RESULTS ON COMPARISON AND SUB/SUPER-STABILIZABILITY OF SOME NEW MEANS

LENKA MIHOKOVIĆ AND MUSTAPHA RAÏSSOULI

Abstract. We present analysis of some new means recently introduced by M. Raïssouli and A. Rezgui. We establish comparison relations and results on (K, N) -sub/super-stabilizability where K and N belong to the class of power means, denoted by B_n , and M is one of the classical or recently studied new means. Assuming that means K , M and N have asymptotic expansions, we present the complete asymptotic expansion of the resultant mean-map. As an application of the obtained asymptotic expansions and the asymptotic inequality between M and $\mathcal{R}(B_p, M, B_q)$, we show how to find the optimal parameters p and q for which M is (B_p, B_q) -sub/super-stabilizable.

1. INTRODUCTION

Through this paper we consider *bivariate mean*, i.e. a function $M: \mathbb{R}^+ \times \mathbb{R}^+ \to$ \mathbb{R}^+ such that $\min(s,t) \leq M(s,t) \leq \max(s,t)$. We say that mean M is symmetric if $M(s,t) = M(t,s)$ for all $s,t > 0$, and homogeneous (of degree 1) if $M(\lambda s, \lambda t) =$ $\lambda M(s,t)$ for all $\lambda, s, t, > 0$. For three homogeneous symmetric bivariate means K, M and N, we define the so-called resultant mean-map

$$
\mathcal{R}(K, M, N)(s, t) = K\Big(M\big(s, N(s, t)\big), M\big(N(s, t), t\big)\Big).
$$

A symmetric mean M is said to be *stable (balanced)*, if $\mathcal{R}(M, M, M) = M$.

Definition 1.1 ([\[9,](#page-18-0) [11\]](#page-18-1)). Let K, N be two nontrivial stable means. Mean M is called

(1) (K, N) -stabilizable, if the following relation is satisfied:

$$
M(s,t) = \mathcal{R}(K,M,N)(s,t) = K\Big(M\big(s,N(s,t)\big),M\big(N(s,t),t\big)\Big).
$$

- (2) (K, N) -sub-stabilizable, if $\mathcal{R}(K, M, N) \leq M$ and M is between K and N,
- (3) (K, N) -super-stabilizable, if $M \leq R(K, M, N)$ and M is between K and N.

Some interesting results regarding these notions can also be found in papers [\[1,](#page-17-0) [12\]](#page-18-2). Motivated by the results on sub/super-stabilizability for bivariate means studied in the paper [\[11\]](#page-18-1) in combination with the general algorithms for the coefficients in the asymptotic expansion of the resultant mean-map and consequently of the stabilizable mean, obtained in the paper [\[8\]](#page-18-3), we present the results on sub/superstabilizability in the context of the asymptotic expansions.

²⁰²⁰ Mathematics Subject Classification. 26E60; 41A60; 39B62.

Key words and phrases. Resultant mean-map; Stable mean; Sub-stabilizable mean; Superstabilizable mean; Asymptotic expansion; Inequalities; Power means.

Definition 1.2. For an asymptotic sequence of functions $(\varphi_n)_{n \in \mathbb{N}_0}$ the (formal) series $\sum_{n=0}^{\infty} a_n \varphi_n(x)$ is said to be an *asymptotic expansion* of a function $f(x)$ as $x \to x_0$ if for each $N \in \mathbb{N}_0$

$$
f(x) = \sum_{n=0}^{N} a_n \varphi_n(x) + o(\varphi_N(x)).
$$

Theoretical background from theory of asymptotic expansions can be found in [\[7\]](#page-17-1). Through this paper, we use asymptotic expansion with respect to the asymptotic sequence $\varphi_n(x) = x^{1-n}, n \in \mathbb{N}_0$, as $x \to \infty$. Based on the sign of the first term in such asymptotic expansion we introduce the notion of the asymptotic inequality.

Definition 1.3 ([\[13\]](#page-18-4)). Let $F(s, t)$ be any homogeneous bivariate function such that

$$
F(x + s, x + t) = c_k(t, s)x^{-k+1} + \mathcal{O}(x^{-k}).
$$

If $c_k(s, t) > 0$ for all s and t, then we say F is asymptotically greater than zero, and write

$$
F\succ 0.
$$

Asymptotic inequality is the necessary condition for the proper inequality, i.e. if $F \geq 0$, then $F \succ 0$.

The subject of study in this paper are means from [\[10\]](#page-18-5) alone for themselves and also in combinations with power means. Recall the definition of the r-th power mean

$$
B_r(a,b) = \begin{cases} \left(\frac{a^r + b^r}{2}\right)^{1/r}, & r \neq 0, \\ \sqrt{ab}, & r = 0. \end{cases}
$$

This class of means covers some well known classical means such as arithmetic mean $A = B_1$, geometric mean $G = B_0$ and harmonic mean $H = B_{-1}$. It has been proved ([\[9\]](#page-18-0)) that power means B_p are stable for all real values of parameter p. Regarding its asymptotic expansion which has been studied in [\[6\]](#page-17-2), one may find that

$$
B_p(x-t, x+t) \sim x + \frac{1}{2}(p-1)t^2 x^{-1} + \frac{1}{24}(p-1)(3+p-2p^2)t^4 x^{-3} + \mathcal{O}(x^{-5}). \tag{1.1}
$$

Remark 1.4. This so-called one variable asymptotic expansion, i.e. asymptotic expansion of a mean in variables $(x - t, x + t)$, is sufficient to determine completely the two variable asymptotic expansion, i.e. asymptotic expansion in variables $(x+s, x+t)$ as it was proved in [\[2,](#page-17-3) Lemma 2.1.]. Following the same procedure as in the mentioned Lemma, we may conclude that the first non-zero coefficient in both of those expansions is the same.

For the convenience of the reader, let us list all the means from [\[10\]](#page-18-5) which will be involved in analysis in this paper. Means from the mentioned paper can be written in a form

$$
m_f(a, b) = \frac{2(a - b)}{f(\frac{a}{b}) - f(\frac{b}{a})}
$$
, $a \neq b$, $m_f(a, a) = a$.

We are interested in some of the special cases. Let

$$
g(x) = \begin{cases} \mu(\ln x), & x \ge 1, \\ -\mu(\ln \frac{1}{x}), & 0 < x < 1, \end{cases} \quad \mu(x) = \int_0^x u(t) dt \text{ odd function,}
$$

and

$$
m_g(a,b) = \frac{|a-b|}{\mu(||\ln \frac{a}{b}|)} = M_u(a,b). \tag{1.2}
$$

With $u(x) = \cosh(\alpha x)$, and for $|\alpha| \leq 1$, we have

$$
L_{\alpha}(a,b) := M_u(a,b) = \frac{2\alpha a^{\alpha}b^{\alpha}(b-a)}{b^{2\alpha} - a^{2\alpha}}.
$$
\n(1.3)

When $u(x) = \frac{1}{\cosh(\alpha x)}$, and for $|\alpha| \leq 1$, we have

$$
S_{\alpha}(a,b) := M_u(a,b) = \frac{2\alpha(b-a)}{4\arctan(\frac{b}{a})^{\alpha} - \pi} = \frac{\alpha(b-a)}{2\arctan\frac{b^{\alpha}-a^{\alpha}}{b^{\alpha}+a^{\alpha}}}.
$$
 (1.4)

Remark that $L_{-\alpha} = L_{\alpha}$ and $S_{-\alpha} = S_{\alpha}$. We can then assume that $0 \le \alpha \le 1$. The means L_{α} and S_{α} cover some well-known standard means and their properties are embodied in the following result. The first part can be seen by a simple verification and for the second part see [\[10\]](#page-18-5).

Proposition 1.5. The following statements are valid:

- (i) $L_0 = S_0 = L$, $L_{1/2} = G$, $L_1 = H$, $S_{1/2} = P$, $S_1 = T$ and $L_{1/4} = HZ_{1/4}$, where L, G, H denote logarithmic, geometric and harmonic mean, P and T refer to the first and the second Seiffert mean respectively, and $HZ_{1/4}$ is the Heinz mean.
- (ii) For fixed $a \neq b$, the map $\alpha \mapsto L_{\alpha}(a, b)$ is strictly decreasing in $\alpha \in [0, 1]$ while $\alpha \mapsto S_{\alpha}(a, b)$ is strictly increasing in $\alpha \in [0, 1]$.

We also recall the definitions of the following means which were introduced in [\[10,](#page-18-5) Examples 4.4–4.7], where $a, b > 0, a \neq b, r > 0, |\alpha| \leq 1$:

$$
M_1(a,b) := \frac{|b-a|}{\ln(1+|\ln b - \ln a|)},
$$

\n
$$
M_2(a,b) := \frac{b-a}{\sqrt{2}\arctan\frac{\ln b - \ln a}{\sqrt{2}}},
$$

\n
$$
M_3(a,b) := \frac{|b-a|}{2\arctan(\left|\ln b - \ln a\right| + 1) - \frac{\pi}{2}},
$$

\n
$$
M_4(a,b) := \frac{b-a}{\sqrt{2}\arcsinh\frac{\ln b - \ln a}{\sqrt{2}}},
$$

\n
$$
M_5(a,b) := \frac{|b-a|}{\sqrt{2}\left(\arcsinh(1 + |\ln b - \ln a|\right) - \arcsinh 1)},
$$

\n
$$
M_{\alpha,r}(a,b) := \frac{(r+\alpha)|b-a|}{(1+r|\ln b - \ln a|)^{\frac{r+\alpha}{r}} - 1}.
$$

\n(1.5)

The (double) parameterized mean $M_{\alpha,r}$ includes some of the known classical means as well as the other means that appear to be new. The following result, which is a simple exercise of Real Analysis, clarifies this claim.

Proposition 1.6. Let $r > 0$ and $|\alpha| \leq 1$. Then the following statements hold:

(i) $M_{\alpha,0}(a, b) = \frac{\alpha |b-a|}{\exp(\alpha |\ln b - \ln a|) - 1}$, with $M_{0,0} = L$. (ii) $M_{0,r} = L$ for any $r > 0$, and $M_{\alpha,\infty} = L$ for any $|\alpha| \leq 1$.

(iii)
$$
M_{-r,r}(a,b) = \frac{r|b-a|}{\ln(1+r|\ln b - \ln a|)}
$$
 for $0 < r \le 1$. In particular, $M_{-1,1} = M_1$.

Let M be a (bivariate) mean. For $a, b > 0$ we can set $a = e^{-x}G$ and $b = e^xG$ for some $x \in \mathbb{R}$, where $G := G(a, b) = \sqrt{ab}$. If M is homogeneous, then

$$
M(a,b) = G M(e^{-x}, e^x) =: G f_M(x). \tag{1.6}
$$

If moreover M is symmetric we can assume that $x \geq 0$. Relying on the associated functions f_{m_i} we may express the characterization of the comparability condition $m_1 < m_2$ for any two symmetric homogeneous means m_1 and m_2 . Namely, it is obvious that

$$
m_1(a, b) < m_2(a, b), \ \forall a \neq b \iff f_{m_1}(x) < f_{m_2}(x), \ \forall x > 0. \tag{1.7}
$$

For example, for the standard means, we have

$$
f_G(x) = 1, \quad f_A(x) = \cosh x, \quad f_H(x) = \frac{1}{\cosh x}, \quad f_L(x) = \frac{\sinh x}{x},
$$

$$
f_P(x) = \frac{\sinh x}{2 \arctan \left(\tanh(x/2)\right)}, \quad f_T(x) = \frac{\sinh x}{\arctan(\tanh x)},\tag{1.8}
$$

from which we easily deduce the well-known chain of inequalities $H < G < L <$ $P < A < T$.

For the means L_{α} and S_{α} , defined by [\(1.3\)](#page-2-0) and [\(1.4\)](#page-2-1), we have for any $x \neq 0$

$$
f_{L_{\alpha}}(x) = 4\alpha \frac{\sinh x}{\sinh(2\alpha x)}, \quad f_{S_{\alpha}}(x) = \alpha \frac{\sinh x}{\arctan(\tanh \alpha x)}, \tag{1.9}
$$

which implies that $L_{\alpha} < S_{\alpha}$ for any $0 < \alpha \leq 1$.

For the means [\(1.5\)](#page-2-2), we can easily find that (for $x > 0$)

$$
f_{M_1}(x) = \frac{2 \sinh x}{\ln(1+2x)}, \qquad f_{M_2}(x) = \frac{\sqrt{2} \sinh x}{\arctan(x\sqrt{2})},
$$

\n
$$
f_{M_3}(x) = \frac{\sinh x}{\arctan(1+2x) - \pi/4}, \qquad f_{M_4}(x) = \frac{\sqrt{2} \sinh x}{\arcsinh(x\sqrt{2})},
$$

\n
$$
f_{M_5}(x) = \frac{\sqrt{2} \sinh x}{\arcsinh(1+2x) - \arcsinh 1}, \qquad f_{M_{\alpha,r}}(x) = \frac{2(r+\alpha)\sinh x}{(1+2rx)^{\frac{r+\alpha}{r}} - 1}.
$$
\n(1.10)

The rest of the paper is organized as follows. In Section 2, we establish comparison relations involving means from [\[10\]](#page-18-5) and some other well-known means. We examine the possibility of being (A, G) - or (G, A) -sub/super-stabilizable for means defined in (1.3) , (1.4) and (1.5) . In Section 3, we present complete asymptotic expansions of the above mentioned means. We also extend the result from [\[8\]](#page-18-3) in order to find the complete asymptotic expansion of the resultant mean-map of means whose asymptotic expansion may include all terms x^{1-n} , $n \in \mathbb{N}_0$, which then could be applied on means which are subject of study in this paper. As a consequence of the coefficient comparison, we find parameters for which means L_{α} are stable and disprove the stability for other means. With use of the coefficients in the asymptotic expansion of power means ([\[6\]](#page-17-2)) we present the coefficients in the asymptotic expansion of the resultant mean-map $\mathcal{R}(B_p, M, B_q)(x - t, x + t)$ as $x \to \infty$. In Section 4, we show some of the applications of the obtained results. We analyze the behaviour of the difference $M - \mathcal{R}(B_p, M, B_q)$, for each of the means defined in [\(1.3\)](#page-2-0), [\(1.4\)](#page-2-1) and [\(1.5\)](#page-2-2). We examine when each of these means can be (B_p, B_q) $sub/super-stabilizable$ and, when possible, how to find such optimal parameters p and q .

2. Comparison of means and sub/super-stabilizability

We start this section by stating some results about comparison between the bivariate means mentioned in the previous section.

Proposition 2.1. Let $\alpha \in [0,1]$. The following statements hold:

(i) If $0 < \alpha < 1/2$ then $G < L_{\alpha} < L < S_{\alpha} < A$. (ii) If $1/2 < \alpha < 1$ then $H < L_{\alpha} < G < L < S_{\alpha} < T$. (iii) If $1/2 \le \alpha \le \sqrt{2}/2$ then $S_{\alpha} < A$.

Proof. For proving [\(i\)](#page-4-0) and [\(ii\)](#page-2-3) we use the statement (ii) of Proposition [1.5](#page-2-4) with the help of [\(i\)](#page-2-5). The details are straightforward and therefore are omitted here.

To show [\(iii\)](#page-4-2) we use [\(1.7\)](#page-3-0) with f_A defined in [\(1.8\)](#page-3-1) and f_{S_α} defined in the second formula in [\(1.9\)](#page-3-2). Thus, we have to establish that

$$
\alpha \frac{\sinh x}{\arctan(\tanh \alpha x)} < \cosh x, \quad \forall x > 0,
$$

or equivalently,

$$
g(x) := \alpha \tanh x - \arctan(\tanh \alpha x) < 0, \quad \forall x > 0.
$$

Simple computation leads to

$$
g'(x) = \frac{\alpha}{(\cosh x)^2} - \frac{1}{1 + (\tanh \alpha x)^2} \frac{\alpha}{(\cosh \alpha x)^2}
$$

$$
= \frac{\alpha}{(\cosh x)^2} - \frac{\alpha}{(\cosh \alpha x)^2 + (\sinh \alpha x)^2}.
$$

We need to study the sign of

$$
h(x) := (\cosh \alpha x)^2 + (\sinh \alpha x)^2 - (\cosh x)^2,
$$

for which we have

$$
h'(x) = 2\alpha(\sinh 2\alpha x) - \sinh 2x \text{ and } h''(x) = 2(2\alpha^2 \cosh 2\alpha x - \cosh 2x).
$$

If $\alpha \leq \sqrt{2}/2$, it is easy to see that $h''(x) < 0$ for all $x > 0$. Thus, h' is strictly decreasing for $x > 0$ and so, $h'(x) < h'(0) = 0$ for all $x > 0$. By the same arguments, we deduce that $h(x) < h(0) = 0$ and therefore $g(x) < g(0) = 0$, for all $x > 0$. The proof is finished. \square

Remark 2.2. Numerical computations show that if $\sqrt{2}/2 < \alpha < 1$ then S_{α} and A are not comparable.

The following Proposition is an extension of the result from [\[10\]](#page-18-5).

Proposition 2.3. We have the following assertions:

- (i) $M_4 < M_5 < M_1 < M_3$ and $L < M_4 < A$. The mean A is not comparable to either one of M_1, M_2, M_3 and M_5 .
- (ii) $L < M_2 < M_3$. The mean M_2 is not comparable to either one of M_1, M_4 and M_5 .

Proof. To prove $M_4 < M_5 < M_1 < M_3$ in [\(i\)](#page-4-3), relying on [\(1.7\)](#page-3-0) and [\(1.10\)](#page-3-3), it is equivalent to show that for all $x > 0$ the following chain of inequalities holds

$$
\frac{\sqrt{2}\sinh x}{\operatorname{arcsinh}(x\sqrt{2})} < \frac{\sqrt{2}\sinh x}{\operatorname{arcsinh}(1+2x) - \operatorname{arcsinh}1} < \frac{2\sinh x}{\ln(1+2x)} < \frac{\sinh x}{\operatorname{arctan}(1+2x) - \pi/4}.
$$
 (2.1)

For the first inequality of [\(2.1\)](#page-5-0) we consider

$$
g(x) := \operatorname{arcsinh}(1 + 2x) - \operatorname{arcsinh}(x\sqrt{2}) - \operatorname{arcsinh} 1,
$$

and then

$$
g'(x) = \frac{2}{\sqrt{1 + (1 + 2x)^2}} - \frac{\sqrt{2}}{\sqrt{1 + 2x^2}},
$$

for which it is easy to see that $g'(x) < 0$ and so $g(x) < g(0) = 0$, for all $x > 0$.

The proof of the two other inequalities in [\(2.1\)](#page-5-0) as well as the proof of the inequalities $L < M_4 < A$ and also $L < M_2 < M_3$ from part [\(ii\)](#page-4-4) is similar. The details are therefore omitted here.

To show that, for example, A is not comparable with M_1 we proceed as follows: we compare $f_A(x)$ and $f_{M_1}(x)$ for $x > 0$, or equivalently, we study the sign of $g(x) := \ln(1 + 2x) - 2 \tanh x$. It is easy to check that $\lim_{x \uparrow \infty} g(x) = +\infty$ and $g(2) = \ln 5 - 2 \tanh 2 < 0$. We then have the conclusion. $g(2) = \ln 5 - 2 \tanh 2 < 0$. We then have the conclusion.

The following result concerns comparison of $M_{\alpha,r}$ with A and G.

Proposition 2.4. Let $r > 0$. If $\alpha \leq 0$ then $M_{\alpha,r} > G$ and, if $\alpha \geq 0$ then $M_{\alpha,r} < A$. *Proof.* Assume that $\alpha \leq 0$. We want to show that $f_{M_{\alpha}}(x) > f_G(x)$ for any $x > 0$. We consider

$$
g(x) := \frac{h(x)}{\left(1+2rx\right)^{\frac{\alpha+r}{r}}-1},
$$

with

$$
h(x) = 2(\alpha + r) \sinh x - (1 + 2rx) \frac{\alpha + r}{r} + 1.
$$

It is clear that $h'(x) = 2(r + \alpha) \left(\cosh x - (1 + 2rx)^{\frac{\alpha}{r}}\right)$. We have the following situations:

- If $\alpha + r > 0$ then $\left(1 + 2rx\right)^{\frac{\alpha + r}{r}} 1 > 0$ and if moreover $\alpha \leq 0$ then $cosh x > 1 \ge (1 + 2rx)^{\alpha/r}$ for any $x > 0$. In this case, $h'(x) > 0$ and so $h(x) > h(0) = 0$, for all $x > 0$. We then deduce that $g(x) > 0$ for all $x > 0$.
- If $\alpha + r < 0$ then $\left(1 + 2rx\right)^{\frac{\alpha + r}{r}} 1 < 0$ and if moreover $\alpha \le 0$ then $h'(x) < 0$ and so $h(x) < h(0) = 0$, for all $x > 0$. We then infer that $g(x) > 0$ for any $x > 0$.

Summarizing, we have shown the desired result.

The inequality $M_{\alpha,r} < A$, for $\alpha \geq 0$, can be established in a similar way, and we ve it to the reader leave it to the reader.

Remark 2.5. Numerical computations show that, if $\alpha > 0$ (resp. $\alpha < 0$) then $M_{\alpha,r}$ is not comparable with G (resp. A).

We will now study the sub/super-stabilizability of some of the above means. We recall the following result.

Proposition 2.6 ([\[11\]](#page-18-1)). Let M be a continuous symmetric mean. Then

- (i) If M is (A, G) -sub-stabilizable then $L \leq M \leq A$.
- (ii) If M is (A, G) -super-stabilizable then $G \leq M \leq L$.

Combining Proposition [2.6](#page-6-0) with Proposition [2.1](#page-4-5) and Proposition [2.3](#page-4-6) we immediately deduce the following corollary.

Corollary 2.7. Let $\alpha \in [0,1]$. The following statements hold:

- (i) L_{α} is not (A, G) -sub-stabilizable and S_{α} is not (A, G) -super-stabilizable.
- (ii) If $1/2 < \alpha \leq 1$ then L_{α} is not (A, G) -super-stabilizable.
- (iii) The three means M_1, M_3 and M_5 are neither (A, G) -sub/super-stabilizable nor (G, A) -sub/super-stabilizable.

To giving more results about sub/super-stabilizability we need the following lemma.

Lemma 2.8 ([\[9\]](#page-18-0)). Let m be a symmetric homogeneous mean. Then, for any $a, b >$ 0, we have

$$
\mathcal{R}(A,m,G)(a,b) = A(\sqrt{a},\sqrt{b}) m(\sqrt{a},\sqrt{b}).
$$

Now, we may state the following result.

Theorem 2.9. Let $0 < \alpha \leq \sqrt{2}/2$. Then, S_{α} is strictly (A, G) -sub-stabilizable. In particular, the first Seiffert mean $P = S_{1/2}$ is strictly (A, G) -sub-stabilizable.

Proof. Firstly, by following Proposition [2.1](#page-4-5) we have $G \leq S_\alpha \leq A$ for any $0 \leq$ $\alpha \leq \sqrt{2}/2$. According to Definition [1.1](#page-0-0) we have to prove that the inequality $\mathcal{R}(A, S_{\alpha}, G)(a, b) < S_{\alpha}(a, b)$ holds for any $a > b$. By Lemma [2.8,](#page-6-1) with substitution [\(1.6\)](#page-3-4), it is equivalent to show that the inequality

$$
f_A(\frac{x}{2})f_{S_\alpha}(\frac{x}{2}) < f_{S_\alpha}(x)
$$

holds for any $x > 0$, and in combination with the corresponding relations in [\(1.8\)](#page-3-1) and in [\(1.9\)](#page-3-2), this is equivalent to

$$
\alpha \cosh(x/2) \frac{\sinh(x/2)}{\arctan(\tanh(\alpha x/2))} < \alpha \frac{\sinh x}{\arctan(\tanh(\alpha x))},
$$

for $x > 0$. Using formula $\sinh x = 2 \sinh(x/2) \cosh(x/2)$ and setting $q(x) :=$ $\arctan\left(\tanh(\alpha x)\right) - 2\arctan\left(\tanh(\alpha x/2)\right)$ we easily verify that $g'(x) < 0$ and so $g(x) < g(0) = 0$, for all $x > 0$. We then deduce the desired result.

Theorem 2.10. Let $0 < \alpha < 1/2$. Then L_{α} is strictly (A, G) -super-stabilizable. In particular, the Heinz mean $HZ_{1/4} = L_{1/4}$ is strictly (A, G) -super-stabilizable.

Proof. By similar way and similar arguments as in the proof of the previous theorem, we are here reduced to study the sign of $g(x) := \sinh(2\alpha x) - 2\sinh(\alpha x)$ for $x > 0$. Obviously, $g'(x) := 2\alpha \cosh(2\alpha x) - 2\alpha \cosh(\alpha x) > 0$ and so $g(x) > g(0) = 0$, for any $x > 0$ so concluding the proof.

Theorem 2.11. The means M_2 and M_4 are both strictly (A, G) -sub-stabilizable.

Proof. It is also similar to the previous proofs. The details are straightforward and therefore omitted here. $\hfill \square$

3. Asymptotic analysis of new means

3.1. Asymptotic expansions of means from [\[10\]](#page-18-5). Let us find asymptotic expansions of means from the Section 1.

The most used result is the expansion for the power of an asymptotic series, which we recall here.

Lemma 3.1 ([\[3\]](#page-17-4)). Let

$$
g(x) \sim \sum_{n=0}^{\infty} a_n x^{-n}
$$

be a given asymptotic expansion (for $x \to \infty$) of $g(x)$ with $a_0 \neq 0$. Then for all real r it holds

$$
[g(x)]^r \sim \sum_{n=0}^{\infty} P[n, r, \mathbf{a}] x^{-n},
$$

where $P[0, r, \mathbf{a}] = a_0^r$ and

$$
P[n,r,\mathbf{a}] = \frac{1}{na_0} \sum_{k=1}^{n} [k(1+r) - n] a_k P[n-k,r,\mathbf{a}], \quad n \in \mathbb{N}.
$$

We assume all sequences are enumerated from 0. Here $P[n, r, a]$ denotes the coefficient by the x^{-n} in the r-th power of series assigned to a sequence $\mathbf{a} = (a_i)_{i \in \mathbb{N}_0}$.

Proposition 3.2. Complete asymptotic expansion of mean L_{α} , $|\alpha| \leq 1$, is given by:

$$
L_{\alpha}(x-t, x+t) \sim x \sum_{n=0}^{\infty} 2\alpha \sum_{k=0}^{n} { \alpha \choose n-k} (-1)^{n-k} P[k, -1, (\binom{2\alpha}{2i+1})_{i \in \mathbb{N}_0}] t^{2n} x^{-2n}.
$$

Proof.

$$
L_{\alpha}(x-t,x+t) = \frac{4\alpha t(x-t)^{\alpha}(x+t)^{\alpha}}{(x+t)^{2\alpha} - (x-t)^{2\alpha}}
$$

\n
$$
= 4\alpha t \left(1 - \frac{t}{x}\right)^{\alpha} \left(1 + \frac{t}{x}\right)^{\alpha} \left[\left(1 + \frac{t}{x}\right)^{2\alpha} - \left(1 - \frac{t}{x}\right)^{2\alpha}\right]^{-1}
$$

\n
$$
\sim 4\alpha t \left(1 - \frac{t^2}{x^2}\right)^{\alpha} \left[\sum_{k=0}^{\infty} {2\alpha \choose k} t^k x^{-k} - \sum_{k=0}^{\infty} {2\alpha \choose k} (-1)^k t^k x^{-k}\right]^{-1}
$$

\n
$$
\sim 2\alpha t \left(1 - \frac{t^2}{x^2}\right)^{\alpha} \left[\sum_{k=0}^{\infty} {2\alpha \choose 2k+1} t^{2k+1} x^{-(2k+1)}\right]^{-1}
$$

\n
$$
\sim 2\alpha t \sum_{j=0}^{\infty} {(\alpha \choose j)} (-1)^j t^{2j} x^{-2j} \cdot \frac{x}{t} \sum_{k=0}^{\infty} P[k, -1, ((\frac{2\alpha}{2i+1}))_{i \in \mathbb{N}_0}] t^{2k} x^{-2k}
$$

\n
$$
\sim 2\alpha x \sum_{n=0}^{\infty} \sum_{k=0}^{n} {(\alpha \choose n-k} (-1)^{n-k} P[k, -1, ((\frac{2\alpha}{2i+1}))_{i \in \mathbb{N}_0}] t^{2n} x^{-2n}.
$$

The beginning of the asymptotic expansion:

$$
L_{\alpha}(x - t, x + t) \sim x - \frac{1}{3}(2\alpha^{2} + 1)t^{2}x^{-1} + \frac{2}{45}(\alpha - 1)(\alpha + 1)(7\alpha^{2} + 2)t^{4}x^{-3}
$$

$$
- \frac{2}{945}(\alpha - 1)(\alpha + 1)(62\alpha^{4} - 85\alpha^{2} - 22)t^{6}x^{-5}
$$

$$
+ \frac{2}{14175}(\alpha - 1)(\alpha + 1)(381\alpha^{6} - 1169\alpha^{4} + 889\alpha^{2} + 214)t^{8}x^{-7} + ...
$$
\n(3.1)

For $\alpha = 1$ and $\alpha = \frac{1}{2}$ these coefficients coincide with coefficients obtained in paper [\[5\]](#page-17-5) for harmonic and geometric mean respectively.

Proposition 3.3. Complete asymptotic expansion of mean S_{α} , $|\alpha| \leq 1$, is given $\mathit{by:}$

$$
S_{\alpha}(x-t, x+t) \sim x \sum_{n=0}^{\infty} \alpha P[n, -1, (D_i)_{i \in \mathbb{N}_0}] t^{2n} x^{-2n},
$$

where

$$
D_n = \sum_{m=0}^{n} \frac{(-1)^m}{2m+1} P[n-m, 2m+1, (C_i)_{i \in \mathbb{N}_0}],
$$

and

$$
C_n = \sum_{k=0}^n \binom{\alpha}{2k+1} P[n-k, -1, \left(\begin{smallmatrix} \alpha \\ 2i \end{smallmatrix}\right))_{i \in \mathbb{N}_0}].
$$

Proof.

$$
S_{\alpha}(x-t, x+t) = \alpha t \left[\arctan \frac{(x+t)^{\alpha} - (x-t)^{\alpha}}{(x+t)^{\alpha} + (x-t)^{\alpha}} \right]^{-1}
$$

\n
$$
= \alpha t \left[\arctan \frac{\left(1 + \frac{t}{x}\right)^{\alpha} - \left(1 - \frac{t}{x}\right)^{\alpha}}{\left(1 + \frac{t}{x}\right)^{\alpha} + \left(1 - \frac{t}{x}\right)} \right]^{-1}
$$

\n
$$
\sim \alpha t \left[\arctan \frac{\sum_{k=0}^{\infty} \binom{\alpha}{k} t^{k} x^{-k} - \sum_{k=0}^{\infty} \binom{\alpha}{k} (-1)^{k} t^{k} x^{-k}}{\sum_{k=0}^{\infty} \binom{\alpha}{k} (-1)^{k} t^{k} x^{-k}} \right]^{-1}
$$

\n
$$
\sim \alpha t \left[\arctan \frac{\sum_{k=0}^{\infty} \binom{\alpha}{k} t^{k} x^{-k} + \sum_{k=0}^{\infty} \binom{\alpha}{k} (-1)^{k} t^{k} x^{-k}}{\sum_{k=0}^{\infty} \binom{\alpha}{2k} t^{2k} x^{-2k}} \right]^{-1}
$$

\n
$$
\sim \alpha t \left[\arctan \left(\frac{t}{x} \sum_{k=0}^{\infty} \binom{\alpha}{2k+1} t^{2k} x^{-2k} \cdot \sum_{j=0}^{\infty} P[j, -1, (\binom{\alpha}{2i})_{i \in \mathbb{N}_0}] t^{2j} x^{-2j} \right) \right]^{-1}
$$

\n
$$
\sim \alpha t \left[\arctan \left(\frac{t}{x} \sum_{n=0}^{\infty} \sum_{k=0}^{n} \binom{\alpha}{2k+1} P[n-k, -1, (\binom{\alpha}{2i})_{i \in \mathbb{N}_0}] t^{2n} x^{-2n} \right) \right]^{-1}
$$

\n
$$
\sim \alpha t \left[\sum_{m=0}^{\infty} \frac{(-1)^m}{2m+1} \left(\frac{t}{x} \sum_{n=0}^{\infty} C_n t^{2n} x^{-2n} \right)^{2m+1} \right]^{-1}
$$

\n
$$
\sim \alpha t \left[\sum_{m
$$

$$
\sim \alpha t \left[\frac{t}{x} \sum_{n=0}^{\infty} \sum_{m=0}^{n} \frac{(-1)^m}{2m+1} P[n-m, 2m+1, (C_i)_{i \in \mathbb{N}_0}] t^{2n} x^{-2n} \right]^{-1}
$$

$$
\sim \alpha x \sum_{n=0}^{\infty} P[n, -1, (D_i)_{i \in \mathbb{N}_0}] t^{2n} x^{-2n}.
$$

Although computed using several recursively defined sequences, the coefficients have a nice form. The beginning of the asymptotic expansion is given by

$$
S_{\alpha}(x-t, x+t) \sim x + \frac{1}{3}(2\alpha^2 - 1)t^2 x^{-1} - \frac{2}{45}(5\alpha^4 - 5\alpha^2 + 2)t^4 x^{-3}
$$

+
$$
\frac{2}{945}(86\alpha^6 - 105\alpha^4 + 63\alpha^2 - 22)t^6 x^{-5}
$$

-
$$
\frac{2}{14175}(214 + 5\alpha^2(\alpha - 1)(\alpha + 1)(135 - 159\alpha^2 + 271\alpha^4))t^8 x^{-7} + ...
$$
\n(3.2)

For $\alpha = \frac{1}{2}$ and $\alpha = 1$ these coefficients coincide with the coefficients from [\[13\]](#page-18-4) for the first and the second Seiffert mean.

We may state more general result for means of the type (1.2) .

Theorem 3.4. Assume that the odd function $\mu: \mathbb{R} \to \mathbb{R}$ has the following expansion

$$
\mu(x) \sim \sum_{n=0}^{\infty} c_n x^{2n+1}, \text{ as } x \to 0,
$$
\n(3.3)

with $c_0 = 1$. Then mean m_g defined in [\(1.2\)](#page-2-6) has the following expansion

$$
m_g(x - t, x + t) \sim x \sum_{m=0}^{\infty} P[m, -1, (E_i)_{i \in \mathbb{N}_0}] t^{2m} x^{-2m}, \text{ as } x \to \infty,
$$
 (3.4)

where

$$
E_m = \sum_{n=0}^{m} c_n 2^{2n} P[m - n, 2n + 1, (\frac{1}{2i+1})_{i \in \mathbb{N}_0}].
$$

Proof. Observe the expression $\ln \frac{a}{b}$ when $a = x - t$ and $b = x + t$, which under assumption $t > 0$ and for x large enough, is equal to

$$
\ln \frac{x+t}{x-t} \sim \ln \left(1 + \frac{t}{x} \right) - \ln \left(1 - \frac{t}{x} \right) = \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} t^k x^{-k} + \sum_{k=1}^{\infty} \frac{1}{k} t^k x^{-k}
$$

$$
\sim 2 \sum_{k=0}^{\infty} \frac{1}{2k+1} t^{2k+1} x^{-(2k+1)}.
$$

Now we have, for any t ,

$$
m_g(x - t, x + t) \sim 2t \left[\mu \left(2 \sum_{k=0}^{\infty} \frac{1}{2k+1} t^{2k+1} x^{-(2k+1)} \right) \right]^{-1}
$$

$$
\sim 2t \left[\sum_{n=0}^{\infty} c_n \left(2 \sum_{k=0}^{\infty} \frac{1}{2k+1} t^{2k+1} x^{-(2k+1)} \right)^{2n+1} \right]^{-1}
$$

$$
\sim 2t \left[\sum_{n=0}^{\infty} c_n t^{2n+1} x^{-2n-1} 2^{2n+1} \sum_{k=0}^{\infty} P[k, 2n+1, (\frac{1}{2i+1})_{i \in \mathbb{N}_0}] t^{2k} x^{-2k} \right]^{-1}
$$

$$
\sim 2t \left[2tx^{-1} \sum_{m=0}^{\infty} \sum_{n=0}^{m} c_n 2^{2n} P[m-n, 2n+1, (\frac{1}{2i+1})_{i \in \mathbb{N}_0}] t^{2m} x^{-2m} \right]^{-1}
$$

$$
\sim x \sum_{m=0}^{\infty} P[m, -1, (E_i)_{i \in \mathbb{N}_0}] t^{2m} x^{-2m}.
$$

- Example 3.5. (1) Asymptotic expansion of mean L_{α} can also be deduced form Theorem [3.4.](#page-9-0) Let $\mu(x) = \frac{1}{\alpha} \sinh(\alpha x)$, i.e. let $c_n = \frac{\alpha^{2n}}{(2n+1)!}$ in [\(3.3\)](#page-9-1). Then using formula (3.4) we may also obtain the coefficients from (3.1) .
	- (2) Let $u(y) = \frac{1}{\cosh(\alpha y)}$ whose asymptotic expansion as $y \to 0$ is $u(y) \sim$ $\sum_{n=0}^{\infty} P[n,-1,(\frac{\alpha^{2i}}{(2i)!})_{i\in\mathbb{N}_0}]y^{2n}$. Then, with $c_n = \frac{1}{2n+1}P[n,-1,(\frac{\alpha^{2i}}{(2i)!})_{i\in\mathbb{N}_0}]$ in (3.3) , formula (3.4) gives the coefficients as in (3.2) .

Regarding the rest of the means from Section 1, similar computations lead to their asymptotic expansions. With prior use of the arctangent addition formula in the denominator of M_3 and the inverse hyperbolic sine addition (subtraction) formula for the denominator of M_5 , with application of Lemma [3.1](#page-7-0) and Taylor series expansion of logarithmic, arctangent or inverse hyperbolic sine function in variable t/x , we obtain the following expansions.

Proposition 3.6. As $x \to \infty$, for means defined in [\(1.5\)](#page-2-2), the following expansions hold:

$$
M_{1}(x-t,x+t) \sim x+|t| - \frac{2}{3}t^{2}x^{-1} + \frac{1}{3}|t|^{3}x^{-2} - \frac{28}{45}t^{4}x^{-3} + \frac{37}{45}|t|^{5}x^{-4}
$$

\n
$$
- \frac{1369}{945}t^{6}x^{-5} + \dots
$$

\n
$$
M_{2}(x-t,x+t) \sim x + \frac{1}{3}t^{2}x^{-1} - \frac{2}{9}t^{4}x^{-3} + \frac{14}{135}t^{6}x^{-5} - \frac{122}{945}t^{8}x^{-7} + \dots
$$

\n
$$
M_{3}(x-t,x+t) \sim x + |t| - \frac{1}{3}|t|^{3}x^{-2} + \frac{4}{15}t^{4}x^{-3} - \frac{13}{45}|t|^{5}x^{-4} + \frac{1}{9}t^{6}x^{-5} + \dots
$$

\n
$$
M_{4}(x-t,x+t) \sim x - \frac{1}{6}t^{4}x^{-3} + \frac{8}{315}t^{6}x^{-5} - \frac{367}{4536}t^{8}x^{-7} + \dots
$$

\n
$$
M_{5}(x-t,x+t) \sim x + \frac{1}{2}|t| - \frac{1}{4}t^{2}x^{-1} - \frac{1}{6}|t|^{3}x^{-2} + \frac{5}{48}t^{4}x^{-3} - \frac{7}{3}360|t|^{5}x^{-4}
$$

\n
$$
+ \frac{1033}{10080}t^{6}x^{-5} + \dots
$$

\n
$$
M_{\alpha,r}(x-t,x+t) \sim x + \alpha|t| + \frac{1}{3}(\alpha^{2} + 2r\alpha - 1)t^{2}x^{-1} - \frac{1}{3}r\alpha(2r+\alpha)|t|^{3}x^{-2}
$$

\n
$$
- \frac{1}{45}(\alpha(\alpha + 2r)(\alpha^{2} - 18r^{2} + 2\alpha r - 5) + 4) t^{4}x^{-3}
$$

\n
$$
- \frac{1}{45}\alpha r(\alpha + 2r)(3(2r - \alpha)(\alpha + 4r) + 10)|t|^{5}x^{-4} + \dots
$$

3.2. Asymptotic expansion of the resultant mean-map. Assuming that all means involved were bivariate, symmetric, homogeneous and had the asymptotic expansions as $x \to \infty$ of the following type

$$
K(x - t, x + t) \sim \sum_{n=0}^{\infty} a_n^K t^{2n} x^{-2n+1},
$$

$$
M(x - t, x + t) \sim \sum_{n=0}^{\infty} a_n^M t^{2n} x^{-2n+1},
$$
 (3.5)

$$
N(x - t, x + t) \sim \sum_{n=0}^{\infty} a_n^N t^{2n} x^{-2n+1},
$$

in [\[8\]](#page-18-3) we found the complete asymptotic expansion of the resultant mean–map:

$$
R(x - t, x + t) = \mathcal{R}(K, M, N)(x - t, x + t) \sim \sum_{n=0}^{\infty} a_n^R t^{2n} x^{-2n+1}.
$$
 (3.6)

Using coefficients of the resultant mean–map of the corresponding means, we found formula for calculating the coefficients in the asymptotic power series expansion of type (3.5) of stable mean, which, up to five terms reads as $([8, formula (34)]):$ $([8, formula (34)]):$ $([8, formula (34)]):$

$$
M(x - t, x + t) = x + a_1 t^2 x^{-1} + \frac{1}{6} a_1 (1 + a_1) (1 - 4a_1) t^4 x^{-3}
$$

+ $\frac{1}{90} a_1 (1 + a_1) (6 - 31a_1 + 36a_1^2 + 64a_1^3) t^6 x^{-5}$
+ $\frac{1}{2520} a_1 (1 + a_1) (90 - 531a_1 + 937a_1^2 + 568a_1^3 - 3088a_1^4 - 2176a_1^5) t^8 x^{-7}$ (3.7)
+ $\mathcal{O}(x^{-9}).$

With $K = B_p$ and $N = B_q$, we obtain coefficients in the asymptotic expansion (3.6) of the resultant mean–map (consequence of $[8, \text{formula } (26)]$):

$$
\mathcal{R}(B_p, M, B_q)(x - t, x + t) = x + \frac{1}{8}(2a_1^M + p + 2q - 3)t^2 x^{-1}
$$

+
$$
\frac{1}{384} \left(24a_2^M + 12a_1^M(-4pq + p + 2q(q + 1) + 1) -2p^3 + 3p^2 + 2p(7 - 6q) + 4q(-4q^2 + 6q + 7) - 39\right)t^4 x^{-3} + \mathcal{O}(x^{-5}).
$$
\n(3.8)

As a consequence of the results of the above mentioned paper, specially relying on the form of the stable mean coefficients given in [\(3.7\)](#page-11-1), in combination with the asymptotic expansion given in Proposition [3.2](#page-7-1) we obtain the following.

Corollary 3.7. Means L_{α} are stable iff $\alpha \in {\{\pm \frac{1}{2}, \pm 1\}}$.

Proof. By comparing coefficients [\(3.1\)](#page-8-0) in the asymptotic expansion of mean L_{α} with the stable mean coefficients [\(3.7\)](#page-11-1) by the powers x^{-1} and x^{-3} we obtain the following equations

$$
-\frac{1}{3}(2\alpha^2 + 1) = a_1,
$$

\n
$$
\frac{2}{45}(\alpha - 1)(\alpha + 1)(7\alpha^2 + 2) = \frac{1}{6}a_1(1 + a_1)(1 - 4a_1)
$$

\n
$$
= \frac{1}{6}(-\frac{1}{3}(2\alpha^2 + 1))(1 - \frac{1}{3}(2\alpha^2 + 1))(1 + \frac{4}{3}(2\alpha^2 + 1)),
$$

which combined give the following

$$
18(\alpha^2 - 1)(7\alpha^2 + 2) = 5(2\alpha^2 + 1)(\alpha^2 - 1)(8\alpha^2 + 7).
$$

The only real solutions are $\alpha \in \{\pm \frac{1}{2}, \pm 1\}$. For either of those values of α the stable mean is obtained. Namely, $L_{\pm 1} = H$ and $L_{\pm \frac{1}{2}} = G$.

The similar procedure as in Corollary [3.7,](#page-11-2) involving coefficients given in [\(3.2\)](#page-9-3), gives no solutions for $|\alpha| \leq 1$, so we have the following conclusion.

Corollary 3.8. Means S_{α} , $|\alpha| \leq 1$, are not stable.

Not all of the means [\(1.3\)](#page-2-0), [\(1.4\)](#page-2-1), [\(1.5\)](#page-2-2) have asymptotic expansion of the form [\(3.5\)](#page-10-0). In order to obtain the asymptotic expansion of the resultant mean–map for all of the means mentioned in the introduction, and afterwards find which of them

are stable and study asymptotic expansion of $\mathcal{R}(B_p, M, B_q)$, we need to adjust the algorithm obtained in the paper [\[8\]](#page-18-3).

Theorem 3.9. Assume that means K , M and N have expansions

$$
K(x - t, x + t) \sim \sum_{n=0}^{\infty} a_n^K t^n x^{-n+1},
$$
\n(3.9)

$$
M(x-t, x+t) \sim \sum_{n=0}^{\infty} a_n^M t^n x^{-n+1},
$$
\n(3.10)

$$
N(x - t, x + t) \sim \sum_{n=0}^{\infty} a_n^N t^n x^{-n+1},
$$
\n(3.11)

as $x \to \infty$, such that $a_1^N \neq \pm 1$. Then the coefficients in the asymptotic expansion of the resultant mean–map

$$
R(x - t, x + t) = \mathcal{R}(K, M, N)(x - t, x + t) \sim \sum_{n=0}^{\infty} a_n^R t^n x^{-n+1},
$$
 (3.12)

can be calulated by the recursive formula

$$
a_m^R = \frac{1}{4} \sum_{n=0}^m \sum_{k=0}^{m-n} a_n^K P[k, n, \mathbf{d}] P[m - n - k, -n + 1, \mathbf{s}], \quad m \in \mathbb{N}_0,
$$
 (3.13)

where

$$
d_{m-1} = \sum_{n=0}^{m} a_n^M \sum_{k=0}^{m-n} (P[k, n, \tilde{\mathbf{g}}] P[m - n - k, -n + 1, \tilde{\mathbf{h}}]
$$

- $P[k, n, \mathbf{g}] P[m - n - k, -n + 1, \mathbf{h}]), \quad m \in \mathbb{N},$

$$
s_m = \sum_{n=0}^{m} a_n^M \sum_{k=0}^{m-n} (P[k, n, \tilde{\mathbf{g}}] P[m - n - k, -n + 1, \tilde{\mathbf{h}}]
$$

+ $P[k, n, \mathbf{g}] P[m - n - k, -n + 1, \mathbf{h}]), \quad m \in \mathbb{N}_0,$

and

$$
\mathbf{g} = (1 + a_1^N, a_2^N, a_3^N, \ldots), \qquad \mathbf{h} = (2, a_1^N - 1, a_2^N, a_3^N, \ldots),
$$

\n
$$
\tilde{\mathbf{g}} = (1 - a_1^N, -a_2^N, -a_3^N, \ldots), \quad \tilde{\mathbf{h}} = (2, 1 + a_1^N, a_2^N, a_3^N, \ldots).
$$
\n(3.14)

Proof. The proof goes by the similar procedure as the proof of the somewhat specific analogue form paper [\[8\]](#page-18-3). With N being the abbreviated version of $N(x - t, x + t)$, the following holds:

$$
M(x-t, N(x-t, x+t)) \sim \sum_{n=0}^{\infty} a_n^M \left(\frac{1}{2}(N-x+t)\right)^n \left(\frac{1}{2}(N+x-t)\right)^{-n+1}
$$

$$
\sim \frac{1}{2} \sum_{n=0}^{\infty} a_n^M \left(t\left(1+a_1^N\right) + \sum_{k=2}^{\infty} a_k^N t^k x^{-k+1}\right)^n \left(2x + \left(a_1^N - 1\right)t + \sum_{j=2}^{\infty} a_j^N t^j x^{-j+1}\right)^{-n+1}
$$

$$
\sim \frac{1}{2} \sum_{n=0}^{\infty} a_n^M t^n \sum_{k=0}^{\infty} P[k, n, \mathbf{g}] t^k x^{-k} \cdot x^{-n+1} \sum_{j=0}^{\infty} P[j, -n+1, \mathbf{h}] t^j x^{-j}
$$

$$
\sim \frac{1}{2} \sum_{m=0}^{\infty} \sum_{n=0}^{m} a_n^M \sum_{k=0}^{m-n} P[k, n, \mathbf{g}] P[m - n - k, -n + 1, \mathbf{h}] t^m x^{-m+1}.
$$

Similarly, the second component to be composed with the mean K has the following expansion:

$$
M(N(x-t, x+t), x+t) \sim \sum_{n=0}^{\infty} a_n^M \left(\frac{1}{2}(x+t-N)\right)^n \left(\frac{1}{2}(x+t+N)\right)^{-n+1}
$$

$$
\sim \frac{1}{2} \sum_{n=0}^{\infty} a_n^M \left(t(1-a_1^N) - \sum_{k=2}^{\infty} a_k^N t^k x^{-k+1}\right)^n \left(2x + \left(a_1^N + 1\right)t + \sum_{j=2}^{\infty} a_j^N t^j x^{-j+1}\right)^{-n+1}
$$

$$
\sim \frac{1}{2} \sum_{n=0}^{\infty} a_n^M t^n \sum_{k=0}^{\infty} P[k, n, \tilde{\mathbf{g}}] t^k x^{-k} \cdot x^{-n+1} \sum_{j=0}^{\infty} P[j, -n+1, \tilde{\mathbf{h}}] t^j x^{-j}
$$

$$
\sim \frac{1}{2} \sum_{m=0}^{\infty} \sum_{n=0}^m a_n^M \sum_{k=0}^{m-n} P[k, n, \tilde{\mathbf{g}}] P[m - n - k, -n+1, \tilde{\mathbf{h}}] t^m x^{-m+1}.
$$

In order to obtain the desired formula, we need to calculate one half of the difference of the previous two expressions

$$
T = \frac{1}{2} \big(M(N(x-t, x+t), x+t) - M(x-t, N(x-t, x+t)) \big) \sim \frac{1}{4} \sum_{m=1}^{\infty} d_{m-1} t^m x^{-m+1},
$$

and one half of the sum of those two expressions

$$
X = \frac{1}{2} \big(M(N(x-t, x+t), x+t) + M(x-t, N(x-t, x+t)) \big) \sim \frac{1}{4} \sum_{m=0}^{\infty} s_m t^m x^{-m+1}.
$$

The resultant mean-map can then be written in a following way

$$
R = K(X - T, X + T)
$$

\n
$$
\sim \sum_{n=0}^{\infty} a_n^K \left(\frac{1}{4} \sum_{k=1}^{\infty} d_{k-1} t^k x^{-k+1}\right)^n \left(\frac{1}{4} \sum_{j=0}^{\infty} s_j t^j x^{-j+1}\right)^{-n+1}
$$

\n
$$
\sim \frac{1}{4} \sum_{n=0}^{\infty} a_n^K t^n \sum_{k=0}^{\infty} P[k, n, \mathbf{d}] t^k x^{-k} \cdot x^{-n+1} \sum_{j=0}^{\infty} P[j, -n+1, \mathbf{s}] t^j x^{-j}
$$

\n
$$
\sim \frac{1}{4} \sum_{m=0}^{\infty} \sum_{n=0}^m \sum_{k=0}^{m-n} a_n^K P[k, n, \mathbf{d}] P[m - n - k, -n+1, \mathbf{s}] t^m x^{-m+1}
$$

\n
$$
\sim \sum_{m=0}^{\infty} a_m^R t^m x^{-m+1}.
$$

 \Box

With assumptions from Theorem [3.9,](#page-12-0) the first few coefficients in the expansion [\(3.12\)](#page-12-1) of the resultant mean are given by:

$$
a_0^R = 1,
$$

\n
$$
a_1^R = \frac{1}{2} \left(a_1^K + a_1^M + a_1^N - a_1^K a_1^M a_1^N \right),
$$

\n
$$
a_2^R = \frac{1}{4} \left(a_2^N (2 - 2a_1^K a_1^M) + a_2^K (-1 + a_1^M a_1^N)^2 + a_2^M + a_2^M a_1^N (a_1^N - 2a_1^K) \right),
$$

\n
$$
a_3^R = \frac{1}{64} \left(-32a_3^N (a_1^K a_1^M - 1) - 8a_3^K (a_1^M a_1^N - 1)^3 - 8a_2^K (-1 + a_1^M a_1^N) ((a_1^N)^2 a_1^M - a_1^M (4a_2^N + 1) + a_1^N ((a_1^M)^2 - 4a_2^M - 1)) + 8a_2^M (a_1^K - a_1^N) ((a_1^N)^2 - 4a_2^N - 1) - 8a_3^M (a_1^K a_1^N (3 + (a_1^N)^2) - 3(a_1^N)^2 - 1)).
$$

\n(3.15)

Remark 3.10. In the previous Theorem formula for the Case I: $a_1^N \neq \pm 1$ was presented. In other cases, first few terms in the sequence g or \tilde{g} is equal to 0 and therefore the expression $P[\cdot,\cdot,\mathbf{g}]$ or $P[\cdot,\cdot,\tilde{\mathbf{g}}]$ is not well defined. In order to complement the result of Theorem [3.9,](#page-12-0) let z be the minimal integer, greater or equal to 2, such that $a_z^N \neq 0$. In Case II: $a_1^N = -1$, let us redefine $\mathbf{g} = (a_z^N, a_{z+1}^N, \ldots)$ and let $\mathbf{h}, \tilde{\mathbf{g}}, \tilde{\mathbf{h}}$ be the same as in [\(3.14\)](#page-12-2). Then

$$
d_{m-1} = \sum_{n=0}^{m} a_n^M \sum_{k=0}^{m-n} P[k, n, \tilde{\mathbf{g}}] P[m - n - k, -n + 1, \tilde{\mathbf{h}}]
$$

$$
- \sum_{n=0}^{\lfloor \frac{m}{z} \rfloor} a_n^M \sum_{k=0}^{m-nz} P[k, n, \mathbf{g}] P[m - nz - k, -n + 1, \mathbf{h}], \quad m \in \mathbb{N},
$$

$$
s_m = \sum_{n=0}^{m} a_n^M \sum_{k=0}^{m-n} P[k, n, \tilde{\mathbf{g}}] P[m - n - k, -n + 1, \tilde{\mathbf{h}}]
$$

$$
+ \sum_{n=0}^{\lfloor \frac{m}{z} \rfloor} a_n^M \sum_{k=0}^{m-nz} P[k, n, \mathbf{g}] P[m - nz - k, -n + 1, \mathbf{h}], \quad m \in \mathbb{N}_0,
$$

and the coefficients of the resultant mean-map can be calculated by the formula [\(3.13\)](#page-12-3). In Case III: $a_1^N = 1$, the sequence \tilde{g} needs to be redefined in a similar manner and the rest of the procedure goes analogously.

Remark 3.11. Once calculated by the procedure given in Theorem [3.9,](#page-12-0) the coefficients behave well for the special values mentioned in Cases II and III from Remark [3.10](#page-14-0) where the existence has been shown. As it was proved in [\[6,](#page-17-2) Lemma 2.2] we may use coefficients from the list (3.15) with $a_1^N = \pm 1$ and $a_2^N = \ldots = a_{z-1}^N = 0$ to obtain the coefficients in Cases II and III as well.

Remark 3.12. Regarding the stability examination, when equating the coefficients a_m^R from [\(3.15\)](#page-14-1) with $a_m^M = a_m^N = a_m^K$, for $m = 1$ we obtain that $a_1^M(1 + a_1^M)(1 - a_1^K)$ a_1^M) = 0. The first possibility $(a_1^M = 0)$ implies (inductively) that also $a_{2m+1} = 0$, for all $m \in \mathbb{N}$, which means that M has the asymptotic expansion of the type [\(3.5\)](#page-10-0). Each of other two possibilities $(a_1^M = \pm 1)$ implies that $a_m = 0$, for all $m \ge 2$, which correspond to the first and the second projection.

As a consequence of the coefficients comparison, using the stable mean expansion [\(3.7\)](#page-11-1), for means whose asymptotic expansion does not contain even powers of x, on the one side and the corresponding expansions given in Proposition [3.6](#page-10-1) on the other side, and employing Remark [3.12](#page-14-2) for those means whose asymptotic expansions contain all powers x^{-n+1} , $n \in \mathbb{N}_0$, we have the following conclusion.

Corollary 3.13. None of the means form the list [\(1.5\)](#page-2-2) is stable.

We finish this Section with a result based on Theorem [3.12](#page-12-1) which will be used in sequel. With $K = B_p$ and $N = B_q$, by incorporating coefficients [\(1.1\)](#page-1-0) into the expansions [\(3.9\)](#page-12-4) and [\(3.11\)](#page-12-5) with parameters p and q respectively, we obtain coefficients in the asymptotic expansion [\(3.12\)](#page-12-1) of the resultant mean–map:

$$
\mathcal{R}(B_p, M, B_q)(x - t, x + t) = x + \frac{1}{2}a_1^M t + \frac{1}{8}(2a_2^M + p + 2q - 3)t^2 x^{-1} + \frac{1}{16}(2a_3^M - (p - 1)(2q - 1)a_1^M)t^3 x^{-2} + \mathcal{O}(x^{-3}).
$$
\n(3.16)

4. Sub-stabilizability and super-stabilizability with power means

In this Section we present some of the possible applications of the previously obtained results. We will show the use of asymptotic expansions when examining the possibility for a given mean to be (B_p, B_q) -sub- or super-stabilizable and when possible, how to find optimal parameters p and q .

Since the asymptotic inequality is the necessary condition for the proper inequality, we analyse when

$$
M - \mathcal{R}(B_p, M, B_q) \succ 0. \tag{4.1}
$$

The best approximation is obtained when as many as possible first coefficients are equal to 0. See [\[13\]](#page-18-4) for detailed analysis. Because of the reasoning in Remark [1.4,](#page-1-1) it is sufficient to use the expansions in variables $(x - t, x + t)$.

Asymptotic inequality corresponds the proper inequality for means when variables s and t are close enough to each other. In order to complement the information obtained form asymptotic side, it is often useful to observe relation between means about the point $(0, 1)$. For more details we refer to paper [\[4\]](#page-17-6).

Example 4.1. Let $M = L_{\alpha}$. Observe the difference between [\(3.1\)](#page-8-0) and [\(3.8\)](#page-11-3). When equating the coefficient by x^{-1} with 0, we have that $q = \frac{1}{2}(1-p) - 2\alpha^2$ and then $(L_{\alpha} - \mathcal{R}(B_p, L_{\alpha}, B_q))(x - t, x + t) \sim -\frac{1}{384}(4\alpha^2 - 1)(p^2 - 1 - 16\alpha^2 + 16\alpha^4)t^4x^{-3} + \dots$ and if also $p = \pm \sqrt{1 + 16\alpha^2 - 16\alpha^4}$ then

$$
(L_{\alpha} - \mathcal{R}(B_{p}, L_{\alpha}, B_{q}))(x - t, x + t) \sim -\frac{1}{720} \alpha^{2} (\alpha^{2} - 1)(4\alpha^{2} - 1)^{3} t^{6} x^{-5} + \dots
$$

Hence, for such p and q, [\(4.1\)](#page-15-0) holds for $\frac{1}{2} < |\alpha| < 1$, and we have the opposite inequality sign in [\(4.1\)](#page-15-0) for $0 < |\alpha| < \frac{1}{2}$.

Additionally, when we equate coefficient by x^{-5} with 0, we obtain the following trivial (and already known) cases: $\alpha = \pm \frac{1}{2}, q = -p; \alpha = 0, p = -1, q = 1; \alpha =$ $0, p = 1, q = 0; \alpha = \pm 1, p = 1, q = -2;$ where the stabilizability is achieved.

On the other side, values of limits $\lim_{s\to 0} L_{\alpha}(s, 1-s) = 0$ and $\lim_{s\to 0} \mathcal{R}(B_p, L_{\alpha}, B_q)(s, 1-s)$ s) > 0 for any p and q, imply that L_{α} could only be (B_p, B_q) -super-stabilizable.

Combining the observations about the sign of the difference L_{α} − $\mathcal{R}(B_p, L_{\alpha}, B_q)$ in two limiting points, with the intent to achieve the best possible order of inequality, we see that for $\frac{1}{2} < |\alpha| < 1$ and above mentioned specific values of p and q mean L_{α} cannot be either sub- or super-stabilizable with the pair of power means (B_p, B_q) .

Motivated by the Example [4.1,](#page-15-1) numerical experiments indicate that the following statement should be true.

Conjecture 4.2. Let $q = \frac{1}{2}(1-p) - 2\alpha^2$ and $p = \pm\sqrt{1+16\alpha^2-16\alpha^4}$. Then L_{α} $\mathcal{R}(B_p, L_\alpha, B_q) < 0$ for $|\alpha| < \frac{1}{2}$.

Example 4.3. Let $M = S_{\alpha}$. Observe the difference between [\(3.2\)](#page-9-3) and [\(3.8\)](#page-11-3). The best order of approximation as $x \to \infty$ is obtained when $q = \frac{1}{2}(1-p) + 2\alpha^2$ and $p^2 = \frac{1-12\alpha^2+112\alpha^4-64\alpha^6}{1+4\alpha^2}$. For such values p and q we have

$$
(S_{\alpha} - \mathcal{R}(B_p, S_{\alpha}, B_q))(x - t, x + t) \sim \frac{1}{720} \alpha^2 (1 + \alpha^2)(1 - 16\alpha^2 + 16\alpha^4)^2 (1 + 4\alpha^2)^{-1} t^6 x^{-5} + \dots
$$

Then, [\(4.1\)](#page-15-0) holds when $|\alpha| \neq \frac{1}{2}\sqrt{2 \pm \sqrt{3}}$ and $\alpha \neq 0$. Examining the sign of the coefficient by x^{-7} for the corresponding values of α we may see that the opposite sign in [\(4.1\)](#page-15-0) holds for $|\alpha| = \frac{1}{2}\sqrt{2 \pm \sqrt{3}}$. When additionally equating the coefficient by x^{-5} with 0 we obtain trivial cases: $\alpha = 0, p = 1, q = 0$ and $\alpha = 0, p = -1, q = 1$, where the stabilizability is achieved.

Conjecture 4.4. Let $q = \frac{1}{2}(1-p) + 2\alpha^2$ and $p^2 = \frac{1-12\alpha^2+112\alpha^4-64\alpha^6}{1+4\alpha^2}$. Then S_α $\mathcal{R}(B_p, S_\alpha, B_q) < 0$ for $|\alpha| = \frac{1}{2}\sqrt{2 \pm \sqrt{3}}$.

Example 4.5. Let $M = M_1$. Mean M has the asymptotic expansion of the form [\(3.10\)](#page-12-6) and hence from [\(3.16\)](#page-15-2) follows that

$$
(M_1 - \mathcal{R}(B_p, M_1, B_q))(x - t, x + t) \sim \frac{1}{2}a_1^M t + \mathcal{O}(x^{-1}).
$$

From Proposition [3.6](#page-10-1) we see that $a_1^M = \frac{|t|}{t}$ so the expansion of the difference starts with coefficient $\frac{1}{2}a_1^M t = \frac{1}{2}|t|$ by x^0 , and the asymptotic inequality [\(4.1\)](#page-15-0) holds, which means that for s and t close enough and $s \neq t$, the difference $M_1(s,t)$ – $\mathcal{R}(B_p, M_1, B_q)(s, t)$ is greater than 0. On the other side, observing the values of limits $\lim_{s\to 0} M_1(s, 1-s) = 0$ and $\lim_{s\to 0} \mathcal{R}(B_p, M_1, B_q)(s, 1-s) = 2^{-1/p} \cdot |2^{-1/q} 1/ \ln(1 + \ln 2^{-1/q}) > 0$, $\forall p, q \in \mathbb{R}$, implies that there are s and t for which the difference between M_1 and $\mathcal{R}(B_p, M_1, B_q)$ is negative. We may conclude that M_1 cannot be either sub- or super-stabilizable with power means.

Example 4.6. Let $M = M_2$. Then from Proposition [3.6](#page-10-1) and [\(3.8\)](#page-11-3) we have

$$
(M_2-\mathcal{R}(B_p, M_2, B_q))(x-t, x+t) \sim \frac{1}{8}(5-p-2q)t^2x^{-1}
$$

+ $\frac{1}{384}(2p^3-3p^2+2p(14q-9)+4q(4(q-2)q-9)-45) t^4x^{-3}+\dots$

For $p = 5 - 2q$ and $q = \frac{1}{2}(5 \pm \sqrt{17})$, coefficients by x^{-1} and x^{-3} become equal to 0, and coefficient by x^{-5} is equal to $-\frac{11}{180}t^6$. Therefore,

$$
M_2-\mathcal{R}(B_p, M_2, B_q) \prec 0.
$$

Conjecture 4.7. Let $p = 5 - 2q$ and $q = \frac{1}{2}(5 \pm \sqrt{17})$. Then $M_2 - \mathcal{R}(B_p, M_2, B_q) < 0$.

Example 4.8. Let $M = M_3$. Then we have the similar situation as in the Example [4.5](#page-16-0) with the same value of the coefficient a_1^M , and therefore asymptotic inequality [\(4.1\)](#page-15-0) holds. On the other side, as $(s, t) \rightarrow (0, 1)$, values of the difference M_3 – $\mathcal{R}(B_p, M_3, B_q)$ may be positive or negative, depending on p and q. Hence, M_3 could only be sub-stabilizable with power means.

Example 4.9. Let $M = M_4$. Then

$$
(M_4 - \mathcal{R}(B_p, M_4, B_q))(x - t, x + t) \sim \frac{1}{8}(3 - p - 2q)t^2 x^{-1}
$$

+ $\frac{1}{384} (p(2 + p)(-7 + 2p) + 12pq - 24q^2 + 16q^3 - 7(3 + 4q)) t^4 x^{-3} + ...$

For $p = 3 - 2q$ and $q = \frac{1}{2}(3 \pm \sqrt{21})$ coefficients by x^{-1} and x^{-3} become equal to 0, the coefficient by x^{-5} is equal to $\frac{13}{120}t^6$ and therefore the asymptotic inequality [\(4.1\)](#page-15-0) holds.

Conjecture 4.10. Let $p = 3 - 2q$ and $q = \frac{1}{2}(3 \pm \sqrt{21})$. Then $M_4 - \mathcal{R}(B_p, M_4, B_q) > 0$.

Example 4.11. Mean $M = M_5$, when compared with the resultant mean-map behaves similarly as mean M_1 . Namely, asymptotic inequality (4.1) holds, and further, $\lim_{s\to 0} M_5(s, 1-s) = 0$ and $\lim_{s\to 0} \mathcal{R}(B_p, M_5, B_q)(s, 1-s) > 0$, $\forall p, q \in \mathbb{R}$. Analogously as in the Example [4.5,](#page-16-0) we may conclude that mean M_5 cannot be either (B_n, B_q) -sub or super-stabilizable.

Example 4.12. Let $M = M_{\alpha,r}$, defined in [\(1.5\)](#page-2-2) with special and limit cases described in Proposition [1.6.](#page-2-7) From [\(3.16\)](#page-15-2) we find the asymptotic expansion of the difference

$$
(M_{\alpha,r}-\mathcal{R}(B_p,M_{\alpha,r},B_q))(x-t,x+t)\sim \frac{1}{2}a_1^Mt+\ldots,
$$

where from Proposition [3.6](#page-10-1) we see that $a_1^M = \alpha \frac{|t|}{t}$. Therefore, asymptotic inequality [\(4.1\)](#page-15-0) holds when $\alpha > 0$, and the opposite asymptotic inequality holds when $\alpha < 0$.

Observing the limit of $M_{\alpha,r}(s,1-s)$ as $s \to 0$, we see that it is equal to 0 for $r + \alpha \geq 0$ and $-(r + \alpha)$ for $r + \alpha < 0$.

If $\alpha > 0$, then by the same argument as in the Example [4.5,](#page-16-0) mean $M_{\alpha,r}$ cannot be either (B_p, B_q) -sub- or super-stabilizable. If $\alpha < 0$, then mean $M_{\alpha,r}$ can only be (B_p, B_q) -super-stabilizable. If $\alpha = 0$, then $M_{\alpha,r}$ corresponds the logarithmic mean L , whose asymptotic expansion can be found in [\[5\]](#page-17-5):

$$
L(x-t, x+t) \sim x - \frac{1}{3}t^2 x^{-1} - \frac{4}{45}t^4 x^{-3} - \frac{44}{945}t^6 x^{-5} + \mathcal{O}(x^{-7}).
$$

Now we have

$$
(L - \mathcal{R}(B_p, L, B_q))(x - t, x + t) \sim \frac{1}{8}(1 - p - 2q)t^2 x^{-1} + \frac{1}{384}(2p^3 - 3p^2 - 2p(5 + 2q) + 4q(4q(q - 1) - 5) + 11)t^4 x^{-3} + \mathcal{O}(x^{-5}).
$$

If we set $p = 1 - 2q$, then

$$
(L - \mathcal{R}(B_{1-2q}, L, B_q))(x - t, x + t) \sim \frac{1}{96}q(q - 1)t^4x^{-3} + \mathcal{O}(x^{-5}),
$$

and if, additionally, $q = 0$ or $q = 1$, we obtain two pairs of power means for which the resultant mean $\mathcal{R}(B_p, L, B_q)$ is equal to L:

$$
\mathcal{R}(A, L, G) = L, \quad \mathcal{R}(H, L, A) = L,
$$

which can be easily verified by direct computation.

REFERENCES

- [1] M.C. Anisiu, V. Anisiu, The first Seiffert mean is strictly (G, A)-super-stabilizable, J. Inequal. Appl. 185, 1–7, 2014.
- [2] T. Burić, L. Mihoković, *Complete asymptotic expansion of the compound means with appli*cations, J. Class. Anal., 22(1), 67–79, 2023.
- [3] C.-P. Chen, N. Elezović, L. Vukšić, Asymptotic formulae associated with the Wallis power function and digamma function, J. Class. Anal., $2(2)$, $151-166$, 2013 .
- [4] N. Elezović, Asymptotic inequalities and comparison of classical means, J. Math. Inequal., 9(1), 177–196, 2015.
- [5] N. Elezović, L. Vukšić, Asymptotic expansions of bivariate classical means and related in $equalities, J. Math. Inequal., 8(4), 707-724, 2014.$
- [6] N. Elezović, L. Vukšić, Asymptotic expansions and comparison of bivariate parameter means, Math. Inequal. Appl., 17(4), 1225–1244, 2014.
- [7] A. Erdélyi, "Asymptotic expansions", Dover Publications, New York, 1956.
- [8] L. Mihoković, Asymptotic expansions of stable, stabilizable and stabilized means with applications, Appl. Anal. Discrete Math.
- [9] M. Raïssouli, Stability and stabilizability for means, Applied Mathematics, $11(2011)$, 159– 174, 2011.
- [10] M. Raïssouli, A. Rezgui, On a class of bivariate means including a lot of old an new means, Commun. Korean Math. Soc., 34(1), 239–251, 2019.
- [11] M. Raïssouli, J. Sándor, Sub-stabilizability and super-stabilizability for bivariate means, J. Inequal. Appl., 2014(28), 2014.
- [12] M. Raïssouli, J. Sándor, Sub-super-stabilizability of certain bivariate means via meanconvexity, J. Inequal. Appl., 2016(273), 2016.
- [13] L. Vukšić, Seiffert means, asymptotic expansions and related inequalities, Rad Hrvat. Akad. Znan. Umjet. Mat. Znan., 19, 129–142, 2015.

LENKA MIHOKOVIĆ, UNIVERSITY OF ZAGREB, FACULTY OF ELECTRICAL ENGINEERING AND COMputing, Unska 3, 10000 Zagreb, Croatia

Email address: lenka.mihokovic@fer.hr

MUSTAPHA RAÏSSOULI, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, MOULAY ISMAIL University, Meknes, Morocco.

Email address: raissouli.mustapha@gmail.com