

# Dominating $K_t$ -Models

Freddie Illingworth\*     David R. Wood†

May 24, 2024

## Abstract

A *dominating  $K_t$ -model* in a graph  $G$  is a sequence  $(T_1, \dots, T_t)$  of pairwise disjoint non-empty connected subgraphs of  $G$ , such that for  $1 \leq i < j \leq t$  every vertex in  $T_j$  has a neighbour in  $T_i$ . Replacing “every vertex in  $T_j$ ” by “some vertex in  $T_j$ ” retrieves the standard definition of  $K_t$ -model, which is equivalent to  $K_t$  being a minor of  $G$ . We explore in what sense dominating  $K_t$ -models behave like (non-dominating)  $K_t$ -models. The two notions are equivalent for  $t \leq 3$ , but are already very different for  $t = 4$ , since the 1-subdivision of any graph has no dominating  $K_4$ -model. Nevertheless, we show that every graph with no dominating  $K_4$ -model is 2-degenerate and 3-colourable. More generally, we prove that every graph with no dominating  $K_t$ -model is  $2^{t-2}$ -colourable. Motivated by the connection to chromatic number, we study the maximum average degree of graphs with no dominating  $K_t$ -model. We give an upper bound of  $2^{t-2}$ , and show that random graphs provide a lower bound of  $(1 - o(1))t \log t$ , which we conjecture is asymptotically tight. This result is in contrast to the  $K_t$ -minor-free setting, where the maximum average degree is  $\Theta(t\sqrt{\log t})$ . The natural strengthening of Hadwiger’s Conjecture arises: is every graph with no dominating  $K_t$ -model  $(t - 1)$ -colourable? We provide two pieces of evidence for this: (1) It is true for almost every graph, (2) Every graph  $G$  with no dominating  $K_t$ -model has a  $(t - 1)$ -colourable induced subgraph on at least half the vertices, which implies there is an independent set of size at least  $\frac{|V(G)|}{2^{t-2}}$ .

---

\*Department of Mathematics, University College London, United Kingdom (f.illingworth@ucl.ac.uk). Research supported by EPSRC grant EP/V521917/1 and the Heilbronn Institute for Mathematical Research.

†School of Mathematics, Monash University, Melbourne, Australia (david.wood@monash.edu). Research supported by the Australian Research Council.

# 1 Introduction

A *dominating  $K_t$ -model* in a graph  $G$  is a sequence  $(T_1, \dots, T_t)$  of pairwise disjoint non-empty connected subgraphs of  $G$ , such that for  $1 \leq i < j \leq t$  every vertex in  $T_j$  has a neighbour in  $T_i$ . Each  $T_i$  dominates  $T_{i+1} \cup \dots \cup T_t$ , hence the name<sup>1</sup>. Contracting each  $T_i$  into a single vertex and deleting vertices not in  $T_1 \cup \dots \cup T_t$  gives  $K_t$  as a minor. Indeed, in the definition of dominating  $K_t$ -model, replacing “every vertex in  $T_j$ ” by “some vertex in  $T_j$ ” retrieves the standard definition of  $K_t$ -model, which is equivalent to  $K_t$  being a minor of  $G$ .

At first glance, the definition of dominating  $K_t$ -model might seem very restrictive. This paper explores in what sense dominating  $K_t$ -models behave like (non-dominating)  $K_t$ -models. We show that several proof methods in the literature regarding  $K_t$ -models in fact work with dominating  $K_t$ -models. On the other hand, we show some significant differences.

For  $t \leq 3$  it is easily seen that a graph  $G$  has a  $K_t$ -minor if and only if  $G$  has a dominating  $K_t$ -model (Observation 1). However, for  $t = 4$ , the behaviour changes dramatically: there are graphs that contain arbitrarily large complete graph minors, but contain no dominating  $K_4$ -model (Corollary 6). Nevertheless, we show that every graph with minimum degree at least 3 contains a dominating  $K_4$ -model (Theorem 7), thus strengthening a classical result of Hadwiger [10] and Dirac [6].

Now consider the chromatic number  $\chi(G)$ . Hadwiger [10] famously conjectured that every  $K_t$ -minor-free graph is  $(t - 1)$ -colourable. This is widely considered to be one of the most important open problems in combinatorics (see [22] for a survey). The best upper bound on the chromatic number is  $\mathcal{O}(t \log \log t)$  due to Delcourt and Postle [5]. It is open whether  $K_t$ -minor-free graphs are  $\mathcal{O}(t)$ -colourable.

The following natural question arises: what is the maximum chromatic number of a graph with no dominating  $K_t$ -model? We prove that this maximum exists, and in particular, the answer is at most  $3 \cdot 2^{t-4}$  for  $t \geq 4$  (see Theorem 22). Complete graphs provide a lower bound of  $t - 1$ . It is possible that every graph with no dominating  $K_t$ -model is  $(t - 1)$ -colourable, which would be a considerable strengthening of Hadwiger’s Conjecture. We prove this is true for  $t \leq 4$  (see Corollary 9).

We show that two pieces of evidence for Hadwiger’s Conjecture also hold for its dominating version. First, consider independent sets. Let  $\alpha(G)$  be the maximum size of an independent set in a graph  $G$ . Hadwiger’s Conjecture would imply that  $\alpha(G) \geq \frac{n}{t-1}$  for every  $n$ -vertex  $K_t$ -minor-free graph  $G$ . Duchet and Meyniel [7] proved that  $\alpha(G) \geq \frac{n}{2t-2}$  for such  $G$ . We extend this result for graphs with no dominating  $K_t$ -model (Theorem 14).

The second piece of evidence concerns random graphs. The *Hadwiger number* of a

---

<sup>1</sup>For a graph  $G$ , a set  $A \subseteq V(G)$  is *dominating* in  $G$  if every vertex in  $V(G) \setminus A$  has a neighbour in  $A$ . A set  $A \subseteq V(G)$  is *connected* if  $G[A]$  is connected.

graph  $G$ , denoted by  $\text{had}(G)$ , is the maximum integer  $t$  such that  $K_t$  is a minor of  $G$ . Hadwiger's Conjecture asserts that  $\chi(G) \leq \text{had}(G)$  for every graph  $G$ . Similarly, define the *dominating Hadwiger number* of a graph  $G$ , denoted by  $\text{domhad}(G)$ , to be the maximum integer  $t$  such that  $G$  contains a dominating  $K_t$ -model.

Bollobás, Catlin, and Erdős [3] showed that almost every graph  $G$  satisfies  $\chi(G) \leq \text{had}(G)$ . This means that the probability that a random  $n$ -vertex graph satisfies  $\chi(G) \leq \text{had}(G)$  tends to 1 as  $n \rightarrow \infty$ . Strengthening the result of Bollobás et al. [3], we show that almost every graph  $G$  satisfies  $\chi(G) \leq \text{domhad}(G)$  (Corollary 23). The proof of this result shows that the Hadwiger number and the dominating Hadwiger number behave differently for the Erdős-Rényi<sup>2</sup> random graph  $G(n, p)$ . In particular,  $\text{domhad}(G(n, \frac{1}{2})) = (1 + o(1)) \frac{n}{\log_2 n}$  while  $\text{had}(G(n, \frac{1}{2})) = (1 + o(1)) \frac{n}{\sqrt{\log_2 n}}$ .

Random graphs and the chromatic number both relate to the natural extremal questions: What is the maximum average degree of a  $K_t$ -minor-free graph? Kostochka [13, 14] and Thomason [23] independently answered this question, by showing that the maximum average degree of a  $K_t$ -minor-free graph is in  $\Theta(t\sqrt{\log t})$ , where random graphs provide the extremal examples. Later, Thomason [24] determined the leading constant asymptotically. Consider the analogous question for dominating models: What is the maximum average degree of a graph with no dominating  $K_t$ -model? We show that the answer is at most  $2^{t-2}$  (Theorem 16) and that random graphs provide a lower bound of  $(1 - o(1))t \log t$ . Closing this gap is an interesting open problem. We establish a number of results that suggest a  $\mathcal{O}(t \log t)$  upper bound.

## 2 Basics

First note that  $K_t$ -models and dominating  $K_t$ -models are equivalent for  $t \leq 3$ .

**Observation 1.** *For  $t \in \{1, 2, 3\}$ , a graph  $G$  has a dominating  $K_t$ -model if and only if  $G$  has a  $K_t$ -model.*

*Proof.* The  $t = 1$  case holds since a graph has a dominating  $K_1$ -model if and only if  $V(G) \neq \emptyset$  if and only if  $G$  has a  $K_1$ -model. The  $t = 2$  case holds since a graph has a dominating  $K_2$ -model if and only if  $E(G) \neq \emptyset$  if and only if  $G$  has a  $K_2$ -model. The  $t = 3$  case holds since a graph has dominating  $K_3$ -model if and only if  $G$  has a cycle if and only if  $G$  has a  $K_3$ -model. In particular, if  $vw$  is an edge of a cycle  $C$ , then  $(C - v - w, \{v\}, \{w\})$  is a dominating  $K_3$ -model.  $\square$

Note the following elementary observations.

---

<sup>2</sup>The *Erdős-Rényi random graph*  $G(n, p)$  is the  $n$ -vertex graph in which each edge is present with probability  $p$  independently of all other edges.

**Observation 2.** For  $t \geq 2$ , a graph  $G$  has a dominating  $K_t$ -model if and only if  $G$  has a non-empty connected subgraph  $T$  such that  $N_G(T)$  has a dominating  $K_{t-1}$ -model.

**Observation 3.** For  $t \geq 2$ , if a graph  $G$  has a dominating  $K_t$ -model, then  $G$  has a  $K_t$ -model  $(T_1, \dots, T_t)$  where each of  $T_{t-1}$  and  $T_t$  have exactly one vertex.

*Proof.* This follows from Observation 2 and the fact that a graph has a dominating  $K_2$ -model if and only if it contains an edge.  $\square$

**Lemma 4.** If a graph  $G$  has a dominating  $K_t$ -model, then  $G$  has a path  $v_1, \dots, v_t$  where  $\deg_G(v_t) \geq t - 1$ , and  $\deg_G(v_i) \geq i$  for  $1 \leq i \leq t - 1$ .

*Proof.* Say  $(T_1, \dots, T_t)$  is a dominating  $K_t$ -model in  $G$ . Let  $v_t$  be any vertex in  $T_t$ . Then  $v_t$  has a neighbour in each of  $T_1, \dots, T_{t-1}$ , so  $\deg_G(v_t) \geq t - 1$ , as claimed. For  $i = t - 1, t - 2, \dots, 1$ , let  $v_i$  be a vertex in  $T_i$  adjacent to  $v_{i+1}$ . Each  $v_i$  has a neighbour in each of  $T_1, \dots, T_{i-1}, T_{i+1}$ , so  $\deg_G(v_i) \geq i$ . By construction,  $v_1, \dots, v_t$  is a path.  $\square$

Lemma 4 implies the following.

**Corollary 5.** Every graph with maximum degree  $\Delta$  contains no dominating  $K_{\Delta+2}$ -model.

Corollary 5 highlights a big difference between  $K_t$ -models and dominating  $K_t$ -models, since there are  $n$ -vertex graphs with maximum degree 3 that contain  $K_{c\sqrt{n}}$ -models, but contain no dominating  $K_5$ -model by Corollary 5. This difference is also highlighted by the following result, which is an immediate consequence of Lemma 4.

**Corollary 6.** For  $n \geq 4$ , if  $G$  is any graph obtained from  $K_n$  by subdividing each edge at least once, then  $G$  contains no dominating  $K_4$ -model.

We conclude this section by mentioning a curious consequence of Corollary 5. If  $\chi(G) \leq \text{domhad}(G)$  for every graph  $G$ , then, Corollary 5 would imply that every graph with maximum degree  $\Delta$  is  $(\Delta + 1)$ -colourable, which is true since such graphs are  $\Delta$ -degenerate.

### 3 Dominating $K_4$ -Models

Hadwiger [10] and Dirac [6] proved that every graph with minimum degree at least 3 has a  $K_4$ -model. Here we strengthen this result.

**Theorem 7.** Every graph  $G$  with minimum degree at least 3 has a dominating  $K_4$ -model.

*Proof.* We may assume that  $G$  is connected. Since  $\delta(G) \geq 3$ ,  $G$  has a cycle. Choose a pair  $(C, H)$  where  $C$  is a cycle of  $G$  and  $H$  is a component of  $G - C$  such that  $|V(H)|$  is maximum (over all choices of  $(C, H)$ ) and, subject to this,  $|V(C)|$  is minimum. By minimality,  $C$  is an induced cycle of  $G$ . Since  $\delta(G) \geq 3$ ,  $H$  is non-empty. Since  $G$  is connected, there is a vertex  $z$  of  $C$  with at least one neighbour in  $H$ .

Suppose that  $G - (V(H) \cup \{z\})$  contains a cycle  $C'$ . Then  $G - C'$  has a component containing  $\{z\} \cup V(H)$  which contradicts the choice of  $(C, H)$ . Thus

$$G - (V(H) \cup \{z\}) \text{ is a forest.} \quad (\dagger)$$

Suppose that  $G - C$  has a component  $H' \neq H$ . If  $H'$  is a single vertex  $a$ , then  $a$  must have at least three neighbours in  $C$ , and so has neighbours  $u, v \in C - z$ . Let  $P_{uv}$  be the path in  $C$  from  $u$  to  $v$  avoiding  $z$ . Then  $va, au$ , and  $P_{uv}$  form a cycle contradicting  $(\dagger)$ . Thus  $H'$  has at least two vertices. By  $(\dagger)$ ,  $H'$  induces a tree and so has at least two leaves. Each leaf must have at least two neighbours in  $C$  and so has a neighbour in  $C - z$ . Let  $a_1$  and  $a_2$  be leaves of  $H'$ , and  $v_1, v_2 \in C - z$  be neighbours of  $a_1$  and  $a_2$ , respectively. Let  $P_{a_2a_1}$  be the path in  $H'$  from  $a_2$  to  $a_1$  and  $P_{v_1v_2}$  be the path in  $C$  from  $v_1$  to  $v_2$  avoiding  $z$ . Then  $P_{a_2a_1}, a_1v_1, P_{v_1v_2}$ , and  $v_2a_2$  form a cycle contradicting  $(\dagger)$ . Thus  $H$  is the only component of  $G - C$ .

Since  $C$  is an induced cycle of  $G$  and  $\delta(G) \geq 3$ , every vertex of  $C$  has a neighbour in  $H$ . Let  $xy$  be an edge of  $C$  and  $P$  be the path  $C - x - y$ . Then  $(H, P, \{x\}, \{y\})$  is a dominating  $K_4$ -model in  $G$ .  $\square$

Theorem 7 implies the following results.

**Corollary 8.** *For  $n \geq 2$ , every  $n$ -vertex graph with no dominating  $K_4$ -model has at most  $2n - 3$  edges.*

Hadwiger [10] and Dirac [6] proved that every  $K_4$ -minor-free graph is 3-colourable. Theorem 7 implies the following strengthening.

**Corollary 9.** *Every graph with no dominating  $K_4$ -model is 2-degenerate and 3-colourable.*

These two results suggest that dominating  $K_4$ -models behave like  $K_4$ -models, although we emphasise that there are graphs with no dominating  $K_4$ -model that contain arbitrarily large complete graph minors (Corollary 6).

## 4 Random Graphs

The main result of this section (Theorem 11) asymptotically determines the dominating Hadwiger number of  $G(n, p)$ . We will ignore ceilings and floors in this section. We first need the following technical lemma.

**Lemma 10.** For fixed  $p \in (0, 1)$ ,

$$\mathbb{P}(G(n, p) \text{ is not connected}) = \mathcal{O}(n(1-p)^n),$$

where the implied constant may depend on  $p$  but not on  $n$ .

*Proof.* Say a set  $S$  of vertices in  $G(n, p)$  is *isolated* if there are no edges between  $S$  and the rest of the vertices. For a fixed set  $S$  of  $k$  vertices,

$$\mathbb{P}(S \text{ is isolated}) = (1-p)^{k(n-k)}.$$

If  $G(n, p)$  is not connected, then there is some non-empty set of at most  $n/2$  vertices that is isolated. Thus, taking a union bound and using the inequality  $\binom{n}{k} \leq n^k$  gives

$$\begin{aligned} \mathbb{P}(G(n, p) \text{ is not connected}) &\leq \sum_{k=1}^{n/2} \binom{n}{k} (1-p)^{k(n-k)} \\ &\leq n(1-p)^{n-1} + n^2(1-p)^{2(n-2)} + \sum_{k=3}^{n/2} (n(1-p)^{n-k})^k \\ &\leq \mathcal{O}(n(1-p)^n) + \sum_{k=3}^{n/2} (n(1-p)^{n/2})^k \\ &\leq \mathcal{O}(n(1-p)^n) + \frac{(n(1-p)^{n/2})^3}{1 - n(1-p)^{n/2}} \\ &= \mathcal{O}(n(1-p)^n). \quad \square \end{aligned}$$

**Theorem 11.** For constant  $p \in (0, 1)$ , asymptotically almost surely,

$$\text{domhad}(G(n, p)) = (1 + o(1)) \frac{n}{\log_b n},$$

where  $b = \frac{1}{1-p}$ .

*Proof.* We first prove the lower bound. Fix  $\varepsilon > 0$  and let  $t = \frac{n}{(1+\varepsilon)\log_b n}$ . Partition  $V(G(n, p))$  into  $t$  parts,  $V_1, \dots, V_t$ , each of size  $n/t = (1+\varepsilon)\log_b n$ . We call a part  $V_i$  *good* if the subgraph of  $G(n, p)$  induced by  $V_i$  is connected, and a pair  $(V_i, V_j)$  (with  $i < j$ ) *good* if  $V_j \subseteq N(V_i)$ .

Note that  $(1-p)^{n/t} = n^{-1-\varepsilon}$ . Since the graph induced by  $V_i$  is  $G(n/t, p)$ , Lemma 10 gives

$$\mathbb{P}(V_i \text{ is bad}) = \mathcal{O}((1+\varepsilon)(\log_b n)n^{-1-\varepsilon}).$$

By the union bound, the probability that some  $V_i$  is bad is  $\mathcal{O}((1+\varepsilon)(\log_b n)n^{-\varepsilon}) = o(1)$ . Hence, with high probability, there are no bad parts.

The probability that a fixed vertex in  $V_j$  has no neighbour in  $V_i$  is  $(1-p)^{|V_i|}$ . Thus,

$$\mathbb{P}((V_i, V_j) \text{ is good}) = (1 - (1-p)^{|V_i|})^{|V_j|}. \quad (1)$$

Using the inequality  $(1+x)^r \geq 1+rx$  for  $r \geq 1$  and  $x \geq -1$  (this is Bernoulli's inequality) we obtain

$$\mathbb{P}((V_i, V_j) \text{ is bad}) = 1 - (1 - n^{-1-\varepsilon})^{n/t} \leq n^{-\varepsilon} t^{-1}.$$

Hence, the expected number of bad pairs of parts is at most

$$\binom{t}{2} \cdot n^{-\varepsilon} t^{-1} \leq t n^{-\varepsilon}.$$

By Markov's inequality, the probability that there are at least  $\varepsilon t$  bad pairs is at most  $\varepsilon^{-1} n^{-\varepsilon} = o(1)$ . Thus, with high probability, the number of bad pairs is at most  $\varepsilon t$  and there are no bad parts. In this case, deleting one part from each bad pair leaves a dominating clique model. Thus, with high probability,  $G(n, p)$  contains a dominating  $K_s$ -model where

$$s \geq (1-\varepsilon)t = \frac{1-\varepsilon}{1+\varepsilon} \cdot \frac{n}{\log_b n} \geq (1-2\varepsilon) \frac{n}{\log_b n}.$$

We now prove the upper bound. Define a *dominating pseudo- $K_t$ -model* in a graph  $G$  to be a sequence  $(V_1, \dots, V_t)$  of pairwise disjoint non-empty subsets of  $V(G)$ , such that  $V_j \subseteq N(V_i)$  for  $1 \leq i < j \leq t$  (this is just a dominating  $K_t$ -model without the connectedness condition).

Fix  $\varepsilon > 0$  and take  $t-1 = (1+\varepsilon) \frac{n-1}{\log_b n}$ . It suffices to show that  $G(n, p)$  does not contain a dominating pseudo- $K_t$ -model with high probability. If  $(V_1, \dots, V_t)$  is a dominating pseudo- $K_t$ -model and  $v \in V_t$ , then  $(V(G) \setminus (V_2 \cup \dots \cup V_{t-1} \cup \{v\}), V_2, \dots, V_{t-1}, \{v\})$  is also a dominating pseudo- $K_t$ -model. Thus it suffices to show that, with high probability,  $G(n, p)$  does not contain a dominating pseudo- $K_t$ -model where  $|V_t| = 1$  and  $V_1 \cup \dots \cup V_t = V(G(n, p))$ .

Fix a list of non-empty vertex sets  $(V_1, \dots, V_t)$  that partition  $V(G(n, p))$  and such that  $|V_t| = 1$ . Then  $(V_1, \dots, V_t)$  is a dominating pseudo- $K_t$ -model if and only if  $(V_i, V_j)$  is good for all  $1 \leq i < j \leq t$ . By (1), and since  $1-x \leq e^{-x}$  for  $x \geq 0$ ,

$$\begin{aligned} \mathbb{P}((V_1, \dots, V_t) \text{ is a dominating pseudo-}K_t\text{-model}) &= \prod_{j=1}^t \prod_{i=1}^{j-1} (1 - (1-p)^{|V_i|})^{|V_j|} \\ &\leq \prod_{j=1}^t \prod_{i=1}^{j-1} \exp\{-|V_j|(1-p)^{|V_i|}\} \\ &= \exp\left\{-\sum_{1 \leq i < j \leq t} |V_j|(1-p)^{|V_i|}\right\}. \end{aligned}$$

We now prove a lower bound for the sum in the parentheses. Since each  $V_j$  has size at least 1,

$$\sum_{1 \leq i < j \leq t} |V_j|(1-p)^{|V_i|} \geq \sum_{1 \leq i < j \leq t} (1-p)^{|V_i|} = \sum_{i=1}^{t-1} (t-i) \cdot (1-p)^{|V_i|}.$$

The AM-GM inequality and the fact that  $|V_t| = 1$  gives

$$\begin{aligned} \sum_{i=1}^{t-1} (t-i)(1-p)^{|V_i|} &\geq (t-1) \left[ \prod_{i=1}^{t-1} (t-i)(1-p)^{|V_i|} \right]^{\frac{1}{t-1}} \\ &= (t-1) \left[ (t-1)! (1-p)^{\sum_{i=1}^{t-1} |V_i|} \right]^{\frac{1}{t-1}} \\ &= (t-1) [(t-1)!]^{\frac{1}{t-1}} (1-p)^{\frac{n-1}{t-1}} \\ &\geq \frac{(t-1)^2}{e} (1-p)^{\frac{n-1}{t-1}}, \end{aligned}$$

where the final inequality used the lower bound  $k! > \left(\frac{k}{e}\right)^k$ , which is valid for all positive integers  $k$ . In particular,

$$\mathbb{P}((V_1, \dots, V_t) \text{ is a dominating pseudo-}K_t\text{-model}) \leq \exp\left\{-\frac{(t-1)^2}{e} (1-p)^{\frac{n-1}{t-1}}\right\}.$$

Since the number of partitions of  $V(G(n, p))$  into  $t$  parts is  $t^n$ ,

$$\begin{aligned} \mathbb{E}(\# \text{ of dominating pseudo-}K_t\text{-models}) &\leq t^n \exp\left\{-\frac{(t-1)^2}{e} (1-p)^{\frac{n-1}{t-1}}\right\} \\ &= \exp\left\{n \log t - \frac{(t-1)^2}{e} (1-p)^{\frac{n-1}{t-1}}\right\}. \end{aligned}$$

Recall that  $t-1 = (1+\varepsilon)\frac{n-1}{\log_b n}$ . Thus  $(1-p)^{\frac{n-1}{t-1}} = n^{-1/(1+\varepsilon)}$  and

$$n \log t - \frac{(t-1)^2}{e} (1-p)^{\frac{n-1}{t-1}} \leq n \log n - \frac{(1+\varepsilon)^2}{e(\log_b n)^2} (n-1)^2 n^{-\frac{1}{1+\varepsilon}}.$$

Since  $2 - \frac{1}{1+\varepsilon} > 1 + \frac{\varepsilon}{2}$  (assuming  $\varepsilon < 1$ ), the right-hand side tends to  $-\infty$  as  $n \rightarrow \infty$ . Thus, the expected number of dominating pseudo- $K_t$ -models, and so the probability that there is a dominating pseudo- $K_t$ -model is  $o(1)$ , as required.  $\square$

## 5 Structure

This section explores the structure of graphs with no dominating  $K_t$ -model, and concludes a lower bound on the independence number of such graphs. The proof follows that of Duchet and Meyniel [7] for  $K_t$ -minor-free graphs.

**Lemma 12** ([7]). *Every non-empty connected graph has a connected dominating set  $D$  and an independent set  $I \subseteq D$  with  $|D| = 2|I| - 1$ .*



*Proof.* Let  $D$  be the largest connected set in  $G$  such that  $D$  contains an independent set  $I$  with  $|D| = 2|I| - 1$ . This is well-defined since  $D = I = \{v\}$  satisfies these properties for any vertex  $v$ . Suppose for the sake of contradiction that  $D$  is not dominating. Since  $G$  is connected, there exists an edge  $xy$  in  $G$  with  $\text{dist}_G(x, D) = 1$  and  $\text{dist}_G(y, D) = 2$ . Thus  $D' := D \cup \{x, y\}$  is a connected set and  $I' := I \cup \{y\}$  is an independent set in  $D'$  with  $|D'| = 2|I'| - 1$ , contradicting the choice of  $D$  and  $I$ . Hence  $D$  is dominating.  $\square$

A *partition* of a graph  $G$  is a partition of  $V(G)$  into non-empty sets. Each element of a partition is called a *part*. Let  $\mathcal{P}$  be a partition of a graph  $G$ . The *quotient* of  $\mathcal{P}$  is the graph with vertex-set  $\mathcal{P}$  where distinct  $A, B \in \mathcal{P}$  are adjacent if and only if there is an edge of  $G$  between  $A$  and  $B$ . If  $H$  is isomorphic to the quotient of  $\mathcal{P}$ , then  $\mathcal{P}$  is called an  *$H$ -partition* of  $G$ . We say  $\mathcal{P}$  is *connected* if each part induces a connected subgraph of  $G$ , in which case the quotient is a minor of  $G$ .

The *depth* of a vertex  $v$  in a tree rooted at  $r$  is the number of vertices in the  $vr$ -path in  $T$ . The *closure* of a rooted tree  $T$  is the graph  $\widehat{T}$  with vertex-set  $V(T)$ , where  $vw$  is an edge of  $\widehat{T}$  if and only if  $v$  is an ancestor or descendent of  $w$  in  $T$ . A graph that is isomorphic to the closure of a tree is called *trivially perfect*, a *comparability graph of a tree*, an *arborescent comparability graph*, or a *quasi-threshold graph* [26]. They are the graphs that contain neither  $P_4$  nor  $C_4$  as induced subgraphs.

**Theorem 13.** *For every connected graph  $G$  there is a tree  $T$  rooted at  $r$  such that  $G$  has a connected  $\widehat{T}$ -partition  $\mathcal{P}$  where:*

- *for every part  $P \in \mathcal{P}$  there is an independent set  $I$  in  $G[P]$  with  $|P| = 2|I| - 1$ , and*
- *for every leaf  $v$  of  $T$ , the sequence of parts in  $\mathcal{P}$  from  $r$  to  $v$  define a dominating complete graph model in  $G$ .*

*Proof.* We proceed by induction on  $|V(G)|$ . The case  $|V(G)| = 1$  is trivial. By Lemma 12,  $G$  has a connected dominating set  $D$  and an independent set  $I \subseteq D$  with  $|D| = 2|I| - 1$ . Let  $G_1, \dots, G_c$  be the components of  $G - D$ . By induction, for each  $i \in \{1, \dots, c\}$ , there is a tree  $T_i$  rooted at  $r_i$  such that  $G_i$  has a  $\widehat{T}_i$ -partition  $\mathcal{P}_i$  such that:

- *for every part  $P \in \mathcal{P}_i$  there is an independent set  $I$  in  $G_i[P]$  with  $|P| = 2|I| - 1$ , and*
- *for every leaf  $v$  of  $T_i$ , the sequence of parts in  $\mathcal{P}_i$  from  $r$  to  $v$  define a dominating complete graph model in  $G_i$ .*

Let  $T$  be the tree obtained from the disjoint union of  $T_1, \dots, T_c$  by adding a new root vertex  $r$  adjacent to  $r_1, \dots, r_c$ . So  $\widehat{T}$  is obtained from the disjoint union of  $\widehat{T}_1, \dots, \widehat{T}_c$  by adding vertex  $r$ , adjacent to every other vertex. Let  $\mathcal{P}$  be the  $\widehat{T}$ -partition of  $G$  obtained by associating  $D$  with  $r$ . Since  $D$  is dominating, for every leaf  $v$  of  $T$ , the sequence of parts in  $\mathcal{P}$  from  $r$  to  $v$  define a dominating complete graph model in  $G$ .  $\square$

**Theorem 14.** For  $t \geq 2$  and every graph  $G$  with no dominating  $K_t$ -model, there is an integer  $h \leq t-1$  and an  $h$ -colourable induced subgraph of  $G$  on at least  $\frac{|V(G)|+h}{2}$  vertices. In particular,

$$\alpha(G) \geq \frac{|V(G)|+t-1}{2t-2}.$$

*Proof.* We may assume  $G$  is connected. Apply Theorem 13 to  $G$ . Let  $T$  be the resulting tree, and let  $\mathcal{P}$  be the resulting  $\hat{T}$ -partition of  $G$ . Let  $h$  be the maximum depth of a node in  $T$ . Since  $G$  has no dominating  $K_t$ -model,  $h \leq t-1$ . Each part  $A \in \mathcal{P}$  contains an independent set  $I_A$  of size  $(|A|+1)/2$ . Let  $X$  be the union of the  $I_A$ . Then

$$|X| \geq \sum_{A \in \mathcal{P}} \frac{|A|+1}{2} = \frac{|V(G)|+|\mathcal{P}|}{2} \geq \frac{|V(G)|+h}{2}.$$

Colour each vertex in  $I_A$  by the depth of the corresponding node in  $T$ . We obtain a proper  $h$ -colouring of  $X$ . Taking the largest colour class in  $X$ ,

$$\alpha(G) \geq \frac{|V(G)|+h}{2h} = \frac{|V(G)|}{2h} + \frac{1}{2} \geq \frac{|V(G)|}{2t-2} + \frac{1}{2} = \frac{|V(G)|+t-1}{2t-2}. \quad \square$$

## 6 Average Degree

This section considers the maximum average degree of a graph with no dominating  $K_t$ -model. Theorem 11 implies the following.

**Corollary 15.** For any  $\varepsilon > 0$  and for sufficiently large  $t$ , there is a graph  $G$  with no dominating  $K_t$ -model and average degree at least  $(1-\varepsilon)t \ln t$ .

*Proof.* Note that  $\ln(\frac{1}{1-x})/x \rightarrow 1$  as  $x \rightarrow 0$ . Fix  $\varepsilon > 0$  small and let  $p > 0$  be small enough so that  $\ln(\frac{1}{1-p}) < (1+\frac{\varepsilon}{3})p$ . For large  $n$ , with high probability,  $G(n, p)$  has average degree at least  $(1-\varepsilon^2)pn$ , and by Theorem 11,

$$\text{domhad}(G(n, p)) \leq (1+\frac{\varepsilon}{3}) \frac{n}{\log_b n} = (1+\frac{\varepsilon}{3}) \frac{n \ln b}{\ln n} < (1+\frac{\varepsilon}{3})^2 \frac{pn}{\ln n} < (1+\varepsilon) \frac{pn}{\ln n},$$

Pick an instance  $G$  of  $G(n, p)$  satisfying both these conditions. Then  $G$  has no dominating  $K_t$ -model where  $t := (1+\varepsilon) \frac{pn}{\log n}$ . Finally,

$$(1-\varepsilon)t \ln t \leq (1-\varepsilon)(1+\varepsilon) \frac{pn}{\ln n} \cdot \ln n = (1-\varepsilon^2)pn$$

which is at most the average degree of  $G$ , as required.  $\square$

Mader [16] first showed that every graph with sufficiently high average degree contains a  $K_t$ -minor. The proof method actually shows the following:

**Theorem 16.** For  $t \geq 2$  every graph with average degree at least  $2^{t-2}$  has a dominating  $K_t$ -model.

*Proof.* We proceed by induction on  $n + t$  with the following hypothesis: for every connected  $n$ -vertex graph  $G$  with average degree at least  $2^{t-2}$  and for every vertex  $v$  of  $G$ , there exists a dominating  $K_t$ -model  $(T_1, \dots, T_t)$  in  $G$  with  $v \in V(T_1)$ . This implies the claim since every graph with average degree at least  $2^{t-2}$  has a connected component with average degree at least  $2^{t-2}$ .

The  $t = 2$  case holds with  $T_1 = G[\{v\}]$  and  $T_2 = G[\{w\}]$  where  $w$  is any neighbour of  $v$  (which exists since  $G$  is connected with average degree at least 1). If  $n \leq 2^{t-2} + 1$  then  $G$  is complete, and the result holds trivially. Now assume that  $t \geq 3$  and  $n > 2^{t-2} + 1$ . Let  $G$  be a connected  $n$ -vertex  $m$ -edge graph  $G$  with average degree at least  $2^{t-2}$ , and let  $v$  be a vertex of  $G$ . So  $2m \geq 2^{t-2}n$ .

For each neighbour  $w$  of  $v$ , let  $d_w$  be the number of common neighbours of  $v$  and  $w$ . Suppose that  $d_w \leq 2^{t-3} - 1$  for some neighbour  $w$  of  $v$ . Let  $G'$  be the graph obtained from  $G$  by contracting  $vw$  into a new vertex  $v'$ . So  $G'$  is connected,  $|E(G')| = m - 1 - d_w$  and  $|V(G')| = n - 1$ . Thus  $G'$  has average degree

$$\frac{2(m - 1 - d_w)}{n - 1} \geq \frac{2^{t-2}n - 2(1 + d_w)}{n - 1} \geq \frac{2^{t-2}n - 2^{t-2}}{n - 1} = 2^{t-2}.$$

By induction, there exists a dominating  $K_t$ -model  $(T_1, \dots, T_t)$  in  $G'$  with  $v' \in V(T_1)$ . Let  $T'_1$  be the tree obtained from  $T_1$  by replacing  $v'$  by  $vw$ , and replacing each edge  $v'x$  in  $T_1$  by  $vx$  or  $wx$  (one of which must exist). By construction, every vertex adjacent to  $v'$  in  $G'$  is adjacent to  $v$  or  $w$  in  $G$ . Thus  $(T'_1, T_2, \dots, T_t)$  is a dominating  $K_t$ -model in  $G$  with  $v \in V(T'_1)$ , as desired.

Now assume that every edge incident to  $v$  is in at least  $2^{t-3}$  triangles. Since  $G$  is connected and with at least two vertices, there is at least one edge incident to  $v$ . Thus  $G[N_G(v)]$  has minimum degree at least  $2^{t-3}$ . By induction,  $G[N_G(v)]$  has a dominating  $K_{t-1}$ -model  $(T_1, \dots, T_{t-1})$ . Thus  $(\{v\}, T_1, \dots, T_{t-1})$  is the desired dominating  $K_t$ -model in  $G$ .  $\square$

We conjecture that average degree  $\Omega(t \log t)$  is enough to force a dominating  $K_t$ -model (and Corollary 15 shows this would be best possible). As a step towards this conjecture, we show it is true for graphs with linear minimum degree. The following proofs make no attempt to optimise constants. The first lemma is well-known.

**Lemma 17.** *Every connected graph  $G$  with  $n$  vertices and minimum degree  $\delta > \ln(2n)$  has a connected dominating set on at most  $\frac{6n \ln(2n)}{\delta}$  vertices.*

*Proof.* Let  $p := \ln(2n)/\delta < 1$ . For each vertex  $v$ , choose  $v$  independently at random with probability  $p$ . Let  $A$  be the set of chosen vertices. Let  $B$  be the set of vertices  $v$  such that  $N_G(v) \cap A = \emptyset$ . Thus  $\mathbb{P}(v \in B) = (1 - p)^{\deg(v)} < e^{-p\delta} = \frac{1}{2n}$ . Thus  $\mathbb{E}(|B|) < \frac{1}{2}$ , and by Markov's Inequality,  $\mathbb{P}(|B| \geq 1) \leq \frac{\mathbb{E}(|B|)}{1} < \frac{1}{2}$ . Now consider  $|A|$ :  $\mathbb{E}(|A|) = pn$  and so, by Markov's Inequality,  $\mathbb{P}(|A| \geq 2pn) \leq \frac{1}{2}$ . A union bound gives

$\mathbb{P}(|B| \geq 1 \text{ or } |A| \geq 2pn) < 1$ . Hence, there exists  $A \subseteq V(G)$  with  $|A| < 2pn = \frac{2n \ln(2n)}{\delta}$  and  $B = \emptyset$ . Since  $B = \emptyset$ ,  $A$  is a dominating set. The result follows, since Duchet and Meyniel [7] showed that if a connected graph has a dominating set on  $k$  vertices, then it has a connected dominating set on at most  $3k - 2$  vertices.  $\square$

The next lemma is similar to several results in the literature on  $K_t$ -minors [11, 15].

**Lemma 18.** *Fix  $c \in (0, 1)$  and  $n$  such that  $cn > \ln(2n)$ . Then every graph  $G$  with at most  $n$  vertices and minimum degree at least  $cn + 6c^{-1}t \ln(2n)$  has a dominating  $K_t$ -model.*

*Proof.* We proceed by induction on  $t$ . The case  $t = 1$  is trivial. We may assume that  $G$  is connected, since any component  $G'$  of  $G$  has  $\delta(G') \geq \delta(G) \geq cn + 6c^{-1}t \ln(2n)$  and  $|V(G')| \leq |V(G)| \leq n$ . Since  $G$  has minimum degree at least  $cn$ , by Lemma 17,  $G$  has a connected dominating set  $A$  with  $|A| \leq 6c^{-1} \ln(2n)$ . Let  $G' := G - A$ . So  $G'$  has at most  $n$  vertices and minimum degree at least  $cn + 6c^{-1}(t - 1) \ln(2n)$ . By induction,  $G'$  has a dominating  $K_{t-1}$ -model  $(A_1, \dots, A_{t-1})$ . Since  $A$  is dominating,  $(A, A_1, \dots, A_{t-1})$  is a dominating  $K_t$ -model in  $G$ .  $\square$

**Proposition 19.** *For any  $c \in (0, 1)$  and sufficiently large integers  $n, t$  with  $n \geq c^{-2}t \log_2 t$ , every graph with  $n$  vertices and minimum degree at least  $2cn$  has a dominating  $K_t$ -model.*

*Proof.* We may assume that  $t \geq (2c^{-2} \log_2 t)^6$ . Thus

$$\log_2 t \geq \ln t \geq 6 \ln(2c^{-2} \log_2 t).$$

Since  $\frac{n}{\ln(2n)}$  is increasing and  $n \geq c^{-2}t \log_2 t$ ,

$$\frac{c^2 n}{\ln(2n)} \geq \frac{t \log_2 t}{\ln(2c^{-2}t \log_2 t)} \geq 6t.$$

Therefore

$$2cn \geq cn + 6c^{-1}t \ln(2n).$$

The result follows from Lemma 18 with minimum degree  $2cn$ , since we may assume that  $cn > \ln(2n)$ .  $\square$

Here is further evidence for the conjecture.

**Proposition 20.** *For sufficiently large  $t$  and  $d = 4t \ln t$ , every  $d$ -regular graph  $G$  contains a dominating pseudo- $K_t$ -model.*

*Proof.* Uniformly and randomly put each vertex of  $G$  into one of  $t$  parts. For each vertex  $v$ , let  $B_v$  be the event that  $v$  has no neighbour in some part. Note that  $\mathbb{P}(B_v)$  equals the probability that a coupon collector has not succeeded by time  $4t \ln t$ , which is at most  $t^{-3}$  by standard tail estimate for the coupon collector problem [25]. Note that  $B_v$  depends

only on the parts where the neighbours of  $v$  were placed, and so  $B_v$  is independent of  $(B_u: \text{dist}(u, v) \geq 3)$ . There are at most  $d^2$  vertices within distance 2 of  $v$  (other than  $v$ ). Since  $4t^{-3}d^2 < 1$  for sufficiently large  $t$ , the Lovász Local Lemma [8] implies that no  $B_v$  occurs with positive probability. That is, there exists a partition of  $V(G)$  such that every vertex in  $G$  has a neighbour in every part. This partition defines a dominating pseudo- $K_t$ -model.  $\square$

This proof still works if all degrees are close to each other, but doesn't work if the degrees vary wildly.

## 7 Colouring

This section considers the chromatic number of graphs with no dominating  $K_t$ -model. Mader [16] showed that every  $K_t$ -minor-free graph is  $2^{t-2}$ -colourable. The proof generalises as follows:

**Theorem 21.** *For  $t \geq 2$ , every graph with no dominating  $K_t$ -model is  $2^{t-2}$ -colourable.*

*Proof.* We proceed by induction on  $t$ . Every graph  $G$  with no dominating  $K_2$ -model has no edges, implying  $G$  is 1-colourable. Now assume that  $t \geq 3$ , and the result holds for  $t - 1$ . Let  $G$  be a graph with no dominating  $K_t$ -model. We may assume that  $G$  is connected. Let  $r$  be any vertex in  $G$ . For  $i \geq 0$ , let  $V_i := \{v \in V(G) : \text{dist}_G(v, r) = i\}$ . So  $V_0 = \{r\}$ , and for  $i \geq 1$ , every vertex in  $V_i$  has a neighbour in  $V_{i-1}$ . If for some  $i \geq 1$ , there is a dominating  $K_{t-1}$ -model  $(T_1, \dots, T_{t-1})$  in  $G[V_i]$ , then  $(G[V_0 \cup \dots \cup V_{i-1}], T_1, \dots, T_{t-1})$  is a dominating  $K_t$ -model in  $G$ . Thus for every  $i \geq 0$ , there is no dominating  $K_{t-1}$ -model in  $G[V_i]$ . By induction, each  $G[V_i]$  is  $2^{t-3}$ -colourable. Use the same set of  $2^{t-3}$  colours for  $\bigcup(V_i : i \text{ even})$  and use a disjoint set of  $2^{t-3}$  colours for  $\bigcup(V_i : i \text{ odd})$ . Since there is no edge between  $V_i$  and  $V_j$  with  $j \geq i + 2$ , we obtain a  $2^{t-2}$ -colouring of  $G$ .  $\square$

Note that Theorem 16 implies that every graph with no dominating  $K_t$ -model is  $(2^{t-2} - 1)$ -degenerate and thus  $2^{t-2}$ -colourable, which provides an alternative proof of Theorem 21.

The proof of Theorem 21 in conjunction with Corollary 9 in the base case shows:

**Theorem 22.** *For  $t \geq 4$ , every graph with no dominating  $K_t$ -model is  $3 \cdot 2^{t-4}$ -colourable.*

Now consider the chromatic number of random graphs. For constant  $p \in (0, 1)$ , the asymptotic value of  $\chi(G(n, p))$  was determined independently by Bollobás [2] and Matula and Kučera [17]. They proved that, with high probability,

$$\chi(G(n, p)) = \left(\frac{1}{2} + o(1)\right) \frac{n}{\log_b n}, \quad (2)$$

where  $b := \frac{1}{1-p}$ . See Heckel [12] for the most precise estimates currently known.

Bollobás, Catlin, and Erdős [3] showed that for fixed  $p \in (0, 1)$ , with high probability,

$$\text{had}(G(n, p)) = (1 + o(1)) \frac{n}{\sqrt{\log_b n}}.$$

It follows that Hadwiger's conjecture comfortably holds for almost every graph.

Theorem 11 and (2) imply that:

**Corollary 23.** *Almost every graph  $G$  satisfies  $\chi(G) \leq \text{domhad}(G)$ .*

It is curious that the chromatic number and dominating Hadwiger number asymptotically differ only by a factor of 2, compared to a factor of  $\sqrt{\log_2 n}$  in the case of the Hadwiger number.

We finish this section with the following amusing observation. Let  $G$  be a graph with  $\chi(G) = t$ . Choose a proper colouring of  $G$  with colours  $1, \dots, t$  minimising  $\sum_{v \in V(G)} \text{col}(v)$ . Then for  $1 \leq i < j \leq t$ , every vertex of colour  $j$  is adjacent to a vertex of colour  $i$ . In particular, the colour classes form a dominating pseudo- $K_t$ -model. This says that the 'dominating pseudo-Hadwiger conjecture' is true!

## 8 Open Problems

We finish with a number of open problems.

- Several authors [1, 4, 18, 19] have noted that graphs  $G$  with independence number  $\alpha(G) = 2$  are a key unsolved case of Hadwiger's Conjecture. Plummer, Stiebitz, and Toft [19] showed that Hadwiger's Conjecture holds for this class if and only if every  $n$ -vertex graph  $G$  with  $\alpha(G) = 2$  has a  $K_t$ -minor with  $t \geq \lceil \frac{n}{2} \rceil$ . The following natural question arises: Does every  $n$ -vertex graph with  $\alpha(G) = 2$  have a dominating  $K_t$ -model with  $t \geq \lceil \frac{n}{2} \rceil$ ? Note that if  $\alpha(G) = 2$  and  $G$  has no dominating  $K_t$ -model, then  $2 \geq \alpha(G) \geq \frac{n+t-1}{2t-2}$  by Theorem 14, implying  $n \leq 3t - 3$ . That is, if  $\alpha(G) = 2$  then  $G$  has a dominating  $K_t$ -model, where  $t \geq \lceil \frac{n}{3} \rceil$ .
- The following potential strengthening of the 4-Colour Theorem is open: Is every graph with no dominating  $K_5$ -model 4-colourable?
- Does every graph with no dominating  $K_t$ -model have fractional chromatic number at most  $2t - 2$ ? Reed and Seymour [20] proved this for  $K_t$ -minor-free graphs.
- Is there a rough structure theorem for graphs with no dominating  $K_t$ -model (in the spirit of Robertson and Seymour's rough structure theorem for  $K_t$ -minor-free graphs [21] and Grohe and Marx's rough structure theorem for  $K_t$ -topological-minor-free graphs [9])?

## References

- [1] JONAH BLASIAK. [A special case of Hadwiger’s conjecture](#). *J. Combin. Theory Ser. B*, 97(6):1056–1073, 2007.
- [2] BÉLA BOLLOBÁS. [The chromatic number of random graphs](#). *Combinatorica*, 8(1):49–55, 1988.
- [3] BÉLA BOLLOBÁS, PAUL A. CATLIN, AND PAUL ERDŐS. [Hadwiger’s conjecture is true for almost every graph](#). *European J. Combin.*, 1(3):195–199, 1980.
- [4] MARIA CHUDNOVSKY AND PAUL SEYMOUR. [Packing seagulls](#). *Combinatorica*, 32(3):251–282, 2012.
- [5] MICHELLE DELCOURT AND LUKE POSTLE. [Reducing linear Hadwiger’s conjecture to coloring small graphs](#). 2021, arXiv:2108.01633.
- [6] GABRIEL A. DIRAC. [A property of 4-chromatic graphs and some remarks on critical graphs](#). *J. London Math. Soc.*, 27:85–92, 1952.
- [7] PIERRE DUCHET AND HENRI MEYNIEL. [On Hadwiger’s number and the stability number](#). *Annals of Discrete Mathematics*, 13:71–73, 1982.
- [8] PAUL ERDŐS AND LÁSZLÓ LOVÁSZ. [Problems and results on 3-chromatic hypergraphs and some related questions](#). In *Infinite and Finite Sets*, vol. 10 of *Colloq. Math. Soc. János Bolyai*, pp. 609–627. North-Holland, 1975.
- [9] MARTIN GROHE AND DÁNIEL MARX. [Structure theorem and isomorphism test for graphs with excluded topological subgraphs](#). *SIAM J. Comput.*, 44(1):114–159, 2015.
- [10] HUGO HADWIGER. [Über eine Klassifikation der Streckenkomplexe](#). *Vierteljahrsschr. Naturforsch. Ges. Zürich*, 88:133–142, 1943.
- [11] DANIEL J. HARVEY AND DAVID R. WOOD. [Average degree conditions forcing a minor](#). *Electron. J. Combin.*, 23(1):#P1.42, 2016.
- [12] ANNIKA HECKEL. [The chromatic number of dense random graphs](#). *Random Structures Algorithms*, 53(1):140–182, 2018.
- [13] ALEXANDR V. KOSTOCHKA. [The minimum Hadwiger number for graphs with a given mean degree of vertices](#). *Metody Diskret. Analiz.*, 38:37–58, 1982.
- [14] ALEXANDR V. KOSTOCHKA. [Lower bound of the Hadwiger number of graphs by their average degree](#). *Combinatorica*, 4(4):307–316, 1984.
- [15] ALEXANDR V. KOSTOCHKA AND NOAH PRINCE. [On  \$K\_{s,t}\$ -minors in graphs with given average degree](#). *Discrete Math.*, 308(19):4435–4445, 2008.
- [16] WOLFGANG MADER. [Homomorphieeigenschaften und mittlere Kantendichte von Graphen](#). *Math. Ann.*, 174:265–268, 1967.
- [17] DAVID MATULA AND LUDĚK KUČERA. [An expose-and-merge algorithm and the chromatic number of a random graph](#). In *Random graphs ’87 (Poznań, 1987)*, pp. 175–187. Wiley, 1990.
- [18] SERGEY NORIN AND PAUL SEYMOUR. [Dense minors of graphs with independence number two](#). 2022, arXiv:2206.00186.
- [19] MICHAEL D. PLUMMER, MICHAEL STIEBITZ, AND BJARNE TOFT. [On a special case of Hadwiger’s conjecture](#). *Discuss. Math. Graph Theory*, 23(2):333–363, 2003.
- [20] BRUCE A. REED AND PAUL SEYMOUR. [Fractional colouring and Hadwiger’s conjecture](#). *J. Combin. Theory Ser. B*, 74(2):147–152, 1998.
- [21] NEIL ROBERTSON AND PAUL SEYMOUR. [Graph minors. XVI. Excluding a non-planar graph](#). *J. Combin. Theory Ser. B*, 89(1):43–76, 2003.
- [22] PAUL SEYMOUR. [Hadwiger’s conjecture](#). In JOHN FORBES NASH JR. AND MICHAEL TH. RASSIAS, eds., *Open Problems in Mathematics*, pp. 417–437. Springer, 2016.
- [23] ANDREW THOMASON. [An extremal function for contractions of graphs](#). *Math. Proc. Cambridge Philos. Soc.*, 95(2):261–265, 1984.
- [24] ANDREW THOMASON. [The extremal function for complete minors](#). *J. Combin. Theory Ser. B*, 81(2):318–338, 2001.
- [25] WIKIPEDIA. [Coupon collector’s problem](#). 2024.
- [26] WIKIPEDIA. [Trivially perfect graph](#). 2024.