Dunkl symmetric coherent pairs of measures. An application to Fourier Dunkl-Sobolev expansions.

Mabrouk Sghaier¹

Higher Institute of Computer Medenine, Medenine - 4119, Tunisia. Francisco Marcellán, ²

Department of Mathematics, Universidad Carlos III de Madrid, 28911-Leganés, Spain Sabrine Hamdi³

Faculty of Sciences of Gabes, Laboratory of Mathematics and Applications (LR17ES11), 6072 Gabes, Tunisia.

Abstract Let \mathcal{T}_{μ} be the Dunkl operator. A pair of symmetric measures (u, v) supported on a symmetric subset of the real line is said to be a symmetric Dunkl-coherent pair if the corresponding sequences of monic orthogonal polynomials $\{P_n\}_{n\geq 0}$ and $\{R_n\}_{n\geq 0}$ (resp.) satisfy

$$
R_n(x) = \frac{\mathcal{T}_{\mu} P_{n+1}(x)}{\mu_{n+1}} - \sigma_{n-1} \frac{\mathcal{T}_{\mu} P_{n-1}(x)}{\mu_{n-1}}, \quad n \ge 2,
$$

where ${\lbrace \sigma_n \rbrace_{n\geq 1}}$ is a sequence of non-zero complex numbers and $\mu_{2n} = 2n, \mu_{2n-1} = 2n - 1 +$ $2\mu, n \geq 1$.

In this contribution we focus the attention on the sequence ${S_n^{(\lambda,\mu)}}_{n\geq 0}$ of monic orthogonal polynomials with respect to the Dunkl-Sobolev inner product

$$
\langle p, q \rangle_{s,\mu} = \langle u, pq \rangle + \lambda \langle v, \mathcal{T}_{\mu} p \mathcal{T}_{\mu} q \rangle, \quad \lambda > 0, \quad p, q \in \mathcal{P}.
$$

An algorithm is stated to compute the coefficients of the Fourier–Sobolev type expansions with respect to $\langle \,\cdot\,,\cdot\rangle_{s,\mu}$ for suitable smooth functions f such that $f \in \mathcal{W}_2^1(\mathbb{R},u,v,\mu) =$ $\{f; ||f||_u^2 + \lambda ||\mathcal{T}_{\mu}f||_v^2 < \infty\}$. Finally, two illustrative numerical examples are presented.

Keywords Orthogonal polynomials, symmetric linear functionals, Dunkl operator, generalized Gegenbauer polynomials, generalized Hermite polynomials, Dunkl coherent pairs of symmetric linear functionals, Dunkl-Sobolev orthogonal polynomials, Fourier expansions.

Mathematics Subject Classification Primary 42C05; Secondary 33C45, 42C10.

1 Introduction

Let P be the linear space of polynomials with complex coefficients. Let u and v denote two symmetric measures supported on a symmetric subset of the real line and let ${P_n}_{n>0}$ and ${R_n}_{n\geq 0}$ be the corresponding sequences of monic orthogonal polynomials (MOPS in short). The pair (u, v) is said to be a symmetric \mathcal{T}_{μ} -coherent pair if there exists a sequence

of non-zero complex numbers $\{\sigma_n\}_{n\geq 1}$ such that

$$
R_n(x) = \frac{\mathcal{T}_{\mu} P_{n+1}(x)}{\mu_{n+1}} - \sigma_{n-1} \frac{\mathcal{T}_{\mu} P_{n-1}(x)}{\mu_{n-1}}, \quad n \ge 2.
$$

Here \mathcal{T}_{μ} is the Dunkl operator introduced in [\[11\]](#page-20-0) that is defined as

$$
\mathcal{T}_{\mu}(f) = f' + 2\mu \mathcal{H}_{-1}f, \quad (\mathcal{H}_{-1}f)(x) = \frac{f(x) - f(-x)}{2x}, \quad f \in \mathcal{P}, \ \mu \neq -n - \frac{1}{2},
$$

and $\mu_{2n} = 2n, \mu_{2n-1} = 2n - 1 + 2\mu, n \ge 1$.

Notice that in a more general framework the concept of symmetric \mathcal{T}_{μ} -coherent pair of linear forms is introduced in [\[23\]](#page-20-1), where a description of them is stated.

In this paper, we consider the sequence ${S_n^{(\lambda,\mu)}}_{n\geq 0}$ of monic Dunkl-Sobolev polynomials orthogonal with respect to a special Sobolev inner product $(\mathcal{T}_{\mu}$ -Sobolev inner product)

$$
\langle p, q \rangle_{s,\mu} = \langle u, pq \rangle + \lambda \langle v, \mathcal{T}_{\mu} p \mathcal{T}_{\mu} q \rangle, \quad \lambda > 0, \quad p, q \in \mathcal{P}.
$$

Our results constitute an extension in the Dunkl framework of the paper [\[12\]](#page-20-2), where coherent and symmetrically coherent pairs of measures supported on the real line are introduced. Indeed, a pair of positive and symmetric measures (u, v) supported on a symmetric subset of the real line is said to be a symmetrically coherent pair if the corresponding MOPS ${P_n(x; u)}_{n>0}, {P_n(x; v)}_{n>0}$ satisfy

$$
P_n(x; v) = \frac{1}{n+1} \left[P'_{n+1}(x; u) - \varrho_{n-1} P'_{n-1}(x; u) \right], \quad \varrho_{n-1} \neq 0, \quad n \ge 2.
$$

The pairs of symmetrically coherent positive measures were completely determined in [\[20\]](#page-20-3), where it is proved that one of the measures must be classical i. e., either Gegenbauer or Hermite, and the other one is a linear spectral transform (see $[24]$). The motivations in $[12]$ for introducing these coherent pairs of measures are their applications in Fourier analysis in terms of sequences of orthogonal polynomials with respect to the Sobolev inner products associated with the vector of measures (u, v) . In the Gegenbauer case, analytic properties of Sobolev-Gegenbauer polynomials were analyzed in [\[3,](#page-19-0) [7,](#page-20-4) [14,](#page-20-5) [19\]](#page-20-6). In the Hermite case, analytic properties of Sobolev-Hermite polynomials were analyzed in [\[4,](#page-19-1) [8,](#page-20-7) [16\]](#page-20-8). An exhaustive information can be found in the survey paper [\[17\]](#page-20-9).

When we deal with symmetric \mathcal{T}_{μ} -coherent pairs of measures, one of our main purposes is the effective computation of the Fourier coefficients in the expansions of functions f living in appropriate function spaces

$$
f(x) \sim \sum_{n=0}^{\infty} \frac{f_n^{(\lambda,\mu)}}{s_n^{(\lambda,\mu)}} S_n^{(\lambda,\mu)}(x),
$$

in terms of the polynomials $\{S_n^{(\lambda,\mu)}\}_{n\geq 0}$ orthogonal with respect to the corresponding Sobolev inner product. This kind of Fourier expansions will be called Dunkl Fourier-Sobolev expansions. Here $s_n^{(\lambda,\mu)} = \langle S_n^{(\lambda,\mu)}, S_n^{(\lambda,\mu)} \rangle_{s,\mu}$ and $f_n^{(\lambda,\mu)} \leq f, S_n^{(\lambda,\mu)} \rangle_{s,\mu}$.

The structure of the paper is as follows. In Section 2, we recall some basic definitions and results on the theory of orthogonal polynomials and the definition of the Dunkl operator, which will be used in the sequel. In Section 3 we assume u (resp. v) is a \mathcal{T}_{μ} -classical

symmetric measure, i. e., the family $\{P_n\}_{n>0}$ (resp. $\{R_n\}_{n>0}$) is either the generalized Hermite polynomial sequence or the generalized Gegenbauer polynomial sequence. The corresponding companion measure is also obtained in terms of either a quadratic Chistoffel or a quadratic Geronimus transformation. In Section 4, we consider Dunkl-Sobolev orthogonal polynomials arising from a symmetric \mathcal{T}_{μ} -coherent pair of measures. In Section 5 an algorithm to compute Fourier Dunkl-Sobolev coefficients is described. Finally, in Section 6, a numerical example is discussed.

2 Preliminaries

Let P be the linear space of polynomials with complex coefficients and let P' be its algebraic dual space. We denote by $\lt u, f >$ the duality bracket for $u \in \mathcal{P}'$ and $f \in \mathcal{P}$. In particular, we denote by $(u)_n := , n \geq 0$, the moments of u. For a form u, any $a \in \mathbb{C}^*, c \in \mathbb{C}$ and a polynomial g, let $Du = u'$, gu, $h_a u$, δ_c and $(x-c)^{-1}u$ be the forms defined, respectively, by

$$
\langle u', f \rangle := -\langle u, f' \rangle, \langle gu, f \rangle := \langle u, gf \rangle, \langle h_a u, f(x) \rangle := \langle u, h_a f(x) \rangle = \langle u, f(ax) \rangle,
$$

$$
\langle \delta_c, f \rangle := f(c), \langle (x - c)^{-1} u, f(x) \rangle := \langle u, \theta_c f(x) \rangle = \langle u, \frac{f(x) - f(c)}{x - c} \rangle, \ f \in \mathcal{P}.
$$

For c, $d \in \mathbb{C}$, $c \neq d$, $u \in \mathcal{P}'$ and $f \in \mathcal{P}$, it is s straightforward to prove that

$$
(x-c)^{-1}(x-c)u = u - (u)_0 \delta_c.
$$
\n(2.1)

Let us recall some basic background on orthogonal polynomials and linear forms (see [\[6\]](#page-19-2)). A form u is called regular if there exists a sequence of polynomials $\{P_n\}_{n>0}$, deg $P_n \leq n$, such that

$$
\langle u, P_n P_m \rangle = p_n \delta_{nm}, \quad p_n \neq 0, \quad n \geq 0,
$$

where δ_{nm} is the Kronecker symbol. Then, deg $P_n = n, n \geq 0$, and we can always assume each P_n is monic. In such a case, the sequence $\{P_n\}_{n\geq 0}$ is unique. It is said to be the sequence of monic orthogonal polynomials (MOPS) with respect to u .

There exist a complex sequence $\{\beta_n\}_{n\geq 0}$ and a non zero complex sequence $\{\gamma_n\}_{n\geq 1}$ such that the MOPS $\{P_n\}_{n\geq 0}$ fulfils the following three-term recurrence relation (TTRR, in short)

$$
P_0(x) = 1, \quad P_1(x) = x - \beta_0,
$$

\n
$$
P_{n+2}(x) = (x - \beta_{n+1})P_{n+1}(x) - \gamma_{n+1}P_n(x), \quad n \ge 0.
$$
\n(2.2)

By convention, we set $\gamma_0 = (u)_0$. The form u is said to be normalized if $(u)_0 = 1$. In this paper, we assume that any form will be normalized.

A form u is said to be symmetric if and only if $(u)_{2n+1} = 0$, $n \ge 0$, or, equivalently, in [\(2.2\)](#page-2-0) $\beta_n = 0, n \geq 0.$

Proposition 2.1. ([\[6\]](#page-19-2)) Let $\{P_n\}_{n>0}$ be the MOPS with respect to a regular form u. Then the following statements are equivalent.

- (i) u is symmetric.
- (ii) $P_n(-x) = (-1)^n P_n(x), n \ge 0.$

(iii) There exist MOPS ${A_n}_{n>0}$, ${B_n}_{n>0}$, orthogonal with respect to linear forms w, xw, respectively, such that $P_{2n}(x) = A_n(x^2), P_{2n+1}(x) = xB_n(x^2), n \ge 0$. On the other hand, w is the linear form with moments $\langle w, x^n \rangle = \langle u, x^{2n} \rangle$.

Let us introduce the Dunkl operator

$$
\mathcal{T}_{\mu}(f) = f' + 2\mu \mathcal{H}_{-1}f, \quad (\mathcal{H}_{-1}f)(x) = \frac{f(x) - f(-x)}{2x}, \quad f \in \mathcal{P}, \ \mu \neq -n - \frac{1}{2}.
$$

This operator was introduced and studied for first time by Dunkl [\[11\]](#page-20-0). Notice that \mathcal{T}_0 is the derivative operator D. The transposed $t_{\mathcal{T}_{\mu}}$ of \mathcal{T}_{μ} is $t_{\mathcal{T}_{\mu}} = -D - H_{-1} = -\mathcal{T}_{\mu}$. Thus, we have

 $\langle \mathcal{T}_{\mu} u, f \rangle = - \langle u, \mathcal{T}_{\mu} f \rangle, \quad u \in \mathcal{P}', \quad f \in \mathcal{P}.$

In particular, this yields $\langle \mathcal{T}_{\mu} u, x^n \rangle = -\mu_n \langle u, x^{n-1} \rangle = -\mu_n(u)_{n-1}, n \geq 0$, where $(u)_{-1} = 0$ and $\mu_n = n + \mu(1 - (-1)^n)$, $n \ge 0$. It is easy to see that

$$
\mathcal{T}_{\mu}(fu) = f\mathcal{T}_{\mu}u + f'u + 2\mu(H_{-1}f)(h_{-1}u), \ \ f \in \mathcal{P}, \ \ u \in \mathbb{P}'. \tag{2.3}
$$

Remark 2.1. Notice that if u is a symmetric form, then (2.3) becomes

$$
\mathcal{T}_{\mu}(fu) = f\mathcal{T}_{\mu}u + \mathcal{T}_{\mu}fu, \quad f \in \mathcal{P}.
$$
\n(2.4)

Now, let consider a sequence of monic polynomials $\{P_n\}_{n>0}$ and let denote by

$$
P_n^{[1]}(x,\mu) = \frac{\mathcal{T}_\mu P_{n+1}(x)}{\mu_{n+1}}, \quad \mu \neq -n - \frac{1}{2}
$$
\n(2.5)

a new sequence of monic polynomials.

Definition 2.1. An MOPS ${P_n}_{n\geq 0}$ is said to be \mathcal{T}_{μ} -classical if ${P_n^{[1]}(.,\mu)}_{n\geq 0}$ is also an MOPS. In this case, the associated form u is called \mathcal{T}_{μ} -classical form.

Theorem 2.1. ([2]) The only \mathcal{T}_{μ} -classical symmetric MOPS are

- The generalized Hermite polynomials $\{\mathrm{H}_n^{\mu}\}_{n\geq 0}$ for $\mu \neq -n-\frac{1}{2}$ $\frac{1}{2}$, $n \geq 1$.
- The generalized Gegenbauer polynomials $\{G_n^{(\alpha,\mu-\frac{1}{2})}\}_{n\geq 0}$ for $\alpha\neq -n$, $\alpha+\mu\neq -n+\frac{1}{2}$ $\frac{1}{2}$, $\mu \neq -n + \frac{1}{2}$ $\frac{1}{2}$, $n \geq 1$.

Remark 2.2. Generalized Hermite polynomials were introduced in [\[5\]](#page-19-4). Generalized Gegenbauer polynomials were studied in $\vert 1 \vert$. For more information about these two families the reader can check [\[6\]](#page-19-2)

Proposition 2.2. 1. The MOPS $\{G_n^{(\alpha,\mu-\frac{1}{2})}\}_{n\geq 0}$ satisfies the TTRR with

$$
\hat{\beta}_n = 0, \quad \gamma_{n+1} = \frac{(n+1+\varrho_n)(n+1+2\alpha+\varrho_n)}{4(n+\alpha+\mu+\frac{1}{2})(n+\alpha+\mu+\frac{3}{2})}, \quad n \ge 0,
$$

$$
\varrho_n = 2\mu \frac{1+(-1)^n}{2}, \quad n \ge 0.
$$

2. The MOPS ${H_n^{\mu}}_{n\geq 0}$ satisfies the TTRR with

$$
\hat{\beta}_n = 0
$$
, $\gamma_{n+1} = \frac{1}{2}(n+1+\mu(1+(-1)^n)), \quad n \ge 0.$

In [\[22\]](#page-20-10) a characterization of \mathcal{T}_{μ} -classical symmetric forms is given in terms of a distributional equation of Pearson type as

Theorem 2.2. A symmetric form u is said to be a \mathcal{T}_{μ} -classical form if and only if u is regular and there exist polynomials Φ and Ψ , $deg(\Phi) \leq 2$ and $deg(\Psi) = 1$, such that

$$
\mathcal{T}_{\mu}(\Phi u) = \Psi u,\tag{2.6}
$$

$$
\Psi'(0) - \frac{1}{2}\Phi''(0)\mu_n \neq 0, \quad n \ge 0.
$$
\n(2.7)

The corresponding MOPS $\{P_n\}_{n>0}$ is called \mathcal{T}_{μ} -classical.

Remark 2.3. Notice that this is the T_{μ} analogue of the Pearson equation for the classical linear forms with respect to the derivative operator (see $[13]$).

We state now the definition of the quasi-orthogonality.

Definition 2.2. Let $u \in \mathcal{P}'$ and s be a nonnegative integer number. A MPS $\{P_n\}_{n\geq 0}$ is said to be quasi-orthogonal of order s with respect to u if

$$
\langle u, P_m P_n \rangle = 0, \quad 0 \le m \le n - s - 1, \quad n \ge s + 1,
$$

$$
\exists r \ge s; \langle u, P_{r-s} P_r \rangle \neq 0.
$$
 (2.8)

If $\langle u, P_{r-s}P_r \rangle \neq 0$ for any $r \geq s$, then $\{P_n\}_{n \geq 0}$ is said to be strictly quasi-orthogonal of order s with respect to u.

3 Symmetric \mathcal{T}_{μ} -Coherent Pairs

Symmetric \mathcal{T}_{μ} -coherent pairs of linear forms have been introduced by M. Sghaier and S. Hamdi in [\[23\]](#page-20-1) as

Definition 3.1. Let u and v denote two symmetric forms and $\{P_n\}_{n\geq 0}$ and $\{R_n\}_{n\geq 0}$ will denote their respective SMOP. Assume that there exists a sequence of non-zero complex numbers $\{\sigma_n\}_{n\geq 1}$ such that

$$
R_n(x) = \frac{\mathcal{T}_{\mu} P_{n+1}(x)}{\mu_{n+1}} - \sigma_{n-1} \frac{\mathcal{T}_{\mu} P_{n-1}(x)}{\mu_{n-1}}, \quad n \ge 2,
$$
\n(3.1)

holds. Then the pair (u, v) is said to be a symmetric \mathcal{T}_{μ} -coherent pair.

In [\[23\]](#page-20-1) the symmetric \mathcal{T}_{μ} -coherent pairs of linear forms are completely described and determined. Let us recall the important result.

Theorem 3.1. If (u, v) denote a symmetric \mathcal{T}_{μ} -coherent pair of linear forms, then one of them is a symmetric Dunkl-classical form, i. e., generalized Hermite or generalized Gegenbauer linear forms.

In the sequel we will assume that (u, v) is a symmetric \mathcal{T}_{μ} -coherent pair of positive Borel measures. We will analyze two cases

3.1 u is symmetric \mathcal{T}_{μ} -classical

Let u be a symmetric \mathcal{T}_{μ} -classical measure, i. e., $\{P_n\}_{n\geq 0}$ is the sequence of generalized Hermite or generalized Gegenbauer polynomials. So, if

$$
T_n(x) = \frac{\mathcal{T}_{\mu} P_{n+1}(x)}{\mu_{n+1}}, \ \ n \ge 0,
$$

and taking into account that $\{T_n\}_{n\geq 0}$ is again a \mathcal{T}_{μ} -classical sequence of polynomials, of the same type, orthogonal with respect to a symmetric positive measure w , then (3.1) becomes

$$
R_n(x) = T_n(x) + \varepsilon_{n-2} T_{n-2}(x), \quad n \ge 2,
$$
\n(3.2)

with $\varepsilon_{n-2} = -\sigma_{n-1}$ for all $n \geq 2$. Furthermore, the coefficients of the TTRR for the sequence ${\{\mathcal{T}_n\}}_{n\geq 0}, \gamma_n$, are known.

Proposition 3.1. ${R_n}_{n>0}$ is an MOPS if and only if

(i) $\gamma_n + \varepsilon_{n-2} - \varepsilon_{n-1} > 0$, $\forall n \ge 1$. By convention, $\varepsilon_{-1} = 0$.

$$
(ii) \varepsilon_n \gamma_n = (\gamma_{n+2} + \varepsilon_n - \varepsilon_{n+1}) \varepsilon_{n-1}, n \ge 1.
$$

Proof. Since

$$
R_n(x) = xT_n(x) + \varepsilon_{n-2}T_{n-2}(x), \quad n \ge 1,
$$
\n(3.3)

from the TTRR that the MOPS ${R_n}_{n>0}$ satisfies, you get

$$
xR_n(x) = R_{n+1}(x) + (\gamma_n + \varepsilon_{n-2} - \varepsilon_{n-1})R_{n-1}(x) + (\varepsilon_{n-2}\gamma_{n-2} - (\gamma_n + \varepsilon_{n-2} - \varepsilon_{n-1})\varepsilon_{n-3}))T_{n-3}(x), \quad n \ge 1.
$$
\n(3.4)

Thus, ${R_n}_{n>0}$ satisfies a TTRR if and only if (i) and (ii) hold.

In the next result, we compute the coefficients $\{\tilde{\gamma}_n\}_{n\geq 1}$ of the TTRR for $\{R_n\}_{n\geq 0}$ in the terms of γ_n and ε_n .

Corollary 3.1. The following relations hold:

(i) $\tilde{\gamma}_n = \gamma_n + \varepsilon_{n-2} - \varepsilon_{n-1}, \forall n \geq 1$. By convention, $\varepsilon_{-1} = 0$. (ii) $\varepsilon_n \gamma_n = \varepsilon_{n-1} \tilde{\gamma}_{n+2}, n > 1.$

Proof. Consider first the TTRR for the sequence ${R_n}_{n>0}$:

$$
\begin{cases} R_0(x) = 1, & R_1(x) = x, \\ R_{n+1}(x) = xR_n(x) - \tilde{\gamma}_n R_{n-1}(x), & n \ge 1. \end{cases}
$$
 (3.5)

Here $\tilde{\gamma}_n > 0, n \geq 1$. Substituting (3.2) into (3.5) , we get

$$
\begin{cases}\nT_{n+1}(x) + \varepsilon_{n-1}T_{n-1}(x) = x(T_n(x) + \varepsilon_{n-2}T_{n-2}(x)) - \tilde{\gamma}_n(T_{n-1}(x) + \varepsilon_{n-3}T_{n-3}(x)), & \forall n \ge 3, \\
T_3(x) + \varepsilon_1T_1(x) = x(T_2(x) + \varepsilon_0) - \tilde{\gamma}_2T_1(x), \\
T_2(x) + \varepsilon_0 = xT_1(x) - \tilde{\gamma}_1, \\
T_1(x) = x.\n\end{cases}
$$

(3.6)

Consider now the TTRR for the sequence $\{\mathrm{T}_n\}_{n\geq 0}$:

$$
\begin{cases} T_0(x) = 1, & T_1(x) = x, \\ xT_n(x) = T_{n+1}(x) + \gamma_n T_{n-1}(x), & n \ge 1. \end{cases}
$$
 (3.7)

Here $\gamma_n > 0, n \geq 1$.

Using [\(3.7\)](#page-6-0) and expanding $xT_i(x)$ in [\(3.6\)](#page-5-2) as a linear combination of the polynomials ${T_n}_{n\geq 0}$, we obtain

$$
\begin{cases}\nT_{n+1}(x) + \varepsilon_{n-1}T_{n-1}(x) = T_{n+1}(x) + \gamma_n T_{n-1}(x) + \varepsilon_{n-2}T_{n-1}(x) + \varepsilon_{n-2}\gamma_{n-2}T_{n-3}(x) \\
- \tilde{\gamma}_n (T_{n-1}(x) + \varepsilon_{n-3}T_{n-3}(x)), \ \forall \ n \ge 3, \\
T_3(x) + \varepsilon_1 T_1(x) = T_3(x) + \gamma_2 T_1(x) + \varepsilon_0 T_1(x) - \tilde{\gamma}_2 T_1(x), \\
T_2(x) + \varepsilon_0 = T_2(x) + \gamma_1 - \tilde{\gamma}_1.\n\end{cases}
$$
\n(3.8)

Since $\{T_n\}_{n\geq 0}$ is a basis in P , after identification of coefficients in [\(3.8\)](#page-6-1) we find

$$
\varepsilon_{n-1} = \gamma_n + \varepsilon_{n-2} - \tilde{\gamma}_n, \quad n \ge 3,
$$
\n(3.9)

$$
\varepsilon_1 = \gamma_2 + \varepsilon_0 - \tilde{\gamma}_2,\tag{3.10}
$$

$$
\varepsilon_0 = \gamma_1 - \tilde{\gamma}_1,\tag{3.11}
$$

$$
\varepsilon_{n-2}\gamma_{n-2} - \varepsilon_{n-3}\tilde{\gamma}_n, \quad n \ge 3. \tag{3.12}
$$

From $(3.9)-(3.11)$ $(3.9)-(3.11)$, we get

$$
\varepsilon_{n-1} = \gamma_n + \varepsilon_{n-2} - \tilde{\gamma}_n, \quad n \ge 1, \quad \varepsilon_{-1} = 0.
$$

Thus (i) holds. (ii) can be deduced in a straightforward way from (3.12) .

Proposition [3.1](#page-5-3) requires the study of properties of the sequence $\{\varepsilon_n\}_{n\geq 0}$. Next, we can prove that this sequence is a solution of a non-linear difference equation.

Proposition 3.2. The sequence $\{\varepsilon_n\}_{n\geq 0}$ satisfies a quadratic difference equation

$$
\varepsilon_{n+1} + \frac{\gamma_n \gamma_{n+1}}{\varepsilon_{n-1}} = \gamma_{n+1} + \gamma_{n+2} + D, \quad n \ge 1,
$$
\n(3.13)

where

$$
D = (\varepsilon_1 - \gamma_2)(1 - \frac{\gamma_1}{\varepsilon_0}).
$$

Proof. From [3.1,](#page-5-3) we have

$$
\varepsilon_n \gamma_n = \varepsilon_{n-1}(\gamma_{n+2} + \varepsilon_n - \varepsilon_{n+1}), \quad n \ge 1.
$$

As a consequence,

$$
\varepsilon_{n+1} - \varepsilon_n = \gamma_{n+2} - \frac{\varepsilon_n}{\varepsilon_{n-1}} \gamma_n, \quad n \ge 1.
$$

We sum from $k = 1$ to $k = n$ and we get

$$
\varepsilon_{n+1} - \varepsilon_1 = \sum_{k=1}^n \gamma_{k+2} - \sum_{k=1}^n \frac{\varepsilon_k}{\varepsilon_{k-1}} \gamma_k, \quad n \ge 1.
$$
 (3.14)

 \blacksquare

On the other hand, again from Corollary [3.1,](#page-5-3) one can see that

$$
\frac{\varepsilon_{k-1}}{\varepsilon_{k-2}}\gamma_{k-1}=\gamma_{k+1}+\varepsilon_{k-1}-\varepsilon_k,\ \ k\geq 2.
$$

Thus

$$
\frac{\varepsilon_k}{\varepsilon_{k-1}} = \frac{\gamma_{k+1}}{\varepsilon_{k-1}} - \frac{\gamma_{k-1}}{\varepsilon_{k-2}} + 1, \quad k \ge 2.
$$

Substitution of the previous equation into [\(3.14\)](#page-6-5) yields

$$
\varepsilon_{n+1} - \varepsilon_1 = \sum_{k=1}^n \gamma_{k+2} - \frac{\varepsilon_1 \gamma_1}{\varepsilon_0} - \sum_{k=2}^n \left(\frac{\gamma_k \gamma_{k+1}}{\varepsilon_{k-1}} - \frac{\gamma_{k-1} \gamma_k}{\varepsilon_{k-2}} + \gamma_k \right)
$$

= $\gamma_{n+1} + \gamma_{n+2} - \gamma_2 - \frac{\gamma_n \gamma_{n+1}}{\varepsilon_{n-1}} + \frac{\gamma_1 \gamma_2}{\varepsilon_0} - \frac{\varepsilon_1 \gamma_1}{\varepsilon_0}, \quad n \ge 2.$

We denote by D the term which does not depend on n in the last relation, then

$$
D = \varepsilon_1 - \frac{\varepsilon_1 \gamma_1}{\varepsilon_0} + \frac{\gamma_1 \gamma_2}{\varepsilon_0} - \gamma_2 = (\varepsilon_1 - \gamma_2)(1 - \frac{\gamma_1}{\varepsilon_0}),
$$

and we obtain (3.13) .

Notice that we can split the above difference equation for even and odd subindices, respectively. Indeed, for $n = 2m$ [\(3.13\)](#page-6-6) reads

$$
\varepsilon_{2m+1} + \frac{\gamma_{2m}\gamma_{2m+1}}{\varepsilon_{2m-1}} = \gamma_{2m+1} + \gamma_{2m+2} + D, \quad m \ge 1.
$$
 (3.15)

If we denote $\varepsilon_{2m+1} = \frac{\rho_{m+1}}{\rho_m}$ $\frac{m+1}{\rho_m}, m \geq 0, \rho_0 = 1$, then the previous equation becomes

$$
\rho_{m+1} + \gamma_{2m}\gamma_{2m+1}\rho_{m-1} = (\gamma_{2m+1} + \gamma_{2m+2} + D)\rho_m, \quad m \ge 1.
$$
\n(3.16)

Here the initial conditions are $\rho_0 = 1, \rho_1 = \varepsilon_1 = \gamma D + \gamma - 2$, with $\gamma = (1 - \frac{\gamma_1}{\varepsilon_0})$ $\frac{\gamma_1}{\varepsilon_0}$)⁻¹.

Thus ρ_m is a polynomial of degree m in D.

For $n = 2m - 1$, [\(3.13\)](#page-6-6) reads

$$
\varepsilon_{2m} + \frac{\gamma_{2m-1}\gamma_{2m}}{\varepsilon_{2m-2}} = \gamma_{2m} + \gamma_{2m+1} + D, \quad m \ge 1.
$$
 (3.17)

If we denote $\varepsilon_{2m} = \frac{\tau_{m+1}}{\tau_m}$ $\frac{m+1}{\tau_m}, m \geq 0, \tau_0 = 1$, then the previous equation becomes

$$
\tau_{m+1} + \gamma_{2m-1} \gamma_{2m} \tau_{m-1} = (\gamma_{2m} + \gamma_{2m+1} + D) \tau_m, \quad m \ge 1.
$$
\n(3.18)

Here the initial conditions are $\tau_0 = 1, \tau_1 = \varepsilon_0$. In both cases, you have a TTRR.

Remark 3.1. Since the measure u is known, we can compute the values of the coefficients ε_n for $n \geq 2$, once the initial values ε_0 and ε_1 are fixed.

Remark 3.2. Notice that (3.2) means that the linear forms w, v associated with the MOPS ${\{\rm T}_n\}_{n\geq 0}, {\{\rm R}_n\}_{n\geq 0}$ are related by $M_2(x)v = w$, where M_2 is a positive quadratic even polynomial. In other words, v is a Geronimus transformation of w. When u is the generalized Hermite form, then $w = u$ and, as a consequence, $M_2(x)v = u$, with $M_2(x) = \lambda(x^2 + \zeta^2), \lambda > 0$. When u is the generalized Gegenbauer form, then $w = (1-x^2)u$ and $M_2(x) = \lambda(\zeta^2 - x^2), |\zeta| \ge$ $1, \lambda > 0$, or $M_2(x) = \lambda(x^2 + \zeta^2), \zeta \in \mathbb{R}, \lambda > 0$.

An alternative way to get the sequence $\{\varepsilon_n\}_{n\geq 0}$ is the following one.

Let consider $T_{2n}(x) = A_n(x^2), T_{2n+1}(x) = xB_n(x^2), R_{2n}(x) = C_n(x^2), R_{2n+1}(x) = xD_n(x^2), n \ge$ 0. Thus [\(3.2\)](#page-5-0) reads

$$
C_n(x) = A_n(x) + \varepsilon_{2n-2}A_{n-1}(x), \quad n \ge 1, D_n(x) = B_n(x) + \varepsilon_{2n-1}B_{n-1}(x), \quad n \ge 1. \tag{3.19}
$$

Taking into account that according to the Christoffel formula (see [\[5\]](#page-19-4)) you get

$$
xB_n(x) = A_{n+1}(x) + a_n A_n(x), xD_n(x) = C_{n+1}(x) + b_n C_n(x), \quad n \ge 0,
$$
 (3.20)

then

$$
A_{n+1}(x) + a_n A_n(x) + \varepsilon_{2n-1}(A_n(x) + a_{n-1} A_{n-1}(x)) = A_{n+1}(x) + \varepsilon_{2n} A_n(x) + b_n (A_n(x) + \varepsilon_{2n-2} A_{n-1}(x)), \quad n \ge 1.
$$
\n(3.21)

As a consequence, identifying coefficients,

$$
a_n + \varepsilon_{2n-1} = \varepsilon_{2n} + b_n, \quad n \ge 1, \varepsilon_{2n-1} a_{n-1} = b_n \varepsilon_{2n-2}, \quad n \ge 1.
$$
 (3.22)

This is a coupled system for the sequence $\{\varepsilon_n\}_{n>0}$.

On the other hand, since $C_n(0) = A_n(0) + \varepsilon_{2n-2}A_{n-1}(0)$, $n \ge 1$, $A_{n+1}(0) + a_nA_n(0) =$ 0, $n \geq 0$, we immediately get

$$
a_n = -\frac{A_{n+1}(0)}{A_n(0)}, \quad b_n = -\frac{C_{n+1}(0)}{C_n(0)} = -\frac{A_{n+1}(0) + \varepsilon_{2n}A_n(0)}{A_n(0) + \varepsilon_{2n-2}A_{n-1}(0)}.
$$
(3.23)

Initial conditions for ε_0 , ε_1 are needed in order to generate the full sequence, Indeed,

$$
\varepsilon_{2n} = a_n + \varepsilon_{2n-1} - \frac{\varepsilon_{2n-1} a_{n-1}}{\varepsilon_{2n-2}}.
$$
\n(3.24)

As a conclusion,

$$
\varepsilon_{2n} = a_n + \varepsilon_{2n-1} (1 - \frac{a_{n-1}}{\varepsilon_{2n-2}}), \quad n \ge 1.
$$
\n(3.25)

Remark 3.3. Notice that the MOPS $\{C_n\}_{n\geq 0}$ is a canonical Geronimus transformation of the MOPS $\{A_n\}_{n\geq 0}$ as well as the MOPS $\{D_n\}_{n\geq 0}$ is a canonical Geronimus transformation of the MOPS ${B_n}_{n>0}$, see [\[18\]](#page-20-12).

3.2 v is symmetric \mathcal{T}_{μ} -classical

Let v be a symmetric \mathcal{T}_{μ} -classical positive measure, i. e., $\{R_n\}_{n\geq 0}$ is the sequence of generalized Hermite or generalized Gegenbauer polynomials. So, if

$$
R_n(x) = \frac{\mathcal{T}_{\mu} Q_{n+1}(x)}{\mu_{n+1}}, \quad n \ge 0,
$$

and taking into account that $\{Q_n\}_{n>0}$ is again a \mathcal{T}_{μ} -classical sequence of polynomials, orthogonal with respect to a symmetric measure \hat{v} , then (3.1) becomes

$$
Q_{n+1}(x) = P_{n+1}(x) - \tau_n P_{n-1}(x) - a_n, \quad n \ge 1,
$$
\n(3.26)

with $\tau_n = \sigma_{n-1} \frac{\mu_{n+1}}{\mu_{n-1}}$ $\frac{\mu_{n+1}}{\mu_{n-1}}$, for all $n \geq 2$, and $\{a_n\}_{n\geq 1}$ is a sequence of real numbers. Furthermore, the coefficients $\hat{\gamma}_n$ of the TTRR for the sequence $\{Q_n\}_{n\geq 0}$ are known.

Notice that $a_{2n} = 0, n \ge 0$, as a straightforward consequence of the fact that P_{2n-1} and Q_{2n-1} are odd functions.

Remark 3.4. If v is the generalized Hermite measure, then $\hat{v} = v$. If v is the generalized Gegenbauer measure, then $(1 - x^2)\hat{v} = v$.

Let consider $Q_{2n}(x) = C_n(x^2), Q_{2n+1}(x) = xD_n(x^2)$. Since $\{Q_n\}_{n\geq 0}$ is an MOPS, then ${C_n}_{n\geq 0}$ and ${D_n}_{n\geq 0}$ are MOPS and $xD_n(x) = C_{n+1}(x) + b_nC_n(x)$, $n \geq 0$. On the other hand, $P_{2n}(x) = A_n(x^2), P_{2n+1} = xB_n(x^2)$. Notice that ${P_n}_{n\geq 0}$ is an MOPS if and only if ${A_n}_{n>0}$ and ${B_n}_{n>0}$ are MOPS. Thus for $n = 2m$ [\(3.26\)](#page-9-0) reads

$$
D_m(x) = B_m(x) - \tau_{2m} B_{m-1}(x), \quad m \ge 1.
$$
\n(3.27)

Assuming $\tau_2 \neq 0$, if $xD_m(x) = D_{m+1}(x) + \beta_m D_m(x) + \alpha_m D_{m-1}(x)$, $m \geq 1$, and ac-cording to item 2, Theorem 1 in [\[15\]](#page-20-13), then ${B_m}_{m\geq 0}$ is an MOPS if and only if τ_{2m} = $\alpha_m \frac{D_{m-1}(a)}{D_m(a)}$ $\frac{D_{m-1}(a)}{D_m(a)}, \quad D_m(a) \neq 0, n \geq 1$, where $a = \beta_0 + \frac{\alpha_1}{\tau_2}$ $\frac{\alpha_1}{\tau_2}.$

If ${D_n}_{n\geq 0}$ is the MOPS associated with the linear form ω , then ${B_m}_{m\geq 0}$ is an MOPS with respect to the linear form $\tilde{\omega} = (x - a)\omega$, see [\[15\]](#page-20-13).

On the other hand, for $n = 2m - 1$ [\(3.26\)](#page-9-0) reads

$$
C_m(x) = A_m(x) - \tau_{2m-1}A_{m-1}(x) - a_{2m-1}, \quad m \ge 1.
$$
\n(3.28)

Assuming $\tau_1 + a_1 \neq 0$, if $xC_m(x) = C_{m+1}(x) + \tilde{\beta}_m C_m(x) + \tilde{\alpha}_m C_{m-1}(x)$, $m \geq 1$, and accord-ing to item 2, Theorem 1 in [\[15\]](#page-20-13), then $\{A_m\}_{m>0}$ is an MOPS if and only if $a_{2m-1} = 0$, $m \ge 1$, and $\tau_{2m-1} = \tilde{\alpha}_m \frac{C_{m-1}(b)}{C_m(b)}$ $C_{m}(b), \quad C_{m}(b) \neq 0, \quad n \geq 1$, where $b = \tilde{\beta}_{0} + \frac{\tilde{\alpha}_{1}}{\tau_{1} + \alpha_{2}}$ $\frac{\alpha_1}{\tau_1+a_1}$.

If ${C_n}_{n>0}$ is the MOPS associated with the linear form ν , then ${A_m}_{m>0}$ is an MOPS with respect to the linear form $\tilde{\nu} = (x - b)\nu$, see item 2, Theorem 1 in [\[15\]](#page-20-13).

Notice that $\tilde{\omega} = x\tilde{\nu}$, and $\omega = x\nu$. This means that $(x - a)x\nu = x\tilde{\nu}$. But $\tilde{\nu} = (x - b)\nu$. As a consequence, $x(x-a)\nu = x(x-b)\nu$. Thus $x(b-a)\nu = 0$. In other words, $(b-a)\nu = M\delta_0$. Taking into account ν is regular, then $a = b$.

Remark 3.5. If $a_1 = -\tau_1$, then $\{A_m\}_{m\geq 0}$ is an MOPS if and only if $a_{2m-1} = 0$ and $\tau_{2m-1}=0$, $m\geq 2$, a contradiction, see Theorem 1, item 1 in [\[15\]](#page-20-13).

Taking into account that $\hat{v}_{2n} = \nu_n$ and $u_{2n} = \tilde{\nu}_n$, as a summary of the previous analysis, we get

Proposition 3.3. If [\(3.26\)](#page-9-0) holds, then $a_n = 0, n \ge 2$. Moreover, $\langle \tilde{\nu}, x^n \rangle = \langle (x - b)\nu, x^n \rangle$. This means that $u_{2n} = \hat{v}_{2n+2} - b\hat{v}_{2n}$. In other words, $u = (x^2 - b)\hat{v}$, i. e., u is a quadratic Christoffel transformation of the measure \hat{v} . In the generalized Hermite case, you have $(x^2 +$ ζ^2) $v = u$, i. e., b must be a nonpositive real number. In the generalized Gegenbauer case, $(1-x^2)u = (x^2+b)v$ holds, i. e., either b is a positive real number or $(x^2-1)u = (x^2-b^2)v$, where $|b| > 1$.

The sequence $\{\tau_n\}_{n\geq 0}$ can be recursively obtained in the following way. According to the above notation [\(3.2\)](#page-5-0) reads

$$
C_{n+1}(x) = A_{n+1}(x) - \tau_{2n+1}A_n(x), n \ge 0, D_n(x) = B_n(x) - \tau_{2n}B_{n-1}(x), n \ge 1.
$$
 (3.29)

Taking into account that according to the Christoffel formula (see [\[5\]](#page-19-4)) you get

$$
xB_n(x) = A_{n+1}(x) + \hat{a}_n A_n(x), x D_n(x) = C_{n+1}(x) + \hat{b}_n C_n(x), n \ge 0,
$$
\n(3.30)

then

$$
A_{n+1}(x) + \hat{a}_n A_n(x) - \tau_{2n}(A_n(x) + \hat{a}_{n-1} A_{n-1}(x)) = A_{n+1}(x) - \tau_{2n+1} A_n(x) + \hat{b}_n (A_n(x) - \tau_{2n-1} A_{n-1}(x)), n \ge 1.
$$
\n(3.31)

As a consequence, identifying coefficients,

$$
\hat{a}_n - \tau_{2n} = -\tau_{2n+1} + \hat{b}_n, n \ge 1, \tau_{2n}\hat{a}_{n-1} = \hat{b}_n\tau_{2n-1}, n \ge 1.
$$
\n(3.32)

This is a coupled system for the sequence ${\tau_n}_{n\geq 0}$. Initial conditions for τ_1 are needed in order to generate the full sequence,

4 Dunkl-Sobolev Orthogonal Polynomials

Let (u, v) be a \mathcal{T}_{μ} -symmetric coherent pair of positive measures and let $\{P_n\}_{n>0}$ and $\{R_n\}_{n>0}$ be the corresponding MOPS, i. e., (3.1) holds. We consider the Dunkl-Sobolev $(\mathcal{T}_{\mu}$ -Sobolev) inner product as

$$
\langle p, q \rangle_{s,\mu} = \langle u, pq \rangle + \lambda \langle v, \mathcal{T}_{\mu} p \mathcal{T}_{\mu} q \rangle, \quad \lambda > 0, \quad p, q \in \mathcal{P}.
$$
 (4.1)

Let $\{S_n^{(\lambda,\mu)}\}_{n\geq 0}$ be the sequence of monic Dunkl-Sobolev polynomials orthogonal with respect to [\(4.1\)](#page-10-0). Notice that $P_k = S_k$ for $k = 0, 1, 2$.

Put $\langle S_n^{(\lambda,\mu)}, S_n^{(\lambda,\mu)} \rangle_{s,\mu} = s_n^{(\lambda,\mu)}, p_n = \langle u, P_n^2 \rangle \text{ and } r_n = \langle v, R_n^2 \rangle.$

The following theorem gives a fundamental algebraic property for \mathcal{T}_{μ} -symmetric coherent pairs of measures.

Theorem 4.1. Let (u, v) be a \mathcal{T}_u -symmetric coherent pair of positive measures and let ${P_n}_{n\geq 0}$ and ${R_n}_{n\geq 0}$ be the corresponding MOPS satisfying [\(3.1\)](#page-4-0). Then the Dunkl-Sobolev polynomials $\{S_n^{(\lambda,\mu)}\}_{n\geq 0}$ are related to $\{P_n\}_{n\geq 0}$ and $\{R_n\}_{n\geq 0}$ by

$$
P_{n+1}(x) + a_{n-2}P_{n-1}(x) = S_{n+1}^{(\lambda,\mu)}(x) + \eta_{n-2}^{\mu}(\lambda)S_{n-1}^{(\lambda,\mu)}(x), \quad n \ge 2,
$$
\n(4.2)

$$
\mu_{n+1} \mathcal{R}_n(x) = \mathcal{T}_{\mu} S_{n+1}^{(\lambda,\mu)}(x) + \eta_{n-2}^{\mu}(\lambda) \mathcal{T}_{\mu} S_{n-1}^{(\lambda,\mu)}(x), \quad n \ge 2,
$$
\n(4.3)

where

$$
a_{n-2} = \frac{\mu_{n+1}}{\mu_{n-1}} \varepsilon_{n-2}, \quad \eta_{n-2}^{\mu}(\lambda) = \frac{p_{n-1}}{s_{n-1}^{(\lambda,\mu)}} a_{n-2}, \quad n \ge 2.
$$
 (4.4)

The coefficients $\eta_n^{\mu}(\lambda)$, $n \geq 0$, will be called Dunkl-Sobolev coefficients.

Proof. Since $\{S_n^{(\lambda,\mu)}\}_{n\geq 0}$ is a basis in \mathcal{P} , then

$$
\frac{P_{n+1}(x)}{\mu_{n+1}} - \sigma_{n-1} \frac{P_{n-1}(x)}{\mu_{n-1}} = \frac{S_{n+1}^{(\lambda,\mu)}(x)}{\mu_{n+1}} + \sum_{k=0}^{n} c_k^{(n)} S_k^{(\lambda,\mu)}(x),
$$

for some constants $c_k^{(n)}$ $k^{(n)}$. So, from (4.1) and (3.1) , we get

$$
c_k^{(n)} \langle S_k^{(\lambda,\mu)}, S_k^{(\lambda,\mu)} \rangle_{s,\mu} = \langle u, \left(\frac{P_{n+1}}{\mu_{n+1}} - \sigma_{n-1} \frac{P_{n-1}}{\mu_{n-1}} \right) S_k^{(\lambda,\mu)} \rangle + \lambda \langle v, R_n \mathcal{T}_{\mu} S_k^{(\lambda,\mu)} \rangle.
$$

Thus,

$$
c_k^{(n)} = 0, \ \ \forall \ 0 \le k \le n - 2.
$$

So,

$$
\frac{P_{n+1}(x)}{\mu_{n+1}} - \sigma_{n-1} \frac{P_{n-1}(x)}{\mu_{n-1}} = \frac{S_{n+1}^{(\lambda,\mu)}(x)}{\mu_{n+1}} + c_n^{(n)} S_n^{(\lambda,\mu)}(x) + c_{n-1}^{(n)} S_{n-1}^{(\lambda,\mu)}(x).
$$

Taking into account the fact that the polynomials P_{n+1} and P_{n-1} are both either even or odd functions, then $c_n^{(n)} = 0$. Therefore,

$$
\frac{P_{n+1}(x)}{\mu_{n+1}} - \sigma_{n-1} \frac{P_{n-1}(x)}{\mu_{n-1}} = \frac{S_{n+1}^{(\lambda,\mu)}(x)}{\mu_{n+1}} + c_{n-1}^{(n)} S_{n-1}^{(\lambda,\mu)}(x).
$$

Furthermore,

$$
c_{n-1}^{(n)} = -\frac{\sigma_{n-1}}{\mu_{n-1}} \frac{p_{n-1}}{s_{n-1}^{(\lambda,\mu)}}.
$$

Thus [\(4.2\)](#page-11-0) holds with

$$
a_{n-2} = \frac{\mu_{n+1}}{\mu_{n-1}} \varepsilon_{n-2}, \quad \eta_{n-2}^{\mu}(\lambda) = \frac{p_{n-1}}{s_{n-1}^{(\lambda,\mu)}} a_{n-2}, \quad n \ge 2.
$$

Applying now the operator \mathcal{T}_{μ} to [\(4.2\)](#page-11-0) and using [\(3.1\)](#page-4-0), we obtain [\(4.3\)](#page-11-1).

Notice that the converse result is not true. Indeed,

Theorem 4.2. Let u and v be two symmetric positive measures and define the Dunkl-Sobolev inner product by [\(4.1\)](#page-10-0). We assume that the monic polynomials $\{P_n\}_{n\geq 0}$ are orthogonal with respect to u and let $\{S_n^{(\lambda,\mu)}\}_{n\geq 0}$ be the sequence of Dunkl-Sobolev monic polynomials orthogonal with respect to [\(4.1\)](#page-10-0). Assume that they are related by [\(4.2\)](#page-11-0), with $\{a_n\}_{n\geq 1}$ and $\{\eta_n^{\mu}(\lambda)\}_{n\geq 1}$ nonzero sequences of real numbers. Then, there exists a sequence of real numbers $\{e_n\}_{n\geq 2}$ such that

$$
\mathcal{T}_{\mu} \mathcal{P}_{n+1}(x) + a_{n-2} \mathcal{T}_{\mu} \mathcal{P}_{n-1}(x) = \mathcal{R}_n(x) + e_n \mathcal{R}_{n-2}(x), \quad n \ge 2,
$$

where ${R_n}_{n\geq 0}$ denotes the MOPS with respect to the measure v.

 \blacksquare

 \blacksquare

Proof. Notice that

$$
\mathcal{T}_{\mu} \mathcal{P}_{n+1}(x) + a_{n-2} \mathcal{T}_{\mu} \mathcal{P}_{n-1}(x) = \mu_{n+1} \mathcal{R}_{n}(x) + \sum_{i=0}^{n-1} e_{n,k} \mathcal{R}_{k}(x),
$$

where

$$
e_{n,k} = \frac{< v, (\mathcal{T}_{\mu}P_{n+1} + a_{n-2}\mathcal{T}_{\mu}P_{n-1})R_k>}{r_k}
$$

.

On the other hand, from [\(4.1\)](#page-10-0) and [\(4.2\)](#page-11-0), for $0 \le k \le n-2$, we get

$$
0 = \langle S_{n+1}^{(\lambda,\mu)} + \eta_{n-2}^{\mu}(\lambda)S_{n-1}^{(\lambda,\mu)}, x^k \rangle_{s,\mu} = \lambda \mu_k \langle v, \left(\mathcal{T}_{\mu}P_{n+1} + a_{n-2}\mathcal{T}_{\mu}P_{n-1}\right)x^{k-1} \rangle.
$$

Then,

$$
< v, (\mathcal{T}_{\mu}P_{n+1} + a_{n-2}\mathcal{T}_{\mu}P_{n-1})x^{k-1} > = 0
$$
, for all $0 \le k \le n-3$.

So, we can write $\mathcal{T}_{\mu}P_{n+1} + a_n \mathcal{T}_{\mu}P_{n-1}$ as

$$
\mathcal{T}_{\mu}P_{n+1} + a_{n-2}\mathcal{T}_{\mu}P_{n-1} = \mu_{n+1}R_n + e_{n,n-1}R_{n-1} + e_{n,n-2}R_{n-2}, \quad n \ge 2.
$$

Because of the symmetry of the polynomials, the coefficient $e_{n,n-1}$ most be zero. Let denote $e_{n,n-2}=e_n$. Then

$$
\mathcal{T}_{\mu}P_{n+1(x)} + a_{n-2}\mathcal{T}_{\mu}P_{n-1}(x) = \mu_{n+1}R_n(x) + e_nR_{n-2}(x), \quad n \ge 2.
$$

Now we give an explicit expression for the coefficients e_n . Using the orthogonality of the sequence $\{S_n^{(\lambda,\mu)}\}_{n\geq 0}$ with respect to (4.1) , we have

$$
\langle S_{n+1}^{(\lambda,\mu)} + \eta_{n-2}^{\mu}(\lambda)S_{n-1}^{(\lambda,\mu)}, x^{n-1} \rangle_{s,\mu} = \eta_{n-2}^{\mu}(\lambda)s_{n-1}^{(\lambda,\mu)}.
$$

On the other hand, by [\(4.2\)](#page-11-0), we get

$$
\langle S_{n+1}^{(\lambda,\mu)} + \eta_{n-2}^{\mu}(\lambda)S_{n-1}^{(\lambda,\mu)}, x^{n-1} \rangle_{s,\mu} = a_{n-2}p_{n-1} + \lambda\mu_{n-1}e_n r_{n-2}, \quad n \ge 2.
$$

Therefore,

$$
e_n = \frac{\eta_{n-2}^{\mu}(\lambda)s_{n-1}^{(\lambda,\mu)} - a_{n-2}p_{n-1}}{\lambda\mu_{n-1}r_{n-2}}, \quad n \ge 2.
$$

Remark 4.1. This is the analogue in the Dunkl framework of a result stated in [\[9\]](#page-20-14).

From the previous theorem, we see that (u, v) is a symmetric \mathcal{T}_{μ} -coherent pair of positive measures if and only if $e_n = 0$ for all $n \geq 2$, which is equivalent to

$$
\eta_{n-2}^{\mu}(\lambda) = \frac{p_{n-1}}{s_{n-1}^{(\lambda,\mu)}} a_{n-2}, \quad n \ge 2.
$$

The following lemma provides a recurrence relation for the sequence $\{s_n^{(\lambda,\mu)}\}_{n\geq 0}$.

Lemma 4.1. For $n \geq 2$

$$
s_{n+1}^{(\lambda,\mu)} = p_{n+1} + a_{n-2}^2 p_{n-1} + \lambda \mu_{n+1}^2 r_n - (\eta_{n-2}^\mu(\lambda))^2 s_{n-1}^{(\lambda,\mu)}.
$$
 (4.5)

 \blacksquare

Proof. For $n \geq 2$, we define

$$
Q_{n+1}(x) = P_{n+1}(x) + a_{n-2}P_{n-1}(x).
$$

Then, from (4.2) ,

$$
Q_{n+1}(x) = S_{n+1}^{(\lambda,\mu)}(x) + \eta_{n-2}^{\mu}(\lambda)S_{n-1}^{(\lambda,\mu)}(x). \tag{4.6}
$$

We first prove that for all $n\geq 4$

$$
\eta_{n-2}^{\mu}(\lambda) = \frac{Q_{n+1}, Q_{n-1} >_{s,\mu}}{Q_{n-1}, Q_{n-1} >_{s,\mu} - \eta_{n-4}^{\mu}(\lambda) < Q_{n-1}, Q_{n-3} >_{s,\mu}}.\tag{4.7}
$$

As a straightforward consequence of [\(4.6\)](#page-13-0) we get

$$
\langle \mathcal{Q}_{n+1}, \mathcal{Q}_{n-1} \rangle_{s,\mu} = \eta_{n-2}^{\mu}(\lambda) \langle S_{n-1}^{(\lambda,\mu)}, \mathcal{Q}_{n-1} \rangle_{s,\mu} \tag{4.8}
$$

and

$$
\langle Q_{n-1}, Q_{n-1} - \eta_{n-4}^{\mu}(\lambda)Q_{n-3} >_{s,\mu} \n= \langle S_{n-1}^{(\lambda,\mu)} + \eta_{n-4}^{\mu}(\lambda)S_{n-3}^{(\lambda,\mu)}, Q_{n-1} - \eta_{n-4}^{\mu}(\lambda)Q_{n-3} >_{s,\mu} \n= \langle S_{n-1}^{(\lambda,\mu)}, Q_{n-1} >_{s,\mu} + \eta_{n-4}^{\mu}(\lambda) < S_{n-3}^{(\lambda,\mu)}, Q_{n-1} >_{s,\mu} - (\eta_{n-4}^{\mu}(\lambda))^2 < S_{n-3}^{(\lambda,\mu)}, Q_{n-3} >_{s,\mu} \n= \langle S_{n-1}^{(\lambda,\mu)}, Q_{n-1} >_{s,\mu} + \eta_{n-4}^{\mu}(\lambda) < S_{n-3}^{(\lambda,\mu)}, S_{n-1}^{(\lambda,\mu)} + \eta_{n-4}^{\mu}(\lambda)S_{n-3}^{(\lambda,\mu)} >_{s,\mu} - (\eta_{n-4}^{\mu}(\lambda))^2 < S_{n-3}^{(\lambda,\mu)}, Q_{n-3} >_{s,\mu} \n= \langle S_{n-1}^{(\lambda,\mu)}, Q_{n-1} >_{s,\mu} + (\eta_{n-4}^{\mu}(\lambda))^2 < S_{n-3}^{(\lambda,\mu)}, Q_{n-3} >_{s,\mu} - (\eta_{n-4}^{\mu}(\lambda))^2 < S_{n-3}^{(\lambda,\mu)}, Q_{n-3} >_{s,\mu} \n= \langle S_{n-1}^{(\lambda,\mu)}, Q_{n-1} >_{s,\mu} .
$$
\n(4.9)

Replacing [\(4.9\)](#page-13-1) in [\(4.8\)](#page-13-2)

$$
{s,\mu}=\eta^{\mu}{n-2}(\lambda)_{s,\mu}
$$

and (4.7) holds.

Now, from the definition,

$$
\langle Q_{n-1}, Q_{n-1} \rangle_{s,\mu} = \langle u, Q_{n-1}Q_{n-1} \rangle + \lambda \langle v, \mathcal{T}_{\mu}Q_{n-1}\mathcal{T}_{\mu}Q_{n-1} \rangle
$$

=\langle u, Q_{n-1}Q_{n-1} \rangle + \lambda \mu_{n-1}^2 \langle v, R_{n-2}(x)R_{n-2}(x) \rangle
=\langle u, (P_{n-1} + a_{n-4}P_{n-3})(P_{n-1} + a_{n-4}P_{n-3}) \rangle + \lambda \mu_{n-1}^2 r_{n-2}
= p_{n-1} + a_{n-4}^2 p_{n-3} + \lambda \mu_{n-1}^2 r_{n-2},

$$
\langle Q_{n+1}, Q_{n-1} \rangle_{s,\mu} = \langle u, Q_{n+1}Q_{n-1} \rangle + \lambda \langle v, \mathcal{T}_{\mu}Q_{n+1}\mathcal{T}_{\mu}Q_{n-1} \rangle
$$

=\langle u, Q_{n+1}Q_{n-1} \rangle + \lambda \mu_{n+1}\mu_{n-1} \langle v, R_n(x)R_{n-2}(x) \rangle
=\langle u, (P_{n+1} + a_{n-2}P_{n-1})(P_{n-1} + a_{n-4}P_{n-3}) \rangle
= a_{n-2}p_{n-1}

and

$$
{s,\mu}=a{n-4}p_{n-3}.
$$

So, [\(4.7\)](#page-13-3) becomes

$$
\eta_{n-2}^{\mu}(\lambda) = \frac{a_{n-2}p_{n-1}}{p_{n-1} + a_{n-4}^2 p_{n-3} + \lambda \mu_{n-1}^2 r_{n-2} - \eta_{n-4}^{\mu}(\lambda) a_{n-4} p_{n-3}}.
$$
(4.10)

According to (4.4) , (4.5) holds.

Remark 4.2. [\(4.5\)](#page-12-0) is useful in order to compute the norms $s_n^{(\lambda,\mu)}$ if the Dunkl-Sobolev coefficients are known, with some initial conditions. First, from their definitions it is easy to see that

$$
s_1^{(\lambda,\mu)} = \lambda + p_1, \quad \eta_0^{\mu}(\lambda) = \frac{a_0 p_1}{\lambda + p_1}.
$$
 (4.11)

Now, for $n=1$,

$$
\eta_1^{\mu}(\lambda) = \frac{a_1 p_2}{s_2^{(\lambda,\mu)}}.
$$

Since $S_2^{(\lambda,\mu)}$ $P_2^{(\lambda,\mu)}(x) = P_2(x) = x^2 - \gamma_1$, then

$$
s_2^{(\lambda,\mu)} = p_2 + 4\lambda r_1, \quad \eta_1^{\mu}(\lambda) = \frac{a_1 p_2}{p_2 + 4\lambda r_1}.
$$
 (4.12)

Now, using [\(4.11\)](#page-14-0) and from [\(4.5\)](#page-12-0) with $n = 2$ we obtain $s_3^{(\lambda,\mu)}$ $\frac{(\lambda,\mu)}{3}$. As a consequence, from (4.4) we get η_2^{μ} $\frac{\mu}{2}(\lambda)$.

In an analogue way, for $n = 2k$, where k is an integer number greater than or equal to 2, in [\(4.5\)](#page-12-0) we obtain $s_{2k+1}^{(\lambda,\mu)}$ and then η_{2k+1}^{μ} $\frac{\mu}{2k}(\lambda)$.

Similarly, using (4.12) and from (4.5) with $n = 2k + 1$, where k is an integer number greater than or equal to 2, we can get $s_{2k+2}^{(\lambda,\mu)}$ and then $\eta_{2k+1}^{\mu}(\lambda)$.

5 Algorithm for the Fourier Dunkl-Sobolev coefficients

Let us define the Dunkl-Sobolev space as

$$
\mathcal{W}_{2}^{1}(\mathbb{R}, u, v, \mu) = \{f; \ ||f||_{u}^{2} + \lambda ||\mathcal{T}_{\mu}f||_{v}^{2} < \infty\}.
$$

If $f \in \mathcal{W}_2^1(\mathbb{R}, u, v, \mu)$, then its Fourier expansion in terms of Dunkl-Sobolev orthogonal polynomials $\{S_n^{\lambda_\mu}\}_{n\geq 0}$ is

$$
f(x) \sim \sum_{n=0}^{\infty} \frac{_{s,\mu}}{s_n^{(\lambda,\mu)}} S_n^{(\lambda,\mu)}(x).
$$

We denote $f_n^{(\lambda,\mu)} \ll f, S_n^{(\lambda,\mu)} >_{s,\mu}$ and $\mathcal{F}_n^{(\lambda,\mu)} = \frac{f_n^{(\lambda,\mu)}}{S_n^{(\lambda,\mu)}}$ $\frac{f_n^{(\lambda,\mu)}}{s_n^{(\lambda,\mu)}}$. $\mathcal{F}_n^{(\lambda,\mu)}$ is said to be the *n*-th Dunkl Fourier-Sobolev coefficient.

Now, take $f \in \mathcal{W}_2^1(\mathbb{R}, u, v, \mu)$ and let consider a sequence of real numbers $\{w_n^{\mu}(f)\}_{n \geq 0}$ such that

$$
w_n^{\mu}(f) = \langle u, (P_{n+2} + a_{n-1}P_n)f \rangle + \lambda \langle v, (\mathcal{T}_{\mu}P_{n+2} + a_{n-1}\mathcal{T}_{\mu}P_n)\mathcal{T}_{\mu}f \rangle. \tag{5.1}
$$

Lemma 5.1. The following relation holds

$$
f_{n+2}^{(\lambda,\mu)} + \eta_{n-1}^{\mu}(\lambda)f_n^{(\lambda,\mu)} = w_n^{\mu}(f), \quad n \ge 0,
$$
\n(5.2)

with initial conditions

$$
\eta_{-1}^{\mu}(\lambda) = 0, \quad f_0^{(\lambda,\mu)} = \langle f, 1 \rangle_{s,\mu} = \langle u, f \rangle, \quad f_1^{(\lambda,\mu)} = \langle u, xf \rangle + \lambda \mu_1 \langle v, \mathcal{T}_{\mu} f \rangle
$$

and

$$
w_0^\mu(f)=+ \lambda .
$$

Proof. Using (4.2) , we get

$$
\langle f, S_{n+2}^{(\lambda,\mu)} \rangle_{s,\mu} = -\eta_{n-1}^{\mu}(\lambda) \langle f, S_{n}^{(\lambda,\mu)} \rangle_{s,\mu} + \langle f, (\mathbf{P}_{n+2} + a_{n-1}\mathbf{P}_{n}) \rangle_{s,\mu}
$$

= $-\eta_{n-1}^{\mu}(\lambda) \langle f, S_{n}^{(\lambda,\mu)} \rangle_{s,\mu} + \langle u, (\mathbf{P}_{n+2} + a_{n-1}\mathbf{P}_{n})f \rangle$
+ $\lambda \langle v, (\mathcal{T}_{\mu}\mathbf{P}_{n+2} + a_{n-1}\mathcal{T}_{\mu}\mathbf{P}_{n})\mathcal{T}_{\mu}f \rangle$

and the result follows.

We will now describe an algorithm to compute the Fourier Dunkl-Sobolev coefficients in the Fourier expansions in terms of Dunkl-Sobolev orthogonal polynomials. We assume that ε_0 and ε_1 are known.

Algorithm 1. Even order Fourier Dunkl-Sobolev coefficients

For *n* even, the Fourier Dunkl-Sobolev coefficients $\mathcal{F}_n^{(\lambda,\mu)}$ can be computed as follows.

Step 0. Initial conditions λ , μ , ε_0 , ε_1 , η_-^{μ} $\frac{\mu}{-1}(\lambda), s_0^{(\lambda,\mu)}$ $\binom{(\lambda,\mu)}{0},\ f_0^{(\lambda,\mu)}$ $v_0^{(\lambda,\mu)},\,w_0^\mu$ $\frac{\mu}{0}(f)$, $s_2^{(\lambda,\mu)}$ and η_1^{μ} $\frac{\mu}{1}(\lambda)$.

Step 1. Using step 0 and from (5.2) when $n = 0$ you get $f_2^{(\lambda,\mu)}$ $F_2^{(\lambda,\mu)}$. Then calculate $F_2^{(\lambda,\mu)}$ $2^{(\lambda,\mu)}$.

Step 2. Using step 0 and the information in step 1 to compute: w_2^{μ} $\frac{\mu}{2}(f)$ with (5.1) and $n = 2, f_4^{(\lambda,\mu)}$ $f_4^{(\lambda,\mu)}$ from [\(5.2\)](#page-14-2) with $n=2$ and $s_4^{(\lambda,\mu)}$ $\chi_4^{(\lambda,\mu)}$ taking $n=3$ in [\(4.5\)](#page-12-0). Then compute $F_4^{(\lambda,\mu)}$ $\frac{1}{4}$.

Step k. For $k > 3$, using the step 0 and the information in steps 1 to $k - 1$ to compute: ε_{2k-3} from [\(3.13\)](#page-6-6) and $n = 2k - 4$, a_{2k-3} from [\(4.4\)](#page-11-2) with $n = 2k - 1$, η_{2k}^{μ} $\int_{2k-3}^{\mu}(\lambda)$ from (4.4) with $n = 2k - 1$, w_2^{μ} $a_{2k-2}^{\mu}(f)$ with [\(5.1\)](#page-14-3) and $n = 2k - 2$, $f_{2k}^{(\lambda,\mu)}$ $\sum_{2k}^{(\lambda,\mu)}$ from (5.2) with $n = 2k - 2$ and, finally, $s_{2k}^{(\lambda,\mu)}$ $\chi_{2k}^{(\lambda,\mu)}$ taking $n = 2k - 1$ in [\(4.5\)](#page-12-0). Then, compute $F_{2k}^{(\lambda,\mu)}$ $\frac{1}{2k}$.

Algorithm 2. Odd order Fourier Dunkl-Sobolev coefficients

For *n* odd, the Fourier Dunkl-Sobolev coefficients $\mathcal{F}_n^{(\lambda,\mu)}$ can be computed as follows.

Step 0. Initial conditions λ , μ , ε_0 , ε_1 , η_0^{μ} $g_0^\mu(\lambda),\,s_1^{(\lambda,\mu)}$ $j_1^{(\lambda,\mu)}, f_1^{(\lambda,\mu)}$ and w_1^{μ} $\frac{\mu}{1}(f).$

Step 1. Using step 0 to compute $f_2^{(\lambda,\mu)}$ $f_2^{(\lambda,\mu)}$ from [\(5.2\)](#page-14-2) with $n=0$ and $s_3^{(\lambda,\mu)}$ $_{3}^{\left(\lambda,\mu\right) }$ from $\left(4.5\right)$ with $n=2$. Then, calculate $F_3^{(\lambda,\mu)}$ $3^{(\lambda,\mu)}$.

Step 2. Using step 0 and the information in step 1 to compute: ε_2 from [\(3.13\)](#page-6-6) and $n = 1$, a_2 from [\(4.4\)](#page-11-2) with $n = 4, \eta_2^{\mu}$ $\binom{\mu}{2}(\lambda)$ from [\(4.4\)](#page-11-2) with $n=4$, w_3^{μ} $j_3^{\mu}(f)$ with [\(5.1\)](#page-14-3) and $n = 3$, $f_5^{(\lambda,\mu)}$ 5 from [\(5.2\)](#page-14-2) with $n=3$ and $s_5^{(\lambda,\mu)}$ $\binom{(\lambda,\mu)}{5}$ taking $n = 4$ in [\(4.5\)](#page-12-0). Then compute $F_5^{(\lambda,\mu)}$ $5^{(\lambda,\mu)}$.

Step k. For $k \geq 3$, using the step 0 and the information in steps 1 to $k-1$ to compute: ε_{2k-2} from [\(3.13\)](#page-6-6) and $n = 2k - 3$, a_{2k-2} from [\(4.4\)](#page-11-2) with $n = 2k$, η_{2k}^{μ} $a_{2k-2}^{\mu}(\lambda)$ from (4.4) with $n = 2k, w^{\mu}_{2k}$ $_{2k-1}^{(\mu}(f)$ with (5.1) and $n = 2k - 1$, $f_{2k+1}^{(\lambda,\mu)}$ from (5.2) with $n = 2k - 1$ and finally $s_{2k+1}^{(\lambda,\mu)}$ taking $n=2k$ in [\(4.5\)](#page-12-0). Then, compute $F_{2k+1}^{(\lambda,\mu)}$.

Remark 5.1. Notice that an algorithm for generate Fourier-Sobolev coefficients when you

deal with $(1, 1)$ coherent pairs of measures with respect to the derivative operator has been presented in [\[10\]](#page-20-15).

6 Numerical examples

In order to illustrate the algorithm presented above, in this section we give an example involving the construction of the Fourier Dunkl–Sobolev series with respect to \mathcal{T}_{μ} – Sobolev inner products of the type (4.1) where (u, v) is a \mathcal{T}_{μ} -symmetric coherent pair of measures and u is the classical one, i.e. u is the generalized Hermite weight \mathcal{H}^{μ} or the generalized Gegenbauer weight $\mathcal{G}^{(\alpha,\mu-\frac{1}{2})}$.

6.1 The generalized Hermite case

The sequence of generalized Hermite polynomials ${H_n^{\mu}}_{n\geq 0}$ is an MOPS with respect to the positive definite linear form \mathcal{H}^{μ} defined by the weight function

$$
\omega(x) = \frac{1}{\Gamma(\alpha + \frac{1}{2})}|x|^{2\mu}e^{-x^2}, \mu > -1/2,
$$

supported on the real line.

This positive measure is \mathcal{T}_{μ} -classical and satisfies the \mathcal{T}_{μ} -Pearson equation

$$
\mathcal{T}_{\mu}(\mathcal{H}^{\mu})=-2x\mathcal{H}^{\mu}.
$$

If we apply \mathcal{T}_{μ} to \mathcal{H}_{n}^{μ} , we obtain [\[21\]](#page-20-16)

$$
\mathcal{T}_{\mu}H_{n}^{\mu}(x) = \mu_{n}H_{n-1}^{\mu}(x).
$$
\n(6.1)

This means that $\{H_n^{\mu}\}_{n\geq 0}$ is an \mathcal{T}_{μ} -Appell sequence.

Now, let $\{u, v\}$ be a \mathcal{T}_{μ} -symmetrically coherent pair of positive measures. Suppose that u is \mathcal{T}_{μ} -classical with $u = \mathcal{H}^{\mu}$. From [\[23\]](#page-20-1), we obtain

$$
\mathcal{H}^{\mu} = 2\frac{\sigma_1 \sigma_3}{r_2 r_4} (x^2 + \xi^2)v.
$$

Using (3.2) , we get

$$
R_n(x) = H_n^{\mu}(x) + \varepsilon_{n-2} H_{n-2}^{\mu}(x), \quad n \ge 2.
$$
 (6.2)

On the other hand,

$$
\begin{cases} H_0^{\mu}(x) = 1, & H_1^{\mu}(x) = x, \\ H_{n+1}^{\mu}(x) = x H_n^{\mu}(x) - \gamma_n H_{n-1}^{\mu}(x), & n \ge 1. \end{cases}
$$

with

$$
\gamma_{2n} = n, \gamma_{2n-1} = n - 1/2 + \mu), \ n \ge 1,
$$

Notice that the positive definite condition yields $\mu > -n - \frac{1}{2}$ $\frac{1}{2}$, $n \geq 0$.

We denote the TTRR relation for the MOPS $\{R_n\}_{n\geq 0}$

$$
R_{n+1}(x) = xR_n(x) - \tilde{\gamma}_n R_{n-1}(x), \quad n \ge 1, \quad R_1(x) = x, \quad R_0(x) = 1.
$$

From Proposition [3.1,](#page-5-4)

$$
\tilde{\gamma}_1 = \frac{1}{2} + \mu - \varepsilon_0,
$$

$$
\tilde{\gamma}_n = \frac{1}{2}\mu_n + \varepsilon_{n-2} - \varepsilon_{n-1}, \quad n \ge 2,
$$

and

$$
\frac{1}{2}\mu_n \varepsilon_n = \varepsilon_{n-1} \left(\frac{1}{2}\mu_{n+2} + \varepsilon_n - \varepsilon_{n+1}\right), \quad n \ge 1.
$$

In addition, by [\(3.13\)](#page-6-6), the sequence $\{\varepsilon_n\}_{n>0}$ satisfies the nonlinear quadratic difference equation

$$
\varepsilon_{n+1} = \frac{1}{2}(\mu_{n+1} + \mu_{n+2}) + (\varepsilon_1 - 1)(1 - \frac{1+2\mu}{2\varepsilon_0}) - \frac{1}{4}\frac{\mu_n\mu_{n+1}}{\varepsilon_{n-1}}, \quad n \ge 1.
$$

Therefore, for fixed initial values ε_0 and ε_1 we can compute the sequences of parameters $\{\varepsilon_n\}_{n\geq 0}$ and $\{\tilde{\delta}_n\}_{n\geq 1}$.

As an illustrative example, if $\mu = 5$, $\lambda = 0.1$, $\varepsilon_0 = 1.2$, $\varepsilon_1 = 1.3$ and $\xi = 0$, then $\gamma_n = \frac{1}{2}$ $rac{1}{2}\mu_n,$ $s_0^{(\lambda,\mu)} = 1, s_1^{(\lambda,\mu)} = 5.6, s_2^{(\lambda,\mu)} = 7.22, \eta_{-}^{\mu}$ $_{-1}^{\mu}(\lambda) = 0, \eta_0^{\mu}$ $j_0^{\mu}(\lambda) = 1.39, \eta_1^{\mu}$ $j_1^{\mu}(\lambda) = 1.98.$ Furthermore,

$$
u = \mathcal{H}^5 = 0.02x^{10}e^{-x^2}
$$

.

As a consequence,

$$
v = 0.45x^8e^{-x^2} + \delta_0.
$$

Next, let consider the function

$$
f(x) = x(10 - x).
$$

Obviously $f \in \mathcal{W}_2^1(\mathbb{R}, u, v, \mu)$. In this case $f_0^{(\lambda, \mu)} = -5.758$, $f_1^{(\lambda, \mu)} = 287.886$, w_0^{μ} $_0^{\mu}(f) = -15.18$ and w_1^{μ} $_{1}^{\mu}(f) = 80.551.$

Now, applying the algorithm introduced in Section 6, we find the Fourier \mathcal{T}_{μ} –Sobolev coefficients of f .

6.2 The generalized Gegenbauer case

The sequence of generalized Gegenbauer polynomials $\{G_n^{(\alpha,\mu-\frac{1}{2})}\}_{n\geq 0}$ is an MOPS with respect to the positive definite linear form $\mathcal{G}^{(\alpha,\mu-\frac{1}{2})}$ defined by the weight function

$$
\omega(x) = \frac{\Gamma(\alpha + \mu + \frac{3}{2})}{\Gamma(\alpha + 1)\Gamma(\mu + \frac{1}{2})}|x|^{2\mu}(1 - x^2)^{\alpha}, \alpha > -1,
$$

supported on the interval $(-1, 1)$.

The corresponding form is a positive definite \mathcal{T}_{μ} -classical linear form and satisfies the \mathcal{T}_{μ} -Pearson equation

$$
\mathcal{T}_{\mu}((x^2-1)\mathcal{G}^{(\alpha,\mu-\frac{1}{2})})=2(\alpha+1)x\mathcal{G}^{(\alpha,\mu-\frac{1}{2})}.
$$

The MOPS $\{G_n^{(\alpha,\mu-\frac{1}{2})}\}_{n\geq 0}$ satisfies the TTRR [\(2.2\)](#page-2-0) (see [\[1\]](#page-19-5))

$$
\hat{\beta}_n = 0, \quad \gamma_{n+1} = \frac{(n+1+\varrho_n)(n+1+2\alpha+\varrho_n)}{4(n+\alpha+\mu+\frac{1}{2})(n+\alpha+\mu+\frac{3}{2})}, \quad n \ge 0,
$$

$$
\varrho_n = 2\mu \frac{1+(-1)^n}{2}, \quad n \ge 0.
$$

If we apply \mathcal{T}_{μ} to $\mathcal{G}_n^{(\alpha,\mu-\frac{1}{2})}$, then we obtain [\[2\]](#page-19-3)

$$
\mathcal{T}_{\mu}(\mathcal{G}_n^{(\alpha,\mu-\frac{1}{2})}) = \mu_n \mathcal{G}_{n-1}^{(\alpha+1,\mu-\frac{1}{2})}.
$$
\n(6.3)

Let v be a symmetric positive measure and denote by $\{R_n\}_{n\geq 0}$ the corresponding MOPS. Assume that $(\mathcal{G}^{(\alpha,\mu-\frac{1}{2})},v)$ is a \mathcal{T}_{μ} -symmetric coherent pair. Thus, from [\[23\]](#page-20-1), we get

$$
(x^{2} - 1)\mathcal{G}^{(\alpha,\mu-\frac{1}{2})} = 2\frac{\sigma_{1}\sigma_{3}}{r_{2}r_{4}}(x^{2} - \xi^{2})v, |\xi| \ge 1.
$$

Since

$$
(1 - x^2) \mathcal{G}^{(\alpha, \mu - \frac{1}{2})} = \frac{\alpha + 1}{\alpha + \mu + \frac{3}{2}} \mathcal{G}^{(\alpha + 1, \mu - \frac{1}{2})},
$$

then

$$
\mathcal{G}^{(\alpha+1,\mu-\frac{1}{2})} = 2\frac{\alpha+\mu+\frac{3}{2}}{\alpha+1}\frac{\sigma_1\sigma_3}{r_2r_4}(\xi^2-x^2)v, \xi \ge 1.
$$

Now, from [\(3.2\)](#page-5-0) and [\(6.3\)](#page-18-0)

$$
R_n(x) = G_n^{(\alpha+1,\mu-\frac{1}{2})}(x) + \varepsilon_{n-2} G_{n-2}^{(\alpha+1,\mu-\frac{1}{2})}(x), n \ge 2,
$$

where ε_n is non-zero constant for all $n \geq 0$.

The TTRR for the MOPS $\{R_n\}_{n\geq 0}$ reads as

$$
R_{n+1}(x) = xR_n(x) - \tilde{\gamma}_n R_{n-1}(x), \quad n \ge 1, \quad R_0(x) = 1, \quad R_1(x) = x,
$$

with $\tilde{\gamma}_n > 0$ for all $n \geq 1$. Then, by Corollary [3.1](#page-5-3)

$$
\tilde{\gamma}_n = \frac{(n + \varrho_{n-1})(n + 2\alpha + \varrho_{n-1})}{4(n + \alpha + \mu - \frac{1}{2})(n + \alpha + \mu + \frac{1}{2})} + \varepsilon_{n-2} - \varepsilon_{n-1}, \quad n \ge 1.
$$

In addition, according to [\(3.13\)](#page-6-6), we can write the parameters $\{\varepsilon_n\}_{n\geq 0}$ as

$$
\varepsilon_{n+1} = \frac{1}{4(n+\alpha+\mu+\frac{3}{2})} \left(\frac{(n+1+\varrho_n)(n+1+2\alpha+\varrho_n)}{n+\alpha+\mu+\frac{1}{2}} + \frac{(n+2+\varrho_{n+1})(n+2+2\alpha+\varrho_{n+1})}{n+\alpha+\mu+\frac{5}{2}} \right)
$$

$$
+ \Lambda - \frac{(n+\varrho_{n-1})(n+2\alpha+\varrho_{n-1})(n+1+\varrho_n)(n+1+2\alpha+\varrho_n)}{16(n+\alpha+\mu+\frac{1}{2})^2(n+\alpha+\mu-\frac{1}{2})(n+\alpha+\mu+\frac{3}{2})\varepsilon_{n-1}},
$$

where

$$
\Lambda = \left(\varepsilon_1 - \frac{1+\alpha}{(\frac{3}{2}+\alpha+\mu)(\frac{5}{2}+\alpha+\mu)}\right)\left(1 - \frac{\mu_1(1+2\alpha+2\mu)}{4\varepsilon_0(\frac{1}{2}+\alpha+\mu)(\frac{3}{2}+\alpha+\mu)}\right).
$$

Therefore, for fixed initial values ε_0 and ε_1 , we can compute the sequences of parameters $\{\varepsilon_n\}_{n\geq 0}$ and $\{\tilde{\gamma}_n\}_{n\geq 1}$.

We choose $\alpha = 5$, $\lambda = 0.5$, $\mu = 1$, $\varepsilon_0 = 0.1$, $\varepsilon_1 = 0.15$ and $\xi = 1$. In this case,

$$
\gamma_{n+1} = \frac{(n+1+\varrho_n)(n+11+\varrho_n)}{4(n+\frac{13}{2})(n+\frac{15}{2})}, \quad \varrho_n = 1 + (-1)^n, \quad n \ge 0,
$$

 $\Lambda = 0.0822, s_0^{(\lambda,\mu)} = 1, s_1^{(\lambda,\mu)} = 0.7, s_2^{(\lambda,\mu)} = 0.219, \eta_{-}^{\mu}$ $_{-1}^{\mu}(\lambda) = 0, \eta_0^{\mu}$ $_0^{\mu}(\lambda) = 0.049, \eta_1^{\mu}$ $j_1^{\mu}(\lambda) = 0.0258.$ Furthermore $u = \mathcal{G}^{(5,\frac{1}{2})} = 17.6x^2(1-x^2)^5$, $-1 < x < 1$.

$$
v = -4.3325 \cdot 10^{-6} (1 - x^2)^{-1} \mathcal{G}^{(6,\frac{1}{2})} + \frac{1}{2} (\delta_1 + \delta_{-1}),
$$

where

$$
\mathcal{G}^{(6,\frac{1}{2})} = 22x^2(1-x^2)^6, \quad -1 < x < 1.
$$

So

$$
v = -0.953 \cdot 10^{-4} x^2 (1 - x^2)^5 + \frac{1}{2} (\delta_1 + \delta_{-1}), \quad -1 < x < 1.
$$

We will use the function

$$
f(x) = xe^{-(x-0.2)^2}.
$$

It is clear that $f \in \mathcal{W}_2^1(\mathbb{R}, u, v, \mu)$. Then $f_0^{(\lambda,\mu)} = 0.0583, f_1^{(\lambda,\mu)} = 1.338, w_0^{\mu}$ $_0^{\mu}(f) = 0.081$ and w_1^{μ} $_1^{\mu}(f) = 1.768.$

These initial conditions and the algorithm introduced in the previous section yield the Fourier Dunkl–Sobolev coefficients of f.

7 Acknowledgements

The work of Francisco Marcellán has been supported by the research project [PID2021- $122154NB-I00$, Ortogonalidad y Aproximación con Aplicaciones en Machine Learning y Teoría de la Probabilidad funded by MICIU/AEI/10.13039/501100011033 and by "ERDF A Way of making Europe"and the Madrid Government (Comunidad de Madrid-Spain) under the Multiannual Agreement with UC3M in the line of Excellence of University Professors, grant EPUC3M23 in the context of the V PRICIT (Regional Program of Research and Technological Innovation).

References

- [1] S. Belmehdi, Generalized Gegenbauer orthogonal polynomials. Proceedings of the Fifth International Symposium on Orthogonal Polynomials, Special Functions and their Applications (Patras, 1999) J. Comput. Appl. Math. 133 (2001), no. 1-2, 195-205.
- [2] Y. Ben Cheikh, M. Gaied, Characterization of the Dunkl-classical symmetric orthogonal polynomials. Appl. Math. Comput. 187 (2007), 105-114.
- [3] C. F. Bracciali, L. Castaño-García, J. J. Moreno-Balcázar, *Some asymptotics for Sobolev* orthogonal polynomials involving Gegenbauer weights. J. Comput. Appl. Math. 235 (2010), no. 4, 904-915.
- [4] L. Castaño-García, J. J. Moreno-Balcázar, A Mehler-Heine-type formula for Hermite-Sobolev orthogonal polynomials. J. Comput. Appl. Math. 150 (2003), no. 1, 25-35.
- [5] T. S. Chihara, Generalized Hermite polynomials. Doctoral Dissertation, Purdue, 1955.
- [6] T. S. Chihara, An Introduction to Orthogonal Polynomials. Gordon and Breach, New York, 1978.
- [7] O. Ciaurri, J. M´ınguez, Fourier series of Gegenbauer-Sobolev polynomials. SIGMA Symmetry Integrability Geom. Methods Appl. 14 (2018), Paper no. 024, 11 pp.
- [8] M. G. De Bruin, W. G. M. Groenevelt, F. Marcellán, H. G. Meijer, J. J. Moreno-Balcázar, Asymptotics and zeros of symmetrically coherent pairs of Hermite type. In Difference Equations, Special Functions and Orthogonal Polynomials, 378-393, World Sci. Publ., Hackensack, NJ, 2007.
- [9] A. M. Delgado, F. Marcellán, On an extension of symmetric coherent pairs of orthogonal polynomials. J. Comput. Appl. Math. 178 (2005), no. 1-2, 155–168.
- [10] H. Dueñas Ruiz, F. Marcellán, A. Molano, On symmetric $(1,1)$ -coherent pairs and Sobolev orthogonal polynomials: an algorithm to compute Fourier coefficients. Rev. Colombiana Mat. 53 (2019), no. 2, 139-164.
- [11] C. F. Dunkl, Integral kernels reflection group invariance. Canad. J. Math. 43 (1991), 1213-1227.
- [12] A. Iserles, P. E. Koch, S. P. Nørsett, J. M. Sanz-Serna, On polynomials orthogonal with respect to certain Sobolev inner products. J. Approx. Theory 65 (1991), no. 2, 151–175.
- [13] F. Marcell´an, A. Branquinho, J. Petronilho, Classical orthogonal polynomials: A functional approach. Acta Appl. Math. 34 (1994), 283-303.
- [14] F. Marcellán, T. E. Pérez, M. A. Piñar, *Gegenbauer-Sobolev orthogonal polynomials.* Math. Appl., 296. Kluwer Academic Publishers Group, Dordrecht, 1994, 71–82.
- [15] F. Marcellán, J. Petronilho, *Orthogonal polynomials and coherent pairs: the classical* case. Indag. Math. (N. S.) 6 (1995), no. 3, 287-307.
- [16] F. Marcellán, J. J. Moreno-Balcázar, Asymptotics and zeros of Sobolev orthogonal polynomials on unbounded supports. Acta Appl. Math. **94** (2006), no. 2, 163-192.
- [17] F. Marcell´an, Yuan Xu, On Sobolev orthogonal polynomials. Expo. Math. 33 (2015), no. 3, 308-352.
- [18] P. Maroni, Sur la suite de polynômes orthogonaux associée à la forme $u = \delta_c + \lambda(x (c)^{-1}L$. Period. Math. Hungar. 21 (1990), no. 3, 223-248.
- [19] A. Martínez-Finkelshtein, J. J. Moreno-Balcázar, H. Pijeira-Cabrera, *Strong asymp*totics for Gegenbauer-Sobolev orthogonal polynomials. J. Comput. Appl. Math. 81 (1997), no. 2, 211-216.
- [20] H. G. Meijer, Determination of all coherent pairs. J. Approx. Theory 89 (1997), no. 3, 321–343.
- [21] M. Rosenblum, Generalized Hermite polynomials and the Bose-like oscillator calculus. Oper. Theory Adv. Appl. 73 (1994), 369-396.
- [22] M. Sghaier A note on the Dunkl-classical orthogonal polynomials. Integral Transforms Spec. Funct. 23 (2012), 753–760.
- [23] M. Sghaier, S. Hamdi, Some results on Dunkl-coherent pairs. Integral Transforms Spec. Funct. 32 (2021), no. 11, 863-882.

[24] A. Zhedanov, Rational spectral transformations and orthogonal polynomials. J. Comput. Appl. Math. 85 (1997), 67-86.