

On Bivariate Jacobi Konhauser Polynomials

Mehmet Ali Özarslan¹ and İlkay Onbaşı Elidemir ^{*1}

¹Department of Mathematics, Faculty of Arts and Sciences, Eastern Mediterranean University, Gazimagusa, TRNC, Mersin 10, Turkey

Abstract

Recently, Özarslan and Elidemir (2023) introduced a methodology for constructing two-variable biorthogonal polynomial families with the help of one-variable biorthogonal and orthogonal polynomial families. The primary objective of the paper is to introduce novel class of two-variable biorthogonal polynomials namely bivariate Jacobi Konhauser polynomials. We investigate several fundamental properties of these polynomials including their biorthogonality property, operational formula, generating function, and integral representation. Furthermore, we investigate their images under the Laplace transform, fractional integral and derivative operators. Corresponding to these polynomials, we define the new type bivariate Jacobi Konhauser Mittag Leffler (JKML) function and obtain the similar properties for them. We introduce an integral operator containing the bivariate JKML function in the kernel.

1 Introduction

The general method for the construction of bivariate biorthogonal polynomials introduced in [1].

Remark 1.1. Let $w(x)$ be an admissible weight function over (a, b) , let $R_n(x)$ and $S_n(x)$ are biorthogonal polynomials of the basic polynomials $r(x)$ and $s(x)$, respectively. In an other words, they satisfy the biorthogonality condition

$$\int_a^b w(x)R_m(x)S_n(x)dx = J_{n,m} := \begin{cases} 0, & \text{if } n \neq m \\ J_{n,n}, & \text{if } n = m \end{cases}.$$

Also,

$$K_n(x) = \sum_{i=0}^n D_{n,i}[r(x)]^i$$

with

$$\int_c^d w(x)K_m(x)K_n(x)dx = \|K_n\|^2 \delta_{nm}.$$

Then the bivariate polynomials

$$P_n(x, y) = \sum_{s=0}^n \frac{D_{n,s}}{J_{n-s,n-s}} [r(x)]^s R_{n-s}(y) \quad \text{and} \quad Q_n(x, y) = K_n(x) \sum_{j=0}^n S_j(y) \quad (1)$$

are biorthogonal with respect to the weight function $w(x)w(y)$ over $(a, b) \times (c, d)$.

Bin-Saad introduced a class of 2D-Laguerre-Konhauser polynomials in [2] by

$${}_{\kappa}L_n^{(\alpha, \beta)}(x, y) = n! \sum_{s=0}^n \sum_{r=0}^{n-s} \frac{(-1)^{s+r} x^r y^{\kappa s + \beta}}{s! r! (n-s-r)! \Gamma(\alpha+r+1) \Gamma(\kappa s + \beta + 1)} \quad (\alpha, \beta, \kappa = 1, 2, \dots).$$

*Corresponding author. Email: ilkay.onbasi@emu.edu.tr

Note that ${}_{\kappa}L_n^{(\alpha,\beta)}(x,y)$ can be written in the following forms:

$${}_{\kappa}L_n^{(\alpha,\beta)}(x,y) = n! \sum_{s=0}^n \frac{(-1)^s x^{s+\alpha} y^{\beta} Z_{n-s}^{\beta}(y; \kappa)}{s! \Gamma(\alpha + s + 1) \Gamma(\kappa n - \kappa s + \beta + 1)}$$

and

$${}_{\kappa}L_n^{(\alpha,\beta)}(x,y) = n! \sum_{s=0}^n \frac{(-1)^s x^{\alpha} y^{\kappa s + \beta} L_{n-s}^{\alpha}(x)}{s! \Gamma(\alpha + n - s + 1) \Gamma(\kappa s + \beta + 1)},$$

where $L_n^{\alpha}(x)$ is the Laguerre polynomial and $Z_n^{\beta}(y; \kappa)$ is the Konhauser polynomial.

The polynomials $Z_n^{\beta}(x; \kappa)$ and $Y_n^{\beta}(x; \kappa)$ are the pair of biorthogonal polynomials that are suggested by the Laguerre polynomials where,

$$Z_n^{\beta}(x; \kappa) = \frac{\Gamma(1 + \beta + \kappa n)}{n!} \sum_{r=0}^n \frac{(-n)_r x^{\kappa r}}{r! \Gamma(\beta + 1 + \kappa r)},$$

$$Y_n^{(\beta)}(y; \kappa) = \frac{1}{n!} \sum_{i=0}^n \frac{x^i}{i!} \sum_{j=0}^i (-1)^j \binom{i}{j} \left(\frac{j + \beta + 1}{\kappa} \right)_n.$$

In [3] Özarslan and Kürt introduced another set of polynomials ${}_{\kappa}\mathfrak{L}_n^{(\alpha,\beta)}(x,y)$, by

$${}_{\kappa}\mathfrak{L}_n^{(\alpha,\beta)}(x,y) = L_n^{\alpha}(x) \sum_{s=0}^n Y_s^{(\beta)}(y; \kappa).$$

By using the orthogonality relations,

$$\int_0^{\infty} e^{-x} x^{\alpha} L_n^{\alpha}(x) L_m^{\alpha}(x) dx = \frac{\Gamma(n + \alpha + 1)}{n!} \delta_{nm}$$

and

$$\int_0^{\infty} e^{-x} x^{\beta} Z_n^{\beta}(x; \kappa) Y_m^{\beta}(x; \kappa) dx = \frac{\Gamma(\kappa n + \beta + 1)}{n!} \delta_{nm},$$

it can be easily seen that, two polynomial sets ${}_{\kappa}\mathfrak{L}_n^{(\alpha,\beta)}(x,y)$ and ${}_{\kappa}L_n^{(\alpha,\beta)}(x,y)$ are bi-orthonormal with respect to the weight function e^{-x-y} over the interval $(0, \infty) * (0, \infty)$. In fact,

$$\int_0^{\infty} \int_0^{\infty} e^{-x-y} {}_{\kappa}\mathfrak{L}_m^{(\alpha,\beta)}(x,y) {}_{\kappa}L_n^{(\alpha,\beta)}(x,y) dy dx = \delta_{nm},$$

where δ_{nm} is the Kronecker's delta.

Motivated by the above result, in [1] the authors introduced new classes of bivariate biorthogonal polynomial families, where the first family named as, Hermite-Konhauser polynomials.

$${}_v H_n^{\varrho}(x,y) = \sum_{s=0}^{\lfloor \frac{n}{2} \rfloor} \sum_{r=0}^{n-s} \frac{(-1)^s (-n)_{2s} (-n)_{s+r}}{(-n)_s \Gamma(\varrho + 1 + vr) s! r!} (2x)^{n-2s} y^{vr}$$

where $\varrho > -1$ and $v = 1, 2, \dots$

And the second class of polynomial families defined as,

$$Q_m(x,y) = H_m(x) \sum_{j=0}^m Y_j^{(\varrho)}(y; \kappa).$$

where $H_m(x)$ is the Hermite polynomials. It can be easily seen that, two polynomial sets ${}_v H_n^{\varrho}(x,y)$ and $Q_m(x,y)$ are biorthogonal with respect to the weight function $e^{-x^2-y} y^{\varrho}$ over the interval $(0, \infty) \times (-\infty, \infty)$. In fact,

$$\int_0^{\infty} \int_{-\infty}^{\infty} e^{-x^2-y} y^{\varrho} {}_v H_n^{\varrho}(x,y) Q_m(x,y) dx dy = \begin{cases} 0, & \text{if } n \neq m \\ 2^n n! \sqrt{\pi} & \text{if } n = m \end{cases},$$

Orthogonal polynomials are very useful in several fields of mathematics, and therefore physics and engineering, thanks to the properties and relations they provide. The theory of orthogonal polynomials has been developed by studying with different methods. Orthogonal polynomials have important applications in different fields such as numerical analysis, number theory, approximation theory, probability theory. Orthogonal polynomials in two variables are studied as the natural generalization of orthogonal polynomials in one variable. In 1967, Krall and Sheffer extended the concept of classical sequence of orthogonal polynomials to the bivariate case (see [4]). In 1975, Koornwinder studied examples of two-variables analogues of Jacobi polynomials, and he introduced seven classes of orthogonal polynomials which he considered to be the bivariate analogues of the Jacobi polynomials (see [5]). Some of these examples are classical polynomials as defined by Krall and Sheffer. We should also mention the book 'Orthogonal polynomials in two Variables' [6].

Corresponding to the polynomials ${}_rL_n^{(\varkappa, \varrho)}(x, y)$, Özarslan and Kürt [3] introduced a bivariate Mittag Leffler functions $E_{\varkappa, \varrho, v}^{(\gamma)}(x, y)$ by

$$E_{\varkappa, \varrho, v}^{(\gamma)}(x, y) = \sum_{r=0}^{\infty} \sum_{s=0}^{\infty} \frac{(\gamma)_{r+s} x^r y^s}{r! s! \Gamma(\varkappa + r) \Gamma(\varrho + vs)},$$

($\varkappa, \varrho, \gamma \in \mathbb{C}, \operatorname{Re}(\varkappa), \operatorname{Re}(\varrho), \operatorname{Re}(v) > 0$), where

$${}_rL_n^{(\varkappa, \varrho)}(x, y) = x^\varkappa y^\varrho E_{\varkappa+1, \varrho+1, r}^{(-n)}(x, y).$$

And corresponding to the polynomials ${}_vH_n^\varrho(x, y)$, Özarslan and Elidemir [1] introduced a bivariate Mittag Leffler functions $E_{\varrho, v}^{(\gamma_1; \gamma_2; \gamma_3)}(x, y)$ by

$$E_{\varrho, v}^{(\gamma_1; \gamma_2; \gamma_3)}(x, y) = \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{2s} (\gamma_2)_{s+r} x^s y^{vr}}{(\gamma_3)_s \Gamma(\varrho + vr) r! s!},$$

($v, \varrho, \gamma_1, \gamma_2, \gamma_3 \in \mathbb{C}, \operatorname{Re}(\gamma_1) > 0, \operatorname{Re}(\gamma_2) > 0, \operatorname{Re}(\gamma_3) > 0, \operatorname{Re}(\varrho) > 0, \operatorname{Re}(v) > 0$), where

$${}_vH_n^\varrho(x, y) = (2x)^n E_{\varrho+1, v}^{(-n; -n; -n)}\left(\frac{-1}{4x^2}, y\right).$$

In 1903 Gösta Mittag Leffler defined the classical Mittag Leffler function. Many generalisations exist for the function $E_\varkappa(x)$ (see [7], [8], [9]). Some generalised forms consist two or more variables instead of single variable. Mittag Leffler function is the important type of the special functions, in connection with fractional calculus. After the work by Prabhakar [10], these functions appeared in the kernel of many fractional calculus operators [11]. In recent years bivariate versions of these functions have shown to have interesting applications in applied science [12]. There have been a severe interest on the investigation of different variant of bivariate Mittag Leffler functions and fractional calculus operators having these functions in the kernel especially in the recent years. (see [13], [14], [15], [16], [17], [18], [19], [20], [21], [22], [23], [24], [25], [26], [27].)

The organization of this study is as follows: In Section 2, by use of Remark1.1 we define new bivariate polynomial which is called bivariate Jacobi Konhauser polynomial and we introduce the corresponding bivariate Jacobi Konhauser Mittag-Leffler (JKML) function. In Section 3, we compute the Riemann-Liouville double fractional integral and derivative of bivariate JKML function and bivariate Jacobi Konhauser polynomials. In section 4, we obtain the double Laplace transforms and give the operational representations for bivariate JKML function and bivariate Jacobi Konhauser polynomials. In section 5, the generating functions for bivariate Jacobi Konhauser polynomials obtained and integral representations of bivariate Jacobi Konhauser polynomials and bivariate JKMLfunction derived. In section 6, we introduce an integral operator containing the bivariate JKML function in the kernel.

2 Bivariate Jacobi Konhauser Polynomials

In this section, firstly we remember some properties of Jacobi polynomials and then we define the 2D- Jacobi Konhauser polynomials.

There are several hypergeometric representations for the one variable Jacobi polynomial,

$$P_n^{(\alpha, \beta)}(x) = \frac{(1+\alpha)_n}{n!} {}_2F_1 \left[-n, 1+\alpha+\beta+n, 1+\alpha, \frac{1-x}{2} \right], \quad (2)$$

$$P_n^{(\alpha, \beta)}(x) = \frac{(1+\alpha)_n}{n!} \left(\frac{x+1}{2} \right)^n {}_2F_1 \left[-n, -\beta-n, 1+\alpha, \frac{x-1}{x+1} \right], \quad (3)$$

$$P_n^{(\alpha, \beta)}(x) = \frac{(-1)^n (1+\beta)_n}{n!} {}_2F_1 \left[-n, 1+\alpha+\beta+n, 1+\beta, \frac{1+x}{2} \right], \quad (4)$$

$$P_n^{(\alpha, \beta)}(x) = \frac{(1+\beta)_n}{n!} \left(\frac{x-1}{2} \right)^n {}_2F_1 \left[-n, -\alpha-n, 1+\beta, \frac{x+1}{x-1} \right], \quad (5)$$

where $\alpha > -1$, $\beta > -1$ and ${}_2F_1[a, b, c, x] = \sum_{s=0}^{\infty} \frac{(a)_s (b)_s x^s}{s! (c)_s}$.

The Jacobi polynomials satisfy the following orthogonality condition

$$\int_{-1}^1 (1-x)^\alpha (1+x)^\beta P_n^{(\alpha, \beta)}(x) P_m^{(\alpha, \beta)}(x) dx = \begin{cases} 0, & \text{if } n \neq m \\ \frac{2^{1+\alpha+\beta} \Gamma(1+\alpha+n) \Gamma(1+\beta+n)}{(2n+\alpha+\beta+1) \Gamma(1+\alpha+\beta+n) n!} & \text{if } n = m \end{cases}.$$

In the next definition we define the explicit form for the bivariate Jacobi Konhauser polynomials by the help of Remark 1.1. To obtain the explicit form we choose $r(x) = \frac{1-x}{2}$, $R_n(x) = Z_n^\beta(x; \kappa)$, $S_n(x) = Y_n^{(\beta)}(x; \kappa)$ and $K_n(x) = P_n^{(\alpha, \beta)}(x)$ with the hypergeometric representation (2).

Definition 2.1. The bivariate Jacobi Konhauser polynomials are defined by the following formula;

$${}_\kappa P_n^{(\alpha, \beta)}(x, y) = \frac{\Gamma(1+\alpha+n)}{n!} \sum_{s=0}^n \sum_{r=0}^{n-s} \frac{(-n)_{s+r} (1+\alpha+\beta+n)_s}{s! r! \Gamma(1+\alpha+s) \Gamma(\beta+1+\kappa r)} \left(\frac{1-x}{2} \right)^s y^{\kappa r} \quad (6)$$

where $\alpha > -1$, $\beta > -1$ and $\kappa = 1, 2, \dots$

Remark 2.1. For the bivariate Jacobi Konhauser polynomials the following representations also hold:

- choosing $R(x) = \frac{x-1}{x+1}$ and using (3) gives

$${}_\kappa P_n^{(\alpha, \beta)}(x, y) = \frac{\Gamma(1+\alpha+n)}{n!} \left(\frac{x+1}{2} \right)^n \sum_{s=0}^n \sum_{r=0}^{n-s} \frac{(-n)_{s+r} (-\beta-n)_s}{s! r! \Gamma(1+\alpha+s) \Gamma(\beta+1+\kappa r)} \left(\frac{x-1}{x+1} \right)^s y^{\kappa r}; \quad (7)$$

- choosing $R(x) = \frac{1+x}{2}$ and using (4) gives

$${}_\kappa P_n^{(\alpha, \beta)}(x, y) = \frac{(-1)^n \Gamma(1+\beta+n)}{n!} \sum_{s=0}^n \sum_{r=0}^{n-s} \frac{(-n)_{s+r} (1+\alpha+\beta+n)_s}{s! r! \Gamma(1+\beta+s) \Gamma(\beta+1+\kappa r)} \left(\frac{1+x}{2} \right)^s y^{\kappa r}; \quad (8)$$

- choosing $R(x) = \frac{x+1}{x-1}$ and using (5) gives

$${}_\kappa P_n^{(\alpha, \beta)}(x, y) = \frac{\Gamma(1+\beta+n)}{n!} \left(\frac{x-1}{2} \right)^n \sum_{s=0}^n \sum_{r=0}^{n-s} \frac{(-n)_{s+r} (-\alpha-n)_s}{s! r! \Gamma(1+\beta+s) \Gamma(\beta+1+\kappa r)} \left(\frac{x+1}{x-1} \right)^s y^{\kappa r}; \quad (9)$$

Theorem 2.1. Bivariate Jacobi Konhauser polynomials, ${}_\kappa P_n^{(\alpha, \beta)}(x, y)$, can be written in the following form:

$${}_\kappa P_n^{(\alpha, \beta)}(x, y) = \Gamma(1+\alpha+n) \sum_{s=0}^n \frac{(-1)^s (1+\alpha+\beta+n)_s}{s! \Gamma(1+\alpha+s) \Gamma(1+\beta+\kappa n - \kappa s)} \left(\frac{1-x}{2} \right)^s Z_{n-s}^\beta(y; \kappa). \quad (10)$$

Proof. Proof directly follows from the representations of the polynomials ${}_\kappa P_n^{(\alpha, \beta)}(x, y)$ and $Z_n^\beta(x; \kappa)$. \square

Theorem 2.2. *Bivariate Jacobi Konhauser polynomials satisfy the following biorthogonality condition;*

$$\int_0^\infty \int_{-1}^1 (1-x)^\alpha (1+x)^\beta e^{-y} y^\beta {}_\kappa P_n^{(\alpha, \beta)}(x, y) Q_m(x, y) dx dy = \begin{cases} 0, & \text{if } n \neq m \\ \frac{2^{1+\alpha+\beta} \Gamma(1+\alpha+n) \Gamma(1+\beta+n)}{(2n+\alpha+\beta+1) \Gamma(1+\alpha+\beta+n) n!} & \text{if } n = m \end{cases}, \quad (11)$$

where

$$Q_m(x, y) = P_m^{(\alpha, \beta)}(x) \sum_{j=0}^m Y_j^{(\beta)}(y; \kappa). \quad (12)$$

Proof. Replacing ${}_\kappa P_n^{(\alpha, \beta)}(x, y)$ from (10) and $Q_m(x, y)$ from (12) in the left side, we get

$$\begin{aligned} & \int_0^\infty \int_{-1}^1 (1-x)^\alpha (1+x)^\beta e^{-y} y^\beta \Gamma(1+\alpha+n) \sum_{s=0}^n \frac{(-1)^s (1+\alpha+\beta+n)_s}{s! \Gamma(1+\alpha+s) \Gamma(1+\beta+\kappa n - \kappa s)} \left(\frac{1-x}{2}\right)^s Z_{n-s}^\beta(y; \kappa) \\ & P_m^{(\alpha, \beta)}(x, y) \sum_{j=0}^m Y_j^{(\beta)}(y; \kappa) dx dy \\ &= \int_{-1}^1 (1-x)^\alpha (1+x)^\beta \Gamma(1+\alpha+n) \sum_{s=0}^n \frac{(-1)^s (1+\alpha+\beta+n)_s}{s! \Gamma(1+\alpha+s) (n-s)!} \left(\frac{1-x}{2}\right)^s P_m^{(\alpha, \beta)}(x, y) dx \\ &= \int_{-1}^1 (1-x)^\alpha (1+x)^\beta P_n^{(\alpha, \beta)}(x, y) P_m^{(\alpha, \beta)}(x, y) dx. \end{aligned}$$

Whence we obtain the result from the orthogonality relation of the Jacobi polynomials. \square

From the view of the definition of the Kampe de Fariet's double hypergeometric series:

$$F_{k,l,m}^{n,p,q} \left[\begin{matrix} (a_n) : (b_p); (c_q); \\ (d_k) : (f_l); (g_m); \end{matrix} ; x, y \right] = \sum_{r,s=0}^{\infty} \frac{\prod_{i=1}^n (a_i)_{r+s} \prod_{i=1}^p (b_i)_r \prod_{i=1}^q (c_i)_s x^s y^r}{\prod_{i=1}^k (d_i)_{r+s} \prod_{i=1}^l (f_i)_r \prod_{i=1}^m (g_i)_s s! r!},$$

we obtain the following series representation for the polynomials ${}_\kappa P_n^{(\alpha, \beta)}(x, y)$.

Theorem 2.3. *The following Kampe de Fariet's double hypergeometric representation holds true for bivariate Jacobi Konhauser polynomials:*

$${}_\kappa P_n^{(\alpha, \beta)}(x, y) = \frac{(1+\alpha)_n}{\Gamma(1+\beta)n!} F_{0,1,\kappa}^{1,1,0} \left[\begin{matrix} -n : (1+\alpha+\beta+n); -; \\ - : (1+\alpha); \Delta(\kappa; \beta+1); \end{matrix} ; \frac{1-x}{2}; \left(\frac{y}{\kappa}\right)^\kappa \right], \quad (13)$$

where $\Delta(k; \sigma)$ denotes the k parameters $\frac{\sigma}{k}, \frac{\sigma+1}{k}, \dots, \frac{\sigma+k-1}{k}$.

Proof. By multiplying explicit form of bivariate Jacobi Konhauser polynomials with $\frac{\Gamma(1+\alpha)\Gamma(1+\beta)}{\Gamma(1+\alpha)\Gamma(1+\beta)}$ we get

$${}_\kappa P_n^{(\alpha, \beta)}(x, y) = \frac{(1+\alpha)_n}{\Gamma(1+\beta)n!} \sum_{s=0}^n \sum_{r=0}^{n-s} \frac{(-n)_{s+r} (1+\alpha+\beta+n)_s}{s! r! (1+\alpha)_s (\beta+1)_{\kappa r}} \left(\frac{1-x}{2}\right)^s y^{\kappa r}, \quad (14)$$

where $(1+\beta)_{\kappa r} = \kappa^{\kappa r} \prod_{j=0}^{\kappa-1} \left(\frac{\beta+1+j}{\kappa}\right)_r$.

Whence we obtain the result. \square

Corollary 2.1. *From the explicit forms of bivariate Jacobi Konhauser polynomials (7), (8), (9) the following Kampe de Fariet's double hypergeometric representations are also hold respectively,*

$${}_\kappa P_n^{(\alpha, \beta)}(x, y) = \frac{(-1)^n (1+\beta)_n}{n!} F_{0,1,\kappa}^{1,1,0} \left[\begin{matrix} -n : (1+\alpha+\beta+n); -; \\ - : (1+\beta); \Delta(\kappa; \beta+1); \end{matrix} ; \frac{1+x}{2}; \left(\frac{y}{\kappa}\right)^\kappa \right], \quad (15)$$

$${}_{\kappa}P_n^{(\alpha,\beta)}(x,y) = \frac{(1+\alpha)_n}{\Gamma(1+\beta)n!} \left(\frac{x+1}{2}\right)^n F_{0,1,\kappa}^{1,1,0} \left[\begin{matrix} -n : (-\beta-n); -; \\ - : (1+\alpha); \Delta(\kappa; \beta+1); \end{matrix} \right. \left. \frac{x-1}{x+1}; \left(\frac{y}{\kappa}\right)^\kappa \right], \quad (16)$$

$${}_{\kappa}P_n^{(\alpha,\beta)}(x,y) = \frac{(1+\beta)_n}{n!} \left(\frac{x-1}{2}\right)^n F_{0,1,\kappa}^{1,1,0} \left[\begin{matrix} -n : (-\alpha-n); -; \\ - : (1+\beta); \Delta(\kappa; \beta+1); \end{matrix} \right. \left. \frac{x+1}{x-1}; \left(\frac{y}{\kappa}\right)^\kappa \right], \quad (17)$$

Definition 2.2. Bivariate Jacobi Konhauser Mittag Leffler functions $E_{\alpha,\beta,\kappa}^{(\gamma_1;\gamma_2)}(x,y)$ are defined by the formula,

$$E_{\alpha,\beta,\kappa}^{(\gamma_1;\gamma_2)}(x,y) = \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{r+s}(\gamma_2)_s x^s y^{\kappa r}}{r!s!\Gamma(\alpha+s)\Gamma(\beta+\kappa r)}, \quad (\alpha, \beta, \gamma_1, \gamma_2 \in \mathbb{C}, Re(\alpha), Re(\beta), Re(\kappa) > 0). \quad (18)$$

Remark 2.2. According to the convergence conditions for the double series by using ratio test the series in (18) converge absolutely for $Re(\kappa) > 0$.

Corollary 2.2. Comparing (6) and (18) it easily seen that;

$${}_{\kappa}P_n^{(\alpha,\beta)}(x,y) = \frac{\Gamma(1+\alpha+n)}{n!} E_{\alpha+1,\beta+1,\kappa}^{(-n;1+\alpha+\beta+n)}\left(\frac{1-x}{2}, y\right). \quad (19)$$

3 Fractional Calculus for $E_{\alpha,\beta,\kappa}^{(\gamma_1;\gamma_2)}(x,y)$ and ${}_{\kappa}P_n^{(\alpha,\beta)}(x,y)$

In this section for the bivariate JKML function $E_{\alpha,\beta,\kappa}^{(\gamma_1;\gamma_2)}(x,y)$ and for the polynomial ${}_{\kappa}P_n^{(\alpha,\beta)}(x,y)$, we compute their images under the action of the Riemann-Liouville double fractional integral and derivative.

Definition 3.1. The Riemann-Liouville fractional integral of order $\zeta \in \mathbb{C}$ ($Re(\zeta) > 0$) is defined by

$$({}_x I_{a+}^{\zeta} g)(x) = \frac{1}{\Gamma(\zeta)} \int_a^x (x-t)^{\zeta-1} g(t) dt \quad (x > a),$$

The double fractional integral of the function $g(x,y)$ is defined by

$$({}_y I_{b+}^{\zeta} {}_x I_{a+}^{\mu} g)(x,y) = \frac{1}{\Gamma(\zeta)\Gamma(\mu)} \int_b^y \int_a^x (x-\epsilon)^{\mu-1} (y-\tau)^{\zeta-1} g(\epsilon,\tau) d\epsilon d\tau. \quad (20)$$

($x > a, y > b, Re(\zeta) > 0, Re(\mu) > 0$)

Definition 3.2. The Riemann-Liouville fractional derivative of order $\zeta \in \mathbb{C}$ ($Re(\zeta) > 0$) is defined by

$$({}_x D_{a+}^{\zeta} g)(x) = \left(\frac{d}{dx}\right)^n ({}_x I_{a+}^{n-\zeta} g)(x), \quad (x > a, n = [Re(\zeta)] + 1)$$

where $[Re(\zeta)]$ is the integral part of $Re(\zeta)$.

The partial Riemann-Liouville fractional derivative define with the following equation,

$$({}_x D_{a+}^{\zeta} {}_y D_{b+}^{\tau} g)(x,y) = \frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m} ({}_x I_{a+}^{n-\zeta} {}_y I_{b+}^{m-\tau} g)(x,y).$$

($x > a, y > b, m = [Re(\tau)] + 1, n = [Re(\zeta)] + 1$)

Theorem 3.1. Bivariate JKML function have the following double fractional integral representation;

$$\begin{aligned} & \left({}_y I_{b+}^{\zeta} {}_x I_{a+}^{\mu} \right) \left[(x-a)^{\alpha} (y-b)^{\beta} E_{\alpha+1,\beta+1,\kappa}^{(\gamma_1;\gamma_2)}(\omega_1(x-a), \omega_2(y-b)) \right] \\ & = (x-a)^{\alpha+\mu} (y-b)^{\beta+\zeta} E_{1+\alpha+\mu,1+\beta+\zeta,\kappa}^{(\gamma_1;\gamma_2)}(\omega_1(x-a), \omega_2(y-b)). \end{aligned}$$

Proof. By replacing $E_{\alpha+1,\beta+1,\kappa}^{(\gamma_1;\gamma_2)}(\omega_1(x-a), \omega_2(y-b))$ with its explicit form in the left hand side and interchanging the order of series and fractional integral operators, we obtain,

$$\begin{aligned} & \left({}_y I_{b^+}^\zeta {}_x I_{a^+}^\mu \right) \left[(x-a)^\alpha (y-b)^\beta E_{\alpha+1,\beta+1,\kappa}^{(\gamma_1;\gamma_2)}(\omega_1(x-a), \omega_2(y-b)) \right] \\ &= \frac{1}{\Gamma(\zeta)\Gamma(\mu)} \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{s+r}(\gamma_2)_s \omega_1^s \omega_2^{\kappa r}}{s!r!\Gamma(\alpha+s+1)\Gamma(\beta+\kappa r+1)} \int_a^x (x-t)^{\mu-1} (t-a)^{\alpha+s} dt \int_b^y (y-\tau)^{\zeta-1} (\tau-b)^{\beta+\kappa r} d\tau \\ &= (x-a)^{\alpha+\mu} (y-b)^{\beta+\zeta} \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{s+r}(\gamma_2)_s [\omega_1(x-a)]^s [\omega_2(y-b)]^{\kappa r}}{s!r!\Gamma(\alpha+\mu+s+1)\Gamma(\beta+\zeta+\kappa r+1)} \end{aligned}$$

whence we get the result. \square

Corollary 3.1. *Bivariate Jacobi Konhauser polynomials have the following double fractional integral representation;*

$$\begin{aligned} & \left({}_y I_{b^+}^\zeta {}_x I_{a^+}^\mu \right) \left[(x-a)^\alpha (y-b)^\beta {}_\kappa P_n^{(\alpha,\beta)}(1-2\omega_1(x-a), \omega_2(y-b)) \right] \\ &= \frac{\Gamma(1+\alpha+n)}{\Gamma(1+\alpha+n+\mu)} (x-a)^{\alpha+\mu} (y-b)^{\beta+\zeta} {}_\kappa P_n^{(\alpha+\mu,\beta+\zeta)}(1-2\omega_1(x-a), \omega_2(y-b)) \end{aligned}$$

Proof. Proof follows from (19) by using Theorem 3.1. \square

Theorem 3.2. *Bivariate JKML function have the following partial fractional derivative representation;*

$$\begin{aligned} & \left({}_y D_{b^+}^\zeta {}_x D_{a^+}^\mu \right) \left[(x-a)^\alpha (y-b)^\beta E_{\alpha+1,\beta+1,\kappa}^{(\gamma_1;\gamma_2)}(\omega_1(x-a), \omega_2(y-b)) \right] \\ &= (x-a)^{\alpha-\mu} (y-b)^{\beta-\zeta} E_{1+\alpha-\mu,1+\beta-\zeta,\kappa}^{(\gamma_1;\gamma_2)}(\omega_1(x-a), \omega_2(y-b)) \end{aligned}$$

Proof. By replacing $E_{\alpha+1,\beta+1,\kappa}^{(\gamma_1;\gamma_2)}(\omega_1(x-a), \omega_2(y-b))$ with its explicit form in the left hand side and interchanging the order of series and fractional integral operators we obtain,

$$\begin{aligned} & \left({}_y D_{b^+}^\zeta {}_x D_{a^+}^\mu \right) \left[(x-a)^\alpha (y-b)^\beta E_{\alpha+1,\beta+1,\kappa}^{(\gamma_1;\gamma_2)}(\omega_1(x-a), \omega_2(y-b)) \right] \\ &= \frac{1}{\Gamma(m-\zeta)\Gamma(n-\mu)} \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{s+r}(\gamma_2)_s \omega_1^s \omega_2^{\kappa r}}{s!r!\Gamma(\alpha+s+1)\Gamma(\beta+\kappa r+1)} \\ & D_x^n \int_a^x (x-t)^{n-\mu-1} (t-a)^{\alpha+s} dt D_y^m \int_b^y (y-\tau)^{m-\zeta-1} (\tau-b)^{\beta-1+\kappa r} d\tau \\ &= (x-a)^{\alpha-\mu} (y-b)^{\beta-\zeta} \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{s+r}(\gamma_2)_s [\omega_1(x-a)]^s [\omega_2(y-b)]^{\kappa r}}{s!r!\Gamma(\alpha-\mu+s+1)\Gamma(\beta-\zeta+\kappa r+1)} \end{aligned}$$

whence we get the result. \square

Corollary 3.2. *Bivariate Jacobi Konhauser polynomials have the following partial fractional derivative representation;*

$$\begin{aligned} & \left({}_y D_{b^+}^\zeta {}_x D_{a^+}^\mu \right) \left[(x-a)^\alpha (y-b)^\beta {}_\kappa P_n^{(\alpha,\beta)}(1-2\omega_1(x-a), \omega_2(y-b)) \right] \\ &= \frac{\Gamma(1+\alpha+n)}{\Gamma(1+\alpha+n-\mu)} (x-a)^{\alpha-\mu} (y-b)^{\beta-\zeta} {}_\kappa P_n^{(\alpha-\mu,\beta-\zeta)}(1-2\omega_1(x-a), \omega_2(y-b)) \end{aligned}$$

Proof. Proof follows from (19) by using Theorem 3.2. \square

4 Laplace transform and Operational representations for $E_{\alpha,\beta,\kappa}^{(\gamma_1;\gamma_2)}(x,y)$ and ${}_n P_n^{(\alpha,\beta)}(x,y)$

In this section, firstly we obtain the double Laplace transforms and then give the operational representations for $E_{\alpha,\beta,\kappa}^{(\gamma_1;\gamma_2)}(x,y)$ and ${}_n P_n^{(\alpha,\beta)}(x,y)$. And also we write bivariate Jacobi Konhauser polynomials in terms of Bessel functions.

Now, recall the bivariate Laplace transform

$$\mathbb{L}_2[h(\bar{x}, \bar{y})](\bar{p}, \bar{q}) = \int_0^\infty \int_0^\infty e^{-(\bar{p}\bar{x} + \bar{q}\bar{y})} h(\bar{x}, \bar{y}) d\bar{x} d\bar{y}, \quad (Re(\bar{p}), Re(\bar{q}) > 0).$$

Theorem 4.1. For $\alpha, \beta, \gamma_1, \gamma_2, \kappa \in \mathbb{C}$, $Re(\alpha), Re(\beta), Re(\gamma_1), Re(\gamma_2), Re(w_1), Re(w_2), Re(p_1), Re(p_2) > 0$, $|\frac{w_2^\kappa}{p_2^\kappa}| < 1$ and $|\frac{w_1 p_2^\kappa}{p_1(p_2^\kappa - w_2^\kappa)}| < 1$ the following double Laplace transformation holds

$$\mathbb{L}_2[x^\alpha y^\beta E_{\alpha+1,\beta+1,\kappa}^{(\gamma_1;\gamma_2)}(w_1 x, w_2 y)](p_1, p_2) = p_1^{-(1+\alpha)} p_2^{-(1+\beta)} \left(1 - \frac{w_2^\kappa}{p_2^\kappa}\right)^{-\gamma_1} {}_2F_0\left[\gamma_1, \gamma_2; -; \frac{w_1 p_2^\kappa}{p_1(p_2^\kappa - w_2^\kappa)}\right].$$

Proof. Because of the hypothesis of the Theorem, we have a right to interchange the order of series and fractional integral operators, which yields

$$\begin{aligned} & \mathbb{L}_2[x^\alpha y^\beta E_{\alpha+1,\beta+1,\kappa}^{(\gamma_1;\gamma_2)}(w_1 x, w_2 y)](p_1, p_2) \\ &= \sum_{s=0}^\infty \sum_{r=0}^\infty \frac{(\gamma_1)_{s+r} (\gamma_2)_s \omega_1^s \omega_2^{\kappa r}}{s! r! \Gamma(\alpha + s + 1) \Gamma(\beta + \kappa r + 1)} \int_0^\infty \int_0^\infty e^{-p_1 x - p_2 y} x^{\alpha+s} y^{\beta+\kappa r} dx dy \\ &= \sum_{s=0}^\infty \sum_{r=0}^\infty \frac{(\gamma_1)_{s+r} (\gamma_2)_s \omega_1^s \omega_2^{\kappa r}}{s! r! \Gamma(\alpha + s + 1) \Gamma(\beta + \kappa r + 1) p_1^{1+\alpha} p_2^{1+\beta}} \int_0^\infty \int_0^\infty e^{-u-v} u^{\alpha+s} v^{\beta+\kappa r} du dv \\ &= \frac{1}{p_1^{1+\alpha} p_2^{1+\beta}} \sum_{s=0}^\infty \frac{(\gamma_1)_s (\gamma_2)_s}{s!} \left(\frac{w_1}{p_1}\right)^s \sum_{r=0}^\infty \frac{(\gamma_1 + s)_r}{r!} \left(\frac{w_2}{p_2}\right)^{\kappa r} \\ &= p_1^{-(1+\alpha)} p_2^{-(1+\beta)} \left(1 - \frac{w_2^\kappa}{p_2^\kappa}\right)^{-\gamma_1} \sum_{s=0}^\infty \frac{(\gamma_1)_s (\gamma_2)_s}{s!} \left(\frac{w_1 p_2^\kappa}{p_1(p_2^\kappa - w_2^\kappa)}\right)^s \end{aligned}$$

Whence the result. □

Corollary 4.1. For the bivariate Jacobi Konhauser polynomials ${}_n P_n^{(\alpha,\beta)}(x,y)$, we have

$$\mathbb{L}_2[x^\alpha y^\beta {}_n P_n^{(\alpha,\beta)}(1 - 2w_1 x, w_2 y)](p_1, p_2) = \frac{\Gamma(1 + \alpha + n)}{n! p_1^{1+\alpha} p_2^{1+\beta}} \left(1 - \frac{w_2^\kappa}{p_2^\kappa}\right)^n {}_2F_0\left[-n, 1 + \alpha + \beta + n; -; \frac{w_1 p_2^\kappa}{p_1(p_2^\kappa - w_2^\kappa)}\right]$$

Theorem 4.2. For the bivariate JKML functions $E_{\alpha,\beta,\kappa}^{(\gamma_1;\gamma_2)}(x,y)$ the following operational representation holds

$$E_{\alpha,\beta,\kappa}^{(\gamma_1;\gamma_2)}(x,y) = x^{1-\alpha} y^{1-\beta} \sum_{s=0}^\infty \sum_{r=0}^\infty \frac{(\gamma_1)_{s+r} (\gamma_2)_s}{s! r!} D_x^{-s} D_y^{-\kappa r} \left\{ \frac{x^{\alpha-1} y^{\beta-1}}{\Gamma(\alpha) \Gamma(\beta)} \right\},$$

where D_x^{-1} is the inverse operator of the derivative.

Proof. Proof directly follows from the explicit form of the bivariate JKML functions. □

Corollary 4.2. For the bivariate JKML functions $E_{\alpha,\beta,\kappa}^{(\gamma_1;\gamma_2)}(x,y)$ the following operational representation also holds

$$E_{\alpha,\beta,\kappa}^{(\gamma_1;\gamma_2)}(x,y) = x^{1-\alpha} y^{1-\beta} \left[\frac{1}{1 - D_y^{-\kappa}} \right]^{-\gamma_1} {}_2F_0\left[\gamma_1, \gamma_2; -; \frac{1}{D_x(1 - D_y^{-\kappa})}\right] \left\{ \frac{x^{\alpha-1} y^{\beta-1}}{\Gamma(\alpha) \Gamma(\beta)} \right\}.$$

Theorem 4.3. For the bivariate Jacobi Konhauser polynomials the following operational representation hold

$$\begin{aligned} {}_{\kappa}P_n^{(\alpha,\beta)}(x,y) &= \frac{\Gamma(1+\alpha+n)}{n!} \left(\frac{1-x}{2}\right)^{-\alpha} y^{-\beta} (1-D_y^{-\kappa})^n {}_2F_0 \left[-n, 1+\alpha+\beta+n; -; \frac{-2}{D_x(1-D_y^{-\kappa})} \right] \\ &\times \left\{ \frac{\left(\frac{1-x}{2}\right)^{\alpha} y^{\beta}}{\Gamma(\alpha+1)\Gamma(\beta+1)} \right\} \\ &= \frac{\Gamma(1+\alpha+n)}{n!} \left(\frac{1-x}{2}\right)^{-\alpha} (1-D_y^{-\kappa})^n y_n \left(\frac{1}{2D_x(1-D_y^{-\kappa})}; 2+\alpha+\beta, 1 \right) \\ &\times \left\{ \frac{\left(\frac{1-x}{2}\right)^{\alpha} y^{\beta}}{\Gamma(\alpha+1)\Gamma(\beta+1)} \right\}, \end{aligned}$$

where $y_n(x; a, b) = {}_2F_0[-n, a-1+n; -, \frac{-x}{b}]$ are the Bessel polynomials.

5 Various properties of ${}_{\kappa}P_n^{(\alpha,\beta)}(x, y)$ and $E_{\alpha,\beta,\kappa}^{(\gamma_1;\gamma_2)}(x, y)$

In this section, firstly we obtain a generating function for the bivariate Jacobi Konhauser polynomials by means of Kampe de Ferit's double hypergeometric functions. Then we obtain Schalfi's contour integral representations of $E_{\alpha,\beta,\kappa}^{(\gamma_1;\gamma_2)}(x, y)$ and ${}_{\kappa}P_n^{(\alpha,\beta)}(x, y)$. The integral representations for the product of $E_{\alpha,\beta,\kappa}^{(\gamma_1;\gamma_2)}(x, y)$, and of course, in its special case to ${}_{\kappa}P_n^{(\alpha,\beta)}(x, y)$ are also derived.

From the view of the definition of the double hypergeometric series [34]

$$\begin{aligned} S_{C:D;D'}^{A:B;B'} &\left[\begin{matrix} [(a) : \theta, \phi] : [(b) : \psi] : [(b') : \psi']; \\ [(c) : \delta, \epsilon] : [(d) : \xi] : [(d') : \xi']; \end{matrix} \right. \\ &\left. \begin{matrix} x; y \end{matrix} \right] \\ &= \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{\prod_{j=0}^A \Gamma(a_j + s\theta + r\phi) \prod_{j=0}^B \Gamma(b_j + s\psi) \prod_{j=0}^{B'} \Gamma(b'_j + r\psi')}{\prod_{j=0}^C \Gamma(c_j + s\delta + r\epsilon) \prod_{j=0}^D \Gamma(d_j + s\xi) \prod_{j=0}^{D'} \Gamma(d'_j + r\xi')} \frac{x^s y^r}{s!r!}. \end{aligned}$$

we obtain the following generating function for the polynomials ${}_{\kappa}P_n^{(\alpha,\beta)}(x, y)$.

Theorem 5.1. Bivariate Jacobi Konhauser polynomials have the following generating function

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(1+\alpha+\beta)_n \Gamma(\beta+1)}{(1+\alpha)_n} {}_{\kappa}P_n^{(\alpha,\beta)}(x, y) t^n &= \frac{1}{(1-t)^{1+\alpha+\beta}} \frac{\Gamma(1+\alpha)\Gamma(1+\beta)}{\Gamma(1+\alpha+\beta)} \\ &\times S_{0,1,1}^{1,0,0} \left[\begin{matrix} [1+\alpha+\beta : 2, 1] : - : -; \\ - : \alpha+1 : 1] : [\beta+1 : \kappa]; \end{matrix} \right. \\ &\left. \frac{t(x-1)}{2(1-t)^2}; \frac{ty^{\kappa}}{(1-t)\kappa^{\kappa}} \right]. \end{aligned} \quad (21)$$

Proof. In left hand side by replacing ${}_{\kappa}P_n^{(\alpha,\beta)}(x, y)$ with (6) and using the properties of series we get,

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(1+\alpha+\beta)_n \Gamma(\beta+1)}{(1+\alpha)_n} {}_{\kappa}P_n^{(\alpha,\beta)}(x, y) t^n &= \sum_{n=0}^{\infty} \sum_{s=0}^n \sum_{r=0}^{n-s} \frac{(-n+s)_r (1+\alpha+\beta)_{n+s}}{(n-s)!s!r!(1+\alpha)_s(1+\beta)_{\kappa r}} \left(\frac{x-1}{2}\right)^2 y^{\kappa r} t^n \\ &= \sum_{n=0}^{\infty} \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(1+\alpha+\beta)_{n+2s+r}}{s!r!(1+\alpha)_s(1+\beta)_{\kappa r}} \left(\frac{tx-t}{2}\right)^2 (-ty)^{\kappa r} t^n. \end{aligned}$$

Since $(1+\alpha+\beta)_{n+2s+r} = (1+\alpha+\beta)_{2s+r} (1+\alpha+\beta+2s+r)_n$ and $\sum_{n=0}^{\infty} \frac{(1+\alpha+\beta+2s+r)_n t^n}{n!} = (1-t)^{-(1+\alpha+\beta+2s+r)}$ we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(1+\alpha+\beta)_n \Gamma(\beta+1)}{(1+\alpha)_n} {}_{\kappa}P_n^{(\alpha,\beta)}(x, y) t^n &= \frac{1}{(1-t)^{1+\alpha+\beta}} \\ &\times \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(1+\alpha+\beta)_{2s+r}}{s!r!(1+\alpha)_s(1+\beta)_{\kappa r}} \left(\frac{t(x-1)}{2(1-t)^2}\right)^s \left(\frac{ty^{\kappa}}{(1-t)\kappa^{\kappa}}\right)^r. \end{aligned}$$

Consequently, we derive the result. \square

Before presenting the integral representations, it is necessary to recall the integral representations of the Gamma function, where the second relation is referred to as Hankel's representation [33].

$$\Gamma(z) = \int_0^\infty e^{-u} u^{z-1} du \quad (\operatorname{Re}(z) > 0), \quad (22)$$

$$\frac{1}{\Gamma(z)} = \frac{1}{2\pi i} \int_{-\infty}^{0+} e^u u^{-z} du \quad (\arg|u| \leq \pi). \quad (23)$$

Theorem 5.2. *The following integral representation holds for the bivariate JKML functions*

$$E_{\alpha, \beta, \kappa}^{(\gamma_1; \gamma_2)}(x, y) = \frac{-1}{4\pi^2 \Gamma(\gamma_2)} \int_0^\infty \int_{-\infty}^{0+} \int_{-\infty}^{0+} t^{\gamma_2-1} u^{-\alpha} w^{-\beta} e^{u+w-t} \left[1 - \left(\frac{y}{w} \right)^\kappa - \frac{tx}{u} \right]^{-\gamma_1} dw du dt.$$

$(\alpha, \beta, \kappa, \gamma_1, \gamma_2 \in \mathbb{C}, \operatorname{Re}(\alpha), \operatorname{Re}(\beta), \operatorname{Re}(\kappa) > 0, |\frac{y^\kappa}{w^\kappa}| < 1 \text{ and } |\frac{txw^\kappa}{u(w^\kappa - y^\kappa)}| < 1)$

Proof. Using formulas (22), (23) and the relation $(a)_{s+r} = (a)_s (a+s)_r$ we obtain that

$$\begin{aligned} E_{\alpha, \beta, \kappa}^{(\gamma_1; \gamma_2)}(x, y) &= \frac{-1}{4\pi^2 \Gamma(\gamma_2)} \int_0^\infty \int_{-\infty}^{0+} \int_{-\infty}^{0+} t^{\gamma_2-1} u^{-\alpha} w^{-\beta} e^{u+w-t} \sum_{s=0}^\infty \frac{(\gamma_1)_s}{s!} \left(\frac{tx}{u} \right)^s \sum_{r=0}^\infty \frac{(\gamma_1+s)_r}{r!} \left(\frac{y}{w} \right)^{\kappa r} dw du dt \\ &= \frac{-1}{4\pi^2 \Gamma(\gamma_2)} \int_0^\infty \int_{-\infty}^{0+} \int_{-\infty}^{0+} t^{\gamma_2-1} u^{-\alpha} w^{-\beta} e^{u+w-t} \left[1 - \left(\frac{y}{w} \right)^\kappa \right]^{-\gamma_1} \sum_{s=0}^\infty \frac{(\gamma_1)_s}{s!} \left(\frac{txw^\kappa}{u(w^\kappa - y^\kappa)} \right)^s. \end{aligned}$$

Whence the result. \square

Corollary 5.1. *The following integral representation holds for the bivariate Jacobi Konhauser polynomials,*

$$\begin{aligned} {}_\kappa P_n^{(\alpha, \beta)}(x, y) &= \frac{-\Gamma(1+\alpha+n)}{4\pi^2 \Gamma(1+\alpha+\beta+n)n!} \int_0^\infty \int_{-\infty}^{0+} \int_{-\infty}^{0+} t^{\alpha+\beta+n} u^{-(1+\alpha)} w^{(1+\beta)} e^{u+w-t} \\ &\quad \times \left[1 - \left(\frac{y}{w} \right)^\kappa - \frac{t}{u} \left(\frac{1-x}{2} \right) \right]^n dw du dt. \end{aligned}$$

Now, we state a double integral representation for the product of $E_{\alpha_1, \beta_1, \kappa}^{(\gamma_1; \gamma_2)}(x, y)$. We will use the following integral representation for the proof of the following theorem.

$$B(\epsilon, \xi) = \frac{\Gamma(\epsilon)\Gamma(\xi)}{\Gamma(\epsilon+\xi)} = \int_0^1 u^{\epsilon-1} (1-u)^{\xi-1} \quad (24)$$

$(\min\{\operatorname{Re}(\epsilon), \operatorname{Re}(\xi)\} > 0)$

Theorem 5.3. *The following integral representation holds true for the multiplication of the bivariate JKML functions*

$$\begin{aligned} E_{\alpha_1, \beta_1, \kappa}^{(\gamma_1; \gamma_2)}(x, y) E_{\alpha_2, \beta_2, \kappa}^{(\sigma_1; \sigma_2)}(x, y) &= \frac{1}{16\pi^4 \Gamma(\gamma_1)\Gamma(\gamma_2)\Gamma(\sigma_1)\Gamma(\sigma_2)} \int_0^1 t^{\gamma_1} (1-t)^{\sigma_1-1} \int_0^1 u^{\gamma_2} (1-u)^{\sigma_2-1} \\ &\quad \times \left(\int_{-\infty}^{0+} \int_{-\infty}^{0+} \int_{-\infty}^{0+} \int_{-\infty}^{0+} e^{\epsilon+\delta+w+\xi} \epsilon^{-\alpha_1} \delta^{-\beta_1} \xi^{-\alpha_2} w^{-\beta_2} \right. \\ &\quad \times S_{0,0,0}^{1,1,0} \left[\begin{matrix} [\gamma_1 + \sigma_1 : 1 : 1] : [\gamma_2 + \sigma_2 : 1 : 0] : -; \\ - : - : -; \end{matrix} \right. \\ &\quad \left. \left. x[(1-t)(1-u)\xi^{-1} + t\epsilon^{-1}]; y^\kappa[(1-t)w^{-\kappa} + t\delta^{-\kappa}] \right] \right. \\ &\quad \left. d\epsilon d\sigma dw d\xi \right) du dt, \end{aligned}$$

$(|\arg(\epsilon)|, |\arg(\sigma)|, |\arg(w)|, |\arg(\xi)| \leq \pi; \min\{\operatorname{Re}(\alpha_1), \operatorname{Re}(\alpha_2)\} > 0; \min\{\operatorname{Re}(\kappa), \operatorname{Re}(\gamma_1), \operatorname{Re}(\gamma_2), \operatorname{Re}(\sigma_1), \operatorname{Re}(\sigma_2)\} > 0)$

Proof. By using the definition (18), we obtain

$$\begin{aligned} E_{\alpha_1, \beta_1, \kappa}^{(\gamma_1; \gamma_2)}(x, y) E_{\alpha_2, \beta_2, \kappa}^{(\sigma_1; \sigma_2)}(x, y) &= \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{r+s} (\gamma_2)_s x^r y^{\kappa s}}{r! s! \Gamma(\alpha_1 + r) \Gamma(\beta_1 + \kappa s)} \sum_{p=0}^{\infty} \sum_{q=0}^{\infty} \frac{(\sigma_1)_{q+p} (\sigma_2)_p x^q y^{\kappa p}}{p! q! \Gamma(\alpha_2 + q) \Gamma(\beta_2 + \kappa p)} \\ &= \frac{1}{\Gamma(\gamma_1) \Gamma(\gamma_2) \Gamma(\sigma_1) \Gamma(\sigma_2)} \sum_{p=0}^{\infty} \sum_{q=0}^{\infty} \frac{\Gamma(\gamma_1 + \sigma_1 + p + q) \Gamma(\gamma_2 + \sigma_2 + q) x^q y^{\kappa p}}{p! q!} \\ &\quad \times \sum_{r=0}^q \sum_{s=0}^p \binom{q}{r} \binom{p}{s} \frac{B(\gamma_1 + s + r, \sigma_1 + p + q - s - r) B(\gamma_2 + r, \sigma_2 + q - r)}{\Gamma(\alpha_1 + r) \Gamma(\beta_1 + \kappa s) \Gamma(\alpha_2 + q - r) \Gamma(\beta_2 + \kappa(p - s))} \end{aligned}$$

in terms of Beta function $B(\epsilon, \xi)$ defined by (24).

Now, by applying the integral formulas (23) and (24) we derive the result of the theorem. \square

Corollary 5.2. *The following integral representation holds true*

$$\begin{aligned} {}_{\kappa}P_n^{(\alpha_1, \beta_1)}(x, y) {}_{\kappa}P_m^{(\alpha_2, \beta_2)}(x, y) &= \frac{\Gamma(1 + \alpha_1 + n) \Gamma(1 + \alpha_2 + m)}{16\pi^4 \Gamma(1 + \alpha_1 + \beta_1 + n) \Gamma(1 + \alpha_2 + \beta_2 + m)} \int_0^{\infty} \int_0^{\infty} e^{-t-u} t^{\alpha_1 + \beta_1 + n} u^{\alpha_2 + \beta_2 + m} \\ &\quad \times \left(\int_{-\infty}^{0+} \int_{-\infty}^{0+} \int_{-\infty}^{0+} \int_{-\infty}^{0+} e^{\epsilon + \delta + w + \xi} \epsilon^{-(1 + \alpha_1)} \delta^{-(1 + \beta_1)} \xi^{-(1 + \alpha_2)} w^{-(1 + \beta_2)} \right. \\ &\quad \left. \times \left[\frac{\delta^{\kappa} - y^{\kappa}}{\delta^{\kappa}} - \frac{t(1-x)}{2\epsilon} \right]^n \left[\frac{w^{\kappa} - y^{\kappa}}{w^{\kappa}} - \frac{u(1-x)}{2\xi} \right]^m d\epsilon d\sigma dw d\xi \right) dudt. \end{aligned}$$

$(|\arg(\epsilon)|, |\arg(\sigma)|, |\arg(w)|, |\arg(\xi)| \leq \pi; \min\{Re(\alpha_1 + 1), Re(\alpha_2 + 1)\} > 0;$
 $\min\{Re(\kappa)\} > 0; \min\{Re(-n), Re(-m)\} > 0)$

6 Integral operator involving $E_{\alpha, \beta, \kappa}^{(\gamma_1; \gamma_2)}(x, y)$ in the kernel

In this section, we consider the following double (fractional) integral operator

$$\begin{aligned} \left(\xi_{\alpha, \beta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)} f \right)(x, y) &= \int_d^y \int_b^x (x-t)^{\alpha-1} (y-u)^{\beta-1} E_{\alpha, \beta, \kappa}^{(\gamma_1; \gamma_2)}(w_1(x-t), w_2(y-u)) f(t, u) dt du. \\ (x > b, y > d) \end{aligned} \quad (25)$$

In the case $\gamma_1 = \gamma_2 = 0$, the integral operator (25) reduces to the Riemann-Liouville double fractional integral operator defined in (20),

$$\left(\xi_{\alpha, \beta, \kappa; w_1, w_2, b^+, d^+}^{(0; 0)} f \right)(x, y) = \left({}_x I_{b^+}^{\alpha} {}_y I_{d^+}^{\beta} f \right)(x, y).$$

The space $L((b, a) \times (d, c))$ of absolutely integrable functions is defined as follows

$$L((b, a) \times (d, c)) = \left\{ g : \|g\|_1 = \int_b^a \int_d^c |g(x, y)| dy dx < \infty \right\}.$$

In the following theorem, we demonstrate that the operator $\xi_{\alpha, \beta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)}$ constitutes a transformation from $L((b, a) \times (d, c))$.

Theorem 6.1. *The double integral operator $\xi_{\alpha, \beta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)}$ is bounded in the space $L((b, a) \times (d, c))$,*

$$\left\| \xi_{\alpha, \beta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)} f \right\|_1 \leq K \|f\|_1, \quad (26)$$

where the constant K is independent of f given by

$$K = (a-b)^{Re(\alpha)} (c-d)^{Re(\beta)} \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{|(\gamma_1)_{r+s}| |(\gamma_2)_s| |w_1(a-b)|^s |w_2(c-d)|^{\kappa r}}{r! s! (Re(\alpha) + s) (Re(\beta) + \kappa r) |\Gamma(\alpha + s)| |\Gamma(\beta + \kappa r)|}.$$

Proof. By applying Fubini's Theorem, we obtain

$$\begin{aligned}
\left\| \xi_{\alpha, \beta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)} f \right\|_1 &\leq \int_b^a \int_d^c |f(t, u)| \\
&\quad \times \left(\int_t^a \int_u^c (x-t)^{\operatorname{Re}(\alpha)-1} (y-u)^{\operatorname{Re}(\beta)-1} \left| E_{\alpha, \beta, \kappa}^{(\gamma_1; \gamma_2)}(w_1(x-t), w_2(y-u)) \right| dy dx \right) dt du \\
&= \int_b^a \int_d^c |f(t, u)| \left(\int_0^{a-t} \int_0^{c-u} \nu^{\operatorname{Re}(\alpha)-1} \tau^{\operatorname{Re}(\beta)-1} \left| E_{\alpha, \beta, \kappa}^{(\gamma_1; \gamma_2)}(w_1 \nu, w_2 \tau) \right| d\tau d\nu \right) dt du \\
&\leq \int_b^a \int_d^c |f(t, u)| \left(\int_0^{a-b} \int_0^{c-d} \nu^{\operatorname{Re}(\alpha)-1} \tau^{\operatorname{Re}(\beta)-1} \left| E_{\alpha, \beta, \kappa}^{(\gamma_1; \gamma_2)}(w_1 \nu, w_2 \tau) \right| d\tau d\nu \right) dt du \\
&\leq \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{|\gamma_1|_{r+s} |(\gamma_2)_s| |w_1|^s |w_2|^{\kappa r}}{r! s! |\Gamma(\alpha+s)| |\Gamma(\beta+\kappa r)|} \\
&\quad \times \int_0^{a-b} \int_0^{c-d} \nu^{\operatorname{Re}(\alpha)+s-1} \tau^{\operatorname{Re}(\beta)+\kappa r-1} d\tau d\nu \|f\|_1 \\
&= K \|f\|_1.
\end{aligned}$$

□

Remark 6.1. The constant K is finite, because the series (18) is absolutely convergent for all x and y , since $\operatorname{Re}(\kappa) > 0$.

The next result of this section is a theorem that establishes a fundamental relationship between the integral operator, defined by equation (25) and the Riemann Liouville fractional calculus, expressed in the form of an infinite series.

Theorem 6.2. For any $\alpha, \beta, \gamma_1, \gamma_2, \kappa, w_1, w_2 \in \mathbb{C}$, $\operatorname{Re}(\alpha), \operatorname{Re}(\beta), \operatorname{Re}(\kappa) > 0$ and $f(x, y) \in L((b, a) \times (d, c))$, the representation of the operator given by equation (25) is as follows:

$$\xi_{\alpha, \beta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)} f(x, y) = \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{r+s} (\gamma_2)_s w_1^s w_2^{\kappa r}}{r! s!} {}_y I_{d^+}^{\beta+\kappa r} {}_x I_{b^+}^{\alpha+s} f(x, y). \quad (27)$$

Proof. The local uniform convergence of the series (18) permits the interchange of summation and integration in formula (25), as follows

$$\begin{aligned}
&\left(\xi_{\alpha, \beta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)} f \right)(x, y) \\
&= \int_d^y \int_b^x (x-t)^{\alpha-1} (y-u)^{\beta-1} E_{\alpha, \beta, \kappa}^{(\gamma_1; \gamma_2)}(w_1(x-t), w_2(y-u)) f(t, u) dt du \\
&= \int_d^y \int_b^x (x-t)^{\alpha-1} (y-u)^{\beta-1} \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{r+s} (\gamma_2)_s [w_1(x-t)]^s [w_2(y-u)]^{\kappa r}}{r! s! \Gamma(\alpha+s) \Gamma(\beta+\kappa r)} f(t, u) dt du \\
&= \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{r+s} (\gamma_2)_s w_1^s w_2^{\kappa r}}{r! s!} \frac{1}{\Gamma(\alpha+s) \Gamma(\beta+\kappa r)} \int_d^y \int_b^x (x-t)^{\alpha+s-1} (y-u)^{\beta+\kappa r-1} f(t, u) dt du \\
&= \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{r+s} (\gamma_2)_s w_1^r w_2^{\kappa s}}{r! s!} {}_y I_{d^+}^{\beta+\kappa r} {}_x I_{b^+}^{\alpha+s} f(x, y).
\end{aligned}$$

□

In the following corollary, we demonstrate the composition of RL fractional integrals and derivatives with the integral operator (25).

Corollary 6.1. *The following composition relationships are valid for all functions $f(x, y) \in L((b, a) \times (d, c))$.*

$$\begin{aligned} ({}_y I_{d^+}^\zeta {}_x I_{b^+}^\mu \xi_{\alpha, \beta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)} f)(x, y) &= (\xi_{\alpha + \mu, \beta + \zeta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)} f)(x, y) \\ &= (\xi_{\alpha, \beta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)} {}_y I_{d^+}^\zeta {}_x I_{b^+}^\mu f)(x, y) \\ ({}_y D_{d^+}^\zeta {}_x D_{b^+}^\mu \xi_{\alpha, \beta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)} f)(x, y) &= (\xi_{\alpha - \mu, \beta - \zeta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)} f)(x, y) \\ &= (\xi_{\alpha, \beta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)} {}_y D_{d^+}^\zeta {}_x D_{b^+}^\mu f)(x, y) \end{aligned}$$

To highlight the effectiveness of the proposed series formula (27), we present a straightforward approach for computing and applying the integral operator (25) to a power function and an exponential function. This alternative method offers a substantial improvement over the use of the original definition (25).

Theorem 6.3. *The integral operator $\xi_{\alpha, \beta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)}$ applied to a power function is expressed as follows:*

$$\begin{aligned} \xi_{\alpha, \beta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)} [(x - b)^\mu (y - d)^\zeta] &= \Gamma(\mu + 1) \Gamma(\zeta + 1) (x - b)^{\mu + \alpha} (y - d)^{\zeta + \beta} \\ &\quad \times E_{\alpha + \mu + 1, \beta + \zeta + 1, \kappa}^{(\gamma_1; \gamma_2)} [w_1(x - b), w_2(y - d)]. \end{aligned} \quad (28)$$

Proof. The Riemann Liouville fractional integral of a power function is given by

$$\begin{aligned} {}_y I_{d^+}^\beta (y - d)^\zeta &= \frac{\Gamma(\zeta + 1)}{\Gamma(\zeta + \beta + 1)} (y - d)^{\zeta + \beta}, \quad \operatorname{Re}(\zeta) > -1, \quad \operatorname{Re}(\beta) > 0, \\ {}_x I_{b^+}^\alpha (x - b)^\mu &= \frac{\Gamma(\mu + 1)}{\Gamma(\mu + \alpha + 1)} (x - b)^{\mu + \alpha}, \quad \operatorname{Re}(\mu) > -1, \quad \operatorname{Re}(\alpha) > 0. \end{aligned}$$

For $\operatorname{Re}(\zeta) > -1$ and $\operatorname{Re}(\mu) > -1$ by using above relations with series formula (27) we obtain

$$\begin{aligned} \xi_{\alpha, \beta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)} [(x - b)^\mu (y - d)^\zeta] &= \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{r+s} (\gamma_2)_s w_1^s w_2^{\kappa r}}{r! s!} {}_y I_{d^+}^{\beta + \kappa r} {}_x I_{b^+}^{\alpha + s} [(x - b)^\mu (y - d)^\zeta] \\ &= \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{r+s} (\gamma_2)_s w_1^s w_2^{\kappa r}}{r! s!} \frac{\Gamma(\zeta + 1) \Gamma(\mu + 1)}{\Gamma(\zeta + \beta + \kappa r + 1) \Gamma(\mu + \alpha + r + 1)} \\ &\quad \times (x - b)^{\alpha + \mu + s} (y - d)^{\zeta + \beta + \kappa s} \\ &= \Gamma(\mu + 1) \Gamma(\zeta + 1) (x - b)^{\mu + \alpha} (y - d)^{\zeta + \beta} \\ &\quad \times \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{r+s} (\gamma_2)_s [w_1(x - b)]^s [w_2(y - d)]^{\kappa r}}{\Gamma(\mu + \alpha + 1) \Gamma(\zeta + \beta + 1) r! s!} \end{aligned}$$

Thus, the result in equation (28) follows. \square

Theorem 6.4. *The integral operator $\xi_{\alpha, \beta, \kappa; w_1, w_2, b^+, d^+}^{(\gamma_1; \gamma_2)}$ applied to an exponential function, with left limits $b = -\infty$ and $d = -\infty$ is given by the formula:*

$$\xi_{\alpha, \beta, \kappa; w_1, w_2, -\infty, -\infty}^{(\gamma_1; \gamma_2)} [e^{\delta x + \sigma y}] = \delta^{-\alpha} \sigma^{-\beta} e^{\delta x + \sigma y} \left[1 - \left(\frac{w_2}{\sigma} \right)^\kappa \right]^{-\gamma_1} {}_2F_0 \left[\gamma_1, \gamma_2; -; \left(\frac{\sigma^\kappa w_1}{\delta(\sigma^\kappa - w_2^\kappa)} \right) \right],$$

where $|\frac{w_1}{\delta}| < 1$.

Proof. The Riemann-Liouville fractional integral of an exponential function is given by

$${}_y I_{-\infty}^\beta e^{\sigma y} = \sigma^{-\beta} e^{\sigma y}, \quad \operatorname{Re}(\sigma) > 0, \operatorname{Re}(\beta) > 0. \quad (29)$$

Using equation (29) together with the series formula (27), and assuming $Re(\sigma) > 0$ and $Re(\delta) > 0$, it follows that:

$$\begin{aligned}
\xi_{\alpha,\beta,\kappa;w_1,w_2,-\infty,-\infty}^{(\gamma_1;\gamma_2)}[e^{\delta x+\sigma y}] &= \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{r+s}(\gamma_2)_s w_1^s w_2^{\kappa r}}{r!s!} {}_y I_{d^+}^{\beta+\kappa r} {}_x I_{b^+}^{\alpha+s} [e^{\delta x+\sigma y}] \\
&= \delta^{-\alpha} \sigma^{-\beta} e^{\delta x+\sigma y} \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{r+s}(\gamma_2)_s \left(\frac{w_1}{\delta}\right)^s \left(\frac{w_2}{\sigma}\right)^{\kappa r}}{r!s!} \\
&= \delta^{-\alpha} \sigma^{-\beta} e^{\delta x+\sigma y} \sum_{s=0}^{\infty} \frac{(\gamma_1)_s (\gamma_2)_s \left(\frac{w_1}{\delta}\right)^s}{s!} \sum_{r=0}^{\infty} \frac{(\gamma_1+s)_r \left(\frac{w_2}{\sigma}\right)^{\kappa r}}{r!} \\
&= \delta^{-\alpha} \sigma^{-\beta} e^{\delta x+\sigma y} \left[1 - \left(\frac{w_2}{\sigma}\right)^{\kappa}\right]^{-\gamma_1} \sum_{s=0}^{\infty} \frac{(\gamma_1)_s (\gamma_2)_s}{s!} \left(\frac{\sigma^{\kappa} w_1}{\delta(\sigma^{\kappa} - w_2^{\kappa})}\right)^s.
\end{aligned}$$

□

Next, we focus on the Laplace transform. Since the operator (25) involves a double integral, it is necessary to consider a two-variable Laplace transform.

Theorem 6.5. *Under the conditions $\gamma_1, \gamma_2, \kappa, \alpha, \beta, w_1, w_2 \in \mathbb{C}, Re(\beta) > 0, Re(\alpha) > 0, Re(\kappa) > 0$ and for any function $\psi \in L((0, a) \times (0, c))$, we have*

$$\mathbb{L}_2 \left(\xi_{\alpha,\beta,\kappa;w_1,w_2,0,0}^{(\gamma_1;\gamma_2)} f \right) (p, q) = \frac{1}{p^{\alpha} q^{\beta}} \left[1 - \left(\frac{w_2}{q} \right)^{\kappa} \right]^{-\gamma_1} {}_2 F_0 \left[\gamma_1, \gamma_2; -; \left(\frac{q^{\kappa} w_1}{p(q^{\kappa} - w_2^{\kappa})} \right) \right] \mathbb{L}_2 f(p, q),$$

where $p, q \in \mathbb{C}$ such that $Re(q) > 0, Re(p) > 0$ and $|\frac{w_1}{q}| < 1$.

Proof. By using (27) we obtain,

$$\begin{aligned}
\mathbb{L}_2 \left(\xi_{\alpha,\beta,\kappa;w_1,w_2,0,0}^{(\gamma_1;\gamma_2)} f \right) (p, q) &= \mathbb{L}_2 \left[\sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{r+s}(\gamma_2)_s w_1^s w_2^{\kappa r}}{r!s!} {}_y I_0^{\beta+\kappa r} {}_x I_0^{\alpha+s} f \right] (p, q) \\
&= \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{r+s}(\gamma_2)_s w_1^s w_2^{\kappa r}}{r!s!} \mathbb{L}_2 {}_y I_0^{\beta+\kappa r} {}_x I_0^{\alpha+s} f (p, q) \\
&= \frac{1}{p^{\alpha} q^{\beta}} \sum_{s=0}^{\infty} \sum_{r=0}^{\infty} \frac{(\gamma_1)_{r+s}(\gamma_2)_s \left(\frac{w_1}{p}\right)^s \left(\frac{w_2}{q}\right)^{\kappa r}}{r!s!} \mathbb{L}_2 f(p, q)
\end{aligned}$$

the remaining manipulation of the series follows precisely the same steps as in the proof of above Theorem. We have utilized the standard result [35] that the double Riemann-Liouville integral has a double Laplace transform given by:

$$\mathbb{L}_2 \left[{}_y I_0^{\beta} {}_x I_0^{\alpha} f(x, y) \right] = \frac{1}{p^{\alpha} q^{\beta}} \mathbb{L}_2 f(x, y), \quad Re(\alpha) > 0, Re(\beta) > 0, Re(p) > 0, Re(q) > 0.$$

□

References

- [1] Özarslan MA, Elidemir İO. Fractional Calculus Operator Emerging from the 2D Biorthogonal Hermite Konhauser Polynomials, 2023 (arXiv:2404.00035).
- [2] Bin-Saad Maged G. Associated Laguerre-Konhauser polynomials quasi-monomiality and operational identities. J. Math. Anal.Appl. 2006;324:1438-1448.
- [3] Özarslan MA, Kürt C. Bivariate Mittag-Leffler functions arising in the solutions of convolution integral equation with 2D-Laguerre-Konhauser polynomials in the kernel. Appl. Math. Comp. 2019;347:631-644.

- [4] Krall HL, Sheffer IM. Orthogonal polynomials in two variables. *Annali di Matematica Pura ed Applicata*, 1967;76: 325-376.
- [5] Koornwinder T. Two-variable analogues of the classical orthogonal polynomials. In *theory and application of special functions*. Academic Press. 1975;435-495.
- [6] Suetin, PK (1999). Orthogonal polynomials in two variables. CRC Press. 1999;3.
- [7] Gorenflo R, Kilbas AA, Mainardi F, Rogosin S V. Mittag-Leffler functions, related topics and applications. 2nd edition: Berlin: Springer;2020.
- [8] Kiryakova VS, Luchko Y F. (2010, November). The Multi-index Mittag-Leffler Functions and their Applications for Solving Fractional Order Problems in Applied Analysis. In *AIP Conference Proceedings*. American Institute of Physics. 2010;1301: 597 - 613.
- [9] Luchko Y. The four-parameters Wright function of the second kind and its applications in FC. *Mathematics*. 2020;8: 970.
- [10] Prabhakar TR. A singular integral equation with a generalized Mittag-Leffler function in the kernel. *Yokohama math. J.* 1971;19:7-15.
- [11] Kilbas AA, Saigo M, Saxena R K. Generalized Mittag-Leffler function and generalized fractional calculus operators. *Integral transforms and Special Functions*. 2004;15:31-49.
- [12] Fernandez A, Kürt C, Özarslan MA. A naturally emerging bivariate Mittag-Leffler function and associated fractional-calculus operators. *Computational and Applied Mathematics*. 2020;39:1-27.
- [13] Ng YX, Phang C, Loh JR, Isah A, MalaYsia S. Analytical solutions of incommensurate fractional differential equation systems with fractional order $1 < \gamma, \beta < 2$ via bivariate Mittag-Leffler functions. *behaviour*. 2022;13:14.
- [14] Garg M, Manohar P, Kalla S L. (2013). A Mittag-Leffler-type function of two variables. *Integral transforms and Special Functions*. 2013;24: 934-944.
- [15] Saigo M, Kilbas AA. (1998). On Mittag-Leffler type function and applications. *Integral transforms and Special Functions*. 1998;7: 97-112.
- [16] Kilbas AA, Saigo M. On Mittag-Leffler type function, fractional calculus operators and solutions of integral equations. *Integral transforms and Special Functions*. 1996;4: 355-370.
- [17] Shukla A K, Prajapati JC. On a generalization of Mittag-Leffler function and its properties. *Journal of mathematical analysis and applications*. 2007;336: 797-811.
- [18] Kilbas AA, Srivastava HM, Trujillo JJ. *theory and Applications of Fractional Differential Equations*. Elsevier: Amsterdam, the Netherlands. 2006.
- [19] Saxena RK, Kalla SL, Saxena R. Multivariate analogue of generalized Mittag-Leffler function, *Int. trans. Special Funct.* 2011;22: 533-548.
- [20] Özarslan MA. On a singular integral equation including a set of multivariate polynomials suggested by Laguerre polynomials. *Applied Mathematics and Computation*. 2014;229: 350-358.
- [21] Özarslan MA, Fernandez A. On the fractional calculus of multivariate Mittag-Leffler functions. *International Journal of Computer Mathematics*. 2022;99:247-273.
- [22] Kürt C, Fernandez A, Özarslan MA. Two unified families of bivariate Mittag-Leffler functions. *Applied Mathematics and Computation*. 2023;443:127785.
- [23] Kürt C, Özarslan MA, Fernandez A. On a certain bivariate Mittag-Leffler function analysed from a fractional-calculus point of view. *Mathematical Methods in the Applied Sciences*. 2021;44:2600-2620.

- [24] Özarslan MA, Fernandez A. On a five-parameter Mittag-Leffler function and the corresponding bivariate fractional operators. *Fractal and Fractional*. 2021;5:45.
- [25] Isah SS, Fernandez A, Özarslan MA. (2023). On bivariate fractional calculus with general univariate analytic kernels. *Chaos, Solitons Fractals*. 2023;171: 113495.
- [26] Isah SS, Fernandez A, Özarslan, MA. On univariate fractional calculus with general bivariate analytic kernels. *Computational and Applied Mathematics*, 2023;42(5):228.
- [27] Shahwan MJ , Bin-Saad M G, Al-Hashami A, Some properties of bivariate Mittag-Leffler function. *the Journal of Analysis*, 2023; 1-21.
- [28] Carlitz L. A note on certain biorthogonal polynomials. *Pac.J. Math.*1968;24:425-430.
- [29] Konhauser JDE. Biorthogonal polynomials suggested by the Laguerre polynomials. *Pacific J. Math.*1967;21: 303-314.
- [30] Konhauser JDE. Some properties of biorthogonal polynomials. *J. Math. Anal.Appl.* 1965;11: 242-260.
- [31] Madhekar HC, thakare NK. Biorthogonal polynomials suggested by the Jacobi polynomials. *Pacific J. Math.* 1982;100:417-424.
- [32] Srivastava HM , Daust MC. A note on the convergence of Kampê de fêriet's double hypergeometric series. *Math. Nachr.* 1972;53: 151-159.
- [33] Whittaker ET, Watson GN. *A course of modern analysis: An introduction to the general theory of infinite process an of analytic functions; with an account of the principal transcendental functions* ed. Cambridge, London and New York. 1927.
- [34] Srivastava HM, Daoust MC. On Eulerian integrals associated with Kampê de Fêriet's function. *Publications de l'Institut Mathématique [Elektronische Ressource]*, 1969;23:199-202.
- [35] Samko SG, Kilbas AA, Marichev OI. *Fractional Integrals and Derivatives: Theory and Applications*, Taylor and Francis, London, 2002.