

The generalized 4-connectivity of godan graphs*

Jing Wang¹, Yuanqiu Huang², Zhangdong Ouyang³

¹ School of Mathematics, Changsha University, Changsha 410022, China

² School of Mathematics and Statistics, Hunan Normal University, Changsha 410081, China

³ School of Mathematics and Statistics, Hunan First Normal University, Changsha 410205, China

Abstract The generalized k -connectivity of a graph G , denoted by $\kappa_k(G)$, is the minimum number of internally edge disjoint S -trees for any $S \subseteq V(G)$ and $|S| = k$. The generalized k -connectivity is a natural extension of the classical connectivity and plays a key role in applications related to the modern interconnection networks. The godan graph EA_n is a kind of Cayley graphs which possesses many desirable properties. In this paper, we shall study the generalized 4-connectivity of EA_n and show that $\kappa_4(EA_n) = n - 1$ for $n \geq 3$.

Keywords interconnection network, godan graph, generalized k -connectivity, tree
MR(2000) Subject Classification 05C40, 05C05

1 Introduction

A network is usually represented by a connected graph $G = (V(G), E(G))$, where $V(G)$ is the set of processors and $E(G)$ is the set of communication links between processors. Fault tolerance has become increasingly significant nowadays since multiprocessor systems failure is inevitable. It is important to consider how the network performs in the event of a certain number of nodes of failure and/or links failure in the network topology. The connectivity is an important parameter for measuring the fault tolerance of the network. A subset $S \subseteq V(G)$ of a connected graph G is called a *vertex-cut* if $G \setminus S$ is disconnected or trivial. The *connectivity* $\kappa(G)$ of G is defined as the minimum cardinality over all vertex-cuts of G . Note that the larger $\kappa(G)$ is, the more reliable the network is.

Let P be a path in G with x and y as its two terminal vertices, then P is called an (x, y) -*path*. Two (x, y) -paths P_1 and P_2 are *internally disjoint* if they have no common internal vertices, that is, $V(P_1) \cap V(P_2) = \{x, y\}$. A well known theorem of Whitney [20] provides an equivalent definition of connectivity. For each 2-subset $S = \{x, y\} \subseteq V(G)$, let $\kappa(S)$ denote the maximum number of internally disjoint

*This work was supported by National Science Foundation of China(No.12271157 & No.12371346) and Natural Science Foundation of Hunan Province (No.2022JJ30028 & No.2023JJ30072).

(x, y) -paths in G . Then

$$\kappa(G) = \min\{\kappa(S) \mid S \subseteq V(G) \text{ and } |S| = 2\}.$$

The generalized k -connectivity, which was introduced by Chartrand et al. [2], is a strengthening of connectivity and can be served as an essential parameter for measuring reliability and fault tolerance of the network. Let $G = (V(G), E(G))$ be a simple graph, S be a subset of $V(G)$. A tree T in G is called an S -tree, if $S \subseteq V(T)$. The trees T_1, T_2, \dots, T_r are called *internally edge disjoint S -trees* if $V(T_i) \cap V(T_j) = S$ and $E(T_i) \cap E(T_j) = \emptyset$ for any integers $1 \leq i < j \leq r$. Throughout this paper, we will use IDSTs to denote internally edge disjoint S -trees for short. Let $\kappa_G(S)$ denote the maximum number of IDSTs.

For an integer k with $2 \leq k \leq |V(G)|$, the *generalized k -connectivity* of G , denoted by $\kappa_k(G)$, is defined as

$$\kappa_k(G) = \min\{\kappa_G(S) \mid S \subseteq V(G) \text{ and } |S| = k\}.$$

The generalized 2-connectivity is exactly the classical connectivity. Over the past few years, research on the generalized connectivity has received meaningful progress. Li et al. [12] derived that it is NP-complete for a general graph G to decide whether there are l internally edge disjoint trees connecting S , where l is a fixed integer and $S \subseteq V(G)$. Authors in [7, 11] investigated the upper and lower bounds of the generalized connectivity of a general graph G .

For $3 \leq k \leq |V(G)|$, Li gave an upper bound of $\kappa_k(G)$ for a general graph G in her Ph.D. thesis [9].

Lemma 1.1 ([9]) *Let G be a connected graph with minimum degree $\delta(G)$. If there are two adjacent vertices of degree $\delta(G)$, then $\kappa_k(G) \leq \delta(G) - 1$ for $3 \leq k \leq |V(G)|$.*

The following result is about the relationship between $\kappa_k(G)$ and $\kappa_{k-1}(G)$ of a regular graph G .

Lemma 1.2 ([15]) *Let G be an r -regular graph. If $\kappa_k(G) = r - 1$, then $\kappa_{k-1}(G) = r - 1$, where $k \geq 4$.*

Many authors tried to study exact values of the generalized connectivity of graphs. The generalized k -connectivity of the complete graphs, $\kappa_k(K_n)$, was determined in [3] for every pair k, n of integers with $2 \leq k \leq n$. The generalized k -connectivity of the complete bipartite graphs $K_{a,b}$ was obtained in [10] for all $2 \leq k \leq a + b$. Apart from these results, the generalized k -connectivity of hypercubes [8, 15], the generalized hypercubes [17], balanced hypercubes [19], exchanged hypercubes [22], dual cubes [26], star graphs and bubble-sort graphs [14], (n, k) -star graphs [6, 24], bubble-sort star graphs [23], pancake graphs [25], Cayley graphs

generated by trees and cycles [13], et al, have drawn many scholars' attention. So far, as we can see, the results on the generalized k -connectivity of networks were almost about $k \leq 4$.

Ren and Wang [16] introduced the n -dimensional godan graph EA_n as a topology structure of interconnection networks, they proved that EA_n poses many desirable properties, such as regularity, high connectivity and vertex-transitivity. The authors in [16] also gave various connectivity, such as nature connectivity, 2-good-neighbour connectivity, 2-extra connectivity, of EA_n . The component connectivity of EA_n was studied in [21]. In this paper, we try to calculate the generalized 4-connectivity of EA_n and obtain the following main result.

Theorem 1.1 *Let EA_n be the n -dimensional godan graph for $n \geq 3$. Then $\kappa_4(EA_n) = n - 1$.*

We end this section with some necessary preliminaries. We only consider a simple, connected graph $G = (V(G), E(G))$ with vertex set $V(G)$ and edge set $E(G)$. For $x \in V(G)$, the *degree* of x in G , denoted by $\deg_G(x)$, is the number of edges of G incident with x . Denote $\delta(G)$ the *minimum degree* of vertices of G . A graph is *d -regular* if $\deg_G(x) = d$ for every $x \in V(G)$. For $x \in V(G)$, we use $N_G(x)$ to denote the set of neighbours of x in G . Let $V' \subseteq V(G)$, denote by $G \setminus V'$ the graph obtained from G by deleting all the vertices in V' together with their incident edges. Denote by $G[V']$ the subgraph of G induced on V' .

The following Lemma 1.3, Lemma 1.4 and Lemma 1.5 are results on the connectivity of a graph that are well-known in the literature.

Lemma 1.3 ([1]) *Let G be a k -connected graph, and let x and y be a pair of distinct vertices of G . Then there exist k internally disjoint (x, y) -paths in G .*

Lemma 1.4 ([1]) *Let G be a k -connected graph, let x be a vertex of G and let $Y \subseteq V(G) \setminus \{x\}$ be a set of at least k vertices of G . Then there exists a k -fan in G from x to Y , that is, there exists a family of k internally disjoint (x, Y) -paths whose terminal vertices are distinct in Y .*

Lemma 1.5 ([1]) *Let G be a k -connected graph, and let X and Y be subsets of $V(G)$ of cardinality at least k . Then there exists a family of k pairwise disjoint (X, Y) -paths in G .*

This paper is organized as follows. Section 2 presents the definition and properties of godan graphs. We note that the lower bound of $\kappa_4(EA_n)$ is critical to obtain the main result. Therefore, some lemmas which will be used to obtain the lower bound of $\kappa_4(EA_n)$ are given in Section 3. Then Theorem 1.1 is proved in Section 4. Finally, the paper is concluded in Section 5.

2 Definition and properties of godan graph

Let $[n] = \{1, 2, \dots, n\}$. For convenience, we denote the permutation $\begin{pmatrix} 1 & 2 & \dots & n \\ p_1 & p_2 & \dots & p_n \end{pmatrix}$ by $p_1 p_2 \dots p_n$ and the permutation $\begin{pmatrix} 1 & \dots & i & \dots & j & \dots & n \\ 1 & \dots & j & \dots & i & \dots & n \end{pmatrix}$ by (ij) , while the latter is called a transposition. The composition $\sigma \circ \tau$ of two permutations σ and τ is the function that maps any element i to $\sigma(\tau(i))$.

Let S_n be the symmetric group on $[n]$. The alternating group A_n ($n \geq 3$) is the subgroup of S_n containing all even permutations. Let $\Omega = \{(123), (132)\} \cup \{(12)(3i) | 4 \leq i \leq n\}$. For $n \geq 3$, the n -dimensional alternating group network AN_n ([5]) is a graph with vertex set $V(AN_n) = A_n$ and edge set $E(AN_n) = \{uv | u = v \circ s, \text{ where } \{u, v\} \subseteq V(AN_n) \text{ and } s \in \Omega\}$.

The alternating group network AN_3 is depicted in Figure 1. It is worth mentioning that AN_n is in fact isomorphic to the $(n, n-2)$ -star $S_{n,n-2}$ [4], whose generalized 4-connectivity is proved to be $(n-2)$ in [6].




Figure 1: The alternating group network AN_3




Figure 2: The godan graph EA_3

Lemma 2.1 ([6, 27]) *For the n -dimensional alternating group network AN_n ($n \geq 3$), the following properties hold:*

- (1) AN_n is $(n-1)$ -regular with $\frac{n!}{2}$ vertices and $\frac{n!(n-1)}{4}$ edges;
- (2) $\kappa(AN_n) = n-1$;
- (3) $\kappa_4(AN_n) = \kappa_4(S_{n,n-2}) = n-2$.

In [16], Ren and Wang introduced the n -dimensional godan graph EA_n as a topology structure of interconnection networks.

Definition 2.1 ([16]) *For $n \geq 3$, let $\Omega^* = \{(12), (123), (132)\} \cup \{(12)(3i) | 4 \leq i \leq n\}$. The n -dimensional godan graph EA_n is the graph with vertex set $V(EA_n) = S_n$ in which two vertices u and v are adjacent in EA_n if and only if $u = v \circ s^*$, where $s^* \in \Omega^*$.*

The godan graphs EA_3 and EA_4 are depicted in Figure 2 and Figure 3, respectively. Some properties of EA_n are listed in Lemma 2.2.




Figure 3: The godan graph EA_4

Lemma 2.2 ([16, 21]) For $n \geq 3$, the n -dimensional godan graph EA_n has the following properties:

- (1) EA_n is n -regular with $n!$ vertices and $\frac{n \times n!}{2}$ edges;
- (2) $EA_n[A_n] \cong AN_n$, $EA_n[S_n \setminus A_n] \cong AN_n$;
- (3) $E(A_n, S_n \setminus A_n)$ is a perfect matching of EA_n , where $E(A_n, S_n \setminus A_n)$ denotes all cross edges between A_n and $S_n \setminus A_n$;
- (4) $\kappa(EA_n) = n$.

It is seen that both $EA_n[A_n]$ and $EA_n[S_n \setminus A_n]$ are isomorphic to AN_n . For simplicity, we denote $EA_n[A_n]$ by AN_n^1 and denote $EA_n[S_n \setminus A_n]$ by AN_n^2 , respectively. Then we can write the construction of EA_n symbolically as

$$EA_n = AN_n^1 \otimes AN_n^2.$$

For $1 \leq i \leq 2$, let x be a vertex in AN_n^i , it is clear that x has a unique neighbor $x \circ (12)$ in $V(AN_n^{3-i})$. Denote $x \circ (12)$ by \tilde{x} . We say that \tilde{x} is the *parity neighbour* of x . Due to Definition 2.1, the following result is an easy observation.

Lemma 2.3 For $n \geq 3$, let $EA_n = AN_n^1 \otimes AN_n^2$ and x be a vertex of AN_n^i , $1 \leq i \leq 2$. If $N_{AN_n^i}(x) = \{b_1, b_2, \dots, b_{n-1}\}$. Then $N_{AN_n^{3-i}}(\tilde{x}) = \{\tilde{b}_1, \tilde{b}_2, \dots, \tilde{b}_{n-1}\}$.

Lemma 2.4 For $n \geq 4$, let x be a vertex of EA_n with $N_{EA_n}(x) = \{b_1, b_2, \dots, b_{n-1}, \tilde{x}\}$. Set $B = \{b_1, \dots, b_{n-2}, \tilde{x}, v\}$, where v is a neighbour of \tilde{x} with $v \neq x$. Then $EA_n \setminus B$ is connected.

Proof Remind that $EA_n = AN_n^1 \otimes AN_n^2$. By symmetry, assume that $x \in V(AN_n^1)$. Let $B_1 = \{b_1, \dots, b_{n-2}\}$ and $B_2 = \{\tilde{x}, v\}$.

By Definition 2.1, for $1 \leq i \leq 2$, $B_i = B \cap V(AN_n^i)$ because $v \neq x$.

By Lemma 2.1(2), for $1 \leq i \leq 2$, $AN_n^i \setminus B_i$ is connected since $|B_1| = n - 2$ and $|B_2| = 2$. Since $\frac{n!}{2} > n$ for $n \geq 4$ and $E(A_n, S_n \setminus A_n)$ is a perfect matching of EA_n , $EA_n \setminus B$ is still connected. \square

Now, we shall investigate the structure of EA_n from another perspective. Let $EA_n^{m:i}$, called a *cluster*, be a subgraph of EA_n induced on vertices with the m th symbol is i , where $4 \leq m \leq n$ and $i \in [n]$. Note that each cluster $EA_n^{m:i}$ is isomorphic to EA_{n-1} , $4 \leq m \leq n$ and $i \in [n]$. Thus, the construction of EA_n may be described as

$$EA_n = EA_n^{m:1} \oplus EA_n^{m:2} \oplus \cdots \oplus EA_n^{m:n}, \quad 4 \leq m \leq n.$$

For simplicity, we use EA_n^i to denote $EA_n^{n:i}$ in the rest of this paper, $i \in [n]$. Thus, we may write the construction of EA_n symbolically as

$$EA_n = EA_n^1 \oplus EA_n^2 \oplus \cdots \oplus EA_n^n.$$

For $i \in [n]$, let x be a vertex in EA_n^i . Clearly, x has $(n - 1)$ neighbours in EA_n^i and exactly one neighbor $x' = x \circ (12)(3n)$ out of EA_n^i . We say that x' is the *out neighbour* of x .

By Definition 2.1, some properties of EA_n can be derived directly, which are presented in Lemma 2.5.

Lemma 2.5 *For $n \geq 3$, let $EA_n = EA_n^1 \oplus EA_n^2 \oplus \cdots \oplus EA_n^n$. The following properties hold:*

- (1) $|E(EA_n^i, EA_n^j)| = (n - 2)!$, where i and j are distinct integers in $[n]$.
- (2) Let $x \in V(EA_n^i)$, $i \in [n]$. Then $|N_{EA_n^i}(x)| = n - 1$. Furthermore, the vertices in $N_{EA_n^i}(x)$ can be ordered in an unique way $x_1, x_2, \dots, x_{i-1}, x_{i+1}, \dots, x_n$ such that $x'_j \in V(EA_n^j)$ for $j \in [n] \setminus \{i\}$. We use $P[x, x'_j]$ to denote the path $\{xx_j, x_jx'_j\}$, where $j \in [n] \setminus \{i\}$.
- (3) Let $x \in V(EA_n^i)$ and x_j be a neighbour of x in EA_n^i , $\{i, j\} \subseteq [n]$ and $j \neq i$. Then x' and x'_j belong to a same cluster of EA_n if and only if $x_j = x \circ (12)$. Moreover, $x'_j = x' \circ (12)$.
- (4) Let x, y and z be three distinct vertices of EA_n . The subgraph induced on $\{x, y, z\}$ is a 3-cycle if and only if $\{y, z\} = \{x \circ (123), x \circ (132)\}$.

We end this section with Lemma 2.6 which determines the lower bound of the connectivity of a subgraph of EA_n .

Lemma 2.6 *For $n \geq 4$, let $EA_n = EA_n^1 \oplus EA_n^2 \oplus \cdots \oplus EA_n^n$. Denote $H = \bigcup_{i=1}^h EA_n^{k_i}$ be a subgraph of EA_n , where $k_i \in [n]$ and $h \geq 2$. Then $\kappa(H) \geq n - 2$.*

Proof Let x and y be any two distinct vertices in H . It suffices to prove that there are $(n - 2)$ internally disjoint (x, y) -paths in H .

Case 1. Both x and y belong to a same cluster, say $EA_n^{k_1}$.

By Lemma 2.2(4), there are $(n - 1)$ internally disjoint (x, y) -paths in $EA_n^{k_1}$ since $\kappa(EA_n^{k_1}) = \kappa(EA_{n-1}) = n - 1$.

Case 2. x and y belong to different clusters of EA_n .

W.l.o.g., assume that $x \in V(EA_n^{k_1})$ and $y \in V(EA_n^{k_2})$. Based on Lemma 2.5(1), $|E(EA_n^{k_1}, EA_n^{k_2})| = (n - 2)! \geq n - 2$ for $n \geq 4$. We may choose $(n - 2)$ different vertices $u_i \in V(EA_n^{k_1})$ such that $u'_i \in V(EA_n^{k_2})$, $i \in [n - 2]$. Combined with Lemma 1.4 and Lemma 2.2(4), there is an $(n - 2)$ -fan L_1, L_2, \dots, L_{n-2} in $EA_n^{k_1}$ from x to $\{u_1, u_2, \dots, u_{n-2}\}$ such that $u_i \in V(L_i)$, where $i \in [n - 2]$. Likewise, there is an $(n - 2)$ -fan Q_1, Q_2, \dots, Q_{n-2} in $EA_n^{k_2}$ from y to $\{u'_1, u'_2, \dots, u'_{n-2}\}$ such that $u'_i \in V(Q_i)$, $i \in [n - 2]$.

For $i \in [n - 2]$, let $P_i = L_i \cup Q_i \cup \{u_i u'_i\}$. Then P_1, \dots, P_{n-2} are $(n - 2)$ internally disjoint (x, y) -paths in H . \square

3 Some lemmas

The lower bound of $\kappa_4(EA_n)$ is the key to the proof of Theorem 1.1. We shall make great efforts to provide several lemmas to obtain the lower bound of $\kappa_4(EA_n)$ for $n \geq 4$.

Lemma 3.1 *For $n \geq 4$, let $EA_n = AN_n^1 \otimes AN_n^2$ and $S = \{x, y, z, w\}$ be any 4-subset of $V(EA_n)$. If there exists an integer i ($1 \leq i \leq 2$) such that $S \subseteq V(AN_n^i)$. Then there are $(n - 1)$ IDSTs in EA_n .*

Proof W.l.o.g., assume that $S \subseteq V(AN_n^1)$. Then there are $(n - 2)$ IDSTs T_1, \dots, T_{n-2} in AN_n^1 by Lemma 2.1(3) and Lemma 2.2(2).

Note that $\{\tilde{x}, \tilde{y}, \tilde{z}, \tilde{w}\} \subseteq V(AN_n^2)$. There is a $\{\tilde{x}, \tilde{y}, \tilde{z}, \tilde{w}\}$ -tree T'_{n-1} in AN_n^2 . Let $T_{n-1} = T'_{n-1} \cup \{x\tilde{x}, y\tilde{y}, z\tilde{z}, w\tilde{w}\}$. Clearly, T_1, \dots, T_{n-1} are $(n - 1)$ IDSTs in EA_n . \square

Lemma 3.2 *For $n \geq 4$, let $EA_n = AN_n^1 \otimes AN_n^2$ and $S = \{x, y, z, w\}$ be any 4-subset of $V(EA_n)$ satisfying the following three conditions:*

- (1) $\{x, y, z\} \subseteq V(AN_n^1)$ and $w \in V(AN_n^2)$;
- (2) $\tilde{w} \notin \{x, y, z\}$;
- (3) $|E(\tilde{w}, \{x, y, z\})| \leq 1$.

Then there are $(n - 1)$ IDSTs in EA_n .

Proof Based on Lemma 2.1(3), there are $(n - 2)$ internally edge disjoint $\{x, y, z, \tilde{w}\}$ -trees T'_1, \dots, T'_{n-2} in AN_n^1 . Furthermore, it has $1 \leq \deg_{T'_i}(\tilde{w}) \leq 2$ for each

$i \in [n-2]$ since $\deg_{AN_n^1}(\tilde{w}) = n-1$. For convenience, we may assume that $\deg_{T'_i}(\tilde{w}) = 1$ and \tilde{u}_i be a neighbour of \tilde{w} in T'_i , where $i \in [n-3]$.

Let $\tilde{U} = \{\tilde{u}_1, \dots, \tilde{u}_{n-3}\}$ and $U = \{u_i \mid \tilde{u}_i \in \tilde{U}, i \in [n-3]\}$.

Note that $U \subseteq V(AN_n^2)$ and $|U| = n-3$. Thus $AN_n^2 \setminus U$ is still connected since $\kappa(AN_n^2) = n-1$. The following two cases are considered.

Case 1. $E(\tilde{w}, \{x, y, z\}) = \emptyset$.

Then $\{\tilde{x}, \tilde{y}, \tilde{z}\} \cap \{w, u_1, \dots, u_{n-3}\} = \emptyset$. Moreover, $wu_i \in E(AN_n^2)$ for $i \in [n-3]$ according to Lemma 2.3.

There is a $\{\tilde{x}, \tilde{y}, \tilde{z}, w\}$ -tree T'_{n-1} in $AN_n^2 \setminus U$. Let $T_{n-1} = T'_{n-1} \cup \{x\tilde{x}, y\tilde{y}, z\tilde{z}\}$ and $T_{n-2} = T'_{n-2} \cup \{w\tilde{w}\}$. For $i \in [n-3]$, let $T_i = (T'_i \setminus \{\tilde{w}\}) \cup \{wu_i, u_i\tilde{u}_i\}$.

Then T_1, \dots, T_{n-1} are $(n-1)$ IDSTs in EA_n .

Case 2. $|E(\tilde{w}, \{x, y, z\})| = 1$.

Assume that $x\tilde{w} \in E(AN_n^1)$. If $x \notin \tilde{U}$, then we can construct $(n-1)$ IDSTs by similar methods in Case 1.

Hereafter, we consider the case that $x \in \tilde{U}$. For simplicity, set $x = \tilde{u}_1$. We only discuss the case that $\deg_{T'_{n-2}}(\tilde{w}) = 2$, since $(n-1)$ IDSTs can be obtained just by replacing T_1 and T_{n-2} in Case 1 if $\deg_{T'_{n-2}}(\tilde{w}) = 1$.

Let \tilde{u}_{n-2} and \tilde{u}_{n-1} be two neighbours of \tilde{w} in T'_{n-2} . Moreover, let $\tilde{U}_1 = \{\tilde{u}_2, \dots, \tilde{u}_{n-1}\}$ and $U_1 = \{u_i \mid \tilde{u}_i \in \tilde{U}_1, 2 \leq i \leq n-1\}$. Then $U_1 \subseteq V(AN_n^2)$ and $|U_1| = n-2$. Therefore, $AN_n^2 \setminus U_1$ is connected and there is a $\{\tilde{x}, \tilde{y}, \tilde{z}, w\}$ -tree T'_{n-1} in $AN_n^2 \setminus U_1$.

Let

$$T_{n-1} = T'_{n-1} \cup \{x\tilde{x}, y\tilde{y}, z\tilde{z}\},$$

$$T_{n-2} = (T'_{n-2} \setminus \{\tilde{w}\}) \cup \{wu_{n-2}, u_{n-2}\tilde{u}_{n-2}, wu_{n-1}, u_{n-1}\tilde{u}_{n-1}\}$$

and $T_1 = T'_1 \cup \{w\tilde{w}\}$. For $2 \leq i \leq n-3$, let T_i be the same as in Case 1. Then T_1, \dots, T_{n-1} are $(n-1)$ IDSTs in EA_n . \square

For $n \geq 4$ and $4 \leq m \leq n$, recall that $EA_n = EA_n^{m:1} \oplus EA_n^{m:2} \oplus \dots \oplus EA_n^{m:n}$. Let $S = \{x, y, z, w\}$ be any 4-subset of $V(EA_n)$. In order to obtain the lower bound of $\kappa_4(EA_n)$, the following subsections are distinguished based on the value of $\max_{i \in [n]} |S \cap V(EA_n^{m:i})|$.

3.1 $\max_{i \in [n]} |S \cap V(EA_n^{m:i})| = 3, 4 \leq m \leq n$.

Lemma 3.3 *For $n \geq 4$ and $4 \leq m \leq n$, let $EA_n = EA_n^{m:1} \oplus EA_n^{m:2} \oplus \dots \oplus EA_n^{m:n}$ and $S = \{x, y, z, w\}$ be any 4-subset of $V(EA_n)$. If there exists an integer $i \in [n]$ such that $|S \cap V(EA_n^{m:i})| = 3$. Then there are $(n-1)$ IDSTs in EA_n .*

Proof W.l.o.g., we assume that $\{x, y, z\} \subseteq V(EA_n^n)$ and $w \in V(EA_n^{n-1})$.

By Lemma 2.5(2), for $i \in [n-2]$, there is a $\{x'_i, y'_i, z'_i, w'_i\}$ -tree T'_i in EA_n^i since EA_n^i is connected, let

$$T_i = T'_i \cup P[x, x'_i] \cup P[y, y'_i] \cup P[z, z'_i] \cup P[w, w'_i].$$

It is possible that $x_i = y_i$ for some $i \in [n-2]$. This possibility will not affect the result of this lemma.

Let $W = \{w_1, \dots, w_{n-2}\}$. By Lemma 2.2(4), $EA_n^{n-1} \setminus W$ is connected since $|W| = n-2$ and $\kappa(EA_n^{n-1}) = n-1$. Note that there is a $\{x'_{n-1}, y'_{n-1}, z'_{n-1}, w\}$ -tree T'_{n-1} in $EA_n^{n-1} \setminus W$. Let

$$T_{n-1} = T'_{n-1} \cup P[x, x'_{n-1}] \cup P[y, y'_{n-1}] \cup P[z, z'_{n-1}].$$

Denote F the subgraph of EA_n^n induced on $\{x, y, z\}$. Clearly, $|E(F)| \leq 3$. The following cases need to be considered.

Case 1. $|E(F)| = 0$.

Clearly, T_1, \dots, T_{n-1} are $(n-1)$ desired IDSTs in EA_n .

Case 2. $|E(F)| = 1$.

For simplicity, assume that $E(F) = \{xy\}$. Consider the out-neighbours of x and y . The following discussions are based on whether x' and y' belong to a same cluster of EA_n or not.

Subcase 2.1. There exists an integer $i \in [n-1]$ with $\{x', y'\} \subseteq V(EA_n^i)$.

We discuss the case that $i = n-2$. For the other cases, $(n-1)$ IDSTs can be obtained by similar arguments.

It is inferred that $x = y_{n-2}$ and $y = x_{n-2}$. There is a $\{x', z'_{n-2}, w'_{n-2}\}$ -tree T''_{n-2} in EA_n^{n-2} . Let

$$\widehat{T}_{n-2} = T''_{n-2} \cup P[z, z'_{n-2}] \cup P[w, w'_{n-2}] \cup \{xx', xy\}.$$

See Figure 4. Then $T_1, \dots, T_{n-3}, \widehat{T}_{n-2}, T_{n-1}$ are $(n-1)$ IDSTs in EA_n .

Subcase 2.2. x' and y' belong to different clusters of EA_n .

By symmetry, assume that $y' \in V(EA_n^{n-1})$ and $x' \in V(EA_n^{n-2})$. That means $x = y_{n-2}$ and $y = x_{n-1}$.

Recall that $x'_{n-2} \in V(EA_n^{n-2})$. Based on Lemma 2.5(3) and (4), either $x'w'_{n-2} \notin E(EA_n^{n-2})$ or $x'_{n-2}w'_{n-2} \notin E(EA_n^{n-2})$. We only consider the case that $x'w'_{n-2} \notin E(EA_n^{n-2})$. For the other case, the proof is similar. By Lemma 2.5(2), there is a neighbour of x' in EA_n^{n-2} , say u , such that $u' \in V(EA_n^{n-1})$. Clearly, $u \notin \{x'_{n-2}, z'_{n-2}, w'_{n-2}\}$.

There exists a $\{u', y', z'_{n-1}, w\}$ -tree T''_{n-1} in $EA_n^{n-1} \setminus W$. Meanwhile, there is a $\{x'_{n-2}, z'_{n-2}, w'_{n-2}\}$ -tree T''_{n-2} in $EA_n^{n-2} \setminus \{x', u\}$ since $EA_n^{n-2} \setminus \{x', u\}$ is connected.

Let

$$\widehat{T}_{n-2} = T''_{n-2} \cup P[x, x'_{n-2}] \cup P[z, z'_{n-2}] \cup P[w, w'_{n-2}] \cup \{xy\}$$




Figure 4: Illustration for Subcase 2.1




Figure 5: Illustration for Subcase 2.2

and

$$\widehat{T}_{n-1} = T''_{n-1} \cup P[z, z'_{n-1}] \cup \{xx', x'u, uu', yy'\}.$$

See Figure 5. Then $T_1, \dots, T_{n-3}, \widehat{T}_{n-2}, \widehat{T}_{n-1}$ are $(n-1)$ IDSTs in EA_n .

Case 3. $|E(F)| = 2$.

For simplicity, assume that $E(F) = \{xy, xz\}$. By Lemma 2.5(2), y' and z' belong to different clusters of EA_n since both y and z are neighbours of x in EA_n^n .

Subcase 3.1. $\{y', z'\} \cap V(EA_n^{n-1}) \neq \emptyset$.

Assume that $z' \in V(EA_n^{n-1})$ and $y' \in V(EA_n^{n-2})$. That means $z = x_{n-1}$ and $y = x_{n-2}$. Consider now the location of x' .




Figure 6: Illustration for Subcase 3.1.1




Figure 7: Illustration for Subcase 3.1.2

Subcase 3.1.1. $x' \in V(EA_n^{n-1})$.

Then $x = y_{n-1} = z_{n-1}$. There is a $\{x', z', w\}$ -tree T''_{n-1} in $EA_n^{n-1} \setminus W$ and a $\{y', z'_{n-2}, w'_{n-2}\}$ -tree T''_{n-2} in EA_n^{n-2} , respectively. Let

$$\widehat{T}_{n-2} = T''_{n-2} \cup P[z, z'_{n-2}] \cup P[w, w'_{n-2}] \cup \{xz, yy'\}$$

and

$$\widehat{T}_{n-1} = T''_{n-1} \cup \{xy, xx', zz'\}.$$

See Figure 6. It is seen that $T_1, \dots, T_{n-3}, \widehat{T}_{n-2}, \widehat{T}_{n-1}$ are $(n-1)$ IDSTs in EA_n .

Subcase 3.1.2. $x' \in V(EA_n^{n-2})$.

That means $x = y_{n-2} = z_{n-2}$. There is a $\{x', y', w'_{n-2}\}$ -tree T''_{n-2} in EA_n^{n-2} and a $\{y'_{n-1}, z', w\}$ -tree T''_{n-1} in $EA_n^{n-1} \setminus W$, respectively. Let

$$\widehat{T}_{n-2} = T''_{n-2} \cup P[w, w'_{n-2}] \cup \{xx', xz, yy'\}$$

and

$$\widehat{T}_{n-1} = T''_{n-1} \cup P[y, y'_{n-1}] \cup \{xy, zz'\}.$$

See Figure 7. Clearly, $T_1, \dots, T_{n-3}, \widehat{T}_{n-2}, \widehat{T}_{n-1}$ are $(n-1)$ IDSTs in EA_n .

Subcase 3.1.3. $x' \in \bigcup_{i=1}^{n-3} V(EA_n^i)$.

W.l.o.g., assume that $x' \in V(EA_n^{n-3})$, which means that $x = y_{n-3} = z_{n-3}$.

Recall that $x'_{n-3} \in V(EA_n^{n-3})$. According to Lemma 2.5(3) and (4), either $x'w'_{n-3} \notin E(EA_n^{n-3})$ or $x'_{n-3}w'_{n-3} \notin E(EA_n^{n-3})$. For convenience, we only consider the case that $x'w'_{n-3} \notin E(EA_n^{n-3})$. Let u be a neighbour of x' in EA_n^{n-3} such that $u' \in V(EA_n^{n-1})$ and v be a neighbour of y' in EA_n^{n-2} such that $v' \in V(EA_n^{n-3})$. It can be inferred that $u \neq w'_{n-3}$.

Let $F_1 = EA_n^{n-2} \setminus \{y', v\}$ and $F_2 = EA_n^{n-3} \setminus \{x', u\}$.

Clearly, both F_1 and F_2 are connected. Thus, there is a $\{y'_{n-2}, z'_{n-2}, w'_{n-2}\}$ -tree T''_{n-2} in F_1 and a $\{x'_{n-3}, v', w'_{n-3}\}$ -tree T''_{n-3} in F_2 , respectively. Moreover, there is a $\{y'_{n-1}, z', w, u'\}$ -tree T''_{n-1} in $EA_n^{n-1} \setminus W$. Let

$$\widehat{T}_{n-3} = T''_{n-3} \cup P[x, x'_{n-3}] \cup P[w, w'_{n-3}] \cup \{xz, yy', y'v, vv'\},$$

$$\widehat{T}_{n-2} = T''_{n-2} \cup P[y, y'_{n-2}] \cup P[z, z'_{n-2}] \cup P[w, w'_{n-2}] \cup \{xy\}$$

and

$$\widehat{T}_{n-1} = T''_{n-1} \cup P[y, y'_{n-1}] \cup \{xx', x'u, uu', zz'\}.$$

The trees $\widehat{T}_{n-3}, \widehat{T}_{n-2}$ and \widehat{T}_{n-1} are depicted in Figure 8. Thus, $T_1, \dots, T_{n-4}, \widehat{T}_{n-3}, \widehat{T}_{n-2}, \widehat{T}_{n-1}$ are $(n-1)$ IDSTs in EA_n .

Subcase 3.2. $\{y', z'\} \cap V(EA_n^{n-1}) = \emptyset$.

For simplicity, let $y' \in V(EA_n^{n-2})$ and $z' \in V(EA_n^{n-3})$. The next discussions are based on the location of x' .

Subcase 3.2.1. Either $x' \in V(EA_n^{n-2})$ or $x' \in V(EA_n^{n-3})$.




Figure 8: Illustration for Subcase 3.1.3




Figure 9: Illustration for Subcase 3.2.1

By symmetry, assume that $x' \in V(EA_n^{n-2})$. There is a $\{x', y', w'_{n-2}\}$ -tree T''_{n-2} in EA_n^{n-2} and a $\{y'_{n-3}, z', w'_{n-3}\}$ -tree T''_{n-3} in EA_n^{n-3} , respectively. Let

$$\widehat{T}_{n-3} = T''_{n-3} \cup P[y, y'_{n-3}] \cup P[w, w'_{n-3}] \cup \{xy, zz'\}$$

and

$$\widehat{T}_{n-2} = T''_{n-2} \cup P[w, w'_{n-2}] \cup \{xx', xz, yy'\}.$$

See Figure 9. Thus, $T_1, \dots, T_{n-4}, \widehat{T}_{n-3}, \widehat{T}_{n-2}, T_{n-1}$ are $(n-1)$ IDSTs in EA_n .

Subcase 3.2.2. $x' \in V(EA_n^i)$, where $i \in [n-4]$ or $i = n-1$.

We consider the case that $i = n-1$. For the other cases, the proofs are similar.

By similar arguments in Subcase 2.2 and Lemma 2.5, either $y'w'_{n-2} \notin E(EA_n^{n-2})$ or $y'_{n-2}w'_{n-2} \notin E(EA_n^{n-2})$. For simplicity, assume that $y'w'_{n-2} \notin E(EA_n^{n-2})$. Likewise, we may assume that $z'w'_{n-3} \notin E(EA_n^{n-3})$.

Let u be a neighbour of y' in EA_n^{n-2} such that $u' \in V(EA_n^{n-1})$ and v be a neighbour of z' in EA_n^{n-3} such that $v' \in V(EA_n^{n-1})$, respectively. It has $u \neq w'_{n-2}$ and $v \neq w'_{n-3}$ by former analysis.

Let $F_1 = EA_n^{n-2} \setminus \{y', u\}$ and $F_2 = EA_n^{n-3} \setminus \{z', v\}$.

We may obtain a $\{y'_{n-2}, z'_{n-2}, w'_{n-2}\}$ -tree T''_{n-2} in F_1 and a $\{y'_{n-3}, z'_{n-3}, w'_{n-3}\}$ -tree T''_{n-3} in F_2 since both F_1 and F_2 are connected. Moreover, there is a $\{x', u', v', w\}$ -tree T''_{n-1} in $EA_n^{n-1} \setminus W$. Let

$$\widehat{T}_{n-3} = T''_{n-3} \cup P[y, y'_{n-3}] \cup P[z, z'_{n-3}] \cup P[w, w'_{n-3}] \cup \{xz\},$$

$$\widehat{T}_{n-2} = T''_{n-2} \cup P[y, y'_{n-2}] \cup P[z, z'_{n-2}] \cup P[w, w'_{n-2}] \cup \{xy\}$$

and

$$\widehat{T}_{n-1} = T''_{n-1} \cup \{xx', yy', y'u, uu', zz', z'v, vv'\}.$$

See Figure 10. Therefore, $T_1, \dots, T_{n-4}, \widehat{T}_{n-3}, \widehat{T}_{n-2}, \widehat{T}_{n-1}$ are $(n-1)$ IDSTs in EA_n .




Figure 10: Illustration for Subcase 3.2.2




Figure 11: Illustration for Case 1

Case 4. $|E(F)| = 3$.

That is to say, $E(F) = \{xy, xz, yz\}$. Remind that $EA_n = AN_n^1 \otimes AN_n^2$. Either $\{x, y, z\} \subseteq V(AN_n^1)$ or $\{x, y, z\} \subseteq V(AN_n^2)$ by Lemma 2.5(4). By symmetry, assume that $\{x, y, z\} \subseteq V(AN_n^1)$. With the help of Lemma 3.1, we only need to consider the possibility that $w \in V(AN_n^2)$.

Consider now \tilde{w} , the parity neighbour of w . Claim 1 can be obtained easily.

Claim 1. $\tilde{w} \notin \{x, y, z\}$.

Proof of Claim 1. Suppose to contrary that $\tilde{w} = x$. Then $w = \tilde{x} \in V(EA_n^n)$. A contradiction to the assumption of this lemma. \square

Based on Lemma 2.5(4), Claim 2 may be followed immediately.

Claim 2. $|E(\tilde{w}, \{x, y, z\})| \leq 1$.

Therefore, Lemma 3.2 implies that there are $(n-1)$ IDSTs in EA_n .

In all, the proof is completed. \square

3.2 $\max_{i \in [n]} \{|S \cap V(EA_n^{m:i})|\} = 2, 4 \leq m \leq n$.

Lemma 3.4 For $n \geq 4$ and $4 \leq m \leq n$, let $EA_n = EA_n^{m:1} \oplus EA_n^{m:2} \oplus \dots \oplus EA_n^{m:n}$ and $S = \{x, y, z, w\}$ be any 4-subset of $V(EA_n)$. If there exist different integers i and j in $[n]$ such that $|S \cap V(EA_n^{m:i})| = |S \cap V(EA_n^{m:j})| = 2$. Then there are $(n-1)$ IDSTs in EA_n .

Proof W.l.o.g., assume that $\{x, y\} \subseteq V(EA_n^n)$ and $\{z, w\} \subseteq V(EA_n^{n-1})$.

By Lemma 2.2(4), $\kappa(EA_n^i) = \kappa(EA_{n-1}) = n-1$ for each $i \in [n]$. Combined with Lemma 1.3, there are $(n-1)$ internally disjoint (x, y) -paths P_1, \dots, P_{n-1} in

EA_n^n and $(n-1)$ internally disjoint (z, w) -paths Q_1, \dots, Q_{n-1} in EA_n^{n-1} , respectively. For convenience, let $x_i \in V(P_i)$ for $i \in [n-1]$.

Since EA_n^{n-1} is connected, there is a (x'_{n-1}, z) -path \tilde{P} in EA_n^{n-1} . Let u be the first vertex in $V(\tilde{P}) \cap (\bigcup_{i=1}^{n-1} V(Q_i))$ when \tilde{P} starts from x'_{n-1} . For convenience, assume that $u \in V(Q_{n-1})$.

Let $z(i)$ be the neighbour of z such that $z(i) \in V(Q_i)$, $i \in [n-1]$. Set

$$X = \{x'_1, \dots, x'_{n-2}\} \quad \text{and} \quad Z = \{z(1)', \dots, z(n-2)'\}.$$

According to Lemma 2.5(2), $X \cap V(EA_n^{n-1}) = \emptyset$ and $|Z \cap V(EA_n^n)| \leq 1$.

Case 1. $Z \cap V(EA_n^n) = \emptyset$.

By Lemma 2.6, $\kappa(\bigcup_{i=1}^{n-2} EA_n^i) \geq n-2$. With the help of Lemma 1.5, there are $(n-2)$ internally disjoint (X, Z) -paths L_1, \dots, L_{n-2} in $\bigcup_{i=1}^{n-2} EA_n^i$ such that $\{x'_i, z(i)'\} \subseteq V(L_i)$ for $i \in [n-2]$. For $i \in [n-2]$, let

$$T_i = P_i \cup Q_i \cup L_i \cup \{x_i x'_i, z(i) z(i)'\}.$$

Let $T_{n-1} = P_{n-1} \cup Q_{n-1} \cup \tilde{P}(x'_{n-1}, u) \cup \{x_{n-1} x'_{n-1}\}$. See Figure 11. Then T_1, \dots, T_{n-1} are $(n-1)$ IDSTs in EA_n .

Case 2. $|Z \cap V(EA_n^n)| = 1$.

For simplicity, we may assume that $z(n-2)' \in V(EA_n^n)$.

Subcase 2.1. $z(n-2)' \in \bigcup_{i=1}^{n-1} V(P_i)$.

Subcase 2.1.1. $z(n-2)' \in V(P_{n-1})$.

Let $Z_1 = \{z(1)', \dots, z(n-3)', z(n-1)'\}$. Note that $Z_1 \cap V(EA_n^n) = \emptyset$. By similar arguments in Case 1, there are $(n-2)$ internally disjoint (X, Z_1) -paths L_1, \dots, L_{n-2} in $\bigcup_{i=1}^{n-2} EA_n^i$ such that $\{x'_i, z(i)'\} \subseteq V(L_i)$ for $i \in [n-3]$ and $\{x'_{n-2}, z(n-1)'\} \subseteq V(L_{n-2})$.

Let $T_{n-2} = P_{n-2} \cup Q_{n-1} \cup L_{n-2} \cup \{x_{n-2} x'_{n-2}, z(n-1) z(n-1)'\}$ and $T_{n-1} = P_{n-1} \cup Q_{n-2} \cup \{z(n-2) z(n-2)'\}$. For $i \in [n-3]$, we may construct T_i be the same as in Case 1. Then T_1, \dots, T_{n-1} are $(n-1)$ IDSTs in EA_n .

Subcase 2.1.2. $z(n-2)' \in \bigcup_{i=1}^{n-2} V(P_i)$.

For convenience, assume that $z(n-2)' \in V(P_{n-2})$. Let $T_{n-2} = P_{n-2} \cup Q_{n-2} \cup \{z(n-2) z(n-2)'\}$ and T_i be the same as in Case 1 for $i \in [n-3]$ or $i = n-1$. Then T_1, \dots, T_{n-1} are $(n-1)$ IDSTs in EA_n .

Subcase 2.2. $z(n-2)' \notin \bigcup_{i=1}^{n-1} V(P_i)$.

Let u_1, \dots, u_{n-1} be $(n-1)$ neighbours of $z(n-2)'$ in EA_n^n . Firstly, we consider the case that $u_1 \in \bigcup_{i=1}^{n-1} V(P_i)$. By similar analysis in Subcase 2.1, we can obtain $(n-1)$ IDSTs in EA_n by replacing the edge $z(n-2) z(n-2)'$ by the path $\{u_1 z(n-2)', z(n-2) z(n-2)'\}$ in T_{n-1} when $u_1 \in V(P_{n-1})$ or in T_{n-2} when $u_1 \in V(P_{n-2})$.

Then we consider the case that $\{u_1, \dots, u_{n-1}\} \cap (\bigcup_{i=1}^{n-1} V(P_i)) = \emptyset$. By Lemma 2.5(2), there is a vertex in $\{u_1, \dots, u_{n-1}\}$, say u_1 , such that $u'_1 \in \bigcup_{i=1}^{n-2} V(EA_n^i)$. Let

$$Z_2 = \{z(1)', \dots, z(n-3)', u'_1\}.$$

Clearly, $|Z_2| = n - 2$ and $Z_2 \cap V(EA_n^n) = \emptyset$.

Combined with Lemma 1.5 and Lemma 2.6, there are $(n-2)$ internally disjoint (X, Z_2) -paths L_1, \dots, L_{n-2} in $\bigcup_{i=1}^{n-2} EA_n^i$ such that $\{x'_i, z(i)'\} \subseteq V(L_i)$ for $i \in [n-3]$ and $\{x'_{n-2}, u'_1\} \subseteq V(L_{n-2})$. Let

$$T_{n-2} = P_{n-2} \cup Q_{n-2} \cup L_{n-2} \cup \{x_{n-2}x'_{n-2}, z(n-2)z(n-2)', z(n-2)'u_1, u_1u'_1\}$$

and T_i be the same as in Case 1 for $i \in [n-3]$ or $i = n-1$. Then T_1, \dots, T_{n-1} are $(n-1)$ IDSTs in EA_n . The proof is done. \square

Lemma 3.5 *For $n \geq 4$ and $4 \leq m \leq n$, let $EA_n = EA_n^{m:1} \oplus EA_n^{m:2} \oplus \dots \oplus EA_n^{m:n}$ and $S = \{x, y, z, w\}$ be any 4-subset of $V(EA_n)$. If there exist different integers i, j and k in $[n]$ such that $|S \cap V(EA_n^{m:i})| = 2$ and $|S \cap V(EA_n^{m:j})| = |S \cap V(EA_n^{m:k})| = 1$. Then there are $(n-1)$ IDSTs in EA_n .*

Proof W.l.o.g., assume that $\{x, y\} \subseteq V(EA_n^n)$, $z \in V(EA_n^{n-1})$ and $w \in V(EA_n^{n-2})$.

There are $(n-1)$ internally disjoint (x, y) -paths P_1, \dots, P_{n-1} in EA_n^n . For simplicity, let $x_i \in V(P_i)$, $i \in [n-1]$.

For $i \in [n-3]$, there is a $\{x'_i, z'_i, w'_i\}$ -tree T'_i in EA_n^i since $\{x'_i, z'_i, w'_i\} \subseteq V(EA_n^i)$. Therefore, let $T_i = T'_i \cup P_i \cup P[z, z'_i] \cup P[w, w'_i] \cup \{x_i x'_i\}$ for $i \in [n-3]$.

Let $Z = \{z_1, \dots, z_{n-2}\}$ and $W = \{w_1, \dots, w_{n-3}, w_{n-1}\}$.

Clearly, $|Z| = |W| = n - 2$. Both $EA_n^{n-1} \setminus Z$ and $EA_n^{n-2} \setminus W$ are connected since $\kappa(EA_n^i) = n - 1$ for $i \in [n]$.

The following discussions are based on whether $w_{n-1} = z'_{n-2}$ or not.

Case 1. $w_{n-1} \neq z'_{n-2}$.

There is a $\{x'_{n-1}, z, w'_{n-1}\}$ -tree T'_{n-1} in $EA_n^{n-1} \setminus Z$ and a $\{x'_{n-2}, z'_{n-2}, w\}$ -tree T'_{n-2} in $EA_n^{n-2} \setminus W$, respectively. Let

$$T_{n-1} = T'_{n-1} \cup P_{n-1} \cup P[w, w'_{n-1}] \cup \{x_{n-1}x'_{n-1}\}$$

and

$$T_{n-2} = T'_{n-2} \cup P_{n-2} \cup P[z, z'_{n-2}] \cup \{x_{n-2}x'_{n-2}\}.$$

See Figure 12. Then T_1, \dots, T_{n-1} are $(n-1)$ IDSTs in EA_n .

Case 2. $w_{n-1} = z'_{n-2}$.

Now consider the location of z' .

Subcase 2.1. $z' \in \bigcup_{i=1}^{n-3} V(EA_n^i)$.




Figure 12: Illustration for Case 1




Figure 13: Illustration for Subcase 2.1

W.l.o.g., assume that $z' \in V(EA_n^{n-3})$. By similar analysis in Subcase 2.2 of Lemma 3.3, either $z'w'_{n-3} \notin E(EA_n^{n-3})$ or $z'_{n-3}w'_{n-3} \notin E(EA_n^{n-3})$. We only consider the possibility that $z'w'_{n-3} \notin E(EA_n^{n-3})$ for convenience. Thus, there is a neighbour of z' in EA_n^{n-3} , say u , such that $u' \in V(EA_n^{n-2})$.

There is a $\{x'_{n-1}, z, w'_{n-1}\}$ -tree T'_{n-1} in $EA_n^{n-1} \setminus \{z_1, \dots, z_{n-3}\}$, a $\{x'_{n-2}, w, u'\}$ -tree T'_{n-2} in $EA_n^{n-2} \setminus W$ and a $\{x'_{n-3}, z'_{n-3}, w'_{n-3}\}$ -tree T''_{n-3} in $EA_n^{n-3} \setminus \{z', u\}$, respectively.

Let

$$T_{n-1} = T'_{n-1} \cup P_{n-1} \cup P[w, w'_{n-1}] \cup \{x_{n-1}x'_{n-1}\},$$

$$T_{n-2} = T'_{n-2} \cup P_{n-2} \cup \{x_{n-2}x'_{n-2}, zz', z'u, uu'\}$$

and

$$\widehat{T}_{n-3} = T''_{n-3} \cup P_{n-3} \cup P[z, z'_{n-3}] \cup P[w, w'_{n-3}] \cup \{x_{n-3}x'_{n-3}\}.$$

See Figure 13. Then $T_1, \dots, T_{n-4}, \widehat{T}_{n-3}, T_{n-2}, T_{n-1}$ are $(n-1)$ IDSTs in EA_n .

Subcase 2.2. $z' \in V(EA_n^{n-2})$.

There is a $\{x'_{n-2}, z', w\}$ -tree T'_{n-2} in $EA_n^{n-2} \setminus W$. Let $T_{n-2} = T'_{n-2} \cup P_{n-2} \cup \{x_{n-2}x'_{n-2}, zz'\}$. For $i \in [n-3]$ or $i = n-1$, let T'_i and T_i be the same as in Case 1. Then T_1, \dots, T_{n-1} are $(n-1)$ IDSTs in EA_n .

Subcase 2.3. $z' \in V(EA_n^n)$.

Subcase 2.3.1. $z' \in \bigcup_{i=1}^{n-1} V(P_i)$.

By Lemma 2.5(3), it has $z_n = z \circ (12)$. There is a neighbour of z_n in EA_n^{n-1} , say v , such that $v' \in V(EA_n^{n-2})$. According to Lemma 2.5(4), $v \neq z_{n-2}$.

Subcase 2.3.1.1. $z' \in V(P_{n-1})$.

There is a $\{x'_{n-2}, v', w\}$ -tree T'_{n-2} in $EA_n^{n-2} \setminus W$. Let $T_{n-2} = T'_{n-2} \cup P_{n-2} \cup \{x_{n-2}x'_{n-2}, zz_n, z_nv, vv'\}$. Let $T_{n-1} = P_{n-1} \cup P[w, w'_{n-1}] \cup \{zz', zw'_{n-1}\}$ since $w'_{n-1} =$

z_{n-2} . Then T_1, \dots, T_{n-1} are $(n-1)$ IDSTs in EA_n .

Subcase 2.3.1.2. $z' \in V(P_{n-2})$.

There is a $\{x'_{n-1}, z, z_n, v\}$ -tree T'_{n-1} in $EA_n^{n-1} \setminus Z$ and a (w, v') -path \tilde{P} in $EA_n^{n-2} \setminus W$. Let $T_{n-2} = P_{n-2} \cup P[w, w'_{n-1}] \cup \{zz', zw'_{n-1}\}$ and $T_{n-1} = T'_{n-1} \cup P_{n-1} \cup \tilde{P} \cup \{x_{n-1}x'_{n-1}, vv'\}$. Then T_1, \dots, T_{n-1} are $(n-1)$ IDSTs in EA_n .

Subcase 2.3.1.3. $z' \in \bigcup_{i=1}^{n-3} V(P_i)$.

W.l.o.g., assume that $z' \in V(P_{n-3})$. Recall that $z_n = \tilde{z}$.

Let $F_1 = EA_n^{n-1} \setminus \{z_1, \dots, z_{n-4}, z_{n-2}, z_n, v\}$. According to Lemma 2.4, F_1 is still connected. There is a $\{x'_{n-1}, z_{n-3}, z\}$ -tree T'_{n-1} in F_1 and a $\{x'_{n-2}, v', w\}$ -tree T'_{n-2} in $EA_n^{n-2} \setminus W$. Moreover, there is a (z'_{n-3}, w'_{n-3}) -path \tilde{P} in EA_n^{n-3} .

Let

$$\begin{aligned}\hat{T}_{n-3} &= P_{n-3} \cup P[w, w'_{n-1}] \cup \{zz', zw'_{n-1}\}, \\ T_{n-2} &= T'_{n-2} \cup P_{n-2} \cup \{x_{n-2}x'_{n-2}, zz_n, z_nv, vv'\}\end{aligned}$$

and

$$T_{n-1} = T'_{n-1} \cup P_{n-1} \cup \tilde{P} \cup P[w, w'_{n-3}] \cup \{x_{n-1}x'_{n-1}, z_{n-3}z'_{n-3}\}.$$

See Figure 14. Then $T_1, \dots, T_{n-4}, \hat{T}_{n-3}, T_{n-2}, T_{n-1}$ are $(n-1)$ IDSTs in EA_n .





Figure 14: Illustration for Subcase 2.3.1.3 Figure 15: Illustration for Subcase 1.1.1

Subcase 2.3.2. $z' \notin \bigcup_{i=1}^{n-1} V(P_i)$.

Let u_1, \dots, u_{n-1} be $(n-1)$ neighbours of z' in EA_n^n .

Firstly, we consider the case that $u_1 \in \bigcup_{i=1}^{n-1} V(P_i)$. By similar methods in Subcase 2.3.1, we may construct $(n-1)$ IDSTs in EA_n just by replacing the edge zz' by the path $\{zz', z'u_1\}$ in T_i when $u_1 \in V(P_i)$, $n-3 \leq i \leq n-1$.

Then we move to the case that $\{u_1, \dots, u_{n-1}\} \cap (\bigcup_{i=1}^{n-1} V(P_i)) = \emptyset$. By Lemma 2.5(2), let u_{n-2} be a neighbour of z' in EA_n^n that $u'_{n-2} \in V(EA_n^{n-2})$. There is a

$\{x'_{n-1}, z, w'_{n-1}\}$ -tree T'_{n-1} in $EA_n^{n-1} \setminus \{z_1, \dots, z_{n-3}\}$ and a $\{x'_{n-2}, u'_{n-2}, w\}$ -tree T'_{n-2} in $EA_n^{n-2} \setminus W$, respectively.

Let

$$T_{n-1} = T'_{n-1} \cup P_{n-1} \cup P[w, w'_{n-1}] \cup \{x_{n-1}x'_{n-1}\}$$

and

$$T_{n-2} = T'_{n-2} \cup P_{n-2} \cup \{x_{n-2}x'_{n-2}, zz', z'u_{n-2}, u_{n-2}u'_{n-2}\}.$$

Then T_1, \dots, T_{n-1} are $(n-1)$ IDSTs in EA_n . □

3.3 $\max_{i \in [n]} \{|S \cap V(EA_n^{m:i})|\} = 1, 4 \leq m \leq n.$

Lemma 3.6 For $n \geq 4$ and $4 \leq m \leq n$, let $EA_n = EA_n^{m:1} \oplus EA_n^{m:2} \oplus \dots \oplus EA_n^{m:n}$ and $S = \{x, y, z, w\}$ be any 4-subset of $V(EA_n)$. If $|S \cap V(EA_n^{m:i})| \leq 1$ for each $i \in [n]$. Then there are $(n-1)$ IDSTs in EA_n .

Proof For simplicity, we assume that $x = 12 \dots n \in V(EA_n^n)$. We need to consider whether $\{y'_n, z'_n, w'_n\} \cap N_{EA_n^n}(x) = \emptyset$ or not.

Case 1. $y'_n \in N_{EA_n^n}(x)$.

Since $|S \cap V(EA_n^{m:i})| \leq 1$ for $4 \leq m \leq n$ and each $i \in [n]$, Claim 1-Claim 3 may be inferred from Definition 2.1.

Claim 1. $n \leq 6$.

Proof of Claim 1. Suppose to contrary that $n \geq 7$. We will conclude that there exists an integer j ($4 \leq j \leq n-1$) such that the j th symbol of y is j .

For $y'_n \in \{x \circ (12), x \circ (123), x \circ (132)\}$, the j th symbol of y_n is j for each $4 \leq j \leq n-1$ due to Definition 2.1. Then there exists an integer j ($4 \leq j \leq n-1$) such that the j th symbol of y is j no matter $y_n \in \{y \circ (12), y \circ (123), y \circ (132)\}$ or $y_n \in \{y \circ (12)(3i) | 4 \leq i \leq n-1\}$.

For $y'_n \in \{x \circ (12)(3i) | 4 \leq i \leq n-1\}$, the j th symbol of y_n is j for each $j \in [n-1] \setminus \{1, 2, 3, i\}$. Then there always exists an integer j ($4 \leq j \leq n-1$) such that the j th symbol of y is j when $n \geq 7$.

The above analysis means that $\{x, y\} \subseteq V(EA_n^{j:j})$, which is a contradiction to the condition of this Lemma 3.6. □

Claim 2. If $n = 6$. Then $y \in \{215364, 214635\}$.

Proof of Claim 2. By similar analysis in the proof of Claim 1, it has $y'_n \notin \{x \circ (12), x \circ (123), x \circ (132)\}$. Moreover, $y_n \notin \{y \circ (12), y \circ (123), y \circ (132)\}$. That means either $y'_n = 214356$ or $y'_n = 215436$. Then $y_n \in \{126354, 126435\}$ and $y \in \{215364, 214635\}$. □

Claim 3. If $n = 5$. Then $y \in \{21453, 31452, 23451, 21534, 51234, 25134, 21354\}$.

Proof of Claim 3. By similar above analysis, $y_n \notin \{y \circ (12), y \circ (123), y \circ (132)\}$ when $y'_n \in \{x \circ (12), x \circ (123), x \circ (132)\}$. Otherwise, $\{x, y\} \subseteq V(EA_5^{4:4})$, a

contradiction to the assumption of this lemma. Therefore, we have $y = 21453$ if $y'_n = x \circ (12)$, $y = 31452$ if $y'_n = x \circ (132)$ and $y = 23451$ if $y'_n = x \circ (123)$.

For $y'_n = x \circ (12)(34) = 21435$. Then $y_n = 12534$. Furthermore, we have $y \in \{21534, 51234, 25134, 21354\}$. \square

Based on Claim 1, we only need to focus on the case that $4 \leq n \leq 6$.

Subcase 1.1. $n = 6$.

According to Claim 2, $|\{z'_n, w'_n\} \cap N_{EA_n^n}(x)| \leq 1$.

Subcase 1.1.1. Either $z'_n \in N_{EA_n^n}(x)$ or $w'_n \in N_{EA_n^n}(x)$.

By symmetry, assume that $z'_n \in N_{EA_n^n}(x)$. Then $y = 214635$ and $z = 215364$ by Claim 2.

Note that $x_5 = y'_6 = 215436$, $y'_4 = 215634 \in V(EA_6^4)$, $x_4 = z'_6$ and $z_5 = 125364$. Clearly, $y'_4 \neq z_5$. Moreover, $w'_6 \notin N_{EA_6^6}(x)$. By Table 1, it is not difficult to check that, for any $w \in V(EA_n^i)$ ($1 \leq i \leq 3$), w cannot be adjacent to y'_i and z'_i at the same time.

Table 1:

vertex a	a_1	a'_1	a_2	a'_2	a_3	a'_3
$y = 214635$	421635	245631	142635	415632	123645	215643
$z = 215364$	521364	254361	152364	514362	123564	214563

W.l.o.g., assume that $w \in V(EA_6^3)$ and $wy'_3 \notin E(EA_6^3)$. That means $w'_5 \neq y_3$.

Let $F_6 = EA_6^6 \setminus \{x_1, x_2, x_3, x_5\}$, $F_5 = EA_6^5 \setminus \{y_1, y_2, y_3, y_4\}$, $F_4 = EA_6^4 \setminus \{z_1, z_2, z_3, z_5\}$ and $F_3 = EA_6^3 \setminus \{w_1, w_2, w_5, w_6\}$.

For $3 \leq i \leq 6$, F_i is connected. Hence, there is a $\{x'_3, y'_3, z'_3, w\}$ -tree T'_3 in F_3 , a $\{x'_4, y'_4, z\}$ -tree T'_4 in F_4 , a $\{x'_5, y, z'_5, w'_5\}$ -tree T'_5 in F_5 and a $\{x, x_4, w'_6\}$ -tree T'_6 in F_6 , respectively.

Let $T_3 = T'_3 \cup P[x, x'_3] \cup P[y, y'_3] \cup P[z, z'_3]$, $T_4 = T'_4 \cup T'_6 \cup P[y, y'_4] \cup P[w, w'_6] \cup \{x_4 x'_4\}$ and $T_5 = T'_5 \cup P[x, x'_5] \cup P[z, z'_5] \cup P[w, w'_5]$. See Figure 15.

For $1 \leq i \leq 2$, there is a $\{x'_i, y'_i, z'_i, w'_i\}$ -tree T'_i in EA_6^i , hereafter, let $T_i = T'_i \cup P[x, x'_i] \cup P[y, y'_i] \cup P[z, z'_i] \cup P[w, w'_i]$. Then T_1, \dots, T_5 are five IDSTs in EA_6 .

Subcase 1.1.2. $z'_n \notin N_{EA_n^n}(x)$ and $w'_n \notin N_{EA_n^n}(x)$.

W.l.o.g., assume that $y \in V(EA_6^5)$, $z \in V(EA_6^4)$ and $w \in V(EA_6^3)$.

Then $z'_6 \neq x_4$. We only need to consider the case that $z'_5 \neq y_4$ and $w'_5 \neq y_3$. Otherwise, the proof is similar to that of Subcase 1.1.1.

Under this circumstance, we may obtain five IDSTs T_1, \dots, T_5 again by the same method in Subcase 1.1.1.

Subcase 1.2. $n = 5$.

By Claim 3, it has $|\{y'_5, z'_5, w'_5\} \cap N_{EA_5^5}(x)| \leq 2$. Otherwise, either $|\{y, z, w\} \cap V(EA_5^{4:3})| \geq 2$ or $|\{y, z, w\} \cap V(EA_5^{4:5})| \geq 2$. Hence, assume that $w'_5 \notin N_{EA_5^5}(x)$.

Subcase 1.2.1. $z'_5 \in N_{EA_5^5}(x)$.

By Claim 3 and the fact that $|S \cap V(EA_5^{m:i})| \leq 1$ for $4 \leq m \leq 5$ and $i \in [5]$, it has $y \in \{21453, 31452, 23451\}$ and $z \in \{21534, 51234, 25134\}$. Note that $\{y', z\} \subseteq V(EA_5^4)$. Moreover, it has $z_5 = x'_4 = 12534$.

Subcase 1.2.1.1. $y = 21453$.

W.l.o.g., assume that $w \in V(EA_5^2)$. Note that $x' \in V(EA_5^3)$ and $y_5 = x'_3 = 12543$. Furthermore, $z_3 \neq y'_4$ when $z \in \{21534, 51234, 25134\}$.

Let $F_5 = EA_5^5 \setminus \{x_1, x_2, x_3\}$, $F_4 = EA_5^4 \setminus \{z_1, z_2, z_3\}$, $F_3 = EA_5^3 \setminus \{y_1, y_4, y_5\}$ and $F_2 = EA_5^2 \setminus \{w_1, w_3, w_5\}$.

Clearly, F_i is connected for $2 \leq i \leq 5$. Thus, there is a $\{x, x_4, w'_5\}$ -tree T'_5 in F_5 , a $\{x'_4, y'_4, z\}$ -tree T'_4 in F_4 , a $\{x', y, z'_3, w'_3\}$ -tree T'_3 in F_3 and a $\{x'_2, z'_2, w\}$ -tree T'_2 in F_2 , respectively.

Let $T_2 = T'_2 \cup P[x, x'_2] \cup P[x, x'_3] \cup P[z, z'_2] \cup \{x'_3 y\}$, $T_3 = T'_3 \cup P[z, z'_3] \cup P[w, w'_3] \cup \{x x'\}$ and $T_4 = T'_4 \cup T'_5 \cup P[y, y'_4] \cup P[w, w'_5] \cup \{x_4 x'_4\}$. See Figure 16. Let T'_1 and T_1 be the same as in Subcase 1.1.1. Then T_1, \dots, T_4 are four desired IDSTs in EA_5 .




Figure 16: Illustration for Subcase 1.2.1.1 Figure 17: Illustration for Subcase 1.2.1.2.1

Subcase 1.2.1.2. $y = 31452$.

It is seen from Table 2 that the 3rd and 4th symbols of y'_1 are different from those of z'_1 no matter $z = 25134$ or $z = 51234$ or $z = 21534$. By Definition 2.1, for any vertex $u \in V(EA_5^1)$, it cannot be adjacent to y'_1 and z'_1 at the same time.

When $z = 25134$, note that $z'_3 = 25413$ and $y'_3 = 41253$. For any vertex $u \in V(EA_5^3)$, it is has $|\{y'_3, z'_3\} \cap N_{EA_5^3}(u)| \leq 1$. When $z = 51234$, a vertex $u \in V(EA_5^3)$ may be adjacent to y'_3 and z'_3 at the same time if and only if $u = 14523$.

Subcase 1.2.1.2.1. $w \in V(EA_5^i)$ is a common neighbour of y'_i and z'_i , $i \in \{1, 3\}$.

Table 2:

vertex a	a_1	a'_1	a_3	a'_3
$y = 31452$	43152	34251	14352	41253
$z = 25134$	52134	25431	52314	25413
$z = 51234$	25134	52431	15324	51423
$z = 21534$	52134	25431	12354	21453

Table 3:

vertex a	a_2	a'_2	a_3	a'_3
$y = 23451$	34251	43152	42351	24153
$z = 25134$	51234	15432	52314	25413
$z = 51234$	15234	51432	15324	51423
$z = 21534$	15234	51432	12354	21453

By above analysis, there are only two possibilities: (1) $z = 51234$ and $w = 14523$; (2) $z = 21534$ and $w \in V(EA_5^3)$.

Note that $\{x, y, z, w\} \in V(AN_5^1)$ when $z = 51234$ and $w = 14523$. Then there are four IDSTs in EA_5 by Lemma 3.1.

Now we consider the possibility that $z = 21534$ and $w \in V(EA_5^3)$. Remind that $w'_5 \neq x_3$. Note that $z' \in V(EA_5^5)$, $x'_2 = y_5$, $x'_4 = z_5$ and $z'_3 = w_4$.

Let $F_5 = EA_5^5 \setminus \{x_1, x_3, x_4\}$, $F_4 = EA_5^4 \setminus \{z_1, z_2\}$, $F_3 = EA_5^3 \setminus \{w_1, w_4, w_5\}$ and $F_2 = EA_5^2 \setminus \{y_1, y_5\}$. Note that F_i is connected for $2 \leq i \leq 5$. Thus, there is a $\{x, y'_5, z', w'_5\}$ -tree T'_5 in F_5 , a $\{x'_4, y', z, w'_4\}$ -tree T'_4 in F_4 , a $\{x'_3, w, w_2\}$ -tree T'_3 in F_3 and a $\{y, z'_2, w'_2\}$ -tree T'_2 in F_2 .

Let $T_2 = T'_2 \cup T'_3 \cup P[x, x'_3] \cup P[z, z'_2] \cup \{w_2 w'_2\}$, $T_4 = T'_4 \cup P[x, x'_4] \cup P[w, w'_4] \cup \{yy'\}$ and $T_5 = T'_5 \cup P[y, y'_5] \cup P[w, w'_5] \cup \{zz'\}$. See Figure 17. Let T_1 be the same as in Subcase 1.1.1. Then T_1, T_2, T_4 and T_5 are four IDSTs in EA_5 .

Subcase 1.2.1.2.2. Either $wy'_i \notin E(EA_5^i)$ or $wz'_i \notin E(EA_5^i)$, where $w \in V(EA_5^i)$ and $i \in \{1, 3\}$.

W.l.o.g., assume that $w \in V(EA_5^3)$ and $wy'_3 \notin E(EA_5^3)$. That means $y'_3 \neq w_2$.

Remind that $x'_3 \neq w_5$, $z \in V(EA_5^4)$ and $x_4 = z'_5$. Moreover, it has $y' \neq z_2$ when $z \in \{51234, 25134, 21534\}$.

There is a $\{x, z'_5, w'_5\}$ -tree T'_5 in $EA_5^5 \setminus \{x_1, x_2, x_3\}$, a $\{y', z, z_5\}$ -tree T'_4 in $EA_5^4 \setminus \{z_1, z_2, z_3\}$, a $\{x'_3, y'_3, z'_3, w\}$ -tree T'_3 in $EA_5^3 \setminus \{w_1, w_2, w_5\}$ and a $\{x'_2, y, z'_2, w'_2\}$ -tree T'_2 in $EA_5^2 \setminus \{y_1, y_3\}$, respectively.

Let $T_4 = T'_4 \cup T'_5 \cup P[w, w'_5] \cup \{yy', z_5 z'_5\}$, $T_3 = T'_3 \cup P[x, x'_3] \cup P[y, y'_3] \cup P[z, z'_3]$ and $T_2 = T'_2 \cup P[x, x'_2] \cup P[z, z'_2] \cup P[w, w'_2]$. See Figure 18. Let T_1 be the same as in Subcase 1.1.1. Then T_1, T_2, T_3 and T_4 are four IDSTs in EA_5 .

Subcase 1.2.1.3. $y = 23451$.

Based on Table 3 and similar arguments in Subcase 1.2.1.2, for $2 \leq i \leq 3$ and any vertex $w \in V(EA_5^i)$, w is adjacent to both y'_i and z'_i if and only if one of the two possibilities: (1) $z = 25134$ and $w = 42513$; (2) $z = 21534$ and $w \in V(EA_5^3)$.

Note that $\{x, y, z, w\} \subseteq V(AN_5^1)$ when $z = 25134$ and $w = 42513$, then there are four IDSTs in EA_5 according to Lemma 3.1. Hence, we only need to consider the possibility that $z = 21534$ and $w \in V(EA_5^3)$ if $w \in V(EA_5^i)$ is a common




Figure 18: Illustration for Subcase 1.2.1.2.2




Figure 19: Three IDSTs in EA_4

neighbour of y'_i and z'_i , $2 \leq i \leq 3$. Note that $y' \in V(EA_5^4)$ and $y' \neq z_1$ when $z = 21534$, moreover, $z' \in V(EA_5^5)$. By similar analysis in Subcase 1.2.1.2.1, four IDSTs can be obtained in EA_5 and the proof is omitted.

If $wy'_i \notin E(EA_5^i)$ or $wz'_i \notin E(EA_5^i)$, where $w \in V(EA_5^i)$ and $i \in \{2, 3\}$. Then we may obtain four IDSTs by similar discussions in Subcase 1.2.1.2.2.

Subcase 1.2.2. $z'_5 \notin N_{EA_5^5}(x)$.

W.l.o.g., assume that $z \in V(EA_5^4)$, $w \in V(EA_5^3)$ and $y \in V(EA_5^2)$. Moreover, we may assume that $\{z'_2, w'_2\} \cap N_{EA_5^2}(y) = \emptyset$. Otherwise, four IDSTs can be obtained by similar method in Subcase 1.2.1.

Note that there is a $\{x'_2, y, z'_2, w'_2\}$ -tree T'_2 in $EA_5^2 \setminus \{y_1, y_3, y_4\}$ and a $\{y'_4, z, z_5\}$ -tree T'_4 in $EA_5^4 \setminus \{z_1, z_2, z_3\}$. Let T'_i ($i \in \{1, 3, 5\}$) and T_i ($i \in [3]$) be the same as in Subcase 1.2.1.2.2. Set $T_4 = T'_4 \cup T'_5 \cup P[w, w'_5] \cup P[y, y'_4] \cup \{z_5 z'_5\}$. Then T_1, \dots, T_4 are four IDSTs in EA_5 .

Subcase 1.3. $n = 4$.

W.l.o.g., assume that $y \in V(EA_4^3)$, $z \in V(EA_4^2)$ and $w \in V(EA_4^1)$. Then $y \in \{2143, 4123, 2413\}$ by Figure 3. By symmetry, we only need to consider the case that $y = 2143$ or $y = 2413$.

Subcase 1.3.1. $y = 2143$.

Recall that $EA_4 = AN_4^1 \otimes AN_4^2$ and $\{x, y\} \subseteq V(AN_4^1)$. We conclude from Lemma 3.1 that three IDSTs may be constructed if $\{z, w\} \subseteq V(AN_4^1)$. Moreover, if $\{z, w\} \cap V(AN_4^1) = \{z\}$, then three IDSTs may be obtained by Lemma 3.2 since $\tilde{w} \in V(EA_4^1)$ and $|E(\tilde{w}, \{x, y, z\})| \leq 1$. Therefore, we only need to consider the case that $\{z, w\} \subseteq V(AN_4^2)$.

Let $V_1 = \{1432, 4312, 3142\}$ and $V_2 = \{2341, 3421, 4231\}$. Then $z \in V_1$ and $w \in V_2$.

Note that the subgraph induced on V_i is a clique, $1 \leq i \leq 2$. By viewing V_i ($1 \leq i \leq 2$) as a vertex, Figure 19 shows that there are three internally edge

disjoint $\{x, y, V_1, V_2\}$ -trees in EA_4 , where three trees are represented by black solid lines, blue dotted lines and red dashed lines, respectively.

From the trees shown in Figure 19, we can obtain three desired IDSTs in EA_4 by adding two edges $z(z \circ (123))$ and $z(z \circ (132))$ in V_1 and by adding two edges $w(w \circ (123))$ and $w(w \circ (132))$ in V_2 .




Figure 20: Three IDSTs in EA_4




Figure 21: Illustration for Case 2

Subcase 1.3.2. $y = 2413$.

By similar analysis in Subcase 1.3.1, we only need to consider the possibility that $z \in V(AN_4^1)$ and $w \in V(AN_4^2)$. Figure 20 shows that we can find three IDSTs in EA_4 by adding four corresponding edges when $z \in \{1342, 4132, 3412\}$ and $w \in \{2341, 3421, 4231\}$.

Case 2. $\{y'_n, z'_n, w'_n\} \cap N_{EA_n^n}(x) = \emptyset$.

W.l.o.g., assume that $y \in V(EA_n^{n-1})$, $z \in V(EA_n^{n-2})$ and $w \in V(EA_n^{n-3})$. Furthermore, we may assume that $\{z'_{n-1}, w'_{n-1}\} \cap N_{EA_n^{n-1}}(y) = \emptyset$ and $z_{n-3} \neq w'_{n-2}$. Otherwise, similar arguments in Case 1 may be applied to obtain $(n-1)$ IDSTs in EA_n .

For $i \in [n-4]$, there is a $\{x'_i, y'_i, z'_i, w'_i\}$ -tree T'_i in EA_n^i since $\{x'_i, y'_i, z'_i, w'_i\} \subseteq V(EA_n^i)$. Let

$$T_i = T'_i \cup P[x, x'_i] \cup P[y, y'_i] \cup P[z, z'_i] \cup P[w, w'_i], \quad i \in [n-4].$$

Denote

$$F_{n-1} = EA_n^{n-1} \setminus \{y_1, \dots, y_{n-2}\}, \quad F_{n-2} = EA_n^{n-2} \setminus \{z_1, \dots, z_{n-3}, z_{n-1}\}$$

and $F_{n-3} = EA_n^{n-3} \setminus \{w_1, \dots, w_{n-4}, w_{n-2}, w_{n-1}\}$.

Clearly, the subgraph F_i is connected for $n-3 \leq i \leq n-1$. Hereafter, there is a $\{x'_{n-1}, y, z'_{n-1}, w'_{n-1}\}$ -tree T'_{n-1} in F_{n-1} , a $\{x'_{n-2}, y'_{n-2}, z, w'_{n-2}\}$ -tree T'_{n-2} in F_{n-2} and a $\{x'_{n-3}, y'_{n-3}, z'_{n-3}, w\}$ -tree T'_{n-3} in F_{n-3} , respectively.

Let

$$T_{n-1} = T'_{n-1} \cup P[x, x'_{n-1}] \cup P[z, z'_{n-1}] \cup P[w, w'_{n-1}],$$

$$T_{n-2} = T'_{n-2} \cup P[x, x'_{n-2}] \cup P[y, y'_{n-2}] \cup P[w, w'_{n-2}]$$

and

$$T_{n-3} = T'_{n-3} \cup P[x, x'_{n-3}] \cup P[y, y'_{n-3}] \cup P[z, z'_{n-3}].$$

See Figure 21. Then T_1, \dots, T_{n-1} are $(n-1)$ desired IDSTs in EA_n . \square

4 Proof of Theorem 1.1

In this section, we are going to prove Theorem 1.1 by induction on n . Firstly, we focus on the induction basis for $n = 3$.

Lemma 4.1 $\kappa_4(EA_3) \geq 2$.

Proof It suffices to prove that there are two IDSTs in EA_3 for any 4-subset $S = \{x, y, z, w\} \subseteq V(EA_3)$. Note that $EA_3 = AN_3^1 \otimes AN_3^2$ and $|V(AN_3^i)| = 3$ for $1 \leq i \leq 2$. The following two cases are discussed.

Case 1. $|S \cap V(AN_3^i)| = 3$ for $1 \leq i \leq 2$.

By symmetry, we may assume that $x = 123$, $y = 231$, $z = 312$ and $w = 213$. Then two IDSTs are shown in Figure 22, where two trees are represented by black solid lines and red dashed lines, respectively.




Figure 22: Two IDSTs in EA_3




Figure 23: Two IDSTs in EA_3




Figure 24: Two IDSTs in EA_3

Case 2. $|S \cap V(AN_3^1)| = |S \cap V(AN_3^2)| = 2$.

W.l.o.g., we may assume that $x = 123$ and $y = 231$. By investigating the structure of EA_3 , two IDSTs are shown in Figure 23 if $\{z, w\} = \{213, 321\}$ and in Figure 24 if $\{z, w\} = \{213, 132\}$, respectively. \square

Now, we are in a position to prove our main result.

Proof of Theorem 1.1. Combined with Lemma 1.1 and Lemma 2.2(1), $\kappa_4(EA_n) \leq \delta(EA_n) - 1 = n - 1$ for $n \geq 3$. We shall prove the reverse inequality by induction on n . First of all, $\kappa_4(EA_3) \geq 2$ by Lemma 4.1. Now suppose that $n \geq 4$ and the result is true for any integer $l < n$, i.e., $\kappa_4(EA_l) \geq l - 1$. Let $S = \{x, y, z, w\}$ be any 4-subset of $V(EA_n)$.

Remind that $EA_n = EA_n^{m:1} \oplus EA_n^{m:2} \oplus \cdots \oplus EA_n^{m:n}$, $4 \leq m \leq n$. The following cases are distinguished.

Case 1. There exists an integer $i \in [n]$ such that $S \subseteq V(EA_n^{m:i})$.

W.l.o.g., we may assume that $S \subseteq V(EA_n^n)$. According to the fact that EA_n^n is isomorphic to EA_{n-1} and our induction hypothesis, there are $(n-2)$ IDSTs T_1, \dots, T_{n-2} in EA_n^n .

By Lemma 2.6, there exists a $\{x', y', z', w'\}$ -tree T'_{n-1} in $\bigcup_{i=1}^{n-1} EA_n^i$. Let $T_{n-1} = T'_{n-1} \cup \{xx', yy', zz', ww'\}$. Clearly, $T_1, \dots, T_{n-2}, T_{n-1}$ are $(n-1)$ IDSTs in EA_n .

Case 2. There exists an integer $i \in [n]$ such that $|S \cap V(EA_n^{m:i})| = 3$.

With the help of Lemma 3.3, we may get $(n-1)$ IDSTs in EA_n .

Case 3. There exist different integers i and j in $[n]$ such that $|S \cap V(EA_n^{m:i})| = |S \cap V(EA_n^{m:j})| = 2$.

Then $(n-1)$ IDSTs in EA_n can be constructed due to Lemma 3.4.

Case 4. There exist different integers i, j and k in $[n]$ such that $|S \cap V(EA_n^{m:i})| = 2$ and $|S \cap V(EA_n^{m:j})| = |S \cap V(EA_n^{m:k})| = 1$.

Based on Lemma 3.5, there are $(n-1)$ IDSTs in EA_n .

Case 5. $|S \cap V(EA_n^{m:i})| \leq 1$ for each $i \in [n]$.

By Lemma 3.6, $(n-1)$ IDSTs in EA_n can be obtained. The proof is done. \square

5 Conclusion

The generalized k -connectivity is a natural generalization of the classical connectivity and can serve for measuring the capability of a network G to connect any k vertices in G . In this paper, we focus on the generalized 4-connectivity of the n -dimensional godan graph EA_n . Together with Theorem 1.1 and Lemma 1.2, the generalized 3-connectivity of EA_n can be derived immediately. This result was also investigated in [18].

References

- [1] J.A. Bondy, U.S.R. Murty, Graph theory, Springer, New York, 2007
- [2] G. Chartrand, S.F. Kapoor, L. Lesniak, D.R. Lick, Generalized connectivity in graphs, Bombay Math., 2 (1984) 1-6
- [3] G. Chartrand, F. Okamoto, P. Zhang, Rainbow trees in graphs and generalized connectivity, Networks, 55 (4) (2010) 360-367
- [4] E. Cheng, K. Qiu, Z. Shen, A note on the alternating group network, J. Supercomput., 59 (2012) 246-248

- [5] Y. Ji, A class of Cayley networks based on the alternating groups, *Adv. Math. (Chinese)*, 4 (1998) 361-362
- [6] C. Li, S. Lin, S. Li, The 4-set tree connectivity of (n, k) -star networks, *Theoret. Comput. Sci.*, 844 (2020) 81-86
- [7] H. Li, X. Li, Y. Mao, Y. Sun, Note on the generalized connectivity, *Ars Combin.*, 114 (2014) 193-202
- [8] H. Li, X. Li, Y. Sun, The generalized 3-connectivity of Cartesian product graphs, *Discrete Math. Theor. Comput. Sci.*, 14 (1) (2012) 43-54
- [9] S. Li, Some topics on generalized connectivity of graphs (Ph.D. thesis), Nankai University, (2012)
- [10] S. Li, W. Li, X. Li, The generalized connectivity of complete bipartite graphs, *Ars Combin.*, 104 (2012) 65-79
- [11] S. Li, X. Li, W. Zhou, Sharp bounds for the generalized connectivity $\kappa_3(G)$, *Discrete Math.*, 310 (2010) 2147-2163
- [12] S. Li, X. Li, Note on the hardness of generalized connectivity, *J. Comb. Optim.*, 24 (2012) 389-396
- [13] S. Li, Y. Shi, J. Tu, The generalized 3-connectivity of Cayley graphs on symmetric groups generated by trees and cycles, *Graphs and Combin.*, 33 (2017) 1195-1209
- [14] S. Li, J. Tu, C. Yu, The generalized 3-connectivity of star graphs and bubble-sort graphs, *Appl. Math. Comput.*, 274 (2016) 41-46
- [15] S. Lin, Q. Zhang, The generalized 4-connectivity of hypercubes, *Discrete Appl. Math.*, 220 (2017) 60-67
- [16] Y. Ren, S. Wang, Reliability analysis of godan graphs, *Discrete Appl. Math.*, 307 (2022) 180-190
- [17] J. Wang, The generalized 3-connectivity of two kinds of regular networks, *Theoret. Comput. Sci.*, 893 (2021) 183-190
- [18] J. Wang, Z. Zhang, Y. Huang, The generalized 3-connectivity of burnt pancake graphs and godan graphs, *AKCE International J. of Graphs and Combin.*, 20 (2023) 98-103
- [19] C. Wei, R.-X. Hao, J.-M. Chang, The reliability analysis based on the generalized connectivity in balanced hypercubes, *Discrete Appl. Math.*, 292 (2021) 19-32

- [20] H. Whitney, Congruent graphs and connectivity of graphs, *J. Amer. Math. Soc.*, 54 (1932) 150-168
- [21] H. Zhang, S. Zhou, Q. Zhang, Component connectivity of alternating group networks and godan graphs, *International J. Foundations of Comput. Sci.*, 34 (2023) 395-410
- [22] S.-L. Zhao, R.-X. Hao, The generalized 4-connectivity of exchanged hypercubes, *Applied Math. Comput.*, 347 (2019) 342-353
- [23] S.-L. Zhao, R.-X. Hao, The generalized connectivity of bubble-sort star graphs, *International J. Foundations of Comput. Sci.*, 30 (2019) 793-809
- [24] S.-L. Zhao, R.-X. Hao, The generalized connectivity of alternating group graphs and (n, k) -star graphs, *Discrete Appl. Math.*, 251 (2018) 310-321
- [25] S.-L. Zhao, J.-M. Chang, H.-Z. Li, The generalized 4-connectivity of pancake graphs, *Discrete Appl. Math.*, 327 (2023) 77-86
- [26] S.-L. Zhao, R.-X. Hao, E. Cheng, Two kinds of generalized connectivity of dual cubes, *Discrete Appl. Math.*, 257 (2019) 306-316
- [27] S.-L. Zhao, R.-X. Hao, S.-L. Peng, Reliability assessment of some regular networks, *The Computer Journal*, 64 (2021) 1-6