

MODULE-THEORETIC APPROACH TO DUALIZABLE GROTHENDIECK CATEGORIES

RYO KANDA

ABSTRACT. We prove that every dualizable Grothendieck category whose dual is again a Grothendieck category satisfies Grothendieck's conditions Ab6 and Ab4*, by taking a module-theoretic approach based on the Gabriel-Popescu embedding. Combining this with a result by Stefanich, we conclude that the class of dualizable linear cocomplete categories is precisely the class of linear Grothendieck category satisfying Ab6 and Ab4*. This provides a complete answer to a modified conjecture on the dualizability, originally posed by Brandenburg, Chirvasitu, and Johnson-Freyd.

CONTENTS

1. Introduction	1
Acknowledgments	2
2. Preliminaries	2
2.1. Dualizability of locally presentable categories	3
2.2. Gabriel-Popescu embedding and tensor products	3
2.3. Roos categories	5
3. Main theorem	6
References	8

1. INTRODUCTION

In [Roo65], Jan-Erik Roos proved the following result:

Theorem 1.1 ([Roo65]). *Let \mathcal{C} be a Grothendieck category. Then the following are equivalent:*

- (1) \mathcal{C} satisfies Grothendieck's conditions Ab6 and Ab4*.
- (2) There is a pair of a ring Λ and an idempotent ideal I of Λ such that \mathcal{C} is the quotient category of $\text{Mod } \Lambda$ by the bilocalizing subcategory $\text{Mod}(\Lambda/I)$.

In this paper, we refer to a Grothendieck category that satisfies Ab6 and Ab4* as a *Roos category*.

Grothendieck's condition Ab6 asserts the distributive property for intersections of filtered unions (see Definition 2.8), and Ab4* asserts that direct products are exact. The category of modules over an arbitrary ring satisfies Ab6 and Ab4*. Every locally noetherian Grothendieck category satisfies Ab6 ([Roo67]; see also [Kan19, Example 2.27]) but does not necessarily satisfy Ab4*. So Theorem 1.1 can be regarded as a characterization of Grothendieck categories that have exact direct products, under the mild assumption Ab6. Theorem 1.1 has been used to investigate the (non-)exactness of direct products for various classes of Grothendieck categories, including the author's result that asserts that the category of quasi-coherent sheaves over a divisorial noetherian scheme has exact direct products if and only if X is an affine scheme ([Kan19, Theorem 1.1]).

2020 *Mathematics Subject Classification*. 18E10 (Primary), 16D90, 18C35, 18E20 (Secondary).

Key words and phrases. Dualizability; Grothendieck category; Roos category; Gabriel-Popescu embedding; idempotent ring.

On the other hand, the class of Grothendieck categories satisfying the condition in Theorem 1.1 (2) is fundamental in the study of (non-unital) *idempotent rings*, that is, rings A that do not necessarily have identities but satisfy $A^2 = A$. For an idempotent ring A , the category of all A -modules is not really the category of interest because it is equivalent to the category of modules over the Dorroh overring of A , which is a unital ring (see [Wis91, sections 1.5 and 6.3]). Instead, one of the categories of interest is the category of *firm A -modules*, which are right A -modules M such that the canonical homomorphism $M \otimes_A A \rightarrow M$ is an isomorphism. It is known that the categories arising as the category of firm modules over an idempotent ring are precisely the Grothendieck categories satisfying the condition in Theorem 1.1 (2) (see [Qui96, section 7] or [Mar97]). Thus, Theorem 1.1 can also be regarded as a characterization of the category of firm modules over an idempotent ring by a purely categorical property.

In [BCJF15], Brandenburg, Chirvasitu, and Johnson-Freyd studied the reflexivity and dualizability of locally presentable linear categories (over a field). It is known that every Grothendieck category is locally presentable. They provide several examples and non-examples of reflexive categories and dualizable categories, and in [BCJF15, Remark 3.6], they conjectured that every dualizable locally presentable linear category is strongly generated by compact projective objects. Although there has been a result that partially confirms the conjecture ([Chi22]), there is a counterexample ([Ste23, Example 3.1.24]). Indeed, as a consequence of their own result in the same paper ([BCJF15, Corollary 3.4]), it follows that every Roos category is dualizable (see Corollary 3.2), and there is a nonzero Roos category that has no nonzero projective objects, constructed by Roos ([Roo66, Example 3]). Therefore, the conjecture should be modified as follows:

Conjecture 1.2. Every dualizable locally presentable linear category is a Roos category.

The aim of this paper is to prove the following result by taking a module-theoretic approach based on the Gabriel-Popescu embedding:

Theorem 1.3 (Theorem 3.5). *Let R be a commutative ring, and let \mathcal{C} be an R -linear Grothendieck category. If \mathcal{C} is dualizable and its dual is again a Grothendieck category, then \mathcal{C} is a Roos category.*

Our theorem only deals with Grothendieck categories whose duals are Grothendieck categories, but in fact, Stefanich [Ste23, Corollary 3.1.19] has already proved that every dualizable linear *cocomplete* category is a Grothendieck category satisfying $\text{Ab}4^*$, and thus its dual is again a Grothendieck category. So, combined with our result, we conclude the following, confirming Conjecture 1.2:

Corollary 1.4 (Corollary 3.7). *Let R be a commutative ring, and let \mathcal{C} be an R -linear cocomplete category. Then the following are equivalent:*

- (1) \mathcal{C} is dualizable.
- (2) \mathcal{C} is a Roos category.

Acknowledgments. The author was supported by JSPS KAKENHI Grant Numbers JP21H04994 and JP24K06693, and Osaka Central Advanced Mathematical Institute: MEXT Joint Usage/Research Center on Mathematics and Theoretical Physics JPMXP0723833165.

2. PRELIMINARIES

Convention 2.1. We use the following convention in this paper: In an abelian category, *direct sums* mean coproducts, *direct products* mean products. All colimits and limits are assumed to be small.

A *ring* means an associative ring with identity, unless otherwise specified. An associative ring that may not have identity will be referred to as a *non-unital ring*. An *idempotent ring* is a non-unital ring A such that $A^2 = A$.

Throughout the paper, we fix a commutative ring R , which will be used as a base ring. An *R -algebra* A means a ring A equipped with a ring homomorphism from R to the center of A , which is equivalent to an R -linear category with a single object. $\text{Mod } A$ denotes the category of right A -modules.

2.1. Dualizability of locally presentable categories. We recall the notion of dualizability and some related results; see [BCJF15] for more details. [BCJF15] only dealt with the case where R is a field, but the results used in this paper are also valid over an arbitrary commutative ring R .

For locally presentable R -linear categories \mathcal{C} and \mathcal{D} , denote by $\text{Cocont}_R(\mathcal{C}, \mathcal{D})$ the category of R -linear cocontinuous functors from \mathcal{C} to \mathcal{D} , with morphisms being the natural transformations. Recall that a functor is *cocontinuous* if it preserves all (small) colimits. It is known that $\text{Cocont}_R(\mathcal{C}, \mathcal{D})$ is again a locally presentable R -linear categories. If $R = \mathbb{Z}$, then $\text{Cocont}_{\mathbb{Z}}(\mathcal{C}, \mathcal{D})$ consists of all cocontinuous functors (which are necessarily additive) functors.

There is a locally presentable R -linear category $\mathcal{C} \boxtimes_R \mathcal{D}$ and a bifunctor $\otimes_R: \mathcal{C} \times \mathcal{D} \rightarrow \mathcal{C} \boxtimes_R \mathcal{D}$ that is cocontinuous and R -linear in each variable satisfying the following universal property: For an R -linear cocomplete category \mathcal{E} , the R -linear functor

$$\begin{array}{ccc} \text{Cocont}_R(\mathcal{C} \boxtimes_R \mathcal{D}, \mathcal{E}) & \rightarrow & \text{Cocont}_R(\mathcal{C}, \mathcal{D}; \mathcal{E}) \\ \downarrow \Psi & & \downarrow \Psi \\ F & \mapsto & F \circ \otimes_R \end{array}$$

is an equivalence, where $\text{Cocont}_R(\mathcal{C}, \mathcal{D}; \mathcal{E})$ denotes the category of bifunctors $\mathcal{C} \times \mathcal{D} \rightarrow \mathcal{E}$ that are cocontinuous and R -linear in each variable.

Define the *dual* of \mathcal{C} to be $\mathcal{C}^* := \text{Cocont}_R(\mathcal{C}, \text{Mod } R)$. Since $\text{Cocont}_R(\mathcal{C}, \text{Mod } R) = \text{Cocont}_{\mathbb{Z}}(\mathcal{C}, \text{Mod } \mathbb{Z})$, the definition of \mathcal{C}^* does not depend on the choice of the base ring R . There is a canonical functor

$$\begin{array}{ccc} \mathcal{C}^* \boxtimes_R \mathcal{D} & \rightarrow & \text{Cocont}_R(\mathcal{C}, \mathcal{D}) \\ \downarrow \Psi & & \downarrow \Psi \\ F \otimes_R \mathcal{D} & \mapsto & (\mathcal{C} \mapsto F(\mathcal{C}) \otimes_R \mathcal{D}) \end{array}$$

where the functor $-\otimes_R \mathcal{D}: \text{Mod } R \rightarrow \mathcal{D}$ is defined to be the left adjoint of $\text{Hom}_{\mathcal{D}}(\mathcal{D}, -): \mathcal{D} \rightarrow \text{Mod } R$ (the left adjoint exists because the functor preserves all limits).

Definition 2.2. A locally presentable R -linear category \mathcal{C} is called *dualizable* if the canonical functor $\mathcal{C}^* \boxtimes_R \mathcal{C} \rightarrow \text{Cocont}_R(\mathcal{C}, \mathcal{C})$ is an equivalence.

Every Grothendieck category is locally presentable. For an R -algebra A , the category $\text{Mod } A$ of right A -modules is an R -linear Grothendieck category, hence locally presentable. For R -algebras A and Γ , the Eilenberg-Watts theorem asserts that there is an R -linear equivalence

$$\begin{array}{ccc} \text{Mod}(A^{\text{op}} \otimes_R \Gamma) & \xrightarrow{\simeq} & \text{Cocont}_R(\text{Mod } A, \text{Mod } \Gamma) \\ \downarrow \Psi & & \downarrow \Psi \\ M & \mapsto & - \otimes_A M. \end{array}$$

Note that $\text{Mod}(A^{\text{op}} \otimes_R \Gamma)$ is canonically equivalent to the category of A - Γ -bimodules over R . In particular, $(\text{Mod } A)^* = \text{Cocont}_R(\text{Mod } A, \text{Mod } R) \cong \text{Mod}(A^{\text{op}})$.

It is shown in [BCJF15, Lemma 3.5] that every locally presentable R -linear category that admits a strongly generating set consisting of compact projective objects is dualizable. In particular, $\text{Mod } A$ is dualizable since $A \in \text{Mod } A$ is a compact projective object and the singleton $\{A\}$ is a strongly generating set.

2.2. Gabriel-Popescu embedding and tensor products.

Definition 2.3. Let \mathcal{C} be a Grothendieck category.

- (1) A *localizing subcategory* of \mathcal{C} is a full subcategory that is closed under subobjects, quotient objects, extensions, and direct sums.
- (2) A *closed subcategory* of \mathcal{C} is a full subcategory that is closed under subobjects, quotient objects, direct sums, and direct products.
- (3) A *bilocalizing subcategory* of \mathcal{C} is a full subcategory that is both localizing and closed.

	sub	quotient	extension	direct sum	direct product
localizing	✓	✓	✓	✓	
closed	✓	✓		✓	✓
bilocalizing	✓	✓	✓	✓	✓

The following result is well-known:

Theorem 2.4 ([Gab62, Lemma V.2.1] and [Ros95, Proposition III.6.4.1]). *Let A be a ring. Then there is a bijection*

$$\begin{array}{ccc} \{\text{ideals of } A\} & \xrightarrow{\simeq} & \{\text{closed subcategories of } \text{Mod } A\} \\ \cup & & \cup \\ I & \mapsto & \text{Mod}(A/I), \end{array} \quad (2.1)$$

where $\text{Mod}(A/I)$ is identified with the closed subcategory consisting of all right A -modules M with $MI = 0$. For ideals I and J of A , the product IJ is sent to $\text{Mod}(A/J) * \text{Mod}(A/I)$, which is, by definition, the full subcategory consisting of all $M \in \text{Mod } A$ that admit a short exact sequence

$$0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$$

with $L \in \text{Mod}(A/J)$ and $N \in \text{Mod}(A/I)$.

As a consequence, the bijection (2.1) induces a bijection

$$\{\text{idempotent ideals of } A\} \xrightarrow{\simeq} \{\text{bilocalizing subcategories of } \text{Mod } A\}.$$

Here an ideal I is called idempotent if $I^2 = I$.

Theorem 2.5 (Gabriel-Popescu [PG64]; see also [Pop73, Theorem 4.14.2]). *Let \mathcal{C} be a Grothendieck category. Let U be a generator in \mathcal{C} and set $A := \text{End}_{\mathcal{C}}(U)$. Then the functor*

$$S := \text{Hom}_{\mathcal{C}}(U, -): \mathcal{C} \rightarrow \text{Mod } A$$

has a left adjoint $Q: \text{Mod } A \rightarrow \mathcal{C}$. Let

$$\mathcal{X} := \text{Ker } Q = \{M \in \text{Mod } A \mid Q(M) = 0\}.$$

Then Q induces an equivalence

$$\frac{\text{Mod } A}{\mathcal{X}} \xrightarrow{\simeq} \mathcal{C}.$$

Moreover, the following hold:

- (1) The functor Q is exact and dense. $Q(A) = U$.
- (2) The functor $S = \text{Hom}_{\mathcal{C}}(U, -)$ is left exact and fully faithful. $S(U) = \text{Hom}_{\mathcal{C}}(U, U) = A$.
- (3) The essential image of S consists of all $M \in \text{Mod } A$ satisfying

$$\text{Hom}_A(X, M) = 0 \quad \text{and} \quad \text{Ext}_A^1(X, M) = 0 \quad \text{for all } X \in \mathcal{X}.$$

- (4) The counit morphism $Q \circ S \rightarrow \text{id}_{\mathcal{C}}$ is an isomorphism.

Convention 2.6. For a Grothendieck category \mathcal{C} , we will refer to $(\text{Mod } A, Q, S)$ in Theorem 2.4 as a Gabriel-Popescu embedding for \mathcal{C} . Note that the unit morphism $\eta_A: A \rightarrow SQ(A)$ is an isomorphism because $A = S(U)$ and $Q \circ S \cong \text{id}_{\text{Mod } A}$.

Theorem 2.7 ([LRGS18]). *Let \mathcal{C}_1 and \mathcal{C}_2 be R -linear Grothendieck categories. For each $i = 1, 2$, take an R -algebra A_i and a localizing subcategory \mathcal{X}_i of $\text{Mod } A_i$ such that there is an R -linear equivalence $\mathcal{C}_i \cong (\text{Mod } A_i)/\mathcal{X}_i$. Then there is an R -linear equivalence*

$$\mathcal{C}_1 \boxtimes_R \mathcal{C}_2 \cong \frac{\text{Mod}(A_1 \otimes_R A_2)}{\mathcal{X}}$$

where \mathcal{X} is the smallest localizing subcategory of $\text{Mod}(A_1 \otimes_R A_2)$ containing all $A_1 \otimes_R A_2$ -modules B such that $B_{A_1} \in \mathcal{X}_1$ or $B_{A_2} \in \mathcal{X}_2$.

Proof. The localizing subcategory \mathcal{X}_i defines a linear topology on the singleton R -linear category $\{\Lambda_i\}$ as in [LRGS18, section 2], and the description of the tensor product follows from [LRGS18, Theorem 5.4]. \square

2.3. Roos categories.

Definition 2.8. Let \mathcal{C} be a cocomplete abelian category.

- (1) \mathcal{C} is said to satisfy *Ab4* if direct sums are exact, that is, for every small family of short exact sequences, its termwise direct sum is again a short exact sequence.
- (2) \mathcal{C} is said to satisfy *Ab5* if filtered colimits are exact, that is, for every small filtered system of short exact sequences, its termwise filtered colimit is again a short exact sequence.
- (3) \mathcal{C} is said to satisfy *Ab6* if the following condition is fulfilled: Let M be an object in \mathcal{C} and let $\{\{L_i^{(j)}\}_{i \in I_j}\}_{j \in J}$ be a small family of (upward) filtered sets of subobjects of M . Then

$$\bigcap_{j \in J} \left(\bigcup_{i \in I_j} L_i^{(j)} \right) = \bigcup_{\{i(j)\}_j \in \prod_{j \in J} I_j} \left(\bigcap_{j \in J} L_{i(j)}^{(j)} \right),$$

where the symbol \bigcup means the filtered union of subobjects (that is, the sum of subobjects that are filtered).

An abelian category with direct products is said to satisfy *Ab4** (resp. *Ab5**, *Ab6**) if the opposite category satisfies *Ab4* (resp. *Ab5*, *Ab6*).

Recall that a Grothendieck category is a cocomplete abelian category that satisfies *Ab5* and has a generator. It is known that every Grothendieck category is also complete.

Theorem 2.9 (Roos [Roo65]; see also [Pop73, section 4.21]). *Let $(\text{Mod } A, Q, S)$ be a Gabriel-Popescu embedding for a Grothendieck category \mathcal{C} . Then the following are equivalent:*

- (1) \mathcal{C} satisfies *Ab6* and *Ab4**.
- (2) $\mathcal{X} = \text{Ker } Q$ is a bilocalizing subcategory of $\text{Mod } A$.
- (3) $Q: \text{Mod } A \rightarrow \mathcal{C}$ has a left adjoint.
- (4) $Q: \text{Mod } A \rightarrow \mathcal{C}$ preserves direct products.

If these equivalent conditions hold, then the left adjoint H of Q is fully faithful and thus satisfies $H \circ Q \cong \text{id}_{\text{Mod } A}$.

Definition 2.10. We call a Grothendieck category \mathcal{C} a *Roos category* if it satisfies *Ab6* and *Ab4**.

	cocomplete	Ab4	Ab5	Ab6	complete	Ab4*	Ab5*	Ab6*
Grothendieck category	✓	✓	✓	✓	✓			
Roos category	✓	✓	✓	✓	✓	✓		

Remark 2.11. The category of modules over an arbitrary ring is a Roos category.

There exists a nonzero Roos category that admits no projective objects other than zero ([Roo66, Example 3]; see also [Roo06, section 4]).

Remark 2.12. Let \mathcal{C} be a Roos category, and let $(\text{Mod } A, Q, S)$ be a Gabriel-Popescu embedding for \mathcal{C} . By Theorem 2.9, \mathcal{C} is equivalent to $(\text{Mod } A)/\mathcal{X}$, where $\mathcal{X} := \text{Ker } Q$. Moreover, by Theorem 2.4, $\mathcal{X} = \text{Mod}(A/I)$ for some idempotent ideal $I \subset A$. Thus

$$\mathcal{C} \cong \frac{\text{Mod } A}{\text{Mod}(A/I)}.$$

Its dual is obtained by taking the opposite ring ([Qui96, Proposition 8.6]):

$$\mathcal{C}^* \cong \frac{\text{Mod } A^{\text{op}}}{\text{Mod}(A^{\text{op}}/I)}. \quad (2.2)$$

Conversely, for a ring A and an idempotent ideal $I \subset A$, the subcategory $\text{Mod}(A/I) \subset \text{Mod } A$ is bilocalizing, so the quotient category $(\text{Mod } A)/(\text{Mod}(A/I))$ is a Roos category (see [Pop73, Theorem 4.21.6]).

Therefore, if \mathcal{C} is a Roos category, its dual \mathcal{C}^* is also a Roos category (because it is of the form (2.2)).

3. MAIN THEOREM

Let R be a commutative ring. The first observation in this section is that every Roos category is dualizable, and its dual is again a Roos category (hence a Grothendieck category). This follows immediately from the following result in [BCJF15]:

Theorem 3.1 ([BCJF15, Corollary 3.4]). *Let \mathcal{M} and \mathcal{C} be locally presentable R -linear categories and suppose that \mathcal{M} is dualizable. If there are cocontinuous functors $Q: \mathcal{M} \rightarrow \mathcal{C}$ and $H: \mathcal{C} \rightarrow \mathcal{M}$ such that $Q \circ H \cong \text{id}_{\mathcal{C}}$, then \mathcal{C} is also dualizable.*

Corollary 3.2. *Every R -linear Roos category \mathcal{C} is dualizable. The dual of \mathcal{C} is also a Roos category.*

Proof. Let $(\text{Mod } A, Q, S)$ be a Gabriel-Popescu embedding for \mathcal{C} . Then Q has a left adjoint H , and $Q \circ H \cong \text{id}_{\mathcal{C}}$. Therefore, \mathcal{C} is dualizable by Theorem 3.1. The dual \mathcal{C}^* is also a Roos category by Remark 2.12. \square

Next we will show that a dualizable Grothendieck category whose dual is again a Grothendieck category is a Roos category.

Lemma 3.3. *Let \mathcal{C} be a dualizable R -linear Grothendieck category whose dual is again a Grothendieck category. Let $(\text{Mod } A, Q, S)$ be a Gabriel-Popescu embedding for \mathcal{C} . Then there is an R -linear cocontinuous functor $H: \mathcal{C} \rightarrow \text{Mod } A$ such that $Q \circ H \cong \text{id}_{\mathcal{C}}$.*

Proof. We have the commutative diagram

$$\begin{array}{ccc} (\text{Mod } A) \boxtimes_R \mathcal{C}^* & \longrightarrow & \text{Cocont}_R(\mathcal{C}, \text{Mod } A) \\ Q \boxtimes \text{id}_{\mathcal{C}^*} \downarrow & & \downarrow Q_* \\ \mathcal{C} \boxtimes_R \mathcal{C}^* & \longrightarrow & \text{Cocont}_R(\mathcal{C}, \mathcal{C}) \end{array}$$

where the horizontal arrows are canonical functors and Q_* is the induced functor defined by $Q_*(F) := Q \circ F$. If Q_* is dense, then there is $H \in \text{Cocont}_R(\mathcal{C}, \text{Mod } A)$ such that $Q \circ H \cong \text{id}_{\mathcal{C}}$, which is the desired claim. Since $\text{Mod } A$ and \mathcal{C} are dualizable, both horizontal arrows are equivalences by [BCJF15, Lemma 3.1]. Hence, it suffices to prove that $Q \boxtimes \text{id}_{\mathcal{C}^*}$ is dense.

Let $(\text{Mod } \Gamma, Q', S')$ be a Gabriel-Popescu embedding for \mathcal{C}^* , and let $\mathcal{X} := \text{Ker } Q$ and $\mathcal{X}' := \text{Ker } Q'$. By Theorem 2.7,

$$\mathcal{C} \boxtimes_R \mathcal{C}^* \cong \frac{\text{Mod } A}{\mathcal{X}} \boxtimes_R \frac{\text{Mod } \Gamma}{\mathcal{X}'} \cong \frac{\text{Mod}(A \otimes_R \Gamma)}{\mathcal{Y}},$$

where \mathcal{Y} is the smallest localizing subcategory of $\text{Mod}(A \otimes_R \Gamma)$ containing all $B \in \text{Mod}(A \otimes_R \Gamma)$ such that $B_A \in \mathcal{X}$ or $B_\Gamma \in \mathcal{X}'$. Similarly, we have

$$(\text{Mod } A) \boxtimes_R \mathcal{C}^* \cong (\text{Mod } A) \boxtimes_R \frac{\text{Mod } \Gamma}{\mathcal{X}'} \cong \frac{\text{Mod}(A \otimes_k \Gamma)}{\mathcal{Z}},$$

where \mathcal{Z} is the smallest localizing subcategory of $\text{Mod}(A \otimes_R \Gamma)$ containing all $B \in \text{Mod}(A \otimes_R \Gamma)$ such that $B_\Gamma \in \mathcal{X}'$. Obviously, $\mathcal{Y} \subset \mathcal{Z}$, and we obtain the commutative diagram

$$\begin{array}{ccccc} (\text{Mod } A) \boxtimes_R \mathcal{C}^* & \xrightarrow{\sim} & (\text{Mod } A) \boxtimes_R \frac{\text{Mod } \Gamma}{\mathcal{X}'} & \xrightarrow{\sim} & \frac{\text{Mod}(A \otimes_R \Gamma)}{\mathcal{Y}} \\ Q \boxtimes \text{id}_{\mathcal{C}^*} \downarrow & & \downarrow & & \downarrow \\ \mathcal{C} \boxtimes_R \mathcal{C}^* & \xrightarrow{\sim} & \frac{\text{Mod } A}{\mathcal{X}} \boxtimes_R \frac{\text{Mod } \Gamma}{\mathcal{X}'} & \xrightarrow{\sim} & \frac{\text{Mod}(A \otimes_R \Gamma)}{\mathcal{Z}}. \end{array}$$

The canonical functor represented by the right-most vertical arrow is dense, and hence $Q \boxtimes \text{id}_{\mathcal{C}^*}$ is also dense. \square

Lemma 3.4. *Let \mathcal{C} be an R -linear Grothendieck category, and let $(\text{Mod } \Lambda, Q, S)$ be a Gabriel-Popescu embedding for \mathcal{C} . If there is an R -linear cocontinuous functor $H: \mathcal{C} \rightarrow \text{Mod } \Lambda$ such that $Q \circ H \cong \text{id}_{\mathcal{C}}$, then \mathcal{C} is a Roos category.*

Proof. Let $\alpha: Q \circ H \xrightarrow{\cong} \text{id}_{\mathcal{C}}$ be an isomorphism. Since $H \circ Q: \text{Mod } \Lambda \rightarrow \text{Mod } \Lambda$ is cocontinuous, the Eilenberg-Watts theorem implies that there is a Λ - Λ -bimodule B over R such that $H \circ Q \cong - \otimes_{\Lambda} B$. Let $\beta: - \otimes_{\Lambda} B \xrightarrow{\cong} H \circ Q$ be an isomorphism, and let $\psi: \Lambda \otimes_{\Lambda} B \xrightarrow{\cong} B$ be the canonical bimodule isomorphism. Define the right Λ -module homomorphism $\varphi: B \rightarrow \Lambda$ as the composition

$$B \xrightarrow{\eta_B} SQ(B) \xleftarrow{\widetilde{SQ(\psi)}} SQ(\Lambda \otimes_{\Lambda} B) \xrightarrow{\widetilde{SQ(\beta_{\Lambda})}} SQHQ(\Lambda) \xrightarrow{\widetilde{S(\alpha_{Q(\Lambda)})}} SQ(\Lambda) \xleftarrow{\widetilde{\eta_{\Lambda}}} \Lambda$$

where $\eta: \text{id} \rightarrow SQ$ is the unit morphism, and η_{Λ} is an isomorphism because this is a Gabriel-Popescu embedding. To observe that φ is a bimodule homomorphism, fix an element $\lambda \in \Lambda$ and let $m_{\Lambda}: \Lambda \rightarrow \Lambda$ and $m_B: B \rightarrow B$ be the multiplication of λ from the left. Then we have the commutative diagram

$$\begin{array}{ccccccc} B & \xrightarrow{\eta_B} & SQ(B) & \xleftarrow{\widetilde{SQ(\psi)}} & SQ(\Lambda \otimes_{\Lambda} B) & \xrightarrow{\widetilde{SQ(\beta_{\Lambda})}} & SQHQ(\Lambda) & \xrightarrow{\widetilde{S(\alpha_{Q(\Lambda)})}} & SQ(\Lambda) & \xleftarrow{\widetilde{\eta_{\Lambda}}} & \Lambda \\ \downarrow m_B & & \downarrow SQ(m_B) & & \downarrow SQ(m_{\Lambda} \otimes B) & & \downarrow SQHQ(m_{\Lambda}) & & \downarrow SQ(m_{\Lambda}) & & \downarrow m_{\Lambda} \\ B & \xrightarrow{\eta_B} & SQ(B) & \xleftarrow{\widetilde{SQ(\psi)}} & SQ(\Lambda \otimes_{\Lambda} B) & \xrightarrow{\widetilde{SQ(\beta_{\Lambda})}} & SQHQ(\Lambda) & \xrightarrow{\widetilde{S(\alpha_{Q(\Lambda)})}} & SQ(\Lambda) & \xleftarrow{\widetilde{\eta_{\Lambda}}} & \Lambda \end{array}$$

Hence, φ is a bimodule homomorphism. So $I := \text{Im } \varphi$ is a (two-sided) ideal of Λ .

We show that the subcategory $\mathcal{X} := \text{Ker } Q$ is equal to $\text{Mod}(\Lambda/I)$. Since $Q(\eta_B): Q(B) \rightarrow QSQ(B)$ is an isomorphism, $Q(\varphi): Q(B) \rightarrow Q(\Lambda)$ is also an isomorphism. By the exactness of Q , the canonical morphism $Q(I) \xrightarrow{\cong} Q(\Lambda)$ is an isomorphism, and $Q(\Lambda/I) = 0$. Hence $\Lambda/I \in \mathcal{X}$, and thus $\text{Mod}(\Lambda/I) \subset \mathcal{X}$.

Let $M \in \mathcal{X}$. Then $M \otimes_{\Lambda} B \cong HQ(M) = 0$. The canonical epimorphism $B \rightarrow \text{Im } \varphi = I$ induces an epimorphism $M \otimes_{\Lambda} B \rightarrow M \otimes_{\Lambda} I$, so $M \otimes_{\Lambda} I = 0$. This implies that $MI = 0$ and $M \in \text{Mod}(\Lambda/I)$. Therefore $\mathcal{X} \subset \text{Mod}(\Lambda/I)$.

Consequently, the localizing subcategory \mathcal{X} is equal to $\text{Mod}(\Lambda/I)$, which is also closed under direct products in $\text{Mod } \Lambda$. Therefore \mathcal{X} is a bilocalizing subcategory of $\text{Mod } \Lambda$, and $\mathcal{C} \cong (\text{Mod } \Lambda)/\mathcal{X}$ is a Roos category. \square

Theorem 3.5. *For an R -linear Grothendieck category \mathcal{C} , the following are equivalent:*

- (1) \mathcal{C} is a Roos category (Ab6 and Ab4*).
- (2) \mathcal{C} is dualizable and \mathcal{C}^* is a Grothendieck category.
- (3) For every/some Gabriel-Popescu embedding $(\text{Mod } \Lambda, Q, S)$ for \mathcal{C} , the functor Q has a left adjoint.
- (4) For every/some Gabriel-Popescu embedding $(\text{Mod } \Lambda, Q, S)$ for \mathcal{C} , there is a cocontinuous functor $H: \mathcal{C} \rightarrow \text{Mod } \Lambda$ such that $Q \circ H \cong \text{id}_{\mathcal{C}}$.

Proof. (1) \Rightarrow (2) follows from Corollary 3.2. (2) \Rightarrow (1) follows from Lemmas 3.3 and 3.4.

(1) \Leftrightarrow (3) \Rightarrow (4) is part of Theorem 2.9.

(4) \Rightarrow (1) is Lemma 3.4. \square

Remark 3.6. Note that (4) \Rightarrow (3) in Theorem 3.5 does not hold for a cocontinuous functor Q (which is not necessarily a Gabriel-Popescu quotient functor) in general. To see this, let k be a field, and consider the functors

$$Q := - \otimes_{k[x]} (k[x]/(x)): \text{Mod } k[x] \rightarrow \text{Mod } k$$

and

$$H := - \otimes_k k[x]: \text{Mod } k \rightarrow \text{Mod } k[x].$$

Then Q and H are cocontinuous functors, and

$$Q \circ H = - \otimes_k k[x] \otimes_{k[x]} (k[x]/(x)) = - \otimes_k (k[x]/(x)) \cong \text{id}_{\text{Mod } k}.$$

However, Q does not have a left adjoint because Q is not left exact (that is, $k[x]/(x)$ is not a flat $k[x]$ -module).

Stefanich [Ste23, Corollary 3.1.19] proved that every dualizable R -linear *cocomplete* (not necessarily locally presentable) category \mathcal{C} is a Grothendieck category satisfying Ab4^* (using $\mathcal{A} = \text{Mod } R$ as the symmetric monoidal Grothendieck category in the statement). Hence its dual \mathcal{C}^* is also a Grothendieck category because it is dualizable. Therefore, by Theorem 3.5, we obtain a complete characterization of dualizable linear cocomplete categories, confirming Conjecture 1.2:

Corollary 3.7. *Let \mathcal{C} be an R -linear cocomplete category. Then the following are equivalent:*

- (1) \mathcal{C} is dualizable.
- (2) \mathcal{C} is a Roos category.

REFERENCES

- [BCJF15] Martin Brandenburg, Alexandru Chirvasitu, and Theo Johnson-Freyd, *Reflexivity and dualizability in categorified linear algebra*, Theory Appl. Categ. **30** (2015), 808–835. MR 3361309
- [Chi22] Alexandru Chirvasitu, *Higher dualizability and singly-generated Grothendieck categories*, Appl. Categ. Structures **30** (2022), no. 1, 1–12. MR 4366933
- [Gab62] Pierre Gabriel, *Des catégories abéliennes*, Bull. Soc. Math. France **90** (1962), 323–448. MR 0232821
- [Kan19] Ryo Kanda, *Non-exactness of direct products of quasi-coherent sheaves*, Doc. Math. **24** (2019), 2037–2056. MR 4033823
- [LRGS18] Wendy Lowen, Julia Ramos González, and Boris Shoikhet, *On the tensor product of linear sites and Grothendieck categories*, Int. Math. Res. Not. IMRN (2018), no. 21, 6698–6736. MR 3873542
- [Mar97] Leandro Marín, *Categories of modules for idempotent rings and morita equivalences*, Master’s thesis, University of Glasgow, 1997.
- [PG64] Nicolae Popescu and Pierre Gabriel, *Caractérisation des catégories abéliennes avec générateurs et limites inductives exactes*, C. R. Acad. Sci. Paris **258** (1964), 4188–4190. MR 0166241
- [Pop73] N. Popescu, *Abelian categories with applications to rings and modules*, Academic Press, London, 1973, London Mathematical Society Monographs, No. 3. MR 0340375
- [Qui96] Daniel Quillen, *Module theory over nonunital rings*, Preprint (1996).
- [Roo65] Jan-Erik Roos, *Caractérisation des catégories qui sont quotients de catégories de modules par des sous-catégories bilocalisantes*, C. R. Acad. Sci. Paris **261** (1965), 4954–4957. MR 0190207
- [Roo66] ———, *Sur les foncteurs dérivés des produits infinis dans les catégories de Grothendieck. Exemples et contre-exemples*, C. R. Acad. Sci. Paris Sér. A-B **263** (1966), A895–A898. MR 0215895
- [Roo67] ———, *Sur la condition AB 6 et ses variantes dans les catégories abéliennes*, C. R. Acad. Sci. Paris Sér. A-B **264** (1967), A991–A994. MR 0217145
- [Roo06] ———, *Derived functors of inverse limits revisited*, J. London Math. Soc. (2) **73** (2006), no. 1, 65–83. MR 2197371
- [Ros95] Alexander L. Rosenberg, *Noncommutative algebraic geometry and representations of quantized algebras*, Mathematics and its Applications, vol. 330, Kluwer Academic Publishers Group, Dordrecht, 1995. MR 1347919
- [Ste23] Germán Stefanich, *Classification of fully dualizable linear categories*, arXiv:2307.16337.
- [Wis91] Robert Wisbauer, *Foundations of module and ring theory*, german ed., Algebra, Logic and Applications, vol. 3, Gordon and Breach Science Publishers, Philadelphia, PA, 1991, A handbook for study and research. MR 1144522

(Ryo Kanda) DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA METROPOLITAN UNIVERSITY, 3-3-138, SUGIMOTO, SUMIYOSHI, OSAKA, 558-8585, JAPAN

Email address: ryo.kanda.math@gmail.com