# Reality Only Happens Once: Single-Path Generalization Bounds for Transformers

Yannick Limmer Limmery@maths.ox.ac.uk

Department of Mathematics, University of Oxford Oxford-Man Institute for Quantitative Finance

Anastasis Kratsios Kratsioa@mcmaster.ca

Department of Mathematics, McMaster University Vector Institute

Xuwei Yang Yangx212@mcmaster.ca

Department of Mathematics, McMaster University

Raeid Sagur RAEIDSAQUR@CS.TORONTO.EDU

 $Department\ of\ Computer\ Science,\ University\ of\ Toronto\ Vector\ Institute$ 

Blanka Horvath HORVATH@MATHS.OX.AC.UK

Department of Mathematics, University of Oxford Oxford-Man Institute for Quantitative Finance

#### Abstract

One of the inherent challenges in deploying transformers on time series is that reality only happens once; namely, one typically only has access to a single trajectory of the data-generating process comprised of non-i.i.d. observations. We derive non-asymptotic statistical guarantees in this setting through bounds on the generalization of a transformer network at a future-time t, given that it has been trained using  $N \leq t$  observations from a single perturbed trajectory of a Markov process. Under the assumption that the Markov process satisfies a log-Sobolev inequality, we obtain a generalization bound which effectively converges at the rate of  $\mathcal{O}(1/\sqrt{N})$ . Our bound depends explicitly on the activation function (Swish, GeLU, or tanh are considered), the number of self-attention heads, depth, width, and norm-bounds defining the transformer architecture. Our bound consists of three components: (I) The first quantifies the gap between the stationary distribution of the data-generating Markov process and its distribution at time t, this term converges exponentially to 0. (II) The next term encodes the complexity of the transformer model and, given enough time, eventually converges to 0 at the rate  $\mathcal{O}(\log(N)^r/\sqrt{N})$  for any r > 0. (III) The third term guarantees that the bound holds with probability at least  $1-\delta$ , and converges at a rate of  $\mathcal{O}(\sqrt{\log(1/\delta)}/\sqrt{N})$ .

# 1. Introduction

Transformers [84] have become the main architectural building block in deep learning-based state-of-the-art foundation models [18, 98, 88]. Transformers are primarily deployed on sequential learning tasks which have complex temporal relationships, and thus, transformers are trained on non-i.i.d. data. The i.i.d. assumption is typically made (e.g. [69, 9, 97]) to derive theoretical statistical guarantees, but in practice it is rarely satisfied; e.g. in natural language processing (NLP) [99], physics [72], medical research [12], reinforcement learning [79], optimal control [80], and in finance [36]. This creates a mismatch between available statistical guarantees in deep learning (which often rely on the i.i.d. assumption) and how transformers are used in practice.

Thus, this paper fills this gap by guaranteeing that transformers trained on a single time-series trajectory can generalize at future moments in time. We, therefore, consider the learning problem where the user is supplied with N paired samples  $(X_1,Y_1),\ldots,(X_N,Y_N)$ , where each input  $Y_n=f^\star(X_n)$  for a smooth (unknown) target function  $f^\star:\mathbb{R}^{d\times M}\to\mathbb{R}^D$  is to be learned, depending on a history length M, and where the inputs are generated by a time-homogeneous Markov process  $X \stackrel{\text{def}}{=} (X_n)_{n=1}^\infty$ . Note that the assumption  $Y_n=f^\star(X_n)$  results in only a mild loss of generality since if X is a discretized solution to a stochastic differential equation then  $Y_n\approx \text{signal}+\text{additive noise}$  due to stochastic calculus considerations (see Appendix E).

The performance of any transformer model  $\mathcal{T}:\mathbb{R}^{d\times M}\to\mathbb{R}^D$  is quantified via a smooth loss

The performance of any transformer model  $\mathcal{T}: \mathbb{R}^{d \times M} \to \mathbb{R}^D$  is quantified via a smooth loss function  $\ell: \mathbb{R}^D \times \mathbb{R}^D \to \mathbb{R}$ . When M = 1, the generalization of such a  $\mathcal{T}$  is measured by the gap between its *empirical risk*  $\mathcal{R}^{(N)}$ , computed from the single-path training data, and its *(true) t-future risk*  $\mathcal{R}_t$  at a (possibly infinite) future time  $N \leq t \leq \infty$   $(t \in \mathbb{N}_+)$  defined by

$$\mathcal{R}_t(f) \stackrel{\text{def.}}{=} \mathbb{E} \big[ \ell(\mathcal{T}(X_t), f^{\star}(X_t)) \big]$$

where  $\mathcal{R}_t$  (resp.  $\mathcal{R}_{\infty}$ ) is computed with respect to the distribution of  $X_t$  (resp. stationary distribution of X.). The time-t excess-risk  $\mathcal{R}_t$ , which is generally unobservable, is estimated by a single-path estimator known as the empirical risk computed using all the noisy samples observed thus far

$$\mathcal{R}^{(N)}(f) \stackrel{\text{\tiny def.}}{=} \frac{1}{N} \sum_{n=1}^{N} \ell(\mathcal{T}(X_n), f^{\star}(X_n)).$$

Our objective is to obtain a statistical learning guarantee bounding the gap between the empirical risk and the t-future risk of transformer models trained on a single path.

Our main result is a bound on the future-generalization, at any given time  $t \ge N$ , of a transformer trained from N samples collected from an unknown transformation  $(f^*)$  of any suitable unknown Markov process (X). Let  $\mathcal{TC}_{L,H,C,W}^{D,d,0}$  be the class of transformers with L transformer blocks, H attention heads, weights constraint C, and maximum width W; then for respective input and output dimensions d and D, and history length 1 (including the current state), the first takeaway of our main result (Theorem 1) is that with probability of at least  $1-\delta$ 

$$\sup_{\mathcal{T} \in \mathcal{T}_{L,W,H}^{D,d,0}} \left| \mathcal{R}_t(\mathcal{T}) - \mathcal{R}^{(N)}(\mathcal{T}) \right| \in \mathcal{O}\left(\frac{\log(1/\delta) + \log(N)^{1/s}}{\sqrt{N}}\right)$$
(FutureGen)

where s > 0 can be made arbitrarily large and  $\mathcal{O}$  hides a dimensional constant depending on s.

#### 1.1 Contributions

Our first main contribution is the first generalization bound (FutureGen) for transformers in the case of non i.i.d. training data at a future point in time. This provides the first set of statistical guarantees for transformers compatible with many time-series tasks. More broadly speaking, our generalization is the first result which estimates the future-generalization (in (FutureGen)) of a learning model given that its performance has only been measured using present/past historical training data.

Our secondary contribution is a full analysis of the constants under the big  $\mathcal{O}$  in our future-generalization bound via a complete estimation of the higher order sensitivities/derivatives of the transformer network. The computation of these constants in Theorem 5 (and their order estimates in Theorem 6) are explicitly expressed in terms of the number of attention heads, depth and width of the transformed model, and weight and bias restriction, as well as on the activation functions used. Our analysis applies to transformers using any smooth activation function, we perform an in-depth analysis for the Swish [74], GeLU [47], and the tanh activation functions. This allows us to analyze the impact of each of these choices on the future generalization of transformers.

#### 1.2 Related Work

The mathematical foundations of transformer networks have recently come into focus in the deep learning theory community. Most of the available statistical guarantees for transformers either concern: in-context learning for linear transformers [97, 39], transformers [86, 1] trained with gradient descent, or instance-dependant bounds [81] for general transformers. These results, however, do not apply in time series analysis contexts where each training sample is not independent of the others but is rather generated by some recursive stochastic process, e.g. a Markov process.

Analytic counterparts to the statistical guarantees for transformers have also emerged. These include universal approximation theorems for transformer networks [94, 95, 34] and contained universal approximation results for networks leveraging generalized attention mechanisms [59], and the identification of function classes which can be efficiently approximated by transformers special classes [63, 37]. From the computability standpoint, transformers are Turing complete [16].

Generalization bounds for multilayer perceptions (MLPs) have been actively studied for years. For classification problems, these generalization bounds often rely on bounding the VC-dimension of classes of MLPs, depending on their depth, width, norm bounds on their parameters and activation functions [6, 10], or similar quantities. In regression problems, one instead controls the Rademacher complexity of similar classes of MLPs [9, 70, 93], due to the results such as [55, 7], or turns to instance-dependant bounds which control the path-norm of the MLP [69, 40, 38] and local variants of these quantities; e.g. [8] or [49]. Our generalization bounds also partially borrow ideas from both of these directions, but instead, we use high-order sensitivities (partial derivatives) of our transformer networks to obtain tighter bounds for large enough N. This does not yield a faster rate, since the  $\mathcal{O}(1/\sqrt{N})$  rate is generally optimal, by the central limit theorem, but it allows us to better control the constants in the generalization bound and thus yields more precise bounds. Thus, a key part of our technical contributions is the computation of these higher-order derivatives ( $C^s$ -norms) both of the transformer and the MLP models using smooth activation functions.

These statistical learning results assume that the data samples are i.i.d. However, time-series data is rarely i.i.d, they are often generated by Markov processor at least embeddable into a Markovian setting [27, 28]. Though there are generalization bounds for non-i.i.d. relying on martingale arguments e.g. [56] and concentration of measure phenomena for martingale sums e.g. [15, 19] those results primarily focus on Lipschitz functions; thus, they do not consider higher-order derivatives. Our results add to this literature since we rely on the concentration of measure phenomena for Markov processes with respect to smooth counterparts of the 1-Wasserstein distance (a tool used in many martingale arguments, e.g. [57], for Lipschitz classes).

Our main result operates under the assumption that the stationary distribution of the Markov chain and its kernels all satisfy a log-Sobolev inequality (LSI). Since their introduction, LSIs have been heavily studied [45, 62, 100, 51, 22] and have found numerous applications in differential privacy [66, 92], optimization [21], random matrix theory [89, 90], optimal transport [31], since they typically imply [42, 43] and effectively characterizes [41] dimension-free rate for concentration of measure.

We require that the data-generating Markov process has an exponentially contracting Markov kernel [54]. For Markov chains, i.e. finite-state space Markov processes, this means that that the generator (Q-matrix) of the Markov chain has a spectral gap. These spectral gaps are actively studied in the Markov chain literature [68, 58, 3, 73, 53] since these have a finite mixing time, meaning that they distribution of such Markov chains approaches their stationary limit after a large finite time has elapsed; i.e. they have well-behaved (approximate) mixing times [67, 50, 91, 96].

Our generalization bounds relied on concentration of measure arguments for the "smooth" integral probability metrics (IPMs) studied in [54, 75], by refining the arguments of [49, 14, 60] to the non-i.i.d. and smooth setting. In the i.i.d. case, our computation of the maximum  $C^s$ -norm  $(R \ge 0)$  of the class  $\mathcal{TC}_{L,W,H}^{D,d,0}$  (Theorem 5) can be used to relate the rate at which measure concentrates (from i.i.d. data) for these IPMs can be bounded-above via classical quantities such as the Rademacher complexity of the class of  $C^s$ -functions on  $\mathbb{R}^d$  with  $C^s$  norm at-most R; see e.g. [78, Theorem 3.3].

#### 1.3 Further Applications of our Secondary Results

The upper bounds, which we compute for the  $C^s$  norms of the transformers models, can be used in conjunction with classical VC-dimension [83], Rademacher complexity [7], or optimal transport [49] type arguments to obtain generalization bounds in the simpler setting of i.i.d. data where there is no notion of (future) time, not considered here. This can be done using classical tools, e.g. entropy estimates in [83, Theorem 2.7.4] on compact domains due to the Sobolev embedding theorem, applied to the larger class of  $C^s$  on  $\mathbb{R}^{d \times M}$  whose  $C^s$ -norms are all most equal to the one we have computed for  $\mathcal{TC}_{L,H,C,W}^{D,d,M}$  in Theorems 5 and 6.

# 2. Background and Preliminaries

This section overviews the necessary background for a self-contained formulation of our main results. This includes the definition of transformers and probabilistic tools such as log-Sobolev inequalities.

**Log-Sobolev Inequalities and Moment Conditions** Fix dimensions  $d, D \in \mathbb{N}_+$  and let P be a Markov kernel on a non-empty Borel  $\mathcal{X} \subseteq \mathbb{R}^d$  with initial distribution  $X_0 \sim \mu_0 \in \mathcal{P}(\mathbb{R}^d)$ , and with associated Markov process  $X \stackrel{\text{def.}}{=} (X_t)_{t \in \mathbb{N}_0}$  given by  $X_t \sim \mu_t \stackrel{\text{def.}}{=} P^t \mu_0$ . The *entropy functional*  $\mathcal{H}_{\mu}$  associated to any Borel probability measure  $\mu$  on  $\mathbb{R}^d$  acts on smooth functions  $g : \mathbb{R}^d \to \mathbb{R}$  by

$$\mathbb{H}_{\mu}(g) \stackrel{\text{\tiny def.}}{=} \mathbb{E}_{X \sim \mu} \Big[ g(X) \log \Big( \frac{g(X)}{\mathbb{E}_{Z \sim \mu}[g(Z)]} \Big) \Big].$$

The entropy functional can be used to express the log-Sobolev inequalities.

**Definition 1 (Log-Sobolev Inequality)** A probability measure  $\mu$  on  $\mathbb{R}^d$  is said to satisfy a log-Sobolev inequality with constant C > 0 (LSI<sub>C</sub>) if for every smooth function  $g : \mathbb{R}^d \to \mathbb{R}$ 

$$\mathbb{H}_{\mu}(g^2) \leqslant C \ \mathbb{E}_{X \sim \mu}[\|\nabla g(X)\|^2]$$

We require that the Markov process is time-homogeneous to admit a satisfactory measure. Further, we require that its Markov kernel and its stationary measure all satisfy  $LSI_C$ .

Assumption 1 (Satisfactions of the Log-Sobolev Inequality) There exists a C > 0 such that  $\bar{\mu}$ ,  $\mu_0$ , and  $P(x,\cdot)$  all satisfy LSI<sub>C</sub>, for each  $x \in \mathcal{X}$ .

Several examples of Markov processes satisfying LSI inequalities are given in [61] and Gaussian processes satisfy the Exponential Moments Assumption. Lastly, we require that the Markov process X is bounded. This will not be required for our concentration arguments (Proposition 2) but it is needed when controlling the higher order derivatives of the transformer model, which can explode when considered over an unbounded domain; i.e. on all of  $\mathbb{R}^{Md}$ .

**Assumption 2 (Bounded Trajectories)** There is a constant  $c_K > 0$  such that, each  $t \in \mathbb{N}_+$ ,  $||X.|| \leq c_k$  holds with probability 1.

The Transformer Model The overall structure of transformers and their involved layers are summarized in Figure 1c. We, thus, encourage the familiar reader to skip this section, which contains precise technical formulations of transformer classes with precise specifications on weight magnitude, depth, width, and a number of attention heads that can be rather involved.

For any  $F \in \mathbb{N}_+$ , we will consider a weighted (parametric) variant of the *layer normalization* function of [5], which permits a variable level of regularization. Our weighted *layer normalization* is defined by LayerNorm:  $\mathbb{R}^F \to \mathbb{R}^F$  defined for any  $u \in \mathbb{R}^F$  by

$$\mathcal{LN}(u; \gamma, \beta, w) \stackrel{\text{def.}}{=} \gamma \frac{(u - \mu_u^w)}{\sqrt{1 + (\sigma_u^w)^2}} + \beta$$




Figure 1: Multi-Head Attention: Consists of parallel application of the attention mechanism, described by the following steps. 1) Inputs are used three-fold, as keys, queries, and values, all are transformed by distinct linear transformations. 2) Keys and queries are multiplied, scaled, and transformed by a softmax application. 3) This output is combined in a matrix multiplication with the values.

**Transformer Block:** 1) Input features are mapped to contexts via multi-head attention mechanism. 2) The output of the multi-head attention mechanism and the input features (via a skip connection) are normalized by a layer-norm. 3) The normalized contextual features are transformed non-linearly by a SLP. 4) The outputs of the SLP, together with the first set of normalized context (via another skip connection), are normalized by a final layer-norm and returned by the transformer block.

**Transformers:** Iteratively feed input features through a series of transformer blocks before processing their outputs with a (fully connected affine layer).

where  $\mu_u^w \stackrel{\text{def.}}{=} \sum_{i=1}^F \frac{w}{F} u_i$  and  $(\sigma_u^w)^2 \stackrel{\text{def.}}{=} \sum_{i=1}^F \frac{w}{F} \|u_i - \mu_u\|^2$ , splus  $\stackrel{\text{def.}}{=} \ln(1 + \exp(\cdot))$ , parameters  $\beta \in \mathbb{R}^F$  and  $\gamma \in \mathbb{R}$ , and the normalization strength parameter  $w \in [0,1]$  with w=1 being the default choice. Here, we prohibit the layer norm from magnifying the size of its outputs when the layer-wise weighted variance  $\sigma_u^w$  is small.<sup>1</sup>

**Definition 2 (Multi-Head Self-Attention)** Fix  $d_{\text{in}} \in \mathbb{N}$ . For  $x \in \mathbb{R}^{(M \times d_{\text{in}})}$ ,  $Q, K \in \mathbb{R}^{d_K \times d_{\text{in}}}$ , and  $V \in \mathbb{R}^{d_V \times d_{\text{in}}}$ , where we have key-dimension  $d_K \in \mathbb{N}$  and value-dimension  $d_V \in \mathbb{N}$ ; we define

$$\operatorname{Att}(x;Q,K,V) \stackrel{\text{\tiny def.}}{=} \left( \sum_{i=0}^{M} \operatorname{softmax} \left( \left( \frac{\langle Qx_m,Kx_i \rangle}{\sqrt{d_k}} \right)_{i=0}^{M} \right)_{j} Vx_j \right)_{m=1}^{M} \in \mathbb{R}^{M \times d_V}.$$

For  $H \in \mathbb{N}$ , set  $Q \stackrel{\text{def.}}{=} (Q^{(h)})_{h=1}^H$ ,  $K \stackrel{\text{def.}}{=} (K^{(h)})_{h=1}^H \subseteq \mathbb{R}^{d_K \times d_{\text{in}}}$ ,  $V \stackrel{\text{def.}}{=} (V^{(h)})_{h=1}^H \subseteq \mathbb{R}^{d_V \times d_{\text{in}}}$ , as well as  $W \stackrel{\text{def.}}{=} (W^{(h)})_{h=1}^H \subseteq \mathbb{R}^{d_{\text{in}} \times d_V}$ . For  $x \in \mathbb{R}^{M \times d_{\text{in}}}$ , we define

$$\mathcal{MH}(x;Q,K,V,W) \stackrel{\text{\tiny def.}}{=} \left( \sum_{h=1}^{H} W^{(h)} \operatorname{Att}(x;Q^{(h)},K^{(h)},V^{(h)})_{m} \right)_{m=1}^{M} \in \mathbb{R}^{(M)\times d_{\mathrm{in}}}.$$

Each transformer block takes a set of inputs and intersperses normalization (via layer norms), contextual comparisons (via multi-head attention mechanism), and non-linear transformations (via an SLP). We also allow the transformer block to extend or contract the length of the generated sequence.

**Definition 3 (Transformer Block)** Fix a non-affine activation function  $\sigma \in C^{\infty}(\mathbb{R})$ . Fix a dimensional multi-index  $d = (d_{\text{in}}, d_K, d_V, d_{\text{ff}}, d_{\text{out}}) \in \mathbb{N}^5$ , a sequence length  $M \in \mathbb{N}_+$ , and a number of

<sup>1.</sup> Note that this formulation of the layer norm avoids division by 0 when the entries of u are identical.

self-attention heads  $H \in \mathbb{N}_+$ . A transformer block is a permutation equivariant map  $\mathcal{TB} : \mathbb{R}^{M \times d_{\mathrm{in}}} \to \mathbb{R}^{M \times d_{\mathrm{out}}}$  represented for each  $x \in \mathbb{R}^{M \times d_{\mathrm{in}}}$ 

$$\mathcal{TB}(x) \stackrel{\text{def.}}{=} \left( \mathcal{LN} \left( B^{(1)} x'_m + B^{(2)} \left( \sigma \bullet (A x'_m + a) \right); \gamma_2, \beta_2, w_2 \right) \right)_{m=1}^{M}$$

$$x' \stackrel{\text{def.}}{=} \left( \mathcal{LN} \left( x_m + \mathcal{MH}(x; Q, K, V, W)_m; \gamma_1, \beta_1, w_1 \right) \right)_{m=1}^{M}$$

$$(1)$$

for  $\gamma_1, \gamma_2 \in \mathbb{R}$ ,  $w_1, w_2 \in [0, 1]$ ,  $\beta_1 \in \mathbb{R}^{d_{\text{in}}}$ ,  $\beta_2 \in \mathbb{R}^{d_{\text{out}}}$ ,  $A \in \mathbb{R}^{d_{\text{ff}} \times d_{\text{in}}}$ ,  $a \in \mathbb{R}^{d_{\text{ff}}}$ ,  $B^{(1)} \in \mathbb{R}^{d_{\text{out}} \times d_{\text{in}}}$ ,  $B^{(2)} \in \mathbb{R}^{d_{\text{out}} \times d_{\text{ff}}}$ , and Q, K, V, W as in Definition 2.

The class of transformer blocks with representation (1) is denoted by  $\mathcal{T}\mathcal{Z}_{d,M,H,C}$ , where the constants are collected in  $C \stackrel{\text{def.}}{=} (C^{\gamma_1}, C^{\gamma_2}, C^{\beta_1}, C^{\beta_2}, C^a, C^A, C^{B^{(1)}}, C^{B^{(2)}}, C^Q, C^K, C^V, C^W)$ .

A transformer concatenates several transformer blocks before passing their outputs to an affine layer and ultimately outputting its prediction.

**Definition 4 (Transformers)** Fix depth  $L \in \mathbb{N}_+$ , memory  $M \in \mathbb{N}$ , width  $W \in \mathbb{N}_+^5$ , number of heads  $H \in \mathbb{N}_+$ , and input-output dimensions  $D, d \in \mathbb{N}_+$ . A transformer (network) is a map  $\mathcal{T} : \mathbb{R}^{M \times D} \to \mathbb{R}^d$  with representation

$$\mathcal{T}(x) = A\left(\operatorname{vec}_{1+M,d_{\text{out}}^L} \circ \mathcal{TB}_L \circ \cdots \circ \mathcal{TB}_1(x)\right) + b \tag{2}$$

where multi-indices  $d^l = (d_{\rm in}^l, d_K^l, d_V^l, d_{\rm ff}^l, d_{\rm out}^l) \leqslant W$  are such that  $d_{\rm in}^1 = D$ ,  $d_{\rm in}^{l+1} = d_{\rm out}^l$  for each  $l = 1, \ldots, L-1$ , and where  $H \stackrel{\rm def.}{=} (H^l)_{l=1}^L$  are the number of self-attention heads,  $C' \stackrel{\rm def.}{=} (C^l)_{l=1}^L$  the parameter bounds, and for  $l = 1, \ldots, L$  we have  $\mathcal{TB}_l \in \mathcal{TB}_{d^l, M, H^l, C^l}$ . Furthermore,  $A \in \mathbb{R}^{d \times M d_{\rm out}^L}$  and  $b \in \mathbb{R}^d$ .

The set of transformer networks with representation (2) and parameters bound by C is denoted by  $\mathcal{TC}_{L,H,C,W}^{D,d,M}$ , where  $C \stackrel{\text{def.}}{=} (*C',C^A,C^b)$ . If the specification is clear, we write  $\mathcal{TC}_{L,H,C,W}^{D,d,M} \equiv \mathcal{TC}$  for short.

**Notation** We make use of reshape operators (standard in most deep learning software) which, for any  $F_1, F_2 \in \mathbb{N}_+$ , are given by reshape  $F_1 \times F_2 = F_1 \times F_2 = F_2 \times F_2 = F_1 \times F_2 = F_2 \times F_2 = F_1 \times F_2 = F_2 \times F_2 = F_2 \times F_2 = F_1 \times F_2 = F_2 \times$ 

$$reshape_{F_1 \times F_2}(x)_{i,j} \stackrel{\text{def.}}{=} x_{(i-1)F_2+j}.$$

We denote the inverse of the map reshape  $F_1 \times F_2$  by  $\text{vec}_{F_1,F_2} : \mathbb{R}^{F_1 \times F_2} \to \mathbb{R}^{F_1 F_2}$ . We also recall the definition of the softmax function, defined by  $\text{softmax}(x) \stackrel{\text{def.}}{=} (\exp(x_i) / \sum_{j=0}^{F-1} \exp(x_j))_{i=0}^{F-1}$  for each  $F \in \mathbb{N}_+$  and each  $x \in \mathbb{R}^F$ . The maximum size of a norm in any vector in any  $K \subseteq \mathbb{R}^k$  is denoted by  $\|K\| \stackrel{\text{def.}}{=} \sup_{x \in K} \|x\|$ , the  $\ell^{\infty}$ -matrix norm of any  $n \times m$  matrix  $A \in \mathbb{R}^{n \times m}$  is denoted by  $C^A \stackrel{\text{def.}}{=} \max_{i \in \{1, \dots, n\}, j \in \{1, \dots, m\}} |A_{i,j}|$ . Fix  $d, D, s \in \mathbb{N}_+$  and a smooth function  $f : \mathbb{R}^d \to \mathbb{R}^D$  then its  $C^s$ -norm on a (regular closed) subset  $K \subseteq \mathbb{R}^D$  is defined to be

$$||f||_{s:K} \stackrel{\text{\tiny def.}}{=} \max_{\substack{k=1,\ldots,s-1\\\alpha \in \{1,1,d\}^k}} \sup_{u \in K} \left| \left| \frac{\partial^k f}{\partial x_{\alpha_1} \dots \partial x_{\alpha_k}}(u) \right| \right| + \max_{\alpha \in \{1,\ldots,d\}^{s-1}} \operatorname{Lip} \left( \frac{\partial^{s-1} f}{\partial x_{\alpha_1} \dots \partial x_{\alpha_{s-1}}} |K \right),$$

where Lip(g|K) denotes the Lipschitz constant of a function  $g: \mathbb{R}^d \to \mathbb{R}^D$  and  $\alpha$  is a multi-index. All additional notation, not required for our main-text results, is relegated to Appendix A.

#### 3. Main Result

We consider smooth loss and target functions functions that are concentrated on a compact region, along with their derivatives. The growth rate of the  $C^s$  norm of the loss function and its derivatives quantifies the degree of concentration. One easily verifies that any function in the Schwartz class satisfies this former of the following conditions.

**Definition 5 (Polynomial Growth of Derivatives)** Let  $d, D \in \mathbb{R}$ . A smooth function  $g : \mathbb{R}^d \to \mathbb{R}^D$  is in the class  $C^{\infty}_{poly:C,r}([0,1]^d,\mathbb{R}^D)$  if are  $C,r \geq 0$  such that:  $\|g\|_{C^s([0,1]^d)} \leq C$  s<sup>r</sup> for each  $s \in \mathbb{N}_+$ .

In Appendix C.2 we show that, in one dimension, any real analytic function whose power series expansion at 0, has coefficients growing at an  $\mathcal{O}((s+1)^r)$  rate belongs to  $g \in C^{\infty}_{poly:C,r}([0,1],\mathbb{R})$ . One can easily extend this argument to multiple dimensions to obtain further examples.

We consider an realizable PAC learning problem, determined by a smooth 1-Lipschitz target function  $f^*: \mathbb{R}^d \to \mathbb{R}^D$  which we would like to learn using a sequence of random observations  $((X_t, Y_t))_{t \in \mathbb{N}}$  as our training data. That is, for each  $t \in \mathbb{N}_+$ 

$$Y_t \stackrel{\text{def.}}{=} f^{\star}(X_t)$$

We aim to learn f from a *single path*. The ability of a model to reliably recover the function  $f: \mathbb{R}^d \to \mathbb{R}^D$  at time t, given the input  $X_t$ , is quantified by the t-future risk

$$\mathcal{R}_t(f) \stackrel{\text{def.}}{=} \mathbb{E} [\ell(f(X_t), f^{\star}(X_t))].$$

The time-t excess-risk  $\mathcal{R}_t$ , which is generally unobservable, is estimated by a single-path estimator known as the empirical risk computed using all the noisy samples to observed thus far

$$\mathcal{R}^{(N)}(f) \stackrel{\text{\tiny def.}}{=} \frac{1}{N} \sum_{n=1}^{N} \ell(f(X_t), f^{\star}(X_t)).$$

Our objective is to obtain a statistical learning guarantee on the quality of our estimate of the target function given by the time  $t \in \mathbb{N}_+$  generalization gap  $|\mathcal{R}_t(f) - \mathcal{R}^{(N)}(f)|$ .

We now summarize our setting and all parameters defining it, e.g. dimension, number of attention heads in the transformer, growth rate of the derivatives of the target and loss functions, etc.

Setting 3.1 Let  $D, d, L, H, *C', C^A, C^b \in \mathbb{N}_+$ ,  $C \stackrel{\text{def.}}{=} (*C', C^A, C^b)$ ,  $r_f, r_\ell, C_f, C_\ell \geqslant 0$ , and consider the hypothesis class  $\mathcal{TC}_{L,H,C}^{D,d,1}$ . Fix a target function  $f^* \in C_{poly:C_f,r_f}^{\infty}(\mathbb{R}^d, \mathbb{R}^D)$  and a loss function,  $\ell \in C_{poly:C_\ell,r_\ell}^{\infty}(\mathbb{R}^{2D}, \mathbb{R})$  and suppose that Assumptions 1 and 2 hold.

Having formalized our setting, we may now state our main result, which is a version of (FutureGen). This version provides insights on the future-generalization of transformers via: 1) explicit constants and 2) explicit *phase transition times* above which the convergence rate in (FutureGen) accelerates by a polylogarithmic factor. We express these times of convergence rate acceleration are expressed using the following *convergence rate function* 

$$\operatorname{rate}_{s}(N) \stackrel{\text{def.}}{=} \begin{cases} \frac{\log(c\,N)^{d-2s+s/d}}{c_{2}\,N^{s/d}} & \text{if } d > 2s \quad (initial\ phases) \\ \frac{\log(c\,N)}{c\,N^{1/2}} & \text{if } d = 2s \quad (critical\ phase) \\ \frac{\log(c\,N)^{d/(2s+1)}}{c\,N^{1/2}} & \text{if } d < 2s \quad (eventual\ phases) \end{cases} \tag{rate}$$

where  $c \stackrel{\text{def.}}{=} 1 - \kappa$ ,  $c_2 \stackrel{\text{def.}}{=} c^{s/d}$ , and  $0 < \kappa < 1$  are constants depending only on the data-generating Markov process.

Theorem 1 (Pathwise Generalization Bounds for Transformers) In Setting 3.1, there is a  $\kappa \in (0,1)$ , depending only on X., and a  $t_0 \in \mathbb{N}_0$  such that: for each  $t_0 \leqslant N \leqslant t \leqslant \infty$  and  $\delta \in (0,1]$  the following holds with probability at-least  $1-\delta$ 

$$\sup_{\mathcal{T} \in \mathcal{TC}_{L,H,C}^{D,d,0}} \left| \mathcal{R}_{\max\{t,N\}}(\mathcal{T}) - \mathcal{R}^{(N)}(\mathcal{T}) \right| \lesssim \sum_{s=0}^{\infty} I_{N \in [\tau_s,\tau_{s+1})} C_{\ell,\mathcal{T},K,s} \left( \underbrace{\kappa^t}_{(\mathbf{I})} + \underbrace{\frac{\sqrt{2\ln(1/\delta)}}{N^{1/2}}}_{(\mathbf{II})} + \underbrace{\frac{1}{2\ln(1/\delta)}}_{(\mathbf{III})} \right)$$

where  $\operatorname{rate}_s(N)$  is defined in (rate), and the constant  $C_{\ell,\mathcal{T},s,K}$  is Table 1 and the transition times  $(\tau_s)_{s=0}^{\infty}$  are given iteratively by  $\tau_0 \stackrel{\text{def.}}{=} 0$  and for each  $s \in \mathbb{N}_+$ 

$$\tau_s \stackrel{\text{\tiny def.}}{=} \inf\bigg\{t \geqslant \tau_{s-1}: \, C_{\ell,\mathcal{T},K,s}(\kappa^t + \mathrm{rate}_s(N) + \frac{\sqrt{\log(1/\delta)}}{\sqrt{N}}) \leqslant C_{\ell,\mathcal{T},K,s-1}(\kappa^t + \mathrm{rate}_{s-1}(N) + \frac{\sqrt{\log(1/\delta)}}{\sqrt{N}}) \bigg\}.$$

Furthermore,  $c \stackrel{\text{def.}}{=} 1 - \kappa$ ,  $c_2 \stackrel{\text{def.}}{=} c^{s/d}$ ,  $\kappa^{\infty} \stackrel{\text{def.}}{=} \lim_{t \to \infty} \kappa^t = 0$ , and  $\lesssim$  hides an absolute constant.

#### 3.1 Discussion

Theorem 1 implies the order estimate in (FutureGen). This is because  $C_{\ell,\mathcal{T},K,s}$  is constant in N and  $\mathrm{rate}_s(N) < \mathrm{rate}_{s-1}(N)$ ; thus, for every s > 0 the right-hand side our bound is eventually bounded by any  $C_{\ell,\mathcal{T},K,s}(\kappa^t + \sqrt{2\ln(1/\delta)}/N^{1/2} + \mathrm{rate}_s(N))$  for N large enough. However, unlike the order estimate (FutureGen), Theorem 1 provides an explicit description of the actual size of the future-generalization gap in terms of three factors which we now interpret.

Non-Stationarity Term: Term (I) quantifies the rate at which the data-generating Markov process X, becomes stationary. This term only depends on the time t and a constant  $0 < \kappa < 1$  determined only by X. We use the notational convention  $\kappa^{\infty} \stackrel{\text{def.}}{=} \lim_{t \to \infty} \kappa^t = 0$  to describe the limiting case.

Model Complexity Term (Phase Transitions): Term (II) captures the complexity of the transformer network in terms of the number of self-attention heads, depth, width, and the activation function used to define the class  $\mathcal{T}_{L,W,H}^{D,d,0}$ . Each constant  $C_1 \leqslant \ldots \leqslant C_s \leqslant \ldots$  collects the higher-order sensitivities ( $s^{th}$  order partial derivatives; where  $s \in \mathbb{N}_+$ ) of the transformer model. Each  $0 = \tau_0 \leqslant \tau_1 \leqslant \ldots \leqslant \tau_s \leqslant \text{indicates}$ the times at which there is a phase-transition in the convergence rate of the generalization bound accelerates. Once  $t \geq \tau_s$ , then the convergence rate of Term (II) accelerates, roughly speaking, by a reciprocal log-factor of  $1/\log(N)$ . Observe that the rate function is asymptotically equal to the rate function from the central limit theorem, as s tends to infinity; that is,  $\lim \operatorname{rate}_s(N) = 1/(c\sqrt{N})$ . The rate (rate) is the (optimal) rate at which the empirical measure generated by observations from a Markov process converges to its stationary distribution in 1-Wasserstein distance [54, 75]. The polylogarithmic factor is removable if the data is i.i.d; in which the rate cannot be improved further [44, 30].




Figure 2: Transition times: (y-axis) when the future-generalization bound accelerates by a polylogarithmic factor (in N) for a single transformer block in terms of the input dimension d (x axis).

**Probabilistic Validity Term:** Term (III) captures the cost of the bound being valid with probability at least  $1 - \delta$ . The convergence rate of this term cannot be improved due to the central limit theorem. It is responsible for the overall convergence rate of our generalization bound being "stuck" at the optimal rate of  $\mathcal{O}(1/\sqrt{N})$  from the central limit theorem; as the other two terms converge exponentially to 0.

The Constant  $C_{\ell,\mathcal{T},K,s}$ : The constant  $C_{\ell,\mathcal{T},K,s}$ , in Theorem 8, encodes the maximal size of the first s partial derivatives of any transformer in the class  $\mathcal{TC}$ . Thus, it encodes the complexity of the class  $\mathcal{TC}$  (e.g. int terms of number of attention heads, depth, width, etc...), the size of the compact set K, and the smoothness of the loss function and target functions.

We note that, any uniform generalization bound for smooth functions thus necessarily contains constants of the same order *hidden within the big*  $\mathcal{O}$ . See e.g. the entropy bound in [83, Theorem 2.71] which yields VC-dimension bounds via standard Dudley integral estimates in the i.i.d. case.

Critically, when the function class is defined by function composition, i.e. as in deep learning, then these maximal partial derivatives tend to grow factorially in s. This is a feature of the derivatives of composite functions in high dimensions as characterized by the multi-variate chain rule (i.e. the Faá di Bruno formula [33, 26]). The combinatorics of these partial derivatives is encoded by the coefficients in the well-studied bell-polynomials [13, 64, 87] whose growth rate has been recently understood in [52] and contains factors of the order of  $\mathcal{O}(\left(\frac{2s}{e\ln s}(1+o(1))\right)^s)$ .

Remark that, in the feedforward case, i.e. when no layernorms or multihead attention are used, then the s=1 case is bounded above by the well-studied path-norms; see e.g. [9, 69], which are simply the product of the weight matrices of in the network and serve as a simple upper-bound for the largest Lipschitz constant (i.e.  $C^1$  norm) of the class  $\mathcal{TC}$ . These constants are included as very specific cases of our constant bounds; whose general expression is complicated and is relegated to the appendix in Theorems 5 and 3. In the case of a single transformer block, the  $C_{\ell,\mathcal{T},K,s}$  is of the order of

$$\mathcal{O}\Big(\underbrace{C_{\ell,f^{\star}}}_{\text{Loss \& Target}}\underbrace{C_{K^{(3)}}^{\mathcal{LN}}(\leqslant s)^{s}C_{K^{(1)}}^{\mathcal{LN}}(\leqslant s)^{s^{3}}}_{\text{Derceptron}}\underbrace{C_{K^{(2)}}^{\mathcal{PL}}(\leqslant s)^{s^{s}}}_{\text{Multihead Attention dimensions Generic: }s\text{-th order Derivative}}^{d^{2}s^{3}}\Big]$$

where the "generic higher-order derivative constant" is  $C_s \stackrel{\text{def.}}{=} \frac{2s}{e \ln s} (1 + o(1))$ . Order estimates on each of these terms are in Table 1, precise analytical estimates are derived in Appendix D.3, and numerical estimates are given in Appendix B (see also Figure 3). If no layer norms, SLP, or multi head attention mechanisms are included in the class, then their respective terms in our order estimate should be taken to be 1.

Table 1: Bounds on the terms in defining the constant  $C_{\ell,\mathcal{T},K,s}$ , in Theorem 8, for a single attention block.

Term	Bound $(\mathcal{O})$
$c_{\ell,f^{\star}}$	$C_f^s s^{r_\ell + 2s^2} C_s^s$
$\mathcal{L}\mathcal{N}$	$s^{(1+s)/2}C_s^s$
$\mathcal{PL}$	$  B^{(1)}   +   B^{(2)}     A  ^s   \sigma  _{s: \left[\pm   a  _{\infty} \pm \sqrt{d_{\text{in}}}\right]} \text{ Width}(\mathcal{PL}) \tilde{C}_s^s$
$\mathcal{MH}$	$  B^{(1)}   +   B^{(2)}     A  ^{s}   \sigma  _{s: \left[\pm   a  _{\infty} \pm \sqrt{d_{\text{in}}}\right]} \text{Width}(\mathcal{PL})  \tilde{C}_{s}^{s} $ $  W     V   (c_{d}   Q     K  )^{s} \left(s^{2} \left(\frac{s}{e}\right)^{2s} C_{s}^{s}\right)$

Here  $C_s \stackrel{\text{def.}}{=} \frac{2s}{e \ln s} (1 + o(1))$ ,  $\tilde{C}_s \stackrel{\text{def.}}{=} s^{1/2} (\frac{n}{e})^s C_s^s$ ,  $c_d \stackrel{\text{def.}}{=} 2 \max\{d_{\text{in}}, d_K, d_V, d_{\text{ff}}, d_{\text{out}}\}$ , Width( $\mathcal{PL}$ ) is the width of the neural network  $\mathcal{PL}$ , where  $\|\cdot\|$  denotes the componentwise max matrix/vector norm.

Figure 3 illustrates the effect of various building blocks in the construction of a transformer (e.g. activation choice, multihead attention (MHA), layernorms) through their effect on the constants in our generalization bounds. Consequentially, we show the effect on the phase-transition times  $(\tau_t)_{t=0}^{\infty}$ , defined in Theorem 1, dictating when the bound accelerates by a polylogarithmic factor in N.

#### 3.2 Explanation of Proof

The first step in deriving our generalization bounds is to quantify the regularity of the transformer model as a function of its depth, number of attention heads, and norm of its weight matrices. By regularity, we mean the number and size of the continuous partial derivatives admitted by the transformer. To quantify the size of the partial derivatives of the transformer we first remark that it is *smooth*; that is, it admits continuous partial derivatives of all orders (see Theorem 3).

We will uniformly bound the generalization capabilities of the class of transformers  $\mathcal{T} \in \mathcal{TC}_{L,W,H}^{D,d,0}$  by instead uniformly bounding the generalization of any  $C^s$  functions on  $\mathbb{R}^d$  with  $C^s$ -norm at most




Figure 3: Effects of Transformer Components of FutureGen: (left to right.) The first figure shows the  $C^s$  bound of various activation functions according to results in Appendix D.3.3. The second illustrates  $C^s$  bounds for Multi-Head Attention (Definition 2), single-layer perceptrons, and the layer norm. The third shows the  $C^s$ -bound of a transformer block (Definition 3), distinguishing if the bound was computed level-specific (Corollary 1) or type specific (Theorem 4). The parameters used for the above plots are the base cases of Tables 2 to 6.

equal to the largest  $C^s$ -norm in the class  $\mathcal{TC}_{L,W,H}^{D,d,0}$ . That is, we control the right-hand side of

$$\sup_{\mathcal{T} \in \mathcal{T}_{L,W,H}^{D,d,M}} \left| \mathcal{R}_t(\mathcal{T}) - \mathcal{R}^{(N)}(\mathcal{T}) \right| \leq \sup_{\hat{f} \in \mathcal{C}_R^s(\mathbb{R}^d)} \left| \mathcal{R}_t(\hat{f}) - \mathcal{R}^{(N)}(\hat{f}) \right| \tag{3}$$

where  $R \stackrel{\text{def.}}{=} \sup_{\mathcal{T} \in \mathcal{TC}_{L,W,H}^{D,d,M}} \|\ell(\mathcal{T}, f^{\star})\|_{C^s}$  describes the higher-order fluctuations of the "difference" between the target function  $f^{\star}$  and any transformer in  $\mathcal{TC}_{L,H,W,C}^{D,d,0}$ , as quantified by the loss function  $\ell$ . Our first step is thus to bound R by upper-bounding maximal size of the  $s^{th}$  partial derivatives of all transformers in  $\mathcal{TC}_{L,W,H}^{D,d,M}$ . Explicit bounds are computed in Theorem 5, and their order estimates (as a function of s) are given in Theorem 6. Combing these estimates with the maximal  $s^{th}$  partial derivatives of the loss and target function, via a Faá di Bruno-type formula (in Theorem 2 or Lemma 3), which is a multivariant higher-order chain rule, yields our estimate for R in (3).

Now that we have bounded R, appearing in the supremum term in (3), it remains to translate this into a generalization bound. We can do this by relating it to the so-called *smooth Wasserstein distance*  $d_s$  between the distribution of the Markov chain at time  $\mu_t$  and its empirical distribution  $\mu^{(N)} \stackrel{\text{def.}}{=} 1/N \sum_{n=1}^{N} \delta_{X_n}$  obtained by collecting samples up to time N. The *smooth Wasserstein distance*  $d_s$ , studied by [54, 75, 48], is the integral probability metric (IPM)-type distance quantifying the distance between any two Borel probability measures  $\mu, \nu$  on  $\mathbb{R}^d$  as the maximal distance which they can produced when tested on any function in  $C_1^s(\mathbb{R}^d)$ ; that is

$$d_s(\mu,\nu) \stackrel{\text{\tiny def.}}{=} \sup_{g \in \mathcal{C}_s^s(\mathbb{R}^d)} \mathbb{E}_{X \sim \mu}[g(X)] - \mathbb{E}_{Y \sim \nu}[g(Y)].$$

The right-hand side (RHS) of (3) can be expressed as R times the  $d_s$  distance between the (true) distribution  $\mu_t$  of the process X. at time t and the (empirical) distribution  $\mu^{(N)}$  collected from samples

RHS (3) 
$$\leq \sup_{\hat{f} \in C_R^s(\mathbb{R}^d)} \|\ell(\hat{f}, f^*)\|_{C^s} d_s(\mu_t, \mu^{(N)}).$$
 (4)

The  $d_s$  distance between the process X, at time t, i.e.  $\mu_t$ , and the running empirical distribution  $\mu^{(N)}$  can be accomplished in two steps. First, we fast-forward time and bound the  $d_s$ -distance between  $\mu^{(N)}$  and the stationary distribution  $\mu_{\infty}$  of the data-generating Markov chain X. (at time  $t = \infty$ ). We then rewind time and bound the  $d_s$ -distance between the stationary distribution  $\mu_{\infty}$  and the distribution  $\mu_t$  of the Markov process up to time t; by setting up the i.i.d. concentration of measure results of [53, 75]. This last step is possible since our assumptions on X. essentially guarantee that it has a finite (approximate) mixing time.

#### 4. Conclusion

We provided a theoretical foundation for the *future-generalization* of transformer trained on a single perturbed realization of a time-series trajectory (Theorem 1). Our results thus help provide insight on the reliability of LLMs outside the i.i.d. framework and their principled use in time-series analysis.

We obtain explicit estimates on the constants in these generalization bounds which relied on explicitly bounding all the higher-order derivatives of transformers; in terms of their number of attention heads, activation functions, depth, width, and weights constraints (Theorems 5 and 6). These bounds can equally be used in conjunction with classical tools, e.g. Rademacher or VC-type bounds in the i.i.d. setting, or other applications where one needs to understand the higher-order sensitivities of transformers to their inputs.

# 5. Limitations and Future Work

Several dynamical systems and financial markets have long-term memory and thus are non-Markovian. In future work, it would be interesting to extend our results to cover such settings as well.

**Acknowledgements** AK acknowledges financial support from an NSERC Discovery Grant No. RGPIN-2023-04482 and their McMaster Startup Funds. RS is supported by Canada NSERC CGS-D Doctoral Grant. RS and AK acknowledge that resources used in preparing this research were provided, in part, by the Province of Ontario, the Government of Canada through CIFAR, and companies sponsoring the Vector Institute <a href="https://vectorinstitute.ai/partnerships/current-partners/">https://vectorinstitute.ai/partnerships/current-partners/</a>.

#### **Appendix Contents**

A	Notation	12
В	Elucidation of Constants in Theorem 1	13
$\mathbf{C}$	Supporting Technical Results on the $C^s$ -Norms of Smooth Functions C.1 Integral Probability Metrics and Restriction to Compact Sets	17 17 18
D	Proof of Theorem 1	18
	<ul> <li>D.1 Step 0 - Bounds on the C<sup>s</sup> Regularity of Multivariate Composite Functions</li> <li>D.1.1 Multivariate Faà di Bruno formula revisited</li> <li>D.1.2 Universal Bounds</li> <li>D.1.3 Bounds in Derivative Type</li> <li>D.2 Step 1 - Concentration of Measure - Bounding the Right-Hand Side of (3)</li> <li>D.3 Step 2 (A) - Bounding the C<sup>s</sup> Regularity of Transformer Building Blocks</li> <li>D.3.1 The Softmax Function</li> <li>D.3.2 The Multi-Head Self-Attention Mechanism</li> <li>D.3.3 The Activation Functions</li> <li>D.3.4 The Layer Norm</li> <li>D.3.5 The Multilayer Perceptron (Feedforward Neural Network) with Skip Connection</li> <li>D.4 Step 2 (B) - Transformers</li> </ul>	19 19 19 22 23 26 26 27 30 32 33 35
	D.5 Step 2 (C) - Merging the $C^s$ -Norm Bounds for Transformers with the Loss Function D.6 Step 3 - Combining Steps 1 and 2 and Completing The Proof of Theorem 1	36 39
$\mathbf{E}$	Example of Additive Noise Using Stochastic Calculus	40

# A. Notation

Notation 1 (Multi-index Notation) We will fix the following multivariate notation.

- Multi-indices  $\alpha \stackrel{\text{def.}}{=} (\alpha_1, \dots, \alpha_k) \in \mathbb{N}^k, k \in \mathbb{N}$  are denoted by Greek letters.
- The sum of entries is given by  $|\alpha| \stackrel{\text{def.}}{=} \sum_{i=1}^k \alpha_k$ .
- Its faculty is defined by  $\alpha! \stackrel{\text{def.}}{=} \prod_{i=1}^k \alpha_k!$ , We denote the derivative w.r.t.  $\alpha$  by  $D^{\alpha} \stackrel{\text{def.}}{=} \partial^{|\alpha|}/\partial x_1^{\alpha_1} \cdots \partial x_k^{\alpha_k}$  if  $|\alpha| > 0$  else  $D^{\alpha}$  is the identity
- For a vector x ∈ ℝ<sup>k</sup>, we write x<sup>α def</sup> = ∏<sub>i=1</sub><sup>k</sup> x<sub>k</sub><sup>α<sub>k</sub></sup>.
  We define the relation α ≺ β for β ∈ ℕ<sup>k</sup> if one of the three following holds
  - (i)  $|\alpha| < |\beta|$ ;
  - (ii)  $|\alpha| = |\beta|$ , and  $\alpha_1 < \beta_1$ ; or
- (iii)  $|\alpha| = |\beta|$ , and  $\alpha_i = \beta_i$  for  $i \in \{1, ..., j-1\}$  and  $\alpha_j < \beta_j$  for  $j \in \{2, ..., k\}$ .

   Unit vectors  $e_i \in \{0, 1\}^k$  are defined by  $(e_i)_j = 0$  for  $i \neq j$  and  $(e_i)_i = 1$ .

For any s>0, the norm  $\|\cdot\|_{C^s}$  of a smooth function  $f:\mathbb{R}^d\to\mathbb{R}$  is defined by

$$||f||_{C^s} \stackrel{\text{\tiny def.}}{=} \max_{k=1,\dots,s-1} \max_{\alpha \in \{1,\dots,d\}^d} \left\| \frac{\partial^k f}{\partial x_{\alpha_1} \dots \partial x_{\alpha_k}} \right\|_{\infty} + \max_{\alpha \in \{1,\dots,d\}^{s-1}} \operatorname{Lip}\left(\frac{\partial^{s-1} f}{\partial x_{\alpha_1} \dots \partial x_{\alpha_{s-1}}}\right)$$

We use the following notation to streamline the analytic challenges the tacking of  $C^s$ -norms.

Notation 2 (Order operator for multi-indeces) Define the order operator o for multi-indeces by

$$\mathfrak{o}: \mathbb{N}^k \longrightarrow \mathbb{N}^k, \quad \alpha_1, \dots, \alpha_k \longmapsto \alpha_{\tau_{\alpha}(1)}, \dots, \alpha_{\tau_{\alpha}(k)},$$

where  $\tau_{\alpha}: \{1, \ldots, k\} \to \{1, \ldots, k\}$  s.t.  $\alpha_{\tau_{\alpha}(1)} \ge \ldots \ge \alpha_{\tau_{\alpha}(k)}$ . We write  $\alpha \sim \beta$  if  $\mathfrak{o}(\alpha) = \mathfrak{o}(\beta)$  for  $\alpha, \beta \in \mathbb{N}^k$ . Further, denote by  $\mathfrak{D}_n^k$  the set  $\{\mathfrak{o}(\alpha) : \alpha \in \mathbb{N}^k, |\alpha| = n\}$  and write  $\mathfrak{D}_{\leq n}^k \stackrel{\text{def.}}{=} \{\mathfrak{o}(\alpha) : \alpha \in \mathbb{N}^k\}$  $\mathbb{N}^k, |\alpha| \leq n$ . Eventually, define  $N(\alpha) \stackrel{\text{def.}}{=} \# |\{\alpha' \in \mathbb{N}^k : \mathfrak{o}(\alpha') = \alpha\}|$ .

We will use the following notation to tabulate the sizes of a  $C^s$ -norm.

Notation 3 (Derivatives) Let  $k \in \mathbb{N}$ ,  $K \in \mathbb{R}^k$  be a set,  $f: K \to \mathbb{R}^m$  a function and  $\alpha \in \mathfrak{O}_n^k$  an ordered multi-index. Then,

• the uniform bound of  $\alpha$ -like derivatives on K is given by

$$C_K^f(\alpha) \stackrel{\text{def.}}{=} \max_{i \in \{1, \dots, m\}} \max_{\gamma \sim \alpha} \|D^{\gamma} f_i\|_K,$$

• we define the bound at / up to derivative level n by

$$C_K^f(n) \stackrel{\scriptscriptstyle\rm def.}{=} \max_{\alpha \in \mathfrak{O}_n^k} C_K^f(\alpha), \qquad C_K^f({\scriptstyle \leqslant} n) \stackrel{\scriptscriptstyle\rm def.}{=} \max_{\alpha \in \mathfrak{O}_{{\scriptstyle \leqslant} n}^k} C_K^f(\alpha),$$

- we write  $||K|| \stackrel{\text{def.}}{=} \sup_{x \in K} ||x||$ , and for a matrix  $A \in \mathbb{R}^{n \times m}$  we write  $C^A \stackrel{\text{def.}}{=} \max_{i \in \{1,\dots,n\}, j \in \{1,\dots,m\}} |A_{i,j}|$ .

When segmenting, truncating, or manipulating time series we will using the following notation.

Notation 4 (Time Series Notation) The following notation is when indexing paths of any time series.

- Realized Path up to time t is denoted by x<sub>≤t</sub> <sup>def.</sup> = (x<sub>s</sub>)<sub>s∈Z, s≤t</sub>.
  Segment of a Path Given a sequence x ∈ R<sup>Z</sup> and integers s ≤ t, we denote x<sub>[s:t]</sub> <sup>def.</sup> = (x<sub>i</sub>)<sup>t</sup><sub>i=s</sub>.

Lastly, we recorded some additional notations that were required throughout our manuscript.

Notation 5 (Miscellaneous) The following notation is when indexing paths of any time series.

- N-Simplex. For  $N \in \mathbb{N}$  we write  $\Delta_N \stackrel{\text{def.}}{=} \{u \in [0,1]^N : \sum_{i=1}^N u_i = 1\}$ . Infinite powers: For  $c \in (0,1)$ , we define  $c^{\infty} \stackrel{\text{def.}}{=} \lim_{t \to \infty} c^t = 0$ .

#### B. Elucidation of Constants in Theorem 1

The aim of this section is to elucidate the magnitude of the constants appearing in Theorem 1. We aim of to make each of these concrete by numerically estimating them, which we report in a series of tables. Importantly, we see how subtle choices of the activation function used to define the transformer model can have dramatic consequences on the size of these constants, which could otherwise be hidden in big  $\mathcal{O}$  notation.<sup>2</sup>

Interestingly, in Tables 2 and 3, we see that the softplus activation function produces significantly tighter bounds than the tanh activation function through much smaller constants, and the GeLU and SWISH activation functions are a relatively comparable second-place.

The bounds depicted in Table 3 exhibits a notable trait of independence from both input dimension and the compactum they are defined on. Notably, the selection of latent dimensionality demonstrates a relatively minor influence in contrast to the pronounced impact of parameter bounds. This suggests that while adjusting the latent dimension may have some effect, the primary driver of the derivative bound lies within the constraints imposed on the parameters. Despite the seemingly conservative nature of the chosen parameter-bounds, it is important to acknowledge their alignment with the parameter ranges observed in trained transformer-models.

<sup>2.</sup> The source code to compute derivative bounds is available at https://github.com/YannickLimmer/ transformer-bounds.

Note that the latter can be observed as well for Multi-Head attention (Table 6), however, we see that here the input dimension (composed of  $d_{\rm in}$  and M) is of greater importance with respect to the derivative bound.

Table 2:  $C^s$ -bounds of activation functions based on numerical maximization of analytic derivatives in Appendix D.3.3.

Bound	softplus	$\operatorname{GeLU}$	tanh	Swish
$C^1$	0.25	1.12	4.00	1.10
$C^2$	0.10	0.48	8.00	0.50
$C^3$	0.12	0.75	16.00	0.31
$C^4$	0.13	1.66	32.00	0.50
$C^5$	0.25	4.34	156.65	0.66
$C^6$	0.41	12.95	1651.32	1.50
$C^7$	1.06	42.77	20405.43	2.91
$C^8$	2.39	153.76	292561.95	8.50
$C^9$	7.75	594.17	4769038.09	21.76
$C^{10}$	22.25	2445.69	87148321.71	77.50

The bound of the layer-norm (see Table 4) seems to be particularly effected by the domain it is defined on, which can be problematic if it appears in later layers. An immediate solution is the usage of its parameter  $\gamma$ , a more drastic approach would be applications in combination with an upstream sigmoid activation.

Eventually, as also shown in Figure 3, we included in Tables 5 and 6 a comparison of using type-specific bounds (see Theorem 4) or level-specific bounds (Theorem 3) in the computation of the constants. This effect seems to become more evident with higher number of function compositions.

Table 3: Derivative Bounds of the Perceptron Layer by derivative level according to Lemma 12.

	Paran	neters		Ι	Perivative L	evel	
$\sigma$	$d_{ m ff}$	$C^{\{A,B^{(1)},B^{(2)}\}}$	1	2	3	4	5
softmax	64	1.0	17.00	50.47	236.94	1.34E+03	1.33E+04
tanh			129.00	1.02E+03	8.96E+03	9.83E+04	1.34E+06
$\operatorname{GeLU}$			73.25	237.41	1.25E+03	1.16E+04	1.81E+05
SWISH			71.39	236.78	870.75	5.33E+03	4.13E+04
	16		5.00	12.62	59.23	334.15	3.31E+03
	32		9.00	25.24	118.47	668.30	6.63E+03
	128		33.00	100.95	473.87	2.67E+03	2.65E+04
	256		65.00	201.90	947.75	5.35E+03	5.30E+04
		0.01	17.00	44.32	180.94	919.87	6.52E+03
		0.1	17.00	44.38	181.00	919.92	6.52E+03
		10.0	17.00	660.15	5.62E+04	4.17E+06	6.73E+08
		100.0	17.00	6.16E+04	5.60E+07	4.17E+10	6.73E+13

Table 4: Layer Norm

	Paramete	ers		Γ	erivative L	evel	
k	$\ K\ $	$\gamma$	1	2	3	4	5
5	10.0	0.1	18.67	28.56	104.49	1.49E+03	4.93E+03
3			18.67	28.56	104.49	945.21	4.93E+03
10			18.67	28.56	104.49	1.49E+03	4.93E+03
20			18.67	28.56	104.49	1.49E+03	4.93E+03
	0.1		0.17	3.61	5.37	7.05	8.87
	1.0		1.73	5.20	6.95	8.71	10.64
	100.0		321.71	7.68E+03	7.88E+05	1.42E+08	4.39E+09
	1000.0		321.71	7.68E+03	7.88E+05	1.42E+08	4.39E+09
		0.01	1.73	2.07	2.24	2.42	2.59
		1.0	321.71	7.75E+03	7.91E+05	1.42E+08	4.44E+09

Table 5: Derivative Bounds of Transformer Block by derivative level according to Theorem 6.

	Para	ameters			De	erivative Le	vel	
$d_{ m in}$	$_{C}^{\{K,Q,V,W\}}$	$C^{\{A,B^{(1,2)}\}}$	$\gamma$	1	2	3	4	5
5	0.01	0.001	0.01	21.15	1.13E+04	4.81E+06	2.59E+09	2.22E+11
	— using der	ivative level -		212.70	1.53E+06	7.47E+09	4.55E+13	3.75E+16
10				111.32	4.51E+05	1.71E+09	1.45E+13	8.70E+16
20				1.29E+03	1.25E+08	3.47E+13	1.85E+19	2.20E+24
	0.001			21.15	1.13E+04	4.81E+06	2.59E+09	2.22E+11
	0.1			21.16	1.13E+04	4.83E+06	2.61E+09	2.32E+11
	1.0			22.30	4.64E+04	6.96E+08	1.70E+13	1.95E+17
		0.0001		5.05	126.27	1.12E+04	4.94E+06	6.87E+08
		0.01		182.17	1.12E+06	4.66E+09	2.43E+13	1.71E+16
		0.1		1.79E+03	1.12E+08	4.65E+12	2.42E+17	1.70E+21
			0.0001	0.21	108.21	4.44E+04	2.27E+07	1.58E+09
			0.001	2.09	1.09E+03	4.46E+05	2.29E+08	1.60E+10
			0.1	240.09	2.45E+05	7.31E+08	4.96E+12	8.79E+15

Table 6: Derivative Bounds of Multi-Head Attention by derivative level according to Corollary 4.

d. M d.	$C^K$	Parameters $\mathcal{C}^{\mathcal{Q}}$	S. S.	$C_M$	$\parallel K \parallel$	_	n De	Derivative Level	vel 1	л
5 1 3	0.1	0.1	0.1	0.1	1.0	7.67	46.08	184.82	931.81	5.73E+03
	— using	using derivative level	e level —			7.67	46.15	186.90	1.01E+03	7.84E+03
10						7.82E+03	4.87E+04	8.14E+05	2.95E+08	2.38E+11
20						1.57E+04	1.09E+05	1.03E+07	9.25E+09	1.50E+13
57						3.90E+03	2.37E+04	1.34E+05	9.96E+06	3.83E+09
20						4.71E+06	2.85E+07	7.11E+08	2.66E+12	1.40E+16
3						7.67	45.99	183.98	920.53	5.53E+03
10						7.70	46.41	190.35	1.07E+03	1.03E+04
20						7.75	47.52	226.02	2.87E+03	1.23E+05
	0.01					7.90	54.45	740.55	6.51E+04	9.33E+06
	1.0					7.65	45.91	183.61	918.01	5.51E+03
		0.01				7.65	45.91	183.61	918.01	5.51E+03
		1.0				7.90	54.45	740.55	6.51E+04	9.33E+06
			0.01			0.77	4.61	18.48	93.18	573.32
			1.0			76.75	460.80	1.85E+03	9.32E+03	5.73E+04
				0.01		0.77	4.61	18.48	93.18	573.32
				1.0		76.75	460.80	1.85E+03	9.32E+03	5.73E+04
	0.001	0.001	0.001	0.001		1.02E+03	8.06E+04	4.95E+07	5.70E+10	8.04E+13
	0.01	0.01	0.01	0.01		0.08	0.46	1.84	9.18	55.08
	1.0	1.0	1.0	1.0		0.00	0.00	0.02	0.09	0.55
					0.1	0.77	32.14	142.34	752.79	4.68E+03
					10.0	79.00	259.65	5.54E+03	5.73E+05	8.04E+07
					100.0	1.02E+03	7.66E+04	4.88E+07	5.63E+10	7.90E+13

# C. Supporting Technical Results on the $C^s$ -Norms of Smooth Functions

This section contains many of the technical tools on which we build our analysis. Most results concern smooth functions, especially their derivatives and those of compositions thereof. However, the first set of results concerns the integral probability metric  $d_s$ .

#### C.1 Integral Probability Metrics and Restriction to Compact Sets

Fix  $d \in \mathbb{N}_+$  and a non-empty compact subset  $K \subseteq \mathbb{R}^d$ . Observe that any Borel probability measure  $\mu$  on K can be canonically extended to a compactly supported Borel probability measure  $\mu^+$  on all of  $\mathbb{R}^d$  via

$$\mu^+(B) \stackrel{\text{\tiny def.}}{=} \mu(B \cap K),$$

for any Borel subset B of  $\mathbb{R}^d$ ; noting only that  $B \cap K$  is Borel.

Let  $\mathcal{P}(K)$  denote the set of Borel probability measures on K. Suppose that K is a regular compact set, i.e. the closure of its interior is itself. As usual, see [32], for any  $s \in \mathbb{N}_+$ , we denote the set of functions from the interior of K to  $\mathbb{R}$  with s continuous partial derivatives thereon and with a continuous extension to K by  $C^s(K)$ . This space, is a Banach space when equipped with the (semi-)norm

$$\|f\|_{s:K} \stackrel{\text{\tiny def.}}{=} \max_{k=1,\dots,s-1} \max_{\alpha \in \{1,\dots,d\}^k} \sup_{u \in K} \left\| \frac{\partial^k f}{\partial x_{\alpha_1} \dots \partial x_{\alpha_k}}(u) \right\| + \max_{\alpha \in \{1,\dots,d\}^{s-1}} \operatorname{Lip} \left( \frac{\partial^{s-1} f}{\partial x_{\alpha_1} \dots \partial x_{\alpha_{s-1}}} \right).$$

We may define an associated integral probability metric  $d_{s:K}$  on  $\mathcal{P}(K)$  via

$$d_{s:K}(\mu,\nu) \stackrel{\text{def.}}{=} \sup_{f \in C^s(K)} \|\mathbb{E}_{X \sim \mu}[f(X)] - \mathbb{E}_{X \sim \mu}[f(X)]\|$$

for any  $\mu, \nu \in \mathcal{P}(K)$ . The main purpose of this technical subsection is simply to reassure ourselves, and the reader, that quantities  $d_{s:K}(\mu, \nu)$  and  $d_s(\mu^+, \nu^+)$  are equal for any  $\mu, \nu \in \mathcal{P}(K)$ . Therefore, we may use them interchangeably.

Lemma 1 (Consistency of Smooth IMP Extension - Beyond Regular Compact Sets) Fix  $d, s \in \mathbb{N}_+$  and let K be a non-empty regular compact subset of  $\mathbb{R}^d$ . For any  $\mu, \nu \in \mathcal{P}(K)$  the following holds

$$d_{s:K}(\mu, \nu) = d_s(\mu^+, \nu^+).$$

**Proof** Let  $\operatorname{int}(K)$  denote the interior of K, By the Whitney extension theorem, as formulated in [35, Theorem A], for any  $f \in C^s(K)$  there exists a  $C^s$ -extension  $F : \mathbb{R}^d \to \mathbb{R}$  of  $f|_{\operatorname{int}(K)}$  to all of  $\mathbb{R}^d$ ; i.e.  $F|_{\operatorname{int}(K)} = f$  and  $\in C^s(\mathbb{R}^d)$ . Since any continuous function is uniformly continuous on a compact set,  $\operatorname{int}(K)$  is dense in K, and since uniformly continuous functions are uniquely determined by their values on compact sets, then f coincides with F on all of K (not only on  $\operatorname{int}(K)$ ).

For any  $\mu \in \mathcal{P}(K)$ , by definition of  $\mu^+$  we have that

$$\mathbb{E}_{X \sim \mu^+}[F(X)] = \mathbb{E}_{X \sim \mu^+}[F(X)I_{X \in K}] = \mathbb{E}_{X \sim \mu^+}[f(X)I_{X \in K}] = \mathbb{E}_{X \sim \mu}[f(X)].$$

Therefore, for any  $\mu, \nu \in \mathcal{P}(K)$  we conclude that and each  $f \in C^s(K)$  there exists some  $F \in C^s(\mathbb{R}^d)$  such that

$$\mathbb{E}_{Y \sim \mu}[f(Y)] - \mathbb{E}_{Y \sim \nu}[f(Y)] = \mathbb{E}_{X \sim \mu^+}[F(X)] - \mathbb{E}_{X \sim \nu^+}[F(X)].$$

Consequentially,  $d_{s:K}(\mu,\nu) \leq d_s(\mu^+,\nu^+)$ . Conversely, since the restriction of any  $g \in C^s(\mathbb{R}^d)$  to K belongs to  $C^s(K)$  then the reverse inequality holds; namely,  $d_{s:K}(\mu,\nu) \geq d_s(\mu^+,\nu^+)$ .

By Lemma 1 we henceforth may interpret any such  $\mu$  as its extension  $\mu^+$ , without loss of generality.

# C.2 Examples of Functions in The Classes $C^s_{poly:C,r}([0,1]^d,\mathbb{R})$ and $C^s_{exp:C,r}([0,1]^d,\mathbb{R})$

In several learning theory papers, especially in the kernel ridge regression literature e.g. [77, 11, 82, 77, 23, 24], one often quantifies the *learnability* of a target function in terms of some sort of decay/growth rates of its coefficients in an appropriate expansion; e.g. the decay of its coefficients in an eigenbasis associated to a kernel. These decay/growth rates are often equivalent to the smoothness of a function<sup>3</sup>. Therefore, in a like spirit, we unpack the meaning of the smoothness condition in Assumption 5 which impacts the learning rates in Theorem 1 by giving examples of functions in the classes  $C_{solveC,r}^s([0,1]^d,\mathbb{R})$  and  $C_{soveC,r}^s([0,1]^d,\mathbb{R})$ .

classes  $C^s_{poly:C,r}([0,1]^d,\mathbb{R})$  and  $C^s_{exp:C,r}([0,1]^d,\mathbb{R})$ . For brevity and transparency in our illustration, we consider the one-dimensional case. In particular, this shows that the class is far from being void.

**Proposition 1 (Functions with polynomially/exponentially growing**  $C^s$ -norms on [0,1]) Fix  $d \in \mathbb{N}_+$  and let K be a non-empty regular compact subset of  $\mathbb{R}^d$ . If  $f : \mathbb{R} \to \mathbb{R}$  is real-analytic with power-series expansion at 0 given by

$$f(x) = \sum_{i=0}^{\infty} \frac{\beta_i x^i}{i!}.$$

If there are C, r > 0 such that

- (i) **Polynomial Growth:**  $|\beta_i| \leq Ce^{ir} \ (\forall i \in \mathbb{N}) \ then \ f \in C^{\infty}_{poly:C,r}([0,1],\mathbb{R}),$
- (ii) **Exponential Growth:**  $|\beta_i| \leq C(1+i)^r \ (\forall i \in \mathbb{N}) \ then \ f \in C^{\infty}_{poly:C,r}([0,1],\mathbb{R}),$

**Proof** Since f is real-analytic we may consider its Maclaurin-Taylor series expansion which coincides with  $\sum_{i=0}^{\infty} \frac{\beta_i x^i}{i!}$ ; meaning that for each  $i \in \mathbb{N}$  we have  $\beta_i = \partial^i f(0)$ . Therefore, standard analytic estimates and manipulations of the Maclaurin-Taylor series, see e.g. [76, page 173], yield

$$\max_{0 \leqslant x \leqslant 1} \left| \sum_{i=0}^{\infty} \frac{\beta_{i} x^{i}}{i!} - f(x) \right| \leqslant \frac{1}{(s+1)!} \sup_{0 \leqslant x \leqslant 1} \left| \left( \sum_{i=0}^{\infty} \frac{\beta_{i} x^{i}}{i!} \right)^{s+1} - \partial f^{s+1}(x) \right| 
\leqslant \frac{1}{(s+1)!} \beta_{s}(s+1)!.$$
(5)

If (i) holds, then the right-hand side of (5) is bounded above by  $C(s+1)^r$  and  $f \in C^{\infty}_{poly:C,r}([0,1],\mathbb{R})$ . If, instead, (ii) holds then the right-hand side of (5) is bounded above by  $C e^{s\,r}$  and  $f \in C^{\infty}_{exp:C,r}([0,1],\mathbb{R})$ .

#### D. Proof of Theorem 1

Section 3.2, the proof will be largely broken down into two steps. First, we derive our concentration of measure result for the empirical mean compared to the true mean general of an arbitrary  $C^s$  function applied to a random input, where the  $C^s$ -norm of the  $C^s$  function is at most  $R \ge 0$  (in Subsection D.2).

Next, (in Subsection D.2), we use the Faà di Bruno-type results in Section D.1 to bound the maximal  $C^s$  norm over the relevant class of transformer networks. We do this by first individually bounding each of the  $C^s$ -norms of its constituent pieces, namely the multi-head attention layers, the SLP blocks with smooth activation functions, and then ultimately, we bound the  $C^s$ -norms of the composition of transformer blocks using the earlier Faà di Bruno-type results.

<sup>3.</sup> See e.g. [4, page 120-121] for an example between the decay rate of the Laplacian eigenspectrum characterize the smoothness of the functions in the RKHS of radially symmetric kernels.




Figure 4: Workflow of the proof technique used to derive Theorem 1.

Our main result (Theorem 1) is then then obtained upon merging these two sets of estimates. The workflow which we use can be applied to derive generalization bounds for other machine learning, and is summarized in Figure 4.

# D.1 Step 0 - Bounds on the $C^s$ Regularity of Multivariate Composite Functions

In this section, we will derive a bound for the Sobolev norm of multivariate composite functions.

# D.1.1 Multivariate Faà di Bruno formula revisited

We begin by establishing notation and stating the multivariate Faà di Bruno formula from [26].

Theorem 2 (Multivariate Faà di Bruno Formula, [26]) Let  $n, m, k \in \mathbb{N}$ ,  $\alpha \in \mathbb{N}^k$  with  $|\alpha| = n$ , and define

$$h(x_1,\ldots,x_k) \stackrel{\text{def.}}{=} f^{(1)}(g^{(1)}(x_1,\ldots,x_k),\ldots,g^{(m)}(x_1,\ldots,x_k)).$$

Then, using the multivariate notation from Notation 1,

$$D^{\alpha}h(x) = \sum_{1 \leq |\beta| \leq n} (D^{\beta}f)(g(x)) \sum_{\eta, \zeta \in \mathcal{P}(\alpha, \beta)} \alpha! \prod_{j=1}^{n} \frac{[D^{\zeta^{(j)}}g(x)]^{\eta^{(j)}}}{\eta^{(j)}!(\zeta^{(j)}!)^{|\eta^{(j)}|}}.$$

where

$$\mathcal{P}(\alpha, \beta) = \left\{ \eta \stackrel{\text{def.}}{=} (\eta^{(1)}, \dots, \eta^{(n)}) \in (\mathbb{N}^m)^n, \zeta \stackrel{\text{def.}}{=} (\zeta^{(1)}, \dots, \zeta^{(n)}) \in (\mathbb{N}^k)^n : \\ \exists j \leqslant m : \eta^{(i)} = 0, \zeta^{(i)} = 0 \text{ for } i < j, |\eta^{(i)}| > 0 \text{ for } i \geqslant j, \\ 0 \prec \zeta^{(j)} \prec \dots \prec \zeta^{(n)}, \sum_{i=1}^n \eta^{(i)} = \beta \text{ and } \sum_{i=1}^n |\eta^{(i)}| \zeta^{(i)} = \alpha \right\}.$$

Proof See [26].

#### D.1.2 Universal Bounds

**Theorem 3** In the notation of Theorem 2, we have for a compact set  $K \subseteq \mathbb{R}^k$  and an multi-index  $\alpha \in \mathbb{N}^k$ ,  $|\alpha| = n$ ,

$$C_K^h(\alpha) \leqslant \max_{n' \in \{1, \dots, n\}} C_{g[K]}^g(n') C_K^f(\leqslant n)^{n'} \sum_{1 \leqslant |\beta| \leqslant n} \sum_{\eta, \zeta \in \mathcal{P}(\alpha, \beta)} \alpha! \prod_{j=1}^n \frac{1}{\eta^{(j)!} (\zeta^{(j)!})^{|\eta^{(j)}|}}$$

where  $C_K^h(\cdot)$ ,  $C_{q[K]}^f(\cdot)$ ,  $C_K^g(\cdot)$  are defined as in Notation 3.

**Proof** Using Theorem 2,

$$\begin{split} C_K^h(\alpha) &\leqslant \sum_{1 \leqslant |\beta| \leqslant n} \|D^{\beta} f\|_{g[K]} \sum_{\eta, \zeta \in \mathcal{P}(\alpha, \beta)} \alpha! \prod_{j=1}^n \frac{\prod_{i=1}^m \|(D^{\zeta^{(j)}}g)_i\|_K^{\eta_i^{(j)}}}{\eta^{(j)}! (\zeta^{(j)}!)^{|\eta^{(j)}|}} \\ &\leqslant \sum_{1 \leqslant |\beta| \leqslant n} C_{g[K]}^g(|\beta|) \sum_{\eta, \zeta \in \mathcal{P}(\alpha, \beta)} \alpha! \prod_{j=1}^n \frac{\prod_{i=1}^m C_K^f(\leqslant n)^{\eta_i^{(j)}}}{\eta^{(j)}! (\zeta^{(j)}!)^{|\eta^{(j)}|}} \\ &\leqslant \sum_{1 \leqslant |\beta| \leqslant n} C_{g[K]}^g(|\beta|) C_K^f(\leqslant n)^{|\beta|} \sum_{\eta, \zeta \in \mathcal{P}(\alpha, \beta)} \alpha! \prod_{j=1}^n \frac{1}{\eta^{(j)}! (\zeta^{(j)}!)^{|\eta^{(j)}|}} \\ &\leqslant \max_{n' \in \{1, \dots, n\}} C_{g[K]}^g(n') C_K^f(\leqslant n)^{n'} \sum_{1 \leqslant |\beta| \leqslant n} \sum_{\eta, \zeta \in \mathcal{P}(\alpha, \beta)} \alpha! \prod_{j=1}^n \frac{1}{\eta^{(j)}! (\zeta^{(j)}!)^{|\eta^{(j)}|}}. \end{split}$$

Next, we refine the strategy used in [49] to convert our uniform risk-bound to a concentration of measure problem. Once done, the remainder of the proof will be to obtain bounds on the rate at which this measure concentrates.

**Lemma 2** For  $\alpha \in \{1, \dots, k\}^n$ , it satisfies that

$$\sum_{1\leqslant |\beta|\leqslant n}\sum_{\eta,\zeta\in\mathcal{P}(\alpha,\beta)}\alpha!\prod_{j=1}^n\frac{1}{\eta^{(j)}!(\zeta^{(j)}!)^{|\eta^{(j)}|}}=\left[\frac{2m|\alpha|}{e\ln|\alpha|}(1+o(1))\right]^{|\alpha|}$$

where  $\mathcal{P}(\alpha, \beta)$  is as defined in Theorem 2.

**Proof** Consider functions

$$g^{(i)}(x) = g^{(i)}(x_1, \dots, x_d) \stackrel{\text{def.}}{=} \exp\left(\sum_{j=1}^d x_j\right) : \mathbb{R}^d \to \mathbb{R}, \quad i = 1, \dots, 2m,$$

$$f(g^{(1)}, \dots, g^{(2m)}) \stackrel{\text{def.}}{=} \exp\left(\sum_{i=1}^{2m} g^{(i)}\right) : \mathbb{R}^{2m} \to \mathbb{R},$$

Since

$$\frac{\partial}{\partial g^{(i)}} f(g^{(1)}, \dots, g^{(2m)}) = f(g^{(1)}, \dots, g^{(2m)}),$$

it follows that

$$(D^{\beta}f)(q^{(1)}(x), \dots, q^{(2m)}(x)) = f(q^{(1)}(x), \dots, q^{(2m)}(x)), \quad \forall \beta \in \{1, \dots, 2m\}^n.$$

Since

$$\frac{\partial}{\partial x^{j}} f(g^{(1)}(x_{1}, \dots, x_{k}), \dots, g^{(2m)}(x_{1}, \dots, x_{k}))$$

$$= \sum_{i=1}^{2m} \frac{\partial}{\partial g^{(i)}} f(g^{(1)}(x_{1}, \dots, x_{k}), \dots, g^{(2m)}(x_{1}, \dots, x_{k})) \frac{\partial g^{(i)}(x_{1}, \dots, x_{k})}{\partial x_{j}}$$

$$= \sum_{i=1}^{2m} f(g^{(1)}(x_{1}, \dots, x_{k}), \dots, g^{(2m)}(x_{1}, \dots, x_{k})) g^{(i)}(x_{1}, \dots, x_{k})$$

$$= \sum_{i=1}^{2m} \frac{\partial}{\partial g^{(i)}} f(g^{(1)}(x), \dots, g^{(2m)}(x)) \frac{\partial g^{(i)}(x)}{\partial x_{j}}$$

$$= f(g^{(1)}(x), \dots, g^{(2m)}(x)) \sum_{i=1}^{2m} g^{(i)}(x)$$

and

$$\frac{\partial g^{(i)}(x_1,\cdots,x_k)}{\partial x_i} = g^{(i)}(x_1,\cdots,x_k),$$

we can show by the Faà di Bruno formula that

$$\begin{split} &D^{\alpha}f(g^{(1)}(x), \cdots, g^{(2m)}(x)) \\ &= D^{\alpha} \exp\left(\sum_{i=1}^{2m} g^{(i)}(x)\right) \\ &= \sum \frac{|\alpha|!}{\gamma_1!(1!)^{\gamma_1} \cdots \gamma_{|\alpha|}!(|\alpha|!)^{\gamma_{|\alpha|}}} \left(D^{\gamma_1 + \cdots + \gamma_{|\alpha|}} \exp\right) \left(\sum_{i=1}^{2m} g^{(i)}(x)\right) \prod_{i=1}^{|\alpha|} \left[m^j \left(\sum_{i=1}^{2m} g^{(i)}(x)\right)\right]^{\gamma_j}, \end{split}$$

where the summation on the right side of the last equality is over all  $|\alpha|$ -tuples  $(\gamma_1, \dots, \gamma_{|\alpha|}) \ge 0$  such that  $1 \cdot \gamma_1 + 2 \cdot \gamma_2 + \dots + |\alpha| \cdot \gamma_{|\alpha|} = |\alpha|$ .

By the multivariate Faà di Bruno formula. For each n=1, ..., s-1 fixed, and for each  $\alpha \in \{1, \dots, k\}^n$ , we have

$$D^{\alpha} f(g^{(1)}(x), \dots, g^{(2m)}(x)) = \sum_{1 \leq |\beta| \leq n} (D^{\beta} f)(g^{(1)}(x), \dots, g^{(2m)}(x)) \sum_{\eta, \zeta \in \mathcal{P}(\alpha, \beta)} \alpha! \prod_{j=1}^{n} \frac{[D^{\zeta^{(j)}}(g^{(1)}(x), \dots, g^{(2m)}(x))]^{\eta^{(j)}}}{\eta^{(j)!} (\zeta^{(j)!})^{|\eta^{(j)}|}}.$$

Taking  $x = (x_1, \dots, x_k) = 0$ , we have

$$D^{\alpha}f(g^{(1)}(0), \dots, g^{(2m)}(0)) = \sum \frac{|\alpha|!}{\gamma_1!(1!)^{\gamma_1} \dots \gamma_{|\alpha|}!(|\alpha|!)^{\gamma_{|\alpha|}}} \exp(2m) \prod_{j=1}^{|\alpha|} (2m)^{\gamma_j} (m^{\beta}f)(g^{(1)}(0), \dots, g^{(2m)}(0)) = f(g^{(1)}(0), \dots, g^{(2m)}(0)) = \exp(2m),$$

$$D^{\zeta^{(j)}}(g^{(1)}(x), \dots, g^{(2m)}(x)) = (1, \dots, 1).$$

Substituting the above derivatives into the Faà di Bruno formula, we obtain

$$\sum_{1 \leqslant |\beta| \leqslant n} \sum_{\eta, \zeta \in \mathcal{P}(\alpha, \beta)} \alpha! \prod_{j=1}^{k} \frac{1}{\eta^{(j)!} (\zeta^{(j)!})^{|\eta^{(j)}|}} = \sum \frac{|\alpha|!}{\gamma_{1}! (1!)^{\gamma_{1}} \cdots \gamma_{|\alpha|}! (|\alpha|!)^{\gamma_{|\alpha|}}} \prod_{j=1}^{|\alpha|} (2m)^{\gamma_{j}} \\
\leqslant (2m)^{|\alpha|} \sum \frac{|\alpha|!}{\gamma_{1}! (1!)^{\gamma_{1}} \cdots \gamma_{|\alpha|}! (|\alpha|!)^{\gamma_{|\alpha|}}} \\
= (2m)^{|\alpha|} \left(\frac{|\alpha|}{e \ln |\alpha|}\right)^{|\alpha|} (1 + o(1))^{|\alpha|},$$

where the last equality follows from [52, Theorem 2.1],

$$\sum \frac{|\alpha|!}{\gamma_1!(1!)^{\gamma_1}\cdots\gamma_{|\alpha|}!(|\alpha|!)^{\gamma_{|\alpha|}}} = \left(\frac{|\alpha|}{e\ln|\alpha|}\right)^{|\alpha|}(1+o(1))^{|\alpha|}.$$

Corollary 1 (Level Specific  $C^s$ -Norm Bounds for Transformer Blocks) In the notation of Theorem 3, it holds for  $n \in \mathbb{N}, n > 1$  that

$$C_K^h(\leqslant n) \leqslant \max_{n' \in \{1, \dots, n\}} C_{g[K]}^g(n') C_K^f(\leqslant n)^{n'} \left[ \frac{2mn}{e \ln n} (1 + o(1)) \right]^n.$$

and if  $C_K^f(\leqslant n) \geqslant 1$ ,

$$C_K^h(\leqslant n) \leqslant C_{g[K]}^g(\leqslant n) C_K^f(\leqslant n)^n \Big[ \frac{2mn}{e \ln n} (1+o(1)) \Big]^n.$$

**Proof** Follows directly from Theorem 3 and Lemma 2.

#### D.1.3 Bounds in Derivative Type

The goal of this section is to bound the derivative of composite functions by grouping with respect to  $\sim$ , defined in Notation 2.

**Theorem 4** In the notation of Theorem 2, we have for a compact set  $K \subseteq \mathbb{R}^k$  and an ordered multi-index  $\alpha \in \mathfrak{O}_n^k$ 

$$C_K^h(\alpha) \leqslant \alpha! \sum_{\beta \in \mathfrak{D}_{\leqslant n}^m} N(\beta) C_{g[K]}^f(\beta) \sum_{\eta, \zeta \in \mathcal{P}'(\alpha, \beta)} \prod_{j=1}^n \frac{C_K^g(\mathfrak{o}(\zeta^{(j)}))^{|\eta^{(j)}|}}{\eta^{(j)}! (\zeta^{(j)}!)^{|\eta^{(j)}|}},$$

where  $C_K^h(\cdot)$ ,  $C_{g[K]}^f(\cdot)$ ,  $C_K^g(\cdot)$  are defined as in Notation 3; and

$$\begin{split} \mathcal{P}'(\alpha,\beta) &= \Big\{ \eta \stackrel{\text{\tiny def.}}{=} (\eta^{(1)},\dots,\eta^{(n)}) \in (\mathbb{N}^m)^n, \zeta \stackrel{\text{\tiny def.}}{=} (\zeta^{(1)},\dots,\zeta^{(n)}) \in (\mathbb{N}^k)^n : \\ &\exists j \leqslant m : \eta^{(i)} = 0, \zeta^{(i)} = 0 \text{ for } i < j, |\eta^{(i)}| > 0 \text{ for } i \geqslant j, \\ &0 < \zeta^{(j)} \lhd \dots \lhd \zeta^{(n)}, \sum_{i=1}^n \eta^{(i)} = \beta \text{ and } \sum_{i=1}^n |\eta^{(i)}| \zeta^{(i)} = \alpha \Big\}, \end{split}$$

where  $\alpha \lhd \beta$  for  $\alpha, \beta \in \mathbb{N}^k$  if  $|\alpha| \leq |\beta|$  and  $\alpha \neq \beta$ .

**Proof** We have for  $\alpha \in \mathfrak{O}_n^k$ 

$$\begin{split} C_K^h(\alpha) \leqslant \max_{\gamma \sim \alpha} \sum_{1 \leqslant |\beta| \leqslant n} \|D^{\beta} f\|_{g[K]} \sum_{\eta, \zeta \in \mathcal{P}(\gamma, \beta)} \alpha! \prod_{j=1}^n \frac{\prod_{i=1}^m \|(D^{\zeta^{(j)}} g)_i\|_K^{\eta_i^{(j)}}}{\eta^{(j)}! (\zeta^{(j)}!)^{|\eta^{(j)}|}} \\ \leqslant \sum_{1 \leqslant |\beta| \leqslant n} C_{g[K]}^f(\mathfrak{o}(\beta)) \max_{\gamma \sim \alpha} \sum_{\eta, \zeta \in \mathcal{P}(\gamma, \beta)} \alpha! \prod_{j=1}^n \frac{C_K^g(\mathfrak{o}(\zeta^{(j)}))^{|\eta^{(j)}|}}{\eta^{(j)}! (\zeta^{(j)}!)^{|\eta^{(j)}|}}. \end{split}$$

Then

$$\left\{\eta,(\mathfrak{o}(\zeta^{(1)}),...,\mathfrak{o}(\zeta^{(n)}))\Big|(\eta,\zeta)\in\mathcal{P}'(\alpha,\beta)\right\}$$

is invariant in  $\alpha$  with respect to  $\sim$  and thus

$$C_K^h(\alpha) \leqslant \sum_{1 \leqslant |\beta| \leqslant n} C_{g[K]}^f(\mathfrak{o}(\beta)) \sum_{n, \zeta \in \mathcal{P}'(\alpha, \beta)} \alpha! \prod_{j=1}^n \frac{C_K^g(\mathfrak{o}(\zeta^{(j)}))^{|\eta^{(j)}|}}{\eta^{(j)}!(\zeta^{(j)}!)^{|\eta^{(j)}|}}.$$

Further, notice that

$$\left\{ ((|\eta^{(1)}|, \eta^{(1)}!), \dots, (|\eta^{(n)}|, \eta^{(n)}!)), \zeta \middle| (\eta, \zeta) \in \mathcal{P}'(\alpha, \beta) \right\}$$

is invariant in  $\beta$  with respect to  $\sim$  and the assertion follows.

**Corollary 2** In the notation of Theorem 4, if f is affine-linear,

$$C_K^h(\alpha) \leqslant m\alpha! C_{q[K]}^f(e_1) C_K^g(\alpha),$$

where  $C_{q[K]}^f(e_1)$  is the maximum weight of the matrix representing f.

**Proof** Theorem 4 yields

$$C_K^h(\alpha) \leq m\alpha! C_{g[K]}^f(e_1) \sum_{\eta, \zeta \in \mathcal{P}'(\alpha, \beta)} \prod_{j=1}^n \frac{C_K^g(\mathfrak{o}(\zeta^{(j)}))^{|\eta^{(j)}|}}{\eta^{(j)!} |(\zeta^{(j)}!)^{|\eta^{(j)}|}},$$

and since  $\mathcal{P}'(\alpha, e_1) = \{(0, ..., 0, e_1), (0, ..., 0, \alpha)\}$  the result follows.

# D.2 Step 1 - Concentration of Measure - Bounding the Right-Hand Side of (3)

We are now ready to derive our main concentration of measure results used to derive our risk-bound. This corresponds to bounding term 3 by controlling the integral probability term  $d_s(\mu_t, \mu^{(N)})$  in (4), with high probability, where the randomness is due to the randomness of the empirical measure  $\mu^{(N)}$ .

Instead of the compact support Assumption 2, our concentration of measure rate, with the same convergence rate, holds under the following weaker condition.

**Assumption 3 (Exponential Moments)** There exist  $\lambda, \tilde{C} > 0$  and  $\gamma \in (0,1)$  such that: for each  $x \in \mathcal{X}$  we have  $\mathbb{E}_{X \sim P(x,\cdot)}[e^{\lambda|X|}] \leqslant \gamma e^{\lambda|x|} + \tilde{C}$ .

Before formulating our concentration result, we remark that Assumption 2 implies 3, but not conversely.

**Proposition 2 (Excess Risk-Bound)** Under Assumptions 1 and 3, let  $f^*: \mathbb{R}^d \to \mathbb{R}^D$ ,  $\ell: \mathbb{R}^{2D} \to \mathbb{R}$ , and R, r > 0 be a such that the composite map  $\mathbb{R}^d \ni x \mapsto \ell(f^*(x), f(x))$  belongs to  $C_R^s(\mathbb{R}^d)$  for all  $f \in C_r^2(\mathbb{R}^d)$ . Then, there exists some  $\kappa \in (0,1)$  depending only on the Markov chain X. and some  $t_0 \in \mathbb{N}_0$  such that for each  $t_0 \leq N < t \leq \infty$ , each "rate-to-constant-tradeoff parameter"  $s \in \mathbb{N}_+$ , and every "confidence level"  $\delta \in (0,1]$  the following

$$\sup_{g \in \mathcal{C}_{R}^{s}(\mathbb{R}^{d})} \frac{\left| \mathcal{R}_{\max\{t,N\}}(g) - \mathcal{R}^{(N)}(g) \right|}{R} \lesssim \kappa^{t} + \frac{\sqrt{2\ln(1/\delta)}}{N^{1/2}} + \begin{cases} \frac{\log\left(c\,N\right)^{d/(2s+1)}}{c\,N^{1/2}} & \text{if } d < 2s \\ \frac{\log\left(c\,N\right)}{c\,N^{1/2}} & \text{if } d = 2s \\ \frac{\log\left(c\,N\right)^{d-2s+(s/d)}}{c\,N^{1/2}} & \text{if } d = 2s \end{cases}$$

holds with probability at least  $1 - \delta$ ; where  $0 < \kappa < 1$ , and we use the notation  $\kappa^{\infty} \stackrel{\text{def}}{=} 0$ .

**Proof** [Proof of Proposition 2] By hypothesis,  $\tilde{f} \in C_r^s(\mathbb{R}^d)$  the induced map

$$f: \mathbb{R}^d \to \mathbb{R}$$

$$x \mapsto f(x) \stackrel{\text{def.}}{=} \ell(f^*(x), \tilde{f}(x))$$
(6)

belongs to  $C_R^s(\mathbb{R}^d)$ .

Conversion to a Concentration of Measure Problem. Denote the empirical (random) measure associated with the samples  $\{(X_n,Y_n)\}_{n=1}^N$  by  $\mu^{(N)} \stackrel{\text{def.}}{=} \frac{1}{N} \sum_{n=1}^N \delta_{(X_n,Y_n)}$ . Note that the generalization bound is 0 for any constant function; therefore, we consider the bound over  $C_R^s(\mathbb{R}^d) \setminus \text{Lip}_0$  where  $\text{Lip}_0$  denotes the set of constant functions from  $\mathbb{R}^d$  to  $\mathbb{R}$ . Note the bijection between  $C_R^s(\mathbb{R}^d) \setminus \text{Lip}_0$  and  $C_1^s(\mathbb{R}^d) \setminus \text{Lip}_0$  given by  $f \mapsto \frac{1}{\max\{1, \|f\|_{C^s(\mathbb{R}^d)}\}} f$ . Therefore, we compute

$$\left| \mathcal{R}_{t}(f) - \mathcal{R}^{(N)}(g) \right| \leq \sup_{g \in \mathcal{C}_{R}^{s}(\mathbb{R}^{d})} \left| \mathcal{R}_{t}(f) - \mathcal{R}^{(N)}(g) \right|$$

$$\leq R \sup_{g \in \mathcal{C}_{1}^{s}(\mathbb{R}^{d})} \left| \mathcal{R}_{t}(g) - \mathcal{R}^{(N)}(g) \right|$$

$$\leq R d_{C^{s}}(\mu_{\max\{t,N\}}, \mu^{(N)})$$

$$\leq R \left( \underbrace{d_{C^{s}}(\mu_{t}, \bar{\mu})}_{(IV)} + \underbrace{d_{C^{s}}(\bar{\mu}, \mu^{(N)})}_{(V)} \right). \tag{7}$$

Next, we bound terms (I) and (II).

**Bounding Term** (IV). Under the log-Sobolev Assumption 1, [17, Theorem 1.3] can be applied to  $\bar{\mu}$  and  $P(x,\cdot)$  for each  $x \in \mathcal{X}$ , implying that the transport inequalities hold: for each  $\nu \in \mathcal{P}(\mathcal{X})$  and each  $\tilde{\mu} \in \{\bar{\mu}, \mu_0\} \cup \{P(x,\cdot)\}_{x \in \mathcal{X}}$ 

$$W_1(\tilde{\mu}, \nu)^2 \leqslant 2C^2 \text{ KL}(\nu|\tilde{\mu}) \tag{8}$$

where we recall the definition of the Kullback–Leibler divergence  $\mathrm{KL}(\nu|\mu) \stackrel{\mathrm{def.}}{=} \mathbb{E}_{X \sim \nu}[\log(\frac{d\nu}{d\mu}(X))]$ . Thus, (8) implies that the following exponential contractility property of the Markov kernel: there exists some  $\kappa \in (0,1)$  such that for each  $x, \tilde{x} \in \mathcal{X}$  and every  $t \in \mathbb{N}_+$ 

$$W_1(P^t(x,\cdot), P^t(\tilde{x},\cdot)) \leqslant \kappa^t \|x - \tilde{x}\|. \tag{9}$$

Furthermore, (9) implies that the conditions for [75, Theorem 1.5] are met; whence, for every  $\varepsilon \geqslant 0$  and each  $N \in \mathbb{N}$  the following holds with probability at-least  $1 - \exp\left(\frac{-N\,\varepsilon^2(1-\kappa)^2}{2C^2}\right)$ 

$$(IV) = d_s(\bar{\mu}, \mu^{(N)}) \leqslant \mathbb{E}\left[d_s(\bar{\mu}, \mu^{(N)})\right] + \varepsilon, \tag{10}$$

for some C>0. Upon setting  $\varepsilon \stackrel{\text{def.}}{=} \frac{C\sqrt{2\ln(1/\delta)}}{\sqrt{N(1-\kappa^2)}}$ , (10) implies that: for every  $N\in\mathbb{N}$  and each  $\delta\in(0,1]$  the following holds with probability at-least  $1-\delta$ 

$$(IV) = d_s(\bar{\mu}, \mu^{(N)}) \leqslant \underbrace{\mathbb{E}\left[d_s(\bar{\mu}, \mu^{(N)})\right]}_{(VI)} + \frac{C\sqrt{2\ln(1/\delta)}}{\sqrt{N(1-\kappa^2)}}.$$
(11)

It remains to bound the expectation term (VI) in (11) to bound term (IV). Under the exponential moment assumption 3, we have that

$$\mathbb{E}_{X \sim P(x,\cdot)}[e^{\beta|X|} - 1] \le \gamma (e^{\beta|x|} - 1) + (C - 1 + \gamma). \tag{12}$$

Therefore [75, Proposition 1.3], implies that  $\sup_{t\in\mathbb{N}_0}\mathbb{E}[e^{\beta|X_t|}-1]<\infty$ . Whence, [75, Assumption 2] holds with Young function  $\Phi(x)=\frac{1}{\max\{1,\sup_{t\in\mathbb{N}_+}\}\mathbb{E}[e^{\beta|X_t-1}]}(e^{\beta|X_t|}-1)$ ; namely,  $\sup_{t\in\mathbb{N}_0}\mathbb{E}[\Phi(|X_t|)]\leqslant 1$ . Consequentially, [75, Theorem 1.1] applies from which we conclude that there is some  $t_0\in\mathbb{N}_+$  such that for all  $N\geqslant t_0$ 

$$(VI) = \mathbb{E}\left[d_{s}(\bar{\mu}, \mu^{(N)})\right] \lesssim \log\left((1-\kappa)N\right)^{s} \begin{cases} \frac{\log\left((1-\kappa)N\right)^{d/(2s)+1}}{(1-\kappa)^{1/2}N^{1/2}} & \text{if } 1 = d < 2s\\ \frac{\log\left((1-\kappa)N\right)}{(1-\kappa)^{1/2}N^{1/2}} & \text{if } d = 2s\\ \frac{\log\left((1-\kappa)N\right)^{d-2s+s/d}}{(1-\kappa)^{s/d}N^{s/d}} & \text{if } d = 2s \end{cases}$$
(13)

Combining the order estimate of (VI) in (13) with the estimate in (11) implies that: for every  $N \ge t_0$  and each  $\delta \in (0, 1]$  we have

$$(IV) = d_s(\bar{\mu}, \mu^{(N)}) \lesssim \frac{\sqrt{2\ln(1/\delta)}}{N^{1/2}} + \begin{cases} \frac{\log(cN)^{d/(2s)+1}}{cN^{1/2}} & \text{if } 1 = d < 2s \\ \frac{\log(cN)}{cN^{1/2}} & \text{if } d = 2s \\ \frac{\log(cN)^{d-2s+s/d}}{cN^{1/2}} & \text{if } d = 2s \end{cases}$$

$$(14)$$

where  $c \stackrel{\text{def.}}{=} (1 - \kappa), c_2 \stackrel{\text{def.}}{=} c^{s/d} \in (0, 1), \text{ and } \lesssim \text{suppresses the absolute constant } \max\{1, C\} > 0.$ 

Bounding Term (V). Next, we bound (V) by computing

$$(V) = d_{C^s}(\mu_t, \bar{\mu}) \stackrel{\text{def.}}{=} \sup_{g \in \mathcal{C}_1^s(\mathbb{R}^d)} \mu_t[g] - \bar{\mu}[g]$$

$$\leqslant \sup_{g \in \text{Lip}_1(\mathbb{R}^d)} \mu_t[g] - \bar{\mu}[g]$$

$$(15)$$

$$=\mathcal{W}_1(\mu_t, \bar{\mu}) \tag{16}$$

$$= \mathcal{W}_1(P^t \mu_0, \bar{\mu}) \tag{17}$$

$$= \mathcal{W}_1(P^t \mu_0, P^t \bar{\mu}) \tag{18}$$

$$\leq \kappa^t \mathcal{W}_1(\mu_0, \bar{\mu}) \stackrel{\text{def.}}{=} \kappa^t C$$
 (19)

where (15) held by definition of the MMD  $d_{C^s}$  and by the inclusion of  $\mathcal{C}_1^s(\mathbb{R}^d) \subset \operatorname{Lip}_1(\mathbb{R}^d)$ , (16) held by Kantorovich duality (see [85, Theorem 5.10]), (18) held since  $\bar{\mu}$  is the stationary probability measure for the Markov chain X., it is invariant to the action of the Markov kernel, and (19) followed from [71, Corollary 21] since we deduced the exponential contractility property (9) of the Markov kernel. Note that  $C \stackrel{\text{def}}{=} \mathcal{W}_1(\mu_0, \bar{\mu})$  is a constant depending only on the initial and stationary distributions of the Markov chain.

**Conclusion.** Incorporating the estimates for (V) and (IV) into the right-hand side of (7) implies that: for every  $t, N \ge t_0$ ,  $s \in \mathbb{N}_+$ , and each  $\delta \in (0, 1]$  the following holds

$$\sup_{g \in \mathcal{C}_{R}^{s}(\mathbb{R}^{d})} \frac{\left| \mathcal{R}_{\max\{t,N\}}(g) - \mathcal{R}^{(N)}(g) \right|}{R} \lesssim I_{t < \infty} \kappa^{t} + \frac{\sqrt{2 \ln(1/\delta)}}{N^{1/2}} + \begin{cases} \frac{\log\left(cN\right)^{d/(2s+1)}}{c N^{1/2}} & \text{if } 1 = d < 2s \\ \frac{\log\left(cN\right)}{c N^{1/2}} & \text{if } d = 2s \\ \frac{\log\left(cN\right)^{d/2s+1}}{c N^{1/2}} & \text{if } d = 2s \end{cases}$$

with probability at-least  $1 - \delta$ ; where  $c \stackrel{\text{def.}}{=} (1 - \kappa)$  and  $\kappa \in (0, 1)$ ; where  $I_{t < \infty} k^{\infty} \stackrel{\text{def.}}{=} 0$  if  $t = \infty$ .

# D.3 Step 2 (A) - Bounding the $C^s$ Regularity of Transformer Building Blocks

We begin by the following simple remark, that if the activation function used to defined the transformer is smooth, then so must the entire transformer model.

Proposition 3 (Transformers with Smooth Activation Functions are Smooth) Define constants D, d, M, L, H, C, as in Definition 4, then every transformer  $T \in \mathcal{TC}_{L,H,C}^{D,d,M}$  is smooth.

**Proof** [Theorem 3] The smoothness of Att follows directly from the smoothness of softmax, which immediately implies smoothness of  $\mathcal{MH}$  since the operators used for its definition are smooth. Furthermore, the  $\mathcal{LN}$  is smooth due to its smooth and lower-bounded denominator and the activation function  $\sigma$  is smooth by definition, therefore we conclude that  $\mathcal{TB} \in \mathcal{TB}_{d',M',H'}$  is smooth for every  $d',M',H' \in \mathbb{N}$  and we obtain smoothness of  $\mathcal{T} \in \mathcal{T}_{L,W,H}^{D,d,M}$  as a consequence.

#### D.3.1 The Softmax Function

Lemma 3 (Representation of higher-order softmax derivatives) For  $F \in \mathbb{N}$  and

smax: 
$$\mathbb{R}^F \to \mathbb{R}^F$$
,  $x \mapsto \left(\exp(x_i)/\sum_{j=0}^{F-1} \exp(x_j)\right)_{i=1}^F$ .

there exists for any multi-index  $\alpha \in \mathbb{N}^F$  and  $m \in \{1, \dots, F\}$  indicators  $(a_{i,j}^k)_{i,j \in I(\alpha)}^{k \in \{1, \dots, |\alpha|!\}} \subseteq \{0,1\}$  such that

$$smax^{(\alpha)}(x_m) = \sum_{k=1}^{|\alpha|!} smax(x_m) \prod_{i,j \in I(\alpha)} (a_{i,j}^k - smax(x_j)),$$
(20)

where  $I(\alpha) \stackrel{\text{def.}}{=} \{(i,j) : i = 1, \dots, F, j = 1, \dots, \alpha_i\}.$ 

**Proof** For  $|\alpha| = 0$ , we have  $n \in \{1, \dots, F\}$  s.t.  $\alpha_n = 1$ , therfore

$$\operatorname{smax}^{(\alpha)}(x_m) = \frac{\partial \operatorname{smax}}{\partial x_n}(x_m) = \operatorname{smax}(x_m) \left(\delta_{mn} - \operatorname{smax}(x_n)\right),$$

which is of the form (20). Now, let  $\alpha \in \mathbb{N}^F$  arbitrary, therefore, by defining  $\alpha' \in \mathbb{N}^F$  by  $\alpha'_i \stackrel{\text{def.}}{=} \alpha_i$  for  $i \neq n$  and  $\alpha'_n \stackrel{\text{def.}}{=} \alpha_n - 1$  (w.l.o.g.  $\alpha_n > 0$ ). We have

$$\operatorname{smax}^{(\alpha)}(x_m) = \frac{\partial \operatorname{smax}^{(\alpha')}}{\partial x_n}(x_m)$$
$$= \frac{\partial}{\partial x_n} \sum_{k=1}^{|\alpha'|!} \operatorname{smax}(x_m) \prod_{i,j \in I(\alpha')} (a'^k_{i,j} - \operatorname{smax}(x_j)).$$

Since for any k

$$\begin{split} &\frac{\partial}{\partial x_n} \operatorname{smax}(x_m) \prod_{i,j \in I(\alpha')} (a_{i,j}'^k - \operatorname{smax}(x_j)) \\ &= \operatorname{smax}(x_m) \left( \delta_{mn} - \operatorname{smax}(x_n) \right) \prod_{i,j \in I(\alpha')} (a_{i,j}'^k - \operatorname{smax}(x_j)) \\ &+ \operatorname{smax}(x_m) \sum_{i',j' \in I(\alpha')} - \operatorname{smax}(x_{j'}) \left( \delta_{j',n} - \operatorname{smax}(x_n) \right) \prod_{\substack{i,j \in I(\alpha) \\ (i,j) \neq (i',j')}} (a_{i,j}'^k - \operatorname{smax}(x_j)), \end{split}$$

we can define  $(a_{i,j}^k)_{i,j\in I(\alpha)}^{k\in\{1,\dots,|\alpha'|+1\}}\subseteq\{0,1\}$  such that

$$\frac{\partial}{\partial x_n} \operatorname{smax}(x_m) \prod_{i,j \in I(\alpha')} (a'_{i,j}^k - \operatorname{smax}(x_j)) = \sum_{k=1}^{|\alpha|} \operatorname{smax}(x_m) \prod_{i,j \in I(\alpha)} (a_{i,j}^k - \operatorname{smax}(x_j)).$$

Since  $|\alpha|! = |\alpha| \cdot |\alpha'|!$ , this concludes the proof.

**Lemma 4 (Bound of higher-order** softmax **derivatives)** With Notation 3, it holds for any set  $K \in \mathbb{R}^k, k \in \mathbb{N}$  and any  $\alpha \in \mathfrak{O}^k_{<\infty}$  that

$$C^{\rm smax}(\alpha) \leqslant |\alpha|!$$

**Proof** This is a direct consequence of the representation in Lemma 3 together with  $\|\operatorname{smax}\| = 1$ .

#### D.3.2 THE MULTI-HEAD SELF-ATTENTION MECHANISM

**Lemma 5 (Bound of Dot product)** In the notation of Definition 2 and for  $m \in \{1, ..., M\}$ 

$$\mathrm{dp}_m(\cdot;Q,K):\mathbb{R}^{Md_{\mathrm{in}}}\longrightarrow\mathbb{R}^M,\quad x\longmapsto\langle Qx_m,Kx_i\rangle_{i=0}^M$$

we have using Notation 3

- 1.  $C_K^{\mathrm{dp}_m}(e_1) \leq 2d_{\mathrm{in}}d_K \|K\|C^Q C^K$ , where  $C^Q \stackrel{\mathrm{def.}}{=} \max_{i,i' \in \{1,...,d_K\} \times \{1,...,d_{\mathrm{in}}\}} |Q_{i,i'}|$ ,  $C^K$  analogously, and  $\|K\| \stackrel{\mathrm{def.}}{=} \max_{x \in K} \|x\|$ . Additionally,
- 2.  $C_K^{\mathrm{dp}_m}(\alpha) \leqslant 2d_K C^Q C^K$ , for  $|\alpha| = 2$ , and
- 3.  $C_K^{dp_m}(\alpha) = 0 \text{ for } |\alpha| > 2.$

Since all bounds are not dependent on m we write  $C^{dp}$  short for  $C^{dp_m}$ .

# Proof

1. Let  $l = (l_1, l_2) \in \{1, \dots, M\} \times \{1, \dots, d_{in}\}$ . Assume  $l_1 = m$ . If  $j \neq m$ , then

$$D^{e_l} \operatorname{dp}_m(x; Q, K)_j = D^{e_l} \sum_{i=1}^{d_K} (Kx_j)_i \sum_{i'=1}^{d_{\text{in}}} Q_{i,i'}(x_m)_{i'} = \sum_{i=1}^{d_K} \left( \sum_{i'=1}^{d_{\text{in}}} K_{i,i'}(x_j)_{i'} \right) Q_{i,l_2},$$

implying

$$||D^{e_l} \operatorname{dp}_m(x; Q, K)|| \le ||K|| \sum_{i=1}^{d_K} Q_{i, l_2} \sum_{i'=1}^{d_{\text{in}}} K_{i, i'} \le d_{\text{in}} d_K ||K|| C^Q C^K.$$
(21)

If j = m,

$$D^{e_l} \operatorname{dp}_m(x; Q, K)_j = D^{e_l} \sum_{i=1}^{d_K} \left( \sum_{i'=1}^{d_{\text{in}}} K_{i,i'}(x_m)_{i'} \right) \left( \sum_{i'=1}^{d_{\text{in}}} Q_{i,i'}(x_m)_{i'} \right)$$

$$= \sum_{i=1}^{d_K} \left( K_{i,l_2} \sum_{i'=1}^{d_{\text{in}}} Q_{i,i'}(x_m)_{i'} + Q_{i,l_2} \sum_{i'=1}^{d_{\text{in}}} K_{i,i'}(x_m)_{i'} \right)$$

therefore implying

$$||D^{e_l} \operatorname{dp}_m(x; Q, K)|| \le 2d_{\operatorname{in}} d_K ||K|| C^Q C^K.$$

If  $l_1 \neq m$  then for  $j \neq l_1$ ,  $D^{e_l} \operatorname{dp}_m(x; Q, K)_j = 0$ , for  $j = l_1$ 

$$D^{e_l} \operatorname{dp}_m(x; Q, K)_j = D^{e_l} \sum_{i=1}^{d_K} (Qx_m)_i \sum_{i'=1}^{d_{\text{in}}} K_{i,i'}(x_j)_{i'} = \sum_{i=1}^{d_K} \left( \sum_{i'=1}^{d_{\text{in}}} Q_{i,i'}(x_m)_{i'} \right) K_{i,l_2},$$

and we obtain (21) analogously.

2. If  $l_1 = m$  and  $j \neq m$ 

$$D^{e_l}\left((x_m)_{l_2} \sum_{i=1}^{d_K} \left(\sum_{i'=1}^{d_{\text{in}}} k_{i,i'}(x_j)_{i'}\right) q_{i,l_2}\right) = 0,$$

implying  $||D^{2e_l} dp_m(x; Q, K)|| \leq 0$ , what analogously holds for  $l_1 \neq m$ . However, for  $l_1 = m$  and j = m

$$D^{e_l}\left(\sum_{i=1}^{d_K} K_{i,l_2} \sum_{i'=1}^{d_{\text{in}}} Q_{i,i'}(x_m)_{i'} + Q_{i,l_2} \sum_{i'=1}^{d_{\text{in}}} K_{i,i'}(x_m)_{i'}\right) = \sum_{i=1}^{d_K} K_{i,l_2} Q_{i,l_2} + Q_{i,l_2} K_{i,l_2}$$

we have

$$||D^{2e_l} \operatorname{dp}_m(x; Q, K)|| \leqslant 2d_K C^Q C^K.$$

3. Let  $l' = (l'_1, l'_2) \in \{1, \dots, M\} \times \{1, \dots, d_{\text{in}}\}$ . Assume  $l_1 = m, j \neq m$ . If  $l'_1 \neq j$ , it follows that  $D^{e_l + e_{l'}} \operatorname{dp}_m(x; Q, K)_j = 0$ . For  $l'_1 = j$  follows  $D^{e_l + e_{l'}} \operatorname{dp}_m(x; Q, K)_j = \sum_{i=1}^{d_K} K_{i, l'_2} Q_{i, l_2}$ . If  $l_1 = m$ ,  $j \neq m$ , we have  $D^{e_l + e_{l'}} \operatorname{dp}_m(x; Q, K)_j = 0$  in the case that  $l'_1 \neq m$ , and for  $l'_1 \neq m$  we obtain

$$D^{e_l + e_{l'}} \operatorname{dp}_m(x; Q, K)_j = \sum_{i=1}^{d_K} K_{i, l_2} Q_{i, l'_2} + Q_{i, l_2} K_{i, l'_2}.$$

This means, we can use the bound

$$||D^{e_l+e_{l'}} \operatorname{dp}_m(x; Q, K)|| \leq 2d_K C^Q C^K.$$

**Lemma 6 (Bound of Self-Attention for Derivative Type)** Using the notation of Notation 3, Definition 2 and Lemma 5, it holds that

$$C_K^{\text{Att}}(\alpha) \leqslant d_{\text{in}} M C^V \left( \|K\| C_K^{\text{smax} \circ \text{dp}}(\alpha) + \sum_{l=1}^{M d_{\text{in}}} \alpha_l C_K^{\text{smax} \circ \text{dp}}(\alpha - e_l) \right)$$

where

$$C_K^{\operatorname{smax} \circ \operatorname{dp}}(\alpha) \leqslant \alpha! \sum_{\beta \in \mathfrak{D}_{\leqslant n}^M} N(\beta) C_{\operatorname{dp}[K]}^{\operatorname{smax}}(\beta) \sum_{\eta, \zeta \in \mathcal{P}'(\alpha, \beta)} \prod_{j=1}^n \frac{C_K^{\operatorname{dp}}(\mathfrak{o}(\zeta^{(j)}))^{|\eta^{(j)}|}}{\eta^{(j)}! (\zeta^{(j)}!)^{|\eta^{(j)}|}}. \tag{22}$$

**Proof** Fix  $\alpha \in \mathbb{N}^k$ , and note that

$$||D^{\alpha} \operatorname{Att}(x; Q, K, V)|| \le \max_{m \in \{1, \dots, M\}} \max_{i \in \{0, \dots, d_V\}} ||D^{\alpha} \operatorname{Att}(x; Q, K, V)_{m,i}||$$

and

$$\begin{split} \|D^{\alpha} \operatorname{Att}(x; Q, K, V)_{m,i}\| \\ &\leqslant \sum_{j=1}^{M} \sum_{i'=0}^{d_{\text{in}}} \|D^{\alpha} \operatorname{smax} \circ \operatorname{dp}(x; Q, K)_{j} V_{i,i'}(x_{j})_{i'}\| \\ &\leqslant d_{\text{in}} M \max_{j \in \{1, \dots, M\}} \max_{i' \in \{0, \dots, d_{\text{in}}\}} \|D^{\alpha} \operatorname{smax} \circ \operatorname{dp}(x; Q, K)_{j} V_{i,i'}(x_{j})_{i'}\|. \end{split}$$

Due to the extended Leibnitz rule [46], we have

$$||D^{\alpha}\operatorname{smax} \circ \operatorname{dp}(x; Q, K)_{i}V_{i,i'}(x_{i})_{i'}||$$

$$\leqslant \|D^{\alpha}\operatorname{smax} \circ \operatorname{dp}(x; Q, K)_{j} V_{i, i'}(x_{j})_{i'}\| + \sum_{l=1}^{Md_{\operatorname{in}}} V_{i, i'} \alpha_{l} \|D^{\alpha - e_{l}} \operatorname{smax} \circ \operatorname{dp}(x; Q, K)_{j}\|.$$

Equation (22) follows directly from Theorem 4.

Corollary 3 (Bound of Self-Attention for Derivative Level) Using the setting of Lemma 6, for  $n \in \mathbb{N}$ ,

$$C_K^{\text{Att}}(n) \leqslant d_{\text{in}} M C^V C_K^{\text{smax} \circ \text{dp}}(\leqslant n) \left( \|K\| + n d_{\text{in}} M \right)$$
(23)

where

$$C_K^{\operatorname{smax} \circ \operatorname{dp}}(\leqslant n) \leqslant C_{\operatorname{dp}[K]}^{\operatorname{smax}}(\leqslant n) C_K^{\operatorname{dp}}(\leqslant n)^n \left[ \frac{2nM}{e \ln n} (1 + o(1)) \right]^n. \tag{24}$$

**Proof** Equation (23) follows directly from Lemma 6; and (24) is a consequence of Corollary 1. ■

Corollary 4 (Bound of Multi-head Self-Attention) In the notation of Definition 2, Theorem 4 and Lemma 5 it holds that

$$C_K^{\mathcal{MH}}(\alpha) \leqslant \alpha! d_V C^W C_K^{\text{Att}}(\alpha)$$

where

$$C_K^{\text{Att def.}} \stackrel{\text{def.}}{=} \max_{h \in \{1, \dots, H\}} C_K^{\text{Att}(\;\cdot\;;Q^{(h)},K^{(h)},V^{(h)})}, \quad C^W \stackrel{\text{def.}}{=} \max_{h \in \{1, \dots, H\}} W^{(h)}.$$

In particular, we have the following order estimate

$$C_K^{\mathcal{MH}}(\leqslant n) \in \mathcal{O}\left(M^2 \|K\| \|W\| \|V\| (c_{d_{\text{in}}, d_K} \|K\| \|Q\| \|K\|)^n \ n^2 \left(\frac{n}{e}\right)^{2n} C_n^n\right).$$

**Proof** From Corollary 2 and Lemma 6 we directly obtain

$$C_K^{\mathcal{MH}}(\alpha) \leqslant n! d_V C^W d_{\text{in}} M C^V n! (2d_{\text{in}} d_K || K || C^Q C^K)^n$$

$$\times (||K|| + nd_{\text{in}} M) \left[ \frac{2nM}{e \ln n} (1 + o(1)) \right]^n.$$

$$(25)$$

Applying Stirling's approximation, we have that

$$C_K^{\mathcal{MH}}(\alpha) \in \mathcal{O}\left(M^2 \|K\| \|W\| \|V\| (c_{d_{\text{in}}, d_K} \|K\| \|Q\| \|K\|)^n \ n^2 \left(\frac{n}{e}\right)^{2n} C_n^n\right), \tag{26}$$

where 
$$C_n \stackrel{\text{def.}}{=} \frac{2nM}{e \ln n} (1 + o(1))$$
 and  $c_{d_{\text{in}}, d_K} \stackrel{\text{def.}}{=} 2d_{\text{in}} d_K$ .

#### D.3.3 The Activation Functions

# Lemma 7 (Derivatives of splus) For

splus: 
$$\mathbb{R} \to \mathbb{R}$$
,  $x \mapsto \ln(1 + \exp(\cdot))$ 

it holds

$$splus^{(1)}(x) = sig(x) \stackrel{\text{def.}}{=} 1/(1 + exp(-x))$$

and for  $n \in \mathbb{N}$ 

$$splus^{(n+1)}(x) = sig^{(n)}(x) = \sum_{k=0}^{n} (-1)^{n+k} k! S_{n,k} sig(x) (1 - sig(x))^{k},$$

where  $S_{n,k}$  are the Stirling numbers of the second kind,  $S_{n,k} \stackrel{\text{def.}}{=} \frac{1}{k!} \sum_{j=0}^{k} (-1)^{k-j} {k \choose j} j^n$ .

Proof We start with Faà di Bruno's formula,

$$\frac{d^n}{dt^n}\operatorname{sig}(x) = \frac{d^n}{dx^n}\frac{1}{f(x)} = \sum_{k=0}^n (-1)^k k! f^{-(k+1)}(x) B_{n,k}(f(x)),$$

where  $f(x) \stackrel{\text{def.}}{=} 1 + \exp(-x)$  and  $B_{n,k}(f(x))$  denotes the Bell polynomials evaluated on f(x). Next, we know the k-th derivative of f(x) is given by

$$\frac{d^k}{dt^k}f(x) = (1-k)_k + ke^{-x}.$$

Now, using the definition of the Bell polynomials  $B_{n,k}(f(t))$ , we have

$$B_{n,k}(f(x)) = (-1)^n S_{n,k} e^{-kx},$$

where  $S_{n,k}$  represents the Stirling numbers of the second kind. Substituting the expression for  $B_{n,k}(f(x))$  into the derivative of sig(x), we obtain

$$\frac{d^n}{dx^n} \operatorname{sig}(x) = \sum_{k=0}^n (-1)^{n+k} k! S_{n,k} \operatorname{sig}(x) (1 - \operatorname{sig}(x))^k.$$

**Corollary 5** In the setting of Lemma 7, for  $n \in \mathbb{N}$ ,

$$C^{\text{splus}}(n) \leqslant \sum_{k=0}^{n} \frac{k^{k} k! S_{n,k}}{(k+1)^{k+1}}.$$

**Proof** For  $k \in \mathbb{N}$  and  $x \in [0,1]$ , we have

$$f^k(x) \stackrel{\text{def.}}{=} x(1-x)^k, \quad (f^k)'(x) = (1-(k+1)x)(1-x)^{k-1};$$

which amounts to  $(f^k)'(x) = 0$  at 1/(k+1), i.e.

$$\max_{x \in [0,1]} f(x) = \frac{1}{k+1} \left( \frac{k}{k+1} \right)^k = \frac{k^k}{(k+1)^{k+1}}.$$

# Lemma 8 (Derivatives of GELU) For

$$GELU : \mathbb{R} \to \mathbb{R}, \quad x \mapsto x\Phi(x),$$

it holds

$$GELU'(x) = \Phi(x) + x\varphi(x),$$

$$GELU^{(n)}(x) = n\varphi^{(n-2)}(x) + x\varphi^{(n-1)}(x), \quad n \ge 2,$$
(27)

with  $\varphi(x) = \frac{1}{\sqrt{2\pi}}e^{-x^2/2}$ ,  $\Phi(x) = \int_{-\infty}^{x} \varphi(u)du$ . The n-th derivative of  $\varphi(x)$  is given by

$$\frac{d^n}{dx^n}\varphi(x) = \frac{1}{\sqrt{2\pi}}e^{-x^2/2} \Big[ \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} 2^k \frac{\Gamma(\frac{2k+1}{2})}{\Gamma(\frac{1}{2})} x^{n-2k} \Big].$$

**Proof** By induction we can show that (27) holds. And the representation of the *n*-th derivative of  $\varphi(x)$  follows from [29].

**Corollary 6** For  $n \ge 2$ , it holds in the setting of Lemma 8

$$C^{\text{GELU}}(n) \leqslant \frac{1}{\sqrt{2\pi}\Gamma(\frac{1}{2})} \left( na_{n-2}b_{n-2} + a_{n-1}c_{n-1} \right),$$

where

$$a_n \stackrel{\text{\tiny def.}}{=} \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} 2^k \Gamma\left(\frac{2k+1}{2}\right), \quad b_n \stackrel{\text{\tiny def.}}{=} \max_{x \in \mathbb{R}} e^{-\frac{x^2}{2}} \sum_{k=0}^{\lfloor n/2 \rfloor} x^{n-k}, \quad c_n \stackrel{\text{\tiny def.}}{=} \max_{x \in \mathbb{R}} e^{-\frac{x^2}{2}} x \sum_{k=0}^{\lfloor n/2 \rfloor} x^{n-k}.$$

**Lemma 9 (Derivatives of** tanh) For tanh :  $\mathbb{R} \to \mathbb{R}$ , the n-th derivatives have the representation

$$\frac{d^n}{dx^n}\tanh x = C_n(\tanh x),$$

$$C_n(z) = (-2)^n (z+1) \sum_{k=0}^n \frac{k!}{2^k} \binom{n}{k} (z-1)^k, \quad n \geqslant 1.$$

Proof See [20].

Corollary 7 In the setting of Lemma 9, for  $n \in \mathbb{N}$ ,  $C^{\tanh}(n) = \max_{z \in [-1,1]} C_n(z)$ .

Lemma 10 (Derivatives of SWISH) For

SWISH: 
$$\mathbb{R} \to \mathbb{R}$$
,  $x \mapsto \frac{x}{1 + e^{-x}}$ 

it holds for  $n \ge 1$ 

$$\frac{d^n}{dx^n} \text{SWISH}(x) = n \sum_{k=1}^n (-1)^{k-1} (k-1)! S_{n,k} \operatorname{sig}^k(x) + x \sum_{k=1}^{n+1} (-1)^{k-1} (k-1)! S_{n+1,k} \operatorname{sig}^k(x),$$

where  $S_{n,k}$  are Stirling numbers of the second kind, i.e.,

$$S_{n,k} = \frac{1}{k!} \sum_{j=0}^{k} (-1)^j \binom{k}{j} (k-j)^n.$$

**Proof** By induction, we can show that

$$\frac{d^n}{dx^n} SWISH(x) = n \operatorname{sig}^{(n-1)}(x) + x \operatorname{sig}^{(n)}(x), \quad n \geqslant 1.$$

By [65, Theorem 2], the derivatives of the sigmod function can be represented as

$$\operatorname{sig}^{(n)}(x) = \sum_{k=1}^{n+1} (-1)^{k-1} (k-1)! S_{n+1,k} \operatorname{sig}^k(x), \ n \geqslant 1.$$

Combining the above two equations, we obtain the general form of the n-th derivative of the SWISH function.

#### D.3.4 The Layer Norm

**Lemma 11 (Bound of the Layer Norm for Derivative Type)** Fix  $k \in \mathbb{N}$ ,  $\beta \in \mathbb{R}^k$ ,  $\gamma \in \mathbb{R}$ , and  $w \in [0,1]$ . For the layer norm, given by

$$\mathcal{LN}: \mathbb{R}^k \to \mathbb{R}^k, \quad x \mapsto \gamma f(x)g \circ \Sigma(x) + \beta;$$

$$f: \mathbb{R}^k \to \mathbb{R}^k, \quad x \mapsto x - M(x); \qquad g: \mathbb{R} \to \mathbb{R}, \quad u \mapsto \frac{1}{\sqrt{1+u}};$$

$$M: \mathbb{R}^k \to \mathbb{R}, \quad x \mapsto \frac{w}{k} \sum_{i=1}^k x_i; \qquad \Sigma: \mathbb{R}^k \to \mathbb{R}, \quad x \mapsto \frac{w}{k} \sum_{i=1}^k (x_i - M(x))^2;$$

holds for a compact symmetric set K (using Notation 3)

$$\begin{split} C_K^{\mathcal{LN}}(\alpha) \leqslant \alpha! \gamma \sum_{m=1}^{n=1} \frac{(2m+1)!!}{2^{2m}} \bigg( \sum_{\substack{\alpha' \leqslant \alpha \\ |\alpha'| = n-1}} \sum_{\eta, \zeta \in \mathcal{P}'(\alpha', m)} \prod_{j=1}^n \frac{C_K^{\Sigma}(\mathfrak{o}(\zeta^{(j)}))^{|\eta^{(j)}|}}{\eta^{(j)}! (\zeta^{(j)}!)^{|\eta^{(j)}|}} \\ &+ \sum_{\eta, \zeta \in \mathcal{P}'(\alpha, m)} \prod_{j=1}^n \frac{C_K^{\Sigma}(\mathfrak{o}(\zeta^{(j)}))^{|\eta^{(j)}|}}{\eta^{(j)}! (\zeta^{(j)}!)^{|\eta^{(j)}|}} \bigg), \end{split}$$

where  $C_K^{\Sigma}(\alpha) = 2w\|K\|$  for  $|\alpha| = 1$ ,  $C_K^{\Sigma}(\alpha) = 2w$  for  $|\alpha| = 2$ , and  $C_K^{\Sigma}(\alpha) = 0$  otherwise.

**Proof** Note that

$$g^{(n)}(x) = (-1)^n \frac{(2n+1)!}{n!2^{2n}} (1+x)^{-\frac{1}{2}-n},$$

implying  $C_K^g(n) \leq (2n+1)!!2^{-2n}$ , !! denoting the double factorial. We have further  $C_K^f(\alpha) \leq \mathbbm{1}_{|\alpha|=1}$  and a direct computation yields

$$C_K^{\Sigma}(\alpha) \leqslant \begin{cases} 2w||K|| & \text{for } |\alpha| = 1\\ 2w & \text{for } |\alpha| = 2\\ 0 & \text{else.} \end{cases}$$

By Theorem 4,

$$C_K^{g \circ \Sigma}(\alpha) \leqslant \alpha! \sum_{m=1}^n C_{\Sigma[K]}^g(m) \sum_{\eta, \zeta \in \mathcal{P}'(\alpha, m)} \prod_{j=1}^n \frac{C_K^{\Sigma}(\mathfrak{o}(\zeta^{(j)}))^{|\eta^{(j)}|}}{\eta^{(j)}! (\zeta^{(j)}!)^{|\eta^{(j)}|}}.$$

According to the general multivariate Leibnitz rule, it holds that

$$D^{\alpha}(f \cdot (g \circ \Sigma)) = \sum_{\beta \leqslant \alpha} \frac{\alpha!}{\beta!(\alpha - \beta)!} D^{\beta} f \cdot D^{\alpha - \beta}(g \circ \Sigma)$$

which implies

$$C_K^{\mathcal{LN}}(\alpha) \leqslant C_K^{g \circ \Sigma}(\alpha) \|K\| + \sum_{\beta \leqslant \alpha, |\beta| = 1} \frac{\alpha!}{(\alpha - \beta)!} C_K^{g \circ \Sigma}(\alpha - \beta).$$

Corollary 8 (Bound of the Layer Norm for Derivative Level) In the setting of Lemma 11, it holds that

$$C_K^{\mathcal{LN}}(\leqslant n) \leqslant 2w \|K\| (2n+1)!! 2^{-2n} (\|K\| + kn) \left[ \frac{2n}{e \ln n} (1 + o(1)) \right]^n.$$

Furthermore, we have the asymptotic estimate

$$C_K^{g \circ \Sigma}(\leqslant n) \in \mathcal{O}\left(w \|K\| n^{1/2} \left(\frac{n^{5/2}}{e^{3/4} \ln(n)} (1 + o(1))\right)^n\right).$$

**Proof** Analogue to the proof of Lemma 11,

$$C_K^{\mathcal{L}\mathcal{N}}(\leqslant n) \leqslant \|K\|C_K^{g \circ \Sigma}(\leqslant n) + knC_K^{g \circ \Sigma}(\leqslant n-1) \leqslant (\|K\| + kn)C_K^{g \circ \Sigma}(\leqslant n),$$

where we can use Corollary 1 to bound

$$C_K^{g \circ \Sigma}(\leqslant n) \leqslant C_{\Sigma[K]}^g(\leqslant n) C_K^{\Sigma}(\leqslant n)^n \left[ \frac{2n}{e \ln n} (1 + o(1)) \right]^n$$

$$\leqslant 2w \|K\| (2n + 1)!! 2^{-2n} \left[ \frac{2n}{e \ln n} (1 + o(1)) \right]^n.$$
(28)

Since 2n + 1 is odd, for each  $n \in \mathbb{N}_+$ , then sterling approximation for double factorial yields the asymptotic

$$(2n+1)!! \in \mathcal{O}\left(\sqrt{2n}\left(\frac{n}{e}\right)^{n/2}\right). \tag{29}$$

Merging (29) with the right-hand side of (28) yields

$$C_K^{g \circ \Sigma}(\leqslant n) \in \mathcal{O}\left(w \|K\| n^{1/2} \left(\frac{n^{5/2}}{e^{3/4} \ln(n)} (1 + o(1))\right)^n\right).$$

D.3.5 THE MULTILAYER PERCEPTRON (FEEDFORWARD NEURAL NETWORK) WITH SKIP CONNECTION

Definition 6 (Single-Layer Feedforward Neural Network with Skip Connection) Fix any non-affine activation function  $\sigma \in C^{\infty}(\mathbb{R})$  and dimensions  $d_{\rm in}, d_{\rm ff}, d_{\rm out} \in \mathbb{N}$ . A feedforward neural network is a map  $\mathcal{PL}: \mathbb{R}^{d_{\rm in}} \to \mathbb{R}^{d_{\rm out}}$  represented for each  $x \in \mathbb{R}^{d_{\rm in}}$  by

$$\mathcal{PL}(x) \stackrel{\text{def.}}{=} B^{(1)}x + B^{(2)} \left(\sigma \bullet (Ax + a)\right) \tag{30}$$

for  $A \in \mathbb{R}^{d_{\text{ff}} \times d_{\text{in}}}$ ,  $a \in \mathbb{R}^{d_{\text{ff}}}$ ,  $B^{(1)} \in \mathbb{R}^{d_{\text{out}} \times d_{\text{in}}}$ , and  $B^{(2)} \in \mathbb{R}^{d_{\text{out}} \times d_{\text{ff}}}$ .

Lemma 12 (Bound of Neural Networks for Derivative Type) In the notation of Notation 3, Lemma 5, and Definition 6, it holds that

$$C_K^{\mathcal{PL}}(\alpha) \leqslant C^{B^{(1)}} \mathbb{1}_{|\alpha|=1} + d_{\mathrm{ff}}(\alpha!)^2 C^{B^{(2)}} \sum_{m=1}^n C_{h[K]}^{\sigma}(m) \cdot (C^A)^m \sum_{\eta, \zeta \in \mathcal{P}'(\alpha, m)} \prod_{j=1}^n \frac{\mathbb{1}_{|\zeta^{(j)}| \leqslant 1}}{\eta^{(j)}!},$$

where h[K] is defined as the image of  $h(x) \stackrel{\text{def.}}{=} Ax + a$  on K.

**Proof** Write  $\mathcal{PL}(x) = B^{(1)}x + B^{(2)}((g_i(x))_{i=1}^{d_{\mathrm{ff}}})$ , where for  $i \in \{1, \dots, d_{\mathrm{ff}}\}$   $g_i(x) \stackrel{\text{def}}{=} \sigma((Ax + a)_i).$ 

If we define  $h_i(x) \stackrel{\text{def.}}{=} h(x)_i$ , we follow with Theorem 4

$$C_K^{g_i}(\alpha) \leqslant \alpha! \sum_{m=1}^n C_{h[K]}^{\sigma}(m) \cdot (C^A)^m \sum_{\eta, \zeta \in \mathcal{P}'(\alpha, m)} \prod_{j=1}^n \frac{\mathbb{1}_{|\zeta^{(j)}| \leqslant 1}}{\eta^{(j)!}} \stackrel{\text{def.}}{=} C_K^g(\alpha),$$

and due to the component wise application of the activation function it holds that

$$||D^{\alpha} \max_{i \in \{1, \dots, d_{\mathrm{ff}}\}} g_i(x)||_K = \max_{i \in \{1, \dots, d_{\mathrm{ff}}\}} C_K^{g_i}(\alpha) \leqslant C_K^g(\alpha).$$

Using Corollary 2, we obtain

$$C_K^{\mathcal{PL}}(\alpha) \leqslant C^{B^{(1)}} \mathbb{1}_{|\alpha|=1} + d_{\text{ff}} \alpha! C^{B^{(2)}} C_K^g(\alpha).$$

Corollary 9 (Bound of Neural Networks for Derivative Level) In the setting of Lemma 12,

$$C_K^{\mathcal{PL}}(\leqslant n) \leqslant C^{B^{(1)}} + d_{\mathrm{ff}} n! C^{B^{(2)}} C_{h[K]}^{\sigma}(\leqslant n) (C^A)^n \Big[ \frac{2n}{e \ln n} (1 + o(1)) \Big]^n.$$

If, moreover,  $K = [-M_1, M_2]^{d_{in}}$  then

$$C_K^{\mathcal{PL}}(\leqslant n) \in \mathcal{O}\Big(\|B^{(1)}\|_{\infty} + \|B^{(2)}\|_{\infty}\|A\|_{\infty}^n \|\sigma\|_{n: \mathrm{Ball}(a, \sqrt{d_{\mathrm{in}}|M_1 + M_2|})} \operatorname{Width}(\mathcal{PL}) \, n^{1/2} \big(\frac{n}{e}\big)^n C_n^n \Big)$$

**Proof** Arguing analogously to the proof of Lemma 12, barring the usage of Corollary 1, we obtain the estimate

$$C_K^{\mathcal{PL}}(\leqslant n) \leqslant C^{B^{(1)}} + d_{\mathrm{ff}} n! C^{B^{(2)}} C_{h[K]}^{\sigma}(\leqslant n) (C^A)^n \left[ \frac{2n}{e \ln n} (1 + o(1)) \right]^n. \tag{31}$$

Let  $C_n \stackrel{\text{def.}}{=} \frac{2n}{e \ln n} (1 + o(1))$  Using Stirling's approximation and the definition of the component-wise  $\|\cdot\|_{\infty}$  norm of a matrix, (31) becomes

$$C_K^{\mathcal{PL}}(\leqslant n) \in \mathcal{O}\Big(\|B^{(1)}\|_{\infty} + \|B^{(2)}\|_{\infty}\|A\|_{\infty}^n C_{h[K]}^{\sigma}(\leqslant n) d_{\mathrm{ff}} \, n^{1/2} \left(\frac{n}{e}\right)^n C_n^n\Big). \tag{32}$$

If, there is some  $M_1, M_2 \leq 0$ , such that  $K = [0, \beta]^d$  then using the estimate between the  $\|\cdot\|_2$  and  $\|\cdot\|_{\infty}$  norms on  $\mathbb{R}^{d_{\text{in}}}$  and the linearity of A we estimate

$$C_{h[K]}^{\sigma}(\leqslant n) \leqslant C_{\mathrm{Ball}(a,\sqrt{d_{\mathrm{in}}|M_1+M_2|})}^{\sigma}(\leqslant n) \leqslant \|\sigma\|_{n:\mathrm{Ball}(a,\sqrt{d_{\mathrm{in}}|M_1+M_2|})}.$$

Upon Width( $\mathcal{PL}$ )  $\stackrel{\text{def.}}{=}$  max $\{d_{\text{in}}, d_{\text{out}}, d_{\text{ff}}\}$ , the estimate (31) implies that  $C_K^{\mathcal{PL}}(\leqslant n)$  is of the order of

$$\mathcal{O}\Big(\|B^{(1)}\|_{\infty} + \|B^{(2)}\|_{\infty}\|A\|_{\infty}^{n}\|\sigma\|_{n:[-\|a\|_{\infty} - \sqrt{d_{\text{in}}|M_{1} + M_{2}|}, \|a\|_{\infty} + \sqrt{d_{\text{in}}|M_{1} + M_{2}|}]} \times \text{Width}(\mathcal{PL}) n^{1/2} \left(\frac{n}{e}\right)^{n} C_{n}^{n}\Big).$$
(33)

# D.4 Step 2 (B) - Transformers

We may now merge the computations in Subsection D.3, with the Fa'a di Bruno-type from Section D.1 to uniformly bound the  $C^s$ -norms of the relevant class transformer networks. Our results are derived in two verions: the first is of "derivative type" (which is much smaller and more precise but consequentially more complicated) and the second is in "derivative level" form (cruder but simpler but also looser).

**Theorem 5 (By Derivative Type)** Let K be a compact set, TB a transformer block as in Definition 3, and  $\alpha \in \mathfrak{O}_n^{Md_{\mathrm{in}}}, n \in \mathbb{N}$ . Then

$$C_K^{\mathcal{TB}}(\alpha) \leqslant \alpha! \sum_{\beta \in \mathfrak{O}_{\leq n}^{d_{\mathrm{out}}}} N(\beta) C_{K^{(3)}}^{\mathcal{LN}}(\beta) \sum_{\eta, \zeta \in \mathcal{P}'(\alpha, \beta)} \prod_{j=1}^n \frac{C_K^{(3)}(\mathfrak{o}(\zeta^{(j)}))^{|\eta^{(j)}|}}{\eta^{(j)}! (\zeta^{(j)}!)^{|\eta^{(j)}|}},$$

where for all  $\gamma \in \mathfrak{O}^{Md_{\mathrm{in}}}_{\leq n}$ :

$$\begin{split} C_K^{(3)}(\gamma) &\stackrel{\text{def.}}{=} \gamma! \sum_{\beta \in \mathcal{D}_{\leqslant n}^{d_{\text{in}}}} N(\beta) C_{K^{(2)}}^{\mathcal{PL}}(\beta) \sum_{\eta, \zeta \in \mathcal{P}'(\gamma, \beta)} \prod_{j=1}^n \frac{C_K^{(2)}(\mathfrak{o}(\zeta^{(j)}))^{|\eta^{(j)}|}}{\eta^{(j)}! (\zeta^{(j)}!)^{|\eta^{(j)}|}}, \\ C_K^{(2)}(\gamma) &\stackrel{\text{def.}}{=} \gamma! \sum_{\beta \in \mathcal{D}_{\leqslant n}^{d_{\text{in}}}} N(\beta) C_{K^{(1)}}^{\mathcal{LN}}(\beta) \sum_{\eta, \zeta \in \mathcal{P}'(\gamma, \beta)} \prod_{j=1}^n \frac{C_K^{(1)}(\mathfrak{o}(\zeta^{(j)}))^{|\eta^{(j)}|}}{\eta^{(j)}! (\zeta^{(j)}!)^{|\eta^{(j)}|}}, \\ C_K^{(1)}(\gamma) &\stackrel{\text{def.}}{=} \mathbbm{1}_{|\gamma|=1} + C_K^{\mathcal{MH}}(\gamma). \end{split}$$

In the above,  $K^{(1)} = \bigcup_{m=0}^{M} \mathcal{MH}_{m}[K]$ ,  $K^{(2)} = \mathcal{LN}[K^{(1)}]$ , and  $K^{(3)} = \mathcal{PL}[K^{(2)}]$ . For respective multi-indices, a bound for  $C_{K}^{\mathcal{LN}}$ ,  $C_{K}^{\mathcal{LN}}$  is given by Lemma 11,  $C_{K}^{\mathcal{PL}}$  is bounded in Lemma 12, and a bound for  $C_{K}^{\mathcal{MH}}$  is given in Corollary 4.

**Proof** This is a direct consequence of Theorem 4.

Theorem 6 (By Derivative Level) Let K be a compact set, TB a transformer block as in Definition 3, and  $n \in \mathbb{N}$ . Then,

$$C_K^{\mathcal{TB}}(\leqslant n) \leqslant C_{K^{(3)}}^{\mathcal{LN}}(\leqslant n) \left(d_{\text{out}}C_{K^{(2)}}^{\mathcal{PL}}(\leqslant n)\right)^n \left(d_{\text{in}}^2 C_{K^{(1)}}^{\mathcal{LN}}(\leqslant n)\right)^{n^2} \\ \cdot \left(1 + C_K^{\mathcal{MH}}(\leqslant n)\right)^{n^3} \left[\frac{2n}{e \ln n} (1 + o(1))\right]^{n+n^2+n^3}$$

where,  $K^{(1)} = \bigcup_{m=0}^{M} \mathcal{MH}_{m}[K]$ ,  $K^{(2)} = \mathcal{LN}[K^{(1)}]$ , and  $K^{(3)} = \mathcal{PL}[K^{(2)}]$ .

A bound for  $C_{K^{(3)}}^{\mathcal{LN}}$ ,  $C_{K^{(1)}}^{\mathcal{LN}}$  is given by Corollary 8,  $C_{K^{(2)}}^{\mathcal{PL}}$  is bounded in Corollary 9, and a bound for  $C_{K}^{\mathcal{MH}}$  is given in Corollary 4.

**Proof** Corollary 1 yields

$$C_K^{\mathcal{TB}}({\scriptscriptstyle \leqslant} n) \leqslant C_{K^{(3)}}^{\mathcal{LN}}({\scriptscriptstyle \leqslant} n) C_K^{(3)}({\scriptscriptstyle \leqslant} n)^n \Big[ \frac{2d_{\mathrm{out}} n}{e \ln n} (1 + o(1)) \Big]^n.$$

where

$$\begin{split} C_K^{(3)}(\leqslant n) &\stackrel{\text{def.}}{=} C_{K^{(2)}}^{\mathcal{PL}}(\leqslant n) C_K^{(2)}(\leqslant n)^n \Big[ \frac{2d_{\text{in}}n}{e \ln n} (1 + o(1)) \Big]^n, \\ C_K^{(2)}(\leqslant n) &\stackrel{\text{def.}}{=} C_{K^{(1)}}^{\mathcal{LN}}(\leqslant n) C_K^{(1)}(\leqslant n)^n \Big[ \frac{2d_{\text{in}}n}{e \ln n} (1 + o(1)) \Big]^n, \\ C_K^{(1)}(\leqslant n) &\stackrel{\text{def.}}{=} 1 + C_K^{\mathcal{MH}}(\leqslant n), \end{split}$$

which concludes the proof.

Theorem 7 (C<sup>s</sup>-Norm Bound of Transformers) Fix  $n, L, H, C, D, d, M \in \mathbb{N}_+$ . For any  $\mathcal{T} \in \mathcal{TC}_{L,H,C}^{D,d,M}$ , any compact  $K_0 \subset \mathbb{R}^{M \times D}$ , and any  $\alpha \in \mathbb{N}^{M \times D}$ ,  $|\alpha| \stackrel{\text{def.}}{=} n$  we have

$$C_{K_0}^{\mathcal{T}}(\alpha) \leqslant d_{\text{out}}^L M \alpha! \cdot C^A \cdot C^L(\alpha),$$
 (34)

where  $C^1(\alpha) \stackrel{\text{def.}}{=} C_{K_0}^{\mathcal{TB}_1}(\alpha)$  and for  $l \in \{2, \ldots, L\}$ ,

$$C^{l}(\alpha) \stackrel{\text{\tiny def.}}{=} \leqslant \alpha! \sum_{\beta \in \mathfrak{O}_{\leqslant n}^{\bar{d}_{l}}} N(\beta) C_{K_{l-1}}^{\mathcal{TB}_{l}}(\beta) \sum_{\eta, \zeta \in \mathcal{P}'(\mathfrak{o}(\alpha), \beta)} \prod_{j=1}^{n} \frac{C^{l}(\mathfrak{o}(\zeta^{(j)}))^{|\eta^{(j)}|}}{\eta^{(j)}! (\zeta^{(j)}!)^{|\eta^{(j)}|}}$$
(35)

where  $K_l \stackrel{\text{def.}}{=} \mathcal{TB}_l[K_{l-1}]$ ,  $\tilde{d}_l \stackrel{\text{def.}}{=} M_l d_{\text{in}}^l$ , and a bound for  $C_{K_{l-1}}^{\mathcal{TB}_l}(\beta)$  is given by Theorem 5, only depending on the transformer block class  $\mathcal{TBC}_{d^l,M^l,H^l,C^l}$ .

**Proof** [Proof of Theorem 7] The bounds (34) are a direct consequence of Theorem 4 and (35) follows directly from Corollary 2.

# D.5 Step 2 (C) - Merging the $C^s$ -Norm Bounds for Transformers with the Loss Function

In this section, we consider the following generalization of the class in Definition 5.

**Definition 7 (Smoothness Growth Rate)** Let  $d, D \in \mathbb{R}$ . A smooth function  $g : \mathbb{R}^d \to \mathbb{R}^D$  is said to belong to the class  $C^{\infty}_{poly:C,r}(\mathbb{R}^d,\mathbb{R}^D)$  (resp.  $C^{\infty}_{exp:C,r}(\mathbb{R}^d,\mathbb{R}^D)$ ) if there exist  $C, r \geq 0$  such that: for each  $s \in \mathbb{N}_+$ 

- (i) Polynomial Growth  $C^{\infty}_{poly:C,r}(\mathbb{R}^d,\mathbb{R}^D)$ :  $\|g\|_{C^s}\leqslant C\,s^r,$
- (ii) Exponential Growth  $C^{\infty}_{exp:C,r}(\mathbb{R}^d,\mathbb{R}^D)$ :  $\|g\|_{C^s} \leqslant C e^{s \, r}$ ,

The next lemma will help us relate the  $C^s$ -regularity of a model, a target function, and a loss function to their composition and product. We use it to relate the  $C^s$ -regularity of a transformed model  $\mathcal{T}: \mathbb{R}^d \to \mathbb{R}^D$ , the target function  $f^*: \mathbb{R}^d \to \mathbb{R}^D$ , and the loss function  $\ell: \mathbb{R}^{2D} \to \mathbb{R}$  to their composition

$$\ell_{\mathcal{T}}: \mathbb{R}^d \to \mathbb{R}$$

$$x \mapsto \ell(\mathcal{T}(x), f^*(x)). \tag{36}$$

One we computed have the  $C^s$ -regularity of  $\ell_{\mathcal{T}}$ , we can apply a concentration of measure-type argument based on an optimal transport-type duality, as in [2, 49, 14, 60], to obtain our generalization bounds. A key technical point where our analysis largely deviates from the mentioned derivations, is that we are not relying on any i.i.d. assumptions.

More generally, the next lemma allows us to bound the size of  $\|\ell(\hat{f}, f^*)\|_{C^s}$  using bounds on  $C^s$  norms of  $\mathcal{TC}$  computed in Theorem 7, the target function  $f^*$ , and on the loss function  $\ell$ . Naturally, to use this result, we must assume a given level of regularity of the target function, as in Definition 5.

Lemma 13 ( $C^s$ -Norm of loss of between two functions) Let  $d, D, s \in \mathbb{N}_+$ ,  $f_1, f_2 : \mathbb{R}^d \to \mathbb{R}^D$  be of class  $C^s$  and  $\ell : \mathbb{R}^{2D} \to \mathbb{R}$  be smooth. If there are constants  $C_1, C_2, \widetilde{C}_1, \ldots, \widetilde{C}_s \geqslant 0$  such that:

 $||f_i||_{C^s} \leqslant C_i \text{ for } i=1,2 \text{ and for } j=1,\ldots,s \text{ we have } ||\ell||_{C^j} \leqslant \widetilde{C}_j \text{ then for all } s>0 \text{ large it satisfies}$ 

$$\|\ell(f_1, f_2)\|_{C^s} = \begin{cases} \mathcal{O}\left[\left(\frac{2Ds}{e\ln s}(1+o(1))\right)^s\right], & \text{if } \max_{1\leqslant k} \widetilde{C}_k(C_1C_2)^k \text{ is bounded,} \\ \mathcal{O}\left[\widetilde{C}_s\left(C_1C_2\frac{2Ds}{e\ln s}(1+o(1))\right)^s\right], & \text{if } \max_{1\leqslant k} \widetilde{C}_k(C_1C_2)^k \text{ is unbounded.} \end{cases}$$
(37)

Particularly, if  $\ell \in C^{\infty}_{poly:C,r}(\mathbb{R}^{2D},\mathbb{R})$ , i.e.,  $\|\ell\|_{C^j} \leqslant C\,j^r$ , then

$$\|\ell(f_1, f_2)\|_{C^s} = \mathcal{O}\left[Cs^r\left(C_1C_2\frac{2Ds}{e\ln s}(1 + o(1))\right)^s\right];$$
 (38)

if  $\ell \in C^{\infty}_{exp:C,r}(\mathbb{R}^{2D}, \mathbb{R})$ , i.e.,  $\|\ell\|_{C^j} \leqslant C e^{j r}$ , then

$$\|\ell(f_1, f_2)\|_{C^s} = \mathcal{O}\left[Ce^{sr}\left(C_1C_2\frac{2Ds}{e\ln s}(1 + o(1))\right)^s\right]. \tag{39}$$

Lemma 13 allows us to obtain a bound on the term  $\sup_{\hat{f} \in C_R^s(\mathbb{R}^d)} \|\ell(\hat{f}, f^*)\|_{C^s}$  in (4), using Theorem 7 and our assumptions on  $\ell$  and on  $f^*$ .

**Proof** [Proof of Lemma 13] We first derive the general bound; which we then specialize to the case where the growth rate of  $\ell$  is known. We first observe that

$$\begin{aligned} & \|\ell(f_1, f_2)\|_{C^s} = \underbrace{\max_{k=1, \cdots, s-1} \max_{\alpha \in \{1, \cdots, d\}^k} \|D^{\alpha}\ell(f_1(x), f_2(x))\|_{\infty}}_{(\text{VII})} \\ & + \underbrace{\max_{\alpha \in \{1, \cdots, d\}^{s-1}} \text{Lip}\left(D^{\alpha}\ell(f_1(x), f_2(x))\right)}_{(\text{VIII})}. \end{aligned}$$

General Case - Term Term (VII): By Corollary 1, we have

$$\left\| (D^{\alpha}\ell)(f_1(x), f_2(x)) \right\|_{\infty} \leqslant \left[ \max_{1 \le k \le s-1} \widetilde{C}_k(C_1C_2)^k \right] \cdot \mathcal{O}\left[ \left( \frac{2Dk}{e \ln k} (1 + o(1)) \right)^k \right], \tag{40}$$

From (40) we have for all large s > 0 that

$$\max_{k=1,\cdots,s-1} \max_{\alpha \in \{1,\cdots,d\}^k} \|D^{\alpha}\ell(f_1(x),f_2(x))\|_{\infty}$$

$$= \begin{cases} \mathcal{O}\left[\left(\frac{2Ds}{e \ln s}(1+o(1))\right)^s\right], & \text{if } \max_{1 \leqslant k} \widetilde{C}_k(C_1C_2)^k \text{ is bounded,} \end{cases}$$

$$= \begin{cases} \mathcal{O}\left[\widetilde{C}_s\left(C_1C_2\frac{2Ds}{e \ln s}(1+o(1))\right)^s\right], & \text{if } \max_{1 \leqslant k} \widetilde{C}_k(C_1C_2)^k \text{ is unbounded.} \end{cases}$$

General Case - Term Term (VIII): For each  $\alpha \in \{1, \dots, d\}^{s-1}$ , by the multivariate Faà di Bruno formula, we have

$$D^{\alpha}\ell(f_1(x), f_2(x)) = \sum_{1 \leq |\beta| \leq s-1} (D^{\beta}\ell)(f_1(x), f_2(x)) \sum_{\eta, \zeta \in \mathcal{P}(\alpha, \beta)} \alpha! \prod_{j=1}^{s-1} \frac{[D^{\zeta^{(j)}}(f_1(x), f_2(x))]^{\eta^{(j)}}}{\eta^{(j)}! (\zeta^{(j)}!)^{|\eta^{(j)}|}}.$$

The Lipschitz constants of the derivatives satisfy

$$\operatorname{Lip}\left(D^{\alpha}\ell(f_{1}(x), f_{2}(x))\right) = \sum_{1 \leq |\beta| \leq s-1} \operatorname{Lip}\left((D^{\beta}\ell)(f_{1}(x), f_{2}(x))\right) \sum_{\eta, \zeta \in \mathcal{P}(\alpha, \beta)} \alpha! \prod_{j=1}^{s-1} \frac{\operatorname{Lip}\left([D^{\zeta^{(j)}}(f_{1}(x), f_{2}(x))]^{\eta^{(j)}}\right)}{\eta^{(j)!}(\zeta^{(j)!})^{|\eta^{(j)}|}} \\
\leq \sum_{1 \leq |\beta| \leq s-1} \widetilde{C}_{|\beta|+1} \sum_{\eta, \zeta \in \mathcal{P}(\alpha, \beta)} \alpha! \prod_{j=1}^{s-1} \frac{(C_{1}C_{2})^{|\eta(j)|}}{\eta^{(j)!}(\zeta^{(j)!})^{|\eta^{(j)}|}} \\
= \sum_{1 \leq |\beta| \leq s-1} \widetilde{C}_{|\beta|+1}(C_{1}C_{2})^{|\beta|} \sum_{\eta, \zeta \in \mathcal{P}(\alpha, \beta)} \alpha! \prod_{j=1}^{s-1} \frac{1}{\eta^{(j)!}(\zeta^{(j)!})^{|\eta^{(j)}|}} \\
\leq \left[ \max_{1 \leq k \leq s-1} \widetilde{C}_{k+1}(C_{1}C_{2})^{k} \right] \sum_{1 \leq |\beta| \leq s-1} \sum_{\eta, \zeta \in \mathcal{P}(\alpha, \beta)} \alpha! \prod_{j=1}^{s-1} \frac{1}{\eta^{(j)!}(\zeta^{(j)!})^{|\eta^{(j)}|}} \\
= \left[ \max_{1 \leq k \leq s-1} \widetilde{C}_{k+1}(C_{1}C_{2})^{k} \right] \cdot \mathcal{O}\left[\left(\frac{2Ds}{e \ln s}(1 + o(1))\right)^{s}\right], \tag{41}$$

where the last equality is due to Lemma 2.

From (41) we have for all s > 0 large that

$$\max_{\alpha \in \{1, \dots, d\}^{s-1}} \operatorname{Lip} \left( D^{\alpha} \ell(f_1(x), f_2(x)) \right) \\
= \begin{cases}
\mathcal{O} \left[ \left( \frac{2Ds}{e \ln s} (1 + o(1)) \right)^s \right], & \text{if } \max_{1 \leq k} \widetilde{C}_k (C_1 C_2)^k \text{ is bounded,} \\
\mathcal{O} \left[ \widetilde{C}_s \left( C_1 C_2 \frac{2Ds}{e \ln s} (1 + o(1)) \right)^s \right], & \text{if } \max_{1 \leq k} \widetilde{C}_k (C_1 C_2)^k \text{ is unbounded.} 
\end{cases}$$

Completing the General Case: Combining our estimates for terms Term (VIII) and Term (VIII) respectively obtained in (40) and (41), we obtain an upper-bound for  $\|\ell(f_1, f_2)\|_{C^s}$  via

$$\left\|\ell(f_1, f_2)\right\|_{C^s} = \begin{cases} \mathcal{O}\left[\left(\frac{2Ds}{e\ln s}(1+o(1))\right)^s\right], & \text{if } \max_{1\leqslant k} \widetilde{C}_k(C_1C_2)^k \text{ is bounded,} \\ \\ \mathcal{O}\left[\widetilde{C}_s\left(C_1C_2\frac{2Ds}{e\ln s}(1+o(1))\right)^s\right], & \text{if } \max_{1\leqslant k} \widetilde{C}_k(C_1C_2)^k \text{ is unbounded.} \end{cases}$$

**Special Cases of Interest:** In particular, in the case that  $\ell$  belongs either to  $C^{\infty}_{poly:C,r}(\mathbb{R}^d,\mathbb{R}^D)$  or to  $C^{\infty}_{exp:C,r}(\mathbb{R}^d,\mathbb{R}^D)$ , as in Definition (5), then: there exists constants  $C_{\ell}, r_{\ell} > 0$  s.t. for each  $j = 1, \ldots, s$  we have

(i) Polynomial Growth -  $C^{\infty}_{poly:C,r}(\mathbb{R}^{2D},\mathbb{R})$  Case:

$$\|\ell\|_{C^j} \leqslant C j^r \stackrel{\text{def.}}{=} \tilde{C}_j,$$

(ii) Exponential Growth -  $C^{\infty}_{exp:C,r}(\mathbb{R}^{2D},\mathbb{R})$  Case:

$$\|\ell\|_{C^j} \leqslant C e^{j r} \stackrel{\text{def.}}{=} \tilde{C}_j.$$

Consequentially, in cases (i) and (ii), the bound in (37) respectively becomes

(i) Polynomial Growth -  $C^{\infty}_{poly:C,r}(\mathbb{R}^{2D},\mathbb{R})$  Case:

$$\|\ell(f_1, f_2)\|_{C^s} \le \mathcal{O}\left[Cs^r\left(C_1C_2\frac{2Ds}{e\ln s}(1 + o(1))\right)^s\right],$$

(ii) Exponential Growth -  $C^{\infty}_{exp:C,r}(\mathbb{R}^{2D},\mathbb{R})$  Case:

$$\|\ell(f_1, f_2)\|_{C^s} \leq \mathcal{O}\left[Ce^{sr}\left(C_1C_2\frac{2Ds}{e\ln s}(1 + o(1))\right)^s\right].$$

## D.6 Step 3 - Combining Steps 1 and 2 and Completing The Proof of Theorem 1

We are now ready to complete the proof of our main result, namely Theorem 1. Before doing so, we state a more technical and general version, which we instead prove and which directly implies the simpler version found in the main body of our manuscript.

We operate under the following more general, but more technical set of assumptions than those considered in the main body of our text (in Setting 3.1).

Setting D.1 (Generalized Setting) Let  $D, d, L, H, *C', C^A, C^b \in \mathbb{N}_+$ , set  $M \stackrel{\text{def.}}{=} 0$ , as well as  $C \stackrel{\text{def.}}{=} (*C', C^A, C^b)$ ,  $r_f, r_\ell, C_f, C_\ell \geqslant 0$ . Suppose that Assumptions 1 and 3 hold.

Fix a target function  $f^*: \mathbb{R}^d \to \mathbb{R}^D$  and a loss function  $\ell: \mathbb{R}^D \times \mathbb{R}^D \to \mathbb{R}$ . Assume either that:

- (i) **Polynomial Growth:**  $f^* \in C^{\infty}_{poly:C_{\ell},r_{\ell}}(\mathbb{R}^d,\mathbb{R}^D)$  and  $\ell \in C^{\infty}_{poly:C_{\ell},r_{\ell}}(\mathbb{R}^{2D},\mathbb{R})$ ,
- (ii) Exponential Growth:  $f^* \in C^{\infty}_{exp:C_f,r_f}(\mathbb{R}^d,\mathbb{R}^D)$  and  $\ell \in C^{\infty}_{exp:C_\ell,r_\ell}(\mathbb{R}^{2D},\mathbb{R})$ ,
- (iii) No Growth: There is a constant  $\bar{C} \geqslant 0$  such that for all s > 0 we have  $||f^*||_{C^s}, ||\ell||_{C^s} \leqslant \bar{C}$ .

**Example 1 (Example of Generalized Setting (iii))** For every  $d \in \mathbb{R}^d$ , the function  $f : \mathbb{R}^d \ni x \mapsto \cos \bullet x = \left(\cos(x_i)\right)_{i=1}^d satisfies \|\frac{\partial^s}{\partial x_i^s} f\|_{\infty} \leqslant 1$  for each  $s \in \mathbb{N}$  and each  $i = 1, \ldots, d$ . Thus, it is an example of a function satisfying Assumption D.1.

We are now ready to prove our main theorem.

**Theorem 8 (Pathwise Generalization Bounds for Transformers)** In Setting D.1, there is a  $\kappa \in (0,1)$ , depending only on X., and a  $t_0 \in \mathbb{N}_0$  such that: for each  $t_0 \leqslant N \leqslant t \leqslant \infty$  and  $\delta \in (0,1]$  the following holds with probability at-least  $1-\delta$ 

$$\sup_{\mathcal{T} \in \mathcal{T}_{L,H,C}^{D,d,0}} \left| \mathcal{R}_{\max\{t,N\}}(\mathcal{T}) - \mathcal{R}^{(N)}(\mathcal{T}) \right| \lesssim \sum_{s=1}^{\infty} I_{N \in [\tau_s, \tau_{s+1})} C_{\ell,\mathcal{T},K,s-1} \left( I_{t < \infty} \kappa^t + \frac{\sqrt{2 \ln(1/\delta)}}{N^{1/2}} + \text{rate}_s(N) \right)$$

where rate<sub>s</sub>(N) is defined in (rate), and the constant  $C_{\ell,\mathcal{T},s,K}$  is Table 1 and the transition phases  $(\tau_s)_{s=0}^{\infty}$  are given iteratively by  $\tau_0 \stackrel{\text{def.}}{=} 0$  and for each  $s \in \mathbb{N}_+$ 

$$\tau_s \stackrel{\text{\tiny def.}}{=} \inf \bigg\{ t \geqslant \tau_{s-1} : C_{\ell,\mathcal{T},K,s} \big( \kappa^t + \mathrm{rate}_s(N) + \frac{\sqrt{\log(1/\delta)}}{\sqrt{N}} \big) \leqslant C_{\ell,\mathcal{T},K,s-1} \big( \kappa^t + \mathrm{rate}_{s-1}(N) + \frac{\sqrt{\log(1/\delta)}}{\sqrt{N}} \big) \bigg\}.$$

Furthermore,  $c \stackrel{\text{def.}}{=} 1 - \kappa$ ,  $c_2 \stackrel{\text{def.}}{=} c^{s/d}$ ,  $\kappa^{\infty} \stackrel{\text{def.}}{=} \lim_{t \to \infty} \kappa^t = 0$ , and  $\lesssim$  hides an absolute constant.

**Proof** [Proof of Theorem 1] Since N is given, we may pick  $s \in \mathbb{N}_+$  to ensure that  $N \in [\tau_s, \tau_{s+1})$ ; where these are defined as in the statement of Theorem 8.

Since we are in Setting 3.1, then  $\ell \in C^{\infty}_{poly:C_{\ell},r_{\ell}}(\mathbb{R}^{2D},\mathbb{R})$  (resp.  $\ell \in C^{\infty}_{exp:C_{\ell},r_{\ell}}(\mathbb{R}^{2D},\mathbb{R})$ ) and  $f^{\star}: \mathbb{R}^{d} \to \mathbb{R}^{D}$  is smooth. Therefore, Lemma 13 implies that there is an absolute constant  $c_{abs} > 0$  such that for any transformer network  $\mathcal{T} \in \mathcal{TC}^{D,d,0}_{L,H,C}$ , the following bound holds

(i) No Growth Case: Using (37) we find that

$$\|\ell(\mathcal{T}, f^*)\|_{C^s} \le c_{\text{abs}} \left(\frac{2Ds}{e \ln s} (1 + o(1))\right)^s \|\mathcal{T}\|_{C^s}^s$$
 (42)

(ii) Polynomial Growth Case -  $\ell \in C^{\infty}_{poly:C_{\ell},r_{\ell}}(\mathbb{R}^{2D},\mathbb{R})$  Case:

$$\|\ell(\mathcal{T}, f^{\star})\|_{C^{s}} \leqslant c_{\text{abs}} \, s^{r_{\ell}} \left(\frac{2Ds}{e \ln s} (1 + o(1))\right)^{s} \|f^{\star}\|_{C^{s}}^{s} \|\mathcal{T}\|_{C^{s}}^{s} \tag{43}$$

(iii) Exponential Growth -  $C^{\infty}_{exp:C_{\ell},r_{\ell}}(\mathbb{R}^{2D},\mathbb{R})$  Case:

$$\|\ell(\mathcal{T}, f^{\star})\|_{C^{s}} \leqslant c_{\text{abs}} e^{s \, r_{\ell}} \left(\frac{2Ds}{e \ln s} (1 + o(1))\right)^{s} \|f^{\star}\|_{C^{s}}^{s} \|\mathcal{T}\|_{C^{s}}^{s}. \tag{44}$$

Since we have assumed that  $f^* \in C^{\infty}_{poly:C_f,r_f}(\mathbb{R}^d,\mathbb{R}^D)$  (resp.  $C^{\infty}_{exp:C_f,r_f}(\mathbb{R}^d,\mathbb{R}^D)$  or the "no growth condition" in Setting D.1 (iii)) then the bounds in (42), (43), and (44), respectively, imply that

(i) No Growth Case:

$$\left\| \ell(\mathcal{T}, f^{\star}) \right\|_{C^{s}} \leqslant c_{\text{abs}} \left( \frac{2Ds}{e \ln s} (1 + o(1)) \right)^{s} C_{K}^{\mathcal{TC}}(s)^{s} \tag{45}$$

(ii) Polynomial Growth Case -  $\ell \in C^{\infty}_{poly:C_{\ell},r_{\ell}}(\mathbb{R}^{2D},\mathbb{R})$  Case:

$$\|\ell(\mathcal{T}, f^*)\|_{C^s} \leqslant c_{\text{abs}} \, s^{r_\ell + 2s^2} \left(\frac{C_f \, 2D}{e \ln s} (1 + o(1))\right)^s C_K^{\mathcal{TC}}(s)^s$$
 (46)

(iii) Exponential Growth -  $C^{\infty}_{exp:C_{\ell},r_{\ell}}(\mathbb{R}^{2D},\mathbb{R})$  Case:

$$\|\ell(\mathcal{T}, f^{\star})\|_{C^{s}} \leqslant c_{\text{abs}} e^{s \, r_{\ell} + s^{2} r_{f}} \left(\frac{2D \, s}{e \ln s} (1 + o(1))\right)^{s} C_{f}^{s} \, C_{K_{0}}^{\mathcal{TC}}(s)^{s}, \tag{47}$$

where we have used the definition of the constant  $C_K^{\mathcal{TC}}(s)$  as a uniform upper bound of  $\sup_{\mathcal{T} \in \mathcal{TC}_{L,H,C}^{D,d,0}}$ . Using Theorem 5 for the "derivative type estimate" (resp.6 for the "derivative level estimate") concludes the implies yields a uniform upper bound (of "derivative type" or "derivative level" respectively) on  $C_{K_0}^{\mathcal{TC}}(s)$ , i.e. independent of the particular transformer instance  $\mathcal{T} \in \mathcal{TC}_{L,H,C}^{D,d,0}$ . In either case, we respectively define R > 0 to be the right-hand side of (46) or (47) depending on the respective assumptions made on  $\ell$  and on  $f^*$ .

The conclusion now follows upon applying Proposition 2 due to the inequality in (3).

## E. Example of Additive Noise Using Stochastic Calculus

In this appendix, we briefly discuss why the seemingly realizable learning setting which we have placed ourselves in, i.e.  $Y_n = f^*(X_n)$ , does not preclude additive noise. Our illustration considers the class of following Markov processes.

**Assumption 4 (Structure on** X.) Let  $g: \mathbb{R}^d \to [0,1]^d$  be a twice continuously differentiable function. Let  $W: \stackrel{\text{def.}}{=} (W_t)_{t \geq 0}$  be d-dimensional Brownian motion and, for each  $n \in \mathbb{N}$ , define

$$X_n \stackrel{\text{def.}}{=} q(W_n).$$

By construction, the boundedness of the change of variables-type function g in Assumption 4, implies that the process  $X = (X_n)_{n \in \mathbb{N}}$  is bounded (and can easily be seen to be Markovian since Brownian motion has the strong Markov property). However, we can say more, indeed under Assumption 4, the Itô Lemma (see e.g. [25, Theorem 14.2.4]) implies that  $X_n$  is given as the following stochastic differential equation (SDE) evaluated at integer times  $n \in \mathbb{N}$ 

$$X_n = g(0) + \int_0^n \mu_s \, ds + \int_0^n \sigma_t^\top \, dW_s \tag{48}$$

where  $\mu_{\cdot} = (\mu_t)_{t \geqslant 0}$  and  $\sigma_{\cdot} = (\sigma_t)_{t \geqslant 0}$  are given by

$$\mu_t \stackrel{\text{def.}}{=} \frac{1}{2} \operatorname{tr} (H(g)(W_s)) \text{ and } \sigma_t \stackrel{\text{def.}}{=} \nabla g(W_t)$$
 (49)

and H(g) is the Hessian of g and tr is the trace of a matrix.

**Example 2** Set d=1 and  $g(x)=(\sin(x)+1)/2$ . Then, for each  $n\in\mathbb{N}$  we have

$$X_n = \int_0^n -\sin(W_s)/4ds + \int_0^t \cos(W_s)/2 \, dW_s.$$

In particular, the expression (48) shows that the input process X is also defined for all intermediate times between non-negative integer times; i.e. for each  $t \ge 0$  the process

$$X_{t} = g(0) + \int_{0}^{t} \mu_{s} \, ds + \int_{0}^{t} \sigma_{t}^{\top} \, dW_{s} \tag{50}$$

is well-defined and coincides with  $X_n$  whenever  $t = n \in \mathbb{N}$ . We may, therefore, also consider the "continuous-time extension"  $Y \stackrel{\text{def.}}{=} (Y_t)_{t \geqslant 0}$  of the target process defined for all intermediate times using (50) by

$$Y_t \stackrel{\text{def.}}{=} f^{\star}(X_t).$$

Note that  $Y_t$  coincides with the target process on non-negative integer times, as defined in our main text, by definition.

The convenience of these continuous-time extensions, of the discrete versions considered in our main text, is that now Y is the transformation of a continuous-time (Itô) process of satisfying the SDE (50) by a smooth function<sup>4</sup>, namely  $f^*$ . Therefore, we may again apply the Itô Lemma (again see e.g. [25, Theorem 14.2.4]) this time to the process X to obtain the desired signal and noise decomposition of the target process Y. (both in discrete and continuous time). Doing so yields the following decomposition

$$Y_{t} = \underbrace{f^{\star}(X_{0}) + \int_{0}^{t} \left( \left( \nabla f^{\star}(X_{s}) \right)^{\top} \mu_{t} + \frac{1}{2} \operatorname{tr} \left( \sigma_{s}^{\top} H(f^{\star})(X_{s}) \sigma_{s} \right) \right) ds}_{\text{Signal (Target)}} + \underbrace{\int_{0}^{t} \left( \nabla f^{\star} \right)^{\top} \sigma_{s} dW_{s}}_{\text{Additive Noise}}.$$

$$(51)$$

This shows that even if it a priori seemed that we are in the realizable PAC setting due to the structural assumption that  $Y_n = f^*(X_n)$  made when defining the target process, we are actually in the standard setting where the target data  $(Y_n)_{n=0}^{\infty}$  can be written as a signal plus an additive noise term. Indeed, when X. is simply a transformation of a Brownian motion by a bounded  $C^2$ -function, as in Assumption 4, then Assumption 2 held and  $Y_n$  admitted the signal-noise decomposition in (51).

<sup>4.</sup> Note that  $f^*$  was assumed to be smooth in our main result (Theorem 1).

## References

- [1] E. Akyürek, D. Schuurmans, J. Andreas, T. Ma, and D. Zhou. What learning algorithm is incontext learning? investigations with linear models. In *The Eleventh International Conference on Learning Representations*, 2023. URL https://openreview.net/forum?id=0g0X4H8yN4I.
- [2] R. Amit, B. Epstein, S. Moran, and R. Meir. Integral probability metrics pac-bayes bounds. *Advances in Neural Information Processing Systems*, 35:3123–3136, 2022.
- [3] Y. F. Atchadé. Approximate spectral gaps for markov chain mixing times in high dimensions. SIAM Journal on Mathematics of Data Science, 3(3):854–872, 2021.
- [4] K. Atkinson and W. Han. Spherical harmonics and approximations on the unit sphere: an introduction, volume 2044 of Lecture Notes in Mathematics. Springer, Heidelberg, 2012. ISBN 978-3-642-25982-1. doi: 10.1007/978-3-642-25983-8. URL https://doi.org/10.1007/ 978-3-642-25983-8.
- [5] J. L. Ba, J. R. Kiros, and G. E. Hinton. Layer normalization. arXiv preprint arXiv:1607.06450, 2016.
- [6] P. Bartlett, V. Maiorov, and R. Meir. Almost linear vc dimension bounds for piecewise polynomial networks. In M. Kearns, S. Solla, and D. Cohn, editors, Advances in Neural Information Processing Systems, volume 11. MIT Press, 1998. URL https://proceedings.neurips.cc/paper\_files/paper/1998/file/bc7316929fe1545bf0b98d114ee3ecb8-Paper.pdf.
- [7] P. L. Bartlett and S. Mendelson. Rademacher and gaussian complexities: Risk bounds and structural results. *Journal of Machine Learning Research*, 3(Nov):463–482, 2002.
- P. L. Bartlett, O. Bousquet, and S. Mendelson. Local Rademacher complexities. Ann. Statist., 33(4):1497–1537, 2005. ISSN 0090-5364,2168-8966. doi: 10.1214/009053605000000282. URL https://doi.org/10.1214/009053605000000282.
- [9] P. L. Bartlett, D. J. Foster, and M. J. Telgarsky. Spectrally-normalized margin bounds for neural networks. In I. Guyon, U. V. Luxburg, S. Bengio, H. Wallach, R. Fergus, S. Vishwanathan, and R. Garnett, editors, Advances in Neural Information Processing Systems, volume 30. Curran Associates, Inc., 2017. URL https://proceedings.neurips.cc/paper\_files/paper/2017/ file/b22b257ad0519d4500539da3c8bcf4dd-Paper.pdf.
- [10] P. L. Bartlett, N. Harvey, C. Liaw, and A. Mehrabian. Nearly-tight VC-dimension and pseudodimension bounds for piecewise linear neural networks. J. Mach. Learn. Res., 20:Paper No. 63, 17, 2019. ISSN 1532-4435,1533-7928.
- [11] D. Barzilai and O. Shamir. Generalization in kernel regression under realistic assumptions, 2023.
- [12] J. R. Beck and S. G. Pauker. The markov process in medical prognosis. Medical decision making, 3(4):419–458, 1983.
- [13] E. T. Bell. Exponential polynomials. Annals of Mathematics, pages 258–277, 1934.
- [14] J. Benitez, T. Furuya, F. Faucher, A. Kratsios, X. Tricoche, and M. V. de Hoop. Out-of-distributional risk bounds for neural operators with applications to the helmholtz equation. arXiv preprint arXiv:2301.11509, 2023.
- [15] B. Bercu, B. Delyon, and E. Rio. Concentration inequalities for sums and martingales. Springer-Briefs in Mathematics. Springer, Cham, 2015. ISBN 978-3-319-22098-7; 978-3-319-22099-4. doi: 10.1007/978-3-319-22099-4. URL https://doi.org/10.1007/978-3-319-22099-4.

- [16] S. Bhattamishra, A. Patel, and N. Goyal. On the computational power of transformers and its implications in sequence modeling. arXiv preprint arXiv:2006.09286, 2020.
- [17] S. G. Bobkov and F. Götze. Exponential integrability and transportation cost related to logarithmic Sobolev inequalities. *J. Funct. Anal.*, 163(1):1–28, 1999. ISSN 0022-1236,1096-0783. doi: 10.1006/jfan.1998.3326. URL https://doi.org/10.1006/jfan.1998.3326.
- [18] R. Bommasani, D. A. Hudson, E. Adeli, R. Altman, S. Arora, S. von Arx, M. S. Bernstein, J. Bohg, A. Bosselut, E. Brunskill, et al. On the opportunities and risks of foundation models. ArXiv preprint, abs/2108.07258, 2021.
- [19] S. Boucheron, G. Lugosi, and P. Massart. Concentration inequalities, 2013. URL https://doi.org/10.1093/acprof:oso/9780199535255.001.0001. A nonasymptotic theory of independence, With a foreword by Michel Ledoux.
- [20] K. N. Boyadzhiev. Derivative polynomials for tanh, tan, sech and sec in explicit form. *Fibonacci Quarterly*, 45(4):291–303, 2007.
- [21] P. Chaudhari, A. Choromanska, S. Soatto, Y. LeCun, C. Baldassi, C. Borgs, J. Chayes, L. Sagun, and R. Zecchina. Entropy-sgd: biasing gradient descent into wide valleys\*. *Journal of Statistical Mechanics: Theory and Experiment*, 2019(12):124018, dec 2019. doi: 10.1088/1742-5468/ab39d9. URL https://dx.doi.org/10.1088/1742-5468/ab39d9.
- [22] H.-B. Chen, S. Chewi, and J. Niles-Weed. Dimension-free log-Sobolev inequalities for mixture distributions. J. Funct. Anal., 281(11):Paper No. 109236, 17, 2021. ISSN 0022-1236,1096-0783. doi: 10.1016/j.jfa.2021.109236. URL https://doi.org/10.1016/j.jfa.2021.109236.
- [23] T. S. Cheng, A. Lucchi, A. Kratsios, and D. Belius. Characterizing overfitting in kernel ridgeless regression through the eigenspectrum, 2024.
- [24] T. S. Cheng, A. Lucchi, A. Kratsios, and D. Belius. A comprehensive analysis on the learning curve in kernel ridge regression, 2024.
- [25] S. N. Cohen and R. J. Elliott. Stochastic calculus and applications. Probability and its Applications. Springer, Cham, second edition, 2015. ISBN 978-1-4939-2866-8; 978-1-4939-2867-5.
   5. doi: 10.1007/978-1-4939-2867-5. URL https://doi.org/10.1007/978-1-4939-2867-5.
- [26] G. M. Constantine and T. H. Savits. A multivariate Faà di Bruno formula with applications. Trans. Amer. Math. Soc., 348(2):503-520, 1996. ISSN 0002-9947. doi: 10.1090/S0002-9947-96-01501-2. URL https://doi.org/10.1090/S0002-9947-96-01501-2.
- [27] C. Cuchiero and J. Teichmann. Markovian lifts of positive semidefinite affine volterra-type processes. *Decisions in Economics and Finance*, 42:407–448, 2019.
- [28] C. Cuchiero and J. Teichmann. Generalized Feller processes and Markovian lifts of stochastic Volterra processes: the affine case. J. Evol. Equ., 20(4):1301–1348, 2020. ISSN 1424-3199,1424-3202. doi: 10.1007/s00028-020-00557-2. URL https://doi.org/10.1007/s00028-020-00557-2.
- [29] M. A. de Oliveira and R. H. Ikeda. Representation of the n-th derivative of the normal pdf using bernoulli numbers and gamma function. Applied Mathematical Sciences, 6(74):3661–3673, 2012.
- [30] S. Dereich, M. Scheutzow, and R. Schottstedt. Constructive quantization: approximation by empirical measures. *Ann. Inst. Henri Poincaré Probab. Stat.*, 49(4):1183–1203, 2013. ISSN 0246-0203,1778-7017. doi: 10.1214/12-AIHP489. URL https://doi.org/10.1214/12-AIHP489.

- [31] E. Dolera and E. Mainini. Lipschitz continuity of probability kernels in the optimal transport framework. *Ann. Inst. Henri Poincaré Probab. Stat.*, 59(4):1778–1812, 2023. ISSN 0246-0203,1778-7017. doi: 10.1214/23-aihp1389. URL https://doi.org/10.1214/23-aihp1389.
- [32] L. C. Evans. Partial differential equations, volume 19. American Mathematical Society, 2022.
- [33] F. Faa di Bruno. Sullo sviluppo delle funzioni. Annali di scienze matematiche e fisiche, 6(1): 479–480, 1855.
- [34] Z. Fang, Y. Ouyang, D.-X. Zhou, and G. Cheng. Attention enables zero approximation error, 2023. URL https://openreview.net/forum?id=AV\_bv4Ydcr9.
- [35] C. L. Fefferman. A sharp form of Whitney's extension theorem. Ann. of Math. (2), 161 (1):509-577, 2005. ISSN 0003-486X,1939-8980. doi: 10.4007/annals.2005.161.509. URL https://doi.org/10.4007/annals.2005.161.509.
- [36] H. Föllmer and A. Schied. Stochastic finance. Walter de Gruyter & Co., Berlin, extended edition, 2011. ISBN 978-3-11-021804-6. doi: 10.1515/9783110218053. URL https://doi.org/10.1515/9783110218053. An introduction in discrete time.
- [37] S. Frieder, L. Pinchetti, R.-R. Griffiths, T. Salvatori, T. Lukasiewicz, P. Petersen, and J. Berner. Mathematical capabilities of chatgpt. Advances in Neural Information Processing Systems, 36, 2024.
- [38] T. Galanti, M. Xu, L. Galanti, and T. Poggio. Norm-based generalization bounds for sparse neural networks. *Advances in Neural Information Processing Systems*, 36, 2024.
- [39] S. Garg, D. Tsipras, P. S. Liang, and G. Valiant. What can transformers learn in-context? a case study of simple function classes. *Advances in Neural Information Processing Systems*, 35: 30583–30598, 2022.
- [40] N. Golowich, A. Rakhlin, and O. Shamir. Size-independent sample complexity of neural networks. *Information and Inference: A Journal of the IMA*, 2020.
- [41] N. Gozlan. A characterization of dimension free concentration in terms of transportation inequalities. Ann. Probab., 37(6):2480-2498, 2009. ISSN 0091-1798,2168-894X. doi: 10.1214/ 09-AOP470. URL https://doi.org/10.1214/09-AOP470.
- [42] N. Gozlan. Poincaré inequalities and dimension free concentration of measure. Ann. Inst. Henri Poincaré Probab. Stat., 46(3):708-739, 2010. ISSN 0246-0203,1778-7017. doi: 10.1214/09-AIHP209. URL https://doi.org/10.1214/09-AIHP209.
- [43] N. Gozlan, C. Roberto, and P.-M. Samson. From dimension free concentration to the Poincaré inequality. Calc. Var. Partial Differential Equations, 52(3-4):899-925, 2015. ISSN 0944-2669,1432-0835. doi: 10.1007/s00526-014-0737-6. URL https://doi.org/10.1007/ s00526-014-0737-6.
- [44] S. Graf and H. Luschgy. Foundations of quantization for probability distributions, volume 1730 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 2000. ISBN 3-540-67394-6. doi: 10.1007/BFb0103945. URL https://doi.org/10.1007/BFb0103945.
- [45] L. Gross. Logarithmic Sobolev inequalities. Amer. J. Math., 97(4):1061–1083, 1975. ISSN 0002-9327,1080-6377. doi: 10.2307/2373688. URL https://doi.org/10.2307/2373688.
- [46] M. Hardy. Combinatorics of partial derivatives. Electron. J. Combin., 13(1):Research Paper 1, 13, 2006. doi: 10.37236/1027. URL https://doi.org/10.37236/1027.

- [47] D. Hendrycks and K. Gimpel. Gaussian error linear units (GeLUs). arXiv preprint arXiv:1606.08415, 2016.
- [48] S. Hou, P. Kassraie, A. Kratsios, A. Krause, and J. Rothfuss. Instance-dependent generalization bounds via optimal transport. J. Mach. Learn. Res., 24:Paper No. [349], 51, 2023. ISSN 1532-4435,1533-7928.
- [49] S. Hou, P. Kassraie, A. Kratsios, A. Krause, and J. Rothfuss. Instance-dependent generalization bounds via optimal transport. *Journal of Machine Learning Research*, 24:1–50, 2023.
- [50] D. J. Hsu, A. Kontorovich, and C. Szepesvári. Mixing time estimation in reversible markov chains from a single sample path. *Advances in neural information processing systems*, 28, 2015.
- [51] J. Inglis and I. Papageorgiou. Log-Sobolev inequalities for infinite-dimensional Gibbs measures with non-quadratic interactions. *Markov Process. Related Fields*, 25(5):879–897, 2019. ISSN 1024-2953.
- [52] O. Khorunzhiy. On asymptotic properties of bell polynomials and concentration of vertex degree of large random graphs. *Journal of Theoretical Probability*, 35:1–32, 03 2022. doi: 10.1007/s10959-020-01025-w.
- [53] B. Kloeckner. Effective berry–esseen and concentration bounds for markov chains with a spectral gap. *The Annals of Applied Probability*, 29(3):1778–1807, 2019.
- [54] B. R. Kloeckner. Empirical measures: regularity is a counter-curse to dimensionality. ESAIM Probab. Stat., 24:408–434, 2020. ISSN 1292-8100,1262-3318. doi: 10.1051/ps/2019025. URL https://doi.org/10.1051/ps/2019025.
- [55] V. Koltchinskii. Rademacher penalties and structural risk minimization. *IEEE Transactions on Information Theory*, 2001.
- [56] A. Kontorovich. Concentration in unbounded metric spaces and algorithmic stability. In E. P. Xing and T. Jebara, editors, Proceedings of the 31st International Conference on Machine Learning, volume 32 of Proceedings of Machine Learning Research, pages 28–36, Bejing, China, 22–24 Jun 2014. PMLR. URL https://proceedings.mlr.press/v32/kontorovicha14.html.
- [57] A. Kontorovich and M. Raginsky. Concentration of measure without independence: a unified approach via the martingale method. In *Convexity and Concentration*, pages 183–210. Springer, 2017.
- [58] I. Kontoyiannis and S. P. Meyn. Geometric ergodicity and the spectral gap of non-reversible markov chains. *Probability Theory and Related Fields*, 154(1):327–339, 2012.
- [59] A. Kratsios and L. Papon. Universal approximation theorems for differentiable geometric deep learning. *Journal of Machine Learning Research*, 23(196):1–73, 2022.
- [60] A. Kratsios, A. M. Neuman, and G. Pammer. Tighter generalization bounds on digital computers via discrete optimal transport, 2024.
- [61] M. Ledoux. Concentration of measure and logarithmic sobolev inequalities. In Seminaire de probabilites XXXIII, pages 120–216. Springer, 2006.
- [62] M. Ledoux, I. Nourdin, and G. Peccati. Stein's method, logarithmic Sobolev and transport inequalities. Geom. Funct. Anal., 25(1):256-306, 2015. ISSN 1016-443X,1420-8970. doi: 10.1007/s00039-015-0312-0. URL https://doi.org/10.1007/s00039-015-0312-0.

- [63] V. Likhosherstov, K. Choromanski, and A. Weller. On the expressive power of self-attention matrices. arXiv preprint arXiv:2106.03764, 2021.
- [64] M. Mihoubi. Bell polynomials and binomial type sequences. *Discrete Mathematics*, 308(12): 2450–2459, 2008.
- [65] A. A. Minai and R. D. Williams. On the derivatives of the sigmoid. Neural Networks, 6(6): 845-853, 1993. ISSN 0893-6080. doi: https://doi.org/10.1016/S0893-6080(05)80129-7. URL https://www.sciencedirect.com/science/article/pii/S0893608005801297.
- [66] K. Minami, H. Arai, I. Sato, and H. Nakagawa. Differential privacy without sensitivity. Advances in Neural Information Processing Systems, 29, 2016.
- [67] R. Montenegro, P. Tetali, et al. Mathematical aspects of mixing times in markov chains. Foundations and Trends® in Theoretical Computer Science, 1(3):237–354, 2006.
- [68] C. Mufa. Estimation of spectral gap for markov chains. *Acta Mathematica Sinica*, 12(4): 337–360, 1996.
- [69] B. Neyshabur, R. Tomioka, and N. Srebro. Norm-based capacity control in neural networks. In P. Grünwald, E. Hazan, and S. Kale, editors, Proceedings of The 28th Conference on Learning Theory, volume 40 of Proceedings of Machine Learning Research, pages 1376-1401, Paris, France, 03-06 Jul 2015. PMLR. URL https://proceedings.mlr.press/v40/Neyshabur15.html.
- [70] B. Neyshabur, Z. Li, S. Bhojanapalli, Y. LeCun, and N. Srebro. The role of over-parametrization in generalization of neural networks. In *International Conference on Learning Representations*, 2019. URL https://openreview.net/forum?id=BygfghAcYX.
- [71] C. Olivera and C. Tudor. Density for solutions to stochastic differential equations with unbounded drift. *Brazilian Journal of Probability and Statistics*, 33(3):520–531, 2019.
- [72] W. Paul and J. Baschnagel. Stochastic processes: From physics to finance. Springer, Heidelberg, second edition, 2013. ISBN 978-3-319-00326-9; 978-3-319-00327-6. doi: 10.1007/978-3-319-00327-6. URL https://doi.org/10.1007/978-3-319-00327-6.
- [73] D. Paulin. Concentration inequalities for Markov chains by Marton couplings and spectral methods. *Electron. J. Probab.*, 20:no. 79, 32, 2015. ISSN 1083-6489. doi: 10.1214/EJP.v20-4039. URL https://doi.org/10.1214/EJP.v20-4039.
- [74] P. Ramachandran, B. Zoph, and Q. V. Le. Searching for activation functions. arXiv preprint arXiv:1710.05941, 2017.
- [75] A. Riekert. Convergence rates for empirical measures of Markov chains in dual and Wasserstein distances. Statist. Probab. Lett., 189:Paper No. 109605, 8, 2022. ISSN 0167-7152,1879-2103. doi: 10.1016/j.spl.2022.109605. URL https://doi.org/10.1016/j.spl.2022.109605.
- [76] W. Rudin. *Principles of mathematical analysis*. International Series in Pure and Applied Mathematics. McGraw-Hill Book Co., New York-Auckland-Düsseldorf, third edition, 1976.
- [77] J. B. Simon, M. Dickens, D. Karkada, and M. Deweese. The eigenlearning framework: A conservation law perspective on kernel ridge regression and wide neural networks. *Transactions on Machine Learning Research*, 2023.
- [78] B. K. Sriperumbudur, K. Fukumizu, A. Gretton, B. Schölkopf, and G. R. G. Lanckriet. On the empirical estimation of integral probability metrics. *Electron. J. Stat.*, 6:1550–1599, 2012. ISSN 1935-7524. doi: 10.1214/12-EJS722. URL https://doi.org/10.1214/12-EJS722.

- [79] R. S. Sutton and A. G. Barto. Reinforcement learning: an introduction. Adaptive Computation and Machine Learning. MIT Press, Cambridge, MA, second edition, 2018. ISBN 978-0-262-03924-6.
- [80] N. Touzi. Optimal stochastic control, stochastic target problems, and backward SDE, volume 29 of Fields Institute Monographs. Springer, New York; Fields Institute for Research in Mathematical Sciences, Toronto, ON, 2013. ISBN 978-1-4614-4285-1; 978-1-4614-4286-8. doi: 10.1007/ 978-1-4614-4286-8. URL https://doi.org/10.1007/978-1-4614-4286-8. With Chapter 13 by Angès Tourin.
- [81] J. Trauger and A. Tewari. Sequence length independent norm-based generalization bounds for transformers. arXiv preprint arXiv:2310.13088, 2023.
- [82] A. Tsigler and P. L. Bartlett. Benign overfitting in ridge regression. J. Mach. Learn. Res., 24: 123–1, 2023.
- [83] A. W. van der Vaart and J. A. Wellner. Weak convergence and empirical processes—with applications to statistics. Springer Series in Statistics. Springer, Cham, second edition, 2023. ISBN 978-3-031-29038-1; 978-3-031-29040-4. doi: 10.1007/978-3-031-29040-4. URL https://doi.org/10.1007/978-3-031-29040-4.
- [84] A. Vaswani, N. Shazeer, N. Parmar, J. Uszkoreit, L. Jones, A. N. Gomez, L. Kaiser, and I. Polosukhin. Attention is all you need. In I. Guyon, U. von Luxburg, S. Bengio, H. M. Wallach, R. Fergus, S. V. N. Vishwanathan, and R. Garnett, editors, Advances in Neural Information Processing Systems 30: Annual Conference on Neural Information Processing Systems 2017, December 4-9, 2017, Long Beach, CA, USA, pages 5998–6008, 2017.
- [85] C. Villani. Optimal transport, volume 338 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 2009. ISBN 978-3-540-71049-3. doi: 10.1007/978-3-540-71050-9. URL https://doi.org/10.1007/978-3-540-71050-9. Old and new.
- [86] J. Von Oswald, E. Niklasson, E. Randazzo, J. Sacramento, A. Mordvintsev, A. Zhmoginov, and M. Vladymyrov. Transformers learn in-context by gradient descent. In *International Conference on Machine Learning*, pages 35151–35174. PMLR, 2023.
- [87] W. Wang and T. Wang. General identities on bell polynomials. Computers & Mathematics with Applications, 58(1):104–118, 2009.
- [88] J. Wei, Y. Tay, R. Bommasani, C. Raffel, B. Zoph, S. Borgeaud, D. Yogatama, M. Bosma, D. Zhou, D. Metzler, et al. Emergent abilities of large language models. ArXiv preprint, abs/2206.07682, 2022.
- [89] E. P. Wigner. Characteristic vectors of bordered matrices with infinite dimensions. Ann. of Math. (2), 62:548-564, 1955. ISSN 0003-486X. doi: 10.2307/1970079. URL https://doi.org/ 10.2307/1970079.
- [90] E. P. Wigner. Characteristic vectors of bordered matrices with infinite dimensions. II. Ann. of Math. (2), 65:203-207, 1957. ISSN 0003-486X. doi: 10.2307/1969956. URL https://doi.org/ 10.2307/1969956.
- [91] G. Wolfer and A. Kontorovich. Estimating the mixing time of ergodic markov chains. In *Conference on Learning Theory*, pages 3120–3159. PMLR, 2019.
- [92] J. Ye and R. Shokri. Differentially private learning needs hidden state (or much faster convergence). Advances in Neural Information Processing Systems, 35:703–715, 2022.

- [93] D. Yin, R. Kannan, and P. Bartlett. Rademacher complexity for adversarially robust generalization. In K. Chaudhuri and R. Salakhutdinov, editors, Proceedings of the 36th International Conference on Machine Learning, volume 97 of Proceedings of Machine Learning Research, pages 7085-7094. PMLR, 09-15 Jun 2019. URL https://proceedings.mlr.press/v97/ yin19b.html.
- [94] C. Yun, S. Bhojanapalli, A. S. Rawat, S. J. Reddi, and S. Kumar. Are transformers universal approximators of sequence-to-sequence functions? arXiv preprint arXiv:1912.10077, 2019.
- [95] C. Yun, Y.-W. Chang, S. Bhojanapalli, A. S. Rawat, S. Reddi, and S. Kumar. O (n) connections are expressive enough: Universal approximability of sparse transformers. *Advances in Neural Information Processing Systems*, 33:13783–13794, 2020.
- [96] B. Zamanlooy. On the mixing times of contractive markov kernels. ArXiV, 2024.
- [97] R. Zhang, S. Frei, and P. L. Bartlett. Trained transformers learn linear models in-context. Journal of Machine Learning Research, 25(49):1–55, 2024.
- [98] W. X. Zhao, K. Zhou, J. Li, T. Tang, X. Wang, Y. Hou, Y. Min, B. Zhang, J. Zhang, Z. Dong, et al. A survey of large language models. *ArXiv preprint*, abs/2303.18223, 2023.
- [99] H. Zhou, S. Zhang, J. Peng, S. Zhang, J. Li, H. Xiong, and W. Zhang. Informer: Beyond efficient transformer for long sequence time-series forecasting. In *Proceedings of the AAAI conference on artificial intelligence*, volume 35, pages 11106–11115, 2021.
- [100] D. Zimmermann. Logarithmic Sobolev inequalities for mollified compactly supported measures. J. Funct. Anal., 265(6):1064-1083, 2013. ISSN 0022-1236,1096-0783. doi: 10.1016/j.jfa.2013.05.029. URL https://doi.org/10.1016/j.jfa.2013.05.029.