

Series Associated with Harmonic Numbers, Fibonacci Numbers and Central Binomial Coefficients $\binom{2n}{n}$

Akerele Olofin Segun¹ and Salami Olamide Esther²

Department of Mathematics, University of Ibadan, Oyo State, Nigeria.

¹email: akerelelofin@gmail.com

²email: salamiolamideesther03@gmail.com

Abstract

We find various series that involves the central binomial coefficients $\binom{2n}{n}$, harmonic numbers and Fibonacci Numbers.

Contrary to the traditional hypergeometric function ${}_pF_q$ approach, our method utilizes a straightforward transformation to obtain new evaluations linked to Fibonacci numbers and the golden ratio. Before the end of this paper, we also gave a new series representation for $\zeta(2)$.

Keywords: Central Binomial coefficients, Harmonic number, Catalan number, Fibonacci number, Lucas number.

2020 Mathematics Subject Classification: 40A05, 11B39, 11B65, 05A10.

1 Introduction

In an article published in the year 2016, Hongwei [1], gave a generating function for the sequence, $\binom{2n}{n}H_n$, $\binom{2n}{n}(H_{2n} - H_n)$, $C_n(H_{2n} - H_n)$ and few others, where C_n is the nth Catalan number, $C_n = \frac{1}{n+1} \binom{2n}{n}$. In search of interesting series associated with central binomial coefficients and Harmonic numbers, Hongwei established several interesting sums as follows;

$$\sum_{n=1}^{\infty} \frac{1}{4^n(2n+1)} \binom{2n}{n} (H_{2n} - H_n) = \pi \ln 2 - 2G, \quad (1)$$

$$\sum_{n=1}^{\infty} \frac{1}{4^n n(2n+1)} \binom{2n}{n} (H_{2n-1} - H_n) = 2 + 2 \ln 2 + \ln^2 2 + 4G - \pi(1 + 2 \ln 2), \quad (2)$$

$$\sum_{n=1}^{\infty} \frac{1}{4^n(2n+3)} C_n H_n = 2 + 4 \ln 2 - 4G - \pi + \pi \ln 2, \quad (3)$$

where G is the Catalan's constant, which is defined by

$$G := \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)^2}.$$

From [1], we have the generating function;

$$\mathcal{M}(x) := \sum_{n=1}^{\infty} \binom{2n}{n} H_n x^n = \frac{2}{\sqrt{1-4x}} \ln \left(\frac{1 + \sqrt{1-4x}}{2\sqrt{1-4x}} \right) \quad (4)$$

which converges on $[-1/4, 1/4)$.

Now before we continue, let F_n and L_n denote the n -th Fibonacci and Lucas numbers, both satisfying the recurrence relation $\Gamma_n = \Gamma_{n-1} + \Gamma_{n-2}$, $n \geq 2$, with conditions $F_0 = 0, F_1 = 1$ and $L_0 = 2, L_1 = 1$. Also $L_{-m} = (-1)^m L_m$ and $F_{-m} = (-1)^{m-1} F_m$. Through out this paper, we denote the golden ratio $\alpha = \frac{1+\sqrt{5}}{2}$ and write it's conjugate $\beta = \frac{1-\sqrt{5}}{2}$, so that $\alpha\beta = -1$ and $\alpha + \beta = 1$. We have the Binet formulas for Fibonacci and Lucas numbers to be;

$$F_m = \frac{\alpha^m - \beta^m}{\alpha - \beta}, \quad L_m = \alpha^m + \beta^m$$

for any integer m .

We will require the following, which are consequences of the Binet formula and well-known identities which are valid for integers m and n .

$$\alpha^{2m} = \alpha^m F_m \sqrt{5} - (-1)^{m+1},$$

$$\alpha^{2m} = \alpha^m L_m - (-1)^m,$$

$$\beta^{2m} = \beta^m L_m - (-1)^m,$$

$$F_n^2 + (-1)^{n+m-1} F_m^2 = F_{n-m} F_{n+m},$$

$$L_{n+m} + (-1)^m L_{n-m} = L_n L_m.$$

Now setting $x = \frac{1}{4(\alpha^{2r} + (-1)^{r+1})}$ in (4) for $r \in \mathbb{N}$ we get that;

$$\sum_{n=1}^{\infty} \frac{1}{(4\sqrt{5})^n \alpha^{rn} F_r^n} \binom{2n}{n} H_n = \frac{2\sqrt{\alpha^r F_r \sqrt{5}}}{\sqrt{\alpha^r F_r \sqrt{5}} - 1} \ln \left(\frac{\sqrt{\alpha^r F_r \sqrt{5}} + \sqrt{\alpha^r F_r \sqrt{5} - 1}}{2\sqrt{\alpha^r F_r \sqrt{5}} - 1} \right) \quad (5)$$

Evaluation at $r = 1, 2, 3$ in (5) gives;

$$\sum_{n=1}^{\infty} \frac{1}{(2^2\sqrt{5})^n \alpha^n} \binom{2n}{n} H_n = \frac{2\sqrt{\alpha\sqrt{5}}}{\sqrt{\alpha\sqrt{5}} - 1} \ln \left(\frac{\sqrt{\alpha\sqrt{5}} + \sqrt{\alpha\sqrt{5} - 1}}{2\sqrt{\alpha\sqrt{5}} - 1} \right) \quad (6)$$

$$\sum_{n=1}^{\infty} \frac{1}{(2^2\sqrt{5})^n \alpha^{2n}} \binom{2n}{n} H_n = \frac{2\alpha\sqrt[4]{5}}{\sqrt{\alpha^2\sqrt{5}-1}} \ln \left(\frac{\alpha\sqrt[4]{5} + \sqrt{\alpha^2\sqrt{5}-1}}{2\sqrt{\alpha^2\sqrt{5}-1}} \right) \quad (7)$$

$$\sum_{n=1}^{\infty} \frac{1}{(2^3\sqrt{5})^n \alpha^{3n}} \binom{2n}{n} H_n = \frac{2\sqrt{2}\sqrt{\alpha^3\sqrt{5}}}{\sqrt{2\alpha^3\sqrt{5}-1}} \ln \left(\frac{\sqrt{2\alpha^3\sqrt{5}} + \sqrt{2\alpha^3\sqrt{5}-1}}{2\sqrt{2\alpha^3\sqrt{5}-1}} \right) \quad (8)$$

Also from (5), by replacing r with $2r$ we get,

$$\sum_{n=1}^{\infty} \frac{1}{(4\sqrt{5})^n \alpha^{2rn} F_{2r}^n} \binom{2n}{n} H_n = \frac{2\alpha^r \sqrt{F_{2r}\sqrt{5}}}{\sqrt{\alpha^{2r} F_{2r}\sqrt{5}-1}} \ln \left(\frac{\alpha^r \sqrt{F_{2r}\sqrt{5}} + \sqrt{\alpha^{2r} F_{2r}\sqrt{5}-1}}{2\sqrt{\alpha^{2r} F_{2r}\sqrt{5}-1}} \right) \quad (9)$$

In this paper, by playing around the $\mathcal{M}(x)$ in (4), we shall produce more interesting results. To ensure accuracy, all formulas appearing in this paper were verified numerically by Mathematica 13.3.

2 Main Theorems

Theorem 2.1. *If r is a natural number, then*

$$\sum_{n=1}^{\infty} \frac{1}{4^n \alpha^{rn} L_r^n} \binom{2n}{n} H_n = \frac{2\sqrt{\alpha^r L_r}}{\sqrt{\alpha^r L_r - 1}} \ln \left(\frac{\sqrt{\alpha^r L_r} + \sqrt{\alpha^r L_r - 1}}{2\sqrt{\alpha^r L_r - 1}} \right) \quad (10)$$

Proof. Setting $x = \frac{1}{4(\alpha^{2r} + (-1)^r)}$ in (4) and using the second identity from the list of consequences from the Binet's formula (*introduction*). The result follows immediately. \square

Example 2.0.1. *Evaluation at $r = 1, 2, 3$ in (10), gives;*

$$\sum_{n=1}^{\infty} \frac{1}{4^n \alpha^n} \binom{2n}{n} H_n = \frac{2\sqrt{\alpha}}{\sqrt{\alpha-1}} \ln \left(\frac{\sqrt{\alpha} + \sqrt{\alpha-1}}{2\sqrt{\alpha-1}} \right), \quad (11)$$

$$\sum_{n=1}^{\infty} \frac{1}{12^n \alpha^{2n}} \binom{2n}{n} H_n = \frac{2\alpha\sqrt{3}}{\sqrt{3\alpha^2-1}} \ln \left(\frac{\alpha\sqrt{3} + \sqrt{3\alpha^2-1}}{2\sqrt{3\alpha^2-1}} \right), \quad (12)$$

$$\sum_{n=1}^{\infty} \frac{1}{16^n \alpha^{3n}} \binom{2n}{n} H_n = \frac{4\sqrt{\alpha^3}}{\sqrt{4\alpha^3-1}} \ln \left(\frac{2\sqrt{\alpha^3} + \sqrt{4\alpha^3-1}}{2\sqrt{4\alpha^3-1}} \right). \quad (13)$$

Corollary 2.0.1. *If r is a natural number, then*

$$\sum_{n=1}^{\infty} \frac{1}{4^n \alpha^{2rn} L_{2r}^n} \binom{2n}{n} H_n = \frac{2\alpha^r \sqrt{L_{2r}}}{\sqrt{\alpha^{2r} L_{2r} - 1}} \ln \left(\frac{\alpha^r \sqrt{L_{2r}} + \sqrt{\alpha^{2r} L_{2r} - 1}}{2\sqrt{\alpha^{2r} L_{2r} - 1}} \right). \quad (14)$$

Proof. Replace r with $2r$ in (10) \square

Theorem 2.2. *If r is a non-negative integer, then*

$$\sum_{n=1}^{\infty} \frac{1}{4^n \alpha^{rn} L_r^n} \binom{2n}{n} (H_{2n} - H_n) = -\frac{\sqrt{\alpha^r L_r}}{\sqrt{\alpha^r L_r} - 1} \ln \left(\frac{\sqrt{\alpha^r L_r} + \sqrt{\alpha^r L_r - 1}}{2\sqrt{\alpha^r L_r}} \right), \quad r > 0 \quad (15)$$

$$\sum_{n=1}^{\infty} \frac{1}{(4\sqrt{5})^n \alpha^{rn} F_r^n} \binom{2n}{n} (H_{2n} - H_n) = -\frac{\sqrt{\alpha^r F_r \sqrt{5}}}{\sqrt{\alpha^r F_r \sqrt{5}} - 1} \ln \left(\frac{\sqrt{\alpha^r F_r \sqrt{5}} + \sqrt{\alpha^r F_r \sqrt{5} - 1}}{2\sqrt{\alpha^r F_r \sqrt{5}}} \right), \quad r > 0 \quad (16)$$

Proof. Before we establish the following identities above, we shall establish a generating function for the sequence $\binom{2n}{n} (H_{2n} - H_n)$ in the ways of [1]. Now observe,

$$\sum_{n=1}^{\infty} \binom{2n}{n} (H_{2n} - H_n) x^n = \sum_{n=1}^{\infty} \sum_{k=1}^n \frac{1}{k} \binom{2n-k}{n} x^n = \sum_{k=1}^{\infty} \frac{1}{k} \left\{ \sum_{n=k}^{\infty} \binom{2n-k}{n} x^n \right\}$$

By setting $(n = m + k)$, we have that:

$$\begin{aligned} \sum_{k=1}^{\infty} \frac{1}{k} \left\{ \sum_{n=k}^{\infty} \binom{2n-k}{n} x^n \right\} &= \sum_{k=1}^{\infty} \sum_{m=0}^{\infty} \binom{2m+k}{m+k} x^{m+k} \\ &= \sum_{k=1}^{\infty} \frac{x^k}{k} \left\{ \sum_{m=0}^{\infty} \binom{2m+k}{m} x^m \right\} \end{aligned}$$

Since, (from [2, p.203])

$$\sum_{m=0}^{\infty} \binom{2m+k}{m} \vartheta^m = \frac{1}{\sqrt{1-4\vartheta}} \left(\frac{1 - \sqrt{1-4\vartheta}}{2\vartheta} \right)^k, \quad |\vartheta| < 1/4$$

Thus, we have that

$$\sum_{n=1}^{\infty} \binom{2n}{n} (H_{2n} - H_n) x^n = \frac{1}{\sqrt{1-4x}} \sum_{k=1}^{\infty} \frac{1}{k} \left(\frac{1 - \sqrt{1-4x}}{2} \right)^k = -\frac{1}{\sqrt{1-4x}} \ln \left(\frac{1 - \sqrt{1-4x}}{2} \right) \quad (17)$$

Now set $x = \frac{1}{4(\alpha^{2r} + (-1)^{r+1})}$ and $x = \frac{1}{4(\alpha^{2r} + (-1)^r)}$ in (17), then (16) and (15) follows directly. \square

Example 2.0.2. *Evaluation at $r = 1, 2, 3$ in (15) and (16), respectively, gives*

$$\sum_{n=1}^{\infty} \frac{1}{4^n \alpha^n} \binom{2n}{n} (H_{2n} - H_n) = -\frac{\sqrt{\alpha}}{\sqrt{\alpha} - 1} \ln \left(\frac{\sqrt{\alpha} + \sqrt{\alpha - 1}}{2\sqrt{\alpha}} \right) \quad (18)$$

$$\sum_{n=1}^{\infty} \frac{1}{12^n \alpha^{2n}} \binom{2n}{n} (H_{2n} - H_n) = -\frac{\alpha\sqrt{3}}{\sqrt{3\alpha^2 - 1}} \ln \left(\frac{\alpha\sqrt{3} + \sqrt{3\alpha^2 - 1}}{2\alpha\sqrt{3}} \right) \quad (19)$$

$$\sum_{n=1}^{\infty} \frac{1}{16^n \alpha^{3n}} \binom{2n}{n} (H_{2n} - H_n) = -\frac{2\sqrt{\alpha^3}}{\sqrt{4\alpha^3 - 1}} \ln \left(\frac{2\sqrt{\alpha^3} + \sqrt{4\alpha^3 - 1}}{4\sqrt{\alpha^3}} \right) \quad (20)$$

$$\sum_{n=1}^{\infty} \frac{1}{(4\sqrt{5})^n \alpha^n} \binom{2n}{n} (H_{2n} - H_n) = -\frac{\sqrt{\alpha\sqrt{5}}}{\sqrt{\alpha\sqrt{5} - 1}} \ln \left(\frac{\sqrt{\alpha\sqrt{5}} + \sqrt{\alpha\sqrt{5} - 1}}{2\sqrt{\alpha\sqrt{5}}} \right) \quad (21)$$

$$\sum_{n=1}^{\infty} \frac{1}{(4\sqrt{5})^n \alpha^{2n}} \binom{2n}{n} (H_{2n} - H_n) = -\frac{\alpha^{\sqrt[4]{5}}}{\sqrt{\alpha^2\sqrt{5} - 1}} \ln \left(\frac{\alpha^{\sqrt[4]{5}} + \sqrt{\alpha^2\sqrt{5} - 1}}{2\alpha^{\sqrt[4]{5}}} \right) \quad (22)$$

$$\sum_{n=1}^{\infty} \frac{1}{(8\sqrt{5})^n \alpha^{3n}} \binom{2n}{n} (H_{2n} - H_n) = -\frac{\sqrt{2}\sqrt{\alpha^3\sqrt{5}}}{\sqrt{\alpha^3 2\sqrt{5} - 1}} \ln \left(\frac{\sqrt{2}\sqrt{\alpha^3\sqrt{5}} + \sqrt{\alpha^3 2\sqrt{5} - 1}}{2\sqrt{2}\sqrt{\alpha^3\sqrt{5}}} \right) \quad (23)$$

Theorem 2.3.

$$\sum_{n=1}^{\infty} \frac{C_n(H_{2n} - \frac{1}{2}H_n)}{4^n(2n+1)} \left(\frac{\pi}{2} - \frac{(2n)!!}{(2n+1)!!} \right) = 2 \ln(2) + \frac{7}{8}\zeta(3) + \frac{\pi}{12}(-12 + \pi(-1 + \ln(8)))$$

where C_n is the n -th Catalan's number and ,

$$n!! := \prod_{k=0}^{\lceil \frac{n}{2} \rceil - 1} (n - 2k) := \begin{cases} \prod_{k=1}^{\frac{n}{2}} (2k), & \text{if } n \text{ is even,} \\ \prod_{k=1}^{\frac{n+1}{2}} (2k - 1), & \text{if } n \text{ is odd.} \end{cases}$$

Proof. From [1] we have that;

$$\sum_{n=1}^{\infty} \binom{2n}{n} (H_{2n} - \frac{1}{2}H_n) x^n = -\frac{1}{\sqrt{1-4x}} \ln \sqrt{1-4x}$$

Immediately we see that;

$$\sum_{n=1}^{\infty} C_n (H_{2n} - \frac{1}{2}H_n) x^n = \frac{1}{2x} (1 - \sqrt{1-4x} + \sqrt{1-4x} \ln \sqrt{1-4x}), \quad |x| < 1/4$$

For the interesting part, set $x = \frac{1}{4} \sin^2 t$ for $t \in (-\pi/2, \pi/2)$. Then we have that;

$$\sum_{n=1}^{\infty} \frac{C_n (H_{2n} - \frac{1}{2}H_n)}{4^n} \sin^{2n} t = \frac{2}{\sin^2 t} (1 - \cos t + \cos t \ln \cos t). \quad (24)$$

Now multiply (24) by $t \cos t$ and integrating both sides from 0 to $\pi/2$. We have;

$$\sum_{n=1}^{\infty} \frac{C_n (H_{2n} - \frac{1}{2}H_n)}{4^n} \int_0^{\pi/2} t \cos t \sin^{2n} t dt = 2 \int_0^{\pi/2} \frac{t \cos t}{\sin^2 t} (1 - \cos t + \cos t \ln \cos t) dt$$

Thus, we evaluated the integral on the right hand side to $\ln 2 + \frac{\pi}{24}(-12 + \pi(-1 + \ln 8)) + \frac{7}{16}\zeta(3)$ and using integration by part for the integral on the left we have that;

$$\int_0^{\frac{\pi}{2}} \frac{t \cos t}{\sin^2 t} (1 - \cos t + \cos t \ln \cos t) dt = \ln 2 + \frac{\pi}{24}(-12 + \pi(-1 + \ln 8)) + \frac{7}{16}\zeta(3),$$

$$\int_0^{\frac{\pi}{2}} t \cos t \sin^{2n} t dt = \frac{1}{2n+1} \left(\frac{\pi}{2} - \frac{(2n)!!}{(2n+1)!!} \right).$$

So the result follows immediately. \square

Theorem 2.4. (Two Ramanujan-Like Series)

$$\sum_{n=1}^{\infty} \frac{C_n(H_{2n} - H_n)}{4^{2n}} \binom{2n+2}{n+1} = \frac{16}{\pi} \psi,$$

where $\psi = 2G + \pi - 2 - \ln 2 - \pi \ln 2$.

$$\sum_{n=1}^{\infty} \frac{C_n H_{2n}}{16^n} \binom{2n}{n} = \frac{2}{\pi} \psi^*,$$

where $\psi^* = 2 + \pi - 2 \ln 8$.

Proof. Integrating both sides of (17) with respect to x , we obtain the generating function for the sequence $C_n(H_{2n} - H_n)$ as follows;

$$\sum_{n=1}^{\infty} C_n(H_{2n} - H_n)x^n = \frac{1}{2x} \left[(1 - \sqrt{1-4x}) + (1 + \sqrt{1-4x}) \ln \left(\frac{1 + \sqrt{1-4x}}{2} \right) \right] \quad (25)$$

Likewise, from [1] we have that;

$$\sum_{n=1}^{\infty} \binom{2n}{n} H_{2n} x^n = \frac{1}{\sqrt{1-4x}} \left[\ln \left(\frac{1 + \sqrt{1-4x}}{2} \right) - 2 \ln \sqrt{1-4x} \right] \quad (26)$$

For this reason, integrating both sides of (26) we have that,

$$\sum_{n=1}^{\infty} C_n H_{2n} x^n = \frac{1}{2x} [(1 - \sqrt{1-4x}) - (1 + \sqrt{1-4x}) \ln(1 + \sqrt{1-4x}) + \ln 2 + \sqrt{1-4x} \ln(2-8x)] \quad (27)$$

Now set $x = \frac{\sin^2 t}{4}$ in (25) and (27) and integrating both sides with respect to x from 0 to $\pi/2$, while using the following results;

$$\int_0^{\frac{\pi}{2}} \left\{ (1 - \cos t) + (1 + \cos t) \ln \left(\frac{1 + \cos t}{2} \right) \right\} dt = 2G + \pi - 2 - \ln 2 - \pi \ln 2,$$

$$\int_0^{\frac{\pi}{2}} \frac{2}{\sin^2 t} [(1 - \cos t) - (1 + \cos t) \ln(1 + \cos t) + \ln 2 + \cos t \ln(2 \cos^2 t)] dt = 2 + \pi - 2 \ln 8,$$

and,

$$\int_0^{\frac{\pi}{2}} \sin^{2n} t \, dt = \frac{\pi}{2} \frac{(2n)!}{4^n (n!)^2}.$$

Then the series follows directly. \square

Theorem 2.5 (A new series representation for $\zeta(2)$).

$$\zeta(2) = \sum_{n=1}^{\infty} \frac{1024n}{3(2n-1)^2(2n+1)(2n+3)^2} \frac{\binom{2n}{n}}{\binom{2n+2}{n+1}}$$

where $\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$, provided $\Re(s) > 0$.

Proof. From [5], we have that;

$$\sum_{n=1}^{\infty} \frac{nx^{2n}}{4^n(2n-1)^2(2n+1)} \binom{2n}{n} = \frac{1}{8} \left(\sqrt{1-x^2} + 2x \sin^{-1} x - \frac{\sin^{-1} x}{x} \right). \quad (28)$$

Next we multiply (28) by x^2 and integrate both sides with respect to x to get;

$$\sum_{n=1}^{\infty} \frac{nx^{2n+3}}{4^n(2n-1)^2(2n+1)(2n+3)} \binom{2n}{n} = \frac{(8x^4 - 8x^2 + 3) \sin^{-1} x + \sqrt{1-x^2}(6x^3 - 3x)}{128} \quad (29)$$

which converges for $x \in [-1, 1]$. Now, we set $x = \sin t$ in (29) and integrate over the interval 0 to $\pi/2$. The result follows. \square

Theorem 2.6 (A Ramanujan-Like Series involving the ratio G/π).

$$\sum_{n=1}^{\infty} \frac{n^2}{16^n(2n-1)^2(2n+1)} \binom{2n}{n}^2 = \frac{G}{4\pi} + \frac{1}{8\pi}$$

Proof. We begin by differentiating (28) with respect to x to get,

$$\sum_{n=1}^{\infty} \frac{2n^2 x^{2n-1}}{4^n(2n-1)^2(2n+1)} \binom{2n}{n} = \frac{1}{8x^2} \left((2x^2 + 1) \sin^{-1} x - x\sqrt{1-x^2} \right). \quad (30)$$

Now multiply (30) by x and set $x = \sin t$, then we integrate both sides over the interval 0 to $\pi/2$. Finally, using the result;

$$\int_0^{\frac{\pi}{2}} \frac{t(2 \sin^2 t + 1) - \sin t \cos t}{\sin t} dt = \int_0^{\frac{\pi}{2}} (2t \sin t - \cos t) dt + \int_0^{\frac{\pi}{2}} \frac{t}{\sin t} dt = 1 + 2G$$

The Series follows immediately. ¹ \square

¹Now we look at some other interesting Series

3 Some Deluxe Series

Observe that the series in (25) converges on $[-1/4, 1/4)$. Setting $x = -1/8, 1/16$ and $-1/16$ respectively, we obtain the following series;

$$\sum_{n=1}^{\infty} \frac{(-1)^n C_n}{8^n} (H_{2n} - H_n) = -\frac{4}{\sqrt{2}} \left[(\sqrt{2} - \sqrt{3}) + (\sqrt{2} + \sqrt{3}) \ln \left(\frac{\sqrt{2} + \sqrt{3}}{2\sqrt{2}} \right) \right], \quad (31)$$

$$\sum_{n=1}^{\infty} \frac{C_n}{16^n} (H_{2n} - H_n) = 4 \left[(2 - \sqrt{3}) + (2 + \sqrt{3}) \ln \left(\frac{2 + \sqrt{3}}{4} \right) \right], \quad (32)$$

$$\sum_{n=1}^{\infty} \frac{(-1)^n C_n}{16^n} (H_{2n} - H_n) = -4 \left[(2 - \sqrt{5}) + (2 + \sqrt{5}) \ln \left(\frac{2 + \sqrt{5}}{4} \right) \right]. \quad (33)$$

In the same light, recall from (29) converges on $[-1, 1]$. Setting $x = 1$ and -1 respectively, we get two interesting series;

$$\sum_{n=1}^{\infty} \frac{n}{4^n (2n-1)^2 (2n+1) (2n+3)} \binom{2n}{n} = \frac{3\pi}{256}, \quad (34)$$

$$\sum_{n=1}^{\infty} \frac{(-1)^{2n+3n}}{4^n (2n-1)^2 (2n+1) (2n+3)} \binom{2n}{n} = \frac{-3\pi}{256}. \quad (35)$$

From (30) we have;

$$\sum_{n=1}^{\infty} \frac{n^2}{4^n (2n-1)^2 (2n+1)} \binom{2n}{n} = \frac{3\pi}{32}. \quad (36)$$

Also, from (17) we get that;

$$\sum_{n=1}^{\infty} \frac{H_{2n} - H_n}{8^n} \binom{2n}{n} = -\sqrt{2} \ln \left(\frac{\sqrt{2} - 1}{2\sqrt{2}} \right), \quad (37)$$

$$\sum_{n=1}^{\infty} \frac{H_{2n} - H_n}{16^n} \binom{2n}{n} = -\frac{2}{\sqrt{3}} \ln \left(\frac{2 - \sqrt{3}}{2\sqrt{2}} \right), \quad (38)$$

$$\sum_{n=1}^{\infty} \frac{C_n}{8^n} \left(H_{2n} - \frac{1}{2} H_n \right) = 4 \left(1 - \frac{1}{\sqrt{2}} - \frac{\ln 2}{2\sqrt{2}} \right), \quad (39)$$

$$\sum_{n=1}^{\infty} \frac{C_n}{16^n} \left(H_{2n} - \frac{1}{2} H_n \right) = 8 \left(1 - \frac{\sqrt{3}}{2} + \frac{\sqrt{3}}{2} \ln \frac{\sqrt{3}}{2} \right). \quad (40)$$

4 Conclusion

1. To assure accuracy of the results, we verified all the series via *Mathematica 13.3*.
2. In this paper we presented new closed forms for some types of series involving the Cen-

tral Binomial coefficients $\binom{2n}{n}$. To prove our results, we used some generating functions, combined with basic differentiation and integration. Using similar techniques, we established series evaluations involving Harmonic numbers with Fibonacci and Lucas sequences. In addition, readers can exploit (28),(29) and (30) to generate more exotic series.

5 Acknowledgment

We would like to appreciate Prof. U. N. Bassey, Dr. D. A. Dikko and Mr. G. S. Lawal from the Department of Mathematics at the University of Ibadan for their motivations. This helped improve and enrich this article.

References

- [1] H. Chen, Interesting Series Associated with Central Binomial Coefficients, Catalan Numbers and Harmonic Numbers. *Journal of Integer Sequences (2016)*, Article 16.1.5.
- [2] R. Graham, D. Knuth and O. Patashnik, *Concrete Mathematics, 2nd Edition*, Addison Wesley, 1994.
- [3] K. Adegoke, R. Frontczak and T. Goy, On some series involving the binomial coefficients $C(3n, n)$. *Notes on Number Theory and Discrete Mathematics (2024)*, Volume 30, Number 2, 319-334.
- [4] H. Chen, Interesting Ramanujan-Like Series Associated with Powers of Central Binomial coefficients. *Journal of Integer Sequences (2022)*, Article 22.1.8.
- [5] Narendra Bhandari, Binomial Sums resulting to the ratios, G/π and $1/\pi$. *Archimede Mathematical Journal (2022)*, Volume 9, Number 1.