

LOG-NOETHERIAN FUNCTIONS

GAL BINYAMINI

ABSTRACT. We introduce the class of *Log-Noetherian* (LN) functions. These are holomorphic solutions to algebraic differential equations (in several variables) with logarithmic singularities. We prove an upper bound on the number of solutions for systems of LN equations, resolving in particular Khovanskii's conjecture for Noetherian functions. Consequently, we show that the structure \mathbb{R}_{LN} generated by LN-functions, as well as its expansion $\mathbb{R}_{\text{LN,exp}}$, are effectively o-minimal: definable sets in these structures admit effective bounds on their complexity in terms of the complexity of the defining formulas.

We show that $\mathbb{R}_{\text{LN,exp}}$ contains the horizontal sections of regular flat connections with quasiunipotent monodromy over algebraic varieties. It therefore contains the universal covers of Shimura varieties and period maps of polarized variations of \mathbb{Z} -Hodge structures. We also give an effective Pila-Wilkie theorem for $\mathbb{R}_{\text{LN,exp}}$ -definable sets. Thus $\mathbb{R}_{\text{LN,exp}}$ can be used as an effective variant of $\mathbb{R}_{\text{an,exp}}$ in the various applications of o-minimality to arithmetic geometry and Hodge theory.

CONTENTS

1. Introduction	2
2. Log-Noetherian cells and functions	8
3. Background from commutative algebra and differential equations	11
4. Basic theory of LN functions	14
5. Proof of the CPT for the LN category	23
6. Effective model completeness and o-minimality of \mathbb{R}_{LN}	29
7. Effective o-minimality of $\mathbb{R}_{\text{LN,PF}}$	33
8. Geometric constructions with LN cells	38
9. Effective Pila-Wilkie in $\mathbb{R}_{\text{LN,PF}}$	40
10. LN-functions and regular flat connections	41
Appendix A. Noetherian functions over number fields	47
References	49

Date: May 28, 2024.

2020 Mathematics Subject Classification. 03C64, 58A17, 14D07, 14Q201.

Key words and phrases. logic, o-minimality, effectivity, differential equations.

Funded by the European Union (ERC, SharpOS, 101087910), and by the ISRAEL SCIENCE FOUNDATION (grant No. 2067/23) and by the Shimon and Golde Picker - Weizmann Annual Grant.

1. INTRODUCTION

1.1. Synopsis. We introduce the class of *Log-Noetherian* (LN) functions, which are holomorphic solutions for systems of algebraic differential equations with logarithmic singularities. This class strictly contains the *Noetherian* functions considered by Khovanskii (which is the non-singular case).

We show that the structure \mathbb{R}_{LN} generated by LN-functions is an effectively o-minimal reduct of \mathbb{R}_{an} . This means that one can associate a *format* \mathcal{F} to each formula ψ in the first-order language of \mathbb{R}_{LN} , and replace the general finiteness results of o-minimality by an effective bound depending on \mathcal{F} . For instance this implies a Bezout-style theorem bounding the number of isolated points in the set defined by ψ in terms of its format. The format \mathcal{F} generally encodes the degrees and magnitude of the coefficients of the algebraic differential equations defining the LN-functions appearing in ψ . We show moreover that the structure generated by graphs of unrestricted Pfaffian functions over \mathbb{R}_{LN} , denoted $\mathbb{R}_{\text{LN,PF}}$ is effectively o-minimal as well. In particular $\mathbb{R}_{\text{LN,exp}}$ is an effectively o-minimal reduct of $\mathbb{R}_{\text{an,exp}}$.

The key to constructing these new effectively o-minimal structures is a solution to Khovanskii's conjecture from the early eighties on effective bounds for the number of solutions of systems of Noetherian equations (and its generalization to the LN class). The proof of this result is based on a version of the cellular parametrization theorem of [10] for the LN-category. It is crucial for the inductive proof of this result to consider log-Noetherian functions and cells even if one is originally only interested in Khovanskii's conjecture for non-singular Noetherian systems.

We show that horizontal sections of regular flat connections with quasiunipotent monodromy over algebraic varieties (with appropriate branch cuts) are definable in $\mathbb{R}_{\text{LN,exp}}$. In particular we show that the universal covering map of the Siegel modular variety \mathcal{A}_g and the universal abelian scheme $\mathbb{A}_g \rightarrow \mathcal{A}_g$, and the period maps for polarized variations of \mathbb{Z} -Hodge structures (PVHS), are all definable in $\mathbb{R}_{\text{LN,exp}}$. Thus all applications of o-minimality to these areas can in principle be made effective using the theory of this structure. In particular we give an effective Pila-Wilkie theorem (a straightforward generalization of [5] from the Pfaffian context) and a bound for the degree of the Hodge locus (a straightforward consequence of the approach of [1]).

1.2. Noetherian functions and Khovanskii's conjecture. Let $\mathcal{P} \subset \mathbb{C}^n$ be a product of discs in \mathbb{C}^n or intervals in \mathbb{R}^n and $f_1, \dots, f_N : \mathcal{P} \rightarrow \mathbb{C}$ a collection of analytic functions satisfying a system of polynomial differential equations

$$\frac{\partial F_i}{\partial \mathbf{z}_j} = G_{i,j}(F_1, \dots, F_N). \quad (1)$$

In [32] Khovanskii studied the situation where $\mathcal{P} \subset \mathbb{R}^n$, and the equations (1) are *triangular* in the sense that $G_{i,j}$ depends only on F_1, \dots, F_i . Functions satisfying this condition are called a *Pfaffian chain*, and the maximum among the degrees of $G_{i,j}$ is called the degree of the chain. A polynomial combination $G(F_1, \dots, F_N)$ of these functions is called a *Pfaffian function* of degree $\deg G$. Khovanskii proved that the number of connected components of a set defined by a system of Pfaffian equations can be bounded in terms of n, N and the degrees of the chain and the functions. This holds even if some of the intervals defining \mathcal{P} are unbounded.

Without the triangularity condition on (1) the chain F_1, \dots, F_N is called a *Noetherian chain* and their polynomial combinations *Noetherian functions*. This terminology is due to Tougeron [44]. Khovanskii's global bounds on the number of solutions certainly do not generalize to the Noetherian context: for instance, $\sin(x)$ is clearly Noetherian on \mathbb{R} and admits infinitely many isolated roots. Nevertheless Khovanskii conjectured in the early eighties that non-global bounds – restricting either to local germs or to compact domains, should still hold in this class. One instance of this conjecture appeared in [27].

Conjecture 1. *Let $P_1(\mathbf{z}, \varepsilon), \dots, P_n(\mathbf{z}, \varepsilon) : (\mathbb{C}^{n+1}, 0) \rightarrow \mathbb{C}$ be holomorphic functions depending analytically on ε , such that for each fixed ε they are Noetherian of degree at most D over a chain F_1, \dots, F_N of degree D' . Then there exists $\delta > 0$ such that for any sufficiently small ε , the number of isolated points in*

$$\{\mathbf{z} \in \mathbb{C}^n : \|\mathbf{z}\| \leq \delta, \quad P_1(\mathbf{z}, \varepsilon) = \dots = P_n(\mathbf{z}, \varepsilon) = 0\} \quad (2)$$

is bounded by some explicit function of N, n and $D + D'$.

In the *complex Pfaffian* case Conjecture 1 was proved by Gabrielov [19], giving a local complex analog of Khovanskii's (real) Pfaffian theory [32]. In the case $n = 1$ the conjecture is proved in [25]. In [27] the general conjecture was proven under the additional assumption that the intersection for $\varepsilon = 0$ is proper (in which case the number of common roots in any deformation as above is bounded by the local intersection number at the origin). Some result on improper intersections are given in [7, 8] under addition assumptions on the deformation, but Conjecture 1 has remained open in general.

A stronger form of Conjecture 1 is to require an explicit bound on the number of isolated zeros in the domain \mathcal{P} . In this case it is clear that the bounds must also involve the magnitude of the coefficients of the equations $G_{i,j}$ in the Noetherian chain (1), as evidenced by the Noetherian function $\sin(Mx)$ with $M \gg 1$. Accordingly, define the *format* \mathcal{F} of a Noetherian chain (F_1, \dots, F_N) and Noetherian function $F := G(F_1, \dots, F_N)$ by

$$\begin{aligned} \mathcal{F}(F_1, \dots, F_N) &:= n + N + \sum_{i,j} \deg G_{i,j} + \|G_{i,j}\| + \max_{\substack{i=1, \dots, N \\ \mathbf{z} \in \mathcal{P}}} |F_i(\mathbf{z})| \\ \mathcal{F}(F) &:= \mathcal{F}(F_1, \dots, F_N) + \deg G + \|G\| \end{aligned} \quad (3)$$

where the norm of a polynomial is given by the sum of the absolute values of its coefficients. With this definition, Khovanskii's conjecture is as follows.

Conjecture 2. *Let $P_1, \dots, P_n : \mathcal{P} \rightarrow \mathbb{C}$ be Noetherian of format \mathcal{F} . Then the number of isolated points in*

$$\{\mathbf{z} \in \mathcal{P} : P_1(\mathbf{z}) = \dots = P_n(\mathbf{z}) = 0\} \quad (4)$$

is bounded by some explicit function of \mathcal{F} .

To see that Conjecture 2 implies Conjecture 1, note that by rescaling the time variable $\tilde{\mathbf{z}} = R\mathbf{z}$ and the chain functions $\tilde{F}_i = F/r$ with $R \gg r \gg 0$ one can arrange that $\|G_{i,j}\|$, $|F_i(\mathbf{z})|$ and $\|G\|$ are all bounded by 1 when considered on the unit polydisc $\tilde{\mathcal{P}}$. One thus obtains an effective bound depending only on $n, N, \deg G_{i,j}$ and $\deg G$ on the isolated points in a ball whose radius δ (in the original \mathbf{z} -coordinates) depends effectively on \mathcal{F} . Conjecture 1 by comparison asserts the same bound without control on δ .

Conjecture 2 has been resolved in the case $n = 1$ by Novikov and Yakovenko [35]. Their approach is based on a combination of commutative algebraic techniques and results from the theory of analytic ODEs (in one variable). We make extensive use of their results in our treatment of the higher dimensional case. To our knowledge no higher dimensional case of Conjecture 2 has been known prior to this work. Some partial results are given in [9, 4] but with bounds degenerating to infinity near the locus of non-proper intersection. It has generally been clear that Conjectures 1 and 2 are the main obstacle toward developing a theory parallel to Khovanskii's Pfaffian theory in the Noetherian context, as the results of the present paper demonstrate.

1.3. Log-Noetherian functions (a simplified version). In his ICM talk Khovanskii listed as an open problem [33, Section 14, Problem 4] the goal of extending the Pfaffian class while retaining the effective bounds, and more specifically [33, Section 14, Problem 5] the goal of extending the effective bounds to the period integrals of algebraic forms over algebraic families. Period integrals are known to satisfy systems of algebraic differential equations (classically known as Picard-Fuchs system, and in greater generality Gauss-Manin connections) which do not seem to be Pfaffian. This is one of the main motivations for considering the more general class of Noetherian functions. It should be noted however that Gauss-Manin connections generally exhibit (regular) singularities, and the Noetherian class only describes their behavior away from the singular locus. Toward this end we enlarge the class of Noetherian functions to allow certain singularities.

Let $\mathcal{P}^\circ \subset \mathbb{C}^n$ denote the product of n punctured discs of radius 1. An LN-chain over \mathcal{P}° is a system analogous to (1) but allowing for logarithmic-type singularities along the divisors $\mathbf{z}_j = 0$, namely

$$\mathbf{z}_j \frac{\partial F_i}{\partial \mathbf{z}_j} = G_{i,j}(F_1, \dots, F_N). \quad (5)$$

We still assume that the functions F_j are holomorphic and bounded (and thus extend holomorphically over the punctures). Under these conditions we call the sequence F_1, \dots, F_N an LN-chain on \mathcal{P}° . More generally we define LN-chains on more complicated cellular domains, but we postpone this definition to §2.1. We resolve Khovanskii's conjecture in the stronger sense of Conjecture 2 for this wider class of functions. For domains \mathcal{P}° as above the result can be stated as follows.

Theorem. *Let $P_1, \dots, P_n : \mathcal{P}^\circ \rightarrow \mathbb{C}$ be LN-functions of format \mathcal{F} . Then the number of isolated points in*

$$\{\mathbf{z} \in \mathcal{P}^\circ : P_1(\mathbf{z}) = \dots = P_n(\mathbf{z}) = 0\} \quad (6)$$

is bounded by some explicit function of \mathcal{F} .

A much more general form of this result is described in §1.6. The proof of this theorem is based on a cellular parametrization theorem (CPT) in the sense of [10] for the LN-category. This can be seen as a type of local desingularization result for sets defined using LN functions. Since the precise statement of the CPT requires some technical setup we postpone it to §2.7.

1.4. Notation for effectivity. When speaking of *effective bounds* in this paper, we mean bounds that are given by primitive-recursive functions that can be fully written out explicitly in principle. We use the notation $\text{Eff}(\dots)$ to denote such

a primitive recursive function, which may be different for each occurrence of this notation.

We define the *format* of various objects such as LN-functions, LN-cells, definable sets in \mathbb{R}_{LN} , etc. To make the notations less cumbersome, when X is one of these objects we sometimes write $\text{Eff}(X)$ to mean $\text{Eff}(\mathcal{F})$ where $\mathcal{F} \in \mathbb{N}$ is the format associated to X .

1.5. Effectively o-minimal structures. Let \mathcal{S} be an o-minimal structure, viewed as a collection of definable subsets of \mathbb{R}^n for $n \in \mathbb{N}$. An *effectively o-minimal structure* on \mathcal{S} consists of a collection $\{\Omega_{\mathcal{F}} \subset \mathcal{S}\}_{\mathcal{F} \in \mathbb{N}}$ called the *format filtration* and a primitive-recursive function $\mathcal{E} : \mathbb{N} \rightarrow \mathbb{N}$ such that

- (1) The filtration is increasing $\Omega_{\mathcal{F}} \subset \Omega_{\mathcal{F}+1}$ and exhaustive $\cup_{\mathcal{F}} \Omega_{\mathcal{F}} = \mathcal{S}$.
- (2) For every $A, B \subset \mathbb{R}^n$ with $A, B \in \Omega_{\mathcal{F}}$, we have

$$A \cup B, A \cap B, \mathbb{R}^n \setminus A, A \times B, \pi_k^n(A) \in \Omega_{\mathcal{F}+1} \quad (7)$$

where $\pi_k^n : \mathbb{R}^n \rightarrow \mathbb{R}^k$ denotes the projection to the first k coordinates.

- (3) If $A \subset \mathbb{R}$ and $A \in \Omega_{\mathcal{F}}$ then A has at most $\mathcal{E}(\mathcal{F})$ connected components.

When we say that a structure is effectively o-minimal in this paper we always mean that the function \mathcal{E} can be explicitly presented in principle, although for brevity we do not compute it exactly. We will usually treat \mathcal{E} as implicit and speak simply of an effectively o-minimal structure (\mathcal{S}, Ω) .

When a structure is effectively o-minimal one has effective variants of all the standard theorems of o-minimality such as cell-decomposition, existence of stratification, definable triangulation, etc. We state the effective cell decomposition theorem as a representative example. The proof follows from the standard proof of cell decomposition verbatim.

Theorem 1 (Effective cell decomposition). *Let $X_1, \dots, X_k \subset \mathbb{R}^n$ be definable subsets. Then there exists a cylindrical decomposition $\mathbb{R}^n = \cup_{\alpha} C_{\alpha}$ compatible with X_1, \dots, X_k such that*

$$\#\{C_{\alpha}\} < \text{Eff}(X_1, \dots, X_k), \quad \forall \alpha : \mathcal{F}(C_{\alpha}) < \text{Eff}(X_1, \dots, X_k). \quad (8)$$

The notion of effective o-minimality is defined analogously to *sharp o-minimality* ($\#$ o-minimality) introduced recently in [11], see also [6, 13]. However unlike in the case of $\#$ o-minimality, here we do not require polynomial growth with respect to a separate *degree* parameter and are content with asserting some explicit bound depending on the format. If $\Omega_{\mathcal{F}, D}$ is a $\#$ o-minimal filtration then $\Omega_{\mathcal{F}, \mathcal{F}}$ is essentially an effectively o-minimal filtration (up to some minor changing of the indices). Berarducci and Servi have considered in [2] a similar notion of effective o-minimality. Our formalism is slightly different but up to some technical differences it amounts to roughly the same concept.

Remark 3. *In $\#$ o-minimality, since the filtration $\Omega_{\mathcal{F}, D}$ involves two indices, there is no canonical way of associating a single format and degree to a given definable set. In effective o-minimality on the other hand one can define the format of a definable set $A \in \mathcal{S}$ to be $\min\{\mathcal{F} : A \in \Omega_{\mathcal{F}}\}$.*

If some collection of sets $\{A_{\alpha} \in \mathcal{S}\}$ generates the structure \mathcal{S} and one assigns a format to each A_{α} , then one can always define a filtration Ω generated by these sets by taking transitive closure under axiom 2 above. The statement that the resulting

structure is effectively o-minimal then means that axiom 3 holds for some suitable choice of \mathcal{E} .

1.6. The structures \mathbb{R}_{LN} and $\mathbb{R}_{\text{LN,exp}}$. Turning back to LN-function, we define the structure \mathbb{R}_{LN} generated by the graphs of LN-functions (for the precise definition see §6). We associate to the graph of each LN-function the format of the function as defined above and generate a format-filtration.

Theorem 2. *The structure \mathbb{R}_{LN} is effectively o-minimal and effectively model-complete.*

Theorem 2 is proved in §6. Model completeness is the statement that every definable set in the structure can be expressed using a purely existential formula. Together with o-minimality it is one of the hallmarks of tame geometry, established for \mathbb{R}_{an} by Gabrielov [24] and for \mathbb{R}_{exp} by Wilkie [47]. Effective model completeness here means that the complexity of the existential formula can be effectively controlled in terms of the complexity of the original formula. We also prove an effective Lojasiewicz inequality for \mathbb{R}_{LN} .

With Theorem 2 in hand, we proceed to show that Khovanskii's theory of Pfaffian functions can be carried out over the structure \mathbb{R}_{LN} . That is, one can consider Pfaffian systems of equations with coefficients in \mathbb{R}_{LN} , and add the graphs of the resulting functions to obtain a larger structure $\mathbb{R}_{\text{LN,PF}}$. As a consequence we prove the following.

Theorem 3. *The structure $\mathbb{R}_{\text{LN,PF}}$ is effectively o-minimal.*

Theorem 3 is proved in §7. As a special case of particular interest for the applications, it follows that the structure $\mathbb{R}_{\text{LN,exp}}$ is effectively o-minimal. We note that this type of "Pfaffian closure" is well known in o-minimality (e.g. by Wilkie [48] and Speissegger [43]), and the effective nature of the construction has also been studied by Berarducci-Servi [2] and Gabrielov-Vorobjov [23]. We give the construction using their different approaches in §7.

1.7. Abelian integrals and the Varchenko-Khovanskii theorem. In the specific case of abelian integrals

$$I(t) := \oint_{\delta(t)} \omega, \quad \text{where } \begin{cases} \omega = P dx + Q dy \\ H \in \mathbb{R}[x, y] \\ \delta(t) \in H_1(\{H = t\}, \mathbb{Z}) \end{cases} \quad (9)$$

the problem of finding an explicit upper bound for the number of zeros of $I(t)$ depending only on $\deg H, \deg \omega$ is known as the *infinitesimal Hilbert sixteenth problem* because of its relation to the limit cycles of a perturbed Hamiltonian differential equation

$$dH + \varepsilon\omega = 0, \quad \varepsilon \ll 1. \quad (10)$$

It can be shown that when the perturbation is non-conservative, the zeros of $I(t)$ correspond to the limit cycles of the perturbed system (10).

The infinitesimal Hilbert problem has been resolved in [12], and we refer the reader to this paper for a review of the long history of this problem. Prior to this effective solution, a uniform (but ineffective) bound depending only on the $\deg H, \deg \omega$ has been obtained by Varchenko and Khovanskii in the papers [46, 30]. Their proof predates modern o-minimality, but can be described as showing that

the period integrals of algebraic families lie in the structure $\mathbb{R}_{\text{an,PF}}$ generated by Pfaffian functions over the globally subanalytic structure.

Going beyond the original infinitesimal Hilbert problem, Varchenko [46, Section 4] also proves a “Bezout” theorem for period integrals, showing that systems of equations in several variables involving period integrals over algebraic families of arbitrary dimension admit finite and uniform bounds for the number of isolated solutions. No effective extension of [12] to this multivariable context has been known.

1.8. Periods and other examples of LN functions. In §10 we show that horizontal sections of regular flat connections with quasiunipotent monodromy over algebraic varieties lie in $\mathbb{R}_{\text{LN,exp}}$ (after making appropriate branch cuts). Applying this result to the Gauss-Manin connection of an algebraic family we obtain an effective form of the Varchenko-Khovanskii theorem from the effective o-minimality of $\mathbb{R}_{\text{LN,exp}}$. This also implies that covering maps of Shimura varieties and period maps for PVHS are definable in $\mathbb{R}_{\text{LN,exp}}$ in the sense of [1].

More specifically, suppose that S is a quasi-projective variety and $\Phi : S \rightarrow D/\Gamma$ is a period map associated to a PVHS (see §10.3 for details). If $Y \subset D/\Gamma$ is a special subvariety then a theorem by Cattani, Deligne and Kaplan [16] states that $S_Y := \Phi^{-1}(Y)$ is algebraic. An alternative proof of this theorem has been given in [1]. Recall that [1, Theorem 1.1] introduces an \mathbb{R}_{alg} -structure on D/Γ such that every special subvariety $Y \subset D/\Gamma$ is \mathbb{R}_{alg} -definable. Below we understand $\mathcal{F}(Y)$ to be taken with respect to this structure with the standard effectively o-minimal structure on \mathbb{R}_{alg} . To deduce the algebraicity of S_Y , [1] proves that Φ is definable in $\mathbb{R}_{\text{an,exp}}$ and appeals to the definable Chow theorem [36]. We show that Φ is actually definable in $\mathbb{R}_{\text{LN,exp}}$ (although see Remark 53 for a discussion of issues around the computation of the format of Φ). Modulo effective computation of the format $\mathcal{F}(\Phi)$ we deduce the following more effective form of the algebraicity of the Hodge locus.

Theorem 4. *Let $Y \subset D/\Gamma$ be a special subvariety. Then $S_Y := \Phi^{-1}(Y)$ is an algebraic subvariety of S and*

$$\deg S_Y = \text{Eff}(\Phi, Y) \tag{11}$$

where \deg denotes the sum of the degrees of the pure-dimensional parts of S_Y .

We also show in §10.5 that the universal covering map for the universal abelian scheme over the Siegel modular variety $\mathbb{A}_g \rightarrow \mathcal{A}_g$ is definable in $\mathbb{R}_{\text{LN,exp}}$ when restricted to an appropriate fundamental domain. Combined with the effective Pila-Wilkie statements in §9 this allows one in principle to effectivize (at least the point counting aspect of) the many applications of the Pila-Wilkie theorem to unlikely intersection problems such as (relative) Manin-Mumford, André-Oort, and Zilber-Pink. We refer the reader to [41, 50] for an introduction to the vast literature around this topic.

In a related direction, [28, Section 6] gives numerous examples of functions arising in quantum systems that are shown to be Noetherian, at least when considered away from their singularities. The question is posed there whether these functions in fact live in an #o-minimal structure, which would allow one to quantify their complexity. While we do not establish #o-minimality of $\mathbb{R}_{\text{LN,exp}}$ in this paper,

it seems that effective o-minimality could provide a replacement suitable for the purposes of [28] and containing many of the examples considered there.

Finally we note that while the examples above mainly come from linear systems, the Noetherian class contains the solutions of essentially arbitrary non-linear systems of ODEs in one or several variables, at least away from their singularities. The theory of \mathbb{R}_{LN} can therefore be used for the quantitative study of geometric complexity in a wide range of examples arising for instance in classical mechanics.

1.9. Conjecture on #o-minimality. In [11, Conjecture 29] we conjectured that the structure generated by restricted Noetherian functions is #o-minimal. We also conjectured [11, Conjecture 33] that the structure generated by Q-functions, which are horizontal sections of regular flat connections (defined over \mathbb{Q}) with quasiunipotent monodromy, is #o-minimal. Since $\mathbb{R}_{\text{LN,exp}}$ contains both of these structures, the results of the present paper make a step toward these conjectures by establishing effective o-minimality. It seems natural to extend the conjecture to the larger structures considered in this paper as follows.

Conjecture 4. *The structure $\mathbb{R}_{\text{LN,PF}}$ is #o-minimal.*

It seems likely in light of the material developed in §7.4 and [14] that Conjecture 4 would follow from the #o-minimality of \mathbb{R}_{LN} , though the technical details of this reduction are not yet fully verified.

2. LOG-NOETHERIAN CELLS AND FUNCTIONS

2.1. Basic definition. We introduce the notion of a *Log-Noetherian (LN) cells* $\mathcal{C} \subset \mathbb{C}^\ell$ and *Log-Noetherian (LN) functions* on \mathcal{C} by induction on ℓ . Denote the set of LN functions on \mathcal{C} by $\mathcal{O}_{\text{LN}}(\mathcal{C})$.

In the case $\ell = 0$, we consider \mathbb{C}^0 to be a singleton and the only cell $\mathcal{C} \subset \mathbb{C}^0$ is \mathbb{C}^0 itself. An LN function is any function $F : \mathcal{C} \rightarrow \mathbb{C}$.

For $r \in \mathbb{C}$ (resp. $r_1, r_2 \in \mathbb{C}$) with $|r| > 0$ (resp. $|r_2| > |r_1| > 0$) we denote

$$\begin{aligned} D(r) &:= \{|z| < |r|\} & D_\circ(r) &:= \{0 < |z| < |r|\} \\ A(r_1, r_2) &:= \{|r_1| < |z| < |r_2|\} & * &:= \{0\}. \end{aligned} \tag{12}$$

We also set $S(r) := \partial D(r)$.

Recall the following notation from [10]. Let \mathcal{X}, \mathcal{Y} be sets and $\mathcal{F} : \mathcal{X} \rightarrow 2^{\mathcal{Y}}$ be a map taking points of \mathcal{X} to subsets of \mathcal{Y} . Then we denote

$$\mathcal{X} \odot \mathcal{F} := \{(x, y) : x \in \mathcal{X}, y \in \mathcal{F}(x)\}. \tag{13}$$

In this paper \mathcal{X} will be a subset of \mathbb{C}^n and \mathcal{Y} will be \mathbb{C} . If $r : \mathcal{X} \rightarrow \mathbb{C} \setminus \{0\}$ then for the purpose of this notation we understand $D(r)$ to denote the map assigning to each $x \in \mathcal{X}$ the disc $D(r(x))$, and similarly for D_\circ, A .

We now introduce the notions of LN-cells and LN-functions. We denote the ring of LN-functions on a cell \mathcal{C} by $\mathcal{O}_{\text{LN}}(\mathcal{C})$. Note that these two definitions are by mutual induction on the length of the cell.

Definition 5. *A cell of length $\ell = 0$ is the singleton $\mathcal{C} = \mathbb{C}^0$. An LN cell $\mathcal{C} \subset \mathbb{C}^{\ell+1}$ of length $\ell + 1$ is a set of the form*

$$\mathcal{C} := \mathcal{C}_{1.. \ell} \odot \mathcal{F} \tag{14}$$

where $\mathcal{C}_{1.. \ell} \subset \mathbb{C}^\ell$ is an LN cell of length ℓ , the fiber \mathcal{F} is one of

$$\mathcal{F} := *, D(r), D_\circ(r), A(r_1, r_2) \tag{15}$$

and the radii are as follows:

- If $\mathcal{F} = D(r), D_\circ(r)$ then $r \in \mathcal{O}_{\text{LN}}(\mathcal{C}_{1..l})$ with $r : \mathcal{C}_{1..l} \rightarrow \mathbb{C}^*$.
- If $\mathcal{F} = A(r_1, r_2)$ then $r_1, r_2 \in \mathcal{O}_{\text{LN}}(\mathcal{C}_{1..l})$ with $r_1, r_2 : \mathcal{C}_{1..l} \rightarrow \mathbb{C}^*$ and $|r_1| < |r_2|$ pointwise on $\mathcal{C}_{1..l}$.

Parting with [10], we require that r in the case of $D(r)$ be a constant functions.

We now define the *standard derivatives* $\partial_j^{\mathcal{C}}$ on \mathcal{C} . Define $\partial_j^{\mathcal{C}}$ to be $\partial_j^{\mathcal{C}_{1..l}}$ if $j = 1, \dots, l$ and

$$\partial_{\ell+1}^{\mathcal{C}} := \begin{cases} r \frac{\partial}{\partial \mathbf{z}_{\ell+1}} & \text{if } \mathcal{F} = D(r) \\ \mathbf{z}_{\ell+1} \frac{\partial}{\partial \mathbf{z}_{\ell+1}} & \mathcal{F} = D_\circ(r), A(r_1, r_2). \\ 0 & \mathcal{F} = *. \end{cases} \quad (16)$$

Remark 6. If we had allowed r in the fibers $D(r)$ to be non-constant then the fields $\partial_j^{\mathcal{C}}$ would be non-commuting. For the correct general definition, note that every cell is biholomorphic to a cell where all fibers of type $D(r)$ are taken to be $D(1)$ by a linear rescaling map $z \rightarrow z/r$. We could define $\partial_j^{\mathcal{C}}$ for the general case by pulling back from this standard model. However this makes the fields harder to visualize, so we prefer to stick to the standard model in our definition of LN cells.

An LN chain on \mathcal{C} is a collection of bounded holomorphic functions $F_1, \dots, F_N : \mathcal{C} \rightarrow \mathbb{C}$ such that the ring $\mathbb{C}[F_1, \dots, F_N]$ is closed under the standard derivatives; that is, such that

$$\partial_j^{\mathcal{C}}(F_i) = G_{i,j}(F_1, \dots, F_N) \quad (17)$$

for suitable polynomials $G_{i,j}$ over \mathbb{C} . An LN function $F : \mathcal{C} \rightarrow \mathbb{C}$ is a function of the form $G(F_1, \dots, F_N)$ for some polynomial G and some LN chain F_1, \dots, F_N on \mathcal{C} . This finishes the recursive definition of LN-cells and LN-functions. Whenever we speak of cells below we will implicitly mean LN cells, unless explicitly stated otherwise.

Remark 7. One may be tempted to use a simpler definition for LN (or Noetherian) functions, namely to work with functions $F : \mathcal{C} \rightarrow \mathbb{C}$ such that the ring generated from $\mathbb{C}[F]$ by closing under the $\partial_j^{\mathcal{C}}$ derivatives is finitely generated. However this does not lead to a good notion, as illustrated in §A.4.

2.2. Format. We introduce a measure for the complexity an LN-cell \mathcal{C} and an LN-function F . If $\ell = 0$ then the format of $\mathcal{F}(\mathcal{C})$ is 1, and the format $\mathcal{F}(f)$ of any (constant) function on $f : \mathcal{C} \rightarrow \mathbb{C}$ is the least integer upper bound for $|f|$. In the notations of §2.1, for general $\mathcal{C} \subset \mathbb{C}^{\ell+1}$,

$$\begin{aligned} & \mathcal{F}(\mathcal{C}_{1..l} \odot *), 1 + \mathcal{F}(\mathcal{C}_{1..l}). \\ \mathcal{F}(\mathcal{C}_{1..l} \odot D_\circ(r)), \mathcal{F}(\mathcal{C}_{1..l} \odot D(r)) &= 1 + \mathcal{F}(\mathcal{C}_{1..l}) + \mathcal{F}(r) \\ \mathcal{F}(\mathcal{C}_{1..l} \odot A(r_1, r_2)) &= 1 + \mathcal{F}(\mathcal{C}_{1..l}) + \mathcal{F}(r_1) + \mathcal{F}(r_2). \end{aligned} \quad (18)$$

We define the norm of a polynomial $\|P\|$ to be the sum of the absolute values of its coefficients. If F_1, \dots, F_N is an LN-chain as in (17) and $F = G(F_1, \dots, F_N)$ an LN-function we define their formats by

$$\begin{aligned} \mathcal{F}(F_1, \dots, F_N) &:= \mathcal{F}(\mathcal{C}) + n + N + \sum_{i,j} \deg G_{i,j} + \|G_{i,j}\| + \sup_{\substack{i=1,\dots,N \\ \mathbf{z} \in \mathcal{C}}} |F_i(\mathbf{z})| \\ \mathcal{F}(F) &:= \mathcal{F}(F_1, \dots, F_N) + \deg G + \|G\|. \end{aligned} \quad (19)$$

To be perfectly accurate, we take the least integer upper bound for the numbers above, and the definition above defines a filtration – so the actual format is defined by the minimum \mathcal{F} over all possible representations of the LN-chain or LN-function.

2.3. Extensions of cells. We now introduce the notion of δ extensions of cells. We largely follow [10] here, but reproduce the definitions for the convenience of the reader.

For any $0 < \delta < 1$ we define the δ -extensions, denoted by superscript δ , by

$$\begin{aligned} D^\delta(r) &:= D(\delta^{-1}r) & D_\circ^\delta(r) &:= D_\circ(\delta^{-1}r) \\ A^\delta(r_1, r_2) &:= A(\delta r_1, \delta^{-1}r_2) & *^\delta &:= *. \end{aligned} \quad (20)$$

We also set $S^\delta(r) = A(\delta r, \delta^{-1}r)$.

Next, we define the notion of a δ -extension of a cell of length ℓ where $\delta \in (0, 1)$.

Definition 8. *The cell of length zero is defined to be its own δ -extension. A cell \mathcal{C} of length $\ell + 1$ admits a δ -extension*

$$\mathcal{C}^\delta := \mathcal{C}_{1..\ell}^\delta \odot \mathcal{F}^\delta \quad (21)$$

if $\mathcal{C}_{1..\ell}$ admits a δ -extension, and if the function r (resp. r_1, r_2) involved in \mathcal{F} admits holomorphic continuation to an LN function on $\mathcal{C}_{1..\ell}^\delta$ and satisfies $|r(\mathbf{z}_{1..\ell})| > 0$ (resp. $0 < |r_1(\mathbf{z}_{1..\ell})| < |r_2(\mathbf{z}_{1..\ell})|$) in this larger domain.

As a shorthand, when say that \mathcal{C}^δ is an LN cell we mean that \mathcal{C} is an LN cell admitting a δ -extension.

The following is a simple exercise in the definitions.

Lemma 9. *If \mathcal{C}^δ is an LN cell then $\mathcal{F}(\mathcal{C}) \leq \mathcal{F}(\mathcal{C}^\delta)$.*

2.4. Maps and cellular maps. A map $f : \mathcal{C} \rightarrow \hat{\mathcal{C}}$ between two LN cells is called an LN map if each coordinate is an LN function. The format $\mathcal{F}(f)$ is defined to be the sum of the formats of $\mathcal{C}, \hat{\mathcal{C}}$ and the coordinate functions.

An LN map $f : \mathcal{C} \rightarrow \hat{\mathcal{C}}$ between two cells of the same length ℓ is called *cellular* if it has the form $\mathbf{w}_j = \phi_j(\mathbf{z}_{1..j})$ where each ϕ_j is an LN function and $\frac{\partial \phi_j}{\partial \mathbf{z}_j}$ is non-vanishing for $j = 1, \dots, \ell$.

Remark 10. *Note that in [10] cellular maps were defined with the additional assumption that ϕ_j is a monic polynomial in \mathbf{z}_j .*

We show in Theorem 11 that the pullback f^*F of an LN function F under an LN map f is again LN and $\mathcal{F}(f^*F) < \text{Eff}(f, F)$, but note that this is not a trivial statement and in involves a slight shrinking of the domain.

2.5. The real setting. We inductively define the notion of a *real* LN cell and a *real* LN function. A cell $\mathcal{C} \odot \mathcal{F}$ is real if \mathcal{C} is real and the radii involved in the definition \mathcal{F} are real LN functions. The *real part* of the cell denoted $\mathbb{R}_+\mathcal{C}$ is defined to be $\mathbb{R}_+\mathcal{C} \odot \mathbb{R}_+\mathcal{F}$, where

$$\begin{aligned} \mathbb{R}_+* &:= * & \mathbb{R}_+D(r) &:= D(r) \cap \mathbb{R} \\ \mathbb{R}_+D_\circ(r) &:= D_\circ(r) \cap \mathbb{R}_{>0} & \mathbb{R}_+A(r_1, r_2) &:= A(r_1, r_2) \cap \mathbb{R}_{>0}. \end{aligned} \quad (22)$$

Note that we took $\mathbb{R}_+D(r)$ to be the whole interval, parting with the convention from [10], as this seems somewhat more natural. However the difference is merely technical.

An LN function $f : \mathcal{C} \rightarrow \mathbb{C}$ is called *real* if \mathcal{C} is real and f is real on $\mathbb{R}_+\mathcal{C}$. A map $f : \mathcal{C} \rightarrow \hat{\mathcal{C}}$ is called real if its coordinate functions are real and $f(\mathbb{R}_+\mathcal{C}) \subset \mathbb{R}_+\hat{\mathcal{C}}$.

2.6. Cellular covers. We will often be interested in covering a cell by cellular images of other cells. Toward this end we introduce the following definition.

Definition 11. Let \mathcal{C}^δ be an LN cell and $\{f_j : \mathcal{C}_j^{\delta'} \rightarrow \mathcal{C}^\delta\}$ be a finite collection of LN cellular maps. We say that this collection is an LN (δ', δ) -cellular cover of \mathcal{C} if

$$\mathcal{C} \subset \cup_j (f_j(\mathcal{C}_j)). \quad (23)$$

Similarly, when $f, \mathcal{C}_j^{\delta'}, \mathcal{C}^\delta$ are all real this is called a *real cover* if

$$\mathbb{R}_+\mathcal{C} \subset \cup_j (f_j(\mathbb{R}_+\mathcal{C}_j)). \quad (24)$$

When (δ', δ) are clear from the context we will speak simply of cellular covers.

The number of maps f_j in a cellular cover is called the *size* of the cover.

Remark 12. We remark that if $\{f_j : \mathcal{C}_j^{\delta'} \rightarrow \mathcal{C}^\delta\}$ is an LN (real) cellular cover of \mathcal{C} and $\{f_{jk} : \mathcal{C}_{jk}^{\delta''/2} \rightarrow \mathcal{C}_j^{\delta'}\}$ is an LN (real) cellular cover of \mathcal{C}_j then $\{f_j \circ f_{jk}\}$ is an LN (real) (δ'', δ) -cover of \mathcal{C} . Note that we need a slightly larger extension $\delta''/2$ in the f_{jk} cover because of the slight shrinking of domains in Theorem 11.

2.7. The cellular parametrization theorem. We say that $f \in \mathcal{O}_{\text{LN}}(\mathcal{C})$ is compatible with \mathcal{C} if F is either identically vanishing or nowhere vanishing on \mathcal{C} . If $f : \mathcal{C} \rightarrow \hat{\mathcal{C}}$ is a cellular map and $F \in \mathcal{O}_{\text{LN}}(\hat{\mathcal{C}})$ we say that f is compatible with F if the pullback f^*F is compatible with \mathcal{C} .

Our main technical tool in this paper is the following cellular parametrization theorem (CPT) in the LN category. In the analytic and algebraic categories, this theorem is the main result of [10].

Theorem 5 (Cellular Parametrization Theorem – LN category). *Let \mathcal{C}^δ be an LN cell and $F_1, \dots, F_M \in \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta)$. Then there exists a cellular cover $\{f_j : \mathcal{C}_j^\delta \rightarrow \mathcal{C}^\delta\}$ such that each f_j is compatible with each F_k . Moreover*

$$\#\{f_j\} < \text{Eff}(F_1, \dots, F_M, 1/(1-\delta)) \quad \text{and} \quad \mathcal{F}(f_j) < \text{Eff}(F_1, \dots, F_M). \quad (25)$$

If $\mathcal{C}, F_1, \dots, F_M$ are real then the cover can be chosen to be real.

The CPT is proven in §5.

3. BACKGROUND FROM COMMUTATIVE ALGEBRA AND DIFFERENTIAL EQUATIONS

In this section we recall some facts about polynomial rings with derivations and about oscillation of solutions of complex ODEs.

3.1. Ascending chains of ideals. Let $R = \mathbb{C}[x_1, \dots, x_n]$ or $R := \mathbb{R}[x_1, \dots, x_n]$ be a polynomial ring and $D : R \rightarrow R$ a derivation of R . Explicitly,

$$D = \sum p_i \frac{\partial}{\partial x_i} \quad \text{where } p_i \in R. \quad (26)$$

For $P \in R$ we write $\mathcal{F}(P) = \deg P + \|P\|$ with $\|P\|$ defined in §2.2 and

$$\mathcal{F}(D) := \sum_i \mathcal{F}(p_i). \quad (27)$$

Define an ascending chain of ideals I_P^k by

$$I_P^0 := (P), \quad I_P^{k+1} := \langle I_P^k, DI_P^k \rangle. \quad (28)$$

The following result of Novikov and Yakovenko plays a key role in our approach.

Theorem 6 ([35]). *The chain I_P^k stabilizes, i.e. $I_P^\ell = I_P^{\ell+1}$ with $\ell \leq \text{Eff}(D, P)$. Moreover,*

$$D^{\ell+1}P = \sum_{j=0}^{\ell} c_j D^j P, \quad c_j \in R \quad (29)$$

with $\sum \mathcal{F}(c_j) < \text{Eff}(D, P)$.

In fact [35] gives more specific bounds, exponential in the degrees and triply exponential in the dimension n . This result is used in loc. cit. to bound the number of intersections between a trajectory of D and the hypersurface defined by P using certain non-oscillation results for solutions of linear ODEs. We use it for similar purposes but require slightly different complex estimates that we recall below.

3.2. Total variation of argument for solutions of linear ODEs. Let $U \subset \mathbb{C}$ be a bounded domain and $f : \bar{U} \rightarrow \mathbb{C}$ be a holomorphic function. The *total variation of argument* [31], also called *Voorhoeve index*, of a holomorphic function $f : U \rightarrow \mathbb{C}$ along a piecewise-smooth curve $\gamma \subset U$ is defined by

$$V_\gamma(f) := \frac{1}{2\pi} \int_\gamma |\text{d Arg } f(z)|. \quad (30)$$

The following theorem, a complex analog of the classical result of de la Vallé Poussin [17], shows that the total variation of argument of a solution of a scalar ODE with analytic coefficients can be explicitly bounded in terms of the order of the equation and the upper bounds for its coefficients.

Theorem 7 ([49, Corollary 2.7]). *Let $\gamma \subset \mathbb{C}$ be either an interval or a circular arc of length ℓ . Suppose $w(t)$ is analytic on γ and satisfies the equation*

$$w^{(n)}(t) + c_1(t)w^{(n-1)}(t) + \cdots + c_n(t)w(t) = 0 \quad (31)$$

where $c_j(t)$ are all analytic on I and bounded in absolute value by $M > 1$. Then

$$V_I(w) \leq C \cdot \ell \cdot n \cdot M \quad (32)$$

where C is some absolute constant.

We remark that in loc. cit. this is stated for straight lines, but the proof is easily seen to apply also for circular arcs. Alternatively one can change variables to turn a circular arc into an interval and this only affects M by some constant factor.

3.3. Total variation of argument for LN functions. We show that the total variation of argument of an LN function along suitable curves in the domain can be bounded in terms of $\mathcal{F}(f)$.

Supposes $\mathcal{C} := \mathcal{C}_{1,\ell} \odot \mathcal{F}$. Fix a point $\mathbf{z} \in \mathcal{C}$ and let γ be a circular arc in the fiber $\{\mathbf{z}\} \odot \mathcal{F}(\mathbf{z})$. Define the standard length of γ , denoted $\text{length}(\gamma)$, to be the length of γ with respect to the $\partial_{\ell+1}^{\mathcal{C}}$ chart. In other words,

$$\text{length}(\gamma) = \begin{cases} \frac{1}{r} \times (\text{Euclidean length of } \gamma) & \mathcal{F} = D(r) \\ \text{Euclidean length of } \log(\gamma) & \mathcal{F} = D_\circ(r), A(r_1, r_2). \end{cases} \quad (33)$$

In this situation we have the following.

Proposition 13. *Let \mathcal{C} be an LN cell and $f \in \mathcal{O}_{\text{LN}}(\mathcal{C})$ and $\gamma \subset \mathcal{C}$ as above. Then*

$$V_\gamma(f) \leq \text{Eff}(f) \cdot \text{length}(\gamma). \quad (34)$$

Proof. According to Theorem 6 applied to f and the derivation $D := \partial_j^{\mathcal{C}}$ we have a differential equation

$$D^{N+1}f = \sum_{j=0}^N c_j D^j f, \quad c_j \in R \quad (35)$$

where R is the ring generated by our LN chain with N , the degrees of the c_j , and their norms, bounded by $\text{Eff}(f)$. Along with the bounds on the format of the LN chain, this implies that all coefficients c_j are bounded in absolute value by $\text{Eff}(f)$ everywhere in \mathcal{C} . Therefore Theorem 7 implies that the total variation of argument of f along γ is bounded as claimed, noting that the time parametrization with respect to D agrees with our notion of standard length. \square

3.4. Taylor and Laurent domination.

Definition 14 (Laurent domination). *A holomorphic function $f : A(r_1, r_2) \rightarrow \mathbb{C}$ is said to possess the (p, q, M) Laurent domination property where $p \leq q$ are integers and $M \in \mathbb{R}_+$ if its Laurent expansion $f(z) = \sum a_k (z - z_0)^k$ satisfies the estimates*

$$\begin{aligned} |a_k| r_2^k &< M \max_{j=p, \dots, q} |a_j| r_2^j, & k = q+1, q+2, \dots \\ |a_k| r_1^k &< M \max_{j=p, \dots, q} |a_j| r_1^j, & k = p-1, p-2, \dots \end{aligned} \quad (36)$$

If $f : D(r) \rightarrow \mathbb{C}$ we formally put $r_1 = 0$ above, and refer to the $(0, q, M)$ Laurent domination property as the (q, M) Taylor domination property.

In the class of LN functions we have the following effective form of Taylor and Laurent domination.

Proposition 15. *Let $\mathcal{C} := \mathcal{C}_{1..l} \odot \mathcal{F}$ be an LN cell and $f \in \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta)$. Then $f(\mathbf{z}_{1..l}, \cdot)$ has the (p, q, M) Laurent domination property for every $\mathbf{z}_{1..l} \in \mathcal{C}_{1..l}$ with $|p| + |q| + M < \text{Eff}(f)$.*

Proof. This is essentially already contained in [10, Corollary 67]. The constants there depend on the constant in [10, Lemma 66], which is derived directly from the total variation of argument of f along the boundary of a certain disc D , which is easily seen to have effectively bounded standard length. The claim thus follows from Proposition 13. Also note that since LN functions are always bounded, if $\mathcal{F} = D_\circ(r)$ then f automatically extends to a holomorphic function over $\mathcal{C}_{1..l} \odot D(r)$ (although not as an LN function) so we have in fact Taylor domination in this case. \square

Recall the following simple consequence of the domination property.

Proposition 16 ([10, Proposition 68]). *Let $f : C^\delta \rightarrow \mathbb{C}$ where $C = D(r), D_\circ(r)$ or $A(r_1, r_2)$ and assume that it has the (p, q, M) domination property. Then for the Taylor or Laurent expansion*

$$f(z) = \sum_{j=p}^q a_j z^j + R(z) \quad (37)$$

we have for every $z \in C$ a bound

$$|R(z)| \leq \frac{2\delta M}{1-\delta} \max_{j=p,\dots,q} |a_j z^j|. \quad (38)$$

We also record two basic propositions on Taylor domination that will be useful later.

Proposition 17. *Suppose*

$$f : D(1) \rightarrow \mathbb{C}, \quad f(z) = \sum a_j z^j \quad (39)$$

has the (q, M) Taylor domination property with $q > 0$, and

$$|a_q| \geq \max_{j=0, q-1} |a_j|. \quad (40)$$

Then $f'(z)$ has the $(q-1, M)$ domination property on $D(1/2)$.

Proof. For every $k > q$ we have $|a_k| < M|a_q|$ by assumption. Then

$$f'(z) = \sum k a_k z^{k-1}. \quad (41)$$

One easily checks that for $k > q$ we have $k/q \leq 2^{k-q}$ so

$$k|a_k|(1/2)^k \leq M \cdot q|a_q|(1/2)^q \quad (42)$$

which proves the claim. \square

Proposition 18. *Suppose $f : D(1) \rightarrow \mathbb{C}^*$ has the (q, M) Taylor domination property and $\delta > 0$. Then there exists $\varepsilon = \varepsilon(q, M, \delta)$ with $1/\varepsilon = \text{Eff}(q, M, 1/\delta)$ such that $f|_{D(\varepsilon)}$ has the $(0, \delta)$ -domination property.*

Proof. For each $j, k = 0, \dots, q$ consider the radius

$$r_{j,k} := \sqrt[k-j]{\frac{|a_j|}{|a_k|}} \quad (43)$$

where $|a_j z^j| = |a_k z^k|$. Outside the annulus $A_{j,k} = A(r_{j,k}/4, 4r_{j,k})$ the ratio $|a_j z^j|/|a_k z^k|$ is either smaller than $1/4$ or larger than 4 . Thus for z outside all of these annuli, it is easy to see that for the maximal among these terms $a_m z^m$ we have

$$|a_m z^m| > 2 \sum_{m \neq j=0,\dots,q} |a_j z^j|. \quad (44)$$

By Proposition 16 we also have that if $|z| < 1/(2+8M)$ then

$$|a_m z^m| > 2|R(z)|. \quad (45)$$

Choose z satisfying $|z| < 1/(2+8M)$ and $z \notin A_{j,k}$ for every j, k . Then $a_m z^m$ dominates all other terms in the Taylor expansion on the circle of radius $|z|$ and it follows by Rouché's theorem that f has m zeros in the disc $D(|z|)$. Hence by assumption $m = 0$, and f has the $(0, M)$ domination property in $D(|z|)$. It is clear that $|z|$ can be chosen with absolute value r satisfying $1/r < \text{Eff}(q, M)$. Setting now $\varepsilon = \delta r/M$ we get the $(0, \delta)$ -domination property on $D(\varepsilon)$. \square

4. BASIC THEORY OF LN FUNCTIONS

In this section we establish some theory concerning LN functions on LN cells.

4.1. Fundamental lemmas from hyperbolic geometry. For any hyperbolic Riemann surface X we denote by $\text{dist}(\cdot, \cdot; X)$ the hyperbolic distance on X . We use the same notation when $X = \mathbb{C}$ and $X = \mathbb{R}$ to denote the usual Euclidean distance, and when $X = \mathbb{C}P^1$ to denote the Fubini-Study metric normalized to have diameter 1. For $x \in X$ and $r > 0$ we denote by $B(x, r; X)$ the open r -ball centered x in X . For $A \subset X$ we denote by $B(A, r; X)$ the union of r -balls centered at all points of A .

The notion of δ -extension is naturally associated with the Euclidean geometry of the complex plane. However, in many cases it is more convenient to use a different normalization associated with the hyperbolic geometry of our domains. For any $0 < \rho < \infty$ we define the $\{\rho\}$ -extension $\mathcal{F}^{\{\rho\}}$ of \mathcal{F} to be \mathcal{F}^δ where δ satisfies the equations

$$\begin{aligned} \rho &= \frac{2\pi\delta}{1-\delta^2} && \text{for } \mathcal{F} \text{ of type } D, \\ \rho &= \frac{\pi^2}{2|\log \delta|} && \text{for } \mathcal{F} \text{ of type } D_\circ, A. \end{aligned} \tag{46}$$

The motivation for this notation comes from the following fact, describing the hyperbolic-metric properties of a domain \mathcal{F} within its $\{\rho\}$ -extension.

Fact 19 ([10, Fact 6]). *Let \mathcal{F} be a domain of type A, D, D_\circ and let S be a component of the boundary of \mathcal{F} in $\mathcal{F}^{\{\rho\}}$. Then the length of S in $\mathcal{F}^{\{\rho\}}$ is at most ρ .*

We define the $\{\rho\}$ -extension $\mathcal{C}^{\{\rho\}}$ by analogy with the δ -extension, $(\mathbb{C} \odot \mathcal{F})^{\{\rho\}} := \mathcal{C}^{\{\rho\}} \odot \mathcal{F}^{\{\rho\}}$. The lemmas below are more natural to state in terms of the $\{\rho\}$ -extension. However, since in the present paper we only care about effectivity of the constants and make no attempt to optimize the asymptotic dependence on parameters, it will make essentially no difference which type of extension we use. We have mostly kept to δ -extensions to simplify the presentation.

Recall the three fundamental lemmas from [10].

Lemma 20 (Fundamental Lemma for \mathbb{D}). *Let $\mathcal{C}^{\{\rho\}}$ be a complex cell. Let $f : \mathcal{C}^{\{\rho\}} \rightarrow \mathbb{D}$ be holomorphic. Then*

$$\text{diam}(f(\mathcal{C}); \mathbb{D}) = O_\ell(\rho). \tag{47}$$

Lemma 21 (Fundamental Lemma for $\mathbb{D} \setminus \{0\}$). *Let $\mathcal{C}^{\{\rho\}}$ be a complex cell and $0 < \rho < 1$. Let $f : \mathcal{C}^{\{\rho\}} \rightarrow \mathbb{D} \setminus \{0\}$ be holomorphic. Then one of the following holds:*

$$f(\mathcal{C}) \subset B(0, e^{-\Omega_\ell(1/\rho)}; \mathbb{C}) \quad \text{or} \quad \text{diam}(f(\mathcal{C}); \mathbb{D} \setminus \{0\}) = O_\ell(\rho). \tag{48}$$

In particular, one of the following holds:

$$\log |f(\mathcal{C})| \subset (-\infty, -\Omega_\ell(1/\rho)) \quad \text{or} \quad \text{diam}(\log |f(\mathcal{C})|; \mathbb{R}) = O_\ell(\rho). \tag{49}$$

Lemma 22 (Fundamental Lemma for $\mathbb{C} \setminus \{0, 1\}$). *Let $\mathcal{C}^{\{\rho\}}$ be a complex cell and let $f : \mathcal{C}^{\{\rho\}} \rightarrow \mathbb{C} \setminus \{0, 1\}$ be holomorphic. Then one of the following holds:*

$$f(\mathcal{C}) \subset B(\{0, 1, \infty\}, e^{-\Omega_\ell(1/\rho)}; \mathbb{C}P^1) \quad \text{or} \quad \text{diam}(f(\mathcal{C}); \mathbb{C} \setminus \{0, 1\}) = O_\ell(\rho). \tag{50}$$

4.2. Bounds for standard derivatives.

Lemma 23. *Let $f \in \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta)$ and $k \in \mathbb{N}$. Then*

$$\|(\partial_i^{\mathbb{C}})^k f\| \leq \rho^k \|f\|_{\mathcal{C}^\delta}, \quad \rho = O\left(\frac{1}{1-\delta}\right) \tag{51}$$

for $j = 1, \dots, \ell$.

Proof. First, we can assume without loss of generality that there are no D_\circ fibers. Indeed, we can replace each $D_\circ(r)$ fiber by $A(\varepsilon r, r)$ for $\varepsilon \ll 1$. Since (51) is independent of ε the claim for the original cell will follow from the claim for these modified cells.

Recall [10, Definition 39] that the skeleton of \mathcal{C} is defined as the set of points in \mathcal{C} where each coordinate is in the boundary of its corresponding fiber. By the maximum principle it is enough to check the maximum of $\partial_i^{\mathcal{C}}$ on the skeleton of \mathcal{C} . Suppose it is obtained at a point p . Moreover, since $\partial^{\mathcal{C}}$ is invariant by rescaling of each fiber and by inversion $z \rightarrow r/z$ in the case of annuli fibers, there is also no harm in assuming that p is in the component of the skeleton given by $S(1)$ in D , A fibers and by $\{0\}$ in $*$ fibers. Then $B_\rho(p) \subset \mathcal{C}^\delta$ where ρ is as given in the statement.

Let $\phi : (\mathbb{C}, 0) \rightarrow \mathcal{C}^\delta$ be the flow chart of $\partial_j^{\mathcal{C}}$ with $\phi(0) = p$. Then we are interested in evaluating $|(f \circ \phi)^{(k)}(0)|$. It is easy to see that ϕ extends to a disc of radius $O(\rho)$, and the claim follows from the Cauchy estimates. \square

4.3. Four basic theorems. In this section we prove four basic theorems for LN functions: monomialization, restricted division, logarithmic derivation, and stability under pullback. We will establish all of these by concurrent induction, so we assume throughout that \mathcal{C}^δ is an LN cell of length $\ell + 1$ and that all statements have been established for cells of length ℓ .

4.3.1. Monomialization. We state an analog of the monomialization lemma of [10], with effective control over the constants in terms of $\mathcal{F}(f)$. We begin by recalling some notation.

A cell \mathcal{C} is homotopy equivalent to a product of points (for fibers $*$, D) and circles (for fibers D_\circ , A) by the map $\mathbf{z}_i \rightarrow \text{Arg } \mathbf{z}_i$. Thus $\pi_1(\mathcal{C}) \simeq \prod G_i$ where G_i is trivial for $*$, D and \mathbb{Z} for D_\circ , A . Let γ_i denote the generator of G_i chosen with positive complex orientation for $G_i = \mathbb{Z}$ and $\gamma_i = e$ otherwise.

Definition 24. Let $f : \mathcal{C} \rightarrow \mathbb{C}^*$ be continuous. Define the monomial associated with f to be $\mathbf{z}^{\alpha(f)}$ where

$$\alpha_i(f) = f_* \gamma_i \in \mathbb{Z} \simeq \pi_1(\mathbb{C}^*). \quad (52)$$

It is easy to verify that $f \mapsto \mathbf{z}^{\alpha(f)}$ is a group homomorphism from the multiplicative group of continuous maps $f : \mathcal{C} \rightarrow \mathbb{C}^*$ to the multiplicative group of monomials, which sends each monomial to itself.

Theorem 8. Let $f : \mathcal{C}^\delta \rightarrow \mathbb{C}^*$ with $f \in \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta)$. Then we have a decomposition $f = m_f(\mathbf{z})U(\mathbf{z})$ where

$$m_f(\mathbf{z}) = \lambda \mathbf{z}^{\alpha(f)}, \quad \lambda \in \mathbb{R}_{>0}, \quad (53)$$

and the branches of $\log U : \mathcal{C}^\delta \rightarrow \mathbb{C}$ are univalued. We have

$$\|\alpha(f)\| + \mathcal{F}(U|_{\mathcal{C}}) \leq \text{Eff}(f), \quad (54)$$

and for all $\mathbf{z} \in \mathcal{C}$,

$$\frac{1}{\text{Eff}(f)} \leq |U(\mathbf{z})| \leq 1. \quad (55)$$

If f is real then U is real.

Proof. We assume that the fiber is of type $A(r_1, r_2)$, the other cases being similar. We begin by bounding $\alpha(f)_{\ell+1}$ by $\text{Eff}(f)$. Let γ denote the curve given by a circle of radius r_2 in the $\mathbf{z}_{\ell+1}$ -coordinate and some point $p \in \mathcal{C}_{1..\ell}$ in the $\mathbf{z}_{1..\ell}$ coordinates. Then $\gamma \subset \mathcal{C}^\delta$, and clearly $\alpha(f)_{\ell+1}$ is bounded by the total variation of argument of f along γ . This circle has standard length 2π , and the bound therefore follows by Theorem 13.

Now consider the cell $\hat{\mathcal{C}} := \mathcal{C}_{1..\ell} \odot *$ and the cellular map $\phi : \hat{\mathcal{C}}^\delta \rightarrow \mathcal{C}^\delta$ given by

$$\phi(\mathbf{z}_{1..\ell}) = (\mathbf{z}_{1..\ell}, r_2(\mathbf{z}_{1..\ell})). \quad (56)$$

By pullback stability (Theorem 11), the function $\hat{f} := \phi^* f$ is LN on the cell $\hat{\mathcal{C}}^{\sqrt{\delta}}$ with effectively bounded format. By induction we have an effective bound on $\|\alpha(\hat{f})\|$. By a similar reasoning we have an effective bound on $\|\alpha(r_2)\|$. We remark that even though $\hat{\mathcal{C}}$ is formally of length $\ell + 1$, in the proof of pullback stability one can identify $\hat{\mathcal{C}}$ with $\mathcal{C}_{1..\ell}$ so our induction is in fact well-founded.

In the notation of Definition 24, the map ϕ takes the fundamental loop $\gamma_i \in \pi_1(\hat{\mathcal{C}})$ to $\gamma_i + \alpha(r)_i \cdot \gamma_{\ell+1}$ in $\pi_1(\mathcal{C})$. Thus

$$\alpha(\hat{f})_i = \alpha(f)_i + \alpha(r)_i \cdot \alpha(f)_{\ell+1}. \quad (57)$$

Solving for $\alpha(f)_i$ we see that it is also effectively bounded.

We proceed with the bounds for the unit U . Fix a branch of $\log U$. We will show that the diameter of $\log U \subset \mathbb{C}$ is effectively bounded. One can then choose the (positive real) constant λ to normalize its maximum to 1. The bound on $\mathcal{F}(U)$ follows from the restricted division theorem (Theorem 9) to be proved later.

First apply the inductive hypothesis to $\hat{U} := \phi^* U$. Note that $\alpha(\hat{U}) = 0$, so this indeed plays the role of a unit in the induction. Thus $\log \hat{U}$ has diameter bounded by $\text{Eff}(f)$. We will thus finish the proof by showing that for any $\mathbf{z}_{1..\ell} \in \mathcal{C}$ we have a bound for the diameter of $\log U$ in the fiber over $\mathbf{z}_{1..\ell}$. Set $r_1 = r_1(\mathbf{z}_{1..\ell})$ and $r_2 = r_2(\mathbf{z}_{1..\ell})$ and

$$U' : A(r_1, r_2)^\delta \rightarrow \mathbb{C}^*, \quad U'(t) = U(\mathbf{z}_{1..\ell}, t). \quad (58)$$

By Theorem 13 we have an effective bound for the total variation of argument of U' along the two boundary components of $A(r_1, r_2)^{\sqrt{\delta}}$.

Let $L := \log U'$ and consider $X := L(A(r_1, r_2)^{\sqrt{\delta}})$. By the open mapping theorem

$$\partial X \subset L(\{|t| = \sqrt{\delta}r_1\}) \cup L(\{|t| = r_2/\sqrt{\delta}\}). \quad (59)$$

In particular,

$$\text{diam Im } X \leq \text{diam Im } L(\{|t| = \sqrt{\delta}r_1\}) + \text{diam Im } L(\{|t| = r_2/\sqrt{\delta}\}) \quad (60)$$

where Im denotes the imaginary part. The two summands on the right hand side are bounded by the two total variations of argument above, so $\text{diam Im } X$ is effectively bounded, say by some constant V .

Now we can finish as in [10]. Set $\tilde{U} := (U')^{1/(4V)}$ and assume further, by multiplying by a scalar, that the image of \tilde{U} contains i . Then the total variation of argument of \tilde{U} is bounded by $\pi/4$, and it follows that $\tilde{U} : A(r_1, r_2)^{\sqrt{\delta}} \rightarrow \mathbb{H}$. Then the fundamental lemma for \mathbb{D} (applied in this case to the upper half space model) implies that the diameter of $\tilde{U}(\mathcal{C})$ is bounded by a constant depending only on $\sqrt{\delta}$. Recovering U' as \tilde{U}^{4V} and recalling that V was effectively bounded finishes the proof. \square

4.3.2. *Restricted division.* The following theorem plays a fundamental role in our approach as it allows us to form an analog of the strict transform used in resolution of singularities.

Theorem 9. *Let \mathcal{C}^δ be an LN cell and $f, g \in \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta)$. Suppose further that*

$$g : \mathcal{C}^\delta \rightarrow \mathbb{C}^* \quad (61)$$

and that f/g is bounded in \mathcal{C}^δ . Then

$$\frac{f}{g} \in \mathcal{O}_{\text{LN}}(\mathcal{C}) \quad \text{and} \quad \mathcal{F}\left(\left(\frac{f}{g}\right)|_{\mathcal{C}}\right) < \text{Eff}(f, g, \|f/g\|_{\mathcal{C}^\delta}). \quad (62)$$

If f, g are real then f/g is real.

Proof. We begin with the case where $g = \lambda \mathbf{z}^\alpha$. Write $\partial_j := \partial_j^{\mathcal{C}^\delta}$. By Theorem 6 there exists $N = \text{Eff}(f)$ such that for each $j = 1, \dots, \ell$ there is an equation

$$\partial_j^{N+1} f = \sum_{i=0}^N c_{j,i} \partial_j^i f \quad (63)$$

where $c_{j,i} \in \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta)$ have format $\text{Eff}(f)$. We claim that adding the functions

$$(\partial^\beta f)/g \text{ for every } \beta \in \{0, \dots, N-1\}^{\ell+1} \quad (64)$$

to the LN chain defining f, g and $c_{j,i}$ gives an LN chain of effectively bounded format on \mathcal{C} .

We start by showing that these new functions are effectively bounded on \mathcal{C} . For this, note that

$$\partial_j(\partial^\beta f/g) = (\partial^{\beta+e_j} f)/g + \partial^\beta f \partial_j(1/g) = (\partial^{\beta+e_j} f)/g - \alpha_j(\partial^\beta f/g) \quad (65)$$

where e_j is the j -th standard basis vector. Here we use the fact that when the j -th fiber is of type $D, *$ we always have $\alpha_j = 0$. From this it easily follows by induction that

$$\partial^\beta(f/g) = (\partial^\beta f)/g + \sum_{\beta' < \beta} a_{\beta'}(\partial^{\beta'} f)/g \quad (66)$$

where $\beta' < \beta$ means $\beta_i \leq \beta_i$ for every i and $\beta \neq \beta'$. Here a_β are effectively bounded integer coefficients computed in terms of α .

By Lemma 23, the norm of the right hand side of (66) is effectively bounded in terms of the norm $\|f/g\|_{\mathcal{C}^\delta}$, and by induction on β with respect to the ordering above we then obtain effective bounds for $\|(\partial^\beta f)/g\|_{\mathcal{C}}$.

We now show how to write an LN chain for our functions. From (65) we already have an expression for $\partial_j(\partial^\beta f/g)$ in terms of our LN chain with effectively bounded coefficients, unless $\beta_j = N$. In this case write $\beta = Ne_j + \hat{\beta}$ and compute

$$\begin{aligned} \partial_j(\partial^\beta f/g) &= (\partial^{\hat{\beta}} \partial_j^{N+1} f)/g - \alpha_j(\partial^\beta f/g) = \\ &= (\partial^{\hat{\beta}} \sum_{i=0}^N c_{j,i} \partial_j^i f)/g - \alpha_j(\partial^\beta f/g) = \\ &= -\alpha_j(\partial^\beta f/g) + \sum_{i=0}^N c_{j,i} \partial^{\hat{\beta}+ie_j} f/g. \end{aligned} \quad (67)$$

The expression in the right hand side is a polynomial in our LN chain with effectively bounded coefficients, thus finishing the proof in the case $g = \lambda \mathbf{z}^\alpha$. Note that the proof in this case does not rely on the monomialization lemma.

In the general case, write $g = m_g U$ using the monomialization lemma. Since U on $\mathcal{C}^{\sqrt{\delta}}$ is effectively bounded above and below in terms of $\mathcal{F}(g)$, we have an effective bound

$$\|f/m_g\|_{\mathcal{C}^{\sqrt{\delta}}} < \text{Eff}(f, g, \|f/g\|_{\mathcal{C}^\delta}). \quad (68)$$

By the preceding case, f/m_g is LN with the desired format bound. The proof will be finished if we show that $1/U$ is also LN with effectively bounded format on \mathcal{C} . For this we add $1/U$ to our LN chain, and note that

$$\partial_j(1/U) = -\partial_j(U)(1/U)^2 \quad (69)$$

and this gives an LN chain of bounded format because we indeed have an effective upper bound for $\|1/U\|_{\mathcal{C}}$.

When f, g are real then clearly f/g is real on $\mathbb{R}_+\mathcal{C}$, and the chain constructed above is readily seen to be real. \square

Remark 25. *Theorem 9 does not hold for Noetherian functions. For example, even though $e^z - 1$ is Noetherian in $D(1)$ and the division $(e^z - 1)/z$ is restricted, it is not possible to write a Noetherian chain for it in $D(1)$. This is not trivial to check, and is proved in §A.3. This explains why the generalization to LN-functions is crucial for our approach even if one is initially interested only in classical Noetherian function*

4.3.3. Logarithmic derivation.

Theorem 10. *Let \mathcal{C}^δ be an LN cell and $f \in \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta)$. Suppose further that*

$$f : \mathcal{C}^\delta \rightarrow \mathbb{C}^*. \quad (70)$$

Then for every $j = 1, \dots, \ell + 1$ we have

$$\partial_j f/f \in \mathcal{O}_{\text{LN}}(\mathcal{C}) \quad \text{and} \quad \mathcal{F}((\partial_j f/f)|_{\mathcal{C}}) < \text{Eff}(f). \quad (71)$$

If f is real then $\partial_j f/f$ is real.

Proof. Using the monomialization theorem write $f = m_f(\mathbf{z})U(\mathbf{z})$ on the cell $\mathcal{C}^{\sqrt{\delta}}$. Then

$$\partial_j f/f = (\alpha(f)_j f + m_f \partial_j U)/f = \alpha(f)_j + \partial_j U/U. \quad (72)$$

Now $\|\partial_j U\|_{\mathcal{C}} < \text{Eff}(f)$ by Lemma 23 and $1/U < \text{Eff}(f)$ by monomialization, so $\|\partial_j f/f\|_{\mathcal{C}} < \text{Eff}(f)$. The claim now follows by restricted division. The real case follows easily. \square

4.3.4. Stability under pullback.

Theorem 11. *Let $\phi : \mathcal{C}^\delta \rightarrow \hat{\mathcal{C}}$ be an LN map between LN cells. Let $F \in \mathcal{O}_{\text{LN}}(\hat{\mathcal{C}})$. Then*

$$\phi^* F \in \mathcal{O}_{\text{LN}}(\mathcal{C}) \quad \text{and} \quad \mathcal{F}(\phi^* F|_{\mathcal{C}}) < \text{Eff}(\phi, F). \quad (73)$$

If ϕ, F are real then $\phi^ F$ is real.*

Proof. Let F_1, \dots, F_N be the LN chain for F . We claim that adding the pullbacks ϕ^*F_i to the LN chain for ϕ gives an LN chain for ϕ^*F on \mathcal{C} with the effective format bound.

Let $\mathbf{z}_{1..\ell}$ denote the coordinates on \mathcal{C} and $\mathbf{w}_{1..\ell}$ the coordinates on $\hat{\mathcal{C}}$, so that ϕ takes the form $\mathbf{w}_j = \phi_j(\mathbf{z})$. Write Σ for the set of indices j where the j -th fiber of $\hat{\mathcal{C}}$ is of type D_\circ, A and Σ' for the disc fibers. When $j \in \Sigma'$ write r_j for the (constant) radius of the j -th disc fiber. Compute

$$\begin{aligned} \partial_k^{\mathcal{C}^\delta}(F_i \circ \phi) &= \sum_{j=1, \dots, \ell} \frac{\partial F_i}{\partial w_j} \circ \phi \cdot (\partial_k^{\mathcal{C}^\delta} \phi_j) = \\ & \sum_{j \in \Sigma'} \frac{\partial_k^{\mathcal{C}^\delta} \phi_j}{r_j} \cdot (\partial_j^{\hat{\mathcal{C}}} F_i) \circ \phi + \\ & \sum_{j \in \Sigma} \frac{\partial_k^{\mathcal{C}^\delta} \phi_j}{\phi_j} \cdot (\partial_j^{\hat{\mathcal{C}}} F_i) \circ \phi. \end{aligned} \quad (74)$$

Now note that for $j \in \Sigma'$ the map ϕ_j maps into $D(r)$, so $\partial_k^{\mathcal{C}^\delta} \phi_j / r$ is bounded effectively by Lemma 23 on $\mathcal{C}^{\sqrt{\delta}}$, and is therefore LN of effectively bounded format on \mathcal{C} by restricted division. For $j \in \Sigma$ the map ϕ_j is non-vanishing since it maps into a fiber of type D_\circ, A , and the logarithmic derivative $\partial_k^{\mathcal{C}^\delta} \phi_j / \phi_j$ is thus LN with effectively bounded format on \mathcal{C} by logarithmic derivation. Putting these together finishes the proof. Note that it is important here that as a real cellular map, $f(\mathbb{R}_+ \mathcal{C}^\delta) \subset \mathbb{R}_+ \hat{\mathcal{C}}$ so the pullback of F is indeed real valued on $\mathbb{R}_+ \mathcal{C}^\delta$. \square

Stability under pullback implies for example that if $f : D^{1/2} \rightarrow \mathcal{C}^{1/2}$ is an LN-map then the diameter of $f(D)$ with respect to the $\partial^{\mathcal{C}}$ -parametrization is bounded in terms of $\mathcal{F}(f)$. Indeed, for the coordinates of type $D, *$ this is obvious. For D_\circ, A coordinates the pullback is a non-vanishing LN-function and therefore, by monomialization, a unit of bounded logarithmic variation.

Remark 26. *We remark that this metric restriction is not hyperbolic, i.e. it is not a consequence of Schwarz-Pick. For example, the map*

$$f : D(1)^{1/2} \rightarrow A(1, \varepsilon)^{1/2}, \quad f(z) := e^{-\frac{1}{2} \log \varepsilon + \frac{1}{4} z \log \varepsilon} \quad (75)$$

has image of logarithmic width proportional to $|\log \varepsilon|$, which is not uniformly bounded independently of the cell.

Stability under pullback is rather a consequence of metric restrictions related to valency of holomorphic maps. Suppose for example that $f : D^{1/2} \rightarrow A(1, \varepsilon)^{1/2}$ is a univalent map from a disc to an annulus. Then it can be lifted under the exponential cover of $A(1, \varepsilon)^{1/2}$ to a map

$$\tilde{f} : D^{1/2} \rightarrow \{\log \varepsilon - \log 2 < \operatorname{Re} z < \log 2\}. \quad (76)$$

The derivative of \tilde{f} at every point of D is bounded by a constant, because otherwise by Koebe's $\frac{1}{4}$ -theorem theorem it would contain a disc of diameter greater than 2π , in which case $f = e^{\tilde{f}}$ would not be univalent. In general the valency of LN functions is always bounded in terms of the format, and this prevents counterexamples like (75) where the valency is indeed proportional to $|\log \varepsilon|$.

4.4. Root extraction.

Proposition 27. *Let \mathcal{C}^δ be an LN cell and $f \in \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta)$. Suppose further that*

$$f : \mathcal{C}^\delta \rightarrow \mathbb{C}^* \quad (77)$$

and that $f^{1/N}$ has a univalued branch for $N \in \mathbb{N}$. Then

$$f^{1/N} \in \mathcal{O}_{\text{LN}}(\mathcal{C}) \quad \text{and} \quad \mathcal{F}(f^{1/N}) < \text{Eff}(f). \quad (78)$$

If f is real and positive on $\mathbb{R}_+\mathcal{C}^\delta$ then $f^{1/N}$ can be chosen to be real.

Proof. By monomialization write $f = m_f U$ on $\mathcal{C}^{\sqrt{\delta}}$. The root $f^{1/N}$ is univalued if and only if $\alpha(f) = \mathbf{0}$ modulo N . In particular note that this implies $N < \text{Eff}(f)$. We will show that both $m_f^{1/N}$ and $U^{1/N}$ are LN with effectively bounded format on \mathcal{C} .

Starting with $m_f^{1/N}$, by assumption it is a monomial effectively bounded on $\mathcal{C}^{\sqrt{\delta}}$. Any such monomial is LN with bounded format. Indeed, it satisfies

$$\partial_j^{\mathcal{C}}(m_f^{1/N}) = \frac{\alpha_j}{N} m_f^{1/N}. \quad (79)$$

As for U , we write

$$\partial_j^{\mathcal{C}}(U^{1/N}) = \frac{U^{1/N} \partial_j^{\mathcal{C}} U}{NU}. \quad (80)$$

Since U is effectively bounded from above and below and has a well-defined root $U^{1/N}$, the function $1/U$ is LN with bounded format by restricted division and the equation above gives an LN chain for $U^{1/N}$.

In the real case it is clear that $f^{1/N}$ defined above is real if one chooses $U^{1/N}$ to be the (unique) positive branch. \square

4.5. Removable singularities. The following is an LN version of the Riemann removable singularity theorem.

Proposition 28. *Let $F \in \mathcal{O}_{\text{LN}}(\mathcal{C} \odot D_\circ(r))$ for some LN cell \mathcal{C} of length ℓ . Then $F_0 := F(\mathbf{z}, 0) \in \mathcal{O}_{\text{LN}}(\mathcal{C})$ and $\mathcal{F}(F_0) \leq \mathcal{F}(F)$. If F is real then F_0 is real.*

Proof. Suppose the LN chain of F is given by functions F_1, \dots, F_N . By assumption these functions are all bounded on \mathcal{C} , and by the Riemann removable singularity theorem they all extend to well-defined functions $F_{0,i}(\mathbf{z}, 0)$. In particular the derivation rules

$$\partial_j^{\mathcal{C}}(F_i) = G_{i,j}(F_1, \dots, F_N), \quad j = 1, \dots, \ell \quad (81)$$

continue holomorphically to the same identities on $F_{0,i}$. The statement follows.

When F is real on $\mathcal{C} \odot D_\circ(r)$ then its limit F_0 is real on \mathcal{C} , and the chain above is real by definition. \square

Remark 29. *We cannot claim in Proposition 28 that F extends to an LN function over $\mathcal{C} \odot D(r)$ in general, because there is no clear way to extend the derivation rules in the $\partial_{\ell+1}^{\mathcal{C}}$ direction. For example $\sin(z)/z$ is LN in $D_\circ(1)$, as one may easily verify (or by restricted division). But we do not know whether it is LN in $D(1)$.*

Corollary 30. *Let $F \in \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta \odot D_\circ(r))$ for some LN cell \mathcal{C} of length ℓ . Write a Taylor expansion*

$$F(\mathbf{z}) = \sum_{k=0}^{\infty} a_j(\mathbf{z}_{1..\ell}) \mathbf{z}_{\ell+1}^k. \quad (82)$$

Then $a_j \in \mathcal{O}_{\text{LN}}(\mathcal{C})$ and $\mathcal{F}(a_j|_{\mathcal{C}}) \leq \text{Eff}(F, j)$.

Proof. First note that we have a Taylor instead of Laurent expansion because of the removable singularity theorem. For a_0 the claim is Proposition 28. Now write

$$\frac{\partial F}{\partial \mathbf{z}_{\ell+1}} = \frac{\partial^{\mathcal{C}^\delta \odot D_\circ(1)} F}{\mathbf{z}_{\ell+1}} \quad (83)$$

and note that this is a restricted division on $\mathcal{C}^\delta \odot D_\circ(1)$. Thus $\frac{\partial F}{\partial \mathbf{z}_{\ell+1}} \in \mathcal{O}_{\text{LN}}(\mathcal{C} \odot D_\circ(r))$ with an effective bound on the format. The claim for a_1 thus follows from Proposition 28 again, and repeating this argument we get a similar bound for a_j . \square

4.6. The ν -cover of a cell. Recall the notion of a ν -cover of a cell from [10, Section 2.6] making small adaptations for the LN setting.

Definition 31 (The ν -cover of a cell). *Let \mathcal{C} be an LN cell of length 1. For $\mathcal{C} = D_\circ, A$ and $\nu \in \mathbb{Z}$ we define the ν -cover $\mathcal{C}_{\times\nu}$ by*

$$D_\circ(r)_{\times\nu} := D_\circ(r^{1/\nu}) \quad A(r_1, r_2)_{\times\nu} := A(r_1^{1/\nu}, r_2^{1/\nu}) \quad (84)$$

For $\mathcal{C} = D(r), *$ the cover $\mathcal{C}_{\times\nu}$ is defined only for $\nu = 1$. In all cases we define $R_\nu : \mathcal{C}_{\times\nu} \rightarrow \mathcal{C}$ by $R_\nu(z) = z^\nu$.

Let \mathcal{C} be a cell of length ℓ and let $\boldsymbol{\nu} = (\nu_1, \dots, \nu_\ell) \in \pi_1(\mathcal{C})$ be such that $\nu_j | \nu_k$ whenever $j > k$ and $G_j = G_k = \mathbb{Z}$. Define the $\boldsymbol{\nu}$ -cover $\mathcal{C}_{\times\boldsymbol{\nu}}$ of \mathcal{C} and the associated cellular map $R_\boldsymbol{\nu} : \mathcal{C}_{\times\boldsymbol{\nu}} \rightarrow \mathcal{C}$ by induction on ℓ . For $\mathcal{C} = \mathcal{C}_{1..\ell-1} \odot \mathcal{F}$ we let

$$\mathcal{C}_{\times\boldsymbol{\nu}} := (\mathcal{C}_{1..\ell-1})_{\times\nu_{1..\ell-1}} \odot (R_{\boldsymbol{\nu}_{1..\ell-1}}^* \mathcal{F}_{\times\nu_\ell}) \quad (85)$$

We define $R_\boldsymbol{\nu}(\mathbf{z}_{1..\ell}) := \mathbf{z}^\boldsymbol{\nu}$.

As explained in [10, Section 2.6], the divisibility conditions on $\boldsymbol{\nu}$ ensure that the radii of $\mathcal{F}_{\times\nu_\ell}$, which are a-priori multivalued roots, are in fact univalued. We will usually consider the ν -cover with $\nu \in \mathbb{N}$, meaning that we take $\boldsymbol{\nu}$ with $\nu_i = \nu$ when $G_i = \mathbb{Z}$ and $\nu_i = 1$ otherwise.

The pullback to a $\boldsymbol{\nu}$ -cover will be used in our treatment to resolve the ramification of multivalued cellular maps. We record a simple proposition concerning the interaction between extensions and ν -covers.

Proposition 32. *Let \mathcal{C}^δ be an LN cell and $\nu \in \mathbb{N}$. Then $\mathcal{C}_{\times\nu}$ is LN and admits a $2\delta^{1/\nu}$ -extension, and the covering map R_ν extends to an LN map $R_\nu : (\mathcal{C}_{\times\nu})^{2\delta^{1/\nu}} \rightarrow \mathcal{C}^\delta$. Moreover*

$$\mathcal{F}(R_\nu|_{(\mathcal{C}_{\times\nu})^{2\delta^{1/\nu}}}) < \text{Eff}(\mathcal{C}^\delta, \nu). \quad (86)$$

When \mathcal{C}^δ is real the cover $\mathcal{C}_{\times\nu}$ can also be chosen to be real.

Proof. The main difference compared with the usual complex cellular case is that we get a slightly weaker extension $2\delta^{1/\nu}$ instead of $\delta^{1/\nu}$. The proposition is proved by using stability under pullbacks and Proposition 27 for extracting univalued roots; since each of these steps involves passing to some (arbitrarily small) extension we only get the LN-ness of R_ν on a slightly smaller extension than in the complex case. In fact 2 is an arbitrary choice and any constant larger than 1 would have sufficed.

For the real case, note that whenever we have a real cell $\mathcal{C} \odot D_\circ(r)$ (or similarly with other fiber types) the radius r must have a constant sign on $\mathbb{R}_+\mathcal{C}$, since it is non-vanishing and real there. Up to changing r with $-r$, which does not affect the cell itself, we can assume that the radius is positive and can therefore choose $r^{1/\nu}$ in the definition of $(D_\circ(r))_{\times\nu}$ to be real as well. \square

4.7. Refinement. The following theorem shows that one can always cover a cell with a given extension by cells with a larger extension. Following [10] we state this with $\{\rho\}$ -extensions because in this natural choice of the parameter the asymptotic bounds is more precise. However in the present paper we will not make use of these precise bounds.

Theorem 12 (Refinement theorem). *Let $\mathcal{C}^{\{\rho\}}$ be a (real) cell and $0 < \sigma < \rho$. Then there exists a (real) cellular cover $\{f_j : \mathcal{C}_j^{\{\sigma\}} \rightarrow \mathcal{C}^{\{\rho\}}\}$ of size $\text{poly}_\ell(\rho, 1/\sigma)$. Moreover $\mathcal{F}(f_j) < \text{Eff}(\mathcal{C}^{\{\rho\}})$.*

The proof of this theorem is the same as in [10, Theorem 9], using stability under pullback of LN maps. We leave the straightforward verification of the details to the reader.

5. PROOF OF THE CPT FOR THE LN CATEGORY

We will prove the CPT for a single function F . The general case follows from this in a straightforward way by first finding cells compatible with F_1 , then covering them by cells compatible with F_2 and so on. Note that the proof of [10] does not seem to extend to the LN category as it requires working with various holomorphic functions that are not LN.

We proceed by induction on the length of the cell \mathcal{C} . So suppose the theorem is established for cells of length ℓ , and we will prove the statement for cells of the form $\mathcal{C} \odot \mathcal{F}$ where \mathcal{C} is of length ℓ . To simplify the notation we will write $\mathbf{z} = \mathbf{z}_{1..\ell}$ for the coordinates on \mathcal{C} and w for the coordinate on \mathcal{F} .

In the case $\mathcal{F} = *$ the map F pulls back to a map $F' \in \mathcal{O}_{\text{LN}}(\mathcal{C})$ and the claim follows from the CPT for \mathcal{C} . For the remaining cases, we will first reduce the cases $\mathcal{F} = D_\circ, A$ to the case $\mathcal{F} = D(r)$ and then prove the disc case at the end.

By applying the refinement theorem we can assume that $\mathcal{C} \odot \mathcal{F}$ admits δ -extension with some small δ to be chosen later. Note that in this reduction the format of $\mathcal{C}^\delta \odot \mathcal{F}^\delta$ remains independent of δ .

5.1. Reducing $\mathcal{F} = A(r_1, r_2)$ to $\mathcal{F} = D(r)$. Write a Laurent expansion for F ,

$$F(\mathbf{z}, w) = \sum_{j=-\infty}^{\infty} a_j(\mathbf{z})w^j. \quad (87)$$

Using Proposition 15 we have integers p, q with $|p| + |q|$ effectively bounded, and some effectively bounded $M > 0$, such that $F(\mathbf{z}, \cdot)$ has the (p, q, M) Laurent domination property for every $\mathbf{z} \in \mathcal{C}^\delta$. Write

$$F(\mathbf{z}, w) = \sum_{j=p}^q a_j(\mathbf{z})w^j + R(\mathbf{z}, w). \quad (88)$$

Unfortunately, even though $a_j(\mathbf{z})$ are bounded holomorphic on \mathcal{C} , we do not know if they are LN. We will approximate them by a slightly more complicated argument.

Write $\partial_w := \partial_{\ell+1}^{\mathcal{C} \odot \mathcal{F}}$. Then

$$\partial_w F(\mathbf{z}, w) = \sum_{j=p}^q j a_j(\mathbf{z})w^j + \partial_w R(\mathbf{z}, w). \quad (89)$$

Define the constant-coefficient differential operators D_k by

$$D_k = \prod_{j=p, \dots, k-1, k+2, \dots, q} \frac{\partial_w - j \cdot \mathbb{I}}{k - j} \quad (90)$$

so that

$$A_k := D_k F(\mathbf{z}, w) = a_k(\mathbf{z})w^k + D_k R(\mathbf{z}, w). \quad (91)$$

Claim 1. *We have*

$$D_k R(\mathbf{z}, w) \leq M' \delta \max_{j=p, \dots, q} |a_j(\mathbf{z})w^j| \quad (92)$$

for every $(\mathbf{z}, w) \in \mathcal{C}^\delta \odot \mathcal{F}$, where $M' < \text{Eff}(F)$.

Proof. Since $|p|, |q|$ are bounded effectively in F it is enough to prove the claim with ∂_w^l instead of D_k , with the bound depending effectively on l . The case $l = 0$ is Proposition 16, and the case of general l is proved in essentially the same way. \square

In particular, let $\varepsilon > 0$ be some constant to be chosen later. Since M depends only on the format of F which is independent of our choice of δ , we can choose $\delta < \varepsilon/(2M)$ and in this case

$$1 - \varepsilon < \frac{\max_{j=p, \dots, q} |a_j(\mathbf{z})w^j|}{\max_{j=p, \dots, q} |A_j(\mathbf{z}, w)|} < 1 + \varepsilon \quad (93)$$

holds uniformly for $(\mathbf{z}, w) \in \mathcal{C}^\delta \odot \mathcal{F}$.

Note that $A_k \in \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta \odot \mathcal{F}^\delta)$. Consider two pullbacks

$$A_k^1(\mathbf{z}) := A_k(\mathbf{z}, r_1(\mathbf{z})) \quad A_k^2(\mathbf{z}) := A_k(\mathbf{z}, r_2(\mathbf{z})) \quad (94)$$

which are both LN functions of effectively bounded format. Apply the CPT inductively to \mathcal{C}^δ and the collection of functions

$$\{A_k^b, A_k^b - A_j^b : j, k = p, \dots, q \text{ and } b = 1, 2\}. \quad (95)$$

For each of the resulting maps $f_j : \mathcal{C}^\delta \rightarrow \mathcal{C}^\delta$ we pull back \mathcal{F} along f_j to give a cell $\mathcal{C}_j^\delta \odot (f_j^* \mathcal{F}^\delta)$. In order to get a cellular cover for the original cell $\mathcal{C} \odot \mathcal{F}$ it will suffice to prove the CPT for each of these cells and the pullback of F separately. In other words, we may replace \mathcal{C} by each of the cells \mathcal{C}_j , and simply assume without loss of generality below that \mathcal{C}^δ is already compatible with the functions (95).

Assume for simplicity of the notation that none of the A_k^b are identically vanishing on \mathcal{C} . If they are then they should simply be removed from consideration below, which would only make the notation slightly more cumbersome. By (95) we have

$$A_k^1/A_j^1 : \mathcal{C}^\delta \rightarrow \mathbb{C} \setminus \{0, 1\}. \quad (96)$$

The fundamental lemma then implies that these ratios “do not move much” on $\mathcal{C}^{\sqrt{\delta}}$.

We claim first that we may assume $A_p^1/A_j^1 > N$ for some large N to be chosen later, for every $j > p$, uniformly over $\mathcal{C}^{\sqrt{\delta}}$. Indeed, suppose this fails for some j at some point in $\mathcal{C}^{\sqrt{\delta}}$. Then by the fundamental lemma with a suitable choice of δ we will have $A_p^1/A_j^1 < N + 1$ uniformly over $\mathcal{C}^{\sqrt{\delta}}$. Then (93) also implies that

$$\frac{\max_{j=p+1, \dots, q} |a_j(\mathbf{z})r_1(\mathbf{z})^j|}{a_p(\mathbf{z})r_1(\mathbf{z})^p} < 2N + 2 \quad (97)$$

Then in fact F has the $(p+1, q, \tilde{M})$ Laurent domination property on $\mathcal{C}^{\sqrt{\delta}} \odot \mathcal{F}^\delta$ for some slightly larger but still effective \tilde{M} . Indeed, whenever the maximum in the domination property is achieved at index p the inequality above implies that the same maximum is achieved at $k > p$ with $\tilde{M} = (2N+2)M$. In this case we can finish by our induction on $q-p$.

Applying a similar reasoning to A_k^2/A_j^2 we conclude that we may assume $A_q^1/A_j^1 > N$ for every $j < q$, uniformly over \mathcal{C}^δ . Under these conditions Claim 1 also implies, with suitable choice of δ , that

$$\begin{aligned} 1 - \frac{1}{N} &< \frac{A_p^1(\mathbf{z})}{a_p(\mathbf{z})r_1(\mathbf{z})^p} < 1 + \frac{1}{N} \\ 1 - \frac{1}{N} &< \frac{A_q^2(\mathbf{z})}{a_q(\mathbf{z})r_2(\mathbf{z})^q} < 1 + \frac{1}{N}. \end{aligned} \quad (98)$$

Consider the quotient

$$\tilde{s}_{p,q} := \frac{A_p^1(\mathbf{z})r_2(\mathbf{z})^q}{A_q^2(\mathbf{z})r_1(\mathbf{z})^p}. \quad (99)$$

We claim that $\tilde{s}_{p,q} \in \mathcal{O}_{\text{LN}}(\mathcal{C}^{\sqrt{\delta}})$ has effectively bounded format. It is enough to show that it is a restricted division with a bound on the norm. And indeed, by (98) we have

$$(1 - 1/N)^2 < \tilde{s}_{p,q}(\mathbf{z})/r_{p,q}(\mathbf{z})^{q-p} < (1 + 1/N)^2 \quad (100)$$

where

$$r_{p,q} = \sqrt[q-p]{\frac{a_p(\mathbf{z})}{a_q(\mathbf{z})}}. \quad (101)$$

To see that this latter $r_{p,q}$ is effectively bounded, note that it is the radius $|w| = r_{p,q}$ where $|a_p w^p| = |a_q w^q|$. Since by our assumption $a_p w^p$ is dominant on $|w| = r_1$ and $a_q w^q$ is dominant on $|w| = r_2$, we must have $r_1 < r_{p,q} < r_2$.

Finally, we wish to extract the $(q-p)$ -th root of $\tilde{s}_{p,q}$. For this purpose consider the covering cell $\hat{\mathcal{C}} := \mathcal{C}_{\times(q-p)}$ and pull back our fiber \mathcal{F} to $\hat{\mathcal{C}}$. This is again an LN cell of effectively bounded format, and as before it will suffice to prove the claim for this new cell (note that we get a slightly smaller extension for $\hat{\mathcal{C}}$ but it is easy to compensate, e.g. by refining $\hat{\mathcal{C}}$). Crucially on $\hat{\mathcal{C}}$ the LN function $\tilde{s}_{p,q}$ admits a $(q-p)$ -th root. Without loss of generality we replace \mathcal{C} by $\hat{\mathcal{C}}$ and simply assume below that $\tilde{s}_{p,q}$ admits a $(q-p)$ -th root $\tilde{r}_{p,q} \in \mathcal{O}_{\text{LN}}(\mathcal{C}^{\sqrt{\delta}})$.

Now cover our original fiber $A(r_1, r_2)$ as

$$A(r_1, r_2) \subset A(r_1, c\tilde{r}_{p,q}) \cup A(c^{-1}\tilde{r}_{p,q}, r_2) \quad (102)$$

where $c > 1$ is arbitrarily close to 1, taken simply to cover the circle of radius $\tilde{r}_{p,q}$. It will suffice to prove the claim for the pullback of F to

$$\mathcal{C}^{\sqrt{\delta}} \odot A(r_1, c\tilde{r}_{p,q}) \text{ and } \mathcal{C}^{\sqrt{\delta}} \odot A(c^{-1}\tilde{r}_{p,q}, r_2) \quad (103)$$

We proceed with the first of these, the other being analogous. On the circle $|w| = c\tilde{r}_{p,q}$ we already have for an appropriate choice of N the estimate

$$\frac{|a_q w^q|}{|a_p w^p|} < 2. \quad (104)$$

Indeed, on the circle of radius $r_{p,q}$ this ratio is 1, and by (100) the ratio $\tilde{r}_{p,q}/r_{p,q}$ is arbitrarily close to 1 for suitable N and c is arbitrarily close to 1. It follows on

$\mathcal{C}^{\sqrt{\delta}} \odot A(r_1, c\tilde{r}_{p,q})$ we have the $(p, q-1, 2M)$ domination property and we can finish as before by induction on $q-p$. This finishes the proof.

In the real case, the functions A_j^b defined above are also real. Then $\tilde{s}_{p,q}$ are real, and the roots $\tilde{r}_{p,q}$ can be chosen to be real, so the new annuli we produce are again real.

5.2. Reducing $\mathcal{F} = D_\circ(r)$ to $\mathcal{F} = D(r)$. Without loss of generality by pulling back along a rescaling map we may assume $\mathcal{F} = D_\circ(1)$. Write a Taylor expansion for F ,

$$F(\mathbf{z}, w) = \sum_{j=0}^{\infty} a_j(\mathbf{z})w^j. \quad (105)$$

where according to Corollary 30 we have $a_j \in \mathcal{O}(\mathcal{C}^\delta)$ with $\mathcal{F}(a_j) \leq \text{Eff}(F, j)$. (Formally we start with a slightly larger extension than δ to get the format bounded on \mathcal{C}^δ .) The proof is now quite similar to the one given in [10, Section 7.2.2] so we outline the argument briefly only to show that the process is indeed effective.

Using Proposition 15 we have an integer q , effectively bounded, and some effectively bounded $M > 0$, such that $F(\mathbf{z}, \cdot)$ has the (q, M) Taylor domination property for every $\mathbf{z} \in \mathcal{C}^\delta$. First apply the CPT to \mathcal{C}^δ with the functions $a_j, a_j - a_k$ for $j, k = 0, \dots, q$. As before, pulling back the fiber to each of the resulting cells we may assume without loss of generality that these functions are all non-vanishing (or some are identically vanishing, which does not affect the arguments below). Then

$$a_j/a_k : \mathcal{C}^\delta \rightarrow \mathbb{C} \setminus \{0, 1\} \quad (106)$$

and the fundamental lemma then implies that these ratios “do not move much” on $\mathcal{C}^{\sqrt{\delta}}$. In particular, we can divide the pairs $(j, k) \in \{0, \dots, q\}^2$ with $j > k$ into a set Σ where a_j/a_k is uniformly bounded by 2, and its complement where a_j/a_k is uniformly bounded below by 1. For $(j, k) \in \Sigma$ we define the radii

$$r_{j,k}(\mathbf{z}) := \sqrt[j-k]{\frac{a_k(\mathbf{z})}{a_j(\mathbf{z})}}. \quad (107)$$

Note that the divisions here are restricted by our definition of Σ . As in §5.1 we may, after pulling back to a $q!$ -cover of \mathcal{C} and refining, assume that $r_{j,k} \in \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta)$ with effectively bounded format.

By definition $r_{j,k}$ is the circle where $|a_j w^j| = |a_k w^k|$. It is easy to see that if $|w|$ has distance at least $\log 3$ to each of the radii $r_{j,k}$ then F is non-vanishing, because one term $a_t w^t$ would dominate all other terms up to order q as well as the Taylor residue. Since the radii $r_{j,k}$ are “almost constant” on \mathcal{C} one can group them into at most q^2 disjoint “special annuli” of effectively bounded logarithmic width, such that every radius along with the annulus of logarithmic width $\log 3$ around it is contained in one of these annuli. The annuli that remain between adjacent pairs of special annuli cannot contain zeros, so they are already compatible with F . We cover each of the special annuli by discs (with their number effectively bounded in terms of the logarithmic width), and thus reduce the problem to the case $\mathcal{F} = D(r)$. For more details on this clustering construction see [10, Section 7.2.2].

In the real case the functions a_j are real, and the radii $r_{j,k}$ can be chosen to be real. Therefore the “special annuli” above are real, and we can cover their real part by real discs centered over the positive real line. Thus we obtain a real cover.

5.3. **The case $\mathcal{F} = D(r)$.** We start similarly to the case of $D_\circ(r)$. Without loss of generality by pulling back along a rescaling map we may assume $\mathcal{F} = D(1)$. It will also suffice to prove that there exists some $\varepsilon > 0$ with $1/\varepsilon < \text{Eff}(F)$ and a covering for $\mathcal{C} \odot D(\varepsilon)$ compatible with F . This simply amounts to choosing our cells (by refinement) to have $\varepsilon\delta$ -extension instead of a δ -extension.

Write a Taylor expansion for F ,

$$F(\mathbf{z}, w) = \sum_{j=0}^{\infty} a_j(\mathbf{z})w^j. \quad (108)$$

We see again that $F(\mathbf{z}, \cdot)$ has the (q, M) Taylor domination property with effective q, M in $D(1)$ for every $\mathbf{z} \in \mathcal{C}^\delta$. After suitable application of the CPT we have

$$a_j/a_k : \mathcal{C}^\delta \rightarrow \mathbb{C} \setminus \{0, 1\} \quad (109)$$

and the fundamental lemma then implies that these ratios do not move much on $\mathcal{C}^{\sqrt{\delta}}$. We proceed by induction on q , so suppose for functions with (q', M') domination and $q' < q$ the CPT is already proved.

For the base case $q = 0$, the free term $a_0(\mathbf{z})$ dominates the Taylor residue on a ball of radius ε with $1/\varepsilon = \text{Eff}(F)$ by Proposition 16 so F is already non-vanishing in $\mathcal{C} \odot D(\varepsilon)$ and the claim is proved.

Proceeding now with general $q > 0$, we may as well assume that at some point $\mathbf{z} \in \mathcal{C}^\delta$ we have $|a_q(\mathbf{z})| \geq |a_j(\mathbf{z})|$ for all j , because otherwise we have the $(q-1, M)$ domination property and are done by induction. Since the ratios are nearly constant, we have $|a_q(\mathbf{z})| \geq |a_j(\mathbf{z})|/2$ uniformly in $\mathcal{C}^{\sqrt{\delta}}$. It follows using Proposition 17 that $\partial_w F(\mathbf{z}, w)$ has the $(q-1, M)$ domination property on the slightly smaller fiber $D(1/2)$.

Apply the inductive case to $\partial_w F$. We get a covering of $\mathcal{C} \odot \mathcal{F}$ and as usual it will suffice to now prove the CPT for the pullback of F to each of the resulting cells. Crucially, since all the maps in the CPT are affine in the final variable (see Remark 33), these pullbacks still the derivative in the ∂_w direction either non-vanishing or identically zero. In other words we are reduced to proving the CPT under the additional assumption that $\partial_w F$ is already compatible with $\mathcal{C}^\delta \odot D(1)$.

If $\partial_w F \equiv 0$ then we can find a covering of \mathcal{C}^δ compatible with $F(\mathbf{z}, 0)$ by induction on ℓ and then multiply each cell by a constant $D(1)$. So assume $\partial_w F$ is nowhere vanishing. By Proposition 18 there is an $\varepsilon > 0$ with $1/\varepsilon = \text{Eff}(F)$ such that $\partial_w F(\mathbf{z}, \cdot)$ has the $(0, 1/4)$ -domination property in $D(\varepsilon)$ for every $\mathbf{z} \in \mathcal{C}^\delta$. To simplify the notation by rescaling $D(\varepsilon)$ we may as well assume that $\partial_w F$ has the $(0, 1/4)$ -domination property in the fiber $D(1)$. Then F has the $(1, 1/4)$ -domination property. Write

$$F(\mathbf{z}, w) = a_0(\mathbf{z}) + a_1(\mathbf{z})w + R(\mathbf{z}, w). \quad (110)$$

Perform an inductive CPT for \mathcal{C}^δ and the functions $a_0(\mathbf{z}), a_1(\mathbf{z}), a_0(\mathbf{z}) - a_1(\mathbf{z})$. By the same reduction used before, after pulling back to the resulting cells we may as well assume that

$$a_1/a_0 : \mathcal{C}^\delta \rightarrow \mathbb{C} \setminus \{0, 1\}. \quad (111)$$

By the fundamental lemma, on $\mathcal{C}^{\sqrt{\delta}}$ we uniformly have either

$$|a_1/a_0| > 100 \quad \text{or} \quad |a_1/a_0| < 101. \quad (112)$$

In the latter case, the $(1, 1/4)$ -domination property implies that in the disc of radius $D(1/200)$ the term $a_0(\mathbf{z})$ already dominates $a_1(\mathbf{z}) + R(\mathbf{z}, w)$, so F has no zeros. In

this case we can just use a cover $\mathcal{C}^{\sqrt{\delta}} \odot D(1/10)$. So assume we are in the case $|a_1/a_0| > 100$.

In this case by a similar reasoning on $\mathcal{C}^{\sqrt{\delta}} \odot D(1/2)$ the term $a_1(\mathbf{z})$ dominates $a_0(\mathbf{z})$ as well as $R(\mathbf{z}, w)$. Taking a restricted division

$$\tilde{F} := \frac{F}{a_1(\mathbf{z})} = a_0(\mathbf{z}) + w + \tilde{R}(\mathbf{z}, w) \quad (113)$$

we have an effective bound for the format of \tilde{F} , and it will be enough to find a covering compatible with \tilde{F} since its zeros agree with those of F . Below we simply replace F by \tilde{F} and assume

$$a_1(\mathbf{z}) \equiv 1 \quad |a_0(\mathbf{z})| < \frac{1}{100} \quad |R(z, w)| < \frac{1}{2} \quad (114)$$

uniformly on $\mathcal{C}^{\sqrt{\delta}} \odot D(1/2)$. Under these conditions it is clear that $F(\mathbf{z}, w)$ has a single zero

$$w = w(\mathbf{z}) \in D(1/2) \quad (115)$$

for each $\mathbf{z} \in \mathcal{C}^{\sqrt{\delta}}$ because the term w in the Taylor expansion is dominant over the circle of radius $1/2$, and in fact the zeros lies in $D(1/10)$ because w is also dominant there. Moreover $w(\mathbf{z})$ is holomorphic on $\mathcal{C}^{\sqrt{\delta}}$. If we show that it is LN with format $\text{Eff}(F)$ then we can finish the proof. Indeed, a covering for $\mathcal{C} \odot D(1/2)$ can then be obtained using the cell $\mathcal{C} \odot *$ with the map $(\mathbf{z}, *) \rightarrow (\mathbf{z}, w(\mathbf{z}))$ to cover the zeros of F and the cell $\mathcal{C} \odot D_\circ(1/5)$ with the map $(\mathbf{z}, w) \rightarrow (\mathbf{z}, w + w(\mathbf{z}))$ to cover $\mathcal{C} \odot D(1/10)$. Note that this map is compatible with F because F has no zeros other than $w(\mathbf{z})$ in $D(1/2)$.

Note that $|\partial_w F|$ is bounded below by $1/2$ on $\mathcal{C}^\delta \odot D(1/2)$. To see this write

$$\partial_w F = 1 + \partial_w R(\mathbf{z}, w) \quad (116)$$

and use the $(0, 1/4)$ -domination property to bound $\partial_w R(\mathbf{z}, w)$ on $D(2/3)$. Therefore the functions

$$D_j \in \mathcal{O}(\mathcal{C}^{\sqrt{\delta}} \odot D(1/2)), \quad D_j := -\frac{\partial_j F}{\partial_w F} \quad (117)$$

are given by restricted division and hence have an effectively bounded format. Moreover,

$$0 = \partial_j F(\mathbf{z}, w(\mathbf{z})) = (\partial_j F)(\mathbf{z}, w(\mathbf{z})) + (\partial_w F)(\mathbf{z}, w(\mathbf{z})) \cdot \partial_j w(\mathbf{z}) \quad (118)$$

where we crucially used the fact that the fiber is of type D , so $\partial_w = \frac{\partial}{\partial w}$. Concluding,

$$\partial_j w(\mathbf{z}) = D_j(\mathbf{z}, w(\mathbf{z})) \quad (119)$$

for $\mathbf{z} \in \mathcal{C}^{\sqrt{\delta}}$.

We are finally in position to construct an LN chain for the function $w(\mathbf{z})$. Let F_1, \dots, F_N be a LN chain containing the functions D_i and the coordinate $F_1 \equiv w$. Now consider the system of equations

$$\partial_j F_i(\mathbf{z}, w(\mathbf{z})) = (\partial_j F_i + \partial_w F_i \cdot D_j)(\mathbf{z}, w(\mathbf{z})). \quad (120)$$

Since $\partial_j F_i, \partial_w F_i$ and D_j are polynomials in the F_i , this is an LN chain of format $\text{Eff}(F)$ for $F_i(\mathbf{z}, w(\mathbf{z}))$ on $\mathcal{C}^{\sqrt{\delta}}$. In particular we got chain for $F_1(\mathbf{z}, w(\mathbf{z})) \equiv w(\mathbf{z})$, finishing the proof.

In the real case, the reduction up to (114) preserves the realness of F . (If we are in the other case where F has no zeros then the cell we use is clearly real.) At this point the real function F admits a unique root, so it follows by symmetry that

the root $w(\mathbf{z})$ must be real. It then easily follows that the two cells used above and corresponding maps are indeed real.

5.4. Simple cellular maps. It is sometimes useful to know that the maps produced by the CPT are of a somewhat special form. By inspection of the proof of the CPT we may assume in the CPT that all maps are simple in the following sense.

Say that a cellular map $f : \mathcal{C} \rightarrow \hat{\mathcal{C}}$ is *basic simple* if each coordinate $w_j = f_j(\mathbf{z}_{1..j})$ is either

- (1) An affine map $\mathbf{z}_j \rightarrow \rho \mathbf{z}_j + \phi_j(\mathbf{z}_{1..j-1})$ with $\rho \in \mathbb{R}_+$, or
- (2) A covering maps $\mathbf{z}_j \rightarrow \mathbf{z}_j^k$ for $k \in \mathbb{N}$, only in the case that the j -th fiber is of type A, D_\circ .

A simple map is a composition of basic simple maps. Note that simple maps are always bijective from $\mathbb{R}_+\mathcal{C}$ to $f(\mathbb{R}_+\mathcal{C})$, as they are monotone on each coordinate.

Remark 33. *With respect to the final variable \mathbf{z}_ℓ we only get affine maps regardless of the fiber type.*

6. EFFECTIVE MODEL COMPLETENESS AND O-MINIMALITY OF \mathbb{R}_{LN}

In this section we establish the effective model completeness and o-minimality of the structure \mathbb{R}_{LN} , proving Theorem 2.

For each real LN-cell \mathcal{C}^δ and real LN-function $F \in \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta)$ let \mathbb{R}_+F denote the restriction $\mathbb{R}_+F : \mathbb{R}_+\mathcal{C}^\delta \rightarrow \mathbb{R}$. Define the structure \mathbb{R}_{LN} to be the structure generated by the relations $=, <$ and the graphs $\text{Gr } \mathbb{R}_+F|_{\mathcal{C}}$ whenever $F \in \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta)$ is real. We declare the $\text{Gr } \mathbb{R}_+F|_{\mathcal{C}}$ to have format $\mathcal{F}(F) + 1/(1 - \delta)$. Consider the language \mathcal{L}_{LN} having relation symbols $=, <$ and relation symbols for the graphs above and for the graphs of $+, *$. Note that while the restrictions of $+, *$ are LN, here we add the full graphs on $\mathbb{R}^2 \rightarrow \mathbb{R}$. We declare the formats of $=, <, +, *$ to be 1. We also include constants for all real numbers in our language and define the format of the singletons to be 1. Finally let $\Omega_{\mathcal{F}}$ be the filtration generated by all these sets with their respective formats.

Remark 34. *Note that even though we define \mathbb{R}_{LN} to be the structure generated by the graphs of real LN-functions restricted to the real part of the cell, we later prove in Proposition 48 that \mathbb{R}_{LN} also contains the graphs of complex LN-functions on complex LN-cells.*

6.1. Boundary equations. Write $\mathcal{C}_F := \mathcal{C} \times D(\|F\|_{\mathcal{C}})$ so

$$\text{Gr } F \subset \mathcal{C}_F \quad \text{and} \quad \text{Gr } \mathbb{R}_+F \subset \mathbb{R}_+\mathcal{C}_F. \quad (121)$$

We introduce a set of equations for the walls of the cell \mathcal{C} and the graph of F .

Definition 35. *Let \mathcal{C}^δ be a real LN-cell. Define the boundary equations*

$$B(\mathcal{C}) \subset \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta) \quad (122)$$

inductively as follows. If \mathcal{C} is of length zero $B(\mathcal{C}) = 0$. If $\mathcal{C} = \mathcal{C}_{1..l} \odot \mathcal{F}$ we define $B(\mathcal{C}) := B(\mathcal{C}_{1..l}) \cup B(\mathcal{F})$ where

$$\begin{aligned} B(*) &:= \{\mathbf{z}_{\ell+1}\} \\ B(D(r)) &:= \{\mathbf{z}_{\ell+1} - r, \mathbf{z}_{\ell+1} + r\} \\ B(D_\circ(r)) &:= \{\mathbf{z}_{\ell+1}, \mathbf{z}_{\ell+1} - r, \mathbf{z}_{\ell+1} + r\} \\ B(A(r_1, r_2)) &:= \{\mathbf{z}_{\ell+1} - r_1, \mathbf{z}_{\ell+1} + r_1, \mathbf{z}_{\ell+1} - r_2, \mathbf{z}_{\ell+1} + r_2\} \end{aligned} \quad (123)$$

If $F : \mathcal{C}^\delta \rightarrow \mathbb{C}$ is a real LN-function on a cell of length ℓ we define $B(F|_e) \subset \mathcal{O}_{\text{LN}}(\mathcal{C}_F^\delta)$ to be

$$B(F|_e) := B(\mathcal{C}) \cup \{\mathbf{z}_{\ell+1} - F(\mathbf{z}_{1..\ell})\}. \quad (124)$$

Lemma 36. *Let \mathcal{C}^δ be a real LN-cell and $\{f_j : \mathcal{C}_j^\delta \rightarrow \mathcal{C}^\delta\}$ a real cover compatible with $B(\mathcal{C})$. Then every image $f_j(\mathbb{R}_+\mathcal{C}_j^\delta)$ is either contained in or disjoint from $\mathbb{R}_+\mathcal{C}$.*

Similarly if $F : \mathcal{C}^\delta \rightarrow \mathbb{C}$ is a real LN-function and $\{f_j : \mathcal{C}_j^\delta \rightarrow \mathcal{C}_F\}$ a real cover compatible with $B(F|_e)$ then every image $f_j(\mathbb{R}_+\mathcal{C}_j^\delta)$ is either contained or disjoint from $\text{Gr } \mathbb{R}_+F|_e$.

Proof. The first statement follows since $\mathbb{R}_+\mathcal{C}$ is given by some sign conditions on the boundary equations $B(\mathcal{C})$ and these signs remain constant on each $f_j(\mathbb{R}_+\mathcal{C}_j^\delta)$ by compatibility. The second statement follows similarly. \square

6.2. Effective model completeness. We say that a formula $\psi(\mathbf{x})$ is *existential* if it is of the form $\exists \mathbf{y} : \psi_0(\mathbf{x}, \mathbf{y})$ where ψ_0 is quantifier-free. Our goal in this section is to prove the effective model completeness part of Theorem 2. More explicitly we prove the following.

Claim 2. *The structure \mathbb{R}_{LN} is effectively model complete. That is, every \mathcal{L}_{LN} -formula ψ is equivalent to an existential formula ψ_e and $\mathcal{F}(\psi_e) < \text{Eff}(\psi)$.*

We make several reductions. First, we can assume that all relations corresponding to real LN-functions $F : \mathcal{C}^\delta \rightarrow \mathbb{C}$ appearing in ψ have $\text{Gr } \mathbb{R}_+F \subset I^n$ where $I := [-1, 1]$ (with different n s). Indeed one can always choose some $M = \text{Eff}(F)$ so that $\tilde{F} := F(M\mathbf{z})/M$ satisfies $\text{Gr } \mathbb{R}_+\tilde{F} \subset I^n$. One can then replace each relation $\mathbf{x} \in \text{Gr } \mathbb{R}_+F$ in ψ by

$$\exists \mathbf{y} : (M\mathbf{y} = \mathbf{x}) \wedge \mathbf{y} \in \text{Gr } \mathbb{R}_+\tilde{F}. \quad (125)$$

Repeating this for each LN-graph in ψ finishes the reduction.

Next, it is enough to prove the case when $\psi(\mathbb{R}^n) \subset I^n$. Let us demonstrate this reduction for the $n = 1$ case. For general ψ we can write

$$\begin{aligned} \psi(x) &\iff (-1 \leq x \leq 1 \wedge \psi(x)) \vee ((x < -1 \vee x > 1) \wedge \psi(x)) \\ &= \psi_1(x) \vee \psi_2(x) \end{aligned} \quad (126)$$

Then $\psi_1(x)$ defines a subset of I . For ψ_2 write the formula for the ‘‘inverse’’ of $\psi_2(\mathbb{R}^n)$, namely

$$\psi_2^i(y) := (-1 \leq y \leq 1) \wedge \exists x : xy = 1 \wedge \psi_2(x). \quad (127)$$

Then ψ_2^i defines a subset of I , so by our assumption is equivalent to an existential formula $\psi_{2e}^i(y)$. Then finally

$$\psi_2(x) \iff \exists y : (xy = 1) \wedge \psi_{2e}^i(y). \quad (128)$$

The case of general n is the same, except one should consider 2^n cases for the potential inversions of each coordinate.

Next, it is well known that by a simple induction over the quantification depth, it is enough to prove the claim for the negation of an existential formula ψ . We can also reduce to the case where all quantifiers in ψ are of the form $\exists y \in I$. Indeed, suppose $\psi(\mathbf{x})$ is of the form $\exists y : \psi_1(\mathbf{x}, y)$. Then it is equivalent to

$$(\exists y \in I : \psi_1(\mathbf{x}, y)) \vee (\exists y' \in I : \psi'(\mathbf{x}, y')) \quad (129)$$

where we will define $\psi'(\mathbf{x}, y')$ satisfying

$$\forall y' \in I : (\psi'(\mathbf{x}, y') \iff \psi_1(\mathbf{x}, 1/y')). \quad (130)$$

We do this as follows. First, if ψ contains a predicate $\mathbf{z} \in \text{Gr } \mathbb{R}_+ F$ and y is one of the components of \mathbf{z} then we just replace this by $0 = 1$ in ψ' , because by our assumption all graphs are contained in I^n . Next, if y appears in an equality $y = T$ where T is a variable or constant we replace it by $y'T = 1$. Similarly we replace $y > T$ by

$$(y' > 0 \wedge y'T < 1) \vee (y' < 0 \wedge y'T > 1). \quad (131)$$

Finally, it remains to treat the case where y appears in the relations $+$, $*$. Since these are semialgebraic we can also express them using polynomial inequalities in $y' = 1/y$, for instance

$$\begin{aligned} x_1 + x_2 = y &\iff y'(x_1 + x_2) = 1 & x_1 + y = x_2 &\iff y'(x_2 - x_1) = 1 \\ x_1 x_2 = y &\iff y' x_1 x_2 = 1 & x_1 y = x_2 &\iff x_1 = x_2 y' \end{aligned} \quad (132)$$

Repeating this for each quantifier in ψ finishes the reduction.

Finally after these reductions, write $\psi(\mathbf{x}, \mathbf{y}) := \exists y \in I^m \psi'(\mathbf{x}, \mathbf{y})$. Consider a cell

$$\hat{\mathcal{C}} = D(2)^\ell \times \prod_{F \in \psi'} \mathcal{C}_F \quad (133)$$

where the product ranges over all function graphs appearing as relation symbols in ψ' , once for each appearance. Denote the variables of \mathcal{C}_F by $\mathbf{z}_{1.. \ell(F)}^F$. Apply the CPT to the following set of equations. First we include all boundary equations $B(F)$ for $F \in \psi'$, and for each relation symbol $\mathbf{x} \in \text{Gr } \mathbb{R}_+ F$ we include $\mathbf{x}_i = \mathbf{z}_i^{\mathcal{C}}$ for $i = 1, \dots, \ell(F)$; for each $+$ relation $x + y = z$ we include $x + y - z$ and similarly with $*$; and for each $x = y$ or $x < y$ we include $x - y$. Note for that all of these relations, even if one of x, y, z is a large constant rather than a variable this does not affect the format of our equations (as LN-functions) because we may just divide by some large constant to make sure all coefficients are bounded by 1. Thus the set of all equations we use have format $\text{Eff}(\psi)$.

Let $\{f_j : \mathcal{C}_j^\delta \rightarrow \hat{\mathcal{C}}^\delta\}$ be the real cover obtained from the CPT. Consider only the maps f_j for which the relations $\mathbf{x}_i = \mathbf{z}_i^{\mathcal{C}}$ vanish identically – denote these $j \in \Sigma$. Then for each of the relations $R(\mathbf{x})$ in ψ' , it is either uniformly true on $f_j(\mathcal{C}_j^\delta)$ or uniformly false when $j \in \Sigma$. Indeed, for the LN-relations this is exactly Lemma 36 and for the remaining relations it true for the same reason.

Since the $f_j(\mathbb{R}_+ \mathcal{C}_j)$ cover $\mathbb{R}_+ \hat{\mathcal{C}}$, we conclude that the set $\psi'(I^{n+m})$ is given by a union of $f_j(\mathbb{R}_+ \mathcal{C}_j)$ for j in a subset $\Sigma' \subset \Sigma$ given by those $j \in \Sigma$ over which ψ' holds true. The cellular structure of f_j then implies that $\psi(I^n)$, given by the projection of $\psi'(I^{n+m})$ to I^n , is given by the images $(f_j)_{1.. \ell}((\mathcal{C}_j)_{1.. \ell})$ with $j \in \Sigma'$. Claim 2 now follows from the following lemma, showing that the complement of each of these sets (and therefore also the complement of their union) is given by an existential formula of effectively bounded format.

Lemma 37. *Let \mathcal{C}^δ be a real LN-cell and $f : \mathcal{C}^\delta \rightarrow D(2)^\ell$ a simple cellular map as described in §5.4. Then $f(\mathbb{R}_+ \mathcal{C})$ and its complement are given by an existential \mathcal{L}_{LN} -formula of format $\text{Eff}(f)$.*

Proof. By assumption f is a composition of affine and covering maps. If we write $\mathbf{w}_{1..l} = f(\mathbf{z}_{1..l})$ then we have equations

$$\begin{aligned} \mathbf{w}_1 &= P_1(\mathbf{z}_1), & P_1 &\in \mathbb{R}[\mathbf{z}_1] \\ \mathbf{w}_2 &= P_2(\mathbf{z}_2), & P_2 &\in \mathcal{O}_{\text{LN}}(\mathcal{C}_1^\delta)[\mathbf{z}_2] \\ & & & \vdots \\ \mathbf{w}_\ell &= P_\ell(\mathbf{z}_\ell), & P_\ell &\in \mathcal{O}_{\text{LN}}(\mathcal{C}_{1..l-1}^\delta)[\mathbf{z}_\ell] \end{aligned} \quad (134)$$

where for each fixed $\mathbf{z}_{1..j-1}$ the polynomials $P_j(\mathbf{z}_{1..j-1}, \cdot)$ are injective on \mathbb{R} for fibers of type D and on \mathbb{R}_+ for fibers of type D_\circ, A . Write $\mathcal{C} = \mathcal{F}_1 \odot \cdots \odot \mathcal{F}_\ell$ and set

$$\mathbb{R}^\mathcal{C} := \prod_{j=1}^{\ell} R_j, \quad R_j := \begin{cases} \{0\} & \mathcal{F}_j = * \\ \mathbb{R} & \mathcal{F}_j = D \\ \mathbb{R}_+ & \mathcal{F}_j = D_\circ, A \end{cases} \quad (135)$$

Then we can express $f(\mathbb{R}_+\mathcal{C})$ with the formula

$$\begin{aligned} \psi(\mathbf{w}) &:= \exists \mathbf{z} \in \mathbb{R}^\mathcal{C} : (134) \text{ holds} \wedge \\ &\quad \mathbf{z}_1 \in \mathcal{F}_1 \wedge \mathbf{z}_2 \in \mathcal{F}_2(\mathbf{z}_1) \wedge \cdots \wedge \mathbf{z}_\ell \in \mathcal{F}_\ell(\mathbf{z}_{1..l-1}) \end{aligned} \quad (136)$$

where the conditions $\mathbf{z}_j \in \mathcal{F}_j(\mathbf{z}_{1..j-1})$ are clearly expressible in \mathcal{L}_{LN} , for instance if $\mathcal{F}_j = A(r_1, r_2)$ then we write this as

$$r_1(\mathbf{z}_{1..j-1}) < \mathbf{z}_j < r_2(\mathbf{z}_{1..j-1}) \quad (137)$$

where r_1, r_2 are by definition real LN-functions on $\mathcal{C}_{1..j-1}^\delta$.

Now consider the complement of $f(\mathbb{R}_+\mathcal{C})$ in \mathbb{R}^ℓ . This can be expressed as follows. We first show that

$$\neg \exists \mathbf{z} \in \mathbb{R}^\mathcal{C} : (134) \text{ holds} \quad (138)$$

is existential. This is equivalent to the disjunction of the following:

$$\begin{aligned} &\neg \exists \mathbf{z}_1 : \mathbf{w}_1 = P_1(\mathbf{z}_1) \\ &\exists \mathbf{z}_1 : \mathbf{w}_1 = P_1(\mathbf{z}_1), \neg \exists \mathbf{z}_2 : \mathbf{w}_2 = P_2(\mathbf{z}_1, \mathbf{z}_2) \\ &\quad \vdots \\ &\exists \mathbf{z}_{1..l-1} : \mathbf{w}_{1..l-1} = P_{1..l-1}(\mathbf{z}_{1..l-1}), \neg \exists \mathbf{z}_\ell : \mathbf{w}_\ell = P_\ell(\mathbf{z}_{1..l-1}, \mathbf{z}_\ell). \end{aligned} \quad (139)$$

where above each $\exists \mathbf{z}_j$ is shorthand for $\exists \mathbf{z}_j \in \mathbb{R}_j^\mathcal{C}$. We claim that the negated formulas

$$\neg \exists \mathbf{z}_j \in \mathbb{R}_j^\mathcal{C} : \mathbf{w}_j = P_j(\mathbf{z}_{1..j-1}, \mathbf{z}_j) \quad (140)$$

can be rewritten as existential formulas. Indeed, for $\mathcal{F}_j = *$ this just means $\mathbf{w}_j \neq P_j(\mathbf{z}_{1..j-1}, 0)$. For $\mathcal{F}_j = D(r)$ this condition is empty because $P_j(\mathbf{z}_{1..j-1}, \cdot)$ is onto \mathbb{R} , being an affine translate. And for $\mathcal{F}_j = D_\circ, A$ this condition is equivalent to $\mathbf{w}_j < P(\mathbf{z}_{1..j-1}, 0)$ because $P_j(\mathbf{z}_{1..j-1}, \mathbf{z}_j)$ is monotone and tends to infinity as $\mathbf{z}_j \rightarrow \infty$.

Having established that (138) is existential, we can finish writing the negation of $\psi(\mathbf{x})$ as

$$\begin{aligned} \psi_n(\mathbf{w}) &:= (\neg \exists \mathbf{z} \in \mathbb{R}^\mathcal{C} : (134) \text{ holds}) \vee \exists \mathbf{z} \in \mathbb{R}^\mathcal{C} : (134) \text{ holds} \wedge \\ &\quad (\mathbf{z}_1 \notin \mathcal{F}_1 \vee \mathbf{z}_2 \notin \mathcal{F}_2(\mathbf{z}_1) \vee \cdots \vee \mathbf{z}_\ell \notin \mathcal{F}_\ell(\mathbf{z}_{1..l-1})) \end{aligned} \quad (141)$$

where this is now an existential formula. \square

6.3. Effective o-minimality. We now finish the proof of Theorem 2 by showing the effective o-minimality of \mathbb{R}_{LN} . More specifically we prove an $\text{Eff}(\psi)$ bound for the number of connected components for a set $A \subset \mathbb{R}^n$ defined by a formula ψ . By effective model completeness we may assume that ψ is existential. Making the same reductions as in §6.2 we may reduce to bounding the case

$$\psi(\mathbf{x}) := \exists \mathbf{y} \in I^m : \psi'(\mathbf{x}, \mathbf{y}) \quad (142)$$

where ψ' is quantifier free. This is certainly bounded by the number of connected components of $\psi'(I^{n+m})$. But we have already seen at the end of §6.2 that this set is given by the union of images $f_j(\mathbb{R}_+ \mathcal{C}_j)$ where $\#\{f_j\} < \text{Eff}(\psi)$. Since each image is connected this gives the required bound.

6.4. Effective polynomial boundedness. Another key feature of \mathbb{R}_{an} that is often used in application is polynomial boundedness. Say that an effectively o-minimal structure is *effectively polynomially bounded* if for every definable map $f : \mathbb{R} \rightarrow \mathbb{R}$ there exists $N = \text{Eff}(f)$ such that $f(t) < t^N$ for all $t \gg 1$.

Proposition 38. \mathbb{R}_{LN} is effectively polynomially bounded.

Proof. It is equivalent to prove that for every definable $f : (0, 1) \rightarrow (0, 1)$ we have $f(t) > t^N$ with $N = \text{Eff}(f)$ and all sufficiently small t . Consider the graph G of f . As in §6.2 we can cover G by images $f_j(\mathbb{R}_+ \mathcal{C}_j)$. In particular, the projection to x of one of these images must contain an interval $(0, \varepsilon)$ for some $\varepsilon > 0$. Denote this cell \mathcal{C} and the corresponding map f . Since $f(\mathcal{C}^\delta) \subset G$ one easily sees that the base of \mathcal{C} must be a punctured disc $D_\circ(r)$, and the map f therefore takes the form

$$f(\mathbf{z}_1, *) = (\mathbf{z}_1^k, f(\mathbf{z}_1)) \quad (143)$$

where $f \in \mathcal{O}_{\text{LN}}(D_\circ(r))$. Both k and the order of zero of f at $z_1 = 0$ are bounded by $\text{Eff}(f)$ for instance by effective o-minimality (or directly by bounds on the variation of argument), and the claim follows. \square

As a consequence of polynomial boundedness we can deduce an effective Łojasiewicz inequality. Below definability can be taken with respect to any effectively o-minimal structure which is effectively polynomially bounded.

Theorem 13 (Effective Łojasiewicz inequality). *Let $X \subset \mathbb{R}^n$ be closed and bounded and $f, g : X \rightarrow \mathbb{R}$ two definable continuous functions with $f^{-1}(0) \subset g^{-1}(0)$. Then there exists $N = \text{Eff}(f, g)$ and $C > 0$ such that $|g(x)|^N < C|f(x)|$ for all $x \in X$.*

Proof. Set

$$\lambda(\varepsilon) := \min\{|f(x)| : x \in X \text{ and } |g(x)| = \varepsilon\}. \quad (144)$$

Note that the set on the left hand side is compact and does not contain 0 by assumption, so $\lambda(\varepsilon) > 0$ for all $\varepsilon > 0$. The claim now easily follows from polynomial boundedness, i.e. $\lambda(\varepsilon) \gg \varepsilon^N$. \square

7. EFFECTIVE O-MINIMALITY OF $\mathbb{R}_{\text{LN,PF}}$

In this section we prove Theorem 3: that the extension of \mathbb{R}_{LN} by (unrestricted) Pfaffian functions is effectively o-minimal. It has long been known [48, 43] that adding Pfaffian functions (or even general Rolle leaves) to an o-minimal structure preserves o-minimality. It has also been shown in [2] that the approach of [48] to

this problem can be made effective, and we explain how this can be used to deduce the effective o-minimality of $\mathbb{R}_{\text{LN,PF}}$ in §7.3. In another direction, Gabrielov and Vorobjov have developed in a series of papers a more concrete approach to this effectivity for Pfaffian functions over the algebraic structure. We show in §7.4 how their approach can be carried over to include Pfaffian functions over \mathbb{R}_{LN} . While this requires somewhat more work than the approach of [2], it seems more likely to play a role in an approach toward Conjecture 4.

7.1. Extension by Pfaffian functions. Let (\mathcal{S}, Ω) be an effective o-minimal structure. We will write $\mathcal{F}^{\mathcal{S}}(\cdot)$ for the format of definable sets in \mathcal{S} to avoid confusion (as we will be defining another type of format).

Let $G \subset \mathbb{R}^n$ be an open cell in \mathcal{S} ,

$$G = I_1 \odot \cdots \odot I_n, \quad \text{where } I_k := (a_k(\mathbf{x}_{1..k-1}), b_k(\mathbf{x}_{1..k-1})) \quad (145)$$

such that the walls a_k, b_k are real analytic on $G_{1..k-1}$ and satisfy $a_k < b_k$ everywhere. We also allow $a_k = \infty$ and $b_k = \infty$.

Let $f_1, \dots, f_\ell : G \rightarrow \mathbb{R}$ be an ℓ -tuple of real analytic functions. Write $X \subset \mathbb{R}^\ell$ for the image of G under (f_1, \dots, f_ℓ) . Say that these functions form a *restricted Pfaffian chain* over \mathcal{S} if they satisfy a system of differential equations

$$\frac{\partial f_j}{\partial x_k} = P_{jk}(f_1, \dots, f_j) \quad (146)$$

where the functions P_{jk} are real analytic on X . Define the format of the Pfaffian chain to be

$$\mathcal{F}(f_1, \dots, f_\ell) := \ell + \mathcal{F}^{\mathcal{S}}(G) + \sum_{j,k} \mathcal{F}^{\mathcal{S}}(P_{jk}|_U). \quad (147)$$

A *Pfaffian function* is a function of the form $f := P(f_1, \dots, f_\ell)$ where P is real analytic on X . Define the format of f to be

$$\mathcal{F}(f) := \mathcal{F}(f_1, \dots, f_\ell) + \mathcal{F}^{\mathcal{S}}(P). \quad (148)$$

Denote by $(\mathcal{S}_{\text{PF}}, \Omega_{\text{PF}})$ the structure generated by all restricted Pfaffian functions over \mathcal{S} with the filtration generated by the formats defined above. Denote by $\mathcal{L}_{\mathcal{S}, \text{PF}}$ the language containing constants for all real numbers and a function symbol for each Pfaffian function over \mathcal{S} .

7.2. Khovanskii's bound over \mathcal{S} . Let (\mathcal{S}, Ω) be an effective o-minimal structure.

Theorem 14 (Khovanskii's bound). *Let F_1, \dots, F_k be Pfaffian over \mathcal{S} , all defined on a common domain G . Then the number of connected components of the set*

$$\{\mathbf{x} \in G : F_1(\mathbf{x}) = \cdots = F_k(\mathbf{x}) = 0\} \quad (149)$$

is bounded by $\text{Eff}(F_1, \dots, F_k)$.

Proof. This is the main result of [32]. The main case is when $k = n$ and one counts isolated solutions of the system of equations. In this case Khovanskii proves the theorem by defining the $\tilde{*}$ -sequence of F_1, \dots, F_n , which turns out to be a sequence of polynomials P_1, \dots, P_n of bounded degrees. In our more general case these will now be functions definable in \mathcal{S} with formats bounded in terms of the format of F_1, \dots, F_k .

Khovanskii shows that the number of isolated solutions of (149) is bounded by the “virtual number of zeros” of the $\tilde{*}$ -sequence. This is bounded [32, Section 3.10, Corollary 3] by the supremum for the number of isolated points in any

fiber of the map $(P_1, \dots, P_n) : G \rightarrow \mathbb{R}^n$ taken over all fibers. When P_1, \dots, P_n this supremum is bounded by Bezout; when P_1, \dots, P_n are definable in \mathcal{S} a bound of the form $\text{Eff}(F_1, \dots, F_n)$ follows trivially by effective o-minimality. \square

Corollary 39. *Let $\psi(\mathbf{x})$ be a quantifier-free $\mathcal{L}_{\mathcal{S}, \text{PF}}$ -formula. Then the number of connected components of $\psi(\mathbb{R}^n)$ is bounded by $\text{Eff}(\psi)$.*

Proof. We can write ψ as a disjunction of basic sets of the form

$$\{\mathbf{x} : P_1(\mathbf{x}) = \dots = P_k(x) = 0, Q_1(\mathbf{x}) > 0, \dots, Q_j(\mathbf{x}) > 0\}. \quad (150)$$

Each connected component of this corresponds to at least one connected component of the following set by projection:

$$\{(\mathbf{x}, \mathbf{y}, \mathbf{z}) : P_1(\mathbf{x}) = \dots = P_k(x) = 0, Q_1(\mathbf{x}) = \mathbf{y}_1^2, \dots, Q_j(\mathbf{x}) = \mathbf{y}_j^2, \mathbf{y}_1 \mathbf{z}_1 = 1, \dots, \mathbf{y}_j \mathbf{z}_j = 1\}. \quad (151)$$

Since this latter set involves only equalities the bound on the number of connected components follows from Theorem 14. \square

7.3. Effective o-minimality following Wilkie and Berarducci-Servi. The structure $\mathbb{R}_{\text{LN}, \text{PF}}$ is generated by Pfaffian functions over \mathbb{R}_{LN} as defined in §7.1. Indeed, every complex cell \mathcal{C} is a domain of the form G (up to a minor technicality with $*$ -fibers which is of no significance), and every LN function $f \in \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta)$ is Pfaffian by definition, taking the Pfaffian chain to consist of just the coordinate variables and P to be f .

The effective o-minimality of $\mathbb{R}_{\text{LN}, \text{PF}}$ now essentially follows from the main result of [2] modulo some small remarks. Indeed, [2, Theorem 2.2] shows that given an effective bound for the number of connected components of quantifier-free formulas, one can obtain effective bounds for the number of connected components for arbitrary formulas.

A few remarks are in order. First, [2, Theorem 2.2] is formulated for recursive bounds rather than primitive-recursive as we have insisted on, but by the remark following the theorem the primitive-recursive analog holds with the same proof. Second, unlike us [2] allows an extension by only finitely many Pfaffian functions (and no constants), but this makes no difference as one can always consider the restriction of $\mathcal{L}_{\text{LN}, \text{PF}}$ that includes the relevant functions or constants used in a given formula. Finally, the expansion considered in [2] is by functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$ rather than our more general cellular domains G . This can easily be circumvented as follows. Consider some map $\phi_{a,b}(x) : \mathbb{R} \rightarrow (a, b)$, algebraic and real analytic in a, b, x for all $a < b$, which restricts to a bijection for every fixed a, b . Then, for any Pfaffian function $f = (f_1, \dots, f_n) : G \rightarrow \mathbb{R}$ we can consider a pullback

$$\tilde{f} : \mathbb{R}^n \rightarrow \mathbb{R}, \quad \tilde{f} = (\phi_{a_1, b_1}^* f_1, \dots, \phi_{a_n(\mathbf{x}_{1..n-1}), b_n(\mathbf{x}_{1..n-1})}^* f_n). \quad (152)$$

Theorem 14 applies to formulas defined with these pullbacks as well – in fact they are also easily seen to be Pfaffian over \mathcal{S} . Then we can consider the structure defined by these \tilde{f} functions, and it is easy to see that the structure generated by these functions contains the graphs of the original f .

7.4. Effective o-minimality following Gabrielov-Vorobjov. In this section we indicate how the approach of Gabrielov-Vorobjov can also be generalized to work over \mathbb{R}_{LN} . The bounds obtained by Gabrielov-Vorobjov are generally more explicit, and often polynomial with respect to degrees, so this approach may offer a more plausible line of attack toward Conjecture 4.

7.4.1. Restricted Pfaffian functions. We start by introducing *restricted* Pfaffian functions. We assume that the domain G in (145) is precompact in \mathbb{R}^n , and also that the cell walls are real analytic on $\bar{G}_{1..k-1}$.

Now let $f_1, \dots, f_\ell : \bar{G} \rightarrow \mathbb{R}$ be an ℓ -tuple of real analytic functions. Write $X \subset \mathbb{R}^\ell$ for the image of \bar{G} under (f_1, \dots, f_ℓ) . Say that these functions form a *restricted* Pfaffian chain over \mathcal{S} if they satisfy a system of differential equations (146) where the functions P_{jk} are real analytic on X , and if there exists some domain $U \subset \mathbb{C}^n$ containing X such that P_{jk} extends holomorphically to U and this extension is definable in \mathcal{S} . Define the format of the Pfaffian chain to be

$$\mathcal{F}(f_1, \dots, f_\ell) := \ell + \mathcal{F}^{\mathcal{S}}(G) + \sum_{j,k} \mathcal{F}^{\mathcal{S}}(P_{jk}|_U). \quad (153)$$

A *restricted Pfaffian function* is a function of the form $f := P(f_1, \dots, f_\ell)$ where P again P extends holomorphically to some complex domain U containing X . Define the format of f to be

$$\mathcal{F}(f) := \mathcal{F}(f_1, \dots, f_\ell) + \mathcal{F}^{\mathcal{S}}(P|_U). \quad (154)$$

Denote by $(\mathcal{S}_{\text{rPF}}, \Omega_{\text{rPF}})$ the structure generated by all restricted Pfaffian functions over \mathcal{S} with the filtration generated by the formats defined above.

Remark 40. *An LN-cell of the form $\mathcal{C} := D_o(1) \odot A(x, 2)$ admits an LN function x/y which does not extend real-analytically to $\bar{\mathcal{C}} \subset \mathbb{R}^4$ as it is not analytic in a neighborhood of the origin. This function would not be restricted Pfaffian by our definition. We will overcome this difficulty with another construction at the end to obtain an honest expansion of \mathbb{R}_{LN} .*

7.4.2. Effective o-minimality of \mathcal{S}_{rPF} . Our goal in this section is to prove the following theorem.

Theorem 15. *The structure $(\mathcal{S}_{\text{rPF}}, \Omega_{\text{rPF}})$ is effectively o-minimal.*

The proof consists in a routine generalization of various results by Gabrielov and Vorobjov from the case where the coefficients P_{ij} in (146) are polynomials to the general case considered above. We outline the main results.

The key ingredient in Gabrielov-Vorobjov's effective theory of Pfaffian structures, in addition to Theorem 14, is the following local complex analog of due to Gabrielov.

Definition 41. *A deformation of format \mathcal{F} of a restricted Pfaffian function F_0 is a function germ $F(\mathbf{z}, \varepsilon) : (\mathbb{C}^{n+1}, 0) \rightarrow \mathbb{C}$ analytic at the origin such that $F(\mathbf{z}, 0) = F_0$ and for every sufficiently small ε we have $F(\cdot, \varepsilon) \in L$ where L is a linear space of functions satisfying $L \subset \Omega_{\mathcal{F}}^{\mathcal{S}}$.*

With this definition we can state Gabrielov's result.

Theorem 16 (Gabrielov's bound). *Let $F_1, \dots, F_n : (\mathbb{C}^{n+1}, 0) \rightarrow \mathbb{C}$ be restricted Pfaffian deformations of format at most \mathcal{F} . Then there exists $r > 0$ such that for all sufficiently small ε the number of isolated points in*

$$\{\mathbf{z} \in \mathbb{C}^n : \|\mathbf{z}\| < r \text{ and } F_1(\mathbf{z}, \varepsilon) = \dots = F_n(\mathbf{z}, \varepsilon) = 0\} \quad (155)$$

is bounded by $\text{Eff}(\mathcal{F})$.

Proof. This is quite similar to the proof given above for Theorem 14. Indeed, Gabrielov [19, Theorem 2.1] defines a sequence much like Khovanskii's \star -sequence, which in the classical case is a polynomial sequence and in our case is a sequence of definable functions of format $\text{Eff}(\mathcal{F})$. The bound is then given by the number of isolated solutions of this sequence – which is bounded by Bezout in Gabrielov's case, and by effective o-minimality in our cases.

We remark that the main difference compared to Khovanskii's result is that Gabrielov needs to involve a derivative in the ε -direction in the definition of the \star -sequence. To accommodate this we had to include the assumption that the deformations $F(\cdot, \varepsilon)$ all belong to a fixed linear space independent of ε , so that the derivative remains in this space and has the same bound on the format. In Gabrielov's case this is automatic (because polynomials degree at most \mathcal{F} form a linear space). \square

We call the sets generated by quantifier free formulas in the language of restricted Pfaffian functions over \mathcal{S} the “semi-Pfaffian sets” over \mathcal{S} . We endow this collection of sets with a filtration $\Omega_{\text{rPF}}^{\text{qf}}$, where we associate format to sets as described at the end of §1.5 excluding the projection axiom.

Theorem 17 ([20]). *Let X be a semi-Pfaffian set over \mathcal{S} with quantifier free format \mathcal{F} . Then \bar{X} is semi-Pfaffian over \mathcal{S} of quantifier free format $\text{Eff}(\mathcal{F})$.*

Proof. The proof is the same as in the original, using Theorem 16 to replace the classical version over the algebraic structure. \square

A *stratification* of a semi-Pfaffian set $X \subset \mathbb{R}^n$ is a partition of X as a disjoint union of smooth, not necessarily connected, semi-Pfaffian subsets. The following is a generalization of a theorem by Gabrielov and Vorobjov.

Theorem 18 ([21]). *Let X be a semi-Pfaffian set over \mathcal{S} with quantifier free format \mathcal{F} . Then there exists a stratification $X = \cup_j X_i$ where the quantifier free formats of the X_i strata, and their number, is bounded by $\text{Eff}(\mathcal{F})$.*

Proof. The proof the same is in the original, using Theorems 14 and 16 to replace the classical versions. \square

Finally we can state the theorem of the complement for \mathcal{S}_{rPF} .

Theorem 19 ([22]). *Let X be semi-Pfaffian over \mathcal{S} with quantifier-free format \mathcal{F} . Then $\mathbb{R}^n \setminus \pi_n(X)$ is the projection $\pi_n(Y)$ of a semi-Pfaffian set $Y \subset \mathbb{R}^N$ and the quantifier-free format of Y is bounded by $\text{Eff}(\mathcal{F})$.*

Proof. This theorem is proved by showing that cell-decomposition is possible in the class of sub-Pfaffian sets (i.e., projections of semi-Pfaffian sets). The proof is based on Theorems 14, 17 and 18. See also [14] for another proof that isolates the dependence of the argument on these three results. \square

As a consequence of Theorem 19, we deduce that a every \mathcal{L}_{rPF} -formula ψ is equivalent to an existential formula $\exists y : \psi'(x, y)$ with $\mathcal{F}(\psi') < \text{Eff}(\psi)$. Since the number of connected components of a set defined by ψ' is bounded by Theorem 14, we obtain Theorem 15.

7.4.3. *The unrestricted case: limit sets.* In [26] Gabrielov develops an effective approach to unrestricted Pfaffian functions. We define the analog of this construction over \mathbb{R}_{LN} .

In the following definition we fix some domain G as in (145) and a Pfaffian chain over it, and consider the formats of restricted semi-Pfaffian sets with respect to this chain. If $X \subset \mathbb{R}^n \times \mathbb{R}$ and $\lambda \in \mathbb{R}$ we denote

$$X_\lambda := \{\mathbf{x} \in \mathbb{R}^n : (\mathbf{x}, \lambda) \in X\}. \quad (156)$$

Definition 42 (A-Limit sets). *Let $X \subset \mathbb{R}^n \times \mathbb{R}_{>0}$ be semi-Pfaffian over \mathcal{S} , and suppose for every $\lambda \in \mathbb{R}_{>0}$ the set X_λ is restricted semi-Pfaffian of format at most \mathcal{F} . Then $(\bar{X})_0$ is called an A-limit. The format of $(\bar{X})_0$ is defined to be \mathcal{F} .*

Denote by $A(\mathbb{R}_{\text{LN},\text{rPF}})$ the structure generated by the A-limit sets and by Ω_A the filtration generated by the formats above. The main result of [26] is that $A(\mathbb{R}_{\text{LN},\text{rPF}})$ is effectively o-minimal. This is done by showing that every set in this structure admits a representation as a union of certain simple sets (*relative closures*), and showing that the format of this representation remains effectively bounded under all boolean operations and projection, and that the format effectively bounds the number of connected components.

Remark 43. *In fact Gabrielov defines the notion “limit set” in a slightly different manner, but such that it is clear that they generate the same structure as the A-limit sets above. We introduced the non-standard terminology of A-limits to avoid potential confusion with his terminology.*

The proof in [26] goes through essentially unchanged for semi-Pfaffian sets over \mathbb{R}_{LN} . The only difference is in [26, Proposition 2.12] which establishes the exponential Lojasiewicz inequality for \mathbb{R}_{PF} . But since \mathbb{R}_{LN} is polynomially bounded (being a reduct of \mathbb{R}_{an}) the same inequality also holds in $\mathbb{R}_{\text{LN},\text{PF}}$ by [34, Theorem 3].

We claim that $A(\mathbb{R}_{\text{LN},\text{rPF}})$ is in fact $\mathbb{R}_{\text{LN},\text{PF}}$. First, we show that all LN-functions are A-definable. Indeed, if \mathcal{C}^δ is an LN-cell and $f \in \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta)$ then for every $0 < \varepsilon < 1$ we can define an LN-cell $\mathcal{C}(\varepsilon)$ by replacing every $D_o(r)$ factor by $A(\varepsilon r, r)$. Then the $\mathcal{C}(\varepsilon)$ are relatively compact in \mathcal{C}^δ and are thus definable in $\mathbb{R}_{\text{LN},\text{rPF}}$. Moreover, their format is independent of ε . Then \mathcal{C} is the A-limit of the family $\mathcal{C}(\varepsilon)$ and is thus contained in $A(\mathbb{R}_{\text{LN},\text{rPF}})$. A similar reasoning applies to the graph of $f|_{\mathcal{C}}$.

In the same way one can show that the graphs of unrestricted Pfaffian functions over \mathbb{R}_{LN} are A-limits. One slightly shrinks the domain G in (145), say by taking I_k to be the interval with the same center and length equal to $(1 - \varepsilon)$ times the length of I_k (or some similar construction if the intervals are infinite). Then the Pfaffian chain restricted to these precompact domains is a restricted Pfaffian chain with format independent of ε , and the A-limit set recovers the graph of the full unrestricted Pfaffian function.

In conclusion the A-limit sets contain all functions generating $\mathbb{R}_{\text{LN},\text{PF}}$ and thus in fact $\mathbb{R}_{\text{LN},\text{PF}} = A(\mathbb{R}_{\text{LN},\text{rPF}})$ and the effective o-minimality of this structure follows from Gabrielov’s result for A-limit sets.

8. GEOMETRIC CONSTRUCTIONS WITH LN CELLS

In this section we collect some useful geometric constructions with LN cells. Our main motivation, which we establish at the end of the section, is to show that complex LN cells and the LN functions on them are definable in \mathbb{R}_{LN} .

8.1. Monomialization of cells. Following [10] we define monomial LN cells. Let \mathcal{C} be an LN cell of length ℓ . An *admissible monomial* on \mathcal{C} is a function of the form $c \cdot \mathbf{z}^\alpha$ where $c \in \mathbb{C}$ and $\alpha = \alpha(f)$ is the associated monomial of some function $f \in \mathcal{O}_{\text{LN}}(\mathcal{C})$.

Definition 44 (Monomial cell). *An LN cell \mathcal{C} is monomial if $\mathcal{C} = *$ or if $\mathcal{C} = \mathcal{C}_{1..l} \odot \mathcal{F}$ where $\mathcal{C}_{1..l}$ is monomial and the radii involved in \mathcal{F} are admissible monomials on $\mathcal{C}_{1..l}$.*

We remark that it is always possible to choose all the constants in the admissible monomials to be real. Therefore, every monomial cell is a real cell.

The following proposition extends from [10] to the LN category with the same proof. Essentially, by the monomialization lemma every radius is given by a monomial times a unit. If we choose the extensions sufficiently large (using the refinement theorem) then the unit becomes nearly constant and one can just replace it by (say) its maximum on the cell.

Proposition 45. *Let \mathcal{C}^δ be a (real) cell. Then there exists a (real) cellular cover $\{f_j : \mathcal{C}_j^\delta \rightarrow \mathcal{C}^\delta\}$ where each \mathcal{C}_j is a monomial cell. Moreover*

$$\#\{f_j\}, \mathcal{F}(f_j) < \text{Eff}(\mathcal{C}^\delta, 1/(1 - \delta)). \quad (157)$$

8.2. Symmetrization.

Proposition 46. *Let \mathcal{C} be a real cell and $F \in \mathcal{O}_{\text{LN}}(\mathcal{C})$, not necessarily real. Then there exist real LN functions $F_R, F_I \in \mathcal{O}_{\text{LN}}(\mathcal{C})$ with $\mathcal{F}(F_R), \mathcal{F}(F_I) < \text{Eff}(F)$ and*

$$F_R|_{\mathbb{R}_+ \mathcal{C}} \equiv \text{Re } F|_{\mathbb{R}_+ \mathcal{C}}, \quad F_I|_{\mathbb{R}_+ \mathcal{C}} \equiv \text{Im } F|_{\mathbb{R}_+ \mathcal{C}}. \quad (158)$$

Proof. Let F_1, \dots, F_N be the LN chain for F . Since \mathcal{C} is real, its radii are all real and it follows that it is symmetric under the map $\mathbf{z} \rightarrow \bar{\mathbf{z}}$. If we define $F_1^\dagger, \dots, F_N^\dagger$ by

$$F_j^\dagger(\mathbf{z}) = \overline{F_j(\bar{\mathbf{z}})} \quad (159)$$

then $F_1^\dagger, \dots, F_N^\dagger$ are again bounded holomorphic functions on \mathcal{C} . By symmetry we have LN chain equations for these function where the polynomials G_{ij} in (17) are replaced by G_{ij}^\dagger (which just means taking complex conjugates of the coefficients).

If $F = G(F_1, \dots, F_N)$ then we put $F^\dagger := G^\dagger(F_1^\dagger, \dots, F_N^\dagger)$ and this is again an LN function. Then setting

$$F_R := \frac{F + F^\dagger}{2}, \quad F_I := \frac{F - F^\dagger}{2} \quad (160)$$

we obtain real LN functions satisfying the required conditions. \square

8.3. Real covers of complex cells.

Proposition 47. *Let \mathcal{C}^δ be an LN cell. Then there exist LN maps $\{f_j : \mathcal{C}_j^\delta \rightarrow \mathcal{C}^\delta\}$ with \mathcal{C}_j^δ real and $\mathcal{C} \subset \cup_j f_j(\mathbb{R}_+ \mathcal{C}_j)$. Moreover*

$$\#\{f_j\}, \mathcal{F}(f_j) < \text{Eff}(\mathcal{C}^\delta, 1/(1 - \delta)). \quad (161)$$

Proof. By refinement and Proposition 45 it is enough to prove the claim for monomial cells. Write $\mathcal{C} := \mathcal{F}_1 \odot \dots \odot \mathcal{F}_\ell$ and let

$$\hat{\mathcal{C}} := A(\frac{1}{2}, 2)^{\odot \ell} \odot \mathcal{C}. \quad (162)$$

Denote the coordinates on $\hat{\mathcal{C}}$ by (\mathbf{x}, \mathbf{z}) . We have an LN map $\phi : \hat{\mathcal{C}}^{2\delta} \rightarrow \mathcal{C}^\delta$ given by

$$\phi(\mathbf{x}, \mathbf{z}) = (\mathbf{x}_1 \mathbf{z}_1, \dots, \mathbf{x}_\ell \mathbf{z}_\ell) \quad (163)$$

where we use the fact that monomial cells are invariant under rotation in each of the coordinates. Then

$$\mathcal{C} \subset \phi(A(\frac{1}{2}, 2)^{\odot \ell} \odot \mathbb{R}_+ \mathcal{C}) \quad (164)$$

and it will therefore suffice to find maps $f_j : \mathcal{C}_j^\delta \rightarrow \hat{\mathcal{C}}^{2\delta}$ such that

$$A(\frac{1}{2}, 2)^{\odot \ell} \odot \mathbb{R}_+ \mathcal{C} \subset \cup_j f_j(\mathbb{R}_+ \mathcal{C}_j). \quad (165)$$

Reducing further, it will suffice to find $g_j : \mathcal{C}'_j \rightarrow A(1/2, 2)^{\odot \ell}$ with $A(1/2, 2)^{\odot \ell} \subset \cup_j g_j(\mathbb{R}_+ \mathcal{C}'_j)$ and then take $C_j := \mathcal{C}'_j \odot \mathcal{C}$ and $f_j := g_j \times \text{id}$. Finally finding g_j as above is elementary. We can first cover each $A(1/2, 2)$ by discs, and then use maps $D(1)^{\odot 2} \rightarrow D(1)^\delta$ given by $(z, w) \rightarrow (z + iw)$. \square

Finally we have the following.

Proposition 48. *Let \mathcal{C}^δ be an LN-cell and $f \in \mathcal{O}_{\text{LN}}(\mathcal{C}^\delta)$. Then $f|_{\mathcal{C}}$ is definable in \mathbb{R}_{LN} under the identification $\mathcal{C}^\ell \simeq \mathbb{R}^{2\ell}$ and its format is bounded by $\text{Eff}(f)$.*

Proof. Let $\{f_j : \mathcal{C}_j^\delta \rightarrow \mathcal{C}^\delta\}$ be as in Proposition 47. For each f_j apply Proposition 46 to the coordinates of f_j and to $f_j^* f$. Then on $\mathbb{R}_+ \mathcal{C}_j^{\sqrt{\delta}}$ the real and imaginary parts of all of these maps are given by LN functions of effectively bounded format. Thus the set

$$\cup_j \{(f_j(\mathbf{z}), f(f_j(\mathbf{z}))) : \mathbf{z} \in \mathcal{C}_j^{\sqrt{\delta}}\} \quad (166)$$

is definable in \mathbb{R}_{LN} . This is in fact the graph of f on some set S with $\mathcal{C} \subset S \subset \mathcal{C}^\delta$. In other words, we have proven the proposition (for all \mathcal{C}^δ, f) with $f|_{\mathcal{C}}$ in the conclusion replaced by $f|_S$.

We now deduce the original statement from this modified statement by induction on ℓ . Suppose the statement is true for $\ell-1$. In particular the cell $\mathcal{C}_{1.. \ell-1}$ is definable in \mathbb{R}_{LN} . By the modified statement, the radii of the fiber over $\mathcal{C}_{1.. \ell-1}$ are definable restricted to some set containing $\mathcal{C}_{1.. \ell-1}$, and since $\mathcal{C}_{1.. \ell-1}$ is definable as well this implies that \mathcal{C} is definable. By the modified statement f is definable on some set containing \mathcal{C} , so it is also definable restricted to \mathcal{C} . The bound on the format follows easily from this. \square

9. EFFECTIVE PILA-WILKIE IN $\mathbb{R}_{\text{LN}, \text{PF}}$

For $\mathbf{x} \in \bar{\mathbb{Q}}^n$ denote

$$H(\mathbf{x}) := \max_j H(\mathbf{x}_j) \quad (167)$$

where $H(\mathbf{x}_j)$ denotes the absolute multiplicative Weil height. For $X \subset \mathbb{R}^n$ and $g \in \mathbb{N}$ denote

$$X(g, H) := \{\mathbf{x} \in X \cap \bar{\mathbb{Q}}^n : [\mathbb{Q}(\mathbf{x}) : \mathbb{Q}] \leq g \text{ and } H(\mathbf{x}) \leq H\}. \quad (168)$$

We also denote by X^{alg} the union of all connected positive-dimensional \mathbb{R}^{alg} -definable subsets of X , and $X^{\text{trans}} := X \setminus X^{\text{alg}}$.

The Pila-Wilkie theorem [38] states that for X definable in an o-minimal structure and $\varepsilon > 0$, there exists a constant $C(X, \varepsilon)$ such that $\#X^{\text{trans}}(1, H) < C(X, \varepsilon)H^\varepsilon$ for every $H \in \mathbb{N}$. This has been extended by Pila [39] to a bound $\#X^{\text{trans}}(g, H) < C(X, g, \varepsilon)H^\varepsilon$. When X is definable in \mathbb{R}_{PF} , it was shown by Jones, Thomas, Schmidt and the author [5] that the constant $C(X, g, \varepsilon)$ can be bounded effectively

in terms of the format of X . The proofs in loc. cit. extend to general effectively o -minimal structures verbatim, although we note that much of the work there is related to obtaining polynomial bounds with respect to degrees in the restricted Pfaffian context. The proofs would be somewhat simpler if one only wants effective bounds. The effective analog of the Pila-Wilkie theorem is as follows.

Theorem 20. *Let $X \subset \mathbb{R}^n$ be definable in an effectively o -minimal structure. Then for every $\varepsilon > 0$ and every $g, H \in \mathbb{N}$ we have*

$$\#X^{\text{trans}}(g, H) \leq C(X, g, \varepsilon)H^\varepsilon, \quad \text{where } C(X, g, \varepsilon) = \text{Eff}(X, g, 1/\varepsilon). \quad (169)$$

We also give a “blocks” version following [39]. Say that a definable set $B \subset \mathbb{R}^n$ is a *basic block* if it is a smooth k -dimensional manifold contained in a semialgebraic set of dimension k .

Theorem 21. *Let $X \subset \mathbb{R}^m \times \mathbb{R}^n$ be definable in an effectively o -minimal structure. Then for every $\varepsilon > 0$ and $g \in \mathbb{N}$ there is a definable family $Y \subset \mathbb{R}^p \times \mathbb{R}^m \times \mathbb{R}^n$ of format $\text{Eff}(X, g, 1/\varepsilon)$ such that for every $(p, y) \in \mathbb{R}^p \times \mathbb{R}^m$ the fiber $Y_{p,y}$ is a basic block contained in X_y .*

Moreover, for every $y \in \mathbb{R}^m$ and $H \in \mathbb{N}$ there is a set $P_y \subset \mathbb{R}^p$ such that

$$\#P_y \leq C(X, g, \varepsilon)H^\varepsilon, \quad \text{where } C(X, g, \varepsilon) = \text{Eff}(X, g, 1/\varepsilon) \quad (170)$$

and

$$X_y(g, H) \subset \bigcup_{p \in P_y} Y_{p,y}. \quad (171)$$

An effective Pila-Wilkie theorem for semi-Noetherian sets was given in [3]. This result was restricted to sets defined by quantifier-free formulas using Noetherian functions in compact domains (in particular, not allowing unrestricted exponentiation). Another result establishing sharper, polylogarithmic bounds in the Noetherian category is given in [4]. However this result is again restricted to compact domains, and also has technical conditions related to the absence of unlikely intersections. It was observed already in [4, Section 1.5] that this limitation involving unlikely intersections is related to Khovanskii’s conjecture.

10. LN-FUNCTIONS AND REGULAR FLAT CONNECTIONS

In this section we show that holomorphic horizontal sections of regular meromorphic connections with LN coefficients are LN-functions, and that if the connection has quasiunipotent monodromy then all sections are definable in $\mathbb{R}_{\text{LN,exp}}$ after making an appropriate branch cut. As a consequence of Deligne’s Riemann-Hilbert correspondence we deduce that $\mathbb{R}_{\text{LN,exp}}$ contains every monodromic tuple of functions with moderate growth and locally quasiunipotent monodromy. We also show as a consequence of these constructions that period maps for PVHS, and the universal covering map for the universal abelian scheme $\mathbb{A}_g \rightarrow \mathcal{A}_g$, are definable in $\mathbb{R}_{\text{LN,exp}}$.

10.1. Connections with log-singularities on LN-cells. Let \mathcal{C}^δ be an LN-cell. Consider a connection on the trivial \mathbb{C}^ℓ -bundle on \mathcal{C}^δ given by

$$\nabla := d - \sum_{j=1}^{\ell} A_j \partial_j^*, \quad A_j \in \text{Mat}_{l \times l}(\mathcal{O}_{\text{LN}}(\mathcal{C}^\delta)). \quad (172)$$

where ∂_j^* is the one-form dual to ∂_j , i.e.

$$\partial_j^* := \begin{cases} d\mathbf{z}_j & \mathcal{F}_j = D \\ \frac{d\mathbf{z}_j}{\mathbf{z}_j} & \mathcal{F}_j = D_\circ, A \end{cases} \quad (173)$$

where \mathcal{F}_j denotes the j -th fiber of \mathcal{C} . We say that the format of a matrix with LN entries is the sum of the formats of the entries, and the format of ∇ is the sum of the formats of A_j for $j = 1, \dots, \ell$. In the case where \mathcal{C} is a product of discs and punctured discs this agrees with the usual notion of a connection with log-singularities.

10.1.1. *Holomorphic sections.* We will consider horizontal sections of ∇ . We start with the holomorphic case.

Proposition 49. *Suppose $X(\mathbf{z})$ is holomorphic and bounded in \mathcal{C}^δ and $\nabla X(\mathbf{z}) = 0$, where either $X(\mathbf{z}) \in \mathbb{C}^l$ or $X(\mathbf{z}) \in \text{End}(\mathbb{C}^n)$. Then X is LN and*

$$\mathcal{F}(X) = \text{Eff}(\nabla, \|X\|_{\mathcal{P}^{1/2}}). \quad (174)$$

Proof. This is essentially just rewriting the connection equation in terms of the standard vector fields,

$$\partial_j X = A_j \partial_j^* X = A_j X \quad (175)$$

and the entries of X indeed form an LN chain over the LN chain defining ∇ . \square

Corollary 50. *Suppose $X(\mathbf{z})$ is holomorphic and bounded in \mathcal{C}^δ and $\nabla X(\mathbf{z}) = 0$. Then X is definable in \mathbb{R}_{LN} and its format is bounded by $\text{Eff}(\nabla)$.*

Proof. Since $\nabla X = 0$ is linear in X we may as well rescale it and assume $\|X\|_{\mathcal{C}^\delta} \leq 1$. We then have a bound $\text{Eff}(\nabla)$ on the format of this rescaled X , and now scaling back gives the result. \square

10.1.2. *The quasiunipotent monodromy case.* We say that the *standard branch cut* on \mathcal{C} is the simply connected domain $\mathcal{C}_{\text{SC}} \subset \mathcal{C}$ given by removing the negative real line from every D_\circ and A fiber in \mathcal{C} .

Theorem 22. *Suppose ∇ has quasiunipotent monodromy. Let $X : \mathcal{C}_{\text{SC}}^\delta \rightarrow \text{End}(\mathbb{C}^l)$ be a horizontal section of ∇ , i.e. $\nabla X = 0$. Then X is definable in $\mathbb{R}_{\text{LN,exp}}$ and its format is bounded by $\text{Eff}(\nabla)$.*

Proof. Fix some base point $s \in \mathcal{C}$ and let $M_1, \dots, M_\ell \in \text{End}(\mathbb{C}^n)$ where M_j denotes the monodromy operator of ∇ along a simple loops around the divisor $\mathbf{z}_j = 0$ (trivial for D or $*$ fibers). Denote

$$t(\mathbf{z}) := \prod_{j: \mathcal{F}_j = D_\circ, A} \mathbf{z}_j. \quad (176)$$

Note that since $\pi_1(\mathcal{C}, s)$ is commutative the M_j s commute. Moreover since $\mathcal{F}(\nabla)$ bounds the norm of the matrices A_j , it is easy to see using Grönwall's inequality that $\|M_j\| < \text{Eff}(\nabla)$. Also by Grönwall's inequality, for every $\mathbf{z} \in \mathcal{C}_{\text{SC}}^\delta$ we have

$$\|X(\mathbf{z})\| \leq t(\mathbf{z})^{-N} \quad (177)$$

where $N = \text{Eff}(\nabla)$. Indeed, working for example with respect to \mathbf{z}_j and assuming it is of type D_\circ, A , we have an equation

$$\partial_j X = A_j X \quad (178)$$

where A_j is bounded on \mathcal{C}^δ , say by some $M = \text{Eff}(\nabla)$. In the logarithmic chart $w = \log \mathbf{z}_j$ this amounts to $\frac{\partial}{\partial w} X = A_j X$ and Grönwall then gives

$$\|X(w)\| < e^{M|w|} \|X(0)\| = |\mathbf{z}_j|^{-M} \|X(0)\|. \quad (179)$$

Repeating this for each coordinate proves the claim.

By [15, Lemma IV.4.5] we may choose commuting logarithms L_j such that

$$e^{2\pi i L_j} = M_j \quad (180)$$

and $\|L_j\| < \text{Eff}(\nabla)$. Now consider the matrix function

$$Y(\mathbf{z}) := X(\mathbf{z}) \cdot \mathbf{z}_1^{-L_1} \cdots \mathbf{z}_\ell^{-L_\ell}. \quad (181)$$

Since the monodromy of $\mathbf{z}_j^{-L_j}$ around $\mathbf{z}_j = 0$ is given by M_j^{-1} and this matrix commutes with every $\mathbf{z}_j^{-L_j}$, we see that $Y(\mathbf{z})$ is univalued on \mathcal{C} . Adding an appropriate integer multiple of the identity (depending on N above) to L_j we can also arrange the $Y(\mathbf{z})$ is bounded on $\mathcal{C}_{\text{SC}}^\delta$, and still $\|L_j\| < \text{Eff}(\nabla)$.

Now we think of $Y(\mathbf{z})$ as a flat holomorphic section of the linear connection

$$dY = d(X(\mathbf{z}) \cdot \mathbf{z}_1^{-L_1} \cdots \mathbf{z}_\ell^{-L_\ell}) = AY - Y\left(\sum_{j=1}^{\ell} L_j \frac{d\mathbf{z}_j}{\mathbf{z}_j}\right) \quad (182)$$

where we think of Y as a vector in \mathbb{C}^{l^2} . In this sense (182) is a connection equation with logarithmic singularities. By Corollary 50 we conclude that $Y(\mathbf{z})$ is definable in \mathbb{R}_{LN} with format bounded by $\text{Eff}(\nabla)$.

Finally the entries of $\mathbf{z}_j^{L_j}$ are each of the form $\mathbf{z}_j^\lambda P(\log \mathbf{z}_j)$ where λ ranges over the spectrum of L_j (modulo $2\pi i$) and P is a polynomial of degree at most l . Since we assume the monodromy is quasiunipotent these λ can be assumed real, and the functions above are all definable in \mathbb{R}_{exp} with format depending only on l . We recover X as

$$X(\mathbf{z}) = Y(\mathbf{z}) \cdot \mathbf{z}_1^{L_1} \cdots \mathbf{z}_\ell^{L_\ell}. \quad (183)$$

which finishes the proof. \square

10.2. Riemann-Hilbert correspondence and monodromic functions. Let M be a smooth quasi-projective variety, and suppose $\bar{M} \setminus M$ is a normal-crossings divisor. Let $U \subset M$ a semialgebraic simply-connected subset. Let \mathcal{V} be a local system on M , by which we always mean a local system of finite-dimensional \mathbb{C} -vector spaces.

Definition 51. *We say that \mathcal{V} has locally quasiunipotent monodromy if for any map $\phi : D_\circ(1) \rightarrow M$ the monodromy of $\phi^* \mathcal{V}$ is quasi-unipotent.*

By Kashiwara's theorem [29] a system \mathcal{V} is locally quasiunipotent if and only if it has quasiunipotent monodromy around every smooth point of the boundary $\bar{M} \setminus M$.

According to Deligne's Riemann-Hilbert correspondence [18], every local system \mathcal{V} on M arises as the horizontal system $\mathcal{V} \simeq V^\nabla$ of an algebraic vector bundle V with a flat connection ∇ with logarithmic singularities along the boundary. The results of the previous sections imply the following.

Proposition 52. *The following two statements hold:*

- (1) If U is precompact in M then the constant sections of $V^\nabla|_U$ are definable in \mathbb{R}_{LN} .
- (2) If \mathcal{V} has locally quasiunipotent monodromy then the constant sections of $V^\nabla|_U$ are definable in $\mathbb{R}_{\text{LN,exp}}$.

Proof. We can cover M by (finitely many) neighborhoods given by products of discs and punctured discs. By a simple covering argument it is enough to consider the intersection of U with each of these separately. Then the first claim is just a reformulation of Proposition 49 and the second a reformulation of Theorem 22. \square

We reformulate this as a result about the $\mathbb{R}_{\text{LN,exp}}$ -definability of multivalued holomorphic functions in terms of growth and monodromy.

Let $f = (f_1, \dots, f_n) : M \rightarrow \mathbb{C}^n$ be a holomorphic map (formally a map on the universal cover of M). We say that f is *monodromic* if fixed $s_0 \in M$ and any $\gamma \in \pi_1(M, s_0)$ we have

$$\Delta_\gamma(f_1, \dots, f_n) = (f_1, \dots, f_n)M_\gamma, \quad M_\gamma \in \text{GL}_n(\mathbb{C}) \quad (184)$$

where Δ_γ is the analytic continuation operator along γ . We say that f has *moderate growth* if whenever $u : D_\circ(1) \rightarrow M$ there are C, N such that

$$\|f \circ u(t)\| < Ct^{-N} \quad \forall t \in (D_\circ)_\text{SC}. \quad (185)$$

Theorem 23. *Let $f : M \rightarrow \mathbb{C}^n$ be a monodromic tuple with locally quasiunipotent monodromy and moderate growth. Then $f|_U$ is definable in $\mathbb{R}_{\text{LN,exp}}$.*

Proof. Let \mathcal{V} be the local system corresponding to the monodromy representation of f . Let (V, ∇) be the corresponding connection as in Proposition 52. It is enough to prove the claim with M replaced by an affine cover, so we may assume without loss of generality that V is trivial over M . In a trivializing chart a fundamental solution matrix $\nabla Y = 0$ satisfies

$$\Delta_\gamma Y = Y M_\gamma \quad (186)$$

so that $f \cdot Y^{-1}$ is univalued. Since it also has moderate growth, it is algebraic (and hence definable in \mathbb{R}_{alg}) by GAGA. Since Y is definable in $\mathbb{R}_{\text{LN,exp}}$ by Proposition 52 we conclude that f is definable in $\mathbb{R}_{\text{LN,exp}}$. \square

10.3. Period maps are definable in $\mathbb{R}_{\text{LN,exp}}$. Let S be a smooth quasi-projective variety over \mathbb{C} , $\mathcal{V}_\mathbb{Z}$ a local system of free \mathbb{Z} -modules, \mathbb{V} the corresponding algebraic vector bundle, and $\mathbb{F}^\bullet \subset \mathbb{V}$ a filtration forming a polarized variation of \mathbb{Z} -Hodge structures. By resolution of singularities we may assume that $\bar{S} \setminus S$ is a normal crossings divisor. By compactness of \bar{S} we can cover \bar{S} by local charts of the form

$$\mathcal{P} := D_\circ(1)^{\odot n} \odot D(1)^{\odot m}. \quad (187)$$

with $\bar{S} \setminus S$ given in \mathcal{P} by $\mathbf{z}_1 \cdots \mathbf{z}_n = 0$. By Borel's monodromy theorem [42, (4.5)] the connection ∇ corresponding to $\mathcal{V}_\mathbb{Z}$ has quasiunipotent monodromy in \mathcal{P} , and by a theorem [42, (4.13)] of Griffiths ∇ has logarithmic singularities on $\bar{S} \setminus S$.

Pick a basepoint $s_0 \in \mathcal{P}$ and let $b_1, \dots, b_l \in (\mathcal{V}_\mathbb{Z})_{s_0}$ be a basis for the free \mathbb{Z} -module $\mathcal{V}_\mathbb{Z}$ at s_0 . Let v_1, \dots, v_l denote a basis of algebraic sections of \mathbb{V} over \mathcal{P} compatible with the filtration \mathbb{F}^\bullet , i.e. such that the first vectors form a basis for the first filtered piece, etc. Then the matrix

$$X : \mathcal{P}_{\text{SC}} \rightarrow \text{GL}_l(\mathbb{C}), \quad X(s) := (v_i(b_j))_{i,j=1,\dots,l} \quad (188)$$

forms a horizontal section of ∇ as we analytically continue from s_0 to $s \in \mathcal{P}_{\text{SC}}$. By Theorem 22 the map $X(s)$ is definable in $\mathbb{R}_{\text{LN,exp}}$.

Denote by \check{D} the variety corresponding to the Hodge flags of dimension $\dim \mathbb{F}^\bullet$, and $q : \text{GL}_m \rightarrow \check{D}$ the quotient map sending the matrix X to the flag with the k -th piece given by the span of the first $\dim \mathbb{F}^k$ columns. Clearly q is definable even in \mathbb{R}_{alg} , so the map $q \circ X$ is also definable in $\mathbb{R}_{\text{LN,exp}}$.

Since the VHS $(S, \mathcal{V}_{\mathbb{Z}}, \mathbb{F}^\bullet)$ is polarized the map above restricts to a map $q \circ X : \mathcal{P}_{\text{SC}} \rightarrow D$ where $D \subset \check{D}$ denotes the open subspace of Hodge flags polarized by the given polarization on \mathbb{V}_{s_0} . The monodromy of ∇ on S induces a monodromy subgroup $\Gamma \subset \text{GL}(\mathbb{V}_{s_0}, \mathbb{Z})$ and factoring $q \circ X$ modulo this monodromy we obtain a well-defined map

$$\Phi : \mathcal{P} \rightarrow D/\Gamma. \quad (189)$$

In [1] it is shown that this map is definable when one chooses an appropriate \mathbb{R}_{alg} -structure on D/Γ . Essentially this is the mod- Γ quotient of the standard \mathbb{R}_{alg} -structure on Siegel domains in D . Accordingly, the statement that Φ is definable in $\mathbb{R}_{\text{LN,exp}}$ follows from the fact that $q \circ X : \mathcal{P}_{\text{SC}} \rightarrow D$ is definable in $\mathbb{R}_{\text{LN,exp}}$, plus the fact that the image of $q \circ X$ lies in the union of finitely many Siegel domains. This latter fact is established in [1, Theorem 1.5], which thus proves the definability of Φ in $\mathbb{R}_{\text{LN,exp}}$.

Remark 53 (The format of Φ). *The computation of the format of Φ in $\mathbb{R}_{\text{LN,exp}}$ involves two separate steps. First, one should effectively determine the Gauss-Manin connection in order to compute $\mathcal{F}(X)$. If the PVHS is given by the Gauss-Manin connection of a projective smooth family of $f : V \rightarrow S$ then this computation is carried out in [45]. In order to deduce a bound for $\mathcal{F}(\Phi)$ one should then also produce an effective bound on the Siegel domains needed in [1, Theorem 1.5], which appears to be a non-trivial task.*

In the rare case that the base S of the PVHS is projective, there are no singularities and Φ is in fact definable in \mathbb{R}_{LN} . In this case it is also far less difficult to estimate the number of Siegel domains that the image of $q \circ X$ meets in terms of the format of the Gauss-Manin connection.

We are now in position to finish the proof of Theorem 4.

Proof of Theorem 4. By [1, Theorem 1.1] the set Y is \mathbb{R}_{alg} -definable, so S_Y above is $\mathbb{R}_{\text{LN,exp}}$ -definable with the format depending effectively on Φ, Y . The algebraicity of S_Y then follows from the definable Chow theorem of Peterzil-Starchenko [36, Theorem 4.4] as explained in [1, Theorem 1.6]. We can now define the irreducible components of S_Y (as an algebraic variety) as the closures of the connected components of $S_Y \setminus (S_Y)_{\text{sing}}$. By effective o-minimality the number of these irreducible components, and the format of each component, are effectively bounded. Finally for each irreducible component of dimension k the degree is given by the number of intersections with k generic hyperplane sections, which is again bounded by effective o-minimality. \square

10.4. The Siegel modular variety. As a special case of the construction of §10.3, the universal covering map of the Siegel modular variety \mathcal{A}_g of principally polarized abelian varieties of genus g , restricted to an appropriate fundamental domain $F \subset \mathbb{H}_g$, is definable in $\mathbb{R}_{\text{LN,exp}}$. To see this let $\mathbb{A}_g \rightarrow \mathcal{A}_g$ be the universal abelian variety over \mathcal{A}_g and denote by ∇ the corresponding Gauss-Manin connection. Choose

a compactification $\bar{\mathcal{A}}_g$ such that $\bar{\mathcal{A}}_g \setminus \mathcal{A}_g$ is a normal crossings divisor. In a neighborhood U of every point $p \in \bar{\mathcal{A}}_g$ one can choose a basis of Shimura differentials $\omega_1, \dots, \omega_g$ generating a basis for the holomorphic differentials over U , and complete it to a basis of $H^1(\mathbb{A}_g/U)$ using meromorphic differentials of the 2nd kind $\omega_{g+1}, \dots, \omega_{2g}$. If one chooses a symplectic basis $\delta_1, \dots, \delta_{2g}$ with respect to the principal polarization for the local system $H_1(\mathbb{A}_g/U, \mathbb{Z})$ then the extended period matrix

$$X(q) := \begin{pmatrix} \oint_{\delta_1} \omega_1 & \cdots & \oint_{\delta_{2g}} \omega_1 \\ \vdots & \ddots & \vdots \\ \oint_{\delta_1} \omega_{2g} & \cdots & \oint_{\delta_{2g}} \omega_{2g} \end{pmatrix} \quad (190)$$

is a horizontal section of the Gauss-Manin connection and therefore definable in $\mathbb{R}_{\text{LN,exp}}$. Writing

$$X(q) = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \quad (191)$$

where A, B, C, D are $g \times g$ blocks, the inverse of the universal cover $e : \mathbb{H}_g \rightarrow \mathcal{A}_g$ is given locally by $B^{-1}A : \mathcal{A}_g \rightarrow \mathbb{H}_g$. It is known, for instance by [37] that the image of this map, after making suitable algebraic branch cuts in \mathcal{A}_g , meets finitely many translates of the standard fundamental domain $F \subset \mathbb{H}_g$. It follows that the graph of $\pi : F \rightarrow \mathcal{A}_g$ can be defined in $\mathbb{R}_{\text{LN,exp}}$ by gluing together finitely many translates of the (inverted) graph of $B^{-1}A$. We remark on issues related to effectivity of this construction at the end of §10.5.

In the special case $g = 1$ one obtains from the construction above the definability of the modular λ -function $\lambda : \mathbb{H} \rightarrow \mathbb{C}$ associated to the Legendre family

$$E_\lambda := \{y^2 = x(x-1)(x-\lambda)\} \quad (192)$$

as the inverse to the ratio $I_1(\lambda)/I_2(\lambda)$ of the two elliptic integrals

$$I_j := \oint_{\delta_j(\lambda)} \frac{dx}{y}, \quad j = 1, 2 \quad (193)$$

where δ_1, δ_2 form a symplectic basis for $H_1(E_\lambda, \mathbb{Z})$. In this case there is no difficulty working out an upper bound for the format of the λ -function restricted to its standard fundamental domain by hand. One should examine the asymptotics of $I_1/I_2(\cdot)$ to determine how many translates of the fundamental domain meet the image of a ball around $\lambda = 0, 1, \infty$ with a branch cut along the negative real axis, and cover the rest of $\mathbb{P}^1 \setminus \{0, 1, \infty\}$ with finitely many discs. Using

$$j = \frac{256(1 - \lambda + \lambda^2)^3}{\lambda^2(1 - \lambda)^2} \quad (194)$$

it then follows that the Klein modular invariant $j : \mathbb{H} \rightarrow \mathbb{C}$ is also definable in $\mathbb{R}_{\text{LN,exp}}$ with effectively bounded format. Thus the structure \mathbb{R}_j generated by the modular invariant is effectively o-minimal, as a substructure of $\mathbb{R}_{\text{LN,exp}}$. This answers a question raised by Pila [40, Section 13.3], who asked for the effective o-minimality of the structure \mathbb{R}_j and noted some implications for effective Andre-Oort.

10.5. The universal abelian variety over \mathcal{A}_g . In this section we show that the universal cover for the universal abelian variety over \mathcal{A}_g restricted to an appropriate fundamental domain is definable in $\mathbb{R}_{\text{LN,exp}}$. It is possible to compute this directly similarly to the construction of the previous section, but for variation we demonstrate this definability using the material of §10.2.

Let $\pi : \mathbb{A}_g \rightarrow \mathcal{A}_g$ be the universal abelian variety over \mathcal{A}_g . Denote by $*$: $\mathcal{A}_g \rightarrow \mathbb{A}_g$ the identity section. Choose a basis $\omega_1, \dots, \omega_g \in \Omega(\mathbb{A}_g/U)$ over an affine open $U \subset \mathcal{A}_g$. Define maps I_1, \dots, I_g by

$$I_j : \mathbb{A}_g \rightarrow \mathbb{C}^{2g+1}, \quad I_j(p) = \left(\oint_{\gamma_1(\pi(p))} \omega_j, \dots, \oint_{\gamma_{2g}(\pi(p))} \omega_j, \int_*^p \omega_j \right) \quad (195)$$

where the path of integration in the first integrand is chosen inside $\pi^{-1}(\pi(p))$ and

$$\gamma_1, \dots, \gamma_{2g} \in H_1(\pi^{-1}(\pi(p)), \mathbb{Z}) \quad (196)$$

form a basis, chosen over some basepoint and then analytically continued.

It is classical that the maps I_j have moderate growth. Moreover they are monodromic with locally quasiunipotent monodromy: the first $2g$ coordinates realize the monodromy of the Gauss-Manin connection of π , and the last coordinate is univalued modulo the previous coordinates. The results of §10.2 then imply that the restriction of I_j to a simply-connected semialgebraic subset $(\mathbb{A}_g)_{\text{SC}}$ of \mathbb{A}_g is $\mathbb{R}_{\text{LN,exp}}$ definable. Consider the universal covering map

$$e : \mathbb{H}_g \times \mathbb{C}^g \rightarrow \mathbb{A}_g. \quad (197)$$

As before, we can describe the inverse of e on $(\mathbb{A}_g)_{\text{SC}}$ in terms of the coordinates of I_1, \dots, I_g : on \mathbb{H}_g it is given as in §10.4, and on \mathbb{C}^g it is given by the last coordinates of I_1, \dots, I_g . It follows from [37] that $e^{-1}((\mathbb{A}_g)_{\text{SC}})$ meets finitely many translates of the standard fundamental domain $\mathcal{F} \subset \mathbb{H}_g \times \mathbb{C}^g$. Gluing together finitely many translates we deduce the following.

Theorem 24. *The restriction $e|_{\mathcal{F}}$ is definable in $\mathbb{R}_{\text{LN,exp}}$.*

Note that the computation of $\mathcal{F}(e|_{\mathcal{F}})$ involves two non-trivial tasks. First one should show how to explicitly compute (or estimate from above) the differential equations for I_j ; this is an additive extension of the Gauss-Manin connection and should be algebraically computable in principle. One should then also effectivize the Peterzil-Starchenko bound on the number of fundamental domains, which is an interesting problem that we plan to return to.

APPENDIX A. NOETHERIAN FUNCTIONS OVER NUMBER FIELDS

In this appendix we consider Noetherian functions “defined over a number field”. We use this in particular to produce some natural examples of LN-functions that are *not* Noetherian, thus demonstrating the necessity of considering the larger LN class.

A.1. Geometric formulation of Noetherian functions. For the purposes of this section we reformulate the notion of a Noetherian chain (1) in a more geometric language. We will say that a ring R is a ring of Noetherian functions in n variables if:

- (1) R is finitely generated over \mathbb{Z} by elements $F_1, \dots, F_N \in R$;
- (2) R is an integral domain.

(3) R is equipped with n commuting derivations $\xi_1, \dots, \xi_n \in \text{Der } R$.

Suppose R satisfies these conditions and let $p \in \text{Spec } R \otimes_{\mathbb{Z}} \mathbb{C}$, thought of as a ring homomorphism $p : R \rightarrow \mathbb{C}$. Then we can form a Noetherian chain (1) given by $F_1, \dots, F_N : \mathcal{P} \rightarrow \mathbb{C}$ on some sufficiently small polydisc \mathcal{P} with

$$\frac{\partial F_i}{\partial \mathbf{z}_j} = \xi_j(F_i) \quad (198)$$

such that $F_i(0) = p(F_i)$. This is just the Frobenius integrability theorem for commuting vector fields. We denote the germs of these functions by $F_1^p, \dots, F_N^p : (\mathbb{C}^n, 0) \rightarrow \mathbb{C}$.

Conversely, let $F_1, \dots, F_N : \mathcal{P} \rightarrow \mathbb{C}$ be a Noetherian chain (1). Adding all coefficients of the G_{ij} to the chain (and defining their derivatives to be zero), we may assume without loss of generality that the coefficients of G_{ij} are in \mathbb{Z} . Then the subring R of the ring of holomorphic functions on \mathcal{P} generated by F_1, \dots, F_N is a finitely generated integral domain equipped with the n commuting derivations $\xi_j = \frac{\partial}{\partial z_j}$.

A.2. Noetherian functions over a field. Let \mathbb{K} be a field of characteristic zero. We are mostly interested in the case $[\mathbb{K} : \mathbb{Q}] < \infty$ but the results of this subsection do not require this assumption. If $p \in \text{Spec } R$ is a \mathbb{K} -point then we say that the functions F_1^p, \dots, F_N^p are defined over \mathbb{K} . We say that F is defined over \mathbb{K} if it is a polynomial combination of F_1^p, \dots, F_N^p with integer coefficients.

Proposition 54. *Suppose that F is a Noetherian function and its Taylor coefficients at the origin are all in \mathbb{K} . Then it is defined over $\overline{\mathbb{K}}$ as a Noetherian function.*

Proof. Let R be a ring of Noetherian functions generated by F_1, \dots, F_N and $F = P(F_1, \dots, F_N)$. We may assume without loss of generality that the coefficients of P are integers, by including any non-integer coefficient as additional functions in the chain F_1, \dots, F_N . For $q \in \text{Spec } R$ write F^q for $P(F_1^q, \dots, F_N^q)$.

We have $F \equiv F^q$ for some $q \in \text{Spec } R$ by definition. The set of all $p \in \text{Spec } R$ such that $F \equiv F^p$ is a \mathbb{K} -variety: indeed, it is given by the conditions

$$\frac{\partial^\alpha}{\partial z^\alpha} F(0) = \frac{\partial^\alpha}{\partial z^\alpha} F^p(0) = p(\xi^\alpha F), \quad \forall \alpha \in \mathbb{N}^n. \quad (199)$$

The expressions $\xi^\alpha F$ on the right hand side are polynomials over \mathbb{Z} in F_1, \dots, F_N , and the expressions on the left hand side are in \mathbb{K} , so this is a collection of polynomial equations over \mathbb{K} . Since the solution set is non-empty, it must also contain a solution p over $\overline{\mathbb{K}}$ by the Nullstellensatz. This finishes the proof. \square

We also record a simple consequence of the definitions.

Proposition 55. *Let F be a Noetherian function over \mathbb{K} , that is $F = P(F_1^p, \dots, F_N^p)$ where P has integer coefficients and $p \in \text{Spec } R$ is a \mathbb{K} -point. Let ν be a valuation on \mathbb{K} with*

$$\nu(F_j^p(0)) \leq 0, \quad \text{for } j = 1, \dots, N. \quad (200)$$

Then

$$\nu\left(\frac{\partial^\alpha}{\partial z^\alpha} F(0)\right) \leq 0 \quad \forall \alpha \in \mathbb{N}^n. \quad (201)$$

Proof. This is simply because all derivatives of F_j^p are given by polynomials in F_1^p, \dots, F_N^p with integer coefficients, and at the origin these all belong to the valuation ring of ν . \square

A.3. An example of an LN function that is not Noetherian. Let \mathbb{K} be a number field, and

$$f(\mathbf{z}) = \sum \frac{c_{\alpha}}{\alpha!} \mathbf{z}^{\alpha} \quad (202)$$

be a Noetherian function with $c_{\alpha} \in \mathbb{K}$ for every $\alpha \in \mathbb{N}^n$. Then by Proposition 54, f is Noetherian over some finite extension $\mathbb{K}' \supset \mathbb{K}$. By Proposition 55 we have $f = P(F_1^p, \dots, F_M^p)$ where P has integer coefficients and $p \in \text{Spec } R$ is a \mathbb{K} -point. Denote by Σ' the finite set of places of \mathbb{K}' corresponding to a valuation ν such that $\nu(F_j^p(0)) > 0$ for some $j = 1, \dots, M$. Denote by Σ_f the set of restrictions of these places to \mathbb{K} . Then we have the following corollary.

Corollary 56. *For every valuation ν on \mathbb{K} corresponding to a place outside the finite set Σ_f ,*

$$\nu(c_{\alpha}) \leq 0 \quad \forall \alpha \in \mathbb{N}^n. \quad (203)$$

As a consequence, the function

$$f(z) = \frac{e^z - 1}{z} = \sum_{j=0}^{\infty} \frac{1/(j+1)}{j!} z^j \quad (204)$$

is not Noetherian: the coefficients $c_j = 1/(j+1)$ lie outside the valuation ring for infinitely many (in fact all) p -adic valuations on \mathbb{Q} . In particular, the Noetherian class is not closed under restricted divisions (or strict transforms). On the other hand f is certainly LN on $D_{\circ}(1)$, for instance by closedness under restricted division, or directly by the Noetherian chain f, z, e^z satisfying

$$\partial_z(f) = e^z - f, \quad \partial_z(z) = z, \quad \partial_z(e^z) = ze^z, \quad (205)$$

with $\partial_z = z \frac{\partial}{\partial z}$.

A.4. An LN function that does not generate a finitely-generated ring.

We close with a simple example showing that an LN function need not generate, by closing under derivatives, a finitely generated ring. In other words, to check that a function is not LN it does not suffice to check that the ring it generates is not finitely generated. Consider

$$f(z) := e^{z^2} \quad (206)$$

on $D(1)$. By straightforward computation

$$\mathbb{C}[f, \partial_z f, \dots, \partial_z^k f] = \mathbb{C}[f, zf, \dots, z^k f]. \quad (207)$$

Since z, f are algebraically independent over \mathbb{C} the chain of rings above does not stabilize as k grows, since $z^{k+1}f$ is clearly not a polynomial in $f, zf, \dots, z^k f$.

REFERENCES

- [1] B. Bakker, B. Klingler, and J. Tsimerman. Tame topology of arithmetic quotients and algebraicity of Hodge loci. *J. Amer. Math. Soc.*, 33(4):917–939, 2020.
- [2] Alessandro Berarducci and Tamara Servi. An effective version of Wilkie’s theorem of the complement and some effective o-minimality results. *Ann. Pure Appl. Logic*, 125(1-3):43–74, 2004.
- [3] Gal Binyamini. Density of algebraic points on Noetherian varieties. *Geometric and Functional Analysis*, 29(1):72–118, 2019.
- [4] Gal Binyamini. Point counting for foliations over number fields. *Forum of Mathematics, Pi*, 10:e6, 2022.

- [5] Gal Binyamini, Gareth O Jones, Harry Schmidt, and Margaret E M Thomas. An effective Pila-Wilkie theorem for sets definable using pfaffian functions, with some diophantine applications. *Preprint*, arXiv:2301.09883, January 2023.
- [6] Gal Binyamini, Dmitri Novikov, and Benny Zack. Sharply o-minimal structures and sharp cellular decomposition. *Preprint*, arXiv:1407.1183, September 2022.
- [7] Gal Binyamini and Dmitri Novikov. Intersection multiplicities of Noetherian functions. *Adv. Math.*, 231(6):3079–3093, 2012.
- [8] Gal Binyamini and Dmitri Novikov. Multiplicities of Noetherian deformations. *Geom. Funct. Anal.*, 25(5):1413–1439, 2015.
- [9] Gal Binyamini and Dmitri Novikov. Multiplicity operators. *Israel Journal of Mathematics*, 210(1):101–124, 2015.
- [10] Gal Binyamini and Dmitri Novikov. Complex cellular structures. *Ann. of Math. (2)*, 190(1):145–248, 2019.
- [11] Gal Binyamini and Dmitri Novikov. Tameness in geometry and arithmetic: beyond o-minimality. In *ICM—International Congress of Mathematicians. Vol. III. Sections 1–4*, pages 1440–1461. EMS Press, Berlin, [2023] ©2023.
- [12] Gal Binyamini, Dmitri Novikov, and Sergei Yakovenko. On the number of zeros of Abelian integrals. *Invent. Math.*, 181(2):227–289, 2010.
- [13] Gal Binyamini, Dmitri Novikov, and Benny Zack. Wilkie’s conjecture for Pfaffian structures. *Accepted to appear in Annals of Math.*, arXiv:2202.05305, 2022.
- [14] Gal Binyamini and Nicolai Vorobjov. Effective cylindrical cell decompositions for restricted sub-Pfaffian sets. *Int. Math. Res. Not. IMRN*, 2022(5):3493–3510, 11 2020.
- [15] A. Borel, P.-P. Grivel, B. Kaup, A. Haefliger, B. Malgrange, and F. Ehlers. *Algebraic D-modules*, volume 2 of *Perspectives in Mathematics*. Academic Press, Inc., Boston, MA, 1987.
- [16] Eduardo Cattani, Pierre Deligne, and Aroldo Kaplan. On the locus of Hodge classes. *J. Amer. Math. Soc.*, 8(2):483–506, 1995.
- [17] C. de la Vallée Poussin. Sur l’équation différentielle linéaire du second ordre. détermination d’une intégrale par deux valeurs assignées. extension aux équations d’ordre n. *Journal de Mathématiques Pures et Appliquées*, 8:125–144, 1929.
- [18] Pierre Deligne. *Équations différentielles à points singuliers réguliers*, volume Vol. 163 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, 1970.
- [19] A. Gabrièlov. Multiplicities of Pfaffian intersections, and the Łojasiewicz inequality. *Selecta Math. (N.S.)*, 1(1):113–127, 1995.
- [20] A. Gabrièlov. Frontier and closure of a semi-Pfaffian set. *Discrete Comput. Geom.*, 19(4):605–617, 1998.
- [21] A. Gabrièlov and N. Vorobjov. Complexity of stratifications of semi-Pfaffian sets. *Discrete Comput. Geom.*, 14(1):71–91, 1995.
- [22] A. Gabrièlov and N. Vorobjov. Complexity of cylindrical decompositions of sub-Pfaffian sets. *J. Pure Appl. Algebra*, 164(1-2):179–197, 2001.
- [23] A. Gabrièlov and N. Vorobjov. Complexity of computations with Pfaffian and Noetherian functions. In *Normal forms, bifurcations and finiteness problems in differential equations*, volume 137 of *NATO Sci. Ser. II Math. Phys. Chem.*, pages 211–250. Kluwer Acad. Publ., Dordrecht, 2004.
- [24] A. M. Gabrièlov. Projections of semianalytic sets. *Funkcional. Anal. i Priložen.*, 2(4):18–30, 1968.
- [25] Andrei Gabrièlov. Multiplicity of a zero of an analytic function on a trajectory of a vector field. In *The Arnoldfest (Toronto, ON, 1997)*, volume 24 of *Fields Inst. Commun.*, pages 191–200. Amer. Math. Soc., Providence, RI, 1999.
- [26] Andrei Gabrièlov. Relative closure and the complexity of Pfaffian elimination. In *Discrete and computational geometry*, volume 25 of *Algorithms Combin.*, pages 441–460. Springer, Berlin, 2003.
- [27] Andrei Gabrièlov and Askold Khovanskii. Multiplicity of a Noetherian intersection. In *Geometry of differential equations*, volume 186 of *Amer. Math. Soc. Transl. Ser. 2*, pages 119–130. Amer. Math. Soc., Providence, RI, 1998.
- [28] Thomas W. Grimm, Lorenz Schlechter, and Mick van Vliet. Complexity in Tame Quantum Theories. 10 2023.
- [29] M. Kashiwara. Quasi-unipotent constructible sheaves. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 28(3):757–773 (1982), 1981.

- [30] A. Khovanskii. Real analytic manifolds with the property of finiteness, and complex abelian integrals. *Funktsional. Anal. i Prilozhen.*, 18(2):40–50, 1984.
- [31] A. Khovanskii and S. Yakovenko. Generalized Rolle theorem in \mathbf{R}^n and \mathbf{C} . *J. Dynam. Control Systems*, 2(1):103–123, 1996.
- [32] A. G. Khovanskii. *Fewnomials*, volume 88 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1991. Translated from the Russian by Smilka Zdravkovska.
- [33] A. G. Khovanskii. Fewnomials and Pfaff manifolds. In *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Warsaw, 1983)*, pages 549–564. PWN, Warsaw, 1984.
- [34] Jean-Marie Lion, Chris Miller, and Patrick Speissegger. Differential equations over polynomially bounded o-minimal structures. *Proc. Amer. Math. Soc.*, 131(1):175–183, 2003.
- [35] D. Novikov and S. Yakovenko. Trajectories of polynomial vector fields and ascending chains of polynomial ideals. *Ann. Inst. Fourier (Grenoble)*, 49(2):563–609, 1999.
- [36] Ya'acov Peterzil and Sergei Starchenko. Complex analytic geometry and analytic-geometric categories. *J. Reine Angew. Math.*, 626:39–74, 2009.
- [37] Ya'acov Peterzil and Sergei Starchenko. Definability of restricted theta functions and families of abelian varieties. *Duke Math. J.*, 162(4):731–765, 2013.
- [38] J. Pila and A. J. Wilkie. The rational points of a definable set. *Duke Math. J.*, 133(3):591–616, 2006.
- [39] Jonathan Pila. On the algebraic points of a definable set. *Selecta Math. (N.S.)*, 15(1):151–170, 2009.
- [40] Jonathan Pila. O-minimality and the André-Oort conjecture for \mathbf{C}^n . *Ann. of Math. (2)*, 173(3):1779–1840, 2011.
- [41] Jonathan Pila. *Point-counting and the Zilber-Pink conjecture*, volume 228 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2022.
- [42] Wilfried Schmid. Variation of Hodge structure: the singularities of the period mapping. *Invent. Math.*, 22:211–319, 1973.
- [43] Patrick Speissegger. The Pfaffian closure of an o-minimal structure. *J. Reine Angew. Math.*, 508:189–211, 1999.
- [44] Jean-Claude Tougeron. Algèbres analytiques topologiquement noethériennes. Théorie de Khovanskii. *Ann. Inst. Fourier (Grenoble)*, 41(4):823–840, 1991.
- [45] David Urbanik. Sets of special subvarieties of bounded degree. *Compos. Math.*, 159(3):616–657, 2023.
- [46] A. N. Varchenko. Estimation of the number of zeros of an abelian integral depending on a parameter, and limit cycles. *Funktsional. Anal. i Prilozhen.*, 18(2):14–25, 1984.
- [47] A. J. Wilkie. Model completeness results for expansions of the ordered field of real numbers by restricted Pfaffian functions and the exponential function. *J. Amer. Math. Soc.*, 9(4):1051–1094, 1996.
- [48] A. J. Wilkie. A theorem of the complement and some new o-minimal structures. *Selecta Math. (N.S.)*, 5(4):397–421, 1999.
- [49] Sergei Yakovenko. On functions and curves defined by ordinary differential equations. In *The Arnoldfest (Toronto, ON, 1997)*, volume 24 of *Fields Inst. Commun.*, pages 497–525. Amer. Math. Soc., Providence, RI, 1999.
- [50] Umberto Zannier. *Some problems of unlikely intersections in arithmetic and geometry*, volume 181 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2012. With appendixes by David Masser.

WEIZMANN INSTITUTE OF SCIENCE, REHOVOT, ISRAEL
 Email address: gal.binyamini@weizmann.ac.il