

THE e -POSITIVITY OF THE CHROMATIC SYMMETRIC FUNCTION FOR TWINNED PATHS AND CYCLES

ESTHER BANAIAN, KYLE CELANO, MEGAN CHANG-LEE, LAURA COLMENAREJO,
OWEN GOFF, JAMIE KIMBLE, LAUREN KIMPEL, JOHN LENTFER, JINTING LIANG,
AND SHEILA SUNDARAM

ABSTRACT. The operation of twinning a graph at a vertex was introduced by Foley, Hoàng, and Merkel (2019), who conjectured that twinning preserves e -positivity of the chromatic symmetric function. A counterexample to this conjecture was given by Li, Li, Wang, and Yang (2021). In this paper, we prove that e -positivity is preserved by the twinning operation on cycles, by giving an e -positive generating function for the chromatic symmetric function, as well as an e -positive recurrence. We derive similar e -positive generating functions and recurrences for twins of paths. Our methods make use of the important triple deletion formulas of Orellana and Scott (2014), as well as new symmetric function identities.

CONTENTS

1. Introduction	1
1.1. Main results	3
2. Preliminaries	4
2.1. Graph operations	4
2.2. Known e -positivity results	6
2.3. A new statistic on partitions	8
3. e -positivity via generating functions	9
3.1. Symmetric function identities and technical lemmas	10
3.2. Generating functions for twinned paths	12
3.3. Generating function for twinned cycles	21
4. e -positivity via recurrences	24
4.1. Recurrences for twinned paths	25
4.2. Recurrence for twinned cycles	29
5. Future directions	33
Acknowledgments	34
References	34

1. INTRODUCTION

The *chromatic symmetric function* of a graph $G = (V, E)$ is defined by Stanley [18] to be

$$X_G(\mathbf{x}) = \sum_{\kappa} \prod_{v \in V} x_{\kappa(v)},$$

2020 *Mathematics Subject Classification.* 05E05, 05C15.

Key words and phrases. chromatic symmetric function, graph twinning, elementary symmetric function, e -positivity.

where the sum is over all proper colorings $\kappa : V \rightarrow \mathbb{Z}_{>0}$ of G by positive integers. The goal of this paper is to study the effect that a small change to the graph G has on $X_G(\mathbf{x})$. Specifically, we look at the change in $X_G(\mathbf{x})$ when one *twins* (or *clones*) a vertex v of a graph G , that is, when one adds a vertex v' incident to v and all its neighbors, to produce a new graph G_v . Precise definitions of this operation and related terms appear in Section 2.

Question 1.1. *For a given vertex v of a graph G , how are the polynomials $X_{G_v}(\mathbf{x})$ and $X_G(\mathbf{x})$ related?*

In the seminal paper [18], Stanley proved that the chromatic symmetric functions of paths and cycles are e -positive, that is, their expansion in the basis of elementary symmetric functions has nonnegative coefficients. As observed in [18], the result for paths is originally due to Carlitz, Scoville, and Vaughan in a different context [2, p.242]. More generally, spurred by the following conjecture of Stanley and Stembridge, much of the research on the chromatic symmetric function has centered around the incomparability graph $\text{Inc}(P)$ of a $(3+1)$ -free poset P , defined as a poset containing no induced subposet isomorphic to the disjoint union of a 3-chain and a 1-chain.

Conjecture 1.2 ([20, 18]). *If P is a $(3+1)$ -free poset, then $X_{\text{Inc}(P)}(\mathbf{x})$ is e -positive.*

To *twinn* a poset P at a vertex v , producing P_v , is to add an element v' such that v' is incomparable to w if and only if either $w = v$ or w is incomparable to v . Note that if $G = \text{Inc}(P)$, then $\text{Inc}(P_v) = G_v$. This next simple lemma is the main motivation for considering the twinning operation. Its proof is immediate from the fact that if $u < v$, then $u < v'$, and if $v < w$ then $v' < w$.

Lemma 1.3. *The twin of a $(3+1)$ -free poset is $(3+1)$ -free.*

One can therefore make a weakened version of the Stanley–Stembridge conjecture, first appearing in the work of Foley, Hoàng, and Merkel [8].

Conjecture 1.4 ([8]). *If P is $(3+1)$ -free and $X_{\text{Inc}(P)}(\mathbf{x})$ is e -positive, then $X_{\text{Inc}(P_v)}(\mathbf{x})$ is e -positive for any $v \in P$.*

Remark 1.5. Li, Li, Wang, and Yang [12, Theorem 3.6] prove that the twinning operation on graphs does not always preserve e -positivity. They give an example of a graph G [12, Theorem 4.1] that is not an incomparability graph of a $(3+1)$ -free poset, but whose chromatic symmetric function $X_G(\mathbf{x})$ is e -positive, and show that for a certain vertex v of G , the chromatic symmetric function for the twinned graph G_v does not expand positively even in the Schur basis, and so it cannot be e -positive. This suggests that there is something special about the twinning operation on $(3+1)$ -free posets.

If Conjecture 1.4 is solved, then the Stanley–Stembridge conjecture can be reduced to the following conjecture. We say that a graph H is *twin-free* if there is no graph G with vertex v such that $H = G_v$.

Conjecture 1.6. *If P is $(3+1)$ -free and twin-free, then $X_{\text{Inc}(P)}(\mathbf{x})$ is e -positive.*

Other than [8, 12], twinning does not seem to have been studied as an operation in its own right. Rather, there has been incidental progress over the years. In 2001, Gebhard and Sagan [10] introduced the stronger notion of (e) -positivity of chromatic symmetric functions in noncommuting variables. This implies e -positivity for chains of complete graphs [10,

Corollary 7.7], and includes twins of paths as a special case. Later, Dahlberg and van Willigenburg [7] gave a direct proof of e -positivity for *lollipop* graphs, which are a special case of [10, Corollary 7.7], and which again include paths twinned at a leaf.

Throughout the paper, we refer to a property of the *chromatic symmetric function of the graph G* as being a property of *the graph G* . For instance, we use interchangeably the phrases “the generating function of the chromatic symmetric function of a graph” and “the generating function of a graph”. Similarly, we use “the chromatic symmetric function of the graph G is e -positive” and “the graph G is e -positive” interchangeably.

1.1. Main results. This paper studies the effect of twinning on the e -expansions of the chromatic symmetric function of certain graphs. We specifically look at twins of path and cycle graphs, a few of which are shown in Figure 2. A summary of our progress on Question 1.1 follows.

Our first main contribution is a series of explicit e -positive formulas for the generating function of the following families of twinned graphs:

- (1) The path twinned at one leaf (Proposition 3.9)
- (2) The path twinned at both leaves (Theorem 3.14)
- (3) The path twinned at an interior vertex (Theorem 3.24)
- (4) The cycle twinned at a vertex (Theorem 3.29)

The fourth family, examined in detail in Section 3.3, and culminating in Theorem 3.29, was not known to be e -positive until now. The first three families appear in [10] and were shown to have the stronger property of (e)-positivity of their chromatic symmetric functions with noncommuting variables [10, Theorems 7.6 and 7.8]. The e -positivity of the first graph was later re-established directly in [7].

However, the explicitly e -positive expressions for the generating functions that we give in Proposition 3.9, Theorem 3.14 and Theorem 3.24 are new. Also, in Corollary 3.4 we provide a new e -positive expression for the generating function of the path that isolates the terms containing e_1 . Our derivations make crucial use of the triple deletion formula of Orellana and Scott [15].

For all but the third family, the expression we obtain for the generating function has the form

$$\sum_{n \geq 0} X_{G_n} z^n = \frac{f_G}{1 - \sum_{i \geq 2} (i-1) e_i z^i} + h_G$$

where f_G and h_G are some e -positive functions depending on the family and h_G has finite degree. The third family has h_G with infinite degree. Note that the denominator in the rational expression above coincides with the denominator in the generating function for both the path and the cycle (see Theorem 3.1). This allows us to easily obtain explicit formulas for the coefficients of the elementary symmetric functions (see Corollary 3.10, Corollary 3.15, Corollary 3.30). We state our formulas for the coefficients using a new statistic $\varepsilon(\lambda)$ associated with a partition λ (see Section 2.3). This statistic appears naturally when computing the e -coefficients of the path and cycle, and appears to be of independent interest.

The identities presented in Section 3.1, particularly in Lemma 3.3 and its proof, are the starting point for our e -positivity results. They also seem interesting in their own right.

Our second main contribution is an e -positive recurrence relation for each of the families listed above, as well as a graph appearing in the computation for the twinned cycle that we call the *moose graph*, which was previously not known to be e -positive:

- (1) The path twinned at one leaf (Proposition 4.2)
- (2) The path twinned at both leaves (Proposition 4.3)
- (3) The path twinned at an interior vertex (Theorem 4.4)
- (4) The cycle twinned at a vertex (Theorem 4.6)
- (5) The moose graph (Theorem 4.7)

2. PRELIMINARIES

In this section, we define the basic notions used throughout the paper, as well as discuss previous results. We assume a familiarity with symmetric functions as in [19, Chapter 7] or [14].

A *graph* G is a pair of sets (V, E) where V is the set of *vertices* and E is a set of 2-element subsets of vertices, called *edges*. We denote edges by $\{u, v\}$ or simply by uv . We assume that V and E are both finite, and that the graph is simple (i.e., there are no loops and no multiple edges). A *leaf* of a graph is a vertex contained within exactly one edge. An *internal vertex* is a vertex contained within at least two edges. Two graphs that are important for this paper are the *path* P_n , which has vertex set $V = [n] = \{1, \dots, n\}$ and edge set $E = \{\{i, i + 1\} \mid i \in [n - 1]\}$, and the *cycle* C_n , which also has vertex set $V = [n]$ and edge set $E = \{\{i, i + 1\} \mid i \in [n - 1]\} \cup \{1, n\}$. We illustrate them in Figure 1.




FIGURE 1. The path and cycle graphs.

A *proper coloring* of a graph $G = (V, E)$ is a function $\kappa : V \rightarrow \mathbb{Z}_{>0}$ such that if $uv \in E$, then $\kappa(u) \neq \kappa(v)$. Let $\mathbf{x} = (x_1, x_2, \dots)$ be an infinite set of commuting variables. The *chromatic symmetric function* of a graph $G = (V, E)$ is defined to be

$$X_G := X_G(\mathbf{x}) = \sum_{\kappa} \prod_{v \in V} x_{\kappa(v)},$$

where the sum is over all proper colorings κ of G . This symmetric function was first introduced by Stanley in [18] and has been studied by numerous authors since then. One central goal has been to characterize graphs G for which X_G is *e*-positive, i.e., X_G has nonnegative coefficients in the elementary symmetric function basis. We give an overview of the previous *e*-positivity results in Section 2.2.

2.1. Graph operations. We begin by defining the two operations on graphs that appear in this paper.

Given an edge ϵ of a graph G , the *deletion of ϵ in G* is the graph, denoted by $G - \epsilon$, obtained by removing the edge ϵ from G . The following formulas of Orellana and Scott are used extensively in our arguments and we refer to them as the *triple deletion arguments*.

Proposition 2.1 (Triple Deletion Formula [15, Theorem 3.1]). *Let G be any graph. Suppose edges $\epsilon_1, \epsilon_2, \epsilon_3$ form a triangle in G . Then,*

$$X_G = X_{G-\epsilon_1} + X_{G-\epsilon_2} - X_{G-\{\epsilon_1, \epsilon_2\}}.$$

Notice that Proposition 2.1 requires the graph to contain a triangle. However, one can use this formula to derive other relations for graphs that do not necessarily contain a triangle. An example of such a relation is the following.

Corollary 2.2 ([15, Corollary 3.2]). *Let $\epsilon_1 = vv_1 \in E$, $\epsilon_2 = vv_2 \in E$ and suppose $\epsilon_3 = v_1v_2 \notin E$. Then*

$$X_G = X_{(G-\epsilon_1) \cup \epsilon_3} + X_{G-\epsilon_2} - X_{(G-\{\epsilon_1, \epsilon_2\}) \cup \epsilon_3}.$$

We now introduce the main operation studied in this paper.

Definition 2.3. Given a graph G and a vertex v , the *twin* of G at v is the graph, denoted by G_v , obtained by adding a new vertex v' and connecting v' to v and to all of its neighbors. We refer to this operation as the *twinning* of a graph and to the resulting graph G_v as the *twinned graph*. By extension, $G_{v,w}$ denotes the graph G twinned at the vertices v and w in succession.

A simple example is the complete graph $G = K_n$ on n vertices. For any vertex v , $(K_n)_v$ is the complete graph K_{n+1} . We illustrate in Figure 2 the twinned path at a leaf and at an interior vertex, and the twinned cycle.




FIGURE 2. Twinning of the path and cycle graphs.

Twinning a non-isolated vertex always produces a triangle, so the triple deletion argument is a natural method to reduce the twinned graph back to the original one, as the following result shows.

Corollary 2.4. *Let H be a graph on n vertices and let u be a vertex of H . Let H' be the graph obtained by adding a new vertex v and the edge uv to H and let H'' be the graph obtained by adding a new vertex w and the edge vw to H' . Finally let H'_v be the graph H' twinned at vertex v , with v' denoting the new vertex. Then*

$$X_{H'_v} = 2(X_{H''} - e_2 X_H).$$

Proof. This is clear by the triple deletion argument using the edges uv, vw of the triangle $\{u, v, v'\}$ as shown in Figure 3. Note also that $X_{P_2} = 2e_2$. \square




FIGURE 3. The triple deletion argument used in Corollary 2.4

2.2. Known e -positivity results. Stanley defined the chromatic symmetric function X_G of a graph G in 1995. Since then, many families of graphs have been examined. We provide an extensive, but by no means exhaustive, list of known e -positivity results in Table 1. We do not define these classes of graphs, but instead provide references containing their definitions as well as proofs of their e -positivity classification. By convention, a family of graphs listed as “not e -positive” means that there is at least one graph in that class that is not e -positive. The table is roughly sorted chronologically by reference, and it is condensed so that some subclasses of other results are omitted.

Graph	Positivity	Reference
Paths	e -positive	[18, Proposition 5.3]
Cycles	e -positive	[18, Proposition 5.4]
Complete graphs	e -positive	[20, Equation 3.1]
Co-triangle-free graphs	e -positive	[20, Theorem 4.3]
K_α -chains	e -positive	[10, Corollary 7.7]
Diamond and path chains	e -positive	[10, Theorem 7.8]
(claw, P_4)-free graphs	e -positive	[23, Theorem 1.4]
(claw, diamond)-free graphs	not e -positive	[11, Lemma 7]
(claw, co-claw)-free graphs	not e -positive	[11, Lemma 7]
(claw, K_4)-free graphs	not e -positive	[11, Lemma 7]
(claw, $4K_1$)-free graphs	not e -positive	[11, Lemma 7]
(claw, $2K_2$)-free graphs	not e -positive	[11, Lemma 7]
(claw, C_4)-free graphs	not e -positive	[11, Lemma 7]
(claw, paw)-free graphs	e -positive	[11, Theorem 3]
(claw, co-paw)-free graphs	e -positive	[11, Theorem 4]
Generalized bull graphs	e -positive	[3, Theorem 3.7]
Lollipops and lariats	e -positive	[7, Theorem 9]
P_3 -free graphs	e -positive	[8, Theorem 5]
(claw, K_3)-free graphs	e -positive	[8, Theorem 5]
(claw, co- P_3)-free graphs	e -positive	[8, Theorem 5]
(co-claw)-free unit interval graphs	e -positive	[8, Theorem 18]
Generalized pyramid graphs	e -positive	[13, Theorem 7]
$2K_2$ -free unit interval graphs	e -positive	[13, Theorem 13]
Triangular ladders	e -positive	[4, Theorem 22]
Star graphs	not e -positive	[6, Example 11]
Spider graphs	e -positive	[5, Theorem 3.2]
Saltire and augmented saltire graphs	not e -positive	[5, Lemmas 4.4, 4.9]
Triangular tower graphs	not e -positive	[5, Lemma 5.4]
Tadpole graphs	e -positive	[12, Theorem 3.1]
Line graphs of tadpole graphs	e -positive	[24, Corollary 3.3]
Cycle-chord graphs $CC_{m,3}$ for any m	e -positive	[24, Corollary 4.6]
Kayak paddle graphs	e -positive	[1, Proposition 6.7]
Generalized nets	not e -positive	[9, Theorem 1]
Melting K_α -chains	e -positive	[22, Theorem 3.19]

TABLE 1. Known e -positivity results and their references.

2.3. A new statistic on partitions. In this subsection, we introduce a new statistic on the set of partitions that will allow us to describe e -coefficients more compactly.

Recall that a *partition* λ of n , written $\lambda \vdash n$, is a weakly decreasing sequence of positive integers $\lambda = (\lambda_1, \dots, \lambda_\ell)$ that sum to n , that is, $\sum_i \lambda_i = n$. We write $\ell = \ell(\lambda)$ for the *length* of the partition, that is, the number of entries in the sequence. A partition λ of n can also be written as $\lambda = \langle 1^{m_1}, 2^{m_2}, \dots, n^{m_n} \rangle$, where $m_i = m_i(\lambda) \geq 0$ denotes the multiplicity of the part i in λ . The *support* of λ , denoted $\text{supp}(\lambda)$, is the set of distinct parts appearing in λ , that is, $\text{supp}(\lambda) := \{i \in \mathbb{Z}_{>0} : m_i(\lambda) \geq 1\}$.

Now we are ready to introduce the new statistic.

Definition 2.5. For a partition λ , define $\varepsilon(\lambda)$ to be the quantity

$$(1) \quad \varepsilon(\lambda) := \prod_{j \in \text{supp}(\lambda)} (j-1)^{m_j(\lambda)} \frac{\ell(\lambda)!}{\prod_j m_j(\lambda)!} \quad \text{with} \quad \varepsilon(\emptyset) = 1.$$

Moreover, for a partition λ of n and a part a such $m_a(\lambda) \geq 1$, let $\lambda - a$ denote the partition of $n - a$ obtained by deleting one part equal to a from λ . By convention, if a is not a part of λ , we set $\varepsilon(\lambda - a) = 0$.

For example, $\varepsilon((n)) = n - 1$ and $\varepsilon((2^n)) = 1$ for any positive integer n . Additionally, $\varepsilon(\lambda) = 0$ if λ contains a 1. In Table 2, we include several examples of partitions λ together with their statistic $\varepsilon(\lambda)$.

λ	(2)	(3)	(4)	(2,2)	(5)	(3,2)	(6)	(4,2)	(3,3)	(2,2,2)	
$\varepsilon(\lambda)$	1	2	3	1	4	4	5	6	4	1	
λ	(7)	(5,2)	(4,3)	(3,2,2)	(8)	(6,2)	(5,3)	(4,4)	(4,2,2)	(3,3,2)	(2,2,2,2)
$\varepsilon(\lambda)$	6	8	12	6	7	10	16	9	9	12	1

TABLE 2. Examples of $\varepsilon(\lambda)$ for some partitions λ .

Remark 2.6. Note that the fraction in (1) is the multinomial coefficient

$$\binom{\ell(\lambda)}{m_1(\lambda), \dots, m_n(\lambda)},$$

which is always a nonnegative integer. Thus, $\varepsilon(\lambda)$ is also a nonnegative integer. In fact, $\varepsilon(\lambda) = 0$ if and only if 1 is a part in λ , i.e., $m_1(\lambda) \geq 1$.

Next, we present other properties of $\varepsilon(\lambda)$.

Lemma 2.7. *Let λ and μ be partitions of n and m , respectively. Then, we have the following:*

$$(a) \text{ For } j \in \text{supp}(\lambda), (j-1)\varepsilon(\lambda - j) = m_j(\lambda) \frac{\varepsilon(\lambda)}{\ell(\lambda)};$$

$$(b) \varepsilon(\lambda) = \sum_{j \in \text{supp}(\lambda)} (j-1)\varepsilon(\lambda - j); \text{ and}$$

$$(c) \varepsilon(\lambda)\varepsilon(\mu) = \varepsilon(\lambda \cup \mu) \binom{\ell(\lambda \cup \mu)}{\ell(\lambda)}^{-1} \prod_{j \in \text{supp}(\lambda \cup \mu)} \binom{m_j(\lambda \cup \mu)}{m_j(\lambda)} \text{ where } \lambda \cup \mu \text{ is the partition of } n + m \text{ formed by listing the parts of } \lambda \text{ and } \mu \text{ together in decreasing order.}$$

Proof. (a) Note first that both sides are identically zero if $1 \in \text{supp}(\lambda)$. For $j \in \text{supp}(\lambda)$ with $j \neq 1$, this identity follows from the definition by noticing that

$$\varepsilon(\lambda) = (j-1) \frac{\ell(\lambda)}{m_j(\lambda)} \left((j-1)^{m_j(\lambda)-1} \prod_{l \in \text{supp}(\lambda), l \neq j} (l-1)^{m_l(\lambda)} \frac{(\ell(\lambda)-1)!}{(m_j(\lambda)-1) \prod_{l \neq j} m_l(\lambda)!} \right).$$

(b) This identity follows from the definition of $\varepsilon(\lambda)$, using $\sum_{j \in \text{supp}(\lambda)} m_j(\lambda) = \ell(\lambda)$.

(c) This identity follows by expanding $\varepsilon(\lambda \cup \mu)$, using $m_j(\lambda \cup \mu) = m_j(\lambda) + m_j(\mu)$ and $\ell(\lambda \cup \mu) = \ell(\lambda) + \ell(\mu)$. \square

Remark 2.8. Intuitively, the formula for $\varepsilon(\lambda)$ can be interpreted as the number of pairs (w, f) of words w on the set $\{1, \dots, \ell(\lambda)\}$ of type λ , i.e. with λ_i occurrences of the letter i , together with a function $f : \{1, \dots, \ell(\lambda)\} \rightarrow \mathbb{Z}$ satisfying $1 \leq f(j) \leq \lambda_j - 1$ for each $j \in \{1, \dots, \ell(\lambda)\}$. These are exactly the *codes* of Stembridge [21] with no fixed points and can be used to prove Lemma 2.7 combinatorially. For example, the right-hand side of part (b) can be interpreted as the number of ways of making a code of type λ from a code whose type has length $\ell(\lambda) - 1$.

3. e -POSITIVITY VIA GENERATING FUNCTIONS

For given family of graphs $G = \{G_n\}_{n \geq 0}$, one can show e -positivity of X_{G_n} by showing that its generating function

$$\mathcal{X}_G(z) = \sum_{n \geq 0} X_{G_n} z^n$$

can be written in the form

$$(2) \quad \mathcal{X}_G(z) = \frac{P(z)}{1 - Q(z)},$$

where $P(z)$ and $Q(z)$ are e -positive formal power series in z . For the path P_n and the cycle C_n , it is known from Stanley's original paper [18] that this can be done. (See also [2, p.242] for paths.)

Theorem 3.1 ([18, Propositions 5.3 and 5.4]).

$$\begin{aligned} \mathcal{X}_P(z) &:= \sum_{n \geq 0} X_{P_n} z^n = \frac{\sum_{i \geq 0} e_i z^i}{1 - \sum_{i \geq 1} (i-1) e_i z^i}, \\ \mathcal{X}_C(z) &:= \sum_{n \geq 2} X_{C_n} z^n = \frac{\sum_{i \geq 2} i(i-1) e_i z^i}{1 - \sum_{i \geq 1} (i-1) e_i z^i}. \end{aligned}$$

In this section, we establish identities of the form (2) for several families of twinned graphs by applying generating function techniques to the relations obtained from the triple deletion argument.

It is useful to convert the preceding result to a recurrence relation for the chromatic symmetric function as follows. We will use this formulation several times in this paper, notably in the proofs of Lemma 3.23 and Proposition 3.25, as well as in Section 4.

Proposition 3.2. *We have the following recurrence relations:*

- (a) For $n \geq 3$, $X_{P_n} = ne_n + \sum_{j=2}^{n-1} (j-1)e_j X_{P_{n-j}} = ne_n + \sum_{i=1}^{n-2} (n-i-1)e_{n-i} X_{P_i}$, with initial conditions $X_{P_1} = e_1$ and $X_{P_2} = 2e_2$.
- (b) For $n \geq 4$, $X_{C_n} = n(n-1)e_n + \sum_{j=2}^{n-2} (j-1)e_j X_{C_{n-j}}$, with initial conditions $X_{C_1} = 0$, $X_{C_2} = 2e_2$, and $X_{C_3} = 6e_3$.

3.1. Symmetric function identities and technical lemmas. In this section, we examine more closely the relationship between the generating function $E(z)$ for the elementary symmetric functions, and the generating functions $\mathcal{X}_P(z)$ and $\mathcal{X}_C(z)$ for the chromatic symmetric functions of the path and the cycle. We also present some formulas for several families of coefficients appearing in the e -expansion of $\mathcal{X}_P(z)$ and $\mathcal{X}_C(z)$. We start by introducing some definitions and notation to facilitate our study.

Let $E(z) := \sum_{i \geq 0} e_i z^i$ be the generating function for the elementary symmetric functions and define

$$D(z) := E(z) - zE'(z) = 1 - \sum_{i \geq 2} (i-1)e_i z^i.$$

Theorem 3.1 can then be rewritten as:

$$(3) \quad \mathcal{X}_P(z) = \frac{E(z)}{D(z)} \quad \text{and} \quad \mathcal{X}_C(z) = \frac{z^2 E''(z)}{D(z)}.$$

It will be useful for our study to collect here the definitions of several e -positive series and their truncations and tails. Considering $k \geq 2$ whenever it appears, we define

$$(4) \quad \begin{aligned} K(z) &= \sum_{i \geq 2} i e_i z^i, & K_{\geq k}(z) &= \sum_{i \geq k} i e_i z^i, \\ G(z) &= 1 - D(z) = \sum_{i \geq 2} (i-1) e_i z^i, & G_{\geq k}(z) &= \sum_{i \geq k} (i-1) e_i z^i, \\ \frac{1}{D(z)} &= \sum_{i \geq 0} G(z)^i, & G_{\leq k}(z) &= \sum_{2 \leq i \leq k} (i-1) e_i z^i = G(z) - G_{\geq k+1}(z). \end{aligned}$$

The next lemma collects some e -positivity results concerning the generating functions introduced above.

Lemma 3.3.

(a) The following expressions are e -positive:

- (i) $z^2 E''(z) - zE'(z) + e_1 z$;
- (ii) $2z^2 E''(z) - 3zE'(z) + 3e_1 z + 2e_2 z^2$; and
- (iii) $z^2 E''(z) - 3zE'(z) + 3E(z) + e_2 z^2$.

(b) The following expressions can be written as rational functions with e -positive numerators:

- (i) $\mathcal{X}_P(z) - (1 + e_1 z)$; and
- (ii) $(1 + e_1 z)\mathcal{X}_C(z) - \mathcal{X}_P(z) + 1 + e_1 z$.

Proof.

(a) The e -positivity results follow, respectively, from the identities:

- (i) $z^2 E''(z) - zE'(z) = -e_1 z + \sum_{i \geq 3} i(i-2)e_i z^i$;
- (ii) $2z^2 E''(z) - 3zE'(z) = -3e_1 z - 2e_2 z^2 + \sum_{i \geq 3} (2i^2 - 5i)e_i z^i$; and
- (iii) $z^2 E''(z) - 3zE'(z) + 3E(z) = 3 - e_2 z^2 + \sum_{i \geq 4} (i-1)(i-3)e_i z^i$.

(b) For the results concerning e -positive numerators, we have that:

(i) By (3),

$$\begin{aligned} \mathcal{X}_P(z) - (1 + e_1 z) &= \frac{zE'(z) + e_1 z[zE'(z) - E(z)]}{D(z)} \\ &= \frac{\sum_{i \geq 2} i e_i z^i + e_1 z \sum_{i \geq 2} (i-1)e_i z^i}{D(z)}. \end{aligned}$$

(ii) By the previous item,

$$\begin{aligned} (1 + e_1 z)\mathcal{X}_C(z) - \mathcal{X}_P(z) + 1 + e_1 z &= \frac{(1 + e_1 z)[z^2 E''(z) - zE'(z)] + e_1 z E(z)}{D(z)} \\ &= \frac{(1 + e_1 z)F_1(z) + e_1 z(E(z) - 1 - e_1 z)}{D(z)}, \end{aligned}$$

where $F_1(z) = \sum_{i \geq 3} i(i-2)e_i z^i$ and $E(z) - 1 - e_1 z = \sum_{i \geq 2} e_i z^i$ are e -positive. \square

Using Lemma 3.3, we get an e -positive expression for the generating function for paths that isolates those terms containing e_1 and that is different from the one in Theorem 3.1.

Corollary 3.4. *We have*

$$\mathcal{X}_P(z) = \frac{K(z)}{D(z)} + e_1 z \frac{G(z)}{D(z)} + (1 + e_1 z).$$

We now analyze these generating functions to extract closed formulas for the coefficients in the e -expansions. Recall the statistic on partitions $\varepsilon(\lambda)$ introduced in Section 2.3.

We start with a result that shows the relation between the coefficients of $G(z)^k$ and $\frac{1}{D(z)}$ in their e -expansion and $\varepsilon(\lambda)$.

Lemma 3.5. *The coefficient of $e_\lambda z^{|\lambda|}$ in $G(z)^k$ is $\varepsilon(\lambda)$ and hence*

$$\frac{1}{D(z)} = \sum_{\lambda} \varepsilon(\lambda) e_\lambda z^{|\lambda|},$$

where the sum is over all partitions λ .

Proof. This follows by manipulating the formal series directly:

$$G(z)^k = \left(\sum_{i \geq 2} (i-1)e_i z^i \right)^k = \sum_{\lambda \vdash k} e_\lambda z^{|\lambda|} \prod_{i \geq 2} (i-1)^{m_i(\lambda)} = \sum_{\lambda \vdash k} \varepsilon(\lambda) e_\lambda z^{|\lambda|}. \quad \square$$

We end this subsection by showing that several families of coefficients in the e -expansion of $\mathcal{X}_P(z)$ and $\mathcal{X}_C(z)$ can be expressed compactly in terms of $\varepsilon(\lambda)$. (See also [25].)

Proposition 3.6. *Given a graph G , let c_λ be the coefficient of $z^{|\lambda|}e_\lambda$ in \mathcal{X}_G , that is, $\mathcal{X}_G = \sum c_\lambda z^{|\lambda|}e_\lambda$. Then, we have the following:*

$$(a) \text{ For } G = P_n, c_\lambda = \varepsilon(\lambda) + \sum_{a \in \text{supp}(\lambda)} \varepsilon(\lambda - a) = \sum_{a \in \text{supp}(\lambda)} a \varepsilon(\lambda - a).$$

$$\text{In particular, if } \lambda = 1 \cup \mu \text{ for some partition } \mu, \text{ then } c_\lambda = \sum_{\substack{a \in \text{supp}(\mu) \\ a \geq 2}} (a-1) \varepsilon(\mu - a).$$

Moreover, we can also extract particular coefficients like

$$c_{(n)} = n, \quad c_{(n-1,1)} = n-2, \quad c_{(2^k)} = 2, \quad \text{and} \quad c_{(2^k,1)} = 1.$$

$$(b) \text{ For } G = C_n, \text{ we have that } c_\lambda = \sum_{a \in \text{supp}(\lambda)} a(a-1) \varepsilon(\lambda - a).$$

Proof. We use the generating functions in (3).

- (a) The first expression comes directly from the path generating function $\mathcal{X}_P(z)$ and the second expression also follows from Lemma 3.3. The equality of the two expressions and the case when $\lambda = 1 \cup \mu$ follow using Lemma 2.7.
- (b) The formula for this coefficient comes directly from the cycle generating function $\mathcal{X}_C(z)$. \square

3.2. Generating functions for twinned paths. In this section, we focus on studying the various ways to twin a path. The following is a key result.

Lemma 3.7. *For $k \geq 2$, the rational function $\frac{1 - G_{\leq k}(z)}{D(z)}$ and the function $\mathcal{X}_P(z)(1 - G_{\leq k}(z))$ are e -positive.*

Proof. For the rational function, we have that

$$(5) \quad \frac{1 - G_{\leq k}(z)}{D(z)} = \frac{1 - G(z) + G_{\geq k+1}(z)}{1 - G(z)} = 1 + \frac{G_{\geq k+1}(z)}{D(z)}.$$

This is e -positive since $G_{\geq k+1}(z) = \sum_{i \geq k+1} (i-1)e_i z^i$ and $\frac{1}{D(z)}$ expands e -positively in powers of $G(z)$. The e -positivity of the second function follows from (3) and the identity

$$\mathcal{X}_P(z) (1 - G_{\leq k}(z)) = \mathcal{X}_P(z) + E(z) G_{\geq k+1}(z). \quad \square$$

3.2.1. Paths twinned at a leaf.

The recurrence for the chromatic symmetric function of twinned paths at a leaf (i.e., a vertex of degree 1) appears in Dahlberg and van Willigenburg [7, Equation 5], where the graph $P_{n,v}$ with v a leaf is called the *lariat graph* L_{n+3} . Its chromatic symmetric function had been considered earlier by Wolfe in [25], and e -positivity was first established by Gebhard and Sagan in [10, Corollary 7.7].

Proposition 3.8. *Let v be a leaf of the path P_n . The generating function for the chromatic symmetric function of the twin $P_{n,v}$ of a path on n vertices satisfies the following identity:*

$$2 + 2e_1 z + \sum_{n \geq 1} X_{P_{n,v}} z^{n+1} = 2(1 - e_2 z^2) \mathcal{X}_P(z).$$

Proof. By [7, Equation 5], the chromatic symmetric function of $P_{n,v}$, with $n \geq 1$, is given by

$$(6) \quad X_{P_{n,v}} = 2X_{P_{n+1}} - X_{P_2}X_{P_{n-1}}.$$

The proof now follows by using the generating function $\mathcal{X}_P(z)$. \square

Now we are ready to derive a generating function for paths twinned at a leaf. Although the e -positivity was established in [10, Corollary 7.7] and again in [7], as mentioned earlier, our contribution here is to give the manifestly e -positive generating function below for $X_{P_{n,v}}$, using only symmetric functions, which enables a more efficient coefficient extraction.

Proposition 3.9. *Let $\mathcal{X}_{P_v}(z)$ be the generating function for the twinned path at a leaf, that is, $\mathcal{X}_{P_v} := \sum_{n \geq 1} X_{P_{n,v}} z^{n+1}$. Then,*

$$\frac{1}{2}\mathcal{X}_{P_v}(z) = K(z)\frac{G_{\geq 3}(z)}{D(z)} + e_1 z G(z)\frac{G_{\geq 3}(z)}{D(z)} + e_2 z^2 + \sum_{i \geq 3} i e_i z^i + e_1 z G_{\geq 3}(z).$$

In particular $X_{P_{n,v}}$ is e -positive, and the initial values are

$$X_{P_{1,v}} = 2e_2, \quad X_{P_{2,v}} = 2(3e_3), \quad X_{P_{3,v}} = 2(4e_4 + 2e_1e_3), \quad X_{P_{4,v}} = 2(4e_2e_3 + 3e_1e_4 + 5e_5).$$

Proof. By Lemma 3.7 and Corollary 3.4, we have that

$$\begin{aligned} \frac{1}{2}\mathcal{X}_{P_v}(z) &= (1 - e_2 z^2)\mathcal{X}_P - (1 + e_1 z) = K(z)\frac{1 - e_2 z^2}{D(z)} + e_1 z G(z)\frac{1 - e_2 z^2}{D(z)} - e_2 z^2(1 + e_1 z) \\ &= (K(z) + e_1 z G(z))\frac{G_{\geq 3}(z)}{D(z)} + (K(z) + e_1 z G(z)) - e_2 z^2(1 + e_1 z) \\ &= (K(z) + e_1 z G(z))\frac{G_{\geq 3}(z)}{D(z)} + (K(z) - e_2 z^2) + e_1 z(G(z) - e_2 z^2). \end{aligned}$$

Since $K(z) - e_2 z^2 = e_2 z^2 + \sum_{i \geq 3} i e_i z^i$ and $G(z) - e_2 z^2 = \sum_{i \geq 3} (i-1) e_i z^i$, the result follows. \square

Corollary 3.10. *Let c_λ be the coefficient of $e_\lambda z^{|\lambda|}$ in \mathcal{X}_{P_v} , that is $\mathcal{X}_{P_v} = \sum c_\lambda e_\lambda z^{|\lambda|}$, where v is a leaf of the path P_n . The following is a list of closed formulas for all the coefficients c_λ involved in the general expression of $\mathcal{X}_{P_v}(z)$:*

(a) $c_{(k)} = 2k$, $k \geq 3$, and $c_{(2)} = 2$;

(b) $c_{(k-1,1)} = 2(k-2)$, $k \geq 4$;

(c) $c_{(k-2,2)} = 4(k-3)$, $k \geq 5$;

(d) $c_{(i,j)} = 4i(j-1) + 4j(i-1) = 4(2ij - i - j)$, $i, j \geq 3$, $i \neq j$;

(e) $c_{(i,i)} = 4i(i-1)$, $i \geq 3$.

(f) $c_{(3,2^k)} = 8$ and $c_{(3,2^k,1)} = 4$, $k \geq 2$.

(g) Assume $c_{1 \cup \mu} \neq 0$ and $\ell(\mu) \geq 2$. Then $1 \notin \text{supp}(\mu)$ and there exists $a \geq 3$ such that $a \in \text{supp}(\mu)$. In particular $c_{(2^k,1)} = 0$. The coefficient of $e_{1 \cup \mu}$ is the coefficient of e_μ in $G(z)G_{\geq 3}(z)G(z)^{\ell(\mu)-2}$, and it equals

$$\sum_{\substack{(a,b) \\ a,b \in \text{supp}(\mu) \\ a \geq 2, b \geq 3}} (a-1)(b-1)\varepsilon((\mu-a)-b).$$

(h) Assume $1 \notin \text{supp}(\lambda)$ and $\ell(\lambda) \geq 2$. If $c_\lambda \neq 0$, then λ contains at least one part of size at least 3. In particular $c_{(2^k)} = 0$. The coefficient c_λ is the coefficient of e_λ in $K(z)G_{\geq 3}(z)G(z)^{\ell(\lambda)-2}$, and it equals

$$\sum_{\substack{(a,b) \\ a,b \in \text{supp}(\lambda) \\ a \geq 2, b \geq 3}} a(b-1)\varepsilon((\lambda-a)-b).$$

Note that cases (c)-(f) are particular cases of (g) and (h).

3.2.2. Paths twinned at both leaves.

In this section, we consider the twinned path $P_{n,w,v}$ at both leaves, which we label with w and v . The e -positivity of its chromatic symmetric function is a consequence of [10, Corollary 7.7], whose proof relies on the theory of symmetric functions in noncommuting variables. Here we derive an e -positive generating function using only symmetric function identities.

Unlike the other families of graphs, here one needs to pay special attention to the smaller values of n . We consider the special case of the path on two vertices first. Twinning both vertices produces the twin of the cycle graph C_3 at one vertex, which is also the complete graph K_4 , as shown in Figure 4, and therefore we have the following.

Lemma 3.11. *For the path P_2 twinned at both vertices, $X_{P_{2,v,w}} = X_{C_{3,v}} = 24e_4$.*




FIGURE 4

For the general case, we start with a consequence of the triple deletion argument.

Corollary 3.12. *Let v, w be the two leaves of the path P_n , and let $P_{n,v,w}$ be the path twinned at both leaves. Then, for $n \geq 3$,*

$$(7) \quad X_{P_{n,v,w}} = 2X_{P_{n+1,v}} - 2e_2X_{P_{n-1,v}} = 4(X_{P_{n+2}} - 2e_2X_{P_n} + e_2^2X_{P_{n-2}}).$$

This relation allows us to give the following generating function identity.

Proposition 3.13. *For the graph $P_{n,v,w}$, we have*

$$(8) \quad \frac{1}{4} \sum_{n \geq 3} X_{P_{n,v,w}} z^{n+2} = (1 - e_2 z^2) \frac{1}{2} \mathcal{X}_{P_v} + \frac{1}{2} \alpha,$$

where $\alpha = 2e_2^2 z^4 - (8e_4 z^4 + 4e_3 e_1 z^4 + 6e_3 z^3 + 2e_2 z^2)$.

Proof. Multiply both sides of the first equality in Corollary 3.12 by z^{n+2} and sum for $n \geq 3$:

$$\begin{aligned}
\sum_{n \geq 3} X_{P_{n,v,w}} z^{n+2} &= 2 \sum_{n \geq 3} X_{P_{n+1,v}} z^{n+2} - 2e_2 \sum_{n \geq 3} X_{P_{n-1,v}} z^{n+2} \\
&= 2 \sum_{n \geq 4} X_{P_{n,v}} z^{n+1} - 2e_2 z^2 \sum_{n \geq 2} X_{P_{n,v}} z^{n+1} \\
&= 2(1 - e_2 z^2) \mathcal{X}_{P_v} - 2(X_{P_{3,v}} z^4 + X_{P_{2,v}} z^3 + X_{P_{1,v}} z^2 - 2e_2 z^2 X_{P_{1,v}} z^2) \\
&= 2(1 - e_2 z^2) \mathcal{X}_{P_v} - 2[(8e_4 + 4e_3 e_1) z^4 + 6e_3 z^3 + 2e_2 z^2 - 2e_2^2 z^4]
\end{aligned}$$

where the computations for $X_{P_{n,v}}$ follow from Proposition 3.9. \square

The next theorem follows from Corollary 3.12 and manipulation of the formal series.

Theorem 3.14. *The generating function $\frac{1}{4} \sum_{n \geq 3} X_{P_{n,v,w}} z^{n+2}$ has the following e -positive expansion:*

$$\begin{aligned}
\frac{1}{4} \sum_{n \geq 3} X_{P_{n,v,w}} z^{n+2} &= (K(z) + e_1 z G(z)) \frac{G_{\geq 3}(z)^2}{D(z)} + e_1 z G_{\geq 3}(z)^2 \\
&\quad + \left(G_{\geq 3}(z) \sum_{i \geq 3} i e_i z^i + e_1 z \sum_{i \geq 4} (i-1) e_i z^i + e_2 z^2 \sum_{i \geq 3} (i-2) e_i z^i \right) + \sum_{i \geq 5} i e_i z^i.
\end{aligned}$$

Proof. We use the generating function in Proposition 3.9 to expand $\frac{1}{2}(1 - e_2 z^2) \mathcal{X}_{P_v}$ as

$$\begin{aligned}
\frac{1}{2}(1 - e_2 z^2) \mathcal{X}_{P_v} &= (1 - e_2 z^2) (K(z) + z e_1 G(z)) \frac{G_{\geq 3}(z)}{D(z)} \\
&\quad + (1 - e_2 z^2) \left(e_2 z^2 + \sum_{i \geq 3} i e_i z^i + z e_1 G_{\geq 3}(z) \right).
\end{aligned}$$

By (5), $\frac{1 - e_2 z^2}{D(z)} = 1 + \frac{G_{\geq 3}(z)}{D(z)}$, and we can rewrite the above expression as

$$\begin{aligned}
\frac{1}{2}(1 - e_2 z^2) \mathcal{X}_{P_v} &= (K(z) + z e_1 G(z)) G_{\geq 3}(z) \left(1 + \frac{G_{\geq 3}(z)}{D(z)} \right) \\
&\quad + (1 - e_2 z^2) \left(e_2 z^2 + \sum_{i \geq 3} i e_i z^i + z e_1 G_{\geq 3}(z) \right).
\end{aligned}$$

Next, we arrange the expression above so that the term $-\frac{1}{2}\alpha$ appears:

$$\begin{aligned}
\frac{1}{2}(1 - e_2 z^2) \mathcal{X}_{P_v} &= (K(z) + z e_1 G(z)) \frac{G_{\geq 3}(z)^2}{D(z)} - \frac{1}{2}\alpha + \sum_{i \geq 5} i e_i z^i + e_1 z \sum_{i \geq 4} (i-1) e_i z^i \\
&\quad + (K(z) + z e_1 G(z)) G_{\geq 3}(z) - e_2 z^2 \sum_{i \geq 3} i e_i z^i - e_2 z^2 (z e_1) G_{\geq 3}(z).
\end{aligned}$$

Thus, we have that

$$\begin{aligned} \frac{1}{4} \sum_{n \geq 3} X_{P_{n,v,w}} z^{n+2} &= \frac{1}{2} (1 - e_2 z^2) \mathcal{X}_{P_v} + \frac{1}{2} \alpha \\ (9) \quad &= (K(z) + z e_1 G(z)) \frac{G_{\geq 3}(z)^2}{D(z)} + \sum_{i \geq 5} i e_i z^i + e_1 z \sum_{i \geq 4} (i-1) e_i z^i \\ (10) \quad &+ (K(z) + z e_1 G(z)) G_{\geq 3}(z) - e_2 z^2 \sum_{i \geq 3} i e_i z^i - e_2 z^2 (z e_1) G_{\geq 3}(z), \end{aligned}$$

where the terms in line (9) are e -positive. Thus, we only need to show that the terms in line (10) are also e -positive. Note that

$$K(z) G_{\geq 3}(z) = \left(2e_2 z^2 + \sum_{i \geq 3} i e_i z^i \right) G_{\geq 3}(z) = 2e_2 z^2 G_{\geq 3}(z) + G_{\geq 3}(z) \sum_{i \geq 3} i e_i z^i.$$

Together with the fact that $G(z) - e_2 z^2 = G_{\geq 3}(z)$, line (10) can be written as

$$\begin{aligned} &(K(z) + z e_1 G(z)) G_{\geq 3}(z) - e_2 z^2 \sum_{i \geq 3} i e_i z^i - e_2 z^2 (z e_1) G_{\geq 3}(z) \\ &= G_{\geq 3}(z) \sum_{i \geq 3} i e_i z^i + e_2 z^2 \left(2 \sum_{i \geq 3} (i-1) e_i z^i - \sum_{i \geq 3} i e_i z^i \right) + e_1 z (G(z) - e_2 z^2) G_{\geq 3}(z) \\ (11) \quad &= G_{\geq 3}(z) \sum_{i \geq 3} i e_i z^i + e_2 z^2 \sum_{i \geq 3} (i-2) e_i z^i + e_1 z G_{\geq 3}(z)^2. \end{aligned}$$

Since the expression in (11) is also e -positive, the result follows. \square

In particular, we can extract the following formulas for the coefficients.

Corollary 3.15. *Let c_λ be the coefficient of $e_\lambda z^{|\lambda|}$ in $\mathcal{X}_{P_{v,w}}$, that is, $\mathcal{X}_{P_{v,w}} = \sum c_\lambda e^\lambda z^{|\lambda|}$. We have the following list of closed formulas:*

- (a) For $k \geq 3$, $c_{(k+2)} = 4(k+2)$, $c_{(k,2)} = 4(k-2)$, and $c_{(k+1,1)} = 4k$.
- (b) For $i \geq 3$, $c_{(i,i)} = 4(i-1)i$, and for $i, j \geq 3$, $i \neq j$, $c_{(i,j)} = 4(j-1)i + 4(i-1)j$.
- (c) For $i, j \geq 3$, $i \neq j$, $c_{(i,i,1)} = 4(i-1)^2$, $c_{(i,j,1)} = 8(i-1)(j-1)$, $i, j \geq 3$, $i \neq j$, and zero otherwise.
- (d) If $c_{1 \cup \mu} \neq 0$, then $1 \notin \text{supp}(\mu)$.
- (e) For all $k \geq 0$, $c_{(3^2, 2^{k+1})} = 32$ and $c_{(3^2, 2^{k+1}, 1)} = 16$.

3.2.3. Paths twinned at an interior vertex.

In this section, we establish an e -positive generating function for the path $P_{n,\ell}$ twinned at an interior vertex ℓ , where we label the vertices of P_n by $1, 2, \dots, n$ from left to right. As stated in the introduction, the e -positivity can also be deduced from [10, Theorem 7.8].

As in the preceding section, we first derive a triple deletion formula for the chromatic symmetric function of $P_{n,\ell}$, (Proposition 3.19), and then deduce an e -positive generating function for its chromatic symmetric function (Theorem 3.24). We begin with some definitions.

Definition 3.16. For $n \geq 2$ and $1 \leq \ell \leq n - 1$, let $\tilde{T}_{n,\ell}$ (T for *triangle*) denote the graph obtained from the path graph P_n by adding a vertex adjacent to both ℓ and $\ell + 1$. For $n \geq 1$ and $1 \leq \ell \leq n$, let $F_{n,\ell}$ (F for *flagpole*) denote the graph obtained from P_n by adding a vertex adjacent to ℓ .

By the triple deletion argument illustrated in Figure 5, we have the following result.

Lemma 3.17. For $n \geq 3$ and $2 \leq \ell \leq n - 1$, we have

$$X_{P_{n,\ell}} = 2X_{\tilde{T}_{n,\ell-1}} - X_{\tilde{T}_{\ell,\ell-1}}X_{P_{n-\ell}}.$$




FIGURE 5. The triple deletion argument applied as in Lemma 3.17.

By carefully applying the triple deletion argument to various $P_{n,\ell}$, we can deal with the triangles $\tilde{T}_{n,\ell}$ by “shifting” them around. Note that $\tilde{T}_{n,\ell}$ has a triangle, and so the triple deletion argument applies to two different sets of edges, to which we refer as *left* and *right shifts*. We illustrate them on the left-hand side and right-hand side of Figure 6, respectively.

Lemma 3.18 (Left and Right Shift Lemma). For $n \geq 3$ and $2 \leq \ell \leq n - 1$, we have

$$\begin{array}{c|c} X_{\tilde{T}_{n,\ell}} = X_{F_{n,\ell}} + X_{P_{n+1}} - X_{P_{\ell+1}}X_{P_{n-\ell}} & X_{\tilde{T}_{n,\ell}} = X_{F_{n,\ell+1}} + X_{P_{n+1}} - X_{P_{\ell}}X_{P_{n-\ell+1}} \\ \text{Left shift} & \text{Right shift} \end{array}$$

Our next step is to use Lemma 3.18 to obtain a formula equivalent to that in Lemma 3.17 which does not involve twinning paths.

Proposition 3.19. For $n \geq 3$ and $2 \leq \ell \leq n - 1$, we have

$$X_{P_{n,\ell}} = -2X_{P_{\ell-1}}X_{P_{n-\ell+2}} + 2e_1X_{P_n} + 4X_{P_{n+1}} - 2X_{P_{\ell}}X_{P_{n-\ell+1}} + 2e_2X_{P_{\ell-1}}X_{P_{n-\ell}} - 2X_{P_{\ell+1}}X_{P_{n-\ell}}.$$

Proof. Applying the Left and Right Shift Lemmas at $\ell - k - 1$ implies that

$$(12) \quad X_{F_{n,\ell-k}} = X_{F_{n,\ell-k-1}} + X_{P_{\ell-k-1}}X_{P_{n-\ell+k+2}} - X_{P_{\ell-k}}X_{P_{n-\ell+k+1}}.$$

By applying (12) repeatedly, we get that

$$(13) \quad X_{F_{n,\ell}} = X_{F_{n,1}} + \sum_{k=0}^{\ell-2} (X_{P_{\ell-k-1}}X_{P_{n-\ell+k+2}} - X_{P_{\ell-k}}X_{P_{n-\ell+k+1}}).$$




FIGURE 6. Illustration of Lemma 3.18.

Since $F_{n,1}$ is precisely P_{n+1} , we can telescope the sum in (13) to obtain

$$(14) \quad X_{F_{n,\ell}} = X_{P_{n+1}} + X_{P_1}X_{P_n} - X_{P_\ell}X_{P_{n-\ell+1}}.$$

Recall the formula in Lemma 3.17:

$$X_{P_{n,\ell}} = 2X_{\tilde{T}_{n,\ell-1}} - X_{\tilde{T}_{\ell,\ell-1}}X_{P_{n-\ell}}.$$

We apply the Left Shift Lemma to $X_{\tilde{T}_{n,\ell-1}}$ and the Right Shift Lemma to $X_{\tilde{T}_{\ell,\ell-1}}$, and obtain

$$X_{P_{n,\ell}} = 2 \left(X_{F_{n,\ell-1}} + X_{P_{n+1}} - X_{P_\ell}X_{P_{n-\ell+1}} \right) - \left(X_{F_{\ell,\ell}} + X_{P_{\ell+1}} - X_{P_{\ell-1}}X_{P_2} \right) X_{P_{n-\ell}}.$$

Since $F_{\ell,\ell}$ is $P_{\ell+1}$, we can rewrite the last equation as:

$$\begin{aligned} X_{P_{n,\ell}} &= 2 \left(X_{F_{n,\ell-1}} + X_{P_{n+1}} - X_{P_\ell}X_{P_{n-\ell+1}} \right) - (2X_{P_{\ell+1}} - X_{P_{\ell-1}}X_{P_2}) X_{P_{n-\ell}} \\ &= 2X_{F_{n,\ell-1}} + [2X_{P_{n+1}} - 2X_{P_\ell}X_{P_{n-\ell+1}} + X_{P_{\ell-1}}X_{P_2}X_{P_{n-\ell}} - 2X_{P_{\ell+1}}X_{P_{n-\ell}}]. \end{aligned}$$

Substituting in (14) for $X_{F_{n,\ell-1}}$, we have:

$$\begin{aligned} X_{P_{n,\ell}} &= 2 \left(X_{P_{n+1}} + X_{P_1}X_{P_n} - X_{P_{\ell-1}}X_{P_{n-\ell}} \right) \\ &\quad + [2X_{P_{n+1}} - 2X_{P_\ell}X_{P_{n-\ell+1}} + X_{P_{\ell-1}}X_{P_2}X_{P_{n-\ell}} - 2X_{P_{\ell+1}}X_{P_{n-\ell}}]. \end{aligned}$$

Finally, the formula in the statement follows by collecting all the terms and evaluating $X_{P_1} = e_1$ and $X_{P_2} = 2e_2$. \square

Now we investigate the generating function of $X_{P_{n,\ell}}$. For this, we introduce two families of polynomials in the variable z with coefficients in the ring of symmetric functions.

Definition 3.20. (a) For $\ell \geq 1$, we define the following polynomial of degree $\ell + 1$:

$$f_\ell(z) := 2 + e_1z - X_{P_{\ell-1}}z^{\ell-1}(1 - e_2z^2) - X_{P_\ell}z^\ell - X_{P_{\ell+1}}z^{\ell+1}.$$

(b) For $\ell \geq 2$, we define the following polynomial of degree ℓ :

$$g_\ell(z) := - \sum_{j=0}^{\ell} X_{P_j} z^j - (1 + e_1 z) \sum_{j=0}^{\ell-2} X_{P_j} z^j.$$

The following result gives an identity for the generating function for the chromatic symmetric function of the twinned path in terms of the generating function for the chromatic symmetric function of the path and the new families of polynomials introduced.

Proposition 3.21. *Let $2 \leq \ell \leq n-1$. The generating function for the chromatic symmetric function of the twinned path $P_{n,\ell}$, twinned at vertex ℓ , can be written in terms of the path generating function \mathcal{X}_P as follows:*

$$(15) \quad \sum_{n \geq \ell+1} X_{P_{n,\ell}} z^{n+1} = 2\mathcal{X}_P(z) f_\ell(z) + 2g_\ell(z).$$

Proof. We reorder the terms appearing in the recurrence in Proposition 3.19 to make the source of the factor $f_\ell(z)$ accompanying \mathcal{X}_P more transparent:

$$(16) \quad X_{P_{n,\ell}} = 4X_{P_{n+1}} + 2e_1 X_{P_n} - 2X_{P_{\ell-1}}(X_{P_{n-\ell+2}} - e_2 X_{P_{n-\ell}}) - 2X_{P_\ell} X_{P_{n-\ell+1}} - 2X_{P_{\ell+1}} X_{P_{n-\ell}}.$$

Multiplying by z^{n+1} and summing over $n \geq \ell+1$ gives

$$\begin{aligned} \sum_{n \geq \ell+1} X_{P_{n,\ell}} z^{n+1} &= 2[2 + e_1 z - X_{P_{\ell-1}} z^{\ell-1} (1 - e_2 z^2) - X_{P_\ell} z^\ell - z^{\ell+1} X_{P_{\ell+1}}] \mathcal{X}_P \\ &\quad - 4 \sum_{j=0}^{\ell} X_{P_j} z^j - 2ze_1 \sum_{j=0}^{\ell-1} X_{P_j} z^j + 2z^{\ell-1} (1 + ze_1) X_{P_{\ell-1}} + 2X_{P_\ell} z^\ell. \end{aligned}$$

The expression for $f_\ell(z)$ follows immediately from the first line above.

Now rewrite the second line as

$$\begin{aligned} -4X_{P_\ell} z^\ell - 4X_{P_{\ell-1}} z^{\ell-1} - 4 \sum_{j=0}^{\ell-2} X_{P_j} z^j - 2z^\ell e_1 X_{P_{\ell-1}} - 2ze_1 \sum_{j=0}^{\ell-2} X_{P_j} z^j \\ + 2X_{P_{\ell-1}} z^{\ell-1} + 2e_1 X_{P_{\ell-1}} z^\ell + 2X_{P_\ell} z^\ell, \end{aligned}$$

which in turn equals the stated expression for $g_\ell(z)$. \square

Although $f_\ell(z)$ is not e -positive, we can conclude the following.

Corollary 3.22. *The e -positivity of $X_{P_{n,\ell}}$ is equivalent to the e -positivity of $\mathcal{X}_P(z) f_\ell(z)$.*

Proof. The degree of $g_\ell(z)$ as a polynomial in z is ℓ , while the left-hand side of (15) has lowest degree $\ell+1$ in z . We conclude that all terms in $g_\ell(z)$ are necessarily canceled out by identical terms in $\mathcal{X}_P(z) f_\ell(z)$. \square

Our next result rewrites $f_\ell(z)$ as a positive expansion of other functions that were introduced in (4).

Lemma 3.23. *For $\ell \geq 2$, we have*

$$(17) \quad f_\ell(z) = \sum_{i=3}^{\ell+1} (i-2)e_i z^i + 2(D + G_{\geq \ell+2}) + \sum_{i=1}^{\ell-2} (D + G_{\geq \ell+2-i}) X_{P_i} z^i.$$

Proof. By Definition 3.20,

$$f_\ell(z) = 2 + e_1z + X_{P_{\ell-1}}e_2z^{\ell+1} - X_{P_{\ell-1}}z^{\ell-1} - X_{P_\ell}z^\ell - X_{P_{\ell+1}}z^{\ell+1}.$$

For $\ell = 2$, recall that $X_{P_3} = e_2e_1 + 3e_3$, and $X_{P_1} = e_1$ and $X_{P_2} = 2e_2$. Then we have

$$\begin{aligned} f_2(z) &= 2 - 2e_2z^2 - 3e_3z^3 = 2(1 - e_2z^2 - 2e_3z^3) + e_3z^3 \\ &= 2(1 - G(z) + G_{\geq 4}(z)) + e_3z^3 = 2(D(z) + G_{\geq 4}(z)) + e_3z^3. \end{aligned}$$

Let φ_ℓ denote the right-hand side of (17). We show that φ_ℓ and f_ℓ satisfy the same recurrence relation. It is straightforward to see that for $\ell \geq 2$,

$$(18) \quad f_{\ell+1} - f_\ell = X_{P_\ell}e_2z^{\ell+2} - X_{P_{\ell+2}}z^{\ell+2} - X_{P_{\ell-1}}e_2z^{\ell+1} + X_{P_{\ell-1}}z^{\ell-1}.$$

Next we look at $\varphi_{\ell+1} - \varphi_\ell$. Observe from (4) that $G_{\geq m+1} - G_{\geq m} = -(m-1)e_mz^m$. We therefore have

$$\begin{aligned} \varphi_{\ell+1} &= \sum_{i=3}^{\ell+2} (i-2)e_iz^i + 2(D + G_{\geq \ell+3}) + \sum_{i=1}^{\ell-1} (D + G_{\ell+3-i})X_{P_i}z^i \\ \varphi_\ell &= \sum_{i=3}^{\ell+1} (i-2)e_iz^i + 2(D + G_{\geq \ell+2}) + \sum_{i=1}^{\ell-2} (D + G_{\ell+2-i})X_{P_i}z^i \end{aligned}$$

and hence we obtain, for $\ell \geq 2$,

$$\varphi_{\ell+1} - \varphi_\ell = \ell e_{\ell+2}z^{\ell+2} - 2(\ell+1)e_{\ell+2}z^{\ell+2} - \sum_{i=1}^{\ell-2} (\ell+1-i)e_{\ell+2-i}z^{\ell+2-i}X_{P_i}z^i + (D + G_{\geq 4})X_{P_{\ell-1}}z^{\ell-1}.$$

The path recurrence relation in Proposition 3.2(a) tells us that

$$X_{P_{\ell+2}} = (\ell+2)e_{\ell+2} + \sum_{j=1}^{\ell-2} (\ell+1-j)e_{\ell+2-j}X_{P_j} + 2e_3X_{P_{\ell-1}} + e_2X_{P_\ell}.$$

Together with $D + G_{\geq 4} = 1 - G_{\leq 3} = 1 - e_2z^2 - 2e_3z^3$, this gives

$$\varphi_{\ell+1} - \varphi_\ell = (-X_{P_{\ell+2}} + 2e_3X_{P_{\ell-1}} + e_2X_{P_\ell})z^{\ell+2} + (1 - e_2z^2 - 2e_3z^3)X_{P_{\ell-1}}z^{\ell-1}.$$

The terms containing $e_3X_{P_{\ell-1}}$ cancel, and the remaining expression coincides with the one for $f_{\ell+1} - f_\ell$ in (18). Hence φ_ℓ and f_ℓ satisfy the same recurrence relation. Since their initial values also coincide, the claim follows. \square

The preceding efforts culminate in the following e -positivity result, as announced at the start of this section.

Theorem 3.24. *For $\ell \geq 2$, we have the e -positive expansion*

$$(19) \quad \mathcal{X}_P f_\ell = \mathcal{X}_P \sum_{i=3}^{\ell+1} (i-2)e_iz^i + 2(E + \mathcal{X}_P G_{\geq \ell+2}) + \sum_{i=1}^{\ell-2} (E + \mathcal{X}_P G_{\geq \ell+2-i})X_{P_i}z^i.$$

Hence the generating function $\sum_{n \geq \ell+1} X_{P_n, \ell} z^{n+1}$ is e -positive.

Proof. The expression for f_ℓ in Lemma 3.23 immediately allows us to conclude (19), using the fact that $\mathcal{X}_P(z)D(z) = E(z)$. It then suffices to observe that the generating function

$\sum_{n \geq \ell+1} X_{P_{n,\ell}} z^{n+1}$ is comprised precisely of all the terms of degree $\geq \ell + 1$ in the e -positive rational expression

$$\begin{aligned} 2\mathcal{X}_P f_\ell &= \frac{2E f_\ell}{D} \\ &= \frac{2E \left(\sum_{i=3}^{\ell+1} (i-2)e_i z^i + G_{\geq \ell+2} + \sum_{i=0}^{\ell-2} G_{\geq \ell+2-i} X_{P_i} z^i \right)}{1 - \sum_{i \geq 2} (i-1)e_i z^i} + 2(1+E) \sum_{i=0}^{\ell-2} X_{P_i} z^i. \quad \square \end{aligned}$$

From Theorem 3.24 and a tedious computation of $\mathcal{X}_P f_\ell + g_\ell$, we obtain the following cancellation-free e -positive expression for $\frac{1}{2} \sum_{n \geq \ell+1} X_{P_{n,\ell}} z^{n+1}$.

Proposition 3.25. *For integers $n \geq 3$ and $2 \leq \ell \leq n-1$, the twin $P_{n,\ell}$ of the path P_n at the degree 2 vertex ℓ is e -positive. In particular, we have*

$$\begin{aligned} \frac{1}{2} \sum_{n \geq \ell+1} X_{P_{n,\ell}} z^{n+1} &= \ell e_{\ell+1} z^{\ell+1} \left(\sum_{i=0}^{\ell-2} X_{P_i} z^i \right) + \sum_{i=3}^{\ell} (i-1) e_i z^i \left(\sum_{j=0}^{i-3} X_{P_{\ell-2-j}} z^{\ell-2-j} \right) + E_{\geq \ell+1} \\ &\quad + E_{\geq \ell+2} \sum_{i=0}^{\ell-2} X_{P_i} z^i + \left(\sum_{i \geq \ell-1} X_{P_i} z^i \right) \left(\sum_{i=2}^{\ell+1} (i-2) e_i z^i \right) \\ &\quad + 2\mathcal{X}_P G_{\geq \ell+2} + \mathcal{X}_P \sum_{i=1}^{\ell-2} G_{\geq \ell+2-i} X_{P_i} z^i. \end{aligned}$$

3.3. Generating function for twinned cycles. In this section, we establish a new result, the e -positivity of the chromatic symmetric function of the twinned cycle. Again, our goal of obtaining an e -positive generating function for the twinned cycle will begin with a formula for the chromatic symmetric function of the twinned cycle, which is derived using the triple deletion formula.

We start by introducing two more families of graphs. Consider the twinned cycle graph $C_{n,v}$ where v' is the twinned vertex of v and u and w are the adjacent vertices to v and v' . Let D_{n+1} be the graph obtained from $C_{n,v}$ by removing the edge uv and let Tad_{n+1} be the graph obtained from $C_{n,v}$ by removing the edges uv and vv' . We illustrate these two definitions in Figure 7.




FIGURE 7

Lemma 3.26. *For $n \geq 3$:*

$$X_{C_{n,v}} = 4X_{C_{n+1}} + 2e_1X_{C_n} - 6X_{P_{n+1}} + 2e_2X_{P_{n-1}}.$$

Proof. Consider $n \geq 3$. By the triple deletion argument applied to $\epsilon_1 = uv$ and $\epsilon_2 = uv'$, we get that

$$(20) \quad X_{C_{n,v}} = 2X_{D_{n+1}} - X_{P_{n,v}} = 2X_{D_{n+1}} - 2X_{P_{n+1}} + X_{P_2}X_{P_{n-1}}.$$

In D_{n+1} , applying the triple deletion argument to $\epsilon_1 = vw$ and $\epsilon_2 = vv'$ gives

$$(21) \quad X_{D_{n+1}} = 2X_{\text{Tad}_{n+1}} - e_1X_{C_n},$$

while applying the triple deletion argument to $\epsilon_1 = vw$ and $\epsilon_2 = v'w$ gives

$$X_{D_{n+1}} = X_{\text{Tad}_{n+1}} + X_{C_{n+1}} - X_{P_{n+1}}.$$

Subtracting both expressions for $X_{D_{n+1}}$ we obtain that

$$X_{\text{Tad}_{n+1}} = X_{C_{n+1}} + e_1X_{C_n} - X_{P_{n+1}},$$

and therefore

$$(22) \quad X_{D_{n+1}} = 2X_{C_{n+1}} + e_1X_{C_n} - 2X_{P_{n+1}}.$$

Finally, putting together (20) and (22), we have

$$X_{C_{n,v}} = 4X_{C_{n+1}} + 2e_1X_{C_n} - 6X_{P_{n+1}} + 2e_2X_{P_{n-1}},$$

as claimed. \square

Let \mathcal{X}_{C_v} be the generating function for the twinned cycle, that is, $\mathcal{X}_{C_v}(z) := \sum_{n \geq 3} X_{C_{n,v}} z^{n+1}$. By Lemma 3.26 we have the following expression for \mathcal{X}_{C_v} .

Corollary 3.27. *The generating function of the twinned cycle can be written as*

$$\mathcal{X}_{C_v}(z) = 2(2 + e_1z)\mathcal{X}_C - 2(3 - e_2z^2)\mathcal{X}_P + 6(1 + e_1z) + 2e_2z^2 - 6e_3z^3.$$

Proof. The generating function follows by multiplying the formula in Lemma 3.26 by z^{n+1} and summing over all $n \geq 3$. In particular, taking into account the initial terms that do not appear and using the initial values $X_{C_1} = 0, X_{P_1} = e_1, X_{C_2} = 2e_2 = X_{P_2}, X_{C_3} = 6e_3$, and $X_{P_3} = e_2e_1 + 3e_3$, we get the following expressions in terms of the generating functions for the cycle and the path:

$$(a) \quad 4 \sum_{n \geq 3} X_{C_{n+1}} z^{n+1} = 4(\mathcal{X}_C - z^2X_{C_2} - z^3X_{C_3}),$$

$$(b) \quad 2(e_1z) \sum_{n \geq 3} X_{C_n} z^n = 2e_1z(\mathcal{X}_C - z^2X_{C_2}),$$

$$(c) \quad 6 \sum_{n \geq 3} X_{P_{n+1}} z^{n+1} = 6(\mathcal{X}_P - 1 - zX_{P_1} - z^2X_{P_2} - z^3X_{P_3}), \text{ and}$$

$$(d) \quad 2e_2z^2 \sum_{n \geq 3} X_{P_{n-1}} z^{n-1} = 2e_2z^2(\mathcal{X}_P - 1 - zX_{P_1}).$$

Putting all this together gives the generating function as stated. \square

Consider the following e -positive generating functions that appear in the proof of Lemma 3.3:

$$F_2 = \sum_{i \geq 3} (2i^2 - 5i)e_i z^i \quad \text{and} \quad F_3 = \sum_{i \geq 4} [(i-1)(i-3)]e_i z^i.$$

Our goal is to show that the expression given in Corollary 3.27 is indeed e -positive.

Lemma 3.28. *The twinned cycle generating function, scaled by $\frac{1}{2}$, can be written as*

$$\frac{1}{2}\mathcal{X}_{C_v} = \frac{1}{D(z)}[F_2 + e_1zF_3 + e_2z^2(E - 1 - e_1z)] - \frac{e_2z^2}{D(z)} + e_2z^2 - 3e_3z^3.$$

Proof. By Corollary 3.27,

$$(23) \quad \frac{1}{2}\mathcal{X}_{C_v} = (2 + e_1z)\mathcal{X}_C - 3(\mathcal{X}_P - 1 - e_1z) + e_2z^2(\mathcal{X}_P + 1) - 3e_3z^3.$$

From the proof of Lemma 3.3, we have the following

$$\mathcal{X}_P - (1 + e_1z) = \frac{z(1 + e_1z)E'(z) - e_1zE(z)}{D(z)} \quad \text{and} \quad \mathcal{X}_C = \frac{z^2E''(z)}{D(z)},$$

and by definition $\mathcal{X}_P(z) = \frac{E(z)}{D(z)}$ and $\mathcal{X}_C(z) = \frac{z^2E''(z)}{D(z)}$. Substituting these into (23), we get

$$\begin{aligned} \frac{1}{2}\mathcal{X}_{C_v} &= \frac{1}{D(z)}[(2 + e_1z)z^2E''(z) - 3(zE'(z) + z^2e_1E'(z) - e_1zE(z)) + e_2z^2E(z)] + e_2z^2 - 3e_3z^3 \\ &= \frac{1}{D(z)}[(2z^2E'' - 3zE') + e_1z(z^2E'' - 3e_1zE' + 3E) + e_2z^2E(z)] + e_2z^2 - 3e_3z^3 \\ &= \frac{1}{D(z)}[F_2 - 3e_1z - 2e_2z^2 + e_1z(F_3 - e_2z^2 + 3) + e_2z^2E(z)] + e_2z^2 - 3e_3z^3, \end{aligned}$$

where the final equality comes from Lemma 3.3. The statement then follows after further algebraic manipulations. \square

Theorem 3.29. *The generating function for $\frac{1}{2}X_{C_{n,v}}$ has the following e -positive rational expression:*

$$\frac{1}{2} \sum_{n \geq 3} X_{C_{n,v}} z^{n+1} = \sum_{i \geq 4} (2i^2 - 5i)e_i z^i + \frac{e_1zF_3 + F_2 \sum_{i \geq 3} (i-1)e_i z^i + e_2z^2 \sum_{i \geq 3} (2i^2 - 6i + 2)e_i z^i}{1 - \sum_{i \geq 2} (i-1)e_i z^i}.$$

Proof. Let $E_{\geq 2} := E - 1 - e_1z = \sum_{i \geq 2} e_i z^i$. Then by Lemma 3.28, we have

$$(24) \quad \begin{aligned} \frac{1}{2}\mathcal{X}_{C_v} &= \frac{1}{D(z)}(F_2 + e_1zF_3 + e_2z^2E_{\geq 2}) - \frac{e_2z^2}{D(z)} + e_2z^2 - 3e_3z^3 \\ &= F_2 - 3e_3z^3 + \frac{e_1zF_3}{D(z)} + \left[F_2 \left(\frac{1}{D(z)} - 1 \right) + \frac{e_2z^2E_{\geq 2}}{D(z)} - \frac{e_2z^2}{D(z)} + e_2z^2 \right]. \end{aligned}$$

The first two terms in (24) can be written as:

$$(25) \quad F_2 - 3e_3z^3 = \sum_{i \geq 4} (2i^2 - 5i)e_i z^i.$$

For the generating function in the brackets of (24), we have

$$\begin{aligned} &F_2 \left(\frac{1}{D(z)} - 1 \right) + \frac{e_2z^2E_{\geq 2}}{D(z)} - \frac{e_2z^2}{D(z)} + e_2z^2 \\ &= F_2 \sum_{k \geq 1} G^k + e_2z^2E_{\geq 2} \sum_{k \geq 0} G^k - e_2z^2 \sum_{k \geq 1} G^k \\ &= F_2 \sum_{k \geq 1} G^k + e_2z^2E_{\geq 2} \sum_{k \geq 1} G^{k-1} - e_2z^2 \sum_{k \geq 1} G^k \end{aligned}$$

$$\begin{aligned}
&= \sum_{k \geq 1} G^{k-1} (GF_2 - e_2 z^2 (G - E_{\geq 2})) \\
&= \sum_{k \geq 1} G^{k-1} \left(GF_2 - e_2 z^2 F_2 + e_2 z^2 F_2 - e_2 z^2 \sum_{i \geq 3} (i-2) e_i z^i \right) \\
&= \sum_{k \geq 1} G^{k-1} \left(F_2 G_{\geq 3} + e_2 z^2 \sum_{i \geq 3} (2i^2 - 6i + 2) e_i z^i \right) \\
(26) \quad &= \frac{F_2 G_{\geq 3} + e_2 z^2 \sum_{i \geq 3} (2i^2 - 6i + 2) e_i z^i}{D(z)},
\end{aligned}$$

where the penultimate equality follows from the definitions of $G_{\geq 3}$ and F_2 . Combining (24), (25), and (26), we obtain the desired expression. \square

From the generating function in Theorem 3.29, we can readily extract the e -coefficients of $X_{C_{n,v}}$.

Corollary 3.30. *Let λ be a partition of k , $\lambda = \langle 1^{m_1}, 2^{m_2}, \dots, k^{m_k} \rangle$, and let c_λ be the coefficient of $e_\lambda z^{|\lambda|}$ in $\frac{1}{2} \mathcal{X}_{C_v}$. We have the following list of expressions for the coefficients:*

- (a) $c_{(k)} = k(2k - 5)$, $k \geq 3$.
- (b) If $m_1 > 1$, then $c_\lambda = 0$.
- (c) If $m_1 = 1$ and $\lambda = \mu \cup 1$ (so that $m_1(\mu) = 0$), then

$$c_\lambda = \sum_{\substack{i \geq 4 \\ i \in \text{supp}(\mu)}} (i-1)(i-3) \varepsilon(\mu - i).$$

Note that this is 0 unless λ has a part of size at least 4.

- (d) If $m_1 = m_2 = 0$, then

$$c_\lambda = \sum_{a \in \text{supp}(\lambda)} (2a^2 - 5a) \varepsilon(\lambda - a).$$

- (e) If $m_1 = 0$ and $m_2 = \ell(\lambda)$, then $c_\lambda = 0$.
- (f) If $m_1 = 0$ and $1 \leq m_2 < \ell(\lambda)$, then

$$c_\lambda = \sum_{\substack{a, b \geq 3 \\ (a, b) \in \text{supp}(\lambda)}} \varepsilon(\lambda - a - b) (2a^2 - 5a)(b-1) + \sum_{\substack{c \geq 3 \\ c \in \text{supp}(\lambda)}} \varepsilon(\lambda - c - 2) (2c^2 - 6c + 2)$$

where $(a, b) \in \text{supp}(\lambda)$ means both a and b are in $\text{supp}(\lambda)$ if $a \neq b$, and $m_a \geq 2$ if $a = b$.

4. e -POSITIVITY VIA RECURRENCES

In this section, we reprove several e -positivity results for certain classes of graphs by exhibiting an e -positive recurrence relation. The recurrence relations for paths and cycles from Proposition 3.2 serve as the model for those of this section, and in fact will play a key role in our derivations. We will also need explicit expressions for some coefficients, which are readily extracted from Proposition 3.2. These are recorded in the next result.

Corollary 4.1. *Given a graph G , let $[e_\lambda]X_G$ denote the coefficient of e_λ in the chromatic symmetric function of G , X_G .*

- For $n \geq 2$, $[e_n]X_{P_n} = n$, $[e_{n-1}e_1]X_{P_n} = n - 2$, and $[e_n]X_{C_n} = n(n - 1)$.
- For $n \geq 5$, $[e_{n-2}e_2]X_{P_n} = 3n - 8$ and $[e_{n-2}e_2]X_{C_n} = n(n - 3)$.
- For $k \geq 2$ and $r \geq 1$, $[(e_k)^r]X_{P_{kr}} = k(k - 1)^{r-1}$ and $[e_{(kr)}]X_{C_{kr}} = k(k - 1)^r$.
- $[e_2^2]X_{C_4} = 2$.

The rest of this section follows the structure of Section 3.

4.1. Recurrences for twinned paths. In this section we derive recurrence formulas for the chromatic symmetric function for a path twinned at one or both leaves or at an internal vertex.

4.1.1. Paths twinned at a leaf.

In the next proposition, we give formulas for the chromatic symmetric function $X_{P_{n,v}}$ of the path twinned at a leaf, one in terms of the path chromatic symmetric function, and the other a recurrence in the spirit of Proposition 3.2. The recurrence given below makes the e -positivity transparent.

Proposition 4.2. *Let v be a leaf of the path P_n . Then, for $n \geq 4$,*

$$X_{P_{n,v}} = 2(n + 1)e_{n+1} + 2 \sum_{j=3}^n (j - 1)e_j X_{P_{n+1-j}}.$$

Thus $X_{P_{n,v}}$ is e -positive. Moreover, for $n \geq 4$, $X_{P_{n,v}}$ satisfies the e -positive recurrence

$$X_{P_{n,v}} = \sum_{j=2}^{n-2} (j - 1)e_j X_{P_{n-j,v}} + 2(n + 1)e_{n+1} + 2(n - 1)e_n e_1 + 2(n - 3)e_{n-1}e_2,$$

with initial values $X_{P_{1,v}} = 2e_2$, $X_{P_{2,v}} = 6e_3$, and $X_{P_{3,v}} = 8e_4 + 4e_3e_1$.

Proof. The first expression follows from Proposition 3.2(a) and Proposition 3.8, noting that

$$\begin{aligned} X_{P_{n,v}} &= 2X_{P_{n+1}} - X_{P_2}X_{P_{n-1}} = 2(n + 1)e_{n+1} + 2 \sum_{j=2}^n (j - 1)e_j X_{P_{n+1-j}} - 2e_2 X_{P_{n-1}} \\ &= 2(n + 1)e_{n+1} + 2 \sum_{j=3}^n (j - 1)e_j X_{P_{n+1-j}} + 2e_2 X_{P_{n-1}} - 2e_2 X_{P_{n-1}}. \end{aligned}$$

For the second recurrence, we apply the triple deletion argument to $P_{n,v}$ followed by Proposition 3.2(a) to both terms. Thus,

$$\begin{aligned} X_{P_{n,v}} &= 2X_{P_{n+1}} - X_{P_2}X_{P_{n-1}} = 2X_{P_{n+1}} - 2e_2 X_{P_{n-1}} \\ &= 2(n + 1)e_{n+1} + 2 \sum_{j=2}^n (j - 1)e_j X_{P_{n+1-j}} - 2e_2 \left[(n - 1)e_{n-1} + \sum_{j=2}^{n-2} (j - 1)e_j X_{P_{n-1-j}} \right] \\ &= 2(n + 1)e_{n+1} - 2(n - 1)e_{n-1}e_2 + 2 \sum_{j=2}^{n-2} (j - 1)e_j [2X_{P_{n+1-j}} - 2e_2 X_{P_{n-1-j}}] \end{aligned}$$

$$\begin{aligned}
& + 2(n-2)e_{n-1}X_{P_2} + 2(n-1)e_nX_{P_1} \\
& = 2(n+1)e_{n+1} + 2 \sum_{j=2}^{n-2} (j-1)e_j X_{P_{n-j,v}} + 2(n-1)e_n e_1 + 2(n-3)e_{n-1}e_2,
\end{aligned}$$

where the final step follows by the triple deletion argument applied to $P_{n-j,v}$. As this is an e -positive recursion with e -positive initial conditions, by induction it follows that $X_{P_{n,v}}$ is e -positive for all n . \square

4.1.2. *Paths twinned at both leaves.* Our new contribution is the e -positive recurrence below.

Proposition 4.3. *For $n \geq 6$, the chromatic symmetric function $X_{P_{n,v,w}}$ for the path P_n twinned at both leaves v, w satisfies the recurrence*

$$\begin{aligned}
\frac{1}{4}X_{P_{n,v,w}} &= \frac{1}{4} \sum_{j=3}^{n-3} (j-1)e_j X_{P_{n-j,v,w}} \\
&+ (n+2)e_{n+2} + n e_{n+1}e_1 + 3(n-2)e_{n-1}e_3 + 2(n-3)e_{n-2}e_3e_1 + 4(n-3)e_{n-2}e_4 \\
&+ e_2 \left[\frac{1}{4}X_{P_{n-2,v,w}} - 2e_n - (n-4)e_{n-2}e_2 - (n-2)e_{n-1}e_1 \right],
\end{aligned}$$

with the initial conditions

$$\begin{aligned}
X_{P_{2,v,w}} &= 24e_4, & X_{P_{4,v,w}} &= 24e_3^2 + 8e_4e_2 + 16e_5e_1 + 24e_6, \\
X_{P_{3,v,w}} &= 4e_3e_2 + 12e_4e_1 + 20e_5, & X_{P_{5,v,w}} &= 16e_3e_3e_1 + 68e_4e_3 + 12e_5e_2 + 20e_6e_1 + 28e_7.
\end{aligned}$$

Moreover, despite the negative terms, the expression is e -positive.

Proof. By the triple deletion argument, we have that

$$X_{P_{n,v,w}} = 2X_{P_{n+1,v}} - 2e_2X_{P_{n-1,v}}.$$

Applying the triple deletion argument again to both twinned terms, we have that $X_{P_{n+1,v}} = 2X_{P_{n+2}} - 2e_2X_{P_n}$ and $X_{P_{n-1,v}} = 2X_{P_n} - 2e_2X_{P_{n-2}}$. Thus, for $n \geq 3$,

$$(27) \quad \frac{1}{4}X_{P_{n,v,w}} = X_{P_{n+2}} + e_2^2X_{P_{n-2}} - 2e_2X_{P_n}.$$

Now we prove the recurrence relation by strong induction on n . The initial conditions are checked directly. For $n \geq 6$, using repeatedly (27) and Proposition 3.2(a), we write

$$\begin{aligned}
\frac{1}{4}X_{P_{n,v,w}} &= X_{P_{n+2}} - 2e_2X_{P_n} + e_2^2X_{P_{n-2}} \\
&= (n+2)e_{n+2} + \sum_{j=2}^{n+1} (j-1)e_j X_{P_{n+2-j}} + (n-2)e_{n-2}e_2^2 + e_2^2 \sum_{j=2}^{n-3} (j-1)e_j X_{P_{n-2-j}} \\
&\quad - 2ne_n e_2 - 2e_2 \sum_{j=2}^{n-1} (j-1)e_j X_{P_{n-j}} \\
&= \frac{1}{4} \sum_{j=2}^{n-3} (j-1)e_j X_{P_{n-j,v,w}} \\
&\quad + (n+2)e_{n+2} + ne_{n+1}e_1 + 3(n-2)e_{n-1}e_3 + 2(n-3)e_{n-2}e_3e_1 + 4(n-3)e_{n-2}e_4 \\
&\quad - 2e_n e_2 - (n-4)e_{n-2}e_2^2 - (n-2)e_{n-1}e_2e_1.
\end{aligned}$$

To rearrange this into the announced form, we peel off the $j = 2$ term from the sum and group it with the negative terms:

$$\begin{aligned} \frac{1}{4}X_{P_{n,v,w}} &= \frac{1}{4} \sum_{j=3}^{n-3} (j-1)e_j X_{P_{n-j,v,w}} \\ &+ (n+2)e_{n+2} + ne_{n+1}e_1 + 3(n-2)e_{n-1}e_3 + 2(n-3)e_{n-2}e_3e_1 + 4(n-3)e_{n-2}e_4 \\ &+ e_2 \left[\frac{1}{4}X_{P_{n-2,v,w}} - 2e_n - (n-4)e_{n-2}e_2 - (n-2)e_{n-1}e_1 \right]. \end{aligned}$$

Next we prove that the term within brackets

$$(28) \quad \left[\frac{1}{4}X_{P_{n-2,v,w}} - 2e_n - (n-4)e_{n-2}e_2 - (n-2)e_{n-1}e_1 \right]$$

is e -positive. This follows by comparing the coefficients of the e -functions involved. Consider (27) applied to $\frac{1}{4}X_{P_{n-2,v,w}}$, and use the coefficients described in Corollary 4.1. We obtain the following formulas for the coefficients:

$$[e_n] \frac{1}{4}X_{P_{n-2,v,w}} = n, \quad [e_{n-1}e_1] \frac{1}{4}X_{P_{n-2,v,w}} = n-2, \quad \text{and} \quad [e_{n-2}e_2] \frac{1}{4}X_{P_{n-2,v,w}} = n-4.$$

In particular, notice that the coefficients of e_n and $e_{n-1}e_1$ are zero and that the coefficient of $e_n e_2$ is $n-2$, which is positive for $n \geq 6$. Thus, the negative terms appearing in (28) are absorbed by terms in $\frac{1}{4}X_{P_{n-2,v,w}}$, and (28) is e -positive.

Finally, as this is an e -positive recurrence with e -positive initial conditions, by induction it follows that $X_{P_{n,v,w}}$ is e -positive for all n . \square

4.1.3. *Paths twinned at an interior vertex.* Next we provide an e -positive recurrence relation for path graphs twinned at an interior vertex.

Theorem 4.4. *The chromatic symmetric function $X_{P_{n,\ell}}$ for the path P_n twinned at the interior vertex ℓ satisfies the e -positive recurrence for $\ell \geq 2$, $n \geq \ell + 1$, and $n \geq 4$,*

$$\begin{aligned} X_{P_{n,\ell}} &= \sum_{j=2}^{n-\ell-1} (j-1)e_j X_{P_{n-j,\ell}} + 4(n+1)e_{n+1} + 2ne_1e_n + 2e_1 \sum_{j=n-\ell+2}^{n-1} (j-1)e_j X_{P_{n-j}} \\ &+ 4 \sum_{j=n-\ell+3}^n (j-1)e_j X_{P_{n+1-j}} + 2 \sum_{j=n-\ell+1}^{n-\ell+2} (j-2)e_j X_{P_{n+1-j}} + (n-\ell-2)e_{n-\ell} X_{P_{\ell,\ell}}. \end{aligned}$$

Thus, for $n \geq 3$ and $2 \leq \ell \leq n-1$, $X_{P_{n,\ell}}$ is e -positive.

Proof. Fix $\ell \geq 2$, and consider $n \geq \ell + 1$ with $n \geq 4$. We start with the recurrence relation in the statement of Proposition 3.19. This can be rewritten as

$$(29) \quad \begin{aligned} X_{P_{n,\ell}} &= 4X_{P_{n+1}} + 2e_1X_{P_n} + 2e_2X_{P_{\ell-1}}X_{P_{n-\ell}} \\ &- 2X_{P_{\ell+1}}X_{P_{n-\ell}} - 2X_{P_\ell}X_{P_{n-\ell+1}} - 2X_{P_{\ell-1}}X_{P_{n-\ell+2}}. \end{aligned}$$

From (29) and using Proposition 3.2(a), we have that

$$X_{P_{n,\ell}} = 4(n+1)e_{n+1} + 4 \sum_{j=2}^n (j-1)e_j X_{P_{n+1-j}} + 2e_1 \left[ne_n + \sum_{j=2}^{n-1} (j-1)e_j X_{P_{n-j}} \right]$$

$$\begin{aligned}
& + 2e_2 X_{P_{\ell-1}} \left[(n-\ell)e_{n-\ell} + \sum_{j=2}^{n-\ell-1} (j-1)e_j X_{P_{n-\ell-j}} \right] \\
& - 2X_{P_{\ell+1}} \left[(n-\ell)e_{n-\ell} + \sum_{j=2}^{n-\ell-1} (j-1)e_j X_{P_{n-\ell-j}} \right] \\
(30) \quad & - 2X_{P_\ell} \left[(n+1-\ell)e_{n+1-\ell} + \sum_{j=2}^{n-\ell} (j-1)e_j X_{P_{n+1-\ell-j}} \right] \\
(31) \quad & - 2X_{P_{\ell-1}} \left[(n+2-\ell)e_{n+2-\ell} + \sum_{j=2}^{n-\ell+1} (j-1)e_j X_{P_{n+2-\ell-j}} \right].
\end{aligned}$$

Notice that, in the six summands above, for each fixed j , the terms attached to the factor $(j-1)e_j$, when collected together, match the six terms in the right-hand side of the recurrence (29) applied to $X_{P_{n-j,\ell}}$. Grouping the remaining terms into an expression Y if they have a positive sign, or Z if they have a negative sign, we obtain

$$X_{P_{n,\ell}} = \sum_{j=2}^{n-\ell-1} (j-1)e_j X_{P_{n-j,\ell}} + Y - Z.$$

The positive terms Y are given by

$$\begin{aligned}
Y &= 4(n+1)e_{n+1} + 2ne_n e_1 + 2e_2 X_{P_{\ell-1}}(n-\ell)e_{n-\ell} \\
&+ 4 \sum_{j=n-\ell}^n (j-1)e_j X_{P_{n+1-j}} + 2e_1 \sum_{j=n-\ell}^{n-1} (j-1)e_j X_{P_{n-j}} \\
&= 4(n+1)e_{n+1} + 2ne_n e_1 + 4 \sum_{j=n-\ell+3}^n (j-1)e_j X_{P_{n+1-j}} + 2e_1 \sum_{j=n-\ell+2}^{n-1} (j-1)e_j X_{P_{n-j}} \\
&+ 4 \underbrace{\sum_{j=n-\ell}^{n-\ell+2} (j-1)e_j X_{P_{n+1-j}} + 2e_1 \sum_{j=n-\ell}^{n-\ell+1} (j-1)e_j X_{P_{n-j}} + 2e_2 X_{P_{\ell-1}}(n-\ell)e_{n-\ell}}_{Y_1},
\end{aligned}$$

where the last line is obtained by splitting the summations. The negative terms Z are given by

$$\begin{aligned}
Z &= 2X_{P_{\ell+1}}(n-\ell)e_{n-\ell} + 2X_{P_\ell}(n+1-\ell)e_{n+1-\ell} + 2X_{P_{\ell-1}}(n+2-\ell)e_{n+2-\ell} \\
&+ 2X_{P_\ell}(n-\ell-1)e_{n-\ell} X_{P_1} + 2X_{P_{\ell-1}} \sum_{j=n-\ell}^{n-\ell+1} (j-1)e_j X_{P_{n+2-\ell-j}},
\end{aligned}$$

where the last two terms come from the sums in (30) and (31). We rewrite Z so that $Y - Z$ is easier to analyze.

$$\begin{aligned} Z &= 2 \sum_{j=n-\ell}^{n-\ell+2} j e_j X_{P_{n+1-j}} \\ &\quad + 2X_{P_\ell}(n-\ell-1)e_{n-\ell}X_{P_1} + 2X_{P_{\ell-1}}(n-\ell)e_{n-\ell+1}X_{P_1} + 2X_{P_{\ell-1}}(n-\ell-1)e_{n-\ell}X_{P_2} \\ &= 2 \sum_{j=n-\ell}^{n-\ell+2} j e_j X_{P_{n+1-j}} + 2e_1 \sum_{j=n-\ell}^{n-\ell+1} (j-1)e_j X_{P_{n-j}} + 2X_{P_2}X_{P_{\ell-1}}(n-\ell-1)e_{n-\ell} \end{aligned}$$

Using $X_{P_2} = 2e_2$, we have

$$Y_1 - Z = 2 \sum_{j=n-\ell}^{n-\ell+2} (j-2)e_j X_{P_{n+1-j}} - 2X_{P_{\ell-1}}(n-\ell-2)e_{n-\ell}e_2,$$

because the terms with the factor $e_1 = X_{P_1}$ can be seen to vanish identically. By splitting the sum, $Y_1 - Z$ can be rewritten as

$$Y_1 - Z = 2 \sum_{j=n-\ell+1}^{n-\ell+2} (j-2)e_j X_{P_{n+1-j}} + 2(n-\ell-2)e_{n-\ell}(X_{P_{\ell+1}} - e_2X_{P_{\ell-1}}).$$

In the last term on the right-hand side, the factor of $2(X_{P_{\ell+1}} - e_2X_{P_{\ell-1}}) = 2X_{P_{\ell+1}} - X_{P_2}X_{P_{\ell-1}} = X_{P_{\ell,\ell}}$ is precisely the chromatic symmetric function of the path P_ℓ twinned at a leaf. Thus, putting all of this together, we obtain the recurrence relation from the statement.

Finally, we can deduce the e -positivity. For the initial values $n = \ell + 1, \ell + 2, \ell + 3$, $X_{P_{n,\ell}}$ is e -positive by Proposition 3.25. We proceed by strong induction on n to show the claimed e -positivity for $X_{P_{n,\ell}}$ for $n \geq \ell + 4$. Our induction hypothesis is that $X_{P_{m,\ell}}$ is e -positive for all $m < n, m \geq \ell + 1$. We only need to look at $Y_1 - Z$ since that is the part containing negative terms. By Proposition 4.2 we know that $Y_1 - Z$ is e -positive. Hence, by induction the proof is complete. \square

As an application of the preceding recurrence, we have the following corollary.

Corollary 4.5 (See also [10, Theorem 7.8]). *Consider the path P_n on n vertices, with $n \geq 4$. Let $2 \leq \ell \leq n - 2$ and let v be the leaf n . Then the chromatic symmetric function of $P_{n,\ell,v}$ is e -positive.*

Proof. The triple deletion argument implies that

$$X_{P_{n,\ell,v}} = 2X_{P_{n+1,\ell}} - X_{P_2}X_{P_{n-1,\ell}} = 2(X_{P_{n+1,\ell}} - e_2X_{P_{n-1,\ell}}).$$

Examining the recurrence for $X_{P_{n,\ell}}$ in Theorem 4.4 one sees that, when $n - \ell - 1 \geq 2$, the initial term in the first sum in the expression for $X_{P_{n,\ell}}$ is $e_2X_{P_{n-2,\ell}}$, making $X_{P_{n,\ell}} - e_2X_{P_{n-2,\ell}}$ e -positive. Replacing n by $n + 1$ now gives e -positivity of $X_{P_{n+1,\ell}} - e_2X_{P_{n-1,\ell}}$ for $n - \ell \geq 2$. \square

4.2. Recurrence for twinned cycles. In this section we derive an e -positive recursive formula for the twinned cycle, analogous to those for the twinned path from the last section. We give a similar formula for another family of graphs that we call moose graphs.

4.2.1. *Twinned cycles.* We start with the cycle graph.

Theorem 4.6. *The chromatic symmetric function $X_{C_{n,v}}$ for the cycle C_n twinned at a vertex v is e -positive. For $n \geq 5$, it satisfies the e -positive recurrence*

$$X_{C_{n,v}} = \sum_{k=3}^{n-2} (k-1)e_k X_{C_{n-k,v}} + 2(n+1)(2n-3)e_{n+1} + 2(n-1)(n-3)e_n e_1 \\ + e_2 [X_{C_{n-2,v}} - 2(n-3)e_{n-1}],$$

with initial conditions

$$X_{C_{1,v}} = 2e_2, \quad X_{C_{2,v}} = 6e_3, \quad X_{C_{3,v}} = 24e_4, \quad \text{and} \quad X_{C_{4,v}} = 50e_5 + 6e_4 e_1 + 4e_3 e_2.$$

Proof. We proceed by induction on n . The initial cases $n \leq 4$ are verified by direct computation using Theorem 3.1 and Lemma 3.26. Note that these initial terms are all e -positive.

Assume we have shown the claim to be true for $X_{C_{m,v}}$ for all $m < n$. Now we rewrite the expression in Lemma 3.26 using Proposition 3.2 to obtain:

$$(32) \quad X_{C_{n,v}} = 4(n+1)ne_{n+1} + 4 \sum_{k=2}^{n-1} (k-1)e_k X_{C_{n+1-k}} \\ + 2e_1 \left[n(n-1)e_n + \sum_{k=2}^{n-2} (k-1)e_k X_{C_{n-k}} \right] \\ - 6 \left[(n+1)e_{n+1} + \sum_{k=2}^n (k-1)e_k X_{P_{n+1-k}} \right] \\ + 2e_2 \left[(n-1)e_{n-1} + \sum_{k=2}^{n-2} (k-1)e_k X_{P_{n-1-k}} \right].$$

Applying Lemma 3.26 again, we can collect the four summations above into one sum and three additional terms as follows:

$$\sum_{k=2}^{n-2} (k-1)e_k [4X_{C_{n+1-k}} + 2e_1 X_{C_{n-k}} - 6X_{P_{n+1-k}} + 2e_2 X_{P_{n-1-k}}] \\ + 4(n-2)e_{n-1} X_{C_2} - 6(n-2)e_{n-1} X_{P_2} - 6(n-1)e_n X_{P_1} \\ = \sum_{k=2}^{n-2} (k-1)e_k X_{C_{n-k,v}} + 4(n-2)e_{n-1} X_{C_2} - 6(n-2)e_{n-1} X_{P_2} - 6(n-1)e_n X_{P_1}.$$

Combining this expression with the remaining terms from (32), we obtain

$$(33) \quad X_{C_{n,v}} = \sum_{k=2}^{n-2} (k-1)e_k X_{C_{n-k,v}} + 2(n+1)(2n-3)e_{n+1} + 2(n-1)(n-3)e_n e_1 \\ - 2(n-3)e_{n-1} e_2.$$

We isolate the term $k = 2$ from the summation and regroup it with the last term in (33), so that it becomes

$$X_{C_{n,v}} = \sum_{k=3}^{n-2} (k-1)e_k X_{C_{n-k,v}} + 2(n+1)(2n-3)e_{n+1} + 2(n-1)(n-3)e_n e_1 + e_2 [X_{C_{n-2,v}} - 2(n-3)e_{n-1}],$$

as stated in the theorem.

Now we want to show e -positivity. By the induction hypothesis, only the last term requires scrutiny. Using Corollary 4.1 and Lemma 3.26, the coefficient of e_{n-1} in $X_{C_{n-2,v}}$ is $2(n-1)(2n-7)$. Therefore, the coefficient of e_{n-1} in $X_{C_{n-2,v}} - 2(n-3)e_{n-1}$ is $2(n-1)(2n-7) - 2(n-3) = 4(n^2 - 5n + 5)$, which is nonnegative for $n \geq 4$. Thus, by the induction hypothesis, the formula in the statement for $X_{C_{n,v}}$ is indeed an e -positive recurrence for the twinned cycles for $n \geq 5$. \square

4.2.2. *The moose graph.* We define the *moose graph* A_{n+2} to be the graph on $n+2$ vertices and $n+1$ edges, obtained from the cycle graph C_n by attaching a leaf to each of the vertices v, w of an edge vw in C_n ,




FIGURE 8. The moose graph A_{n+2} .

We provide an e -positive recurrence relation for the chromatic symmetric function of the moose graph.

Theorem 4.7. *For $n \geq 2$, the chromatic symmetric function of the moose graph A_{n+2} is e -positive. For $n \geq 4$, it satisfies the e -positive recurrence*

$$X_{A_{n+2}} = \left(\sum_{j=2}^{n-2} (j-1)e_j X_{A_{n+2-j}} \right) + (n+2)(n-1)e_{n+2} + 2e_1 e_{n+1} (n^2 - n - 1) + (n-1)(n-2)e_1^2 e_n + 2e_2 e_n,$$

with initial values

$$\begin{aligned} X_{A_4} &= X_{P_4} = 2e_2^2 + 2e_3 e_1 + 4e_4, \\ X_{A_5} &= 2e_3 e_1^2 + 2e_3 e_2 + 10e_4 e_1 + 10e_5, \\ X_{A_6} &= 2e_2^3 + 2e_3 e_2 e_1 + 6e_4 e_1^2 + 6e_4 e_2 + 22e_5 e_1 + 18e_6. \end{aligned}$$

Proof. We start by applying the triple deletion argument to A_{n+2} . The resulting expression involves two graphs T_{n+2} and Y_{n+2} , to which we apply the triple deletion argument as stated in Corollary 2.2. We depict this triple deletion argument in Figure 9. Using $X_{P_1} = e_1$ and $X_{P_2} = 2e_2$, the resulting expression only involves cycles paths:

$$(34) \quad X_{A_{n+2}} = X_{C_{n+2}} + 2e_1X_{C_{n+1}} + e_1^2X_{C_n} + 2e_2X_{P_n} - 2e_1X_{P_{n+1}} - 2X_{P_{n+2}}.$$




FIGURE 9. Derivation of (34) in the proof of Theorem 4.7, via three applications of the triple deletion argument.

The e -positive initial values follow from the recurrence and the initial values for X_{C_n} and X_{P_n} from Proposition 3.2.

The induction hypothesis is that $X_{A_{m+2}}$ is e -positive for $m < n$. Using the recurrences in Proposition 3.2, we obtain the following recurrence for $X_{A_{n+2}}$, valid for $n \geq 4$:

$$X_{A_{n+2}} = (n+2)(n+1)e_{n+2} + \sum_{j=2}^n (j-1)e_j X_{C_{n+2-j}}$$

$$\begin{aligned}
& + 2e_1 \left((n+1)ne_{n+1} + \sum_{j=2}^{n-1} (j-1)e_j X_{C_{n+1-j}} \right) \\
& + e_1^2 \left(n(n-1)e_n + \sum_{j=2}^{n-2} (j-1)e_j X_{C_{n-j}} \right) + 2e_2 \left(ne_n + \sum_{j=2}^{n-1} (j-1)e_j X_{P_{n-j}} \right) \\
& - 2e_1 \left((n+1)e_{n+1} + \sum_{j=2}^n (j-1)e_j X_{P_{n+1-j}} \right) - 2 \left((n+2)e_{n+2} + \sum_{j=2}^{n+1} (j-1)e_j X_{P_{n+2-j}} \right).
\end{aligned}$$

Collecting terms now gives, for $n \geq 4$,

$$\begin{aligned}
X_{A_{n+2}} &= \sum_{j=2}^{n-2} (j-1)e_j X_{A_{n+2-j}} + (n+2)(n+1)e_{n+2} + (n-1)e_n X_{C_2} + (n-2)e_{n-1} X_{C_3} \\
&+ 2(n+1)ne_1e_{n+1} + 2e_1(n-2)e_{n-1} X_{C_2} \\
&+ e_1^2 n(n-1)e_n \\
&+ 2ne_2e_n + 2e_2(n-2)e_{n-1} X_{P_1} \\
&- 2(n+1)e_1e_{n+1} - 2(n-1)e_1e_n X_{P_1} - 2(n-2)e_1e_{n-1} X_{P_2} \\
&- 2(n+2)e_{n+2} - 2ne_{n+1} X_{P_1} - 2(n-1)e_n X_{P_2} - 2(n-2)e_{n-1} X_{P_3}.
\end{aligned}$$

Now collect the terms outside the summation, using values from Proposition 3.2 to obtain

$$\begin{aligned}
X_{A_{n+2}} &= \sum_{j=2}^{n-2} (j-1)e_j X_{A_{n+2-j}} \\
&+ (n+2)(n-1)e_{n+2} + 2(n^2 - n - 1)e_{n+1}e_1 + (n-1)(n-2)e_n e_1^2 + 2e_n e_2.
\end{aligned}$$

All terms in the right side of this equation are now e -positive, and so by induction the proof is complete. \square

5. FUTURE DIRECTIONS

This work provides explicit e -positive generating functions for the chromatic symmetric function of twinned paths and cycles, and suggests that it may be worthwhile to undertake a similar study for twins of other graph families.

More generally, an examination of Table 1 shows that much of the recent literature focuses on establishing Gebhard and Sagan's (e)-positivity of the chromatic symmetric function in *noncommuting* variables. Although (e)-positivity implies e -positivity as a symmetric function in ordinary commuting variables, in such cases an explicit e -positive generating function or recurrence would be desirable. We propose the following future investigations in this direction:

- (1) For the twinned cycle graph, is the chromatic symmetric function in noncommuting variables (e)-positive?
- (2) Are there pleasing e -positive symmetric function expansions for those families whose e -positivity is known only via the stronger (e)-positivity property? Specific examples that may admit nice generating functions are the triangular ladder [4, 19], the kayak paddle graphs [1] and the tadpole graph [10, 12].

It would also be interesting to examine twinning for the *chromatic quasisymmetric function* of Shareshian and Wachs [16, 17] since the main class of posets of study for these, whose incomparability graphs are unit interval graphs, is also closed under the appropriately defined twinning operation for labeled graphs.

Acknowledgments. This work began at the 2023 Graduate Research Workshop in Combinatorics, which was supported in part by NSF grant #1953445, NSA grant #H98230-23-1-0028, and the Combinatorics Foundation. We would like to thank the organizers for providing a great venue for collaboration.

E. Banaian was supported by Research Project 2 from the Independent Research Fund Denmark (grant no. 1026-00050B). J. Lentfer was supported by the National Science Foundation Graduate Research Fellowship DGE-2146752.

REFERENCES

1. F. Aliniaiefard, V. Wang, and S. van Willigenburg, *The chromatic symmetric function of a graph centred at a vertex*, arXiv:2108.04850 (2021), preprint.
2. L. Carlitz, R. Scoville, and T. Vaughan, *Enumeration of pairs of sequences by rises, falls and levels*, Manuscripta Math. **19** (1976), no. 3, 211–243.
3. S. Cho and J. Huh, *On e -positivity and e -unimodality of chromatic quasisymmetric functions*, SIAM Journal on Discrete Mathematics **33** (2019), 2286–2315.
4. S. Dahlberg, *A new formula for Stanley’s chromatic symmetric function for unit interval graphs and e -positivity for triangular ladder graphs*, Sémin. Lothar. Combin. **82B** (2020), Art. 59, 12.
5. S. Dahlberg, A. Foley, and S. van Willigenburg, *Resolving Stanley’s e -positivity of claw-contractible-free graphs*, Journal of the European Mathematical Society **22** (2020), no. 8, 2673–2696.
6. S. Dahlberg, A. She, and S. van Willigenburg, *Schur and e -positivity of trees and cut vertices*, Electron. J. Combin. **27** (2020), no. 1, Paper No. 1.2, 22. MR 4057182
7. S. Dahlberg and S. van Willigenburg, *Lollipop and lariat symmetric functions.*, SIAM Journal on Discrete Mathematics **32** (2018), no. 2, 1029 – 1039.
8. A. M. Foley, C. T. Hoàng, and O. D. Merkel, *Classes of graphs with e -positive chromatic symmetric function*, Electron. J. Combin. **26** (2019), no. 3, Paper No. 3.51, 19. MR 4014621
9. A. M. Foley, J. Kazdan, L. Kröll, S. Martínez Alberga, O. Melnyk, and A. Tenenbaum, *Spiders and their kin: an investigation of Stanley’s chromatic symmetric function for spiders and related graphs*, Graphs Combin. **37** (2021), no. 1, 87–110. MR 4197368
10. D. D. Gebhard and B. E. Sagan, *A chromatic symmetric function in noncommuting variables*, Journal of Algebraic Combinatorics **13** (2001), no. 3, 227–255.
11. A. M. Hamel, C. T. Hoàng, and J. E. Tuero, *Chromatic symmetric functions and H -free graphs*, Graphs Combin. **35** (2019), no. 4, 815–825. MR 3962011
12. E. Y. H. Li, G. M. X. Li, D. G. L. Wang, and A. L. B. Yang, *The twinning operation on graphs does not always preserve e -positivity*, Taiwanese Journal of Mathematics **25** (2021), no. 6, 1089 – 1111.
13. G. M. X. Li and A. L. B. Yang, *On the e -positivity of $(\text{claw}, 2K_2)$ -free graphs*, Electron. J. Combin. **28** (2021), no. 2, Paper No. 2.40, 14. MR 4281206
14. I. G. Macdonald, *Symmetric functions and Hall polynomials*, Oxford University Press, 1998.
15. R. Orellana and G. Scott, *Graphs with equal chromatic symmetric functions*, Discrete Mathematics **320** (2014), 1–14.
16. J. Shareshian and M. L. Wachs, *Chromatic quasisymmetric functions and Hessenberg varieties*, Configuration spaces, CRM Series, vol. 14, Ed. Norm., Pisa, 2012, pp. 433–460.
17. J. Shareshian and M. L. Wachs, *Chromatic quasisymmetric functions*, Advances in Mathematics **295** (2016), 497–551.
18. R. P. Stanley, *A symmetric function generalization of the chromatic polynomial of a graph*, Adv. Math. **111** (1995), no. 1, 166–194.
19. R. P. Stanley, *Enumerative Combinatorics. volume 2*, Cambridge University Press, 1999.

20. R. P. Stanley and J. R. Stembridge, *On immanants of Jacobi-Trudi matrices and permutations with restricted position*, Journal of Combinatorial Theory, Series A **62** (1993), no. 2, 261–279.
21. J. R. Stembridge, *Eulerian numbers, tableaux, and the Betti numbers of a toric variety*, Discrete Mathematics **99** (1992), no. 1-3, 307–320.
22. F. Tom, *A signed e -expansion of the chromatic symmetric function and some new e -positive graphs*, arXiv:2311.08020 (2023), preprint.
23. S. Tsujie, *The chromatic symmetric functions of trivially perfect graphs and cographs*, Graphs and Combinatorics **34** (2018), no. 5, 1037–1048.
24. D. G. L. Wang and M. M. Y. Wang, *The e -positivity of two classes of cycle-chord graphs*, J. Algebraic Combin. **57** (2023), no. 2, 495–514. MR 4570410
25. M. Wolfe, *Symmetric chromatic functions*, Pi Mu Epsilon Journal **10** (1998), no. 8, 643–657.

DEPARTMENT OF MATHEMATICS, AARHUS UNIVERSITY, AARHUS DK

Email address: banaian@math.au.dk

DEPARTMENT OF MATHEMATICS, WAKE FOREST UNIVERSITY, NC

Email address: celanok@wfu.edu

DEPARTMENT OF MATHEMATICS, BROWN UNIVERSITY, PROVIDENCE, RI

Email address: megan_chang-lee@brown.edu

DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NC

Email address: lcolmen@ncsu.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WI

Email address: ogoff@wisc.edu

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MI

Email address: kimplej2@msu.edu

DEPARTMENT OF MATHEMATICS, JOHNS HOPKINS UNIVERSITY, BALTIMORE, MD

Email address: lkimpel1@jhu.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA

Email address: jlentfer@berkeley.edu

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MI

Email address: liangj26@msu.edu

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MN

Email address: shsund@umn.edu