

On analytic characterization of convex sets in \mathbb{R}^m (a survey)

Nikolay Kuznetsov

Laboratory for Mathematical Modelling of Wave Phenomena
 Institute for Problems in Mechanical Engineering
 Russian Academy of Sciences
 V.O., Bol'shoy pr. 61, St Petersburg 199178, Russia
 nikolay.g.kuznetsov@gmail.com

In the first part of this note, we review results concerning analytic characterization of convexity for planar sets. The second part is devoted to results valid for arbitrary $m \geq 2$.

1 Introduction

The notion of a convex set was introduced by Hermann Minkowski at the close of the 19th century. In 1897–1903, he published four papers treating different aspects of convex sets (see [23, p. 224] for the corresponding references), and his work initiated the modern theory of convexity. Minkowski's definition (it is given in his first paper [17, p. 198], where these sets are studied on their own), is equivalent to the common, modern one based on an intrinsic property of convex sets (cf. [23, Definition 1.3]):

Definition 1.1. A set $S \subset \mathbb{R}^m$ is convex if with every two points x, y belonging to S the entire segment joining x and y lies in S .

It is amazing that Motzkin's theorem that characterizes the convexity in another natural way—in terms of the nearest point property—appeared only in 1935.

Theorem 1.1 (Motzkin, [18]; see also [23], Theorem 7.8). *Let S be a closed set in \mathbb{R}^m . This set is convex if and only if a unique nearest point in S corresponds to every point in \mathbb{R}^m .*

In the 1950s, this result was generalized by using S_z —the set of all points in \mathbb{R}^m having z as a nearest point in S . Namely, the following characterization of convexity is valid.

Theorem 1.2 (Phelps, [20]). *Let S be a closed set in \mathbb{R}^m . This set is convex if and only if S_z is a closed cone in \mathbb{R}^m with vertex z .*

The following characterization of convexity was obtained in 1976. It involves the Steiner symmetrization of a set (see, for example, [23, Definition 12.7]), and its converse is also true; see the monograph [8].

Theorem 1.3 (Falconer, [9]). *Let $S \subset \mathbb{R}^m$ be a compact set of positive Lebesgue measure. If the Steiner symmetrizations of this set about all hyperplanes are convex, then S is convex.*

Further data on history of convexity can be found in an account presented by Fenchel [10] in 1983. However, it was not until the mid-1980s that first results on analytic characterization of convex sets had been published; see [3] and [19]. Almost 40 years past since then, but there is still no summarizing survey on diverse results in this field which has numerous applications; in particular, in the analysis of Hardy-type inequalities [4]. In the present paper, our aim is to fill in this gap, at least partially, by reviewing the mentioned, now classical, results along with new ones that appeared during the past few years; see [15] and [22].

The paper's plan is as follows. Several criteria for the convexity of a planar domain/closed set are considered in Section 2; in particular, a characterization of half-planes in terms of a distance function is given. In Section 3, various results are presented which are valid for arbitrary $m \geq 2$. In particular, the characterization of convexity is considered which involves a bilinear form defined on exterior, unit normals on ∂D . It is valid for a bounded domain D with smooth boundary and its proof is based on the Crofton formula for the surface measure $|\partial D|$. The proof of another characterization of convexity is based on application of another bilinear form; namely, the norm in the Sobolev space $H^{1/2}(\mathbb{R}^m)$.

2 The convexity of planar sets

Throughout this section D will denote a proper subdomain of the Euclidean plane \mathbb{R}^2 , and so the boundary ∂D is not empty. Hence the distance function with respect to ∂D is defined; namely,

$$d(x, \partial D) = \inf_{y \in \partial D} |x - y| \text{ or } d(x) \text{ for brevity;} \quad (2.1)$$

here $x = (x_1, x_2) \in D$ and $|x - y| = \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2}$ is the Euclidean norm in \mathbb{R}^2 . Thus the disc

$$B_r(x) = \{y \in \mathbb{R}^2 : |y - x| < r\} \subset D \text{ provided } r = d(x). \quad (2.2)$$

It occurs that the distance function $d(\cdot, \partial D)$ is a convenient tool for characterization of convexity.

2.1. Convexity of D via the distance function. The following result has a simple proof.

Theorem 2.1 (Armitage and Kuran, [3]). *If $d(\cdot, \partial D)$ is superharmonic in a domain $D \subset \mathbb{R}^2$, then D is convex.*

Proof. Let us assume $D \subset \mathbb{R}^2$ to be nonconvex and show that $d(\cdot, \partial D)$ is not superharmonic in D . According to the assumption, there exist a point $y \in \partial D$, a closed half-plane P with $y \in \partial P$ (without loss of generality, we set $y = (0, 0)$ and $P = \{x \in \mathbb{R}^2 : x_2 \geq 0\}$), and $r > 0$ such that

$$P \cap (\overline{B_r(y)} \setminus \{y\}) \subset D, \text{ whereas } (\partial D \setminus \{y\}) \cap B_r(y) \subset \mathbb{R}^2 \setminus P;$$

cf. [23, Theorem 4.8].

Then for $x \in B = B_{r/8}(x_0)$, where $x_0 = (0, r/4)$, we have that $d(x, \partial D) > x_2$ provided $x_1 \neq 0$. Hence

$$\int_B d(x, \partial D) dx > \int_B x_2 dx = \pi(r/8)^2(r/4) = \pi(r/8)^2 d(x_0, \partial D),$$

and so the area mean-value inequality, guaranteeing the superharmonicity of $d(\cdot, \partial D)$, is violated for B . \square

It should be emphasized that Theorem 2.1 is specifically a two-dimensional result, because neither D nor \overline{D} need be convex in higher dimensions. The example given by Armitage and Kuran [3, Sect. 5] to confirm this assertion involves the domain in \mathbb{R}^3 bounded by a particular torus.

2.2. Characterization of half-planes. Presumably, the first question about an analytic characterization of a convex set appeared as an exercise in the textbook [12]. It concerns a half-plane which is the maximal, so to speak, convex domain in \mathbb{R}^2 . The exceptional analytic property of its distance function d arises from the following observation.

Let a point $x \in \mathbb{R}^2$ belong to a half-plane which can be taken $\{x \in \mathbb{R}^2 : x_2 > 0\}$, without loss of generality. It is obvious that the distance from x to the x_1 -axis bounding this half-plane—the length of the segment orthogonal to the x_1 -axis—is equal to x_2 , and so is a harmonic function in this domain. The converse assertion is far from being trivial to prove; it is referred to as one of “three secrets about harmonic functions” in the interesting note [6], where the characterization of half-planes is as follows.

Theorem 2.2. *If $d(\cdot, \partial D)$ is harmonic in a domain $D \subset \mathbb{R}^2$, then D is a half-plane.*

A sketch of proof. It is clear that (2.2) is valid for every $x \in D$ and $D = d^{-1}(0, +\infty)$. Fixing $x^* \in D$ arbitrarily, we denote by y^* a point on ∂D with $|y^* - x^*| = d(x^*)$. Without loss of generality, y^* can be taken as the origin, whereas $x^* = (0, x_2^*)$ with $x_2^* > 0$. Then $B_{x_2^*}(x^*) \subset D$ and the fact that

$$d(x) = x_2 \text{ for every } x = (0, x_2) \text{ with } x_2 \in (0, 2x_2^*]$$

is a consequence of harmonicity and continuity of d . Hence the equality $d(x) = x_2$ is valid in $B_{x_2^*}(x^*)$ because harmonic functions are real-analytic. In the same way, we have that $B_{2x_2^*}(x_1^*) \subset D$, where $x_1^* = (0, 2x_2^*)$, and so

$$d(x) = x_2 \text{ for every } x = (0, x_2) \text{ with } x_2 \in (0, 2^2x_2^*],$$

in which case $d(x) = x_2$ in $B_{2x_2^*}(x_1^*)$. Iterating this procedure, we obtain that $d(x) = x_2$ in the disc $B_{2^n x_2^*}(x_n^*) \subset D$, where $x_n^* = (0, 2^n x_2^*)$.

It is clear that the union of discs $\cup_{n \in \{1, 2, \dots\}} B_{2^n x_2^*}(x_n^*)$ coincides with the half-plane

$$P_0 = \{x \in \mathbb{R}^2 : x_2 > 0\} \text{ and } d(x) = x_2 \text{ in } P_0,$$

because this equality is valid in each disc. Since the latter is a subset of D , we have that $P_0 \subset D$. On the other hand, d is superharmonic in D , and so this domain is convex by Theorem 2.1. But a convex domain can contain a half-plane only coinciding with it; thus, $D = P_0$. \square

2.3. Convexity of D via solutions to the modified Helmholtz equation. It occurs that it is possible to describe the domain’s convexity by means of these solutions (labelled panharmonic functions in [7]) by imposing some conditions on them. Let ∂D be bounded and sufficiently smooth, say C^2 ; then the Dirichlet problem for the modified Helmholtz equation

$$\nabla^2 v - \mu^2 v = 0 \text{ for } x \in D, \quad \mu \in \mathbb{R}_+, \tag{2.3}$$

has a unique solution satisfying the boundary condition

$$v(x) = 1 \text{ for } x \in \partial D; \tag{2.4}$$

here and below $\nabla = (\partial_1, \partial_2)$ is the gradient operator, $\partial_i = \partial/\partial x_i$. It has long been known (see, for example, [14]) that a solution of this problem, say $v(x, \mu)$, satisfies the inequalities

$$0 < v(x, \mu) \leq 1 \text{ for all } x \in \overline{D} \text{ and any } \mu > 0. \tag{2.5}$$

In this connection, see the recent survey [2, Sect. 2.1], where a detailed account of results concerning the strong maximum principle for elliptic operators is presented.

Let us show that problem’s solutions corresponding to large values of μ characterize the domain D in the following way.

Theorem 2.3. *Let $D \subset \mathbb{R}^2$ be a bounded domain with smooth boundary. If the condition*

$$|\nabla v(x, \mu)| \leq \mu v(x, \mu), \quad x \in D, \quad (2.6)$$

holds for solutions of problem (2.3), (2.4) with all large $\mu > 0$, then D is convex.

Before proving this theorem, let us illustrate it with two elementary examples. In the first one, the domain D is the half-plane P_0 , in which case it is natural to require that $v(x, \mu) \rightarrow 0$ as $x_2 \rightarrow \infty$. Duffin [7] established that $v(x, \mu) = e^{-\mu x_2}$ solves problem (2.3), (2.4) complemented by the latter condition; see his proof of Theorem 5. Condition (2.6) is fulfilled for this function for all $\mu > 0$, and so Theorem 2.3 implies the obvious fact that D is convex. Notice that

$$-\mu^{-1} \log v(x, \mu) = x_2 = d(x, \partial D)$$

in agreement with the assertion of Theorem 2.4 formulated below.

In the second example, the domain D is a disc, and so convex. As usual, I_0 stands for the modified Bessel function of the first kind (see, for example, [7, p. 111]). If $a \in (0, 1)$, then $v(x, \mu) = aI_0(\mu|x|)$ solves problem (2.3), (2.4) in the disc $B_r(0) = \{y \in \mathbb{R}^2 : |y| < r\}$ provided $r > 0$ is such that $aI_0(\mu r) = 1$.

It is clear that properties of the function I_0 (see [1, sect. 9.6]) yield condition (2.6) for $v(\cdot, \mu)$; indeed, it takes the form $I_1(\mu|x|) < I_0(\mu|x|)$, because

$$|\nabla v(x, \mu)| = \mu a I_1(\mu|x|)$$

in view of formula [1, 9.6.27].

Now, we formulate two assertions essential for proving Theorem 2.3; they follow from results obtained in the classical Varadhan's article [24].

Theorem 2.4 ([24], p. 434). *Let $D \subset \mathbb{R}^2$ be a bounded domain with smooth boundary. If $v(x, \mu)$ is a solution of the Dirichlet problem (2.3), (2.4), then*

$$-\mu^{-1} \log v(x, \mu) \rightarrow d(x, \partial D) \quad \text{as } \mu \rightarrow \infty \quad (2.7)$$

uniformly on \overline{D} .

The second assertion is a corollary of Theorem 3.6, [24].

Proposition 2.1. *Let $v(x, \mu)$ be a solution of the Dirichlet problem (2.3), (2.4) in a bounded domain $D \subset \mathbb{R}^2$ with smooth boundary. Then for every $\rho \in (0, 1/2)$, there exists a constant $C_\rho > 1$ such that the estimate*

$$v(x, \mu) \leq C_\rho \exp\{-\mu(1-\rho)d(x, \partial D)\}, \quad x \in \overline{D},$$

is valid for all $\mu > 0$.

Proof of Theorem 2.3. It is sufficient to show that conditions (2.6) imply that d is superharmonic in D ; indeed, this allows us to apply Theorem 2.1, thus demonstrating the convexity of D .

For a nonvanishing $u \in C^2(D)$, one obtains by a straightforward calculation that

$$\nabla^2(\log u) = -\frac{|\nabla u|^2}{u^2} + \frac{\nabla^2 u}{u} \quad \text{in } D.$$

Therefore, if $v(\cdot, \mu) > 0$ is a solution of the Dirichlet problem (2.3), (2.4) with large $\mu > 0$, then

$$-\nabla^2(\log v) = -\mu^2 + |\nabla v|^2/v^2 < 0$$

provided condition (2.6) is valid. Hence, $-\mu^{-1} \log v(\cdot, \mu)$ is superharmonic in D for all large $\mu > 0$.

The next step is to demonstrate that passage to the limit in Theorem 2.4 preserves superharmonicity. In view of inequality (2.5) and condition (2.6), Proposition 2.1 implies that

$$-\mu^{-1} \log v(x, \mu) \geq -\mu^{-1} \log C_\rho + (1 - \rho) d(x, \partial D), \quad x \in \overline{D}. \quad (2.8)$$

Averaging this over an arbitrary open disc $B_r(x)$ such that $\overline{B_r(x)} \subset D$, we obtain

$$-\mu^{-1} \log v(x, \mu) \geq -\mu^{-1} \log C_\rho + \frac{1 - \rho}{\pi r^2} \int_{B_r(x)} d(y, \partial D) dy.$$

Indeed, the function on the left-hand side of (2.8) is superharmonic and the first term on the right is harmonic. Letting $\mu \rightarrow \infty$ first and then $\rho \rightarrow +0$, we see that the last inequality turns into

$$d(x, \partial D) \geq \frac{1}{\pi r^2} \int_{B_r(x)} d(y, \partial D) dy, \quad x \in D,$$

in view of (2.7).

Since the obtained inequality is valid for all $r > 0$ such that $\overline{B_r(x)} \subset D$, the distance function $d(\cdot, \partial D)$ is superharmonic in D . Then Theorem 2.1 yields the assertion of Theorem 2.3, thus completing its proof. \square

Many assertions about convexity (see, in particular, Theorems 1.1 and 1.2) are formulated in the following form: “a set is convex if and only if some property is fulfilled”. Therefore, it is interesting to ascertain whether the converse of Theorem 2.3 is true.

Conjecture 2.1. Let $D \subset \mathbb{R}^2$ be a bounded, convex domain with smooth boundary. Then inequality (2.6) is fulfilled for solutions of problem (2.3), (2.4) provided μ is sufficiently large.

It is not unlikely that this assertion can be true for every $m \geq 2$.

2.4. A local version of Theorem 2.1. Let $F \neq \emptyset$ denote a proper, closed subset of \mathbb{R}^2 ; for every $x \in \mathbb{R}^2$ the distance function from F is

$$d(x, F) = \inf_{y \in F} |x - y|, \quad (2.9)$$

which is analogous to (2.1). It occurs that this function characterizes the convexity of F in the same way as the distance function $d(\cdot, \partial D)$ does it for the domain D ; namely, we have

Theorem 2.5 (Parker, [19]). *Let $D \subset \mathbb{R}^2$ be a domain such that $F \subset D$. Then the distance function $d(\cdot, F)$ is subharmonic in D if and only if F is convex.*

The proof is substantially more complicated than that of Armitage and Kuran’s result. Indeed, along with some simple auxiliary assertions it requires applying the Krein–Milman theorem (see [24, Theorem 11.5]), as well as the following local Motzkin-type theorem.

Theorem 2.6 (Parker, [19]). *Let $F \neq \emptyset$ be proper, closed subset of \mathbb{R}^m , $m \geq 2$. If for $x^* \in \partial F$ and some $r > 0$ every point in $B_r(x^*)$ has a unique nearest point in F , then there exists an open ball of radius r which touches F at x^* ; that is, this ball has no common points with F , but x^* belongs to the ball’s boundary.*

There is a very simple counterexample to Theorem 2.5 in three dimensions, and it can be easily extended to higher dimensions. Indeed; let (ρ, θ, z) be cylindrical coordinates in \mathbb{R}^3 , and let

$$F = \{\rho = 1, z = 0\};$$

that is, F is the unit circumference in the plane $\{z = 0\}$, which is obviously not convex. Calculating the Laplacian of the axisymmetric distance function

$$d(\rho, \theta, z, F) = \sqrt{(\rho - 1)^2 + z^2},$$

one immediately gets

$$d_{\rho\rho} + \rho^{-1}d_{\rho} + d_{zz} = (2\rho - 1)/(\rho d) \text{ for } \rho \neq 0.$$

Hence $d(\cdot, F)$ is subharmonic in the domain $\{\rho > 1/2, z \in \mathbb{R}\}$ containing the nonconvex set F , which demonstrates that Theorem 2.5 is not valid for domains of dimensions higher than two.

3 The convexity of a set in arbitrary dimension

At the beginning of this section, we consider results similar to those presented in Sections 2.1 and 2.4. Thereupon, an inequality characterizing the convexity of a bounded domain D with smooth boundary is described; it involves a bilinear form defined on exterior, unit normals on ∂D .

3.1. The convexity via the subharmonicity of the distance function. Let $F \neq \emptyset$ be a proper, closed subset of \mathbb{R}^m , $m \geq 2$, then the distance $d(x, F)$ from F is defined by (2.9) for every $x = (x_1, \dots, x_m) \in \mathbb{R}^m$. The following assertion extends Theorem 2.5 to higher dimensions.

Theorem 3.1 (Armitage and Kuran, [3]). *Let the closed set $F \subset \mathbb{R}^m$, $m \geq 2$, be proper. Then the distance function $d(\cdot, F)$ is subharmonic in $\mathbb{R}^m \setminus F$ if and only if F is convex.*

A sketch of proof. If F is convex, then the signed distance $d(\cdot, H)$ (it is measured from every its support hyperplane H so that it is negative in the interior of F) is harmonic in \mathbb{R}^m ; cf. Section 2.2. Moreover, we have that $d(\cdot, F) = \sup_H d(\cdot, H)$, and so this function is subharmonic in $\mathbb{R}^m \setminus F$, because $d(\cdot, H)$ is harmonic.

To prove the “only if” part of theorem’s assertion, let us assume that F is nonconvex. Then according to Theorem 1.1, there are two distinct points in F , say y_1 and y_2 , equidistant from some point in $\mathbb{R}^m \setminus F$; without loss of generality, the origin can be taken as this point, and so $|y_1| = |y_2| > 0$. Putting

$$v(x) = \min\{|x - y_1|, |x - y_2|\} \text{ for every } x \in \mathbb{R}^m,$$

it is easy to show that there exists $r_* > 0$ such that

$$v(0) > M(v, r) \text{ for every } r \in (0, r_*).$$

Here $M(v, r)$ stands for the mean value of v over the sphere $\{x \in \mathbb{R}^m : |x| = r\}$. Since

$$v(x) \geq d(x, F) \text{ for every } x \in \mathbb{R}^m \setminus F$$

with equality valid at the origin, we see that

$$d(0, F) = v(0) > M(v, r) \geq M(d(\cdot, F), r),$$

and so $d(\cdot, F)$ is not subharmonic in $\mathbb{R}^m \setminus F$. □

3.2. Characterization of convexity via an inequality for a bilinear form. The recent preprint [22] begins with the following heuristic observation.

Let $D \subset \mathbb{R}^m$, $m \geq 2$, be a bounded C^1 -domain. If two points $x, y \in \partial D$ are close, then the normals n_x, n_y at these points are almost parallel to each other and both of them are roughly orthogonal to $x - y$. However, the normal turns quickly in a region, where the curvature is large, but along with these regions there are “flatter” regions on the boundary of a convex domain. Therefore, it might all average out in this case.

The quantitative result based on this argument (it can be interpreted as a global conservation law for convex domains) is as follows.

Theorem 3.2 (Steinerberger, [22]). *Let $D \subset \mathbb{R}^m$, $m \geq 2$, be a bounded C^1 -domain. Then there exists a constant $C_m > 0$ such that*

$$\int_{\partial D \times \partial D} \frac{|(n_x, y - x)(y - x, n_y)|}{|x - y|^{m+1}} dS_x dS_y \geq C_m |\partial D| \quad (3.1)$$

with equality taking place if and only if D is convex.

Here (\cdot, \cdot) is the inner product in \mathbb{R}^m , whereas dS is the surface measure on ∂D and $|\partial D|$ is its total measure. If D is the unit ball, then the expression on the left-hand side of (3.1) reduces to

$$\frac{\sqrt{2} (\pi/2)^{m/2}}{\Gamma(m/2)} \int_{S^{m-1}} [1 - (x, w)]^{-(m-3)/2} dS_x, \quad w = (1, 0, \dots, 0) \in S^{m-1},$$

which yields that $C_m = 2^{-1} \int_{S^{m-1}} |x_1| dS$; in particular, $C_2 = 2$ and $C_3 = \pi$.

The proof of Theorem 3.2 given in the preprint [22] relies essentially on probabilistic technique; moreover, two assertions are crucial for this proof. The first one is the Crofton formula for a rectifiable hypersurface in \mathbb{R}^m (see, for example, [21]); for our purpose it takes the form:

$$|\partial D| = \alpha_m \int_L N_{\partial D}(\ell) d\varphi dp, \quad \alpha_m = \frac{\Gamma([m+1]/2)}{2 \pi^{(m-1)/2}}, \quad (3.2)$$

Here L is the set of all oriented lines in \mathbb{R}^m , the kinematic measure on $S^{m-1} \times (0, \infty)$ is denoted $d\varphi dp$, and $N_{\partial D}(\ell)$ stands for the number of transversal crossings of ∂D by $\ell \in L$. The second assertion is the following.

Lemma 3.1. *Let $D \subset \mathbb{R}^m$, $m \geq 2$, be a bounded C^1 -domain. Almost every (with respect to the kinematic measure) line $\ell \in L$ either has no common points with ∂D or intersects it transversally at two points exactly if and only if D is convex.*

As a consequence of this lemma and formula (3.2), one obtains the inequality

$$|\partial D| = \alpha_m \int_L N_{\partial D}(\ell) d\varphi dp \leq \frac{\alpha_m}{2} \int_L [N_{\partial D}(\ell)]^2 d\varphi dp,$$

where equality takes place if and only if D is convex. It occurs that (3.1) follows from this inequality through a chain of tricky manipulations. A slight modification of these considerations leads to the following.

Corollary 3.1. *Let $D \subset \mathbb{R}^m$, $m \geq 2$, be a bounded, convex C^1 -domain. Then*

$$\int_{\partial D} \frac{|(n_x, y - x)(y - x, n_y)|}{|x - y|^{m+1}} dS_y = C_m \quad \text{for every } x \in \partial D;$$

here C_m is the same as in (3.1).

Since the Crofton formula is known to be true in a very general setting (see [21]), some conditions imposed on D in this section can be relaxed. Indeed, an analogue of inequality (3.1) was recently obtained [5] under the assumption that D is just a set of finite perimeter. We recall that D is a set of this kind provided the distributional gradient $\nabla\chi_D$ of its characteristic function χ_D defines a finite measure on \mathbb{R}^m ; that is, $\int_{\mathbb{R}^m} |\nabla\chi_D| < \infty$. (An exposition of the theory of these sets is given in the monograph [16].) It occurs that the generalized inequality (see [5, formula (1.3)]) involves the so-called reduced boundary ∂^*D defined for a set of finite perimeter; in particular, integration over this boundary is used instead of the usual domain's boundary ∂D .

Another analytic characterization of convexity was obtained by Figalli and Jerison [11] also for a set F of finite perimeter. Namely, the authors considered the function

$$\{x \in \mathbb{R}^m : (x, v) = 0\} \ni u \mapsto W_v(u) = \int_{\mathbb{R}} \chi_F(u + tv) dt \in \mathbb{R}, \quad \text{where } v \in \mathbb{R}^m \text{ and } |v| = 1,$$

and proved the following.

Theorem 3.3. *Let $F \subset \mathbb{R}^m$, $m \geq 2$, be a bounded set of finite perimeter. If the function W_v is log-concave for almost every v with respect to the $(m - 1)$ -dimensional Hausdorff measure, then F is convex (up to a set of measure zero).*

We recall that W_v is log-concave if it has the form e^{-V} , where V is a convex function mapping $\{x \in \mathbb{R}^m : (x, v) = 0\}$ to $\mathbb{R} \cup \{+\infty\}$. Thus this theorem improves Theorem 1.3, according to which W_v is required to be concave to guarantee that F is convex. The converse of Theorem 3.3 is true as well. Indeed, if F is convex, then the Brunn–Minkowski inequality [13] implies that W_v is concave which is even stronger than log-concave.

The proof of Theorem 3.3 is based on measuring the perimeter of a set through the standard norm in the Sobolev space $H^{1/2}(\mathbb{R}^m)$, and so the assumption that F has finite perimeter is essential. However, the authors point out that the theorem could be true without it. It is also worth noticing that the norm in $H^{1/2}(\mathbb{R}^m)$ is a bilinear form like the left-hand side in (3.1).

Acknowledgement. The author thanks Stefan Steinerberger, whose comments about the original manuscript helped him to improve the presentation.

References

1. M. Abramowitz, I. A. Stegun (eds.), *Handbook of Mathematical Functions*. US National Bureau of Standards, Washington, DC, 1964.
2. D. E. Apushkinskaya, A. I. Nazarov, “The normal derivative lemma and surrounding issues,” *Russian Math. Surveys* **77** (2), 189–249 (2022).
3. D. H. Armitage, Ü. Kuran, “The convexity of a domain and the superharmonicity of the signed distance function,” *Proc. Amer. Math. Soc.* **93**, 598–600 (1985).
4. A. A. Balinsky, W. D. Evans, R. T. Lewis, *The analysis and geometry of Hardy’s inequality*, Springer Cham, Heidelberg-New York-Dordrecht-London, 2015.
5. R. E. G. Bushling, “A singular integral identity for surface measure,” *J. Geom. Anal.* **34**, # 16 (2024).
6. R. B. Burckel, “Three secrets about harmonic functions,” *Amer. Math. Monthly* **104**, 52–56 (1997).

7. R. J. Duffin, “Yukawan potential theory,” *J. Math. Anal. Appl.* **35**, 105–130 (1971).
8. H. G. Eggleston, *Convexity*, Cambridge University Press, Cambridge, 1958.
9. K. J. Falconer, “A result on the Steiner symmetrization of a compact set,” *J. Lond. Math. Soc.* (2) **14**, 385–386 (1976).
10. W. Fenchel, “Convexity through the ages,” in *Convexity and its Applications*, pp. 121–130. P. M. Gruber and J. M. Wills (eds.), Birkhäuser, Basel-Boston-Stuttgart, 1983.
11. A. Figalli, D. Jerison, “How to recognize convexity of a set from its marginals,” *J. Func. Anal.* **266**, 1685–1701 (2014).
12. W. H. J. Fuchs, *Topics in the Theory of Functions of One Complex Variable*, Van Nostrand, Princeton, 1967.
13. R. J. Gardner, “The Brunn–Minkowski inequality,” *Bull. Amer. Math. Soc.* **39**, 355–405 (2002).
14. E. Hopf, “Elementare Bemerkungen über die Lösungen partieller Differentialgleichungen zweiter Ordnung vom elliptischen Typus,” *Sitzungsber. Preuss. Akad. Wiss., Phys.-Math. Kl.* **19**, 147–152 (1927).
15. N. Kuznetsov, “The convexity of a planar domain via properties of solutions to the modified Helmholtz equation,” *J. Math. Sci.* **281**, 607–611 (2024).
16. F. Maggi, *Sets of Finite Perimeter and Geometric Variational Problems: An Introduction to Geometric Measure Theory*, Cambridge University Press, Cambridge, 2012.
17. H. Minkowski, “Allgemeine Lehrsätze über die konvexen Polyeder,” *Nachr. Ges. Wiss. Göttingen, Math.-Phys. Kl.* **1897**, 198–220 (1897); also *Gesammelte Abhandlungen*, II, B. G. Teubner, Leipzig-Berlin, 103–121 (1911).
18. T. Motzkin, “Sur quelques propriétés caractéristiques des ensembles convexes,” *Rend. Acad. Naz. Lincei, Classe Sci. Fis., Mat. Nat.* **21**, 562–567 (1935).
19. M. J. Parker, “Convex sets and subharmonicity of the distance function,” *Proc. Amer. Math. Soc.* **103**, 503–506 (1988).
20. R. R. Phelps, “Convex sets and nearest points,” *Proc. Amer. Math. Soc.* **8**, 790–797 (1957).
21. L. A. Santaló, *Integral Geometry and Geometric Probability*, Cambridge University Press, Cambridge, 1984.
22. S. Steinerberger, “An inequality characterizing convex domains,” preprint arXiv: 2209.14153v2
23. F. A. Valentine, *Convex Sets*, McGraw-Hill, New York, 1964.
24. S. R. S. Varadhan, “On the behavior of the fundamental solution of the heat equation with variable coefficients,” *Comm. Pure Appl. Math.* **20**, 431–455 (1967).