

ON THE LAPLACE OPERATOR WITH A WEAK MAGNETIC FIELD IN EXTERIOR DOMAINS

AYMAN KACHMAR, VLADIMIR LOTOREICHIK, AND MIKAEL SUNDQVIST

ABSTRACT. We study the magnetic Laplacian in a two-dimensional exterior domain with Neumann boundary condition and uniform magnetic field. For the exterior of the disk we establish accurate asymptotics of the low-lying eigenvalues in the weak magnetic field limit. For the exterior of a star-shaped domain, we obtain an asymptotic upper bound on the lowest eigenvalue in the weak field limit, involving the 4-moment, and optimal for the case of the disk. Moreover, we prove that, for moderate magnetic fields, the exterior of the disk is a local maximizer for the lowest eigenvalue under a p -moment constraint.

1. INTRODUCTION

1.1. Background and motivation. Inequalities between eigenvalues in terms of geometric data are central in spectral geometry. Many interesting results are proven for the Laplace operator. We mention the Faber–Krahn inequality [8, 19], that says that among all domains of finite volume, the ball minimizes the first Dirichlet eigenvalue. On the contrary, the ball is a maximizer of the first non-zero eigenvalue in the case of Neumann boundary conditions [25, 27].

L. Erdős extended the Faber–Krahn inequality to the magnetic Laplace operator with a uniform magnetic field, still under a Dirichlet boundary condition [7]. A quantitative version of the isoperimetric inequality by Erdős was recently obtained in [11]. Much less is known for other boundary conditions. It was suggested in [10] that the disk might in fact be a *maximizer* among all bounded two-dimensional simply-connected domains with fixed area. This conjecture is still out of reach, but there is some progress [3, 4, 16]. Related results concern eigenvalue optimization for the magnetic Laplacian with Robin boundary conditions [5, 15].

Optimization of the lowest eigenvalue for the non-magnetic Laplace operator with attractive Robin boundary conditions in exterior domains was analysed in [2, 20, 21], see also the monograph [1]. Less is known in this context for the magnetic Laplace operator in exterior domains, where the optimization of the lowest eigenvalue is meaningful also for Neumann boundary conditions.

We study here the case of magnetic Neumann boundary conditions, and weak magnetic fields, in the exterior of star-shaped domains. The exterior of the disk is the main suspect to be a maximizer of the lowest eigenvalue. Therefore, we develop in Section 2 precise estimates of the eigenvalues of the magnetic Laplacian in the exterior of a disk, and we obtain the eigenvalue splitting in the limit of weak

2020 *Mathematics Subject Classification.* 35P15; 58J50, 34L15, 35Q40.

Key words and phrases. magnetic Laplacian, eigenvalue asymptotics, weak magnetic fields, isoperimetric inequality, Landau levels.

magnetic field. We discuss the exterior of star-shaped domains about the origin in Section 3, and show that the exterior of the disk is indeed a local maximizer of the lowest eigenvalue under a fixed p -th moment of the exterior domain. Moreover, we obtain in the same section an asymptotic upper bound on the lowest eigenvalue in the weak field limit, which yields optimality of the exterior of a disk for sufficiently weak magnetic fields under fixed fourth moment of the exterior domain.

The main results in this paper are stated below in Theorem 1.1 concerning the exterior of the disk, and Theorems 1.3 and 1.5 concerning the exterior of a star-shaped domain about the origin. They highlight two new phenomena, the first is that the weak field limit in exterior domains turns out to have a *semi-classical* character, due to the fact the spectrum of the non-magnetic Laplacian is purely continuous in exterior domains. The second is the emergence of the moment constraint under which we verify the local optimality of the disk's exterior. A part of the novelty in our results, we believe that our analysis paves the way towards understanding whether the exterior of the disk is a maximizer among all exterior domains with fixed area or perimeter of the interior complement.

1.2. The magnetic Laplacian in an exterior domain. Consider a bounded simply connected domain $\Omega \subset \mathbb{R}^2$ with a C^∞ -smooth boundary $\partial\Omega$ and the outer unit normal vector field ν . Throughout this paper, we denote by $\Omega^{\text{ext}} = \mathbb{R}^2 \setminus \overline{\Omega}$, the exterior of the domain Ω . We also let $\mathcal{B}_R = \{x \in \mathbb{R}^2 : |x| < R\}$ denote the open disk of radius $R > 0$, and in particular \mathcal{B}_1 the unit disk.

Let \mathbf{A} be the vector potential defined as

$$\mathbf{A}(x) = \frac{1}{2}(-x_2, x_1)^\top, \quad (1.1)$$

and which generates the unit magnetic field, $\text{curl } \mathbf{A} = 1$. We study the self-adjoint magnetic Neumann Laplacian in $L^2(\Omega^{\text{ext}})$,

$$\mathcal{H}u := (-i\nabla - b\mathbf{A})^2 u,$$

$$\text{dom}(\mathcal{H}) := \{u \in H_{\mathbf{A}}^1(\Omega^{\text{ext}}) : (-i\nabla - b\mathbf{A})^2 u \in L^2(\Omega^{\text{ext}}), \nu \cdot (\nabla - ib\mathbf{A})u = 0 \text{ on } \partial\Omega\},$$

where $b \in \mathbb{R}$ and the first-order magnetic Sobolev space is defined by $H_{\mathbf{A}}^1(\Omega^{\text{ext}}) := \{u \in L^2(\Omega^{\text{ext}}) : (-i\nabla - b\mathbf{A})u \in [L^2(\Omega^{\text{ext}})]^2\}$. In fact, the operator is unitarily equivalent under a sign-flip of the magnetic field, so it is sufficient to consider $b \geq 0$. The operator \mathcal{H} is associated with the quadratic form

$$\mathfrak{q}_b(u) := \int_{\Omega^{\text{ext}}} |(-i\nabla - b\mathbf{A})u|^2 dx, \quad \text{dom}(\mathfrak{q}_b) := H_{\mathbf{A}}^1(\Omega^{\text{ext}}).$$

Contrary to the case of an interior domain, the spectrum of \mathcal{H} depends on the gauge of the vector potential \mathbf{A} , since we can not rule out this dependence by a gauge transformation.

The essential spectrum of \mathcal{H} consists of the Landau levels, $\{(2n-1)b : n \in \mathbb{N}\}$, and the discrete spectrum is infinite and accumulates at each Landau level from below [12]. In particular, the spectrum below b consists of a sequence of discrete eigenvalues

$$\lambda_1(b, \Omega^{\text{ext}}) \leq \lambda_2(b, \Omega^{\text{ext}}) \leq \dots < b,$$

counted with multiplicity.

The eigenvalues can, as usual, be characterized by the variational min-max theorem,

$$\lambda_k(b, \Omega^{\text{ext}}) = \inf_{\substack{M \subset H_{\mathbf{A}}^1(\Omega^{\text{ext}}) \\ \dim M = k}} \left(\max_{\substack{u \in M \\ u \neq 0}} \frac{\mathfrak{q}_b(u)}{\|u\|_{L^2(\Omega^{\text{ext}})}^2} \right).$$

1.3. Exterior of the disk. We will concentrate on the exterior of the disk, and prove that for weak magnetic fields, all the eigenvalues below b are simple, and obey a two-term asymptotics.

Theorem 1.1. *Assume that $\Omega = \mathcal{B}_R$ and that $b > 0$. Then the ground state of \mathcal{H} will be non-radial. Moreover, for any $k \geq 1$,*

$$\lambda_k(b, \mathcal{B}_R^{\text{ext}}) = b - \frac{R^{2k}}{2^{k-1}(k-1)!} b^{k+1} + \mathcal{O}(b^{k+\frac{3}{2}}), \quad (b \rightarrow 0^+).$$

Remark 1.2. (i) By a scaling argument, it suffices to prove Theorem 1.1 in the case of the exterior of the unit disk, $\Omega = \mathcal{B}_1$, which corresponds to $R = 1$.

- (ii) Our proof of Theorem 1.1 relies on separation of variables and the analysis of fibre operators. Even though this method does not directly carry over to general domains, we will give in Theorem 1.3 an asymptotic upper bound on the lowest eigenvalue for star-shaped domains that will enable us to compare with the disk under a certain moment constraint.
- (iii) For the Neumann magnetic operator in the disk \mathcal{B}_R , ground states are radial at least for $bR^2 < 1$ (see [16, Prop. 2.3]). The limit of the lowest eigenvalue as $b \rightarrow 0^+$ follows from standard perturbation theory [9, Prop. 1.5.2]; in fact, as $b \rightarrow 0^+$,

$$\lambda_1(b, \mathcal{B}_R) = \frac{b^2 R^2}{8} + \mathcal{O}(b^3).$$

- (iv) Interestingly, there is an analogy between Theorem 1.1 and the strong field asymptotics for the Dirichlet eigenvalues in the interior of the disk, which highlights an interior-exterior, Dirichlet-Neumann and weak-strong field duality. In fact, [14, Thm. 5.2] yields that,

$$\lambda_k^{\text{D}}(b, \mathcal{B}_R) = b + \frac{R^{2k}}{2^{k-1}(k-1)!} b^{k+1} e^{-bR^2/2} + o(b^{k+1} e^{-bR^2/2}), \quad (b \rightarrow +\infty),$$

where $\lambda_k^{\text{D}}(b, \mathcal{B}_R)$ denotes the k -th eigenvalue of $(-i\nabla - b\mathbf{A})^2$ in $L^2(\mathcal{B}_R)$, with Dirichlet boundary condition on $\partial\mathcal{B}_R$.

1.4. Exterior of a star-shaped domain. In our considerations of general domains we employ a quantity associated merely to the exterior domain itself. Clearly, the area of the exterior domain is always infinite, and we cannot use it in the analysis. However, the following p -th moment

$$\mathcal{I}_p(\Omega^{\text{ext}}) := \int_{\Omega^{\text{ext}}} \frac{1}{|x|^p} dx, \quad p > 2,$$

is finite, provided that the origin belongs to Ω . Such a moment appears in the geometric constraint and in the asymptotic upper bound for the values of p specified below. In some cases, suitable values of p will also depend on the intensity of the magnetic field b . We also remark that the p -th moment of Ω^{ext} is not invariant under translations of the domain and the results below can be further optimized by choosing the origin so that $\mathcal{I}_p(\Omega^{\text{ext}})$ is minimal.

1.4.1. *Asymptotic upper bound.* In the case where the domain Ω is star-shaped with respect to the origin and parametrized by a smooth mapping $\mathbb{R}/[0, 2\pi] \ni \theta \mapsto \rho(\theta)$ as $\Omega = \{(r, \theta) : r < \rho(\theta)\}$, we establish an upper bound on the lowest eigenvalue $\lambda_1(b, \Omega^{\text{ext}})$, valid in the limit of small b .

Theorem 1.3. *Let $\Omega \subset \mathbb{R}^2$ be a bounded C^∞ -smooth domain star-shaped with respect to the origin and parametrized by the C^∞ -smooth function $\rho : \mathbb{R}/[0, 2\pi] \rightarrow \mathbb{R}_+$. Then, as $b \rightarrow 0^+$, we have*

$$\lambda_1(b, \Omega^{\text{ext}}) \leq b - \left(\frac{|\Omega|}{2\pi} + \frac{\pi}{2\mathcal{I}_4(\Omega^{\text{ext}})} \right) b^2 + \mathcal{O}(b^{5/2}).$$

Moreover, for Ω being not a disk, there exists $b_0 = b_0(\Omega) > 0$ such that, for $0 < b < b_0$, we have,

$$\lambda_1(b, \Omega^{\text{ext}}) < \lambda_1(b, \mathcal{B}_{R_\star}^{\text{ext}}),$$

where R_\star denotes the radius of the disk satisfying $\mathcal{I}_4(\mathcal{B}_{R_\star}^{\text{ext}}) = \mathcal{I}_4(\Omega^{\text{ext}})$.

Remark 1.4. (i) The upper bound in Theorem 1.3 is consistent with the asymptotics in Theorem 1.1 if we take $\Omega = \mathcal{B}_R$, the disk of radius R .

(ii) If we choose the radius R so that $|\mathcal{B}_R| = |\Omega|$, the upper bound in Theorem 1.3 does not yield that the exterior of the disk is a maximizer of the lowest eigenvalue for sufficiently small b .

1.4.2. *Moderate field and moment constraint.* We consider next the more general case with a moderate magnetic field and address whether the exterior of the disk is a local maximizer of the lowest magnetic Neumann eigenvalue under suitable geometric constraints. We provide an affirmative answer when imposing a certain constraint and when Ω is a small deformation of the disk in a certain sense, which will be clarified below.

Our result demonstrates that the exterior of the disk is a local optimizer under the p -moment constraint. In the formulation of the theorem below, we use an auxiliary quantity

$$p_\star(b) := 2 + \frac{16b - 4b^2}{4 - 8b + b^2}.$$

Theorem 1.5. *Let $R > 0$, $bR^2 \in (0, 4 - 2\sqrt{3})$ and $p > p_\star(bR^2)$. Let $\Omega \subset \mathbb{R}^2$ be a bounded C^∞ -smooth domain star-shaped with respect to the origin and parametrized by the C^∞ -smooth function $\rho : \mathbb{R}/[0, 2\pi] \rightarrow \mathbb{R}_+$. Then, there exists $r_\star = r_\star(b, p, R) \in (0, R)$ such that under the assumptions $\mathcal{I}_p(\Omega^{\text{ext}}) = \mathcal{I}_p(\mathcal{B}_R^{\text{ext}})$ and $R - r_\star \leq \rho \leq R + r_\star$, the following isoperimetric inequality holds:*

$$\lambda_1(b, \Omega^{\text{ext}}) \leq \lambda_1(b, \mathcal{B}_R^{\text{ext}}).$$

Remark 1.6. Natural constraints under which optimality of the exterior of the disk can be analysed are fixed area or fixed perimeter of the interior domain, as in related analysis on the optimization of the lowest eigenvalue for the Robin Laplacian on exterior domains [2, 20, 21]. The method we propose for the magnetic Neumann Laplacian does not imply local optimality of the exterior disk under such constraints. (Local) optimality of the exterior disk under these constraints remains an open problem. The magnetic problem is subtle due to the fact that the ground state is not a radial function and the problem shares some common features with the optimization of the second Robin eigenvalue in exterior domains [22].

- Remark 1.7.* (i) In the limit $b \rightarrow 0^+$ we have $p_*(b) \rightarrow 2$. Thus, for any $p > 2$, we get by Theorem 1.5 local optimality of the exterior of the disk for a fixed p -moment provided that $b > 0$ is sufficiently small.
- (ii) We also point out that Theorem 1.5 establishes uniform local optimality of the exterior of the disk (under fixed p -moment) within exteriors of all smooth star-shaped domains, whose Hausdorff distance from the disk is smaller than some fixed positive constant, which depends only on the intensity b of the magnetic field, the parameter p , and the radius R of the disk.

We organized the rest of the paper as follows. Section 2 is devoted to the proof of Theorem 1.1. Section 3 is devoted to the proof of Theorems 1.3 and 1.5. Finally, Appendix A contains a result on the intersection of eigenvalue curves of the fibre operators.

2. THE WEAK FIELD LIMIT IN THE EXTERIOR OF THE DISK

In this section we prove the two-term asymptotic formula of the smallest eigenvalue in the exterior of the disk, stated in Theorem 1.1.

2.1. Fibre operators in the exterior of the disk.

2.1.1. *Splitting into fibre operators.* We work in the exterior of the unit disk, $\mathcal{B}_1^{\text{ext}}$. The action of the operator \mathcal{H} reads in polar coordinates as,

$$\mathcal{H} = -\frac{\partial^2}{\partial r^2} - \frac{1}{r} \frac{\partial}{\partial r} + \left(\frac{-i\partial_\theta}{r} - \frac{br}{2} \right)^2.$$

Since $\{e^{im\theta}\}_{m \in \mathbb{Z}}$ is an orthogonal basis for the square integrable functions on the circle, our operator splits naturally. We can work with the family $\{\mathcal{H}^{(m)}\}_{m \in \mathbb{Z}}$ of self-adjoint fibre operators in $L^2((1, +\infty); r dr)$,

$$\begin{aligned} \mathcal{H}^{(m)} = \mathcal{H}^{(m)}(b) &:= -\frac{d^2}{dr^2} - \frac{1}{r} \frac{d}{dr} + V_{m,b}(r), \\ \text{dom}(\mathcal{H}^{(m)}) &:= \left\{ u: u, -u'' - \frac{u'}{r} + V_{m,b}u \in L^2((1, +\infty); r dr), u'(1) = 0 \right\}, \end{aligned} \quad (2.1)$$

where

$$V_{m,b}(r) = \left(\frac{m}{r} - \frac{br}{2} \right)^2.$$

For each $b \in \mathbb{R}$,

$$\sigma(\mathcal{H}(b)) = \overline{\bigcup_{m \in \mathbb{Z}} \sigma(\mathcal{H}^{(m)}(b))},$$

where we use the decomposition $\mathcal{H} \simeq \bigoplus_{m \in \mathbb{Z}} \mathcal{H}^{(m)}$. We let $\lambda_1^{(m)}(b) \leq \lambda_2^{(m)}(b) \leq \dots$ denote the eigenvalues of $\mathcal{H}^{(m)}$. Away from eventual crossings, all $b \mapsto \lambda_k^{(m)}(b)$ are real analytic, see [18, Chap. VII, §3.5, Thm. 3.9].

We remark that the ground state energy of \mathcal{H} can be expressed as

$$\lambda_1(b, \mathcal{B}_1^{\text{ext}}) = \inf_{m \in \mathbb{Z}} \lambda_1^{(m)}(b). \quad (2.2)$$

In fact, we know that the essential spectrum of \mathcal{H} consists of the Landau levels b , $3b$, and so on. We will find eigenvalues of $\mathcal{H}^{(m)}(b)$ that are less than b , so $\lambda_1(b, \mathcal{B}_1^{\text{ext}})$ will indeed be an eigenvalue.

2.1.2. *A first control of each eigenvalue.* We recall that we only consider positive b , and warm up by excluding all $m \leq 0$ from the game. We also show that each $b \mapsto \lambda_1^{(m)}(b)$ will cross the lowest Landau level b exactly once for $b > 0$, and that it will happen for $b = 2m$.

Proposition 2.1. *For $b > 0$ the following holds.*

- (1) *If $m < 0$, then $\lambda_1^{(m)}(b) \geq 2b$.*
- (2) *If $m = 0$ and $b > 0$, then $\lambda_1^{(m)}(b) > b$.*
- (3) *If $m \geq 1$ then*

$$\begin{cases} \lambda_1^{(m)}(b) < b & \text{if } b < 2m, \\ \lambda_1^{(m)}(b) = b & \text{if } b = 2m, \\ \lambda_1^{(m)}(b) > b & \text{if } b > 2m. \end{cases}$$

- (4) *If $m \geq 1$, then the second eigenvalue of $\mathcal{H}^{(m)}$ satisfies $\lambda_2^{(m)}(b) > b$.*

Proof. We show the statements one by one.

- (1) For $m < 0$, we have

$$V_{m,b}(r) \geq 2|m|b$$

which directly yields

$$\lambda_1^{(m)}(b) > 2b.$$

(2) For $m = 0$ and $b > 0$, the eigenvalue equation $\mathcal{H}^{(0)}u = bu$ has the general solution

$$u(r) = u_0(r) := e^{-br^2/4} \left(c_1 + c_2 \int_1^r \rho^{-1} e^{b\rho^2/2} d\rho \right).$$

For the function not to increase exponentially at $+\infty$ we must have $c_2 = 0$. But then we find that $u_0'(r) = -c_1(br/2)e^{-br^2/4}$, and so we cannot satisfy the Neumann boundary condition $u_0'(1) = 0$ unless also $c_1 = 0$. Hence, there is no $b > 0$ such that $\lambda_1^{(0)}(b) = b$. We notice next that $V_{0,b}(r) = b^2r^2/4 \geq b^2/4$, so if $b > 2$ it holds that $\lambda_1^{(0)}(b) > b$. By continuity of $b \mapsto \lambda_1^{(0)}(b)$, this inequality is valid for all $b > 0$.

(3) Let us fix $m \geq 1$, solving again the eigenvalue equation for the lowest Landau level, $\mathcal{H}^{(m)}u = bu$, we get

$$u(r) = u_m(r) := r^m e^{-br^2/4} \left(c_1 + c_2 \int_1^r \rho^{-1-2m} e^{b\rho^2/2} d\rho \right).$$

The term with the coefficient c_2 is not in $L^2((1, +\infty); r dr)$, so we take $c_2 = 0$. We also choose c_1 so that u_m becomes normalized. The solution u_m satisfies the Neumann boundary condition if and only if $b = 2m$. Thus,

$$b \in \sigma(\mathcal{H}^{(m)}) \iff b = 2m. \quad (2.3)$$

Moreover, since $V_{m,b}(r) \geq b^2/4 - mb$ it holds that $V_{m,b}(r) > b$ if b is sufficiently large. This implies that $\lambda_1^{(m)}(b) > b$ if b is sufficiently large. We conclude that $\lambda_1^{(m)}(b) = 2m$ precisely when $b = 2m$ and that $\lambda_1^{(m)}(b) > b$ for all $b > 2m$.

We next show that $(\lambda_1^{(m)})'(2m) > 1$; that implies that we really have a crossing at $b = 2m$, and that $\lambda_1^{(m)}(b) < b$ for $0 < b < 2m$. By the Feynmann–Hellman formula

$$(\lambda_1^{(m)})'(b) = - \int_1^{+\infty} \left(m - \frac{br^2}{2} \right) |u_m|^2 r dr,$$

which can be justified by the fact that the sesquilinear forms associated with $\mathcal{H}^{(m)}(b)$ form a holomorphic family of the type (B) in the sense of Kato with respect to the parameter b and thus the perturbation formulae in [18, Chap. VII, §4.6] apply. Since $1 = \int_1^{+\infty} |u_m|^2 r \, dr$, we get for $b = 2m$,

$$(\lambda_1^{(m)})'(2m) - 1 = c_1^2 \int_1^{+\infty} (m(r^2 - 1) - 1) r^{2m+1} e^{-mr^2} \, dr.$$

Integrating by parts, we get

$$\begin{aligned} \int_1^{+\infty} m(r^2 - 1) r^{2m+1} e^{-mr^2} \, dr &= \int_1^{+\infty} \left(1 + m - \frac{m}{r^2}\right) r^{2m+1} e^{-mr^2} \, dr \\ &> \int_1^{+\infty} r^{2m+1} e^{-mr^2} \, dr, \end{aligned}$$

which yields that $(\lambda_1^{(m)})'(2m) > 1$.

(4) By the previous step we already know that $\lambda_2^{(m)}(b) > b$ for $b > 2m$. The inequality must still hold for $b = 2m$ since the groundstate of $\mathcal{H}^{(m)}$ is simple. In fact, it must also hold for $0 < b < 2m$, since otherwise we would have $\lambda_2^{(m)}(b) = b$ for some $0 < b < 2m$, and that is not possible according to (2.3). \square

Corollary 2.2. *For all $b > 0$ we have $\lambda_1^{(m)}(b) < b$ and \mathcal{H} does not have radially symmetric ground states.*

Proof. By Proposition 2.1, the ground state must be of the form $u_m(r)e^{im\theta}$ with m a positive integer. \square

Corollary 2.3. *For all $0 < b < 2$ and $m \geq 1$, $\lambda_1^{(m)}(b) < b$ is a discrete eigenvalue of \mathcal{H} ; moreover, $\lambda_1^{(m)}(b) \rightarrow b$ as $m \rightarrow +\infty$.*

Proof. Proposition 2.1 tells us that $\{\lambda_1^{(m)}(b)\}_{m \geq 1}$ is the sequence of discrete eigenvalues of \mathcal{H} below the bottom of the essential spectrum, b . Hence, $\lambda_1^{(m)}(b) \rightarrow b$ as $m \rightarrow +\infty$. \square

2.2. An effective operator. We would like to understand the limit behaviour, as $b \rightarrow 0^+$, of the eigenvalues $\lambda_1^{(m)}(b)$ and $\lambda_2^{(m)}(b)$ of the operator $\mathcal{H}^{(m)}(b)$ introduced in (2.1). This will be done by deriving an effective operator, which allows us to prove the following proposition.

Proposition 2.4. *For every $m \geq 1$, we have as $b \rightarrow 0^+$,*

$$\lambda_1^{(m)}(b) = b + o(b), \quad \lambda_2^{(m)}(b) \geq 3b + o(b).$$

The rest of this subsection is devoted to the proof of Proposition 2.4.

2.2.1. Unitary transformation and scaling. In order to transform the weighted space $L^2((1, +\infty); r \, dr)$ to the non-weighted Hilbert space $L^2((1, +\infty); dr)$, we apply the unitary transformation $U: L^2((1, +\infty); dr) \rightarrow L^2((1, +\infty); r \, dr)$

$$U: u(r) \mapsto r^{-1/2} u(r),$$

and then, we do the change of variable

$$r \mapsto \sqrt{\frac{b}{2}} (r - 1)$$

and get

$$U^{-1}\mathcal{H}^{(m)}(b)U \simeq \frac{b}{2}\mathcal{S}_b^{(m)},$$

where the self-adjoint operator $\mathcal{S}_b^{(m)}$ in $L^2(\mathbb{R}_+)$ is given by

$$\begin{aligned} \mathcal{S}_b^{(m)} &= -\frac{d^2}{dr^2} + w_{m,b}(r), \\ \text{dom}(\mathcal{S}_b^{(m)}) &= \left\{ u : u, -u'' + w_{m,b}u \in L^2(\mathbb{R}_+), u'(0) = \sqrt{\frac{1}{2b}}u(0) \right\}, \end{aligned}$$

with the potential $w_{m,b}$ defined by

$$w_{m,b} := \frac{4m^2 - 1}{4\left(r + \sqrt{\frac{b}{2}}\right)^2} + \left(r + \sqrt{\frac{b}{2}}\right)^2 - 2m.$$

2.2.2. *Localizing the spectrum of $\mathcal{S}_b^{(m)}$.* For fixed m , the spectrum of $\mathcal{S}_b^{(m)}$ converges to that of an operator on \mathbb{R}_+ formally defined by

$$\mathcal{S}_\star^{(m)} = -\frac{d^2}{dr^2} + \frac{4m^2 - 1}{4r^2} + r^2 - 2m,$$

with the boundary condition $u(0) = 0$; (we introduce $\mathcal{S}_\star^{(m)}$ rigorously as the self-adjoint operator associated with the closed, densely defined, non-negative quadratic form $\mathbf{q}_\star^{(m)}$ defined below). We will make this convergence precise in Proposition 2.5, but let us first introduce the two eigenvalues

$$\begin{aligned} \mu_1(\mathcal{S}_b^{(m)}) &= \inf_{u \in D(\mathbf{q}_b^{(m)}) \setminus \{0\}} \frac{\mathbf{q}_b^{(m)}(u)}{\|u\|_{L^2(\mathbb{R}_+)}^2}, \\ \mu_1(\mathcal{S}_\star^{(m)}) &= \inf_{u \in D(\mathbf{q}_\star^{(m)}) \setminus \{0\}} \frac{\mathbf{q}_\star^{(m)}(u)}{\|u\|_{L^2(\mathbb{R}_+)}^2}, \end{aligned} \tag{2.4}$$

where

$$\begin{aligned} \mathbf{q}_b^{(m)}(u) &= \int_{\mathbb{R}_+} (|u'(r)|^2 + w_{m,b}(r)|u(r)|^2) dr + \sqrt{\frac{1}{2b}}|u(0)|^2, \\ \mathbf{q}_\star^{(m)}(u) &= \int_{\mathbb{R}_+} (|u'(r)|^2 + w_{m,0}(r)|u(r)|^2) dr, \end{aligned}$$

and¹

$$\begin{aligned} \text{dom}(\mathbf{q}_b^{(m)}) &= \{u \in H^1(\mathbb{R}_+) : ru \in L^2(\mathbb{R}_+)\}, \\ \text{dom}(\mathbf{q}_\star^{(m)}) &= \{u \in H_0^1(\mathbb{R}_+) : ru \in L^2(\mathbb{R}_+)\}. \end{aligned}$$

We also denote by $\{\mu_k(\mathcal{S}_b^{(m)})\}_{k \geq 1}$ and $\{\mu_k(\mathcal{S}_\star^{(m)})\}_{k \geq 1}$ the sequences of eigenvalues of $\mathcal{S}_b^{(m)}$ and $\mathcal{S}_\star^{(m)}$ respectively, numbered in increasing order including multiplicity. Observe that, the scaling and unitary transformation we did, yield the following

$$\lambda_k^{(m)}(b) = \frac{b}{2}\mu_k(\mathcal{S}_b^{(m)}) \quad (k \geq 1).$$

¹By the Hardy inequality, $\int_{\mathbb{R}_+} r^{-2}|u|^2 dr \leq 4 \int_{\mathbb{R}_+} |u'|^2 dr$ for all $u \in H_0^1(\mathbb{R}_+)$, so we get for free that $u/r \in L^2(\mathbb{R}_+)$.

Therefore, we have as immediate consequence of Proposition 2.1,

$$\forall m \geq 1, \forall b \in (0, 2), \quad \mu_1(\mathcal{S}_b^{(m)}) < 2. \quad (2.5)$$

Proposition 2.5. *For all $m \geq 1$, the operator $\mathcal{S}_b^{(m)}$ converges in the strong resolvent sense to $\mathcal{S}_\star^{(m)}$ as $b \rightarrow 0^+$. Moreover, the eigenvalues of $\mathcal{S}_b^{(m)}$ converge to the corresponding eigenvalues of $\mathcal{S}_\star^{(m)}$ as $b \rightarrow 0^+$.*

Proof. To be able to apply the abstract result [23, Theorem S.14] we need a monotone family of quadratic forms. Let us therefore introduce an intermediate self-adjoint operator $\tilde{\mathcal{S}}_b^{(m)}$ in $L^2(\mathbb{R}_+)$ associated with the closed, semi-bounded, symmetric and densely defined quadratic form

$$\begin{aligned} \tilde{\mathbf{q}}_b^{(m)}(u) &:= \int_{\mathbb{R}_+} (|u'(r)|^2 + w_{m,b}^{\text{int}}(r)|u(r)|^2) dr + \sqrt{\frac{1}{2b}}|u(0)|^2, \\ \text{dom}(\tilde{\mathbf{q}}_b^{(m)}) &:= \{u \in H^1(\mathbb{R}_+) : ru \in L^2(\mathbb{R}_+)\}, \end{aligned}$$

where the potential is defined by

$$\tilde{w}_{m,b}(r) := \frac{4m^2 - 1}{4 \left(r + \sqrt{\frac{b}{2}} \right)^2} + r^2 - 2m.$$

It is straightforward to see that the family of quadratic forms $\mathbb{R}_+ \ni b \mapsto \tilde{\mathbf{q}}_b^{(m)}$ is monotonously decreasing in b in the sense of ordering of the quadratic forms. It follows from [23, Theorem S.14] using the identities

$$D(\mathbf{q}_\star^{(m)}) = \left\{ u \in H^1(\mathbb{R}_+) : ru \in L^2(\mathbb{R}_+), \sup_{b>0} \tilde{\mathbf{q}}_b^{(m)}(u) < \infty \right\}$$

and

$$\lim_{b \rightarrow 0^+} \tilde{\mathbf{q}}_b^{(m)}(u) = \mathbf{q}_\star^{(m)}(u), \quad \text{for all } u \in D(\mathbf{q}_\star^{(m)}),$$

that the operators $\tilde{\mathcal{S}}_b^{(m)}$ converge to $\mathcal{S}_\star^{(m)}$ in the strong resolvent sense as $b \rightarrow 0^+$. Moreover, for any $b \in (0, 8)$ and all $u \in D(\mathbf{q}_b^{(m)}) = D(\tilde{\mathbf{q}}_b^{(m)})$, we have

$$\begin{aligned} \left| \mathbf{q}_b^{(m)}(u) - \tilde{\mathbf{q}}_b^{(m)}(u) \right| &= \int_{\mathbb{R}_+} \left(\sqrt{2b}r + \frac{b}{2} \right) |u(r)|^2 dr \\ &\leq \sqrt{2b} \int_{\mathbb{R}_+} (r+1) |u(r)|^2 dr \\ &\leq 2\sqrt{2b} \left(\tilde{\mathbf{q}}_b^{(m)}(u) + (2m+1) \|u\|_{L^2(\mathbb{R}_+)}^2 \right). \end{aligned}$$

Hence, we get by [18, Theorem VI 3.4] that

$$\|(\mathcal{S}_b^{(m)} + 2m+1)^{-1} - (\tilde{\mathcal{S}}_b^{(m)} + 2m+1)^{-1}\| \rightarrow 0 \quad \text{as } b \rightarrow 0^+. \quad (2.6)$$

It follows from the fact that $\tilde{\mathcal{S}}_b^{(m)}$ converge to $\mathcal{S}_\star^{(m)}$ in the strong resolvent sense together with (2.6) that $\mathcal{S}_b^{(m)}$ also converge to $\mathcal{S}_\star^{(m)}$ in the strong resolvent sense as $b \rightarrow 0^+$.

Note that $\mathcal{S}_b^{(m)} \geq \mathcal{T}^{(m)}$ where the semi-bounded self-adjoint operator $\mathcal{T}^{(m)}$ in $L^2(\mathbb{R}_+)$ with purely discrete spectrum, associated with the form

$$\{u \in H^1(\mathbb{R}_+): ru \in L^2(\mathbb{R}_+)\} \ni u \mapsto \int_{\mathbb{R}_+} (|u'(r)|^2 + (r^2 - 2m)|u(r)|^2) dr.$$

Thus, we get by [26] that the eigenvalues of $\mathcal{S}_b^{(m)}$ converge to the corresponding eigenvalues of $\mathcal{S}_\star^{(m)}$. \square

2.2.3. Spectral gap for the effective operator. Now we turn to calculating the lowest eigenvalue of $\mathcal{S}_\star^{(m)}$ and to estimating the spectral gap between the first and second eigenvalues. We will compare with the Landau Hamiltonian to obtain the gap.

Proposition 2.6. *For all $m \geq 1$, we have $\mu_1(\mathcal{S}_\star^{(m)}) = 2$ and $\mu_2(\mathcal{S}_\star^{(m)}) \geq 6$.*

Note that Proposition 2.6 yields the conclusion in Proposition 2.4, thanks to Proposition 2.5. The proof of Proposition 2.6 relies on the following observation.

Lemma 2.7. *Fix a non-negative integer m and suppose that $g \in \text{dom}(\mathcal{S}_\star^{(m)})$, with $g \neq 0$. Let $h \in L^2(\mathbb{R}^2)$ be defined in polar coordinates as*

$$h = r^{-1/2}g(r)e^{im\theta}.$$

Then,

$$h \in H_{2\mathbf{A}}^1(\mathbb{R}^2) := \{u \in L^2(\mathbb{R}^2): (-i\nabla - 2\mathbf{A})u \in L^2(\mathbb{R}^2)\},$$

and

$$\frac{\langle \mathcal{S}_\star^{(m)}g, g \rangle_{L^2(\mathbb{R}_+)}}{\|g\|_{L^2(\mathbb{R}_+)}^2} = \frac{\|(-i\nabla - 2\mathbf{A})h\|_{L^2(\mathbb{R}^2)}^2}{\|h\|_{L^2(\mathbb{R}^2)}^2}. \quad (2.7)$$

Moreover, if g is orthogonal to u in $L^2(\mathbb{R}_+)$, with $u(r) = r^{m+1/2}e^{-r^2/2}$, then

$$\frac{\|(-i\nabla - 2\mathbf{A})h\|_{L^2(\mathbb{R}^2)}^2}{\|h\|_{L^2(\mathbb{R}^2)}^2} \geq 6. \quad (2.8)$$

Proof. Let $\dot{\Omega} = \mathbb{R}^2 \setminus \{0\}$ and consider the magnetic Sobolev space

$$H_{2\mathbf{A}}^1(\dot{\Omega}) = \{u \in L^2(\dot{\Omega}): (-i\nabla - 2\mathbf{A})u \in L^2(\dot{\Omega})\},$$

where the derivative is seen as a distribution on $\dot{\Omega}$, i.e., $(-i\nabla - 2\mathbf{A})u \in \mathcal{D}'(\dot{\Omega})$. Since $\mathbf{A} \in L_{\text{loc}}^2(\mathbb{R}^2)$, we have that $H_{2\mathbf{A}}^1(\dot{\Omega}) = H_{2\mathbf{A}}^1(\mathbb{R}^2)$ (see [17, Rem. 2.2]).

By a straightforward calculation, we have

$$r^{-1/2}\mathcal{S}_\star^{(m)}r^{1/2} = -\frac{d^2}{dr^2} - \frac{1}{r}\frac{d}{dr} + \left(\frac{m}{r} - r\right)^2$$

and

$$\left(-\frac{d^2}{dr^2} - \frac{1}{r}\frac{d}{dr} + \left(\frac{m}{r} - r\right)^2\right)\psi(r) = e^{-im\theta}(-i\nabla - 2\mathbf{A})^2e^{im\theta}\psi(r).$$

Writing $g = r^{1/2}\psi$, we get $h = e^{im\theta}\psi$, $(-i\nabla - 2\mathbf{A})^2h \in L^2(\dot{\Omega})$, and

$$\begin{aligned} \langle \mathcal{S}_\star^{(m)}g, g \rangle_{L^2(\mathbb{R}_+; dr)} &= \langle r^{-1/2}\mathcal{S}_\star^{(m)}r^{1/2}\psi, \psi \rangle_{L^2(\mathbb{R}_+; r dr)} \\ &= \langle (-i\nabla - 2\mathbf{A})^2h, h \rangle_{L^2(\mathbb{R}_+; r dr)}. \end{aligned}$$

Integrating with respect to θ on $[0, 2\pi)$, we get

$$2\pi \langle \mathcal{S}_\star^{(m)}g, g \rangle_{L^2(\mathbb{R}_+; dr)} = \langle (-i\nabla - 2\mathbf{A})^2h, h \rangle_{L^2(\mathbb{R}^2; dx)} = \|(-i\nabla - 2\mathbf{A})h\|_{L^2(\dot{\Omega})}^2.$$

This proves that $h \in H_{2\mathbf{A}}^1(\dot{\Omega}) = H_{2\mathbf{A}}^1(\mathbb{R}^2)$, and yields the identity in (2.7).

Now, we assume that the orthogonality condition on g holds,

$$\int_{\mathbb{R}_+} g(r)r^{m+\frac{1}{2}}e^{-r^2/2} dr = 0. \quad (2.9)$$

Recall a characterization of the lowest Landau level

$$\begin{aligned} \mathcal{L} &:= \{u \in L^2(\mathbb{R}^2) : (-i\nabla - 2\mathbf{A})^2 u = 2u\} \\ &= \{f(z)e^{-|z|^2/2} : f \text{ is holomorphic in } \mathbb{C}\}, \end{aligned}$$

and that

$$\inf_{\substack{v \in \mathcal{L}^\perp \cap H_{2\mathbf{A}}^1(\mathbb{R}^2) \\ v \neq 0}} \frac{\|(-i\nabla - 2\mathbf{A})v\|_{L^2(\mathbb{R}^2)}^2}{\|v\|_{L^2(\mathbb{R}^2)}} = 6.$$

Therefore, to prove (2.8), it suffices to verify that $h \in \mathcal{L}^\perp$. Let f be a holomorphic function, and notice that, for $r > 0$, we have by Cauchy's integral formula,

$$\int_0^{2\pi} r^{-m} e^{-im\theta} f(re^{i\theta}) d\theta = \frac{1}{i} \int_{C_r} \frac{f(z)}{z^{m+1}} dz = \frac{2\pi}{m!} f^{(m)}(0),$$

where C_r is the circle centred at the origin and of radius r . Consequently, with $u = f(z)e^{-|z|^2/2}$, we have

$$\begin{aligned} \langle u, h \rangle_{L^2(\mathbb{R}^2)} &= \int_0^{+\infty} \left(\int_0^{2\pi} r^{-m} e^{-im\theta} f(re^{i\theta}) d\theta \right) r^{m+\frac{1}{2}} e^{-r^2/2} \overline{g(r)} dr \\ &= \frac{2\pi}{m!} f^{(m)}(0) \int_{\mathbb{R}_+} r^{m+\frac{1}{2}} e^{-r^2/2} \overline{g(r)} dr = 0, \end{aligned}$$

where we used (2.9) in the last step. \square

Proof of Proposition 2.6. It is easy to check that $u = r^{m+\frac{1}{2}} \exp(-r^2/2)$ is an eigenfunction of $\mathcal{S}_*^{(m)}$ with eigenvalue 2, so $\mu_1(\mathcal{S}_*^{(m)}) \leq 2$.

By Lemma 2.7 and the min-max principle, we have $\mu_1(\mathcal{S}_*^{(m)}) \geq 2$, and the function u is a ground state of $\mathcal{S}_*^{(m)}$. If $g \in \text{Dom}(\mathcal{S}_*^{(m)})$ is orthogonal to u , we have by (2.8), $\langle \mathcal{S}_*^{(m)} g, g \rangle_{L^2(\mathbb{R}_+)} \geq 6 \|g\|_{L^2(\mathbb{R}_+)}^2$, and by the min-max principle, we get that $\mu_2(\mathcal{S}_*^{(m)}) \geq 6$. \square

2.3. Refined eigenvalue asymptotics. We improve the leading term asymptotics in Proposition 2.4 by calculating the subleading term.

Theorem 2.8. *Let $m \geq 1$. As $b \rightarrow 0^+$,*

$$\lambda_1^{(m)}(b) = b - \frac{1}{2^{m-1}(m-1)!} b^{m+1} + \mathcal{O}(b^{m+\frac{3}{2}}).$$

Remark 2.9. The $\mathcal{O}(b^{m+\frac{3}{2}})$ can likely be improved to $\mathcal{O}(b^{m+2})$.

Proof of Theorem 2.8. We organize our calculations to use the Temple inequality on $\mathcal{H}^{(m)} - b$, to get both the upper and lower bounds needed. We first work out the sizes of the terms involved. Throughout the proof we use the shorthand notations $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$ for the inner product and the norm in $L^2((1, +\infty); r dr)$, respectively. All order calculations are done in the limit $b \rightarrow 0^+$.

Step 1 (Quasimode). Fix m , and let χ be the function defined as

$$\chi(r) = 1 + \frac{m - b/2}{m + b/2} r^{-2m}.$$

We will define a quasimode Ψ in the domain of $\mathcal{H}^{(m)}$,

$$\Psi(r) = \chi(r)f(r), \quad f(r) = r^m e^{-br^2/4},$$

with

$$\Psi'(1) = [\chi'(1) + (m - b/2)\chi(1)]f(1) = 0.$$

We next show that

$$\|\Psi\|^2 = \frac{2^m m!}{b^{m+1}} + \mathcal{O}(1/b). \quad (2.10)$$

Since Ψ is real-valued, we can skip absolute values, and note that the $\Psi(r)^2$ can be written as a sum of the terms

$$A_1(r) = r^{2m} e^{-br^2/2}, \quad A_2(r) = B_m^2 r^{-2m} e^{-br^2/2}, \quad A_3(r) = 2B_m e^{-br^2/2},$$

where $B_m = (m - b/2)/(m + b/2) = 1 + \mathcal{O}(b)$.

We will see that the integrals of A_2 and A_3 are small when compared to the integral of A_1 . We get

$$\int_1^{+\infty} A_1(r)r \, dr = \int_1^{+\infty} r^{2m+1} e^{-br^2/2} \, dr = \frac{2^m}{b^{m+1}} \Gamma(m+1, b/2) = \frac{2^m m!}{b^{m+1}} + \mathcal{O}(1).$$

For the integral of A_3 , a change of variable yields

$$\int_1^{+\infty} A_3(r)r \, dr = \frac{2B_m}{b} \int_{b/2}^{+\infty} e^{-s} \, ds = \frac{2B_m e^{-b/2}}{b} = \mathcal{O}(1/b),$$

and for the integral of A_2 , we write by Hölder's inequality,

$$\int_1^{+\infty} A_2(r)r \, dr \leq B_m^2 \left(\int_1^{+\infty} r^{-4m} r \, dr \right)^{1/2} \left(\int_1^{+\infty} e^{-br^2} r \, dr \right)^{1/2} = \frac{B_m^2 e^{-b/2}}{\sqrt{4(2m-1)b}}.$$

The claimed size of the norm in (2.10) therefore holds.

Step 2. We next want to estimate the size of $\|(\mathcal{H}^{(m)} - b)\Psi\|$. Observing that

$$\mathcal{H}^{(m)} f = bf, \quad f' = \left(\frac{m}{r} - \frac{br}{2} \right) f,$$

we get

$$(\mathcal{H}^{(m)} - b)\Psi = -f \left(\chi'' + \frac{1+2m}{r} \chi' - br\chi' \right).$$

Now, we note that $\chi'' + \frac{1+2m}{r} \chi' = 0$, hence

$$(\mathcal{H}^{(m)} - b)\Psi = -2bmB_m r^{-m} e^{-br^2/4}, \quad (2.11)$$

and, just as we estimated the integral of A_2 in the previous step, we get eventually

$$\|(\mathcal{H}^{(m)} - b)\Psi\| = \mathcal{O}(b^{\frac{3}{4}}). \quad (2.12)$$

Step 3. Now we estimate the inner product $\langle (\mathcal{H}^{(m)} - b)\Psi, \Psi \rangle$. Thanks to (2.11), we have

$$\langle (\mathcal{H}^{(m)} - b)\Psi, \Psi \rangle = -2bmB_m \int_1^{+\infty} \chi(r) e^{-br^2/2} r \, dr.$$

We write the integral on the right-hand side as the sum of the following two integrals

$$\int_1^{+\infty} e^{-br^2/2} r \, dr = e^{-b/2}/b = \mathcal{O}(1/b)$$

$$B_m \int_1^{+\infty} r^{-2m} e^{-br^2/2} r \, dr = \mathcal{O}(1/\sqrt{b}),$$

and consequently, we end up with

$$\langle (\mathcal{H}^{(m)} - b)\Psi, \Psi \rangle = -2m + \mathcal{O}(\sqrt{b}). \quad (2.13)$$

Step 4 (Temple's inequality). As $b \rightarrow 0^+$, we have a spectral gap, thanks to Proposition 2.4 which yields

$$\lambda_2^{(m)}(b) - \lambda_1^{(m)}(b) \geq 2b + o(b).$$

So we can apply Temple's inequality (see, e.g., [13, Thm. 2]) for the operator $\mathcal{H}^{(m)} - b$,

$$\eta - \frac{\epsilon^2}{\beta - \eta} \leq \lambda_1^{(m)}(b) - b \leq \eta,$$

where

$$\eta = \frac{\langle (\mathcal{H}^{(m)} - b)\Psi, \Psi \rangle}{\|\Psi\|^2} \quad \text{and} \quad \epsilon^2 = \frac{\|(\mathcal{H}^{(m)} - b)\Psi\|^2}{\|\Psi\|^2} - \eta^2,$$

and $\beta = 2b$.

Collecting the estimates in (2.10), (2.11) and (2.13), we get positive constants C_1, C_2 and C_3 , as well as $\hat{b} > 0$ so that for $0 < b < \hat{b}$, it holds that $\beta < \lambda_2^{(m)}(b)$,

$$\frac{b^{m+1}}{(m-1)!2^{m-1}} - C_1 b^{m+\frac{3}{2}} \leq \eta \leq \frac{b^{m+1}}{(m-1)!2^{m-1}} + C_2 b^{m+\frac{3}{2}}$$

and $0 \leq \epsilon^2 \leq C_3 b^{m+\frac{5}{2}}$. The promised asymptotic formula follows. \square

2.4. Ordering of the eigenvalues. Our next aim is to prove the existence of $b_0 > 0$ such that, for all $0 < b < b_0$, the eigenvalues are ordered as

$$\lambda_1^{(1)}(b) < \lambda_1^{(2)}(b) < \lambda_1^{(3)}(b) < \dots < b.$$

To achieve this, we will use the asymptotic expansion of each $\lambda_1^{(m)}(b)$ as $b \rightarrow 0^+$, as well as some information about intersections.

2.4.1. Intersection of band functions. With the asymptotic formula in Theorem 2.8 at hand, we can now with the help of Proposition A.2 find that the first eigenvalues $\lambda_1^{(m)}(b)$ do not intersect for small b .

First we note that the asymptotic formula gives the existence of constants \hat{b}_m such that if $0 < b < \hat{b}_m$ then

$$\lambda_1^{(m-1)}(b) < \lambda_1^{(m)}(b).$$

The next proposition states an explicit bound for \hat{b}_m .

Proposition 2.10. *Assume that $m \geq 2$ and $0 < b \leq 2m + 1 - \sqrt{8m + 1}$. Then*

$$\lambda_1^{(m-1)}(b) < \lambda_1^{(m)}(b).$$

Proof. From Proposition A.2 we know that if $\lambda_1^{(m-1)}(b) = \lambda_1^{(m)}(b) = \lambda < b$ then

$$(b/2 - m)(b/2 - (m - 1)) < b.$$

We recall that both eigenvalues $\lambda_1^{(m-1)}(b)$ and $\lambda_1^{(m)}(b)$ are smaller than b only if $b < 2(m - 1)$. For these values of b the displayed inequality above is equivalent to $b > 2m + 1 - \sqrt{8m + 1}$. Hence, if $b \leq 2m + 1 - \sqrt{8m + 1}$ then $\lambda_1^{(m-1)}(b)$ and $\lambda_1^{(m)}(b)$ cannot be equal, and since by Theorem 2.8 it holds that $\lambda_1^{(m-1)}(b) < \lambda_1^{(m)}(b)$ for small b , this must still be true for all b with $0 < b \leq 2m + 1 - \sqrt{8m + 1}$. \square

Corollary 2.11. *If $0 < b < 5 - \sqrt{17}$ then*

$$\lambda_1^{(1)}(b) < \lambda_1^{(2)}(b) < \lambda_1^{(3)}(b) < \dots < b.$$

Proof. This is immediate from Proposition 2.10 and the fact that $m \mapsto 2m + 1 - \sqrt{8m + 1}$ is increasing for $m \geq 2$. \square

2.4.2. *Uniform convergence of eigenvalues.*

Proposition 2.12. *For $0 < b < 5 - \sqrt{17}$, it holds that $\lambda_1^{(m)}(b) \rightarrow b$ uniformly as $m \rightarrow +\infty$.*

Proof. We know that the eigenvalues obey in this interval

$$\lambda_1^{(1)}(b) < \lambda_1^{(2)}(b) < \lambda_1^{(3)}(b) < \dots < b.$$

The eigenvalues must converge to b pointwise as $m \rightarrow +\infty$ since b is the bottom of the essential spectrum of \mathcal{H} . The convergence is uniform by the Dini theorem. \square

Remark 2.13. The convergence will in fact be uniform on any interval $(0, \hat{b})$, $\hat{b} > 0$, since we have a good control on where the intersections occur.

2.5. **Proof of Theorem 1.1.** Assume that $R = 1$. By (2) in Proposition 2.1, the ground state is non-radial. Moreover, by (1) and (4) in Proposition 2.1, and by Corollary 2.11, we get for $0 < b < 5 - \sqrt{17}$ that

$$\lambda_1(b, \mathcal{B}_1^{\text{ext}}) = \lambda_1^{(1)}(b) < \lambda_2(b, \mathcal{B}_1^{\text{ext}}) = \lambda_2^{(2)}(b) < \dots < b.$$

Finally, the announced two term asymptotics of $\lambda_k(b, \mathcal{B}_1^{\text{ext}})$ follows from Theorem 2.8.

3. OPTIMALITY OF THE DISK'S EXTERIOR

The purpose of this section to prove Theorems 1.3 and 1.5. Recall that $\mathcal{B}_1 \subset \mathbb{R}^2$ denotes the unit disk centred at the origin and that $\mathcal{B}_1^{\text{ext}} := \mathbb{R}^2 \setminus \overline{\mathcal{B}_1}$ is the exterior of \mathcal{B}_1 .

Throughout this section, we suppose that $0 < b < 5 - \sqrt{17}$. Thanks to Corollary 2.11, and Proposition 2.1, we know that

$$\lambda_1(b, \mathcal{B}_1^{\text{ext}}) = \lambda_1^{(1)}(b), \tag{3.1}$$

is the lowest eigenvalue of the fibre operator $\mathcal{H}^{(1)}(b)$ introduced in (2.1). In particular, we know that, as $b \rightarrow 0^+$,

$$\lambda_1(b, \mathcal{B}_1^{\text{ext}}) = b - b^2 + o(b^2). \tag{3.2}$$

This is what we need to compare with for general domains, when addressing optimality of the exterior of the unit disk in the limit $b \rightarrow 0^+$.

3.1. Trial function. The ground state of \mathcal{H} on $\mathcal{B}_1^{\text{ext}}$ can be represented in polar coordinates as

$$u(r, \theta) := f(r)e^{i\theta}, \quad (3.3)$$

where f is the positive normalized eigenfunction of the fibre operator $\mathcal{H}^{(1)}(b)$ corresponding to its lowest eigenvalue $\lambda := \lambda_1^{(1)}(b) < b$. In particular, we know that $f \in L^2((1, +\infty); r dr)$ satisfies the ordinary differential equation

$$-f''(r) - \frac{1}{r}f'(r) + \left(\frac{1}{r} - \frac{br}{2}\right)^2 f(r) = \lambda f(r), \quad r > 1, \quad (3.4)$$

and the boundary condition $f'(1) = 0$. By standard properties of ordinary differential equations, we can construct a C^∞ -smooth extension f to $(0, \infty)$, which satisfies the same differential equation (3.4) for all $r > 0$. This function can be expressed in terms of a confluent hypergeometric function (Whittaker function), see Appendix A. Without any danger of confusion we will denote the respective extension again by f . Let us also introduce the extension of u to $\mathbb{R}^2 \setminus \{0\}$,

$$v(r, \theta) = f(r)e^{i\theta}, \quad r > 0, \theta \in [0, 2\pi). \quad (3.5)$$

It is not hard to see that

$$((-\mathbf{i}\nabla - b\mathbf{A})^2 v)(x) = \lambda v(x), \quad \text{for all } x \in \mathbb{R}^2 \setminus \{0\}. \quad (3.6)$$

3.2. Upper bound for general exterior domains. Using the trial function in (3.5), we can prove an upper bound on the lowest eigenvalue for an exterior domain not containing the origin of \mathbb{R}^2 .

Proposition 3.1. *Suppose that $\Omega^{\text{ext}} = \mathbb{R}^2 \setminus \overline{\Omega}$, where Ω a bounded simply connected domain with a C^∞ -smooth boundary and such that $0 \in \Omega$. If $b < 5 - \sqrt{17}$, then the lowest eigenvalue in Ω^{ext} satisfies*

$$\lambda_1(b, \Omega^{\text{ext}}) \leq \lambda_1(b, \mathcal{B}_1^{\text{ext}}) - \frac{\int_{\partial\Omega} (\nu \cdot \nabla v) \bar{v} \, d\sigma}{\int_{\Omega^{\text{ext}}} |v|^2 \, dx}, \quad (3.7)$$

where v is the function introduced in (3.5), and ν is the outer unit normal for Ω .

Proof. By construction, we have $v|_{\Omega^{\text{ext}}} \in H_{\mathbf{A}}^1(\Omega^{\text{ext}})$. For the sake of brevity we will denote the restriction of v to Ω^{ext} again by v as no confusion can arise. Using the min-max principle and integrating by parts we get

$$\begin{aligned} \lambda_1(b, \Omega^{\text{ext}}) &\leq \frac{\int_{\Omega^{\text{ext}}} |(-\mathbf{i}\nabla - b\mathbf{A})v|^2 \, dx}{\int_{\Omega^{\text{ext}}} |v|^2 \, dx} \\ &= \frac{\int_{\Omega^{\text{ext}}} (-\mathbf{i}\nabla - b\mathbf{A})^2 v \bar{v} \, dx}{\int_{\Omega^{\text{ext}}} |v|^2 \, dx} - \frac{\int_{\partial\Omega} (\nu \cdot (\nabla - \mathbf{i}b\mathbf{A})v) \bar{v} \, d\sigma}{\int_{\Omega^{\text{ext}}} |v|^2 \, dx} \\ &= \lambda_1(b, \mathcal{B}_1^{\text{ext}}) - \frac{\int_{\partial\Omega} (\nu \cdot \nabla v) \bar{v} \, d\sigma}{\int_{\Omega^{\text{ext}}} |v|^2 \, dx}, \end{aligned}$$

where we used in the last step that v satisfies the differential equation in (3.6) for the first term, whereas the magnetic contribution in the boundary term vanishes. Indeed, integrating by parts we get

$$\begin{aligned} \int_{\partial\Omega} (\nu \cdot \mathbf{A})|v|^2 d\sigma &= - \int_{\Omega^{\text{ext}}} \operatorname{div}(\mathbf{A}|v|^2) dx \\ &= - \int_{\Omega^{\text{ext}}} (\operatorname{div} \mathbf{A})|v|^2 dx - \int_{\Omega^{\text{ext}}} \mathbf{A} \cdot \nabla(|v|^2) dx = 0, \end{aligned}$$

where in the last step we used that $\operatorname{div} \mathbf{A} = 0$ and that $\mathbf{A}(x)$ is orthogonal to $\nabla(|v|^2)(x)$ for any $x \in \Omega^{\text{ext}}$, since $|v|$ is radial. \square

3.3. Star-shaped domains. When Ω is star-shaped and contains the origin, we can express the boundary term in (3.7) in terms of the following function

$$F_\alpha(r) := r^\alpha f'(r^\alpha) f(r^\alpha), \quad (\alpha \neq 0, r > 0), \quad (3.8)$$

where f is the function appearing in (3.5). Notice that $F_1(r) = F_\alpha(r^{1/\alpha})$, for any $\alpha \neq 0$.

Proposition 3.2. *Suppose that Ω is defined by polar coordinates as $\Omega = \{(r, \theta) : r < \rho(\theta)\}$, where $\rho: \mathbb{R}/[0, 2\pi] \rightarrow \mathbb{R}_+$ is a C^∞ -smooth function. Then,*

$$\int_{\partial\Omega} (\nu \cdot \nabla v) \bar{v} d\sigma = \int_0^{2\pi} F_1(\rho(\theta)) d\theta, \quad (3.9)$$

where v is the function introduced in (3.5), and F_1 is the function introduced in (3.8).

Proof. We can write the outer unit normal vector field to $\partial\Omega$ as a function of the angular variable (cf. [22, Eq. (3.10)])

$$\nu(\theta) = \frac{1}{\sqrt{\rho^2(\theta) + [\dot{\rho}(\theta)]^2}} (\rho(\theta) \cos \theta + \dot{\rho}(\theta) \sin \theta, \rho(\theta) \sin \theta - \dot{\rho}(\theta) \cos \theta)^\top. \quad (3.10)$$

The gradient of the trial function v in polar coordinates can be written as

$$(\nabla v)(r, \theta) = f'(r) e^{i\theta} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} + \frac{i f(r) e^{i\theta}}{r} \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix}. \quad (3.11)$$

Using (3.10) and (3.11) we can express the boundary term as

$$\begin{aligned} \int_{\partial\Omega} (\nu \cdot \nabla v) \bar{v} d\sigma &= \int_0^{2\pi} \left(\rho(\theta) f'(\rho(\theta)) - i \frac{\dot{\rho}(\theta)}{\rho(\theta)} f(\rho(\theta)) \right) f(\rho(\theta)) d\theta \\ &= \int_0^{2\pi} \rho(\theta) f'(\rho(\theta)) f(\rho(\theta)) d\theta. \quad \square \end{aligned}$$

Thanks to Propositions 3.1 and 3.2, we need to determine the sign of the integral involving F_1 in order to finish the proof of Theorem 1.5. Towards that aim, the next proposition will be handy.

Proposition 3.3. *If $\alpha < 0$, then the function F_α introduced in (3.8) satisfies*

$$F_\alpha''(1) \geq |f(1)|^2 \left[2\alpha^2 \left(\frac{b^2}{2} - 2b \right) - \alpha \left(1 - 2b + \frac{b^2}{4} \right) \right].$$

In particular, if

$$0 < b < 4 - 2\sqrt{3} \quad \text{and} \quad \frac{1 - 2b + \frac{b^2}{4}}{b^2 - 4b} < \alpha < 0,$$

then $F''_\alpha(1) > 0$.

Proof. Differentiating F_α twice, inserting $r = 1$, and using $f'(1) = 0$, we get

$$F''_\alpha(1) = (3\alpha^2 - \alpha)f''(1)f(1) + \alpha^2 f'''(1)f(1). \quad (3.12)$$

From the differential equation (3.4) employing that $f'(1) = 0$ we derive

$$f''(1) = \left[\left(1 - \frac{b}{2}\right)^2 - \lambda \right] f(1). \quad (3.13)$$

Differentiating (3.4) with respect to r and inserting $r = 1$, we get

$$f'''(1) + f''(1) = -2 \left(1 - \frac{b^2}{4}\right) f(1). \quad (3.14)$$

Substituting (3.13) and (3.14) into (3.12)

$$F''_\alpha(1) = |f(1)|^2 \left[(2\alpha^2 - \alpha) \left(1 - b + \frac{b^2}{4} - \lambda\right) - 2\alpha^2 \left(1 - \frac{b^2}{4}\right) \right].$$

For $\alpha < 0$ we get using the inequality $\lambda < b$,

$$F''_\alpha(1) \geq |f(1)|^2 \left[2\alpha^2 \left(\frac{b^2}{2} - 2b\right) - \alpha \left(1 - 2b + \frac{b^2}{4}\right) \right].$$

Writing

$$2\alpha^2 \left(\frac{b^2}{2} - 2b\right) - \alpha \left(1 - 2b + \frac{b^2}{4}\right) = -\alpha \left[1 - 2b + \frac{b^2}{4} - 2\alpha \left(\frac{b^2}{2} - 2b\right)\right]$$

then the conditions $0 < b < 4 - 2\sqrt{3}$ and $\frac{1-2b+\frac{b^2}{4}}{b^2-4b} < \alpha < 0$ yield

$$1 - 2b + \frac{b^2}{4} > 0 \quad \text{and} \quad 1 - 2b + \frac{b^2}{4} - 2\alpha \left(\frac{b^2}{2} - 2b\right) > 0,$$

and consequently $F''_\alpha(1) > 0$. \square

3.4. Proof of Theorem 1.5. We will prove the theorem for the unit radius of the disk ($R = 1$). The general case follows by scaling. The star-shaped domain Ω is parametrized by a smooth mapping $\mathbb{R}/[0, 2\pi] \ni \theta \mapsto \rho(\theta)$ as $\Omega = \{(r, \theta) : r < \rho(\theta)\}$, and in this case the p -moment of Ω can be computed as follows:

$$\mathcal{I}_p(\Omega^{\text{ext}}) = \int_0^{2\pi} \int_{\rho(\theta)}^\infty r^{1-p} dr = \frac{1}{p-2} \int_0^{2\pi} (\rho(\theta))^{2-p} d\theta. \quad (3.15)$$

Recall that we impose the constraint $\mathcal{I}_p(\Omega^{\text{ext}}) = \mathcal{I}_p(\mathcal{B}_1^{\text{ext}})$. Let us set

$$\alpha = \frac{1}{2-p}.$$

By the assumptions on p and b in the formulation of Theorem 1.5, the conditions in Proposition 3.3 are fulfilled, and we have $F''_\alpha(1) > 0$. By smoothness of F_α there exists $\varepsilon \in (0, 1)$ such that $F''_\alpha(r) > 0$ for all $r \in [1 - \varepsilon, 1 + \varepsilon]$, hence F_α is convex on $[1 - \varepsilon, 1 + \varepsilon]$.

Thus, there exists $r_\star \in (0, 1)$ such that $\rho^{1/\alpha} \in (1 - \varepsilon, 1 + \varepsilon)$ provided that $1 - r_\star \leq \rho \leq 1 + r_\star$. Using Jensen's inequality and (3.15) we get

$$\begin{aligned} \int_0^{2\pi} F_1(\rho(\theta)) \, d\theta &= \int_0^{2\pi} F_\alpha((\rho(\theta))^{1/\alpha}) \, d\theta \\ &\geq 2\pi F_\alpha\left(\frac{1}{2\pi} \int_0^{2\pi} (\rho(\theta))^{1/\alpha} \, d\theta\right) \\ &= 2\pi F_\alpha\left(\frac{p-2}{2\pi} \mathcal{I}_p(\Omega^{\text{ext}})\right) \\ &= 2\pi F_\alpha\left(\frac{p-2}{2\pi} \mathcal{I}_p(\mathcal{B}_1^{\text{ext}})\right) \\ &= 2\pi F_\alpha(1) = 0. \end{aligned}$$

Hence, the boundary term in (3.7) is positive by Proposition 3.2, and the desired isoperimetric inequality follows from Proposition 3.1. This finishes the proof of Theorem 1.5 for $R = 1$. The general case follows by scaling.

Remark 3.4. The fixed area constraint corresponds to the choice $\alpha = \frac{1}{2}$ in the above proof of Theorem 1.5 (due to the formula $|\Omega| = \frac{1}{2} \int_0^{2\pi} \rho^2(\theta) \, d\theta$ for star-shaped Ω). In this case, we get $F_{1/2}''(1) = -\frac{1}{2}|f(1)|^2(1 - b^2/4)$ is positive provided that $b > 2$. However, for such magnetic fields the ground state of the magnetic Neumann Laplacian on the exterior of the unit disk does not correspond to the fibre labelled by $m = 1$, and we no more have (3.1). Thus, our method is not directly applicable.

3.5. Proof of Theorem 1.3. We can derive an upper bound when $b \rightarrow 0^+$ and $\Omega = \{(r, \theta) : r < \rho(\theta)\}$ is star-shaped and contains 0, by modifying the trial state in (3.5) as follows

$$u(r, \theta) = \Psi(r)e^{i\theta}, \quad \Psi(r) = (r + cr^{-1})e^{-br^2/4}, \quad (3.16)$$

with some parameter $c \in \mathbb{R}_+$ to be determined at a later stage. The function Ψ is reminiscent of the one appearing in the proof of Theorem 2.8 for $m = 1$. It satisfies the differential equation

$$-\Psi'' - \frac{1}{r}\Psi' + \left(\frac{1}{r} - \frac{br}{2}\right)^2 \Psi - b\Psi = -2bcr^{-1}e^{-br^2/4} \leq 0.$$

Fixing a positive constant ε , similar calculations as in the proof of Theorem 2.8 yield the following asymptotics as $b \rightarrow 0^+$,

$$\begin{aligned} \int_\varepsilon^{+\infty} |\Psi|^2 r \, dr &= \frac{2}{b^2} + \mathcal{O}(1/b), \\ \int_\varepsilon^{+\infty} \left[-\Psi'' - \frac{1}{r}\Psi' + \left(\frac{1}{r} - \frac{br}{2}\right)^2 \Psi - b\Psi\right] \Psi r \, dr &= -2c + \mathcal{O}(\sqrt{b}). \end{aligned}$$

Observing that there are positive constants $r_1 < r_2$ such that

$$\mathcal{B}_{r_1} \subset \Omega \subset \mathcal{B}_{r_2},$$

we get by a straightforward comparison with the integrals over $\mathcal{B}_{r_1}^{\text{ext}}$ and $\mathcal{B}_{r_2}^{\text{ext}}$,

$$\|u\|_{L^2(\Omega^{\text{ext}})}^2 = \frac{4\pi}{b^2} + \mathcal{O}(1/b), \quad \langle [(-i\nabla - b\mathbf{A})^2 - b]u, u \rangle_{L^2(\Omega^{\text{ext}})} = -4\pi c + \mathcal{O}(\sqrt{b}).$$

Arguing as in Propositions 3.1 and 3.2, we get by the min-max principle

$$\begin{aligned}\lambda_1(b, \Omega^{\text{ext}}) &\leq \frac{\langle (-i\nabla - b\mathbf{A})^2 u, u \rangle_{L^2(\Omega^{\text{ext}})}}{\|u\|_{L^2(\Omega^{\text{ext}})}^2} - \frac{\int_{\partial\Omega} \nu \cdot \nabla u \bar{u} \, d\sigma}{\|u\|_{L^2(\Omega^{\text{ext}})}^2} \\ &= b - b^2 c - \frac{b^2}{4\pi} \int_0^{2\pi} \rho(\theta) \Psi'(\rho(\theta)) \Psi(\rho(\theta)) \, d\theta + \mathcal{O}(b^{5/2}).\end{aligned}$$

A routine calculation shows that, as $b \rightarrow 0^+$,

$$r\Psi'(r)\Psi(r) = r^2 - c^2 r^{-2} + \mathcal{O}(b)$$

uniformly with respect to r , as long as r is in a compact interval of \mathbb{R}_+ . Consequently,

$$\int_0^{2\pi} \rho(\theta) \Psi'(\rho(\theta)) \Psi(\rho(\theta)) \, d\theta = 2(|\Omega| - c^2 \mathcal{I}_4(\Omega^{\text{ext}})) + \mathcal{O}(b).$$

We therefore have proved that

$$\lambda_1(b, \Omega^{\text{ext}}) \leq b - \left(c + \frac{|\Omega| - c^2 \mathcal{I}_4(\Omega^{\text{ext}})}{2\pi} \right) b^2 + \mathcal{O}(b^{5/2}). \quad (3.17)$$

Optimizing the bound with respect to the parameter c we easily get that the optimal value is $c = \frac{\pi}{2\mathcal{I}_4(\Omega^{\text{ext}})}$ and we obtain the following:

$$\lambda_1(b, \Omega^{\text{ext}}) \leq b - \mathcal{G}(\Omega) b^2 + \mathcal{O}(b^{5/2}), \quad (3.18)$$

where

$$\mathcal{G}(\Omega) = \frac{|\Omega|}{2\pi} + \frac{\pi}{2\mathcal{I}_4(\Omega^{\text{ext}})}.$$

This bound is consistent with the conclusion of Theorem 1.1, since $\mathcal{G}(\Omega) = R^2$ if $\Omega = \mathcal{B}_R$ is the disk of radius R .

In the remaining part of the proof we assume that Ω is not a disk centred at the origin. By Hölder's inequality,

$$2\pi < \left(\int_0^{2\pi} \rho^2 \, d\theta \right)^{1/2} \left(\int_0^{2\pi} \rho^{-2} \, d\theta \right)^{1/2},$$

which reads as $\pi^2 < |\Omega| \mathcal{I}_4(\Omega^{\text{ext}})$, Thus, we always have

$$\frac{\pi}{\mathcal{I}_4(\Omega^{\text{ext}})} < \mathcal{G}(\Omega) < \frac{|\Omega|}{\pi}.$$

In particular, if we fix R_\star such that $\mathcal{I}_4(\mathcal{B}_{R_\star}^{\text{ext}}) = \mathcal{I}_4(\Omega^{\text{ext}})$, we have

$$\mathcal{G}(\Omega) > \mathcal{G}(\mathcal{B}_{R_\star}),$$

and we get from (3.18) and Theorem 1.1 that $\lambda_1(b, \Omega^{\text{ext}}) < \lambda_1(b, \mathcal{B}_{R_\star}^{\text{ext}})$, for b sufficiently small, by which the proof of Theorem 1.3 is complete.

Remark 3.5. However, if we fix the area by choosing R so that $|\mathcal{B}_R| = |\Omega|$, then $\mathcal{G}(\Omega) < \mathcal{G}(\mathcal{B}_R)$ and (3.18) is not sufficient to compare $\lambda_1(b, \Omega^{\text{ext}})$ and $\lambda_1(b, \mathcal{B}_R^{\text{ext}})$.

Acknowledgments. The first listed author (A. K.) is partially supported by The Chinese University of Hong Kong, Shenzhen (grant UDF01003322). The second listed author (V. L.) is supported by the Czech Science Foundation (GAČR) within the project 21-07129S.

We initiated this work when the first listed author visited Lund University and the Nuclear Physics Institute, Czech Academy of Sciences, Řež, Czech Republic. We acknowledge financial support through Knut and Alice Wallenberg Foundation (grant KAW 2021.0259).

APPENDIX A. INTERSECTION OF EIGENVALUES

A.1. Introduction. We consider here the self-adjoint realization of the operator

$$\mathcal{H}^{(m)}(b) = -\frac{d^2}{dr^2} - \frac{1}{r} \frac{d}{dr} + \left(\frac{m}{r} - \frac{br}{2}\right)^2, \quad (\text{A.1})$$

either in $L^2((0, 1), r dr)$ or $L^2((1, +\infty), r dr)$, with Neumann condition at $r = 1$. We denote in both cases by $\lambda_1^{(m)}(b)$ the lowest eigenvalue of $\mathcal{H}^{(m)}(b)$.

A.2. The interior of the disk. We start by working in the interior of the disk. The following result is old, but not so well-known, since it is in the original paper only mentioned as a comment, without a proof.

Theorem A.1 ([24]). *Assume that $b > 0$ and $m \geq 1$. If $\lambda_1^{(m-1)}(b) = \lambda_1^{(m)}(b) < b$ then $\lambda_1^{(m)}(b) = (b/2 - m)[b/2 - (m - 1)]$.*

The condition that the eigenvalues should be less than b is merely to enforce the eigenvalues to be the bottom ones, and to be sure that they are not eigenvalues of the Dirichlet operator.

Proof. The result relies on several identities for confluent hypergeometric functions, we will use the so-called Whittaker functions.

The Whittaker differential equation reads

$$\frac{d^2 y}{dz^2} + \left(-\frac{1}{4} + \frac{\kappa}{z} + \frac{1/4 - \mu^2}{z^2}\right)y = 0. \quad (\text{A.2})$$

It has two linearly independent solutions, denoted by $M(\kappa, \mu, z)$ (well-defined if 2μ is not a negative integer, which it will not be in our case) and $W(\kappa, \mu, z)$. According to [6, (13.14.14)],

$$M(\kappa, \mu, z) = z^{\mu+1/2}(1 + O(z)) \quad (z \rightarrow 0). \quad (\text{A.3})$$

With a change of variables $z = br^2/2$ and with a scaling by r of the independent function, one finds that the general solution to $\mathcal{H}^{(m)}(b)u = \lambda u$ is given by

$$u(r) = c_1 \frac{1}{r} M\left(\Lambda, \frac{m}{2}, \frac{br^2}{2}\right) + c_2 \frac{1}{r} W\left(\Lambda, \frac{m}{2}, \frac{br^2}{2}\right) \quad (\text{A.4})$$

For notational convenience in the continuation, we have introduced the constant

$$\Lambda = \frac{mb + \lambda}{2b}.$$

Since M is bounded at 0 according to (A.3), and since the differential equation has a regular singularity at the origin, W becomes singular as $r \rightarrow 0^+$. We are forced to take $c_2 = 0$. Let us also set $c_1 = 1$, and consider below the eigenfunction

$$u(r) = \frac{1}{r} M\left(\Lambda, \frac{m}{2}, \frac{br^2}{2}\right). \quad (\text{A.5})$$

The Whittaker function M satisfies [6, (13.15.15)]

$$\frac{d}{dz}M(\kappa, \mu, z) = \left(\frac{1}{2} - \frac{\kappa}{z}\right)M(\kappa, \mu, z) + \frac{1 + 2\kappa + 2\mu}{2z}M(\kappa + 1, \mu, z). \quad (\text{A.6})$$

Implementing this, we find that $u'(1) = 0$ precisely when

$$(b^2 - 2b(m+1) - 2\lambda)M\left(\Lambda, \frac{m}{2}, \frac{b}{2}\right) + 2(b + 2bm + \lambda)M\left(\Lambda + 1, \frac{m}{2}, \frac{b}{2}\right) = 0. \quad (\text{A.7})$$

Replacing m by $m - 1$ we get

$$\begin{aligned} & (b^2 - 2bm - 2\lambda)M\left(\Lambda - \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) \\ & + 2(-b + 2bm + \lambda)M\left(\Lambda + \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) = 0. \end{aligned} \quad (\text{A.8})$$

We note that the third arguments in the Whittaker functions in (A.7) and (A.8) are the same. We will ‘massage’ the first equation (A.7) in order to have Whittaker functions with the same arguments in both equations.

In the first step we will make sure that the second arguments become the same. To do this, we will use formula [6, (13.15.4)], which says that

$$M(\kappa, \mu, z) = \frac{2\mu}{\sqrt{z}} \left(M\left(\kappa - \frac{1}{2}, \mu - \frac{1}{2}, z\right) - M\left(\kappa + \frac{1}{2}, \mu - \frac{1}{2}, z\right) \right). \quad (\text{A.9})$$

Implementing this in (A.7), the equation transforms into (here we have skipped the non-zero factor $m/\sqrt{b/2}$ from both terms)

$$\begin{aligned} & (b^2 - 2b(m+1) - 2\lambda) \left[M\left(\Lambda - \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) - M\left(\Lambda + \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) \right] \\ & + 2(b + 2bm + \lambda) \left[M\left(\Lambda + \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) - M\left(\Lambda + \frac{3}{2}, \frac{m-1}{2}, \frac{b}{2}\right) \right] = 0. \end{aligned} \quad (\text{A.10})$$

We need a formula that makes the first argument comparable as well. To this aim, we will use the formula [6, (13.15.1)],

$$\left(\kappa + \mu + \frac{1}{2}\right)M(\kappa + 1, \mu, z) = \left(\mu - \kappa + \frac{1}{2}\right)M(\kappa - 1, \mu, z) + (2\kappa - z)M(\kappa, \mu, z). \quad (\text{A.11})$$

We will soon implement it in (A.10), with

$$\kappa = \Lambda + \frac{1}{2}, \quad \mu = \frac{m-1}{2}, \quad z = \frac{b}{2}.$$

This gives

$$\mu + \kappa + \frac{1}{2} = \Lambda + \frac{m+1}{2}, \quad \mu - \kappa + \frac{1}{2} = \frac{m-1}{2} - \Lambda, \quad 2\kappa - z = 2\Lambda + 1 - \frac{b}{2},$$

and

$$\begin{aligned} \left(\Lambda + \frac{m+1}{2}\right)M\left(\Lambda + \frac{3}{2}, \frac{m-1}{2}, \frac{b}{2}\right) &= \left(\frac{m-1}{2} - \Lambda\right)M\left(\Lambda - \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) \\ &+ \left(2\Lambda + 1 - \frac{b}{2}\right)M\left(\Lambda + \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right). \end{aligned}$$

We are now ready to insert this into (A.10),

$$\begin{aligned}
& (b^2 - 2b(m+1) - 2\lambda) \left(\Lambda + \frac{m+1}{2} \right) \\
& \quad \times \left[\mathbf{M}\left(\Lambda - \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) - \mathbf{M}\left(\Lambda + \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) \right] \\
& \quad + 2(b + 2bm + \lambda) \left(\Lambda + \frac{m+1}{2} \right) \mathbf{M}\left(\Lambda + \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) \\
& \quad - 2(b + 2bm + \lambda) \left(\frac{m-1}{2} - \Lambda \right) \mathbf{M}\left(\Lambda - \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) \\
& \quad - 2(b + 2bm + \lambda) \left(2\Lambda + 1 - \frac{b}{2} \right) \mathbf{M}\left(\Lambda + \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) = 0.
\end{aligned} \tag{A.12}$$

Collecting the two different Whittaker expressions, and cancelling the non-zero common factor $(b + 2bm + \lambda)/2$, we find that

$$(b - 2m) \mathbf{M}\left(\Lambda - \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) + (b + 2m) \mathbf{M}\left(\Lambda + \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) = 0. \tag{A.13}$$

Now assume that we have eigenvalues $\lambda_1^{(m)}(b) = \lambda_1^{(m-1)}(b) = \lambda$. This means that we have

$$\begin{bmatrix} b^2 - 2bm - 2\lambda & 2(-b + 2bm + \lambda) \\ b - 2m & b + 2m \end{bmatrix} \begin{bmatrix} \mathbf{M}\left(\Lambda - \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) \\ \mathbf{M}\left(\Lambda + \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

If it would hold that $\mathbf{M}\left(\Lambda - \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) = 0$, it would according to (A.5) mean that λ is also an eigenvalue of the Dirichlet realization of $\mathcal{H}_{m-1}(b)$. Since the smallest Dirichlet eigenvalue is greater than b , and we assume that $\lambda < b$, this is not possible. Thus, the determinant of the coefficient matrix must be zero,

$$\begin{aligned}
0 &= \det \begin{bmatrix} b^2 - 2bm - 2\lambda & 2(-b + 2bm + \lambda) \\ b - 2m & b + 2m \end{bmatrix} \\
&= -4b \left[\lambda - \left(\frac{b}{2} - m \right) \left(\frac{b}{2} - (m-1) \right) \right].
\end{aligned}$$

We conclude that if $\lambda < b$ is a common eigenvalue of $\mathcal{H}^{(m)}(b)$ and $\mathcal{H}^{(m-1)}(b)$, then

$$\lambda = \left(\frac{b}{2} - m \right) \left(\frac{b}{2} - (m-1) \right). \quad \square$$

The fact that it is the lowest eigenvalues of $\mathcal{H}^{(m-1)}(b)$ and $\mathcal{H}^{(m)}(b)$ that intersect follows from the upper bound $\lambda < b$. The property in the theorem, however, seems also to be true for higher eigenvalues of $\mathcal{H}^{(m-1)}(b)$ and $\mathcal{H}^{(m)}(b)$.

A.3. The exterior of the disk. For the exterior the situation is similar. We use again the same notation as above, but with the operator now acting in $L^2((1, +\infty), r dr)$. Here, the other Whittaker function W is used. Similar formulas exist for it. In fact, the formula (A.6) is replaced by

$$\frac{d}{dz} W(\kappa, \mu, z) = \left(\frac{1}{2} - \frac{\kappa}{z} \right) W(\kappa, \mu, z) - \frac{1}{z} W(\kappa + 1, \mu, z),$$

and then the formula (A.7) is replaced by

$$\left(\frac{\lambda}{b} + m + 1 - \frac{b}{2}\right) \mathbb{W}\left(\Lambda, \frac{m}{2}, \frac{b}{2}\right) + 2\mathbb{W}\left(\Lambda + 1, \frac{m}{2}, \frac{b}{2}\right) = 0,$$

and the formula (A.8) is replaced by

$$\left(\frac{\lambda}{b} + m - \frac{b}{2}\right) \mathbb{W}\left(\Lambda - \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) + 2\mathbb{W}\left(\Lambda + \frac{1}{2}, \frac{m-1}{2}, \frac{b}{2}\right) = 0.$$

The formula (A.9) is replaced by

$$\mathbb{W}(\kappa, \mu, z) = \frac{1}{\sqrt{z}} \left[\mathbb{W}\left(\kappa + \frac{1}{2}, \mu - \frac{1}{2}, z\right) + \left(\kappa + \mu - \frac{1}{2}\right) \mathbb{W}\left(\kappa - \frac{1}{2}, \mu - \frac{1}{2}, z\right) \right],$$

and the formula (A.11) is replaced by

$$\mathbb{W}(\kappa + 1, \mu, z) + (2\kappa - z)\mathbb{W}(\kappa, \mu, z) + \left(\kappa - \mu - \frac{1}{2}\right) \left(\kappa + \mu - \frac{1}{2}\right) \mathbb{W}(\kappa - 1, \mu, z) = 0.$$

Once that is done, the reasoning is similar, and in fact exactly the same formula as for the interior of the disk holds.

Proposition A.2. *Assume that $b > 0$ and $m \geq 2$. If $\lambda_1^{(m-1)}(b) = \lambda_1^{(m)}(b) < b$ then $\lambda_1^{(m)}(b) = (b/2 - m)(b/2 - (m - 1))$.*

REFERENCES

- [1] C. Bandle and A. Wagner. *Shape optimization. Variations of domains and applications*, volume 42 of *De Gruyter Ser. Nonlinear Anal. Appl.* Berlin: De Gruyter, 2023.
- [2] L. Bundrock. Geometric optimization of the first Robin eigenvalue in exterior domains. *arXiv preprint*, 2404.10886, 2024.
- [3] B. Colbois, C. Léna, L. Provenzano, and A. Savo. Geometric bounds for the magnetic Neumann eigenvalues in the plane. *J. Math. Pures Appl. (9)*, 179:454–497, 2023.
- [4] B. Colbois, C. Léna, L. Provenzano, and A. Savo. A reverse Faber–Krahn inequality for the magnetic Laplacian. *arXiv preprint*, 2403.11336, 2024.
- [5] C. Dietze, A. Kachmar, and V. Lotoreichik. Isoperimetric inequalities for inner parallel curves. *arXiv preprint*, 2311.18413, 2023.
- [6] *NIST Digital Library of Mathematical Functions*. <https://dlmf.nist.gov/>, Release 1.2.0 of 2024-03-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [7] L. Erdős. Rayleigh-type isoperimetric inequality with a homogeneous magnetic field. *Calc. Var. Partial Differential Equations*, 4(3):283–292, 1996.
- [8] G. Faber. Beweis, daß unter allen homogenen Membranen von gleicher Fläche und gleicher Spannung die kreisförmige den tiefsten Grundton gibt. *Math.-Nat. Kl. Bayer. Akad. Wiss.*, pages 169–172, 1923.
- [9] S. Fournais and B. Helffer. *Spectral methods in surface superconductivity*, volume 77 of *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser Boston, Inc., Boston, MA, 2010.
- [10] S. Fournais and B. Helffer. Inequalities for the lowest magnetic Neumann eigenvalue. *Lett. Math. Phys.*, 109(7):1683–1700, 2019.
- [11] R. Ghanta, L. Junge, and L. Morin. Quantitative magnetic isoperimetric inequality. *J. Spectr. Theory*, 14(1):185–205, 2024.
- [12] M. Goffeng, A. Kachmar, and M. Persson Sundqvist. Clusters of eigenvalues for the magnetic Laplacian with Robin condition. *J. Math. Phys.*, 57(6):063510, 19, 2016.
- [13] E. M. Harrell. Generalizations of Temple’s inequality. *Proc. Am. Math. Soc.*, 69:271–276, 1978.
- [14] B. Helffer and M. Persson Sundqvist. On the semi-classical analysis of the ground state energy of the Dirichlet Pauli operator. *J. Math. Anal. Appl.*, 449(1):138–153, 2017.

- [15] A. Kachmar and V. Lotoreichik. On the isoperimetric inequality for the magnetic Robin Laplacian with negative boundary parameter. *J. Geom. Anal.*, 32(6):20, 2022. Id/No 182.
- [16] A. Kachmar and V. Lotoreichik. A geometric bound on the lowest magnetic Neumann eigenvalue via the torsion function. *To appear in SIAM J. Math. Anal.*, *arXiv:2312.06161*, 2024.
- [17] A. Kachmar and X.B. Pan. Oscillatory patterns in the Ginzburg-Landau model driven by the Aharonov-Bohm potential. *J. Funct. Anal.*, 279(10):108718, 37, 2020.
- [18] T. Kato. *Perturbation theory for linear operators*. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the 1980 edition.
- [19] E. Krahn. Über eine von Rayleigh formulierte Minimaleigenschaft des Kreises. *Math. Ann.*, 94(1):97–100, 1925.
- [20] D. Krejčířík and V. Lotoreichik. Optimisation of the lowest Robin eigenvalue in the exterior of a compact set. *J. Convex Anal.*, 25(1):319–337, 2018.
- [21] D. Krejčířík and V. Lotoreichik. Optimisation of the lowest Robin eigenvalue in the exterior of a compact set, II: Non-convex domains and higher dimensions. *Potential Anal.*, 52(4):601–614, 2020.
- [22] D. Krejčířík and V. Lotoreichik. Optimisation and monotonicity of the second Robin eigenvalue on a planar exterior domain. *arXiv preprint*, 2307.14286, 2023.
- [23] M. Reed and B. Simon. *Methods of modern mathematical physics. I*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, second edition, 1980. Functional analysis.
- [24] D. Saint-James. Etude du champ critique H_{c_3} dans une geometrie cylindrique. *Physics Letters A*, 15(1):13–15, 1965.
- [25] G. Szegő. Inequalities for certain eigenvalues of a membrane of given area. *J. Rational Mech. Anal.*, 3:343–356, 1954.
- [26] J. Weidmann. Continuity of the eigenvalues of selfadjoint operators with respect to the strong operator topology. *Integral Equations Operator Theory*, 3(1):138–142, 1980.
- [27] H. F. Weinberger. An isoperimetric inequality for the N -dimensional free membrane problem. *J. Rational Mech. Anal.*, 5:633–636, 1956.

(A. Kachmar) THE CHINESE UNIVERSITY OF HONG KONG, SHENZHEN, GUANGDONG, 518172, P.R. CHINA.

Email address: akachmar@cuhk.edu.cn

(V. Lotoreichik) DEPARTMENT OF THEORETICAL PHYSICS, NUCLEAR PHYSICS INSTITUTE, CZECH ACADEMY OF SCIENCES, 25068, ŘEŽ, CZECH REPUBLIC

Email address: lotoreichik@ujf.cas.cz

(M. Sundqvist) DEPARTMENT OF MATHEMATICS, LUND UNIVERSITY, SWEDEN

Email address: mikael.persson.sundqvist@math.lth.se