

# Letter frequency vs factor frequency in pure morphic words

Shuo Li

Department of Mathematics & Statistics, The University of Winnipeg,  
Winnipeg, Canada.

**Abstract.** We prove that, for any pure morphic word  $w$ , if the frequencies of all letters in  $w$  exist, then the frequencies of all factors in  $w$  exist as well. This result answers a question of Saari in his doctoral thesis.

## 1 Introduction

Let  $A$  be a finite alphabet and let  $w = w[1]w[2]w[3]\cdots$  be an infinite word over  $A$ . Let  $v$  be a factor of  $w$  of length  $k$ , the (*ordinary*) frequency of  $v$  in  $w$  is defined as

$$\lim_{n \rightarrow \infty} \frac{\#\{i \mid w[i]w[i+1]\cdots w[i+k-1] = v, 1 \leq i \leq n\}}{n},$$

if the limit exists. For the factors such that  $k = 1$ , the previous frequency is called the (*ordinary*) frequency of letters. The frequency of letters in morphic words has been studied since the early 1970's. It was proved by Cobham in 1972 [3] that the frequency of a letter in an automatic word, if it exists, is rational. Michel [4,5] proved that the frequencies of all letters exist in primitive morphic words. Peter [6] gave a sufficient and necessary condition for the existence of the frequency of a letter in an automatic word. Saari proved in [7] that the frequencies of both letters exist in any pure morphic binary word and gave a sufficient and necessary condition for the existence of the frequency of a letter in a morphic sequence over an arbitrary alphabet [8]. For any infinite word  $w$  and any factor  $v$  of length  $k$  of  $w$ , one can also define the *logarithmic frequency* of  $v$  in  $w$  as

$$\lim_{n \rightarrow \infty} \frac{1}{\log(n)} \sum_{\{i \mid w[i]w[i+1]\cdots w[i+k-1] = v, 1 \leq i \leq n\}} \frac{1}{i},$$

if this limit exists. It is proved in [1, Proposition 8.4.4] that if the frequency of a letter  $a$  in  $w$  exists, then the logarithmic frequency of  $a$  in  $w$  exists as well, and these two frequencies are equal. Allouche and Shallit asked in [1, Section 8.8] if the logarithmic frequency of any letter in a morphic word must exist and Bell gave a positive answer to this question in [2] and generalized it to all factors. Concerning the relation between the frequency of letters and the frequency of factors, Saari proved in his doctoral thesis [9, Proposition 3.1] that, for an arbitrary morphic word  $w$ , the existence of the frequencies of all letters in  $w$  cannot imply the existence of the frequencies of all factors in  $w$  and asked if it

is true for pure morphic words [9, Problem 3.1]. We establish a positive answer to this question.

**Theorem 1** *For any pure morphic word  $w$ , if the frequencies of all letters in  $w$  exist, then the frequencies of all factors in  $w$  exist as well.*

## 2 Definitions and notation

Let  $A$  be a finite set. It will be called an *alphabet* and its elements will be called *letters*. Let  $A^*$  denote the free monoid generated by  $A$  under concatenations having neutral element the empty word  $\varepsilon$ . The elements in  $A^*$  are called the *finite words* with letters in  $A$ . For any finite word  $w = w[1]w[2]w[3] \cdots w[n] \in A^*$ , the *length* of  $w$  is the integer  $|w| = n$ . Let  $A^{\mathbb{N}}$  be the set of infinite concatenations of elements in  $A$ . The elements in  $A^{\mathbb{N}}$  are called the *infinite words* with letters in  $A$ . For any infinite word  $w$ , the length of  $w$ , which is also denoted by  $|w|$ , is infinite. Let  $A^\infty = A^* \cup A^{\mathbb{N}}$ . For any  $w \in A^\infty$  and  $v \in A^*$ , let  $|w|_v$  denote the number of occurrences of  $v$  in  $w$ .

Let  $w = w[1]w[2]w[3] \cdots$  be an element in  $A^\infty$  and let  $v \in A^*$ . We say  $v$  a *prefix* of  $w$  if there exists an integer  $t$  such that  $1 \leq t \leq |w|$  and  $v = w[1]w[2]w[3] \cdots w[t]$  and we say  $v$  a *factor* of  $w$  if there exists a pair of integers  $t, r$  such that  $1 \leq t \leq r \leq |w|$  and  $v = w[t]w[t+1]w[t+2] \cdots w[r]$ . A prefix  $v$  of  $w$  is called *proper* if  $|v| < |w|$ . For any pair of integers  $t, r$  such that  $0 \leq t \leq r \leq |w|$ , let  $w[t, r] = w[t]w[t+1]w[t+2] \cdots w[r]$ .

Let  $A$  and  $B$  be two alphabets. A *morphism*  $\phi$  is a map  $A^* \rightarrow B^*$  satisfying  $\phi(xy) = \phi(x)\phi(y)$  for any pair of elements  $x, y$  in  $A^*$ . The morphism  $\phi$  is called  *$k$ -uniform* for some positive integer  $k$  if for all elements  $a \in A$ ,  $|\phi(a)| = k$ , and it is called *non-uniform* otherwise. A morphism  $\phi$  is called a *coding function* if it is 1-uniform, and it is called *non-erasing* if  $\phi(a) \neq \varepsilon$  for all  $a \in A$ . For any positive integer  $k$ , by  $\phi^k$ , we mean the  $k$ -fold composition of the morphism  $\phi$ . Let  $u \in A^*$  and  $v \in B^*$ , if  $\phi(u) = v$ , then  $v$  is called the *image* of  $u$  and  $u$  is the *pre-image* of  $v$  under  $\phi$ . We write  $u = \phi^{-1}(v)$ . A morphism  $\phi : A^* \rightarrow A^*$  is called *primitive* if there exists an integer  $n \geq 1$  such that for all  $a, b \in A$ ,  $a$  occurs in  $\phi^n(b)$ .

Let  $A$  be an alphabet and let  $\phi : A^* \rightarrow A^*$  be a morphism. A letter  $a \in A$  is called *bounded* if  $|\phi^k(a)|$  is upper bounded by some constant  $C$  for all positive integers  $k$  and it is called *unbounded* otherwise.

Let  $A$  be a finite alphabet, and let  $w$  be an infinite word over  $A$ .  $w$  is called *morphic* if there exists an alphabet  $B$ , a letter  $b \in B$ , a nonempty word  $v \in B^*$ , a non-erasing morphism  $\phi : B^* \rightarrow B^*$ , and a coding function  $\psi : B \rightarrow A$ , such that  $\phi(b) = bv$  and

$$w = \lim_{i \rightarrow \infty} \psi(\phi^i(b)).$$

Moreover, for any positive integer  $k \geq 2$ , the word  $w$  is called  *$k$ -automatic* if  $\phi$  is  $k$ -uniform, it is called *automatic* if  $\phi$  is  $k$ -uniform for some integer  $k \geq 2$ , and it is called *non-automatic* if  $\phi$  is not  $k$ -uniform for any integer  $k \geq 2$ . The word

$w$  is called *pure morphic* if  $A = B$  and  $\psi = Id$  and it is called *primitive* if  $\phi$  is primitive. Remark that if  $w$  is pure morphic, then  $w = \phi(w)$ .

### 3 Proof of the main theorem

Let  $A$  be an alphabet,  $\phi : A^* \rightarrow A^*$  be a non-erasing morphism and  $w$  be a pure morphic word such that  $w = \phi(w)$ . Let  $A_B = \{a \mid a \in A, a \text{ is bounded}\}$  and let  $A_U = \{a \mid a \in A, a \text{ is unbounded}\}$ . Obviously,  $A_B \cup A_U = A$  and  $A_B \cap A_U = \emptyset$ . If all letters in  $A$  have a frequency, then for any  $a \in A$ , let  $\alpha_a = \lim_{n \rightarrow \infty} \frac{|w[1,n]|_a}{n}$  and let  $\alpha = \sum_{a \in A_B} \alpha_a$ . Let  $k_1$  be an integer such that  $|\phi^n(a)| < k_1$  for all  $a \in A_B$  and for all  $n \in \mathbb{N}^+$ .

The idea of the proof is that, for any pure morphic word  $w$ , any factor  $v$  of  $w$  and any prefix  $w[1, m]$  of  $w$  with  $m$  large enough, supposing that there exists the word  $w[1, m''] = \phi^{-k}(w[1, m])$  for some positive integer  $k$ , then one can prove that, when  $k$  is large enough,  $|w[1, m]|_v$  can be approximated by the summation of  $|\phi^k(w[i])|_v$  for all unbounded letters  $w[i]$ ,  $1 \leq i \leq m''$ . Thus, the frequency of  $v$  can be estimated in terms of  $\alpha_a$ ,  $|\phi^k(a)|$  and  $|\phi^k(a)|_v$ , for all  $a \in A_U$ .

The strategy of the proof is as follow: supposing that there exists some factor  $v$  of  $w$  such that  $\limsup \frac{|w[1,n]|_v}{n} - \liminf \frac{|w[1,n]|_v}{n} = \Delta > 0$ , then one can find a real number  $C$ , independent from  $\Delta$ , such that  $\left| \frac{|w[1,n]|_v}{n} - C \right| < \frac{1}{3}\Delta$  when  $n$  is large, which leads to a contradiction.

*Proof (of Theorem 1).* Let  $w$  be a pure morphic word satisfying the hypothesis in Theorem 1 and suppose that there exists a factor  $v$  of  $w$  of length  $L$  such that  $\limsup \frac{|w[1,n]|_v}{n} - \liminf \frac{|w[1,n]|_v}{n} = \Delta > 0$ . Let  $M \in \mathbb{N}$  such that for all  $a \in A_U$

$$|\phi^M(a)| \geq \max \left\{ \frac{6(L + \alpha k_1)}{(1 - \alpha)\Delta}, \frac{k_1 \alpha (6 - \Delta)}{(1 - \alpha)\Delta} \right\}.$$

Let  $k_2$  be an integer such that  $|\phi^M(a)| < k_2$  for all  $a \in A$ .

For any  $m \in \mathbb{N}^+$ , let  $m'$  be the largest integer smaller than or equal to  $m$  such that  $w[1, m']$  is the image of a prefix of  $w$  under  $\phi^M$ . Thus, there exists  $m'' \in \mathbb{N}^+$  such that

$$w[1, m'] = \phi^M(w[1, m'']). \quad (1)$$

From the definition of  $m'$ , if  $w[m' + 1, m]$  is not empty, then it is a proper prefix of  $\phi^M(a)$  for some  $a \in A$ . Thus,

$$m' \leq m < m' + k_2, \quad (2)$$

$$|w[1, m']|_v \leq |w[1, m]|_v < |w[1, m']|_v + k_2. \quad (3)$$

One can estimate  $m'$  as well as  $|w[1, m']|_v$  in terms of  $m''$ .

$$m' = \sum_{a \in A_U} |\phi^M(a)| \cdot |w[1, m'']|_a + \epsilon_1, \quad (4)$$

where

$$\epsilon_1 = \sum_{a \in A_B} |\phi^M(a)| \cdot |w[1, m'']|_a. \quad (5)$$

Since  $|\phi^M(a)| < k_1$  for all  $a \in A_B$  and  $\sum_{a \in A_B} |w[1, m'']|_a = \alpha m'' + o(m'')$ , one has

$$0 \leq \epsilon_1 \leq k_1 m'' \alpha + o(m''). \quad (6)$$

Similarly,

$$|w[1, m']|_v = |\phi^M(w[1, m''])|_v = \sum_{a \in A_U} |\phi^M(a)|_v \cdot |w[1, m'']|_a + \epsilon_2 + \epsilon_3, \quad (7)$$

where

$$\epsilon_2 = \sum_{a \in A_B} |\phi^M(a)|_v \cdot |w[1, m'']|_a. \quad (8)$$

and  $\epsilon_3$  is the number of occurrences of  $v$  which begin in a factor  $\phi^M(w[t])$  for some  $t$  but do not end in the same factor. For any bounded letter  $a$ ,  $|\phi^M(a)|_v \leq |\phi^M(a)| < k_1$ . Thus,

$$0 \leq \epsilon_2 \leq k_1 m'' \alpha + o(m''). \quad (9)$$

For any integer  $i$ ,  $1 \leq i \leq m''$ , the number of occurrences of  $v$  which begin in  $\phi^M(w[i])$  but do not end in the same factor is upper-bounded by the length of  $v$ , which is  $L$ . Thus,

$$0 \leq \epsilon_3 \leq L m'', \quad (10)$$

One can estimate  $\frac{|w[1, m]|_v}{m}$  in terms of  $m''$  using Equations 4, 7.

$$\frac{|w[1, m]|_v}{m} = \frac{\sum_{a \in A_U} |\phi^M(a)|_v \cdot |w[1, m'']|_a + \epsilon_2 + \epsilon_3 + \epsilon_4}{\sum_{a \in A_U} |\phi^M(a)| \cdot |w[1, m'']|_a + \epsilon_1 + \epsilon_5}, \quad (11)$$

where  $\epsilon_4 = |w[1, m]|_v - |w[1, m']|_v$  and  $\epsilon_5 = m - m'$ . From Equations 2, 3,

$$0 \leq \epsilon_4, \epsilon_5 < k_2 = o(m''). \quad (12)$$

To simplify the notation, let

$$S_1(m) = \sum_{a \in A_U} |\phi^M(a)|_v \cdot |w[1, m'']|_a, \quad S_2(m) = \sum_{a \in A_U} |\phi^M(a)| \cdot |w[1, m'']|_a.$$

Since

$$\frac{S_1(m)}{S_2(m)} = \frac{\sum_{a \in A_U} |\phi^M(a)|_v \cdot \frac{|w[1, m'']|_a}{m''}}{\sum_{a \in A_U} |\phi^M(a)| \cdot \frac{|w[1, m'']|_a}{m''}} \quad (13)$$

and  $\lim_{m \rightarrow \infty} \frac{|w[1, m'']|_a}{m''} = \alpha_a$  for all  $a \in A_U$ , one has

$$\lim_{m \rightarrow \infty} \frac{S_1(m)}{S_2(m)} = \frac{\sum_{a \in A_U} |\phi^M(a)|_v \alpha_a}{\sum_{a \in A_U} |\phi^M(a)| \alpha_a} < 1. \quad (14)$$

$$\text{Let } C_{v,M} = \frac{\sum_{a \in A_U} |\phi^M(a)|_v \alpha_a}{\sum_{a \in A_U} |\phi^M(a)|_{\alpha_a}}.$$

To upper bound  $\frac{|w[1,m]|_v}{m}$ , take  $\epsilon_1 = \epsilon_5 = 0$ ,  $\epsilon_2 = k_1 m'' \alpha + o(m'')$ ,  $\epsilon_3 = L m''$  and  $\epsilon_4 = o(m'')$  using Equations 6, 8, 10, 12. One has

$$\frac{|w[1,m]|_v}{m} = \frac{S_1(m) + \epsilon_2 + \epsilon_3 + \epsilon_4}{S_2(m) + \epsilon_1 + \epsilon_5} \leq \frac{S_1(m)}{S_2(m)} + \frac{(k_1 \alpha + L)m'' + o(m'')}{S_2(m)}. \quad (15)$$

Since  $|\phi^M(a)| \geq \frac{6(L+\alpha k_1)}{(1-\alpha)\Delta}$  for all  $a \in A_U$ ,

$$\begin{aligned} \frac{(k_1 \alpha + L)m'' + o(m'')}{S_2(m)} &= \frac{(k_1 \alpha + L)m'' + o(m'')}{\sum_{a \in A_U} |\phi^M(a)| \cdot |w[1,m'']|_a} \\ &\leq \frac{(k_1 \alpha + L)m'' + o(m'')}{\frac{6(L+\alpha k_1)}{(1-\alpha)\Delta} ((1-\alpha)m'' + o(m''))} \\ &\leq \frac{\Delta}{6} + o(1). \end{aligned} \quad (16)$$

Thus, combining Equations 15, 16,

$$\frac{|w[1,m]|_v}{m} \leq \frac{S_1(m)}{S_2(m)} + \frac{\Delta}{6} + o(1). \quad (17)$$

From Equation 14, There exists an integer  $N_1$  such that for all  $m \geq N_1$ ,

$$\frac{|w[1,m]|_v}{m} < C_{v,M} + \frac{\Delta}{6} + \frac{\Delta}{6} = C_{v,M} + \frac{\Delta}{3}. \quad (18)$$

To lower bound  $\frac{|w[1,m]|_v}{m}$ , take  $\epsilon_1 = k_1 m'' \alpha + o(m'')$ ,  $\epsilon_5 = o(m'')$  and  $\epsilon_2 = \epsilon_3 = \epsilon_4 = 0$  using Equations 6, 8, 10, 12. One has

$$\frac{|w[1,m]|_v}{m} = \frac{S_1(m) + \epsilon_2 + \epsilon_3 + \epsilon_4}{S_2(m) + \epsilon_1 + \epsilon_5} \geq \frac{S_1(m)}{S_2(m) + k_1 m'' \alpha + o(m'')} \quad (19)$$

Since  $|\phi^M(a)| \geq \frac{k_1 \alpha (6-\Delta)}{(1-\alpha)\Delta}$  for all  $a \in A_U$ ,

$$\begin{aligned} \frac{|w[1,m]|_v}{m} &\geq \frac{S_1(m)}{S_2(m) + k_1 m'' \alpha + o(m'')} \\ &\geq \frac{S_1(m)}{S_2(m)} \frac{1}{1 + \frac{k_1 m'' \alpha + o(m'')}{S_2(m)}} \\ &\geq \frac{S_1(m)}{S_2(m)} \frac{1}{1 + \frac{k_1 m'' \alpha + o(m'')}{\frac{k_1 \alpha (6-\Delta)}{(1-\alpha)\Delta} ((1-\alpha)m'' + o(m''))}} \\ &\geq \frac{S_1(m)}{S_2(m)} \left(1 - \frac{\Delta}{6}\right) + o(1) \\ &\geq \frac{S_1(m)}{S_2(m)} - \frac{\Delta}{6} + o(1). \end{aligned} \quad (20)$$

The last inequality is from the fact that  $\frac{S_1(m)}{S_2(m)} < 1$  for all  $m$ . From Equation 14, there exists an integer  $N_2$  such that for all  $m \geq N_2$ ,

$$\frac{|w[1, m]|_v}{m} > C_{v, M} - \frac{\Delta}{6} - \frac{\Delta}{6} = C_{v, M} - \frac{\Delta}{3}. \quad (21)$$

In conclusion, let  $N = \max\{N_1, N_2\}$ , then for any  $m \geq N$ ,  $\left| \frac{|w[1, m]|_v}{m} - C_{v, M} \right| < \frac{\Delta}{3}$  which is a contradiction to the hypothesis that  $\limsup \frac{|w[1, n]|_v}{n} - \liminf \frac{|w[1, n]|_v}{n} = \Delta > 0$ . Thus, all factors in  $w$  have a frequency.

**Corollary 2** *Let  $w$  be a pure morphic word over the alphabet  $A$  such that each element in  $A$  has a frequency in  $w$ . If  $\phi$  is the morphism such that  $\phi(w) = w$ , then for any  $v \in A^*$ , with the same notation as above,*

$$\lim_{m \rightarrow \infty} \frac{|w[1, m]|_v}{m} = \lim_{M \rightarrow \infty} C_{v, M}. \quad (22)$$

*Proof.* Once it is proved that each factor  $v$  of  $w$  has a frequency, the argument in the proof of Theorem 1 shows that, for any  $\Delta > 0$ , there exists  $M \in \mathbb{N}^+$  such that for all  $m \geq M$ , there exists  $N_m \in \mathbb{N}^+$  such that for all  $n \geq N_m$ ,

$$\left| C_{v, m} - \frac{|w[1, n]|_v}{n} \right| < \frac{\Delta}{3}. \quad (23)$$

See Equations 21, 18. On the other hand, letting  $\alpha_v$  be the frequency of  $v$  in  $w$ , there exists  $N'_m \in \mathbb{N}^+$  such that for all  $n \geq N'_m$ ,

$$\left| \alpha_v - \frac{|w[1, n]|_v}{n} \right| < \frac{\Delta}{3}. \quad (24)$$

Thus, for  $n$  large,

$$|\alpha_v - C_{v, m}| < \frac{2\Delta}{3}. \quad (25)$$

However, there is no term in Equation 25 involving the variable  $n$ , thus, Equation 25 holds independently from the choice of  $n$ . Consequently,

$$\lim_{M \rightarrow \infty} C_{v, M} = \alpha_v. \quad (26)$$

**Remark 3** *In the proof of Theorem 1, the existence of the frequency of the factor  $v$  is not established in a direct way. The reason is that once Equation 23 is established, it is not trivial to prove that the sequence  $(C_{v, m})_{m \in \mathbb{N}^+}$  converges. However, when it is proved that the frequency of  $v$  exists, then the limit of the above sequence exists as well.*

## 4 Conclusion

The author concludes this note by asking the following question:

**Question 4** *For any pure morphic word  $w$  and any factor  $v$  of  $w$ , is it possible to prove the existence of the frequency of  $v$  and directly compute the value using the incidence matrix of  $w$ ?*

## References

1. Allouche, J.-P., Shallit, J.: *Automatic Sequences: Theory, Applications, Generalizations*. Cambridge University Press (2003)
2. Bell, J.P.: Logarithmic frequency in morphic sequences. *Journal de Théorie des Nombres de Bordeaux* 20(2), 227–241 (2008)
3. Cobham, A.: Uniform tag sequences. *Mathematical Systems Theory* 6, 164–192 (1972)
4. Michel, P.: *Sur les ensembles minimaux engendrés par les substitutions de longueur non constante*. Ph.D. Thesis, Université de Rennes (1975)
5. Michel, P.: Stricte ergodicité d'ensembles minimaux de substitution. In: Conze, J.P., Keane, M.S. (eds.) *Théorie Ergodique: : Actes des Journées Ergodiques, Rennes 1973/1974*. pp. 189–201. Springer Berlin Heidelberg, Berlin, Heidelberg (1976)
6. Peter, M.: The asymptotic distribution of elements in automatic sequences. *Theoretical Computer Science* 301(1), 285–312 (2003)
7. Saari, K.: On the frequency of letters in pure binary morphic sequences. In: De Felice, C., Restivo, A. (eds.) *Developments in Language Theory*. pp. 397–408. Springer Berlin Heidelberg, Berlin, Heidelberg (2005)
8. Saari, K.: On the frequency of letters in morphic sequences. In: Grigoriev, D., Harrison, J., Hirsch, E.A. (eds.) *Computer Science – Theory and Applications*. pp. 334–345. Springer Berlin Heidelberg, Berlin, Heidelberg (2006)
9. Saari, K.: *On the frequency and periodicity of infinite words*. Ph.D. Thesis, University of Turku (2008)