

# LOCAL SUPER-DERIVATIONS OF THE SUPER SCHRÖDINGER ALGEBRAS

ALAUADINOV A.K., YUSUPOV B.B.

ABSTRACT. The present paper is devoted to study local super-derivations of the super Schrödinger algebras. We prove that every local super-derivation on the super Schrödinger algebra is a super-derivation.

## 1. INTRODUCTION

This notion was introduced and investigated independently by R.V. Kadison [18] and D.R. Larson and A.R. Sourour [21]. The above papers gave rise to a series of works devoted to the description of mappings which are close to automorphisms and derivations of  $C^*$ -algebras and operator algebras. R.V. Kadison set out a program of study for local maps in [18], suggesting that local derivations could prove useful in building derivations with particular properties. R.V. Kadison proved in [18, Theorem A] that each continuous local derivation of a von Neumann algebra  $M$  into a dual Banach  $M$ -bimodule is a derivation. This theorem gave way to studies on derivations on  $C^*$ -algebras, culminating with a result due to B.E. Johnson, which asserts that every local derivation of a  $C^*$ -algebra  $A$  into a Banach  $A$ -bimodule is automatically continuous, and hence is a derivation [16, Theorem 5.3].

Let us present a list of finite or infinite dimensional algebras for which all local derivations are derivations:

- $C^*$ -algebras, in particular, the algebra  $M_n(\mathbb{C})$  of all square matrices of order  $n$  over the field of complex numbers [16, 18];
- the complex polynomial algebra  $\mathbb{C}[x]$  [18];
- finite dimensional simple Lie algebras over an algebraically closed field of characteristic zero [6];
- Borel subalgebras of finite-dimensional simple Lie algebras [24];
- infinite dimensional Witt algebras over an algebraically closed field of characteristic zero [11];
- Witt algebras over a field of prime characteristic [25];
- solvable Lie algebras of maximal rank [22];
- Cayley algebras [4];
- finite dimensional semi-simple Leibniz algebras over an algebraically closed field of characteristic zero [23];
- locally finite split simple Lie algebras over a field of characteristic zero [9];
- the Schrödinger algebras [2];
- Lie superalgebra  $q(n)$  [10];
- conformal Galilei algebras [3].

---

2020 *Mathematics Subject Classification.* 08A35, 17A32, 17A36 .

*Key words and phrases.* super Schrödinger algebras, super-derivations, local super-derivations.

On the other hand, some algebras (in most cases close to nilpotent algebras) admit pure local derivations, that is, local derivations which are not derivations. Below a short list of some classes of algebras which admit pure local derivations:

- the algebra  $\mathbb{C}(x)$  of rational functions [18];
- finite dimensional filiform Lie algebras [6];
- $p$ -filiform Leibniz algebras [8];
- solvable Leibniz algebras with abelian nilradicals, which have a one dimensional complementary space [5];
- the algebra of lower triangular  $n \times n$ -matrices [13];
- the ternary Malcev algebra  $M_8$  [14];
- direct sum null-filiform Leibniz algebras [1].

The Schrödinger Lie group describes symmetries of the free particle Schrödinger equation in [20]. The Lie algebra  $\mathbb{S}(n)$  in  $(n + 1)$ -dimensional space-time of the Schrödinger Lie group is called the Schrödinger algebra, see [17]. The Schrödinger algebra is a non-semisimple Lie algebra and plays an important role in mathematical physics. The Lie algebra  $\mathbb{S}(1)$  is one of the most essential case for  $n = 1$  and admits a universal 1-dimensional central extension which is called the centrally extended Schrödinger algebra or, simply, the Schrödinger algebra, abusing the language.

The present paper is devoted to study local super-derivations on the super Schrödinger algebra.

## 2. PRELIMINARIES

In this section we recall definitions, symbols and establish some auxiliary results for later use in this paper.

We begin by providing some conventions in this paper. Denote the degree for  $\mathbb{Z}_2$ -graded vector spaces  $\mathcal{L} = \mathcal{L}_{\bar{0}} \oplus \mathcal{L}_{\bar{1}}$  by

$$|x| = \begin{cases} 0, & \text{if } x \in \mathcal{L}_{\bar{0}}; \\ 1, & \text{if } x \in \mathcal{L}_{\bar{1}}. \end{cases}$$

Elements in  $\mathcal{L}_{\bar{0}}$  or  $\mathcal{L}_{\bar{1}}$  are called homogeneous. We implicitly assume that  $x$  is as a  $\mathbb{Z}_2$ -homogeneous element when  $|x|$  occurs in some expression. A linear map  $g : \mathcal{L} \rightarrow \mathcal{L}$  is a homogeneous linear map of degree  $\alpha \in \mathbb{Z}_2$ , i.e.  $|g| = \alpha$ , provided that

$$g(\mathcal{L}_{\beta}) \subseteq \mathcal{L}_{\alpha+\beta}, \quad \forall \beta \in \mathbb{Z}_2.$$

Similarly, a bilinear map  $\varphi : \mathcal{L} \times \mathcal{L} \rightarrow \mathcal{L}$  is a homogeneous bilinear map of degree  $\alpha \in \mathbb{Z}_2$ , i.e.  $|\varphi| = \alpha$  if it satisfies that

$$\varphi(\mathcal{L}_{\beta}, \mathcal{L}_{\gamma}) \subseteq \mathcal{L}_{\alpha+\beta+\gamma}, \quad \forall \beta, \gamma \in \mathbb{Z}_2.$$

Moreover, if  $|g|$  or  $|\varphi|$  occurs in some expression, we regard it as a homogeneous map.

A Lie superalgebra (see [19]) is a superspace  $\mathcal{L} = \mathcal{L}_{\bar{0}} \oplus \mathcal{L}_{\bar{1}}$  with a bilinear mapping  $[\cdot, \cdot] : \mathcal{L} \times \mathcal{L} \rightarrow \mathcal{L}$  satisfying the following identities:

- (1)  $[\mathcal{L}_{\alpha}, \mathcal{L}_{\beta}] \subseteq \mathcal{L}_{\alpha+\beta}$ , for  $\alpha, \beta \in \mathbb{Z}_2$  ( $\mathbb{Z}_2$ -grading),
- (2)  $[x, y] = -(-1)^{|x||y|}[y, x]$  (graded skew-symmetry),
- (3)  $(-1)^{|x||z|}[x, [y, z]] + (-1)^{|y||x|}[y, [z, x]] + (-1)^{|z||y|}[z, [x, y]] = 0$  (graded Jacobi identity),

for all  $x, y, z \in \mathcal{L}$ . It is clear that the graded Jacobi identity is equivalent to the equation

$$[x, [y, z]] = [[x, y], z] + (-1)^{|z||x+y|} [[z, x], y]. \quad (1)$$

**Definition 2.1.** [12] The  $N = 1$  super Schrödinger algebra in  $(1 + 1)$ -dimensional spacetime  $\mathcal{S} = \mathcal{S}_0 \oplus \mathcal{S}_1$  is a 9-dimensional Lie superalgebra with the even part  $\mathcal{S}_0 = \text{span}_{\mathbb{C}}\{e, f, h, p, q, z\}$  and the odd part  $\mathcal{S}_1 = \text{span}_{\mathbb{C}}\{E, F, G\}$  satisfying the following non-vanishing Lie super brackets:

$$\begin{aligned} [h, e] &= 2e, & [h, f] &= -2f, & [e, f] &= h, \\ [h, p] &= p, & [h, q] &= -q, & [p, q] &= z, \\ [e, q] &= p, & [p, f] &= -q, & [h, E] &= E, \\ [h, F] &= -F, & [e, F] &= -E, & [f, E] &= -F, \\ [p, F] &= G, & [q, E] &= G, & [E, E] &= 2e, \\ [F, F] &= -2f, & [G, G] &= z, & [E, F] &= -h, \\ [E, G] &= -p, & [F, G] &= q. \end{aligned}$$

**Remark 1.** (1) The even part  $\mathcal{S}_0 = \text{span}_{\mathbb{C}}\{e, f, h, p, q, z\}$  is the Schrodinger algebra in  $(1 + 1)$ -dimensional spacetime.

(2) It's easy to see that  $\mathcal{S}$  contains two subalgebras: the super Heisenberg subalgebra  $H = \text{span}_{\mathbb{C}}\{p, q, z, G\}$  and the Lie superalgebra  $osp(1|2)$  spanned by  $\{h, e, f, E, F\}$ . In particular, the super Heisenberg subalgebra  $H$  is also an ideal of  $\mathcal{S}$ . Then,  $\mathcal{S}$  can be viewed as a semidirect product  $\mathcal{S} = H \rtimes osp(1|2)$ .

**Definition 2.2.** Let  $\mathcal{L}$  be a Lie superalgebra. A linear map  $D : \mathcal{L} \rightarrow \mathcal{L}$  is called a super-derivation of  $\mathcal{L}$  if

$$D([x, y]) = [D(x), y] + (-1)^{|D||x|} [x, D(y)], \quad \forall x, y \in \mathcal{L}.$$

Write  $Der_{\alpha}(\mathcal{L})$  for the set of all homogeneous super-derivations of degree  $\alpha \in \mathbb{Z}_2$  of  $\mathcal{L}$ . Then  $Der(\mathcal{L}) = Der_0(\mathcal{L}) \oplus Der_1(\mathcal{L})$ . For arbitrary  $x \in \mathcal{L}$ , it is easy to see that the map  $ad_x : \mathcal{L} \rightarrow \mathcal{L}$  defined by  $ad_x(y) = [x, y]$  for all  $y \in \mathcal{L}$  is a super-derivation of  $\mathcal{L}$ , which is termed an inner super-derivation. The set of all inner super-derivations is denoted by  $IDer(\mathcal{L})$ .

**Definition 2.3.** A linear operator  $\Delta$  is called a local super-derivation if for any  $x \in \mathcal{L}$ , there exists a super-derivation  $D_x : \mathcal{L} \rightarrow \mathcal{L}$  (depending on  $x$ ) such that  $\Delta(x) = D_x(x)$ . The set of all local super-derivations on  $\mathcal{L}$  we denote by  $LocDer(\mathcal{L})$ .

The following theorem gives a description of the super-derivation on the super Schrödinger algebra  $\mathcal{S}$ .

**Theorem 2.4.** [15]

$$\text{Der}(\mathcal{S}) = \text{Ider}(\mathcal{S}) \oplus \mathbb{C}\delta,$$

where  $\delta$  are linear transformations of  $\mathcal{S}$  and satisfy that

$$\begin{aligned} \delta|_{osp(1|2)} &= 0, & \delta(p) &= p, \\ \delta(q) &= q, & \delta(z) &= 2z, & \delta(G) &= G. \end{aligned}$$

### 3. LOCAL SUPER-DERIVATIONS OF THE SUPER SCHRÖDINGER ALGEBRA $\mathcal{S}$

In this section we investigate local super-derivations on the super Schrödinger algebra  $\mathcal{S}$ . The following theorem is the main result of this section.

**Theorem 3.1.** *Every local super-derivations on  $\mathcal{S}$  is a super-derivation.*

For the proof of this theorem we need several lemmas.

For any local super-derivations  $\Delta$  on  $\mathcal{S}$  we have by Theorem 2.4

$$\Delta(x) = [b_x, x] + \lambda_x \delta(x).$$

Put

$$b_x = b_{e,x}e + b_{f,x}f + b_{h,x}h + b_{p,x}p + b_{q,x}q + b_{z,x}z + b_{E,x}E + b_{F,x}F + b_{G,x}G \in \mathcal{S}.$$

Then we obtain

$$\begin{aligned} \Delta(e) &= 2b_{h,e}e - b_{f,e}h - b_{q,e}p + b_{F,e}E, \\ \Delta(f) &= -2b_{h,f}f + b_{e,f}h - b_{p,f}q + b_{E,f}F, \\ \Delta(h) &= -2b_{e,h}e + 2b_{f,h}f - b_{p,h}p + b_{q,h}q - b_{E,h}E + b_{F,h}F, \\ \Delta(p) &= (b_{h,p} + \lambda_p)p + b_{f,p}q - b_{q,p}z - b_{F,p}G, \\ \Delta(q) &= b_{e,q}p + (-b_{h,q} + \lambda_q)q + b_{p,q}z - b_{E,q}G, \\ \Delta(z) &= 2b_{z,z}z, \\ \Delta(E) &= 2b_{E,E}e - b_{F,E}h + b_{G,E}p + b_{h,E}E - b_{f,E}F + b_{q,E}G, \\ \Delta(F) &= -2b_{F,F}f + b_{E,F}h - b_{G,F}q - b_{e,F}E - b_{h,F}F + b_{p,F}G, \\ \Delta(G) &= -b_{E,G}p + b_{F,G}q + b_{G,G}z + \lambda_G G. \end{aligned}$$

For the proof of this Theorem we need several Lemmata.

There exists an element  $a = a_{e,x}e + a_{f,x}f + a_{h,x}h + a_{p,x}p + a_{q,x}q + a_{z,x}z + a_{E,x}E + a_{F,x}F + a_{G,x}G \in \mathcal{S}$  such that

$$\Delta(x) = [a, x] + \mu_x \delta(x).$$

In the following lemmas, we assume that  $\Delta$  be a local super-derivation on  $\mathcal{S}$  such that  $\Delta(h + z) = 0$ .

**Lemma 3.2.**  $\Delta(h) = \Delta(z) = 0$ .

*Proof.* We considering,

$$0 = \Delta(h + z) = \Delta(h) + \Delta(z) = -2b_{e,h}e + 2b_{f,h}f - b_{p,h}p + b_{q,h}q - b_{E,h}E + b_{F,h}F + 2b_{z,z}z,$$

which implies  $b_{e,h} = b_{f,h} = b_{p,h} = b_{q,h} = b_{E,h} = b_{F,h} = b_{z,z} = 0$ . Then  $\Delta(h) = \Delta(z) = 0$ .  $\square$

**Lemma 3.3.**

$$\begin{aligned} \Delta(e) &= 2b_{h,e}e - b_{q,e}p + b_{F,e}E, \\ \Delta(f) &= -2b_{h,f}f - b_{p,f}q + b_{E,f}F. \end{aligned}$$

*Proof.* We consider

$$\begin{aligned}
\Delta(h+e) &= [a, h+e] + \mu_{h+e}\delta(h+e) = \\
&= [a_{e,h+e}e + a_{f,h+e}f + a_{h,h+e}h + a_{p,h+e}p + a_{q,h+e}q + a_{z,h+e}z + \\
&\quad + a_{E,h+e}E + a_{F,h+e}F + a_{G,h+e}G, h+e] + \mu_{h+e}\delta(h+e) = \\
&= -2a_{e,h+e}e + 2a_{f,h+e}f - a_{p,h+e}p + a_{q,h+e}q - a_{E,h+e}E + a_{F,h+e}F + \\
&\quad + 2a_{h,h+e}e - a_{f,h+e}h - a_{q,h+e}p + a_{F,h+e}E.
\end{aligned}$$

On the other hand,

$$\Delta(h+e) = \Delta(h) + \Delta(e) = 2b_{h,e}e - b_{f,e}h - b_{q,e}p + b_{F,e}E.$$

Comparing the coefficients at the basis elements  $f$  and  $h$ , we get  $b_{f,e} = 0$ . Similarly, considering

$$\begin{aligned}
\Delta(h+f) &= [a, h+f] + \mu_{h+f}\delta(h+f) = \\
&= [a_{e,h+f}e + a_{f,h+f}f + a_{h,h+f}h + a_{p,h+f}p + a_{q,h+f}q + a_{z,h+f}z + \\
&\quad + a_{E,h+f}E + a_{F,h+f}F + a_{G,h+f}G, h+f] + \mu_{h+f}\delta(h+f) = \\
&= -2a_{e,h+f}e + 2a_{f,h+f}f - a_{p,h+f}p + a_{q,h+f}q - a_{E,h+f}E + a_{F,h+f}F \\
&\quad - 2a_{h,h+f}f + a_{e,h+f}h - a_{p,h+f}q + a_{E,h+f}F.
\end{aligned}$$

On the other hand,

$$\Delta(h+f) = \Delta(h) + \Delta(f) = -2b_{h,f}f + b_{e,f}h - b_{p,f}q + b_{E,f}F.$$

Comparing the coefficients at the basis elements  $e$  and  $h$ , we get  $b_{e,f} = 0$ . □

**Lemma 3.4.**

$$\begin{aligned}
\Delta(e) &= 2b_{h,e}e - b_{q,e}p, \\
\Delta(f) &= -2b_{h,f}f - b_{p,f}q, \\
\Delta(p) &= (b_{h,p} + \lambda_p)p - b_{q,p}z, \\
\Delta(q) &= (-b_{h,q} + \lambda_q)q + b_{p,q}z.
\end{aligned}$$

*Proof.* We consider

$$\begin{aligned}
\Delta(h+p) &= [a, h+p] + \mu_{h+p}\delta(h+p) = \\
&= [a_{e,h+p}e + a_{f,h+p}f + a_{h,h+p}h + a_{p,h+p}p + a_{q,h+p}q + \\
&\quad + a_{z,h+p}z + a_{E,h+p}E + a_{F,h+p}F + a_{G,h+p}G, h+p] + \mu_{h+p}\delta(h+p) = \\
&= -2a_{e,h+p}e + 2a_{f,h+p}f - a_{p,h+p}p + a_{q,h+p}q - a_{E,h+p}E + a_{F,h+p}F + \\
&\quad + (a_{h,h+p} + \mu_{h+p})p + a_{f,h+p}q - a_{q,h+p}z - a_{F,h+p}G.
\end{aligned}$$

On the other hand,

$$\Delta(h+p) = \Delta(h) + \Delta(p) = (b_{h,p} + \lambda_p)p + b_{f,p}q - b_{q,p}z - b_{F,p}G.$$

Comparing the coefficients at the basis elements  $F$  and  $G$ , we get  $b_{F,p} = 0$ .

We consider

$$\begin{aligned}
\Delta(h+q) &= [a, h+q] + \mu_{h+q}\delta(h+q) = \\
&= [a_{e,h+q}e + a_{f,h+q}f + a_{h,h+q}h + a_{p,h+q}p + a_{q,h+q}q + \\
&\quad + a_{z,h+q}z + a_{E,h+q}E + a_{F,h+q}F + a_{G,h+q}G, h+q] + \mu_{h+q}\delta(h+q) = \\
&= -2a_{e,h+q}e + 2a_{f,h+q}f - a_{p,h+q}p + a_{q,h+q}q - a_{E,h+q}E + a_{F,h+q}F + \\
&\quad + a_{e,h+q}p + (-a_{h,h+q} + \mu_{h+q})q + a_{p,h+q}z - a_{E,h+q}G.
\end{aligned}$$

On the other hand,

$$\Delta(h+q) = \Delta(h) + \Delta(q) = b_{e,q}p + (-b_{h,q} + \lambda_q)q + b_{p,q}z - b_{E,q}G.$$

Comparing the coefficients at the basis elements  $E$  and  $G$ , we get  $b_{E,q} = 0$ .

We consider

$$\begin{aligned}
\Delta(e+p) &= [a, e+p] + \mu_{e+p}\delta(e+p) = \\
&= [a_{e,e+p}e + a_{f,e+p}f + a_{h,e+p}h + a_{p,e+p}p + a_{q,e+p}q + \\
&\quad + a_{z,e+p}z + a_{E,e+p}E + a_{F,e+p}F + a_{G,e+p}G, e+p] + \mu_{e+p}\delta(e+p) = \\
&= 2a_{h,e+p}e - a_{f,e+p}h - a_{q,e+p}p + a_{F,e+p}E + \\
&\quad + (a_{h,e+p} + \lambda_{e+p})p + a_{f,e+p}q - a_{q,e+p}z - a_{F,e+p}G.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
\Delta(e+p) &= \Delta(e) + \Delta(p) = \\
&= 2b_{h,e}e - b_{q,e}p + b_{F,e}E + (b_{h,p} + \lambda_p)p + b_{f,p}q - b_{q,p}z.
\end{aligned}$$

Comparing the coefficients at the basis elements  $h$ ,  $q$ ,  $E$ , and  $G$ , we get  $b_{F,e} = b_{f,p} = 0$ .

We consider

$$\begin{aligned}
\Delta(f+q) &= [a, f+q] + \mu_{f+q}\delta(f+q) = \\
&= [a_{e,f+q}e + a_{f,f+q}f + a_{h,f+q}h + a_{p,f+q}p + a_{q,f+q}q + \\
&\quad + a_{z,f+q}z + a_{E,f+q}E + a_{F,f+q}F + a_{G,f+q}G, f+q] + \mu_{f+q}\delta(f+q) = \\
&= -2a_{h,f+q}f + a_{e,f+q}h - a_{p,f+q}q + a_{E,f+q}F + \\
&\quad + a_{e,f+q}p + (-a_{h,f+q} + \lambda_{f+q})q + a_{p,f+q}z - a_{E,f+q}G.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
\Delta(f+q) &= \Delta(f) + \Delta(q) = \\
&= -2b_{h,f}f - b_{p,f}q + b_{E,f}F + b_{e,q}p + (-b_{h,q} + \lambda_q)q + b_{p,q}z.
\end{aligned}$$

Comparing the coefficients at the basis elements  $h$ ,  $p$ ,  $F$  and  $G$ , we get  $b_{E,f} = b_{e,q} = 0$ .  $\square$

**Lemma 3.5.**

$$\begin{aligned}
\Delta(e) &= 2b_{h,e}e - b_{q,e}p, \\
\Delta(f) &= -2b_{h,f}f - b_{q,e}q, \\
\Delta(E) &= 2b_{E,E}e - b_{F,E}h + b_{G,E}p + b_{h,E}E - b_{f,E}F, \\
\Delta(F) &= -2b_{F,F}f + b_{E,F}h - b_{G,E}q - b_{e,F}E - b_{h,F}F, \\
\Delta(G) &= -b_{E,F}p + b_{F,E}q + b_{G,G}z + \lambda_G G.
\end{aligned}$$

*Proof.* We consider

$$\begin{aligned}
\Delta(h + E) &= [a, h + E] + \mu_{h+E}\delta(h + E) = \\
&= [a_{e,h+E}e + a_{f,h+E}f + a_{h,h+E}h + a_{p,h+E}p + a_{q,h+E}q + a_{z,h+E}z + \\
&\quad + a_{E,h+E}E + a_{F,h+E}F + a_{G,h+E}G, h + E] + \mu_{h+E}\delta(h + E) = \\
&= -2a_{e,h+E}e + 2a_{f,h+E}f - a_{p,h+E}p + a_{q,h+E}q - a_{E,h+E}E + a_{F,h+E}F + \\
&\quad + 2a_{E,h+E}e - a_{F,h+E}h + a_{G,h+E}p + a_{h,h+E}E - a_{f,h+E}F + a_{q,h+E}G.
\end{aligned}$$

On the other hand,

$$\Delta(h + E) = \Delta(h) + \Delta(E) = 2b_{E,E}e - b_{F,E}h + b_{G,E}p + b_{h,E}E - b_{f,E}F + b_{q,E}G.$$

Comparing the coefficients at the basis elements  $q$  and  $G$ , we get  $b_{q,E} = 0$ .

Similarly, considering

$$\begin{aligned}
\Delta(h + F) &= [a, h + F] + \mu_{h+F}\delta(h + F) = \\
&= [a_{e,h+F}e + a_{f,h+F}f + a_{h,h+F}h + a_{p,h+F}p + a_{q,h+F}q + a_{z,h+F}z + \\
&\quad + a_{E,h+F}E + a_{F,h+F}F + a_{G,h+F}G, h + F] + \mu_{h+F}\delta(h + F) = \\
&= -2a_{e,h+F}e + 2a_{f,h+F}f - a_{p,h+F}p + a_{q,h+F}q - a_{E,h+F}E + a_{F,h+F}F \\
&\quad - 2a_{F,h+F}f + a_{E,h+F}h - a_{G,h+F}q - a_{e,h+F}E - a_{h,h+F}F + a_{p,h+F}G.
\end{aligned}$$

On the other hand,

$$\Delta(h + F) = \Delta(h) + \Delta(F) = -2b_{F,F}f + b_{E,F}h - b_{G,F}q - b_{e,F}E - b_{h,F}F + b_{p,F}G.$$

Comparing the coefficients at the basis elements  $p$  and  $G$ , we get  $b_{p,F} = 0$ .

Considering

$$\begin{aligned}
\Delta(e - f + h) &= [a, e - f + h] + \mu_{e-f+h}\delta(e - f + h) = \\
&= [a_{e,e-f+h}e + a_{f,e-f+h}f + a_{h,e-f+h}h + a_{p,e-f+h}p + a_{q,e-f+h}q + a_{z,e-f+h}z + \\
&\quad + a_{E,e-f+h}E + a_{F,e-f+h}F + a_{G,e-f+h}G, e - f + h] + \mu_{e-f+h}\delta(e - f + h) = \\
&= 2a_{h,e-f+h}e - a_{f,e-f+h}h - a_{q,e-f+h}p + a_{F,e-f+h}E + \\
&\quad + 2a_{h,e-f+h}f - a_{e,e-f+h}h + a_{p,e-f+h}q - a_{E,e-f+h}F - \\
&\quad - 2a_{e,e-f+h}e + 2a_{f,e-f+h}f - a_{p,e-f+h}p + a_{q,e-f+h}q - a_{E,e-f+h}E + a_{F,e-f+h}F.
\end{aligned}$$

On the other hand,

$$\Delta(e - f + h) = \Delta(e) - \Delta(f) = 2b_{h,e}e - b_{q,e}p - 2b_{h,f}f - b_{p,f}q.$$

Comparing the coefficients at the basis elements  $p$  and  $q$ , we get  $b_{p,f} = b_{q,e}$ .

We consider

$$\begin{aligned}
\Delta(E + F) &= [a, E + F] + \mu_{E+F}\delta(E + F) = \\
&= [a_{e,E+F}e + a_{f,E+F}f + a_{h,E+F}h + a_{p,E+F}p + a_{q,E+F}q + a_{z,E+F}z + \\
&\quad + a_{E,E+F}E + a_{F,E+F}F + a_{G,E+F}G, E + F] + \mu_{E+F}\delta(E + F) = \\
&= 2a_{E,E+F}e - a_{F,E+F}h + a_{G,E+F}p + a_{h,E+F}E - a_{f,E+F}F + a_{q,E+F}G - \\
&\quad - 2a_{F,E+F}f + a_{E,E+F}h - a_{G,E+F}q - a_{e,E+F}E - a_{h,E+F}F + a_{p,E+F}G.
\end{aligned}$$

On the other hand,

$$\begin{aligned}\Delta(E + F) &= \Delta(E) + \Delta(F) = 2b_{E,E}e - b_{F,E}h + b_{G,EP} + b_{h,E}E - b_{f,E}F - \\ &\quad - 2b_{F,F}f + b_{E,F}h - b_{G,F}q - b_{e,F}E - b_{h,FF}.\end{aligned}$$

Comparing the coefficients at the basis elements  $p$  and  $q$ , we get  $b_{G,E} = b_{G,F}$ .

We consider

$$\begin{aligned}\Delta(E + G) &= [a, E + G] + \mu_{E+G}\delta(E + G) = \\ &= [a_{e,E+G}e + a_{f,E+G}f + a_{h,E+G}h + a_{p,E+G}p + a_{q,E+G}q + a_{z,E+G}z + \\ &\quad + a_{E,E+G}E + a_{F,E+G}F + a_{G,E+G}G, E + G] + \mu_{E+G}\delta(E + G) = \\ &= 2a_{E,E+G}e - a_{F,E+G}h + a_{G,E+G}p + a_{h,E+G}E - a_{f,E+G}F + a_{q,E+G}G - \\ &\quad - a_{E,E+G}p + a_{F,E+G}q + a_{G,E+G}z + \mu_{E+G}G.\end{aligned}$$

On the other hand,

$$\begin{aligned}\Delta(E + G) &= \Delta(E) + \Delta(G) = 2b_{E,E}e - b_{F,E}h + b_{G,EP} + b_{h,E}E - b_{f,E}F - \\ &\quad - b_{E,GP} + b_{F,G}q + b_{G,G}z + \lambda_G G.\end{aligned}$$

Comparing the coefficients at the basis elements  $h$  and  $q$ , we get  $b_{F,E} = b_{F,G}$ .

We consider

$$\begin{aligned}\Delta(F + G) &= [a, F + G] + \mu_{F+G}\delta(F + G) = \\ &= [a_{e,F+G}e + a_{f,F+G}f + a_{h,F+G}h + a_{p,F+G}p + a_{q,F+G}q + a_{z,F+G}z + \\ &\quad + a_{E,F+G}E + a_{F,F+G}F + a_{G,F+G}G, F + G] + \mu_{F+G}\delta(F + G) = \\ &= -2a_{F,F+G}f + a_{E,F+G}h - a_{G,F+G}q - a_{e,F+G}E - a_{h,F+G}F + a_{p,F+G}G - \\ &\quad - a_{E,F+G}p + a_{F,F+G}q + a_{G,F+G}z + \mu_{F+G}G.\end{aligned}$$

On the other hand,

$$\begin{aligned}\Delta(F + G) &= \Delta(F) + \Delta(G) = -2b_{F,F}f + b_{E,F}h - b_{G,F}q - b_{e,F}E - b_{h,FF} - \\ &\quad - b_{E,GP} + b_{F,G}q + b_{G,G}z + \lambda_G G.\end{aligned}$$

Comparing the coefficients at the basis elements  $h$  and  $p$ , we get  $b_{E,F} = b_{E,G}$ . □

### Lemma 3.6.

$$\Delta(e) = 2b_{h,e}e,$$

$$\Delta(f) = -2b_{h,e}f.$$

*Proof.* We consider take an element  $y = e + f + E + F$

$$\begin{aligned}\Delta(y) &= [a, y] + \mu_y\delta(y) = \\ &= [a_{e,y}e + a_{f,y}f + a_{h,y}h + a_{p,y}p + a_{q,y}q + a_{z,y}z + a_{E,y}E + a_{F,y}F + a_{G,y}G, y] + \mu_y\delta(y) = \\ &= 2a_{h,y}e - a_{f,y}h - a_{q,y}p + a_{F,y}E - \\ &\quad - 2a_{h,y}f + a_{e,y}h - a_{p,y}q + a_{E,y}F + \\ &\quad + 2a_{E,y}e - a_{F,y}h + a_{G,y}p + a_{h,y}E - a_{f,y}F + a_{q,y}G - \\ &\quad - 2a_{F,y}f + a_{E,y}h - a_{G,y}q - a_{e,y}E - a_{h,y}F + a_{p,y}G.\end{aligned}$$



On the other hand,

$$\begin{aligned}\Delta(y) &= \Delta(e) + \Delta(f) + \Delta(E) + \Delta(F) = 2b_{h,e}e - b_{q,e}p - 2b_{h,f}f - b_{q,e}q + \\ &+ 2b_{E,E}e - b_{F,E}h + b_{G,EP} + b_{h,E}E - b_{f,E}F - \\ &- 2b_{F,F}f + b_{E,F}h - b_{G,Eq} - b_{e,F}E - b_{h,FF}.\end{aligned}$$

Comparing the coefficients at the basis elements  $p$ ,  $q$  and  $G$ , we get

$$\begin{aligned}-b_{q,e} + b_{G,E} &= -a_{q,y} + a_{G,y}, \\ -b_{q,e} - b_{G,E} &= -a_{p,y} - a_{G,y}, \\ 0 &= a_{q,y} + a_{p,y},\end{aligned}$$

which implies  $b_{q,e} = 0$ .

We consider

$$\begin{aligned}\Delta(e + f) &= [a, e + f] + \mu_{e+f}\delta(e + f) = \\ &= [a_{e,e+f}e + a_{f,e+f}f + a_{h,e+f}h + a_{p,e+f}p + a_{q,e+f}q + \\ &+ a_{z,e+f}z + a_{E,e+f}E + a_{F,e+f}F + a_{G,e+f}G, e + f] + \mu_{e+f}\delta(e + f) = \\ &= 2a_{h,e+f}e - a_{f,e+f}h - a_{q,e+f}p + a_{F,e+f}E - \\ &- 2a_{h,e+f}f + a_{e,e+f}h - a_{p,e+f}q + a_{E,e+f}F.\end{aligned}$$

On the other hand,

$$\Delta(e + f) = \Delta(e) + \Delta(f) = 2b_{h,e}e - 2b_{h,f}f.$$

Comparing the coefficients at the basis elements  $e$  and  $f$ , we get  $b_{h,e} = b_{h,f}$ . □

**Lemma 3.7.**

$$\begin{aligned}\Delta(p) &= (b_{h,p} + \lambda_p)p, \\ \Delta(q) &= (-b_{h,q} + \lambda_q)q, \\ \Delta(E) &= b_{G,EP} + b_{h,e}E, \\ \Delta(F) &= -b_{G,Eq} - b_{h,e}F, \\ \Delta(G) &= b_{G,Gz} + \lambda_G G.\end{aligned}$$

*Proof.* We consider

$$\begin{aligned}\Delta(e + G) &= [a, e + G] + \mu_{e+G}\delta(e + G) = \\ &= [a_{e,e+G}e + a_{f,e+G}f + a_{h,e+G}h + a_{p,e+G}p + a_{q,e+G}q + \\ &+ a_{z,e+G}z + a_{E,e+G}E + a_{F,e+G}F + a_{G,e+G}G, e + G] + \mu_{e+G}\delta(e + G) = \\ &= 2a_{h,e+G}e - a_{f,e+G}h - a_{q,e+G}p + a_{F,e+G}E - \\ &- a_{E,e+G}p + a_{F,e+G}q + a_{G,e+G}z + \mu_{e+G}G.\end{aligned}$$

On the other hand,

$$\Delta(e + G) = \Delta(e) + \Delta(G) = 2b_{h,e}e - b_{E,FP} + b_{F,Eq} + b_{G,Gz} + \lambda_G G.$$

Comparing the coefficients at the basis elements  $E$  and  $q$ , we get  $b_{F,E} = 0$ .

We consider

$$\begin{aligned}
\Delta(f + G) &= [a, f + G] + \mu_{f+G}\delta(f + G) = \\
&= [a_{e,f+G}e + a_{f,f+G}f + a_{h,f+G}h + a_{p,f+G}p + a_{q,f+G}q + \\
&+ a_{z,f+G}z + a_{E,f+G}E + a_{F,f+G}F + a_{G,f+G}G, f + G] + \mu_{f+G}\delta(f + G) = \\
&= -2a_{h,f+G}f + a_{e,f+G}h - a_{p,f+G}q + a_{E,f+G}F \\
&- a_{E,f+G}p + a_{F,f+G}q + a_{G,f+G}z + \mu_{f+G}G.
\end{aligned}$$

On the other hand,

$$\Delta(f + G) = \Delta(f) + \Delta(G) = -2b_{h,f}f - b_{E,F}p + b_{G,G}z + \lambda_G G.$$

Comparing the coefficients at the basis elements  $F$  and  $p$ , we get  $b_{E,F} = 0$ .

We consider take an element  $y = E + F + G$

$$\begin{aligned}
\Delta(y) &= [a, y] + \mu_y\delta(y) = \\
&= [a_{e,y}e + a_{f,y}f + a_{h,y}h + a_{p,y}p + a_{q,y}q + a_{z,y}z + a_{E,y}E + a_{F,y}F + a_{G,y}G, y] + \mu_y\delta(y) = \\
&= 2a_{E,y}e - a_{F,y}h + a_{G,y}p + a_{h,y}E - a_{f,y}F + a_{q,y}G - \\
&- 2a_{F,y}f + a_{E,y}h - a_{G,y}q - a_{e,y}E - a_{h,y}F + a_{p,y}G - \\
&- a_{E,y}p + a_{F,y}q + a_{G,y}z + \mu_y G.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
\Delta(y) &= \Delta(E) + \Delta(F) + \Delta(G) = 2b_{E,E}e + b_{G,EP} + b_{h,E}E - b_{f,E}F - \\
&- 2b_{F,F}f - b_{G,Eq} - b_{e,F}E - b_{h,FF} + b_{G,G}z + \lambda_G G.
\end{aligned}$$

Comparing the coefficients at the basis elements  $e$ ,  $h$  and  $f$ , we get  $b_{E,E} = b_{F,F}$ .

We consider

$$\begin{aligned}
\Delta(e + p + E) &= [a, e + p + E] + \mu_{e+p+E}\delta(e + p + E) = \\
&= [a_{e,e+p+E}e + a_{f,e+p+E}f + a_{h,e+p+E}h + a_{p,e+p+E}p + a_{q,e+p+E}q + \\
&+ a_{z,e+p+E}z + a_{E,e+p+E}E + a_{F,e+p+E}F + a_{G,e+p+E}G, e + p + E] + \\
&+ \mu_{e+p+E}\delta(e + p + E) = 2a_{h,e+p+E}e - a_{f,e+p+E}h - \\
&- a_{q,e+p+E}p + a_{F,e+p+E}E + (a_{h,e+p+E} + \mu_{e+p+E})p + \\
&+ a_{f,e+p+E}q - a_{q,e+p+E}z - a_{F,e+p+E}G + 2a_{E,e+p+E}e - a_{F,e+p+E}h + \\
&+ a_{G,e+p+E}p + a_{h,e+p+E}E - a_{f,e+p+E}F + a_{q,e+p+E}G.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
\Delta(e + p + E) &= \Delta(e) + \Delta(p) + \Delta(E) = 2b_{h,e}e + \\
&+ (b_{h,p} + \lambda_p)p - b_{q,p}z + 2b_{E,E}e + b_{G,EP} + b_{h,E}E - b_{f,E}F.
\end{aligned}$$

Comparing the coefficients at the basis elements  $h$ ,  $q$ ,  $z$ ,  $E$ ,  $F$  and  $G$  we get  $b_{f,E} = b_{q,p} = 0$ .

We consider

$$\begin{aligned}
\Delta(q + F) &= [a, q + F] + \mu_{q+F}\delta(q + F) = \\
&= [a_{e,q+F}e + a_{f,q+F}f + a_{h,q+F}h + a_{p,q+F}p + a_{q,q+F}q + \\
&+ a_{z,q+F}z + a_{E,q+F}E + a_{F,q+F}F + a_{G,q+F}G, q + F] + \mu_{q+F}\delta(q + F) = \\
&= a_{e,q+F}p + (-a_{h,q+F} + \mu_{q+F})q + a_{p,q+F}z - a_{E,q+F}G - \\
&- 2a_{F,q+F}f + a_{E,q+F}h - a_{G,q+F}q - a_{e,q+F}E - a_{h,q+F}F + a_{p,q+F}G.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
\Delta(q + F) &= \Delta(q) + \Delta(F) = (-b_{h,q} + \lambda_q)q + b_{p,q}z + \\
&- 2b_{F,F}f - b_{G,E}q - b_{e,F}E - b_{h,F}F.
\end{aligned}$$

Comparing the coefficients at the basis elements  $h$ ,  $p$ ,  $z$ ,  $E$  and  $G$ , we get  $b_{e,F} = b_{p,q} = 0$ .

We consider take an element  $\nu = f + p + E$

$$\begin{aligned}
\Delta(\nu) &= [a, \nu] + \mu_\nu\delta(\nu) = \\
&= [a_{e,\nu}e + a_{f,\nu}f + a_{h,\nu}h + a_{p,\nu}p + a_{q,\nu}q + a_{z,\nu}z + a_{E,\nu}E + a_{F,\nu}F + a_{G,\nu}G, \nu] + \mu_\nu\delta(\nu) = \\
&= -2a_{h,\nu}f + a_{e,\nu}h - a_{p,\nu}q + a_{E,\nu}F + \\
&+ (a_{h,\nu} + \lambda_\nu)p + a_{f,\nu}q - a_{q,\nu}z - a_{F,\nu}G + \\
&+ 2a_{E,\nu}e - a_{F,\nu}h + a_{G,\nu}p + a_{h,\nu}E - a_{f,\nu}F + a_{q,\nu}G.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
\Delta(\nu) &= \Delta(f) + \Delta(p) + \Delta(E) = -2b_{h,e}f + \\
&+ (b_{h,p} + \lambda_p)p - b_{q,p}z + 2b_{E,E}e + b_{G,E}p + b_{h,E}E.
\end{aligned}$$

Comparing the coefficients at the basis elements  $e$ ,  $q$  and  $F$ , we get  $b_{E,E} = 0$ .

We consider

$$\begin{aligned}
\Delta(e + F) &= [a, e + F] + \mu_{e+F}\delta(e + F) = \\
&= [a_{e,e+F}e + a_{f,e+F}f + a_{h,e+F}h + a_{p,e+F}p + a_{q,e+F}q + \\
&+ a_{z,e+F}z + a_{E,e+F}E + a_{F,e+F}F + a_{G,e+F}G, e + F] + \mu_{e+F}\delta(e + F) = \\
&= 2a_{h,e+F}e - a_{f,e+F}h - a_{q,e+F}p + a_{F,e+F}E - \\
&- 2a_{F,e+F}f + a_{E,e+F}h - a_{G,e+F}q - a_{e,e+F}E - a_{h,e+F}F + a_{p,e+F}G.
\end{aligned}$$

On the other hand,

$$\Delta(e + F) = \Delta(e) + \Delta(F) = 2b_{h,e}e - b_{G,E}q - b_{h,F}F.$$

Comparing the coefficients at the basis elements  $e$ ,  $f$ ,  $h$ ,  $E$  and  $F$ , we get  $b_{h,F} = b_{h,e}$ .

We consider

$$\begin{aligned}
\Delta(f + E) &= [a, f + E] + \mu_{f+E}\delta(f + E) = \\
&= [a_{e,f+E}e + a_{f,f+E}f + a_{h,f+E}h + a_{p,f+E}p + a_{q,f+E}q + \\
&+ a_{z,f+E}z + a_{E,f+E}E + a_{F,f+E}F + a_{G,f+E}G, f + E] + \mu_{f+E}\delta(f + E) = \\
&= -2a_{h,f+E}f + a_{e,f+E}h - a_{p,f+E}q + a_{E,f+E}F + \\
&+ 2a_{E,f+E}e - a_{F,f+E}h + a_{G,f+E}p + a_{h,f+E}E - a_{f,f+E}F + a_{q,f+E}G.
\end{aligned}$$

On the other hand,

$$\Delta(f + E) = \Delta(f) + \Delta(E) = -2b_{h,e}f + b_{G,EP} + b_{h,E}E.$$

Comparing the coefficients at the basis elements  $e$ ,  $f$ ,  $h$ ,  $E$  and  $F$ , we get  $b_{h,E} = b_{h,e}$ .  $\square$

**Lemma 3.8.**

$$\begin{aligned}\Delta(p) &= b_{h,e}p, \\ \Delta(q) &= -b_{h,e}q, \\ \Delta(E) &= b_{G,EP} + b_{h,e}E, \\ \Delta(F) &= -b_{G,EQ} - b_{h,e}F, \\ \Delta(G) &= b_{G,E}z.\end{aligned}$$

*Proof.* We consider take an element  $y = E - F + G$

$$\begin{aligned}\Delta(y) &= [a, y] + \mu_y \delta(y) = \\ &= [a_{e,y}e + a_{f,y}f + a_{h,y}h + a_{p,y}p + a_{q,y}q + a_{z,y}z + a_{E,y}E + a_{F,y}F + a_{G,y}, y] + \mu_y \delta(y) = \\ &= 2a_{E,y}e - a_{F,y}h + a_{G,y}p + a_{h,y}E - a_{f,y}F + a_{q,y}G + \\ &+ 2a_{F,y}f - a_{E,y}h + a_{G,y}q + a_{e,y}E + a_{h,y}F - a_{p,y}G - \\ &- a_{E,y}p + a_{F,y}q + a_{G,y}z + \mu_y G.\end{aligned}$$

On the other hand,

$$\begin{aligned}\Delta(y) &= \Delta(E) - \Delta(F) + \Delta(G) = b_{G,EP} + b_{h,e}E + \\ &+ b_{G,EQ} + b_{h,e}F + b_{G,G}z + \lambda_G G.\end{aligned}$$

Comparing the coefficients at the basis elements  $e$ ,  $p$  and  $z$ , we get  $b_{G,G} = b_{G,E}$ .

We consider take an element  $\nu = f + q - \frac{1}{2}z$

$$\begin{aligned}\Delta(\nu) &= [a, \nu] + \mu_\nu \delta(\nu) = \\ &= [a_{e,\nu}e + a_{f,\nu}f + a_{h,\nu}h + a_{p,\nu}p + a_{q,\nu}q + a_{z,\nu}z + a_{E,\nu}E + a_{F,\nu}F + a_{G,\nu}G, \nu] + \mu_\nu \delta(\nu) = \\ &= -2a_{h,\nu}f + a_{e,\nu}h - a_{p,\nu}q + a_{E,\nu}F + \\ &+ a_{e,\nu}p + (-a_{h,\nu} + \mu_\nu)q + a_{p,\nu}z - a_{E,\nu}G - \mu_\nu z.\end{aligned}$$

On the other hand,

$$\Delta(\nu) = \Delta(f) + \Delta(p) - \frac{1}{2}\Delta(z) = -2b_{h,e}f + (-b_{h,q} + \lambda_q)q.$$

Comparing the coefficients at the basis elements  $f$ ,  $q$  and  $z$ , we get  $-b_{h,q} + \lambda_q = -b_{h,e}$ .

We consider take an element  $\tau = e + p + \frac{1}{2}z$

$$\begin{aligned}\Delta(\tau) &= [a, \tau] + \mu_\tau \delta(\tau) = \\ &= [a_{e,\tau}e + a_{f,\tau}f + a_{h,\tau}h + a_{p,\tau}p + a_{q,\tau}q + a_{z,\tau}z + a_{E,\tau}E + a_{F,\tau}F + a_{G,\tau}G, \tau] + \mu_\tau \delta(\tau) = \\ &= 2a_{h,\tau}e - a_{f,\tau}h - a_{q,\tau}p + a_{F,\tau}E + \\ &+ (a_{h,\tau} + \mu_\tau)p + a_{f,\tau}q - a_{q,\tau}z - a_{F,\tau}G + \mu_\tau z.\end{aligned}$$

On the other hand,

$$\Delta(\tau) = \Delta(e) + \Delta(p) + \frac{1}{2}\Delta(z) = 2b_{h,e}e + (b_{h,p} + \lambda_p)p.$$

Comparing the coefficients at the basis elements  $e$ ,  $p$  and  $z$ , we get  $b_{h,p} + \lambda_p = b_{h,e}$ .

We consider take an element  $t = p + q + E - F + G$

$$\begin{aligned}\Delta(t) &= [a, t] + \mu_t \delta(t) = \\ &= [a_{e,t}e + a_{f,t}f + a_{h,t}h + a_{p,t}p + a_{q,t}q + a_{z,t}z + a_{E,t}E + a_{F,t}F + a_{G,t}t] + \mu_t \delta(t) = \\ &= (a_{h,t} + \mu_t)p + a_{f,t}q - a_{q,t}z - a_{F,t}G + \\ &+ a_{e,t}p + (-a_{h,t} + \mu_t)q + a_{p,t}z - a_{E,t}G + \\ &+ 2a_{E,t}e - a_{F,t}h + a_{G,t}p + a_{h,t}E - a_{f,t}F + a_{q,t}G - \\ &+ 2a_{F,t}f - a_{E,t}h + a_{G,t}q + a_{e,t}E + a_{h,t}F - a_{p,t}G - \\ &- a_{E,t}p + a_{F,t}q + a_{G,t}z + \mu_t G.\end{aligned}$$

On the other hand,

$$\begin{aligned}\Delta(t) &= \Delta(p) + \Delta(q) + \Delta(E) - \Delta(F) + \Delta(G) = b_{h,e}p - \\ &- b_{h,e}q + b_{G,E}p + b_{h,e}E - b_{G,E}q - b_{h,e}F + b_{G,G}z + \lambda_G G.\end{aligned}$$

Comparing the coefficients at the basis elements  $e, f, p, z, E$  and  $G$ , we get  $\lambda_G = 0$ .  $\square$

**Lemma 3.9.**  $\Delta$  is an inner super-derivation on  $\mathcal{S}$

*Proof.* Let  $x = x_e e + x_f f + x_h h + x_p p + x_q q + x_z z + x_E E + x_F F + x_G G \in \mathcal{S}$ .

We consider, by Lemma 3.2, Lemma 3.6 and Lemma 3.8

$$\begin{aligned}\Delta(x) - [b_{h,e}h + b_{G,E}G, x] &= \Delta(x_e e + x_f f + x_h h + x_p p + x_q q + x_z z + x_E E + x_F F + x_G G) - \\ &- [b_{h,e}h + b_{G,E}G, x_e e + x_f f + x_h h + x_p p + x_q q + x_z z + x_E E + x_F F + x_G G] = \\ &= (2x_e e - 2x_f f + x_p p - x_q q + x_E E - x_F F)b_{h,e} + (x_E p - x_F q + x_G z)b_{G,E} - \\ &- (2x_e e - 2x_f f + x_p p - x_q q + x_E E - x_F F)b_{h,e} - (x_E p - x_F q + x_G z)b_{G,E} = 0.\end{aligned}$$

Then  $\Delta$  is an inner super-derivation.  $\square$

Now we are in position to prove Theorem 3.1.

**Proof of Theorem 3.1.** Let  $\nabla$  be a local super-derivation of  $\mathcal{S}$ . For the element  $h + z$  exists there a super-derivation  $D_{h+z}$  such that

$$\nabla(h + z) = D_{h+z}(h + z).$$

Set  $\Delta = \nabla - D_{h+z}$ . Then  $\Delta$  is a local super-derivation such that  $\Delta(h + z) = 0$ . By Lemma 3.9,  $\Delta(x) = ad_{b_{h,e}h + b_{G,E}G}(x)$ . Thus  $\nabla = D_{h+z} + ad_{b_{h,e}h + b_{G,E}G}$ . Then  $\nabla$  is a super-derivation. The proof is complete.

### Declarations

Conflict of interest The author declares no conflict of interest.

### REFERENCES

- [1] J. Adashev and B. Yusupov, *Local derivations and automorphisms of direct sum null-filiform Leibniz algebras*, Lobachevskii Journal of Mathematics, **43**:12, (2022) 1–7.
- [2] A. K. Alauadinov, B. B. Yusupov, *Local derivations of the Schrödinger algebras*, Algebra Colloquium, (2024) in press.
- [3] A. K. Alauadinov, B. B. Yusupov, *Local derivations of conformal Galilei algebra*, Communications in Algebra, Communications in Algebra, **52**:6 (2024), 2489–2508.
- [4] Sh. Ayupov, A. Elduque and K. Kudaybergenov, *Local derivations and automorphisms of Cayley algebras*, Journal of Pure and Applied Algebra, **227**:5, (2023) 107277.
- [5] Sh.A. Ayupov, A.Kh. Khudoyberdiyev, B.B. Yusupov, *Local and 2-local derivations of solvable Leibniz algebras*, Internat. J. Algebra Comput., **30**:6 (2020) 1185–1197.

- [6] Sh.A. Ayupov, K.K. Kudaybergenov, *Local derivations on finite-dimensional Lie algebras*, Linear Algebra Appl., **493** (2016) 381-398.
- [7] Sh. A. Ayupov, K. K. Kudaybergenov, A. Allambergenov, *Local and 2-local derivations on octonion algebras*, Journal of algebra and its applications, Vol. 22, No. 07 (2024), 2350147.
- [8] Sh. A. Ayupov, K. K. Kudaybergenov, B. B. Yusupov, *Local and 2-local derivations of  $p$ -filiform Leibniz algebras*, Journal of Mathematical Sciences, **245** (3), (2020) 359–367.
- [9] Sh. A. Ayupov, K. K. Kudaybergenov, B. B. Yusupov, *Local and 2-Local Derivations of Locally Simple Lie Algebras*, Journal of Mathematical Sciences, **278**: 4 (2024) 613–622.
- [10] H. Chen, Y. Wang, *Local superderivation on Lie superalgebra  $q(n)$* , Czechoslovak Mathematical Journal **68** (143), (2018) 661–675.
- [11] Y. Chen, K. Zhao, Y. Zhao, *Local derivations on Witt algebras*, Linear and multilinear algebra, **70**:6 (2022) 1159–1172.
- [12] C. Duval, P. Horvathy, *On Schrodinger superalgebras*, Journal of Mathematical Physics, **35**:5, 2516–2538, (1994).
- [13] A.P. Elisova, I.N. Zotov, V.M. Levchuk and G.S. Suleymanova, *Local automorphisms and local derivations of nilpotent matrix algebras*, Izv. Irkutsk Gos. Univ., **4**:1 (2011) 9-19.
- [14] B. Ferreira, I. Kaygorodov, K. Kudaybergenov, *Local and 2-local derivations of simple  $n$ -ary algebras*, Ricerche di Matematica, (2021).
- [15] Y. He, *Structures and representation theory for  $sl_{n+1}(\mathbb{C})$  and related Lie (super) algebra*, Thesis (Ph.D.)- Soochow University, 66 pp, (2019).
- [16] B.E. Johnson, *Local derivations on  $C^*$ -algebras are derivations*, Transactions of the American Mathematical Society, **353** (2001) 313–325.
- [17] V. Dobrev, H. Doebner, C. Mrugalla, *Lowest weight representations of the Schrodinger algebra and generalized heat/Schrödinger equations*, Rep. Math. Phys. 39 (1997) 201–218.
- [18] R.V. Kadison, *Local derivations*, J. Algebra, **130** (1990) 494–509.
- [19] V. G. Kac, *Lie superalgebras*, Adv. Math., **26**:1, (1977) 8–96.
- [20] M. Perroud, *Projective representations of the Schrödinger group*, Helv. Phys. Acta., **50** (1997) 233–252.
- [21] D. R. Larson, A. R. Sourour, *Local derivations and local automorphisms of  $B(X)$* , Proc. Sympos. Pure Math., **51** (1990) 187-194.
- [22] K.K. Kudaybergenov, B.A. Omirov, T.K. Kurbanbaev, *Local derivations on solvable Lie algebras of maximal rank*, Communications in Algebra, **50**:9 (2022), 1-11.
- [23] K.K. Kudaybergenov, I. Kaygorodov, I. Yuldashev, *Local derivations of semisimple Leibniz algebras*, Communications in Mathematics, **30**:2 (2022), 1–12.
- [24] Y. Yu, Zh. Chen, *Local derivations on Borel subalgebras of finite-dimensional simple Lie algebras*, Comm. Algebra, **48**:1 (2020) 1-10.
- [25] Y. F. Yao, *Local derivations on the Witt algebra in prime characteristic*, Linear and multilinear algebra, **70**(15) (2022), 2919-2933.

(Amir Alauadinov) CH.ABDIROV 1, DEPARTMENT OF MATHEMATICS, KARAKALPAK STATE UNIVERSITY, NUKUS 230113, UZBEKISTAN

AND

V.I.ROMANOVSKIY INSTITUTE OF MATHEMATICS, UZBEKISTAN ACADEMY OF SCIENCES, UNIVERSITY STREET, 9, OLMAZOR DISTRICT, TASHKENT, 100174, UZBEKISTAN

*Email address:* amir\_t85@mail.ru

(Bakhtiyor Yusupov) V.I.ROMANOVSKIY INSTITUTE OF MATHEMATICS, UZBEKISTAN ACADEMY OF SCIENCES, UNIVERSITY STREET, 9, OLMAZOR DISTRICT, TASHKENT, 100174, UZBEKISTAN

AND

DEPARTMENT OF PHYSICS AND MATHEMATICS, URGENCH STATE UNIVERSITY, H. ALIMDJAN STREET, 14, URGENCH 220100, UZBEKISTAN

*Email address:* baxtiyor\_yusupov\_93@mail.ru b.yusupov@mathinst.uz