

# Full Asymptotic Expansion of Monodromy Data for the First Painlevé Transcendent: Applications to Connection Problems

Yun-Jiang Jiang<sup>†</sup>, Yu-Tian Li<sup>‡</sup>, and Wen-Gao Long<sup>\*†</sup>

<sup>†</sup>School of Mathematics and Computational Science, Hunan University of Science and Technology, Xiangtan, Hunan, 411201, PR China

<sup>‡</sup>School of Science and Engineering, The Chinese University of Hong Kong, Shenzhen, Guangdong, 518172, PR China.

## Abstract

We study the full asymptotic expansion of the monodromy data (*i.e.*, Stokes multipliers) for the first Painlevé transcendent (PI) with large initial data or large pole parameters. Our primary approach involves refining the complex WKB method, also known as the method of uniform asymptotics, to approximate the second-order ODEs derived from PI's Lax pair with higher-order accuracy. As an application, we provide a rigorous proof of the full asymptotic expansion of the nonlinear eigenvalues proposed numerically by Bender, Komijani, and Wang. Additionally, we present the full asymptotic expansion for the pole parameters  $(p_n, H_n)$  corresponding to the  $n$ -th pole of the real tritronquée solution of the PI equation as  $n \rightarrow +\infty$ .

*Keywords:* First Painlevé transcendent, connection formula, Stokes multiplier, uniform asymptotics, Airy function,

*MSC2020:* 34M40, 33E17, 34A12, 34E05, 33C10

## 1 Introduction

Painlevé equations are a group of six nonlinear second-order ordinary differential equations that possess the Painlevé property. This property requires that the movable singularities of the solutions must be poles, not branch points or essential singularities. In general, the solutions of these equations do not have explicit expressions. Consequently, asymptotic analysis is commonly used to study the behavior of solutions to Painlevé equations. This leads to a

---

\*Corresponding author: longwg@hnust.edu.cn

natural question: if the asymptotic approximations of a Painlevé function are given in different regimes, such as the variable  $t \rightarrow -\infty$  and  $t \rightarrow \infty$ , what is the relation between these asymptotic formulas? This is known as the connection formula problem in Painlevé equation theory, which has received significant attention in recent research; see for example [1–3, 7, 9, 10, 13–16, 19, 20, 22].

In this work, we consider the connection problem for the first Painlevé equation:

$$\frac{d^2y}{dt^2} = 6y^2 + t. \quad (1.1)$$

This problem has been previously studied by Joshi and Kruskal [10], as well as by Kapaev and Kitaev [13], yielding significant progress. However, the problem is not yet fully solved and has been highlighted by Clarkson as an open problem on several occasions [5, 6]. Since then, further investigations have been conducted in this direction [2, 14, 15].

The asymptotic behaviors of the PI solutions can be derived in several interesting regimes:

$$(i) \ t \rightarrow -\infty, \quad (ii) \ t \rightarrow 0, \quad (iii) \ t \rightarrow p, \quad (1.2)$$

where  $p$  is a pole of the PI solution.

The situation (i) has been extensively studied by Kapaev [11] using isomonodromy theory. There are three types of solutions for the PI equation according to the asymptotic behavior of  $y(t)$  as  $t \rightarrow -\infty$ . They are classified as Type A (oscillatory solutions), Type B (separatrix solutions) and Type C (singular solutions). The classification rules and details of the asymptotic behaviors are provided in Appendix B, for structural considerations. Situation (ii) is the initial value problem, specifically where the solution has a MacLaurin series expansion:

$$y(t) = a + bt + O(t^2), \quad t \rightarrow 0, \quad (1.3)$$

with the initial data  $(y(0), y'(0)) = (a, b)$  uniquely determining the solution. For the situation (iii), as  $t \rightarrow p$ , where  $p$  is a pole of  $y(t)$ , the solution has a Laurent series expansion:

$$y(t) = \frac{1}{(t-p)^2} - \frac{p}{10}(t-p)^2 - \frac{1}{6}(t-p)^3 + H(t-p)^4 + \mathcal{O}((t-p)^5), \quad t \rightarrow p. \quad (1.4)$$

Here,  $H$  is a free parameter, and the pair  $(p, H)$  uniquely determines a PI solution.

The monodromy theory plays a major role in the study of the connection problem of Painlevé transcendents. For structural considerations again, we describe the monodromy theory for PI in Appendix A. In principle, there is a one-to-one correspondence between PI solutions and Stokes multipliers: a pair of Stokes multipliers determines a unique PI solution and vice versa; see Appendix A. Therefore, one often-used approach in the study of connection problems is first to establish the relation between the Stokes multipliers and particular asymptotic approximations, just as Kapaev [11] did for situation (i). Then, using the Stokes multipliers as a bridge, the behaviors of a particular PI solution in different regimes can be connected.

For PI, we are particularly interested in two connection problems: (I)connection problem between the asymptotic behavior at negative infinity and the initial data; and (II)connection problem between the asymptotic behaviors at negative infinity and the pole. To this end, one need to calculate the Stokes multipliers for situations (ii) and (iii).

In situation (ii), the Stokes multipliers are studied in [14]. Although it is very challenging to derive the exact expression of the Stokes multipliers in terms of  $a$  and  $b$ , the leading asymptotic approximations of the Stokes multipliers are calculated when  $a$  or  $b$  is large. The corresponding results are applied to build some limiting-form connection formulas between the initial data and the parameters in the asymptotic expansions at negative infinity.

In situation (iii), i.e., when  $t \rightarrow p$ , the PI solution is expressed as a Laurent series (1.4). In [15], the authors study the leading asymptotic behavior of the Stokes multipliers  $s_k$ 's as  $p$  or  $H$  tends to infinity and provide an asymptotic classification of the PI solutions with respect to  $(p, H)$ .

In the present work, we aim to achieve a deep understanding and refined results of the above two connection problems for PI. The major step is to derive the full asymptotic expansion of the Stokes multipliers in situations (ii) and (iii). One of our motivations comes from [3], where the authors find that there exists a sequence  $\{b_n\}$  such that the PI solutions with  $y(0) = 0$  and  $y'(0) = b_n$  are the separatrix solutions. Similarly, there exists a sequence  $\{a_n\}$  such that the PI solutions with  $y(0) = a_n$  and  $y'(0) = 0$  are also the separatrix solutions. The values  $a_n$  and  $b_n$  are termed as the *nonlinear eigenvalues* for the PI equation. It is stated in [3], based on numerical simulations, that

$$\begin{aligned} a_n &\sim - \left[ \frac{\sqrt{3\pi}\Gamma\left(\frac{11}{6}\right)}{\Gamma\left(\frac{1}{3}\right)} \right]^{\frac{2}{5}} \left( n - \frac{1}{2} \right)^{\frac{2}{5}} \left[ 1 - \frac{0.0096518}{\left(n - \frac{1}{2}\right)^2} + \frac{0.0240}{\left(n - \frac{1}{2}\right)^4} \right], \\ b_n &\sim 2 \left[ \frac{\sqrt{3\pi}\Gamma\left(\frac{11}{6}\right)}{\Gamma\left(\frac{1}{3}\right)} \right]^{\frac{3}{5}} \left( n - \frac{1}{6} \right)^{\frac{3}{5}} \left[ 1 - \frac{0.00551328}{\left(n - \frac{1}{6}\right)^2} + \frac{0.29334}{\left(n - \frac{1}{6}\right)^3} \right] \end{aligned} \quad (1.5)$$

as  $n \rightarrow \infty$ . A natural problem is to obtain the full asymptotic expansions of  $a_n$  and  $b_n$  as  $n \rightarrow \infty$ . Moreover, equation (1.5) suggests that all the odd terms vanish and that  $a_n$  has a full asymptotic expansion of the form:

$$a_n \sim - \left[ \frac{\sqrt{3\pi}\Gamma\left(\frac{11}{6}\right)}{\Gamma\left(\frac{1}{3}\right)} \right]^{\frac{2}{5}} \left( n - \frac{1}{2} \right)^{\frac{2}{5}} \sum_{k=0}^{\infty} \frac{A_k}{\left(n - \frac{1}{2}\right)^{2k}}, \quad n \rightarrow \infty. \quad (1.6)$$

It turns out that the answer is affirmative. To provide a rigorous proof of (1.6), we need to calculate the full asymptotic expansion of the Stokes multipliers corresponding to the PI equation as  $a \rightarrow -\infty$ .

A similar question can be proposed in the connection problem of the PI equation between negative infinity and the pole. In [15], the authors obtain the leading asymptotic approximation of the pole parameter  $(p_n, H_n)$  for the  $n$ -th pole of the tritronquée solution of the PI equation as  $n \rightarrow \infty$ . Numerical evidence also shows that

$$\begin{aligned} p_n &\sim 2C_0 \left( \frac{4n-2}{\kappa^2(C_0)} \right)^{\frac{2}{5}} \left[ 1 - \frac{0.0045}{\left(n - \frac{1}{2}\right)^2} \right], \\ H_n &\sim -\frac{1}{7} \left( \frac{4n-2}{\kappa^2(C_0)} \right)^{\frac{6}{5}} \left[ 1 + \frac{0.0081}{\left(n - \frac{1}{2}\right)^2} \right], \end{aligned} \quad (1.7)$$

as  $n \rightarrow \infty$ , where  $C_0$  and  $\kappa^2(C_0)$  are defined in [15, Lemma 2.1]. A natural question is whether the asymptotic behavior of  $p_n$  and  $H_n$  both possess full asymptotic expansions with the odd terms vanishing. To answer this question, the full asymptotic expansion of the Stokes multipliers as  $p$  or  $H$  tends to infinity is needed.

In the study of connection problems of Painlevé equations, an often-used approach is the complex WKB method (also known as the method of *uniform asymptotics*) introduced by Bassom *et al.* [1] and developed by [15, 16, 19, 20, 22]. There are two steps in this approach. The first step is to approximate a second-order linear ODE (derived from the Lax pair) uniformly on two adjacent Stokes lines. These uniform asymptotics usually involve special functions like Airy or parabolic cylinder functions. The second step is to derive the Stokes multipliers of particular Painlevé functions by using the Stokes phenomena of the special functions used in the first step.

For the first Painlevé equation, when considering the cases as  $t \rightarrow 0$  and  $t \rightarrow p$ , although it is very difficult to calculate the explicit representation of the Stokes multipliers  $\{s_k\}$ , it is possible to obtain the asymptotic behavior of  $\{s_k\}$  when the initial data or pole parameter is large. However, since [1, Theorems 1 and 2] only give the leading approximation of the solutions of the Lax pair, we can only obtain the leading asymptotic behavior of the Stokes multipliers. To get a full asymptotic expansion of the Stokes multipliers, we need to improve the complex WKB method to obtain a higher-order approximation of the second-order ODEs derived from the Lax pair. This is another motivation for this paper.

The rest of this paper is organized as follows. In Sec. 2, we state the main results, which are divided into three parts: (1) Higher-order approximations of the Lax pair; (2) Full asymptotic expansions of the Stokes multipliers; (3) Full asymptotic expansions of the nonlinear eigenvalues and pole parameters. The proof of the third part is also included in this section, based on the results in the first two parts. In Sec. 3 and Sec. 4, we provide the proofs of the first and second parts, respectively.

## 2 Main results

The main results of this paper consist of three parts. First, we obtain the higher-order approximation of the second-order ODEs derived from the Lax pair for the PI equation in case (ii) [*resp.* case (iii)] stated in (1.2). Second, we obtain the full asymptotic expansion of the Stokes multipliers in case (ii) when the initial data [*resp.* the pole parameters] tend to infinity. The third part is an application of the full asymptotic expansions of the Stokes multipliers. We derive the full asymptotic expansions of  $a_n$ ,  $p_n$ , and  $H_n$  as  $n \rightarrow \infty$ , namely eqs. (1.6) and (1.7).

### 2.1 Higher-order approximations of the Lax pair

If we regard (A.10) as a linear system of ODEs for  $\hat{\Phi} = (\phi_1, \phi_2)^T$ , where  $T$  denotes the transpose, and eliminate the equation for  $\phi_2$ , then system (A.10) is equivalent to a second-order ODE for  $\phi_1$ . We then apply the method of uniform asymptotics developed by Bassom *et al.* [1] to obtain asymptotic approximations for  $\phi_1(\lambda)$  with a large parameter. To obtain

full asymptotic expansions, rather than just the leading term, we incorporate the method of uniform asymptotics with an idea developed by Dunster [8].

We first work on case (ii) with large initial data.

Assume  $t \neq p$ . It follows from (A.10) that  $Y$  satisfies a second-order ODE:

$$\frac{d^2\phi_1}{d\lambda^2} = \left[ y_t^2 + 4\lambda^3 + 2\lambda t - 2yt - 4y^3 - \frac{yt}{\lambda - y} + \frac{3}{4} \frac{1}{(\lambda - y)^2} \right] \phi_1. \quad (2.1)$$

When  $t = 0$ , we introduce a large parameter  $\xi$  defined by

$$\xi^{\frac{6}{5}} = y(0)^3 - \frac{y'(0)^2}{4} = \frac{b^2}{4} - a^3$$

and set

$$a = A(\xi)\xi^{\frac{2}{5}} \quad \text{and} \quad b = B(\xi)\xi^{\frac{3}{5}},$$

where  $A(\xi), B(\xi)$  are both bounded. Then  $\frac{B(\xi)^2}{4} - A(\xi)^3 = 1$ . Under the scaling transformation  $\lambda = \xi^{\frac{2}{5}}z, Y(z) = \phi_1(\xi^{\frac{2}{5}}z)$ , eq. (2.1) becomes

$$\frac{d^2Y}{dz^2} = \xi^2 \left[ f(z) + \frac{g(z)}{\xi} + \frac{h(z)}{\xi^2} \right] Y, \quad (2.2)$$

where

$$f(z) = 4(z - z_0)(z - z_1)(z - z_2), \quad g(z) = -\frac{B(\xi)}{z + A(\xi)}, \quad h(z) = \frac{3}{4(z + A(\xi))^2}$$

with  $z_0 = -1, z_1 = e^{\frac{\pi i}{3}}$  and  $z_{-1} = e^{-\frac{\pi i}{3}}$ . It is clear that  $f(z), g(z)$ , and  $h(z)$  are all analytic in a neighbourhood of  $z = z_1$ , and that  $f(z)$  has a simple zero at  $z = z_1$ . Define a transformation  $\zeta := \zeta(z)$  by

$$\frac{2}{3}\zeta^{\frac{3}{2}} = \int_{z_1}^z f(t)^{\frac{1}{2}} dt. \quad (2.3)$$

Then  $\zeta(z)$  is a conformal mapping in the neighbourhood of  $z = z_1$  and the two adjacent Stokes lines  $\{\arg \zeta = \frac{\pi}{3}, \pi\}$  emanating from  $z = z_1$  to infinity. Moreover, a direct calculation of the integral in (2.3) yields

$$\frac{2}{3}\zeta^{\frac{3}{2}} = \frac{4}{5}z^{\frac{5}{2}} + E_0 + \mathcal{O}(z^{-\frac{1}{2}}) \quad (2.4)$$

as  $z \rightarrow \infty$  uniformly for  $\arg z \in [0, \frac{4\pi}{5}]$ , where  $E_0 = \frac{3}{5}B(\frac{1}{2}, \frac{1}{3}) - \frac{\sqrt{3}i}{5}B(\frac{1}{2}, \frac{1}{3})$ . Set  $W(\zeta, \xi) = \left(\frac{f(z)}{\zeta}\right)^{\frac{1}{4}} Y$ , then (2.2) is transformed into

$$\frac{d^2W}{d\zeta^2} = \xi^2 \left[ \zeta + \frac{\varphi(\zeta)}{\xi} + \frac{\psi(\zeta)}{\xi^2} \right] W, \quad (2.5)$$

where

$$\begin{aligned} \varphi(\zeta) &= \frac{\zeta g(z)}{f(z)}, \\ \psi(\zeta) &= \frac{5}{16\zeta^2} + \frac{\zeta[4f(z)f''(z) - 5f'(z)^2]}{16f(z)^3} + \frac{\zeta h(z)}{f(z)} \end{aligned} \quad (2.6)$$

are both analytic at  $\zeta = 0$ . Moreover, inspired by the work of Dunster [8], we have the following Lemma.

**Lemma 1.** *Let  $\delta$  and  $\epsilon$  be fixed small positive constants and denote*

$$\mathbb{S} := \{\zeta \in \mathbb{C} : \arg \zeta = \frac{k\pi}{3}, k = 1, 3\}$$

and

$$\mathbb{D} := \left\{ \zeta \in \mathbb{C} : |\zeta| \leq \delta \text{ or } \frac{\pi}{3} - \epsilon < \arg \zeta < \pi + \epsilon \right\}.$$

For any  $n \in \mathbb{N}$ , define  $\hat{\zeta} = \zeta + \mathcal{A}_n(\zeta, \xi)$  with

$$\mathcal{A}_n(\zeta, \xi) = \sum_{s=1}^{2n} \frac{a_s(\zeta)}{\xi^s} \quad (2.7)$$

satisfying

$$\xi^2 \{ \mathcal{A}_n + (\zeta + \mathcal{A}_n)(2 + \mathcal{A}'_n) \mathcal{A}'_n \} + \frac{3\mathcal{A}_n''^2 - 2(1 + \mathcal{A}'_n)\mathcal{A}_n'''}{4(1 + \mathcal{A}'_n)^2} = \xi\varphi(\zeta) + \psi(\zeta) + \mathcal{O}(\xi^{-2n+1}) \quad (2.8)$$

as  $\xi \rightarrow +\infty$  uniformly for all  $\zeta \in \mathbb{D}$ . Assume also that each  $a_s(\zeta)$  is analytic in  $\mathbb{D}$ . Then for all  $s \in \mathbb{N}$ , the limits

$$\alpha_s(A(\xi), B(\xi)) := \lim_{\zeta \rightarrow \infty} \zeta^{\frac{1}{2}} a_s(\zeta)$$

exist and depend only on  $A(\xi)$  and  $B(\xi)$ . Moreover, for any solution  $W(\zeta, \xi)$  of (2.18), there exist two constants  $C_1$  and  $C_2$  such that

$$W(\zeta, \xi) = [C_1 + r_1(\zeta, \xi)]\text{Ai}_n(\zeta, \xi) + [C_2 + r_2(\zeta, \xi)]\text{Bi}_n(\zeta, \xi), \quad (2.9)$$

where

$$\text{Ai}_n(\zeta, \xi) = \left( \frac{d\hat{\zeta}}{d\zeta} \right)^{-\frac{1}{2}} \text{Ai} \left( \xi^{\frac{2}{3}} \hat{\zeta} \right) \quad \text{and} \quad \text{Bi}_n(\zeta, \xi) = \left( \frac{d\hat{\zeta}}{d\zeta} \right)^{-\frac{1}{2}} \text{Bi} \left( \xi^{\frac{2}{3}} \hat{\zeta} \right), \quad (2.10)$$

and  $r_{1,2}(\zeta, \xi) = \mathcal{O}(\xi^{-2n})$  as  $\xi \rightarrow +\infty$  uniformly for all  $\zeta \in \mathbb{D} \cap \mathbb{S}$ .

**Remark 1.** *Combining (2.7), (2.8) and the analyticity of  $a_s(\zeta)$ , we see that  $a_s(\zeta)$  are uniquely determined in a recursive manner. Specifically*

$$\begin{aligned} a_1(\zeta) &= \frac{1}{2\zeta^{\frac{1}{2}}} \int_0^\zeta \frac{\varphi(t)}{t^{\frac{1}{2}}} dt, \\ a_2(\zeta) &= \frac{1}{2\zeta^{\frac{1}{2}}} \int_0^\zeta \frac{\psi(t) - 2a_1(t)a_1'(t) - t(a_1'(t))^2}{t^{\frac{1}{2}}} dt, \\ a_s(\zeta) &= \frac{1}{2\zeta^{\frac{1}{2}}} \int_0^\zeta \frac{F_s(t)}{t^{\frac{1}{2}}} dt, \quad s > 2, \end{aligned} \quad (2.11)$$

where  $F_s(t)$  is can be expressed in terms of  $a_1(t), a_2(t), \dots, a_{s-1}(t)$ . If  $B(\xi) = 0$ , then  $g(z) = 0$  which implies  $a_1(\zeta) = 0$ . By induction, we can show that  $a_{2s+1}(\zeta) = 0$  for all  $s \in \mathbb{N}$ . Moreover, in this case, the estimates of  $r_{1,2}(\zeta, \xi)$  can be improved as

$$r_{1,2}(\zeta, \xi) = \mathcal{O}(\xi^{-2n-1}), \quad \xi \rightarrow +\infty. \quad (2.12)$$

**Remark 2.** *The proof of Lemma 1 is provided in Sec. 3, where we show that the limits of  $r_1(\zeta, \xi)$  and  $r_2(\zeta, \xi)$  as  $\zeta \rightarrow \infty$  with  $\arg \zeta \sim \frac{k\pi}{3}$ ,  $k = 1, 3$  exist. The four limit values may not be equal to each other, but they all have an estimate of  $\mathcal{O}(\xi^{-2n-1})$  as  $\xi \rightarrow +\infty$ . For convenience, hereafter, we set  $r_{j,k}(\xi) = \lim_{\zeta \rightarrow \infty} r_i(\zeta, \xi)$  with  $\arg \zeta \sim \frac{k\pi}{3}$ , for  $k = 1, 3$ , and  $j = 1, 2$ .*

By comparing Lemma 1 with [1, Theorem 2], one may observe that Lemma 1 is a refined version of [1, Theorem 2], as the error bounds in (2.9) are expressed in higher order terms. It is noteworthy that the special case of Lemma 1 for real  $\zeta$  has been considered by Dunster [8], which inspires our extension of this result to the complex plane for  $\zeta \in \mathbb{D} \cap \mathbb{S}$ .

Next, we work on case (iii) with the pole parameters  $(p, H)$  being large.

Assume  $t = p$ . Recall that  $\phi_1(\lambda)$  is the first component of  $\hat{\Phi}(\lambda)$  in (A.10). It follows from eq. (A.10) that  $\phi_1(\lambda)$  satisfies the following reduced triconfluent Heun equation (RTHE; see [21, eq. (6)] or [18, p. 108])

$$\frac{d^2 \phi_1}{d\lambda^2} = [4\lambda^3 + 2p\lambda - 28H] \phi_1. \quad (2.13)$$

Let  $\xi$  be a large parameter and write

$$p = 2C_0 \xi^{\frac{2}{5}} \quad \text{and} \quad H = -\frac{1}{7} \xi^{\frac{6}{5}}.$$

Under the scaling transformation  $\lambda = \xi^{\frac{2}{5}} z$  and  $\phi_1(\xi^{\frac{2}{5}} z) = Y(z)$ , eq. (2.13) becomes

$$\frac{d^2 Y}{dz^2} = \xi^2 \tilde{f}(z) Y, \quad (2.14)$$

where

$$\tilde{f}(z) = 4(z^3 + C_0 z + 1) = 4(z - \tilde{z}_0)(z - \tilde{z}_1)(z - \tilde{z}_2)$$

with  $\tilde{z}_0 < 0$ , and  $\tilde{z}_1, \tilde{z}_2$  being a conjugate pair. We shall assume  $\text{Im } \tilde{z}_1 \geq \text{Im } \tilde{z}_2$  without loss of generality. Define the transformation  $\eta := \eta(z)$  by

$$\frac{2}{3} \eta^{\frac{3}{2}} = \int_{\tilde{z}_1}^z \tilde{f}(t)^{\frac{1}{2}} dt. \quad (2.15)$$

Then  $\eta(z)$  is a conformal mapping in a neighbourhood of  $z = \tilde{z}_1$  and the two adjacent Stokes lines emanating from  $z = \tilde{z}_1$  to infinity. Moreover, a direct calculation of the integral in (2.15) yields

$$\frac{2}{3} \eta^{\frac{3}{2}} = \frac{4}{5} z^{\frac{5}{2}} + 2C_0 z^{\frac{1}{2}} + \tilde{E}_0 + \mathcal{O}(z^{-\frac{1}{2}}) \quad (2.16)$$

as  $z \rightarrow \infty$  uniformly for  $\arg z \in [0, \frac{4\pi}{5}]$ , where

$$\tilde{E}_0 = \int_{\tilde{z}_1}^{\infty e^{i\theta}} \left[ 2(t^3 + C_0 t + 1)^{\frac{1}{2}} - \left( 2t^{\frac{3}{2}} + C_0 t^{-\frac{1}{2}} \right) \right] dt - \left( \frac{4}{5} \tilde{z}_1^{\frac{5}{2}} + 2C_0 \tilde{z}_1^{\frac{1}{2}} \right) \quad (2.17)$$

with  $\theta \in [0, \frac{4\pi}{5}]$ . Set  $\tilde{W}(\eta, \xi) = \left(\frac{\tilde{f}(z)}{\eta}\right)^{\frac{1}{4}} Y$ , then (2.2) becomes

$$\frac{d^2 \tilde{W}}{d\eta^2} = \xi^2 \left[ \eta + \frac{\tilde{\psi}(\eta)}{\xi^2} \right] \tilde{W}, \quad (2.18)$$

where

$$\tilde{\psi}(\eta) = \frac{5}{16\eta^2} + \frac{\eta[4\tilde{f}(z)\tilde{f}''(z) - 5\tilde{f}'(z)^2]}{16\tilde{f}(z)^3} \quad (2.19)$$

is analytic at  $\eta = 0$ . Similar to Lemma 1, we have the following result.

**Lemma 2.** *Let  $\delta$  and  $\epsilon$  be fixed small positive constants and denote*

$$\tilde{\mathbb{S}} := \left\{ \eta \in \mathbb{C} : \arg \eta = \frac{k\pi}{3}, k = 1, 3 \right\}$$

and

$$\tilde{\mathbb{D}} := \left\{ \eta \in \mathbb{C} : |\eta| \leq \delta \text{ or } \frac{\pi}{3} - \epsilon < \arg \eta < \pi + \epsilon \right\}.$$

For any  $n \in \mathbb{N}$ , define  $\hat{\eta} = \eta + \tilde{\mathcal{A}}_n(\eta, \xi)$  with

$$\tilde{\mathcal{A}}_n(\eta, \xi) = \sum_{s=1}^n \frac{\tilde{a}_s(\eta)}{\xi^{2s}} \quad (2.20)$$

satisfying

$$\xi^2 \left\{ \tilde{\mathcal{A}}_n + (\eta + \tilde{\mathcal{A}}_n)(2 + \tilde{\mathcal{A}}_n') \mathcal{A}'_n \right\} + \frac{3(\tilde{\mathcal{A}}_n'')^2 - 2(1 + \tilde{\mathcal{A}}_n') \tilde{\mathcal{A}}_n'''}{4(1 + \tilde{\mathcal{A}}_n')^2} = \tilde{\psi}(\eta) + \mathcal{O}(\xi^{-2n}) \quad (2.21)$$

as  $\xi \rightarrow +\infty$  uniformly for all  $\eta \in \tilde{\mathbb{D}}$ . Assume also that each  $\tilde{a}_s(\eta)$  is analytic at  $\eta = 0$ . Then for all  $s \in \mathbb{N}$ , the limits

$$\beta_s = \lim_{\eta \rightarrow \infty} \eta^{\frac{1}{2}} \tilde{a}_s(\eta)$$

exist. Moreover, for any solution of (2.18), there exist two constants  $\tilde{C}_1$  and  $\tilde{C}_2$  such that

$$\tilde{W}(\eta, \xi) = [\tilde{C}_1 + \tilde{r}_1(\eta, \xi)] \text{Ai}_n(\eta, \xi) + [\tilde{C}_2 + \tilde{r}_2(\eta, \xi)] \text{Bi}_n(\eta, \xi), \quad (2.22)$$

where

$$\text{Ai}_n(\eta, \xi) = \left( \frac{d\hat{\eta}}{d\eta} \right)^{-\frac{1}{2}} \text{Ai} \left( \xi^{\frac{2}{3}} \hat{\eta} \right) \quad \text{and} \quad \text{Bi}_n(\eta, \xi) = \left( \frac{d\hat{\eta}}{d\eta} \right)^{-\frac{1}{2}} \text{Bi} \left( \xi^{\frac{2}{3}} \hat{\eta} \right), \quad (2.23)$$

and  $\tilde{r}_{1,2}(\eta, \xi) = \mathcal{O}(\xi^{-2n-1})$  as  $\xi \rightarrow +\infty$  uniformly for all  $\eta \in \tilde{\mathbb{D}} \cap \tilde{\mathbb{S}}$ .

**Remark 3.** *Similar to  $a_s(\zeta)$  in Lemma 1, the coefficients  $\tilde{a}_s(\eta)$  can be determined in a recursive manner by (2.21) and the fact that  $\tilde{a}_s(\eta)$  are analytic in  $\tilde{\mathbb{D}}$ . Specifically,*

$$\tilde{a}_1(\eta) = \frac{1}{2\eta^{\frac{1}{2}}} \int_0^\eta \frac{\tilde{\psi}(t)}{t^{\frac{1}{2}}} dt \quad (2.24)$$



and

$$\tilde{a}_s(\eta) = \frac{1}{2\eta^{\frac{1}{2}}} \int_0^\eta \frac{\tilde{F}_s(t)}{t^{\frac{1}{2}}} dt \quad (2.25)$$

for all  $s \geq 2$ , where  $\tilde{F}_s(t)$  are polynomials of  $\tilde{a}_1(t), \tilde{a}_2(t), \dots, \tilde{a}_{s-1}(t)$  and their first three derivatives. For example,  $\tilde{F}_1(t) = \tilde{a}_1'''(t) - 4\tilde{a}_1(t)\tilde{a}_1'(t) - 2t\tilde{a}_1'(t)^2$ .

**Remark 4.** We may regard eq. (2.14) as a special case of eq. (2.2), and so that Lemma 2 as a special case of Lemma 1. Similar to  $r_{1,2}(\zeta)$  in Lemma 1, the limits of  $\tilde{r}_1(\eta, \xi)$  and  $\tilde{r}_2(\eta, \xi)$  as  $\eta \rightarrow \infty$  with  $\arg \eta \sim \frac{k\pi}{3}$ ,  $k = 1, 3$ , exist. We shall denote  $\tilde{r}_{j,k}(\xi) = \lim_{\eta \rightarrow \infty} \tilde{r}_j(\eta, \xi)$  with  $\arg \eta \sim \frac{k\pi}{3}$ ,  $k = 1, 3$ , and  $j = 1, 2$ .

**Remark 5.** It can be shown by induction that, for all  $s \in \mathbb{N}$ , the values  $\beta_s$  are purely imaginary; see Appendix C.

## 2.2 Full asymptotic expansion of the Stokes multipliers

Using Lemma 1 and the Stokes phenomenon of the Airy functions, we obtain the full asymptotic expansion of the Stokes multipliers corresponding to the PI solution with  $y(0) = a$  and  $y'(0) = b$  as  $\xi = \left(\frac{b^2}{4} - a^3\right)^{\frac{5}{6}} \rightarrow +\infty$ .

**Theorem 1.** Assume that  $a = A(\xi)\xi^{\frac{2}{3}}$  and  $b = B(\xi)\xi^{\frac{3}{5}}$  with  $\frac{B(\xi)^2}{4} - A(\xi)^3 = 1$ . Then the Stokes multipliers corresponding to  $y(t; a, b)$  have the following asymptotic expansions:

$$\begin{aligned} s_0 &\sim 2i \exp \left\{ - \sum_{s=0}^{\infty} \frac{2 \operatorname{Re} \alpha_s(A(\xi), B(\xi))}{\xi^{s-1}} \right\} \left\{ \cos \left[ \sum_{s=0}^{\infty} - \frac{2 \operatorname{Im} \alpha_s(A(\xi), B(\xi))}{\xi^{s-1}} \right] \right\}, \\ s_1 &= -\overline{s_{-1}} \sim i \exp \left\{ \sum_{s=0}^{\infty} \frac{2\alpha_s(A(\xi), B(\xi))}{\xi^{s-1}} \right\}, \\ s_2 &= -\overline{s_3} \sim -i \exp \left\{ \sum_{s=0}^{\infty} \frac{-4i \operatorname{Im} \alpha_s(A(\xi), B(\xi))}{\xi^{s-1}} \right\} \end{aligned} \quad (2.26)$$

as  $\xi \rightarrow +\infty$ , where

$$\alpha_0(A(\xi), B(\xi)) = \frac{3}{5} \operatorname{B} \left( \frac{1}{2}, \frac{1}{3} \right) - \frac{\sqrt{3}i}{5} \operatorname{B} \left( \frac{1}{2}, \frac{1}{3} \right) \quad (2.27)$$

and  $\alpha_s(A(\xi), B(\xi))$ ,  $s = 1, 2, \dots$ , are given in Lemma 1.

**Remark 6.** When  $A(\xi)$  and  $B(\xi)$  are both fixed,  $\alpha_s(A(\xi), B(\xi))$  are all fixed and independent of  $\xi$ . In particular, when  $B(\xi) = 0$ , we have  $A(\xi) = -1$ , then according to Remark 1, we have

$$\alpha_{2s+1}(-1, 0) = 0 \quad \text{for all } s = 0, 1, 2, \dots \quad (2.28)$$

When  $A(\xi) = 0$ , we have  $B(\xi) = \pm 2$ . Hence, from (2.8), we obtain

$$\alpha_1(0, \pm 2) = \lim_{\zeta \rightarrow \infty} \frac{1}{2} \int_0^\zeta \frac{\pm \varphi(t)}{t^{\frac{1}{2}}} dt = \frac{1}{2} \int_{e^{\frac{\pi i}{3}}}^\infty \frac{g(z)}{f(z)^{\frac{1}{2}}} dz = \mp \int_{e^{\frac{\pi i}{3}}}^\infty \frac{1}{2z(z^3 + 1)^{\frac{1}{2}}} dz = \pm \frac{\pi i}{6}. \quad (2.29)$$

However, when  $A(\xi)$  or  $B(\xi)$  is not fixed, we find that  $\alpha_s(A(\xi), B(\xi))$  all depend on  $\xi$ . For instance, when  $a \neq 0$  is fixed,  $A(\xi) = a\xi^{-\frac{2}{5}}$  and  $B(\xi) = \pm \left(2 + a^3\xi^{-\frac{6}{5}}\right) + \mathcal{O}(\xi^{-\frac{12}{5}})$  as  $\xi \rightarrow +\infty$ . This implies that  $\alpha_1(A(\xi), B(\xi)) = \pm \frac{\pi i}{6} + \mathcal{O}(\xi^{-\frac{2}{5}})$  in this case. Similarly, when  $b \neq 0$  is fixed, we have  $B(\xi) = b\xi^{-\frac{3}{5}}$  and  $A(\xi) = -1 + \frac{b^2}{12}\xi^{-\frac{6}{5}} + \mathcal{O}(\xi^{-\frac{12}{5}})$  as  $\xi \rightarrow +\infty$ . This implies that  $\alpha_1(A(\xi), B(\xi)) = \mathcal{O}(\xi^{-\frac{3}{5}})$  in this case.

Next, let us consider the PI solution with large pole parameters  $(p, H) = \left(2C_0\xi^{\frac{2}{5}}, -\frac{1}{7}\xi^{\frac{6}{5}}\right)$ . Using Lemma 2 and the Stokes phenomenon of the Airy functions, we obtain the full asymptotic expansion of the Stokes multipliers corresponding to such PI solution as  $\xi \rightarrow +\infty$ .

**Theorem 2.** *Let  $C_0$  be the constant given in [15, Lemma 1.2]. Then the full asymptotic expansion of the Stokes multipliers corresponding to the PI solutions with  $(p, H) = \left(2C_0\xi^{\frac{2}{5}}, -\frac{1}{7}\xi^{\frac{6}{5}}\right)$  are given by*

$$\begin{aligned} s_0 &\sim i \left\{ 2 \cos \left[ \sum_{s=0}^{\infty} -\frac{2i\beta_s}{\xi^{2s-1}} \right] - 1 \right\}, \\ s_1 &\sim -\overline{s_{-1}} \sim i \exp \left\{ \sum_{s=0}^{\infty} \frac{2\beta_s}{\xi^{2s-1}} \right\}, \\ s_2 &\sim -\overline{s_{-2}} \sim 2 \exp \left\{ -\sum_{s=0}^{\infty} \frac{3\beta_s}{\xi^{2s-1}} \right\} \sin \left\{ \sum_{s=0}^{\infty} \frac{i\beta_s}{\xi^{2s-1}} \right\} \end{aligned} \quad (2.30)$$

as  $\xi \rightarrow +\infty$ , where  $\beta_0 = \tilde{E}_0$  defined in (2.17) and  $\beta_s, s = 1, 2, \dots$ , are given in Lemma 2.

### 2.3 Full asymptotic expansions of the nonlinear eigenvalues and pole parameters

An asymptotic expansion of the nonlinear eigenvalues of PI is proposed in [3] based on numerical simulations. Now, we can show that such a full asymptotic expansion is indeed valid as a direct consequence of Theorem 1.

**Corollary 1.** *For  $a = 0$ , there exist two sequences  $\{b_n^+\}$  and  $\{b_n^-\}$  such that the PI solutions  $y(t; a, b_n^\pm)$  are separatrix solutions, and the following asymptotic expansions hold:*

$$b_n^+ \sim 2 \left[ \frac{\sqrt{3\pi}\Gamma\left(\frac{11}{6}\right)}{\Gamma\left(\frac{1}{3}\right)} \right]^{\frac{3}{5}} \left( n - \frac{1}{6} \right)^{\frac{3}{5}} \left[ 1 + \sum_{s=2}^{\infty} \frac{B_s^+}{\left(n - \frac{1}{6}\right)^s} \right] \quad (2.31)$$

and

$$b_n^- \sim -2 \left[ \frac{\sqrt{3\pi}\Gamma\left(\frac{11}{6}\right)}{\Gamma\left(\frac{1}{3}\right)} \right]^{\frac{3}{5}} \left( n - \frac{5}{6} \right)^{\frac{3}{5}} \left[ 1 + \sum_{s=2}^{\infty} \frac{B_s^-}{\left(n - \frac{5}{6}\right)^s} \right] \quad (2.32)$$

as  $n \rightarrow \infty$ , where  $B_s^\pm, s = 2, 3, \dots$ , can be expressed in terms of  $\alpha_k(0, \pm 2), k = 0, 1, 2, \dots, s$ .

*Proof.* When  $a = 0$ , we have  $B(\xi) = \pm 2$ . Regard the Stokes multipliers  $s_k$  as functions of  $\xi$ . According to (2.26), it is readily seen that there exist two sequences  $\{\xi_n^+\}$  and  $\{\xi_n^-\}$  such that  $s_0(\xi_n^\pm) = 0$  and

$$-2 \operatorname{Im} \alpha_0(0, \pm 2) \xi_n^\pm - 2 \operatorname{Im} \alpha_1(0, \pm 2) - \sum_{s=1}^{\infty} \frac{2 \operatorname{Im} \alpha_{s+1}(0, \pm 2)}{(\xi_n^\pm)^s} \sim \left(n - \frac{1}{2}\right) \pi \quad (2.33)$$

as  $n \rightarrow \infty$ . This implies that

$$\xi_n^\pm \sim -\frac{\left(n - \frac{1}{2} \pm \frac{1}{3}\right) \pi}{2 \operatorname{Im} \alpha_0(0, \pm 2)} - \sum_{s=1}^{\infty} \frac{2 \operatorname{Im} \alpha_{s+1}(0, \pm 2)}{2 \operatorname{Im} \alpha_0(0, \pm 2) (\xi_n^\pm)^s} \quad (2.34)$$

as  $n \rightarrow \infty$ . Noting that  $b = B(\xi) \xi^{\frac{3}{5}} = \pm 2 \xi^{\frac{3}{5}}$ , then  $b_n^\pm = \pm 2 (\xi_n^\pm)^{\frac{3}{5}}$ . We further obtain (2.31) and (2.32) respectively, where

$$B_2^\pm = -\frac{4 \operatorname{Im} \alpha_0(0, \pm 2) \operatorname{Im} \alpha_2(0, \pm 2)}{\pi^2} = \frac{4\sqrt{3}}{5} B\left(\frac{1}{2}, \frac{1}{3}\right) \operatorname{Im} \alpha_2(0, \pm 2). \quad (2.35)$$

and for  $s > 2$ ,  $B_s^\pm$  can be expressed in term of  $\alpha_0(0, \pm 2), \alpha_1(0, \pm 2), \dots, \alpha_s(0, \pm 2)$ .  $\square$

**Remark 7.** In Corollary 1, if  $a \neq 0$  is fixed, according to the analysis in Remark 6, we can also obtain the following asymptotic expansion of  $(b_n^\pm)^2 - 4a^3$  as  $n \rightarrow +\infty$

$$\left((b_n^\pm)^2 - 4a^3\right)^{\frac{1}{2}} \sim \pm 2 \left[ \frac{\sqrt{3\pi} \Gamma\left(\frac{11}{6}\right)}{\Gamma\left(\frac{1}{3}\right)} \right]^{\frac{3}{5}} \left(n - \gamma_n^\pm\right)^{\frac{3}{5}} \left[ 1 + \sum_{s=2}^{\infty} \frac{B_s^\pm(n)}{(n - \gamma_n^\pm)^s} \right], \quad (2.36)$$

which is similar to (2.31) and (2.32). Here,  $\gamma_n^\pm = -\frac{1}{2} \pm \frac{1}{3} + \mathcal{O}(n^{-\frac{2}{5}})$  as  $n \rightarrow +\infty$  and  $B_s^\pm(n)$  are all bounded but depend on  $n$ .

When  $b = 0$ , then  $A(\xi) = -1$  and  $B(\xi) = 0$ . Noting that  $\alpha_{2s+1}(-1, 0) = 0$ , we have the following result.

**Corollary 2.** For  $b = 0$ , there exists a sequence  $\{a_n\}$  such that the PI solutions  $y(t; a_n, b)$  are separatrix solutions and

$$a_n \sim -\left[ \frac{\sqrt{3\pi} \Gamma\left(\frac{11}{6}\right)}{\Gamma\left(\frac{1}{3}\right)} \right]^{\frac{2}{5}} \left(n - \frac{1}{2}\right)^{\frac{2}{5}} \left[ 1 + \sum_{s=1}^{\infty} \frac{A_s}{\left(n - \frac{1}{2}\right)^{2s}} \right] \quad (2.37)$$

as  $n \rightarrow \infty$ , where  $A_s$  can be expressed in terms of  $\alpha_{2k}(-1, 0), k = 1, 2, \dots, s$ .

**Remark 8.** In Corollary 2, if  $b \neq 0$  is fixed, according to the analysis in Remark 6, we can also obtain the following asymptotic expansion of  $a_n$  as  $n \rightarrow \infty$

$$\left(a_n^3 - \frac{b^2}{4}\right)^{\frac{1}{3}} \sim -\left[ \frac{\sqrt{3\pi} \Gamma\left(\frac{11}{6}\right)}{\Gamma\left(\frac{1}{3}\right)} \right]^{\frac{2}{5}} \left(n - \mu_n\right)^{\frac{2}{5}} \left[ 1 + \sum_{s=1}^{\infty} \frac{A_s(n)}{(n - \mu_n)^{2s}} \right], \quad (2.38)$$

which is an analogue to (2.37). Here,  $\mu_n = \frac{1}{2} + \mathcal{O}(n^{-\frac{3}{5}})$  as  $n \rightarrow \infty$  and  $A_s(n)$  are all bounded but depend on  $n$ .

Let us turn to the real tritronquée solution of PI. We can obtain the full asymptotic expansions for the pole parameters  $(p_n, H_n)$  as  $n \rightarrow \infty$ , where  $p_n$  denotes the  $n$ -th pole of this solution. It is a direct consequence of the combination of Theorem 2 and the fact that the Stokes multipliers corresponding to PI's real tritronquée solution are  $s_1 = s_{-1} = i, s_2 = s_{-2} = 0, s_0 = i$ .

**Corollary 3.** *Let  $(p_n, H_n)$  be the pole parameters of the  $n$ -th real pole of PI's real tritronquée solution, then*

$$\begin{aligned} p_n &\sim 2C_0 \left( \frac{4n-2}{\kappa^2(C_0)} \right)^{\frac{2}{5}} \left[ 1 + \sum_{s=1}^{\infty} \frac{\rho_s}{\left(n - \frac{1}{2}\right)^{2s}} \right], \\ H_n &\sim -\frac{1}{7} \left( \frac{4n-2}{\kappa^2(C_0)} \right)^{\frac{6}{5}} \left[ 1 + \sum_{s=1}^{\infty} \frac{\mathcal{H}_s}{\left(n - \frac{1}{2}\right)^{2s}} \right] \end{aligned} \quad (2.39)$$

as  $n \rightarrow \infty$ , where  $\rho_s$  and  $\mathcal{H}_s$  can be expressed in terms of  $\beta_1, \dots, \beta_s$ . Here,  $C_0$  and  $\kappa^2(C_0)$  are given in [15, Lemma 2.1].

### 3 Uniform asymptotics and proof of Lemma 1

To prove Lemma 1, we first use induction to show that

$$a_s(\zeta) = \mathcal{O}(\zeta^{-\frac{1}{2}}), \quad a_s^{(k)}(\zeta) = \mathcal{O}(\zeta^{-\frac{3}{2}}), \quad k = 1, 2, \dots, \quad (3.1)$$

as  $\zeta \rightarrow \infty$  in  $\mathbb{D}$  for all  $s \in \mathbb{N}$ .

Let  $s = 1, 2$ . It follows from (2.4) and (2.6) that

$$\varphi(\zeta) = \mathcal{O}(\zeta^{-\frac{7}{5}}), \quad \psi(\zeta) = \mathcal{O}(\zeta^{-2}), \quad \text{as } \zeta \rightarrow \infty, \quad (3.2)$$

which, together with the explicit representations of  $a_1(\zeta)$  and  $a_2(\zeta)$  in (2.11), leads to (3.1).

Suppose (3.1) holds for  $s = 1, 2, \dots, m-1$  with  $m > 2$ . It then follows from (2.8) and the analyticity of  $a_m(\zeta)$  in  $\mathbb{D}$  that

$$a_m(\zeta) = \frac{1}{2\zeta^{\frac{1}{2}}} \int_0^\zeta \frac{F_m(t)}{t^{\frac{1}{2}}} dt, \quad (3.3)$$

where  $F_m(\zeta)$  is a polynomial of  $a'_s(\zeta), a''_s(\zeta), a'''_s(\zeta), s = 1, 2, \dots, m-1$ . Therefore,  $F_m(\zeta)$  and its derivatives all have an estimate of  $\mathcal{O}(\zeta^{-\frac{3}{2}})$  as  $\zeta \rightarrow \infty$ . Hence, the integral in (3.3) is convergent when  $\zeta \rightarrow \infty$ , and then  $a_m(\zeta) = \mathcal{O}(\zeta^{-\frac{1}{2}})$ . According to (2.8) [or (3.3)], we also have

$$a_m(\zeta) + 2\zeta a'_m(\zeta) = F_s(\zeta) = \mathcal{O}(\zeta^{-\frac{3}{2}}), \quad (3.4)$$

which implies that the derivative of  $a_m(\zeta)$  of any order has the estimate of  $\mathcal{O}(\zeta^{-\frac{3}{2}})$  as  $\zeta \rightarrow \infty$ . Therefore, (3.1) holds for any  $s = 1, 2, \dots$  by induction.

It follows from (2.7) and (3.1) that  $\zeta^{\frac{1}{2}}\mathcal{A}_n(\zeta, \xi) = \mathcal{O}(\xi^{-1})$  as  $\xi \rightarrow +\infty$  uniformly for all  $\zeta \in \mathbb{D}$ . Hence,

$$e^{\xi\hat{\zeta}^{\frac{3}{2}}} = \exp\{\xi\zeta^{\frac{3}{2}}(1 + \zeta^{-1}\mathcal{A}_n(\zeta, \xi))^{\frac{3}{2}}\} = \exp\{\xi\zeta^{\frac{3}{2}}\} \cdot \mathcal{O}(\xi\zeta^{\frac{1}{2}}\mathcal{A}_n(\zeta, \xi)) \quad (3.5)$$

as  $\xi \rightarrow +\infty$ , where  $\zeta \in \mathbb{D}$ . Note that  $\zeta^{\frac{3}{2}}$  is purely imaginary when  $\zeta \in \mathbb{D} \cap \mathbb{S}$ . Therefore,  $e^{\xi\hat{\zeta}^{\frac{3}{2}}} = \mathcal{O}(1)$  as  $\xi \rightarrow +\infty$  uniformly for  $\zeta \in \mathbb{D} \cap \mathbb{S}$ .

Now, we are in a position to prove Lemma 1.

**Proof of Lemma 1** It is clear from (3.1) that the limits  $\lim_{\zeta \rightarrow \infty} a_s(\zeta)$  exist for all  $s = 1, 2, \dots$ .

It is also evident that  $\text{Ai}_n(\zeta, \xi)$  and  $\text{Bi}_n(\zeta, \xi)$  satisfy the equation

$$\frac{d^2 w}{d\zeta^2} = \xi^2 [\zeta + Q_n(\zeta, \xi)] w, \quad (3.6)$$

where

$$Q_n(\zeta, \xi) = \{\mathcal{A}_n + (\zeta + \mathcal{A}_n)(2 + \mathcal{A}'_n)\mathcal{A}'_n\} + \frac{3\mathcal{A}_n''^2 - 2(1 + \mathcal{A}'_n)\mathcal{A}_n'''}{4\xi^2(1 + \mathcal{A}'_n)^2}. \quad (3.7)$$

It follows from (2.7) and (3.4) that  $\mathcal{A}_n + 2\zeta\mathcal{A}'_n = \mathcal{O}(\zeta^{-\frac{3}{2}})$ , which, together with (3.1), imply that  $Q_n(\zeta, \xi) = \mathcal{O}(\zeta^{-\frac{3}{2}})$  as  $\zeta \rightarrow \infty$  in  $\mathbb{D}$ . For any solution  $W(\zeta, \xi)$  of (2.18), we have

$$\frac{d^2 W}{d\zeta^2} = \xi^2 [\zeta + Q_n(\zeta, \xi)] W + [\xi\varphi(\zeta) + \psi(\zeta) - \xi^2 Q_n(\zeta, \xi)] W. \quad (3.8)$$

Then there exist two constants  $C_1$  and  $C_2$  such that for all  $\zeta \in \mathbb{D} \cap \mathbb{S}$

$$\begin{aligned} W(\zeta, \xi) = & C_1 \text{Ai}_n(\zeta, \xi) + C_2 \text{Bi}_n(\zeta, \xi) \\ & + \int_0^\zeta \frac{\text{Ai}_n(t, \xi)\text{Bi}_n(\zeta, \xi) - \text{Ai}_n(\zeta, \xi)\text{Bi}_n(t, \xi)}{\mathcal{W}(\text{Ai}_n(t, \xi), \text{Bi}_n(t, \xi))} W(t, \xi) R(t, \xi) dt \end{aligned} \quad (3.9)$$

where

$$R(\zeta, \xi) = [\xi\varphi(\zeta) + \psi(\zeta) - \xi^2 Q_n(\zeta, \xi)] = \mathcal{O}(\zeta^{-\frac{3}{2}}), \quad \zeta \rightarrow \infty, \zeta \in \mathbb{D} \quad (3.10)$$

and

$$R(\zeta, \xi) = [\xi\varphi(\zeta) + \psi(\zeta) - \xi^2 Q_n(\zeta, \xi)] = \mathcal{O}(\xi^{-2n+1}), \quad \xi \rightarrow +\infty \quad (3.11)$$

uniformly for all  $\zeta \in \mathbb{D} \cap \mathbb{S}$ . Applying the iterative argument used in the proof of [1, Theorem 2], we see that  $W(\zeta, \xi)$  is bounded on  $\mathbb{D} \cap \mathbb{S}$ . A combination of this fact and (3.10) yields that the last term in (3.9) is integrable. Using the Wronskian for Airy functions ([17, eq. (9.2.7)]), we have

$$\mathcal{W}(\text{Ai}_n(t, \xi), \text{Bi}_n(t, \xi)) \sim \frac{1}{\pi} \xi^{\frac{2}{3}}, \quad \xi \rightarrow +\infty, \quad (3.12)$$

we obtain (2.9), and

$$r_{1,2}(\zeta, \xi) = \mathcal{O}(\xi^{-2n+\frac{1}{3}}) \quad (3.13)$$

as  $\xi \rightarrow +\infty$  uniformly for all  $\zeta \in \mathbb{D} \cap \mathbb{S}$ .

To get the desired approximations of  $r_{1,2}(\zeta, \xi)$  as stated in Lemma 1, we need the following approximation of  $W(\zeta, \xi)$

$$W(\zeta, \xi) \leq M(|C_1||\text{Ai}_n(\zeta, \xi)| + |C_2||\text{Bi}_n(\zeta, \xi)|) \quad (3.14)$$

which can be obtained as a direct consequence of (2.9) and (3.13). Let  $\zeta^*$  be a point on the integration contour in (3.9) satisfying  $|\zeta^*| \sim \xi^{-\frac{1}{3}}$ . A comparison of (3.9) and (2.9) leads to

$$\begin{aligned} r_1(\zeta, \xi) &= - \int_0^\zeta \frac{\text{Bi}_n(t, \xi)W(t, \xi)R(t, \xi)}{\mathcal{W}(\text{Ai}_n(t, \xi), \text{Bi}_n(t, \xi))} dt \\ &= - \int_0^{\zeta^*} \frac{\text{Bi}_n(t, \xi)W(t, \xi)R(t, \xi)}{\mathcal{W}(\text{Ai}_n(t, \xi), \text{Bi}_n(t, \xi))} dt - \int_{\zeta^*}^\zeta \frac{\text{Bi}_n(t, \xi)W(t, \xi)R(t, \xi)}{\mathcal{W}(\text{Ai}_n(t, \xi), \text{Bi}_n(t, \xi))} dt \\ &= I_1 + I_2. \end{aligned} \quad (3.15)$$

The integrand of  $I_1$  is bounded by  $\mathcal{O}(\xi^{-2n+\frac{1}{3}})$ , then

$$I_1 = \mathcal{O}(\xi^{-2n}) \quad \text{as } \xi \rightarrow +\infty. \quad (3.16)$$

To estimate  $I_2$ , we use (3.11), (3.12) and (3.14) to get

$$|I_2| \leq \frac{4\pi M^2}{\xi} \int_{\zeta^*}^\zeta \left| t^{-\frac{1}{2}} R(t, \xi) \right| dt \leq \tilde{M} \xi^{-2n} \quad (3.17)$$

where  $\tilde{M} > 0$  is a constant. This fact, together with (3.10), (3.16) and (3.17), lead to  $r_1(\zeta, \xi) = \mathcal{O}(\xi^{-2n})$  as  $\xi \rightarrow +\infty$ . In a similar manner, we can show that  $r_2(\zeta, \xi)$  has the same estimate as  $r_1(\zeta, \xi)$ ; completing the proof of Lemma 1.

## 4 Asymptotic matching and proof of Theorems 1 and 2

In this section, we derive full asymptotic expansions of the Stokes multipliers  $s_k$ 's in case (ii) [*resp.* case (iii)], which is done by matching the uniform asymptotic approximation of  $W(\zeta, \xi)$  stated in Lemma 1 [*resp.* Lemma 2] and the asymptotic expansions of the canonical solutions  $\hat{\Phi}_k$  in (A.11). The approach is similar to that used in [14, 15], and hence we only state some crucial steps, referring to these two articles for details.

### Proof of Theorem 1

Since the Stokes multipliers  $s_k$ 's depend on  $\xi$ , hence we will only take the limit  $\lambda \rightarrow \infty$  when we do the asymptotic matching between the Airy functions and  $\Phi_k$ 's. Recall the asymptotics for the Airy functions for large arguments (see [17, eqs. (9.2.12), (9.7.5), (9.2.10)])

$$\begin{cases} \text{Ai}(Z) \sim \frac{1}{2\sqrt{\pi}} Z^{-\frac{1}{4}} e^{-\frac{2}{3}Z^{\frac{3}{2}}}, & \arg Z \in (-\pi, \pi), \\ \text{Ai}(Z) \sim \frac{1}{2\sqrt{\pi}} Z^{-\frac{1}{4}} e^{-\frac{2}{3}Z^{\frac{3}{2}}} + \frac{i}{2\sqrt{\pi}} Z^{-\frac{1}{4}} e^{\frac{2}{3}Z^{\frac{3}{2}}}, & \arg Z \in \left(\frac{\pi}{3}, \frac{5\pi}{3}\right), \\ \text{Ai}(Z) \sim \frac{1}{2\sqrt{\pi}} Z^{-\frac{1}{4}} e^{-\frac{2}{3}Z^{\frac{3}{2}}} - \frac{i}{2\sqrt{\pi}} Z^{-\frac{1}{4}} e^{\frac{2}{3}Z^{\frac{3}{2}}}, & \arg Z \in \left(-\frac{5\pi}{3}, -\frac{\pi}{3}\right) \end{cases} \quad (4.1)$$

and

$$\begin{cases} \text{Bi}(Z) \sim \frac{i}{2\sqrt{\pi}} Z^{-\frac{1}{4}} e^{-\frac{2}{3}Z^{\frac{3}{2}}} + \frac{1}{\sqrt{\pi}} Z^{-\frac{1}{4}} e^{\frac{2}{3}Z^{\frac{3}{2}}}, & \arg Z \in \left(-\frac{\pi}{3}, \pi\right), \\ \text{Bi}(Z) \sim \frac{i}{2\sqrt{\pi}} Z^{-\frac{1}{4}} e^{-\frac{2}{3}Z^{\frac{3}{2}}} + \frac{1}{2\sqrt{\pi}} Z^{-\frac{1}{4}} e^{\frac{2}{3}Z^{\frac{3}{2}}}, & \arg Z \in \left(\frac{\pi}{3}, \frac{5\pi}{3}\right), \\ \text{Bi}(Z) \sim -\frac{i}{2\sqrt{\pi}} Z^{-\frac{1}{4}} e^{-\frac{2}{3}Z^{\frac{3}{2}}} + \frac{1}{\sqrt{\pi}} Z^{-\frac{1}{4}} e^{\frac{2}{3}Z^{\frac{3}{2}}}, & \arg Z \in \left(-\pi, \frac{\pi}{3}\right). \end{cases} \quad (4.2)$$

When  $\lambda \rightarrow \infty$  with  $\arg \lambda \sim \frac{\pi}{5}$ , it follows from  $\lambda = \xi^{\frac{2}{5}}z$  and (2.4) that  $\arg \zeta \sim \frac{\pi}{3}$ . Substituting (2.4) into (4.1) and (4.2), and using  $\lambda = \xi^{\frac{2}{5}}z$  again, we get

$$\begin{cases} \left(\frac{\zeta}{f(z)}\right)^{\frac{1}{4}} \text{Ai}_n(\xi^{\frac{2}{3}}\zeta) = \left(\frac{\zeta}{f(z)}\right)^{\frac{1}{4}} \left(\frac{d\hat{\zeta}}{d\zeta}\right)^{-\frac{1}{2}} \text{Ai}\left(\xi^{\frac{2}{3}}\hat{\zeta}\right) \sim c_1 \frac{-1}{\sqrt{2}} \lambda^{-\frac{3}{4}} e^{-\frac{4}{5}\lambda^{\frac{5}{2}}}, \\ \left(\frac{\zeta}{f(z)}\right)^{\frac{1}{4}} \text{Bi}_n(\xi^{\frac{2}{3}}\zeta) = \left(\frac{\zeta}{f(z)}\right)^{\frac{1}{4}} \left(\frac{d\hat{\zeta}}{d\zeta}\right)^{-\frac{1}{2}} \text{Bi}\left(\xi^{\frac{2}{3}}\hat{\zeta}\right) \sim ic_1 \frac{-1}{\sqrt{2}} \lambda^{-\frac{3}{4}} e^{-\frac{4}{5}\lambda^{\frac{5}{2}}} + 2c_2 \frac{1}{\sqrt{2}} \lambda^{-\frac{3}{4}} e^{\frac{4}{5}\lambda^{\frac{5}{2}}} \end{cases} \quad (4.3)$$

as  $\lambda \rightarrow \infty$  (and  $\zeta \rightarrow \infty$  accordingly), where

$$\begin{aligned} c_1 &= \frac{-1}{2\sqrt{\pi}} \xi^{-\frac{2}{15}} \exp \left\{ -\sum_{s=0}^n \alpha_s(A(\xi), B(\xi)) \xi^{-s+1} \right\}, \\ c_2 &= \frac{1}{2\sqrt{\pi}} \xi^{-\frac{2}{15}} \exp \left\{ \sum_{s=0}^n \alpha_s(A(\xi), B(\xi)) \xi^{-s+1} \right\}. \end{aligned} \quad (4.4)$$

From (A.4), a straightforward calculation yields

$$\left( (\hat{\Phi}_k)_{21}, (\hat{\Phi}_k)_{22} \right) \sim \left( \frac{1}{\sqrt{2}} \lambda^{-\frac{3}{4}} e^{\frac{4}{5}\lambda^{\frac{5}{2}}}, \frac{-1}{\sqrt{2}} \lambda^{-\frac{3}{4}} e^{-\frac{4}{5}\lambda^{\frac{5}{2}}} \right), \quad k \in \mathbb{Z} \quad (4.5)$$

as  $\lambda \rightarrow \infty$ . Substituting (4.3) into (2.9), noting that  $Y = \left(\frac{\zeta}{f(z)}\right)^{\frac{1}{4}}$  and  $Y$  is the first component of  $\hat{\Phi}$ , and then comparing the resulting equation with (4.5), we get

$$\begin{aligned} W &= [C_1 + r_{1,1}(\xi)]c_1(\hat{\Phi}_1)_{22} + [C_2 + r_{2,1}(\xi)][ic_1(\hat{\Phi}_1)_{22} + 2c_2(\hat{\Phi}_1)_{21}] \\ &= [C_2 + r_{2,1}(\xi)]2c_2(\hat{\Phi}_1)_{21} + \{[C_1 + r_{1,1}(\xi)]c_1 + [C_2 + r_{2,1}(\xi)]ic_1\}(\hat{\Phi}_1)_{22} \end{aligned} \quad (4.6)$$

with  $\lambda \in \Omega_1$ .

When  $\lambda \rightarrow \infty$  with  $\arg \lambda \sim \frac{3\pi}{5}$ , one has  $\arg \zeta \sim \pi$ . Using the asymptotics of the Airy functions with  $\arg Z \sim \pi$  in (4.1) and (4.2), we have

$$\begin{cases} \left(\frac{\zeta}{f(z)}\right)^{\frac{1}{4}} \text{Ai}_n(\xi^{\frac{2}{3}}\zeta) = \left(\frac{\zeta}{f(z)}\right)^{\frac{1}{4}} \left(\frac{d\hat{\zeta}}{d\zeta}\right)^{-\frac{1}{2}} \text{Ai}\left(\xi^{\frac{2}{3}}\hat{\zeta}\right) \sim c_1 \frac{-1}{\sqrt{2}} \lambda^{-\frac{1}{4}} e^{-\frac{4}{5}\lambda^{\frac{5}{2}}} + ic_2 \frac{1}{\sqrt{2}} \lambda^{-\frac{1}{4}} e^{\frac{4}{5}\lambda^{\frac{5}{2}}}, \\ \left(\frac{\zeta}{f(z)}\right)^{\frac{1}{4}} \text{Bi}_n(\xi^{\frac{2}{3}}\zeta) = \left(\frac{\zeta}{f(z)}\right)^{\frac{1}{4}} \left(\frac{d\hat{\zeta}}{d\zeta}\right)^{-\frac{1}{2}} \text{Ai}\left(\xi^{\frac{2}{3}}\hat{\zeta}\right) \sim ic_1 \frac{-1}{\sqrt{2}} \lambda^{-\frac{1}{4}} e^{-\frac{4}{5}\lambda^{\frac{5}{2}}} + c_2 \frac{1}{\sqrt{2}} \lambda^{-\frac{1}{4}} e^{\frac{4}{5}\lambda^{\frac{5}{2}}} \end{cases} \quad (4.7)$$

as  $\lambda \rightarrow \infty$  (and  $\zeta \rightarrow \infty$  accordingly). In a similar manner for deriving (4.6), we get

$$\begin{aligned} W &= \{[C_1 + r_{1,2}(\xi)]ic_2 + [C_2 + r_{2,2}(\xi)]c_2\}(\Phi_2)_{21} \\ &\quad + \{[C_1 + r_{1,2}(\xi)]c_1 + [C_2 + r_{2,2}(\xi)]ic_1\}(\Phi_2)_{22} \end{aligned} \quad (4.8)$$

$\lambda \in \Omega_2$ . Combining (4.6) with (4.8), and noting that

$$((\Phi_2)_{21}, (\Phi_2)_{22}) = ((\Phi_1)_{21}, (\Phi_1)_{22}) \begin{pmatrix} 1 & s_1 \\ 0 & 1 \end{pmatrix},$$

we further obtain

$$r_{1,1}(\xi) - r_{1,2}(\xi) = i(r_{2,2}(\xi) - r_{2,1}(\xi)) \quad (4.9)$$

and

$$\begin{aligned} s_1 &= \frac{(C_2 - iC_1)c_2 + (2r_{2,1}(\xi) - ir_{1,2}(\xi) - r_{2,2}(\xi))c_2}{(C_1 + iC_2)c_1 + r_{1,2}(\xi)c_1 + r_{2,2}(\xi)ic_1} \\ &= -\frac{ic_2}{c_1} + \frac{2(r_{2,1}(\xi) - r_{2,2}(\xi))c_2}{(C_1 + iC_2 + r_{1,2}(\xi) + ir_{2,2}(\xi))c_1}. \end{aligned} \quad (4.10)$$

By Lemma 1,  $r_{j,k}(\xi) = \mathcal{O}\left(\frac{|C_1|+|C_2|}{\xi^n}\right)$  as  $\xi \rightarrow +\infty$  for  $j = 1, 2$  and  $k = 1, 2$ . Hence,

$$s_1 = i \exp \left\{ \sum_{s=0}^{2n} 2\alpha_s(A(\xi), B(\xi))\xi^{-s+1} \right\} (1 + \mathcal{O}(\xi^{-2n})) \quad (4.11)$$

as  $\xi \rightarrow +\infty$ .

The asymptotic expansion of  $s_{-1}$  in (2.26) follows from the conjugate relation  $s_{-1} = \overline{s_1}$ , and the asymptotic expansions for other Stokes multipliers follow from the constraint  $s_k = i(1 + s_{k+2}s_{k+3})$ .

### Proof of Theorem 2

The full asymptotic expansion of  $s_1$  and  $s_{-1}$  in Theorem 2 can be derived similarly to those in Theorem 1. To obtain the full asymptotic expansion of the other Stokes multipliers, we need to use the constraint  $s_k = i(1 + s_{k+2}s_{k+3})$  and note that  $\beta_s(A(\xi), B(\xi))$  are purely imaginary for all  $s \in \mathbb{N}$ .

## A Monodromy theory of PI

Painlevé equations can be expressed as compatibility conditions of Lax pairs. In particular, consider the Lax pair of PI (see [13])

$$\begin{cases} \frac{\partial \Psi}{\partial \lambda} = \left\{ (4\lambda^4 + t + 2y^2)\sigma_3 - i(4y\lambda^2 + t + 2y^2)\sigma_2 - \left(2y_t\lambda + \frac{1}{2\lambda}\right)\sigma_1 \right\} \Psi := \mathcal{A}(\lambda)\Psi, \\ \frac{\partial \Psi}{\partial t} = \left\{ \left(\lambda + \frac{y}{\lambda}\right)\sigma_3 - \frac{iy}{\lambda}\sigma_2 \right\} \Psi := \mathcal{B}(\lambda)\Psi, \end{cases} \quad (A.1)$$

where

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

are the Pauli matrices and  $y_t = \frac{dy}{dt}$ . If the Lax pair satisfies the compatibility condition  $\frac{\partial^2 \Psi}{\partial t \partial \lambda} = \frac{\partial^2 \Psi}{\partial \lambda \partial t}$ , then  $y = y(t)$  is a solution of the PI equation (1.1).



Under the transformation

$$\Phi(\lambda) = \lambda^{\frac{1}{4}\sigma_3} \frac{\sigma_3 + \sigma_1}{\sqrt{2}} \Psi(\sqrt{\lambda}), \quad (\text{A.2})$$

the first equation of (A.1) becomes

$$\frac{\partial \Phi}{\partial \lambda} = \begin{pmatrix} y_t & 2\lambda^2 + 2y\lambda - t + 2y^2 \\ 2(\lambda - y) & -y_t \end{pmatrix} \Phi. \quad (\text{A.3})$$

Following [13] (see also [12]), the only singularity of equation (A.3) is the irregular singular point at  $\lambda = \infty$ , and there exist canonical solutions  $\Phi_k(\lambda)$ ,  $k \in \mathbb{Z}$ , of (A.3) with the following asymptotic expansion

$$\Phi_k(\lambda, t) = \lambda^{\frac{1}{4}\sigma_3} \frac{\sigma_3 + \sigma_1}{\sqrt{2}} \left( I + \frac{\mathcal{H}}{\lambda} + \mathcal{O}\left(\frac{1}{\lambda^2}\right) \right) e^{(\frac{1}{5}\lambda^{\frac{5}{2}} + t\lambda^{\frac{1}{2}})\sigma_3}, \quad \lambda \rightarrow \infty, \quad \lambda \in \Omega_k, \quad (\text{A.4})$$

uniformly for all  $t$  bounded away from poles, where  $\mathcal{H} = -(\frac{1}{2}y_t^2 - 2y^3 - ty)\sigma_3$ , and the canonical sectors are

$$\Omega_k = \left\{ \lambda \in \mathbb{C} : \arg \lambda \in \left( -\frac{3\pi}{5} + \frac{2k\pi}{5}, \frac{\pi}{5} + \frac{2k\pi}{5} \right) \right\}, \quad k \in \mathbb{Z}.$$

These canonical solutions are related by

$$\Phi_{k+1} = \Phi_k S_k, \quad S_{2k-1} = \begin{pmatrix} 1 & s_{2k-1} \\ 0 & 1 \end{pmatrix}, \quad S_{2k} = \begin{pmatrix} 1 & 0 \\ s_{2k} & 1 \end{pmatrix}, \quad (\text{A.5})$$

where  $s_k$ 's are the *Stokes multipliers*, and independent of  $\lambda$  and  $t$  according to the isomonodromy condition. The Stokes multipliers are subject to the constraints

$$s_{k+5} = s_k \quad \text{and} \quad s_k = i(1 + s_{k+2}s_{k+3}), \quad k \in \mathbb{Z}. \quad (\text{A.6})$$

Moreover, regarding  $s_k$ 's as functions of  $(t, y(t), y'(t))$ , they also satisfy (see [11, eq. (13)])

$$s_k(t, y(t), y'(t)) = \overline{-s_{-k}(\bar{t}, \overline{y(t)}, \overline{y'(t)})}, \quad k \in \mathbb{Z}, \quad (\text{A.7})$$

where  $\bar{z}$  stands for the complex conjugate of  $z$ . From (A.6), it is readily seen that, generically, two of the Stokes multipliers determine all others. The derivation of (A.4), (A.5), and (A.6), along with more details about the Lax pairs, can be found in [9].

For simplicity, we make a further transformation:

$$\hat{\Phi}(\lambda, t) = G(\lambda, t)\Phi(\lambda, t) \quad (\text{A.8})$$

with

$$G(\lambda, t) = \begin{bmatrix} 0 & 1 \\ 1 & -\frac{1}{2} \left( -y_t + \frac{1}{2(\lambda-y)} \right) \end{bmatrix} (\lambda - y)^{\frac{\sigma_3}{2}}. \quad (\text{A.9})$$

Then,  $\hat{\Phi}(\lambda, t)$  satisfies

$$\frac{d}{d\lambda}\hat{\Phi}(\lambda, t) = \begin{bmatrix} 0 & 2 \\ V(\lambda, t) & 0 \end{bmatrix} \hat{\Phi}(\lambda, t), \quad (\text{A.10})$$

where

$$2V(\lambda, t) = \begin{cases} y_t^2 + 4\lambda^3 + 2\lambda t - 2yt - 4y^3 - \frac{yt}{\lambda - y} + \frac{3}{4} \frac{1}{(\lambda - y)^2}, & t \neq p, \\ 4\lambda^3 + 2p\lambda - 28H, & t = p, \end{cases}$$

$p$  is a pole of the PI solution and  $H$  is the parameter in the Laurent series (1.4).

Now, the approximation of  $\Phi(\lambda, t)$  in (A.4) is transformed to

$$\hat{\Phi}_k(\lambda, t) = \frac{\lambda^{-\frac{3}{4}\sigma_3}}{\sqrt{2}} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \left( I + \mathcal{O}(\lambda^{-\frac{1}{2}}) \right) e^{(\frac{4}{5}\lambda^{\frac{5}{2}} + t\lambda^{\frac{1}{2}})\sigma_3} \quad (\text{A.11})$$

as  $\lambda \rightarrow \infty$  with  $\lambda \in \Omega_k$  when  $t$  is bounded away from  $p$ . When  $t \rightarrow p$ , according to [4], (A.11) should be replaced by

$$\hat{\Phi}_k(\lambda, p) = \frac{\lambda^{-\frac{3}{4}\sigma_3}}{\sqrt{2}} \begin{bmatrix} -i & -i \\ -i & i \end{bmatrix} \left( I + \mathcal{O}(\lambda^{-\frac{1}{2}}) \right) e^{(\frac{4}{5}\lambda^{\frac{5}{2}} + p\lambda^{\frac{1}{2}})\sigma_3} \quad (\text{A.12})$$

as  $\lambda \rightarrow \infty$  with  $\lambda \in \Omega_k$ ; see [4, Corollary A.8].

## B Asymptotics of PI solutions as $t \rightarrow -\infty$

The asymptotic behavior of the PI solution as  $t \rightarrow -\infty$  has been extensively studied by Kapaev [11] using isomonodromy theory. There are three types of solutions for the PI equation according to the asymptotic behavior of  $y(t)$  as  $t \rightarrow -\infty$  as follows:

(A) a two-parameter family of solutions, oscillating about the parabola  $y = -\sqrt{-t/6}$  and satisfying

$$y = -\sqrt{-\frac{t}{6}} + d(-t)^{-\frac{1}{8}} \cos \left[ 24^{\frac{1}{4}} \left( \frac{4}{5}(-t)^{\frac{5}{4}} - \frac{5}{8}d^2 \log(-t) + \varphi \right) \right] + \mathcal{O}\left(t^{-\frac{5}{8}}\right) \quad (\text{B.1})$$

as  $t \rightarrow -\infty$ , where

$$\begin{cases} 24^{\frac{1}{4}}d^2 = -\frac{1}{\pi} \log |s_0|, \\ 24^{\frac{1}{4}}\theta = -\arg s_3 - 24^{\frac{1}{4}}d^2 \left( \frac{19}{8} \log 2 + \frac{5}{8} \log 3 \right) - \frac{\pi}{4} - \arg \Gamma \left( -i \frac{24^{\frac{1}{4}}}{2} d^2 \right); \end{cases} \quad (\text{B.2})$$

(B) a one-parameter family of solutions (termed *separatrix solutions*), satisfying

$$y(t) = y_0(t) - \frac{h}{4\sqrt{\pi}} 24^{-\frac{1}{8}} (-t)^{-\frac{1}{8}} \exp \left\{ -\frac{4}{5} 24^{\frac{1}{4}} (-t)^{\frac{5}{4}} \right\} \left( 1 + \mathcal{O}\left(|t|^{-\frac{5}{4}}\right) \right) \quad (\text{B.3})$$

as  $t \rightarrow -\infty$ , where  $y_0(t) = \sqrt{\frac{-t}{6}} \left[ 1 + \mathcal{O}\left((-t)^{-\frac{5}{2}}\right) \right]$  and

$$h = s_1 - s_4; \quad (\text{B.4})$$

(C) a two-parameter family of solutions, having infinitely many double poles on the negative real axis and satisfying

$$\frac{1}{y(t) + \sqrt{-t/6}} \sim \frac{\sqrt{6}}{2} \sin^2 \left\{ \frac{2}{5} 24^{1/4} (-t)^{5/4} + \frac{5}{8} \rho \log(-t) + \sigma \right\} \quad \text{as } t \rightarrow -\infty, \quad (\text{B.5})$$

where

$$\begin{cases} \rho = \frac{1}{2\pi} \log(|s_2|^2 - 1) = \frac{1}{2\pi} \log(|1 + s_2 s_3|) = \frac{1}{2\pi} \log |s_0|, \\ \sigma = \frac{19}{8} \rho \log 2 + \frac{5}{8} \rho \log 3 + \frac{1}{2} \arg \Gamma \left( \frac{1}{2} - i\rho \right) - \frac{\pi}{4} + \frac{1}{2} \arg s_2. \end{cases} \quad (\text{B.6})$$

The above classification is characterized by

$$\begin{cases} \text{Type A (oscillatory solutions)} : & 1 + s_2 s_3 > 0; \\ \text{Type B (separatrix solutions)} : & 1 + s_2 s_3 = 0; \\ \text{Type C (singular solutions)} : & 1 + s_2 s_3 < 0. \end{cases} \quad (\text{B.7})$$

## C Proof of Remark 5

Note that  $\beta_0 = \tilde{E}_0$ , and from [15, eqs. (2.2),(2.7)], we conclude that  $\beta_0$  is purely imaginary. In this section, we continue to show that  $\beta_s$  are purely imaginary for all  $s = 1, 2, \dots$ . To this end, we first prove the following lemma.

**Lemma 3.** *Assume that*

$$\tilde{f}(z) = 4(z^3 + Cz + 1) = 4(z - z_0)(z - z_1)(z - z_{-1}) \quad (\text{C.1})$$

with  $z_0 < 0$  and  $z_1 = \overline{z_{-1}}$ . Define the Stokes line  $\Gamma$  by

$$\Gamma = \left\{ z \in \mathbb{C} : \operatorname{Re} \left( \int_{z_1}^z \tilde{f}(t)^{1/2} dt \right) = 0 \right\}. \quad (\text{C.2})$$

Then

$$\operatorname{Im} \left( \frac{4\tilde{f}(z)\tilde{f}''(z) - 5\tilde{f}'(z)^2}{16\tilde{f}(z)^3} \right) = 0 \quad \text{for all } z \in \Gamma. \quad (\text{C.3})$$

*Proof.* From (C.2), we find that  $\operatorname{Re} \left( \tilde{f}(z)^{1/2} dz \right) = 0$ , which implies that  $\operatorname{Im} \left( \tilde{f}(z) dz^2 \right) = 0$  for  $z \in \Gamma$ , and that

$$\begin{aligned} \operatorname{Im} \left( \tilde{f}'(z) dz^3 \right) &= 0, \\ \operatorname{Im} \left( \tilde{f}''(z) dz^4 \right) &= 0 \end{aligned} \quad (\text{C.4})$$

on  $\Gamma$ . A combination of the above facts yields (C.3).  $\square$

**Proof of Remark 5:** Since the integrands in (2.24) and (2.25) are analytic in  $\tilde{\mathbb{D}}$  and are bounded by  $\mathcal{O}(t^{-2})$  as  $t \rightarrow \infty$ , we conclude that the limits  $\lim_{\eta \rightarrow \infty} \eta^{\frac{1}{2}} \tilde{a}_s(\eta)$ ,  $s = 1, 2, \dots$ , are independent of the integration contour (or limiting directions). For simplicity, we assume that  $\arg \eta = \pi$ . Now we show by induction that for all  $s = 1, 2, \dots$ ,  $a_s(\eta)$  are real when  $\arg \eta = \pi$ , which shall imply that the limits  $\beta_s = \lim_{\eta \rightarrow \infty} \eta^{\frac{1}{2}} \tilde{a}_s(\eta)$  are all purely imaginary.

For  $s = 1$ , it follows from (C.3) that the integral

$$\int_0^\eta \tilde{\psi}(t) t^{-\frac{1}{2}} dt \quad (\text{C.5})$$

is purely imaginary when  $\arg \eta = \pi$ , where  $\tilde{\psi}(\eta)$  is defined in (2.19). This implies that  $a_1(\eta)$  is real when  $\arg \eta = \pi$ . Moreover, we can conclude that  $a_1'(\eta)$ ,  $a_1''(\eta)$ , and  $a_1'''(\eta)$  are also real when  $\arg \eta = \pi$ .

Suppose that, when  $\arg \eta = \pi$ , all of  $a_k(\eta)$ ,  $a_k'(\eta)$ ,  $a_k''(\eta)$ , and  $a_k'''(\eta)$  are real-valued functions for  $k = 1, 2, \dots, s-1$ . Since  $\tilde{F}_s(\eta)$  is a real-coefficient polynomial of  $a_k(\eta)$ ,  $a_k'(\eta)$ ,  $a_k''(\eta)$ , and  $a_k'''(\eta)$  for  $k = 1, 2, \dots, s-1$ , we conclude that  $\tilde{F}_s(\eta)$  is also real when  $\arg \eta = \pi$ . It follows that the integral

$$\int_0^\eta \frac{\tilde{F}_s(t)}{t^{\frac{1}{2}}} dt, \quad \arg \eta = \pi \quad (\text{C.6})$$

is purely imaginary. Consequently,  $\tilde{a}_s(\eta)$  is real when  $\arg \eta = \pi$ .

## Acknowledgements

The work of Wen-Gao Long was supported in part by the Natural Science Foundation of Hunan Province [Grant no. 2024JJ5131] and the Outstanding Youth Fund of Hunan Provincial Department of Education [Grant no. 23B0454]. The work of Yu-Tian Li was supported in part by the National Natural Science Foundation of China [Grant no. 11801480].

## References

- [1] A. P. Bassom, P. A. Clarkson, C. K. Law and J. B. McLeod, Application of uniform asymptotics to the second Painlevé transcendent, *Arch. Rational Mech. Anal.*, **143** (1998), 241–271.
- [2] C. M. Bender and J. Komijani, Painlevé transcendents and PT-symmetric Hamiltonians, *J. Phys. A: Math. Theor.*, **48** (2015), 475202, 15 pp.
- [3] C. M. Bender, J. Komijani and Q. Wang, Nonlinear eigenvalue problems for generalized Painlevé equations, *J. Phys. A: Math. and Theor.*, **52** (2019), 315202.
- [4] M. Bertola and A. Tovbis, Universality for the focusing nonlinear Schrödinger equation at the gradient catastrophe point: rational breathers and poles of the tritronquée solution to Painlevé I, *Comm. Pure Appl. Math.*, **66** (2013), pp. 678–752.
- [5] P. A. Clarkson, Painlevé equations-nonlinear special functions, *J. Comput. Appl. Math.*, **153** (2003), 127–140.

- [6] P. A. Clarkson, Painlevé equations-nonlinear special functions, in *Orthogonal Polynomials and Special Functions*, Springer, Berlin and Heidelberg, (2006), 331–411.
- [7] D. Dai and W. Hu, Connection formulas for the Ablowitz–Segur solutions of the inhomogeneous Painlevé II equation. *Nonlinearity*, **30** (2017), 2982.
- [8] T. M. Dunster, Olver’s error bound methods applied to linear ordinary differential equations having a simple turning point, *Anal. Appl.*, **12** (2014), 385–402.
- [9] A. S. Fokas, A. R. Its, A. A. Kapaev and V. Y. Novokshenov, *Painlevé Transcendents: the Riemann-Hilbert Approach*, American Mathematical Soc., R.I., 2006.
- [10] N. Joshi and M. D. Kruskal, The Painlevé connection problem: an asymptotic approach. I, *Stud. Appl. Math.*, **86** (1992), 315–376.
- [11] A. A. Kapaev, Asymptotic behavior of the solutions of the Painlevé equation of the first kind, *Differ. Uravn.*, **24** (1988), 1684–1695.
- [12] A. A. Kapaev, Quasi-linear Stokes phenomenon for the Painlevé first equation, *J. Phys. A: Math. Gen.*, **37** (2004), 11149–11167.
- [13] A. A. Kapaev and A. V. Kitaev, Connection formulae for the first Painlevé transcendent in the complex domain, *Lett. Math. Phys.*, **27** (1993), 243–252.
- [14] W.-G. Long and Y.-T. Li Connection problem of the first Painlevé transcendents with large initial data, *J. Phys. A: Math. Theor.*, 56 (2023), 175201.
- [15] W.-G. Long, Y.-T. Li, and Q.-h. Wang, Connection problem of the first Painlevé transcendent between poles and negative infinity, *SIAM J. Math. Anal.*, 55 (2023), 6676–6706.
- [16] W.-G. Long, Z.-Y. Zeng, On the Connection Problem for the Second Painlevé Equation with Large Initial Data, *Constr. Approx.*, **55** (2022), 861–889.
- [17] F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds. *NIST Digital Library of Mathematical Functions*. <https://dlmf.nist.gov/>, Release 1.2.0 of 2024-03-15.
- [18] S. Y. Slavyanov and W. Lay, *Special Functions: A Unified Theory Based on Singularities*, Oxford University Press, Oxford, 2000.
- [19] R. Wong and H.-Y. Zhang, On the connection formulas of the third Painlevé transcendent, *Discrete Contin. Dyn. Syst.*, **23** (2009), 541–560.
- [20] R. Wong and H.-Y. Zhang, On the connection formulas of the fourth Painlevé transcendent, *Anal. Appl.*, **7** (2009), 419–448.
- [21] J. Xia, S.-X. Xu and Y.-Q. Zhao, Isomonodromy sets of accessory parameters for Heun class equations, *Stud. Appl. Math.*, 146 (2021), 901–952.
- [22] Z.-Y. Zeng and Y.-Q. Zhao, Application of uniform asymptotics to the connection formulas of the fifth Painlevé equation, *Appl. Anal.*, **95** (2016), 390–404.