

On reverse Markov-Nikol'skii inequalities for polynomials with restricted zeros¹

Mikhail A. Komarov²

Abstract. Let Π_n be the class of algebraic polynomials P of degree n , all of whose zeros lie on the segment $[-1, 1]$. In 1995, S. P. Zhou has proved the following Turán type reverse Markov-Nikol'skii inequality: $\|P'\|_{L_p[-1,1]} > c(\sqrt{n})^{1-1/p+1/q} \|P\|_{L_q[-1,1]}$, $P \in \Pi_n$, where $0 < p \leq q \leq \infty$, $1 - 1/p + 1/q \geq 0$ ($c > 0$ is a constant independent of P and n). We show that Zhou's estimate remains true in the case $p = \infty$, $q > 1$. Some of related Turán type inequalities are also discussed.

For any $n \in \mathbb{N}$, let Π_n be the class of polynomials P of degree n , all of whose zeros lie in the closed interval $[-1, 1]$, and $\Pi_n(D^+)$ — the class of polynomials P of degree n , all of whose zeros lie in the closed half-disk $D^+ := \{z : |z| \leq 1, \operatorname{Im} z \geq 0\}$. We will use the notation $\|P\|_p := \|P\|_{L_p[-1,1]}$, $0 < p \leq \infty$, where $\|P\|_{L_\infty[-1,1]} = \|P\|_{C[-1,1]}$.

In [6], the author proved the inequality

$$\|P'\|_\infty > A\sqrt{n} \|P\|_\infty, \quad A = \frac{2}{3\sqrt{210e}}, \quad P \in \Pi_n(D^+), \quad (1)$$

that generalizes the well-known *reverse Markov inequality*

$$\|P'\|_\infty > \frac{1}{6}\sqrt{n} \|P\|_\infty, \quad P \in \Pi_n, \quad (2)$$

obtained by P. Turán in 1939. The L_p -version of (2) was obtained for $p = 2$ — by A. K. Varma (1976) and for any $p > 0$ — by S. P. Zhou (1984, 1986), see [11], [13]:

$$\|P'\|_p > c\sqrt{n} \|P\|_p, \quad P \in \Pi_n, \quad 0 < p < \infty, \quad (3)$$

where $c > 0$ is a constant depending only on p . The following more general result (*reverse Markov-Nikol'skii inequality*) is also due to Zhou [14]: suppose that

$$0 < p \leq q \leq \infty, \quad 1 - 1/p + 1/q \geq 0;$$

then

$$\|P'\|_p > c(\sqrt{n})^{1-1/p+1/q} \|P\|_q, \quad P \in \Pi_n, \quad (4)$$

where again $c = c(p) > 0$. Taking the polynomials $Q_n(x) := (1 - x^2)^n$, we see that the order of n in (4) (and hence in (1)-(3)) can not be improved, because

$$\int_{-1}^1 |Q'_n(x)|^p dx \asymp n^{(p-1)/2}, \quad \int_{-1}^1 |Q_n(x)|^q dx \asymp n^{-1/2} \quad (n \rightarrow \infty)$$

for any fixed $p, q > 0$.

¹MSC 41A17; *key words*: Turán inequality, Markov-Nikol'skii inequality, polynomials with restricted zeros

²Vladimir State University, Gor'kogo street 87, 600000 Vladimir, Russia; e-mail: kami9@yandex.ru

For extensions of (1) and (4), see [4] and [12], correspondingly. The limiting case $p = 0$ of (3) was investigated in [7]. We also recall the following weighted Turán type inequalities with the best possible constants: if $P \in \Pi_n$, then

$$\|P'(x)\sqrt{1-x^2}\|_\infty \geq A_\infty(n) \|P\|_\infty, \quad A_\infty(n) := \frac{\sqrt{n}}{\sqrt{2}} \left(1 - \frac{1}{2n}\right)^{n-1/2} \quad (5)$$

(V. F. Babenko and S. A. Pichugov, [1]), and

$$\|P'(x)(1-x^2)^{\frac{p-1}{2p}}\|_p \geq A_p(n) \|P\|_\infty, \quad 1 \leq p < \infty, \quad (6)$$

$$A_p(n) := n \left\{ B \left(\frac{(2n-1)p+1}{2}, \frac{p+1}{2} \right) \right\}^{1/p}$$

(I. Ya. Tyrygin, [10]), where $B(x, y)$ is the Euler beta-function. It can be checked that $A_p(n) \rightarrow A_\infty(n)$ ($p \rightarrow \infty$) and

$$A_p(n) \asymp (\sqrt{n})^{1-1/p}, \quad 1 \leq p \leq \infty \quad (n \rightarrow \infty).$$

In this short note, we show the existence of the estimate of the form (4) in the case when $p = \infty$, $q > 1$ (in contrast to Zhou's case, where $p \leq q$). Namely, for the class $\Pi_n(D^+)$, the following extension of (4), as well as of (1), holds:

Theorem 1 *If $P \in \Pi_n(D^+)$, then for any $1 < q \leq \infty$*

$$\|P'\|_\infty > 6^{1/q} (1 - q^{-1})^{1/q} (A\sqrt{n})^{1+1/q} \|P\|_q, \quad (7)$$

where A is the constant defined in (1).

This bound can be slightly improved for the class Π_n as follows:

Theorem 2 *If $P \in \Pi_n$, then for any $1 < q \leq \infty$*

$$\|P'(x)\sqrt{1-x^2}\|_\infty^{1-1/q} \|P'(x)(1-x^2)\|_\infty^{1/q} \geq c_q n^{1/q} \{A_\infty(n)\}^{1-1/q} \|P\|_q, \quad (8)$$

where $A_\infty(n) > \sqrt{n}/\sqrt{2e}$ is the constant defined in (5), $c_q := 2^{-1/q}(1 - q^{-1})^{1/q}$.

Corollary 1 *If $P \in \Pi_n$, then for any $1 < q \leq \infty$*

$$\|P'(x)\sqrt{1-x^2}\|_\infty > c_q (\sqrt{2e})^{-1+1/q} (\sqrt{n})^{1+1/q} \|P\|_q.$$

Of course, in the case $q = \infty$ we have (7) and (8) directly from (1) and (5).

Next, applying (5), (6) to the derivative $P' \in \Pi_{n-1}$ and combining the result with (8), we get the reverse Markov-Nikol'skii inequality for the second derivative.

Corollary 2 *If $P \in \Pi_n$, $n \geq 2$, then for any $1 \leq p \leq \infty$ and $1 < q \leq \infty$*

$$\|P''(x)(1-x^2)^{\frac{p-1}{2p}}\|_p \geq C_{p,q}(n) \|P\|_q,$$

where $C_{p,q}(n) := c_q n^{1/q} \{A_\infty(n)\}^{1-1/q} \{A_p(n-1)\}$,

$$C_{p,q}(n) \asymp (\sqrt{n})^{2-1/p+1/q} \quad (n \rightarrow \infty).$$

For example, $C_{1,\infty} = A_\infty$, and we have $\|P''\|_1 > (\sqrt{n}/\sqrt{2e}) \|P\|_\infty$.

It should be mentioned that the inequality $\|P''\|_\infty \geq C(n) \|P\|_\infty$ with the best possible constant $C(n) = \min\{n; (n-1)n/4\}$ was constructed by Babenko and Pichugov [2].

Taking again $Q_n(x) = (1-x^2)^n$, we see that Theorems 1, 2 and Corollaries 1, 2 can not be improved in the order of n . In the case of Corollary 2 we use the fact that $Q_n''(x) = 2n(1-x^2)^{n-2}\{x^2(2n-1) - 1\}$, where $|x^2(2n-1) - 1| \leq 1$ ($|x| \leq 1/\sqrt{2n-1}$) and $0 < x^2(2n-1) - 1 < 2nx^2$ ($|x| > 1/\sqrt{2n-1}$). Hence

$$\begin{aligned} \int_{-1}^1 |Q_n''(x)|^p dx &< 2 \int_0^{1/\sqrt{2n-1}} (2n)^p dx + 2 \int_{1/\sqrt{2n-1}}^1 (2nx)^{2p} (1-x^2)^{(n-2)p} dx \\ &< 2^{1+p} n^{p-1/2} + 2^{1+2p} n^{2p} \int_0^1 x^{2p} (1-x^2)^{(n-2)p} dx \asymp n^{p-1/2}. \end{aligned}$$

Proof of Theorems 1 and 2. To prove the theorems, we need the following non-standard Nikol'skii type inequality between different metrics:

Lemma 1 *Let $1 < q < \infty$. Then*

$$\int_{-1}^1 |Q(x)|^q dx \leq K \frac{q}{q-1} \frac{\|Q\|_\infty^{q-1}}{n} \quad (9)$$

with $K := 70e$ for any polynomial Q of degree n such that

$$Q \in \Pi_n(D^+), \quad \|Q'\|_\infty = 1.$$

Moreover, (9) holds true with $K := 2$, if

$$Q \in \Pi_n, \quad \|Q'(x)(1-x^2)\|_\infty = 1. \quad (10)$$

Thus, from Lemma 1 we immediately deduce that for any polynomial $P \in \Pi_n$

$$n 2^{-1} (1-q^{-1}) \int_{-1}^1 |P(x)|^q dx \leq \|P'(x)(1-x^2)\|_\infty \|P\|_\infty^{q-1} \quad (q > 1), \quad (11)$$

and (8) follows by this and $\|P\|_\infty \leq \|P'(x)\sqrt{1-x^2}\|_\infty / A_\infty$ (see (5)). The proof of (7) is similar and uses the inequalities $\|P\|_\infty < \|P'\|_\infty / (A\sqrt{n})$, $P \in \Pi_n(D^+)$ (see (1)), and $K = 70e = 4/(27A^2) < 1/(6A^2)$.

Remark. Some ‘‘quasi’’ Turán type estimates for the class Π_n follow from (11) with (6). For example, we have $c_q n^{1/q} \{A_p(n)\}^{1-1/q} \|P\|_q \leq \|P'\|_\infty^{1/q} \|P'\|_p^{1-1/q}$ ($p \geq 1$, $q > 1$).

Proof of Lemma 1. If a given polynomial Q from $\Pi_n(D^+)$ (or Π_n) satisfies $n\|Q\|_\infty \leq 1$, then (9) obviously holds with $K = 2(q-1)/q < 2$:

$$\int_{-1}^1 |Q(x)|^q dx \leq 2\|Q\|_\infty^q \leq \frac{2}{n} \|Q\|_\infty^{q-1}.$$

Now let $Q \in \Pi_n$ be a polynomial such that (10) holds and $n\|Q\|_\infty > 1$.

For any $\delta > 0$, let us define the set

$$E(\delta) = \{x \in [-1, 1] : n|Q(x)| > \delta\}.$$

The measure $m(E(\delta)) =: h(\delta)$ is zero if $\delta \geq T := n\|Q\|_\infty$, therefore

$$\begin{aligned} \int_{-1}^1 (n|Q(x)|)^q dx &= q \int_0^\infty \delta^{q-1} h(\delta) d\delta = q \int_0^T \delta^{q-1} h(\delta) d\delta \\ &\leq 2q \int_0^1 \delta^{q-1} d\delta + q \int_1^T \delta^{q-1} h(\delta) d\delta = 2 + q \int_1^T \delta^{q-1} h(\delta) d\delta. \end{aligned}$$

By (10), we have $(1-x^2)|Q'(x)| \leq 1$ for $-1 \leq x \leq 1$. Hence

$$E(\delta) \subset \left\{x \in [-1, 1] : \frac{(1-x^2)|Q'(x)|}{n|Q(x)|} < \frac{1}{\delta}\right\}. \quad (12)$$

But for any polynomial $\pi_n \in \Pi_n$ and any $\alpha > 0$, the measure

$$m\left\{x \in [-1, 1] : (1-x^2) \left| \frac{\pi_n'(x)}{n\pi_n(x)} \right| \leq \alpha\right\} \leq 2\alpha;$$

this was shown in [8] (see also [9]) with help of P. Borwein's identity [3, p. 75]. This metric estimate, applied to (12), provides the bound

$$h(\delta) \leq 2/\delta,$$

and we easily get the desired inequality (9):

$$n^q \int_{-1}^1 |Q(x)|^q dx \leq 2 + 2q \int_1^T \delta^{q-2} d\delta = 2 + 2q \frac{T^{q-1} - 1}{q-1} < \frac{2qT^{q-1}}{q-1}, \quad T = n\|Q\|_\infty.$$

To obtain (9) in the remaining case ($Q \in \Pi_n(D^+)$, $\|Q'\|_\infty = 1$, $n\|Q\|_\infty > 1$), we use similar arguments with another metric estimate

$$m\left\{x \in [-1, 1] : \left| \frac{\pi_n'(x)}{n\pi_n(x)} \right| \leq \alpha\right\} < K\alpha, \quad K = 70e, \quad \pi_n \in \Pi_n(D^+)$$

(N. V. Govorov and Yu. P. Lapenko, [5]). Lemma is proved.

Remark. Concerning the case $q = 1$, it seems that (for example) the right version of (7) should be of the form $\|P'\|_\infty > Cn\|P\|_1$. Our approach gives the weaker bound

$$\|P'\|_\infty > (30e)^{-1} \frac{n}{9 + \ln n} \|P\|_1, \quad P \in \Pi_n(D^+).$$

Indeed (see (1) and the proof of Lemma 1), in this case the polynomial $Q(x) := P(x)/\|P'\|_\infty$ satisfies $\|Q'\|_\infty = 1$, $T = n\|Q\|_\infty < n(A\sqrt{n})^{-1} = \sqrt{n}/A$ and

$$n \int_{-1}^1 |Q(x)| dx < 2 + \int_1^{\sqrt{n}/A} \frac{70e}{\delta} d\delta = 2 + 70e \ln(\sqrt{n}/A) < 30e(9 + \ln n).$$

References

- [1] Babenko, V.F., Pichugov, S.A.: An exact inequality for the derivative of a trigonometric polynomial having only real zeros, *Mat. Zametki* **39**(3) (1986), 330–336. (in Russian) [Engl. transl.: *Math. Notes* **39**(3) (1986), 179–182, <https://doi.org/10.1007/BF01170244>]
- [2] Babenko, V.F., Pichugov, S.A.: Inequalities for the derivatives of polynomials with real zeros, *Ukr. Mat. Zh.* **38**(4) (1986), 411–416. (in Russian) [Engl. transl.: *Ukrain. Math. J.* **38**(4) (1986), 347–351, <https://doi.org/10.1007/BF01057288>]
- [3] Borwein, P.: The size of $\{x : r'_n/r_n \geq 1\}$ and lower bounds for $\|e^{-x} - r_n\|$, *J. Approx. Theory* **36**(1) (1982), 73–80. [https://doi.org/10.1016/0021-9045\(82\)90072-7](https://doi.org/10.1016/0021-9045(82)90072-7)
- [4] Erdélyi, T.: Turán-type reverse Markov inequalities for polynomials with restricted zeros, *Constr. Approx.* **54**(1) (2021), 35–48. <https://doi.org/10.1007/s00365-020-09509-y>
- [5] Govorov, N.V., Lapenko, Yu.P.: Lower bounds for the modulus of the logarithmic derivative of a polynomial, *Mat. Zametki* **23**(4) (1978), 527–535. (in Russian) [Engl. transl.: *Math. Notes* **23**(4) (1978), 288–292, <https://doi.org/10.1007/BF01786958>]
- [6] Komarov, M.A.: Reverse Markov inequality on the unit interval for polynomials whose zeros lie in the upper unit half-disk, *Anal. Math.* **45**(4) (2019), 817–821. <https://doi.org/10.1007/s10476-019-0009-y>
- [7] Komarov, M.A.: The Turán-type inequality in the space L_0 on the unit interval, *Anal. Math.* **47**(4) (2021), 843–852. <https://doi.org/10.1007/s10476-021-0097-3>
- [8] Komarov, M.A.: On Borwein’s identity and weighted Turán type inequalities on a closed interval, *Trudy Inst. Mat. Mekh.* **28**(1) (2022), 127–138. (in Russian) <https://doi.org/10.21538/0134-4889-2022-28-1-127-138>
- [9] Komarov, M.A.: On the reverse Dzyadyk inequality for polynomials with zeros on a closed interval, *Mat. Zametki* **112**(6) (2022), 941–946. (in Russian) [Engl. transl.: *Math. Notes* **112**(6) (2022), 1065–1070, <https://doi.org/10.1134/S0001434622110384>]
- [10] Tyrygin, I.Ya.: On inequalities of Turán type in some integral metrics, *Ukr. Mat. Zh.* **40**(2) (1988), 256–260. (in Russian) [Engl. transl.: *Ukrain. Math. J.* **40**(2) (1988), 223–226, <https://doi.org/10.1007/BF01056484>]
- [11] Varma, A.K.: An analogue of some inequalities of P. Turán concerning algebraic polynomials having all zeros inside $[-1, +1]$, *Proc. Amer. Math. Soc.* **55**(2) (1976), 305–309. <https://doi.org/10.1090/S0002-9939-1976-0396878-7>
- [12] Zhao, D., Zhou, S.P., Yu, D.S., Wang, J.L.: Weighted Turán type inequality for rational functions with prescribed poles, *J. Math. Inequal.* **8**(2) (2014), 251–265. <https://doi.org/10.7153/jmi-08-17>
- [13] Zhou, S.P.: An extension of the Turán inequality in L_p -space for $0 < p < 1$, *J. Math. Res. Expos.* **6**(2) (1986), 27–30. <http://dx.doi.org/10.3770/j.issn:1000-341X.1986.02.010>
- [14] Zhou, S.P.: Some remarks on Turán’s inequality. III: The completion, *Anal. Math.* **21**(4) (1995), 313–318. <https://doi.org/10.1007/BF01909153>