

**ABSOLUTELY MONOTONIC FUNCTIONS RELATED TO THE
ASYMPTOTIC FORMULA FOR THE COMPLETE ELLIPTIC
INTEGRAL OF THE FIRST KIND**

TIEHONG ZHAO AND ZHEN-HANG YANG*

ABSTRACT. Let $\mathcal{K}(x)$ be the complete elliptic integral of the first kind and

$$\mathcal{G}_p(x) = e^{\mathcal{K}(\sqrt{x})} - \frac{p}{\sqrt{1-x}}$$

for $p \in \mathbb{R}$ and $x \in (0, 1)$. In this paper we find the necessary and sufficient conditions for the functions $\pm \mathcal{G}_p^{(k)}(x)$ ($k \in \mathbb{N} \cup \{0\}$) to be absolutely monotonic on $(0, 1)$, which extend previous known results and yield several new functional inequalities involving the complete elliptic integral of the first kind. More importantly, we provide a new method to deal with those absolute monotonicity problem by proving the monotonicity of a sequence generated by the coefficients of the power series of $\mathcal{G}_p(x)$.

1. INTRODUCTION

The complete elliptic integral of the first kind $\mathcal{K}(r)$ is defined on $(0, 1)$ by

$$\mathcal{K}(r) = \int_0^{\pi/2} \frac{1}{\sqrt{1-r^2 \sin^2 t}} dt,$$

which can also be expressed by the Gaussian hypergeometric function

$$(1.1) \quad \mathcal{K}(r) = \frac{\pi}{2} F\left(\frac{1}{2}, \frac{1}{2}; 1; r^2\right) = \frac{\pi}{2} \sum_{n=0}^{\infty} \frac{(1/2)_n^2}{(n!)^2} r^{2n},$$

where

$$F(a, b; c; x) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{x^n}{n!}$$

is the Gaussian hypergeometric function defined on $x \in (-1, 1)$, here $-c \notin \mathbb{N} \cup \{0\}$ and $(a)_n = \Gamma(n+a)/\Gamma(a)$ in which $\Gamma(x) = \int_0^{\infty} t^{x-1} e^{-t} dt$ ($x > 0$) is the gamma function. The complete elliptic integral of the first kind $\mathcal{K}(r)$ has the asymptotic formula

$$(1.2) \quad \mathcal{K}(r) \sim \ln \frac{4}{r'} \quad \text{as } r \rightarrow 1^-,$$

where and in what follows $r' = \sqrt{1-r^2}$ (see [1]).

2000 *Mathematics Subject Classification.* Primary 33E05, 26A48 Secondary 40A05, 41A10.

Key words and phrases. Complete elliptic integral of the first kind, absolute monotonicity, hypergeometric function, recurrence formula, functional inequality.

Corresponding Author: Zhen-Hang Yang*.

There are many known properties of $\mathcal{K}(r)$, including monotonicity, convexity, absolute monotonicity and inequalities. Here we mention several properties related to the asymptotic formula (1.2), which are more interesting. Let

$$(1.3) \quad \mathcal{F}_p(x) = (1-x)^p e^{\mathcal{K}(\sqrt{x})} \quad \text{and} \quad \mathcal{G}_p(x) = e^{\mathcal{K}(\sqrt{x})} - \frac{p}{\sqrt{1-x}}.$$

In 1996, Qiu and Vamanamurthy [2, Theorem 1.2] showed that the function $r \mapsto \mathcal{F}_{1/2}(r^2)$ is strictly decreasing and concave from $(0, 1)$ onto $(4, e^{\pi/2})$. Yang and Chu [3, Theorem 3.1] proved that the function $r \mapsto \mathcal{F}_{p/2}(r^2)$ is strictly increasing on $(0, 1)$ if and only if $p \leq \pi/4$ and strictly decreasing on $(0, 1)$ if and only if $p \geq 1$ using a different way. Further, Yang and Tian [4] proved that $-\mathcal{F}_p'$ is absolutely monotonic on $(0, 1)$ if and only if $1/2 \leq p \leq (\pi + 4 + \sqrt{16 - \pi})/8 = 1.340\dots$, and \mathcal{F}_p is absolutely monotonic on $(0, 1)$ if and only if $p \leq \pi/8 = 0.392\dots$. Yang and Chu [3, Theorem 3.3] proved that \mathcal{G}_p is strictly decreasing on $(0, 1)$ if and only if $p \geq 4$ and strictly increasing on $(0, 1)$ if and only if $p \leq \pi e^{\pi/2}/4$.

For the function $f_p(x) = \mathcal{K}(\sqrt{x})/\ln(p/\sqrt{1-x})$ ($p \geq 1$), Anderson, Vamanamurthy and Vuorinen [5, Theorem 2.2] proved that the function $r \mapsto f_c(r^2)$ is strictly decreasing if and only if $1 \leq c \leq 4$ and increasing if and only if $c \geq e^2$. Qiu and Vamanamurthy [6] further verified that $r \mapsto f_4(r^2)$ is strictly concave on $(0, 1)$. Yand and Tian [7] showed that f_p is concave on $(0, 1)$ if and only if $p = e^{4/3}$. After three years, Tian and Yang [8, Theorem 1] proved further that $p = e^{4/3}$ is also the necessary and sufficient condition for the function $-f_p''$ to be absolutely monotonic on $(0, 1)$ by using recurrence technique. Recently, Alzer and Richards [9] found that the function $x \mapsto 1/f_p(x)$ is strictly concave on $(0, 1)$ if and only if $p \geq e^{8/5}$. This condition was proved in [8, Theorem 3] to be also necessary and sufficient such that the function $x \mapsto -[1/f_p(x)]''$ is absolutely monotonic on $(0, 1)$. Moreover, Tian and Yang [10, Theorem 1] discovered the function $x \mapsto 1/f_p(x)$ is strictly convex on $(0, 1)$ if and only if $0 < p \leq 4$. More information about monotonicity, convexity and absolute monotonicity for the functions $f_p(x)$ and its reciprocal can be found in [11, 12, 8, 10], also refer to [13, 14, 15, 16, 17, 18, 19, 20, 21] for their generalizations. Besides, several remarkable inequalities related to the asymptotic formula (1.2) can be seen in [22, 23, 24, 25, 26, 27, 28, 29].

Let us return to (1.3). The absolute monotonicity of the functions \mathcal{F}_p in [4] and the monotonicity of \mathcal{G}_p in [3] encourage us to investigate the absolute monotonicity of the function \mathcal{G}_p on $(0, 1)$. This seems to be not difficult, however, it is actually quite challenging. Fortunately, we succeed in solving the problem using the following new idea:

Step 1: Let

$$(1.4) \quad e^{\mathcal{K}(\sqrt{x})} = \sum_{n=0}^{\infty} b_n x^n \quad \text{and} \quad \frac{1}{\sqrt{1-x}} = \sum_{n=0}^{\infty} W_n x^n,$$

where

$$(1.5) \quad W_n = \frac{(2n-1)!!}{(2n)!!} = \frac{\Gamma(n+1/2)}{\Gamma(1/2)\Gamma(n+1)}$$

which satisfies the recurrence relation

$$(1.6) \quad W_{n+1} = \frac{n+1/2}{n+1} W_n.$$

Then

$$(1.7) \quad G(x) = \left[\frac{\pi}{8} F\left(\frac{1}{2}, \frac{1}{2}; 2; x\right) - \frac{1}{2} \right] e^{\mathcal{K}(\sqrt{x})} = \sum_{n=0}^{\infty} \left[(n+1)b_{n+1} - \left(n + \frac{1}{2}\right)b_n \right] x^n.$$

Differentiation yields

$$G'(x) = \frac{\pi}{64} e^{\mathcal{K}(\sqrt{x})} g(x),$$

where

$$(1.8) \quad g(x) = F\left(\frac{3}{2}, \frac{3}{2}; 3; x\right) + \pi F\left(\frac{3}{2}, \frac{3}{2}; 2; x\right) F\left(\frac{1}{2}, \frac{1}{2}; 2; x\right) - 4F\left(\frac{3}{2}, \frac{3}{2}; 2; x\right).$$

Step 2: We prove that $g(x)$ is absolutely monotonic on $(0, 1)$. Since $e^{\mathcal{K}(\sqrt{x})}$ is obviously absolutely monotonic on $(0, 1)$, so is $G'(x)$.

Step 3: Due to the absolute monotonicity of $G'(x)$, we deduce the positivity of those coefficients of power series of $G(x)$ (1.7), except for the constant term, that is,

$$(1.9) \quad (n+1)b_{n+1} - \left(n + \frac{1}{2}\right)b_n > 0 \quad \text{or equivalently,} \quad \frac{b_{n+1}}{W_{n+1}} > \frac{b_n}{W_n}$$

for $n \geq 1$. Moreover, we also need to prove that $\lim_{n \rightarrow \infty} (b_n/W_n) = 4$.

Step 4: From Step 3, we easily arrive at the absolute monotonicity results for the functions $\pm \mathcal{G}_p^{(k)}(x)$ on $(0, 1)$ for $k \geq 0$, which are contained in the following theorem.

Theorem 1. *Let \mathcal{G}_p be defined on $(0, 1)$ in (1.3). Then the following statements are valid:*

(i) \mathcal{G}_p is absolutely monotonic on $(0, 1)$ if and only if $p \leq \pi e^{\pi/2}/4 = 3.778\dots$, and $-\mathcal{G}_p$ is absolutely monotonic on $(0, 1)$ if and only if $p \geq e^{\pi/2}$.

(ii) For $k \in \mathbb{N}$, $\mathcal{G}_p^{(k)}$ is absolutely monotonic on $(0, 1)$ if and only if $p \leq b_k/W_k$, and $-\mathcal{G}_p^{(k)}$ is absolutely monotonic on $(0, 1)$ if and only if $p \geq 4$.

Remark 1. *The conclusion in Step 3 follows from the following obvious fact:*

A real power series $S(x) = \sum_{n=0}^{\infty} a_n x^n$ converging on $(-R, R)$ ($R > 0$) is absolutely monotonic on $(0, R)$ if and only if $a_k \geq 0$ for all $k \geq 0$.

In fact, if $S(x)$ is absolutely monotonic on $(0, R)$, then $S^{(k)}(0) \geq 0$ for all $k \geq 0$, which means that $a_k \geq 0$ for all $k \geq 0$. Conversely, if $a_k \geq 0$ for all $k \geq 0$, then $S^{(k)}(x) \geq 0$ for $x \in (0, R)$ and all $k \geq 0$.

2. PRELIMINARIES AND PROOF OF THEOREM 1

2.1. Basic knowledge. In order to prove our results, we need several basic properties of the hypergeometric function for real a, b, c , where $-c \notin \mathbb{N} \cup \{0\}$.

(i) Differentiation formula

$$(2.1) \quad F'(a, b; c; x) = \frac{ab}{c} F(a+1, b+1; c+1; x).$$

(ii) The value of $x = 1$:

$$F(a, b; c; 1) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)},$$

where $-c \notin \mathbb{N} \cup \{0\}$, and $c > a + b$. In particular, we have

$$(2.2) \quad F\left(\frac{1}{2}, \frac{1}{2}; 2; 1\right) = \frac{4}{\pi}.$$

(iii) Linear transformation formula

$$(2.3) \quad F(a, b; c; x) = (1-x)^{c-a-b} F(c-a, c-b; c; x).$$

(iv) Asymptotic formula

$$(2.4) \quad F(a, b; a+b; x) = \frac{R(a, b) - \ln(1-x)}{B(a, b)} + O((1-x) \ln(1-x)) \text{ as } x \rightarrow 1^-,$$

where

$$B(a, b) = \frac{\Gamma(a) \Gamma(b)}{\Gamma(a+b)}, \quad a, b > 0$$

is the classical beta function,

$$(2.5) \quad R \equiv R(a, b) = -2\gamma - \psi(a) - \psi(b),$$

here $\psi(x) = \Gamma'(x)/\Gamma(x)$, $x > 0$ is the psi function and γ is the Euler-Mascheroni constant. In particular, we have

$$(2.6) \quad F\left(\frac{3}{2}, \frac{3}{2}; 3; x\right) \sim \frac{8}{\pi} [4 \ln 2 - 4 - \ln(1-x)] \rightarrow \infty \text{ as } x \rightarrow 1^-.$$

Moreover, using the notation W_n , certain specific hypergeometric functions can be simply represented as

$$(2.7) \quad \begin{aligned} F\left(\frac{1}{2}, \frac{1}{2}; 1; x\right) &= \sum_{n=0}^{\infty} W_n^2 x^n, & F\left(\frac{3}{2}, \frac{3}{2}; 2; x\right) &= 4 \sum_{n=0}^{\infty} (n+1) W_{n+1}^2 x^n, \\ F\left(\frac{1}{2}, \frac{1}{2}; 2; x\right) &= \sum_{n=0}^{\infty} \frac{W_n^2}{n+1} x^n, & F\left(\frac{3}{2}, \frac{3}{2}; 3; x\right) &= 8 \sum_{n=0}^{\infty} \frac{n+1}{n+2} W_{n+1}^2 x^n, \end{aligned}$$

2.2. The first three steps. In this subsection, we complete the proofs of Step 1-3 introduced in Section 1.

Lemma 1 (Step 1). *Let b_n be defined as in (1.4) Then the formula (1.7) is valid. Moreover, b_n satisfies the recurrence formula: $b_0 = e^{\pi/2}$ and for $n \geq 0$,*

$$(2.8) \quad b_{n+1} = \frac{n}{n+1} b_n + \frac{\pi}{8(n+1)} \sum_{k=0}^n \frac{W_k^2}{k+1} b_{n-k}.$$

In particular, the first few coefficients are:

$$b_1 = \frac{1}{8} \pi e^{\pi/2}, \quad b_2 = \frac{1}{128} \pi (\pi + 9) e^{\pi/2}, \quad \text{and} \quad b_3 = \frac{\pi}{3072} (\pi^2 + 27\pi + 150) e^{\pi/2}.$$

Proof. Differentiation yields

$$\left[e^{\mathcal{K}(\sqrt{x})} \right]' = \frac{\pi}{8} F\left(\frac{3}{2}, \frac{3}{2}; 2; x\right) e^{\mathcal{K}(\sqrt{x})} = \sum_{n=0}^{\infty} n b_n x^{n-1},$$

which, by (2.3), implies that

$$(2.9) \quad \frac{\pi}{8} F\left(\frac{1}{2}, \frac{1}{2}; 2; x\right) e^{\mathcal{K}(\sqrt{x})} = (1-x) \sum_{n=0}^{\infty} n b_n x^{n-1} = \sum_{n=0}^{\infty} [(n+1) b_{n+1} - n b_n] x^n.$$

Hence,

$$G(x) = \frac{\pi}{8} F\left(\frac{1}{2}, \frac{1}{2}; 2; x\right) e^{\mathcal{K}(\sqrt{x})} - \frac{1}{2} e^{\mathcal{K}(\sqrt{x})} = \sum_{n=0}^{\infty} \left[(n+1) b_{n+1} - \left(n + \frac{1}{2}\right) b_n \right] x^n,$$

which proves the formula (1.7).

On the other hand, applying the third formula of (2.7) and Cauchy product formula to (2.9) gives

$$\frac{\pi}{8} \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \frac{W_k^2}{k+1} b_{n-k} \right) x^n = \sum_{n=0}^{\infty} [(n+1) b_{n+1} - n b_n] x^n,$$

then comparing coefficients of x^n leads to

$$\frac{\pi}{8} \sum_{k=0}^n \frac{W_k^2}{k+1} b_{n-k} = (n+1) b_{n+1} - n b_n,$$

which implies (2.8). Letting $x \rightarrow 0^+$ leads us to $b_0 = e^{\pi/2}$, and the first few coefficients follows from the recurrence formula (2.8), which completes the proof. \square

Lemma 2 (Step 2). *Let $G(x)$ be defined by (1.7). Then $G'(x)$ is absolutely monotonic on $(0, 1)$.*

Proof. Differentiation yields

$$G'(x) = \frac{\pi}{64} e^{\mathcal{K}(\sqrt{x})} g(x),$$

where $g(x)$ is defined by (1.8). Since $e^{\mathcal{K}(\sqrt{x})}$ is obviously absolutely monotonic on $(0, 1)$, if we prove that $g(x)$ is absolutely monotonic on $(0, 1)$, then by Remark 1, so is $G'(x)$.

Let

$$g_0(x) = (1-x) F\left(\frac{3}{2}, \frac{3}{2}; 3; x\right) + \pi F^2\left(\frac{1}{2}, \frac{1}{2}; 2; x\right) - 4F\left(\frac{1}{2}, \frac{1}{2}; 2; x\right).$$

By the third and fourth formulas of (2.7), $g_0(x)$ can be expressed as

$$g_0(x) = \sum_{n=0}^{\infty} u_n x^n,$$

where

$$(2.10) \quad u_n = \pi \sum_{k=0}^n \frac{W_k^2 W_{n-k}^2}{(k+1)(n-k+1)} - 6 \frac{2n+1}{(n+2)(n+1)} W_n^2.$$

It was proved in [3, Lemma 2.5] that $u_n < 0$ for all $n \geq 8$. This together with [3, Eqs. (3.6)–(3.8)] gives that $u_0 = \pi - 3 > 0$, $u_1 = (\pi - 3)/4 > 0$ and $u_n < 0$ for all $n \geq 2$. Moreover, by (2.2) and (2.6), we have

$$g_0(1^-) = \sum_{n=0}^{\infty} u_n = 0.$$

On the other hand, applying the linear transformation formula (2.3) and the Cauchy product formula gives

$$\begin{aligned} g(x) &= F\left(\frac{3}{2}, \frac{3}{2}; 3; x\right) + \pi F\left(\frac{3}{2}, \frac{3}{2}; 2; x\right) F\left(\frac{1}{2}, \frac{1}{2}; 2; x\right) - 4F\left(\frac{3}{2}, \frac{3}{2}; 2; x\right) \\ &= \frac{g_0(x)}{1-x} = \frac{1}{1-x} \sum_{n=0}^{\infty} u_n x^n = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n u_k \right) x^n := \sum_{n=0}^{\infty} v_n x^n. \end{aligned}$$

It is verified that $v_0 = u_0 = \pi - 3 > 0$, $v_1 = u_0 + u_1 = 5(\pi - 3)/4 > 0$. For $n \geq 2$, since

$$v_n - v_{n-1} = \sum_{k=0}^n u_k - \sum_{k=0}^{n-1} u_k = u_n < 0,$$

it is readily deduced that

$$v_n > \lim_{n \rightarrow \infty} v_n = \sum_{n=0}^{\infty} u_n = 0.$$

This proves the absolute monotonicity of $g(x)$ on $(0, 1)$, thereby completing the proof. \square

Lemma 3 (Step 3). *Let b_n be defined by (2.8). Then the sequence $\{b_n/W_n\}_{n \geq 1}$ is increasing with*

$$(2.11) \quad \lim_{n \rightarrow \infty} \frac{b_n}{W_n} = 4.$$

Proof. To prove the increasing property of the sequence $\{b_n/W_n\}_{n \geq 1}$, it suffices to prove the inequality

$$b_{n+1} > \frac{W_{n+1}}{W_n} b_n = \frac{n+1/2}{n+1} b_n,$$

or equivalently,

$$(2.12) \quad (n+1)b_{n+1} - \left(n + \frac{1}{2}\right)b_n > 0$$

for $n \in \mathbb{N}$. By Lemma 2, $G'(x)$ is absolutely monotonic on $(0, 1)$, which together with (1.7) and Remark 1 implies (2.12).

Since the sequence $\{b_n/W_n\}_{n \geq 1}$ is increasing and $b_1/W_1 > 0$, the $\lim_{n \rightarrow \infty} (b_n/W_n) = p_0 > 0$ or tends to ∞ . We first prove that $\lim_{n \rightarrow \infty} (b_n/W_n) \leq 4$. If not, that is, $\lim_{n \rightarrow \infty} (b_n/W_n) > 4$, then there exists $n_0 > 1$ such that $b_n/W_n > 4$ for $n \geq n_0$. Let us denote by $p^* = b_{n_0}/W_{n_0}$. The increasing property of $\{b_n/W_n\}_{n \geq 1}$ makes us to deduce

$$\frac{b_n}{W_n} - p^* = \frac{b_n}{W_n} - \frac{b_{n_0}}{W_{n_0}} \geq 0, \text{ that is, } b_n - p^* W_n \geq 0 \text{ for } n \geq n_0,$$

which in conjunction with (1.4) yields

$$\mathcal{G}_{p^*}(x) = \sum_{n=0}^{\infty} (b_n - p^* W_n) x^n \geq \sum_{n=0}^{n_0-1} (b_n - p^* W_n) x^n,$$

which contradicts with

$$\mathcal{G}_{p^*}(1^-) = \lim_{x \rightarrow 1^-} \left[e^{\ln(4/\sqrt{1-x})} - \frac{p^*}{\sqrt{1-x}} \right] = -\infty.$$

If $\lim_{n \rightarrow \infty} (b_n/W_n) = p_0 < 4$, that is to say, $\sup_{n \geq 1} \{b_n/W_n\} = p_0$, then $b_n/W_n \leq p_0$ for all $n \geq 1$. Similarly,

$$\mathcal{G}_{p_0}(x) = \sum_{n=0}^{\infty} (b_n - p_0 W_n) x^n \leq b_0 - p_0 W_0$$

which is a contradiction with $\mathcal{G}_{p_0}(1^-) = \infty$ due to $p_0 < 4$. This proves $\lim_{n \rightarrow \infty} (b_n/W_n) = 4$, and the proof is done. \square

2.3. Step 4: Proof of Theorem 1. By means of Lemma (Step 3) 3, we can easily complete the proof of our main result.

Proof of Theorem 1. From (1.4), $\mathcal{G}_p(x)$ can be expressed in power series:

$$(2.13) \quad \mathcal{G}_p(x) = \sum_{n=0}^{\infty} b_n x^n - p \sum_{n=0}^{\infty} W_n x^n = \sum_{n=0}^{\infty} (b_n - p W_n) x^n := \sum_{n=0}^{\infty} c_n(p) x^n.$$

As a matter to remember, Lemma 3 states that the sequence $\{b_n/W_n\}$ is strictly increasing for $n \geq 1$ with $\lim_{n \rightarrow \infty} (b_n/W_n) = 4$.

(i) By (2.13), \mathcal{G}_p is absolutely monotonic on $(0, 1)$ if and only if

$$c_n(p) = b_n - p W_n \geq 0 \quad \text{for } n \geq 0,$$

which implies that

$$p \leq \inf_{n \geq 0} \left\{ \frac{b_n}{W_n} \right\} = \min \left\{ \frac{b_0}{W_0}, \frac{b_1}{W_1} \right\} = \min \left\{ e^{\pi/2}, \frac{\pi}{4} e^{\pi/2} \right\} = \frac{\pi}{4} e^{\pi/2}.$$

Similarly, $-\mathcal{G}_p$ is absolutely monotonic on $(0, 1)$ if and only if

$$p \geq \sup_{n \geq 0} \left\{ \frac{b_n}{W_n} \right\} = \max \left\{ \frac{b_0}{W_0}, 4 \right\} = \max \left\{ e^{\pi/2}, 4 \right\} = e^{\pi/2}.$$

(ii) In a fashion to (i), $\mathcal{G}_p^{(k)}$ for $k \geq 1$ is absolutely monotonic on $(0, 1)$ if and only if

$$p \leq \inf_{n \geq k} \left\{ \frac{b_n}{W_n} \right\} = \frac{b_k}{W_k};$$

and $-\mathcal{G}_p^{(k)}$ is absolutely monotonic on $(0, 1)$ if and only if

$$p \geq \sup_{n \geq k} \left\{ \frac{b_n}{W_n} \right\} = 4.$$

This completes the proof. \square

3. NEW FUNCTIONAL INEQUALITIES RELATED TO $\mathcal{K}(r)$

In this Section, Theorem 1 will be applied to establish several functional inequalities involving the complete elliptic integral of the first kind $\mathcal{K}(r)$.

Proposition 1. *Let b_n be defined in (1.4) with $b_0 = e^{\pi/2}$ and $c_n(p) = b_n - p W_n$ for $n \geq 0$. The double inequality*

$$(3.1) \quad \ln \left(\frac{p}{r'} + \sum_{n=0}^m c_n(p) r^{2n} \right) < \mathcal{K}(r) < \ln \left(\frac{q}{r'} + \sum_{n=0}^m c_n(q) r^{2n} \right)$$

holds for $r \in (0, 1)$ if and only if $p \leq b_{m+1}/W_{m+1}$ and $q \geq 4$.

Proof. As shown in (2.13), we denote by

$$(3.2) \quad \mathcal{R}_{p,m}(x) := e^{\mathcal{K}(\sqrt{x})} - \frac{p}{\sqrt{1-x}} - \sum_{n=0}^m c_n(p)x^n = \sum_{n=m+1}^{\infty} c_n(p)x^n.$$

Theorem 1 (ii) enables us to know that $c_n(p) \geq 0$ for $n \geq m+1$ if $p \leq b_{m+1}/W_{m+1}$ and $c_n(q) \leq 0$ for $n \geq m+1$ if $q \geq 4$. This together with (3.2) gives the desired inequality (3.1).

Conversely, the necessary conditions that (3.1) holds for all $r \in (0, 1)$ require to satisfy $\mathcal{R}_{p,m}(0^+) \geq 0$ and $\mathcal{R}_{q,m}(1^-) \leq 0$, which imply $c_{m+1}(p) = b_{m+1} - pW_{m+1} \geq 0$, (i.e. $p \leq b_{m+1}/W_{m+1}$) and

$$\begin{aligned} \mathcal{R}_{q,m}(1^-) &= \lim_{x \rightarrow 1^-} \frac{\sqrt{1-x}e^{\mathcal{K}(\sqrt{x})} - q - \sqrt{1-x} \sum_{n=0}^m c_n(p)x^n}{\sqrt{1-x}} \\ &= \lim_{x \rightarrow 1^-} \frac{4-q}{\sqrt{1-x}} \leq 0, \end{aligned}$$

namely, $q \geq 4$. □

Remark 2. Taking $m = 0$ in the double inequality (3.1) leads to

$$(3.3) \quad \ln \left(e^{\pi/2} - p + \frac{p}{r'} \right) < \mathcal{K}(r) < \ln \left(e^{\pi/2} - q + \frac{q}{r'} \right)$$

for all $r \in (0, 1)$ with the best constants $p = b_1/W_1 = \pi e^{\pi/2}/4$ and $q = 4$. This result was proven in [3, Corollary 3.7]. Clearly, Proposition 1 is a generalization of [3, Corollary 3.7].

Remark 3. Taking $m = 1$ in (3.1) yields

$$(3.4) \quad \ln \left[e^{\pi/2} - p - \left(\frac{p}{2} - \frac{\pi}{8} e^{\pi/2} \right) r^2 + \frac{p}{r'} \right] < \mathcal{K}(r) < \ln \left[e^{\pi/2} - q - \left(\frac{q}{2} - \frac{\pi}{8} e^{\pi/2} \right) r^2 + \frac{q}{r'} \right]$$

for all $r \in (0, 1)$ with the best constants $p = b_2/W_2 = [\pi(\pi+9)e^{\pi/2}]/48$ and $q = 4$. Since

$$\begin{aligned} & \left[e^{\pi/2} - \frac{b_2}{W_2} - \left(\frac{b_2}{2W_2} - \frac{\pi}{8} e^{\pi/2} \right) r^2 + \frac{b_2}{W_2 r'} \right] - \left(e^{\pi/2} - \frac{b_1}{W_1} + \frac{b_1}{W_1 r'} \right) \\ &= \frac{\pi(\pi-3)e^{\pi/2}}{96} \frac{r^4(r^2+3)}{r'[2+(r^2+2)r']} > 0 \end{aligned}$$

and $2 - \pi e^{\pi/2}/8 = 0.110 \dots > 0$, we conclude that the optimal inequality of (3.4) improves the one of (3.3).

For $n \geq 0$, Theorem 1 (ii) demonstrates that the function $x \mapsto \mathcal{R}_{4,n}(x)/x^{n+1}$ is strictly decreasing on $(0, 1)$ together with

$$\lim_{x \rightarrow 0^+} \frac{\mathcal{R}_{4,n}(x)}{x^{n+1}} = c_{n+1}(4) \quad \text{and} \quad \lim_{x \rightarrow 1^-} \frac{\mathcal{R}_{4,n}(x)}{x^{n+1}} = - \sum_{k=0}^n c_k(4).$$

This yields

$$- \left(\sum_{n=0}^m c_n(4) \right) x^{m+1} < e^{\mathcal{K}(\sqrt{x})} - \frac{4}{\sqrt{1-x}} - \sum_{n=0}^m c_n(4)x^n < c_{m+1}(4)x^{m+1}$$

for $x \in (0, 1)$. By replacing $x = r^2$, we obtain therefore the following proposition.

Proposition 2. Let b_n be defined by (1.4) and $c_n = b_n - 4W_n$ for $n \geq 0$. The double inequality

$$(3.5) \quad \ln \left[\frac{4}{r'} + \sum_{n=0}^m c_n r^{2n} - \left(\sum_{n=0}^m c_n \right) r^{2m+2} \right] < \mathcal{K}(r) < \ln \left[\frac{4}{r'} + \sum_{n=0}^{m+1} c_n r^{2n} \right]$$

holds for $r \in (0, 1)$, where the lower and upper bounds are sharp.

Remark 4. Taking $m = 0$ in Proposition 2 yields

$$(3.6) \quad \ln \left[\frac{4}{r'} + \left(e^{\pi/2} - 4 \right) r'^2 \right] < \mathcal{K}(r) < \ln \left[\frac{4}{r'} + \left(e^{\pi/2} - 4 \right) - \left(2 - \frac{\pi}{8} e^{\pi/2} \right) r^2 \right]$$

for $r \in (0, 1)$. It was proved in [24, Corollary 3.5] that

$$(3.7) \quad \ln \left[\frac{4}{r'} + \left(e^{\pi/2} - 4 \right) r' \right] < \mathcal{K}(r) < \ln \left(\frac{4}{r'} + e^{\pi/2} - 4 \right)$$

for $r \in (0, 1)$. In [3, Eq. (3.5)], it was found that

$$(3.8) \quad \ln \left[\frac{4}{r'} + \left(e^{\pi/2} - 4 \right) r' \right] < \mathcal{K}(r) < \ln \left[\frac{(\pi + 4)e^{\pi/2}}{8r'} + \left(\frac{1}{2} - \frac{\pi}{8} \right) e^{\pi/2} r' \right]$$

for $r \in (0, 1)$. The left side inequality of (3.6) looks like the one of (3.7) but a bit worse than the left side of (3.6), while the right side of (3.6) is better than the ones of (3.7) and of (3.8). This can be confirmed by

$$\left[\frac{4}{r'} + \left(e^{\pi/2} - 4 \right) - \left(2 - \frac{\pi}{8} e^{\pi/2} \right) r^2 \right] - \left[\frac{4}{r'} + \left(e^{\pi/2} - 4 \right) \right] < 0$$

and

$$\begin{aligned} & \left[\frac{4}{r'} + e^{\pi/2} - 4 - \left(2 - \frac{\pi}{8} e^{\pi/2} \right) r^2 \right] - \left[\frac{(\pi + 4)e^{\pi/2}}{8r'} + \left(\frac{1}{2} - \frac{\pi}{8} \right) e^{\pi/2} r' \right] \\ &= -\frac{(1-r')^2}{8r'} \left[\left((\pi + 4) e^{\pi/2} - 32 \right) - \left(16 - \pi e^{\pi/2} \right) r' \right] \\ &< -\frac{(\pi + 2) e^{\pi/2} - 24 (1-r')^2}{4} \frac{1}{r'} < 0 \end{aligned}$$

for $r \in (0, 1)$.

It was shown in [30] that every convex function $f : [0, \infty) \mapsto \mathbb{R}$ satisfies a functional inequality

$$(3.9) \quad f(x) + f(y) \leq f(0) + f(x + y)$$

for $x, y \in [0, \infty)$. In particular, $f(x)$ is superadditive if $f(0) \leq 0$. As a matter to remind, a function $f(x)$ defined on $\mathbb{R}_+ \setminus (0, a]$ is said to be superadditive if

$$(3.10) \quad f(x) + f(y) \leq f(x + y)$$

for $x, y \in (a, \infty)$, while it is called subadditive if the inequality (3.10) reverses.

Let $\mathcal{R}_{p,m}$ be defined as in (3.2). Then it is clear that $\mathcal{R}_{p,m}(0^+) = 0$ for $m \geq 0$. Since

$$\mathcal{R}_{p,m}''(x) = \left[\sum_{n=m+1}^{\infty} c_n(p) x^n \right]'' = \begin{cases} \sum_{n=2}^{\infty} n(n-1) c_n(p) x^{n-2} & \text{if } m = 0, \\ \sum_{n=m+1}^{\infty} n(n-1) c_n(p) x^{n-2} & \text{if } m \geq 1, \end{cases}$$

it follows from Theorem 1 (ii) that $c_n(p) \geq 0$ for $n \geq 2$ if $p \leq b_2/W_2 = [\pi(\pi + 9)e^{\pi/2}]/48$, while $c_n(p) \leq 0$ for $n \geq 2$ if $p \geq 4$. This together with (3.9) deduces that the double inequality

$$2\mathcal{R}_{p,m}\left(\frac{x+y}{2}\right) < \mathcal{R}_{p,m}(x) + \mathcal{R}_{p,m}(y) < \mathcal{R}_{p,m}(x+y)$$

holds for $x, y, x+y \in (0, 1)$ if $p \leq [\pi(\pi + 9)e^{\pi/2}]/48$ and it reverses if $p \geq 4$.

Proposition 3. *Let b_n be defined in (1.4) and $c_n(p) = b_n - pW_n$. If $p \leq [\pi(\pi + 9)e^{\pi/2}]/48$, then the double inequality*

$$\begin{aligned} & 2e^{\mathcal{K}(\sqrt{\frac{x+y}{2}})} + p \left(\frac{1}{\sqrt{1-x}} + \frac{1}{\sqrt{1-y}} - \frac{2}{\sqrt{1-\frac{x+y}{2}}} \right) + \sum_{n=0}^m c_n(p) \left[x^n + y^n - 2 \left(\frac{x+y}{2} \right)^n \right] \\ & < e^{\mathcal{K}(\sqrt{x})} + e^{\mathcal{K}(\sqrt{y})} \\ & < e^{\mathcal{K}(\sqrt{x+y})} + p \left(\frac{1}{\sqrt{1-x}} + \frac{1}{\sqrt{1-y}} - \frac{1}{\sqrt{1-x-y}} \right) + \sum_{n=0}^m c_n(p) [x^n + y^n - (x+y)^n] \end{aligned}$$

holds for $x, y, x+y \in (0, 1)$, and it reverses if $p \geq 4$.

Taking $m = 0$ into Proposition 3, we obtain

Corollary 1. *If $p \leq [\pi(\pi + 9)e^{\pi/2}]/48$, then the double inequality*

$$\begin{aligned} & 2e^{\mathcal{K}(\sqrt{\frac{x+y}{2}})} + p \left(\frac{1}{\sqrt{1-x}} + \frac{1}{\sqrt{1-y}} - \frac{2}{\sqrt{1-\frac{x+y}{2}}} \right) < e^{\mathcal{K}(\sqrt{x})} + e^{\mathcal{K}(\sqrt{y})} \\ & < e^{\mathcal{K}(\sqrt{x+y})} + p \left(\frac{1}{\sqrt{1-x}} + \frac{1}{\sqrt{1-y}} - \frac{1}{\sqrt{1-x-y}} - 1 \right) + e^{\pi/2} \end{aligned}$$

holds for $x, y, x+y \in (0, 1)$, and it is reversed if $p \geq 4$.

Remark 5. *Let $p = 4$, $y = 1 - r^2$ and $x \rightarrow r^2$ in Corollary 1. Then we obtain*

$$e^{\pi/2} - 4 < e^{\mathcal{K}(r)} + e^{\mathcal{K}(r')} - 4 \left(\frac{1}{r} + \frac{1}{r'} \right) \leq 2e^{\mathcal{K}(1/\sqrt{2})} - 8\sqrt{2}$$

for $r \in (0, 1)$, where

$$\mathcal{K}\left(\frac{1}{\sqrt{2}}\right) = \frac{\pi}{2} F\left(\frac{1}{2}, \frac{1}{2}; 1; \frac{1}{2}\right) = \frac{\Gamma(1/4)^2}{4\sqrt{\pi}},$$

which can be computed by the formula (see [31, Eq. (15.1.26)])

$$(3.11) \quad F\left(a, 1-a; b; \frac{1}{2}\right) = \frac{2^{1-b}\sqrt{\pi}\Gamma(b)}{\Gamma(a/2 + b/2)\Gamma(1/2 + b/2 - a/2)}.$$

Clearly, the lower and upper bounds are sharp by (1.2).

The following lemma was proved in [8, Lemma 4], where h should be infinitely differentiable instead of “thrice differentiable”.

Lemma 4. *Let the function h be defined on \mathbb{R} which is infinitely differentiable. If $h^{(2k+1)}(x) > (<) 0$ for $x \in \mathbb{R}$ and $k \in \mathbb{N}$ then the function*

$$\phi(x) = \frac{h(2c-x) - h(x)}{2c-2x} \text{ if } x \neq c \text{ and } \phi(c) = h'(c)$$

satisfies that $\phi^{(2k-2)}(x)$ is strictly convex (concave) on \mathbb{R} , and is decreasing (increasing) on $(-\infty, c)$ and increasing (decreasing) on (c, ∞) . Furthermore, if $h'(x)$ is absolutely monotonic on \mathbb{R} then $\phi(x)$ is completely monotonic on $(-\infty, c)$ and absolutely monotonic on (c, ∞) .

Applying Lemma 4 and the absolute monotonicity of $-\mathcal{G}'_4(x)$, we have the following statement.

Proposition 4. *The function*

$$\mathcal{H}(x) = \frac{\mathcal{G}_4(x) - \mathcal{G}_4(1-x)}{1-2x} = \frac{1}{1-2x} \left[e^{\mathcal{K}(\sqrt{x})} - e^{\mathcal{K}(\sqrt{1-x})} + \frac{4}{\sqrt{x}} - \frac{4}{\sqrt{1-x}} \right]$$

is completely monotonic on $(0, 1/2)$ and absolutely monotonic on $(1/2, 1)$. Consequently, the double inequality

$$(3.12) \quad \beta(1-2r^2) < e^{\mathcal{K}(r)} - e^{\mathcal{K}(r')} + \frac{4}{r} - \frac{4}{r'} < \alpha(1-2r^2)$$

holds for $r \in (0, 1/\sqrt{2})$ with the best constants $\alpha = e^{\pi/2} - 4 \approx 0.810$ and

$$\beta = 4\sqrt{2} - \frac{1}{\sqrt{\pi}}\Gamma^2\left(\frac{3}{4}\right) \exp\left(\frac{\Gamma(1/4)^2}{4\sqrt{\pi}}\right) \approx 0.246.$$

The double inequality (3.12) reverses for $r \in (1/\sqrt{2}, 1)$.

Proof. Taking $h(x) = -\mathcal{G}_4(x)$ and $c = 1/2$ in Lemma 4 yields the complete and absolute monotonicity of $\mathcal{H}(x)$ on $(0, 1/2)$ and $(1/2, 1)$, respectively. The decreasing property of $\mathcal{H}(x)$ on $(0, 1/2)$ implies that

$$(3.13) \quad -\mathcal{G}'_4\left(\frac{1}{2}\right) = \mathcal{H}\left(\frac{1}{2}\right) < \mathcal{H}(x) = \mathcal{H}(0) = e^{\pi/2} - 4$$

for $x \in (0, 1/2)$. Using the formula (3.11) we have

$$\begin{aligned} F\left(\frac{1}{2}, \frac{1}{2}; 1; \frac{1}{2}\right) &= \frac{1}{2\pi^{3/2}}\Gamma^2\left(\frac{1}{4}\right), \\ F\left(\frac{1}{2}, \frac{1}{2}; 2; \frac{1}{2}\right) &= \frac{4}{\pi^{3/2}}\Gamma^2\left(\frac{3}{4}\right). \end{aligned}$$

Then

$$\begin{aligned} -\mathcal{G}'_4\left(\frac{1}{2}\right) &= \left[\frac{2}{(1-x)^{3/2}} - \frac{\pi}{8(1-x)} F\left(\frac{1}{2}, \frac{1}{2}; 2; x\right) e^{\mathcal{K}(\sqrt{x})} \right]_{x=1/2} \\ &= 4\sqrt{2} - \frac{1}{\sqrt{\pi}}\Gamma^2\left(\frac{3}{4}\right) \exp\left(\frac{\Gamma^2(1/4)}{4\sqrt{\pi}}\right). \end{aligned}$$

Let $x = r^2$. Then (3.13) implies (3.12). In a similar way, we can prove that (3.12) reverses for $r \in (1/\sqrt{2}, 1)$. \square

4. CONCLUDING REMARKS

In this paper, we investigated the absolute monotonicity of the functions $\pm\mathcal{G}_p(x)$ and $\pm\mathcal{G}_p^{(k)}(x)$ for $k \in \mathbb{N}$ on $(0, 1)$, where

$$\mathcal{G}_p(x) = e^{\mathcal{K}(\sqrt{x})} - \frac{p}{\sqrt{1-x}} = \sum_{n=0}^{\infty} b_n x^n - p \sum_{n=0}^{\infty} W_n x^n.$$

These absolute monotonicity results yield several new functional inequality related to $\mathcal{K}(r)$. It is worth empathizing our approach used in this paper is very novel and interesting. Follow the usual method, ones should determine the signs of those coefficients of power series of $\pm\mathcal{G}_p(x)$ and $\pm\mathcal{G}_p^{(k)}(x)$ by directly using the recurrence formula (1.9) and W_n defined by (1.5), which is very complicated and hard. By our new idea (Steps 1 and 2), we found a surprising property about b_n and W_n , that is, the sequence $\{b_n/W_n\}_{n \geq 1}$ is increasing with $\lim_{n \rightarrow \infty} (b_n/W_n) = 4$. It is with this property that we can conveniently prove the desired absolute monotonicity results. Such new idea may contribute to deal with certain problems on monotonicity, convexity and absolute monotonicity occurred in special functions.

Finally, we present two remarks and a problem.

Remark 6. *It was proved in [4] that*

$$\mathcal{F}_{1/2}(x) = \sqrt{1-x}e^{\mathcal{K}(\sqrt{x})} = \sum_{n=0}^{\infty} a_n x^n,$$

where $a_0 = e^{\pi/2}$ and $a_n < 0$ for all $n \geq 1$. Differentiation yields

$$\sqrt{1-x}\mathcal{F}'_{1/2}(x) = \left[\frac{\pi}{8}F\left(\frac{1}{2}, \frac{1}{2}; 2; x\right) - \frac{1}{2} \right] e^{\mathcal{K}(\sqrt{x})} = G(x).$$

Can prove that $G'(x)$ is absolutely monotonic on $(0, 1)$ by the absolute monotonicity of $-\mathcal{F}'_{1/2}(x)$? Or vice versa?

Remark 7. *Since $b_0/W_0 > b_1/W_1$, we see that the sequence $\{b_n/W_n\}_{n \geq 0}$ is decreasing for $0 \leq n \leq 1$ and increasing for $n \geq 1$. Making use of the unimodal type monotonicity rule of the ratio of power series given in [32, Theorem 2.1], we can only deduce that*

$$\mathcal{F}_{1/2}(x) = \frac{e^{\mathcal{K}(\sqrt{x})}}{1/\sqrt{1-x}} = \frac{\sum_{n=0}^{\infty} b_n x^n}{\sum_{n=0}^{\infty} W_n x^n}$$

is decreasing on $(0, 1)$. Employing the recurrence technique, however, Yang and Tian have shown that $-\mathcal{F}'_{1/2}(x)$ is absolutely monotonic on $(0, 1)$. In a similar way, by the monotonicity rule of the ratio of power series given in [33], we can draw a conclusion that

$$x \mapsto \mathcal{J}(x) = \frac{e^{\mathcal{K}(\sqrt{x})} - e^{\pi/2}}{1/\sqrt{1-x} - 1} = \frac{\sum_{n=1}^{\infty} b_n x^n}{\sum_{n=1}^{\infty} W_n x^n}$$

is increasing on $(0, 1)$. However, by the increasing property of $\{b_n/W_n\}_{n \geq 1}$ and recurrence technique, we can prove that this function $\mathcal{J}(x)$ is absolutely monotonic on $(0, 1)$, and the proof of which is left to the reader.

Remark 8. *So far, several interesting questions remain unresolved involving the absolutely monotonic functions related to the asymptotic formula (1.2), for example,*

(i) *In [8, Conjecture 1], it was conjectured that the function $-f_4'''(x)$ is absolutely monotonic on $(0, 1)$, where $f_p(x)$ is defined in Introduction of this paper.*

(ii) *In [8, Conjecture 1], it was conjectured that the function $1/f_4(x)$ is absolutely monotonic on $(0, 1)$.*

(iii) *Is the function $(f_4(x) - 1)/(1 - x)$ absolutely monotonic on $(0, 1)$? This is a special of the open problem proposed in [34, Appendix G, Open Problems (2)], which is still open.*

Acknowledgements

We would like to thank the three referees for the careful review on the manuscript. This research was supported by the National Natural Science Foundation of China (11971142) and the Natural Science Foundation of Zhejiang Province (LY24A010011).

Data availability

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Declarations

Conflict of interest The authors declare that they have no conflict of interest.

REFERENCES

- [1] P. F. Byrd and M. D. Friedman, *Handbook of Elliptic Integrals for Engineers and Scientists*, Springer-Verlag, New York, 1971.
- [2] S.-L. Qiu and M. K. Vamanamurthy, Sharp estimates for complete elliptic integrals, *SIAM J. Math. Anal.* **27** (1996), no. 3, 823–834. <https://doi.org/10.1137/052704>
- [3] Z.-H. Yang, W.-M. Qian and Y.-M. Chu, Monotonicity properties and bounds involving the complete elliptic integrals of the first kind, *Math. Inequal. Appl.* **21** (2018), no. 4, 1185–1199. doi:10.7153/mia-2018-21-82
- [4] Z.-H. Yang and J. Tian, Absolutely monotonic functions involving the complete elliptic integrals of the first kind with applications, *J. Math. Inequal.* **15** (2021), no. 3, 1299–1310. DOI: 10.7153/jmi-2021-15-87
- [5] G. D. Anderson, M. K. Vamanamurthy and M. Vuorinen, Functional inequalities for complete elliptic integrals and their ratios, *SIAM J. Math. Anal.* **21** (1990), no. 2, 536–549. DOI: 10.1137/0521029
- [6] S.-L. Qiu and M. K. Vamanamurthy, Elliptic integrals and the modulus of Grötzsch ring, *Panamer. Math. J.* **5** (1995), 41–60.
- [7] Z.-H. Yang and J.-F. Tian, Convexity and monotonicity for elliptic integrals of the first kind and applications, *Appl. Anal. Discrete Math.* **13** (2019), 240–260. <https://doi.org/10.2298/AADM171015001Y>
- [8] J.-F. Tian and Z.-H. Yang, Several absolutely monotonic functions related to the complete elliptic integral of the first kind, *Results Math.* **77** (2022), Art. no. 109. <https://doi.org/10.1007/s00025-022-01641-4>
- [9] H. Alzer and K. C. Richards, A concavity property of the complete elliptic integral of the first kind, *Integral Transforms Spec. Funct.* **31** (2020), no. 9, 758–768. <https://doi.org/10.1080/10652469.2020.1738423>
- [10] J.-F. Tian and Z.-H. Yang, Convexity and monotonicity involving the complete elliptic integral of the first kind, *Results Math.* **78** (2023), Art. no. 29. <https://doi.org/10.1007/s00025-022-01799-x>
- [11] M.-K. Wang, H.-H. Chu, Y.-M. Li, and Y.-M. Chu, Answers to three conjectures on convexity of three functions involving complete elliptic integrals of the first kind, *Appl. Anal. Discrete Math.* **14** (2020), 255–271. <https://doi.org/10.2298/AADM190924020W>
- [12] Z. Yang and J. Tian, Absolute monotonicity involving the complete elliptic integrals of the first kind with applications, *Acta Math. Sci.* **42** (2022), 847–864. <https://doi.org/10.1007/s10473-022-0302-x>
- [13] S.-L. Qiu, M. Vuorinen, Infinite products and the normalized quotients of hypergeometric functions, *SIAM J. Math. Anal.* **30** (1999), no. 5, 1057–1075. <https://doi.org/10.1137/S003614109732680>
- [14] Z.-H. Yang, Y.-M. Chu, A monotonicity property involving the generalized elliptic integrals of the first kind, *Math. Inequal. Appl.* **20** (2017), no. 3, 729–735. doi:10.7153/mia-20-46
- [15] M.-K. Wang, W. Zhang and Y.-M. Chu, Monotonicity, convexity and inequalities involving the generalized elliptic integrals, *Acta Math. Sci.* **39B** (2019), no. 5, 1440–1450. <https://doi.org/10.1007/s10473-019-0520-z>
- [16] Z.-H. Yang and J.-F. Tian, Sharp inequalities for the generalized elliptic integrals of the first kind, *Ramanujan J.* **48** (2019), 91–116. <https://doi.org/10.1007/s11139-018-0061-4>

- [17] K. C. Richards and J. N. Smith, A concavity property of generalized complete elliptic integrals, *Integral Transforms Spec. Funct.* **32** (2020), no. 3, 1–13. DOI: 10.1080/10652469.2020.1815726
- [18] T.-H. Zhao, M.-K. Wang and Y.-M. Chu, Monotonicity and convexity involving generalized elliptic integral of the first kind, *Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat.* **115** (2021): 46. <https://doi.org/10.1007/s13398-020-00992-3>
- [19] T.-H. Zhao, M.-K. Wang and Y.-M. Chu, Concavity and bounds involving generalized elliptic integral of the first kind, *J. Math. Inequal.* **15** (2021), no. 2, 701–724. doi:10.7153/jmi-2021-15-50
- [20] Y.-J. Chen and T.-H. Zhao, On the monotonicity and convexity for generalized elliptic integral of the first kind, *Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat.* **116** (2022): 77. <https://doi.org/10.1007/s13398-022-01211-x>
- [21] M.-K. Wang, T.-H. Zhao, X.-J. Ren, Y.-M. Chu and Z.-Y. He, Monotonicity and concavity properties of the Gaussian hypergeometric functions, with applications, *Indian J. Pure Appl. Math.* **54** (2023), 1105–1124. <https://doi.org/10.1007/s13226-022-00325-7>
- [22] B. C. Carlson, J. L. Guatafson, Asymptotic expansion of the first elliptic integral, *SIAM J. Math. Anal.* **16** (1985), no. 5, 1072–1092. <https://doi.org/10.1137/0516080>
- [23] G. D. Anderson, M. K. Vamanamurthy and M. Vuorinen, Functional inequalities for hypergeometric functions and complete elliptic integrals, *SIAM J. Math. Anal.* **23** (1992), no. 2, 512–524. DOI 10.1137/0523025
- [24] S.-L. Qiu, M. K. Vamanamurthy and M. Vuorinen, Some inequalities for the growth of elliptic integrals, *SIAM J. Math. Anal.* **29** (1998), no. 5, 1224–1237. <https://doi.org/10.1137/S0036141096310491>
- [25] H. Alzer, Sharp inequalities for the complete elliptic integral of the first kind, *Math. Proc. Camb. Phil. Soc.* **124** (1998), no. 2, 309–314. DOI: <https://doi.org/10.1017/S0305004198002692>
- [26] H. Alzer and S.-L. Qiu, Monotonicity theorems and inequalities for the complete elliptic integrals, *J. Comput. Appl. Math.* **172** (2004), 289–312. <https://doi.org/10.1016/j.cam.2004.02.009>
- [27] Z.-H. Yang, Y.-Q. Song and Y.-M. Chu, Sharp bounds for the arithmetic-geometric mean, *J. Inequal. Appl.* **2014** (2014): 192. DOI: 10.1186/1029-242X-2014-192
- [28] Z.-H. Yang, W.-M. Qian, W. Zhang and Y.-M. Chu, Notes on the complete elliptic integral of the first kind, *Math. Inequal. Appl.* **23** (2020), no. 1, 77–93. DOI: 10.7153/mia-2020-23-07
- [29] Z.-H. Yang, J.-F. Tian and Y.-R. Zhu, A sharp lower bound for the complete elliptic integrals of the first kind, *Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat.* **115** (2021): 8. <https://doi.org/10.1007/s13398-020-00949-6>
- [30] M. Petrović, Sur une équation fonctionnelle, *Publ. Math. Univ. Belgrade* **1** (1932), 149–156.
- [31] M. Abramowitz and I. A. Stegun (Eds), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Applied Mathematics Series 55, 10th printing, Dover Publications, New York and Washington, 1972.
- [32] Z.-H. Yang, Y.-M. Chu, and M.-K. Wang, Monotonicity criterion for the quotient of power series with applications, *J. Math. Anal. Appl.* **428** (2015), no. 1, 587–604. <https://doi.org/10.1016/j.jmaa.2015.03.043>
- [33] M. Biernacki, J. Krzyz, On the monotonicity of certain functionals in the theory of analytic functions, *Annales Universitatis Mariae Curie-Sklodowska*, **9** (1955), 135–147.
- [34] G. D. Anderson, M. K. Vamanamurthy and M. Vuorinen, *Conformal Invariants, Inequalities, and Quasiconformal Maps*, John Wiley & Sons, New York, 1997.

TIEHONG ZHAO, SCHOOL OF MATHEMATICS, HANGZHOU NORMAL UNIVERSITY, HANGZHOU, ZHEJIANG, CHINA, 311121

Email address: tiehong.zhao@hznu.edu.cn

URL: <https://orcid.org/0000-0002-6394-1049>

ZHEN-HANG YANG, STATE GRID ZHEJIANG ELECTRIC POWER COMPANY RESEARCH INSTITUTE, HANGZHOU, ZHEJIANG, CHINA, 310014

Email address: yzhkm@163.com

URL: <https://orcid.org/0000-0002-2719-4728>