# Symmetries in Overparametrized Neural Networks: A Mean-Field View

Javier Maass Martínez \* Center for Mathematical Modeling University of Chile Joaquín Fontbona<sup>†</sup> Center for Mathematical Modeling University of Chile

## Abstract

We develop a Mean-Field (MF) view of the learning dynamics of overparametrized Artificial Neural Networks (NN) under distributional symmetries of the data w.r.t. the action of a general compact group G. We consider for this a class of generalized shallow NNs given by an ensemble of N multi-layer units, jointly trained using stochastic gradient descent (SGD) and possibly symmetry-leveraging (SL) techniques, such as Data Augmentation (DA), Feature Averaging (FA) or Equivariant Architectures (EA). We introduce the notions of weakly and strongly invariant laws (WI and SI) on the parameter space of each single unit, corresponding, respectively, to G-invariant distributions, and to distributions supported on parameters fixed by the group action (which encode EA). This allows us to define symmetric models compatible with taking  $N \rightarrow \infty$  and give an interpretation of the asymptotic dynamics of DA, FA and EA in terms of Wasserstein Gradient Flows describing their MF limits. When activations respect the group action, we show that, for symmetric data, DA, FA and freely-trained models obey the exact same MF dynamic, which stays in the space of WI parameter laws and attains therein the population risk's minimizer. We also provide a counterexample to the general attainability of such an optimum over SI laws. Despite this, and quite remarkably, we show that the space of SI laws is also preserved by these MF distributional dynamics even when freely trained. This sharply contrasts the finite-N setting, in which EAs are generally not preserved by unconstrained SGD. We illustrate the validity of our findings as N gets larger, in a teacher-student experimental setting, training a student NN to learn from a WI, SI or arbitrary teacher model through various SL schemes. We last deduce a data-driven heuristic to discover the largest subspace of parameters supporting SI distributions for a problem, that could be used for designing EA with minimal generalization error.

# 1 Introduction

Learning in complex tasks, employing ever larger datasets, has strongly benefited from the implementation and training of Artificial Neural Networks (NN) with a huge number of parameters; as well as from training schemes or architectures that can leverage underlying symmetries of the data in order to reduce the problem's complexity (see [27, 28] for general reference). This raises questions, on one hand, of understanding the puzzling generalizability in overparametrized NN; and on the other, of when and how symmetry-leveraging (SL) techniques (such as Data Augmentation, Feature Averaging or Equivariant Architectures), can induce useful biases towards learning with symmetries, without hindering approximation and generalization properties. The recent Mean-Field (**MF**) theory of NN (see [14] and further references below) provides a partial, yet promissory, viewpoint to address the

<sup>\*</sup>javier.maass@gmail.com

<sup>&</sup>lt;sup>†</sup>fontbona@dim.uchile.cl

first question for shallow NN: in the Mean-Field *Limit* (**MFL**) of an infinitely wide hidden layer, stochastic gradient descent (SGD) training dynamics approximates the Wasserstein Gradient Flow (**WGF**) of certain *convex population risk* on the space of distributions on parameters. Confluently, the incorporation of combined algebraic and probabilistic viewpoints have yielded a more complete view of the benefits of SL techniques under symmetry (see e.g. [11, 22, 47] and further references below); however, it is not clear if and how those findings can scale to overparametrized NN and their **MFL**.

In this work we develop a systematic **MF** analysis of the limiting learning dynamics of a class of generalized shallow NNs, under distributional symmetries of the data w.r.t. the action of a compact group, and including the possible effects of employing some of the most popular SL techniques. The effect of symmetries on the **WGF** dynamics was already studied in [29], in the particular case of two-layer ReLU networks, under data generated by a function symmetric w.r.t. a single orthogonal transformation. We consider our (independent <sup>3</sup>) work to largely broaden the scope and applicability of such initial contributions, as it provides a unified **MF** interpretation for both the use of SL techniques under general distributional invariances, and the interplay of such symmetries at the levels of data, architectures and training dynamics. The paper unfolds as follows:

In Section 2 we introduce a class of generalized shallow models with multi-layer units on which we will focus, we recall **WGFs** and their role in the **MFL** of NN training dynamics, and review the SL techniques to be studied. Section 3 contains the bulk of our contributions, as we study how SL techniques applied on these models can be interpreted in terms of their limiting **WGFs**, how they relate to each other in terms of the optima of their corresponding population risks, and how their limiting **MF** training dynamics behave with or without symmetric data. Finally, Section 4 presents the empirical *validation* of our main theoretical results through some empirical simulations; it also suggests a potential heuristic for *architecture-discovery* in ML problems. Proofs and complements to our results can be found in the Supplementary Material (henceforth SuppMat for short).

# 2 Preliminaries

## 2.1 Supervised learning with generalized shallow neural networks

Let  $\mathcal{X}$ ,  $\mathcal{Y}$  and  $\mathcal{Z}$  be separable Hilbert Spaces, termed as the *feature*, *label* and *parameter* spaces respectively. Typically, these are finite-dimensional, e.g.  $\mathcal{X} = \mathbb{R}^d$  and  $\mathcal{Y} = \mathbb{R}^c$  (for  $c, d \in \mathbb{N}^*$ ) with  $\mathcal{Z}$  the space of affine transformations between hidden layers. We write  $\mathcal{P}(\cdot)$  for the space of Borel probability measure on a metric space ( $\cdot$ ). Let  $\pi \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$  denote the data distribution from which i.i.d. samples  $(X, Y) \in \mathcal{X} \times \mathcal{Y}$  will be drawn, and  $\ell : \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$  be a **convex** loss function. Consider also an *activation function*  $\sigma_* : \mathcal{X} \times \mathcal{Z} \to \mathcal{Y}$ . We introduce a general class of shallow NN:

**Definition 1.** A shallow neural network model of parameter  $\theta := (\theta_i)_{i=1}^N \in \mathbb{Z}^N$  is the function  $\Phi_{\theta}^N : \mathcal{X} \to \mathcal{Y}$  given by  $\Phi_{\theta}^N(x) := \frac{1}{N} \sum_{i=1}^N \sigma_*(x;\theta_i), \forall x \in \mathcal{X}$ . Equivalently, if  $\nu_{\theta}^N := \frac{1}{N} \sum_{i=1}^N \delta_{\theta_i}$  is the empirical measure associated with  $\theta \in \mathbb{Z}^N$ , we can write  $\forall x \in \mathcal{X}, \Phi_{\theta}^N(x) = \langle \sigma_*(x; \cdot), \nu_{\theta}^N \rangle$  or, abusing notation, simply  $\Phi_{\theta}^N = \langle \sigma_*, \nu_{\theta}^N \rangle$ .

In the setting where  $\mathcal{X} = \mathbb{R}^d$ ,  $\mathcal{Y} = \mathbb{R}^c$  and  $\mathcal{Z} = \mathbb{R}^{c \times b} \times \mathbb{R}^{d \times b} \times \mathbb{R}^b$  (for  $b \in \mathbb{N}^*$ ) this corresponds exactly to a single-hidden-layer neural network with N hidden units, by defining, for  $z = (W, A, B) \in \mathcal{Z}$  and  $\sigma : \mathbb{R}^b \to \mathbb{R}^b$ ,  $\sigma_*(x, z) := W\sigma(A^Tx + B)$ . Depending on  $\sigma_*$ , however, these shallow NN models can go way beyond this example. In fact,  $\sigma_*$  can be taken to be an entire Multi-Layer NN model, in which case  $\Phi^N_{\theta}$  will represent an **ensemble** of N such units trained simultaneously (see SuppMat-A.4). They can also model the deconvolution of sparse spikes, RBF networks, density estimation via MMD minimization, among many others (see [14, 50, 56]).

This class thus allows for non-trivial internal units, while enabling the width  $N \to \infty$  consistently, and regardless of the possible underlying structure of the (fixed size) units represented by  $\sigma_*$ . Inspired by this possibility, and by our writing of *shallow NN models*, we define a more general notion:

**Definition 2** (Shallow Model). A shallow model is any function of the form  $\Phi_{\mu}(x) := \langle \sigma_*(x; \cdot), \mu \rangle$ for some  $\mu \in \mathcal{P}(\mathcal{Z})$  (whenever the integral makes sense for all  $x \in \mathcal{X}$ ). We write  $\Phi_{\mu} := \langle \sigma_*, \mu \rangle$  and denote the space of such models as  $\mathcal{F}_{\sigma_*}(\mathcal{P}(\mathcal{Z}))$ .

<sup>&</sup>lt;sup>3</sup>We became aware of that work in [29] after a first version of this paper was posted.

Classically, we want to find a NN model that performs well with respect to  $\pi$  and  $\ell$ . More precisely, having fixed an *architecture* (given here by N and  $\sigma_*$ ), we consider the *generalization error* or *population risk* given by  $R(\theta) = \mathbb{E}_{\pi} \left[ \ell(\Phi_{\theta}^N(X), Y) \right]$ , and look for a vector of parameters  $\theta \in \mathbb{Z}^N$  attaining  $\min_{\theta \in \mathbb{Z}^N} R(\theta)$ . However, not only is this function highly non-convex and hard to optimize; but in practice we generally don't have access to  $\pi$  (and thus R) and we have to solve this problem only with a set of i.i.d. data samples  $\{(X_k, Y_k)\}_{k \in \mathbb{N}}$  drawn from  $\pi$ . The usual way of *training* a *NN model*  $\Phi_{\theta}^N$  thus is by following an SGD scheme (see e.g. [5]) of the type :

- First, initialize  $\theta_i^0, \forall i \in \{1, \dots, N\}$ , i.i.d. from a fixed distribution  $\mu_0 \in \mathcal{P}(\mathcal{Z})$ .
- Iterate, for  $k \in \mathbb{N}$ , defining  $\forall i \in \{1, \dots, N\}$ :

$$\theta_i^{k+1} = \theta_i^k - s_k^N \left( \nabla_z \sigma_*(X_k, \theta_i^k) \cdot \nabla_1 \ell(\Phi_{\theta^k}^N(X_k), Y_k) + \tau \nabla r(\theta_i^k) \right) + \sqrt{2\beta s_k^N} \xi_i^k.$$
(1)

Here,  $s_k^N = \varepsilon_N \varsigma(k\varepsilon_N)$  is the *step-size* (or *learning rate*), parametrized in terms of  $\varsigma : \mathbb{R}_+ \to \mathbb{R}_+$ a regular function and  $\varepsilon_N > 0$ . Also, we have a penalization function  $r : \mathbb{Z} \to \mathbb{R}$ , regularizing gaussian noise  $\xi_i^k \stackrel{i.i.d.}{\sim} \mathcal{N}(0, \operatorname{Id}_{\mathbb{Z}})$  independent from the initialization, and  $\tau, \beta \ge 0$ . When  $\tau, \beta > 0$ , the method is called stochastic gradient Langevin dynamics or SGLD ([60]) or simply *noisy* SGD, which we will use for simplicity. Note that requiring an *infinite* and *i.i.d.* sample from  $\pi$  is key when letting later  $N \to \infty$ . This is not truly a restriction, since one can always sample from the empirical distribution of finite data points. We assume *i.i.d.* parameter initialization for simplicity.

In principle, there are no guarantees that this training procedure will be truly optimizing  $R(\theta)$  let alone approaching its minimum. However, by extending the definition of the **generalization error** to models in  $\mathcal{F}_{\sigma_*}(\mathcal{P}(\mathcal{Z}))$ , one gets the convex functional  $R: \mathcal{P}(\mathcal{Z}) \to \mathbb{R}$  given by  $R(\mu) := \mathbb{E}_{\pi} \left[ \ell(\Phi_{\mu}(X), Y) \right]$ . The problem on  $\mathcal{Z}^N$  is thus lifted to the **convex** optimization problem on  $\mathcal{P}(\mathcal{Z})$ :

$$\min_{\mu \in \mathcal{P}(\mathcal{Z})} R(\mu).$$
<sup>(2)</sup>

Accordingly, this motivates looking at the evolution of empirical measures  $(\nu_k^N)_{k \in \mathbb{N}} := (\nu_{\theta^k}^N)_{k \in \mathbb{N}} \subseteq \mathcal{P}(\mathcal{Z})$  instead of that of the specific parameters  $(\theta^k)_{k \in \mathbb{N}} \subseteq \mathcal{Z}^N$ . The **MF** approach to NNs (see [14, 42, 50, 55]) aims at providing theoretical guarantees for problem (2), justifying that a *global optimum* of the population risk can be approximated by training a NN with SGD for large N. We next provide some necessary background on **WGF**s and on the **MF** theory of shallow NN models.

## 2.2 Wasserstein gradient flow and mean-field limit of shallow models

We briefly recall some elements of Optimal Transport and Wasserstein Gradient Flows, referring to [1, 51, 58] for further background. Let  $\mathcal{Z}$  be a Hilbert space with norm  $\|\cdot\|$  and, for  $p \in [1, \infty)$ , let  $\mathcal{P}_p(\mathcal{Z}) := \{\mu \in \mathcal{P}(\mathcal{Z}) : \int_{\mathcal{Z}} \|\theta\|^p \mu(d\theta) < +\infty\}$  be the space of probability measures on  $\mathcal{Z}$  with finite *p*-th moment. We endow this space with the *p*-th *Wasserstein distance*, defined as:  $W_p(\mu, \nu) := \left[\inf_{\gamma \in \Pi(\mu,\nu)} \mathbb{E}_{\gamma}[\|X - Y\|^p]\right]^{\frac{1}{p}}, \forall \mu, \nu \in \mathcal{P}_p(\mathcal{Z}) \text{ with } \Pi(\mu,\nu) \text{ being the set of$ *couplings between* $<math>\mu$  and  $\nu$  (the infimum is always attained). The metric space  $(\mathcal{P}_p(\mathcal{Z}), W_p)$  is Polish and called the *p*-th Wasserstein Space. In the remainder of this section we consider p = 2 and  $\mathcal{Z} = \mathbb{R}^D$ .

We recall important objects for the sequel, including the Lions derivative [7, 37] popularized in the area of mean-field games (see e.g. [8, 10, 13, 31]), also shown in [26] to coincide with the Wassertein gradient ([1]):

**Definition 3** (Linear Functional Derivative and Intrinsic Derivative). Given  $F : \mathcal{P}_2(\mathcal{Z}) \to \mathbb{R}$ , its linear functional derivative is the function (if it exists)  $\frac{\partial F}{\partial \mu} : Dom(F) \times \mathcal{Z} \to \mathbb{R}$  such that  $\forall \mu, \nu \in Dom(F)$ ,  $\lim_{h\to 0} \frac{F((1-h)\mu+h\nu)-F(\mu)}{h} = \int_{\mathcal{Z}} \frac{\partial F}{\partial \mu}(\mu, z)d(\nu-\mu)(z)$  and  $\int_{\mathcal{Z}} \frac{\partial F}{\partial \mu}(\mu, z)d\mu(z) = 0$ . The function  $F' : \mu \in \mathcal{P}_2(\mathcal{Z}) \mapsto \frac{\partial F}{\partial \mu}(\mu, \cdot)$  is known as the first variation of F at  $\mu$ . Moreover, if  $\frac{\partial F}{\partial \mu}$  exists and is differentiable in its second argument, we define the intrinsic derivative of F at  $\mu$  to be:  $D_{\mu}F(\mu, z) = \nabla_z \left(\frac{\partial F}{\partial \mu}(\mu, z)\right)$ . Abusing notation, we will write  $\frac{\partial F}{\partial \mu} : \mathcal{P}_2(\mathcal{Z}) \times \mathcal{Z} \to \mathbb{R}$  and  $D_{\mu}F : \mathcal{P}_2(\mathcal{Z}) \times \mathcal{Z} \to \mathcal{Z}$ , even if they are only partially defined.

This allows us to define next a Wasserstein Gradient Flow (following e.g. [1, 14]):

**Definition 4.** [Wasserstein Gradient Flow (WGF)] Let  $\varsigma : \mathbb{R}_+ \to \mathbb{R}_+$  be a regular scalar function and  $F : \mathcal{P}_2(\mathcal{Z}) \to \mathbb{R}$  be a convex functional for which the intrinsic derivative  $D_\mu F$  is defined. We define a Wasserstein Gradient Flow (WGF) for F (shortened WGF(F)) as any absolutely continuous trajectory  $(\mu_t)_{t>0}$  in  $\mathcal{P}_2(\mathcal{Z})$  that satisfies, distributionally on  $[0, \infty) \times \mathcal{Z}$ :

$$\partial_t \mu_t = \varsigma(t) \operatorname{div} \left( D_\mu F(\mu_t, \cdot) \mu_t \right).$$

Several authors ([1, 14, 51, 58], among others) have proven under various sets of assumptions that, given an initial condition  $\mu_0 \in \mathcal{P}_2(\mathcal{Z})$ , the **WGF**(*F*) admits a unique (weak) solution,  $(\mu_t)_{t\geq 0}$ . In a sense, **WGF**(*F*) "follows the negative gradient" of *F*. Unfortunately, even for convex *F*, stationary points of **WGF**(*F*) need not be global minima, see [14].

We are interested in the case where F is the following convex, **entropy-regularized population risk**:  $R^{\tau,\beta}(\mu) := R(\mu) + \tau \int r d\mu + \beta H_{\lambda}(\mu)$ , where  $\tau, \beta \ge 0, \lambda$  is the Lebesgue Measure on Z,  $r: Z \to \mathbb{R}_+$  is a *penalization*, and  $H_{\lambda}$  defined as  $H_{\lambda}(\mu) := \int \log(\frac{d\mu}{d\lambda}(z))d\mu(z)$  if  $\mu \ll \lambda$  or  $+\infty$  otherwise, is the *Boltzmann entropy* of  $\mu$ . In this case, **WGF** $(R^{\tau,\beta})$  reads as the PDE:

$$\partial_t \mu_t = \varsigma(t) \left[ \operatorname{div} \left( \left( D_\mu R(\mu_t, \cdot) + \tau \nabla_\theta r) \, \mu_t \right) + \beta \Delta \mu_t \right], \tag{3}$$

known as McKean-Vlasov equation in the probability and PDE communities (see the classic references [43, 57], and the recent review [9]) and popularized as "distributional dynamics" in NN literature (e.g. [42]). When  $\beta > 0$ , a solution to (3) has a density w.r.t.  $\lambda$  and is actually strong. Under rather simple technical assumptions (see SuppMat-B.3, or [10, 13, 31, 45, 56]), when  $\tau$ ,  $\beta > 0$  it is known that the **WGF**( $R^{\tau,\beta}$ )  $W_2$ -converges to a (unique) minimizer. When  $\tau$ ,  $\beta = 0$  a sort of converse holds (see [14]): if **WGF**(R) converges in  $W_2$ , then the limit minimizes R.

Proven by [14, 42, 50, 55] and later refined e.g. by [12, 19, 20, 40, 54, 56], the main result in the **MF** Theory of overparametrized shallow NNs states that SGD *training* for a shallow NN, in the right *scaling* limit as  $N \to \infty$ , approximates a **WGF** :

**Theorem 1** (Mean-Field limit, sketch). For each T > 0, under relevant technical assumptions including regularity of  $\sigma_*$  and a proper asymptotic behaviour of  $\varepsilon_N \to 0$  as  $N \to \infty$ , the rescaled empirical process given by  $\mu^N := (\nu_{\lfloor t/\varepsilon_N \rfloor}^N)_{t \in [0,T]}$  converges in law (in the Skorokhod space  $D_{\mathcal{P}(\mathcal{Z})}([0,T])$ ) to  $\mu := (\mu_t)_{t \in [0,T]}$  given by the **unique WGF** $(R^{\tau,\beta})$  starting at  $\mu_0$ .

### 2.3 Symmetry-leveraging techniques

We next discuss mathematical formulations of the main techniques to leverage posited distributional symmetries of the data at the training or architecture levels. We henceforth fix a *compact group* G of normalized Haar measure  $\lambda_G$ , acting on  $\mathcal{X}$  and  $\mathcal{Y}$ , which we denote  $G \ominus_{\rho} \mathcal{X}$ ,  $G \ominus_{\hat{\rho}} \mathcal{Y}$ . <sup>4</sup> A function  $f: \mathcal{X} \to \mathcal{Y}$  is termed *equivariant* if  $\forall g \in G$ ,  $\hat{\rho}_{g^{-1}}.f(\rho_g.x) = f(x) \ d\pi_{\mathcal{X}}(x)$ -a.s. We further say that the data  $(X, Y) \sim \pi$  is *equivariant*, and write  $\pi \in \mathcal{P}^G(\mathcal{X} \times \mathcal{Y})$ , if  $\forall g \in G$ ,  $(\rho_g.X, \hat{\rho}_g.Y) \sim \pi$  (this is not enforced unless stated). The space of functions  $f: \mathcal{X} \to \mathcal{Y}$  square-integrable (in Bochner sense) w.r.t  $\pi_{\mathcal{X}} = Law(X)$  is called  $L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})$ . Further relevant concepts are introduced as needed.

**Data Augmentation (DA):** This training scheme considers  $\{g_k\}_{k\in\mathbb{N}} \stackrel{i.i.d.}{\sim} \lambda_G$  independent from the  $\{(X_k, Y_k)\}_{k\in\mathbb{N}}$  in (1), and carries out SGD on samples  $\{(\rho_{g_k}.X_k, \hat{\rho}_{g_k}.Y_k)\}_{k\in\mathbb{N}}$ . **DA** and the *vanilla* training scheme would thus be equivalent if  $\pi \in \mathcal{P}^G(\mathcal{X} \times \mathcal{Y})$ . One can show (see [11, 38]) that, performing SGD with **DA**, results in an optimization scheme for the *symmetrized population risk*,  $R^{DA}(\theta) := \mathbb{E}_{\pi} \left[ \int_G \ell \left( \Phi_{\theta}^N(\rho_g.X), \hat{\rho}_g.Y \right) d\lambda_G(g) \right]$ . Despite being effective in practice, **DA** gives no guarantee that the resulting model will be equivariant. For deeper insights, see [11, 18, 32, 36, 38, 41].

Feature Averaging (FA): Instead of focusing on the data, FA works with symmetrized versions of the vanilla NN models  $\Phi_{\theta}^{N}$  at hand, averaging model copies over all possible translations through the group action. This amounts to constructing (or approximating) the symmetrization operator over  $L^{2}(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})$  defined as  $(\mathcal{Q}_{G}.f)(x) := \int_{G} \hat{\rho}_{g^{-1}}.f(\rho_{g}.x)d\lambda_{G}(g)$  (see [22]), and trying to minimize  $R^{FA}(\theta) := \mathbb{E}_{\pi} \left[ \ell \left( (\mathcal{Q}_{G}.\Phi_{\theta}^{N})(X), Y \right) \right]$  (see [11, 18, 36, 38]). The resulting model will be equivariant, however, FA is inefficient, as  $\approx |G|$  times more evaluations are needed for training and inference.

Equivariant Architectures (EA): Following [6], EA in multilayer NNs are configurations yielding models equivariant between each of the hidden layers (where G is assumed to act). As stated in

<sup>&</sup>lt;sup>4</sup>w.l.o.g. by compactness, all *G*-actions are assumed to be via orthogonal representations

[24, 25, 49, 52, 53, 61] and [2, 15, 34, 35, 59], once the (equivariant) activation functions between the different layers have been fixed, **EAs** are plainly parameter-sharing schemes (determined by the space of *intertwinners/group convolutions* between layers). In our context, assuming that  $G \odot_M Z$ is some group action, we require that  $\sigma_* : \mathcal{X} \times Z \to \mathcal{Y}$  is *jointly equivariant*, namely,  $\forall (g, x, z) \in$  $G \times \mathcal{X} \times Z$ ,  $\sigma_*(\rho_g.x, M_g.z) = \hat{\rho}_g \sigma_*(x, z)$ ; to ensure *G*-actions over different spaces are properly related. Introducing the set of fixed points for  $G \odot_M Z$ ,  $\mathcal{E}^G := \{z \in Z : \forall g \in G, M_g.z = z\}$ , a shallow NN model  $\Phi^N_{\theta}$  thus has an **EA** if  $\theta \in (\mathcal{E}^G)^N$ . Under the right choices of  $\sigma_*$  and *M*, the obtained **EAs** can encode interesting and widely applied architectures, such as CNNs [16] and DeepSets [63] (see SuppMat-A.4 for further discussion). We call  $\mathcal{E}^G$  the **subspace of invariant parameters**, which is a closed linear subspace of Z, with unique *orthogonal projection*  $P_{\mathcal{E}^G}: Z \to$  $\mathcal{E}^G$ , explicitly given by  $P_{\mathcal{E}^G.z} := \int_G M_g.z d\lambda_G(g)$  for  $z \in Z$  (see [25]). We are thus led to solve:  $\min_{\theta \in (\mathcal{E}^G)^N} R(\theta)$  or, equivalently, to find the best *projected* model  $\Phi^{N,EA}_{\theta} := \langle \sigma_*, P_{\mathcal{E}^G} \# \nu^N_{\theta} \rangle$ , by minimizing  $R^{EA}(\theta) := \mathbb{E}_{\pi} \left[ \ell \left( \Phi^{N,EA}_{\theta}(X), Y \right) \right]$ . This can considerably reduce the parameter space dimension; however **EA** might generally yield a decreased expressivity or approximation capacity.

## **3** Symmetries in overparametrized neural networks: main results

## 3.1 Two notions of symmetries for parameter distributions

The following notions regarding distributions from  $\mathcal{P}(\mathcal{Z})$  are central to our work:

**Definition 5.** Given  $\mu \in \mathcal{P}(\mathcal{Z})$ , we respectively define its symmetrized and projected versions as  $\mu^G := \int_G (M_g \# \mu) d\lambda_G$  and  $\mu^{\mathcal{E}^G} := P_{\mathcal{E}^G} \# \mu$ . Moreover, we introduce two subspaces of  $\mathcal{P}(\mathcal{Z})$ :  $\mathcal{P}^G(\mathcal{Z}) := \{\mu \in \mathcal{P}(\mathcal{Z}) : \forall g \in G, M_g \# \mu = \mu\}$  and  $\mathcal{P}(\mathcal{E}^G) := \{\mu \in \mathcal{P}(\mathcal{Z}) : \mu(\mathcal{E}^G) = 1\}.$ 

**Definition 6** (Invariant Probability Measures). We say that  $\mu \in \mathcal{P}(\mathcal{Z})$  is:

Weakly-Invariant (WI) if  $\mu = \mu^G$  and Strongly-Invariant (SI) if  $\mu = \mu^{\mathcal{E}^G}$ .

We notice that:  $\mathcal{P}(\mathcal{E}^G) \subseteq \mathcal{P}^G(\mathcal{Z}), \ \mu^G \in \mathcal{P}^G(\mathcal{Z}) \text{ and } \mu^{\mathcal{E}^G} \in \mathcal{P}(\mathcal{E}^G).$  Thus, **SI** implies **WI**. Next result relates the symmetrization operation on  $\mathcal{P}(\mathcal{Z})$  with the one on shallow models  $\mathcal{F}_{\sigma_*}(\mathcal{P}(\mathcal{Z}))$ : **Proposition 1.** Let  $\Phi_{\mu} \in \mathcal{F}_{\sigma_*}(\mathcal{P}(\mathcal{Z}))$  with  $\sigma_* : \mathcal{X} \times \mathcal{Z} \to \mathcal{Y}$  jointly equivariant. Then:

$$(\mathcal{Q}_G \Phi_\mu) = \Phi_{\mu^G}$$

That is to say, the closest equivariant function (in  $L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})$ ) to  $\Phi_{\mu}$  is given by the shallow model associated to the symmetrized version of  $\mu$ .

**Remark.** In particular,  $\Phi_{\mu}$  is equivariant as soon as  $\mu$  is **WI** only. Conversely, if  $\Phi_{\mu} : \mathcal{X} \to \mathcal{Y}$  is an equivariant function, then  $\Phi_{\mu} = \Phi_{\mu^{G}}$ , i.e. it can be expressed in terms of a **WI** distribution. This highlights a priority role of **WI** distributions on  $\mathcal{Z}$  in representing invariant shallow models.

The alternative, "projected model"  $\Phi_{\mu^{\mathcal{E}G}}$ , in turn, is never the closest equivariant shallow model, to  $\Phi_{\mu}$  in  $L^2(\mathcal{X}, \mathcal{Y}, \pi_{\mathcal{X}})$ , unless equal to  $\Phi_{\mu^G}$ . The latter rarely is the case (unlike commonly implied in the literature). In fact, the symmetrized version  $\mathcal{Q}_G \Phi_{\theta}^N$  of a shallow NN model  $\Phi_{\theta}^N$  involves  $(\nu_{\theta}^N)^G = \frac{1}{N} \sum_{i=1}^N \varphi_{\theta_i}$ , where  $\forall z \in \mathcal{Z}, \varphi_z$  is the orbit measure of the action,<sup>5</sup> and has  $N \cdot |G|$  $\mathcal{Z}$ -valued parameters (possibly with  $|G| = \infty$ ). This sharply contrasts  $(\nu_{\theta}^N)^{\mathcal{E}^G} = \frac{1}{N} \sum_{i=1}^N \delta_{P_{\mathcal{E}^G},\theta_i}$ , which has  $\leq N$  distinct parameters, all living in  $\mathcal{E}^G$ . So, in general, depending on  $\sigma_*$  and G, one might have  $\langle \sigma_*, (\nu_{\theta}^N)^{\mathcal{E}^G} \rangle \neq \mathcal{Q}_G \Phi_{\theta}^N$ . A notable case in which the equality holds is the class of linear models, which is discussed in SuppMat-C.1.

## **3.2** Invariant functionals on $\mathcal{P}(\mathcal{Z})$ and their optima

In the same spirit as when defining the population risk  $R : \mathcal{P}(\mathcal{Z}) \to \mathbb{R}$  in (2), the risk functions associated with SL-techniques from Section 2.3 can be lifted to functionals over  $\mathcal{P}(\mathcal{Z})$ , namely

<sup>&</sup>lt;sup>5</sup>defined as:  $\varphi_z = T_z \# \lambda_G$ , where  $T_z := [g \in G \mapsto M_g . z \in \mathcal{Z}]$ 

to:  $R^{DA}(\mu) := \mathbb{E}_{\pi} \left[ \int_{G} \ell \left( \Phi_{\mu}(\rho_{g}.X), \hat{\rho}_{g}.Y \right) d\lambda_{G}(g) \right], R^{FA}(\mu) := \mathbb{E}_{\pi} \left[ \ell \left( \mathcal{Q}_{G}(\Phi_{\mu})(X), Y \right) \right]$  and  $R^{EA}(\mu) := \mathbb{E}_{\pi} \left[ \ell \left( \Phi_{\mu^{\mathcal{E}G}}(X), Y \right) \right]$ , respectively. This will allow us to study these SL-techniques, in the overparametrized regime, under a common **MF** framework. We need the following assumption:

**Assumption 1.**  $\pi \in \mathcal{P}_2(\mathcal{X} \times \mathcal{Y}); \ell : \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$  is convex, jointly invariant and differentiable with  $\nabla_1 \ell$  linearly growing; and  $\sigma_* : \mathcal{X} \times \mathcal{Z} \to \mathcal{Y}$  is bounded, jointly equivariant and differentiable.

The quadratic loss  $\ell(y, \hat{y}) = \frac{1}{2}||y - \hat{y}||^2$  is an example of such  $\ell$ . Now, having bounded and differentiable  $\sigma_*$  is generally very restrictive since, in practice, many commonly used activations don't satisfy either. Assuming this is, however, standard in the **MF** literature (see e.g. [10, 31, 42]), and crucial for establishing insightful key results such as global convergence. Overcoming these limitations to make **MF** theory more broadly applicable is an active area of research. We also define:

**Definition 7.** A functional  $F : \mathcal{P}(\mathcal{Z}) \to \mathbb{R}$  is invariant if  $F(M_g \# \mu) = F(\mu) \forall (g, \mu) \in G \times \mathcal{P}(\mathcal{Z})$ ; equivalently, if it equals its symmetrized version  $F^G(\mu) := \int_G F(M_g \# \mu) d\lambda_G(g)$ .

**Proposition 2.** Under Assumption 1,  $R^{DA}$ ,  $R^{FA}$  and  $R^{EA}$  are invariant (and convex) and we have:

$$R^{DA}(\mu) = R^{G}(\mu), \ R^{FA}(\mu) = R(\mu^{G}) \ and \ R^{EA}(\mu) = R(\mu^{\mathcal{E}^{G}}).$$

In particular,  $R = R^{DA}$  if R is invariant. Moreover,  $\forall \mu \in \mathcal{P}^G(\mathcal{Z})$ ,  $R(\mu) = R^{DA}(\mu) = R^{FA}(\mu)$ . Last, if  $\pi \in \mathcal{P}^G(\mathcal{X} \times \mathcal{Y})$  (the data distribution is equivariant), then R is invariant.

The proof relies on Proposition 1 and calculations as in [25], see SuppMat-C.2. Next result is a general property of functionals over  $\mathcal{P}(\mathcal{Z})$ , which is key for the forthcoming analysis:

**Proposition 3** (Optimality for Invariant Functionals). Let  $F : \mathcal{P}(\mathcal{Z}) \to \mathbb{R}$  be convex,  $\mathcal{C}^1$  and invariant. Then:  $\forall \mu \in \mathcal{P}(\mathcal{Z}), F(\mu^G) \leq F(\mu)$ ; and so,  $\inf_{\mu \in \mathcal{P}^G(\mathcal{Z})} F(\mu) = \inf_{\mu \in \mathcal{P}(\mathcal{Z})} F(\mu)$ . In particular, if F has a unique minimizer over  $\mathcal{P}(\mathcal{Z})$ , it must be WI.

The proof relies on an ad-hoc version of Jensen's inequality. Next, we state that optimizing under **DA** and **FA** is essentially equivalent, and corresponds to optimizing R exclusively over **WI** measures:

Theorem 2 (Equivalence of DA and FA). Under assumption 1, we have:

$$\inf_{\mu \in \mathcal{P}(\mathcal{Z})} R^{DA}(\mu) = \inf_{\mu \in \mathcal{P}^G(\mathcal{Z})} R^{DA}(\mu) = \inf_{\mu \in \mathcal{P}^G(\mathcal{Z})} R(\mu) = \inf_{\mu \in \mathcal{P}^G(\mathcal{Z})} R^{FA}(\mu) = \inf_{\mu \in \mathcal{P}(\mathcal{Z})} R^{FA}(\mu).$$

Note that, on the other hand,  $R^{EA}$  only satisfies:  $\inf_{\mu \in \mathcal{P}(\mathcal{Z})} R^{EA}(\mu) = \inf_{\mu \in \mathcal{P}(\mathcal{E}^G)} R(\mu)$ . In the case of the quadratic loss, Theorem 2 can be made more explicit:

**Corollary 1.** Under Assumption 1, when the loss is quadratic and  $\pi_{\chi}$  is invariant, we have:

$$\inf_{\mu\in\mathcal{P}^G(\mathcal{Z})} R(\mu) = R_* + \inf_{\mu\in\mathcal{P}^G(\mathcal{Z})} \|\Phi_\mu - f_*\|_{L^2(\mathcal{X},\mathcal{Y};\pi_{\mathcal{X}})}^2 = \bar{R}_* + \inf_{\mu\in\mathcal{P}^G(\mathcal{Z})} \|\Phi_\mu - \mathcal{Q}_G \cdot f_*\|_{L^2(\mathcal{X},\mathcal{Y};\pi_{\mathcal{X}})}^2.$$

where  $f_* = \mathbb{E}_{\pi}[Y|X = \cdot]$ , and  $R_*$ ,  $\tilde{R}_*$  are constants only depending on  $\pi$  and  $f_*$ . That is, optimizing under **DA** and **FA** corresponds to approximating the **symmetrized** version of  $f_*$ .

Under equivariance of the data distribution  $\pi$ , the following general result also holds:

**Corollary 2.** Let Assumption 1 hold and suppose  $\pi \in \mathcal{P}^G(\mathcal{X} \times \mathcal{Y})$ . Then, R is invariant and therefore:  $\inf_{\mu \in \mathcal{P}(\mathcal{Z})} R(\mu) = \inf_{\mu \in \mathcal{P}^G(\mathcal{Z})} R(\mu) = \inf_{\mu \in \mathcal{P}(\mathcal{Z})} R^{DA}(\mu) = \inf_{\mu \in \mathcal{P}(\mathcal{Z})} R^{FA}(\mu)$ .

**Remark.** Consequently, equivariant data allow us to globally optimize the population risk by only considering WI measures. It also shows that DA and FA provide no advantage for this optimization.

The same unfortunately is not true for SI measures (answering a question in [22]), as shown by the following result, which constructs a simple example in which  $\mathcal{E}^G$  is trivial:

**Proposition 4.** Even with a finite group G acting orthogonally on  $\mathcal{X} = \mathbb{R}^d$ ,  $\mathcal{Y} = \mathbb{R}$  and  $\mathcal{Z} = \mathbb{R}^{(d+2)}$ ; with  $\pi$  being compactly-supported and equivariant; with  $\ell$  being quadratic; and with  $\sigma_*$  being  $\mathcal{C}^{\infty}$ , bounded and jointly equivariant; we can have:  $\inf_{\mu \in \mathcal{P}(\mathcal{Z})} R(\mu) < \inf_{\nu \in \mathcal{P}(\mathcal{E}^G)} R(\nu)$ .

In fact, even if R is invariant, when  $\mathcal{E}^G$  is *too restrictive*, it might become impossible to globally optimize R over **SI** measures (which would correspond to using  $R^{EA}$  as a proxy for R). Nevertheless, if  $\mathcal{E}^G$  has good *universality* properties, a true **SI** solution to the learning problem can be sought for:

**Proposition 5.** Let Assumption 1 hold,  $\ell$  be quadratic and  $\pi \in \mathcal{P}_2^G(\mathcal{X} \times \mathcal{Y})$ . If  $\mathcal{F}_{\sigma_*}(\mathcal{P}(\mathcal{E}^G))$  is dense in  $L^2_G(\mathcal{X}, \mathcal{Y}; \pi|_{\mathcal{X}}) := \mathcal{Q}_G(L^2(\mathcal{X}, \mathcal{Y}; \pi|_{\mathcal{X}}))$ , then:  $\inf_{\mu \in \mathcal{P}(\mathcal{Z})} R(\mu) = \inf_{\nu \in \mathcal{P}(\mathcal{E}^G)} R(\nu) = R_*$ .

**Remark.** Conditions on  $\mathcal{E}^G$  and  $\sigma_*$  allowing us to guarantee this "restricted" universality on  $L^2_G(\mathcal{X}, \mathcal{Y}; \pi|_{\mathcal{X}})$ , are provided e.g. in [39, 48, 62, 63]. These allow for effectively solving the problem in fewer dimensions, which is key in successful **EA** like CNNs and DeepSets. See SuppMat-C.2.5 for a deeper discussion.

## 3.3 Symmetries and SL training dynamics in the overparametrized regime

We now study the **MFL** of the various training dynamics when  $\mathcal{Z} = \mathbb{R}^D$ . We begin with the general: **Theorem 3.** Let  $F : \mathcal{P}(\mathcal{Z}) \to \overline{\mathbb{R}}$  be an invariant functional such that **WGF**(F) is well defined and has a unique (weak) solution  $(\mu_t)_{t\geq 0}$ . If  $\mu_0 \in \mathcal{P}_2^G(\mathcal{Z})$ , then, for dt-a.e.  $t \geq 0$ ,  $\mu_t \in \mathcal{P}_2^G(\mathcal{Z})$ .

The proof of Theorem 3 relies on  $D_{\mu}F$  being equivariant (in a suitable sense) and  $(M_g\mu_t)_{t\geq 0}$ satisfying also, as a consequence, **WGF**(F) (See SuppMat-C.3 for the details). Note that  $\mu_0 \in \mathcal{P}_2^G(\mathcal{Z})$  is simply verified, e.g. by a standard Gaussian in  $\mathcal{Z}$ . Specializing this result, we get:

**Corollary 3.** Let Assumption 1 and technical assumptions (as in [14]) hold. Then, if R and r are invariant, **WGF**( $R^{\tau,\beta}$ ) starting from  $\mu_0 \in \mathcal{P}_2^G(\mathcal{Z})$  satisfies: for dt-a.e.  $t \ge 0$ ,  $\mu_t \in \mathcal{P}_2^G(\mathcal{Z})$ . If moreover  $\beta > 0$ , each  $\mu_t$  has an invariant density function.

**Remark.** If  $\pi$  is equivariant, R is invariant, and this result is valid for a **freely-trained NN**, without employing SL-techniques. In a way, MFL incorporates these symmetries from infinite SGD iterations.

Theorem 3 (and Corollary 3) can thus be seen as significant generalizations of Proposition 2.1 from [29], which addresses the case of wide 2-layer ReLU networks with a target function that's symmetric w.r.t. a single orthogonal transformation. The fact that strong solutions to  $WGF(R^{\tau,\beta})$  can be sought among invariant functions to reduce the complexity when  $\pi$  is equivariant, was also first hinted in [42]. The natural domain of invariant functions is in fact the quotient space of  $G \bigcirc_M \mathcal{Z}$  (and not  $\mathcal{E}^G$ , which is strictly embedded in it).

Comparing the different training dynamics at the MF level and applying Proposition 2, we also get:

**Theorem 4.** Under assumptions of Corollary 3, if  $\mu_0 \in \mathcal{P}_2^G(\mathcal{Z})$ ,  $WGF(\mathbb{R}^{DA})$  and  $WGF(\mathbb{R}^{FA})$  solutions are equal. If further  $\mathbb{R}$  is invariant, the  $WGF(\mathbb{R})$  solution coincides with them too.

**Remark.** In particular, with equivariant data (i.e. invariant R), training with **DA** or **FA** is essentially the same, at least at the **MF** level, as **using no SL-technique whatsoever**. Hence, a relevant, practical open question, is: how do the convergence rates to the **MFL** compare in all three cases, as  $N \to \infty$ ?

We wil now see that similar results hold for  $\mathcal{P}(\mathcal{E}^G)$  instead of  $\mathcal{P}^G(\mathcal{Z})$ . Notice that the *entropy-regularized* risk forces each  $\mu_t$  to have a density w.r.t.  $\lambda$  in  $\mathcal{Z}$ . Therefore, if  $G \odot_M \mathcal{Z}$  is non-trivial (thus  $\mathcal{E}^G$  is a strict subspace), we always have  $\mu_t \notin \mathcal{P}(\mathcal{E}^G)$ . It seems natural then to *force* the noise in equation (1) to stay within  $\mathcal{E}^G$ ; namely, to consider the following *projected noisy SGD* dynamic:

$$\theta_i^{k+1} = \theta_i^k - s_k^N \left( \nabla_z \sigma_*(X_k, \theta_i^k) \cdot \nabla_1 \ell(\Phi_{\theta^k}^N(X_k), Y_k) + \tau \nabla r(\theta_i^k) \right) + \sqrt{2\beta s_k^N P_{\mathcal{E}^G} \xi_i^k}, \quad (4)$$

which should approximate the **WGF** of the **projected-regularized functional**:  $R_{\mathcal{E}^G}^{\tau,\beta}(\mu) := R(\mu) + \tau \int r d\mu + \beta H_{\lambda_{\mathcal{E}^G}}(\mu)$ , with  $\lambda_{\mathcal{E}^G} := P_{\mathcal{E}^G} \# \lambda$  the Lebesgue Measure over  $\mathcal{E}^G$ . We indeed have :

**Theorem 5.** Let Assumption 1 and technical assumptions on R hold. Suppose that R and r are invariant. Then, if  $\mu_0 \in \mathcal{P}_2(\mathcal{E}^G)$ ,  $(\mu_t)_{t\geq 0}$  solution of  $WGF(R_{\mathcal{E}^G}^{\tau,\beta})$  satisfies  $\forall t \geq 0$ ,  $\mu_t \in \mathcal{P}_2(\mathcal{E}^G)$ .

The proof of this result is based on pathwise properties of the McKean-Vlasov stochastic differential equation associated with the  $WGF(R_{\mathcal{E}^G}^{\tau,\beta})$  (studied e.g. in [19]), see SuppMat-B.2. Notice that  $\mu_0 \in \mathcal{P}_2(\mathcal{E}^G)$  can be obtained e.g. with  $\mu_0 = \delta_{\vec{0}}$ ; though calculating  $\mathcal{E}^G$  is usually hard.

**Remark.** Theorem 5 is significantly stronger than Corollary 3: it implies that, for  $\pi \in \mathcal{P}^G(\mathcal{X} \times \mathcal{Y})$ , the flow will remain in the set of **SI** distributions all throughout its evolution, despite there being **no** constraint whatsoever on the network parameters during training (they can all be freely updated), nor any SL-technique being used. This even holds for  $\beta = 0$ , so the noise projection is not "forcing" the result. This is a large N exclusive phenomenon, as our numerical experiments will show.

**Remark.** Theorem 5 holds too for the invariant functionals  $R^{DA}$ ,  $R^{FA}$  and  $R^{EA}$  in the role of R, even if  $\pi$  is not equivariant. Notably, **DA**, **FA** and **EA** procedures starting on a **SI** distribution, despite being free to involve all parameters, will keep the distribution **SI** all throughout training.

Last, we also have:

**Theorem 6.** Let the conditions for Theorem 5 hold. If  $\mu_0 \in \mathcal{P}_2(\mathcal{E}^G)$ , the WGF( $\mathbb{R}^{FA}$ ), WGF( $\mathbb{R}^{DA}$ ) and WGF( $\mathbb{R}^{EA}$ ) solutions coincide. If  $\mathbb{R}$  is invariant, WGF( $\mathbb{R}$ ) solution coincides with them too.

# 4 Numerical experiments and architecture-discovery heuristic

To empirically illustrate some of our results from the previous section, we consider synthetic data produced in a **teacher-student** setting (see e.g. [12, 14]). Code necessary for replicating the obtained results, as well as a detailed description of our experimental setting, can be sought in the SuppMat.

We study a simple setting with:  $\mathcal{X} = \mathcal{Y} = \mathbb{R}^2$ ,  $\mathcal{Z} = \mathbb{R}^{2 \times 2} \cong \mathbb{R}^4$ , and  $G = C_2$  acting on  $\mathcal{X}$  and  $\mathcal{Y}$  by *permuting the coordinates*; and on  $\mathcal{Z}$  via the natural *intertwining* action (for which  $\mathcal{E}^G$  is explicit). We take the jointly equivariant activation  $\sigma_*(x, z) = \sigma(z \cdot x)$ ,  $\forall (x, z) \in \mathcal{X} \times \mathcal{Z}$  with  $\sigma : \mathbb{R} \to \mathbb{R}$  a sigmoidal function applied pointwise; and consider *normally* distributed features and labels produced from a **teacher model**  $f_* = \Phi_{\theta^*}^{N_*}$  with  $N_* = 5$  particles. Notice that the data will be equivariant only if  $f_*$  is. We try to **mimic** the **teacher** with a **student model**,  $\Phi_{\theta}^N$ , with the same  $\sigma_*$ , but different particles  $\theta \in \mathcal{Z}^N$  that will be trained to minimize the regularized population risk  $R^{\tau,\beta}$  (with quadratic loss and penalization). For this we employ the SGD dynamic given by Equation (1) (or projected, if required, as in Equation (4)), possibly involving **DA**, **FA** or **EA** techniques. We refer to *free training*, with no SL-techniques involved, as **vanilla** training. Each experiment was repeated  $N_r = 10$  times to ensure consistency. Explicit values of the fixed training parameters are found in SuppMat-D.

#### 4.1 Study for varying N

We will demonstrate how properties of  $\mathbf{WGF}(\mathbb{R}^{\tau,\beta})$  stated in Section 3.3 become apparent as N grows. We consider here a teacher with  $\nu_{\theta^*}^{N_*}$  either **arbitrary** or **WI**, and different training schemes performed over  $N_e$  epochs, all initialized with the same particles drawn from given  $\mu_0 \in \mathcal{P}_2(\mathcal{Z})$  that is either **SI** or **WI**. Figure 1 displays the behavior of the student's particle distribution,  $\nu_{N_e}^N$ , after training, in terms of certain "normalized version" of the  $W_2$ -distance, which we call simply Relative Measure Distance, or **RMD** for short. <sup>6</sup> We refer to SuppMat-D for further insights and, additionally: a deeper analysis of the case of  $\nu_{\theta^*}^{N_*}$  being **SI**, comparisons between different techniques and **EA**, and  $L^2$  comparisons between  $\Phi_{\theta}^N$  and both  $f_*$  and  $\mathcal{Q}_G.f_*$  (to illustrate Corollary 1).

We first look at the **SI**-initialized training. Though from [25] we know that (exact) **DA** or **FA** during training will *respect*  $\mathcal{E}^G$  without needing to pass to the **MFL**. This is certainly not true in general for the **vanilla** scheme, where the symmetry is never *explicit* for the model. We notice in Figure 1, however, that, as N grows big, the **SI**-initialized **vanilla** training scheme, under only a **WI** teacher, does **remain SI** throughout training, as predicted in Theorem 5. This is absolutely remarkable, since there is no intuitive reason why the **vanilla** scheme (were parameters can be updated *freely*) *shouldn't escape*  $\mathcal{E}^G$  to better approximate  $f_*$ . For instance, for an **arbitrary teacher** (with the same particles, but *un-symmetrized*) **vanilla** training readily *leaves*  $\mathcal{E}^G$  to better approximate  $f_*$ . Though this isn't a *predicted behaviour* from our theory, it motivates an heuristic we present in the upcoming section. On the other hand, and as expected, both **DA** and **FA** consistently remain within  $\mathcal{E}^G$  almost independently of N, and even if  $f_*$  isn't equivariant. Finally, as N grows bigger, the end-of-training distribution of the **vanilla** scheme *approaches* that of **DA** and **FA** (as expected from Theorem 4).

Regarding the WI-initialized training, unlike the SI case, particles sampled *i.i.d.* from a WI distribution don't necessarily yield a WI empirical distribution  $\nu_0^N$ . On the one hand, this means we require large N to see  $\nu_{N_e}^N$  being (approx.) WI; and on the other hand, it means we have no guarantee that DA and FA will be close unless we look at larger N (where Theorem 4 applies). The second column of Figure 1 precisely shows these behaviours as N grows: both a trend of  $\nu_{N_e}^N$  towards becoming WI, and a clear coincidence between the DA, FA and vanilla schemes (the latter only for equivariant  $f_*$ ).

<sup>&</sup>lt;sup>6</sup>It is roughly equivalent to  $W_2$  when far from the  $\delta_0$  measure, see SuppMat-D for details




Figure 1: **RMD**s, at the end of training, for N = 5, 10, 50, 100, 500, 1000, 5000 and the **vanilla** (**V**), **DA**, **FA** and **EA** schemes. Each column corresponds to a different training initialization (**SI** and **WI** resp.), while each row corresponds to a different teacher distribution (**arbitrary** and **WI** resp.). Each position has two plots: the first displays  $\mathbf{RMD}^2(\nu_{N_e}^N, (\nu_{N_e}^N)^{\mathcal{E}^G})$  or  $\mathbf{RMD}^2(\nu_{N_e}^N, (\nu_{N_e}^N)^G)$  depending on initialization (either **SI** or **WI**), and the second the **RMD** between **DA**, **FA** and **vanilla** schemes.

## 4.2 Heuristic algorithm for architechture discovery

From these experiments, for non-equivariant  $f_*$ , the **SI**-initialized WGF is seen to eventually **escape**  $\mathcal{E}^G$ . In turn, for equivariant  $f_*$ , Figure 2 shows that a training scheme initialized at  $E \subsetneq \mathcal{E}^G$  (i.e.  $\nu_{\theta_0}^N \in \mathcal{P}(E)$ ), eventually **leaves** E, but **stays within**  $\mathcal{E}^G$  (as expected from Theorem 5). These empirical observations hint to an heuristic for **discovering the "good" EAs for the task at hand**.

Assume indeed  $\pi$  equivariant w.r.t. G. We want to find the unknown, largest (i.e. most expressive) subspace of  $\mathcal{Z}$  supporting **SI** measures. We hence consider some (large) N, a shallow NN model with e.g.  $\sigma_*(x, z) = \sigma(z.x)$ , and numerical thresholds  $(\delta_j)_{j \in \mathbb{N}} \subseteq \mathbb{R}_+$ . We define  $E_0 = \{0\} \leq \mathcal{E}^G$  as an initial subspace and initialize  $\nu_{\theta_0}^N = \nu_{\overline{0}}^N \in \mathcal{P}(E_0)$ . Then, we iteratively proceed as follows:

For j = 0, 1, ..., initialize a model at  $\nu_{\theta_0}^N \in \mathcal{P}(E_j)$ , train it for  $N_e$  epochs, and check whether  $\mathbf{RMD}^2(\nu_{N_e}^N, P_{E_j} \# \nu_{N_e}^N) \leq \delta_j$ . If that is the case, the training didn't **escape**  $E_j$ , and one could suppose  $\mathcal{E}^G := E_j$ . Otherwise, it **left**  $E_j$  (so  $\mathcal{E}^G \neq E_j$ ) and one can set e.g.  $E_{j+1} := E_j \oplus v_{E_j}$ , with  $v_{E_j} := \frac{1}{N} \sum_{i=1}^N (\theta_i^{N_e} - P_{E_j}, \theta_i^{N_e})$ . Allegedly, this scheme would eventually leave all strict subspaces to reach the "right"  $\mathcal{E}^G$ . Figure 2 shows indeed this scheme in our simple **teacher-student** example (see SuppMat-D.2 for its details). This idea might have potential for real world applications, yet a larger scale experimental analysis and rigorous theoretical guarantees need to be provided.

## 5 Conclusion

In the light of theoretical guarantees given by the **MF** theory of overparametrized shallow NN, we explored their training dynamics when data is possibly equivariant for some group action and/or SL techniques are employed. We thus described how **DA**, **FA** and **EA** schemes can be understood in the limit of infinite internal units, and studied in that setting the qualitative advantages that can be attired from their use. In this **MFL**, **DA** and **FA** are essentially equivalent in terms of the optimization problem they solve and the trajectory of their associated **WGFs**. Moreover, for equivariant data, *freely*-trained NN, in the **MFL**, obey the same **WGF** as **DA/FA**. They also "respect" symmetries during training, as **WI** and **SI** initializations (corresponding to symmetric parameter distributions and **EA** configurations) are preserved throughout, even if potentially all NN parameters can be updated. We also highlighted the prominent role of **WI** laws in representing equivariant models. We illustrated our results with appropriate numerical experiments, which in turn suggested a data-driven heuristic to find appropriate parameter subspaces for **EAs** in a given task. Providing theoretical guarantees for this heuristic is an interesting problem left for future work. A further relevant question to address, is to quantify and compare the speeds at which all studied training schemes approach the **MFL**, as




Figure 2: Application of our heuristic method on **student** particles. Squares represent the *teacher* particles, dots represent the student particles, and the hyperplane is  $\mathcal{E}^G$ . Each row displays a different viewpoint of the situation (*Row 2* being *parallel* to  $\mathcal{E}^G$ ). Column 1 shows the particles after training on step j = 0. Column 2 shows the j = 1 step initialization on  $E_1 = \langle v_{E_1} \rangle$ . Column 3 shows the particles after training on step j = 1; they **leave**  $E_1$  (see Row 1), but stay in  $\mathcal{E}^G$  (see Row 2).

this would a provide a full comparative picture of their performances. Extending the **MF** analysis of symmetries to NNs with more complex inner structures is another interesting line of work.

## Acknowledgements

JM thanks partial financial support from ANID Magister Fellowship 22231325 and CMM-Basal Grant FB210005, ANID-Chile. JF thanks partial financial support from Fondecyt Regular Grants 1201948/1242001 and CMM-Basal Grant FB210005, ANID-Chile . Both authors thank Roberto Cortez, Daniel Remenik and Felipe Tobar for comments and discussions which motivated part of the numerical experiments developed in the paper, as well as the refinement of some ideas. The authors also thank Daniela Bellott for her assistance in running part of the numerical experiments.

## References

- Luigi Ambrosio, Nicola Gigli, and Giuseppe Savaré. Gradient Flows in Metric Spaces and in the Space of Probability Measures. Lectures in Mathematics ETH Zürich. Birkhäuser, 2. ed edition, 2008. OCLC: 254181287.
- [2] Jimmy Aronsson. Homogeneous vector bundles and g-equivariant convolutional neural networks. *Sampling Theory, Signal Processing, and Data Analysis,* 20(2), jul 2022.
- [3] Dominique Bakry, Ivan Gentil, Michel Ledoux, et al. *Analysis and geometry of Markov diffusion operators*, volume 103. Springer, 2014.
- [4] Andrew Barron. Universal approximation bounds for superpositions of a sigmoidal function. ieee trans. on information theory 39, 930-945. *Information Theory, IEEE Transactions on*, 39:930 945, 06 1993.
- [5] Léon Bottou, Frank E Curtis, and Jorge Nocedal. Optimization methods for large-scale machine learning. *SIAM review*, 60(2):223–311, 2018.
- [6] Michael M Bronstein, Joan Bruna, Taco Cohen, and Petar Veličković. Geometric deep learning: Grids, groups, graphs, geodesics, and gauges. *arXiv preprint arXiv:2104.13478*, 2021.
- [7] Pierre Cardaliaguet. Notes on mean-field games (from P.-L. Lions lectures at Collège de France). 2013. Available at: https://www.ceremade.dauphine.fr/~cardaliaguet/ MFG20130420.pdf.

- [8] R. Carmona and F. Delarue. Probabilistic Theory of Mean Field Games with Applications I: Mean Field FBSDEs, Control, and Games. Probability Theory and Stochastic Modelling. Springer International Publishing, 2018.
- [9] Louis-Pierre Chaintron and Antoine Diez. Propagation of chaos: A review of models, methods and applications. I. models and methods. *Kinetic and Related Models*, 15(6):895–1015, 2022.
- [10] Fan Chen, Yiqing Lin, Zhenjie Ren, and Songbo Wang. Uniform-in-time propagation of chaos for kinetic mean field langevin dynamics. *Electronic Journal of Probability*, 29:1–43, 2024.
- [11] Shuxiao Chen, Edgar Dobriban, and Jane H Lee. A group-theoretic framework for data augmentation. *Journal of Machine Learning Research*, 21(245):1–71, 2020.
- [12] Zhengdao Chen, Grant Rotskoff, Joan Bruna, and Eric Vanden-Eijnden. A dynamical central limit theorem for shallow neural networks. *Advances in Neural Information Processing Systems*, 33:22217–22230, 2020.
- [13] Lénaïc Chizat. Mean-field langevin dynamics : Exponential convergence and annealing. *Transactions on Machine Learning Research*, 2022.
- [14] Lenaic Chizat and Francis Bach. On the global convergence of gradient descent for overparameterized models using optimal transport. *Advances in neural information processing systems*, 31, 2018.
- [15] Taco S Cohen, Mario Geiger, and Maurice Weiler. A general theory of equivariant cnns on homogeneous spaces. Advances in neural information processing systems, 32, 2019.
- [16] Y. Le Cun, B. Boser, J. S. Denker, R. E. Howard, W. Habbard, L. D. Jackel, and D. Henderson. *Handwritten digit recognition with a back-propagation network*, page 396–404. Morgan Kaufmann Publishers Inc., San Francisco, CA, USA, 1990.
- [17] George V. Cybenko. Approximation by superpositions of a sigmoidal function. *Mathematics of Control, Signals and Systems*, 2:303–314, 1989.
- [18] Tri Dao, Albert Gu, Alexander Ratner, Virginia Smith, Chris De Sa, and Christopher Ré. A kernel theory of modern data augmentation. In *International conference on machine learning*, pages 1528–1537. PMLR, 2019.
- [19] Valentin De Bortoli, Alain Durmus, Xavier Fontaine, and Umut Simsekli. Quantitative propagation of chaos for sgd in wide neural networks. *Advances in Neural Information Processing Systems*, 33:278–288, 2020.
- [20] Arnaud Descours, Arnaud Guillin, Manon Michel, and Boris Nectoux. Law of large numbers and central limit theorem for wide two-layer neural networks: the mini-batch and noisy case. *arXiv preprint arXiv:2207.12734*, 2022.
- [21] Paul Dupuis and Richard S Ellis. A weak convergence approach to the theory of large deviations. John Wiley & Sons, 2011.
- [22] Bryn Elesedy and Sheheryar Zaidi. Provably strict generalisation benefit for equivariant models. In *International conference on machine learning*, pages 2959–2969. PMLR, 2021.
- [23] Stewart N Ethier and Thomas G Kurtz. Markov processes: characterization and convergence. John Wiley & Sons, 2009.
- [24] Marc Finzi, Max Welling, and Andrew Gordon Wilson. A practical method for constructing equivariant multilayer perceptrons for arbitrary matrix groups. In *International conference on machine learning*, pages 3318–3328. PMLR, 2021.
- [25] Axel Flinth and Fredrik Ohlsson. Optimization dynamics of equivariant and augmented neural networks. *arXiv preprint arXiv:2303.13458*, 2023.
- [26] Wilfrid Gangbo and Adrian Tudorascu. On differentiability in the wasserstein space and wellposedness for hamilton–jacobi equations. *Journal de Mathématiques Pures et Appliquées*, 125:119–174, 2019.

- [27] Ian Goodfellow, Yoshua Bengio, and Aaron Courville. *Deep Learning*. MIT Press, 2016. http://www.deeplearningbook.org.
- [28] Philipp Grohs and Gitta Kutyniok. *Mathematical aspects of deep learning*. Cambridge University Press, 2022.
- [29] Karl Hajjar and Lénaïc Chizat. On the symmetries in the dynamics of wide two-layer neural networks. *Electron. Res. Arch.*, 31(4):2175–2212, 2023.
- [30] Kurt Hornik, Maxwell Stinchcombe, and Halbert White. Multilayer feedforward networks are universal approximators. *Neural Networks*, 2(5):359–366, 1989.
- [31] Kaitong Hu, Zhenjie Ren, David Siska, and Lukasz Szpruch. Mean-field langevin dynamics and energy landscape of neural networks. In *Annales de l'Institut Henri Poincare (B) Probabilites et statistiques*, volume 57, pages 2043–2065. Institut Henri Poincaré, 2021.
- [32] Kevin H Huang, Peter Orbanz, and Morgane Austern. Quantifying the effects of data augmentation. arXiv preprint arXiv:2202.09134, 2022.
- [33] Ningyuan Huang, Ron Levie, and Soledad Villar. Approximately equivariant graph networks. *Advances in Neural Information Processing Systems*, 36, 2024.
- [34] Risi Kondor and Shubhendu Trivedi. On the generalization of equivariance and convolution in neural networks to the action of compact groups. In *International conference on machine learning*, pages 2747–2755. PMLR, 2018.
- [35] Leon Lang and Maurice Weiler. A wigner-eckart theorem for group equivariant convolution kernels. In *International Conference on Learning Representations*, 2020.
- [36] Zhiyuan Li, Ruosong Wang, Dingli Yu, Simon S Du, Wei Hu, Ruslan Salakhutdinov, and Sanjeev Arora. Enhanced convolutional neural tangent kernels. arXiv preprint arXiv:1911.00809, 2019.
- [37] Pierre Luis Lions. Cours au College de France. 2008.
- [38] Clare Lyle, Mark van der Wilk, Marta Kwiatkowska, Yarin Gal, and Benjamin Bloem-Reddy. On the benefits of invariance in neural networks. *arXiv preprint arXiv:2005.00178*, 2020.
- [39] Haggai Maron, Ethan Fetaya, Nimrod Segol, and Yaron Lipman. On the universality of invariant networks. In *International conference on machine learning*, pages 4363–4371. PMLR, 2019.
- [40] Song Mei, Theodor Misiakiewicz, and Andrea Montanari. Mean-field theory of two-layers neural networks: dimension-free bounds and kernel limit. In *Conference on Learning Theory*, pages 2388–2464. PMLR, 2019.
- [41] Song Mei, Theodor Misiakiewicz, and Andrea Montanari. Learning with invariances in random features and kernel models. In *Conference on Learning Theory*, pages 3351–3418. PMLR, 2021.
- [42] Song Mei, Andrea Montanari, and Phan-Minh Nguyen. A mean field view of the landscape of two-layer neural networks. *Proceedings of the National Academy of Sciences*, 115(33):E7665– E7671, 2018.
- [43] Sylvie Méléard. Asymptotic behaviour of some interacting particle systems; McKean-Vlasov and Boltzmann models, pages 42–95. Springer Berlin Heidelberg, Berlin, Heidelberg, 1996.
- [44] Tomohiro Nishiyama. Convex optimization on functionals of probability densities. *arXiv* preprint arXiv:2002.06488, 2020.
- [45] Atsushi Nitanda, Denny Wu, and Taiji Suzuki. Convex analysis of the mean field langevin dynamics. In *International Conference on Artificial Intelligence and Statistics*, pages 9741–9757. PMLR, 2022.
- [46] F. Otto and C. Villani. Generalization of an inequality by talagrand and links with the logarithmic sobolev inequality. *Journal of Functional Analysis*, 173(2):361–400, 2000.

- [47] Mircea Petrache and Shubhendu Trivedi. Approximation-generalization trade-offs under (approximate) group equivariance. Advances in Neural Information Processing Systems, 36, 2024.
- [48] Siamak Ravanbakhsh. Universal equivariant multilayer perceptrons. In Hal Daumé III and Aarti Singh, editors, *Proceedings of the 37th International Conference on Machine Learning*, volume 119 of *Proceedings of Machine Learning Research*, pages 7996–8006. PMLR, 13–18 Jul 2020.
- [49] Siamak Ravanbakhsh, Jeff Schneider, and Barnabas Poczos. Equivariance through parametersharing. In *International conference on machine learning*, pages 2892–2901. PMLR, 2017.
- [50] Grant Rotskoff and Eric Vanden-Eijnden. Trainability and accuracy of artificial neural networks: An interacting particle system approach. *Communications on Pure and Applied Mathematics*, 75(9):1889–1935, jul 2022.
- [51] Filippo Santambrogio. Optimal transport for applied mathematicians. *Birkäuser, NY*, 55(58-63):94, 2015.
- [52] John Shawe-Taylor. Threshold network learning in the presence of equivalences. In J. Moody, S. Hanson, and R.P. Lippmann, editors, *Advances in Neural Information Processing Systems*, volume 4. Morgan-Kaufmann, 1991.
- [53] John Shawe-Taylor. Sample sizes for threshold networks with equivalences. *Information and Computation*, 118(1):65–72, 1995.
- [54] Justin Sirignano and Konstantinos Spiliopoulos. Mean field analysis of neural networks: A central limit theorem. *Stochastic Processes and their Applications*, 130(3):1820–1852, 2020.
- [55] Justin Sirignano and Konstantinos Spiliopoulos. Mean field analysis of neural networks: A law of large numbers. *SIAM Journal on Applied Mathematics*, 80(2):725–752, 2020.
- [56] Taiji Suzuki, Denny Wu, and Atsushi Nitanda. Convergence of mean-field langevin dynamics: time-space discretization, stochastic gradient, and variance reduction. In *Thirty-seventh Conference on Neural Information Processing Systems*, 2023.
- [57] Alain-Sol Sznitman. Topics in propagation of chaos. In Paul-Louis Hennequin, editor, *Ecole d'Eté de Probabilités de Saint-Flour XIX 1989*, pages 165–251, Berlin, Heidelberg, 1991. Springer Berlin Heidelberg.
- [58] Cédric Villani et al. Optimal transport: old and new, volume 338. Springer, 2009.
- [59] Maurice Weiler and Gabriele Cesa. General e (2)-equivariant steerable CNNs. Advances in neural information processing systems, 32, 2019.
- [60] Max Welling and Yee W Teh. Bayesian learning via stochastic gradient langevin dynamics. In Proceedings of the 28th international conference on machine learning (ICML-11), pages 681–688. Citeseer, 2011.
- [61] Jeffrey Wood and John Shawe-Taylor. Representation theory and invariant neural networks. *Discrete Applied Mathematics*, 69(1):33–60, 1996.
- [62] Dmitry Yarotsky. Universal approximations of invariant maps by neural networks. *Constructive Approximation*, 55(1):407–474, 2022.
- [63] Manzil Zaheer, Satwik Kottur, Siamak Ravanbakhsh, Barnabas Poczos, Russ R Salakhutdinov, and Alexander J Smola. Deep sets. *Advances in neural information processing systems*, 30, 2017.

## A Symmetries in functions, measures, data and shallow models

In this section we state some useful, basic results on the effect of symmetries acting on measures, functions and data, that will be used in the sequel. We also explain how symmetries of interest can be incorporated into the generalized shallow NN setting from Definition 1, complementing also the discussions given in Section 2.1 and Section 2.3 in that regard.

Recall  $\mathcal{X}, \mathcal{Y}$  and  $\mathcal{Z}$  are (separable) Hilbert Spaces and G a compact group with Haar measure  $\lambda_G$ , such that  $G \ominus_{\rho} \mathcal{X}, G \ominus_{\hat{\rho}} \mathcal{Y}$  and  $G \ominus_M \mathcal{Z}$ . Also, let  $\mathcal{E}^G \subseteq \mathcal{Z}$  be the linear subspace of parameters that are *fixed points of the action of G over*  $\mathcal{Z}$ , and  $P_{\mathcal{E}^G}$  the orthogonal projection onto it.

#### A.1 Differentials and integrals of equivariant functions

The following lemma characterizes the differential of *jointly equivariant* functions with respect to the action of some group G. Here we can assume G be a lcsH group w.l.o.g. and consider representations that aren't necessarily orthogonal (we denote them differently to avoid confusion).

**Proposition 6.** Let  $G \circ_{\chi} \mathcal{X}, G \circ_{\tilde{\chi}} \mathcal{Z}, G \circ_{\tilde{\chi}} \mathcal{Y}$  via some representations  $\chi, \tilde{\chi}$  and  $\tilde{\chi}$  respectively (not necessarily orthogonal). Let  $f : \mathcal{X} \times \mathcal{Z} \to \mathcal{Y}$  be jointly *G*-equivariant with respect to these actions (i.e.  $\forall g \in G, \forall x \in \mathcal{X}, \forall z \in \mathcal{Z}, \tilde{\chi}_g.f(x, z) = f(\chi_g.x, \tilde{\chi}_g.z)$ ) and Fréchet-differentiable on its first argument. Then:

$$\forall g \in G, \ \forall x \in \mathcal{X}, \ \forall z \in \mathcal{Z}, \ \ D_x f(\chi_g.x, \tilde{\chi}_g.z) = \check{\chi}_g.D_x f(x, z)\chi_q^{-1}$$

*Proof of Proposition 6.* Indeed, we know that  $\forall z \in \mathbb{Z} D_x f(\cdot, z)$  is the unique function that satisfies,  $\forall \tilde{x} \in \mathcal{X}$ :

$$\lim_{h \to 0} \frac{\|f(\tilde{x}+h,z) - f(\tilde{x}) - D_x f(\tilde{x},z)h\|}{\|h\|} = 0$$

Since we want to prove that  $\forall \tilde{x} \in \mathcal{X}, \forall z \in \mathcal{Z}, \forall g \in G : D_x f(\chi_g.\tilde{x}, \tilde{\chi}_g.z) = \check{\chi}_g D_x f(\tilde{x}, z) \chi_g^{-1}$ , it will be enough to check that:

$$\lim_{h \to 0} \frac{\|f(\chi_g.\tilde{x} + h, \tilde{\chi}_g z) - f(\chi_g.\tilde{x}, \tilde{\chi}_g.z) - \check{\chi}_g D_x f(\tilde{x}, z) \chi_g^{-1} h\|}{\|h\|} = 0$$

which by uniqueness implies the result. Thanks to the joint equivariance of f, we have  $\forall h \neq 0$ :

$$\frac{\|f(\chi_g.\tilde{x}+h,\tilde{\chi}_gz) - f(\chi_g.\tilde{x},\tilde{\chi}_g.z) - \check{\chi}_g D_x f(\tilde{x},z)\chi_g^{-1}h\|}{\|h\|} = \frac{\|f(\chi_g.(\tilde{x}+\chi_g^{-1}.h),\tilde{\chi}_gz) - f(\chi_g.\tilde{x},\tilde{\chi}_g.z) - \check{\chi}_g D_x f(\tilde{x},z)\chi_g^{-1}h\|}{\|h\|} = \frac{\|\check{\chi}_g.f(\tilde{x}+\chi_g^{-1}.h,z) - \check{\chi}_g.f(\tilde{x},z) - \check{\chi}_g D_x f(\tilde{x},z)\chi_g^{-1}h\|}{\|h\|} = \frac{\|\check{\chi}_g.\left[f(\tilde{x}+\chi_g^{-1}.h,z) - f(\tilde{x},z) - D_x f(\tilde{x},z)(\chi_g^{-1}h)\right]\|}{\|\chi_g.\chi_g^{-1}.h\|}$$

Now, recall that for every  $g \in G$ , the operator  $\check{\chi}_g$  is bounded, i.e. it has finite operator norm  $0 < \|\check{\chi}_g\| < \infty$  (non-zero as  $\check{\chi}_g$  is invertible). By defining  $\tilde{h} := \chi_g^{-1} h$ , we have:

$$\frac{\|\check{\chi}_g.\left[f(\tilde{x}+\tilde{h},z)-f(\tilde{x},z)-D_xf(\tilde{x},z)\tilde{h}\right]\|}{\|\chi_g\tilde{h}\|} \leq \frac{\|\check{\chi}_g\|\|f(\tilde{x}+\tilde{h},z)-f(\tilde{x},z)-D_xf(\tilde{x},z)\tilde{h}\|}{\|\chi_g.\tilde{h}\|}$$

Multiplying by  $1 = \frac{\|\chi_g^{-1}\chi_g\tilde{h}\|}{\|\tilde{h}\|}$  the last term is seen to be bounded by

$$\|\check{\chi}_{g}\|\|\chi_{g}^{-1}\| \cdot \frac{\|f(\tilde{x}+\tilde{h},z) - f(\tilde{x},z) - D_{x}f(\tilde{x},z)\tilde{h}\|}{\|\tilde{h}\|}$$

Since  $\chi_g$  and  $\chi_g^{-1}$  are bounded operators, we have that:  $h \to 0 \iff \tilde{h} = \chi_g^{-1}h \to 0$ . Thus

$$\lim_{h \to 0} \frac{\|f(\chi_g.\tilde{x} + h, \tilde{\chi}_g z) - f(\chi_g.\tilde{x}, \tilde{\chi}_g.z) - \tilde{\chi}_g D_x f(\tilde{x}, z) \chi_g^{-1} h\|}{\|h\|} \le \lim_{h \to 0} \|\tilde{\chi}_g\| \|\chi_g^{-1}\| \cdot \frac{\|f(\tilde{x} + \tilde{h}, z) - f(\tilde{x}, z) - D_x f(\tilde{x}, z) \tilde{h}\|}{\|\tilde{h}\|} = 0$$

This, in particular, allows us to characterize the differential of equivariant functions. **Corollary 4.** If  $G \ominus_{\chi} \mathcal{X}$ ,  $G \ominus_{\tilde{\chi}} \mathcal{Y}$ , and  $f : \mathcal{X} \to \mathcal{Y}$  is a *G*-equivariant and Fréchet-differentiable function, then:

$$\forall g \in G, \ \forall x \in \mathcal{X}, \ \ D_x f(\chi_g.x) = \tilde{\chi}_g D_x f(x) \chi_g^T$$

Proof of Corollary 4. Direct.

We can also get some interesting *integral* properties of *jointly* equivariant functions.

**Proposition 7.** Let  $\mathcal{X}, \mathcal{Y}$  and  $\mathcal{Z}$  be (separable) Hilbert Spaces and G be a lcsH group. Let  $G \ominus_{\chi} \mathcal{X}$ ,  $G \ominus_{\tilde{\chi}} \mathcal{Z}, G \ominus_{\tilde{\chi}} \mathcal{Y}$  via some representations  $\chi, \tilde{\chi}$  and  $\check{\chi}$  respectively (not necessarily orthogonal).

Let  $f : \mathcal{X} \times \mathcal{Z} \to \mathcal{Y}$  be a jointly *G*-equivariant function (with respect to these actions). Consider a measure  $\mu \in \mathcal{P}(\mathcal{Z})$  and let f be Bochner integrable on its second argument with respect to  $\mu$ . Then:

$$\forall x \in \mathcal{X}, \ \forall g \in G, \ \check{\chi}_g \langle f(x; \cdot), \mu \rangle = \langle f(\chi_g x; \cdot), \check{\chi}_g \# \mu \rangle$$

*Proof of Proposition 7.* Let  $\mu \in \mathcal{P}(\mathcal{Z})$  and f be as stated. Notice that,  $\forall x \in \mathcal{X}, \forall g \in G$ , we have:

$$\check{\chi}_g \langle f(x, \cdot), \mu \rangle = \check{\chi}_g \int_{\mathcal{Z}} f(x, \theta) d\mu(\theta) = \int_{\mathcal{Z}} \check{\chi}_g f(x, \theta) d\mu(\theta),$$

where we've used the linearity of the Bochner integral under continuous linear operators (as is  $\check{\chi}_g$ ). It follows, from the joint *G*-equivariance of *f* and the definition of the *pushforward measure*, that:

$$\int_{\mathcal{Z}} \check{\chi}_g f(x,\theta) d\mu(\theta) = \int_{\mathcal{Z}} f(\chi_g x, \check{\chi}_g \theta) d\mu(\theta) = \int_{\mathcal{Z}} f(\chi_g x, \tilde{\theta}) d(\check{\chi}_g \# \mu)(\tilde{\theta})$$

We conclude the desired result.

#### A.2 Properties of symmetric measures

Consider again compact G acting orthogonally over the different spaces. Recall that, given  $\mu \in \mathcal{P}(\mathcal{Z})$ , we defined  $\mu^G := \int_G (M_g \# \mu) d\lambda_G$  as its **symmetrized** version and  $\mu^{\mathcal{E}^G} := P_{\mathcal{E}^G} \# \mu$  as its **projected** version. The following two results assumed in the core of the paper are elementary, but we provide their detailed proofs for completeness:

**Lemma 1.** We have the following inclusion:  $\mathcal{P}(\mathcal{E}^G) \subseteq \mathcal{P}^G(\mathcal{Z})$ . Also, for any  $\mu \in \mathcal{P}(\mathcal{Z})$  the following equalities hold  $\forall g \in G: \mu^{\mathcal{E}^G} = (M_g \# \mu)^{\mathcal{E}^G} = (\mu^G)^{\mathcal{E}^G} = (\mu^{\mathcal{E}^G})^G$  and  $(M_g \# \mu)^G = \mu^G$ .

*Proof of Lemma 1.* Let  $\mu \in \mathcal{P}(\mathcal{E}^G)$  (i.e.  $\mu(\mathcal{E}^G) = 1$ ),  $g \in G$  and consider any positive measurable  $f : \mathcal{Z} \to \mathbb{R}$ . We can see:

$$\int_{\mathcal{Z}} f(M_g z) \mu(dz) = \int_{\mathcal{E}^G} f(M_g z) \mu(dz) = \int_{\mathcal{E}^G} f(z) \mu(dz) = \int_{\mathcal{Z}} f(z) \mu(dz).$$

So, that  $\forall g \in G$ ,  $\mu = M_g \# \mu$ , and thus  $\mu \in \mathcal{P}^G(\mathcal{Z})$ . Regarding the equalities, consider  $\mu \in \mathcal{P}(\mathcal{Z})$ and  $A \in \mathcal{B}_{\mathcal{Z}}$  some borel set. Since the  $\lambda_G$  is right-invariant, we have  $\forall g \in G, \forall z \in \mathcal{Z}$ :

$$P_{\mathcal{E}^G}M_g z = \int_G M_h(M_g z) d\lambda_G(h) = \int_G (M_h M_g z) d\lambda_G(h) = \int_G M_{\tilde{h}} z) d\lambda_G(\tilde{h}) = P_{\mathcal{E}^G} z d\lambda_G(h) = \int_G M_{\tilde{h}} z d\lambda_G(h)$$

Then, for  $g \in G$ ,  $M_g^{-1}P_{\mathcal{E}^G}^{-1}(A) = (P_{\mathcal{E}^G}M_g)^{-1}(A) = (P_{\mathcal{E}^G})^{-1}(A)$  and so:

$$(M_g \# \mu)^{\mathcal{E}^G}(A) = \mu(M_g^{-1}P_{\mathcal{E}^G}^{-1}(A)) = \mu(P_{\mathcal{E}^G}^{-1}(A)) = \mu^{\mathcal{E}^G}(A),$$

and:

$$(\mu^G)^{\mathcal{E}^G}(A) = \int_G \mu(M_g^{-1}P_{\mathcal{E}^G}^{-1}(A))d\lambda_G(g) = \int_G \mu(P_{\mathcal{E}^G}^{-1}(A))d\lambda_G(g) = \mu(P_{\mathcal{E}^G}^{-1}(A)) = \mu^{\mathcal{E}^G}(A).$$

On the other hand, since  $\mu^{\mathcal{E}^G} \in \mathcal{P}^{\mathcal{E}^G}(\mathcal{Z}) \subseteq \mathcal{P}^G(\mathcal{Z}), (\cdot)^G$  leaves it unchanged:  $(\mu^{\mathcal{E}^G})^G = \mu^{\mathcal{E}^G}$ . For the last equality, let  $g \in G$  and  $f : \mathcal{Z} \to \mathbb{R}_+$  be measurable. We have:

$$\langle f, (M_g \# \mu)^G \rangle = \int_G \langle f, M_h \# (M_g \# \mu) \rangle d\lambda_G(h) = \int_G \langle f, (M_{\tilde{h}}) \# \mu \rangle d\lambda_G(\tilde{h}) = \langle f, \mu^G \rangle,$$

once again by the right-invariance of  $\lambda_G$ . Namely,  $(M_g \# \mu)^G = \mu^G$ .

**Proposition 8.** For  $\mu \in \mathcal{P}(\mathcal{Z})$ , we have:  $\mu^G \in \mathcal{P}^G(\mathcal{Z})$  and  $\mu^{\mathcal{E}^G} \in \mathcal{P}(\mathcal{E}^G)$ .

*Proof of Proposition 8.* Indeed, let  $h \in G$  and  $B \in \mathcal{B}_{\mathcal{Z}}$  (borel set of  $\mathcal{Z}$ ), using the properties of M and the left-invariance of  $\lambda_G$ , we get that:

$$\mu^{G}(M_{h}^{-1}(B)) = \int_{G} \mu(M_{g}^{-1}(M_{h}^{-1}(B))) d\lambda_{G}(g) = \int_{G} \mu(M_{\tilde{g}}^{-1}(B)) d\lambda_{G}(\tilde{g}) = \mu^{G}(B)$$

So,  $\forall g \in G, \ \mu^G = M_g \# \mu^G$ , implying that  $\mu^G \in \mathcal{P}^G(\mathcal{Z})$ . On the other hand, by definition we have (as the projection is surjective)  $\mu^{\mathcal{E}^G}(\mathcal{E}^G) = \mu(P_{\mathcal{E}^G}^{-1}(\mathcal{E}^G)) = \mu(\mathcal{Z}) = 1$ , so that  $\mu^{\mathcal{E}^G} \in \mathcal{P}(\mathcal{E}^G)$ .  $\Box$ 

**Remark.** It's not hard to notice that, on  $\mathcal{Z} = \mathbb{R}^D$  and with  $\lambda$  being the lebesgue measure, if  $\mu \in \mathcal{P}(\mathcal{Z})$  is such that  $\mu \ll \lambda$  and has density  $u : \mathcal{Z} \to \mathbb{R}_+$ , then:  $\mu^G \in \mathcal{P}^G(\mathcal{Z})$  has density  $u^G := \int_G u \circ M_g d\lambda_G(g)$  w.r.t.  $\lambda$  (whereas  $\mu^{\mathcal{E}^G}$  doesn't have a density w.r.t.  $\lambda$  unless the action is trivial). This follows from the O(D)-invariance of  $\lambda$  and some standard calculations.

Since we will be working on  $\mathcal{P}_2(\mathcal{Z})$  on Section 3.3, it's useful to also notice that:

**Remark.** If  $\mu \in \mathcal{P}_p(\mathcal{Z})$ , then  $\mu^G, \mu^{\mathcal{E}^G} \in \mathcal{P}_p(\mathcal{Z})$ . Indeed, it follows from the fact that  $||M_g|| \le 1$   $\forall g \in G$  (since the representation is orthogonal) and  $||P_{\mathcal{E}^G}|| \le 1$  (since  $P_{\mathcal{E}^G}$  is a projection).

Also, we have that:

**Proposition 9.**  $\mathcal{P}_p(\mathcal{E}^G) := \mathcal{P}(\mathcal{E}^G) \cap \mathcal{P}_p(\mathcal{Z}) \text{ and } \mathcal{P}_p^G(\mathcal{Z}) := \mathcal{P}^G(\mathcal{Z}) \cap \mathcal{P}_p(\mathcal{Z}) \text{ are closed and convex subspaces of } \mathcal{P}_p(\mathcal{Z}) \text{ (under the topology induced by } W_p).$ 

Proof of Proposition 9. Convexity is direct by definition of the involved spaces. For closedness under the Wasserstein topology, recall (see e.g. [1]) that  $W_p(\mu_n, \mu) \xrightarrow[n \to \infty]{} 0$  is equivalent to:  $\mu_n \xrightarrow[n \to \infty]{} \mu$ (weak convergence) and  $\int_{\mathcal{Z}} \|\theta\|^p d\mu_n(\theta) \xrightarrow[n \to \infty]{} \int_{\mathcal{Z}} \|\theta\|^p d\mu(\theta)$ . Since for  $f \in C_b(\mathcal{Z})$ ,  $f \circ M_g$  (for  $g \in G$ ) is continuous and bounded,  $(\mu_n)_{n \in \mathbb{N}} \subseteq \mathcal{P}^G(\mathcal{Z})$  implies  $\forall g \in G$ ,  $M_g \# \mu = \mu$  (namely,  $\mu \in \mathcal{P}^G(\mathcal{Z})$ ). Similarly,  $f \circ P_{\mathcal{E}^G}$  is continuous and bounded, and so if  $(\mu_n)_{n \in \mathbb{N}} \subseteq \mathcal{P}(\mathcal{E}^G)$ , then  $P_{\mathcal{E}^G} \# \mu = \mu$  (i.e.  $\mu \in \mathcal{P}(\mathcal{E}^G)$ ).

## A.3 Properties of equivariant data

We are representing the idea of "data being symmetric" by assuming the data distribution  $\pi$  to be equivariant. This may however seem "unintuitive", as one might imagine equivariant data to be actually given by an invariant r.v. X on  $\mathcal{X}$  and  $Y = f^*(X) + \xi$  for some equivariant function  $f^* : \mathcal{X} \to \mathcal{Y}$  and some centered independent noise  $\xi$ . The following property tells us that, assuming  $\pi \in \mathcal{P}^G(\mathcal{X} \times \mathcal{Y})$  implies such behaviour, but with a  $\xi = \xi_X$ , "conditionally" centered given X. This result will be required in proving Proposition 5.

**Proposition 10.** Let  $\pi \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$  be an equivariant data distribution such that  $\mathbb{E}_{\pi}[||Y||^2] < \infty$ . Then  $f^* = \mathbb{E}_{\pi}[Y|X = \cdot]$  is an equivariant function.

Proof of Proposition 10. Indeed, as  $\mathbb{E}[||Y||^2] < \infty$ , we know the conditional expectation  $\mathbb{E}[Y|X]$ is well defined and there exists a measurable  $f^* : \mathcal{X} \to \mathcal{Y}$  s.t.  $f^*(X) = \mathbb{E}[Y|X]$ . Now, by properties of the conditional expectation): Given any  $h : \mathcal{X} \to \mathbb{R}$  square integrable, we will show that:  $\mathbb{E}_{\pi}[Yh(X)] = \mathbb{E}_{\pi}[\int_{G} \hat{\rho}_{g}^{-1} f^*(\rho_{g}.X) d\lambda_{G}(g)h(X)]$ . Indeed, notice that by *Fubini's theorem*(as  $f^* \in L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})$  and h square integrable), linearity of the integral and G-invariance of  $\pi$ :

$$\mathbb{E}_{\pi} \left[ (\mathcal{Q}_G \cdot f^*)(X)h(X) \right] = \mathbb{E}_{\pi} \left[ \int_G \hat{\rho}_g^{-1} \cdot f^*(\rho_g \cdot X) d\lambda_G(g)h(X) \right]$$
$$= \int_G \hat{\rho}_g^{-1} \cdot \mathbb{E}_{\pi} [f^*(X)h(\rho_g \cdot f^{-1} \cdot X)] d\lambda_G(g)$$
$$= \int_G \mathbb{E}_{\pi} [\hat{\rho}_g^{-1} \cdot Yh(\rho_g \cdot f^{-1} \cdot X)] d\lambda_G(g)$$
$$= \int_G \mathbb{E}_{\pi} [Yh(X)] d\lambda_G(g) = \mathbb{E}_{\pi} [Yh(X)]$$

By uniqueness of the conditional expectation, we know therefore that  $f^*(X) \stackrel{a.s.}{=} \int_G \hat{\rho}_g^{-1} f^*(\rho_g X) \lambda_G(g)$ . In particular,  $\pi_{\mathcal{X}}$ -a.e.  $f^* = \mathcal{Q}_G(f^*)$ , making  $f^*$  *G*-equivariant.

We notice that Proposition 10 can also be used to recover a celebrated result from [22] (later generalized by [33]), in the general setting where data symmetry is encoded by the condition that  $\pi \in \mathcal{P}_2(\mathcal{X} \times \mathcal{Y})$ . Define the **symmetrization gap** of a learning problem with quadratic loss as:

$$\Delta(f, \mathcal{Q}_G f) := \mathbb{E}_{\pi}[\|Y - f(X)\|_{\mathcal{Y}}^2] - \mathbb{E}_{\pi}[\|Y - (\mathcal{Q}_G f)(X)\|_{\mathcal{Y}}^2]$$

The following extension of mentioned statements from [22, 33] is not needed for our results, but we provide a proof of it in SuppMat-E, in view of its potential, independent interest:

**Lemma 2** (Symmetrization Gap Characterization). Consider the quadratic loss and  $\pi \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$  such that  $\mathbb{E}_{\pi}[||Y||^2] < \infty$ . Also, assume that  $\pi|_{\mathcal{X}}$  is *G*-invariant, but  $\pi$  is only *H*-invariant with respect to some  $H \leq G$  (closed). Then, the generalization gap satisfies:

$$\Delta(f, \mathcal{Q}_G f) = -2\langle f^*, f_G^{\perp} \rangle_{L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})} + \|f_G^{\perp}\|_{L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})}^2$$

where  $f^*(x) = \mathbb{E}_{\pi}[Y|X=x]$  is the conditional expectation function, and  $f_G^{\perp} := f - \mathcal{Q}_G f$ .

In particular, if  $\pi$  is G-invariant as well, we get  $\Delta(f, \mathcal{Q}_G f) = \|f_G^{\perp}\|_{L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})}^2$ 

#### A.4 Shallow models with symmetries

The class of models we have introduced in Definition 1 allows for taking an arbitrary, common parameter space Z for all hidden units, as well as an arbitrary function  $\sigma_* : \mathcal{X} \times Z \to \mathcal{Y}$ . As noted in the description of **EA**s in Section 2.3, by requiring only that  $\sigma_*$  is jointly-equivariant, we moreover ensure that  $G \oplus_M Z$  is *properly related* to the actions  $G \oplus_\rho \mathcal{X}$  and  $G \oplus_\rho \mathcal{Y}$ . This might seem like an abstract idea, but it is heavily inspired from what happens in the concrete examples of *single-hidden-layer* shallow NN setting.

Indeed, recall the finite-dimensional setting of single-hidden-layer neural networks. We considered  $\mathcal{X} = \mathbb{R}^d$ ,  $\mathcal{Y} = \mathbb{R}^c$  and  $\mathcal{Z} = \mathbb{R}^{c \times b} \times \mathbb{R}^{d \times b} \times \mathbb{R}^b$ , and defined, for  $z = (W, A, B) \in \mathcal{Z}$  and  $\sigma : \mathbb{R}^b \to \mathbb{R}^b$ ,  $\sigma_*(x, z) := W\sigma(A^Tx + B)$ . This allows us to express a *shallow* NN with N hidden units, of parameters  $\theta = (\theta_i)_{i=1}^N \in \mathcal{Z}^N$ , with  $\theta_i = (W_i, A_i, B_i)$ , as the function  $\Phi^N_{\theta} : \mathcal{X} \to \mathcal{Y}$  given by:

$$\forall x \in \mathcal{X}, \ \Phi^N_{\theta}(x) := \frac{1}{N} \sum_{i=1}^N W_i \sigma(A_i^T . x + B_i) = \frac{1}{N} \overline{W} . \sigma(\overline{A}^T . x + \overline{B}),$$

where we write the expression by blocks, considering

$$\overline{W} = (W_1, \dots, W_N) \in \mathbb{R}^{c \times bN}, \ \overline{A} = (A_1, \dots, A_N) \in \mathbb{R}^{d \times bN}, \text{ and } \overline{B} = \begin{pmatrix} B_1 \\ \vdots \\ B_N \end{pmatrix} \in \mathbb{R}^{bN}.$$

This corresponds exactly to the usual *single-hidden-layer* setting (see e.g. [20, 42, 50, 55]), only that we allow for the structure to involve the use of *block matrices*. This allows us to translate many relevant **EAs**, such as CNNs (see [16]) or DeepSets (see [63]) into the *shallow NN* framework that we propose.

We now also consider a *G*-action on the *intermediate layer*,  $G \,_{\eta} \mathbb{R}^{b}$  (so that  $G \,_{\eta \otimes \mathrm{Id}_{N}} (\mathbb{R}^{b})^{N}$ ). We could ask, for instance, for  $\sigma : \mathbb{R}^{b} \to \mathbb{R}^{b}$  to be equivariant with respect to this action. This is relatively reasonable, since for actions via *permutations*, it's enough to consider a  $\sigma$  that is the *pointwise* application of a scalar function (see [25]).<sup>7</sup>

With this in place, we consider the *natural intertwinning* action of G on Z, which is given by:  $M_g.z = (\hat{\rho}_g.W\eta_g^T, \rho_g.A\eta_g^T, \eta_g.B)$ . This is exactly the action under which the fixed points (i.e.  $\mathcal{E}^G$ ) correspond exactly to **EAs** in the traditional sense for this architechture (i.e. each layer is an equivariant function). Also, having an equivariant  $\sigma : \mathbb{R}^b \to \mathbb{R}^b$ , ensures that  $\sigma_*$  is *jointly equivariant*, and so all other results from our work can be applied.

This readily generalizes to the multilayer case. Namely, if  $\sigma_*$  encodes a Multi-Layer Perceptron (MLP) with multiple hidden layers whose parameters live in some linear space  $\mathcal{Z}$ , we can define  $G \odot_M \mathcal{Z}$  as the *intertwining* action between each successive layer (similar to the previous example).  $\sigma_*$  can be made jointly-equivariant by making every activation function on each hidden layer equivariant (see [25]). Then,  $\mathcal{E}^G$  corresponds exactly to the parameters that make the entire MLP an equivariant architechture (in the sense that every layer is an equivariant function). With this,  $\Phi^N_{\theta}$  is an ensemble of N such MLPs trained in parallel, to which our results would also apply.

# **B** Further elements from the MF theory of shallow neural networks

In this section we study several theoretical notions required in the MF approach to overparametrized shallow NN. For the purpose of our results, Subsections B.1 and B.2 are the most relevant ones, as we establish therein some useful properties or formula that will be explicitly required. Subsections B.3 and B.4 review some results and recent literature regarding well-posedness and long-time convergence of WGFs, which are relevant to the MF interpretation of the training dynamics of shallow models.

#### **B.1** Linear functional derivatives and intrinsic derivatives

Let  $\mathcal{X}, \mathcal{Y}$  and  $\mathcal{Z}$  be separable Hilbert spaces. Recall the definitions of the linear functional derivative and intrinsic derivatives:

**Definition 8** (Linear Functional Derivative (First Variation)). For a functional  $F : \mathcal{P}_2(\mathcal{Z}) \to \mathbb{R}$ , its linear functional derivative (lfd) is a function:  $\frac{\partial F}{\partial \mu} : \mathcal{P}_2(\mathcal{Z}) \times \mathcal{Z} \to \mathbb{R}$  such that  $\forall \mu, \nu \in \mathcal{P}_2(\mathcal{Z})$ :

$$\lim_{h \to 0} \frac{F((1-h)\mu + h\nu) - F(\mu)}{h} = \int_{\mathcal{Z}} \frac{\partial F}{\partial \mu}(\mu, z) d(\nu - \mu)(z), \quad and \quad \int_{\mathcal{Z}} \frac{\partial F}{\partial \mu}(\mu, z) d\mu(z) = 0$$

The function  $F': \mu \in \mathcal{P}_2(\mathcal{Z}) \mapsto \frac{\partial F}{\partial \mu}(\mu, \cdot)$  is also known as the first variation of R at  $\mu$ .

**Definition 9** (Intrinsic Derivative). Whenever  $\frac{\partial F}{\partial \mu} : \mathcal{P}_2(\mathcal{Z}) \times \mathcal{Z} \to \mathbb{R}$  exists and is differentiable on its second argument, the intrinsic derivative of F is defined as:  $D_{\mu}F(\mu, z) = \nabla_z \left(\frac{\partial F}{\partial \mu}(\mu, z)\right)$ .

**Example.** To better illustrate the notion of the linear functional derivative and the intrinsic derivative, consider the following usual examples:

1. An important example is that of the **KL Divergence**. Let  $\mu \ll \nu$  and  $\frac{d\mu}{d\nu}$  be the corresponding Radon-Nykodym derivative; the KL Divergence between  $\mu$  and  $\nu$  is defined as:  $D(\mu||\nu) := \int \log(\frac{d\mu}{d\nu}(z))d\mu(z)$ . Fixing  $\nu \in \mathcal{P}(\mathcal{Z})$  and working with  $F(\mu) = D(\mu||\nu)$ , we have that (modulo a constant that doesn't depend on z, see [44]):

$$\frac{\partial F}{\partial \mu}(\mu, z) = \log\left(\frac{d\mu}{d\nu}(z)\right) + 1 \text{ and } D_{\mu}F(\mu, z) = \frac{1}{\frac{d\mu}{d\nu}(z)}\nabla_{z}\frac{d\mu}{d\nu}(z)$$

<sup>&</sup>lt;sup>7</sup>For infinite groups this might be more challenging, and we leave its exploration as future work.

2. Whenever  $F(\mu) := \int_{\mathcal{Z}} \phi(z) d\mu(z)$  for some bounded continuously differentiable function  $\phi : \mathcal{Z} \to \mathbb{R}$ , it is well known that :

$$\frac{\partial F}{\partial \mu}(\mu,z) = \phi(z) - \int \phi d\mu \text{ and } D_{\mu}F(\mu,z) = \nabla_{z}\phi(z)$$

The most relevant example, in our case, regards the linear functional and intrinsic derivatives of the **population risk** functional,  $R(\mu) = \mathbb{E}_{\pi}[\ell(\Phi_{\mu}(X), Y)]$ . We can consider the general setting presented in [14], where some Hilbert Space  $\mathcal{H}$  is considered, and it is assumed that  $F : \mathcal{P}(\mathcal{Z}) \to \mathbb{R}$ can be written as  $F(\mu) = L(\langle \Phi, \mu \rangle)$ , where  $\Phi : \mathcal{Z} \to \mathcal{H}$  is a parametrization of elements in  $\mathcal{H}$ ;  $L : \mathcal{H} \to \mathbb{R}$  is some *loss* functional, and the integral  $\langle \Phi, \mu \rangle$  is a **Bochner integral** on  $\mathcal{H}$ . This generalizes the *shallow NN* setting, as one might consider  $\mathcal{H} = L^2(\mathcal{X}, \mathcal{Y}, \pi_{\mathcal{X}}), L : \mathcal{H} \to \mathbb{R}$  given by  $L(f) = \mathbb{E}_{\pi}[\ell(f(X), Y)]$  and  $\Phi : \mathcal{Z} \to \mathcal{H}$  defined as  $\forall z \in \mathcal{Z}, \quad \Phi(z) = \sigma_*(\cdot; z)$ ; so that  $R(\mu) = L(\langle \Phi, \mu \rangle)$ . In this setting, we can prove the following result:<sup>8</sup>

**Proposition 11.** Let  $\mathcal{H}$  be a separable Hilbert Space and  $F(\mu) := L(\langle \Phi, \mu \rangle)$ , for some function that's Gateaux-differentiable  $L : \mathcal{H} \to \mathbb{R}$  on every direction and of continuous differential; and  $\Phi : \mathcal{Z} \to \mathcal{H}$  such that  $\forall \mu \in \mathcal{P}(\mathcal{Z}), ||\langle \Phi, \mu \rangle||_{\mathcal{H}} < \infty$ .

Then  $\forall z \in \mathcal{Z}, \ \forall \mu \in \mathcal{P}(\mathcal{Z})$ :

$$\frac{\partial F}{\partial \mu}(\mu, z) = D_h L(\langle \Phi, \mu \rangle)(\Phi(z)) = \langle \nabla_h L(\langle \Phi, \mu \rangle), \Phi(z) \rangle_{\mathcal{H}} - C_{F,\mu}$$
$$D_\mu F(\mu, z) = (D_h L(\langle \Phi, \mu \rangle)(D_z \Phi(z)))^* = \nabla_z \Phi(z)(\nabla_h L(\langle \Phi, \mu \rangle)).$$

Here,  $C_{F,\mu} := \langle \nabla_h L(\langle \Phi, \mu \rangle), \langle \Phi, \mu \rangle \rangle_{\mathcal{H}}$  is exactly the constant needed to avoid ambiguity in the definition;  $(\cdot)^*$  denotes the adjoint operator and, in particular,  $\nabla_z \Phi(z) = (D_z \Phi(z))^* : \mathcal{H} \to \mathcal{Z}$ . When  $\mathcal{Z} = \mathbb{R}^D$  this corresponds to the usual definition of the gradient.

*Proof of Proposition 11.* We know that,  $\forall \mu, \nu \in \mathcal{P}(\mathcal{Z}), h \in [0, 1]$ :

$$\frac{F((1-h)\mu + h\nu) - F(\mu)}{h} = \frac{L(\langle \Phi, (1-h)\mu + h\nu \rangle) - L(\langle \Phi, \mu \rangle)}{h}$$
$$= \frac{L(\langle \Phi, \mu \rangle + h\langle \Phi, \nu - \mu \rangle) - L(\langle \Phi, \mu \rangle)}{h}$$

Let's denote by  $q_{\mu} := \langle \Phi, \mu \rangle$  (analogously  $q_{\nu-\mu} := \langle \Phi, \nu - \mu \rangle$ ) and  $s_{\mu,\nu} := hq_{\nu-\mu}$ , so we can write:

$$\frac{F((1-h)\mu + h\nu) - F(\mu)}{h} = \frac{L(q_{\mu} + s_{\mu,\nu}) - L(q_{\mu})}{h}.$$

As L is Gateaux differentiable, we have the following first order Taylor expansion  $\forall x, s \in \mathcal{H}, \forall t \in \mathbb{R}$ :

$$L(x + ts) = L(x) + t D_h L(x) \cdot s + o(|t|||s||),$$

which allows us to write:

$$\frac{F((1-h)\mu + h\nu) - F(\mu)}{h} = \frac{L(q_{\mu} + h q_{\nu-\mu}) - L(q_{\mu})}{h} = \frac{h \cdot D_h L(q_{\mu}) \cdot q_{\nu-\mu} + o(|h| ||q_{\nu-\mu}||)}{h}$$

As  $||q_{\nu-\mu}|| < \infty$  by hypothesis, we can say that:  $o(|h|||q_{\nu-\mu}||) = o(h)$ . Therefore, taking the limit with  $h \to 0$ , we get that:

$$\lim_{h \to 0} \frac{F((1-h)\mu + h\nu) - F(\mu)}{h} = D_h L(q_\mu) \cdot q_{\nu-\mu} + \lim_{h \to 0} \frac{o(h)}{h} = D_h L(q_\mu) \cdot q_{\nu-\mu}$$

Now, developping this last term (using, for instance, the linearity of the Bochner integral, as we know  $D_h L(x, \cdot).(\cdot)$  to be linear and bounded), we get that:

$$D_h L(q_\mu).q_{\nu-\mu} = D_h L(q_\mu).\langle \Phi, \nu - \mu \rangle = \langle D_h L(q_\mu)(\Phi), \nu - \mu \rangle,$$

<sup>&</sup>lt;sup>8</sup>Where the **gradient**  $\nabla_x f(x)$  is the unique vector in  $\mathcal{H}$  representing the action of  $D_x f(x) : \mathcal{H} \to \mathbb{R}$ 

and so by definition of the gradient of L:

$$\lim_{h \to 0} \frac{F((1-h)\mu + h\nu) - F(\mu)}{h} = \int_{\mathcal{Z}} \langle \nabla_h L(\langle \Phi, \mu \rangle), \Phi(z) \rangle_{\mathcal{H}} d(\nu - \mu)(z).$$

From here we deduce that:

$$\frac{\partial F}{\partial \mu}(\mu, z) = \langle \nabla_h L(\langle \Phi, \mu \rangle), \Phi(z) \rangle_{\mathcal{H}} - C_{F, \mu},$$

where  $C_{F,\mu}$  is a fixed constant, given by:

$$C_{F,\mu} = \int_{\mathcal{Z}} \frac{\partial F}{\partial \mu}(\mu, z) d\mu(z) = \int_{\mathcal{Z}} \langle \nabla_h L(\langle \Phi, \mu \rangle), \Phi(z) \rangle_{\mathcal{H}} d(\mu)(z) = \langle \nabla_h L(\langle \Phi, \mu \rangle), \langle \Phi, \mu \rangle \rangle_{\mathcal{H}}$$

On the other hand, for the intrinsic derivative, notice that  $D_z(\frac{\partial F}{\partial \mu}(\mu, z)): \mathbb{Z} \to \mathbb{R}$  is a bounded linear functional over  $\mathbb{Z}$ , so (by Riesz Representation)  $\exists D_\mu F(\mu, z) := \nabla_z(\frac{\partial F}{\partial \mu}(\mu, z)) \in \mathbb{Z}$  such that:

$$\forall z \in \mathcal{Z}, \ \left\langle \nabla_z \left( \frac{\partial F}{\partial \mu}(\mu, z) \right), z \right\rangle_{\mathcal{Z}} = D_z \left( \frac{\partial F}{\partial \mu}(\mu, z) \right)(z)$$

However, we can develop the RHS, and as the constant  $C_{F,\mu}$  doesn't depend on z, we get that:

$$D_z\left(\frac{\partial F}{\partial \mu}(\mu, z)\right)(z) = D_z\left(\langle \nabla_h L(\langle \Phi, \mu \rangle), \Phi(z) \rangle_{\mathcal{H}}\right)(z)$$

Now, by the chain rule and the definition of the *adjoint operator* of  $D_z \Phi(z)$ :

$$D_{z}\left(\frac{\partial F}{\partial \mu}(\mu, z)\right)(z) = \left\langle \nabla_{h}L(\langle \Phi, \mu \rangle), D_{z}\Phi(z)(z) \right\rangle_{\mathcal{H}} = \left\langle \left(D_{z}\Phi(z)\right)^{*}\left(\nabla_{h}L(\langle \Phi, \mu \rangle)\right), z \right\rangle_{\mathcal{Z}}$$

So, as they coincide for every  $z \in \mathcal{Z}$ , we conclude that:

$$D_{\mu}F(\mu, z) = (D_{z}\Phi(z))^{*} \left(\nabla_{h}L(\langle \Phi, \mu \rangle)\right)$$

Proposition 11 applies directly to our population risk functional  $R(\mu) := \mathbb{E}_{\pi} \left[ \ell(\Phi_{\mu}(X), Y) \right]$ , by considering:

- The Hilbert space:  $\mathcal{H} = L^2(\mathcal{X}, \mathcal{Y}, \pi_{\mathcal{X}})$
- $L : \mathcal{H} \to \mathbb{R}$  given by  $L(f) = \mathbb{E}_{\pi}[\ell(f(X), Y)]$ , which is Gateaux-differentiable on every direction in  $\mathcal{H}$  if we assume  $\ell : \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$  to be continuously differentiable on its first argument, with  $\nabla_1 \ell$  linearly growing. The differential (which is continuous) can be explicitly computed to be:

$$D_h L(f)(h) = \mathbb{E}_{\pi} \left| \left\langle \nabla_1 \ell \left( \left( f(X), Y \right), h(X) \right\rangle_{\mathcal{Y}} \right| \right|$$

•  $\Phi : \mathcal{Z} \to \mathcal{H}$  defined as  $\forall z \in \mathcal{Z}, \quad \Phi(z) = \sigma_*(\cdot; z)$ , which satisfies  $\forall \mu \in \mathcal{P}(\mathcal{Z}), \|\langle \Phi, \mu \rangle\|_{\mathcal{H}} < \infty$  under the assumption of  $\sigma_*$  being **bounded** and continuous.

**Corollary 5.** We can explicitly compute the linear functional derivative and the intrinsic derivative for the learning problem's population risk:

$$\begin{split} &\frac{\partial R}{\partial \mu}(\mu,z) = \mathbb{E}_{\pi}\left[ \left\langle \nabla_{1}\ell\left( \left\langle \sigma_{*}(X;\cdot),\mu\right\rangle,Y\right),\sigma_{*}(X;z)\right\rangle_{\mathcal{Y}}\right] + (\textit{constant not depending on }z) \\ &D_{\mu}R(\mu,z) = \mathbb{E}_{\pi}\left[ \nabla_{z}\sigma_{*}(X;z).\nabla_{1}\ell(\left\langle \sigma_{*}(X;\cdot),\mu\right\rangle,Y)\right] \end{split}$$

Beyond this particular example, the linear functional derivative and the intrinsic derivative behave well when the underlying functional is invariant, as shown by the following result:

**Proposition 12.** Let  $F : \mathcal{P}(\mathcal{Z}) \longrightarrow \mathbb{R}$  be invariant and of class  $\mathcal{C}^1$ . Then:  $\forall z \in \mathcal{Z}, \forall \mu \in \mathcal{P}(\mathcal{Z}), \forall g \in G$ :

$$\frac{\partial F}{\partial \mu}(M_g \# \mu, M_g.z) = \frac{\partial F}{\partial \mu}(\mu, z) \quad and \quad D_{\mu}F(M_g \# \mu, M_g.z) = M_g.D_{\mu}F(\mu, z)$$

*i.e.*  $\frac{\partial F}{\partial \mu}$  *is jointly invariant and*  $D_{\mu}F$  *jointly equivariant.* 

*Proof of Proposition 12.* To prove this, recall that the linear functional derivative of F is the only function  $\frac{\partial F}{\partial \mu} : \mathcal{P}(\mathcal{Z}) \times \mathcal{Z} \to \mathbb{R}$  satisfying  $\forall \mu, \nu \in \mathcal{P}(\mathcal{Z})$ :

$$\lim_{h\to 0}\frac{F((1-h)\mu+h\nu)-F(\mu)}{h}=\int_{\mathcal{Z}}\frac{\partial F}{\partial \mu}(\mu,z)d(\nu-\mu)(z) \ \text{ and } \ \int_{\mathcal{Z}}\frac{\partial F}{\partial \mu}(\mu,z)d\mu(z)=0.$$

In particular, as F is G-invariant (and  $M_g$  linear), we can write  $\forall \mu, \nu \in \mathcal{P}(\mathcal{Z}), \forall h \neq 0 \text{ and } g \in G$ :

$$\frac{F((1-h)\mu + h\nu) - F(\mu)}{h} = \frac{F((1-h)(M_g \# \mu) + h(M_g \# \nu)) - F(M_g \# \mu)}{h}.$$

Taking the limit as  $h \rightarrow 0$  on both sides, we get:

$$\lim_{h \to 0} \frac{F((1-h)\mu + h\nu) - F(\mu)}{h} = \int_{\mathcal{Z}} \frac{\partial F}{\partial \mu} (M_g \# \mu, z) d(M_g \# \nu - M_g \# \mu)(z)$$
$$= \int_{\mathcal{Z}} \frac{\partial F}{\partial \mu} (M_g \# \mu, M_g. z) d(\nu - \mu)(z),$$

and also:

$$\int_{\mathcal{Z}} \frac{\partial F}{\partial \mu} (M_g \# \mu, M_g. z) d\mu(z) = \int_{\mathcal{Z}} \frac{\partial F}{\partial \mu} (M_g \# \mu, z) dM_g \# \mu(z) = 0.$$

So, by uniqueness, we get  $\forall g \in G$ :

$$\frac{\partial F}{\partial \mu}(\mu,z) = \frac{\partial F}{\partial \mu}(M_g \# \mu, M_g.z),$$

and, from proposition 6 (since  $\frac{\partial F}{\partial \mu}$  is jointly invariant), we get  $\forall g \in G$ :

$$D_{\mu}F(M_{g}\#\mu, M_{g}.z) = \nabla_{z} \left(\frac{\partial F}{\partial \mu}\right) (M_{g}.\mu, M_{g}.z) = M_{g}.\nabla_{z} \left(\frac{\partial F}{\partial \mu}\right) (\mu, z) = M_{g}.D_{\mu}F(\mu, z)$$

## B.2 Expression for the WGF of the regularized population risk

In the case of the **regularized** population risk functional  $R^{\tau,\beta} : \mathcal{P}(\mathcal{Z}) \to \mathbb{R}$ , we can explicitly write its intrinsic derivative. Consider a slightly more *general* functional, denoted by  $R_{\nu}^{\tau,\beta}$ , where the entropy is calculated against a **Gibbs measure**  $\nu \ll \lambda$  such that  $\nu(dz) = e^{-U(z)}\lambda(dz)$  for some function  $U : \mathcal{Z} \to \mathbb{R}$  (as in [31]). We have  $\forall \mu \in \mathcal{P}(\mathcal{Z})$  s.t.  $\mu \ll \nu, \forall z \in \mathcal{Z}$ :

$$D_{\mu}R_{\nu}^{\tau,\beta}(\mu,z) = D_{\mu}R(\mu,z) + \tau\nabla_{z}r(z) + \beta\nabla_{z}U(z) + \beta\left(\frac{1}{\mu(z)}\nabla_{z}\mu(z)\right),$$

so that **WGF** $(R_{\nu}^{\tau,\beta})$  as in definition 4 reads:

$$\partial_t \mu_t = \varsigma(t) \left[ \operatorname{div} \left( D_\mu R_\nu^{\tau,\beta}(\mu_t, \cdot) \, \mu_t \right) \right] = \varsigma(t) \left[ \operatorname{div} \left( \left( D_\mu R(\mu_t, \cdot) + \tau \nabla_z r + \beta \nabla_z U \right) \, \mu_t \right) + \beta \Delta \mu_t \right].$$

We recover the expression for  $WGF(R^{\tau,\beta})$  in Equation (3) by considering  $U \equiv 0$ :

$$\partial_t \mu_t = \varsigma(t) \left[ \operatorname{div} \left( \left( D_\mu R(\mu_t, \cdot) + \tau \nabla_z r \right) \mu_t \right) + \beta \Delta \mu_t \right] \right]$$

We can see that Equation (3) corresponds to a Fokker-Planck equation, which can be rewritten into a **non-linear SDE** system representing the behaviour of the *type* parameter: the **McKean-Vlasov SDE** [57, 43, 9] (also known as *the Mean Field Langevin Dynamics (MFLD*) in the NN literature). In the case of  $R^{\tau,\beta}$  it reads:

$$dZ_t = \varsigma(t) \left[ -\left( D_\mu R(\mu_t, Z_t) + \tau \nabla_\theta r(Z_t) \right) dt + \sqrt{2\beta} dB_t \right] \text{ with } \mu_t = \mathbf{Law}(Z_t),$$
 (5)

Wwere  $(B_t)_{t\geq 0}$  is a *D*-dimensional standard Brownian Motion. Both formulations are equivalent (see [57], Theorem 1.1 for a reference result), but this SDE will prove useful to establish some of the relevant results of the paper.

#### **B.3** Global convergence in the regularized case

For the example we just presented of the entropy-regularized population risk, multiple authors (see [10, 13, 31, 45, 56] among many others) have studied the properties of  $WGF(R^{\tau,\beta})$ , particularly, the global convergence results that can be obtained. For instance, consider the following results from [31] (where we look at  $R_{\nu}^{\tau,\beta}$  for generality). First, define:

**Definition 10.** We say that a functional  $R : \mathcal{P}_p(\mathcal{Z}) \to \mathbb{R}$  is of class  $\mathcal{C}^1$  if  $\frac{\partial R}{\partial \mu}(\mu, \cdot)$  is well defined and bounded for every  $\mu \in \mathcal{P}_p(\mathcal{Z})$ , and the function  $(\mu, z) \in \mathcal{P}_p(\mathcal{Z}) \times \mathcal{Z} \mapsto \frac{\partial R}{\partial \mu}(\mu, z)$  is continuous.

Now, from [13, 31], we get the following key result. We include the proof for completeness:

**Lemma 3** (as in [31, 13]). Assume that  $R : \mathcal{P}_p(\mathcal{Z}) \to \mathbb{R}$  is convex and of class  $\mathcal{C}^1$ . Then, for any  $\mu, \mu' \in P_p(\mathcal{Z})$ , we have:

$$R(\mu') - R(\mu) \ge \int_{\mathcal{Z}} \frac{\partial R}{\partial \mu}(\mu, z) d(\mu' - \mu)(z)$$

*Proof of Lemma 3 (from [31]).* Define  $\mu^{\epsilon} := (1 - \epsilon)\mu + \epsilon \mu'$ . Since R is convex, we have

$$\epsilon \left( R(\mu') - R(\mu) \right) \ge R(\mu^{\epsilon}) - R(\mu) = \int_0^{\epsilon} \int_{\mathcal{Z}} \frac{\partial R}{\partial \mu} (\mu^s, z) d(\mu' - \mu) (dz) \, ds.$$

Since the map  $s \in [0, 1] \mapsto \mu^s$  is continuous, it is of **compact image** (denoted  $[\mu, \mu']$ ). In particular, as  $\frac{\partial R}{\partial \mu}$  is continuous and bounded on its second argument, we get that, it is bounded on  $[\mu, \mu'] \times \mathcal{Z}$ . The dominated convergence and *Lebesgue differentiation* theorems (as  $\varepsilon \to 0$ ) allow us to conclude.  $\Box$ 

Consider now the following assumption:

**Assumption 2** (As in [31]).  $U : \mathbb{Z} \to \mathbb{R}$  is assumed to be  $C^{\infty}$ , with  $\nabla U$  Lipschitz continuous, and such that  $\exists C_U > 0$ ,  $\exists C'_U \in \mathbb{R}$  such that  $\forall x \in \mathbb{Z} : \nabla U(x) \cdot x \ge C_U ||x||^2 + C'_U$ . When required, we will also assume that  $r : \mathbb{Z} \to \mathbb{R}$  satisfies these conditions.

Notice that these conditions imply that  $\exists 0 \leq C' \leq C$  s.t.  $\forall x \in \mathcal{Z}, C' ||x||^2 - C \leq U(x) \leq C(1 + ||x||^2)$  (i.e. U has quadratic growth) and  $|\Delta U(x)| \leq C$ 

Since  $R_{\nu}^{\tau,\beta}$  includes an **entropy term**, it guarantees **strict convexity, weak lower semicontinuity and compact sublevel sets** (see e.g. [31] or [21]). On the other hand, assumption 2 implies that U (or r) will have quadratic growth. Namely, we get (see [31] for a detailed proof):

**Proposition 13** (Existence and Uniqueness of the minimizer (regularized case)). Let R be **convex**, of class  $C^1$  and bounded from below. Let  $\nu$  be the Gibbs measure with potential U. Then,  $R_{\nu}^{\tau,\beta}$  has a **unique minimizer**,  $\mu^{*,\tau,\beta,\nu} \in \mathcal{P}(\mathcal{Z})$ , absolutely continuous with respect to Lebesgue measure  $\lambda$ . When either U or r satisfies assumption 2, this minimizer also belongs to  $\mathcal{P}_2(\mathcal{Z})$ .

For establishing global convergence results further assumptions are requred

**Assumption 3** (Assumptions for well definedness (from [10] and [31])). Assume that the intrinsic derivative  $D_{\mu}R : \mathcal{P}(\mathcal{Z}) \times \mathcal{Z} \to \mathcal{Z}$  of the functional  $R : \mathcal{P}(\mathcal{Z}) \to \mathbb{R}$  exists and satisfies either one of the following:

- 1. (From [31]). Assume:
  - $D_{\mu}R$  is bounded and Lipschitz continuous, i.e.  $\exists C_R > 0$  s.t.  $\forall z, z' \in \mathcal{Z}, \ \forall \mu, \mu' \in \mathcal{P}_2(\mathcal{Z}),$

 $|D_{\mu}R(\mu, z) - D_{\mu}R(\mu', z')| \le C_R[|z - z'| + W_2(\mu, \mu')]$ 

- $\forall \mu \in \mathcal{P}(\mathcal{Z}), \ D_{\mu}R(\mu, \cdot) \in C^{\infty}(\mathcal{Z}).$
- $\nabla D_{\mu}R : \mathcal{P}(\mathcal{Z}) \times \mathcal{Z} \to \mathcal{Z} \times \mathcal{Z}$  is jointly continuous.
- 2. (From [10], who relax some differentiability conditions at the cost of boundedness assumptions; this allows them to avoid altogether the coercivity condition from assumption 2, which is used in [31]):

- $\forall x \in \mathcal{Z}, \forall m, m' \in \mathcal{P}_2(\mathcal{Z}), |D_\mu R(m, x) D_\mu R(m', x)| \leq M_{mm}^R W_1(m, m')$  for some constant  $M_{mm}^R \geq 0$  (i.e. it is lipschitz on the measure argument).
- Suppose that

$$\sup_{\mu \in \mathcal{P}_2(\mathcal{Z})} \sup_{x \in \mathcal{Z}} |\nabla D_{\mu} R(\mu, x)| \le M_{mx}^R$$

for some constant  $M_{mx}^R \ge 0$  i.e.  $\nabla D_{\mu} R(\mu, x)$  is uniformly bounded.

This allows to establish a traditional global convergence result from the MF Theory of NNs:

**Theorem 7** (from [10] and [31]). Let  $\mu_0 \in \mathcal{P}_2(\mathcal{Z})$ , and let assumption 2 and 3 hold; then:

$$\forall t > 0, \quad \frac{d}{dt} (R_{\nu}^{\tau,\beta}(\mu_t)) = -\varsigma(t) \int_{\mathcal{Z}} \left| D_{\mu} R(\mu_t, z) + \tau \nabla r(z) + \beta \frac{\nabla u_t}{u_t}(z) + \beta \nabla U(z) \right|^2 \, d\mu_t(z)$$

where  $u_t$  denotes the density of  $\mu_t := Law(X_t)$ , the solution to equation (3). i.e. following the WGF makes the regularized risk decrease at a known rate. This is known as the energy dissipation equation.

**Remark.** Notice that this equation can be rewritten using the **Fisher divergence** (or relative Fisher Information) between two measures. This quantity is defined as:

$$I(\mu||\nu) := \int_{\mathcal{Z}} \left\| \nabla \log(\frac{d\mu}{d\nu}(z)) \right\|^2 d\mu(z)$$

Then, almost by definition, we get:

$$\frac{d}{dt}(R_{\nu}^{\tau,\beta}(\mu_t)) = -\beta^2 \varsigma(t) I(\mu_t || \hat{\mu}_t)$$

*This allows us to characterize the stationary points of the dynamic explicitly, as done in [10, 31, 45, 56].* 

Theorem 7 implies that the WGF converges to the unique global optimizer of the regularized problem: **Theorem 8** (from [31]). Let R be convex, bounded from below and  $C^1$ ; also assume that assumption 2 and 3 hold. Consider  $\mu_0 \in \bigcup_{p>2} \mathcal{P}_p(\mathcal{Z})$  and let  $(\mu_t)_{t\geq 0}$  be the **WGF** $(R_{\nu}^{\tau,\beta})$  starting from  $\mu_0$ . Then, the equation has a stationary distribution,  $\mu_{\infty}$ , that satisfies:

$$\mu_{\infty} := \arg\min_{\mu \in \mathcal{P}(\mathcal{Z})} R_{\nu}^{\tau, \beta}(\mu) \text{ and } \lim_{t \to \infty} W_2(\mu_t, \mu_{\infty}) = 0$$

**Remark.** Global Convergence Results such as Theorem 7 or Theorem 8 have been established as early as in [42] (for the quadratic loss). However, settings such as those of [10, 31, 45, 56, 13] are of notorious interest to establish essentially the same results under fundamentally more general assumptions.

Making further *technical* assumptions on our regularized functionals leads to better *convergence results*. Namely, consider the following definition:

**Definition 11.** We say  $\mu \in \mathcal{P}(\mathcal{Z})$  satisfies the Log-Sobolev Inequality with constant  $\vartheta > 0$  (in short,  $LSI(\vartheta)$ ), if for any  $\nu \in \mathcal{P}(\mathcal{Z})$  such that  $\nu \ll \mu$ , we have:

$$D(\nu||\mu) := \int_{\mathcal{Z}} \log(\frac{d\nu}{d\mu}(z)) d\nu(z) \le \frac{1}{2\vartheta} \int_{\mathcal{Z}} \left\| \nabla \log(\frac{d\nu}{d\mu}(z)) \right\|^2 d\nu(z) =: \frac{1}{2\vartheta} I(\nu||\mu)$$

where  $D(\nu || \mu)$  is the KL divergence and  $I(\mu || \nu)$  is the **Fisher divergence**.

(see [3, 46] for background on functional inequalities and [13, 10, 45, 56] for applications of it to the NN context). In our setting, as done by most authors in recent years to achieve the desired global convergence results, the following "uniform-LSI" on the functional  $R : \mathcal{P}(\mathcal{Z}) \to \mathcal{R}$  is assumed to hold:

**Assumption 4** (Uniform LSI from [13, 10, 45, 56]). There exists  $\vartheta > 0$  such that  $\forall \mu \in \mathcal{P}_2(\mathcal{Z})$ ,  $\hat{\mu}$  satisfies LSI( $\vartheta$ ). Here  $\hat{\mu}$  is the probability measure with density w.r.t.  $\lambda$  given by (slightly abusing notation, and considering U the potential of a Gibbs measure  $\nu$  used in the entropy, which is 0 if  $\nu = \lambda$ ):

$$\hat{\mu}(z) \propto \exp\left(-\frac{1}{\beta}\frac{\partial R}{\partial \mu}(\mu, z) - \frac{\tau}{\beta}r(z) - U(z)\right)$$

**Remark.** This LSI is a recurrent element in the literature of WGF and Optimal Transport in general. In particular, it implies (see [3]) Poincaré Inequality:  $\forall \phi \in C_b^1(\mathcal{Z})$ ,  $\operatorname{Var}_{\hat{\mu}}(\phi) \leq \frac{1}{2\vartheta} \mathbb{E}_{\hat{\mu}}[|\nabla \phi|^2]$ , and ([46]) the Talagrand's  $T_2$ -transport inequality as well:  $\forall \nu \in \mathcal{P}_2(\mathcal{Z})$ ,  $\vartheta W_2^2(\nu, \hat{\mu}) \leq D(\nu || \hat{\mu})$ 

Beyond the *characterization* of the decay (from [31]), we have the following guarantee: **Theorem 9** (from [10, 13]). Let R be convex,  $C^1$  and bounded from below, and let assumptions 3 and 4 hold. Then, if for some  $t_0 \ge 0$ ,  $\mu_{t_0}$  has finite entropy and finite second moment; then  $\forall t \ge t_0$ ,

$$D(\mu_t || \mu_{\infty}) \leq R_{\nu}^{\tau,\beta}(\mu_t) - R_{\nu}^{\tau,\beta}(\mu_{\infty}) \leq (R_{\nu}^{\tau,\beta}(\mu_{t_0}) - R_{\nu}^{\tau,\beta}(\mu_{\infty}))e^{-2\beta\vartheta \int_{t_0}^{\tau} \varsigma(s)ds}$$

where  $\mu_{\infty} = \mu^{\tau, \beta, \nu} = \arg \min_{\mu \in \mathcal{P}(\mathcal{Z})} R_{\nu}^{\tau \beta}(\mu)$ . That is, the value function following the WGF thus converges exponentially fast to the optimum value of the problem, and this implies an exponential convergence in relative entropy.

One thus recovers, under the right technical assumptions, a version of Theorem 4 from [42] and actually a quantitative improvement of it. By Talagrand's inequality this also implies *exponential*  $W_2$  convergence of  $\mu_t$  to  $\mu_{\infty}$ . We note that the result in [10] is established in the setting with  $\tau = 0, \beta = 1$  and  $\varsigma \equiv 1$ ; however, one can show that the result holds as stated by using standard arguments.

## B.4 Conditions for well-posedness of WGF

Most of the technical conditions that will be here presented are directly taken from both [14] and [19]. We only adapt them slightly to fit into our notation.

Regarding the existence of **weak** solutions to the WGF presented in equation (3), [14] are able to guarantee it under the following assumptions (more general assumptions might be sought in [1, 51], but these are relatively standard in the MF context):

Assumption 5 (Assumptions for existence and uniqueness of the WGF solutions (taken from [14])). Consider a setting as described in proposition 11, with  $R(\mu) := L(\langle \Phi, \mu \rangle) + V(\mu)$ , with  $V(\mu) = \tau \int_{\mathbb{Z}} r d\mu$ .

- 1. Let Z to be the closure of a convex open set within some finite-dimensional euclidean space.
- 2. Let  $L : \mathcal{H} \to \mathbb{R}^+$  be differentiable, with a differential dL that is Lipschitz on bounded sets and bounded on sublevel sets.
- 3. Let  $\Phi : \mathbb{Z} \to \mathcal{H}$  be differentiable and  $V : \mathbb{Z} \to \mathbb{R}^+$  be semiconvex (i.e.  $\exists \lambda \in \mathbb{R} : V + \lambda | \cdot |^2$  is convex).
- 4. There exists a family  $(Q_r)_{r>0}$  of nested nonempty closed convex subsets of  $\mathcal{Z}$  such that:
  - (a)  $\{u \in \Omega; dist(u, Q_{r'}) \leq r\} \subset Q_{r+r'}$  for all r, r' > 0,
  - (b)  $\Phi$  and V are bounded, and  $d\Phi$  is Lipschitz on each  $Q_r$
  - (c)  $\exists C_1, C_2 > 0$  such that  $\sup_{u \in Q_r} (\|d\Phi(u)\| + \|\partial V(u)\|) \le C_1 + C_2 r$  for all r > 0, where  $\|\partial V(u)\|$  stands for the maximal norm of an element in  $\partial V(u)$ .

On the other hand, [19] are able to prove (based on Theorem 1.1. from [57]) the existence of **strong** solutions with pathwise uniqueness for McKean-Vlasov SDE given by

$$dZ_t = b(t, Z_t, \mu_t)dt + \Sigma(t, Z_t, \mu_t)dB_t$$

where b and  $\Sigma$  satisfies the conditions of **B2** (presented below) and for all  $t \ge 0$ ,  $\mu_t = \text{Law}(Z_t) \in \mathcal{P}_2(\mathbb{R}^D)$ ,  $(B_t)_{t\ge 0}$  is an r-dimensional Brownian motion (with  $r \in \mathbb{N}^*$  potentially different from  $D \in \mathbb{N}^*$ ), and  $Z_0$  has the (fixed) law  $\mu^0 \in \mathcal{P}_2(\mathbb{R}^D)$ . For this, consider the following technical assumptions (**B1** and **B2**) which have been taken directly from [19]:

Assumption 6 (Assumptions for the existence and uniqueness of solutions in [19]). Consider:

**B1.** There exist a measurable function  $g : \mathbb{R}^D \times \mathcal{W} \to \mathbb{R}$ ,  $M_1 \ge 0$  and  $\mu_0 \in \mathcal{P}_2(\mathbb{R}^D)$  such that for any  $N \in \mathbb{N}$ , the following hold.

(a) For any 
$$w_1, w_2 \in \mathbb{R}^D$$
 and  $z \in W$  we have  
 $\|g(w_1, z) - g(w_2, z)\| \le \zeta(z) \|w_1 - w_2\|$ , and  $\|g(w_1, z)\| \le \zeta(z)$   
with  $\int_{\mathcal{W}} \zeta^2(z) \, d\pi_{\mathcal{W}}(z) < +\infty$ 

(b)  $b_N \in C(\mathbb{R}_+ \times \mathbb{R}^D \times \mathcal{P}_2(\mathbb{R}^D), \mathbb{R}^D)$  and  $\Sigma_N \in C(\mathbb{R}_+ \times \mathbb{R}^D \times \mathcal{P}_2(\mathbb{R}^D), \mathbb{R}^{D \times r})$ . (c) For any  $w_1, w_2 \in \mathbb{R}^D$  and  $\mu_1, \mu_2 \in \mathcal{P}_2(\mathbb{R}^D)$ 

$$\sup_{t\geq 0} \left\{ \|b_N(t,w_1,\mu_1) - b_N(t,w_2,\mu_2)\| + \|\Sigma_N(t,w_1,\mu_1) - \Sigma_N(t,w_2,\mu_2)\| \right\} \\
\leq M_1 \left( \|w_1 - w_2\| + \left( \int_{\mathcal{W}} \int_{\mathbb{R}^D} |\langle g(\cdot,z),\mu_1 \rangle - \langle g(\cdot,z),\mu_2 \rangle|^2 \, d\pi_{\mathcal{W}}(z) \right)^{1/2} \right\} \right), \\
\sup_{t\geq 0} \{ \|b_N(t,0,\mu_0)\| + \|\Sigma_N(t,0,\mu_0)\| \} \leq M_1.$$

**B2.** There exist  $M_2 \ge 0$ ,  $\kappa > 0$ ,  $b \in C(\mathbb{R}_+ \times \mathbb{R}^D \times \mathcal{P}_2(\mathbb{R}^D), \mathbb{R}^D)$  and  $\Sigma \in C(\mathbb{R}_+ \times \mathbb{R}^D \times \mathcal{P}_2(\mathbb{R}^D), \mathbb{R}^{D \times r})$  such that

$$\sup_{t \ge 0, w \in \mathbb{R}^{D}, \mu \in \mathcal{P}_{2}(\mathbb{R}^{D})} \{ \|b_{N}(t, w, \mu) - b(t, w, \mu)\| + \|\Sigma_{N}(t, w, \mu) - \Sigma(t, w, \mu)\| \} \le M_{2} N^{-\kappa}.$$

**Proposition 14** (Proposition 11 in [19]). Assuming **B1** and **B2**. Given  $\mu^0 \in \mathcal{P}_2(\mathbb{R}^D)$  as a fixed initial condition; then, there exists an  $(\mathcal{F}_t)_{t\geq 0}$ -adapted process  $(Z_t)_{t\geq 0}$  that is the **unique (pathwise)** strong solution of the McKean-Vlasov SDE:

 $dZ_t = b(t, Z_t, \mu_t)dt + \Sigma(t, Z_t, \mu_t)dB_t$ 

Additionally, it satisfies for each  $T \ge 0$ :  $\sup_{t \in [0,T]} \mathbb{E}[||Z_t||^2] < \infty$ 

## **C** Proofs and discussions of main results

#### C.1 Proofs of Section 3.1

*Proof of Proposition 1.* By definition of the symmetrization operator, we know that  $\forall x \in \mathcal{X}$ :

$$(\mathcal{Q}_G \Phi_\mu)(x) = \int_G \hat{\rho}_{g^{-1}} \Phi_\mu(\rho_g x) d\lambda_G(g)$$

For  $g \in G$ , since  $\sigma_*$  is equivariant and  $M_g$  is invertible, we can write:

$$\Phi_{\mu}(\rho_g x) = \langle \sigma_*(\rho_g x, \cdot), \mu \rangle = \langle \sigma_*(\rho_g x, \cdot), M_g \# (M_g^{-1} \# \mu) \rangle = \hat{\rho}_g \langle \sigma_*(x, \cdot), M_g^{-1} \# \mu \rangle$$

where we've used Proposition 7 in the last equality. In turn, we can write (via the inversion-invariance of  $\lambda_G$ ):

$$\begin{aligned} (\mathcal{Q}_G \Phi_\mu)(x) &= \int_G \hat{\rho}_{g^{-1}} \hat{\rho}_g \langle \sigma_*(x, \cdot), M_g^{-1} \# \mu \rangle d\lambda_G(g) = \int_G \langle \sigma_*(x, \cdot), M_{g^{-1}} \# \mu \rangle d\lambda_G(g) \\ &= \int_G \langle \sigma_*(x, \cdot), M_g \# \mu \rangle d\lambda_G(g) = \langle \sigma_*(x, \cdot), \mu^G \rangle = \Phi_{\mu^G}(x) \end{aligned}$$

As mentioned in Section 3.1, a simple case where we will have  $\Phi_{\mu^G} = \Phi_{\mu^{\mathcal{E}^G}}$  is when  $\sigma_*$  is *linear*: **Proposition 15.** If  $\sigma_* : \mathcal{X} \times \mathcal{Z} \to \mathcal{Y}$  is jointly equivariant and  $\pi_{\mathcal{X}}$ -c.s.  $\forall x \in \mathcal{X}, [z \mapsto \sigma_*(x; z)]$  is a bounded linear operator, then, for any  $\mu \in \mathcal{P}(\mathcal{Z})$ :  $\Phi_{\mu^G} = \langle \sigma_*, \mu^G \rangle = \langle \sigma_*, \mu^{\mathcal{E}^G} \rangle = \Phi_{\mu^{\mathcal{E}^G}}$ .

*Proof of Proposition 15.* A straightforward computation yields (using Fubini's theorem and the linearity of integrals and  $\sigma_*$ ),  $\forall \mu \in \mathcal{P}(\mathcal{Z})$ :

$$\langle \sigma_*, \mu^G \rangle = \int_G \int_{\mathcal{Z}} \sigma_*(x, M_g.z) d\mu(z) d\lambda_G(g) = \int_{\mathcal{Z}} \int_G \sigma_*(x, M_g.z) d\lambda_G(g) d\mu(z)$$
  
= 
$$\int_{\mathcal{Z}} \sigma_*(x, \int_G M_g.z \, d\lambda_G(g)) d\mu(z) = \int_{\mathcal{Z}} \sigma_*(x, P_{\mathcal{E}^G}.z) d\mu(z) = \langle \sigma_*, \mu^{\mathcal{E}^G} \rangle$$

#### C.2 Proofs of results in Section 3.2

#### C.2.1 Proof of Proposition 2

Now consider, as a shorthand notation,  $\forall x \in \mathcal{X}, \forall y \in \mathcal{Y}$  the functional  $L_{x,y} : \mathcal{P}(\mathcal{Z}) \to \mathbb{R}$  given by  $\forall \mu \in \mathcal{P}(\mathcal{Z})$ :  $L_{x,y}(\mu) = \ell (\Phi_{\mu}(x), y)$ . The following lemma that shall be useful for later stages. **Lemma 4.** Let  $\sigma_*$  be jointly equivariant and  $\ell$  be invariant. Then,  $\forall g \in G, \forall x \in \mathcal{X}, \forall y \in \mathcal{Y}, \forall \mu \in \mathcal{P}(\mathcal{Z})$ ,

$$L_{\rho_g.x,\hat{\rho}_g.y}(M_g \# \mu) = L_{x,y}(\mu)$$

Equivalently, the map  $L : \mathcal{P}(\mathcal{Z}) \to L^2(\mathcal{X} \times \mathcal{Y}, \pi)$  given by  $L(\mu) \mapsto [(x, y) \mapsto L_{x,y}(\mu)]$  is equivariant (under the appropriate<sup>9</sup> G-actions).

*Proof of Lemma 4.* Using the joint equivariance of  $\sigma_*$  (via proposition 7) and the invariance of  $\ell$ , a straightforward computation yields, for all  $x \in \mathcal{X}, y \in \mathcal{Y}$ , and  $g \in G$ :

$$L_{\rho_g.x,\hat{\rho}_g.y}(M_g \# \mu) = \ell \left( \langle \sigma_*(\rho_g.x,\cdot), M_g \# \mu \rangle, \hat{\rho}_g.y \right)$$
  
=  $\ell \left( \hat{\rho}_g. \langle \sigma_*(x,\cdot), \mu \rangle, \hat{\rho}_g.y \right)$   
=  $\ell \left( \langle \sigma_*(x,\cdot), \mu \rangle, y \right) = L_{x,y}(\mu)$ 

With this we can prove Proposition 2. Notice that we will basically utilize the equivariance properties of  $\sigma_*$  and  $\ell$  in Assumption 1 (the convexity of the functions comes directly from the convexity of R when  $\ell$  is convex, together with the linearity of  $(\cdot)^G$ ,  $(\cdot)^{\mathcal{E}^G}$  and  $\int_G (\cdot) d\lambda_G$ .).

Proof of Proposition 2. We can readily see that:  $R^{EA}(\mu) = \mathbb{E}_{\pi} \left[ \ell \left( \Phi_{\mu^{\mathcal{E}G}}(X), Y \right) \right] = R(\mu^{\mathcal{E}^G}).$ On the other hand, as  $\sigma_*$  is jointly equivariant, from Proposition 1, we have:

$$R^{FA}(\mu) = \mathbb{E}_{\pi} \left[ \ell \left( \mathcal{Q}_G(\Phi_{\mu})(X), Y \right) \right] = \mathbb{E}_{\pi} \left[ \ell \left( \Phi_{\mu^G}(X), Y \right) \right] = R(\mu^G)$$

Next, using Lemma 4, Fubini's theorem and the inversion-invariance of  $\lambda_G$ , we get:

$$\begin{split} R^{DA}(\mu) &= \mathbb{E}_{\pi} \left[ \int_{G} \ell \left( \Phi_{\mu}(\rho_{g}.X), \hat{\rho}_{g}.Y \right) d\lambda_{G}(g) \right] = \mathbb{E}_{\pi} \left[ \int_{G} L_{\rho_{g}.X, \hat{\rho}_{g}.Y}(\mu) d\lambda_{G}(g) \right] \\ &= \int_{G} \mathbb{E}_{\pi} \left[ L_{\rho_{g}.X, \hat{\rho}_{g}.Y}(\mu) \right] d\lambda_{G}(g) = \int_{G} \mathbb{E}_{\pi} \left[ L_{\rho_{g}.X, \hat{\rho}_{g}.Y}(M_{g} \# M_{g}^{-1} \# \mu) \right] d\lambda_{G}(g) \\ &= \int_{G} \mathbb{E}_{\pi} \left[ L_{X,Y}(M_{g^{-1}} \# \mu) \right] d\lambda_{G}(g) = \int_{G} \mathbb{E}_{\pi} \left[ L_{X,Y}(M_{g} \# \mu) \right] d\lambda_{G}(g) \\ &= \int_{G} R(M_{g} \# \mu) d\lambda_{G}(g) = R^{G}(\mu) \end{split}$$

From these expressions we can quickly verify that  $R^{DA}$ ,  $R^{FA}$  and  $R^{EA}$  are invariant. Namely, by Lemma 1, for  $g \in G$  and  $\mu \in \mathcal{P}(\mathcal{Z})$  we have:  $R^{FA}(M_g \# \mu) = R((M_g \# \mu)^G) = R(\mu^G) = R^{FA}(\mu)$ , and:  $R^{EA}(M_g \# \mu) = R((M_g \# \mu)^{\mathcal{E}^G}) = R(\mu^{\mathcal{E}^G}) = R^{EA}(\mu)$ . On the other hand, the right-invariance of  $\lambda_G$  implies:

$$R^{DA}(M_g \# \mu) = \int_G R((M_h \# (M_g \# \mu)) d\lambda_G(h) = \int_G R((M_{\tilde{h}} \# \mu)) d\lambda_G(\tilde{h}) = R^{DA}(\mu).$$

We can also see that whenever R is invariant, we have that, for  $\mu \in \mathcal{P}(\mathcal{Z})$  and  $g \in G$ ,  $R(M_g \# \mu) = R(\mu)$ , so that:  $R^{DA}(\mu) = R^G(\mu) = \int_G R(M_g \# \mu) d\lambda_G(g) = \int_G R(\mu) d\lambda_G(g) = R(\mu)$ .

<sup>&</sup>lt;sup>9</sup>In this case,  $\forall g \in G$ , let  $g.\mu = M_g \# \mu$  and  $g.f = f^g$  given by  $\forall x \in \mathcal{X}, \ \forall y \in \mathcal{Y}, \ f^g(x,y) = f(\rho_g^{-1}.x, \hat{\rho}_g^{-1}.y)$ 

Also, if  $\mu \in \mathcal{P}^G(\mathcal{Z})$ , we have, for all  $g \in G$ ,  $\mu = \mu^G = M_g \# \mu$ , so that:  $R^{DA}(\mu) = \int_G R(M_g \# \mu) d\lambda_G(g) = \int_G R(\mu) d\lambda_G(g) = R(\mu) = R(\mu^G) = R^{FA}(\mu)$ .

Finally, we verify that our population risk  $R : \mathcal{P}(\mathcal{Z}) \to \mathbb{R}$  is invariant whenever  $\pi \in \mathcal{P}^G(\mathcal{X} \times \mathcal{Y})$ . Indeed,  $\forall g \in G$  and  $\forall \mu \in \mathcal{P}(\mathcal{Z})$ , by the invariance of  $\ell$  and  $\pi$ , together with Proposition 7 from the equivariance of  $\sigma_*$ , we get:

$$R(M_g \# \mu) = \mathbb{E}_{\pi} \left[ \ell(\langle \sigma_*(X; \cdot), M_g \# \mu \rangle, Y) \right] = \mathbb{E}_{\pi} \left[ \ell(\hat{\rho}_g \langle \sigma_*(\rho_g^{-1}X; \cdot), \mu \rangle, \hat{\rho}_g \hat{\rho}_g^{-1}Y) \right]$$
$$= \mathbb{E}_{\pi} \left[ \ell(\langle \sigma_*(\rho_g^{-1}X; \cdot), \mu \rangle, \hat{\rho}_g^{-1}Y) \right] = \mathbb{E}_{\pi} \left[ \ell(\langle \sigma_*(X; \cdot), \mu \rangle, Y) \right] = R(\mu)$$

That is, R is invariant.

Notice that the equivariance of the data distribution  $\pi$  can also make the *regularized* population risk be invariant, under the right choice of r. Namely:

**Corollary 6.** If  $R : \mathcal{P}(\mathcal{Z}) \to \mathbb{R}$  and  $r : \mathcal{Z} \to \mathbb{R}$  are invariant (in their respective sense), then  $R^{\tau,\beta}$  is invariant.

The result can be proven for  $R_{\nu}^{\tau,\beta}$  with  $\nu$  some *G*-invariant measure (such as  $\lambda$  for orthogonal representations). Notice that  $r(\theta) = \|\theta\|^2$  is an example of invariant function for *orthogonal representations*.

Proof of Corollary 6. It is enough to notice that  $V(\mu) = \int_{\mathcal{Z}} r(\theta) d\mu(\theta)$  and  $H_{\nu}(\mu) = \int_{\mathcal{Z}} \log(\frac{d\mu}{d\nu}) d\mu$ (with  $\mu \ll \nu$ ) are invariant when  $r : \mathcal{Z} \to \mathbb{R}$  and  $\nu \in \mathcal{P}(\mathcal{Z})$  are invariant (in their respective sense):

1. For V, notice that for  $g \in G$ :

$$V(M_g \# \mu) = \int_{\mathcal{Z}} r(\theta) d(M_g \# \mu)(\theta) = \int_{\mathcal{Z}} r(M_g \theta) d\mu(\theta) = \int r(\theta) d\mu(\theta) = V(\mu)$$

thanks to the invariance of r. i.e. V is invariant.

2. For  $H_{\nu}$ , notice that, for  $g \in G$ , as  $\nu$  is invariant, we know that  $\frac{d(M_g \# \mu)}{d\nu}(x) = \frac{d\mu}{d\nu}(M_g^{-1}x)$ . Therefore:

$$H_{\nu}(M_g \# \mu) = \int \log\left(\frac{d(M_g \# \mu)}{d\nu}(\theta)\right) d(M_g \# \mu)(\theta)$$
$$= \int \log\left(\frac{d\mu}{d\nu}(M_g^{-1}\theta)\right) d(M_g \# \mu)(\theta)$$
$$= \int \log\left(\frac{d\mu}{d\nu}(M_g^{-1}M_g\theta)\right) d\mu(\theta) = H_{\nu}(\mu)$$

Which proves that  $H_{\nu}$  is invariant.

We can readily conclude, since:  $R^{\tau,\beta}(\mu) = R(\mu) + \tau \int r d\mu + \beta H_{\lambda}(\mu)$ , for all  $\mu \in \mathcal{P}(\mathcal{Z})$ .

## C.2.2 Proof of Proposition 3

In order to prove Proposition 3, we require a version of Jensen's inequality that's suited for our context. Such a result might exist in the literature, but since we couldn't find a complete proof under our assumptions, we provide our own.

**Proposition 16 (Jensen's Inequality).** Let  $F : \mathcal{P}(\mathcal{Z}) \longrightarrow \mathbb{R}$  be such that Lemma 3 holds. Let S be some measurable space,  $\lambda \in \mathcal{P}(S)$  and  $s \in S \mapsto \mu_s \in \mathcal{P}(\mathcal{Z})$  a measurable function. Define  $\tilde{\mu} \in \mathcal{P}(\mathcal{Z})$  as the intensity measure:  $\tilde{\mu} = \int_S \mu_s d\lambda(s) \in \mathcal{P}(\mathcal{Z})$ . Then, Jensen's inequality holds:

$$F(\tilde{\mu}) \leq \int_{S} F(\mu_s) d\lambda(s)$$

*Proof of Proposition 16.* Since Lemma 3 holds, we have that  $\forall \mu_1, \mu_2 \in \mathcal{P}(\mathcal{Z})$ :

$$F(\mu_1) \ge F(\mu_2) + \int \frac{\partial F}{\partial \mu}(\mu_2, z) d(\mu_1 - \mu_2)(z)$$

Let  $\tilde{s} \in S$  be arbitrary and consider  $\mu_2 = \tilde{\mu} := \int \mu_s d\lambda(s)$ ; and  $\mu_1 = \mu_{\tilde{s}}$ . Then:

$$F(\left(\int \mu_s d\lambda(s)\right) \le F(\mu_{\tilde{s}}) - \int \frac{\partial F}{\partial \mu} \left(\int \mu_s d\lambda(s), z\right) d\left(\mu_{\tilde{s}} - \int \mu_s d\lambda(s)\right)(z)$$

Integrating the inequality with respect to  $\lambda$  (on  $\tilde{s}$ ):

$$\int_{S} F\left(\int \mu_{s} d\lambda(s)\right) d\lambda(\tilde{s})$$
  
$$\leq \Delta := \left[\int_{S} F(\mu_{\tilde{s}}) d\lambda(\tilde{s}) - \int_{S} \left(\int_{\mathcal{Z}} \frac{\partial F}{\partial \mu} \left(\int_{S} \mu_{s} d\lambda(s), \cdot\right) d\left(\mu_{\tilde{s}} - \int_{S} \mu_{s} d\lambda(s)\right)\right) d\lambda(\tilde{s})\right]$$

We notice that the LHS doesn't depend on  $\tilde{s}$ , so that  $\int_{S} F\left(\int \mu_s d\lambda(s)\right) d\lambda(\tilde{s}) = F\left(\int \mu_s d\lambda(s)\right)$ . On the other hand, the right-most term in  $\triangle$  can be developed as:

$$\begin{split} \star &:= \int_{S} \left( \int_{\mathcal{Z}} \frac{\partial F}{\partial \mu} \left( \int_{S} \mu_{s} d\lambda(s), \cdot \right) d \left( \mu_{\tilde{s}} - \int_{S} \mu_{s} d\lambda(s) \right) \right) d\lambda(\tilde{s}) \\ &= \int_{S} \left( \int_{\mathcal{Z}} \frac{\partial F}{\partial \mu} \left( \int_{S} \mu_{s} d\lambda(s), \cdot \right) d\mu_{\tilde{s}} - \int_{\mathcal{Z}} \frac{\partial F}{\partial \mu} \left( \int_{S} \mu_{s} d\lambda(s), \cdot \right) d \left( \int_{S} \mu_{s} d\lambda(s) \right) \right) d\lambda(\tilde{s}) \\ &= \int_{S} \left( \int_{\mathcal{Z}} \frac{\partial F}{\partial \mu} \left( \int \mu_{s} d\lambda(s), \cdot \right) d(\mu_{\tilde{s}}) \right) d\lambda(\tilde{s}) \\ &- \int_{S} \left( \int_{\mathcal{Z}} \frac{\partial F}{\partial \mu} \left( \int_{S} \mu_{s} d\lambda(s), \cdot \right) d \left( \int_{S} \mu_{s} d\lambda(s) \right) \right) d\lambda(\tilde{s}) \end{split}$$

Notice that the *linear functional derivative* is chosen in such a way so that it satisfies  $\forall \nu \in \mathcal{P}(\mathcal{Z}), \int_{\mathcal{Z}} \frac{\partial F}{\partial \mu}(\nu, z) d\nu(z) = 0$ . In particular, the second term of the previous expression vanishes. We get that

$$\star = \int_{S} \left( \int_{\mathcal{Z}} \frac{\partial F}{\partial \mu} \left( \int_{S} \mu_{s} d\lambda(s), z \right) d\mu_{\tilde{s}}(z) \right) d\lambda(\tilde{s})$$

But, by definition:  $\forall f : \mathbb{Z} \to \mathbb{R}$  integrable,

$$\langle f, \int_{S} \mu_{s} d\lambda(s) \rangle = \int_{S} \langle f, \mu_{s} \rangle d\lambda(s) = \int_{S} \left( \int_{\mathcal{Z}} f(z) d\mu_{s}(z) \right) d\lambda(s)$$

So this is, by definition, and applying the same convention on the definition of the linear functional derivative<sup>10</sup>:

$$\star = \int_{\mathcal{Z}} \frac{\partial F}{\partial \mu} \left( \int \mu_s, z \right) d\left( \int \mu_{\tilde{s}} d\lambda(\tilde{s}) \right)(z) = 0$$

With this, we conclude that  $\triangle = \int_{S} F(\mu_s) d\lambda(s)$ , and so, we get that:

$$F(\tilde{\mu}) = F\left(\int_{S} \mu_s d\lambda(s)\right) \le \int_{S} F(\mu_s) d\lambda(s)$$

which corresponds to Jensen's inequality.

**Remark.** We believe that the  $C^1$  hypothesis can be lifted. Understanding what happens when equality holds should be of interest in our context and more generally.

Thanks to this *Jensen inequality*, we readily get the following result: **Corollary 7.** If  $F : \mathcal{P}(\mathcal{Z}) \longrightarrow \mathbb{R}$  is convex,  $\mathcal{C}^1$  and invariant, then  $\forall \mu \in \mathcal{P}(\mathcal{Z}): F(\mu^G) \leq F(\mu)$ 

*Proof.* Direct from the definition of  $(\cdot)^G$  and Proposition 16.

With these results in place, we are ready to prove Proposition 3:

<sup>&</sup>lt;sup>10</sup>Notice that the *convention* on the linear functional derivative's definition isn't truly important for the proof, since in the end  $\star$  simply corresponds to the term  $\int_{\mathcal{Z}} \frac{\partial F}{\partial \mu} \left( \int \mu_s, z \right) d\left( \int \mu_{\tilde{s}} d\lambda(\tilde{s}) \right)(z)$  being substracted to itself.

*Proof of Proposition 3.* Evidently, since  $\mathcal{P}^G(\mathcal{Z}) \subseteq \mathcal{P}(\mathcal{Z})$ , we have:

$$\inf_{\mu \in \mathcal{P}^G(\mathcal{Z})} F(\mu) \ge \inf_{\mu \in \mathcal{P}(\mathcal{Z})} F(\mu)$$

For the other inequality, take  $(\mu_n)_{n\in\mathbb{N}} \subseteq \mathcal{P}(\mathcal{Z})$  to be an *infinizing sequence* for F; i.e. such that  $F(\mu_n) \ge F(\mu_{n+1})$  and  $F(\mu_n) \xrightarrow[n\to\infty]{} \inf_{\mu\in\mathcal{P}(\mathcal{Z})} F(\mu)$ ). Such a sequence always exists. By Corollary 7, we have  $\forall n \in \mathbb{N}, \ F(\mu_n^G) \le F(\mu_n)$ ; thus,  $\forall n \in \mathbb{N}$ :

$$\inf_{\mu \in \mathcal{P}^G(\mathcal{Z})} F(\mu) \le F(\mu_n^G) \le F(\mu_n)$$

Which allows us to infer, by taking  $n \to \infty$ , that:  $\inf_{\mu \in P^G(\mathcal{Z})} F(\mu) \leq \inf_{\mu \in P(\mathcal{Z})} F(\mu)$ . In turn, we can conclude that:

$$\inf_{\in \mathcal{P}^G(\mathcal{Z})} F(\mu) = \inf_{\mu \in \mathcal{P}(\mathcal{Z})} F(\mu)$$

Notice that if there was some minimizer  $\mu_* \in \arg \min_{\mu \in \mathcal{P}(\mathcal{Z})} F(\mu)$ , then by Corollary 7 we would also have  $\mu^G_* \in \arg \min_{\mu \in \mathcal{P}(\mathcal{Z})} R(\mu)$ . Namely, if such a minimizer was **unique**, then it would satisfy:  $\mu_* = \mu^G_* \in \mathcal{P}^G(\mathcal{Z})$ . That is, the unique solution would be **WI**.

# C.2.3 Proof of Theorem 2 and Corollary 1

Notice that, under Assumption 1, from Corollary 5 we know R is of class  $C^1$  (as well as convex). This properties actually transfers to the functionals  $R^{DA}$ ,  $R^{FA}$  and  $R^{EA}$ , as shown by the following result:

**Proposition 17.** If  $R : \mathcal{P}(\mathcal{Z}) \to \mathbb{R}$  is a convex and  $\mathcal{C}^1$  functional, then  $R^{DA}$ ,  $R^{FA}$  and  $R^{EA}$  are convex and  $\mathcal{C}^1$  as well, with linear functional derivatives given by:

$$\begin{aligned} \frac{\partial R^{DA}}{\partial \mu}(\mu,z) &= \int_{G} \frac{\partial R}{\partial \mu} (M_{g} \# \mu, M_{g}.z) d\lambda_{G}(g), \quad \frac{\partial R^{FA}}{\partial \mu}(\mu,z) = \int_{G} \frac{\partial R}{\partial \mu} (\mu^{G}, M_{g}.z) d\lambda_{G}(g) \\ and \quad \frac{\partial R^{EA}}{\partial \mu}(\mu,z) &= \frac{\partial R}{\partial \mu} (\mu^{\mathcal{E}^{G}}, P_{\mathcal{E}^{G}}.z) \end{aligned}$$

And intrinsic derivatives given by (when well defined):

$$D_{\mu}R^{DA}(\mu,z) = \int_{G} M_g^T D_{\mu}R(M_g \# \mu, M_g.z) d\lambda_G(g)$$

$$D_{\mu}R^{FA}(\mu,z) = \int_{G} M_{g}^{T} D_{\mu}R(\mu^{G}, M_{g}.z)d\lambda_{G}(g) \text{ and } D_{\mu}R^{EA}(\mu,z) = P_{\mathcal{E}^{G}}^{*} D_{\mu}R(\mu^{\mathcal{E}^{G}}, P_{\mathcal{E}^{G}}.z)$$
  
In particular, from Proposition 16, we have that:  $\forall \mu \in \mathcal{P}(\mathcal{Z}), \ R^{FA}(\mu) \leq R^{DA}(\mu)$ 

*Proof of Proposition 17.* We can calculate the linear functional derivatives (l.d.f. for short) as follows. Let  $\mu, \nu \in \mathcal{P}(\mathcal{Z})$ , and consider:

$$\begin{split} \lim_{h \to 0} \frac{R^{DA}((1-h)\mu + h\nu) - R^{DA}(\mu)}{h} \\ &= \lim_{h \to 0} \frac{\int_G R(M_g \# ((1-h)\mu + h\nu))d\lambda_G(g) - \int_G R(M_g \# \mu)d\lambda_G(g)}{h} \\ &= \lim_{h \to 0} \int_G \frac{R((1-h)M_g \# \mu + hM_g \# \nu) - R(M_g \# \mu)}{h}d\lambda_G(g) \\ &= \int_G \lim_{h \to 0} \frac{R((1-h)M_g \# \mu + hM_g \# \nu) - R(M_g \# \mu)}{h}d\lambda_G(g) \\ &= \int_G \int_Z \frac{\partial R}{\partial \mu} (M_g \# \mu, z) d(M_g \# \nu - M_g \# \mu)(z) d\lambda_G(g) \\ &= \int_G \int_Z \frac{\partial R}{\partial \mu} (M_g \# \mu, M_g.z) d(\nu - \mu)(z) d\lambda_G(g) \\ &= \int_Z \int_G \frac{\partial R}{\partial \mu} (M_g \# \mu, M_g.z) d\lambda_G(g) d(\nu - \mu)(z). \end{split}$$

We have used Fubini's theorem, which is applicable<sup>11</sup> thanks to the fact that R is of class  $C^1$ , and we've used the definition of the linear functional derivative for R. Also, we see that (using Fubini's theorem once again, as well as the definition of the linear functional derivative of R):

$$\int_{\mathcal{Z}} \int_{G} \frac{\partial R}{\partial \mu} (M_g \# \mu, M_g. z) d\lambda_G(g) d\mu(z) = \int_{G} \int_{\mathcal{Z}} \frac{\partial R}{\partial \mu} (M_g \# \mu, z) d(M_g \# \mu)(z) d\lambda_G(g) = 0.$$

We can then identify:

$$\frac{\partial R^{DA}}{\partial \mu}(\mu, z) = \int_{G} \frac{\partial R}{\partial \mu} (M_g \# \mu, M_g. z) d\lambda_G(g),$$

and, by taking the gradient:

$$D_{\mu}R^{DA}(\mu,z) = \int_{G} M_{g}^{T}.D_{\mu}R(M_{g}\#\mu,M_{g}.z)d\lambda_{G}(g).$$

We analogously calculate the expression for the l.f.d. of  $R^{FA}$ ; let  $\mu, \nu \in \mathcal{P}(\mathcal{Z})$ :

$$\lim_{h \to 0} \frac{R^{FA}((1-h)\mu + h\nu) - R^{FA}(\mu)}{h} = \lim_{h \to 0} \frac{R((1-h)\mu^G + h\nu^G) - R(\mu^G)}{h}$$
$$= \int_{\mathcal{Z}} \frac{\partial R}{\partial \mu} (\mu^G, z) d(\nu^G - \mu^G)(z)$$
$$= \int_{\mathcal{Z}} \int_{G} \frac{\partial R}{\partial \mu} (\mu^G, M_g.z) d\lambda_G(g) d(\nu - \mu)(z),$$

and also:

$$\begin{split} \int_{\mathcal{Z}} \int_{G} \frac{\partial R}{\partial \mu} (\mu^{G}, M_{g}.z) d\lambda_{G}(g) d\mu(z) &= \int_{G} \int_{\mathcal{Z}} \frac{\partial R}{\partial \mu} (\mu^{G}, z) d(M_{g} \# \mu)(z) d\lambda_{G}(g) \\ &= \int_{\mathcal{Z}} \frac{\partial R}{\partial \mu} (\mu^{G}, z) d\mu^{G}(z) = 0. \end{split}$$

So, by the definition of the lineal functional derivative, we identify:

$$\frac{\partial R^{FA}}{\partial \mu}(\mu, z) = \int_{G} \frac{\partial R}{\partial \mu}(\mu^{G}, M_{g}.z) d\lambda_{G}(g),$$

and taking the gradient we get:

$$D_{\mu}R^{FA}(\mu,z) = \int_{G} M_{g}^{T}.D_{\mu}R(\mu^{G},M_{g}.z)d\lambda_{G}(g)$$

Lastly, the l.f.d. of  $R^{EA}$  is calculated similarly, noticing that:

$$\lim_{h \to 0} \frac{R^{EA}((1-h)\mu + h\nu) - R^{EA}(\mu)}{h} = \lim_{h \to 0} \frac{R(P_{\mathcal{E}^G} \# ((1-h)\mu + h\nu)) - R(P_{\mathcal{E}^G} \mu)}{h}$$
$$= \int_{\mathcal{Z}} \frac{\partial R}{\partial \mu} (\mu^{\mathcal{E}^G}, z) d(P_{\mathcal{E}^G} \# \nu - P_{\mathcal{E}^G} \# \mu)(z)$$
$$= \int_{\mathcal{Z}} \frac{\partial R}{\partial \mu} (\mu^{\mathcal{E}^G}, P_{\mathcal{E}^G} z) d(\nu - \mu)(z),$$

and that:

$$\int_{\mathcal{Z}} \frac{\partial R}{\partial \mu} (\mu^{\mathcal{E}^G}, P_{\mathcal{E}^G}.z) d\mu(z) = \int_{\mathcal{Z}} \frac{\partial R}{\partial \mu} (\mu^{\mathcal{E}^G}, z) d(\mu^{P_{\mathcal{E}^G}})(z) = 0$$

We can thus identify:

$$\frac{\partial R^{EA}}{\partial \mu}(\mu,z) = \frac{\partial R}{\partial \mu}(\mu^{\mathcal{E}^{G}},P_{\mathcal{E}^{G}}.z) \quad \text{and} \quad D_{\mu}R^{EA}(\mu,z) = P_{\mathcal{E}^{G}}^{T}.D_{\mu}R(\mu^{\mathcal{E}^{G}},P_{\mathcal{E}^{G}}.z)$$

The last remark is direct from proposition 16.

<sup>&</sup>lt;sup>11</sup>In particular, as for any fixed  $\mu \in \mathcal{P}(\mathcal{Z})$  the function  $g \in G \mapsto M_g \# \mu$  is continuous (thus, of **compact** image), then the function  $(g, z) \in G \times \mathcal{Z} \mapsto \frac{\partial R}{\partial \mu}(M_g \# \mu, M_g.z)$  is **bounded** 

With all of these different elements in place, we are ready to prove Theorem 2.

*Proof of Theorem* 2. Under Assumption 1, R is **convex** and of **class**  $C^1$  from Corollary 5; and so are  $R^G$ ,  $R^{FA}$  and  $R^{EA}$ , from Proposition 17. Since Proposition 2 ensures that the latter are always invariant, Proposition 3 implies that  $R^{DA}$ ,  $R^{FA}$  and  $R^{EA}$  can all be optimized by only considering **weakly equivariant models** (explaining the first and last equalities). The two middle equalities follow directly from Proposition 2, since R,  $R^{DA}$  and  $R^{FA}$  coincide over  $\mathcal{P}^G(\mathcal{Z})$ .

In the case of the quadratic loss, one can employ the properties of  $Q_G$  from [22] to show Corollary 1.

Proof of Corollary 1. Notice that, for  $\mu \in \mathcal{P}^G(\mathcal{Z})$ ,  $\Phi_\mu$  is a *G*-invariant function (i.e.  $\Phi_\mu \in L^2_G(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})$ ). Also, a simple calculation (see e.g. [4, 40]) allows us to write:  $R(\mu) = \mathbb{E}_{\pi}[\|\Phi_{\mu}(X) - Y\|^2_{\mathcal{Y}}] = R_* + \mathbb{E}_{\pi_{\mathcal{X}}}[\|\Phi_{\mu}(X) - f_*(X)\|^2_{\mathcal{Y}}] = R_* + \|\Phi_{\mu} - f_*\|^2_{L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})}$  with  $R_* = \mathbb{E}_{\pi}[\|Y - f^*(X)\|^2_{\mathcal{Y}}]$  being independent of  $\mu$ . We can thus write (simplifying subscripts for simplicity):

$$\begin{aligned} R(\mu) &= R_* + \|\Phi_{\mu} - \mathcal{Q}_G.f_* + \mathcal{Q}_G.f_* - f_*\|_{L^2(\mathcal{X},\mathcal{Y};\pi_{\mathcal{X}})}^2 \\ &= R_* + \|\Phi_{\mu} - \mathcal{Q}_G.f_*\|_{L^2}^2 + \|(f_*)_G^{\perp}\|_{L^2}^2 - 2\langle \Phi_{\mu} - \mathcal{Q}_G.f_*, (f_*)_G^{\perp} \rangle_{L^2} \end{aligned}$$

where  $(f_*)_G^{\perp} := f_* - \mathcal{Q}_G \cdot f_*$ . We notice that, since  $\Phi_{\mu}$  and  $\mathcal{Q}_G \cdot f_*$  are *G*-equivariant functions, we have that  $\langle (\Phi_{\mu} - \mathcal{Q}_G \cdot f_*), (f_*)_G^{\perp} \rangle_{L^2} = 0$ . That is, for any  $\mu \in \mathcal{P}^G(\mathcal{Z})$ , we have  $R(\mu) = \tilde{R}_* + \|\Phi_{\mu} - \mathcal{Q}_G \cdot f_*\|_{L^2}^2$ , where  $\tilde{R}_* := R_* + \|(f_*)_G^{\perp}\|_{L^2(\mathcal{X},\mathcal{Y};\pi_{\mathcal{X}})}^2$  is independent of  $\mu$  (and doesn't intervene in the optimization). Finally, we get:

$$\inf_{\mu \in \mathcal{P}^G(\mathcal{Z})} R(\mu) = \ddot{R}_* + \inf_{\mu \in \mathcal{P}^G(\mathcal{Z})} \|\Phi_\mu - \mathcal{Q}_G \cdot f_*\|_{L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})}^2$$

#### C.2.4 Proof of Corollary 2

When  $\pi$  is assumed to be equivariant, we can summon our previous results to prove Corollary 2.

*Proof of Corollary 2.* From Proposition 2 we know that equivariant data implies  $R : \mathcal{P}(\mathcal{Z}) \to \mathbb{R}$  is invariant, and also that this makes  $R = R^{DA}$ . We conclude using Theorem 2.

We can readily extend this to the regularized case by recalling Corollary 6:

**Corollary 8.** When  $R : \mathcal{P}(\mathcal{Z}) \to \mathbb{R}$  and  $r : \mathcal{Z} \to \mathbb{R}$  are *G*-invariant, a minimum for the regularized population risk  $R^{\tau,\beta}$  can be found within  $\mathcal{P}^G(\mathcal{Z})$ . When  $\beta > 0$  such **WI** minimum is **unique**.

*Proof of Corollary* 8. Direct from Corollary 6, together with Proposition 3 (as in Corollary 2). The uniqueness comes from the strict convexity of the entropy term (see proposition 13).  $\Box$ 

## C.2.5 Proof of Proposition 4 and Proposition 5

*Proof of Proposition 4.* Consider the group  $G = C_4$  acting on  $\mathbb{R}^2$  via 90° rotations. Let  $K = B(0,1) \subseteq \mathbb{R}^2$  be a compact set. Consider a random variable  $X \sim \mathcal{N}(0, \mathrm{Id}_2)|_K$  (i.e. given by  $X = Z \mathbb{1}_{Z \in K}$  for  $Z \sim \mathcal{N}(0, \mathrm{Id}_2)$ ) and set  $Y = ||X||^2$ . Notice that  $\pi$  defined this way is compactly supported.

Clearly G is finite (thus compact) and it can be seen as its ortogonal representation:

$$\rho_G = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right\} \subseteq O(2) , \ \hat{\rho}_G = \{ \mathrm{Id}_1 \} \subseteq O(1) \ \text{(trivial repr.)}$$

By the definition of our r.v.s, it is clear that:

•  $X \stackrel{(d)}{=} \rho_g X \ \forall g \in G \text{ because } X \sim N(0, \mathrm{Id}_2)$ 

•  $\forall g \in G, (X,Y) \stackrel{(d)}{=} (\rho_g X, \hat{\rho}_g Y)$  (since  $\hat{\rho}$  is the trivial representation, it is enough to notice that  $\|\rho_g X\|^2 = \|X\|^2$  for all  $g \in G$ .

Therefore,  $\pi = \text{Law}(X, Y)$  is *G*-invariant (and compactly supported). Consider a *shallow NN* given by:  $\Phi_{\theta}^{N} : \mathbb{R}^{2} \longrightarrow \mathbb{R}^{N \times b} \longrightarrow \mathbb{R}$  (with  $b \in \mathbb{N}$  and some action  $G \ominus_{\eta} \mathbb{R}^{b}$ ) as:  $\Phi_{\theta}^{N}(x) = \frac{1}{N} \sum_{i=1}^{N} W_{i} \sigma(A_{i}^{T}x + B_{i}), \quad \forall x \in \mathbb{R}^{d}$ ; where  $\theta_{i} = (W_{i}, A_{i}, B_{i}) \in \mathcal{Z} := \mathbb{R}^{1 \times b} \times \mathbb{R}^{2 \times b} \times \mathbb{R}^{b} \cong \mathbb{R}^{D}$ . We let  $G \ominus_{M} \mathcal{Z}$  as described in appendix A.4:

$$M_g.\theta_i = (\hat{\rho}_g W_i \ \eta_g^T, \rho_g \ A_i \ \eta_g^T, \eta_g.B_i) = (W_i \ \eta_g^T, \rho_g \ A_i \ \eta_g^T, \eta_g.B_i)$$

We can assume, for instance, that b = 1 and  $\eta$  is the trivial representation (making it so no condition is required for  $\sigma_*$  to be jointly *G*-equivariant) and recall that:  $\theta_i \in \mathcal{E}^G \iff \forall g \in G, M_g \theta_i = \theta_i$ . However, if we assume that:  $\forall g \in G, \rho_g A_i = A_i$ , then, in particular:  $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} A_i^1 \\ A_i^2 \end{pmatrix} = \begin{pmatrix} A_i^1 \\ A_i^2 \end{pmatrix}$ . This in turn implies, as  $A_i^1 = -A_i^1$  and  $A_i^2 = -A_i^2$  that  $A_i^1 = A_i^2 = 0$ . i.e.  $A_i \equiv 0$ . Thus, any  $\theta_i = \begin{pmatrix} w_i \\ A_i \end{pmatrix} \in \mathcal{E}^G$  has  $A_i = 0$ . Therefore, if we choose any activation  $\sigma$  (e.g the *sigmoid* activation or  $\sigma = \tanh$ , both  $\mathcal{C}^\infty$  and bounded) and we choose  $N \in \mathbb{N}^*$  and  $\theta_i \in \mathcal{E}^G \forall i = 1, ..., N$ ; then:

$$\forall x \in \mathbb{R}^2, \ \Phi_{\theta}^{N,\mathcal{E}^G}(x) = \frac{1}{N} \sum_{i=1}^N W_i \sigma(0^T \cdot x + B_i) = \frac{1}{N} \sum_{i=1}^N W_i \sigma(B_i)$$

which is a constant independent of x. i.e. **any equivariant architecture in this context** is a constant function (whereas  $Y = ||X||^2$  is not). Notice, in particular, that any shallow model  $\Phi_{\nu}$  with  $\nu \in \mathcal{P}(\mathcal{E}^G)$  will also be a constant function. In particular, notice that we will never do "better" than minimizing over all possible constants:

$$\inf_{\substack{\theta_i \in \mathcal{E}^G \\ i=1,\dots,N \\ N \in \mathbb{N}}} R(\Phi_{\theta}^{\mathcal{E}^G}) \ge \inf_{\nu \in \mathcal{P}(\mathcal{E}^G)} R(\nu) \ge \inf_{C \in \mathbb{R}} \mathbb{E}[|Y - C|^2] = \inf_{C \in \mathbb{R}} \mathbb{E}[|\|X\|^2 - C|^2].$$

The problem on the right has a known answer, which is  $C^* = \mathbb{E}_{\pi}[||X||^2] > 0$ . On the other hand, consider a fully conected neuronal network. By the **universal approximation theorem** (which applies for the chosen  $\sigma$ , as in [30, 17, 4]), as  $\pi$  is **compactly supported** (in particular,  $\pi_{\mathcal{X}}(K) = 1$ ); we consider the **parameters** that approximate the function  $f(x) = ||x||^2$  in K = B(0, 1) to precision  $\varepsilon > 0$ . i.e. For  $\varepsilon \in (0, \sqrt{C^*})$ , we know:  $\Longrightarrow \exists N \in \mathbb{N}, \exists a_1, ..., a_N \in \mathbb{R}^2, \exists w_1, ..., w_N \in \mathbb{R}^1$  such that:

$$\|\Phi_{\theta}^{N} - f\|_{\infty,K} = \sup_{x \in K} |\Phi_{\theta}^{N}(x) - f(x)| < \varepsilon < \sqrt{C^{*}}$$

Then:

$$\mathbb{E}[|Y - \Phi^N_{\theta}(x)|^2] \le \mathbb{E}[(\sup_{x \in K} |\Phi^N_{\theta}(x) - f(x)|)^2] < \mathbb{E}[C^*] = C^*$$

But, in particular,  $\exists \nu_{\theta}^{N} \in \mathcal{P}(\mathcal{Z})$  such that:

$$\mathbb{E}[|Y - \Phi^N_\theta(x)|^2] < C^*$$

and so:

$$\inf_{\mu \in \mathcal{P}(\mathcal{Z})} R(\mu) \leq \inf_{\theta \in \mathcal{Z}^N} \mathbb{E}[|Y - \Phi^N_{\theta}(x)|^2] < C^* \leq \inf_{\nu \in \mathcal{P}(\mathcal{E}^G)} R(\nu)$$

In particular, we **can't expect** an optimum of the learning problem to be achieved within  $\mathcal{P}(\mathcal{E}^G)$ .  $\Box$ 

To overcome situations as in Proposition 4, the usual setting is to assume some **universality** condition. This leads to Proposition 5, which we will now prove:

*Proof of Proposition 5.* A standard calculation from the quadratic loss case (see the proof of Corollary 1 for details) yields that, for any  $\mu \in \mathcal{P}(\mathcal{Z})$ :

$$R(\mu) = \mathbb{E}_{\pi}[\|Y - \Phi_{\mu}(X)\|_{\mathcal{Y}}^{2}] = R_{*} + \mathbb{E}_{\pi}[\|f^{*}(X) - \Phi_{\mu}(X)\|_{\mathcal{Y}}^{2}]$$

where  $f^*(x) := \mathbb{E}_{\pi}[Y|X = x]$  and  $R_*$  is the *Bayes risk* of the problem. From Proposition 10, we know that  $f^* \in L^2_G(\mathcal{X}, \mathcal{Y}; \pi|_{\mathcal{X}})$ , and so by universality of  $\mathcal{F}_{\sigma_*}(\mathcal{P}(\mathcal{E}^G))$  onto that space (as well as that of  $\mathcal{F}_{\sigma_*}(\mathcal{P}(\mathcal{Z}))$ ), we conclude directly that:  $\inf_{\nu \in \mathcal{P}(\mathcal{E}^G)} R(\nu) = R_* = \inf_{\mu \in \mathcal{P}(\mathcal{Z})} R(\mu)$ .  $\Box$ 

**Remark.** Works such as [39, 48, 62, 63] precisely provide conditions under which such **universality** condition can be guaranteed. Particularly, as noted in [39], equivariant NNs of tensor order 1 (as are the ones presented in our single-hidden-layer example) are unable to achieve universality for certain types of group actions (see Theorem 2 from [39]). Bridging this gap (eventually allowing for arbitrary order tensors in our MF formulation) is part of our future challenges to make our approach more broadly applicable. In any case, first order universality (which is what we hope to have in our setting) has been established for relevant examples such as Deep Sets ([62, 63]) and CNNs ([62]).

#### C.3 Proofs of results in Section 3.3

Having laid out all the different relevant elements for our work, we can now procede with the proofs of some of our main results.

## C.3.1 Proof of Theorem 3, Corollary 3 and Theorem 4

We start by proving Theorem 3 on the general case.

*Proof of Theorem 3.* We know that a family  $(\mu_t)_{t\geq 0} \subseteq \mathcal{P}_2(\mathcal{Z})$  satisfies **WGF**(*F*) in the weak sense if  $\forall \varphi \in C_c^{\infty}(\mathcal{Z} \times (0,T))$ :

$$\int_0^T \int_{\mathcal{Z}} \left( \partial_t \varphi(z,t) - \langle \varsigma(t) D_\mu F(\mu_t,z), \nabla_z \varphi(z,t) \rangle \right) d\mu_t(z) \, dt = 0$$

Now, profiting from the **uniqueness** of the solutions of this equation, it will be enough to show that, given a solution  $(\mu_t)_{t\geq 0} \subseteq \mathcal{P}_2(\mathcal{Z})$  of **WGF**(F), then  $(\mu_t^G)_{t\geq 0} \subseteq \mathcal{P}_2^G(\mathcal{Z})$  is also a solution. Indeed, consider, for  $g \in G$ ,  $\tilde{\mu}_t = M_g \# \mu_t$ , and notice that for  $\varphi \in C_c^{\infty}(\mathcal{Z} \times (0,T))$ :

$$\begin{split} \int_0^T \int_{\mathcal{Z}} (\partial_t \varphi(z,t) - \langle \varsigma(t) D_\mu F(\tilde{\mu}_t,z), \nabla_z \varphi(z,t) \rangle) d\tilde{\mu}_t(z) \, dt \\ &= \int_0^T \int_{\mathcal{Z}} \left( \partial_t \varphi(M_g.z,t) - \langle \varsigma(t) D_\mu F(M_g \# \mu_t, M_g.z), \nabla_z \varphi(M_g.z,t) \rangle \right) d\mu_t(z) \, dt =: \star \end{split}$$

Now, we can define  $\varphi^g \in C_c^{\infty}(\mathcal{Z} \times (0,T))$  given by  $\forall (z,t) \in \mathcal{Z} \times (0,T) \varphi^g(z,t) = \varphi(M_g.z,t)$ , which satisfies:

$$\partial_t \varphi^g(z,t) = \partial_t \varphi(M_g.z,t) \text{ and } \nabla_z \varphi^g(z,t) = M_g^T \nabla_z \varphi(M_g.z,t)$$

So that, by also using proposition 12 and the orthogonality of the group action, we get:

$$\star = \int_0^T \int_{\mathcal{Z}} \left( \partial_t \varphi^g(z, t) - \langle M_g.\varsigma(t) D_\mu F(\mu_t, z), M_g \nabla_z \varphi^g(z, t) \rangle \right) d\mu_t(z) dt$$
$$= \int_0^T \int_{\mathcal{Z}} \left( \partial_t \varphi^g(z, t) - \langle \varsigma(t) D_\mu F(\mu_t, z), \nabla_z \varphi^g(z, t) \rangle \right) d\mu_t(z) dt = 0$$

Where the last equality comes from the fact that  $(\mu_t)_{t>0}$  is a solution to the WGF.

In particular, as we also have that  $\tilde{\mu}_0 = M_g \# \mu_0 = \mu_0$  (because  $\mu_0 \in \mathcal{P}^G(\mathcal{Z})$ ), by uniqueness we can conclude that this means that  $\forall g \in G, \forall t \in (0, T) \lambda$ -a.e.,  $\mu_t = M_g \# \mu_t$ .

This may seem *weaker* that what we want to prove. Nevertheless, as our group is compact and has a unique normalized Haar measure, we can proceed as follows: let  $f : [0,T] \times \mathbb{Z} \to \mathbb{R}_+$  be any positive and measurable function. Given  $g \in G$ , take  $\Omega_g \subseteq [0,T]$  a full measure set where it holds that  $\mu_t = M_g \# \mu_t$ . In particular,  $f_t = f(t, \cdot) : \mathbb{Z} \to \mathbb{R}$  is positive and measurable, so that:

$$\forall t \in \Omega_g, \ \langle f_t, \mu_t \rangle = \langle f_t, M_g \# \mu_t \rangle = \langle f_t \circ M_g, \mu_t \rangle$$

and we can integrate this equality to get:  $\int_0^T \langle f_t, \mu_t \rangle dt = \int_0^T \langle f_t \circ M_g, \mu_t \rangle dt$  Now, by integrating both sides with respect to the Haar measure, and applying Fubini's theorem (because everything is positive) we get:

$$\int_0^T \langle f_t, \mu_t \rangle dt = \int_G \int_0^T \langle f_t, \mu_t \rangle dt d\lambda_G(g) = \int_G \int_0^T \langle f_t \circ M_g, \mu_t \rangle dt d\lambda_G(g) = \int_0^T \langle f_t, \mu_t^G \rangle dt$$

Implying (by a standard argument) that  $\forall t \in [0,T]$  a.e.  $\mu_t = \mu_t^G$ , and therefore:  $\forall t \in [0,T]$  a.e.  $\mu_t \in \mathcal{P}^G(\mathcal{Z})$ .

*Proof of Corollary 3.* From Corollary 2, Corollary 6 and Corollary 8, we know that  $R^{\tau,\beta}$  is invariant.

On the other hand, from [31] (or [57]) we know that, under our assumptions, a **unique** weak solution to the Fokker-Planck equation exists. Furthermore, this solution is known to be strong if  $\beta > 0$ . In particular, theorem 3 applies and allows us to conclude that if  $\mu_0 \in \mathcal{P}_2^G(\mathcal{Z})$ , then  $\forall t \ge 0$  (a.e.)  $\mu_t \in \mathcal{P}_2^G(\mathcal{Z})$ .

When  $\beta > 0$ , since solutions are **strong**, we conclude that the densities  $(u_t)_{t\geq 0}$  are all *G*-invariant functions ( $\lambda$ -a.e.). This follows from the remark about *densities of invariant measures* provided in SuppMat-A.2.

**Remark.** When  $\beta > 0$ , we have a unique weakly-invariant minimizer (from proposition 3 and/or corollary 8); and also, under mild assumptions, a **global convergence** result. That is, independently of the network's initialization, we will converge to the *G*-invariant solution. An interesting question in this setting is then: At which point does the **WGF** enter the space  $\mathcal{P}_2^G(\mathcal{Z})$ ?

We can also prove Theorem 4:

*Proof of Theorem 4.* This proof follows from the fact that  $\forall z \in \mathcal{Z}, \forall \mu \in \mathcal{P}^G(\mathcal{Z})$ :

$$D_{\mu}R^{DA}(\mu, z) = D_{\mu}R^{FA}(\mu, z).$$

Indeed, notice that, from proposition 17:

$$D_{\mu}R^{FA}(\mu,z) = \int_{G} M_{g}^{T} \cdot D_{\mu}R(\mu^{G}, M_{g}.z)d\lambda_{G}(g) = \int_{G} M_{g}^{T} \cdot D_{\mu}R(\mu, M_{g}.z)d\lambda_{G}(g),$$

while also:

$$D_{\mu}R^{DA}(\mu, z) = \int_{G} M_{g}^{T} \cdot D_{\mu}R(M_{g} \# \mu, M_{g} \cdot z)d\lambda_{G}(g) = \int_{G} M_{g}^{T} \cdot D_{\mu}R(\mu, M_{g} \cdot z)d\lambda_{G}(g)$$

Now, let  $(\mu_t^{FA})_{t\geq 0}$  and  $(\mu_t^{DA})_{t\geq 0}$  be the WGF solutions starting from  $\mu_0$  for  $R^{FA}$  and  $R^{DA}$  respectively. As  $R^{FA}$  is *G*-invariant, by corollary 3,  $(a.e.)\forall t\geq 0$ ,  $\mu_t^{FA}\in \mathcal{P}^G(\mathcal{Z})$ . Now, let's see that this process actually **also satisfies WGF** $(R^{DA})$ , forcing both processes to coincide by uniqueness.

Indeed, we know that  $(\mu_t^{FA})_{t\geq 0}$  satisfies:  $\forall \varphi \in C_c^{\infty}(\mathcal{Z} \times (0,T))$ :

$$\int_0^T \int_{\mathcal{Z}} \left( \partial_t \varphi(z,t) - \langle \varsigma(t) D_\mu R^{FA}(\mu_t^{FA},z), \nabla_z \varphi(z,t) \rangle \right) d\mu_t^{FA}(z) \, dt = 0$$

Now, as  $(a.e.)\forall t \ge 0$ ,  $\mu_t^{FA} \in \mathcal{P}^G(\mathcal{Z})$ , we have  $\forall z \in \mathcal{Z}$ :  $D_\mu R^{FA}(\mu_t^{FA}, z) = D_\mu R^{DA}(\mu_t^{FA}, z)$ . In particular,  $(\mu_t^{FA})_{t\ge 0}$  satisfies  $\forall \varphi \in C_c^{\infty}(\mathcal{Z} \times (0, T))$ :

$$\int_0^T \int_{\mathcal{Z}} \left( \partial_t \varphi(z,t) - \langle \varsigma(t) D_\mu R^{DA}(\mu_t^{FA},z), \nabla_z \varphi(z,t) \rangle \right) d\mu_t^{FA}(z) \, dt = 0$$

Implying that  $(\mu_t^{FA})_{t\geq 0}$  solves **WGF** $(R^{DA})$  starting from  $\mu_0$ ; thus by uniqueness:  $(\mu_t^{FA})_{t\geq 0} = (\mu_t^{DA})_{t\geq 0}$ .

The last part of the theorem comes from Proposition 2, since if R is invariant, its WGF will exactly coincide with that of  $R^{DA}$  (they are the same functional).

**Remark.** This results tells us the ultimate bottom line: at the MF level, training with **DA** or **FA** results in the exact same dynamic. Furthermore, whenever data is equivariant, they are both essentially equivalent to **applying no technique whatsoever**. This result, despite the obvious limitation of considering infinitely wide NNs, still provides some meaningful practical insights (as shown Appendix D) for large NNs, that could be used in applications.

**Remark.** Since  $R^{DA}$ ,  $R^{FA}$  and R all coincide on  $\mathcal{P}^G(\mathcal{Z})$ , one could expect Theorem 4 to hold for R even without assuming  $\pi$  to be equivariant. However, the invariance of R is **crucial** for such a result: if R isn't invariant, nothing guarantees that its WGF process will stay within  $\mathcal{P}_G(\mathcal{Z})$ , whereas  $WGF(R^{DA})$  and  $WGF(R^{FA})$  always do so.

#### C.3.2 Proof of Theorem 5 and Theorem 6

We can now provide the proof for the *stronger result*, Theorem 5, stating that the WGF of an invariant functional will *respect*  $\mathcal{E}^G$  all along training. This proof uses the **McKean-Vlasov non-linear SDE** (Equation (5)) presented in Appendix B.2 (see [19] for a reference). Namely, we will consider the following **projected McKean-Vlasov SDE** (with  $(B_t)_{t>0}$  a BM on  $\mathcal{Z}$ ), given by:

$$dZ_t = \varsigma(t) \left[ -\left( D_\mu R(\mu_t, \cdot) + \tau \nabla_\theta r(Z_t) \right) dt + \sqrt{2\beta} P_{\mathcal{E}^G} dB_t \right] \text{ with } \operatorname{Law}(Z_t) = \mu_t,$$
 (6)

which corresponds to the MF limit dynamics arising from performing the **projected noisy SGD** scheme from Equation (4). Namely, it corresponds to the WGF of  $R_{\mathcal{E}^G}^{\tau,\beta}$  presented in Section 3.3.

*Proof of Theorem 5.* Consider the (**pathwise unique**) solution of the **projected McKean-Vlasov SDE** (6),  $Z = (Z_t)_{t \ge 0}$ . This means that it satisfies:

$$Z_t \stackrel{a.s.}{=} Z_0 - \int_0^t \varsigma(s) D_\mu R^\tau(\mu_s, Z_s) ds + \sqrt{2\beta} \int_0^t \varsigma(s) P_{\mathcal{E}^G} dB_s, \text{ and } Z_0 = \xi_0 \text{ (initial condition)}$$
(7)

With  $\xi_0$  such that  $Law(\xi_0) = \mu^0$ , and  $R^{\tau} := R + \langle r, \cdot \rangle$  being used as shorthand notation.

We first let  $g \in G$  be an arbitrary group element, and we study how the process  $\tilde{Z} = (\tilde{Z}_t)_{t \ge 0} := (M_g Z_t)_{t \ge 0}$  satisfies this same equation (7).

Denote  $\nu_s := M_g \# \mu_s$  as the law of  $\tilde{Z}_s$ , we want to show that for all  $t \ge 0$ :

$$\tilde{Z}_t \stackrel{a.s.}{=} \tilde{Z}_0 - \int_0^t \varsigma(s) D_\mu R^\tau(\nu_s, \tilde{Z}_s) ds + \sqrt{2\beta} \int_0^t \varsigma(s) P_{\mathcal{E}^G} dB_s \tag{8}$$

Indeed, first notice that:

- 1. Let  $\Omega$  be the full measure set where  $\xi_0 \in \mathcal{E}^G$  (which we can do since  $\mu_0 \in \mathcal{P}(\mathcal{E}^G)$ , or, equivalently:  $\mathbb{P}(\xi_0 \in \mathcal{E}^G) = 1$ ). Then,  $\forall \omega \in \Omega$ ,  $Z_0(\omega) = \xi_0(\omega) \in \mathcal{E}^G$ . In particular,  $\forall \omega \in \Omega$ ,  $\forall g \in G, \tilde{Z}_0(\omega) = M_g Z_0(\omega) = M_g \xi_0(\omega) = \xi_0(\omega) = Z_0(\omega)$ . That is,  $\tilde{Z}_0 \stackrel{a.s.}{=} Z_0$ .
- 2. Now, the equation is satisfied by  $(Z_t)_{t>0}$  and therefore, for  $t \ge 0$ , we have:

$$\begin{split} \tilde{Z}_t &= M_g Z_t = M_g Z_0 - M_g \left( \int_0^t \varsigma(s) D_\mu R^\tau(\mu_s, Z_s) ds \right) + \sqrt{2\beta} M_g \int_0^t \varsigma(s) P_{\mathcal{E}^G} dB_s \\ &= \tilde{Z}_0 - \int_0^t \varsigma(s) M_g D_\mu R^\tau(\mu_s, Z_s)) ds + \sqrt{2\beta} \int_0^t \varsigma(s) M_g. P_{\mathcal{E}^G} dB_s \\ &= \tilde{Z}_0 - \int_0^t \varsigma(s) D_\mu R^\tau(M_g \# \mu_s, M_g. Z_s) ds + \sqrt{2\beta} \int_0^t \varsigma(s) P_{\mathcal{E}^G} dB_s \\ &= \tilde{Z}_0 - \int_0^t \varsigma(s) D_\mu R^\tau(\nu_s, \tilde{Z}_s) ds + \sqrt{2\beta} \int_0^t \varsigma(s) P_{\mathcal{E}^G} dB_s \end{split}$$

Here, we used the linearity of the integral (and the stochastic integral), the fact that  $\forall g \in G$ ,  $M_g P_{\mathcal{E}^G} = P_{\mathcal{E}^G}$ , and Proposition 12, which holds for  $\forall \theta \in \mathcal{Z}, \forall \mu \in \mathcal{P}(\mathcal{Z})$  (in particular for  $\theta = Z_s(\omega)$ ,  $\forall \omega \in \Omega$  and  $\mu_s = \text{Law}(Z_s)$ ). Thus,  $\forall g \in G$ , (8) holds.

By the **pathwise uniqueness** of the solution  $(Z_t)_{t\geq 0}$ , we have (following, for instance, [23]):

$$\mathbb{P}\left(\sup_{t\geq 0} \|Z_t - \tilde{Z}_t\| = 0\right) = 1$$

In particular, as  $g \in G$  was arbitrary, we have that:

$$\forall g \in G, \ \sup_{t \ge 0} \|Z_t - M_g Z_t\| \stackrel{a.s.}{=} 0 \tag{9}$$

We now want to be able to **interchange** the  $\forall g \in G$  with the probability measure. Fortunately, we are dealing with a compact group with a normalized Haar measure  $\lambda_G$ . Indeed, from equation (9) we deduce that  $\forall g \in G, \forall t \geq 0, \mathbb{P}(||Z_t - M_g Z_t|| = 0) = 1$ .

Now, notice that, for any  $t \ge 0$  and  $\omega \in \Omega$ :

$$\left\|Z_t(\omega) - P_{\mathcal{E}^G} Z_t(\omega)\right\| = \left\|Z_t(\omega) - \int_G M_g Z_t(\omega) d\lambda_G(g)\right\| \le \int_G \left\|Z_t(\omega) - M_g Z_t(\omega)\right\| d\lambda_G(g)$$

We can integrate both sides by  $\mathbb{P}$  to get (using Fubini as functions are positive and measurable):

$$0 \leq \int_{\Omega} \|Z_t(\omega) - P_{\mathcal{E}^G} Z_t(\omega)\| d\mathbb{P}(\omega) \leq \int_{\Omega} \int_{G} \|Z_t(\omega) - M_g Z_t(\omega)\| d\lambda_G(g) d\mathbb{P}(\omega)$$
$$\leq \int_{G} \int_{\Omega} \|Z_t(\omega) - M_g Z_t(\omega)\| d\mathbb{P}(\omega) d\lambda_G(g) = 0$$

where in the last step we have used the fact that  $\forall g \in G, \ \forall t \geq 0, \ \mathbb{P}(\|Z_t - M_g Z_t\| = 0) = 1$ , so that  $\forall t \geq 0, \ \forall g \in G, \ \int_{\Omega} \|Z_t(\omega) - M_g Z_t(\omega)\| \ d\mathbb{P}(\omega) = 0$ .

This implies that  $\forall t \geq 0 \mathbb{P}$ -a.s.  $Z_t = P_{\mathcal{E}^G} Z_t$ , i.e.  $\mathbb{P}(Z_t \in \mathcal{E}^G) = \mu_t(\mathcal{E}^G) = 1$ , or, in other words,  $\forall t \geq 0, \ \mu_t \in \mathcal{P}(\mathcal{E}^G)$  as required.  $\Box$ 

**Remark.** Notice that, by a.s. continuity of the McKean-Vlasov diffusion (6) and the fact that  $\mathcal{E}^G$  is closed, the previous proof actually shows that  $\mathbb{P}(Z_t \in \mathcal{E}^G, \forall t \ge 0) = 1$ .

Notice also that Theorem 5 bears some resemblance to *Corollary 1* in [25], which states that  $\mathcal{E}^G$  is *stable* under the traditional gradient flow of the *augmented risk* ( $[\theta \in \mathbb{Z}^N \mapsto R^{DA}(\theta) \in \mathbb{R}]$ , as in Section 2.3). Our result shares a similar flavor, but for the MF dynamics of freely-trained NNs with equivariant data.

**Remark.** Unlike with **WI** distributions, initializing a shallow NN with  $\mu_0 \in \mathcal{P}(\mathcal{E}^G)$  isn't as straightforward as using a normal distribution. Effectively (and efficiently) computing the space  $\mathcal{E}^G$  is actually quite challenging (as noted in [24]).

A natural way to ensure that  $\mu_0 \in \mathcal{P}(\mathcal{E}^G)$ , independently of the form of  $\mathcal{E}^G$ , is to initialize all parameters to be 0. The question of whether under such initialization the parameters will eventually exit {0} (or some larger, strict subspace  $E \subsetneq \mathcal{E}^G$ ) and find values over the entire space  $\mathcal{E}^G$  is left for future work. Some insights on this behaviour can be sought in our experimental results, see Section 4. If true, this behavior could point towards some type of underlying hypoellipticity of the McKean-Vlasov dynamics (6) (or variants) on  $\mathcal{E}^G$ , which would be interesting to analyze, in particular in view of potential theoretical guarantees for architecture-discovering heuristics as suggested in Section 4.2.

Note that there is no need for seeing  $\mathcal{E}^G$  as a subspace of an ambient space  $\mathcal{Z}$ . When training with **EAs**, we simply *force* our parameters to *live on*  $\mathcal{E}^G$ , since we fix the architechture beforehand. Namely, our "whole space" is  $\tilde{\mathcal{Z}} = \mathcal{E}^G$  (regarded directly as a vector space  $\mathcal{E}^G \cong \mathbb{R}^{\tilde{D}}$ ) rather than  $\mathcal{Z}$ . Thus, the relevant *population risk* is the *restricted* version of the original:  $\tilde{R} := R|_{\mathcal{P}(\tilde{\mathcal{Z}})} : \mathcal{P}(\tilde{\mathcal{Z}}) \to \mathbb{R}$ ; and we can apply the usual results from the MF Theory when the relevant hypothesis are satisfied by  $\tilde{\mathcal{Z}}$  and  $\tilde{R}$ . Notably, we can have global convergence of  $\tilde{R}^{\tau,\beta}$  to  $\inf_{\mu \in \mathcal{P}(\tilde{\mathcal{Z}})} \tilde{R}^{\tau,\beta}(\mu) = \inf_{\mu \in \mathcal{P}(\mathcal{E}^G)} R^{\tau,\beta}_{\mathcal{E}^G}(\mu)$ 

**Remark.** As shown in [31] (see Proposition 18 in Appendix E for the details) the regularized versions of the involved functionals (i.e.  $R^{\tau,\beta}$  and  $R^{\tau,\beta}_{\mathcal{E}^G}$ )  $\Gamma$ -converge to the original R as  $\tau, \beta \to 0$ ; meaning that, for small values of the regularization parameters, we should expect the achieved optima to ressemble  $\inf_{\mu \in \mathcal{P}^G(\mathcal{Z})} R(\mu)$  and  $\inf_{\nu \in \mathcal{P}(\mathcal{E}^G)} R(\nu)$  respectively (or, under Proposition 5, both to  $R_*$ ). We will also leave the exploration of how this approximation behaves as future work.

Finally, we provide a proof for Theorem 6

*Proof of Theorem 6.* The proof structure is very similar to that of Theorem 4. Namely, it comes from noticing that for  $\mu \in \mathcal{P}(\mathcal{E}^G)$  and  $z \in \mathcal{E}^G$ , we have:

$$D_\mu R^{DA}(\mu,z) = D_\mu R^{FA}(\mu,z) = D_\mu R^{EA}(\mu,z). \label{eq:D_matrix}$$

We already know the first equality, as seen in Theorem 4 (since  $\mu \in \mathcal{P}^G(\mathcal{Z})$  from lemma 1). We only need to show the last equality. Indeed, notice that:

$$D_{\mu}R^{FA}(\mu,z) = \int_{G} M_{g}^{T} D_{\mu}R(\mu,z)d\lambda_{G}(g) = P_{\mathcal{E}^{G}} D_{\mu}R(\mu,z),$$

while also, from Proposition 17:

$$D_{\mu}R^{\mathcal{E}A}(\mu,z) = P_{\mathcal{E}^G}^*.D_{\mu}R(\mu^{\mathcal{E}^G},P_{\mathcal{E}^G}.z) = P_{\mathcal{E}^G}.D_{\mu}R(\mu,z).$$

Knowing this, the rest of the proof is analogous to that of Theorem 4. Let  $(\mu_t^{FA})_{t\geq 0}$ ,  $(\mu_t^{DA})_{t\geq 0}$ and  $(\mu_t^{EA})_{t\geq 0}$  be the WGF solutions starting from  $\mu_0$  for  $R^{FA}$ ,  $R^{DA}$  and  $R^{EA}$  respectively. Since  $\mu_0 \in \mathcal{P}(\mathcal{E}^G) \subseteq \mathcal{P}^G(\mathcal{Z})$ , from Theorem 4 we know that  $(\mu_t^{FA})_{t\geq 0}$  and  $(\mu_t^{DA})_{t\geq 0}$  coincide. Let's see that, w.l.o.g.,  $(\mu_t^{FA})_{t\geq 0}$  coincides with  $(\mu_t^{EA})_{t\geq 0}$ .

As  $R^{FA}$  is *G*-invariant, by theorem 5,  $\forall t \geq 0$ ,  $\mu_t^{FA} \in \mathcal{P}(\mathcal{E}^G)$ . Now, let's see that this process also satisfies WGF( $R^{EA}$ ), forcing both processes to coincide by uniqueness.

As before, we know that  $(\mu_t^{FA})_{t\geq 0}$  satisfies:  $\forall \varphi \in C_c^{\infty}(\mathcal{Z} \times (0,T))$ :

$$\int_0^T \int_{\mathcal{Z}} \left( \partial_t \varphi(z,t) - \langle \varsigma(t) D_\mu R^{FA}(\mu_t^{FA},z), \nabla_z \varphi(z,t) \rangle \right) d\mu_t^{FA}(z) \, dt = 0$$

Now, as  $\forall t \geq 0$ ,  $\mu_t^{FA} \in \mathcal{P}(\mathcal{E}^G)$ , we can restrict our integral to  $\mathcal{E}^G$ . Also, we have  $\forall z \in \mathcal{E}^G$ :  $D_{\mu}R^{FA}(\mu_t^{FA}, z) = D_{\mu}R^{EA}(\mu_t^{FA}, z)$ . With these properties,  $(\mu_t^{FA})_{t\geq 0}$  satisfies  $\forall \varphi \in C_c^{\infty}(\mathcal{Z} \times (0,T))$ :

$$\int_0^T \int_{\mathcal{E}^G} \left( \partial_t \varphi(z,t) - \langle \varsigma(t) D_\mu R^{EA}(\mu_t^{FA},z), \nabla_z \varphi(z,t) \rangle \right) d\mu_t^{FA}(z) \, dt = 0$$

Making the integral over  $\mathcal{Z}$  once again (we can since  $\mu_t^{FA} \in \mathcal{P}(\mathcal{E}^G)$  for all  $t \geq 0$ ), we get that  $(\mu_t^{FA})_{t\geq 0}$  solves  $\mathbf{WGF}(\mathbb{R}^{EA})$  starting from  $\mu_0$ ; thus by uniqueness:  $(\mu_t^{FA})_{t\geq 0} = (\mu_t^{EA})_{t\geq 0}$ .

The last part of the theorem comes, once again, from Proposition 2, since if R is invariant, its WGF exactly coincides with that of  $R^{DA}$ .

## **Experimental setting and further experiments** D

All the different experiments were run on Python 3.10, on a Google Colab session consisting (by default) of 2 Intel Xeon virtual CPUs (2.20GHz) and with 13GB of RAM.

In order to obtain results that can be visualized, we consider a simple setting where  $\mathcal{X} = \mathcal{Y} = \mathbb{R}^2$ and  $\mathcal{Z} = \mathbb{R}^{2 \times 2} \cong \mathbb{R}^4$ . We let  $G = C_2$  acting on  $\mathcal{X}$  and  $\mathcal{Y}$  via the coordinate transposition action (i.e. the group generated by the orthogonal matrix  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ ); and on  $\mathcal{Z}$  via the natural intertwining action (i.e.  $M_g \cdot z = \hat{\rho}_g \cdot z \cdot \rho_q^T$ ). We also consider the jointly equivariant activation given by  $\sigma_*(x,z) = \sigma(z \cdot x) \ \forall x \in \mathbb{R}^2, \ \forall z \in \mathbb{R}^{2 \times 2} \ \text{with } \sigma : \mathbb{R} \to \mathbb{R} \ \text{a sigmoidal activation function}$ (which is  $\mathcal{C}^{\infty}$  and bounded) applied pointwise. Under this setting,  $\mathcal{E}^G$  can be explicitly computed as  $\mathcal{E}^G = \left\langle \begin{pmatrix} \frac{1}{\sqrt{2}} & 0\\ 0 & \frac{1}{\sqrt{2}} \end{pmatrix}, \begin{pmatrix} 0 & \frac{1}{\sqrt{2}}\\ \frac{1}{\sqrt{2}} & 0 \end{pmatrix} \right\rangle$ , which is a 2-dimensional subspace of the ambient 4-dimensional space. It's projection operator  $P_{\mathcal{E}^G}$  is also explicitly known.

We consider a **teacher** model  $f_* = \Phi_{\theta^*}^{N_*}$  with  $N_* = 5$  fixed particles, such that  $\nu_{\theta^*}^{N_*}$  is either **arbitrary**, **WI** or **SI**. Let  $\vartheta = 0.5$  be a scale parameter. The **arbitrary** particles were chosen to be:

$$\theta_1^* = \vartheta.(-1, 0, 0, 0.5)^T$$
  

$$\theta_2^* = \vartheta.(0.5, 1, 0, 1)^T$$
  

$$\theta_3^* = \vartheta.(-0.5, 0.3, 1, 0)^T$$
  

$$\theta_4^* = \vartheta.(0, -1, -0.5, 1)^T$$
  

$$\theta_5^* = \vartheta.(0.7, -0.7, 0.5, 0.7)^T$$

This was fixed in order to make the task non-trivial and interesting. The WI teacher distribution was simply chosen to be  $(\nu_{\theta^*}^{N_*})^G$ , with  $\theta^*$  as just described. The **SI** particles were also fixed, but their chosen coordinates had to be expressed in terms of the basis vector of  $\mathcal{E}^{G}$  (i.e. only providing 2 parameters). Particularly, they were fixed to be (denoting then by  $a^* = (a_i^*)_{i=1}^{N_*}$  to avoid confusion):

$$a_1^* = \vartheta.(1,0)^T, \ a_2^* = \vartheta.(0.5,1)^T, \ a_3^* = \vartheta.(-0.5,0.3), \ a_4^* = \vartheta.(0,-1), \ a_5^* = \vartheta.(0.7,0.7).$$

As seen on Section 3.1, the teacher  $f_*: \mathcal{X} \to \mathcal{Y}$  will be an equivariant function as soon as  $\nu_{\theta^*}^{N_*}$  is chosen either **WI** or **SI**. Our data distribution  $\pi$  will be such that  $(X, Y) \sim \pi$  will satisfy  $X \sim \mathcal{N}(0, \sigma_{\pi}^2.\mathrm{Id}_2)$  (with  $\sigma_{\pi} = 4$ ), and  $Y = f_*(X)$ . Namely,  $\pi_{\mathcal{X}}$  will always be *G*-invariant, whereas  $\pi$  will only be *G*-invariant if  $f_*$  is. This setting allows for testing the different results provided, without losing the properties of  $\mathcal{Q}_G$  as a projection (which require  $\pi_{\mathcal{X}}$  to be G-invariant, as shown in [22]).

We will try to **mimic** the teacher network by using **student networks**, which will be given by  $\Phi_{\theta}^{\mathcal{H}}$ ; namely, with the same  $\sigma_*$ , but varying values of N and  $\theta \in \mathbb{Z}^N$ . We will train them to minimize the regularized population risk  $R^{\tau,\beta}$  given by a quadratic loss,  $\ell(y,\hat{y}) = ||y - \hat{y}||_{\mathcal{Y}}^2$ , and a quadratic **penalization**,  $r(z) = ||z||_{\mathcal{Z}}^2$ . For this purpose, we employ a *minibatch* variant of the SGD training scheme provided in Equation (1) (possibly projected, as in Equation (4), when required). We will also employ the different symmetry-leveraging techniques presented in Section 2.3, such as DA, FA and EA. We refer to the *free training* with no SL-techniques whatsoever as the **vanilla** training.

The training parameters were fixed to be (unless explicitly stated otherwise):

- Step Size:  $\varsigma \equiv \alpha > 0$  (with  $\alpha = 50$  in most experiments),  $\varepsilon_N = \frac{1}{N}$ , so that  $s_k^N = \frac{\alpha}{N}$ . This was convenient, since it corresponds to the usual implementation of SGD on most common NN frameworks in Python (namely, **pytorch** and **jax**).
- Regularization parameters:  $\tau = 10^{-4}$  and  $\beta = 10^{-6}$ .
- Batch Size: It was chosen to be B = 20.
- Number of Training Epochs: In line with the statement of Theorem 1, to observe phenomena at a MF scale, we need an amount of iterations (commonly known as *epochs* in the ML literature) that is proportional to the number of particles. For this purpose, we fix an

observation time horizon of T = 20. All training schemes were performed for a total of  $N_e = N \cdot T$  epochs (iterations). An additional "granularity" parameter (usually set to be gr = 5) is introduced to determine how often in the dynamic we will observe and save the training losses and particle positions: we do so every  $\lfloor \frac{N_e}{gr} \rfloor$  steps. Notice that  $N_e$  depends on N, and so models with different values of N were trained for a different amount of epochs.

- Student Initialization: The student's particles, θ ∈ Z<sup>N</sup>, are initialized i.i.d. from some μ<sub>0</sub> that is chosen to be either WI or SI. When WI-initialized, they are sampled from a random gaussian Z ~ N(0, 1/16). When SI-initialized, particles are taken to be P<sub>E<sup>G</sup></sub>.Z with Z as before.
- Number of Repetitions: Each experiment was repeated a total of  $N_r = 10$  times to ensure consistency. Each repetition, a different random *seed* was employed to: initialize the student's particles, generate the training data, and generate the noise for the SGD iteration. In particular, on a fixed repetition, all models were trained with the **same data** and the **same noise** being applied on SGD updates.

**Remark.** As  $N_e$  is chosen to be proportional to the number of particles, N, computational burden and memory requirements quickly became heavy for the simple machines we employed (which didn't even have a dedicated GPU). This is the reason why we don't scale our experiments beyond the N = 5000 case. As reference, for N = 5000 a single training (with the above hyperparameters) of a single model (either of vanilla, DA, FA or EA) took  $\approx 15$  minutes (which quickly amounts to large amounts of running time for the  $N_r = 10$  repetitions, the 4 different training schemes and the 6 possible settings with WI or SI initialization and arbitrary, WI or SI teacher). This is a clear point of improvement and shall be tackled in future work.

**Remark.** As here mentioned, on every fixed "repetition" of the experiments, the same noise was used during the SGD training iterations for the **vanilla**, **DA** and **FA** schemes. The **EA** scheme, despite using the same seeds for the data and student initializations, didn't have the same noise applied during SGD. This was because, despite using the same seed for the noise generator, noise for our EAs was only 2-dimensional (since **EAs** are parametrized by  $\mathcal{E}^G$ ), while it was 4-dimensional for the other schemes. This made the resulting training schemes have an additional layer of noise separating them; and so solving this issue, in order to properly visualize Theorem 6, becomes fundamental.

**Remark.** Notice that the  $N_r = 10$  performed repetitions were largely enough to allow for plots with error bars (actually, we do boxplots which encode the variability of the different quantities more precisely) that allow for significant analysis of the observed phenomena. We do not go beyond  $N_r = 10$  due to the low computational capabilities of our machines, and the already high computational cost of running the experiments (for thousands of hidden units and epochs, in many different settings, and involving the calculation of Wasserstein Distances, as we'll comment below).

To facilitate the implementation of the ideas behind our **EA** models, we use the *group* and *representations* tools from the **emlp** repository provided as part of [24]. This code is openly available and has an MIT License for unrestricted access. We employ it to numerically (and efficiently) determine the space  $\mathcal{E}^G$  (namely, its basis), as well as  $P_{\mathcal{E}^G}$ . We do remark that these calculations were correct only up to a precision of  $10^{-8}$ , which results in a slight burden for our empirical results. On the other hand, regarding the implementation of **EMLP**s provided in the package (**EAs** in our setting), some slight modifications to the source code had to be performed in order to correctly represent our setting.

On a similar note, we can numerically compute the squared Wasserstein-2 distance between two empirical distributions of particles by employing the **pyot** library. This allows us to evaluate *to what extent* our resulting models are close to each other in terms of their particle distribution  $\nu_{\theta}^{N}$  (which is what the **MF** approach suggests). In order to fix a common scale in which the experiments can be compared, for different values of N, and mitigate the effects of fluctuating empirical estimates (mainly for small values of N, and in the low dimensions considered), we consider a natural normalization of the Wasserstein-2 distance, which we refer to as the **RMD** (Relative-Measure-Distance). This is defined as:  $\mathbf{RMD}^2(\mu,\nu) = \frac{W_2^2(\mu,\nu)}{M_{\mu}^2 + M_{\nu}^2}$  where  $M_{\mu}^2 = 2\mathbb{E}[||Z||^2]$  for  $Z \sim \mu$  (so that  $0 \leq \mathbf{RMD} \leq 1$ ). The **RMD** provided a good metric for the experiments here presented. Notice that, as  $N \to \infty$ , by the L.L.N. for empirical distribution following from the **MFL** convergence, the **RMD** is expected to stabilize at the corresponding value of the limiting distributions. Therefore, up to a multiplicative quantity approaching a (finite, non null, in our case) constant, we are observing the behavior of the Wasserstein-2 distance. A drawback from using the Wasserstein-2 metric, is that calculating them




Figure 3: **RMD** comparisons between training regimes, for different values of N, at the end of an **SI**-initialized training for  $N_e$  epochs. Each column corresponds to a teacher with, respectively, an **arbitrary**, **WI** and **SI** distribution. Row 1 displays  $\text{RMD}^2(\nu_{N_e}^N, (\nu_{N_e}^N)^{\mathcal{E}^G})$  for the different regimes, in order to evaluate to what extent the training remained within  $\mathcal{E}^G$ . Row 2 displays the **RMD** between **DA**, **FA** and **vanilla** training regimes; and Row 3 does the same for each of them against **EA**.

can be very expensive computationally. This is another one of the reasons why our experiments only get to N = 5000 particles.

## **D.1** Study for varying N

Beyond the analysis already provided in Section 4, we here provide some meaningful insights. We want to observe to what extent the properties proved in Section 3.3 for the **WGF** of  $R^{\tau,\beta}$  can be observed in practice. For this purpose, we observe, for  $N \in \{5, 10, 50, 100, 500, 1000, 5000\}$ , different relevant quantities to evaluate the different combinations of *teachers* and *students*.

For this set of experiments, **WI**-initialized students were trained with the usual **SGD** scheme from Equation (1); while, **SI**-initialized students, were trained with the **projected SGD dynamics** from Equation (4). Models for the different schemes to be compared, are all initialized with the exact same (random) particles.

Figure 3 displays some comparisons between particle distributions in terms of **RMD** at the end of training, knowing that they were all initialized with the same particles drawn from  $\mu_0 \in \mathcal{P}(\mathcal{E}^G)$ . From this, we can visually see that, when the teacher distribution is either **WI** of **SI**, the resulting distribution from vanilla training *stays* on  $\mathcal{E}^G$  (since **RMD**<sup>2</sup> $(\nu_{NT}^N, (\nu_{NT}^N)^{\mathcal{E}^G})$  is small) increasingly more as N becomes large. This fact is absolutely remarkable, since, for a **WI** teacher there should be no reason why the **vanilla** training (that's completely *free*, in principle) *shouldn't escape*  $\mathcal{E}^G$  to achieve a better approximation of  $f_*$ . On the other hand, in every single teacher setting, almost independently of N, both **DA** and **FA** consistently remain within  $\mathcal{E}^G$  (as expected, even if  $f_*$  isn't equivariant). For an **arbitrary teacher**, we see that the vanilla training distribution readily *leaves*  $\mathcal{E}^G$ 

to better approximate  $f_*$ , which isn't a *predicted behaviour* from our theory (we have no guarantees of "leaving  $\mathcal{E}^G$  when data isn't equivariant"), but motivates the heuristic defined in Section 4.

Still from Figure 3, we can see that, as N grows bigger, the end-of-training distribution of the **vanilla** scheme becomes closer and closer to that of **DA** and **FA** (from Theorem 4 we actually expect them to be equal in the limit). A similar result is obtained relating **vanilla**, **DA** and **FA** to the **EA** scheme; the values are however larger than before and *less significantly close* in general. This is possibly due to the *different noises* employed during training (as mentioned in a remark above). We do however notice that for increasing values of N, the **EA**, **DA**, **FA** and **vanilla** schemes (the latter only under equivariant  $f_*$ ) tend towards coinciding, which serves to illustrate the constatations from Theorem 6. Finally, notice that the results obtained for **WI** and **SI** teachers present almost no quantitative differences whatsoever between them.

In Figure 4 we present a visualization of the final particle distribution, after an **SI**-initialized training under a **WI** teacher  $f_*$ , of the **vanilla**, **DA**, **FA** and **EA** schemes (on a single realization of the experiment). At least visually (and macroscopically), it seems like all these regimes followed (approximately) the same flow, as they end up with an approximately equal particle distribution. This isn't a rigorous comparison at all, and providing better quantitative comparisons between the methods is to be considered for future work. As a counterfactual, we provide in Figure 5 the results of an **SI**-initialized training that is performed under a non-equivariant  $f_*$ . We can see that the vanilla model readily *leaves*  $\mathcal{E}^G$  to achieve a better approximation of  $f_*$ , while the **DA**, **FA** and **EA** schemes "stay inside" (roughly coinciding between them as well).

Figure 6 also displays **RMD** comparisons between particle distributions at the end of training, but for an (identic) initialization with particles drawn from  $\mu_0 \in \mathcal{P}^G(\mathcal{Z})$ . Now, unlike the **SI** case, with particles sampled *i.i.d.* from a **WI** distribution, nothing ensures that the resulting  $\nu_0^N$  will be **WI** as well. Namely, in this case the limit as  $N \to \infty$  becomes significantly more important to visualize the theoretical results. Indeed, since  $\nu_0^N$  isn't necessarily **WI**, we no longer have a guarantee that the finite-N networks trained with **DA**, **FA** or **vanilla** methods will be close to each other in any sense. We do however notice on Figure 6 that, for increasing N, the end-of-training distributions of **DA**, **FA** and **vanilla** schemes (the latter only when  $f_*$  is equivariant) become increasingly closer to their symmetrized versions (namely,  $\mathbf{RMD}^2(\nu_{NT}^N, (\nu_{NT}^N)^G)$  becomes smaller, though never as small as in the **SI**-initialized experiments). Also, as guaranteed by Theorem 4, for large N we see that **DA** and **FA** become indistinguishably close, no matter the teacher's properties; also, when  $f_*$  is equivariant, they both "coincide" with the **vanilla** scheme. A comparison between the **EA** scheme and the result from projecting the **FA** scheme on the last step is also presented. It is used simply to illustrate that, in principle, directly training on  $\mathcal{E}^G$  isn't necessarily comparable to performing "free-training" and *projecting* the resulting particle distribution only on the last step (even when an SL technique such as **FA** is used).

Now, beyond the analysis of the underlying particle distributions after training, we turn our focus to comparisons of the resulting **models**. We measure some distances in  $L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})$ , by approximating  $\|\cdot\|_{L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})}$  with a Monte-Carlo sample of 100 random data points drawn from  $\pi$ .

Figure 7 shows that the observed behaviour for the underlying *particles*,  $\nu_{\theta}^{N}$ , of each model, is consistent with the behaviour of the obtained model  $\Phi_{\theta}^{N}$ . That is, as particles become *close* to being *symmetric* in some sense, the resulting *shallow model* also becomes increasingly equivariant as well.

Finally, Figure 8 and Figure 9 illustrate quite well the observations from Corollary 1. When our teacher **isn't equivariant**, models trained using any kind of SL technique end up suffering from the **inductive bias** introduced by the symmetric assumption (something that's hinted by the symmetrization gap characterization from Lemma 2 presented in SuppMat-A.3). On the other hand, the **vanilla** model thrives in approximating  $f_*$  as it is capable of breaking the assumed symmetry, unlike **DA**, **FA** and **EA**. The training regimes that use SL techniques are effectively approximating  $Q_G.f_*$ , as shown in Figure 9 (and proven in Corollary 1). We also notice that, for the **WI**-initialized experiments, **EA**s end up suffering from their constraint of *staying within*  $\mathcal{E}^G$ , as they can't approximate  $f_*$  (or  $Q_G.f_*$ ) as well as **DA** or **FA** (even when the teacher is **SI**). This isn't the case for the **SI**-initialized experiments, where the performance of **DA**, **FA** and **EA** (and **vanilla** only for equivariant  $f_*$ ) is quite closely comparable (once again, hinting at Theorem 6). We also notice a general trend showing that, for bigger N, the approximations of  $f_*$  (or, eventually,  $Q_G.f_*$ ) become increasingly *better* (specially for the **WI**-initialization).




Figure 4: Visualization of the NN particles after training under the **vanilla**, **DA**, **FA** and **EA** schemes, for a single realization of the experiment. Squares represent the *teacher* particles (which are **WI**), dots represent the student particles, and the hyperplane is  $\mathcal{E}^G$ . The bigger plots show an aerial view of the global particle distribution after training; and the minor plots below them show a viewpoint at the level of (and parallel to)  $\mathcal{E}^G$ . The student particles were all initialized to be **SI** (and to coincide at initialization between the different schemes), and trained with equation (4) correspondingly applying the proper SL technique.




Figure 5: Visualization of the NN particles after training under the **vanilla**, **DA**, **FA** and **EA** schemes for a single realization of the experiment. Squares represent the *teacher* particles (which are **arbitrary**), dots represent the student particles, and the hyperplane is  $\mathcal{E}^G$ . The bigger plots show an aerial view of the global particle distribution after training; and the minor plots below them show a viewpoint at the level of (and parallel to)  $\mathcal{E}^G$ . The student particles were all initialized to be **SI** (and to coincide at initialization between the different schemes), and trained with equation (4) correspondingly applying the proper SL technique. Notice how the particles for the **vanilla** scheme readily **leave**  $\mathcal{E}^G$  (despite the noise being projected onto it) and seem to *approach* the teacher particles.




Figure 6: **RMD** comparisons between training regimes, for different values of N, at the end of a **WI**-initialized training for  $N_e$  epochs. Each column corresponds to a teacher with, respectively, an **arbitrary**, **WI** and **SI** distribution. *Row 1* displays  $\text{RMD}^2(\nu_{N_e}^N, (\nu_{N_e}^N)^G)$  for the different regimes, to evaluate to what extent the training *remained* **WI**. *Row 2* displays the **RMD** between **DA**, **FA** and **vanilla** training regimes; as well as a comparison between **EA** and the projected particles of **FA**.




Figure 7: Approximation of  $L^2$ -distance between each model and its **symmetrized** version for increasing values of N. Each column corresponds to a different *teacher* as before. *Row 1* corresponds to the **SI**-initialized experiment and *Row 2* to the **WI**-initialized one.




Figure 8: Approximation of  $L^2$ -distance between each model and the corresponding **teacher network**  $f_*$ , for increasing values of N. Each column corresponds to a different *teacher* as before; *Row 1* corresponds to the **SI**-initialized experiment and *Row 2* to the **WI**-initialized one.




Figure 9: Approximation of  $L^2$ -distance between each model and the **symmetrized teacher network**  $Q_G.f_*$ , for increasing values of N. Each column corresponds to a different *teacher* as before; *Row 1* corresponds to the **SI**-initialized experiment and *Row 2* to the **WI**-initialized one.






Figure 10: **RMD** comparison between the empirical student particle distribution,  $\nu_{\theta}^{N}$ , to both  $P_{E_j} \# \nu_{\theta}^{N}$  and  $(\nu_{\theta}^{N})^{\mathcal{E}^{G}}$  (where *j* is the heuristic step). These are performed at the beginning and the end of training on every fixed heuristic step. The red line is placed at the value  $10^{-3}$  and represents a possible *threshold*  $\delta$ , to be used in the heuristic to determine whether *training left its corresponding*  $E_j$  or not.

## D.2 Heuristic algorithm for architecture discovery

The proposed heuristic that we infer from the results on the previous experimental setting is quite thoroughly described in Section 4.2. We only notice that, for this particular setting, the *learning rate* was chosen to be  $\alpha = 20$  (to better approximate the **MFL** conditions). Beyond the description of the proposed heuristic and the simple example visualized in Figure 2, we also provide Figure 10 here, illustrating a possible threshold choice in that setting.

Considering  $E_j$  for j = 0, 1, ... as the *spaces* that are discovered on each step of the heuristic, Figure 10 displays the values of:  $\mathbf{RMD}^2(\nu_{\theta}^N, P_{E_j} \# \nu_{\theta}^N)$  and  $\mathbf{RMD}^2(\nu_{\theta}^N, P_{\mathcal{E}^G} \# \nu_{\theta}^N)$ ; both before and after training on a given heuristic step j. The red line simbolizes a possible value of  $\delta$  that could be fixed to **detect** whenever the obtained particle distribution *after training* **stayed** on  $E_j$ . In the case of this example, on steps 0 and 1 we would decide that *the training left the original space*  $E_j$ , but we wouldn't do so on step 2, allowing us to fix  $\mathcal{E}^G := E_2$ . As shown by the values of  $\mathbf{RMD}^2(\nu_{\theta}^N, P_{\mathcal{E}^G} \# \nu_{\theta}^N)$ , we wouldn't be *too far off* with our prediction.

Despite this proposed heuristic being potentially interesting for real-world applications, we acknowledge that the setting where it is applied here might be too *simple*, *synthetic* and *idealized*. On one hand, this provided a *clean-enough* framework, where the underlying phenoma could be easily observed. However, in order to properly validate our heuristic approach, experiments with *more complex settings* (and with larger and more intricate datasets) need to be performed. These should also be coupled with sound theoretical guarantees, whose exploration we leave for future work.

## **E** Further theoretical insights

The following result provides consistency guarantees when the *regularization parameters*  $\tau$  and  $\beta$  are small, and is a slight extension of a result in [31].

**Proposition 18** ( $\Gamma$ -convergence, as in ). Let  $\mathcal{Z} = \mathbb{R}^D$ . If R is  $W_p$ -continuous,  $\nu$  is a Gibbs measure of potential U, and both U and r satisfy assumption 2 (or alternatively, are equal to 0), then  $R_{\nu}^{\tau,\beta}$   $\Gamma$ -converges to R when  $\tau, \beta \downarrow 0$ . Particularly, given  $\mu^{*,\tau,\beta,\nu}$  the minimizer of  $R_{\nu}^{\tau,\beta}$ , we have

$$\overline{\lim_{\tau,\beta\to 0}} R(\mu^{*,\,\tau,\,\beta,\,\nu}) = \inf_{\mu\in\mathcal{P}_2(\mathcal{Z})} R(\mu).$$

In particular, every cluster point of  $(\mu^{*, \tau, \beta, \nu})_{\tau, \beta}$  is a minimizer of R.

*Proof of Proposition 18.* We follow the exact same proof structure as [31], employing essentially their same techniques. However, we do adapt it to the case of taking the simultaneous limit of  $\tau, \beta \to 0$ , and so we do include it for completeness.

Let  $(\tau_n)_{n\in\mathbb{N}}$  and  $(\beta_n)_{n\in\mathbb{N}}$  be two positive sequences decreasing to 0. On the one hand, since R is continuous (weakly if p = 0 or in  $W_p$  for other  $p \ge 1$ ) and  $H_{\nu}(\mu) = D(\mu || \nu) \ge 0$ , for all  $\mu_n \to \mu$  (in the appropriate sense), we have

$$\lim \inf_{n \to +\infty} \inf_{\nu} R_{\nu}^{\tau_n, \beta_n}(\mu_n) \ge \lim_{n \to +\infty} R(\mu_n) = R(\mu).$$

On the other hand, given  $\mu \in \mathcal{P}_p(\mathcal{Z})$ , consider  $\rho$  to be the heat kernel in  $\mathcal{Z} = \mathbb{R}^D$  and  $\rho_n(x) := \beta_n^{-D} \nu(x/\beta_n)$ . In particular, from [1] (as the heat kernel has finite *p*-th moments) we know that  $\mu_n := \mu * \rho_n \xrightarrow[n \to \infty]{} \mu$  in  $W_p$  (or weakly if it is the case).

Now, since the function  $h(x) := x \log(x)$  is convex, from Jensen's inequality we get that

$$\int_{\mathcal{Z}} h(\mu * \rho_n) dx \le \int_{\mathcal{Z}} \int_{\mathcal{Z}} h\left(\rho_n(x-y)\right) \mu(dy) dx = \int_{\mathcal{Z}} h(\rho_n(x)) dx = \int_{\mathcal{Z}} h(\rho(x)) dx - D \log(\sqrt{2\beta_n}),$$

Besides, we have (denoting here  $g(x) = e^{-U(x)}$ ):

$$\int_{\mathcal{Z}} (\mu * \rho_n) \log(g) dx = -\int_{\mathcal{Z}} \mu(dy) \int_{\mathcal{Z}} \rho_n(x) U(x-y) dx \ge -C \left(1 + \int_{\mathcal{Z}} |y|^2 \mu(dy)\right).$$

The last inequality is due to the quadratic growth of U; and by the same argument on r:

$$\int_{\mathcal{Z}} (\mu * \rho_n) r dx = \int_{\mathcal{Z}} \mu(dy) \int_{\mathcal{Z}} \rho_n(x) r(x-y) dx \le C \left( 1 + \int_{\mathcal{Z}} |y|^2 \mu(dy) \right)$$

Notice that whenever  $U \equiv 0$  or  $r \equiv 0$ , despite them not satisfying assumption 2, we still get the same inequalities (since the leftmost term would be 0).

Now, as R is  $W_p$ -continuous,  $R(\mu_n) \xrightarrow[n \to \infty]{} R(\mu)$ , and:

$$\begin{split} \lim \sup_{n \to +\infty} R_{\nu}^{\tau_n, \, \beta_n}(\mu * \nu_n) \\ &\leq R(\mu) + \lim \sup_{n \to +\infty} \tau_n \left( \int_{\mathcal{Z}} (\mu * \rho_n) r dx \right) \\ &+ \lim \sup_{n \to +\infty} \beta_n \left( \int_{\mathcal{Z}} h(\mu * \rho_n) dx - \int_{\mathcal{Z}} (\mu * \rho_n) \log(g) dx \right) \end{split}$$

And, as  $\lim_{n\to\infty} \beta_n \log(\sqrt{2\beta_n}) = 0$  and the rest of the terms are bounded, we conclude that:

$$\lim \sup_{n \to +\infty} R_{\nu}^{\tau_n, \, \beta_n}(\mu * \rho_n) \le R(\mu)$$

In particular, denoting by  $\mu_*^{\tau,\beta,\nu}$  the unique minimizer of  $R_{\nu}^{\tau,\beta}$ , then from the previous expressions we get  $\forall n \in \mathbb{N}$  and  $\forall \mu \in \mathcal{P}_p(\mathcal{Z})$ :

$$R(\mu_*^{\tau_n,\beta_n,\nu}) \le R_{\nu}^{\tau_n,\beta_n}(\mu_*^{\tau_n,\beta_n,\nu}) \le R_{\nu}^{\tau_n,\beta_n}(\mu*\rho_n)$$

So that,

$$\lim \sup_{n \to \infty} R(\mu_*^{\tau_n, \beta_n, \nu}) \le \lim \sup_{n \to +\infty} R_{\nu}^{\tau_n, \beta_n}(\mu * \rho_n) \le R(\mu), \quad \text{for all } \mu \in P_2(\mathcal{Z}).$$

Finally, we provide, for completeness, a proof of Lemma 2 :

Proof of Lemma 2 (based on [22] and [33]). As  $H \leq G$  is a compact group and  $\pi$  is H-invariant, from proposition 10 we know that  $f^* = \mathbb{E}_{\pi}[Y|X = \cdot]$  lives in  $f^* \in L^2_H(\mathcal{X}, \mathcal{Y}; \pi|_{\mathcal{X}})$ . Consider any  $f \in L^2(\mathcal{X}, \mathcal{Y}; \pi|_{\mathcal{X}})$ , by Lemma 1 from [22] (which applies since  $\pi|_{\mathcal{X}}$  is G-invariant), we can decompose f as  $f = \overline{f}_G + f_G^{\perp}$ , where  $\overline{f}_G = \mathcal{Q}_G f$  is its symmetric part and  $f_G^{\perp} = f - \mathcal{Q}_G f$  its antisymmetric part. A standard calculation of the population risk under the quadratic loss setting (see the proof of Corollary 1 for further insight) gives:  $R(f) = R_* + ||f - f^*||^2_{L^2(\mathcal{X},\mathcal{Y};\pi_{\mathcal{X}})}$ , and so:

$$\Delta(f, \mathcal{Q}_G.f) = R(f) - R(\mathcal{Q}_G.f) = \mathbb{E}\left[\|f^*(X) - f(X)\|_{\mathcal{Y}}^2\right] - \mathbb{E}\left[\|f^*(X) - \overline{f}_G(X)\|_{\mathcal{Y}}^2\right],$$

which can be written as:

$$\begin{split} \Delta(f, \mathcal{Q}_G f) &= \mathbb{E}_{\pi} \left[ \| f^*(X) - \overline{f}_G(X) \|_{\mathcal{Y}}^2 - 2 \langle f^*(X) - \overline{f}_G(X), f_G^{\perp}(X) \rangle_{\mathcal{Y}} + \| f_G^{\perp}(X) \|_{\mathcal{Y}}^2 \right] \\ &- \mathbb{E}_{\pi} \left[ \| f^*(X) - \overline{f}_G(X) \|_{\mathcal{Y}}^2 \right] \\ &= -2 \langle f^* - \overline{f}_G, f_G^{\perp} \rangle_{L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})} + \| f_G^{\perp} \|_{L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})}^2 \\ &= -2 \langle f^*, f_G^{\perp} \rangle_{L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})} + \| f_G^{\perp} \|_{L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})}^2 \end{split}$$

Where we used that  $\langle \overline{f}_G, f_G^{\perp} \rangle_{L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})} = 0$ . The first term on the right hand side,  $-2\langle f^*, f_G^{\perp} \rangle_{L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})}$ , is what [33] call the *mismatch* between the real *underlying model* (which is only *H*-equivariant) and the *symmetrized* version of our model (which is made entirely *G*-equivariant).

Now, when  $\pi$  is *G*-equivariant, by proposition 10,  $Q_G f^* = f^*$ , and so:  $-2\langle f^*, f_G^{\perp} \rangle_{L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})} = 0$ , giving us the desired result:

$$\Delta(f, \mathcal{Q}_G f) = \|f_G^{\perp}\|_{L^2(\mathcal{X}, \mathcal{Y}; \pi_{\mathcal{X}})}^2$$

Lemma 2 essentially says that if we *try to symmetrize* a model with respect to a group that has "*more symmetries*" than what are actually observable in our data (i.e.  $\pi$  in itself is only *H*-invariant, but we symmetrize with respect to  $G \ge H$ ); we can either *win* or *lose* generalization power according to the interplay between the two presented terms. In particular, if  $\pi$  is *G* equivariant, there's a *strict generalization benefit* from choosing a *symmetric model* to tackle our *learning problem* (which gives the name to the paper [22]). In particular, whenever  $f_G^{\perp}$  is non-zero (on a strict) positive  $\pi|_{\mathcal{X}}$ -measure set) there's a *strict gain* in generalization power from using the symmetrized version of the model.