

FUNCTIONAL INEQUALITIES IN THE FRAMEWORK OF BANACH SPACES

CONSTANTIN P. NICULESCU

ABSTRACT. A quadrilateral inequality established by Schötz [28] in the context of Hilbert spaces is extended to the framework of Banach spaces. Our approach is based on the theory of majorization and a substitute for the parallelogram law associated with Clarkson's notion of von Neumann-Jordan constant. As a by-product, several functional inequalities that extend classical inequalities from linear algebra and geometry of Banach spaces are also obtained.

1. INTRODUCTION

This paper aims to illustrate the usefulness of majorization theory and higher-order convexity theory in generalizing several classical inequalities as functional inequalities. This includes results due to Ball, Carlen, and Lieb [1], Hanner [8], Popoviciu [25] and Zhang [33].

Our starting point was a recent paper by Schötz [28], that reveals the connection between a class of higher-order convex functions and the following quadruple inequality,

$$(1.1) \quad \|p - x\| + \|y - z\| \leq \|p - y\| + \|x - z\| + \|p - z\| + \|x - y\|,$$

that occurs in any normed vector space. To see that (1.1) indeed works, add side by side the following two inequalities that result from the triangle inequality,

$$2\|p - x\| \leq \|p - y\| + \|p - z\| + \|y - x\| + \|z - x\|$$

and

$$2\|y - z\| \leq \|y\| + \|z\| + \|y - x\| + \|z - x\|.$$

When dealing with inner product spaces, we also have

$$\begin{aligned} \|y - z\|^2 - \|p - y\|^2 - \|x - z\|^2 + \|p - x\|^2 &= 2\langle z - p, x - y \rangle \\ &\leq 2\|p - z\| \|x - y\| \\ &\leq \|p - z\|^2 + \|x - y\|^2, \end{aligned}$$

so the inequality (1.1) remains valid by squaring the norms. Schötz was able to single out an entire class of functions $f : [0, \infty) \rightarrow [0, \infty)$ that extend this inequality by replacing $\|\cdot\|$ with $f(\|\cdot\|)$:

Date: August 15, 2024.

2020 Mathematics Subject Classification. Primary 26A51, 39B62 46 B20; Secondary 26D15.

Key words and phrases. Hanner's inequalities; optimal 2-uniform convexity inequalities; higher order convexity; majorization; von Neumann-Jordan constant; quadruple functional inequalities.

Theorem 1. (Schötz [28], Theorem 3) *Let x, y, z, p be four points in an inner product space V and let $f : [0, \infty) \rightarrow [0, \infty)$ be a nondecreasing, convex, and differentiable function such that $f(0) = 0$ and f' is concave. Then*

$$\begin{aligned} f(\|p - x\|) + f(\|y - z\|) \\ \leq f(\|p - y\|) + f(\|x - z\|) + f(\|p - z\|) + f(\|x - y\|). \end{aligned}$$

For convenience, we will denote by \mathcal{S} the set of all functions $f : [0, \infty) \rightarrow [0, \infty)$ which are nondecreasing, convex and differentiable, and have concave derivatives and by \mathcal{S}_0 the set of functions in \mathcal{S} which vanishes at the origin. Some few examples of functions belonging to \mathcal{S}_0 are

$$\begin{aligned} (1 + \alpha x^2)^{1/2} - 1 \quad (\text{for } \alpha > 0), \\ x^\alpha \quad (\text{for } \alpha \in [1, 2]), \quad x \log(x + 1), \quad \text{and } \log(\cosh x). \end{aligned}$$

The aim of the present paper is to gain more insight into this matter by noticing that the functions considered by Schötz belong to the subject of higher order convexity (briefly summarized in Section 2). Indeed, the set \mathcal{S} coincides with the set of all functions $f : [0, \infty) \rightarrow \mathbb{R}$ which are nondecreasing, convex and 3-concave in the sense of E. Hopf [9] and T. Popoviciu [22], [24]. See Theorem 3, Section 2. Combining this fact with Popoviciu's approximation theorem (Theorem 4 in Section 2) we may reduce the reasoning with functions in \mathcal{S} to the case where they are also three times continuously differentiable.

In Section 3 we establish the "truncated form" of the two basic results of majorization theory, the Hardy-Littlewood-Pólya theorem and the Tomić-Weyl theorem on weak majorization. Their main advantage is the possibility to handle pairs of finite sequences of numbers, of different sizes. See Theorem 7 and Theorem 8. Unlike Sherman's theorem of majorization (see [18]) our results do not make explicit use of the stochastic matrices.

The applications proved in Section 4 represent functional generalization of the determinantal inequalities of Zhang [33] and Popoviciu, of the optimal 2-uniform convexity inequality for L^p spaces (with $p \in (1, 2]$) and of Hanner's inequality. See respectively Theorem 9, Theorem 10, Theorem 11 and 12. The 2-uniform convexity functional inequality for $p \in [2, \infty)$ makes the objective of Theorem 16 in Section 6.

The Section 5 is devoted to a quick presentation of a basic ingredient necessary in our extension of Theorem 1 to the general context of Banach spaces: the von Neumann-Jordan constant. According to Clarkson [5], the *von Neumann-Jordan constant* $C_{NJ}(X)$ of a Banach space X is defined by the formula

$$C_{NJ}(X) = \sup \left\{ \frac{\|u + v\|^2 + \|u - v\|^2}{2\|u\|^2 + 2\|v\|^2} : u, v \in X \text{ and } \|u\| + \|v\| \neq 0 \right\}.$$

He noticed that $C_{NJ}(X) \in [1, 2]$ and the equality $C_{NJ}(X) = 1$ characterizes inner product spaces.

The *von Neumann-Jordan constant* makes possible the following substitute of the parallelogram law in any Banach space X :

$$\|x + y\|^2 + \|x - y\|^2 \leq 2C_{NJ}(X) (\|x\|^2 + \|y\|^2), \quad \text{for all } x, y \in X.$$

A functional inequality generalizing this fact makes the objective of Theorem 13. This theorem is used in Section 6 to prove the following generalization of Theorem 1:

Theorem 2. *Let y, z, q, r be four points in the Banach space X and let f be a nondecreasing, convex and 3-concave function defined on $[0, \infty)$ such that $f(0) = 0$. Then*

$$f(\|y - q\|) + f(\|z - r\|) \leq \frac{N(X)}{2} \{f(\|y - z\|) + f(\|r - q\|) + f(\|z - q\|) + f(\|y - r\|)\}.$$

Here $N(X) = 2C_{NJ}(X)$ if $2C_{NJ}(X)$ is an integer and $N(X) = 4$ otherwise.

Since $N(X) = 2$ if X is an inner product space, in this case Theorem 2 reduces to Theorem 1.

2. SOME BASIC FACTS CONCERNING THE n -CONVEX FUNCTIONS ($n \leq 3$)

Higher order convexity was introduced by Hopf [9] and Popoviciu [22], [24], who defined it in terms of divided differences. Assuming that f is a real-valued function defined on an interval I , its divided differences of order $0, 1, \dots, n$ associated to a family x_0, x_1, \dots, x_n of $n+1$ distinct points are respectively defined by the formulas:

$$\begin{aligned} [x_0; f] &= f(x_0) \\ [x_0, x_1; f] &= \frac{f(x_1) - f(x_0)}{x_1 - x_0} \\ &\dots \\ [x_0, x_1, \dots, x_n; f] &= \frac{[x_1, x_2, \dots, x_n; f] - [x_0, x_1, \dots, x_{n-1}; f]}{x_n - x_0} \\ &= \sum_{j=0}^n \frac{f(x_j)}{\prod_{k \neq j} (x_j - x_k)}. \end{aligned}$$

Notice that all these divided differences are invariant under the permutation of points x_0, x_1, \dots, x_n . As a consequence, we may always assume that $x_0 < x_1 < \dots < x_n$.

A function f is called n -convex (respectively n -concave) if all divided differences $[x_0, x_1, \dots, x_n; f]$ are nonnegative (respectively nonpositive). In particular,

- the convex functions of order 0 are precisely the nonnegative functions;
- the convex functions of order 1 are the nondecreasing functions;
- the convex functions of order 2 are nothing but the usual convex functions since in this case for all $x_0 < x_1 < x_2$ in I ,

$$[x_0, x_1, x_2; f] = \frac{\frac{f(x_0) - f(x_1)}{x_0 - x_1} - \frac{f(x_1) - f(x_2)}{x_1 - x_2}}{x_0 - x_2} \geq 0,$$

that is,

$$(x_2 - x_0) f(x_1) \leq (x_2 - x_1) f(x_0) + (x_1 - x_0) f(x_2).$$

The description of 3-convex functions (as well as of the higher order convex functions) in terms of divided differences is rather intricate. For example, a function f is 3-convex if for every quadruple $x_0 < x_1 < x_2 < x_3$ of elements we have

$$[x_0, x_1, x_2, x_3; f] = \frac{f(x_0)}{(x_0 - x_1)(x_0 - x_2)(x_0 - x_3)} - \frac{f(x_1)}{(x_0 - x_1)(x_1 - x_2)(x_1 - x_3)} \\ + \frac{f(x_2)}{(x_0 - x_2)(x_1 - x_2)(x_2 - x_3)} - \frac{f(x_3)}{(x_0 - x_3)(x_1 - x_3)(x_2 - x_3)} \geq 0,$$

equivalently,

$$(x_2 - x_0)(x_3 - x_0)(x_3 - x_2)f(x_1) + (x_1 - x_0)(x_2 - x_0)(x_2 - x_1)f(x_3) \\ \geq (x_2 - x_1)(x_3 - x_1)(x_3 - x_2)f(x_0) + (x_1 - x_0)(x_3 - x_0)(x_3 - x_1)f(x_2).$$

When the points x_0, x_1, x_2, x_3 are equidistant, that is, when $x_1 = x_0 + h$, $x_2 = x_0 + 2h$, $x_3 = x_0 + 3h$ for some $h > 0$, the last inequality becomes

$$f(x_0 + 3h) - 3f(x_0 + 2h) + 3f(x_0 + h) - f(x_0) \geq 0,$$

equivalently,

$$(2.1) \quad f(x_0) + 3f\left(\frac{x_0 + 2x_3}{3}\right) \leq 3f\left(\frac{2x_0 + x_3}{3}\right) + f(x_3).$$

Fortunately, some others, more convenient, approaches are available.

If f is 3-times differentiable, then a repeated application of the mean value theorem yields the existence of a point $\xi \in (\min_k x_k, \max_k x_k)$ such that

$$[x_0, x_1, x_2, x_3; f] = \frac{f^{(3)}(\xi)}{6}.$$

As a consequence, one obtains the sufficiency part of the following practical criterion of 3-convexity.

Lemma 1. *Suppose that f is a continuous function defined on an interval I which is 3-times differentiable on the interior of I . Then f is 3-convex if and only if its derivative of third order is nonnegative.*

The necessity part is also immediate by using the standard formulas for derivatives via iterated differences,

$$f^{(3)}(x) = \lim_{h \rightarrow 0^+} \frac{f(x_0 + 3h) - 3f(x_0 + 2h) + 3f(x_0 + h) - f(x_0)}{h^3}.$$

According to Lemma 1, the following functions are 3-convex functions on \mathbb{R}_+ :

$$x^\alpha \text{ (for } \alpha \in (0, 1] \cup [2, \infty)), \quad x/(1+x), \\ \log(1+x), \quad -x \log x, \quad \sinh, \quad \cosh, \quad \text{and} \quad -\log(\Gamma(x)).$$

Notice that the polynomials of degree ≤ 2 are both 3-convex and 3-concave functions on the whole real line.

Remark 1. *(Permanence proprieties) The continuous n -convex functions defined on an interval I constitute a convex cone in the vector space $C(I)$, of all continuous functions on I .*

Every continuous function which is n -convex on the interior of I is n -convex on the whole interval.

The limit of a pointwise convergent sequence of n -convex functions is also an n -convex function.

The following characterization of higher order convexity is due to Hopf ([9], p. 24) and Popoviciu ([22], p. 48):

Theorem 3. *Suppose that f is a continuous function defined on an interval I . Then f is 3-convex if and only if it is differentiable on the interior of I and f' is a convex function.*

Corollary 1. *Every function $f \in \mathcal{S}$ is 3-concave and every nondecreasing, convex and 3-concave function $f : [0, \infty) \rightarrow [0, \infty)$ belongs to \mathcal{S} .*

Proof. It suffices to show that every nondecreasing, convex and 3-concave function $f : [0, \infty) \rightarrow [0, \infty)$ is continuously differentiable at the origin. For this, notice first that

$$0 < x < y \quad \text{implies} \quad 0 \leq \frac{f(x) - f(0)}{x} \leq \frac{f(y) - f(0)}{y},$$

since f is nondecreasing and convex. As a consequence,

$$\lim_{x \rightarrow 0+} \frac{f(x) - f(0)}{x} = \inf_{x > 0} \frac{f(x) - f(0)}{x} \geq 0,$$

which assures the differentiability at the origin (and thus everywhere, according to Theorem 3). Since f is convex, its derivative f' is nondecreasing. Therefore,

$$\begin{aligned} \lim_{x \rightarrow 0+} f'(x) &= \inf_{x > 0} f'(x) = \inf_{x, h > 0} \frac{f(x+h) - f(x)}{h} \\ &= \liminf_{h \rightarrow 0+} \frac{f(h) - f(0)}{h} = f'(0), \end{aligned}$$

which means that f' is continuous at the origin. □

An important source of nonnegative, nondecreasing, convex and 3-concave functions on a compact interval $[0, A]$ is that of completely monotone functions. Recall that a function $f : [0, \infty) \rightarrow [0, \infty)$ is *completely monotone* if it is continuous on $[0, \infty)$, indefinitely differentiable on $(0, \infty)$ and

$$(-1)^n f^{(n)}(x) \geq 0 \quad \text{for all } x > 0 \text{ and } n \geq 0.$$

Some simple examples are e^{-x} , $1/(1+x)$, and $(1/x) \log(1+x)$. For more details, see the monograph of Schilling, Song and Vondraček [27]. Every completely monotone function f is nonnegative, nonincreasing, convex and 3-concave, but adding to it linear functions αx with $\alpha \geq -\inf_{x \in [0, A]} f'(x)$ one obtains nonnegative, nondecreasing, convex and 3-concave functions on a given compact interval $[0, A]$.

Popoviciu has characterized the property of n -convexity in terms of higher order differences.

The *difference operator* Δ_h (of step size $h \geq 0$) associates to each function f defined on an interval I the function $\Delta_h f$ defined by

$$(\Delta_h f)(x) = f(x+h) - f(x),$$

for all x such that the right-hand side formula makes sense. Notice that no restrictions are necessary if $I = \mathbb{R}^+$ or $I = \mathbb{R}$. The difference operators are linear and commute to each other,

$$\Delta_{h_1} \Delta_{h_2} = \Delta_{h_2} \Delta_{h_1}.$$

They also verify the following property of invariance under translation:

$$\Delta_h (f \circ T_a) = (\Delta_h f) \circ T_a,$$

where T_a is the translation defined by the formula $T_a(x) = x + a$.

The higher order iterated differences can be introduced via the formulas:

$$\begin{aligned} (\Delta_h)^0 f(x) &= f(x) \\ (\Delta_h)^n f(x) &= \underbrace{\Delta_h \cdots \Delta_h}_{n \text{ times}} f(x) \\ &= \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} f(x + kh) \quad \text{for } n \geq 1. \end{aligned}$$

Their connection with the higher order divided differences is given by

$$(\Delta_h)^n f(x) = h^n [x, x+h, \dots, x+nh; f]$$

and this applies to every function f defined on an interval I of the form $[0, A]$ or $[0, \infty)$, all points $x \in I$ and all steps $h > 0$ such that $x + nh \in I$.

Clearly, if f is an n -convex function ($n \geq 1$) defined on an interval I of the form $[0, A]$ or $[0, \infty)$, then

$$(2.2) \quad (\Delta_h)^n f(x) \geq 0$$

for all $x \in I$ and all $h > 0$ such that $x + nh \in I$. As was noticed by Popoviciu [24] (at the beginning of Section 24, p. 49) this property characterizes the n -convex functions under the presence of continuity. See also [14] and [19] (as well as the references therein).

The inequality (2.2) together with Bernstein's variant of the Weierstrass approximation theorem (see [3], Theorem 8.8.1, p. 256) yields the following shape preserving approximation result.

Theorem 4. (Popoviciu's approximation theorem [23]) *If a continuous function $f : [0, 1] \rightarrow \mathbb{R}$ is k -convex, then so are the Bernstein polynomials associated to it,*

$$B_n(f)(x) = \sum_{i=0}^n \binom{n}{i} x^i (1-x)^{n-i} f\left(\frac{i}{n}\right).$$

Moreover, by the well-known property of simultaneous uniform approximation of a function and its derivatives by the Bernstein polynomials and their derivatives, it follows that $B_n(f)$ and any derivative (of any order) of it converge uniformly to f and to its derivatives, correspondingly.

Using a change of variable, one can easily see that the approximation theorem extends to functions defined on compact intervals $[a, b]$ with $a < b$.

Proof. Using mathematical induction one can easily show that the derivatives of Bernstein's polynomials verify the formula

$$\begin{aligned} B_n^{(k)}(f)(x) &= n(n-1) \cdots (n-k+1) \sum_{i=0}^{n-k} \underbrace{\Delta_{1/n} \cdots \Delta_{1/n}}_{k \text{ times}} f(j/n) \binom{n-k}{i} x^i (1-x)^{n-k-i}. \end{aligned}$$

The proof ends by taking into account the formula (2.2) and Lemma 1. □

When combined with Remark 1, Theorem 4 implies that any result valid for the smooth n -convex functions also works for all n -convex continuous functions.

Corollary 2. *If $f : \mathbb{R}_+ \rightarrow [0, \infty)$ is a continuous 3-convex function which is also nondecreasing and concave, then the same properties hold for f^α if $\alpha \in (0, 1]$.*

Proof. According to Theorem 4, we may reduce the proof to the case where the involved function is of class C^3 , in which case the conclusion follows from Lemma 1. \square

For a second application of Theorem 4 we need the following well known fact on concave functions.

Lemma 2. *If $f : [0, \infty) \rightarrow \mathbb{R}$ is a concave function and $f(0) \geq 0$, then the function $f(x)/x$ is nonincreasing on $(0, \infty)$.*

We are now in a position to state a rather general result concerning the composition of functions with opposite properties of convexity. It extends Lemma 27 in [28].

Theorem 5. *Suppose that $f : [0, \infty) \rightarrow \mathbb{R}$ is a nondecreasing, continuous and 3-concave function. Then the function $g(x) = f(x^\alpha)$ is nondecreasing and concave for every $\alpha \in (0, 1/2]$.*

Proof. Combining Popoviciu’s approximation theorem with Remark 1, we can reduce ourselves to the case where f is of class C^2 . Since f is nondecreasing, it follows that $f' \geq 0$. According to Theorem 3, f' is a concave function, so by Lemma 2 it results that $f'(x)/x$ is nonincreasing on $(0, \infty)$. The fact that g is nondecreasing is clear. To prove that g is also concave it suffices to show that its derivative is nonincreasing. Indeed, g' can be represented as the product of two nonincreasing nonnegative functions,

$$g'(x) = \alpha x^{\alpha-1} f'(x^\alpha) = \alpha x^{2\alpha-1} \cdot \frac{f'(x^\alpha)}{x^\alpha},$$

and the proof is done. \square

Theorem 5 is not valid for $\alpha \in (1/2, \infty)$, a counterexample being provided by the function $f(x) = x^2, x \geq 0$.

Remark 2. (a) *The argument of Theorem 5 also shows that $f(x^\alpha)$ is a nonincreasing convex function if $\alpha \in (0, 1/2]$ and f is a function of the same nature;*

(b) *$f(x^\alpha)$ is a convex function provided that $\alpha \in (0, 1/2]$ and f is a differentiable, convex and 3-convex function such that $f'(0) \leq 0$. Indeed, proceeding as in the proof of Theorem 5 one can assume that f is of class C^2 . According to Theorem 3, the condition of 3-convexity implies that f has a convex derivative on $(0, \infty)$, so taking into account Lemma 2, the function $f'(x)/x$ is nondecreasing, Therefore the derivative of $f'(x)/x$ is nonnegative, a fact that assures that the second derivative of the function $f(x^\alpha)$ is also nonnegative.*

Some few examples of differentiable and 3-convex functions $f : [0, \infty) \rightarrow \mathbb{R}$ such that $f(0) = f'(0) = 0$ are

$$x^\alpha \ (\alpha \geq 2), \ e^x - 1 - x, \ -x \log(x + 1) \text{ and } -\log(\cosh x).$$

More results concerning the 3-convex/3-concave functions are made available by the recent survey of Marinescu and Niculescu [14].

3. SOME CONSEQUENCES OF THE HARDY-LITTLEWOOD-PÓLYA THEOREM OF MAJORIZATION

We start by recalling the Hardy-Littlewood-Pólya theorem of majorization:

Theorem 6. *Let f be a real-valued convex function defined on a nonempty interval I . If $\mathbf{x} = (x_k)_{k=1}^n$ and $\mathbf{y} = (y_k)_{k=1}^n$ are two families of points in I such that*

$$(3.1) \quad x_1 \geq \cdots \geq x_n$$

and

$$(3.2) \quad \sum_{k=1}^m x_k \leq \sum_{k=1}^m y_k \quad \text{for } m = 1, \dots, n,$$

with equality for $m = n$, then

$$(3.3) \quad \sum_{k=1}^n f(x_k) \leq \sum_{k=1}^n f(y_k).$$

When condition (3.1) is replaced by $y_1 \leq \cdots \leq y_n$, then the conclusion (3.3) works in the reverse direction.

Pólya [21] noticed that this result implies the other classical result of majorization theory, precisely, the theorem of Tomić [31] and Weyl [32] on weak majorization:

Corollary 3. (The Tomić-Weyl theorem) *Let f be a real-valued nondecreasing convex function defined on a nonempty interval I . Then the inequality (3.3) holds for every two families $\mathbf{x} = (x_k)_{k=1}^n$ and $\mathbf{y} = (y_k)_{k=1}^n$ of points in I that verify the conditions (3.1) and (3.2).*

When condition (3.1) is replaced by $y_1 \leq \cdots \leq y_n$, then the conclusion (3.3) works in the reverse direction.

The details can be found in [18].

Of special interest for us will be the following "truncated forms" of Corollary 3 and Theorem 6.

Theorem 7. (Truncated majorization: the convex case) *If $f : [0, \infty) \rightarrow \mathbb{R}$ is a nondecreasing convex function, then for all integer numbers $2 \leq m \leq n$ and all strings x_1, \dots, x_n and y_1, \dots, y_m of positive numbers such that*

$$(3.4) \quad \max \left\{ \sum_{p=1}^k x_{i_p} : i_r \neq i_s \right\} \leq \max \left\{ \sum_{p=1}^k y_{j_p} : j_r \neq j_s \right\}$$

for $k = 1, \dots, m - 1$ and

$$(3.5) \quad \sum_{k=1}^n x_k \leq \sum_{k=1}^m y_k,$$

we have

$$(3.6) \quad \sum_{k=1}^n f(x_k) \leq \sum_{k=1}^m f(y_k) + (n - m)f(0).$$

Proof. Apply Corollary 3 to the families

$$\mathbf{x} = (x_1, x_2, \dots, x_n) \quad \text{and} \quad \mathbf{y} = (y_1, \dots, y_m, \underbrace{0, \dots, 0}_{n-m \text{ times}}).$$

□

Corollary 4. *If $f : [0, \infty) \rightarrow \mathbb{R}$ is a nondecreasing convex function and $x_1, x_2, \dots, x_n, y_1, y_2$ ($n \geq 2$) are nonnegative numbers such that*

$$\max\{x_1, x_2, \dots, x_n\} \leq \max\{y_1, y_2\}$$

and

$$\sum_{k=1}^n x_k \leq y_1 + y_2,$$

then

$$\sum_{k=1}^n f(x_k) \leq f(y_1) + f(y_2) + (n-2)f(0).$$

In the case of nonincreasing concave functions defined on $[0, \infty)$, the conclusion works in the reverse direction.

Theorem 8. (Truncated majorization: the concave case) *Let $f : [0, \infty) \rightarrow \mathbb{R}$ be a nondecreasing concave function and let $x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_m$ ($2 \leq m \leq n$) be nonnegative numbers such that*

$$\begin{aligned} \max\{x_1, x_2, \dots, x_n\} &\leq \max\{y_1, y_2, \dots, y_m\}, \\ \max\{x_{i_1} + x_{i_2} : i_1 \neq i_2\} &\leq \max\{y_{j_1} + y_{j_2} : j_1 \neq j_2\} \\ &\dots \\ \max\left\{\sum_{p=1}^k x_{i_p} : i_r \neq i_s\right\} &\leq \max\left\{\sum_{p=1}^k y_{j_p} : j_r \neq j_s\right\} \end{aligned}$$

for $k \leq m-1$ and

$$\sum_{i=1}^n x_i \geq \sum_{j=1}^m y_j.$$

Then

$$(3.7) \quad \sum_{i=1}^n f(x_i) \geq \sum_{j=1}^m f(y_j) + (n-m)f(0).$$

Proof. Without loss of generality we may increase y_k to \tilde{y}_k (for $k \in \{1, \dots, m\}$) so that

$$\sum_{k=1}^n x_k = \sum_{j=1}^m \tilde{y}_j$$

and also we may assume that $x_1 \geq x_2 \geq x_3 \geq \dots \geq x_n$ and $\tilde{y}_1 \geq \tilde{y}_2 \geq \dots \geq \tilde{y}_m$. Due to our hypotheses $\tilde{y}_1 \geq x_1$, $\tilde{y}_1 + \tilde{y}_2 \geq x_1 + x_2$, ..., and $\tilde{y}_1 + \tilde{y}_2 + \dots + \tilde{y}_m = \sum_{k=1}^n x_k \geq x_1 + \dots + x_m$. Therefore the string $\mathbf{x} = (x_1, x_2, x_3, \dots, x_n)$ is majorized by $\mathbf{y} = (\tilde{y}_1 + \tilde{y}_2 + \dots + \tilde{y}_m, \underbrace{0, \dots, 0}_{n-m \text{ times}})$ and the inequality (3.7) follows from the Hardy-

Littlewood-Pólya theorem of majorization and the fact that f is nondecreasing. □

Corollary 5. *If $f : [0, \infty) \rightarrow \mathbb{R}$ is a nondecreasing concave function and $x_1, x_2, \dots, x_n, y_1, y_2$ ($n \geq 2$) are nonnegative numbers such that*

$$\max\{x_1, x_2, \dots, x_n\} \leq \max\{y_1, y_2\}$$

and

$$\sum_{k=1}^n x_k \geq y_1 + y_2,$$

then

$$\sum_{k=1}^n f(x_k) \geq f(y_1) + f(y_2) + (n-2)f(0).$$

In the case of nonincreasing convex functions defined on $[0, \infty)$, the conclusion works in the reverse direction.

Theorem 8 and its Corollary 5 apply to functions such as x^α ($\alpha \in (0, 1]$), $\log(1+x)$, $\frac{x}{x+r}$ ($r \geq 0$) and $1 - e^{-tx}$ ($t > 0$), all defined on $[0, \infty)$ and vanishing at the origin.

A particular case of Corollary 5 was noticed by Schötz [28], Lemma 47, using a different argument.

A nice illustration of Corollary 4 (applied to the function $f(x) = x^{p/q}$) is the following inequality noticed by Enflo [6]: if r_1, r_2, \dots, r_6 are positive real numbers and $\max(r_1, \dots, r_4) \leq \max(r_5, r_6)$ and $r_1^q + \dots + r_4^q = r_5^q + r_6^q$, then

$$r_1^p + \dots + r_4^p \leq r_5^p + r_6^p \quad \text{if } p \geq q.$$

For applications to the functional inequalities see the next section.

4. FUNCTIONAL INEQUALITIES VIA MAJORIZATION

The following inequality can be found in the book of Zhang [33], Problem 36, p. 215: if A, B, C are positive semidefinite matrices of the same dimension, then

$$\det(A + B + C) + \det C \geq \det(A + C) + \det(B + C).$$

Since $A + B + C \geq A + C$, $B + C$ and the function \det is nondecreasing on the cone of positive semidefinite matrices, one can derive via Corollary 4 the following result.

Theorem 9. *If $f : [0, \infty) \rightarrow \mathbb{R}$ is a nondecreasing convex function and A, B and C are positive semidefinite matrices then*

$$f(\det(A + B + C)) + f(\det C) \geq f(\det(A + C)) + f(\det(B + C)),$$

which (by symmetrization) leads to the following inequality,

$$\begin{aligned} & \frac{f(\det A) + f(\det B) + f(\det C)}{3} + f(\det(A + B + C)) \\ & \geq \frac{2}{3}f(\det(A + B)) + f(\det(B + C)) + f(\det(A + C)), \end{aligned}$$

that reminds us of Popoviciu's inequality [25].

The case of the positive semidefinite matrices

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}, \quad C = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$$

and of the convex function $f(x) = x^2$, $x \geq 0$, shows that the conclusion of Theorem 9 cannot be strengthened to

$$\begin{aligned} & \frac{f(\det A) + f(\det B) + f(\det C)}{3} + f\left(\det\left(\frac{A+B+C}{3}\right)\right) \\ & \geq \frac{2}{3} \left(f\left(\det\left(\frac{A+B}{2}\right)\right) + f\left(\det\left(\frac{B+C}{2}\right)\right) + f\left(\det\left(\frac{A+C}{2}\right)\right) \right), \end{aligned}$$

that is, it is not a generalization of Popoviciu's inequality.

A partial generalization of Popoviciu's inequality for functions of a vector variable can be found in [17]. See also [2].

The statement of *the optimal 2-uniform convexity inequality* stated below can be found in the paper of Ball, Carlen and Lieb [1], Proposition 3:

Proposition 1. *If $p \in (1, 2]$ and x and y belong to an L^p space (or to the Schatten space $S^p(H)$), then*

$$\|x\|_p^2 + \|y\|_p^2 \geq 2 \left\| \frac{x+y}{2} \right\|_p^2 + 2(p-1) \left\| \frac{x-y}{2} \right\|_p^2.$$

For $p \in [2, \infty)$, the inequality is reversed.

Here $\|\cdot\|_p$ denotes either the L^p norm or the Schatten norm of index p .

Recall that the Schatten space of index $p \in [1, \infty)$, associated to a Hilbert space H , is the space $S^p(H)$ of all compact linear operators $T : H \rightarrow H$ whose sequences $(s_n(T))_n$ of singular values belong to ℓ^p . $S^p(H)$ is a Banach space with respect to the norm

$$\|T\|_p = \left(\sum_n |s_n(T)|^p \right)^{1/p}.$$

See [30] for a comprehensive presentation of the theory of these spaces.

As noticed that Pisier and Xu [20], Theorem 5.3, the Proposition 1 also works in the context of noncommutative L^p spaces.

Corollary 5 allows us to extend Proposition 1 as follows:

Theorem 10. *(The 2-uniform convexity functional inequality for $p \in (1, 2]$) If $p \in (1, 2]$ and $f : [0, \infty) \rightarrow \mathbb{R}$ is a nondecreasing function such that $f(0) = 0$ and $f(x^{1/2})$ is convex, then*

$$f\left(\|x\|_p\right) + f\left(\|y\|_p\right) \geq 2f\left(\left\|\frac{x+y}{2}\right\|_p\right) + [2(p-1)]f\left(\left\|\frac{x-y}{2}\right\|_p\right),$$

for all elements x and y belonging to an L^p space, to a Schatten space $S^p(H)$ or to a noncommutative L^p space.

Here $[\cdot]$ denotes the floor function ($[x] =$ the greatest integer less than or equal to x).

According to Remark 2 (b), the hypotheses of Theorem 10 are fulfilled by every nondecreasing and differentiable function $f : [0, \infty) \rightarrow \mathbb{R}$ such that $f(0) = 0$, f' is convex and $f'(0) = 0$.

Proof. Clearly

$$\left\| \frac{x \pm y}{2} \right\|_p^2 \leq \left(\frac{\|x\|_p + \|y\|_p}{2} \right)^2 \leq \max\{\|x\|_p^2, \|y\|_p^2\}$$

and

$$\left\| \frac{x+y}{2} \right\|_p^2 + \left\| \frac{x-y}{2} \right\|_p^2 \leq \|x\|_p^2 + \|y\|_p^2,$$

so the conclusion follows from Corollary 4. \square

The case $p > 2$ of Theorem 10 makes the objective of Theorem 16 in Section 6.

The next consequence of Corollary 5 makes use of the following special case of the parallelogram law: for all real numbers a and b ,

$$|a|^2 + |b|^2 = 2 \left| \frac{a-b}{2} \right|^2 + 2 \left| \frac{a+b}{2} \right|^2.$$

Corollary 6. *Suppose that $f : [0, \infty) \rightarrow [0, \infty)$ is a nondecreasing, convex, and 3-concave function such that $f(0) = 0$. Then for all real numbers a and b ,*

$$f(|a|) + f(|b|) \leq 2f\left(\left|\frac{a-b}{2}\right|\right) + 2f\left(\left|\frac{a+b}{2}\right|\right).$$

In particular, for all $\alpha \in [1, 2]$,

$$|a|^\alpha + |b|^\alpha \leq 2 \left| \frac{a-b}{2} \right|^\alpha + 2 \left| \frac{a+b}{2} \right|^\alpha.$$

See Corollary 7 below for an extension to the framework of Banach spaces.

Proof. According to Theorem 5, the function $g(x) = f(x^{1/2})$ is concave. As a consequence, the result of Corollary 6 follows from Corollary 5, when applied to the function g and to the elements

$$\begin{aligned} y_1 &= |a|^2, \quad y_2 = |b|^2, \\ x_1 = x_2 &= \left| \frac{a-b}{2} \right|^2 \quad \text{and} \quad x_3 = x_4 = \left| \frac{a+b}{2} \right|^2. \end{aligned}$$

\square

Hanner's inequalities for L^p spaces with $p \in (1, 2]$ assert that

$$(4.1) \quad (\|u\|_p + \|v\|_p)^p + \left| \|u\|_p - \|v\|_p \right|^p \leq \|u+v\|_p^p + \|u-v\|_p^p,$$

while for $p \in [2, \infty)$ these inequalities work in the reversed direction. See [?], p. 139 for details.

The argument of Corollary 6 can be easily adapted to obtain the following generalization of Hanner's aforementioned inequalities:

Theorem 11. *(The generalization of Hanner's inequalities for $p \in (1, 2]$) Let $f : [0, \infty) \rightarrow \mathbb{R}$ be a nondecreasing convex function such that $f(0) = 0$ and $f(x^{1/p})$ is concave for some $p \in (1, 2]$. Then*

$$f(\|u\|_p + \|v\|_p) + f(\left| \|u\|_p - \|v\|_p \right|) \leq f(\|u+v\|_p) + f(\|u-v\|_p)$$

for all u, v belonging to a Lebesgue space $L^p(\mu)$.

The companion of Theorem 11 for $p \geq 2$ is a direct consequence of Corollary 4:

Theorem 12. (The generalization of Hanner's inequalities for $p \in [2, \infty)$) Let $f : [0, \infty) \rightarrow \mathbb{R}$ be a nondecreasing function such that $f(0) = 0$ and $f(x^{1/p})$ is concave. Then

$$f(\|u\|_p + \|v\|_p) + f(|\|u\|_p - \|v\|_p|) \geq f(\|u + v\|_p) + f(\|u - v\|_p)$$

for all u, v belonging to a Lebesgue space $L^p(\mu)$.

As noticed by Ball, Carlen and Lieb [1], Theorem 2, the inequalities (4.1) also hold in the context of Schatten spaces $S^p(H)$ in the following two cases:

- (HS1) $1 < p \leq 4/3$; and
- (HS2) u and v belong to $S^p(H)$ for some $p \in (1, 2]$ and $u \pm v$ are positive semidefinite.

For $p \geq 2$, the inequalities (4.1) work in the reverse direction and the restriction in (HS1) becomes $p \geq 4$, and the restriction in (HS2) changes to the restriction that u and v are positive semidefinite. Subject to these restrictions, Theorem 11 and Theorem 12 continue to work in the context of Schatten spaces.

5. A SUBSTITUTE OF THE PARALLELOGRAM LAW IN THE CONTEXT OF BANACH SPACES

In connection with the famous work [10] of Jordan and von Neumann concerning the inner product spaces, Clarkson [5] has introduced the *von Neumann-Jordan constant* $C_{NJ}(X)$ of a Banach space X as

$$C_{NJ}(X) = \sup \left\{ \frac{\|u + v\|^2 + \|u - v\|^2}{2\|u\|^2 + 2\|v\|^2} : u, v \in X \text{ and } \|u\| + \|v\| \neq 0 \right\}.$$

We have $1 \leq C_{NJ}(X) \leq 2$ for all Banach spaces X and $C_{NJ}(X) = 1$ if and only if X is a Hilbert space.

In general, $C_{NJ}(X) < 2$ for any uniformly convex space. $C_{NJ}(X) = 2$ in the case of spaces $L^p(\mathbb{R})$ with $p = 1$ or $p = \infty$ and the same is true in the case of Banach spaces of continuous functions endowed with the sup norm.

Let $1 < p < \infty$ and $t = \min\{p, p/(p-1)\}$. Then

$$C_{NJ}(X) = 2^{2/t-1}$$

for each of the following Banach spaces X of dimension at least 2:

- $L^p(\mathbb{R})$ (see Clarkson [5]);
- the Sobolev spaces $X = W^{k,p}(\mathbb{R})$ (see Kato and Miyazaki [11]);
- the Schatten classes of index p , $S_p(H)$ (Kato and Takahashi [12]).

The definition of the von Neumann-Jordan constant gives rise to the following generalization of the parallelogram rule:

$$(5.1) \quad \|u\|^2 + \|v\|^2 \leq 2C_{NJ}(X) \left\| \frac{u-v}{2} \right\|^2 + 2C_{NJ}(X) \left\| \frac{u+v}{2} \right\|^2$$

for all $u, v \in X$.

The next result extends Theorem 1 to the general context of Banach spaces. Its statement makes use of a modification of the von Neumann-Jordan constant, precisely,

$$N(X) = \begin{cases} 2C_{NJ}(X) & \text{if } 2C_{NJ}(X) \text{ is an integer} \\ 4 & \text{otherwise,} \end{cases}$$

motivated by the use of Corollary 5 which deals with strings of elements indexed by integers. Notice that $N(X) = 3$ in the case of L^p spaces with $p = (2 \log 2) / (\log 3) \approx 1.261859507\dots$

Theorem 13. *Let X be a Banach space and let $f : [0, \infty) \rightarrow \mathbb{R}$ be a nondecreasing, convex and 3-concave function such that $f(0) = 0$. Then*

$$(5.2) \quad f(\|u\|) + f(\|v\|) \leq N(X)f\left(\left\|\frac{u-v}{2}\right\|\right) + N(X)f\left(\left\|\frac{u+v}{2}\right\|\right),$$

and

$$f(\|u\|) + f(\|v\|) \leq \frac{N(X)}{2} \{f(\|u+x\|) + f(\|v+x\|) + f(\|x\|) + f(\|u+v+x\|)\},$$

for all $u, v, x \in X$.

Proof. We will apply Corollary 5 to the function $f \circ \sqrt{\cdot}$ and the points

$$\begin{aligned} x_1 = \dots = x_{N(X)} &= \left\|\frac{u-v}{2}\right\|^2, \\ x_{N(X)+1} = \dots = x_{2N(X)} &= \left\|\frac{u+v}{2}\right\|^2, \\ y_1 &= \|u\|^2 \text{ and } y_2 = \|v\|^2. \end{aligned}$$

We have $\max\{x_1, x_2, \dots, x_{2N(X)}\} \leq \max\{y_1, y_2\}$ since

$$\begin{aligned} x_1 = \dots = x_{N(X)} &= \left\|\frac{u-v}{2}\right\|^2 \leq \left(\frac{\|u\| + \|v\|}{2}\right)^2 \\ &\leq \max\{\|u\|^2, \|v\|^2\} = \max\{y_1, y_2\} \end{aligned}$$

and the same is true for $x_{N(X)+1}, \dots, x_{2N(X)}$. The fact that $x_1 + x_2 + x_3 + \dots + x_{2N(X)} \geq y_1 + y_2$ follows from the inequality (5.1), while Theorem 5 assures that $f \circ \sqrt{\cdot}$ is a nondecreasing concave function. The inequality (5.2) is now clear.

The second inequality in the statement of Theorem 13 is a consequence of the inequality (5.2). Indeed, since f is convex and nondecreasing, the function $f \circ \|\cdot\|$ is also convex, which yields

$$\begin{aligned} 2f\left(\left\|\frac{u-v}{2}\right\|\right) &\leq f(\|u+x\|) + f(\|-v-x\|) \\ &= f(\|u+x\|) + f(\|v+x\|), \end{aligned}$$

and

$$2f\left(\left\|\frac{u+v}{2}\right\|\right) \leq f(\|x\|) + f(\|u+v+x\|),$$

for all $u, v, x \in \mathbb{R}^N$. □

In the particular case when f is the function x^α with $\alpha \in [1, 2]$, Theorem 13 yields the following result:

Corollary 7. *Let X be a Banach and $\alpha \in [1, 2]$. Then*

$$\|u\|^\alpha + \|v\|^\alpha \leq N(X) \left(\left\|\frac{u-v}{2}\right\|^\alpha + \left\|\frac{u+v}{2}\right\|^\alpha \right) \leq N(X)^2 (\|u\|^\alpha + \|v\|^\alpha)$$

whenever $u, v \in X$.

6. THE GENERALIZATION OF THE QUADRUPLE INEQUALITY OF SCHÖTZ

We are now in a position to state the following generalization to the context of Banach spaces of the quadruple inequality of Schötz (see Theorem 1 in the Introduction) :

Theorem 14. *Let y, z, q, r be four points in the Banach space X and let f be a nondecreasing, convex and 3-concave function defined on $[0, \infty)$ such that $f(0) = 0$. Then*

$$\begin{aligned} f(\|y - q\|) + f(\|z - r\|) \\ \leq \frac{N(X)}{2} \{f(\|y - z\|) + f(\|r - q\|) + f(\|z - q\|) + f(\|y - r\|)\}. \end{aligned}$$

In the case of inner product spaces, $N(X) = 2$ and we retrieve the quadruple inequality of Schötz.

Proof. Given four points y, z, q, r in the space X , let us denote

$$z - q = x, \quad q - y = u, \quad \text{and} \quad r - z = v.$$

Then

$$z - y = u + x, \quad r - y = u + v + x, \quad \text{and} \quad r - q = v + x,$$

so the proof ends by taking into account the second part of Theorem 13. $\square \quad \square$

In the case of L^p spaces, Clarkson (see [4], Theorem 2) noticed the following two inequalities, usually known as *the easy Clarkson inequalities*:

$$\begin{aligned} 2^{p-1} \left(\|x\|_p^p + \|y\|_p^p \right) &\leq \|x - y\|_p^p + \|x + y\|_p^p \leq 2 \left(\|x\|_p^p + \|y\|_p^p \right), \quad \text{if } p \in (1, 2], \\ 2 \left(\|x\|_p^p + \|y\|_p^p \right) &\leq \|x - y\|_p^p + \|x + y\|_p^p \leq 2^{p-1} \left(\|x\|_p^p + \|y\|_p^p \right), \quad \text{if } p \in [2, \infty). \end{aligned}$$

As a consequence, by replacing the inequalities (5.1) with the easy Clarkson inequalities and using a similar argument to the one that we used for Theorem 13 and Theorem 14, we arrive at the following companion of these theorems.

Theorem 15. *Consider an L^p space X with $p \in (1, 2]$ and let $f : [0, \infty) \rightarrow \mathbb{R}$ be a nondecreasing, convex function such that $f(0) = 0$ and $f(x^{1/p})$ is concave. Then*

$$f(\|u\|_p) + f(\|v\|_p) \leq 2f\left(\left\|\frac{u-v}{2}\right\|_p\right) + 2f\left(\left\|\frac{u+v}{2}\right\|_p\right),$$

and

$$f(\|u\|_p) + f(\|v\|_p) \leq f(\|u + x\|_p) + f(\|v + x\|_p) + f(\|x\|_p) + f(\|u + v + x\|_p),$$

for all $u, v, x \in X$. As a consequence,

$$\begin{aligned} f(\|y - q\|_p) + f(\|z - r\|_p) \\ \leq f(\|y - z\|_p) + f(\|r - q\|_p) + f(\|z - q\|_p) + f(\|y - r\|_p) \end{aligned}$$

for all $q, r, y, z \in X$.

For $p \in [2, \infty)$, these inequalities should be replaced by the following ones:

$$\begin{aligned} f(\|u\|_p) + f(\|v\|_p) &\leq C(p)f\left(\left\|\frac{u-v}{2}\right\|_p\right) + C(p)f\left(\left\|\frac{u+v}{2}\right\|_p\right), \\ f(\|u\|_p) + f(\|v\|_p) &\leq \frac{C(p)}{2} \left\{ f(\|u+x\|_p) + f(\|v+x\|_p) \right. \\ &\quad \left. + f(\|x\|_p) + f(\|u+v+x\|_p) \right\}, \end{aligned}$$

and

$$\begin{aligned} f(\|y-q\|_p) + f(\|z-r\|_p) &\leq \frac{C(p)}{2} \left\{ f(\|y-z\|_p) + f(\|r-q\|_p) \right. \\ &\quad \left. + f(\|z-q\|_p) + f(\|y-r\|_p) \right\}, \end{aligned}$$

where $C(p) = 2^{p-1}$ if 2^{p-1} is an integer and $C(p) = \lfloor 2^{p-1} + 1 \rfloor$ otherwise.

As McCarthy noticed in [16] (see also Simon [30]), the two easy Clarkson inequalities also work in the context of Schatten classes $S^p(H)$ (provided that the L^p norms $\|\cdot\|_p$ are replaced by the Schatten p norms $\|\cdot\|_p$). Accordingly, Theorem 15 still works in the framework of Schatten classes.

We end this section by noticing a quadruple inequality (similar to that stated in Theorem 15) that results in the case $p \in [2, \infty)$ of the optimal 2-uniform convexity inequality. For convenience, we denote

$$\tilde{C}(p) = \begin{cases} (p-1)/2 & \text{if } (p-1)/2 \in \mathbb{Z} \\ \lfloor (p-1)/2 + 1 \rfloor & \text{otherwise.} \end{cases}$$

Theorem 16. (The 2-uniform convexity functional inequality: case $p \in [2, \infty)$) If $p \in [2, \infty)$ and $f : [0, \infty) \rightarrow \mathbb{R}$ is a nondecreasing, convex, and 3-concave function such that $f(0) = 0$. Then

$$f(\|x\|_p) + f(\|y\|_p) \leq 2f\left(\left\|\frac{x+y}{2}\right\|_p\right) + 2\tilde{C}(p)f\left(\left\|\frac{x-y}{2}\right\|_p\right)$$

and

$$\begin{aligned} f(\|x\|_p) + f(\|y\|_p) &\leq f(\|v\|) + f(\|x+y+v\|_p) \\ &\quad + \tilde{C}(p)f(\|x+u\|_p) + \tilde{C}(p)f(\|y+u\|_p) \end{aligned}$$

for all elements x, y , and u belonging to an L^p space, to a Schatten space $S^p(H)$ or to a non-commutative L^p space.

As a consequence,

$$\begin{aligned} f(\|y-q\|_p) + f(\|z-r\|_p) &\leq f(\|z-q\|_p) + f(\|y-r\|_p) \\ &\quad + \tilde{C}(p)f(\|y-z\|_p) + \tilde{C}(p)f(\|r-q\|_p) \end{aligned}$$

for all q, r, y, z in an L^p space (or in a Schatten space $S^p(H)$).

7. FURTHER COMMENTS

Not everything can be obtained via majorization theory.

Here is an example based on Fréchet's identity (see [7]),

$$(Fr) \quad \|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 + \|\mathbf{z}\|^2 + \|\mathbf{x} + \mathbf{y} + \mathbf{z}\|^2 = \|\mathbf{x} + \mathbf{y}\|^2 + \|\mathbf{y} + \mathbf{z}\|^2 + \|\mathbf{z} + \mathbf{x}\|^2,$$

valid in any Euclidean space \mathbb{R}^N . As Ressel noticed in [26], Theorem 2, this identity implies the Hornich-Hlawka inequality in \mathbb{R}^N ,

$$(HH) \quad \|\mathbf{x}\| + \|\mathbf{y}\| + \|\mathbf{z}\| + \|\mathbf{x} + \mathbf{y} + \mathbf{z}\| \geq \|\mathbf{x} + \mathbf{y}\| + \|\mathbf{y} + \mathbf{z}\| + \|\mathbf{z} + \mathbf{x}\|,$$

which in turn yields

$$\begin{aligned} \|\mathbf{x}\|^{1/2^n} + \|\mathbf{y}\|^{1/2^n} + \|\mathbf{z}\|^{1/2^n} + \|\mathbf{x} + \mathbf{y} + \mathbf{z}\|^{1/2^n} \\ \geq \|\mathbf{x} + \mathbf{y}\|^{1/2^n} + \|\mathbf{y} + \mathbf{z}\|^{1/2^n} + \|\mathbf{z} + \mathbf{x}\|^{1/2^n} \end{aligned}$$

for all $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{R}^N$ and all integers $n = 1, 2, 3, \dots$.

One may think of the Hornich-Hlawka inequality as a particular case of a functional inequality attached to Fréchet's identity and including the square root function in its domain. So far, no such functional inequality is known, despite a great deal of attention received by the Hornich-Hlawka inequality over the years. See [15] for details.

To understand why the majorization theorem cannot be helpful for proving the implication $(Fr) \implies (HH)$, let's restrict ourselves to the positive cone \mathbb{R}_+^N . In this case one can add to Fréchet's identity the inequalities

$$\begin{aligned} \max \left\{ \|\mathbf{x}\|^2, \|\mathbf{y}\|^2, \|\mathbf{z}\|^2, \|\mathbf{x} + \mathbf{y} + \mathbf{z}\|^2 \right\} &\geq \max \left\{ \|\mathbf{x} + \mathbf{y}\|^2, \|\mathbf{y} + \mathbf{z}\|^2, \|\mathbf{z} + \mathbf{x}\|^2 \right\}, \\ \|\mathbf{z}\|^2 + \|\mathbf{x} + \mathbf{y} + \mathbf{z}\|^2 &\geq \|\mathbf{y} + \mathbf{z}\|^2 + \|\mathbf{z} + \mathbf{x}\|^2, \end{aligned}$$

and the similar ones obtained by permuting the variables. It becomes clear that none of the strings

$$X = (\|\mathbf{x}\|^2, \|\mathbf{y}\|^2, \|\mathbf{z}\|^2, \|\mathbf{x} + \mathbf{y} + \mathbf{z}\|^2)$$

and

$$Y = (\|\mathbf{x} + \mathbf{y}\|^2, \|\mathbf{y} + \mathbf{z}\|^2, \|\mathbf{z} + \mathbf{x}\|^2, 0)$$

is majorized by the other one in the sense of Hardy, Littlewood and Pólya.

However, some particular results may still be obtained.

For example, from Corollary 5 one can deduce the following fact.

Theorem 17. *For every nondecreasing and concave function $f : [0, \infty) \rightarrow \mathbb{R}$ and every $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{R}_+^N$.*

$$\begin{aligned} f(\|\mathbf{x} + \mathbf{y}\|^2) + f(\|\mathbf{y} + \mathbf{z}\|^2) + f(\|\mathbf{z} + \mathbf{x}\|^2) \\ \geq f(\|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 + \|\mathbf{z}\|^2) + f(\|\mathbf{x} + \mathbf{y} + \mathbf{z}\|^2) + f(0). \end{aligned}$$

The particular case where f is the square root function gives rise to an inequality that complements the Hornich-Hlawka inequality:

$$(7.1) \quad \|\mathbf{x} + \mathbf{y}\| + \|\mathbf{y} + \mathbf{z}\| + \|\mathbf{z} + \mathbf{x}\| \geq \left(\|\mathbf{x}\|^2 + \|\mathbf{y}\|^2 + \|\mathbf{z}\|^2 \right)^{1/2} + \|\mathbf{x} + \mathbf{y} + \mathbf{z}\|$$

for every $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{R}_+^N$.

Notice that the inequality (7.1) does not work for arbitrary elements of \mathbb{R}^N .

The square norm is strongly superadditive on the positive cone of \mathbb{R}^N , that is,

$$\|\mathbf{x} + \mathbf{y} + \mathbf{z}\|^2 + \|\mathbf{z}\|^2 \geq \|\mathbf{x} + \mathbf{z}\|^2 + \|\mathbf{y} + \mathbf{z}\|^2$$

for every $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{R}_+^N$. By proceeding as in the case of Theorem 9 one obtains the following Popoviciu like inequality.

Proposition 2. *For every $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{R}_+^N$ and every nondecreasing function $f : [0, \infty) \rightarrow \mathbb{R}$ such that $f(0) = 0$ and $f(x^{1/2})$ is convex, we have*

$$\begin{aligned} \frac{f(\|\mathbf{x}\|) + f(\|\mathbf{y}\|) + f(\|\mathbf{z}\|)}{3} + f(\|\mathbf{x} + \mathbf{y} + \mathbf{z}\|) \\ \geq \frac{2}{3}f(\|\mathbf{x} + \mathbf{y}\|) + f(\|\mathbf{y} + \mathbf{z}\|) + f(\|\mathbf{z} + \mathbf{x}\|). \end{aligned}$$

Similarly, starting from Serre's determinantal inequality [29],

$$\begin{aligned} \det^{1/2} A + \det^{1/2} B + \det^{1/2} C + \det^{1/2}(A + B + C) \\ \leq \det^{1/2}(A + B) + \det^{1/2}(B + C) + \det^{1/2}(C + A), \end{aligned}$$

valid for all triplets A, B, C of positive semidefinite matrices, one obtains from Corollary 5 the functional inequality

$$\begin{aligned} f\left(\det^{1/2}(A + B)\right) + f\left(\det^{1/2}(B + C)\right) + f\left(\det^{1/2}(C + A)\right) \\ \geq f\left(\det^{1/2} A + \det^{1/2} B + \det^{1/2} C\right) + f\left(\det^{1/2}(A + B + C)\right), \end{aligned}$$

for every nonnegative, nondecreasing and concave function $f : [0, \infty) \rightarrow \mathbb{R}$.

We leave open the problem whether Propositions 17 and 2 still work for all triplets of vectors in \mathbb{R}^N .

Acknowledgement. The author would like to thank Ștefan Cobzaș, Dan-Ștefan Marinescu and Christof Schötz for many useful comments on the subject of this paper.

REFERENCES

- [1] Ball, K., Carlen, E.A., Lieb, E.H.: Sharp uniform convexity and smoothness inequalities for trace norms. *Invent. Math.* **115**(1), 463–482 (1994).
- [2] Bencze, M., Niculescu, C.P., Popovici, F.: Popoviciu's inequality for functions of several variables. *J. Math. Anal. Appl.* **365**(1), 399–409 (2010).
- [3] Choudary, A.D.R., Niculescu, C.P.: *Real Analysis on Intervals*. Springer India (2014).
- [4] Clarkson, J.A.: Uniformly convex spaces. *Trans. Amer. Math. Soc.* **40**(3) 396–414 (1936).
- [5] Clarkson, J.A.: The von Neumann-Jordan constant for the Lebesgue space. *Ann. of Math.* **38**(1), 114–115 (1937).
- [6] Enflo, P.: Uniform structures and square roots in topological groups. II. *Israel J. Math.* **8**, 253–272 (1970).
- [7] Fréchet, M.: Sur la définition axiomatique d'une classe d'espaces vectoriels distanciés applicables vectoriellement sur l'espace de Hilbert. *Ann. of Math.* **36**, 705–718 (1935).
- [8] Hanner, O.: On the uniform convexity of L^p and ℓ^p . *Ark. Mat.* **3**(3), 239–244 (1956).
- [9] Hopf, E.: Über die Zusammenhänge zwischen gewissen höheren Differenzenquotienten reeller Funktionen einer reellen Variablen und deren Differenzierbarkeitseigenschaften. *Dissertation*, Univ. Berlin (1926).
- [10] Jordan, P., von Neumann, J.: On inner products in linear metric spaces. *Ann. of Math.* **36**, 719–723 (1935).

- [11] Kato, M., Miyazaki, K.: On generalized Clarkson's inequalities for $L_p(\mu; L_q(\nu))$ and Sobolev spaces. *Math. Japon.* **43**, 505-515 (1996).
- [12] Kato, M., Takahashi, Y.: Von Neumann–Jordan constant for Lebesgue–Bochner spaces. *J. Inequal. Appl.* **2**(1), 89-97 (1998).
- [13] Kato, M., Takahashi, Y.: On optimal 2-uniform convexity inequalities. *Nonlinear Analysis and Convex Analysis* 1298, 65-69 (2002).
- [14] Marinescu, D-Ş., Niculescu, C.P.: Old and new on the 3-convex functions. *Math. Inequal. Appl.* **26**(4), 911–933 (2023).
- [15] Marinescu, D-Ş., Niculescu, C.P.: A survey of the Hornich-Hlawka inequality. Preprint arXiv <http://arxiv.org/abs/2407.03278> (2024).
- [16] McCarthy, J.: C_p . *Israel J. Math.* **5**, 249-271 (1967).
- [17] Niculescu, C.P.: A new look at the Hardy-Littlewood-Polya inequality of majorization. *J. Math. Anal. Appl.* **501**(2), article no. 125211 (2021).
- [18] Niculescu, C.P., Persson, L.- E.: *Convex Functions and their Applications. A Contemporary Approach.* (Third edition) CMS Books in Mathematics, Springer (2024).
- [19] Niculescu, C.P., Sra, S.: Functional inequalities for functions with positive differences on convex cones. *Math. Results* 78(6), article no. 217 (2023).
- [20] Pisier, G., Xu, Q.: Non-commutative L_p -spaces. In *Handbook of the geometry of Banach spaces 2*, 1459-1517. North-Holland, Amsterdam (2003).
- [21] Pólya, G.: Remark on Weyl's note "Inequalities between the two kinds of eigenvalues of a linear transformation", *Proc. Natl. Acad. Sci. U.S.A.* **36**, 49–51 (1950).
- [22] Popoviciu, T.: Sur quelques propriétés des fonctions d'une ou de deux variables réelles. *Mathematica (Cluj)* **8**, 1-85 (1934).
- [23] Popoviciu, T.: Sur l'approximation des fonctions convexes d'ordre supérieur. *Mathematica (Cluj)* **10**, 49-54 (1935).
- [24] Popoviciu, T.: *Les Fonctions Convexes.* Hermann Cie. Editeurs, Paris (1944).
- [25] Popoviciu, T.: Sur certaines inégalités qui caractérisent les fonctions convexes. *Analele ştiinţifice Univ. "Al. I. Cuza" Iasi, Secţia I-a Mat.* **11**, 155–164 (1965).
- [26] Ressel, P.: The Hornich–Hlawka inequality and Bernstein functions. *J. Math. Inequal.* **9**, 883–888 (2015).
- [27] Schilling, R.L., Song, R., Vondraček, Z.: *Bernstein Functions. Theory and Applications.* De Gruyter, Berlin (2010).
- [28] Schötz, C.: Quadruple inequalities: Between Cauchy-Schwarz and triangle. ArXiv preprint, [arXiv:2307.01361 v.3](https://arxiv.org/abs/2307.01361) (2024).
- [29] Serre, D.: The reverse Hlawka inequality in a Minkowski space. *C. R. Math. Acad. Sci. Paris* **353**(7), 629–633 (2015).
- [30] Simon, B.: *Trace ideals and their applications.* Second edition. *Mathematical Surveys and Monographs* vol. **120**, American Mathematical Society, Providence, RI (2005).
- [31] Tomić, M.: Théorème de Gauss relatif au centre de gravité et son application. *Bull. Soc. Math. Phys. Serbie* **1**, 31–40 (1949)
- [32] Weyl, H.: Inequalities between two kinds of eigenvalues of a linear transformation. *Proc. Nat. Acad. Sci. U. S. A.* **35**, 408–411 (1949)
- [33] Zhang, F.: *Matrix Theory: Basic Results and Techniques.* 2nd ed. Springer, New York (2011).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CRAIOVA, CRAIOVA 200585, ROMANIA
Email address: constantin.p.niculescu@gmail.com