

CLASSIFYING THE POLISH SEMIGROUP TOPOLOGIES ON THE SYMMETRIC INVERSE MONOID

S. BARDYLA, L. ELLIOTT, J. D. MITCHELL, AND Y. PÉRESSE

ABSTRACT. We classify all Polish semigroup topologies on the symmetric inverse monoid $I_{\mathbb{N}}$ on the natural numbers. This result answers a question of Elliott et al. There are countably infinitely many such topologies. Under containment, these Polish semigroup topologies form a join-semilattice with infinite descending chains, no infinite ascending chains, and arbitrarily large finite anti-chains. Also, we show that the monoid $I_{\mathbb{N}}$ endowed with any second countable T_1 semigroup topology is homeomorphic to the Baire space $\mathbb{N}^{\mathbb{N}}$.

1. INTRODUCTION

Recall that the full transformation monoid X^X on a set X consists of all functions from X to X under the operation of composition of functions. The monoid X^X plays the same role in semigroup theory, as the symmetric group S_X does for groups. That is, by an analogue of Cayley's Theorem every semigroup embeds into some full transformation monoid. If X is countable, then X^X naturally has the topology of the Baire space, i.e. the product topology on X^X arising from the discrete topology on X . This topology is compatible with the operation of X^X (the compositions of functions), making X^X a topological semigroup. The Baire space is a Polish space, a separable completely metrizable topological space. Polish spaces and groups are widely studied in descriptive set theory, see [2, 3, 5, 8, 9, 13, 14, 15, 16, 17, 22, 25, 26].

The symmetric group $S_{\mathbb{N}}$ is a G_{δ} subspace of $\mathbb{N}^{\mathbb{N}}$, and hence is a Polish space also. In addition to the multiplication in $S_{\mathbb{N}}$ being continuous, inversion is also continuous, making $S_{\mathbb{N}}$ a Polish topological group. In fact, the topology of the Baire space is intrinsically linked to the monoid $\mathbb{N}^{\mathbb{N}}$ and the group $S_{\mathbb{N}}$. That is, the Baire space topology is the unique Polish semigroup topology on $\mathbb{N}^{\mathbb{N}}$ [7, Theorem 5.4]; and, similarly, the subspace topology inherited from the Baire space is the unique Polish group topology on $S_{\mathbb{N}}$ [10, 18, 27]. For more about Polish topologies on groups and semigroups and their relation to automatic continuity see [1, 4, 6, 7, 11, 19, 21, 23, 28, 29, 30].

Inverse semigroups are semigroups satisfying a "local inverses" axiom: for every $x \in S$ there exists a unique $x^{-1} \in S$ such that $xx^{-1}x = x$ and $x^{-1}xx^{-1} = x^{-1}$. Lying between general semigroups, which are rich in their diversity, and groups, which are rich in structure, inverse semigroups have been widely studied in the literature; see [12, 20, 24] and references therein. The symmetric inverse monoid I_X consists of all bijections between subsets of X under the usual composition of binary relations. By the Wagner-Preston Theorem [12, Theorem 5.1.7], I_X plays the same role for the class of inverse semigroups as X^X for semigroups, or S_X for groups. It was shown in [7, Theorem 5.15] that $I_{\mathbb{N}}$ also possesses a unique Polish semigroup topology \mathcal{I}_4 under which the inversion function $x \mapsto x^{-1}$ is also continuous, and two further Polish semigroup topologies \mathcal{I}_2 and \mathcal{I}_3 , which are defined as follows: For every $x, y \in \mathbb{N}$ let,

$$(1) \quad U_{x,y} = \{h \in I_{\mathbb{N}} : (x,y) \in h\}, \quad W_x = \{h \in I_{\mathbb{N}} : x \notin \text{dom}(h)\}, \quad W_x^{-1} = \{h \in I_{\mathbb{N}} : x \notin \text{im}(h)\}.$$

Then a subbasis for \mathcal{I}_2 is

$$\{U_{x,y} : x, y \in \mathbb{N}\} \cup \{W_x : x \in \mathbb{N}\};$$

The first named author was supported by the Austrian Science Fund FWF (Grant ESP 399).

a subbasis for \mathcal{I}_3 is

$$\{U_{x,y} : x, y \in \mathbb{N}\} \cup \{W_x^{-1} : x \in \mathbb{N}\};$$

and \mathcal{I}_4 is the topology generated by $\mathcal{I}_2 \cup \mathcal{I}_3$.

It was also shown in [7] that every Polish semigroup topology for $I_{\mathbb{N}}$ is contained in \mathcal{I}_4 and contains either \mathcal{I}_2 or \mathcal{I}_3 which led the authors to ask the following question:

Question 1.1 ([7, Question 5.17]). *Are \mathcal{I}_2 , \mathcal{I}_3 , and \mathcal{I}_4 the only Polish semigroup topologies on $I_{\mathbb{N}}$?*

In this paper, we answer Question 1.1 by classifying all of the Polish semigroup topologies on the symmetric inverse monoid $I_{\mathbb{N}}$; Definition 2.2, Theorem 3.10, and Theorem 2.3. In particular, there are countably many such semigroup topologies, forming a join-semilattice with: infinite descending chains, no infinite ascending chains, and arbitrarily long finite anti-chains; see Corollary 2.7. If ω denotes the first infinite ordinal, then each such Polish semigroup topology is characterised by a special kind of function $f : \omega + 1 \rightarrow \omega + 1$ that we will refer to as *waning*; see Definition 2.2. The order on the join-semilattice of the Polish semigroup topologies on $I_{\mathbb{N}}$ is also completely characterised in terms of the corresponding waning functions, see Theorem 2.6. Also, we show that the symmetric inverse monoid $I_{\mathbb{N}}$ endowed with any T_1 second countable semigroup topology is homeomorphic to the Baire space $\mathbb{N}^{\mathbb{N}}$, see Theorem 2.5.

This paper is organised as follows: in Section 2 we provide the necessary definitions, and then state the main theorems; in Section 3 we show how to associate a Polish semigroup topology on $I_{\mathbb{N}}$ to every waning function; and finally in Section 4 we show that every Polish semigroup topology on $I_{\mathbb{N}}$ equals one of the topologies from Section 3.

2. MAIN RESULTS

In this section, after first introducing the required definitions and notation, we state the main theorems of this paper.

The topologies on $I_{\mathbb{N}}$ arising from functions from $\omega + 1$ to $\omega + 1$ are defined as follows.

Definition 2.1 (The topology \mathcal{T}_f). Suppose that $f : \omega + 1 \rightarrow \omega + 1$ is any function. We define \mathcal{T}_f to be the least topology on $I_{\mathbb{N}}$ containing \mathcal{I}_2 and the sets of the form:

$$U_{f,n,X} = \{g \in I_{\mathbb{N}} \mid |\text{im}(g) \setminus X| \geq n \text{ and } |X \cap \text{im}(g)| \leq (n)f\}$$

where $n \in \mathbb{N}$ and $X \subseteq \mathbb{N}$ is finite.

Roughly speaking, the open sets $U_{f,n,X}$ consist of those elements of $I_{\mathbb{N}}$ which try not to map any points into X . An element $g \in U_{f,n,X}$ must map at least n points outside of X and is allowed at most $(n)f$ mistakes, i.e. points in the image of g in X .

It will transpire that every function $f : \omega + 1 \rightarrow \omega + 1$ gives rise to a Polish semigroup topology \mathcal{T}_f on $I_{\mathbb{N}}$ (Theorem 3.10). However, distinct functions can give rise to the same topology. To circumvent this, we require the following definition.

Definition 2.2 (Waning function). We say that a non-increasing function $f : \omega + 1 \rightarrow \omega + 1$ is *waning* if either: f is constant with value ω ; or there exists $i \in \omega$ such that $(i)f \in \omega$, and $(j+1)f < (j)f$ for all $j \in \omega$ such that $0 \neq (j)f \in \omega$.

For example, the constant function $\bar{0}$ with value 0 is a waning function, and so too is the following function:

$$(x)f = \begin{cases} \omega & \text{if } x \leq 42 \\ 1337 - x & \text{if } 42 \leq x < 69 \\ 0 & \text{if } x \geq 69. \end{cases}$$

If f is a waning function, continuing to speak roughly, partial functions in $U_{f,n,X}$ defined on more points are allowed fewer mistakes.

For a topology \mathcal{T} on $I_{\mathbb{N}}$ let $\mathcal{T}^{-1} = \{U^{-1} : U \in \mathcal{T}\}$. It is routine to verify that if \mathcal{T} is a (Polish) semigroup topology on $I_{\mathbb{N}}$, then so is \mathcal{T}^{-1} . The following result answers Question 1.1 and is the main result of this paper.

Theorem 2.3. *If \mathcal{T} is a T_1 second countable semigroup topology on $I_{\mathbb{N}}$, then there exists a waning function f such that either $\mathcal{T} = \mathcal{T}_f$, or $\mathcal{T}^{-1} = \mathcal{T}_f$.*

Conversely, if f and g are distinct waning functions, then $\mathcal{T}_f \neq \mathcal{T}_g$.

Clearly, there are only countably many waning functions, and so Theorem 2.3 has the following corollary.

Corollary 2.4. *The symmetric inverse monoid has countably infinitely many T_1 distinct second countable semigroup topologies.*

As noted earlier, every T_1 and second-countable topology for $I_{\mathbb{N}}$ is Polish. In fact, up to homeomorphism, there is only one such topology:

Theorem 2.5. *The symmetric inverse monoid $I_{\mathbb{N}}$ endowed with any T_1 second countable semigroup topology is homeomorphic to the Baire space $\mathbb{N}^{\mathbb{N}}$.*

Proof. By Corollary 3.9, the spaces $(I_{\mathbb{N}}, \mathcal{T}_f)$ and $(I_{\mathbb{N}}, \mathcal{T}_f^{-1})$ are Polish and zero-dimensional for every waning function f . By the classical Alexandrov-Urysohn Theorem [17, Theorem 7.7], a non-empty Polish zero-dimensional topological space X is homeomorphic to the Baire space $\mathbb{N}^{\mathbb{N}}$ if and only if each compact subset of X has empty interior. Recall that for each waning function f the spaces $(I_{\mathbb{N}}, \mathcal{T}_f)$ and $(I_{\mathbb{N}}, \mathcal{T}_f^{-1})$ are homeomorphic. By Theorem 2.3, it is enough to show that $(I_{\mathbb{N}}, \mathcal{T}_f)$ is homeomorphic to the Baire space for each waning function f . Since all the topologies \mathcal{T}_f fall between \mathcal{I}_2 and \mathcal{I}_4 , it suffices to check that the sets which are compact with respect to \mathcal{I}_2 have empty interior with respect to \mathcal{I}_4 . If this was not the case, then there would be a basic open set U in \mathcal{I}_4 whose closure in $(I_{\mathbb{N}}, \mathcal{I}_2)$ is compact. Then there exist $n \in \mathbb{N}$, a finite subset $X \subset \mathbb{N}$ and a finite partial bijection $h : n \rightarrow \mathbb{N} \setminus X$ such that

$$U = \{f \in I_{\mathbb{N}} \mid f \upharpoonright_n = h \text{ and } \text{im}(f) \cap X = \emptyset\}.$$

It is straightforward to check that the set U is closed with respect to \mathcal{I}_2 and the open cover

$$\mathcal{U} = \{\{f \in I_{\mathbb{N}} \mid (n)f = m\} \mid m \in \mathbb{N}\} \cup \{\{f \in I_{\mathbb{N}} \mid n \notin \text{dom}(f)\}\}$$

has no finite subcover. □

As discussed in the introduction, we now address the question of how the Polish semigroup topologies on $I_{\mathbb{N}}$ are ordered with respect to containment. The set $(\omega + 1)^{(\omega + 1)}$ of all functions from $\omega + 1$ to $\omega + 1$ can be partially ordered by coordinate-wise comparison arising from the reverse of the usual well-ordering on $\omega + 1$. In other words, if $f, g \in (\omega + 1)^{(\omega + 1)}$ we write $f \preceq g$ whenever $(i)f \geq (i)g$ for all $i \in \omega + 1$. The reason that we order $f \preceq g$ when f is coordinatewise greater than g will become apparent shortly. We denote by \mathfrak{W} the set of all waning functions on $\omega + 1$ with the partial order \preceq .

Recall that, by [7], every T_1 second-countable semigroup topology on $I_{\mathbb{N}}$ contains \mathcal{I}_2 or $\mathcal{I}_3 = \mathcal{I}_2^{-1}$ and is contained in $\mathcal{I}_4 = \mathcal{I}_4^{-1}$. It follows that the entire lattice of such semigroup topologies is described by the following theorem.

Theorem 2.6. *The function $f \mapsto \mathcal{T}_f$ is an order-isomorphism from \mathfrak{W} to the interval of the lattice of Polish semigroup topologies on $I_{\mathbb{N}}$ with respect to containment between \mathcal{I}_2 and \mathcal{I}_4 .*

Similarly, $f \mapsto \mathcal{T}_f^{-1}$ is an order-isomorphism from \mathfrak{W} to the interval between \mathcal{I}_3 and \mathcal{I}_4 .

The reason for defining \preceq as we did, should now be clear: to avoid having an ‘‘anti-embedding’’ and ‘‘anti-isomorphism’’ in the statement of Theorem 2.6. It is possible to show that (\mathfrak{W}, \prec) contains infinite descending linear orders, only finite ascending linear orders, and every finite partial order.


FIGURE 1. A Hasse diagram of part of the poset \mathfrak{W} .

Figure 1 shows a Hasse diagram of some of the Polish semigroup topologies ordered by inclusion. We obtain the following corollary to Theorem 2.6.

Corollary 2.7. *The partial order of Polish semigroup topologies on $I_{\mathbb{N}}$ contains the following:*

- (a) *infinite descending linear orders;*
- (b) *only finite ascending linear orders; and*
- (c) *every finite partial order.*

3. FROM WANING FUNCTIONS TO SEMIGROUP TOPOLOGIES

In this section we will prove the following for any waning function $f : \omega + 1 \rightarrow \omega + 1$:

- (a) in Lemma 3.5 we establish a convenient neighbourhood basis for \mathcal{T}_f at every $g \in I_{\mathbb{N}}$;
- (b) in Theorem 3.10 we show that \mathcal{T}_f (defined in Definition 2.2) is a Polish semigroup topology on the symmetric inverse monoid $I_{\mathbb{N}}$;
- (c) in Lemma 3.11 we show that if $f, h : \omega \rightarrow \omega$ are waning functions, then $\mathcal{T}_f \subseteq \mathcal{T}_h$ if and only if $f \preceq h$.

In terms of proving the main results of the paper, (b) above proves one direction of Theorem 2.3, and (c) together with Theorem 2.3 proves Theorem 2.6.

Let $g \in I_{\mathbb{N}}$. We denote by $|g|$ the size of g as a subset of $\mathbb{N} \times \mathbb{N}$ (i.e. $|g| = |\{(x, (x)g) : x \in X\}|$), and by $g \upharpoonright_r := \{(x, y) \in g \mid x \in r\}$ the restriction of g to $\{0, \dots, r-1\}$.

Lemma 3.1. *If f is a waning function and $g \in I_{\mathbb{N}}$, then there exist $r \in \mathbb{N}$ such that $(r)f \leq (|g|)f = (g \upharpoonright_r)f$.*

Proof. If f is constant with value ω , then we define $r = 0$ and so $(r)f = \omega \leq \omega = (|g|)f = (g \upharpoonright_r)f$.

Suppose that f is not a constant function with value ω . If $|g| < \omega$, then we define $r \in \mathbb{N}$ so that $r > \max(\text{dom}(g))$. In this case, $g \upharpoonright_r = g$ and $(r)f \leq (|g|)f$ since $|g| < r$. This implies that $(r)f \leq (|g|)f = (g \upharpoonright_r)f$.

If $|g| = \omega$, then we choose r large enough so that $(g \upharpoonright_r)f = 0$. Since $r \geq |g \upharpoonright_r|$, it follows that $(r)f \leq (g \upharpoonright_r)f = 0 = (\omega)f = (|g|)f$. \square

For any waning function f , any $g \in I_{\mathbb{N}}$, and any $r \in \mathbb{N}$ such that the condition in Lemma 3.1 holds we define:

$$(2) \quad W_{f,g,r} = \{h \in I_{\mathbb{N}} : h \upharpoonright_r = g \upharpoonright_r \text{ and } |\text{im}(h) \cap (r \setminus \text{im}(g))| \leq (|g|)f\}.$$

Lemma 3.2. *If f is a waning function, $g \in I_{\mathbb{N}}$, and $r \in \mathbb{N}$ are such that $W_{f,g,r}$ is defined, then for any $p \geq r$, $W_{f,g,p}$ is defined and $W_{f,g,p} \subseteq W_{f,g,r}$.*

Proof. Assume that $W_{f,g,r}$ is defined, i.e. $(r)f \leq (|g|)f = (|g\upharpoonright_r|)f$. Fix any $p \geq r$. Since f is a waning function and by the choice of r we have the following:

$$(p)f \leq (r)f \leq (|g|)f = (|g\upharpoonright_r|)f \geq (|g\upharpoonright_p|)f \geq (|g|)f.$$

It follows that $(p)f \leq (|g|)f = (|g\upharpoonright_p|)f$ and, consequently, the set $W_{f,g,p}$ is defined. The inclusion $W_{f,g,p} \subseteq W_{f,g,r}$ is trivial. \square

Definition 3.3. For all $f \in (\omega + 1)^{(\omega+1)}$ we define $f' : \omega + 1 \rightarrow \omega + 1$ to be the unique waning function obtained inductively as follows:

- Let $(0)f' = (0)f$.
- If $i \in \omega$ and $(i)f = 0$, then let $(i+1)f' = \min\{(i+1)f, (i)f' - 1\}$.
- If $i \in \omega$ and $(i)f = 0$, then let $(i+1)f = 0$.
- if $(i)f' = \omega$ for all $i \in \omega$, then let $(\omega)f' = \omega$, and otherwise let $(\omega)f' = 0$.

In other words, for every $i \in \omega$ we have

$$(i)f' = \max(0, \min\{(j)f - (i-j) : j \leq i\}),$$

where we use the convention that $\omega - n = \omega$ for every $n \in \omega$. Note that f' is equal to the minimum waning function with respect to the order \leq on $(\omega + 1)^{(\omega+1)}$ such that $(i)f' \leq (i)f$ for all $i \in \omega$. In particular, $f' = f$ whenever f is a waning function.

Lemma 3.4 (Small open sets). *Let $f \in (\omega + 1)^{(\omega+1)}$, $g \in I_{\mathbb{N}}$, and $r \in \mathbb{N}$ satisfies $(r)f' \leq (|g|)f' = (|g\upharpoonright_r|)f'$. Then the set $W_{f',g,r}$ is an open neighborhood of g in $(I_{\mathbb{N}}, \mathcal{T}_f)$.*

Proof. By Definition 3.3, there is some $j \leq |g\upharpoonright_r|$ such that $(|g\upharpoonright_r|)f' \geq (j)f - (|g\upharpoonright_r| - j)$. Put

$$b = (|g\upharpoonright_r|)f' + |g\upharpoonright_r| - (j)f.$$

Observe that if $(|g\upharpoonright_r|)f' \neq 0$, then $(|g\upharpoonright_r|)f' = (j)f - (|g\upharpoonright_r| - j)$ and hence $b = j$, and if $(|g\upharpoonright_r|)f' = 0$, then $b = |g\upharpoonright_r| - (j)f \geq j$. Note that in either case we have the following:

- (i) $j \leq b \leq |g\upharpoonright_r|$.
- (ii) $(|g\upharpoonright_r|)f' = (j)f - (|g\upharpoonright_r| - b)$.

Let $Y = r \setminus \text{im}(g)$ and $Z \subseteq \text{im}(g\upharpoonright_r)$ be such that $|Z| = |g\upharpoonright_r| - b$ and hence

$$(3) \quad (|g\upharpoonright_r|)f' = (j)f - |Z|.$$

Let $X = Y \cup Z$ and note that $b = |\text{im}(g\upharpoonright_r) \setminus Z| = |g\upharpoonright_r| - |Z|$. Then $\text{im}(g\upharpoonright_r) \setminus X = \text{im}(g\upharpoonright_r) \setminus Z$ and so

$$(4) \quad |\text{im}(g) \setminus X| \geq |\text{im}(g\upharpoonright_r) \setminus X| = |\text{im}(g\upharpoonright_r) \setminus Z| = b \geq j,$$

and $X \cap \text{im}(g) = Z \cap \text{im}(g)$ and so

$$|X \cap \text{im}(g)| = |Z \cap \text{im}(g)| = |Z| = |g\upharpoonright_r| - b = (j)f - (|g\upharpoonright_r|)f' \leq (j)f.$$

So $g \in U_{f,j,X} = \{l \in I_{\mathbb{N}} \mid |\text{im}(l) \setminus X| \geq j \text{ and } |X \cap \text{im}(l)| \leq (j)f\}$. Moreover

$$\begin{aligned} g &\in \{l \in I_{\mathbb{N}} \mid l\upharpoonright_r = g\upharpoonright_r\} \cap U_{f,j,X} \\ &= \{l \in I_{\mathbb{N}} \mid l\upharpoonright_r = g\upharpoonright_r \text{ and } |\text{im}(l) \setminus X| \geq j \text{ and } |X \cap \text{im}(l)| \leq (j)f\} \\ &= \{l \in I_{\mathbb{N}} \mid l\upharpoonright_r = g\upharpoonright_r \text{ and } |X \cap \text{im}(l)| \leq (j)f\} && \text{(by 4)} \\ &= \{l \in I_{\mathbb{N}} \mid l\upharpoonright_r = g\upharpoonright_r \text{ and } |Z| + |\text{im}(l) \cap Y| \leq (j)f\} && \text{(as } Z \subseteq \text{im}(g\upharpoonright_r) = \text{im}(l\upharpoonright_r)) \\ &= \{l \in I_{\mathbb{N}} \mid l\upharpoonright_r = g\upharpoonright_r \text{ and } |\text{im}(l) \cap Y| \leq (j)f - |Z|\} \\ &= \{l \in I_{\mathbb{N}} \mid l\upharpoonright_r = g\upharpoonright_r \text{ and } |\text{im}(l) \cap Y| \leq (|g\upharpoonright_r|)f'\} && \text{(by 3)} \\ &= \{l \in I_{\mathbb{N}} \mid l\upharpoonright_r = g\upharpoonright_r \text{ and } |\text{im}(l) \cap Y| \leq (|g|)f'\} \\ &= \{l \in I_{\mathbb{N}} \mid l\upharpoonright_r = g\upharpoonright_r \text{ and } |\text{im}(l) \cap (r \setminus \text{im}(g))| \leq (|g|)f'\} = W_{f',g,r}. \end{aligned} \quad \square$$

Lemma 3.5 (Neighbourhood basis). *If f is a waning function and $g \in I_{\mathbb{N}}$, then the family*

$$\mathcal{B}_f(g) = \{W_{f,g,r} : (r)f \leq (|g|)f = (|g \upharpoonright_r|)f\}$$

forms an open neighbourhood basis at g in $(I_{\mathbb{N}}, \mathcal{T}_f)$.

Proof. Clearly, for each open neighbourhood V of g in $(I_{\mathbb{N}}, \mathcal{I}_2)$, there exists large enough $r \in \mathbb{N}$ such that the set $W_{f,g,r}$ is open in $(I_{\mathbb{N}}, \mathcal{T}_f)$ (see Lemma 3.4) and $g \in W_{f,g,r} \subseteq V$. Let $n \in \mathbb{N}$ and X be a finite subset of \mathbb{N} such that

$$g \in U_{f,n,X} = \{h \in I_{\mathbb{N}} : |\text{im}(h) \setminus X| \geq n \text{ and } |X \cap \text{im}(h)| \leq (n)f\}.$$

Let $r \in \mathbb{N}$ be larger than all the elements of X , and large enough that $|\text{im}(g \upharpoonright_r) \setminus X| \geq n$ and $(r)f \leq (|g|)f = (|g \upharpoonright_r|)f$. By Lemma 3.4, $g \in W_{f,g,r}$ and $W_{f,g,r} \in \mathcal{T}_f$. We show that $W_{f,g,r} \subseteq U_{f,n,X}$.

Let $h \in W_{f,g,r}$. As $h \upharpoonright_r = g \upharpoonright_r$ we have $|\text{im}(h) \setminus X| \geq n$. It remains to show that $|X \cap \text{im}(h)| \leq (n)f$. If $(n)f = \omega$, then there is nothing to show. Assume that $(n)f \in \omega$. By the assumption we have that

$$|\text{im}(h) \cap (r \setminus \text{im}(g))| \leq (|g|)f.$$

So, $|\text{im}(h) \cap (X \setminus \text{im}(g))| \leq (|g|)f$ and hence

$$\begin{aligned} |\text{im}(h) \cap X| &= |\text{im}(h) \cap \text{im}(g) \cap X| + |\text{im}(h) \cap (X \setminus \text{im}(g))| \\ &\leq |\text{im}(g) \cap X| + |\text{im}(h) \cap (r \setminus \text{im}(g))| \\ &\leq |\text{im}(g) \cap X| + (|g|)f \\ &\leq |\text{im}(g) \cap X| + (n + |\text{im}(g) \cap X|)f \quad (\text{as } |\text{im}(g) \setminus X| \geq n). \end{aligned}$$

If $(n + |\text{im}(g) \cap X|)f = 0$, then $|\text{im}(h) \cap X| \leq |\text{im}(g) \cap X| \leq (n)f$, as $g \in U_{f,n,X}$. If $(n + |\text{im}(g) \cap X|)f > 0$, then taking into account that f is a waning function, we get that

$$(n + |\text{im}(g) \cap X|)f \leq (n)f - |\text{im}(g) \cap X|.$$

In the latter case $|\text{im}(h) \cap X| \leq |\text{im}(g) \cap X| + (n)f - |\text{im}(g) \cap X| = (n)f$. Hence $|X \cap \text{im}(h)| \leq (n)f$, witnessing that $W_{f,g,r} \subseteq U_{f,n,X}$. \square

Lemmas 3.4 and 3.5 imply the following.

Corollary 3.6. *If $f \in (\omega + 1)^{(\omega+1)}$ is a function, then $\mathcal{T}_{f'} \subseteq \mathcal{T}_f$.*

Lemma 3.5 implies the following, which will be important in Section 4.

Remark 3.7. For any waning function f the following assertions hold:

- For each $h \in I_{\mathbb{N}}$ such that $(|h|)f = 0$, a set $U \subseteq I_{\mathbb{N}}$ is a neighbourhood of h in \mathcal{T}_f if and only if U is a neighbourhood of h in \mathcal{I}_4 .
- For each $h \in I_{\mathbb{N}}$ such that $(|h|)f = \omega$, a set $U \subseteq I_{\mathbb{N}}$ is a neighbourhood of h in \mathcal{T}_f if and only if U is a neighbourhood of h in \mathcal{I}_2 .

We will now show that the topologies \mathcal{T}_f are Polish semigroup topologies for $I_{\mathbb{N}}$. We first show that the topology generated by \mathcal{I}_2 and any single set $U_{f,n,X}$ is Polish.

Lemma 3.8. *If $f : \omega + 1 \rightarrow \omega + 1$, $n \in \mathbb{N}$ and X is a finite subset of \mathbb{N} , then the topology $\mathcal{T}_{n,X}$ generated by $\mathcal{I}_2 \cup \{U_{f,n,X}\}$ is Polish and zero-dimensional.*

Proof. The set

$$\{g \in I_{\mathbb{N}} \mid |\text{im}(g) \setminus X| \geq n\}$$

is open in $(I_{\mathbb{N}}, \mathcal{I}_2)$. Hence we can find a family of clopen sets $\{U_m \mid m \in \mathbb{N}\} \subseteq \mathcal{I}_2$ with

$$\bigcup_{m \in \mathbb{N}} U_m = \{g \in I_{\mathbb{N}} \mid |\text{im}(g) \setminus X| \geq n\}.$$

For each $m \in \mathbb{N}$, let $U_{f,n,X,m} = U_{f,n,X} \cap U_m$. By definition these sets are open in $\mathcal{T}_{n,X}$. Moreover,

$$U_{f,n,X} = \bigcup_{m \in \mathbb{N}} U_{f,n,X,m}.$$

So the topology $\mathcal{T}_{n,X}$ is generated by $\mathcal{I}_2 \cup \{U_{f,n,X}\}$ is equal to the topology generated by $\mathcal{I}_2 \cup \{U_{f,n,X,m} : m \in \mathbb{N}\}$.

Since \mathcal{I}_2 is Polish, Lemma 13.2 from [17] implies that for each $m \in \mathbb{N}$ the topology generated by $\mathcal{I}_2 \cup \{U_{f,n,X,m}\}$ is Polish if $U_{f,n,X,m}$ is closed in \mathcal{I}_2 . Lemma 13.3 from [17] then implies that the topology $\mathcal{T}_{n,X}$ is Polish as well.

Thus we need only show that for every $m \in \mathbb{N}$ the set $U_{f,n,X,m}$ is closed in $(I_{\mathbb{N}}, \mathcal{I}_2)$. Note that for a given basis \mathcal{B} of \mathcal{I}_2 consisting of clopen sets the family $\mathcal{B} \cup \{U_{f,n,X,m} : m \in \mathbb{N}\}$ will be a subbasis for $\mathcal{T}_{n,X}$ consisting of clopen sets.

Fix an arbitrary $m \in \mathbb{N}$ and $g \in I_{\mathbb{N}} \setminus U_{f,n,X,m}$. If $g \notin U_m$ then we define

$$W_g = I_{\mathbb{N}} \setminus U_m.$$

Since the set U_m is closed in $(I_{\mathbb{N}}, \mathcal{I}_2)$, W_g is an open neighbourhood of g in $(I_{\mathbb{N}}, \mathcal{I}_2)$ which is disjoint from $U_{f,n,X,m}$.

If $g \in U_m \setminus U_{f,n,X,m}$, then $|\text{im}(g) \setminus X| \geq n$ and $|X \cap \text{im}(g)| > (n)f$. We define

$$W_g = \{h \in I_{\mathbb{N}} : h|_{(\text{im}(g) \cap X)g^{-1}} \subseteq g\}.$$

Again, W_g is an open neighbourhood of g in $(I_{\mathbb{N}}, \mathcal{I}_2)$ which is disjoint from $U_{f,n,X,m}$. So the set $U_{f,n,X,m}$ is closed in $(I_{\mathbb{N}}, \mathcal{I}_2)$ as required. \square

Corollary 3.9. *If $f : \omega + 1 \rightarrow \omega + 1$ is any function, then the topology \mathcal{T}_f is Polish and zero-dimensional.*

Proof. Since \mathcal{T}_f is generated by the union of the $\mathcal{T}_{n,X}$ where $n \in \mathbb{N}$ and $X \subset \mathbb{N}$ is finite, Lemma 13.3 from [17] implies that the space $(I_{\mathbb{N}}, \mathcal{T}_f)$ is Polish. \square

Theorem 3.10. *For each waning function f , $(I_{\mathbb{N}}, \mathcal{T}_f)$ is a Polish topological semigroup.*

Proof. By Corollary 3.9, the space $(I_{\mathbb{N}}, \mathcal{T}_f)$ is Polish. Fix an arbitrary $a, b, c \in I_{\mathbb{N}}$ such that $ab = c$. If $(|c|)f = \omega$, then by Remark 3.7 the neighbourhoods of c in \mathcal{T}_f and \mathcal{I}_2 coincide. Since \mathcal{I}_2 is a semigroup topology and $\mathcal{I}_2 \subseteq \mathcal{T}_f$ we get that the semigroup operation is continuous at the point (a, b) . Assume that $(|c|)f \in \omega$. Fix an arbitrary open neighborhood $W_{f,c,r}$ of c (see Lemma 3.5). Choose a positive integer p that satisfies the following conditions:

- (1) $(p)f \leq (|a|)f = (|a|_p)f$;
- (2) $(p)f \leq (|b|)f = (|b|_p)f$;
- (3) $p \geq \max(\{0, \dots, r\}a)$;
- (4) $p \geq \max(\{0, \dots, r\}b^{-1})$.

Lemma 3.5 and conditions (1) and (2) imply that the sets $W_{f,a,p}$ and $W_{f,b,p}$ are open neighborhoods of a and b , respectively. Let us show that $W_{f,a,p} \cdot W_{f,b,p} \subseteq W_{f,c,r}$. Fix any elements $d \in W_{f,a,p}$ and $e \in W_{f,b,p}$. Since $a|_p = d|_p$ and $b|_p = e|_p$, condition (3) yields that $(de)|_r = c|_r$. Let

$$A = \text{im}(b) \cap \text{im}(de) \cap (r \setminus \text{im}(c)) \quad \text{and} \quad B = \text{im}(e) \cap (r \setminus \text{im}(b)).$$

It is easy to check that, $\text{im}(de) \cap (r \setminus \text{im}(c)) \subseteq A \cup B$. In order to show that $de \in W_{f,c,r}$ it suffices to check that $|A| + |B| \leq (|c|)f$.

Consider any $y \in A$. If $(y)b^{-1} \in \text{im}(a)$, then $y \in \text{im}(c)$, which contradicts the choice of y . By condition (4), $(y)b^{-1} \leq p$. Hence $(y)b^{-1} \in \text{dom}(b) \cap (p \setminus \text{im}(a))$, witnessing that $(A)b^{-1} \subseteq \text{dom}(b) \cap (p \setminus \text{im}(a))$. Since b is a partial bijection, $|A| = |(A)b^{-1}| \leq |\text{dom}(b) \cap (p \setminus \text{im}(a))|$. For the sake of brevity we put $t = |\text{dom}(b) \cap (p \setminus \text{im}(a))|$. Since $(y)b^{-1} \leq p$ and $e|_p = b|_p$ we get that $((y)b^{-1}, y) \in e$. Taking into account that $y \in \text{im}(de)$, we obtain $(y)b^{-1} \in \text{im}(d) \cap (p \setminus \text{im}(a))$. It

follows that $(A)b^{-1} \subseteq \text{im}(d) \cap (p \setminus \text{im}(a))$. Since $d \in W_{f,a,p}$ and b is a partial bijection, we get that $|A| = |(A)b^{-1}| \leq (|a|)f$. Hence

$$|A| \leq \min\{t, (|a|)f\}.$$

Since $e \in W_{f,b,p}$, $|B| \leq (|b|)f$.

Finally, let us check that $\min\{t, (|a|)f\} + (|b|)f \leq (|c|)f$. Since for every $x \in \text{dom}(b) \cap (p \setminus \text{im}(a))$, $(x)b \in \text{im}(b) \setminus \text{im}(c)$ we get that $|c| \leq |b| - t$. If $(|b|)f > 0$, then $(|b|)f + t \leq (|c|)f$ as f is a waning function. If $(|b|)f = 0$, then $\min\{t, (|a|)f\} + (|b|)f \leq (|a|)f \leq (|c|)f$, as $|a| \geq |c|$ and f is a waning function. Thus, $de \in W_{f,c,r}$ and, consequently, $W_{f,a,p} \cdot W_{f,b,p} \subseteq W_{f,c,r}$. Hence $(I_{\mathbb{N}}, \mathcal{T}_f)$ is a Polish topological semigroup. \square

Finally, in terms of the promises made at the beginning of this section, we now show how the topologies \mathcal{T}_f are ordered with respect to inclusion.

Lemma 3.11. *If f and g are waning functions, then the following are equivalent:*

- (i) $\mathcal{T}_f \subseteq \mathcal{T}_g$.
- (ii) for all $n \in \omega$, we have $(n)g \leq (n)f$.

Proof. (i) \Rightarrow (ii): Seeking a contradiction, we suppose that $\mathcal{T}_f \subseteq \mathcal{T}_g$ but that there exists $n \in \mathbb{N}$ such that $(n)f < (n)g$.

Let id_n be the identity function on $\{0, 1, \dots, n-1\}$, let $b \in \mathbb{N}$ be such that $(n)f < b - n \leq (n)g$. By Lemma 3.4, the set $W_{f,\text{id}_n,b}$ is open in $(I_{\mathbb{N}}, \mathcal{T}_f)$. Since $W_{f,\text{id}_n,b} \in \mathcal{T}_g$, by Lemma 3.5 there is some $r \in \mathbb{N}$ such that $W_{g,\text{id}_n,r} \subseteq W_{f,\text{id}_n,b}$. Since $W_{g,\text{id}_n,t+1} \subseteq W_{g,\text{id}_n,t}$ for any large enough t , we may choose $r > b$. By assumption $b - n > (n)f \geq 0$, $b > n$ and so we may define:

$$h = \text{id}_n \cup \{(r+i, n+i) \mid i \in \{0, 1, \dots, b-n-1\}\} \in I_{\mathbb{N}}.$$

Then $h \upharpoonright_r = \text{id}_n \upharpoonright_r$ and $|\text{im}(h) \cap (b \setminus \text{im}(\text{id}_n))| = |\text{im}(h) \cap (r \setminus \text{im}(\text{id}_n))| = b - n$. In particular,

$$(n)f < |\text{im}(h) \cap (b \setminus \text{im}(\text{id}_n))| = |\text{im}(h) \cap (r \setminus \text{im}(\text{id}_n))| \leq (n)g = (|\text{id}_n|)g$$

and so $h \in W_{g,\text{id}_n,r}$ but $h \notin W_{f,\text{id}_n,b}$, which is a contradiction.

(ii) \Rightarrow (i): Let $U \in \mathcal{T}_f$. We show that U is an open neighbourhood of all its elements with respect to \mathcal{T}_g . Suppose that $h \in U$. By Lemma 3.5, there is $r \in \mathbb{N}$ such that $h \in W_{f,h,r} \subseteq U$. Enlarging r if necessary we can assume that $h \in W_{g,h,r} \in \mathcal{T}_g$. Since $(|h|)g \leq (|h|)f$,

$$W_{g,h,r} \subseteq \{l \in I_{\mathbb{N}} : l \upharpoonright_r = h \upharpoonright_r \text{ and } |\text{im}(l) \cap (r \setminus \text{im}(h))| \leq (|h|)f\} = W_{f,h,r}.$$

Hence U is an open neighbourhood of h with respect to \mathcal{T}_g . \square

Lemma 3.11 implies the following.

Corollary 3.12. *If f and g are waning functions, then $\mathcal{T}_f = \mathcal{T}_g$ if and only if $f = g$.*

The following corollary will be used in Section 4.

Corollary 3.13. *Let $f, g : \omega + 1 \rightarrow \omega + 1$ be waning functions, and let \mathcal{F} and \mathcal{G} be the sets of neighbourhoods of \emptyset in \mathcal{T}_f and \mathcal{T}_g , respectively. Then $\mathcal{F} \subseteq \mathcal{G}$ if and only if $(0)f \geq (0)g$.*

Proof. By Lemma 3.5, the family $B_f(\emptyset) = \{W_{f,\emptyset,r} : (r)f \leq (0)f\}$ is a filter basis for \mathcal{F} . Since f is a waning function, $(r)f \leq (0)f$ holds for all $r \in \omega + 1$, and so $B_f(\emptyset) = \{W_{f,\emptyset,r} : r \in \omega\}$.

If $(0)f \geq (0)g$, then $W_{f,\emptyset,r} \supseteq W_{g,\emptyset,r}$ for all r by Eq. (2), and so $B_f(\emptyset) = \{W_{f,\emptyset,r} : r \in \omega\}$ is coarser than $\{W_{g,\emptyset,r} : r \in \omega\} = B_g(\emptyset)$. Hence $\mathcal{F} \subseteq \mathcal{G}$.

Conversely, suppose that $\mathcal{F} \subseteq \mathcal{G}$. If we define waning functions f' and g' such that $(0)f' = (0)f$ and $(0)g' = (0)g$ and $(x)f' = (x)g' = 0$ if $x \neq 0$, then the set of neighbourhoods of \emptyset in $\mathcal{T}_{f'}$ is \mathcal{F} . Similarly for $\mathcal{T}_{g'}$ and \mathcal{G} . A subset N of $I_{\mathbb{N}}$ is a neighbourhood in $\mathcal{T}_{f'}$ of $p \in I_{\mathbb{N}} \setminus \{\emptyset\}$ if and only if N is a neighbourhood of p in \mathcal{I}_4 . Similarly, for $\mathcal{T}_{g'}$. Hence every neighbourhood in $\mathcal{T}_{f'}$ of every $p \in I_{\mathbb{N}}$ is also a neighbourhood of p in $\mathcal{T}_{g'}$. So $\mathcal{T}_{f'} \subseteq \mathcal{T}_{g'}$ and so, by Lemma 3.11, $(0)f = (0)f' \geq (0)g' = (0)g$. \square

By Corollary 3.9, every function $f : \omega + 1 \rightarrow \omega + 1$ (not just waning ones) corresponds to a Polish topology \mathcal{T}_f on $I_{\mathbb{N}}$. It is natural (but not necessary) to ask how these compare to the Polish semigroup topologies \mathcal{T}_f arising from waning functions f . We will show in Theorem 3.15 that \mathcal{T}_f is a Polish semigroup topology for every function $f : \omega + 1 \rightarrow \omega + 1$, but that there always exists a waning function f' such that $\mathcal{T}_f = \mathcal{T}_{f'}$ and so non-waning functions do not give rise to any additional topologies.

Lemma 3.14. *Let f be a waning function with $(\omega)f = 0$, $n \in \mathbb{N}$ and $X \subseteq \mathbb{N}$ be finite. If $g \in I_{\mathbb{N}}$ is arbitrary with $|\text{im}(g) \setminus X| \geq n$ and $|X \cap \text{im}(g)| \geq (n)f$, then $(|g|)f = 0$.*

Proof. If g is infinite, then we are done by the assumption that $(\omega)f = 0$. We have

$$(|g|)f \leq \max(0, (|g| - 1)f - 1) \leq \dots \leq \max(0, (|g| - (|g| - n))f - (|g| - n)) = \max(0, (n)f - (|g| - n)).$$

As $|\text{im}(g) \setminus X| \geq n$, it follows that $|g| - n \geq |\text{im}(g) \cap X| \geq (n)f$. So $(n)f - (|g| - n) \leq 0$ and hence $\max(0, (n)f - (|g| - n)) = 0$ as required. \square

Theorem 3.15. *If $f \in (\omega + 1)^{(\omega + 1)}$, then $\mathcal{T}_f = \mathcal{T}_{f'}$.*

Proof. Note that if $f \upharpoonright_{\omega}$ is constant with value ω , then $f \upharpoonright_{\omega} = f' \upharpoonright_{\omega}$ and we are done. So assume that this is not the case which, in particular, implies that $(\omega)f' = 0$.

By Corollary 3.6, we have $\mathcal{T}_f \supseteq \mathcal{T}_{f'}$ and so it only remains to show that $\mathcal{T}_{f'} \supseteq \mathcal{T}_f$. Let $n \in \mathbb{N}$, $X \subseteq \mathbb{N}$ be finite, and

$$U_{f,n,X} = \{g \in I_{\mathbb{N}} \mid |\text{im}(g) \setminus X| \geq n \text{ and } |X \cap \text{im}(g)| \leq (n)f\}.$$

In order to show that $U_{f,n,X} \in \mathcal{T}_{f'}$, let $g \in U_{f,n,X}$ be arbitrary. We show that $U_{f,n,X}$ is a neighbourhood of g with respect to $\mathcal{T}_{f'}$.

First assume that $(|g|)f' > 0$. Since $|\text{im}(g) \setminus X| \geq n$ and f' is a waning function with $(\omega)f' = 0$, it follows from the contra-positive of Lemma 3.14, that $|X \cap \text{im}(g)| < (n)f'$ and so $g \in U_{f',n,X}$. Note that $U_{f',n,X} \subseteq U_{f,n,X}$ so we have that $U_{f,n,X}$ is a neighbourhood of g with respect to $\mathcal{T}_{f'}$.

So now assume that $(|g|)f' = 0$. Let r be larger than all elements of $(X)g^{-1}$ as well as the first n elements of $(\mathbb{N} \setminus X)g^{-1}$, and large enough that $W_{f',g,r} \in \mathcal{T}_{f'}$. It follows that

$$g \in W_{f',g,r} \subseteq \{l \in I_{\mathbb{N}} \mid g \upharpoonright_r = l \upharpoonright_r \text{ and } \text{im}(l) \cap X = \text{im}(g) \cap X\} \subseteq U_{f,n,X}$$

so $U_{f,n,X}$ is a neighbourhood of g with respect to $\mathcal{T}_{f'}$ as required. \square

4. FROM SEMIGROUP TOPOLOGIES TO WANING FUNCTIONS

We have shown in Section 3 that every waning function gives rise to a Polish semigroup topology on $I_{\mathbb{N}}$. In this section we will prove the converse, namely that every Polish (in fact every T_1 and second-countable) semigroup topology on $I_{\mathbb{N}}$ corresponds to a waning function. In other words, the aim of this section is to finish the proof of Theorem 2.3.

Recall that there are two minimal Polish semigroup topologies on $I_{\mathbb{N}}$, namely \mathcal{I}_2 and $\mathcal{I}_3 = \mathcal{I}_2^{-1}$. Reflecting this fact, the correspondence between waning functions and Polish topologies on $I_{\mathbb{N}}$ is “one-to-two” in the following sense. Every waning function f gives rise to a unique Polish topology \mathcal{T}_f on $I_{\mathbb{N}}$ which contains \mathcal{I}_2 and a unique topology \mathcal{T}_f^{-1} containing \mathcal{I}_3 . Due to this duality we will mostly consider an arbitrary T_1 and second-countable topology \mathcal{T} containing \mathcal{I}_2 .

The aim of the next definitions and results (up to and including Lemma 4.8) is to construct a sequence on non-decreasing filters for such a \mathcal{T} which we will call good filters. The good filters will later be used to define the waning function corresponding to \mathcal{T} .

Definition 4.1. If $X \subseteq \mathbb{N}$ is finite, then we define $i_X : \mathbb{N} \rightarrow \mathbb{N} \setminus X$ to be the unique order isomorphism. For $g \in I_{\mathbb{N}}$ let $\uparrow g = \{h \in I_{\mathbb{N}} \mid g \subseteq h\}$. If $g \in I_{\mathbb{N}}$ has finite domain and image, then we define a bijection

$$d_g : \uparrow g \rightarrow I_{\mathbb{N}} \quad \text{by} \quad (h)d_g = i_{\text{dom}(g)} \circ h \circ i_{\text{im}(g)}^{-1}.$$

Note that the function d_g is continuous with respect to any shift continuous topology on $I_{\mathbb{N}}$ and has a clopen domain with respect to \mathcal{I}_1 .

Lemma 4.2. *If g is a finite idempotent of $I_{\mathbb{N}}$, then d_g is an isomorphism.*

Proof. Since g is an idempotent, then $\uparrow g$ is an inverse subsemigroup of $I_{\mathbb{N}}$. As d_g is a bijection it suffices to show that d_g is a homomorphism. Let $f, h \in \uparrow g$. Note that $\text{im}(i_{\text{dom}(g)}f) \cap \text{im}(g) = \emptyset$, as $g \subseteq f$ and f is injective. It follows that $i_{\text{dom}(g)}f i_{\text{im}(g)}^{-1} i_{\text{im}(g)} = i_{\text{dom}(g)}f$. Then

$$(f)d_g(h)d_g = i_{\text{dom}(g)}f i_{\text{im}(g)}^{-1} i_{\text{im}(g)} h i_{\text{im}(g)}^{-1} = i_{\text{dom}(g)}f h i_{\text{im}(g)}^{-1} = (fh)d_g.$$

□

Definition 4.3. Let \mathcal{T} be a second countable semigroup topology on $I_{\mathbb{N}}$ containing \mathcal{I}_2 and $g \in I_{\mathbb{N}}$ be finite. Then let

$$\mathcal{F}_{\mathcal{T},g} := \{(N \cap \uparrow g)d_g \mid N \text{ is a neighbourhood of } g \text{ with respect to } \mathcal{T}\}.$$

Lemma 4.4. *If \mathcal{T} is a second countable semigroup topology on $I_{\mathbb{N}}$ containing \mathcal{I}_2 , and $g, h \in I_{\mathbb{N}}$ are such that $|g| = |h| < \omega$, then $\mathcal{F}_{\mathcal{T},g} = \mathcal{F}_{\mathcal{T},h}$.*

Proof. Let θ_1 and θ_2 be the unique topologies on $I_{\mathbb{N}}$ such that $d_g : (\uparrow g, \mathcal{T}) \rightarrow (I_{\mathbb{N}}, \theta_1)$ and $d_h : (\uparrow h, \mathcal{T}) \rightarrow (I_{\mathbb{N}}, \theta_2)$ are homeomorphisms. Note that $\mathcal{F}_{\mathcal{T},g}$ is the set of neighbourhoods of \emptyset with respect to θ_1 , and similarly $\mathcal{F}_{\mathcal{T},h}$ is the set of neighbourhoods of \emptyset with respect to θ_2 .

It suffices to show that $\theta_1 = \theta_2$. Let $k : \text{dom}(g) \rightarrow \text{dom}(h)$ be a bijection. Let

$$k_d = k \cup (i_{\text{dom}(g)}^{-1} \circ i_{\text{dom}(h)}) \in \text{Sym}(\mathbb{N}).$$

Set

$$k_i = g^{-1}k_d h \cup (i_{\text{im}(g)}^{-1} \circ i_{\text{im}(h)}) \in \text{Sym}(\mathbb{N}).$$

Then $\phi : I_{\mathbb{N}} \rightarrow I_{\mathbb{N}}$ defined by $(x)\phi = k_d^{-1}xk_i$ is a homeomorphism from $(I_{\mathbb{N}}, \mathcal{T})$ to $(I_{\mathbb{N}}, \mathcal{T})$. Observe that

$$(g)\phi = k_d^{-1}gk_i = k^{-1}gg^{-1}k_d h = k^{-1}k_d h = k^{-1}k h = h.$$

Also, for all $f \in I_{\mathbb{N}}$ we have

$$(f)d_g^{-1} = i_{\text{dom}(g)}^{-1} f i_{\text{im}(g)} \cup g.$$

Hence

$$\begin{aligned} (f)d_g^{-1}\phi &= \left((i_{\text{dom}(g)}^{-1} \circ i_{\text{dom}(h)})^{-1} i_{\text{dom}(g)}^{-1} f i_{\text{im}(g)} (i_{\text{im}(g)}^{-1} \circ i_{\text{im}(h)}) \right) \cup (g)\phi \\ &= \left(i_{\text{dom}(h)}^{-1} f i_{\text{im}(h)} \right) \cup h. \end{aligned}$$

Thus

$$(f)d_g^{-1}\phi d_h = i_{\text{dom}(h)} \left(\left(i_{\text{dom}(h)}^{-1} f i_{\text{im}(h)} \right) \cup h \right) i_{\text{im}(h)}^{-1} = f.$$

So $d_g^{-1}\phi d_h : (I_{\mathbb{N}}, \theta_1) \rightarrow (I_{\mathbb{N}}, \theta_2)$ is the identity function. Being a composition of homeomorphisms it is also a homeomorphism and so $\theta_1 = \theta_2$. □

Definition 4.5. From now on we denote the filter $\mathcal{F}_{\mathcal{T},g}$ by $\mathcal{F}_{\mathcal{T},|g|}$ (this is well defined by Lemma 4.4). We say that a filter \mathcal{F} is *good* if $\mathcal{F} = \mathcal{F}_{\mathcal{T},n}$ for some $n \in \mathbb{N}$ and some second countable semigroup topology on $I_{\mathbb{N}}$ containing \mathcal{I}_2 .

Lemma 4.6 (Good filters are zero neighbourhoods). *If \mathcal{F} is a filter, then \mathcal{F} is good if and only if there is a second countable semigroup topology \mathcal{T} on $I_{\mathbb{N}}$ containing \mathcal{I}_2 such that \mathcal{F} is the set of neighbourhoods of \emptyset .*

Proof. (\Leftarrow): In this case $\mathcal{F} = \mathcal{F}_{\mathcal{T}, \emptyset}$.

(\Rightarrow): Suppose that $\mathcal{F} = \mathcal{F}_{\mathcal{T}, n}$ for some $n \in \mathbb{N}$.

Let id_n be the identity function on n . By Lemma 4.2, d_{id_n} is an isomorphism from $\uparrow \text{id}_n$ to $I_{\mathbb{N}}$ which maps id_n to \emptyset . Let the semigroup $\uparrow \text{id}_n$ have the subspace topology inherited from $(I_{\mathbb{N}}, \mathcal{T})$, and θ be the unique topology on $I_{\mathbb{N}}$ such that $d_{\text{id}_n} : \uparrow \text{id}_n \rightarrow (I_{\mathbb{N}}, \theta)$ is a topological isomorphism. Clearly, θ is a second countable semigroup topology. Since $\uparrow \text{id}_n$ is open in $(I_{\mathbb{N}}, \mathcal{T})$ we get that $(\mathcal{F}_{\mathcal{T}, n})d_{\text{id}_n}^{-1}$ is the set of all neighbourhoods of id_n in $\uparrow \text{id}_n$. Thus $\mathcal{F}_{\mathcal{T}, n} = ((\mathcal{F}_{\mathcal{T}, n})d_{\text{id}_n}^{-1})d_{\text{id}_n}$ is the set of all neighbourhoods of 0 in $(I_{\mathbb{N}}, \theta)$. Hence $\mathcal{F}_{\theta, 0} = \mathcal{F}_{\mathcal{T}, n}$. \square

Definition 4.7. If \mathcal{T} is a second countable semigroup topology on $I_{\mathbb{N}}$ containing \mathcal{I}_2 , then we define the filter data of \mathcal{T} to be the sequence

$$\mathfrak{F}_{\mathcal{T}} := (\mathcal{F}_{\mathcal{T}, n})_{n \in \omega}.$$

Lemma 4.8. *If $\mathfrak{F}_{\mathcal{T}} = (\mathcal{F}_{\mathcal{T}, n})_{n \in \omega}$ is the filter data of a second countable semigroup topology $\mathcal{T} \supseteq \mathcal{I}_2$ on $I_{\mathbb{N}}$, then $\mathcal{F}_{\mathcal{T}, n} \subseteq \mathcal{F}_{\mathcal{T}, m}$ for all $n \leq m$.*

Proof. Assume without loss of generality that $n < m$.

Let $k \in I_{\mathbb{N}}$ be such that $\text{id}_n \subseteq k$ and k bijectively maps $\mathbb{N} \setminus (m \setminus n)$ to \mathbb{N} in an order preserving fashion. It follows that the function $\phi_k : \uparrow \text{id}_m \rightarrow \uparrow \text{id}_n$ defined by

$$(x)\phi_k = k^{-1}xk$$

is a continuous bijection with inverse $x \mapsto (kxk^{-1}) \cup \text{id}_{m \setminus n}$. Note that $\text{id}_{\mathbb{N} \setminus (m \setminus n)} \text{id}_m = \text{id}_n$ and $k|_n = \text{id}_n$. Then

$$(\text{id}_m)\phi_k = k^{-1} \text{id}_m k = \text{id}_n \text{id}_m \text{id}_n = \text{id}_n.$$

By θ we denote the subspace topology on $\uparrow \text{id}_n$ inherited from $(I_{\mathbb{N}}, \mathcal{T})$. Hence if $N \in \mathcal{F}_{\mathcal{T}, n}$, then $(N)d_{\text{id}_n}^{-1}$ is a neighbourhood of id_n with respect to θ and hence $(N)d_{\text{id}_n}^{-1}\phi_k^{-1}$ is a neighbourhood of id_m with respect to θ . To show that $N \in \mathcal{F}_{\mathcal{T}, m}$, it therefore suffices to show that $(N)d_{\text{id}_n}^{-1}\phi_k^{-1}d_{\text{id}_m} = N$.

We show in fact that $\phi_k^{-1}d_{\text{id}_m} = d_{\text{id}_n}$. Note that $\phi_k, d_{\text{id}_m}, d_{\text{id}_n}$ are all injections and we have by definition that

$$\text{dom}(\phi_k) = \uparrow \text{id}_m = \{g \in I_{\mathbb{N}} \mid \text{id}_m \subseteq g\}, \quad \text{im}(\phi_k) = \uparrow \text{id}_n = \{g \in I_{\mathbb{N}} \mid \text{id}_n \subseteq g\},$$

$$\text{im}(d_{\text{id}_m}) = \text{im}(d_{\text{id}_n}) = I_{\mathbb{N}}.$$

Thus ϕ_k^{-1} is a bijection from $\uparrow \text{id}_n$ to $\uparrow \text{id}_m$. For all $x \in \uparrow \text{id}_n$ we have

$$(x)d_{\text{id}_n} = i_n x i_n^{-1} \text{ and } (x)\phi_k^{-1}d_{\text{id}_m} = i_m(kxk^{-1} \cup \text{id}_{m \setminus n})i_m^{-1} = (i_m k)x(i_m k)^{-1}.$$

Thus we need only show that $i_n = i_m k$. This is immediate as $i_n : \mathbb{N} \rightarrow \mathbb{N} \setminus n$, $i_m : \mathbb{N} \rightarrow \mathbb{N} \setminus m$, and $k|_{\mathbb{N} \setminus m} : \mathbb{N} \setminus m \rightarrow \mathbb{N} \setminus n$ are all order isomorphisms. \square

The next major step is to show in Theorem 4.13 that every good filter has a convenient basis (we will call it a basis consisting of *wany* sets). Furthermore, we will show that if this basis is particularly nice (consisting of *finitely wany* sets), then the good filter arises from a waning function.

Recall that by $[X]^{<\infty}$ we denote the set of all finite subsets of a set X .

Definition 4.9. If $\mathfrak{W} \subseteq [\mathbb{N}]^{<\infty}$ is non-empty, then we say that a set $S \subseteq I_{\mathbb{N}}$ is \mathfrak{W} -wany if there is $n \in \mathbb{N}$ such that

$$S = N_{n, \mathfrak{W}} := \{f \in I_{\mathbb{N}} \mid \text{dom}(f) \cap n = \emptyset \text{ and there is } Y \in \mathfrak{W} \text{ with } \text{im}(f) \cap Y = \emptyset\}.$$

We say a set is wany if it is \mathfrak{W} -wany for some \mathfrak{W} . We say that a set is finitely wany if it is \mathfrak{W} -wany for some finite \mathfrak{W} .

Remark 4.10. Note that for example if $\mathfrak{W} = \{\emptyset\}$, then

$$N_{n,\mathfrak{W}} := \{f \in I_{\mathbb{N}} \mid \text{dom}(f) \cap n = \emptyset\},$$

if $\mathfrak{W} = \{\{0\}\}$, then

$$N_{n,\mathfrak{W}} := \{f \in I_{\mathbb{N}} \mid \text{dom}(f) \cap n = \emptyset \text{ and } 0 \notin \text{im}(f)\},$$

if $\mathfrak{W} = \{\{0, 1, \dots, k\}\}$, then

$$N_{n,\mathfrak{W}} := \{f \in I_{\mathbb{N}} \mid \text{dom}(f) \cap n = \emptyset \text{ and } \text{im}(f) \cap \{0, 1, \dots, k\} = \emptyset\},$$

if $\mathfrak{W} = \{Y \subseteq r \mid |Y| = r - 3\}$, then

$$N_{n,\mathfrak{W}} := \{f \in I_{\mathbb{N}} \mid \text{dom}(f) \cap n = \emptyset \text{ and } |\text{im}(f) \cap r| \leq 3\},$$

etc. So all the usual sets we've used to define waning function topologies are of this form.

From here onward we denote the set of finite elements of $I_{\mathbb{N}}$ by $I_{\mathbb{N}}^{<\infty}$ (not to be confused with $[I_{\mathbb{N}}]^{<\infty}$ which is the set of finite subsets of $I_{\mathbb{N}}$).

Definition 4.11. We say that a filter \mathcal{F} on $I_{\mathbb{N}}$ is *defined by* a waning function f , if \mathcal{F} is the filter of all neighbourhoods of \emptyset with respect to the topology \mathcal{T}_f (as defined in Definition 2.1).

Definition 4.12. If $X \subseteq Y$ are sets and \mathcal{F} is a filter of subsets of Y , then we define the trace of this filter on X by

$$\mathcal{F}|_X := \{F \cap X \mid F \in \mathcal{F}\}.$$

Theorem 4.13. *If \mathcal{F} is a good filter on $I_{\mathbb{N}}$, then the following hold:*

- (i) *The filter \mathcal{F} has a filter base consisting of wany sets.*
- (ii) *If \mathcal{F} has a filter base consisting of finitely wany sets, then \mathcal{F} can be defined by a waning function.*
- (iii) *If the filter $\mathcal{F}|_{I_{\mathbb{N}}^{<\infty}}$ on $I_{\mathbb{N}}^{<\infty}$ has a base of the form $(S_i \cap I_{\mathbb{N}}^{<\infty})_{i \in \mathbb{N}}$ where each set S_i is finitely wany, then there is a good filter \mathcal{F}' defined by a waning function such that $\mathcal{F}|_{I_{\mathbb{N}}^{<\infty}} = \mathcal{F}'|_{I_{\mathbb{N}}^{<\infty}}$.*

Proof. We start with point (1). Let \mathcal{F} be a good filter and let $N \in \mathcal{F}$ be arbitrary. We will construct a wany set $N'''' \in \mathcal{F}$ with $N'''' \subseteq N$. By Lemma 4.6, \mathcal{F} is the set of neighbourhoods of an idempotent in a semigroup topology \mathcal{T} . Theorem 5.1.5(vii) from [7] implies that $\mathcal{T} \subseteq \mathcal{I}_4$. Then there exists an $N' \in \mathcal{F} \cap \mathcal{I}_4$ such that $N'N' \subseteq N$.

For all $(F, X, Y) \in I_{\mathbb{N}}^{<\infty} \times [\mathbb{N}]^{<\infty} \times [\mathbb{N}]^{<\infty}$, define

$$U_{F,X,Y} := \{f \in I_{\mathbb{N}} \mid F \subseteq f, \text{dom}(f) \cap X = \text{im}(f) \cap Y = \emptyset\}.$$

As $N' \in \mathcal{I}_4$, there exists $D \subseteq I_{\mathbb{N}}^{<\infty} \times [\mathbb{N}]^{<\infty} \times [\mathbb{N}]^{<\infty}$ such that

$$N' = \bigcup_{(F,X,Y) \in D} U_{F,X,Y}.$$

Moreover, there must be some triple $(\emptyset, A, B) \in D$ as $\emptyset \in N'$. Let $n \in \mathbb{N}$ be such that $A \cup B \subseteq n$ and define

$$D' = \{(F, X \cup n, Y) \mid (F, X, Y) \in D \text{ and } U_{F,X \cup n, Y} \neq \emptyset\}.$$

Let

$$N'' := \bigcup_{(F,X,Y) \in D'} U_{F,X,Y} = \bigcup_{(F,X,Y) \in D} U_{\emptyset, n, \emptyset} \cap U_{F,X,Y} = N' \cap U_{\emptyset, n, \emptyset}.$$

Since $U_{\emptyset, n, \emptyset} \in \mathcal{I}_2 \subseteq \mathcal{T}$ we get that $N'' \in \mathcal{F}$. Let $N'''' := N''N'' \subseteq N'N' \subseteq N$.

Claim 4.14. *If $(F, X, Y) \in D'$, then*

$$U_{\emptyset, n, B} U_{F, X, Y} = U_{\emptyset, n, Y}.$$

Proof. The containment \subseteq is immediate. Let $h \in U_{\emptyset, n, Y}$ be arbitrary. Note that $U_{F, X, Y}$ is non-empty so $\text{dom}(F) \cap X = \text{im}(F) \cap Y = \emptyset$.

Let $g_1 = F$ and let g_2 be any partial bijection with image equal to $\text{im}(h) \setminus \text{im}(g_1)$ and domain disjoint from $n \cup X \cup Y \cup \text{dom}(F) \cup \text{im}(F)$. Let $g = g_1 \cup g_2 \in I_{\mathbb{N}}$. Note that $g \in U_{F, X, Y}$ as $\text{dom}(F) \cap X = \text{im}(F) \cap Y = \emptyset$ and $\text{im}(h) \cap Y = \emptyset$.

Let $f = hg^{-1} = h(g_1 \cup g_2)^{-1}$. Note that $\text{dom}(f) \subseteq \text{dom}(h)$, so

$$\text{dom}(f) \cap n = \text{dom}(h) \cap n = \emptyset \quad \text{and} \quad \text{im}(f) \subseteq \text{dom}(F) \cup \text{dom}(g_2).$$

By the definition of D' , $X \supseteq n \supseteq B$, and so

$$\text{im}(f) \cap B \subseteq (\text{dom}(F) \cup \text{dom}(g_2)) \cap B = (\text{dom}(F) \cap B) \cup (\text{dom}(g_2) \cap B) = \text{dom}(F) \cap B \subseteq \text{dom}(F) \cap X = \emptyset.$$

Hence $f \in U_{\emptyset, n, B}$ and $h = fg \in U_{\emptyset, n, B} U_{F, X, Y}$ as required. \square

Let $\mathfrak{Y} := \{Y \subseteq \mathbb{N} \mid Y \text{ appears as the third entry of an element of } D'\}$. Since $(\emptyset, A, B) \in D$, we have $(\emptyset, n, B) \in D'$. Thus by Claim 4.14, we have

$$\bigcup_{Y \in \mathfrak{Y}} U_{\emptyset, n, Y} = \bigcup_{(F, X, Y) \in D'} U_{\emptyset, n, B} U_{F, X, Y} \subseteq \left(\bigcup_{(F, X, Y) \in D'} U_{F, X, Y} \right) \left(\bigcup_{(F, X, Y) \in D'} U_{F, X, Y} \right) = N'' N'' = N'''$$

For convenience denote $N'''' := \bigcup_{Y \in \mathfrak{Y}} U_{\emptyset, n, Y}$. Note that N'''' is \mathfrak{Y} -wamy. As $n \subseteq X$ for all $(F, X, Y) \in D'$ we have $U_{F, X, Y} \subseteq U_{F, n, Y} \subseteq U_{\emptyset, n, Y}$. It follows that $N'' \subseteq N''''$ and subsequently $N'''' \in \mathcal{F}$. Since N was chosen arbitrarily, $N'''' \subseteq N''' \subseteq N$, and N'''' is a wamy set, the result follows.

We now prove the parts (2) and (3) of the theorem together by proving the following statement.

(2)&(3) If $I \in \{I_{\mathbb{N}}, I_{\mathbb{N}}^{<\infty}\}$ and $\mathcal{F}|_I$ has a filter base of the form $(S_i \cap I)_{i \in \mathbb{N}}$ where each set S_i is finitely wamy, then there is a good filter \mathcal{F}' defined by a waning function such that $\mathcal{F}|_I = \mathcal{F}'|_I$.

Suppose that \mathcal{F} and I are as hypothesised. We define

$$K = \{l \in \mathbb{N} \mid \{f \in I \mid \text{im}(f) \not\supseteq (l+1)\} \in \mathcal{F}|_I\}.$$

Claim 4.15. *Let $l, n \in \mathbb{N}$ and $X \in [\mathbb{N}]^{<\infty}$ be arbitrary such that $l < |X|$. Then*

$$l \in K \iff \{f \in I \mid \text{dom}(f) \cap n = \emptyset \text{ and } |\text{im}(f) \cap X| \leq l\} \in \mathcal{F}|_I.$$

Proof. We fix $l, n \in \mathbb{N}$ and $X \in [\mathbb{N}]^{<\infty}$ such that $l < |X|$.

(\Leftarrow): Let $\sigma \in \text{Sym}(\mathbb{N})$ be such that $l+1 \subseteq (X)\sigma$. Then

$$\begin{aligned} (\{f \in I \mid \text{dom}(f) \cap n = \emptyset \text{ and } |\text{im}(f) \cap X| \leq l\})\sigma &\subseteq \{f \in I \mid |\text{im}(f) \cap (X)\sigma| \leq l\} \\ &\subseteq \{f \in I \mid \text{im}(f) \not\supseteq (l+1)\}. \end{aligned}$$

Since \mathcal{F} is a good filter and elements of $\text{Sym}(\mathbb{N})$ act bijectively on $I_{\mathbb{N}}^{<\infty}$ and they are homeomorphisms of $I_{\mathbb{N}}$ endowed with any shift-continuous topology, we get that the filter $\mathcal{F}|_I$ is closed under right multiplication by elements of $\text{Sym}(\mathbb{N})$. It follows that

$$\{f \in I \mid \text{im}(f) \not\supseteq (l+1)\} \in \mathcal{F}|_I,$$

so $l \in K$.

(\Rightarrow) : We have that $\{f \in I \mid |\text{im}(f) \cap (l+1)| \leq l\} \in \mathcal{F}|_I$. Let $r = |X|$. Let G_r be the finite subgroup of $\text{Sym}(\mathbb{N})$ consisting of all permutations with support contained in r . Note that

$$\begin{aligned}
\{f \in I \mid |\text{im}(f) \cap r| \leq l\} &= \{f \in I \mid \text{im}(f) \cap r \text{ has no subset of size } l+1\} \\
&= \{f \in I \mid \text{there is no } \sigma \in G_r \text{ such that } (\text{im}(f))\sigma \supseteq l+1\} \\
&= \{f \in I \mid \text{there is no } \sigma \in G_r \text{ such that } \text{im}(f\sigma) \supseteq l+1\} \\
&= \bigcap_{\sigma \in G_r} \{f \in I \mid \text{im}(f\sigma) \not\supseteq l+1\} \\
&= \bigcap_{\sigma \in G_r} \{f \in I\sigma^{-1} \mid \text{im}(f\sigma) \not\supseteq l+1\} \quad (\text{as } I = I\sigma^{-1}, \text{ because } \sigma \text{ is a bijection}) \\
&= \bigcap_{\sigma \in G_r} \{g\sigma^{-1} \in I\sigma^{-1} \mid \text{im}(g\sigma^{-1}\sigma) \not\supseteq l+1\} \\
&= \bigcap_{\sigma \in G_r} \{g\sigma^{-1} \in I\sigma^{-1} \mid \text{im}(g) \not\supseteq l+1\} \\
&= \bigcap_{\sigma \in G_r} \{f\sigma^{-1} \in I \mid \text{im}(f) \not\supseteq l+1\} \\
&= \bigcap_{\sigma \in G_r} \{f \in I \mid \text{im}(f) \not\supseteq l+1\} \sigma^{-1}.
\end{aligned}$$

So, since the filter $\mathcal{F}|_I$ is closed under right multiplication by elements of $\text{Sym}(\mathbb{N})$ we have that $\{f \in I \mid |\text{im}(f) \cap r| \leq l\} \in \mathcal{F}|_I$. As \mathcal{F} contains all the \mathcal{I}_2 neighbourhoods of \emptyset , we obtain that

$$\begin{aligned}
\{f \in I \mid \text{dom}(f) \cap n = \emptyset \text{ and } |\text{im}(f) \cap r| \leq l\} \\
= \{f \in I \mid \text{dom}(f) \cap n = \emptyset\} \cap \{f \in I \mid |\text{im}(f) \cap r| \leq l\} \in \mathcal{F}|_I.
\end{aligned}$$

Let $\rho \in \text{Sym}(\mathbb{N})$ be such that $(r)\rho = X$. It follows that

$$\begin{aligned}
\{f \in I \mid \text{dom}(f) \cap n = \emptyset \text{ and } |\text{im}(f) \cap X| \leq l\} \\
= \{f \in I \mid \text{dom}(f) \cap n = \emptyset \text{ and } |\text{im}(f) \cap r| \leq l\} \rho \in \mathcal{F}|_I
\end{aligned}$$

as required. □

Claim 4.16. *If $F \in \mathcal{F}|_I$, then there exist $l, n \in \mathbb{N}$ and $X \in [\mathbb{N}]^{<\infty}$ such that*

$$U := \{f \in I \mid \text{dom}(f) \cap n = \emptyset \text{ and } |\text{im}(f) \cap X| \leq l\} \in \mathcal{F}|_I$$

and $U \subseteq F$.

Proof. Let $F \in \mathcal{F}$ be arbitrary. By the assumption, $\mathcal{F}|_I$ has a filter base of the form $(S_i \cap I)_{i \in \mathbb{N}}$ where each set S_i is finitely wany. Hence there is a finite non-empty set \mathfrak{Y} and $n \in \mathbb{N}$ such that $N_{n, \mathfrak{Y}} \cap I \in \mathcal{F}|_I$ and

$$N_{n, \mathfrak{Y}} \cap I = \{f \in I \mid \text{dom}(f) \cap n = \emptyset \text{ and there is } Y \in \mathfrak{Y} \text{ with } \text{im}(f) \cap Y = \emptyset\} \subseteq F.$$

Let l be the largest integer such that we both have $l \leq |\bigcup \mathfrak{Y}|$ and that every subset of $\bigcup \mathfrak{Y}$ of size l is disjoint from at least one element of \mathfrak{Y} (note that 0 satisfies this).

Let G be the finite subgroup of $\text{Sym}(\mathbb{N})$ consisting of all permutations supported on $\bigcup \mathfrak{Y}$. Then define

$$M := I \cap \bigcap_{\sigma \in G} N_{n, \mathfrak{Y}}\sigma.$$

Note that $M \in \mathcal{F}|_I$. Note that if for some $f \in I_{\mathbb{N}}$, $|\text{im}(f) \cap \bigcup \mathfrak{Y}| \leq l$, then $\text{im}(f\sigma)$ must be disjoint from an element of \mathfrak{Y} for all $\sigma \in G$, so $f \in M$. Conversely, if $f \in M$, then $|\text{im}(f) \cap \bigcup \mathfrak{Y}| \leq l$ as otherwise there is $\sigma \in G$ such that $\text{im}(f\sigma)$ intersects every element of \mathfrak{Y} and hence $f \notin N_{n,\mathfrak{Y}}\sigma^{-1}$. So

$$M = \{f \in I \mid \text{dom}(f) \cap n = \emptyset \text{ and } |\text{im}(f) \cap \bigcup \mathfrak{Y}| \leq l\} \in \mathcal{F}|_I$$

choosing $X = \bigcup \mathfrak{Y}$ the claim follows. \square

If for all $F \in \mathcal{F}|_I$, there is $l, n \in \mathbb{N}$ and $X \in [\mathbb{N}]^{<\infty}$ with $l = |X|$ such that

$$\{f \in I \mid \text{dom}(f) \cap n = \emptyset \text{ and } |\text{im}(f) \cap X| \leq l\} \in \mathcal{F}|_I$$

and

$$\{f \in I \mid \text{dom}(f) \cap n = \emptyset \text{ and } |\text{im}(f) \cap X| \leq l\} \subseteq F,$$

then we have that $\mathcal{F}|_I$ is the set of all neighbourhoods of \emptyset in the subspace topology on I inherited from $(I_{\mathbb{N}}, \mathcal{I}_2)$, and we are done. Otherwise it follows from Claim 4.16 and Claim 4.15 that $K \neq \emptyset$. Let $k = \min(K)$. It then follows from the same two claims that the sets of the form

$$\{f \in I \mid \text{dom}(f) \cap n = \emptyset \text{ and } |\text{im}(f) \cap X| \leq k\}$$

for $n \in \mathbb{N}$ and $X \in [\mathbb{N}]^{<\infty}$ with $|X| > k$ all belong to $\mathcal{F}|_I$ and are a basis for this filter. Hence if \mathcal{F}' is any filter defined by a waning function ϕ such that $(0)\phi = k$, then $\mathcal{F}|_I = \mathcal{F}'|_I$. \square

We have shown that every good filter has a basis of wany sets and that a basis of finitely wany sets may be used to construct a waning function. To bridge the remaining gap, we show in Corollary 4.24 that a good filter cannot have a basis of wany sets without having a basis of finitely wany sets.

Lemma 4.17. *If $\mathcal{F}_1, \mathcal{F}_2$ are good filters and $\mathcal{F}_1|_{I_{\mathbb{N}}^{<\infty}} = \mathcal{F}_2|_{I_{\mathbb{N}}^{<\infty}}$, then $\mathcal{F}_1 = \mathcal{F}_2$.*

Proof. Let θ_1, θ_2 be second countable semigroup topologies on $I_{\mathbb{N}}$ containing \mathcal{I}_2 such that \mathcal{F}_1 is the set of θ_1 neighbourhoods of \emptyset and \mathcal{F}_2 is the set of θ_2 neighbourhoods of \emptyset .

Suppose for a contradiction that $\mathcal{F}_1 \not\subseteq \mathcal{F}_2$. By Theorem 4.13, we can fix a wany set $N_{m,\mathfrak{Y}} \in \mathcal{F}_1 \setminus \mathcal{F}_2$. Consider the product $\emptyset \text{id}_{\mathbb{N}} = \emptyset$ in the topological semigroup $(I_{\mathbb{N}}, \theta_1)$. Since $(I_{\mathbb{N}}, \theta_1)$ is a topological semigroup, we can find $W_l \in \mathcal{F}_1 \cap \theta_1$ and $W_r \in \theta_1 \subseteq \mathcal{I}_4$ such that $\text{id}_{\mathbb{N}} \in W_r$ and $W_l W_r \subseteq N_{m,\mathfrak{Y}}$. Since open neighborhood bases at $\text{id}_{\mathbb{N}}$ in $(I_{\mathbb{N}}, \mathcal{I}_2)$ and $(I_{\mathbb{N}}, \mathcal{I}_4)$ coincide, we lose no generality assuming there is $n \geq m$ such that $W_r = \uparrow \text{id}_n$.

As $W_l \cap I_{\mathbb{N}}^{<\infty} \in \mathcal{F}_1|_{I_{\mathbb{N}}^{<\infty}} = \mathcal{F}_2|_{I_{\mathbb{N}}^{<\infty}}$, we can (by Theorem 4.13(1)) find a wany set $N_{k,\mathfrak{Y}'} \in \mathcal{F}_2$ such that $N_{k,\mathfrak{Y}'} \cap I_{\mathbb{N}}^{<\infty} \subseteq W_l$ and $k \geq n$. Since $N_{m,\mathfrak{Y}} \notin \mathcal{F}_2$ and $N_{k,\mathfrak{Y}'} \in \mathcal{F}_2$, we get that $N_{k,\mathfrak{Y}'} \not\subseteq N_{m,\mathfrak{Y}}$. On the other hand, $(N_{k,\mathfrak{Y}'} \cap I_{\mathbb{N}}^{<\infty}) \cdot \uparrow \text{id}_n \subseteq W_l W_r \subseteq N_{m,\mathfrak{Y}}$.

Let $g \in N_{k,\mathfrak{Y}'} \setminus N_{m,\mathfrak{Y}}$. If g is finite, then $g \in (N_{k,\mathfrak{Y}'} \cap I_{\mathbb{N}}^{<\infty}) \cdot \text{id}_{\mathbb{N}} \subseteq (N_{k,\mathfrak{Y}'} \cap I_{\mathbb{N}}^{<\infty}) \cdot \uparrow \text{id}_n \subseteq N_{m,\mathfrak{Y}}$, which is a contradiction. Hence g is infinite. Then $g \notin N_{m,\mathfrak{Y}}$ and $k \geq m \geq n$ implies that $\text{im}(g)$ intersects every element of \mathfrak{Y} . Hence $N_{m,\mathfrak{Y}}$ contains no elements with the same image as g .

Let $g' := g|_{\{0,1,\dots,n-1\}g^{-1}}$. Since $g \in N_{k,\mathfrak{Y}'} = \downarrow N_{k,\mathfrak{Y}'}$, it follows that $g' \in N_{k,\mathfrak{Y}'} \cap I_{\mathbb{N}}^{<\infty} \subseteq W_l \in \mathcal{I}_4$. So $W_l \cap \uparrow g' \in \mathcal{I}_4$ and $(W_l \cap \uparrow g') \cdot \uparrow \text{id}_n \subseteq N_{m,\mathfrak{Y}}$. Since $g' \in W_l \cap \uparrow g' \in \mathcal{I}_4$, there must be some infinite $g'' \in W_l \cap \uparrow g'$ with $\text{im}(g'') \cap n = \text{im}(g') = \text{im}(g) \cap n$. So the set $g'' \cdot \uparrow \text{id}_n$ contains an element with the same image as g . This is a contradiction, as $g'' \cdot \uparrow \text{id}_n \subseteq W_l' \cdot \uparrow \text{id}_n \subseteq N_{m,\mathfrak{Y}}$, and $N_{m,\mathfrak{Y}}$ contains no elements with the same image as g . Therefore $\mathcal{F}_1 \subseteq \mathcal{F}_2$ and by symmetry $\mathcal{F}_1 = \mathcal{F}_2$. \square

Definition 4.18. We say that a non-empty set $\mathfrak{Y} \subseteq [\mathbb{N}]^{<\infty}$ is *bad* if there is a good filter \mathcal{F} and $n \in \mathbb{N}$ such that $N_{n,\mathfrak{Y}} \in \mathcal{F}$ and there is no finitely wany set $S \in \mathcal{F}$ with $S \subseteq N_{n,\mathfrak{Y}}$.

Note that if \mathfrak{Y} is bad and the good filter \mathcal{F} witnesses this, then \mathcal{F} has no filter base consisting of finitely wany sets.

We will show (eventually) that there are, in fact, no bad sets.

Recall that $[\mathbb{N}]^n$ denotes the set of all subsets of \mathbb{N} of cardinality n .

Lemma 4.19. *If there exists a bad set $\mathfrak{Y} \subseteq [\mathbb{N}]^{<\infty}$, then there is a set $\mathfrak{Y}' \subseteq [\mathbb{N}]^{<\infty}$ such that the following hold:*

- (1) *There is a good filter \mathcal{F} such that $N_{0,\mathfrak{Y}'} \in \mathcal{F}$, and there is no finitely many set S , with $S \cap I_{\mathbb{N}}^{<\infty} \in \mathcal{F}|_{I_{\mathbb{N}}^{<\infty}}$ and $S \cap I_{\mathbb{N}}^{<\infty} \subseteq N_{0,\mathfrak{Y}'}$.*
- (2) *The elements of \mathfrak{Y}' are incomparable with respect to containment and if $Y \in [\mathbb{N}]^{<\infty}$ and $\{f \in I_{\mathbb{N}}^{<\infty} \mid \text{im}(f) \cap Y = \emptyset\} \subseteq N_{0,\mathfrak{Y}'}$, then there is $Y' \in \mathfrak{Y}'$ with $Y' \subseteq Y$.*
- (3) *For all $n \in \mathbb{N}$, $[\mathbb{N}]^n \cap \mathfrak{Y}'$ is finite.*

Proof. We first construct a set satisfying (1), we will then modify this set to satisfy (2) while ensuring it still satisfies (1). The resulting set will satisfy (3) as well. Let \mathfrak{Y} be a bad set and let \mathcal{F} be a good filter witnessing that \mathfrak{Y} is bad.

By Theorem 4.13, we can find a sequence $(N_{i,\mathfrak{Y}_i})_{i \in \mathbb{N}}$ of many sets forming a filter base for \mathcal{F} .

For a good filter \mathcal{F}' and a set $\mathfrak{Y}' \subseteq [\mathbb{N}]^{<\infty}$, let $P(\mathcal{F}', \mathfrak{Y}')$ be the statement:

- $N_{0,\mathfrak{Y}'} \in \mathcal{F}'$, and there is no finitely many set S , with $S \cap I_{\mathbb{N}}^{<\infty} \in \mathcal{F}'|_{I_{\mathbb{N}}^{<\infty}}$ and $S \cap I_{\mathbb{N}}^{<\infty} \subseteq N_{0,\mathfrak{Y}'}$.

Seeking a contradiction, assume that for every $i \in \mathbb{N}$ the statement $P(\mathcal{F}, \mathfrak{Y}_i)$ is false. Then for all $i \in \mathbb{N}$, we can find a finitely many set S_i such that $S_i \cap I_{\mathbb{N}}^{<\infty} \in \mathcal{F}|_{I_{\mathbb{N}}^{<\infty}}$ and $S_i \cap I_{\mathbb{N}}^{<\infty} \subseteq N_{0,\mathfrak{Y}_i}$. Since \mathcal{F} contains every \mathcal{I}_2 neighbourhood of \emptyset , for all $i \in \mathbb{N}$ the set $U_i := \{f \in I_{\mathbb{N}} \mid \text{dom}(f) \cap i = \emptyset\}$ belongs to \mathcal{F} . Hence replacing S_i by the finitely many set $S_i \cap U_i$ if needed, we can assume that $S_i \cap I_{\mathbb{N}}^{<\infty} \subseteq N_{i,\mathfrak{Y}_i}$ for all $i \in \mathbb{N}$. Since $\{N_{i,\mathfrak{Y}_i} \mid i \in \mathbb{N}\}$ is a filter base for \mathcal{F} , it follows that $\{S_i \cap I_{\mathbb{N}}^{<\infty} \mid i \in \mathbb{N}\}$ is a filter base for $\mathcal{F}|_{I_{\mathbb{N}}^{<\infty}}$.

By Theorem 4.13(3), there is a filter \mathcal{F}' defined by a waning function such that $\mathcal{F}'|_{I_{\mathbb{N}}^{<\infty}} = \mathcal{F}|_{I_{\mathbb{N}}^{<\infty}}$. By Lemma 4.17, $\mathcal{F}' = \mathcal{F}$. This is a contradiction as by assumption \mathcal{F} witnesses that \mathfrak{Y} is bad, and \mathcal{F} has a filter base consisting of finitely many sets (see Remark 4.10).

Thus we can find a set \mathfrak{A} such that $P(\mathcal{F}, \mathfrak{A})$.

We next define a set \mathfrak{Y}' using \mathfrak{A} to satisfy conditions (1) and (2). First, however, let

$$\mathfrak{A}' := \mathfrak{A} \cup \{Y \in [\mathbb{N}]^{<\infty} \mid \{f \in I_{\mathbb{N}}^{<\infty} \mid \text{im}(f) \cap Y = \emptyset\} \subseteq N_{0,\mathfrak{A}}\}.$$

Note that $N_{0,\mathfrak{A}} \subseteq N_{0,\mathfrak{A}'}$ and $N_{0,\mathfrak{A}} \cap I_{\mathbb{N}}^{<\infty} = N_{0,\mathfrak{A}'} \cap I_{\mathbb{N}}^{<\infty}$ and hence \mathfrak{A}' also satisfies (1). Let \mathfrak{Y}' be the set obtained from \mathfrak{A}' by removing all elements from \mathfrak{A}' which are not minimal with respect to containment. It is easy to see that $N_{0,\mathfrak{A}'} = N_{0,\mathfrak{Y}'}$. Note that \mathfrak{Y}' now satisfies (1) and (2).

In order to show that \mathfrak{Y}' satisfies (3), we need the following auxiliary fact.

Claim 4.20. *There are pairwise disjoint $Y_1, \dots, Y_k \in \mathfrak{Y}'$ such that $Y \cap \left(\bigcup_{1 \leq i \leq k} Y_i\right) \neq \emptyset$ for all $Y \in \mathfrak{Y}'$.*

Proof. Note that if $\emptyset \in \mathfrak{Y}'$, then $N_{0,\mathfrak{Y}'} = I_{\mathbb{N}} = N_{0,\{\emptyset\}}$. Then the finitely many set $N_{0,\mathfrak{Y}'}$ belongs to \mathcal{F} , and so setting $S = N_{0,\mathfrak{Y}'}$, we obtain that \mathfrak{Y}' doesn't satisfy condition (1). The obtained contradiction implies $\emptyset \notin \mathfrak{Y}'$.

Suppose for a contradiction that the claim is false. Let $Y_1 \in \mathfrak{Y}'$ be arbitrary. If $Y_1, \dots, Y_l \in \mathfrak{Y}'$ are defined and pairwise disjoint, then as the claim is false we can find an $Y_{l+1} \in \mathfrak{Y}'$ disjoint from all of them. So we have an infinite sequence of pairwise disjoint sets $Y_1, Y_2, \dots \in \mathfrak{Y}'$. If $f \in I_{\mathbb{N}}^{<\infty}$, then $\text{im}(f)$ intersects at most finitely many of the Y_i , hence $N_{0,\mathfrak{Y}'} \cap I_{\mathbb{N}}^{<\infty} = I_{\mathbb{N}}^{<\infty} = N_{0,\{\emptyset\}} \cap I_{\mathbb{N}}^{<\infty}$. This is a contradiction to the fact that \mathfrak{Y}' satisfies (1). \square

We show inductively that:

- ★ For all $n \in \mathbb{N}$, there is a finite set $B_n \subseteq \mathbb{N}$ such that for all $Y \in \mathfrak{Y}'$, we have $|Y \cap B_n| \geq \min\{|Y|, n\}$.

Note that this implies that if $|Y| \leq n$ then $Y \subseteq B_n$, in particular this will imply (3) since $[\mathbb{N}]^n \cap \mathfrak{Y}' \subseteq \mathcal{P}(B_n)$. For the base case of the induction we set $B_1 = \bigcup_{1 \leq i \leq k} Y_i$ where the sets Y_1, \dots, Y_k are those from Claim 4.20.

Suppose inductively that the statement holds for some $i \in \mathbb{N}$, so the set B_i is defined. For every subset S of B_i , let

$$\mathfrak{Y}'_S := \{Y \setminus B_i \mid Y \in \mathfrak{Y}' \text{ and } Y \cap B_i = S\} = \{Y \setminus S \mid Y \in \mathfrak{Y}' \text{ and } Y \cap B_i = S\}.$$

Let $S \subseteq B_i$ be arbitrary. If $\mathfrak{Y}'_S \neq \emptyset$, let $Y_{1,S} \in \mathfrak{Y}'_S$ (otherwise define $Y_{1,S} := \emptyset$). If $Y_{1,S}, \dots, Y_{j,S}$ are defined, then let $Y_{j+1,S} \in \mathfrak{Y}'_S$ be disjoint from $Y_{1,S} \cup \dots \cup Y_{j,S}$ (if this is impossible define $Y_{j+1,S} := \emptyset$).

Claim 4.21. $\bigcup_{j=1}^{\infty} Y_{j,S}$ is finite.

Proof. Suppose for a contradiction that the set is infinite. It follows that $Y_{j,S} \neq \emptyset$ for infinitely many j . Without loss of generality suppose that every $Y_{j,S}$ is non-empty. Note that, for all j , $Y_{j,S} \cup S \in \mathfrak{Y}'$, and $Y_{j,S} \cap B_i = \emptyset$.

We show that $\{f \in I_{\mathbb{N}}^{<\infty} \mid \text{im}(f) \cap S = \emptyset\} \subseteq N_{0,\mathfrak{Y}'}$ (so we can apply part (2) to S). Let $f \in I_{\mathbb{N}}^{<\infty}$ be such that $\text{im}(f) \cap S = \emptyset$. As $\text{im}(f)$ is finite, and the sets $Y_{j,S}$ are non-empty and pairwise disjoint, and there are infinitely many such sets, there is $j \in \mathbb{N}$ such that $Y_{j,S} \cap \text{im}(f) = \emptyset$. It follows that $(Y_{j,S} \cup S) \cap \text{im}(f) = \emptyset$, so $f \in N_{0,\mathfrak{Y}'}$. By (2) (using S as Y), there is a subset Y' of S belonging to \mathfrak{Y}' .

Let $Y \in \mathfrak{Y}'_S$ be arbitrary. Then $Y' \subseteq Y \cup S \in \mathfrak{Y}'$ and so $Y', Y \cup S \in \mathfrak{Y}'$. But elements of \mathfrak{Y}' are incomparable and so $Y' \subseteq S \subseteq Y \cup S = Y'$. Since Y and S are disjoint, $Y = \emptyset$ which shows that $\mathfrak{Y}'_S \subseteq \{\emptyset\}$. This contradicts the assumption that the sets $Y_{j,S}$ are non-empty, as required. \square

If $Y \in \mathfrak{Y}'$ and $Y \cap B_i = S$, then we have by construction that either $Y \subseteq B_i$ or

$$Y \cap \bigcup_{j=1}^{\infty} Y_{j,S} \neq \emptyset.$$

Thus the set

$$B_{i+1} = B_i \cup \bigcup_{S \subseteq B_i} \bigcup_{j=1}^{\infty} Y_{j,S}$$

satisfies \star . \square

Lemma 4.22. *Bad sets do not exist.*

Proof. Suppose for a contradiction that bad sets do exist, and let \mathfrak{Y} be such a set. Let \mathfrak{Y}' be the set from Lemma 4.19, and let \mathcal{F} be the good filter from Lemma 4.19(1). By Theorem 4.13(1), we may suppose that

$$N_{1,\mathfrak{Y}_1} \supseteq N_{2,\mathfrak{Y}_2} \supseteq \dots$$

is a filter base for \mathcal{F} for some sets $\mathfrak{Y}_i \subseteq [\mathbb{N}]^{<\omega}$. Since \mathcal{F} is the set of neighbourhoods of \emptyset in a semigroup topology on $I_{\mathbb{N}}$ and composition with an element of $\text{Sym}(\mathbb{N})$ is a homeomorphism, it follows that $\mathcal{F}\phi = \mathcal{F}$ for all $\phi \in \text{Sym}(\mathbb{N})$, where

$$\mathcal{F}\phi = \{A\phi : A \in \mathcal{F}\}.$$

In particular, since $N_{0,\mathfrak{Y}'} \in \mathcal{F}$ it follows that $N_{0,\mathfrak{Y}'}\phi \in \mathcal{F}$ for any $\phi \in \text{Sym}(\mathbb{N})$. Since the sets N_{i,\mathfrak{Y}_i} are a base for \mathcal{F} , there exists $i_\phi \in \mathbb{N}$ such that $N_{i_\phi,\mathfrak{Y}_{i_\phi}} \subseteq N_{0,\mathfrak{Y}'}\phi$. For all $j \in \mathbb{N}$, let $S_j := \{\phi \in \text{Sym}(\mathbb{N}) \mid i_\phi = j\}$. Note that

$$\text{Sym}(\mathbb{N}) = \bigcup_{i \in \mathbb{N}} S_i.$$

As $\text{Sym}(\mathbb{N})$ is Baire (under the pointwise topology), at least one of the sets S_j is not nowhere dense. In other words, the closure of S_j contains a neighbourhood of some $\theta \in \text{Sym}(\mathbb{N})$.

If $\phi \in \theta^{-1}S_j$, then $\theta\phi \in S_j$ and so from the definition of S_j ,

$$\begin{aligned}
 N_{j, \mathfrak{Y}_j} \subseteq N_{0, \mathfrak{Y}'\theta\phi} &= \{f \in I_{\mathbb{N}} : \exists Y \in \mathfrak{Y}', \text{im}(f) \cap Y = \emptyset\} \theta\phi \\
 (\star) \qquad \qquad \qquad &= \{f \in I_{\mathbb{N}} : \exists Y \in \mathfrak{Y}', \text{im}(f) \cap Y\theta = \emptyset\} \phi \\
 &= N_{0, \mathfrak{Y}'\theta\phi}.
 \end{aligned}$$

Also $\mathfrak{Y}'\theta$ satisfies the conditions of Lemma 4.19 with respect to the same \mathcal{F} (as $\mathcal{F}\theta = \mathcal{F}$ and $\mathcal{F}|_{I_{\mathbb{N}}^{<\infty}}\theta = \mathcal{F}|_{I_{\mathbb{N}}^{<\infty}}$). Since the closure of S_j contains a neighbourhood of θ , it follows that the closure of $\theta^{-1}S_j$ contains a neighbourhood of $\text{id}_{\mathbb{N}}$. In other words, the closure of $\theta^{-1}S_j$ contains the pointwise stabilizer $\{f \in \text{Sym}(\mathbb{N}) \mid x \in F \Rightarrow (x)f = x\}$ in $\text{Sym}(\mathbb{N})$ of some finite set F .

Claim 4.23. *There exists $Y \in \mathfrak{Y}_j$ such that there is no $Y' \in \mathfrak{Y}'\theta$ with $Y' \subseteq F \cap Y$.*

Proof. If the claim is false, then for all $Y \in \mathfrak{Y}_j$ there is $Y' \in \mathfrak{Y}'\theta$ such that $Y' \subseteq F \cap Y$. Thus

$$N_{j, \mathfrak{Y}_j} \cap I_{\mathbb{N}}^{<\infty} \subseteq \{f \in I_{\mathbb{N}}^{<\infty} : \text{im}(f) \cap Y' = \emptyset \text{ for some } Y' \in \mathfrak{Y}'\theta \cap \mathcal{P}(F)\} \subseteq N_{0, \mathfrak{Y}'\theta \cap \mathcal{P}(F)} \subseteq N_{0, \mathfrak{Y}'\theta}.$$

Then the set $S = N_{0, \mathfrak{Y}'\theta \cap \mathcal{P}(F)}$ is finitely wany, $S \cap I_{\mathbb{N}}^{<\infty} \in \mathcal{F}|_{I_{\mathbb{N}}^{<\infty}}$, and $S \cap I_{\mathbb{N}}^{<\infty} \subseteq N_{0, \mathfrak{Y}'\theta}$. But Lemma 4.19(1) applied to $\mathfrak{Y}'\theta$ states that no such set S exists, a contradiction. \square

Let the set $Y \in \mathfrak{Y}_j$ be as given in Claim 4.23. Let B be the union of all elements of $\mathfrak{Y}'\theta$ of size at most $|Y|$. By Lemma 4.19(3), the set B is finite. Since $Y \in \mathfrak{Y}_j$, Y is also finite. Hence there exists $\phi \in \text{Sym}(\mathbb{N})$ fixing F pointwise and satisfying $(B \setminus F)\phi \cap Y = \emptyset$. Since the closure of $\theta^{-1}S_j$ contains the pointwise stabilizer of F , we can choose $\phi \in \theta^{-1}S_j$. From (\star) , $N_{0, \mathfrak{Y}_j} \subseteq N_{0, \mathfrak{Y}'\theta\phi}$, and Lemma 4.19(2) implies that there is $Y' \in \mathfrak{Y}'\theta\phi$ such that $Y' \subseteq Y$. In particular $Y'\phi^{-1} \in \mathfrak{Y}'\theta$ and has at most $|Y|$ elements. So $Y'\phi^{-1} \subseteq B$ implying that $Y' \subseteq B\phi$. Thus $Y' \subseteq B\phi \cap Y$. But by the choice of ϕ , $B\phi \cap Y \subseteq F$. So $Y' \subseteq F \cap Y$ and since ϕ fixes F pointwise, $Y'\phi^{-1} = Y' \subseteq F \cap Y$. This is a contradiction since, as was mentioned before, $Y'\phi^{-1} \in \mathfrak{Y}'\theta$ and Y was chosen to satisfy the property given in Claim 4.23. \square

Corollary 4.24. *If \mathcal{F} is good, then it can be defined by a waning function.*

Proof. By Theorem 4.13(1), \mathcal{F} has a filter base \mathcal{B} consisting of wany sets. By Lemma 4.22, every set $B \in \mathcal{B}$ is not bad, and so there exists a finitely wany set $A_B \in \mathcal{F}$ such that $A_B \subseteq B$. Hence the collection $\{A_B : B \in \mathcal{B}\}$ of finitely wany sets is a filter base for \mathcal{F} as well. Thus by Theorem 4.13(2), \mathcal{F} can be defined by a waning function. \square

Definition 4.25. Let \mathcal{T} be a second countable semigroup topology on $I_{\mathbb{N}}$ containing \mathcal{I}_2 , with filter data $(\mathcal{F}_{\mathcal{T}, i})_{i \in \omega}$. By Definition 4.7, every $\mathcal{F}_{\mathcal{T}, i}$ is good, and so, by Corollary 4.24, there exists a waning function $f_i : \omega + 1 \rightarrow \omega + 1$ such that $\mathcal{F}_{\mathcal{T}, i}$ is the set of all neighbourhoods of \emptyset in the topology \mathcal{T}_{f_i} .

We define a function $f_{\mathcal{T}} : (\omega + 1) \rightarrow (\omega + 1)$ by

- (1) $(i)f_{\mathcal{T}} = (0)f_i$ for all $i \in \omega$;
- (2) $(\omega)f_{\mathcal{T}} = \min\{(0)f_{\mathcal{T}}, (1)f_{\mathcal{T}}, \dots\}$.

One might think that $f_{\mathcal{T}}$ depends on the choice of the waning functions f_i in Definition 4.25. However, if $f, g : \omega + 1 \rightarrow \omega + 1$ are waning functions, then the topologies \mathcal{T}_f and \mathcal{T}_g have the same neighbourhoods of \emptyset if and only if $(0)f = (0)g$ (by Corollary 3.13). Hence if f_i and g_i are waning functions such that \mathcal{F}_i is the set of neighbourhoods of \emptyset in \mathcal{T}_{f_i} and \mathcal{T}_{g_i} , then $(0)f_i = (0)g_i$. It follows that $f_{\mathcal{T}}$ is independent of the choice of the f_i in Definition 4.25.

Theorem 4.26. *If \mathcal{T} is a second countable semigroup topology on $I_{\mathbb{N}}$ containing \mathcal{I}_2 , then $f_{\mathcal{T}}$ is a waning function.*

Proof. If $(\mathcal{F}_i)_{i \in \omega}$ is the filter data of \mathcal{T} , then by Lemma 4.8, $\mathcal{F}_m \subseteq \mathcal{F}_n$ for all $m \leq n$. So, by Corollary 3.13, if f_m and f_n are waning functions corresponding to \mathcal{F}_m and \mathcal{F}_n (as in Definition 4.25), then $(0)f_m \geq (0)f_n$. In other words, $f_{\mathcal{T}}$ is non-increasing.

It therefore suffices to show that if $i \in \mathbb{N}$ and $(i)f_{\mathcal{T}} \in \mathbb{N} \setminus \{0\}$, then $(i+1)f_{\mathcal{T}} < (i)f_{\mathcal{T}}$.

Claim 4.27. *If $g \in I_{\mathbb{N}}^{<\infty}$ and $N \subseteq I_{\mathbb{N}}$, then N is a neighbourhood of g with respect to \mathcal{T} if and only if N contains the sets*

$$\{h \in I_{\mathbb{N}} \mid h \upharpoonright_r = g \upharpoonright_r \text{ and } |\text{im}(h) \cap (r \setminus \text{im}(g))| \leq (|g|)f_{\mathcal{T}}\}$$

for all but finitely many $r \in \mathbb{N}$.

Proof. Suppose that $g \in I_{\mathbb{N}}^{<\infty}$. Then

$$\mathcal{F}_{|g|} = \mathcal{F}_{\mathcal{T},g} = \{(N \cap \uparrow g)d_g : N \text{ is a neighbourhood of } g \text{ with respect to } \mathcal{T}\}.$$

(by Definition 4.5 and Lemma 4.4). Hence N is a neighbourhood of g if and only if $(N \cap \uparrow g)d_g \in \mathcal{F}_{|g|}$. But $\mathcal{F}_{|g|}$ is the set of all neighbourhoods of \emptyset in $\mathcal{T}_{f_{|g|}}$ for some waning function $f_{|g|}$ (as in Definition 4.25). By the definition of $f_{\mathcal{T}}$, $(|g|)f_{\mathcal{T}} = (0)f_{|g|}$, and so Lemma 3.5 implies that $N \in \mathcal{F}_{|g|}$ if and only if N contains a set of the form

$$W_{f_{|g|},\emptyset,r} = \{h \in I_{\mathbb{N}} \mid h \upharpoonright_r = \emptyset \text{ and } |\text{im}(h) \cap r| \leq (0)f_{|g|} = (|g|)f_{\mathcal{T}}\}$$

for some large enough $r \in \mathbb{N}$. It follows that N is a neighbourhood of g (with respect to \mathcal{T}) if and only if $(N \cap \uparrow g)d_g$ contains a set $W_{f_{|g|},\emptyset,r}$ for large enough r .

We claim that the set $(N \cap \uparrow g)d_g$ contains a set $W_{f_{|g|},\emptyset,r}$ for all but finitely many $r \in \mathbb{N}$ if and only if N contains a set of the form

$$\{h \in I_{\mathbb{N}} \mid h \upharpoonright_r = g \upharpoonright_r \text{ and } |\text{im}(h) \cap (r \setminus \text{im}(g))| \leq (|g|)f_{\mathcal{T}}\}$$

for all but finitely many $r \in \mathbb{N}$, as we can check that if $r > |g|$ then

$$\{h \in I_{\mathbb{N}} \mid h \upharpoonright_r = g \upharpoonright_r \text{ and } |\text{im}(h) \cap (r \setminus \text{im}(g))| \leq (|g|)f_{\mathcal{T}}\} = W_{f_{|g|},\emptyset,r-|g|}d_g^{-1}.$$

Hence

$$(\{h \in I_{\mathbb{N}} \mid h \upharpoonright_r = g \upharpoonright_r \text{ and } |\text{im}(h) \cap (r \setminus \text{im}(g))| \leq (|g|)f_{\mathcal{T}}\})d_g = W_{f_{|g|},\emptyset,r-|g|}.$$

□

By Claim 4.27, the set

$$N := \{f \in I_{\mathbb{N}} \mid \text{id}_i \subseteq f \text{ and } |\text{im}(f) \cap \{i, i+1, \dots, i+(i)f_{\mathcal{T}}\}| \leq (i)f_{\mathcal{T}}\}$$

is a neighbourhood of id_i with respect to \mathcal{T} . Let $U \in \mathcal{T}$ be such that $\text{id}_i \in U \subseteq N$. As U is a neighbourhood of id_i and $(i)f_{\mathcal{T}} \neq 0$, it follows that we can find $g \in U$ such that $\text{im}(g) = i+1$. As U is open, it is also a neighbourhood of g . So N is a neighbourhood of g .

It follows that

$$\begin{aligned} N' &:= \{f \in I_{\mathbb{N}} \mid g \subseteq f \text{ and } |\text{im}(f) \cap \{i, i+1, \dots, i+(i)f_{\mathcal{T}}\}| \leq (i)f_{\mathcal{T}}\} \\ &= \{f \in I_{\mathbb{N}} \mid g \subseteq f \text{ and } |\text{im}(f) \cap \{i+1, \dots, i+(i)f_{\mathcal{T}}\}| \leq (i)f_{\mathcal{T}} - 1\} \end{aligned}$$

is a neighbourhood of g with respect to \mathcal{T} . Since $|g| = i+1$, it follows that $(i+1)f_{\mathcal{T}} \leq (i)f_{\mathcal{T}} - 1$ as required. □

Theorem 4.28. *If \mathcal{T} is a second countable semigroup topology on $I_{\mathbb{N}}$ containing \mathcal{I}_2 , then $\mathcal{T} = \mathcal{T}_{f_{\mathcal{T}}}$. In particular every second countable, semigroup topology on $I_{\mathbb{N}}$ containing \mathcal{I}_2 can be defined by a waning function.*

Proof. By the definition of $f_{\mathcal{T}}$, Lemma 3.5, and Claim 4.27, we know that for all $g \in I_{\mathbb{N}}^{<\infty}$, g has the same neighbourhoods with respect to \mathcal{T} as $\mathcal{T}_{f_{\mathcal{T}}}$. Thus we need only consider the neighbourhoods of the infinite elements of $I_{\mathbb{N}}$.

We split the proof into two cases.

Claim 4.29. *If $(\omega)f_{\mathcal{T}} = 0$, then $\mathcal{T} = \mathcal{T}_{f_{\mathcal{T}}}$.*

Proof. Let $N \in \mathbb{N}$ be such that $(N)f_{\mathcal{T}} = 0$. Let $a \in \mathbb{N}$ be arbitrary. We show that the set

$$\{f \in I_{\mathbb{N}} \mid N \leq |f| \text{ and } a \notin \text{im}(f)\}$$

is open with respect to \mathcal{T} . Note that this is sufficient as the \mathcal{I}_4 neighbourhoods of any infinite element of $I_{\mathbb{N}}$ are generated by these sets and sets from \mathcal{I}_2 .

Let $F_{N,a} := \{f \in I_{\mathbb{N}} \mid |f| = N \text{ and } a \notin \text{im}(f)\}$. Observe that

$$\{f \in I_{\mathbb{N}} \mid N \leq |f| \text{ and } a \notin \text{im}(f)\} = \bigcup_{g \in F_{N,a}} \{f \in I_{\mathbb{N}} \mid g \subseteq f \text{ and } a \notin \text{im}(f)\}.$$

Thus we need only show that the sets in the above union are open.

Let $g \in F_{N,a}$ be arbitrary. Define $\phi : \{f \in I_{\mathbb{N}} \mid g \subseteq f\} \rightarrow \{f \in I_{\mathbb{N}} \mid N \leq |f| < \infty\}$ by

$$(f)\phi = f \circ \text{id}_{\text{im}(g) \cup \{a\}}.$$

Note that ϕ is continuous, well-defined, and the topology on $\{f \in I_{\mathbb{N}} \mid N \leq |f| < \infty\}$ is the subspace topology inherited from $(I_{\mathbb{N}}, \mathcal{I}_4)$. It follows that

$$(\{f \in I_{\mathbb{N}} \mid a \notin \text{im}(f)\})\phi^{-1} = \{f \in I_{\mathbb{N}} \mid g \subseteq f \text{ and } a \notin \text{im}(f)\}$$

is open in $\{f \in I_{\mathbb{N}} \mid g \subseteq f\}$, which is also open in $\mathcal{T} \supseteq \mathcal{I}_2$ so the result follows. \square

Claim 4.30. *If $(\omega)f_{\mathcal{T}} = \omega$, then $\mathcal{T} = \mathcal{T}_{f_{\mathcal{T}}} = \mathcal{I}_2$.*

Proof. Let $W \in \mathcal{T}$, and $g \in W$ be infinite. We need only show that W is a \mathcal{I}_2 neighbourhood of g . Consider the product $\text{id}_{\mathbb{N}}g = g$. As $\mathcal{I}_2 \subseteq \mathcal{T} \subseteq \mathcal{I}_4$ and $\mathcal{B} := \{\uparrow \text{id}_n : n \in \mathbb{N}\}$ is an open neighborhood base of $\text{id}_{\mathbb{N}}$ in both \mathcal{I}_2 and \mathcal{I}_4 , we get the family \mathcal{B} also forms a base at $\text{id}_{\mathbb{N}}$ in \mathcal{T} . Since \mathcal{T} is a semigroup topology, there is $U \in \mathcal{T}$ and $n \in \mathbb{N}$ such that

$$(\uparrow \text{id}_n)U \subseteq W.$$

Since $U \in \mathcal{I}_4$, there exists finite $g' \subseteq g$ such that $g \upharpoonright_n = g' \upharpoonright_n$ and $g' \in U$. As g' is finite and $f_{\mathcal{T}}(|g'|) = \omega$, it follows that U is a \mathcal{I}_2 neighbourhood of g' . So there is some finite $k \in \mathbb{N}$ such that

$$U_{k,g'} := \{f \in I_{\mathbb{N}} \mid f \upharpoonright_k = g' \upharpoonright_k\} \subseteq U.$$

In particular

$$(\uparrow \text{id}_n)U_{k,g'} \subseteq W.$$

Let $m := \max\{\{k, n\} \cup \text{dom}(g') \cup \text{im}(g')\} + 1$. We have

$$(\uparrow \text{id}_n)U_{m,g'} \subseteq W.$$

It therefore suffices to show that $U_{n,g'} \subseteq (\uparrow \text{id}_n)U_{m,g'}$ as this set is open in \mathcal{I}_2 and contains g . Let $h \in U_{n,g'}$ be arbitrary. Put $h_1 = g \upharpoonright_n = g' \upharpoonright_n = h \upharpoonright_n$ and $h_2 = h \setminus h_1$. We have that

$$h = h_1 \cup h_2.$$

Let $l : \text{im}(h_2) \rightarrow \mathbb{N} \setminus m$ be an injective function.

Note that

$$\emptyset = \text{dom}(\text{id}_n) \cap \text{dom}(h_2l) = \text{im}(\text{id}_n) \cap \text{im}(h_2l) = \text{dom}(l^{-1}) \cap \text{dom}(g') = \text{im}(l^{-1}) \cap \text{im}(g').$$

It follows that

$$h = h_1 \cup h_2 = (\text{id}_n \circ g') \cup (h_2l \circ l^{-1}) = (\text{id}_n \cup h_2l) \circ (g' \cup l^{-1}) \in (\uparrow \text{id}_n)U_{m,g'}.$$

\square

The two claims above complete the proof of the theorem. \square

REFERENCES

- [1] S. Bardyla, L. Elliott, J. D. Mitchell and Y. Péresse, *Topological embeddings into transformation monoids*, Forum Mathematicum, 2024 (published online), <https://doi.org/10.1515/forum-2023-0230>
- [2] H. Becker, A. S. Kechris, *The descriptive set theory of Polish group actions*, London Math. Soc. Lecture Notes Ser. 232, Cambridge Univ. Press, 1996.
- [3] I. Yaacov, A. Berenstein and J. Melleray, *Polish topometric groups*, Transactions of the American Mathematical Society, **365** (2013), 3877–3897
- [4] R. M. Bryant, D. M. Evans, *The small index property for free groups and relatively free groups*, J. London Math. Soc. **55**:2 (1997), 363–369.
- [5] M. Cohen, R. R. Kallman, *$PL_+(I)$ is not a Polish group*, Ergodic Theory and Dynamical Systems **36** (2016), 2121–2137.
- [6] R. M. Dudley, *Continuity of homomorphisms*, Duke Mathematical Journal **28** (1961), 587–594.
- [7] L. Elliott, J. Jonušas, Z. Mesyan, J. D. Mitchell, M. Morayne and Y. Péresse, *Automatic continuity, unique Polish topologies, and Zariski topologies on monoids and clones*, Trans. Amer. Math. Soc. **376**:11 (2023), 8023–8093.
- [8] L. Elliott, J. Jonušas, J. D. Mitchell, Y. Péresse, and M. Pinsker, *Polish topologies on endomorphism monoids of relational structures*, Adv. Math. **431** (2023), 109214.
- [9] P. Gartside, B. Pejić, *Uniqueness of Polish group topology*, Topology Appl. **155** (2008), 992–999.
- [10] E. D. Gaughan, *Topological group structures of infinite symmetric groups*, Proceedings of the National Academy of Sciences of the United States of America, **58** (1967), 907–910.
- [11] W. Hodges, I. Hodkinson, D. Lascar and S. Shelah, *The small index property for ω -stable ω -categorical structures and for the random graph*, J. London Math. Soc. **S2-48**:2 (1993), 204–218.
- [12] Howie, J. *Fundamentals of Semigroup Theory*, (Oxford university Press, 1995)
- [13] R. R. Kallman, *A uniqueness result for topological groups*, Proc. Amer. Math. Soc. **54** (1976), 439–440.
- [14] R. R. Kallman, *Uniqueness results for groups of measure preserving transformations*, Proc. Amer. Math. Soc. **95** (1985), 87–90.
- [15] R. R. Kallman, *Uniqueness results for homeomorphism groups*, Trans. Amer. Math. Soc. (1986), 389–396.
- [16] R. R. Kallman, A. P. McLinden, *The Poincaré and related groups are algebraically determined Polish groups*, Collect. Math. **61** (2010), 337–352.
- [17] A. Kechris, *Classical Descriptive Set Theory*, volume 156 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1995. Berlin, 1989.
- [18] A. Kechris, C. Rosendal. *Turbulence, amalgamation and generic automorphisms of homogeneous structures*. Proceedings of the London Mathematical Society, **94**:2 (2007), 302–350.
- [19] D. Lascar, *Autour De La Propriété Du Petit Indice*, Proceedings of the London Mathematical Society **s3-62** (1991), 25–53.
- [20] M. Lawson, *Inverse semigroups, the theory of partial symmetries*, World Scientific Publishing, Singapore, 1998.
- [21] K. Mann, *Automatic continuity for homeomorphism groups and applications*, Geometry & Topology **20** (2016), 3033–3056.
- [22] D. Montgomery, *Continuity in topological groups*, Bull. Amer. Math. Soc. **42** (1936), 879–882.
- [23] J. Pérez, C. Uzcátegui, *Topologies on the symmetric inverse semigroup*, Semigroup Forum **104**:2 (2022), 398–414.
- [24] M. Petrich, *Inverse Semigroups*, John Wiley & Sons, 1984.
- [25] M. Pinsker, C. Schindler, *The semigroup of increasing functions on the rational numbers has a unique Polish topology*, preprint, 2023, 10.48550/arXiv.2305.04921.
- [26] C. Rosendal, *On the non-existence of certain group topologies*, Fund. Math. **187** (2005), 213–228.
- [27] C. Rosendal, S. Solecki, *Automatic continuity of homomorphisms and fixed points on metric compacta*, Israel J. Math. **162** (2007), 349–371.
- [28] M. Sabok, *Automatic continuity for isometry groups*, J. Inst. Math. Jussieu **18** (2019), 561–590.
- [29] J. K. Truss, *Infinite permutation groups II. Subgroups of small index*, J. Algebra **120**:2 (1989), 494–515.
- [30] T. Tsankov, *Automatic continuity for the unitary group*, Proc. Amer. Math. Soc. **141** (2013), 3673–3680.

S. BARDYLA: UNIVERSITY OF VIENNA, INSTITUTE OF MATHEMATICS, VIENNA, AUSTRIA
Email address: sbardyla@gmail.com

L. ELLIOTT: UNIVERSITY OF BINGHAMTON, DEPARTMENT OF MATHEMATICS AND STATISTICS, BINGHAMTON, SUNY, USA
Email address: luna.elliott142857@gmail.com

J. D. MITCHELL: UNIVERSITY OF ST ANDREWS, SCHOOL OF MATHEMATICS AND STATISTICS, SCOTLAND, UK
Email address: jdm3@st-andrews.ac.uk

Y. PÉRESSE: UNIVERSITY OF HERTFORDSHIRE, SCHOOL OF PHYSICS, ENGINEERING AND COMPUTER SCIENCE, UK
Email address: y.peresse@herts.ac.uk