

A Point-cloud Clustering & Tracking Algorithm for Radar Interferometry

Magnus F Ivarsen*

Department of Physics, University of Oslo, Oslo, Norway

Jean-Pierre St-Maurice[†] and Glenn C Hussey

Department of Physics and Engineering Physics, University of Saskatchewan, Saskatoon, Canada

Devin R Huyghebaert

Department of Physics and Technology, University of Tromsø, Tromsø, Norway

Megan D Gilles

Department of Chemistry and Physics, Mount Royal University, Calgary, Canada

(Dated: June 4, 2024)

In data mining, density-based clustering, which entails classifying datapoints according to their distributions in some space, is an essential method to extract information from large datasets. With the advent of software-based radio, ionospheric radars are capable of producing unprecedentedly large datasets of plasma turbulence backscatter observations, and new automatic techniques are needed to sift through them. We present an algorithm to automatically identify and track clusters of radar echoes through time, using **dbscan**, a celebrated density-based clustering method for noisy point-clouds. We demonstrate its efficiency by tracking turbulent structures in the E-region ionosphere, the so-called radar aurora. Through conjugate auroral imagery, as well as *in-situ* satellite observations, we demonstrate that the observed turbulent structures generally track the motion of the aurora. We discuss instances when this prediction is brought to fruition and when it ostensibly is not. Through case studies, we highlight the important instances when the radar echo bulk motions vary considerably around discrete auroral arcs, an effect we argue is produced by strong electric field modulations caused by energetic particle precipitation.

I. INTRODUCTION

In the practice of data mining, ‘data clustering’ attempts to segment a dataset into groups or classes that share common traits and characteristics. Crucially, these characteristics are not immediately discernible in individual datapoints, and their close association is hidden, as it were, in the data [1]. Density-based clustering, in turn, achieves this goal by classifying datapoints based on their density inside a space defined by two or more variables, a point-cloud. In the present study, we apply density-based clustering to interferometry radar data in the field of ionospheric research, an endeavour that will enable key insights into ionospheric physics.

With recent advances in radar and lidar (optical radar) technologies, spatial point-clouds are becoming increasingly common. For example, self-driving cars rely on real-time obstacle-detection using lidar scans, in which fast and accurate point-cloud clustering is vital [2, 3]. Lidar mounted on aircraft are frequently used in geographic surveying [4] and for discovering future archaeological sites [5]. In the survey of Earth’s space environment, the ICEBEAR 3D radar represents a recent advancement in radar technology that is capable of producing

3D point-clouds of plasma turbulence echoes in the lower ionosphere [6]. In a recent paper, [7] demonstrated that such echoes tend to be organized in tight clusters, and the degree of clustering is directly reflecting the filamentation of geomagnetic currents in the ionosphere [8, 24].

Unlike conventional radar and lidar imaging, in which the subject is a coherent man-made or natural object, ionospheric radar images are produced by turbulent, or stochastic, processes [9]. Measurements of stochastic processes are unpredictable and must in general be understood in terms of statistical representation, which is usually obtained through Fourier analysis [10]. The turbulent shapes that are imaged by the ICEBEAR radar are thus viable at any time to bifurcate, merge, dissipate, and grow, making efficient tracking of their bulk motion a challenging endeavour.

A. Plasma Turbulence in the E-region

The ionosphere, a partially ionized layer of gas existing in Earth’s atmosphere at altitudes above ~ 90 km, can at times be highly turbulent, with large portions of the ionosphere’s plasma exhibiting chaotic fluid properties [11]. Collectively and ultimately, the turbulence serves to dissipate the energy imparted to Earth from the solar wind during geomagnetic storms. Field-aligned currents and precipitating particles pass electric fields that stir the plasma, which, together with the interplanetary magnetic field, create a gigantic convection pattern that

* Also at Department of Physics and Engineering Physics, University of Saskatchewan, Saskatoon, Canada

[†] Also at Department of Physics and Astronomy, University of Western Ontario, London, Ontario, Canada

covers Earth’s high-latitude regions [12, 13], as well as direct large amounts of electromagnetic energy into the atmosphere [14]. The ionosphere’s conductivity, its ability to support electrical currents, is central to how these processes unfold, and small-scale turbulence in the ionosphere can cause stochastic changes to this conductivity [15, 16].

At high latitudes, on Earth’s nightside, the particles that precipitate into the atmosphere eventually collide with neutral gas particles, whereby they excite light emissions, creating the aurora [17]. The aurora, appearing to observers on the ground as elongated curtains of shifting, or dancing, light, is accompanied by a conspicuous radio signature in the bottomside (E-region) ionosphere, the radar aurora [18]. Radars, when pointed in the direction perpendicular to Earth’s magnetic field lines, can detect an echo of their outgoing signal, echoes that are produced when that signal is scattered off of sharp density gradients in the plasma. With a 50 MHz radar signal, such sharp gradients are 3-meter waves produced by the Farley-Buneman instability, a modified two-stream instability mechanism [19, 20].

The growth of FB waves in the E-region depends on the electric field being sufficiently strong [21]. Detectability of the structures by radar additionally necessitates a large enough ambient plasma density, without which radar echoes are too weak to exceed the received noise level [22]. As a result, the echo count inside a scattering volume depends both on the electric field strength and on the plasma density field. In addition, as alluded to, FB waves mostly grow in directions perpendicular to Earth’s magnetic field [21, 23]. The ICEBEAR radar, located in Saskatchewan, Canada, beams its signal towards the horizon. Here, given regions of favourable conditions, a continuous distribution of FB turbulence will produce a 3D point cloud every second. These echo point-clouds are frequently observed to cluster around auroral arcs [22, 24], a crucial observation that is valid for the radar aurora in general [25–27]. Since the aurora is an emergent phenomenon that arises through the interaction between the ionosphere, Earth’s magnetosphere, and the solar wind [28], we can infer key characteristics of the behaviour of echo clusters: their motion must be governed by magnetospheric processes, often arising from the complex interplay between Alfvénic and electrodynamic considerations [29].

In the present study, we describe a set of novel methods from data mining that we applied to automatically detect and track dense clusters of coherent backscatter echoes in the ICEBEAR 3D dataset. The ICEBEAR radar routinely detects multitudes of echoes in excess of 10^4 per minute, each of which is associated with a 3D spatial location. In space, then, the E-region turbulence regions frequently form, bifurcate, break up, and reappear, all the while exhibiting some form of motion in the plane perpendicular to Earth’s field lines. This motion is complicated and composite, and must be described as *apparent* by an observer on Earth. However, as we

demonstrate in a companion letter, a clear tendency to follow the flow and propagation of magnetospheric field structures is discernible in the motions of ICEBEAR echo clusters. The present paper is a *method* paper whose intent is to explain, elucidate, and discuss the process of extracting spatial tracks made by distinct echo clusters, as well as to demonstrate a clear and unambiguous tendency in those tracks to follow the motion of auroral arcs

II. METHODOLOGY & DATA

As already mentioned, our dataset consists of E-region radio echoes from 3-meter size irregularities produced by the FB instability, as detected by the ICEBEAR (Ionospheric Continuous-wave E-region Bistatic Experimental Auroral Radar), a 49.5 MHz coherent scatter radar in Saskatchewan, Canada [32]. The ICEBEAR 3D dataset is the result of a reconfiguration so that echo locations can be determined with a higher spatial and temporal resolution [6]. In this reconfiguration, the data processing chain produces a full field-of-view radio image for every range-Doppler time bin, thereby handling some 200,000 such images per second. In each image, a single target radar echo is identified by a single peak in the brightness distribution, at which point its location in latitude, longitude, and altitude is determined from its bearing and heading. This is in addition to the target echo’s spectral properties, particularly its Doppler shift.

The present work has been motivated by recent results showing that ICEBEAR 3D echoes appear to cluster tightly in the plane perpendicular to Earth’s magnetic field lines [7]. Building on this, we have applied the `dbscan` algorithm to automatically detect clusters in two-dimensional point-clouds [33, 34]. To implement the methods, we have tracked clusters through time with a rudimentary test of similarity from one time-step to the next. We sort the data into time-intervals of three seconds, with a one-second cadence. Based on this approach, we were able to systematically identify and track turbulent regions as they appeared in the radar FOV.

The `dbscan` algorithm assigns points to groups, or clusters, by requiring that the k^{th} nearest *neighbour* of a point is less than a distance ϵ away from that point. Points for which this distance is above ϵ are considered *noise* or outliers, and do not belong to any cluster. The threshold distance ϵ is determined by a so-called k -distance plot, which simply sorts the k^{th} nearest neighbour distance metric for each point. The kink or elbow of this curve, where the k^{th} nearest neighbour distance metric flattens to a plateau, identifies the characteristic noise threshold ϵ for a particular point-cloud. Examples of this procedure are shown in Figure 1, where the left column shows ICEBEAR 3D point-clouds, the middle column shows the k^{th} distance plot associated with each cloud, and the right column shows the resulting clusters that were identified.

In the middle column of Figure 1, we observe that the




FIG. 1. Examples of the echo point-cloud clustering categorization using the `dbSCAN` algorithm. Leftmost column: the raw echo locations. Middle column: k -distance plot for all the echoes (with $k = 100^{\text{th}}$ nearest neighbour). A solid red line gives a piecewise linear fit, while a dashed red line shows the determined threshold distance ϵ . Right column: displayed and labeled clusters that were identified. The labels attached to clusters in panels c), f), and i) do not carry over to other panels.

noise threshold ϵ takes on various values, based on the natural tendency for points to cluster in separate point-clouds. Based on extensive testing, we limited the value of ϵ to be between 3 km and 200 km, and we took the average value of ϵ determined within a six-second period to determine the transition to a plateau. In the right column of Figure 1, we illustrate how, based on our selection criteria, the `dbSCAN` algorithm has identified two clusters in each point-cloud (black data), with a smaller number of non-clustered points (green data).

The red 'bounding boxes' that enclose each cluster of

echoes is then used to track echo clusters through time, by requiring that the bounding box size and location of a particular cluster must be reasonably similar to a cluster identified at the previous timestep. The bounding box similarity test consists of minimizing three distance metrics: 1) the longitudinal extent of the bounding box, 2) the latitudinal extent of the bounding box, and 3) the median echo location inside the bounding box. Two consecutive populations that minimize the difference in each of the three distance metrics are determined to be the same population if the metrics are all smaller than




FIG. 2. Two example echo cluster tracks (solid blue lines) and their corresponding linear polynomial fits (dashed blue lines). The linear coefficient (v_E, v_N) is displayed for both clusters, with the 95-percentile (3-sigma) confidence interval for each fit denoting the error margin.

a pre-defined distance D . For the purpose of identifying and tracking large echo populations through time we have found, through extensive testing, that the procedure favours an abnormally high D (e.g., $D = 80$ km), as well as a relatively high k (e.g., $k = 100$). This ensures that echo populations that grow or shrink drastically will still be successfully tracked through time, and echo clusters that consist of very few echoes will be ignored. On the other hand, to accurately determine bulk velocities (which is the main focus of the results of the present paper) the procedure favours relatively low parameters (e.g., $D = 80$ km and $k = 20$). As we shall demonstrate in the two next sections, this scheme will produce a high number of tracks whose average motion will accurately quantify the bulk motion of the radar aurora.

The final product in this process are the *tracks* traced out by each cluster as it moves through space. In Figure 2 we show two such tracks as solid blue lines. We fit a linear polynomial to the longitudinal and latitudinal

positions (dashed blue lines). The coefficient, or slope, in these fits comprise the eastward and poleward bulk velocity respectively, and the 95-percentile (3-sigma) confidence intervals of those fits readily provide errorbars to those bulk velocity estimates. Individual clusters (such as those identified in Figure 1) produce spatial tracks, and the minutae of their time histories represent each cluster’s spatio-temporal evolution. In the next section we shall present extensive results of such bulk motion determination.

III. RESULTS

Figure 3 shows four such time histories, where we plot four equally spaced frames in the life of four selected clusters that were identified on 1 March and 25 August, in 2021, all observed pre-midnight magnetic local time. We display the auroral images (in the green channel) as contour lines of constant emission intensity, with green arrows showing the result of applying the Horn-Schunk optical flow algorithm to the images [35]. The four clusters exhibit bulk motions, and the spatial tracks (solid blue lines) are well approximated by the linear polynomial fits (dashed blue lines), with the notable exception of the bottom row (panels m–p), which we will discuss in the next paragraph. All the clusters of echoes are seen in close association with the spatial distribution of auroral emissions, in accordance with existing observations of auroral turbulence [22, 24, 26]. Likewise, for all the clusters in Figure 3, the direction of the straight, dashed blue lines in point in the overall direction of the aurora. The distribution of specific Doppler shifts, indicated by the color of the echo point-clouds, follow the distribution of auroral emissions, with distinct yellow regions (fast Doppler speeds) appearing close to auroral emissions, where the electric field is expected to maximize [36]. The Doppler speeds of radar echoes in the E-region are expected to correlate with the F-region drifts convection electric field, but are limited in magnitude by instability physics [37].

The first and last row of Figure 3 corresponds to the two tracks on display in Figure 2, showing how echo clusters can drift in a clear and coherent way (Figure 3a–d) or drifting hither and thither (Figure 3m–p), while still tracing out a spatial track that clearly follows the motion of an auroral form. In terms of ionospheric electrodynamics surrounding auroral arcs, we offer the following explanation. In the case of the former, Figure 3a–d), the echo cluster and the auroral form both follow an ambient ionospheric drift, which at that time and location (evening auroral oval) pointed in the south-western direction. In the case of the latter, Figure 3m–p), we likewise observe an auroral arc that is embedded into an ambient, convection electric field, but this time the auroral arc is unusually intense, producing a local electric field that is strong and variable, effectively *overriding* the ambient field (see Figure 7 in [38], Figure 5 in [39], and Figure 7 in [40]). This explanation makes sense if the ICEBEAR




FIG. 3. Detailed spatio-temporal evolution of four ICEBEAR echo clusters (red-yellow-white point-clouds, colorscale indicated), as well as their spatial tracks (solid blue lines). The point-clouds are superposed on optical auroral imaging, represented by green contour lines that trace pixel intensity. Green arrows show the optical flow of pixel intensity. A dashed blue line shows linear polynomials fitted to (magnetic) longitudinal and latitudinal components of the tracks, and display the overall direction travelled. The bulk speed inferred from those fits is displayed above each of the three rows.

echo clusters, which are excited by a strong perpendicular electric field [21], follow the motion of electric field source regions. In the next section, we will discuss and elaborate on the premise for this suggestion.

IV. DISCUSSION

The motion of the aurora follows plasma convection and large-scale instability processes in the magnetosphere [29, 41–43]. The same fields that are involved in those large-scale processes also power the (smaller-scale) wave-particle interactions which ultimately cause particles to precipitate into Earth’s atmosphere [44–46]. Since the local fields are, in turn, responsible for producing auroral plasma turbulence [21, 36], it is not surprising that the latter should ultimately follow the motion of magnetospheric source regions, and hence also the general ionospheric convection. To propose (not demonstrate beyond doubt) that the ICEBEAR bulk motions follow the ionospheric plasma convection, we present in Figure 4 a space-ground conjunction with one of the European Space Agency’s Swarm satellites. The Swarm mission consists of three low-Earth-orbit (450 km – 500 km altitude) with a high inclination (87°), each equipped with an electric field instrument [47, 48]. The latter computes ion drift velocities from an advanced imaging procedure.

Figure 4a–c) show three consecutive three-minute intervals during geomagnetic dawn on 5 September 2022. The auroral images (green contour lines) show a large, roughly north-south oriented diffuse auroral form, itself consisting of numerous flashing or pulsating auroral patches, that drift steadily in the eastward direction. The pulsating aurora has been shown statistically to follow the large-scale convection pattern, to the extent that it has been proposed as its proxy measurement [49]. At first sunlight, on 10:31:15 UT, the TREX RGB camera system was shut off to preserve the light-sensitive sensors. Naturally, ICEBEAR, being a radar, continued unaffected to observe the bulk motion of extant echo clusters, of which the majority were pointing in the direction of the diffuse auroral form. At around 10:36:30 UT Swarm B orbited through ICEBEAR’s field-of-view, with the cross-track *in-situ* ion velocity measurements from the F-region at an altitude 500 km exhibiting a steady eastward drift. A video of the event is provided in the Supporting Information, which clearly shows the inferences written here.

Figure 4d) compares ICEBEAR’s echo-weighted average bulk motion with Swarm B’s average cross-track ion drift for the interval 10:35 UT – 10:38 UT, showing excellent agreement in both direction and magnitude. This observation agrees well with the inferences made from Figure 3a–l), namely that the ICEBEAR echo clusters’ general motion and spatial distribution track the aurora. Next, we shall discuss the prominent and important exceptions to this rule.

The bottom row of Figure 3 presents a curious case where ICEBEAR’s echo cluster track ostensibly vary con-




FIG. 4. **Panels a–c):** An early-morning conjunction between ground-based optical imaging of the aurora (green contour lines, with green arrows denoting auroral optical flow), ICEBEAR echo cluster tracks (red lines), and Swarm B cross-track ion drift observations from the F-region (blue arrows). Each of the three panels display consecutive three-minute intervals from 10:29 UT to 10:38 UT on 5 September 2022, of concurrent observations plotted in an MLON-MLAT plane that take field lines curvature between the E- and F-regions into account. **Panel d)** shows the echo-weighted average ICEBEAR bulk motion (red arrow) as well as the average cross-track Swarm B ion drifts (blue) for the time interval 10:35 UT – 10:38 UT.

siderably in directions perpendicular to the general motion of the aurora. As we suggested in the previous section, if the local field enhancements poleward of the discrete arc in 3m-p) are strong enough, they will cancel out the ambient drift and cause the local $\mathbf{E} \times \mathbf{B}$ -direction to fluctuate. Despite this, the average direction over 51 seconds (the dashed blue lines in Figure 3m-p) is perfectly perpendicular to the discrete arc and pointing in the direction of the auroral flow vectors. In other words, in this case, the average bulk motion still uncovers a general drift direction consistent with the expectations for the large-scale electric field as observed in the pre-midnight auroral region.

Figure 5 shows an even more extreme example of the foregoing, where the average tracks point parallel to an intense, undulating auroral arc, with the astonishing bulk speed of 2200 m/s. Following the explanation laid out in the previous paragraphs, the local electric field around these unusually intense precipitation is strong enough to completely override the ambient drift, causing the average track direction to be parallel to the arc and the bulk motions to be extremely fast. In this case, the local electric field, in principle a linear superposition of that field and the ambient, is almost wholly produced and maintained by the energetic precipitating particles (1 keV – 10 keV [50]), which impart to the plasma charge upon their impact with the atmosphere, an action that ultimately produces a convergence electric field [38–40, 46]. In a letter, a companion to the present paper, we present conjunctions with magnetospheric satellites that elucidate this phenomenon.

Pitfalls Our proposed method of tracking turbulent structures is still in its infancy and has yet to be perfected. The resulting tracks and the bulk velocities they imply must, at present, be carefully assessed before conclusions are drawn about their physical significance. Critical tracking situations may occur when the clustering algorithm decides to drop or include echo regions, effectively causing bifurcation or merging events in the dataset. During such occurrences the median echo location may suddenly shift, causing a jump in the track. This jump will go on to form the track’s subsequent trajectory after merging or bifurcation, and as such is not necessarily unphysical, but it places upon the algorithm what may appear as an arbitrary means of deciding when clusters merge or bifurcate.

A second point of potential improvement lies in the possibility of clustering in higher dimensions. Here, time could be included as a dimension, removing the need for timestep-by-timestep matching of clusters to their preceding form. Doppler speed and signal-to-noise ratio, or even altitude, may also be added to the dimensions to cluster in. This can be done, for example, through dimensionality reduction by principle component analysis.




FIG. 5. **Panels a–c):** ICEBEAR echo tracks (solid red lines) and their linear polynomial fits (dashed red lines), superposed on auroral images, with green arrows denoting auroral optical flow. Each panel encompasses 20 seconds of data. Panel d) shows the echo-weighted average ICEBEAR bulk motion for the entire one-minute interval.

V. CONCLUDING REMARKS

We have applied industry-proven point-cloud analysis techniques to automatically mine large datasets of interferometry radar data. The method entails point-cloud clustering in two dimensions (in our case, the local plane




FIG. 6. 35 clusters of ICEBEAR 3D echoes identified on 25 August 2021. Each panel covers an area of $266 \text{ km} \times 266 \text{ km}$ and shows the cluster as it appeared at its most echo-numerous timestep. The clusters are superposed on the TREX RGB auroral images observed at that time. A red-yellow-white color scale indicates the absolute Doppler speed of the echoes.

perpendicular to Earth’s magnetic field lines). We track clusters of radar echoes through time by a simple similarity test from timestep-to-timestep. The resulting time histories of echo clusters trace out distinct spatial tracks, from which we derive a bulk motion. Through several case studies, as well as fortuitous conjunction studies, we show that this motion tends to follow the motion of auroral forms. Notable exceptions to this rule occur when especially intense auroral arcs cause a fast, peculiar motion parallel to those arcs. This phenomenon is explored further in a companion publication.

Our method, which is fully automatic, can be used to mine point-cloud data for irregular and dynamic clustering and flag this data for subsequent analyses. To illus-

trate the usefulness of our algorithm’s automatic nature, we have produced in Figure 6 a serial collection of 35 ICEBEAR echo clusters that were observed between 04:00 UT and 05:40 UT, each superposed on optical images of the aurora. The echo clusters are represented by snapshots taken at a prominent point during their evolution. For each of the 35 shapes, then, there readily exists a detailed frame-by-frame evolution akin to those presented in Figures 3.

With ICEBEAR’s new observations of the radar aurora bulk motions come new opportunities to study the magnetosphere-ionosphere coupling that lies at the heart of such motions, and to shed further light on the understudied E-region ionosphere.

ACKNOWLEDGEMENTS

This work is supported in part by Research Council of Norway (RCN) grant 324859. We acknowledge the support of the Canadian Space Agency (CSA) [20SUG-OICEB], the Canada Foundation for Innovation (CFI) John R. Evans Leaders Fund [32117], the Natural Science and Engineering Research Council (NSERC), the International Space Mission Training Program supported by the Collaborative Research and Training Experience (CREATE) [479771-2016], the Discovery grants program [RGPIN-2019-19135]; and the Digital Research Alliance of Canada [RRG-FT2109]. DRH was funded during this

study through a UiT The Arctic University of Norway contribution to the EISCAT_3D project funded by RCN grant 245683. MFI is grateful to M. Oppenheim, Y. Dimant, F. Lind, P. Erickson, and Y. Miyashita for stimulating discussions.

DATA AVAILABILITY

ICEBEAR 3D echo data for 2020, 2021 is published with DOI 10.5281/zenodo.7509022. TReX optical data can be accessed at <https://data.phys.ucalgary.ca/>. Data from the European Space Agency's Swarm mission can accessed at <https://swarm-diss.eo.esa.int/>.

-
- [1] D. Birant and A. Kut, ST-DBSCAN: An algorithm for clustering spatial-temporal data, *Data & Knowledge Engineering Intelligent Data Mining*, **60**, 208 (2007).
- [2] M. W. McDaniel, T. Nishihata, C. A. Brooks, P. Salesses, and K. Iagnemma, Terrain classification and identification of tree stems using ground-based LiDAR, *Journal of Field Robotics* **29**, 891 (2012),.
- [3] J. Li, H. Bao, X. Han, F. Pan, W. Pan, F. Zhang, and D. Wang, Real-time self-driving car navigation and obstacle avoidance using mobile 3D laser scanner and GNSS, *Multimedia Tools and Applications* **76**, 23017 (2017).
- [4] M. E. Hodgson, J. Jensen, G. Raber, J. Tullis, B. A. Davis, G. Thompson, and K. Schuckman, An Evaluation of Lidar-derived Elevation and Terrain Slope in Leaf-off Conditions, *Photogrammetric Engineering & Remote Sensing* **71**, 817 (2005).
- [5] J. Schindling and C. Gibbes, LiDAR as a tool for archaeological research: a case study, *Archaeological and Anthropological Sciences* **6**, 411 (2014).
- [6] A. Lozinsky, G. Hussey, K. McWilliams, D. Huyghebaert, and D. Galeschuk, ICEBEAR-3D: A Low Elevation Imaging Radar Using a Non-Uniform Coplanar Receiver Array for E Region Observations, *Radio Science* **57**, e2021RS007358 (2022),.
- [7] M. F. Ivarsen, J.-P. St-Maurice, G. C. Hussey, D. Galeschuk, A. Lozinsky, B. Pitzel, and K. A. McWilliams, An Algorithm to Separate Ionospheric Turbulence Radar Echoes From Those of Meteor Trails in Large Data Sets, *Journal of Geophysical Research: Space Physics* **128**, e2022JA031050 (2023),.
- [8] M. F. Ivarsen, A. Lozinsky, J.-P. St-Maurice, A. Spicher, D. Huyghebaert, G. C. Hussey, D. Galeschuk, B. Pitzel, and J. Vierinen, The Distribution of Small-Scale Irregularities in the E-Region, and Its Tendency to Match the Spectrum of Field-Aligned Current Structures in the F-Region, *Journal of Geophysical Research: Space Physics* **128**, e2022JA031233 (2023),.
- [9] B. Fejer and J. Providakes, High latitude E-region irregularities: New results, *Physica Scripta* **T18**, 167 (1987).
- [10] M. K. Ochi, *Ocean Waves: The Stochastic Approach*, Cambridge Ocean Technology Series (Cambridge University Press, Cambridge, 1998).
- [11] J. D. Huba, A. B. Hassam, I. B. Schwartz, and M. J. Keskinen, Ionospheric turbulence: Interchange instabilities and chaotic fluid behavior, *Geophysical Research Letters* **12**, 65 (1985),.
- [12] J. W. Dungey, Interplanetary Magnetic Field and the Auroral Zones, *Physical Review Letters* **6**, 47 (1961).
- [13] S. W. H. Cowley, TUTORIAL: Magnetosphere-Ionosphere Interactions: A Tutorial Review, Washington DC American Geophysical Union Geophysical Monograph Series **118**, 91 (2000).
- [14] A. Keiling, S. Thaller, J. Wygant, and J. Dornbeck, Assessing the global Alfvén wave power flow into and out of the auroral acceleration region during geomagnetic storms, *Science Advances* **5**, eaav8411 (2019), publisher: American Association for the Advancement of Science.
- [15] M. Wiltberger, V. Merkin, B. Zhang, F. Toffoletto, M. Oppenheim, W. Wang, J. G. Lyon, J. Liu, Y. Dimant, M. I. Sitnov, and G. K. Stephens, Effects of electrojet turbulence on a magnetosphere-ionosphere simulation of a geomagnetic storm, *Journal of Geophysical Research: Space Physics* **122**, 5008 (2017),.
- [16] J.-P. St-Maurice and L. Goodwin, Revisiting the Behavior of the E-Region Electron Temperature During Strong Electric Field Events at High Latitudes, *Journal of Geophysical Research: Space Physics* **126**, 2020JA028288 (2021),.
- [17] G. W. Prölss, Absorption and Dissipation of Solar Wind Energy, in *Physics of the Earth's Space Environment: An Introduction*, edited by G. W. Prölss (Springer, Berlin, Heidelberg, 2004) pp. 349–399.
- [18] D. L. Hysell, The Radar Aurora, in *Auroral Dynamics and Space Weather* (American Geophysical Union (AGU), 2015) pp. 191–209,.
- [19] O. Buneman, Excitation of Field Aligned Sound Waves by Electron Streams, *Physical Review Letters* **10**, 285 (1963), publisher: American Physical Society.
- [20] D. T. Farley, Incoherent Scatter Correlation Function Measurements, *Radio Science* **4**, 935 (1969),.
- [21] B. G. Fejer and M. C. Kelley, Ionospheric irregularities, *Reviews of Geophysics* **18**, 401 (1980),.
- [22] D. Huyghebaert, J.-P. St.-Maurice, K. McWilliams, G. Hussey, A. D. Howarth, P. Rutledge, and S. Erion, The Properties of ICEBEAR E-Region Coherent Radar Echoes in the Presence of Near Infrared Auroral Emis-

- sions, as Measured by the Swarm-E Fast Auroral Imager, *Journal of Geophysical Research: Space Physics* **126**, e2021JA029857 (2021),.
- [23] J. Drexler, J.-P. St. Maurice, D. Chen, and D. R. Moorcroft, New insights from a nonlocal generalization of the Farley-Buneman instability problem at high latitudes, *Annales Geophysicae* **20**, 2003 (2002), publisher: Copernicus GmbH.
- [24] M. F. Ivarsen, D. M. Gilles, D. R. Huyghebaert, J. P. St-Maurice, A. Lozinsky, D. Galeschuk, and G. Hussey, Turbulence Embedded into the Ionosphere by Electromagnetic Waves, Manuscript accepted at JGR: Space Physics (2024).
- [25] G. Hall, D. R. Moorcroft, L. L. Cogger, and D. Andre, Spatial relationship between large aspect angle VHF radio aurora and 557.7-nm emissions: Evidence for refraction, *Journal of Geophysical Research: Space Physics* **95**, 15281 (1990),.
- [26] H. Bahcivan, D. L. Hysell, D. Lummerzheim, M. F. Larsen, and R. F. Pfaff, Observations of colocated optical and radar aurora, *Journal of Geophysical Research: Space Physics* **111**, 10.1029/2006JA011923 (2006),.
- [27] D. L. Hysell, R. Miceli, J. Munk, D. Hampton, C. Heinselman, M. Nicolls, S. Powell, K. Lynch, and M. Lessard, Comparing VHF coherent scatter from the radar aurora with incoherent scatter and all-sky auroral imagery, *Journal of Geophysical Research: Space Physics* **117**, 10.1029/2012JA018010 (2012),.
- [28] J. E. Borovsky, J. Birn, M. M. Echim, S. Fujita, R. L. Lysak, D. J. Knudsen, O. Marghitu, A. Otto, T.-H. Watanabe, and T. Tanaka, Quiescent Discrete Auroral Arcs: A Review of Magnetospheric Generator Mechanisms, *Space Science Reviews* **216**, 1 (2019).
- [29] C. C. Chaston, K. Seki, T. Sakanoi, K. Asamura, and M. Hirahara, Motion of aurorae, *Geophysical Research Letters* **37**, 10.1029/2009GL042117 (2010),.
- [30] M. F. Ivarsen, J.-P. St-Maurice, D. R. Huyghebaert, D. M. Gilles, and G. C. Hussey, The Electric Field Derived from the Bulk Motion of the Radar Aurora, Submitted to JGR: Space Physics (2024).
- [31] M. F. Ivarsen, K. Song, D. Billett, J. P. St-Maurice, P. T. Jayachandran, and G. Hussey, Turbulent E-region Signatures of Poleward-moving Auroral Forms, Observed During the 23 April 2023 Geomagnetic Storm, Manuscript under preparation (2024).
- [32] D. Huyghebaert, G. Hussey, J. Vierinen, K. McWilliams, and J.-P. St-Maurice, ICEBEAR: An all-digital bistatic coded continuous-wave radar for studies of the E region of the ionosphere, *Radio Science* **54**, 349 (2019), conference Name: Radio Science.
- [33] M. Ester, H.-P. Kriegel, J. Sander, and X. Xu, A density-based algorithm for discovering clusters in large spatial databases with noise, in *kdd*, Vol. 96 (1996) pp. 226–231, issue: 34.
- [34] K. Khan, S. U. Rehman, K. Aziz, S. Fong, and S. Saravady, DBSCAN: Past, present and future, in *The Fifth International Conference on the Applications of Digital Information and Web Technologies (ICADIWT 2014)* (2014) pp. 232–238.
- [35] J. L. Barron, D. J. Fleet, and S. S. Beauchemin, Performance of optical flow techniques, *International Journal of Computer Vision* **12**, 43 (1994).
- [36] M. F. Ivarsen, D. R. Huyghebaert, D. M. Gilles, J. P. St-Maurice, D. R. Themens, M. M. Oppenheim, B. Gustavsson, D. Billett, B. Pitzel, E. Donovan, and G. C. Hussey, Turbulence around Auroras, Under revision at JGR: Space Physics (2024).
- [37] J. C. Foster and P. J. Erickson, Simultaneous observations of E-region coherent backscatter and electric field amplitude at F-region heights with the Millstone Hill UHF Radar, *Geophysical Research Letters* **27**, 3177 (2000),.
- [38] G. Marklund, Electric Fields and Plasma Processes in the Auroral Downward Current Region, Below, Within, and Above the Acceleration Region, *Space Science Reviews* **142**, 1 (2009).
- [39] K. Hosokawa, S. E. Milan, M. Lester, A. Kadokura, N. Sato, and G. Bjornsson, Large flow shears around auroral beads at substorm onset, *Geophysical Research Letters* **40**, 4987 (2013), publisher: Wiley Online Library.
- [40] B. Gallardo-Lacourt, Y. Nishimura, L. R. Lyons, J. M. Ruohoniemi, E. Donovan, V. Angelopoulos, K. A. McWilliams, and N. Nishitani, Ionospheric flow structures associated with auroral beading at substorm auroral onset, *Journal of Geophysical Research: Space Physics* **119**, 9150 (2014),.
- [41] V. M. Uritsky, J. Liang, E. Donovan, E. Spanswick, D. Knudsen, W. Liu, J. Bonnell, and K. H. Glassmeier, Longitudinally propagating arc wave in the pre-onset optical aurora, *Geophysical Research Letters* **36**, 10.1029/2009GL040777 (2009),.
- [42] K. Ogasawara, Y. Kasaba, Y. Nishimura, T. Hori, T. Takada, Y. Miyashita, V. Angelopoulos, S. B. Mende, and J. Bonnell, Azimuthal auroral expansion associated with fast flows in the near-Earth plasma sheet: Coordinated observations of the THEMIS all-sky imagers and multiple spacecraft, *Journal of Geophysical Research: Space Physics* **116**, 10.1029/2010JA016032 (2011),.
- [43] Y. Miyashita, T.-F. Chang, Y. Miyoshi, T. Hori, A. Kadokura, S. Kasahara, S.-Y. Wang, K. Keika, A. Matsuoka, Y. Tanaka, Y. Kasahara, M. Teramoto, C.-W. Jun, K. Asamura, Y. Kazama, S. W. Y. Tam, B.-J. Wang, S. Yokota, A. Kumamoto, F. Tsuchiya, M. Shoji, S. Kurita, S. Imajo, and I. Shinohara, Magnetic Field and Energetic Particle Flux Oscillations and High-Frequency Waves Deep in the Inner Magnetosphere During Substorm Dipolarization: ERG Observations, *Journal of Geophysical Research: Space Physics* **126**, e2020JA029095 (2021),.
- [44] S. Kasahara, Y. Miyoshi, S. Yokota, T. Mitani, Y. Kasahara, S. Matsuda, A. Kumamoto, A. Matsuoka, Y. Kazama, H. U. Frey, V. Angelopoulos, S. Kurita, K. Keika, K. Seki, and I. Shinohara, Pulsating aurora from electron scattering by chorus waves, *Nature* **554**, 337 (2018), number: 7692 Publisher: Nature Publishing Group.
- [45] K. Hosokawa, Y. Miyoshi, M. Ozaki, S.-I. Oyama, Y. Ogawa, S. Kurita, Y. Kasahara, Y. Kasaba, S. Yagitani, S. Matsuda, F. Tsuchiya, A. Kumamoto, R. Kataoka, K. Shiokawa, T. Raita, E. Turunen, T. Takashima, I. Shinohara, and R. Fujii, Multiple time-scale beats in aurora: precise orchestration via magnetospheric chorus waves, *Scientific Reports* **10**, 3380 (2020), number: 1 Publisher: Nature Publishing Group.
- [46] S. Imajo, Y. Miyoshi, Y. Kazama, K. Asamura, I. Shinohara, K. Shiokawa, Y. Kasahara, Y. Kasaba, A. Matsuoka, S.-Y. Wang, S. W. Y. Tam, T.-F. Chang, B.-J. Wang, V. Angelopoulos, C.-W. Jun, M. Shoji, S. Naka-

- mura, M. Kitahara, M. Teramoto, S. Kurita, and T. Hori, Active auroral arc powered by accelerated electrons from very high altitudes, *Scientific Reports* **11**, 1610 (2021), publisher: Nature Publishing Group.
- [47] E. Friis-Christensen, H. Lühr, and G. Hulot, Swarm: A constellation to study the Earth's magnetic field, *Earth, Planets and Space* **58**, BF03351933 (2006).
- [48] D. J. Knudsen, J. K. Burchill, S. C. Buchert, A. I. Eriksson, R. Gill, J.-E. Wahlund, L. Åhlen, M. Smith, and B. Moffat, Thermal ion imagers and Langmuir probes in the Swarm electric field instruments, *Journal of Geophysical Research: Space Physics* **122**, 2016JA022571 (2017).
- [49] B. Yang, E. Donovan, J. Liang, and E. Spanswick, A statistical study of the motion of pulsating aurora patches: using the THEMIS All-Sky Imager, *Annales Geophysicae* **35**, 217 (2017), publisher: Copernicus GmbH.
- [50] P. T. Newell, K. M. Lyons, and C.-I. Meng, A large survey of electron acceleration events, *Journal of Geophysical Research: Space Physics* **101**, 2599 (1996),.