

Bridging the micro-Hz gravitational wave gap via Doppler tracking with the Uranus Orbiter and Probe Mission: Massive black hole binaries, early universe signals and ultra-light dark matter

Lorenz Zwick^{1,*}, Deniz Soyuer², Daniel J. D’Orazio¹, David O’Neill¹, Andrea Derdzinski^{3,4}, Prasenjit Saha⁵, Diego Blas^{6,7}, Alexander C. Jenkins⁸, and Luke Zoltan Kelley⁹

¹*Niels Bohr International Academy, The Niels Bohr Institute, Blegdamsvej 17, DK-2100, Copenhagen, Denmark*

²*Department of Astrophysics, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland*

³*Department of Life and Physical Sciences, Fisk University, 1000 17th Avenue N., Nashville, TN 37208, USA*

⁴*Department of Physics & Astronomy, Vanderbilt University, 2301 Vanderbilt Place, Nashville, TN 37235, USA*

⁵*Physik-Institut, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland*

⁶*Institut de Física d’Altes Energies (IFAE), The Barcelona Institute of Science and Technology, Campus UAB, 08193 Bellaterra (Barcelona), Spain*

⁷*Institució Catalana de Recerca i Estudis Avançats (ICREA), Passeig Lluís Companys 23, 08010 Barcelona, Spain*

⁸*Department of Physics and Astronomy, University College London, London WC1E 6BT, UK*

⁹*Department of Astronomy, University of California, Berkeley, California 94720, USA*

(Dated: June 5, 2024)

With the recent announcement by NASA’s *Planetary Science and Astrobiology Decadal Survey 2023-2032*, a priority flagship mission to the planet Uranus is anticipated. Here, we explore the prospects of using the mission’s radio Doppler tracking equipment to detect gravitational waves (GWs) and other analogous signals related to dark matter (DM) over the duration of its interplanetary cruise. By employing a methodology to stack tracking data in combination with Monte-Carlo Markov-Chain parameter recovery tests, we show that the mission will be sensitive to GWs over the wide frequency range of 3×10^{-9} Hz to 10^{-1} Hz, provided that tracking data is taken consistently over a large fraction of the cruise duration. Thus, the mission has the potential to fill the gap between pulsar timing and space-based-interferometry GW observatories. Within this assumption, we forecast the detection of $\mathcal{O}(1 - 100)$ individual massive black hole binaries using two independent population models. Additionally, we determine the mission’s sensitivity to both astrophysical and primordial stochastic gravitational wave backgrounds, as well as its capacity to test, or even confirm via detection, ultralight DM models. In all these cases, the tracking of the spacecraft over its interplanetary cruise would enable coverage of unexplored regions of parameter space, where signals from new phenomena in our Universe may be lurking.

I. INTRODUCTION

Almost half a century after the launch of the *Voyager 2* space probe, the prospect of a new visit to the ice giant Uranus is finally crystallizing again: A flagship mission to the planet Uranus (hereafter referred to as “UOP”, standing for Uranus Orbiter and Probe) has been declared a priority, according to the announcement by NASA’s *Planetary Science and Astrobiology Decadal Survey 2023-2032*.¹ The announcement is timely; numerous publications have underlined the rich potential of such a mission in terms of planetary science over the past few years [42, 43, 49, 51, 54, 58, 59, 63, 93]. Yet, any mission to the outer Solar system must undergo a long cruise in interplanetary space before reaching its desti-

nation. According to a notional, short duration trajectory of the UOP (see Fig. 1), just the transfer between Jupiter and Uranus is expected to require around 9 years, during which the planetary science yield is virtually non-existent.

Several works have highlighted how the UOP’s long cruise may instead provide unique opportunities for *non-Uranian science*. Prospects include the possible measurement of the local dark matter content in the Solar system [89, 109], as well as the localisation of the hypothetical Planet 9 on the sky [24]. The opportunity to take such measurements would be afforded by the presence of a radio link between the spacecraft and the Earth, which would allow one to closely monitor changes in the spacecraft’s velocity and reconstruct deviations from the expected mission trajectory. Among all, the most timely application of interplanetary radio tracking is arguably the detection of gravitational wave (GW) signals. The idea of detecting GWs in the Doppler tracking data of planetary missions has a rich history and has been attempted previously with *Pioneer 11* [8], the *Galileo*–

* lorenz.zwick@nbi.ku.dk

¹ Survey provided in: [Origins, Worlds, and Life: A Decadal Strategy for Planetary Science and Astrobiology 2023-2032 \(2022\)](#)

Ulysses–Mars Observer coincidence experiment [5, 14], and *Cassini* [9, 13, 30], albeit without any successful detection candidates due to insufficient sensitivity. As first highlighted in Ref. [97], the prospective UOP offers an exceptional opportunity compared to the previous attempts due to a longer spacecraft cruise, a longer detector arm, and potentially improved radio technology.²

In this work, we consider how the collection of tracking data over the entire interplanetary cruise can increase the sensitivity of the mission to GWs from many sources, allowing us to probe the yet poorly covered micro-Hz regime. Additionally, we outline the mission’s capability to greatly expand the range of ultra-light dark matter (ULDM) models [21, 57] that can be directly detected, including the possibility of a first detection of DM in the Solar System. The key for these results is a method developed here to stack individual short-duration tracking runs, showing how the mission’s GW frequency sensitivity band and DM mass sensitivity are in principle only limited by the total observation time and the data cadence, in analogy to the Pulsar Timing Array analysis for nano-Hz GW detection [hereafter PTA, see, *e.g.*, 2, 6, 46, 53, 90]. For the UOP, this places the frequency limits to be between 3×10^{-9} Hz ($\sim 1/(10$ yr); the interplanetary cruise time) and 10^{-1} Hz ($1/(10$ s); the resolution time), respectively, and the DM mass limits in the range 10^{-23} eV/ c^2 to 10^{-15} eV/ c^2 .

In summary, we show that, beyond its Uranian science goals, the UOP could double as a sensitive GW detector in the micro-Hz band, probing the astrophysics of massive black holes or signals from the early Universe, and as a dark matter detector for models not accessible today by any other probe. This should be achievable without any sacrifice to planetary science goals by tracking the mission consistently over its interplanetary cruise. We also remark that the mission is expected to fly in concomitance with the Laser Interferometer Space Antenna (LISA [29]). The opportunities for complementarity between these two missions were briefly discussed in Ref. [97]. They deserve a more thorough analysis, which we leave for future work.

This paper is structured as follows: In Section II we introduce the basics of GW and ULDM detection via Doppler tracking and detail our strategy to track signals over the entire mission duration. In Section III, we construct GW sensitivity curves and define three possible mission configurations with increasing levels of technological development. In Section IV we estimate the number of detections of individual massive black hole binaries with two independent population models, as well as the associated stochastic gravitational wave background.

² Recently, there was another proposal to use tracking of satellites to Mars to study GWs of frequencies from 10^{-4} Hz to 10^{-1} Hz [15]. As compared to our proposal, these constraints are too weak to be relevant for known sources, while they will be largely superseded in the future by LISA [29].




FIG. 1. One of the many proposed mission plans of a prospective Uranus mission that involves a Jupiter Gravity Assist and a subsequent cruise to Uranus. This specific mission timeline corresponds to the shortest proposed cruise time and is available in the public [mission document](#).

Section V and Section VI include our forecast constraints on GW signals from the early universe, as well as various DM models, respectively. Lastly, we discuss the prospects to further improve the sensitivity of the mission in Section VII and present our concluding remarks in Section VIII.

II. DOPPLER TRACKING OF GWS AND OTHER SIGNALS

A. Response of Doppler tracking to a passing GW

The passage of a GW between the Earth and the spacecraft modifies the propagation time of photons, inducing small timing shifts in the tracking system. These shifts are best described by defining the dimensionless fractional frequency fluctuation of a two-way Doppler system:

$$y_2(t) = \Delta\nu/\nu_0, \quad (1)$$

where ν_0 is the tracking system’s carrier frequency, most often in the Ka-band (~ 30 GHz) or the X-band (~ 8 GHz), while $\Delta\nu$ is the measured Doppler shift with respect to the carrier frequency. Following Ref. [7], we can write down an explicit formula for the frequency fluctuation due to a passing GW:

$$y_2^{\text{GW}}(t) = \frac{\mu - 1}{2} \bar{\Psi}(t) - \mu \bar{\Psi} \left(t - \frac{\mu + 1}{2} T_2 \right) + \frac{\mu + 1}{2} \bar{\Psi}(t - T_2), \quad (2)$$

where T_2 is the two-way light travel time between the Earth and the spacecraft, $\mu = \hat{\mathbf{k}} \cdot \hat{\mathbf{n}}$ is the projection of the

unit wave vector $\hat{\mathbf{k}}$ of the GW onto the unit vector connecting the Earth and the spacecraft $\hat{\mathbf{n}}$, and $\bar{\Psi}$ is the projection of the GW amplitude onto the Doppler link [105]. The latter is given by $\bar{\Psi}(t) = (\hat{\mathbf{n}} \cdot \mathbf{h}(t) \cdot \hat{\mathbf{n}})/(1 - \mu^2)$, where $\mathbf{h}(t) = h_+(t) \mathbf{e}_+ + h_\times(t) \mathbf{e}_\times$ is the GW amplitude (*i.e.*, the strain) and $\mathbf{e}_{+, \times}$ are the usual ‘‘plus’’ and ‘‘cross’’ polarization states of a transverse, traceless plane GW [22, 82]. While Eq. (2) can be used to describe any incident GW, in this Section we will focus on coherent, monochromatic GWs with the form:

$$h_+(t) = A \frac{1 + \cos^2 \iota}{2} \cos(f_{\text{GW}} t + \phi_0), \quad (3a)$$

$$h_\times(t) = A \cos \iota \sin(f_{\text{GW}} t + \phi_0), \quad (3b)$$

where f_{GW} is the frequency of the GW, A its amplitude, ι is the binary’s inclination, and ϕ_0 is an initial phase [see, *e.g.*, 69, 72, 74]. In particular, Eqs. (3) can be used to describe signals produced by supermassive BH binaries in the early inspiral stages, the most promising astrophysical source of GWs in the nano-Hz to micro-Hz frequency band, and will be discussed in detail in Section IV.

B. Response of Doppler tracking to a SGWB

Following [9], the results from the previous section can be immediately used to derive a bound on stochastic GW backgrounds (SGWBs). Recall that many sources of GWs produce a signal that is comprised of several independent waves that interfere incoherently. These are the kind of signals expected from early Universe physics [25], and many astrophysical sources (see, *e.g.* [91] for examples in the micro-Hz band). SGWBs are modelled by considering the power spectrum of the sky- and polarization-average of the signal given in Eq. (2): $S_{y_2}(f)$. Following [37], this quantity can be expressed as $S_{y_2}(f) = \bar{R}_2(f) S_{\text{GW}}(f)$, where $S_{\text{GW}}(f)$ is the GW power spectrum, and the transfer function $\bar{R}_2(f)$ can be found in [9]. In the limit of low frequencies ($2\pi f T_2 \ll 1$), it reads simply $\bar{R}_2(f) \approx (8/15) (\pi f T_2)^2$, while at higher frequencies it displays some oscillatory behaviour. In order to investigate the sensitivity of Doppler tracking to SGWBs, it is customary to express the constraints in terms of the normalised energy density of the SGWB for logarithmic frequency unit:

$$\Omega_{\text{GW}}(f) \equiv \frac{8\pi G}{3H_0^2} \frac{d\rho_{\text{GW}}}{d \log f} = \frac{8\pi^2 f^3 S_{y_2}(f)}{3\bar{R}_2(f) H_0^2}, \quad (4)$$

where we will consider the Hubble parameter to be $H_0 \approx 67 \text{ km/s/Mpc}$ [4]. In practice, we will impose that $\Omega_{\text{GW}}(f) < \frac{8\pi^2 f^3 S_{y_2}(f)}{3\bar{R}_2(f) H_0^2}$, where $S_{y_2}(f)$ represents the power spectrum of the noise of the experiment.

C. Response of Doppler tracking to ultra-light dark matter

Dark matter (DM) particles of masses $m_{\text{DM}} \lesssim 1 \text{ eV}/c^2$ are distributed in the Milky Way at distances smaller than their de Broglie wavelength (associated with the ‘size’ of the particle). As a result, DM can be approximated in this range of masses as a classical non-relativistic massive field ϕ_{DM} [57]. From the virialised properties of DM in the Milky Way, this field will be distributed in coherent patches of characteristic length

$$l \approx 0.9 \times 10^6 \text{ AU} \left(\frac{10^{-3}}{\sigma_0} \right) \left(\frac{10^{-20} \text{ eV}/c^2}{m_{\text{DM}}} \right), \quad (5)$$

where σ_0 is the DM velocity dispersion, generated by virialization in the Milky Way. In these regions, the DM field can be approximated as a monochromatic function oscillating at a frequency

$$f_{\text{DM}} \approx 2.4 \times 10^{-6} \left(\frac{m_{\text{DM}}}{10^{-20} \text{ eV}/c^2} \right) \text{ Hz}, \quad (6)$$

and coherent over c^2/σ_0^2 oscillations. As a result, the gravitational potentials generated by the galactic DM (derived from solving the Poisson equation, which is quadratic in ϕ_{DM}) inherit the coherent features of the DM field, and oscillate with frequency $f_{\text{grav}} \approx 2f_{\text{DM}}$ in these patches (*cf.* Eq. (8) below). When approaching the lowest viable masses, these DM models are known as fuzzy dark matter [56] or ultra-light dark matter (ULDM) [57].

The Doppler tracking suggested above for GWs can be easily adapted to search for ULDM in different ways. First, notice that, at leading order, the fluctuations of the gravitational potential $g_{ij} \equiv -(1 - 2\psi)\delta_{ij}$ modify the relative frequency of the radio signal as [21, 62, 83, 101]

$$\Delta\nu/\nu_0 = \psi(t) - \psi(t - T_2). \quad (7)$$

For ULDM, the fluctuating piece of the gravitational potential ψ behaves (at leading order in $\sigma_0 \approx 10^{-3}$) as

$$\psi(t) \approx \psi_g(t) \equiv 6 \cdot 10^{-16} \left(\frac{\rho_{\text{DM}}}{0.3 \text{ GeV}/\text{cm}^3} \right) \left(\frac{10^{-23} \text{ eV}/c^2}{m_{\text{DM}}} \right)^2 \cos(4\pi f_{\text{DM}} t + 2\alpha_0), \quad (8)$$

where ρ_{DM} is the dark matter energy density and α_0 is a random phase.³ This immediately allows one to use the bounds on $y_2^{\text{GW}}(t)$ to constrain the mass of dark matter

³ Technically, the interference of the different classical waves that comprise the ULDM implies that the local density has a degree of randomness over this value, which corresponds to the average DM energy density [45]. We will assume that the DM energy density in the Solar System is 1/2 of the averaged one (expected to be of order $\rho_{\text{DM}} \approx 0.3 \text{ GeV}/\text{cm}^3$) to provide our estimates.

candidates (it is enough to use (7) in (1), and connect it to (2)). To find the precise bound, it is important to realise that from the lack of polarization and directional dependence, a factor of $\sqrt{15}$ appears when comparing the bounds on rms of $h(t)$ vs $\psi(t)$ [21].

Second, it is normally expected that DM is not completely ‘dark’, and couples, though very weakly, to matter or light. As a result, the fluctuations of ϕ_{DM} can affect the tracking data in new ways. For instance, new forces may affect the trajectory of gravitating probes or the properties of the tracking beam, see, *e.g.*, [1, 40, 47, 71, 77, 81]. For our current study, we will focus on two simple, though representative possibilities of *direct coupling* of DM to ordinary matter and light. In the first model, dark matter is coupled to the constituents of the detectors, the spacecraft, and the tracking beam through a *universal coupling*, common to all particles of the standard model of particle physics. This universality is similar to the coupling of the metric $g_{\mu\nu}$ in general relativity. As a consequence, in this model the coupling of matter and light to gravity ($g_{\mu\nu}$) and dark matter (ϕ_{DM}) happens through the interaction with an effective metric $\bar{g}_{\mu\nu} = g_{\mu\nu}(1 + 2\alpha(\phi_{\text{DM}}))$. In this situation, matter and light will move in geodesics of $\bar{g}_{\mu\nu}$, implying that the expression (7) will still be valid, identifying⁴ $\psi \equiv \psi_g(t) + \alpha(\phi_{\text{DM}})$. The function $\alpha(\phi)$ is model-dependent. To illustrate the potential of UOP to test DM models, we will choose $\alpha(\phi_{\text{DM}}) = \phi_{\text{DM}}^2/\Lambda_2^2$, where Λ_2 parameterises the intensity of the coupling. The effect of ULDM is described in this case through (7), after the identification

$$\psi(t) \approx \left(1 + 40 \left(\frac{10^{18}\text{GeV}}{\Lambda_2}\right)^2\right) \psi_g(t). \quad (9)$$

To illustrate other possibilities opened from the coupling of the tracking (light) beam to DM, we will consider a model where DM is an axion-light particle [28]. In this case, a coupling of the form

$$g_{a\gamma\gamma}\phi_{\text{DM}}\vec{E} \cdot \vec{B}, \quad (10)$$

is expected, where \vec{E} and \vec{B} are the electric and magnetic fields respectively, and $g_{a\gamma\gamma}$ represents the intensity of the coupling. Following [17], for the ULDM case one expects frequency oscillations in the tracking beam of order

$$\frac{\Delta\nu}{\nu_0} \sim 10^{-16} \left(\frac{g_{a\gamma\gamma}}{10^{-10}\text{GeV}^{-1}}\right) \left(\frac{\text{GHz}}{\nu/2\pi}\right) \sqrt{\frac{\rho_{\text{DM}}}{0.3\text{GeV}/\text{cm}^3}} \quad (11)$$

at frequency f_{DM} , where we have normalised $g_{a\gamma\gamma}$ to a value representative of current sensitivity [50, 80].

The previous three effects, summarised by Eqs. (8), (9) and (11) are far from being comprehensive (for instance,

we are not discussing the possibility of other possibilities for $\alpha(\phi_{\text{DM}})$, beyond the quadratic case of (9), or models where DM is a vector field, as in the case of dark photons). Still, they represent three families of ULDM models: universal gravitational coupling; direct coupling to matter; and direct coupling to light, which illustrate the potential of UOP to explore ULDM models.

D. Noise in a Doppler tracking system

The response of a Doppler tracking system to GWs and other oscillatory signals must be compared to its intrinsic noise. Noise sources range from mechanical (caused, *e.g.*, by flexibility in the antennae), to astrophysical (*e.g.*, scintillation due to propagation through the interplanetary plasma). These are thoroughly compiled and discussed in the extensive review by Ref. [7], and further discussed in Section VII. In general, the total noise profile can be described in either the time domain or the frequency domain. In the former, the size of typical noise fluctuations for every data-point is characterised by the system’s Allan deviation, σ_A , which is a function of the chosen averaging time τ . In the latter, noise is best described by its one-sided power spectrum S_n , which is a function of frequency f . These two crucial quantities are related by the following equation:

$$\sigma_A(\tau) = \sqrt{4 \int_0^\infty S_n(f) \frac{\sin(f\tau)^4}{(\pi f\tau)^2} df} \approx \sqrt{\frac{S_n(1/\tau)}{\tau}}, \quad (12)$$

where the latter approximation can be verified numerically, and is exact in the case of white noise [13]. Eq. (12) essentially states that the typical noise fluctuations given an averaging time τ are determined by the noise power spectral density at the reciprocal frequency $f \sim 1/\tau$.

The dependence of the Allan deviation on the averaging time τ is crucial when attempting to reconstruct a signal: On the one hand, a longer averaging time generally leads to reduced noise and improves the quality of each data point. On the other hand, the signal-to-noise ratio (SNR) also scales as the square root of the total amount of collected data points $N_s(\tau)$ which is inherently limited by the minimum sampling rate $1/\tau$ ($N \sim 1/\tau$). These competing effects define a sweet spot for the averaging time, which compromises between the quantity and quality of tracking data. It can be found by minimising the function:

$$\frac{\sigma_A(\tau)}{\sqrt{N_s(\tau)}} \sim \sqrt{S_n(1/\tau)}, \quad (13)$$

where we neglect an arbitrary normalisation given by the total observation time.

A simple parameterisation of a noise power spectral density is shown in Fig. 2, based on theoretical work by Ref. [13] and extensive analysis of the *Ulysses* and *Cassini* tracking data [14, 30]. The broad features of the

⁴ More specifically, at leading order in perturbations over the flat background, $\bar{g}_{ij} = -(1 - 2\psi - 2\alpha(\phi_{\text{DM}}))\delta_{ij}$.




FIG. 2. The blue curve shows the noise power spectral density based on theoretical work by Ref. [14] and extensive analysis of the *Ulysses* and *Cassini* tracking data [30]. The gray dashed curve is a visualisation of the trade-off between the total amount of data-points N_d and the quality of each data-point for the purposes of reconstructing the trajectory, where the absolute values for the latter curve have no significance. The curve is proportional to $\sqrt{fS_n} \times \sqrt{N_d}$, where $N_d \sim 1/\tau$ and τ is the averaging time of the Doppler link. The lowest point on the gray dashed curve therefore represents the sampling rate at which optimal SNR is achieved.

power spectral density can be captured by a simple three power law structure. At frequencies higher than $f \sim 0.1$ Hz, the noise is dominated by thermal fluctuations and steeply increases as $\sim f^2$. This limit tends to constrain the minimum averaging time of tracking systems to $\tau \sim 10$ s, beyond which the Allan deviation rapidly deteriorates. A central region of optimal sensitivity spans between $\sim 10^{-4}$ Hz and $\sim 10^{-1}$ Hz and is characterised by a red spectrum $\sim f^{-1/2}$ [30]. The upturn in noise below 10^{-4} Hz is an attempt to model less constrained low-frequency noise sources, which range from uncertainties in the dynamical modelling of the spacecraft orbits to the effects of atmospheric turbulence. They are conservatively estimated to generate a slope of f^{-2} , as has been originally observed in the *Ulysses* tracking data. However, it is important to note that these estimates are purely phenomenological. The effects of achieving a flatter low-frequency slope for the noise, and the resulting improvements in sensitivity, have been explored in [30]. Ultimately, the behaviour of the power spectral density below frequencies of $\sim 10^{-4}$ Hz cannot be studied straightforwardly owing to the rotation of the Earth: The spacecraft will fall below the horizon of any given tracking station within a fraction of a day, severing the Doppler link and preventing the continuous tracking of a coherent signal.

To circumvent this complication we consider a typical tracking run of a single station to have a maximal duration of approximately 8 hours, after which the link may only be reinstated after a clock reset. This amounts to either switching tracking stations or waiting for the next day. For the purposes of this work, we will con-

servatively assume that the noise of the *Cassini* Doppler tracking system is representative of modern interplanetary spacecraft within the relevant frequency range. We also discuss some more advanced technological prospects in Section VII B. With this choice, we are assuming that noise components at lower frequencies do not bleed into the sampled range of the power spectral density. Such components take the form of unmodelled accelerations due to small massive bodies and/or variations in the solar radiation pressure. Typically, they result in a subdominant contribution to the Allan deviation at 10 s of $\sim 10^{-16}$ [7].

As mentioned above, the noise power spectral density is specified by two crucial parameters, the optimal sampling rate and its corresponding Allan deviation. Given the *Cassini* mission specifications, the two parameters take the values:

$$\tau^{\text{Cass}} \sim 10 \text{ s}, \quad \sigma_A^{\text{Cass}}(10 \text{ s}) \sim 10^{-14}, \quad (14)$$

which are found by looking for the minimum of Eq. (13). For the purposes of this work, we fix the optimal sampling rate at 10 s, but allow the Allan deviation to vary slightly from its *Cassini*-era value in order to reflect technological advancements in the last four decades. The effects of these choices will be further discussed in Section VII.

E. Strategy for the tracking of low-frequency GWs

Measuring a low-frequency gravitational wave requires a long observation time, at least of the order $T \sim 1/f_{\text{GW}}$. Here we devise a strategy to integrate gravitational wave signals over the total mission duration of ~ 10 years. Previous attempts to estimate the sensitivity of Doppler tracking to such GWs have typically assumed that the noise power spectral density may be extended to lower frequencies following the phenomenological slope of $\sim f^{-2}$ [13]. Here we take a different approach, which consists of stacking many independent tracking runs in the time domain with a maximal duration of 8 hours, taken repeatedly in the entirety of the expected Uranus mission duration.

The first step for the stacking is to be able to generate noise realisations for single 8-hour tracking passes as a time series, while preserving the properties defined by the power spectral density. We construct the hypothetical noise time series y_2^{noise} of an entire 10 year tracking run by performing a discrete inverse Fourier transform of $\sqrt{fS_n}$, where we take the *Cassini* noise spectral density as shown in Fig. 2:

$$y_2^{\text{noise}}(t) = \frac{\sigma_A(10 \text{ s})}{\mathcal{N}} \times \sum_{f_i \in [f_{\min}, f_{\max}]} \sqrt{f_i S_n(f_i)} [\sin(f_i t + \alpha_i) + \cos(f_i t + \beta_i)]. \quad (15)$$

Here $f_{\min} = (10\text{yr})^{-1}$ is the inverse of the cruise time duration and $f_{\max} = \tau^{-1} = (10\text{s})^{-1}$ is the sampling rate. The phases α_i and β_i are uniformly sampled from $[0, 2\pi)$, resulting in an approximately Gaussian noise distribution. Additionally, the normalisation factor \mathcal{N} simply assures that the standard deviation of the noise takes the correct value $\sigma_A(10\text{s})$, regardless of the specific discretisation.

We can simulate tracking data from a virtual mission by averaging evaluations of Eq. (15) and adding it to the coherent Doppler response to a GW signal over the time segments during which the Doppler tracking is active. Thus, we model the mission's total tracking data as being composed of a sum of the GW response and the noise realisation, stacked over N consecutive tracking runs:

$$y_2^{\text{tot}} = \bigoplus_{n \leq N} (y_2^{\text{GW}}(t_n; t) + y_2^{\text{noise}}(t_n; t)), \quad (16)$$

where the tensor sum symbol describes the stacking of several, independent 8 h passes with arbitrary gaps between them. Here t_n corresponds to the timestamp at which the n -th tracking run is initiated, and $0 < t < 8$ hours follows a clock that is started anew at the beginning of every tracking run. A possible realisation of the virtual tracking data, where the gaps have been removed, is visualised in Fig. 3. We note that from here on, we will assume that the tracking runs are spaced regularly over the mission duration (see Section VII). While this is most likely not realistic, investigating the distribution of tracking runs is only useful once an official mission trajectory is determined. To summarise, in addition to the Allan deviation, each realisation of y_2^{tot} is thus determined by the parameters characterising both the GW and the spacecraft trajectory. Schematically, they are given by the following:

$$\begin{aligned} \text{GW} &\rightarrow \begin{cases} \hat{\mathbf{k}}; & \text{Propagation vector} \\ f_{\text{GW}}; & \text{Frequency} \\ A; & \text{Amplitude} \\ \phi_0; & \text{Phase} \end{cases} \\ \text{Trajectory} &\rightarrow \begin{cases} \hat{\mathbf{n}}(t_n; t); & \text{Link unit vector} \\ T_2(t_n; t); & \text{Two-way travel time} \end{cases} \end{aligned}$$

III. GW SENSITIVITY CURVES FOR THE PROSPECTIVE URANUS MISSION

A. Detection thresholds and null hypothesis

We shall estimate how well the GW signal can be extracted from the tracking noise. In a Bayesian statement of the problem, we are looking to evaluate the posterior probability of a GW with parameters $[\hat{\mathbf{k}}, f_{\text{GW}}, A, \phi_0]$ to be present in a given realisation of y_2^{tot} . Or in other words, we wish to evaluate and maximise a likelihood

function \mathcal{L} of the form:

$$\log(\mathcal{L}) \propto - \sum (y_2^{\text{tot}} - y_2^{\text{trial}})^2, \quad (17)$$

where y_2^{trial} is a trial GW response. To this end, we employ a Markov Chain-Monte Carlo (MCMC) framework, where we use the Monte Carlo sampling package `emcee` [44] running 32 parallel walkers in order to recover the posteriors of the GW parameters $[\hat{\mathbf{k}}, f_{\text{GW}}, A, \phi_0]$. Note that, since the trajectory parameters $[\hat{\mathbf{n}}(t_n; t), T_2(t_n; t)]$ only evolve very slowly, we can essentially treat them as constant over a single 8-hour tracking run. Therefore, we only have to specify the mission's trajectory at the timestamps t_n :

$$\hat{\mathbf{n}}(t_n; t) \rightarrow \hat{\mathbf{n}}(t_n), \quad (18)$$

$$T_2(t_n; t) \rightarrow T_2(t_n). \quad (19)$$

This simplification not only drastically expedites the likelihood evaluations, but also easily allows to efficiently repeat the analysis with different trajectories. This will become especially convenient once the official mission trajectory becomes available.

Our ultimate goal is to obtain a GW sensitivity curve for the mission which can be easily compared to the sensitivity curves of other GW observatories. A possible way to define such a curve is to attempt to maximise the likelihood of a tracking time series that does not contain any GW signal. The MCMC walkers will be able to freely sample regions in which the trial GW amplitude A^{trial} is well below the noise. However, they will be strongly discouraged to sample large values of A^{trial} , since they are inconsistent with the absence of a signal. Thus, the posterior probability for the GW amplitude A will sharply decrease above a certain threshold A_s :

$$\mathcal{P}(A) \sim \begin{cases} \text{const.} & \text{for } A < A_s \left(\hat{\mathbf{k}}, f_{\text{GW}}, \phi_0 \right), \\ 0 & \text{for } A > A_s \left(\hat{\mathbf{k}}, f_{\text{GW}}, \phi_0 \right), \end{cases} \quad (20)$$

After marginalising over the initial phase ϕ_0 and wave vector $\hat{\mathbf{k}}$, we can specify the threshold to the desired confidence level and obtain a curve $A_s(f_{\text{GW}})$, purely as a function of GW frequency. An example of this procedure is visualised in the left panel of Fig. 4, where one can clearly see a transition between allowed and disallowed regions of parameter space in the 2-dimensional posteriors of the GW frequency and amplitude.

As seen in the bottom panel of Fig. 5, the boundary between the allowed and disallowed regions is closely approximated by a half-Gaussian. This allows us to easily find the 1σ confidence threshold, above which signals are distinguishable from the null hypothesis. Note that this does not necessarily imply that the individual signal parameters are well constrained, only that the overall signal is distinguishable from its absence. This corresponds to the analogous definition of a sensitivity curve for other




FIG. 3. The n^{th} (with $n = \{1, \dots, 9\}$) two-way frequency fluctuation $y_{2,n}$ due to the total signal (black lines), due to the noise (salmon lines) and due to the GW signal (dashed lines) for 9 different 8-hour measurements conducted with a 1-year separation. The 9 equally separated sections between two-way light travel times of 6000s and 20000s, represent roughly Jupiter's and Uranus' semi-major axes, respectively. The horizontal axis has the units of the resolution time (10s), such that a single 8-hour observation lasts $8 \times 60 \times 60\text{s}/(10\text{s}) = 2880$ units. The observations are then glued together in the time axis, hence the discontinuities in the waveform of the GW. The change in the shape of the GW signal is due to the variation of the spacecraft-Earth vector with respect to the GW source as the spacecraft cruises interplanetary space. The mock amplitude of the GW ($A = 10^{-15}$) is exaggerated to guide the eye (see Eqs. (3)).




FIG. 4. We show the two-dimensional marginalised posterior distributions in amplitude and frequency for a set of three different injected GW amplitudes, corresponding to an SNR of 0, 3, and 12 from left to right. Here the SNR is calculated according to Eq. (24) with respect to the threshold amplitude shown in Fig. 5. As we can see, increasing SNR produces better parameter estimation in both frequency and amplitude of the GW, where sources with $SNR=3$ can already be constrained. The upper bound on the GW amplitude suffers from a degeneracy with the source's position in the sky (not shown) and therefore stops improving significantly after an $SNR \sim 8$ for this particular mission trajectory.

GW detectors (see, *e.g.*, [87]). For the purposes of defining a simple sensitivity curve, we fit the one sigma boundary with a broken power law, shown in the top panel of Fig. 5.

The value of the 1σ threshold A_s depends on the noise power spectral density of the tracking runs, the specified mission trajectory, as well as the total number of tracking runs N_t . By performing several numerical MCMC tests, we find, as one would expect, that its overall normalisation scales linearly with the Allan deviation and scales as

the square root of the total number of samples:

$$A_s \sim \sigma_A(\tau) \times \frac{\sqrt{\tau}}{\sqrt{N_t}}. \quad (21)$$

The effect of the spacecraft's trajectory is also folded into the specific realisation of A_s , as it modulates the GW response function y_2^{GW} over the 10-year observation window. In particular the relative orientation of the wave vector \hat{k} and the link vector \hat{n} can change significantly for elliptical trajectories. The presence of curvature in




FIG. 5. In the top panel we show the two-dimensional marginalised posterior samples in amplitude and frequency for a run with no injected signal. We highlight the threshold at which the absence of a signal is excluded at 1σ confidence, which is well fit by a broken power law. In the lower panel, we showcase the sample count in 50 frequency bins, as a function of amplitude. We normalise the latter with the 1σ threshold.

the trajectory is crucial, as it allows one to break the sky localisation degeneracy that would be present for a single, straight detector arm.

B. Defining the GW sensitivity curve

To reiterate, the recovered 1σ threshold A_s , which corresponds to the definition of a sensitivity curve, results from distributing N_t tracking runs over the entire mission duration of $T = 10$ years. For a monochromatic signal, this results in tracking a total of $n \sim (f_{\text{GW}} \times N_t \times 8 \text{ hour})$ cycles. This is reminiscent of matched filtering techniques, which enable one to compare the sensitivity curves of GW observatories to the characteristic strain of a signal, rather than its instantaneous amplitude [see, e.g., 87]. Thus, we adjust the 1σ detection threshold by heuristically factoring out the number of detected cycles:

$$h_{\text{sens}}(f_{\text{GW}}) = A_s(f_{\text{GW}}) \times (f_{\text{GW}} \times N_t \times 8 \text{ hr})^{1/2}, \quad (22)$$

where we define $h_{\text{sens}}(f_{\text{GW}})$ the GW sensitivity curve of the mission, given a trajectory, an Allan deviation and a set of tracking day timestamps t_n :

$$h_{\text{sens}}(f_{\text{GW}}) = h_{\text{sens}}(f_{\text{GW}}; \sigma_A(10 \text{ s}), t_n). \quad (23)$$

Then, we can also define the SNR of a GW detected with Doppler tracking in complete analogy with other gravitational wave detectors:

$$\text{SNR} = \sqrt{4 \times \int \frac{h_c(f')^2}{h_{\text{sens}}(f')^2 f'} df'}, \quad (24)$$

where h_c is the GW's characteristic strain, and the factor 4 comes from the normalisation of the one-sided-power spectrum. Here, an SNR of 1 corresponds to a barely detectable GW that can be distinguished from the absence of signal with 1σ confidence. As shown in Fig. 4, higher SNRs result in both better parameter estimation and in reduced false alarm rates, also in complete analogy with GW detectors. We find that the frequency and amplitude of sources with an $\text{SNR} \sim 3$ can be recovered within ~ 0.02 and ~ 0.5 dex, respectively. The bounds improve by a factor ~ 3 for $\text{SNR} = 12$ sources, though a residual degeneracy between the source's amplitude and optimal localisation on the sky becomes manifest, inducing a skew in the recovered posterior distribution.

C. Baseline, Priority and Optimistic scenarios

The sensitivity curves constructed above span the interesting frequency range of few $\sim 10^{-9}$ Hz to $\sim 10^{-5}$ Hz with a slope of approximately $f^{-1/3}$, before a turning point at $f \sim 1/(8 \text{ hours})$. They then rise with a slope of $\sim f^{1/3}$ until it is truncated by the Nyquist frequency at $\sim 10^{-1}$ Hz. Crucially, strain sensitivities of the order $\sim 10^{-15}$ in the nano-Hz to milli-Hz band can be achieved without requiring significant improvements in the Allan deviation, *provided that tracking data is accumulated over a substantial amount of the entire mission duration*. This is in stark contrast with previous analyses, which did not include the stacking procedure detailed in Section II E. We note that this amount of tracking is not typical for deep interplanetary missions. Here, however, we argue that it should be prioritised in parallel with Allan deviation improvements in order to achieve the scientific objectives detailed in this manuscript. Assuming this level of commitment to the tracking, we define three possible future scenarios different technological improvements are implemented with respect to Cassini era ranging.

In the *baseline* scenario, we do not assume any improvement with respect to *Cassini*-era tracking technology. We assume the tracking of the spacecraft for 8 hours each day over the mission's 10-year interplanetary cruise. The Allan deviation of this scenario is then $\sigma_A(10 \text{ s}) = 10^{-14}$, at a sampling rate of 10 seconds. The total number of 8-hour \times tracking runs is approximately 3650.

In the *priority* scenario, we assume that optimising currently available tracking technology becomes one of

the explicit targets of the mission before launch. We assume that the Allan deviation at 10 seconds is reduced to the optimal *Cassini*-era value of 3×10^{-15} , which was achieved with a longer averaging time of $\tau = 10^4$ s.

In the *optimistic* scenario, we assume that technological improvements in tracking technology (see, *e.g.*, Sec. VII) will have acquired an appropriate readiness level for an interplanetary mission. As a proxy for such technological improvements, we assume an Allan deviation at 10 seconds of 10^{-15} . Additionally, we assume that several tracking stations distributed across the globe be utilised, assuring 24 hours of tracking per day. We discuss the prospects to achieve such low levels of noise in Section VII B. Here we simply note that by the time the UOP is operational, almost half a century of technological improvement will have taken place since the launch of the *Cassini* mission.

IV. DETECTION FORECAST FOR BLACK HOLE BINARIES

Supermassive black hole binaries (SMBHBs) are among the loudest sources in the low-frequency GW sky. They are expected as a consequence of galaxy mergers, although how frequently they form and how they evolve in their environments is still not fully understood [see, *e.g.*, the seminal papers 12, 55, 64, 99, 104]. Recent constraints on a GW background signal by pulsar timing arrays (PTA) provide evidence of an SMBHB population in the mass range $\sim 10^8 - 10^{10} M_\odot$ [2, 6, 85, 108] while the direct detections of lighter SMBHB mergers below a few $10^7 M_\odot$ is expected in the 2030s with the advent of space-based interferometers such as LISA [29] and TianQin [68]. As we will show, the data stacking method in Section III facilitates the detection of SMBHBs that are evolving beyond the nano-Hz band and into the milli-Hz regime. As an illustration, we show the hypothetical sensitivity of the mission to chirping, equal-mass BH binaries as a waterfall plot in the bottom panel of Fig. 6. The mission's peak sensitivity lies in the total mass range of 10^8 to $10^{10} M_\odot$. Such heavy, chirping binaries would in principle be detectable up to redshifts of $z \sim 10$ for an Allan deviation of $\sigma_A = 10^{-14}$ and essentially over the whole cosmological volume for $\sigma_A = 3 \times 10^{-15}$.

However, a detection forecast requires us to consider the actual population of sources we expect to fall in this frequency band, and whether the majority will be detectable as a monochromatic signal rather than a chirp. Here we adopt two separate approaches to estimate the redshift-dependent merger rate of such sources.




FIG. 6. In the top panel, we compare the GW sensitivity of the prospective Uranus mission to the current sensitivity of PTAs [36] and the sensitivity of the recently adopted LISA mission. The three blue lines correspond to the scenarios detailed in Section III. In the bottom panel, we show a waterfall plot of the mission's hypothetical detection horizon for *chirping* equal mass ratio massive black hole binaries, assuming a maximum of 10 years for the signal duration and the *priority* scenario, as described in Section III C. We compare it to the same plot for PTAs and for LISA (assuming 4 years of maximal signal duration). Note that monochromatic sources, expected for PTAs and from our population models in §IV, will have lower SNR for the same mass and redshift in the bottom panel.

A. Model 1: Millennium simulation

Here we detail a simple prescription that links the massive black hole merger rate to the much more established halo merger rate, based on the two Millennium simulations [38]. The latter works provide a convenient fit characterising the differential halo merger rate:

$$\frac{d^2\Gamma}{d\xi dz} = B_1 \left(\frac{M_{\text{halo}}}{10^{12} M_\odot} \right)^{b_1} \xi^{b_2} \exp \left[\left(\frac{\xi}{B_2} \right)^{b_3} \right] (1+z)^{b_4}, \quad (25)$$

where Γ is the total number of mergers that a halo of mass M_{halo} experiences over cosmic time, $\xi \leq 1$ is

the halo merger mass ratio and the best-fit parameters are given by $(B_1, B_2, b_1, b_2, b_3, b_4) = (0.0104, 9.72 \times 9.72, 0.133, -1.995, 0.263, 0.0993)$. We link Eq. (25) to the SMBH merger rate $\dot{N}_{\bullet\bullet}$ by multiplying the halo merger rate with the black hole mass function:

$$\frac{d^3 \dot{N}_{\bullet\bullet}}{dM_{\bullet} d\xi dz} = P_{\text{occ}}(M_{\text{halo}}, z) \frac{4\pi c D_{\text{com}}^2(z)}{(1+z)^3} \times \frac{dn_{\bullet}}{dM_{\bullet}} \frac{d^2 \Gamma}{d\xi dz}(\xi, z_{\text{del}}), \quad (26)$$

where D_{com} is the comoving distance at redshift z and we must additionally supply an occupation fraction P_{occ} of black holes in halos and a delay prescription z_{del} between the nominal halo merger time and the actual black hole merger time. Here we use the SMBH mass function $dn_{\bullet}/dM_{\bullet}$ as reported in [92] and adopt a simple relation between the halo and SMBH mass from [31]:

$$M_{\bullet} = \left[\frac{M_{\text{halo}}(1+z)}{2 \times 10^7 M_{\odot}} \right]^{3/2} M_{\odot}, \quad (27)$$

which is consistent with both simulations and observations of massive galaxies [10, 70].

We can now establish the number of detectable sources of gravitational radiation by integrating the differential SMBH merger rate. We define an SNR threshold C and count all SMBH binaries that exceed it:

$$N_{\bullet\bullet}^{\text{det}} = \int \int \int \int \frac{d^3 \dot{N}_{\bullet\bullet}}{dM_{\bullet} d\xi dz} \frac{1}{\dot{f}_{\text{GW}}} \Theta(\text{SNR} - C) \times df_{\text{GW}} d\xi dM_{\bullet} dz, \quad (28)$$

where Θ is a Heaviside function, and we distribute all binaries in frequency bins according to their residence times \dot{f}_{GW}^{-1} . Here it is necessary to make two additional simplifications. Firstly, we model the occupation fraction P_{occ} as an unspecified constant. For massive black holes $M > 10^7 M_{\odot}$, simulations [86] and observations of AGN [48] show that the occupation fraction P_{occ} is a fraction of order unity. Secondly, we also neglect the effect of time delays between the halo merger and the BH merger, as they typically only amount to 10^8 – 10^9 years for the mass range we are considering [35, 103]. This simplification is justified a posteriori, given that tracking will only ever be sensitive to binaries at $z \lesssim 1$ for realistic populations,⁵ where a delay of 10^9 years at most corresponds to a small redshift shift $\Delta z_{\text{del}} \sim 0.3$.

B. Model 2: Holodeck

We present another estimate of the detectable SMBHB population via the `holodeck` code [61, and see Ref. [3]].

Importantly, we utilise `holodeck` to build population models that generate nano-Hz-frequency GWBs with an amplitude that is consistent with the GWB for which multiple PTA experiments have recently found strong evidence [2, 6, 85, 108].

We utilise `holodeck`'s semi-analytical SMBHB population models. Specifically, these models combine an observationally constrained galaxy mass function from Ref. [66], with a galaxy merger rate from Ref. [88] to generate galaxy mergers. An SMBH-mass versus bulge-mass relation from Ref. [64] is then used to calculate SMBHB parameters, which are then self-consistently evolved from galaxy-scales to nano-Hz separations using a phenomenological binary evolution model developed for `holodeck` that combines binary decay due to environmental interactions and GWs [3]. The change in source redshift over the course of binary evolution is naturally included. Each of these model components depends on a number of uncertain parameters with measurements and uncertainties described in the cited studies.

To sample SMBHBs that are consistent with PTA observations, we generate 1080 `holodeck` populations by sampling over the uncertainties in each model parameter⁶. We then calculate the resulting PTA-measured GWB spectrum, fit a power-law spectrum to the lowest five frequency bins, and select the populations with amplitudes and spectral indices consistent with recent measurements. In particular, we require the GWB amplitude (characteristic strain at a frequency of 1 yr^{-1}) that is $A_{\text{yr}} \in [10^{-15}, 10^{-14}]$, and a power-law index (in power-spectral-density of timing residuals) that is $\gamma \in [-5.0, -2.5]$ [see, *e.g.*, 100]. This procedure is repeated for each UOP sensitivity scenario and threshold SNR, drawing independent populations for each combination. Out of the 1080 populations, 130 ± 10 pass the above selection cuts. Those populations are then integrated over with a cut in detection SNR similarly to Eq. (28) to determine detectable SMBHB systems.

C. Detection forecast

The results of our detection forecast are summarised in Table I and Fig. 7. Regarding the former, we compute the number of expected detections of individual binary SMBHBs at three separate SNR thresholds (1, 3, 8), for the three mission scenarios of baseline, priority, and optimistic (§III C). For the `holodeck` results we quote the median, 25%, and 75% quantiles derived from the ~ 130 GWB-selected populations described above. For the Millennium results, we quote expectation values in terms of the occupation fraction of BHs in halos. The expected number of detections is broadly consistent for the two

⁵ The seeming discrepancy with the detection horizon in Fig. 6 derives from the detectability of monochromatic sources vs. the hypothetical chirping sources drawn for consistent comparison between bands in Fig. 6.

⁶ 29 parameters are marginalised over in this way, sampling from the `PS_Astro_AStrong_AA11-v1.0` parameter space.

adopted population models; the Millennium expectation values fall within the inner 50% range of `holodeck` values for $P_{\text{occ}} \gtrsim 0.4$. Even in the baseline scenario, which does not assume any technological improvement with respect to *Cassini*-era technology, we would typically expect the detection of $\mathcal{O}(1)$ binaries at $\text{SNR} = 3$. A mild improvement in the Allan deviation, *e.g.*, to 3×10^{-15} , boosts the number of detections to $\mathcal{O}(10)$. Overall, the results shown in Table I suggest that a mild improvement in Allan deviation over the baseline mission would guarantee the detection of several well-constrained monochromatic signals from SMBH binaries, in a mass range that is yet to be explored. Indeed, as seen in Fig. 7, the majority of detected sources would lie around a mass of $10^9 M_\odot$, being detectable out to redshifts of $z \lesssim 1$. The majority of these are detected as almost monochromatic sources at frequencies of tens to hundreds of nHz, *i.e.*, above the PTA band (see bottom panel of Fig. 7). A more sensitive mission can pick up more monochromatic signals at lower frequencies, where a larger number of binaries reside.

The main difference between the Millennium and the `holodeck` population models lies in the redshift distribution of sources, as can be seen in the middle panel of Fig. 7. As seen in Fig. 8, we also note that in the priority scenario, the UOP will be sensitive to the stochastic background of SMBH binaries over the entire frequency range of 10^{-8} Hz to 10^{-6} , provided that the latter follows the scaling of $f^{-2/3}$ that is reported by, *e.g.*, Ref. [2].

As a final comment on the topic of SMBHBs, we briefly mention that tracking the UOP would provide stringent bounds on the existence of an intermediate-mass companion to the main SMBH of the Milky Way, *i.e.*, SgrA*. Current constraints arise from a range of techniques, see, *e.g.*, [78, 106]. Quite relevant for this work are the models where the GWs emitted fall in the micro-Hz band. In [78], it was shown that several of these models generate GWs with strains at the Solar system of up to 10^{-12} . From Fig. 6, it can be easily extrapolated that all hidden companions that would produce strains of order $\sim 10^{-15}$ over the frequency range of 10^{-8} Hz to 10^{-5} Hz would be detectable.

Combining all of these prospects, we have demonstrated that Doppler tracking with the UOP is uniquely suited to detect GWs in the frequency regime between PTAs and space-based interferometry. Sources in this frequency range comprise heavy SMBH binaries of several $10^9 M_\odot$ approaching coalescence, inspiraling pre-LISA systems and intermediate-mass black holes in the Milky Way. Such detections would unveil a more complete picture of SMBH formation, growth, and interaction, complementing both existing and planned GW detectors in the low-frequency regime. In the event of a concurrent detection with PTAs or LISA, a UOP detection may also aid in the difficult task of localising SMBHB sources [*e.g.*, 97].




FIG. 7. The differential distribution of detections (*i.e.*, the probability density function) shown with the dashed curves and the cumulative distribution function with the solid curves, for both the Millennium and `holodeck` (here chosen to be the median percentile realisation) population models, normalised with their respective maximum values. The top panel shows the distribution in mass decades $dN/d\log_{10}(M)$, while the middle panel shows the distribution in redshift dN/dz . The curves are displayed for the optimistic scenario with an SNR threshold of 3, but apply with little variation for all scenarios and SNR thresholds. The lowest panel shows the distribution of sources in $dN/d\log_{10}(f_{\text{GW}})$, showing how a more sensitive mission is capable of detecting more monochromatic signals at lower frequencies, shifting the PDFs towards the left. The highest SNR sources are therefore always detected at ~ 100 nHz, above the PTA sensitivity band. Note that all the displayed curves are normalised.

Scenario 1: Baseline			Scenario 2: Priority			Scenario 3: Optimistic		
Pop.	SNR	Nr. Det	Pop.	SNR	Nr. Det	Pop.	SNR	Nr. Det
Millennium	1	$12 \times P_{\text{occ}}$	Millennium	1	$71 \times P_{\text{occ}}$	Millennium	1	$1049 \times P_{\text{occ}}$
Millennium	3	$2 \times P_{\text{occ}}$	Millennium	3	$12 \times P_{\text{occ}}$	Millennium	3	$197 \times P_{\text{occ}}$
Millennium	8	0	Millennium	8	$2 \times P_{\text{occ}}$	Millennium	8	$42 \times P_{\text{occ}}$
holodeck	1	$6.8^{+20.0}_{-2.0}$	holodeck	1	$58.2^{+193.1}_{-21.4}$	holodeck	1	1375^{+2582}_{-584}
holodeck	3	$1.0^{+1.8}_{-0.3}$	holodeck	3	$6.5^{+14.8}_{-1.7}$	holodeck	3	190^{+497}_{-68}
holodeck	8	$0.15^{+0.28}_{-0.06}$	holodeck	8	$1.2^{+3.1}_{-0.4}$	holodeck	8	$30.1^{+67.2}_{-10.6}$

TABLE I. Three tables summarising the expected number of detections of SMBHBs for the UOP scenarios detailed in Section III C, over the entire mission duration. The detection estimates are performed with the two population models explained in Section IV and reported for three different SNR thresholds (see Fig. 4 for a visualisation of the typical parameter posteriors given these SNRs). The Millennium results scale with the occupation fraction of SMBHBs in massive galaxies, P_{occ} . The holodeck numbers are quoted as median \pm 25% and 75% quantile values.

V. FORECAST CONSTRAINTS ON EARLY UNIVERSE SIGNALS

As well as filling the gap in our sensitivity to SMBHBs, our proposed GW searches are also sensitive to potential cosmological signals from the early Universe. There is a huge range of mechanisms for generating GWs at early times [25], each of which, if detected, would give valuable new insights into fundamental physics. In the context of Doppler tracking with the UOP however, the most interesting prospects are for signals that are peaked in a relatively small range of frequencies, which risk going undetected by other GW experiments if there is not sufficient coverage of the micro-Hz frequency band.

The quintessential example of such a peaked cosmological GW signal is that generated by a first-order phase transition (FOPT) in the early Universe. These transitions, in which a fundamental quantum field escapes from a metastable state by nucleating ‘bubbles’ of a new phase, are a generic prediction of many extensions to the Standard Model of particle physics [25–27], notably beyond the reach of even the most powerful particle accelerators on Earth. Collisions between nucleated bubbles, and the subsequent acoustic and turbulent motion of the thermal plasma, can produce strong SGWB signals that peak at a frequency set by the energy scale of the transition, with higher frequencies corresponding to higher energies and thus earlier cosmic epochs [60]. This peak frequency is commonly approximated as

$$f_* \approx 19 \mu\text{Hz} \times \frac{k_B T_*}{100 \text{ GeV}} \frac{\beta/H_*}{v_w/c} \left(\frac{g_*}{106.75} \right)^{1/6}, \quad (29)$$

where T_* is the temperature of the thermal plasma at the time when the GWs are generated, β is the duration of the transition, measured in units of the Hubble rate H_* at that epoch, v_w is the terminal expansion velocity of the bubble walls, and g_* is the number of relativistic degrees of freedom in the plasma, normalised here to the Standard Model value. Focusing for simplicity on the contribution from sound waves in the plasma [52], which is dominant in many scenarios, the SGWB spectrum can

be modelled as a broken power law,

$$\Omega_{\text{GW}}(f) \approx \Omega_{\text{GW}}(f_*) \times (f/f_*)^3 \left[\frac{7}{4 + 3(f/f_*)^2} \right]^{7/2}, \quad (30)$$

where the peak intensity is

$$\Omega_{\text{GW}}(f_*) \approx 5.7 \times 10^{-6} \times \frac{v_w/c}{\beta/H_*} \left(\frac{\kappa\alpha}{1+\alpha} \right)^2 \left(\frac{g_*}{106.75} \right)^{-1/3} \times \left[1 - (1 + 2\tau_{\text{sw}} H_*)^{-1/2} \right], \quad (31)$$

with κ an efficiency parameter and τ_{sw} the lifetime of the sound-wave source, both of which can be modelled as functions of the parameters α , β , and v_w [26, 27].

Following [18, 19], we carry out a scan over the 4D parameter space $(T_*, \alpha, \beta, v_w)$ to assess the sensitivity of our proposed searches to FOPT signals. As shown in Fig. 9, there is a broad swathe of parameter space for these models that is accessible with Doppler tracking searches in the micro-Hz band that would go undetected by pulsar timing arrays in the nano-Hz band and LISA in the milli-Hz band. These constraints are complementary to those that have been forecasted for ‘binary resonance’ searches for micro-Hz GWs using Lunar laser ranging data [18, 19], allowing us to potentially detect or rule out a much larger family of FOPT models.

VI. FORECAST ON ULDM CONSTRAINTS.

The expected constraints on h_c from the top panel of Fig. 6 already give an idea of the reach of UOP to detect ULDM (compare Eqs. (1) and (7)). This suggests that for masses $m_{\text{DM}} \gtrsim 10^{-22} \text{ eV}/c^2$, UOP may represent the most sensitive probe for large regions in parameter space of dark matter models, even in the baseline scenario. Notably, this mass range is particularly relevant for ULDM phenomenology [41, 57]. To illustrate this reach, let us consider the equivalent h_c that would be produced by the three families of models we introduced in Section II C,




FIG. 8. Forecast sensitivity to the SGWB spectrum $\Omega_{\text{GW}}(f)$ for our baseline, priority, and optimistic scenarios (navy solid, dashed, and dotted curves), along with existing sensitivities of the NANOGrav PTA (green) [3] and Doppler tracking with *Cassini* (blue), as well as forecast sensitivities of LISA (brown) [87] and binary resonance searches with lunar laser ranging data (LR, light blue). The black horizontal line (N_{eff} constraint) shows the indirect constraint on the *integrated* SGWB intensity from measurements of the Hubble rate during radiation domination. The pink shaded region indicates the astrophysical SGWB from unresolved SMBHBs (with amplitude inferred from NANOGrav and fixed $\propto f^{2/3}$ spectral tilt, as appropriate in the absence of environmental or relativistic effects). The yellow shaded region indicates a potential cosmological signal from a first-order phase transition in the early Universe.

and that could be summarised by Eqs. (8), (9) and (11). Recall that $f_{\text{grav}} \approx 2f_{\text{DM}}$, where f_{DM} can be found in Eq. (6) for the purely gravitational coupling and the quadratic direct coupling $\alpha(\phi) = \phi^2/\Lambda^2$. For the linear coupling, such as the one in Eq. (10), $f = f_{\text{DM}}$.

The purely gravitational case represented by Eq. (8) (which is independent of other couplings) generates the black line in Fig. 10 as a function of the ULDM mass (horizontal axes). We notice that the priority UOP scenario may reach a sensitivity that would directly detect dark matter in the Solar System for masses below $\sim 10^{-23}$ eV. While these low masses are already in tension with other astrophysical observations (see, *e.g.*, [11, 41]), this is a very interesting result since, as compared to other bounds, it would be based on *direct* sensitivity to the ULDM field.

Regarding direct coupling, from Eq. (9) and the bounds on the gravitational effect from Fig. 10, it is easy to understand the constraining power on Λ_2 for different masses. In the upper panel of Fig. 10 we show the equivalent h_c predicted for two values of Λ_2 : one corresponding to the typical minimum value UOP could constrain (around $\Lambda_2 = 10^{-18}$ GeV) and another one that improves on the best bounds from *Cassini* ($\Lambda_2 = 10^{-16}$ GeV). We see clearly that in both cases, even the baseline UOP will produce the most stringent results for ULDM in a large range of masses. To make this more explicit, in




FIG. 9. Forecast constraints on cosmological SGWB signals from early Universe phase transitions. These are shown using 2D projections of the 4D space of parameters describing the transition: the temperature of the Universe at the time of the transition, T_* , the fractional energy density released by the transition, α , the rate of the transition, β , and the terminal velocity of the bubble walls, v_w .

Fig. 10, lower panel, we show the constraints that can be achieved on Λ_2 as a function of the DM mass for different configurations and compare them to *Cassini* and PTA [21] data (see also [20] and the recent [65, 95] for related bounds). It is clear that the UOP will enormously extend the searches for these models, with the potential to generate the first direct detection of dark matter.

Finally, we also show in Fig. 10 the equivalent h_c from the effect from Eq. (11) representing axion-like particles. Recall that this effect also arises from an oscillating part of the ULDM with frequency f_{DM} . We have used a value of the coupling $g_{a\gamma\gamma}$ characteristic of current bounds at the relevant masses, see [80]. From this plot, we conclude that the priority or optimistic cases of the UOP mission would also have a unique legacy in the search for axion-like particles.




FIG. 10. Upper plot: Expected sensitivity of the UOP for ULDM. The effect of ULDM is represented as a corresponding h_c that can be directly compared with Fig. 6. We plot the gravitational case (Eq. (8)), two cases where ULDM couples to matter (Eq. (9)), and a case where ULDM couples to light (Eqs. (11)). Lower plot: Expected sensitivity of FUM for ULDM for the model with direct coupling of matter to $\alpha(\phi) = \phi^2/\Lambda_2^2$. We compare with bounds from *Cassini* [9] and the EPTA [94] (see also [83]).

VII. BEYOND THE BASELINE: REDUCING THE ALLAN DEVIATION

A. Noise sources

Predicting realistic improvements in the Allan deviation requires considering recent advancements in a wide range of technologies. Here we base some qualitative considerations on the thorough discussion in Ref. [7]. *Cassini*-era noise was dominated by 3 major components; the antenna mechanical noise, the plasma interference noise, and the tropospheric scintillation noise. The first is simply a mechanical limitation, which can be addressed by implementing a complementary smaller and stiffer antenna in combination with the main dish, as well as via three-point antenna calibration, which carries the potential to reduce σ_A of Ka-band tracking missions by one

order of magnitude.⁷

The noise of plasma scintillation, the variation of the refractive index of the interplanetary medium, is localised around 3×10^{-3} Hz and then steadily drops down for lower frequencies [73]. It is astrophysical in origin, and therefore cannot be easily addressed with technological improvements. One possibility is to prioritise tracking measurements at optimal Sun-Earth-spacecraft configuration, as well as by upgrading the Doppler link to higher frequencies, as discussed in Section VII B.

Tropospheric noise is thought to dominate the noise power spectral density frequency below 10^{-4} Hz. Since the *Cassini* era, corrective measures based on water-vapor-radiometers have been shown to be able to reduce the Allan deviation up to a factor of 10, down to $1.5 \times 10^{-15} - 3 \times 10^{-15}$ [7]. Furthermore, drastic improvements in tropospheric noise would be achievable by tracking the spacecraft with high-altitude facilities, either ground-based or with balloons. Finally, an ulterior opportunity to reduce tropospheric noise would be using multiple measurement points via a radio telescope array [107].

Glitches and unmodelled accelerations, *e.g.*, from solar winds or imprecisions in the planetary ephemeris, may also leave residual imprints on the Doppler time series. However, these outliers will not influence the coherent stacking of the GW signal in the experiment. A more quantitative treatment of their influence can be addressed by a more detailed simulation of the mission's trajectory, once the satellite specifications are known. Connected to this, one must also consider the noise from asteroids in the Solar system. A simple estimate using Brownian-motion argument [67] suggests a noise level in the solar system barycenter of order 10^{-15} in the 10 nHz range due to asteroids smaller than 80 km. Furthermore, the results in [39] indicate a lower level of 10^{-18} for the potential generated by asteroids on a satellite at \sim 1 AU from the Sun (with typical frequencies around and below μ Hz), which would be irrelevant to the scientific outcome we outlined. However, a more detailed analysis confirming this expectation and considering a realistic mission trajectory and possible mitigation measures using updated asteroid ephemerides (see [102] for the current strategy followed by NANOGrav) would be opportune.

B. Breakthrough: Optical links?

A true breakthrough in improving the Allan deviation to $\lesssim 10^{-15}$ could be achieved by upgrading the Doppler link to optical frequencies. Here we briefly review theoretical limits and recent experiments. There are three distinct techniques in optical ranging.

⁷ Private conversation with Sami W. Asmar, NASA JPL.

- The passive-reflector method has reflecting cubes on the target, which reflect light back to the source. This has been very successful for lunar ranging [75] but (as we will see below) is not feasible at interplanetary distances.
- The synchronous transponder method has the target receive a signal and then send another signal back with a fixed delay. This has been recently demonstrated with the Hayabusa2 mission to ~ 15 lunar distances [79], with ground observations even during the day.
- An asynchronous transponder ranging both sides sends signals at pre-designated intervals, and the ranging is computed indirectly [32]. The Deep Space Optical Communications (DSOC) system on the Psyche mission [76, 84] is an example.

We can get an idea of what is and is not feasible using order-of-magnitude arguments. Similar arguments appear in [33]. Consider a signal at wavelength λ from a transmitting telescope of area A_T to a receiving telescope of area A_R at distance R . A diffraction-limited beam will have a solid angle of $\sim \lambda^2/A_T$. Of the photons transmitted from A_T , the fraction received at A_R will be $\sim A_R/R^2$ divided by the solid angle. Thus a fraction

$$\frac{A_T A_R}{\lambda^2 R^2} \quad (32)$$

will be received. For a transmitting power P , the rate of photons received is then

$$\frac{P}{hc\lambda} \frac{A_T A_R}{R^2}. \quad (33)$$

The maximum data rate for a given power and area is thus $\propto \lambda^{-1}$.

For some orders of magnitude, let us put $\lambda = 1 \times 10^{-6}$ m, $A_T = A_R = 1$ m² and $R = 3 \times 10^9$ m. This makes the photon fraction $\sim 10^{-13}$ and the photon rate $\sim 5 \times 10^4$ W⁻¹.

With a passive reflector the photon fraction (32) gets squared through the round trip. At lunar distances ($R \simeq 3 \times 10^9$ m) this is tolerable, but at interplanetary distances, it would become hopeless. Hence an active system is needed.

Ref. [98] discusses optical telecom prospects for ice-giant missions briefly, noting the positives and negatives, which also follow from the above arguments.

- + The data rates can be orders of magnitude higher. Ref. [96] demonstrated uplink and downlink rates of 19 and 38 Mbps respectively from lunar distances. The Psyche mission to the asteroid belt was designed for rates of 250 Mbps from deep space [23], and a demonstration of the data rate (showing a cat pursuing a laser pointer) was widely reported in the media and social media.

- The narrowness of the optical beam requires accurate pointing, including the effect of light travel time.
- To be weather-proof, the system requires redundant ground stations (or balloon or orbiting stations).

Laboratory experiments of asynchronous ranging [16] give precisions of better than 1 mm, which amounts to 3×10^{-16} of the Uranus distance. Reference [34] discuss the error budget for interplanetary ranging, and conclude that there is potential for (sub-)mm range accuracy. That is, accuracies even better than the optimistic scenario considered earlier appear feasible with optical ranging.

VIII. CONCLUSION: MISSION OUTLOOK

We have demonstrated that the prospective flagship mission to Uranus (UOP) has rich scientific opportunities that go beyond its nominal planetary science goals. Doppler tracking data may be accumulated over the entire interplanetary cruise phase, and subsequently stacked according to the methodology showcased in Section II E. This results in the capacity to detect imprints from deterministic GWs from individual SMBH binaries (Section IV), astrophysical as well as early universe stochastic GW backgrounds (Sections IV and V), and finally gravitational and non-gravitational couplings between the spacecraft and dark matter (Section VI). Such varied and timely prospects would greatly expand the scientific yield of the mission, broadening its relevancy to communities beyond the planetary sciences. So the question beckons: what is required to actually achieve what we have shown to be possible in this paper? We note that our analysis of the mission's capacity to distinguish various signals should be repeated with more sophisticated noise models, including, *e.g.*, non-Gaussianities and the time dependence of different noise contributions along the cruise phase. A more thorough treatment of the mission's trajectory is also crucial, in particular considering the influence of unmodelled accelerations due to uncertainties in the masses of planets, asteroids, and solar radiation pressure. Additionally, a more sophisticated data analysis pipeline should take into account the possibility of multiple signals to be present at the same time, including the possibility of stochastic backgrounds overshadowing individual signals. Nevertheless, taking these limitations at face value, we have demonstrated how two crucial factors determine the magnitude of the achievable scientific goals. They are:

- *The total amount of tracking data* collected during the cruise phase, which could range from a few data points to constant tracking at the optimal allowed cadence of 0.1 Hz for the entire duration of the cruise phase.

- *The technological feasibility of reducing the Allan deviation* with respect to its baseline value of $\sim 10^{-14}$ at 10s, as observed for the *Cassini* mission data more than three decades ago, in 1992.

We maintain that these two points should become part of the mission requirements, in light of the numerous scientific opportunities demonstrated in this work. With them, the UOP mission has the potential for groundbreaking discoveries about black holes living in galactic centers, the study of particle physics beyond the standard model and the reach of current particle accelerators, and the nature of dark matter.

ACKNOWLEDGMENTS

L.Z., D.S, D.J.D, D.O., A.D. and P.S. acknowledge the support of the *International Space Science Institute (ISSI)* in Bern, Switzerland. L.Z. and D.O. acknowledge support from ERC Starting Grant No. 121817–Black-

HoleMergs. D.J.D. and D.O. received funding from the Danish Independent Research Fund through Sapere Aude Starting Grant No. 121587. A.D. acknowledges support from NSF grant AST-2319441. We thank Mark Hofstadter, Joseph Lazio and Daniele Durante for useful discussions. D.S. thanks Timothée Schaeffer for continuing moral support and inspiration. L.Z. also acknowledges G.M.’s curiosity. The research leading to these results has received funding from the Spanish Ministry of Science and Innovation (PID2020-115845GB-I00/AEI/10.13039/501100011033). IFAE is partially funded by the CERCA program of the Generalitat de Catalunya. D.B. acknowledges the support from the Departament de Recerca i Universitats de la Generalitat de Catalunya al Grup de Recerca i Universitats from Generalitat de Catalunya to the Grup de Recerca 00649 (Codi: 2021 SGR 00649). A.C.J. is supported by the Science and Technology Facilities Council (STFC) through the UKRI Quantum Technologies for Fundamental Physics Programme [grant number ST/T005904/1]. This work was partly enabled by the UCL Cosmoparticle Initiative.

-
- [1] Abbott, R., et al. 2022, *Phys. Rev. D*, 105, 063030, doi: [10.1103/PhysRevD.105.063030](https://doi.org/10.1103/PhysRevD.105.063030)
- [2] Agazie, G., Anumalapudi, A., Archibald, A. M., et al. 2023, *ApJ*, 951, L8, doi: [10.3847/2041-8213/acdac6](https://doi.org/10.3847/2041-8213/acdac6)
- [3] —. 2023, arXiv e-prints, arXiv:2306.16220, doi: [10.48550/arXiv.2306.16220](https://doi.org/10.48550/arXiv.2306.16220)
- [4] Aghanim, N., et al. 2020, *Astron. Astrophys.*, 641, A6, doi: [10.1051/0004-6361/201833910](https://doi.org/10.1051/0004-6361/201833910)
- [5] Anderson, J. D., Armstrong, J. W., Campbell, J. K., et al. 1992, *Space Sci. Rev.*, 60, 591, doi: [10.1007/BF00216869](https://doi.org/10.1007/BF00216869)
- [6] Antoniadis, J., Arumugam, P., Arumugam, S., et al. 2023, arXiv e-prints, arXiv:2306.16214, doi: [10.48550/arXiv.2306.16214](https://doi.org/10.48550/arXiv.2306.16214)
- [7] Armstrong, J. W. 2006, *Living Reviews in Relativity*, 9, 1, doi: [10.12942/lrr-2006-1](https://doi.org/10.12942/lrr-2006-1)
- [8] Armstrong, J. W., Estabrook, F. B., & Wahlquist, H. D. 1987, *Astrophys. J.*, 318, 536, doi: [10.1086/165390](https://doi.org/10.1086/165390)
- [9] Armstrong, J. W., Iess, L., Tortora, P., & Bertotti, B. 2003, *Astrophys. J.*, 599, 806, doi: [10.1086/379505](https://doi.org/10.1086/379505)
- [10] Bansal, A., Ichiki, K., Tashiro, H., & Matsuoka, Y. 2023, *MNRAS*, 523, 3840, doi: [10.1093/mnras/stad1608](https://doi.org/10.1093/mnras/stad1608)
- [11] Bar, N., Blas, D., Blum, K., & Sibiriyakov, S. 2018, *Phys. Rev. D*, 98, 083027, doi: [10.1103/PhysRevD.98.083027](https://doi.org/10.1103/PhysRevD.98.083027)
- [12] Begelman, M. C., Blandford, R. D., & Rees, M. J. 1980, *Nature (London)*, 287, 307, doi: [10.1038/287307a0](https://doi.org/10.1038/287307a0)
- [13] Bertotti, B., Vecchio, A., & Iess, L. 1999, *Phys. Rev. D*, 59, 082001, doi: [10.1103/PhysRevD.59.082001](https://doi.org/10.1103/PhysRevD.59.082001)
- [14] Bertotti, B., Ambrosini, R., Asmar, S. W., et al. 1992, *A&AS*, 92, 431
- [15] Bi, X., Guo, Z., Zou, X., et al. 2024, arXiv e-prints, arXiv:2402.06096, doi: [10.48550/arXiv.2402.06096](https://doi.org/10.48550/arXiv.2402.06096)
- [16] Birnbaum, K. M., Chen, Y., & Hemmati, H. 2010, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 7587, Free-Space Laser Communication Technologies XXII, ed. H. Hemmati, 75870A, doi: [10.1117/12.843315](https://doi.org/10.1117/12.843315)
- [17] Blas, D., Caputo, A., Ivanov, M. M., & Sberna, L. 2020, *Phys. Dark Univ.*, 27, 100428, doi: [10.1016/j.dark.2019.100428](https://doi.org/10.1016/j.dark.2019.100428)
- [18] Blas, D., & Jenkins, A. C. 2022, *Phys. Rev. D*, 105, 064021, doi: [10.1103/PhysRevD.105.064021](https://doi.org/10.1103/PhysRevD.105.064021)
- [19] —. 2022, *Phys. Rev. Lett.*, 128, 101103, doi: [10.1103/PhysRevLett.128.101103](https://doi.org/10.1103/PhysRevLett.128.101103)
- [20] Blas, D., López Nacir, D., & Sibiriyakov, S. 2020, *Phys. Rev. D*, 101, 063016, doi: [10.1103/PhysRevD.101.063016](https://doi.org/10.1103/PhysRevD.101.063016)
- [21] Blas, D., Nacir, D. L., & Sibiriyakov, S. 2017, *Phys. Rev. Lett.*, 118, 261102, doi: [10.1103/PhysRevLett.118.261102](https://doi.org/10.1103/PhysRevLett.118.261102)
- [22] Bondi, H., Pirani, F. A. E., & Robinson, I. 1959, *Proceedings of the Royal Society of London Series A*, 251, 519, doi: [10.1098/rspa.1959.0124](https://doi.org/10.1098/rspa.1959.0124)
- [23] Buck, B. R., Allen, G. D., Duerr, E. K., et al. 2019, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 10978, *Advanced Photon Counting Techniques XIII*, ed. M. A. Itzler, J. C. Bienfang, & K. A. McIntosh, 1097809, doi: [10.1117/12.2520617](https://doi.org/10.1117/12.2520617)
- [24] Bucko, J., Soyuer, D., & Zwick, L. 2023, *MNRAS*, 524, L32, doi: [10.1093/mnrasl/slad067](https://doi.org/10.1093/mnrasl/slad067)
- [25] Caprini, C., & Figueroa, D. G. 2018, *Classical and Quantum Gravity*, 35, 163001, doi: [10.1088/1361-6382/aac608](https://doi.org/10.1088/1361-6382/aac608)
- [26] Caprini, C., et al. 2016, *JCAP*, 04, 001, doi: [10.1088/1475-7516/2016/04/001](https://doi.org/10.1088/1475-7516/2016/04/001)
- [27] —. 2020, *JCAP*, 03, 024, doi: [10.1088/1475-7516/2020/03/024](https://doi.org/10.1088/1475-7516/2020/03/024)
- [28] Chadha-Day, F., Ellis, J., & Marsh, D. J. E. 2022, *Sci. Adv.*, 8, abj3618, doi: [10.1126/sciadv.abj3618](https://doi.org/10.1126/sciadv.abj3618)
- [29] Colpi, M., Danzmann, K., Hewitson, M., et al. 2024, arXiv e-prints, arXiv:2402.07571, doi: [10.48550/arXiv.2402.07571](https://doi.org/10.48550/arXiv.2402.07571)

- 2402.07571
- [30] Comoretto, G., Bertotti, B., Iess, L., & Ambrosini, R. 1992, *Nuovo Cimento C Geophysics Space Physics C*, 15, 1193, doi: [10.1007/BF02506712](https://doi.org/10.1007/BF02506712)
- [31] Croton, D. J. 2009, *MNRAS*, 394, 1109, doi: [10.1111/j.1365-2966.2009.14429.x](https://doi.org/10.1111/j.1365-2966.2009.14429.x)
- [32] Degnan, J. J. 2002, *Journal of Geodynamics*, 34, 551, doi: [10.1016/S0264-3707\(02\)00044-3](https://doi.org/10.1016/S0264-3707(02)00044-3)
- [33] —. 2023, *Photonics for Solar Energy Systems IX*, 10, 98, doi: [10.3390/photonics10020098](https://doi.org/10.3390/photonics10020098)
- [34] Dirkx, D., Prochazka, I., Bauer, S., et al. 2019, *Journal of Geodesy*, 93, 2405, doi: [10.1007/s00190-018-1171-x](https://doi.org/10.1007/s00190-018-1171-x)
- [35] Dosopoulou, F., & Antonini, F. 2017, *Astrophys. J.*, 840, 31, doi: [10.3847/1538-4357/aa6b58](https://doi.org/10.3847/1538-4357/aa6b58)
- [36] EPTA Collaboration, InPTA Collaboration, Antoniadis, J., et al. 2023, *A&A*, 678, A50, doi: [10.1051/0004-6361/202346844](https://doi.org/10.1051/0004-6361/202346844)
- [37] Estabrook, F. B., & Wahlquist, H. D. 1975, *General Relativity and Gravitation*, 6, 439, doi: [10.1007/BF00762449](https://doi.org/10.1007/BF00762449)
- [38] Fakhouri, O., Ma, C.-P., & Boylan-Kolchin, M. 2010, *MNRAS*, 406, 2267, doi: [10.1111/j.1365-2966.2010.16859.x](https://doi.org/10.1111/j.1365-2966.2010.16859.x)
- [39] Fedderke, M. A., Graham, P. W., & Rajendran, S. 2021, *Phys. Rev. D*, 103, 103017, doi: [10.1103/PhysRevD.103.103017](https://doi.org/10.1103/PhysRevD.103.103017)
- [40] Fedderke, M. A., & Mathur, A. 2023, *Phys. Rev. D*, 107, 043004, doi: [10.1103/PhysRevD.107.043004](https://doi.org/10.1103/PhysRevD.107.043004)
- [41] Ferreira, E. G. M. 2021, *Astron. Astrophys. Rev.*, 29, 7, doi: [10.1007/s00159-021-00135-6](https://doi.org/10.1007/s00159-021-00135-6)
- [42] Fletcher, L. N., Simon, A. A., Hofstadter, M. D., et al. 2020, arXiv e-prints, arXiv:2008.12125. <https://arxiv.org/abs/2008.12125>
- [43] Fletcher, L. N., Helled, R., Roussos, E., et al. 2020, *Planet. Space Sci.*, 191, 105030, doi: [10.1016/j.pss.2020.105030](https://doi.org/10.1016/j.pss.2020.105030)
- [44] Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, 125, 306, doi: [10.1086/670067](https://doi.org/10.1086/670067)
- [45] Foster, J. W., Rodd, N. L., & Safdi, B. R. 2018, *Phys. Rev. D*, 97, 123006, doi: [10.1103/PhysRevD.97.123006](https://doi.org/10.1103/PhysRevD.97.123006)
- [46] Foster, R. S., & Backer, D. C. 1990, *Astrophys. J.*, 361, 300, doi: [10.1086/169195](https://doi.org/10.1086/169195)
- [47] Fukusumi, K., Morisaki, S., & Suyama, T. 2023, *Phys. Rev. D*, 108, 095054, doi: [10.1103/PhysRevD.108.095054](https://doi.org/10.1103/PhysRevD.108.095054)
- [48] Georgakakis, A., Comparat, J., Merloni, A., et al. 2019, *MNRAS*, 487, 275, doi: [10.1093/mnras/sty3454](https://doi.org/10.1093/mnras/sty3454)
- [49] Girija, A. P. 2023, *Acta Astronautica*, 202, 104, doi: [10.1016/j.actaastro.2022.10.005](https://doi.org/10.1016/j.actaastro.2022.10.005)
- [50] Graham, P. W., Irastorza, I. G., Lamoreaux, S. K., Lindner, A., & van Bibber, K. A. 2015, *Ann. Rev. Nucl. Part. Sci.*, 65, 485, doi: [10.1146/annurev-nucl-102014-022120](https://doi.org/10.1146/annurev-nucl-102014-022120)
- [51] Helled, R., & Fortney, J. J. 2020, arXiv e-prints, arXiv:2007.10783. <https://arxiv.org/abs/2007.10783>
- [52] Hindmarsh, M., Huber, S. J., Rummukainen, K., & Weir, D. J. 2014, *Phys. Rev. Lett.*, 112, 041301, doi: [10.1103/PhysRevLett.112.041301](https://doi.org/10.1103/PhysRevLett.112.041301)
- [53] Hobbs, G., Archibald, A., Arzoumanian, Z., et al. 2010, *Classical and Quantum Gravity*, 27, 084013, doi: [10.1088/0264-9381/27/8/084013](https://doi.org/10.1088/0264-9381/27/8/084013)
- [54] Hofstadter, M., Simon, A., Atreya, S., et al. 2019, *Planet. Space Sci.*, 177, 104680, doi: [10.1016/j.pss.2019.06.004](https://doi.org/10.1016/j.pss.2019.06.004)
- [55] Hopkins, P. F., Hernquist, L., Cox, T. J., et al. 2005, *Astrophys. J.*, 630, 705, doi: [10.1086/432438](https://doi.org/10.1086/432438)
- [56] Hu, W., Barkana, R., & Gruzinov, A. 2000, *Phys. Rev. Lett.*, 85, 1158, doi: [10.1103/PhysRevLett.85.1158](https://doi.org/10.1103/PhysRevLett.85.1158)
- [57] Hui, L., Ostriker, J. P., Tremaine, S., & Witten, E. 2017, *Phys. Rev. D*, 95, 043541, doi: [10.1103/PhysRevD.95.043541](https://doi.org/10.1103/PhysRevD.95.043541)
- [58] Iorio, L., Girija, A. P., & Durante, D. 2023, *MNRAS*, 523, 3595, doi: [10.1093/mnras/stad1446](https://doi.org/10.1093/mnras/stad1446)
- [59] Jarmak, S., Leonard, E., Akins, A., et al. 2020, *Acta Astronautica*, 170, 6, doi: [10.1016/j.actaastro.2020.01.030](https://doi.org/10.1016/j.actaastro.2020.01.030)
- [60] Kamionkowski, M., Kosowsky, A., & Turner, M. S. 1994, *Phys. Rev. D*, 49, 2837, doi: [10.1103/PhysRevD.49.2837](https://doi.org/10.1103/PhysRevD.49.2837)
- [61] Kelley et. al. In Prep.
- [62] Khmelnskiy, A., & Rubakov, V. 2014, *JCAP*, 02, 019, doi: [10.1088/1475-7516/2014/02/019](https://doi.org/10.1088/1475-7516/2014/02/019)
- [63] Kollmann, P., Cohen, I., Allen, R. C., et al. 2020, *Space Sci. Rev.*, 216, 78, doi: [10.1007/s11214-020-00696-5](https://doi.org/10.1007/s11214-020-00696-5)
- [64] Kormendy, J., & Ho, L. C. 2013, *ARA&A*, 51, 511, doi: [10.1146/annurev-astro-082708-101811](https://doi.org/10.1146/annurev-astro-082708-101811)
- [65] Kûs, P., López Nacir, D., & Urban, F. R. 2024. <https://arxiv.org/abs/2402.04099>
- [66] Leja, J., Speagle, J. S., Johnson, B. D., et al. 2020, *Astrophys. J.*, 893, 111, doi: [10.3847/1538-4357/ab7e27](https://doi.org/10.3847/1538-4357/ab7e27)
- [67] Loeb, A. 2024. <https://arxiv.org/abs/2405.05410>
- [68] Luo, J., Chen, L.-S., Duan, H.-Z., et al. 2016, *Classical and Quantum Gravity*, 33, 035010, doi: [10.1088/0264-9381/33/3/035010](https://doi.org/10.1088/0264-9381/33/3/035010)
- [69] Maggiore, M. 2018, *Gravitational Waves: Volume 2: Astrophysics and Cosmology*, doi: [10.1093/oso/9780198570899.001.0001](https://doi.org/10.1093/oso/9780198570899.001.0001)
- [70] Marasco, A., Cresci, G., Posti, L., et al. 2021, *MNRAS*, 507, 4274, doi: [10.1093/mnras/stab2317](https://doi.org/10.1093/mnras/stab2317)
- [71] Miller, A. L., Badaracco, F., & Palomba, C. 2022, *Phys. Rev. D*, 105, 103035, doi: [10.1103/PhysRevD.105.103035](https://doi.org/10.1103/PhysRevD.105.103035)
- [72] Misner, C. W., Thorne, K. S., & Wheeler, J. A. 1973, *Gravitation*
- [73] Molera Calvés, G., Pogrebenko, S. V., Cimò, G., et al. 2014, *A&A*, 564, A4, doi: [10.1051/0004-6361/201322925](https://doi.org/10.1051/0004-6361/201322925)
- [74] Moore, C. J., Cole, R. H., & Berry, C. P. L. 2015, *Classical and Quantum Gravity*, 32, 015014, doi: [10.1088/0264-9381/32/1/015014](https://doi.org/10.1088/0264-9381/32/1/015014)
- [75] Müller, J., Murphy, T. W., Schreiber, U., et al. 2019, *Journal of Geodesy*, 93, 2195, doi: [10.1007/s00190-019-01296-0](https://doi.org/10.1007/s00190-019-01296-0)
- [76] Mumolo, J. M., Pestana, N., Clark, H., et al. 2021, Evaluation of asynchronous geiger-mode avalanche photodiode arrays for deep-space optical communications, DRAFT VERSION, Pasadena, CA: Jet Propulsion Laboratory, National Aeronautics and Space Administration, 2021, doi: [2014/54760](https://doi.org/2014/54760)
- [77] Nagano, K., Nakatsuka, H., Morisaki, S., et al. 2021, *Phys. Rev. D*, 104, 062008, doi: [10.1103/PhysRevD.104.062008](https://doi.org/10.1103/PhysRevD.104.062008)
- [78] Naoz, S., Will, C. M., Ramirez-Ruiz, E., et al. 2020, *Astrophys. J. Lett.*, 888, L8, doi: [10.3847/2041-8213/ab5e3b](https://doi.org/10.3847/2041-8213/ab5e3b)
- [79] Noda, H., Senshu, H., Osubo, T., et al. 2023, *Advances in Space Research*, 71, 4196, doi: [10.1016/j.asr.2022.06.004](https://doi.org/10.1016/j.asr.2022.06.004)

- 12.057
- [80] O'Hare, C. 2020, *cajohare/AxionLimits: AxionLimits*, v1.0, <https://cajohare.github.io/AxionLimits/>, Zenodo, doi: [10.5281/zenodo.3932430](https://doi.org/10.5281/zenodo.3932430)
- [81] Oshima, Y., Fujimoto, H., Kume, J., et al. 2023, *Phys. Rev. D*, 108, 072005, doi: [10.1103/PhysRevD.108.072005](https://doi.org/10.1103/PhysRevD.108.072005)
- [82] Pirani, F. A. E. 1956, *Acta Physica Polonica*, 15, 389
- [83] Porayko, N. K., et al. 2018, *Phys. Rev. D*, 98, 102002, doi: [10.1103/PhysRevD.98.102002](https://doi.org/10.1103/PhysRevD.98.102002)
- [84] Pretorius, H., Aull, B., Clark, H., et al. 2021, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 11721, *Advanced Photon Counting Techniques XV*, ed. M. A. Itzler, J. C. Bienfang, & K. A. McIntosh, 117210S, doi: [10.1117/12.2588469](https://doi.org/10.1117/12.2588469)
- [85] Reardon, D. J., Zic, A., Shannon, R. M., et al. 2023, *ApJ*, 951, L6, doi: [10.3847/2041-8213/acdd02](https://doi.org/10.3847/2041-8213/acdd02)
- [86] Ricarte, A., & Natarajan, P. 2018, *MNRAS*, 474, 1995, doi: [10.1093/mnras/stx2851](https://doi.org/10.1093/mnras/stx2851)
- [87] Robson, T., Cornish, N. J., & Liu, C. 2019, *Classical and Quantum Gravity*, 36, 105011, doi: [10.1088/1361-6382/ab1101](https://doi.org/10.1088/1361-6382/ab1101)
- [88] Rodriguez-Gomez, V., Genel, S., Vogelsberger, M., et al. 2015, *MNRAS*, 449, 49, doi: [10.1093/mnras/stv264](https://doi.org/10.1093/mnras/stv264)
- [89] Sereno, M., & Jetzer, P. 2006, *MNRAS*, 371, 626, doi: [10.1111/j.1365-2966.2006.10670.x](https://doi.org/10.1111/j.1365-2966.2006.10670.x)
- [90] Sesana, A., Vecchio, A., & Colacino, C. N. 2008, *MNRAS*, 390, 192, doi: [10.1111/j.1365-2966.2008.13682.x](https://doi.org/10.1111/j.1365-2966.2008.13682.x)
- [91] Sesana, A., et al. 2021, *Exper. Astron.*, 51, 1333, doi: [10.1007/s10686-021-09709-9](https://doi.org/10.1007/s10686-021-09709-9)
- [92] Shankar, F., Weinberg, D. H., & Miralda-Escudé, J. 2013, *MNRAS*, 428, 421, doi: [10.1093/mnras/sts026](https://doi.org/10.1093/mnras/sts026)
- [93] Simon, A. A., Fletcher, L. N., Arridge, C., et al. 2020, *Space Sci. Rev.*, 216, 17, doi: [10.1007/s11214-020-0639-1](https://doi.org/10.1007/s11214-020-0639-1)
- [94] Smarra, C., et al. 2023, *Phys. Rev. Lett.*, 131, 171001, doi: [10.1103/PhysRevLett.131.171001](https://doi.org/10.1103/PhysRevLett.131.171001)
- [95] —. 2024. <https://arxiv.org/abs/2405.01633>
- [96] Sodnik, Z., Smit, H., Sans, M., et al. 2014, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 8971, *Free-Space Laser Communication and Atmospheric Propagation XXVI*, ed. H. Hemmati & D. M. Boroson, 89710W, doi: [10.1117/12.2045510](https://doi.org/10.1117/12.2045510)
- [97] Soyuer, D., Zwick, L., D'Orazio, D. J., & Saha, P. 2021, *MNRAS*, 503, L73, doi: [10.1093/mnras1/slab025](https://doi.org/10.1093/mnras1/slab025)
- [98] Spilker, T. R. 2020, *Philosophical Transactions of the Royal Society of London Series A*, 378, 20190488, doi: [10.1098/rsta.2019.0488](https://doi.org/10.1098/rsta.2019.0488)
- [99] Springel, V., Di Matteo, T., & Hernquist, L. 2005, *ApJ*, 620, L79, doi: [10.1086/428772](https://doi.org/10.1086/428772)
- [100] The International Pulsar Timing Array Collaboration, Agazie, G., Antoniadis, J., et al. 2023, arXiv e-prints, arXiv:2309.00693, doi: [10.48550/arXiv.2309.00693](https://doi.org/10.48550/arXiv.2309.00693)
- [101] Unal, C., Urban, F. R., & Kovetz, E. D. 2022. <https://arxiv.org/abs/2209.02741>
- [102] Vallisneri, M., et al. 2020, doi: [10.3847/1538-4357/ab7b67](https://doi.org/10.3847/1538-4357/ab7b67)
- [103] Vasiliev, E., Antonini, F., & Merritt, D. 2015, *Astrophys. J.*, 810, 49, doi: [10.1088/0004-637X/810/1/49](https://doi.org/10.1088/0004-637X/810/1/49)
- [104] Volonteri, M., Haardt, F., & Madau, P. 2003, *Astrophys. J.*, 582, 559, doi: [10.1086/344675](https://doi.org/10.1086/344675)
- [105] Wahlquist, H. 1987, *General Relativity and Gravitation*, 19, 1101, doi: [10.1007/BF00759146](https://doi.org/10.1007/BF00759146)
- [106] Will, C. M., Naoz, S., Hees, A., et al. 2023, *Astrophys. J.*, 959, 58, doi: [10.3847/1538-4357/ad09b3](https://doi.org/10.3847/1538-4357/ad09b3)
- [107] Williams, S., Bock, Y., & Fang, P. 1998, *J. Geophys. Res.*, 103, 27,051, doi: [10.1029/98JB02794](https://doi.org/10.1029/98JB02794)
- [108] Xu, H., Chen, S., Guo, Y., et al. 2023, *Research in Astronomy and Astrophysics*, 23, 075024, doi: [10.1088/1674-4527/acdfa5](https://doi.org/10.1088/1674-4527/acdfa5)
- [109] Zwick, L., Soyuer, D., & Bucko, J. 2022, *A&A*, 664, A188, doi: [10.1051/0004-6361/202243741](https://doi.org/10.1051/0004-6361/202243741)