

# THE SMASHING SPECTRUM OF SHEAVES

KO AOKI

ABSTRACT. For an arbitrary  $\infty$ -topos, we classify the smashing localizations in the  $\infty$ -category of sheaves valued in derived vector spaces: Any of them is the restriction functor to a (unique) closed subtopos. Our proof is based on the existence of a Boolean cover.

This result in particular gives us the first example of a nonzero presentably symmetric monoidal stable  $\infty$ -category whose smashing spectrum has no points.

Combining this with the sheaves-spectrum adjunction, we obtain a Tannaka-type categorical reconstruction result for locales.

## 1. INTRODUCTION

Given a presentably symmetric monoidal stable  $\infty$ -category  $\mathcal{C}$ , understanding its localizations is fundamental. Particularly, *smashing localizations*, i.e., localizations  $L: \mathcal{C} \rightarrow \mathcal{C}$  that are equivalent to  $L(\mathbf{1}) \otimes -$ , are classical and important in stable homotopy theory; cf. [16, Section 1]. A basic observation is that they form a *frame*, i.e., a complete lattice satisfying the distributivity law  $\bigvee_{i \in I} x \wedge y_i = x \wedge \bigvee_{i \in I} y_i$  for any  $x$  and  $(y_i)_{i \in I}$ . The associated locale, for which we write  $\mathrm{Sm}(\mathcal{C})$ , is called the *smashing spectrum* of  $\mathcal{C}$ . See [2] and references therein about this construction.

Computing  $\mathrm{Sm}(\mathcal{C})$  for given  $\mathcal{C}$  is typically extremely difficult; e.g., even the most standard case  $\mathcal{C} = \mathbf{Sp}$ , the  $\infty$ -category of spectra, contains the telescope conjecture. Although the conjecture itself was recently settled in [8], our knowledge of  $\mathrm{Sm}(\mathbf{Sp})$  remains very limited. The case  $\mathcal{C} = \mathbf{D}(R)$  for a commutative ring  $R$  has been actively studied: When  $R$  is noetherian,  $\mathrm{Sm}(\mathbf{D}(R))$  is the Zariski spectrum of  $R$ ; see [15, Section 3] or [5, Section 6]. In general, there are many more smashing localizations; see, e.g., the calculation when  $R$  is a valuation ring in [7, Section 5]. Still, determining  $\mathrm{Sm}(\mathbf{D}(R))$  is unapproachable in general.

In this paper, as  $\mathcal{C}$  we consider  $\mathrm{Shv}(\mathcal{X}; \mathbf{D}(k))$  for an  $\infty$ -topos  $\mathcal{X}$  and a field  $k$ . For example, for a topological space (or a locale)  $X$ , by considering  $\mathcal{X} = \mathrm{Shv}(X)^{\mathrm{hyp}}$ , we get the derived category of sheaves on  $X$  valued in classical  $k$ -vector spaces as  $\mathcal{C}$ . There is an obvious class of smashing localizations on  $\mathcal{C}$ : For any closed subtopos  $\mathcal{Z}$ , the composite

$$\mathrm{Shv}(\mathcal{X}; \mathbf{D}(k)) \xrightarrow{i^*} \mathrm{Shv}(\mathcal{Z}; \mathbf{D}(k)) \xrightarrow{i_*} \mathrm{Shv}(\mathcal{X}; \mathbf{D}(k))$$

is smashing due to the projection formula. The main theorem of this paper states that there are no other smashing localizations:

**Theorem A.** *For an  $\infty$ -topos  $\mathcal{X}$  and a field  $k$ , the smashing spectrum of  $\mathrm{Shv}(\mathcal{X}; \mathbf{D}(k))$  is canonically isomorphic to the locale of subterminal objects.*

We note that our proof uses a nontrivial theorem from (higher) topos theory as an input.

*Remark 1.1.* The phenomenon of the smashing spectrum only depending on the underlying locale in Theorem A is special to our choice of coefficients: An interesting example is given by Clausen–Mathew [9, Theorem 1.9], which says that for the étale  $\infty$ -topos of a nice scheme, the hypercompletion of  $\mathbf{Sp}$ -valued sheaves is smashing. Typically, hypercompletion is nontrivial (even over  $\mathrm{KU}$ ), as demonstrated by the argument of Wieland in [13, Warning 7.2.2.31]

The importance of this theorem is two-fold: First, as already explained in [2, Section 1.2], we can deduce the following Tannaka-type duality for locales by exploiting the main result of [2]:

**Theorem B.** *For a field  $k$ , the functor*

$$\mathrm{Shv}(-; \mathbf{D}(k)): \mathrm{Loc}^{\mathrm{op}} \rightarrow \mathrm{CAlg}_{\mathbf{D}(k)}(\mathrm{Pr})$$

is fully faithful.

*Remark 1.2.* In Theorem B, note that the category of locales has a canonical bicategorical structure; see Theorem 3.8 for a version where this is taken into account.

*Remark 1.3.* In Theorem B, we cannot replace  $k$  with  $\mathbf{Z}$ ; i.e.,

$$(1.4) \quad \mathrm{Shv}(-; \mathbf{D}(\mathbf{Z})) : \mathrm{Loc}^{\mathrm{op}} \rightarrow \mathrm{CAlg}_{\mathbf{D}(\mathbf{Z})}(\mathrm{Pr})$$

is not fully faithful: Consider the Sierpiński space, i.e., the Zariski spectrum of a discrete valuation ring. Then its sheaf  $\infty$ -category is equivalent to  $\mathrm{Fun}(\Delta^1, \mathbf{D}(\mathbf{Z}))$  with the pointwise symmetric monoidal structure. There is a symmetric monoidal functor  $\mathrm{Fun}(\Delta^1, \mathbf{D}(\mathbf{Z})) \rightarrow \mathbf{D}(\mathbf{Z})$  that pointwise carries  $F$  to the limit of  $\mathbf{Q} \otimes F(0) \rightarrow \mathbf{Q} \otimes F(1) \leftarrow F(1)$ , but it does not come from any locale map.

However, we can show that

$$\mathrm{Shv}(-; \mathbf{D}(\mathbf{Z})_{\geq 0}) : \mathrm{Loc}^{\mathrm{op}} \rightarrow \mathrm{CAlg}_{\mathbf{D}(\mathbf{Z})_{\geq 0}}(\mathrm{Pr})$$

is fully faithful by a similar proof to that of Theorem B: For a locale  $X$  we can identify the (unstable) smashing spectrum of  $\mathrm{Shv}(X; \mathbf{D}(\mathbf{Z})_{\geq 0})$  with  $X$  by applying Theorem A for prime fields. Then the desired result follows from considering  $\mathcal{V} = \mathbf{D}(\mathbf{Z})_{\geq 0}$  in the proof of Theorem B.

In future work, we will prove that (1.4) is fully faithful when restricting to compact Hausdorff spaces. Our proof uses a certain “continuous” version of  $\mathrm{Sm}$ .

Second, Theorem A has the following consequence:

**Theorem C.** *Any locale is the smashing spectrum of some presentably symmetric monoidal stable  $\infty$ -category. In particular, there is a nonzero presentably symmetric monoidal stable  $\infty$ -category whose smashing spectrum has no points.*

Recall that a locale is called *spatial* if it is isomorphic to the frame of open subsets of a (unique sober) topological space. We can ask the following:

**Question 1.5.** Under what condition on a presentably symmetric monoidal stable  $\infty$ -category  $\mathcal{C}$ , is the smashing spectrum  $\mathrm{Sm}(\mathcal{C})$  spatial?

This question is interesting because if we know that  $\mathrm{Sm}(\mathcal{C})$  is spatial, we can study smashing localizations of  $\mathcal{C}$  by studying the points and topology of  $\mathrm{Sm}(\mathcal{C})$  separately. In the compactly generated rigid case, people have attempted to prove the spatiality, but it is still open; see [4, Appendix A]. Lastly, we note that even the answer to the following is not known:

**Question 1.6.** Is  $\mathrm{Sm}(\mathbf{D}(R))$  always spatial for a commutative ring  $R$ ?

This paper is organized as follows: We first recall facts on sheaves on Boolean locales in Section 2. Using those, we prove Theorem A and deduce (a refined version of) Theorem B in Section 3.

**Acknowledgments.** I thank Alexander Efimov and Peter Scholze for helpful communications; the spatial case of Theorem A was obtained during a discussion with them. I also thank Scholze for his useful comments on the draft. I thank the Max Planck Institute for Mathematics for its hospitality.

**Conventions.** Regarding smashing spectrum, we continue using the notations from [2].

We do not use  $L$  or  $R$  to signify how functors are derived; e.g.,  $\otimes$  on derived  $\infty$ -categories means the derived tensor product.

## 2. BOOLEAN LOCALES AND SHEAVES THEREON

In this section, we study sheaves on Boolean locales. In Section 2.1, we review the notion of Boolean locales. In Section 2.2, we consider sheaves on them. In Section 2.3, we focus on the case of  $\mathbf{D}(k)$ -valued sheaves for a field  $k$ .

**2.1. Boolean locales.** We recall the following definition:

**Definition 2.1.** For an element  $x$  in a distributive lattice, its *complement* is an element  $x'$  satisfying  $x \wedge x' = \perp$  and  $x \vee x' = \top$ . By distributivity, it is necessarily unique as

$$x' = x' \wedge \top = x' \wedge (x \vee x'') = (x' \wedge x) \vee (x' \wedge x'') = \perp \vee (x' \wedge x'') = x' \wedge x''$$

and similarly  $x'' = x' \wedge x''$  hold when  $x'$  and  $x''$  are complements of  $x$ . A *Boolean lattice* is a distributive lattice in which every element has a complement. A locale is called *Boolean* if its underlying frame is Boolean. In a Boolean lattice, we write  $x \setminus y$  for  $x \wedge y'$  where  $y'$  is the complement of  $y$ .

**Example 2.2.** In the poset of open subsets of a topological space, complementable elements correspond to clopen subsets. Therefore, a spatial locale is Boolean if and only if it is discrete. In other words, a Boolean locale is either discrete or nonspatial.

*Remark 2.3.* A *complete Boolean algebra* is another name for a Boolean locale (or frame).

We here give two typical nondiscrete examples:

**Example 2.4.** Consider the poset of Borel subsets (or Lebesgue-measurable subsets) of the interval  $[0, 1]$ . We take a quotient with respect to the equivalence relation identifying  $B$  and  $B'$  when their symmetric difference  $B \Delta B'$  has measure zero. Then we get a Boolean frame, which is commonly called the *random algebra* by set theorists (cf. [17]). This does not have any points.

**Example 2.5.** We again consider the poset of Borel subsets (or open subsets) of  $[0, 1]$ , but this time we consider the equivalence relation that identifies  $B$  and  $B'$  when  $B \Delta B'$  is a meager set, i.e., a countable union of nowhere dense subsets. The quotient is again a Boolean frame, which is often called the *Cohen algebra* because of its role in forcing. This does not have any points.

*Remark 2.6.* Examples 2.4 and 2.5 are classical; most famously, they appeared in von Neumann's 1936–1937 lectures on continuous geometry [18, Part III], where he showed that these two locales are not isomorphic.

*Remark 2.7.* If a locale does not have any points, neither does its sheaf 1-topos. The example of a nontrivial 1-topos without points given in [3, IV.7.4], which is attributed to Deligne there, is based on this observation; it is simply the sheaf 1-topos of the locale given in Example 2.4.

**2.2. Sheaves on a Boolean locale.** We recall the following from [2, Proposition 3.18]:

**Theorem 2.8.** *Let  $X$  be a locale with frame of opens  $F$  and  $\mathcal{C}$  an  $\infty$ -category with limits. Then a presheaf  $\mathcal{F}: F^{\text{op}} \rightarrow \mathcal{C}$ , is a sheaf if and only if it satisfies the following:*

- (1) *The value  $\mathcal{F}(\perp)$  is final.*
- (2) *The square*

$$\begin{array}{ccc} \mathcal{F}(V \vee V') & \longrightarrow & \mathcal{F}(V) \\ \downarrow & & \downarrow \\ \mathcal{F}(V') & \longrightarrow & \mathcal{F}(V \wedge V') \end{array}$$

*is cartesian for any two opens  $V$  and  $V'$ .*

- (3) *The morphism*

$$\mathcal{F}\left(\bigvee D\right) \rightarrow \varinjlim_{U \in D} \mathcal{F}(U)$$

*is an equivalence for any directed subset  $D \subset F$ .*

We prove the following variant in the Boolean case:

**Theorem 2.9.** *Let  $X$  be a Boolean locale with frame of opens  $F$  and  $\mathcal{C}$  an  $\infty$ -category with limits. Then a presheaf is a sheaf if and only if it carries disjoint joins (see Definition 2.11) to products.*

*Remark 2.10.* By Theorem 2.9, the notion of sheaves on a Boolean locale makes sense when the coefficient  $\infty$ -category only has products and not necessarily all limits.

**Definition 2.11.** Let  $D$  be a Boolean lattice. We say that a family of elements  $(a_i)_{i \in I}$  is (pairwise) *disjoint* if  $a_i \wedge a_j = \perp$  for  $i \neq j$ .

**Proposition 2.12.** *In the statement of Theorem 2.8, when  $X$  is Boolean, (2) can be replaced by the condition that for any disjoint opens  $V$  and  $V'$ , the morphism  $\mathcal{F}(V \amalg V') \rightarrow \mathcal{F}(V) \times \mathcal{F}(V')$  is an equivalence.*

*Proof.* When we consider disjoint opens  $V$  and  $V'$  in (2), we exactly get our condition by (1). On the other hand, if  $\mathcal{F}$  satisfies (1) and our condition, the diagram in (2) is equivalent to

$$\begin{array}{ccc} \mathcal{F}(V \setminus V') \times \mathcal{F}(V \wedge V') \times \mathcal{F}(V' \setminus V) & \longrightarrow & \mathcal{F}(V \setminus V') \times \mathcal{F}(V \wedge V') \\ \downarrow & & \downarrow \\ \mathcal{F}(V \wedge V') \times \mathcal{F}(V' \setminus V) & \longrightarrow & \mathcal{F}(V \wedge V'), \end{array}$$

which is cartesian.  $\square$

*Remark 2.13.* The same argument as in Proposition 2.12 shows a similar statement for sheaves on a Stone space (aka totally disconnected compact Hausdorff space): A presheaf on a Stone space is a sheaf if and only if it carries directed joins to limits and finite disjoint joins of quasicompact opens to products.

We need the following, which is the  $\infty$ -categorical version of [1, Corollary 1.7]:

**Proposition 2.14.** *An  $\infty$ -category  $\mathcal{C}$  has all filtered colimits if and only if it has colimits indexed by ordinals.*

*For an  $\infty$ -category  $\mathcal{C}$  having small filtered colimits, a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$  preserves filtered colimits if and only if it preserves colimits indexed by ordinals.*

*Remark 2.15.* Whereas Proposition 2.14 is true, beware that it is not the case that any directed poset has a cofinal map from an ordinal: The poset of the finite subsets of an uncountable set is such an example. However, we also note that any countable directed poset has a cofinal map from  $\omega$ .

*Remark 2.16.* The same proof shows the refinement of Proposition 2.14 with a bound given by a fixed cardinal  $\kappa$ . For example, an  $\infty$ -category has all  $\kappa$ -small filtered colimits if and only if it has colimits indexed by ordinals in  $\kappa$ .

*Proof.* We prove the first statement since the second one follows from the same argument. The “only if” direction is clear. We prove the “if” direction. Let  $P$  be a directed poset with cardinality  $\kappa$ . We want to show that any diagram  $P \rightarrow \mathcal{C}$  has colimits. We proceed by induction on  $\kappa$ . If  $\kappa$  is finite,  $P$  has a final object and its image in  $\mathcal{C}$  is the colimit. Suppose that  $\kappa$  is infinite and that the statement holds for smaller cardinals. By [1, Lemma 1.6],<sup>1</sup> we can write  $P$  as an increasing of subposets  $P_\lambda$  for  $\lambda < \kappa$  such that each  $P_\lambda$  is directed and of cardinality  $< \kappa$ . By the inductive hypothesis, we can take the colimit of each restriction  $P_\lambda \rightarrow \mathcal{C}$  and then take the colimit of the resulting diagram indexed by  $\kappa$ , which exists by our assumption. This is a colimit of the original diagram by [13, Corollary 4.2.3.10].  $\square$

*Proof of Theorem 2.9.* By Proposition 2.12, it suffices to show that for a presheaf  $\mathcal{F}: F^{\text{op}} \rightarrow \mathcal{C}$  carrying finite disjoint joins to products, (3) of Theorem 2.8 holds if and only if  $\mathcal{F}$  carries arbitrary disjoint joins to products. The “only if” direction is clear since we can write disjoint joins as a filtered colimit of finite disjoint unions. Hence we prove the “if” direction.

By Proposition 2.14, it suffices to show that  $\mathcal{F}$  carries any colimit indexed by an ordinal  $\alpha$  to a limit. Let  $U_\beta$  be such a diagram with join  $U$ . Then we take

$$V_\beta = U_\beta \setminus \bigvee_{\gamma < \beta} U_\gamma.$$

<sup>1</sup>Beware a small error in the proof given there: With their construction,  $I_0 = \emptyset$  is not directed. We can fix this by reindexing.

This family is disjoint and has  $U$  as its join. We have a diagram

$$\mathcal{F}(U) \rightarrow \varprojlim_{\beta < \alpha} \mathcal{F}(U_\beta) \rightarrow \prod_{\beta < \alpha} \mathcal{F}(V_\beta)$$

and want to show that the first map is an equivalence. Since we know that the composite is an equivalence by assumption, it suffices to show that the second map is an equivalence. However, since

$$U_\beta = \bigvee_{\gamma \leq \beta} U_\gamma,$$

by using our assumption again, we can rewrite the second map as

$$\varprojlim_{\beta < \alpha} \prod_{\gamma \leq \beta} \mathcal{F}(V_\gamma) \rightarrow \prod_{\beta < \alpha} \mathcal{F}(V_\beta),$$

which is an equivalence.  $\square$

Theorem 2.9 is useful since it says that the sheaf condition can be checked on homotopy groups. Here we record some immediate consequences:

**Corollary 2.17.** *For a Boolean locale  $X$  and  $n \geq -2$ , the  $(n+1)$ -connective/ $n$ -truncated factorization system (see [13, Example 5.2.8.16]) for  $\mathrm{Shv}(X)$  can be computed pointwise.*

The following were also obtained in [14, Section A.4.1]:

**Corollary 2.18.** *For a Boolean locale  $X$ , the  $\infty$ -topos  $\mathrm{Shv}(X)$  is Postnikov complete.*

**Corollary 2.19.** *For a Boolean locale  $X$ , the final object of  $\mathrm{Shv}(X)$  is projective in the sense that any epimorphism  $\mathcal{F} \rightarrow *$  has a section.*

**2.3. Linear algebra on a Boolean locale.** In this section, we prove the following structure theorem:

**Theorem 2.20.** *For a Boolean locale  $X$  and a field  $k$ , any object of  $\mathrm{Shv}(X; \mathbf{D}(k))$  is (noncanonically) equivalent to a direct sum of objects of the form  $\Sigma^n k_U$  where  $n$  is an integer and  $U$  is an open of  $X$ . Here we write  $k_U$  for the extension by zero of the constant sheaf  $k$  on  $U$ .*

We first prove the following underived variant; note that  $\mathbf{D}(k)^\heartsuit$  denotes the 1-category of discrete  $k$ -vector spaces:

**Proposition 2.21.** *For a Boolean locale  $X$  and a field  $k$ , any object of the 1-category of discrete  $k$ -linear sheaves  $\mathrm{Shv}(X; \mathbf{D}(k)^\heartsuit)$  is (noncanonically) the direct sum of the sheaves of the form  $k_U$  for  $U \subset X$  an open.*

We need several lemmas to prove this:

**Lemma 2.22.** *Let  $X$  be a locale and  $k$  a field. Any subobject of  $k_X \in \mathrm{Shv}(X; \mathbf{D}(k)^\heartsuit)$  is of the form  $k_U \hookrightarrow k_X$  for a unique open  $U$ .*

*Proof.* Let  $\mathcal{F} \hookrightarrow k_X$  be an arbitrary subobject. In the category of set-valued sheaves  $\mathrm{Shv}(X; \mathbf{Set})$ , we base change this morphism along  $1: * \rightarrow k_X$  to obtain a monomorphism to  $*$ . It is written as  $*_U \hookrightarrow *$  for a unique open  $U$ . Hence we have a map  $*_U \rightarrow \mathcal{F}$  and also get a map  $k_U \rightarrow \mathcal{F}$  by adjunction. Therefore, we have a morphism of subobjects  $k_U \hookrightarrow \mathcal{F}$  of  $k_X$  that is an isomorphism after base change along  $1: * \rightarrow k_X$ . By  $k^\times$ -equivariance, this is an isomorphism after base change along  $i: * \rightarrow k_X$  for  $i \in k^\times$ . This is also an isomorphism after base change along  $0: * \rightarrow k_X$  as  $k_U \times_{k_X} * = *$ . Since the map  $\prod_{i \in k} * \rightarrow k_X$  is an isomorphism,  $k_U \hookrightarrow \mathcal{F}$  is an isomorphism. The uniqueness is clear.  $\square$

**Lemma 2.23.** *In the situation of Proposition 2.21, an epimorphism from  $k_X$  in  $\mathrm{Shv}(X; \mathbf{D}(k)^\heartsuit)$  is isomorphic to the projection  $k_X \simeq k_U \oplus k_{X \setminus U} \rightarrow k_U$  for a unique open  $U$ .*

*Proof.* This follows from Lemma 2.22 by taking the kernel of a given epimorphism.  $\square$

**Lemma 2.24.** *In the situation of Proposition 2.21, let  $\mathcal{F} \rightarrow \mathcal{F}'$  be a morphism in  $\mathrm{Shv}(X; \mathbf{D}(k)^\heartsuit)$ . It is an epimorphism if and only if the map  $\mathcal{F}(X) \rightarrow \mathcal{F}'(X)$  is surjective.*

*Proof.* We first prove the “if” direction. Then we can show that it is surjective on local sections by Theorem 2.9, since for any open  $U$ , the map  $\mathcal{F}(U) \rightarrow \mathcal{F}'(U)$  is a retract of

$$\mathcal{F}(X) \simeq \mathcal{F}(U) \oplus \mathcal{F}(X \setminus U) \rightarrow \mathcal{F}'(U) \oplus \mathcal{F}'(X \setminus U) \simeq \mathcal{F}'(X)$$

We prove the “only if” direction, which means that  $k_X$  is projective. Let  $\mathcal{F} \rightarrow k_X$  be an epimorphism in  $\text{Shv}(X; \mathbf{D}(k)^\heartsuit)$ . In  $\text{Shv}(X; \text{Set})$  we take a base change of this map along  $1: * \rightarrow k_X$  to get an epimorphism  $\mathcal{F}' \rightarrow *$ . By Corollary 2.19, we have a section  $* \rightarrow \mathcal{F}'$ . The map  $k_X \rightarrow \mathcal{F}$  corresponding to the composite  $* \rightarrow \mathcal{F}' \rightarrow \mathcal{F}$  by adjunction gives us a desired splitting.  $\square$

*Proof of Proposition 2.21.* Let  $\mathcal{F}$  be a sheaf. First, by recursion, we construct an open  $U_s$  and a map  $a_s: k_{U_s} \rightarrow \mathcal{F}$  for each section  $s \in \mathcal{F}(X)$ : We fix a well ordering on  $\mathcal{F}(X)$ . For each  $s \in \mathcal{F}(X)$ , let  $b_s: \bigoplus_{s' < s} k_{U_{s'}} \rightarrow \mathcal{F}$  be the map obtained by  $(a_{s'})_{s' < s}$ . Then the composite  $k_X \xrightarrow{s} \mathcal{F} \rightarrow \text{coker}(b_s)$  is written as the composite  $k_X \simeq k_U \oplus k_{X \setminus U} \twoheadrightarrow k_U \hookrightarrow \text{coker}(b_s)$  by Lemma 2.23. We declare  $U_s$  to be this open  $U$  and  $a_s$  to be the composite  $k_U \rightarrow k_U \oplus k_{X \setminus U} \simeq k_X \xrightarrow{s} \mathcal{F}$ .

We wish to show that the morphism  $c: \bigoplus_{s \in \mathcal{F}(X)} k_{U_s} \rightarrow \mathcal{F}$  induced by  $(a_s)_{s \in \mathcal{F}(X)}$  is an isomorphism. We first see that  $c$  is a monomorphism by induction: For each  $s$ , we show that the map  $\bigoplus_{s' \leq s} k_{U_{s'}} \rightarrow \mathcal{F}$  induced by  $a_{s'}$  and  $b_s$  is a monomorphism. By the inductive hypothesis,  $b_s$  is a monomorphism. Since  $k_U \xrightarrow{a_s} \mathcal{F} \rightarrow \text{coker}(b_s)$  is a monomorphism, so is the map induced by  $a_s$  and  $b_s$ . Then we show that  $c$  is an epimorphism. By Lemma 2.24, we need to see that for  $s \in \mathcal{F}(X)$ , the corresponding morphism  $k_X \rightarrow \mathcal{F}$  factors through  $c$ . Since  $b_s$  is a monomorphism, we obtain a map

$$k_{X \setminus U_s} \simeq \ker(k_X \rightarrow k_{U_s}) \rightarrow \ker(\mathcal{F} \rightarrow \text{coker}(b_s)) \simeq \bigoplus_{s' < s} k_{U_{s'}}.$$

Then the coproduct of  $a_s$  and this map gives a desired factorization.  $\square$

**Corollary 2.25.** *For a Boolean locale  $X$  and a field  $k$ , the Grothendieck abelian category  $\text{Shv}(X; \mathbf{D}(k)^\heartsuit)$  has homological dimension  $\leq 0$ ; i.e., every object is injective and projective.*

*Proof.* It suffices to show that every object is projective. By Proposition 2.21, it suffices to show that  $k_U$  for any open  $U$  is projective. Since this is a direct summand of  $k_X$ , it is projective by Lemma 2.24.  $\square$

*Proof of Theorem 2.20.* By Corollary 2.18, the  $\infty$ -category  $\text{Shv}(X; \mathbf{D}(k))$  is equivalent to the derived  $\infty$ -category of its heart. Therefore, by Corollary 2.25, any object  $\mathcal{F}$  is equivalent to  $\bigoplus_n \Sigma^n \pi_n \mathcal{F}$  and hence Proposition 2.21 gives the desired result.  $\square$

We also record the following:

**Corollary 2.26.** *For a Boolean locale  $X$  and a field  $k$ , the heart  $\text{Shv}(X; \mathbf{D}(k))^\heartsuit \subset \text{Shv}(X; \mathbf{D}(k))$  is closed under tensor product operations.*

*Proof.* By Proposition 2.21, it suffices to consider the tensor product of finite copies of the unit, which is again the unit.  $\square$

### 3. THE SMASHING SPECTRUM OF $\mathbf{D}(k)$ -VALUED SHEAVES

We first review facts we need from higher topos theory in Section 3.1. We then prove Theorems A and B in Sections 3.2 and 3.3, respectively.

**3.1. The existence of a Boolean cover.** We recall the following from [14, Corollary A.4.3.2]:

**Theorem 3.1** (Lurie). *For any hypercomplete  $\infty$ -topos  $\mathcal{X}$ , there is a Boolean locale  $X$  and a geometric morphism*

$$\text{Shv}(X) \rightarrow \mathcal{X}$$

*such that its inverse image functor is conservative.*



*Remark 3.2.* In the situation of Theorem 3.1, when  $\mathcal{X}$  is the hypercompletion of a 0-localic  $\infty$ -topos, the statement is simple: It says that any frame has a frame injection to a Boolean frame. This theorem was proven by Funayama [10]. His proof is quite involved because of its generality and a simpler proof in this case can be found, e.g., in [11, Corollary II.2.6]. When  $\mathcal{X}$  is the hypercompletion of a 1-localic  $\infty$ -topos, the statement was proven by Barr [6].

This theorem is useful to us through the following observation:

**Lemma 3.3.** *Let  $\mathcal{Y} \rightarrow \mathcal{X}$  be a map of  $\infty$ -toposes whose inverse image functor is conservative. Then the inverse image functor  $\mathrm{Shv}(\mathcal{X}; \mathcal{C}) \rightarrow \mathrm{Shv}(\mathcal{Y}; \mathcal{C})$  for a compactly generated  $\infty$ -category  $\mathcal{C}$  is conservative as well.*

*Proof.* Let  $\mathcal{C}_0$  denote the full subcategory of compact objects. The desired result follows from the identification of  $\mathrm{Shv}(-; \mathcal{C})$  with the  $\infty$ -category of the functors  $\mathcal{C}_0^{\mathrm{op}} \rightarrow -$  preserving finite limits.  $\square$

**3.2. The smashing spectrum of  $D(k)$ -valued sheaves.** In this section, we prove Theorem A using Theorem 3.1, whose usefulness can be seen from the following:

**Lemma 3.4.** *Let  $\mathcal{C} \rightarrow \mathcal{D}$  be a colimit-preserving symmetric monoidal functor between presentably symmetric monoidal  $\infty$ -categories.<sup>2</sup> If it is conservative, the induced map  $\mathrm{Sm}(\mathcal{D}) \rightarrow \mathrm{Sm}(\mathcal{C})$  of locales is an epimorphism.*

*Proof.* It induces a conservative frame map  $F \rightarrow G$ , where  $F$  and  $G$  are  $\mathrm{Sm}(\mathcal{C})$  and  $\mathrm{Sm}(\mathcal{D})$ , respectively. We wish to show that it is injective. Suppose that  $U$  and  $U' \in F$  are mapped to the same open. In that case,  $U \wedge U'$  is also mapped to the same open, but by conservativity, we get  $U = U \wedge U' = U'$ .  $\square$

We do not directly apply Lemma 3.4 in our proof of Theorem A. Instead, we use the following observations:

**Lemma 3.5.** *Let  $\mathcal{X}$  be an  $\infty$ -topos and  $k$  a field. For any  $E_1$ -algebra  $A$  in  $\mathrm{Shv}(\mathcal{X}; D(k))$ , there is a (unique) subterminal object  $U$  such that  $A|_{\mathcal{X}_{/U}}$  is zero and the unit  $k|_{\mathcal{X}_{\setminus U}} \rightarrow \pi_0 A|_{\mathcal{X}_{\setminus U}}$  is a monomorphism in  $\mathrm{Shv}(\mathcal{X}_{\setminus U}; D(k)^{\heartsuit})$ , where  $\mathcal{X}_{\setminus U}$  denotes the closed subtopos complementary to  $\mathcal{X}_{/U}$ .*

*Proof.* In  $\mathrm{Shv}(\mathcal{X}; \mathrm{Set})$ , we base change the monomorphism  $1: * \rightarrow \pi_0 A$  along  $0: * \rightarrow \pi_0 A$  to get a subterminal object, which corresponds to an open  $U$ . We wish to show that  $U$  satisfies the requirement. By the functoriality of this construction, it suffices to consider the cases  $U = *$  and  $U = \emptyset$ .

We first assume  $U = *$ . This means that  $* \xrightarrow{1} k \rightarrow \pi_0 A$  is zero, which implies  $A = 0$  as  $A$  is an  $E_1$ -algebra.

We then assume  $U = \emptyset$  and show that  $k \rightarrow \pi_0 A$  is a monomorphism. By considering the decomposition  $\coprod_{i \in k} * \simeq k$ , it suffices to show that the limit of  $* \xrightarrow{i} \pi_0 A \xleftarrow{i'} *$  is initial when  $i$  and  $i'$  are distinct elements of  $k$ . The assumption is the case  $(i, i') = (0, 1)$ . The other cases are also reduced to this case by considering the automorphism of  $\pi_0 A \in \mathrm{Shv}(\mathcal{X}; D(k)^{\heartsuit})$  given by  $(i - (-))(i - i')^{-1}$ .  $\square$

**Lemma 3.6.** *For a field  $k$  and an  $\infty$ -topos  $\mathcal{X}$ , suppose that  $e: k \rightarrow E$  is an idempotent object in  $\mathrm{Shv}(\mathcal{X}; D(k))$ . If  $\pi_0 e: k \rightarrow \pi_0 E$  is a monomorphism in  $\mathrm{Shv}(\mathcal{X}; D(k)^{\heartsuit})$ ,  $e$  is an equivalence.*

*Proof.* We first prove this under the assumption that  $\mathcal{X} = \mathrm{Shv}(X)$  holds for a Boolean locale  $X$ . By Theorem 2.20 and Corollary 2.25,  $e$  is an inclusion to the direct summand. Hence  $k_X$  is a retract of  $E$ . Since  $\mathrm{id}_E \otimes e$  is an equivalence, so is  $\mathrm{id}_{k_X} \otimes e \simeq e$ .

We consider the general case. By Theorem 3.1, there is a Boolean locale  $Y$  and a geometric morphism  $\mathrm{Shv}(Y) \rightarrow \mathcal{X}^{\mathrm{hyp}}$  whose inverse image is conservative. We write  $f$  for its composite with  $\mathcal{X}^{\mathrm{hyp}} \rightarrow \mathcal{X}$ . Then  $f^* e: k_Y \rightarrow f^* E$  is an idempotent object and hence an equivalence by

<sup>2</sup>Note that here we state this in the unstable setting; see [2] for the definition of  $\mathrm{Sm}$ .

the argument above. Therefore by Lemma 3.3, the composite  $k \xrightarrow{e} E \rightarrow E^{\text{hyp}}$  is an equivalence and thus  $e$  is a split monomorphism in  $\text{CAlg}(\text{Shv}(\mathcal{X}; D(k)))$ . Since  $e$  is also an epimorphism in  $\text{CAlg}(\text{Shv}(\mathcal{X}; D(k)))$ , it must be an equivalence.  $\square$

*Proof of Theorem A.* It is clear that  $\text{Sm}(\mathcal{X}) \rightarrow X$  is an epimorphism of locales. It suffices to show that it induces a surjection of frames. Let  $E$  be an idempotent algebra in  $\text{Shv}(\mathcal{X}; D(k))$ . By Lemma 3.5, we have a subterminal object  $U$  such that  $E|_{\mathcal{X}/U}$  is zero and  $E|_{\mathcal{X} \setminus U}$  satisfies the assumption of Lemma 3.6. By Lemma 3.6, as an idempotent algebra,  $E$  is equivalent to  $\text{cofib}(k_U \rightarrow k)$ .  $\square$

**3.3. Application: Tannaka duality.** We deduce a refined version of Theorem B from Theorem A. First, we explain its namesake:

*Remark 3.7.* Since the work of Lurie [12], Tannaka duality in algebraic geometry means a categorical reconstruction result of certain type. For example, Lurie showed in [14, Corollary 9.6.0.2] that the functor

$$\text{QCoh}: \{\text{quasicompact quasiseparated spectral algebraic spaces}\}^{\text{op}} \rightarrow \text{CAlg}(\text{Pr})$$

is fully faithful; see [14, Remark 9.0.0.5] for other references.

*Proof of Theorem B.* By [2], for a presentably symmetric monoidal  $\infty$ -category  $\mathcal{V}$ , we can compose  $\text{Sm}$  with the functor  $\mathcal{V} \otimes -: \text{CAlg}(\text{Pr}) \rightarrow \text{CAlg}_{\mathcal{V}}(\text{Pr})$  to obtain the adjunction

$$\text{Loc}^{\text{op}} \begin{array}{c} \xrightarrow{\text{Shv}(-; \mathcal{V})} \\ \xleftarrow{\text{Sm}} \end{array} \text{CAlg}_{\mathcal{V}}(\text{Pr}).$$

The desired statement is about the case  $\mathcal{V} = D(k)$ . Theorem A says that the unit is an equivalence so  $\text{Shv}(-; D(k))$  is fully faithful.  $\square$

Note that the 1-category  $\text{Loc}$  can be upgraded to a  $(1, 2)$ -category, i.e., a bicategory whose mapping categories are (essentially) posets. Our convention is that  $\text{Loc}(Y, X)$  denotes the full subcategory of  $\text{Fun}(F, G)$  spanned by frame morphisms where  $F$  and  $G$  denote the frames of  $X$  and  $Y$ , respectively. We can show that this is again fully faithful in the bicategorical sense:

**Theorem 3.8.** *For any field  $k$  and locales  $X$  and  $Y$ , the functor*

$$\text{Loc}(Y, X) \rightarrow \text{Fun}_{D(k)}^{\otimes}(\text{Shv}(X; D(k)), \text{Shv}(Y; D(k)))$$

*is an equivalence.*

*Proof.* Note that a functor of  $\infty$ -categories  $\mathcal{C} \rightarrow \mathcal{D}$  is an equivalence if and only if  $\text{Fun}(\Delta^1, \mathcal{C})^{\simeq} \rightarrow \text{Fun}(\Delta^1, \mathcal{D})^{\simeq}$  is an equivalence. By applying this to our situation, the desired result follows from Theorem B by considering the situation where  $X = X$  and  $Y$  is the locale corresponding to the frame  $\text{Fun}(\Delta^1, G)$  where  $G$  is the frame of  $Y$ .  $\square$

## REFERENCES

- [1] Jiří Adámek and Jiří Rosický. *Locally presentable and accessible categories*, volume 189 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1994.
- [2] Ko Aoki. The sheaves-spectrum adjunction, 2023. [arXiv:2302.04069v1](https://arxiv.org/abs/2302.04069v1).
- [3] M. Artin, A. Grothendieck, and J.-L. Verdier. *Théorie des topos et cohomologie étale des schémas. Tome 1: Théorie des topos*. Lecture Notes in Mathematics, Vol. 269. Springer-Verlag, Berlin-New York, 1972. Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4). with the collaboration of N. Bourbaki, P. Deligne, and B. Saint-Donat.
- [4] Scott Balchin and Greg Stevenson. Big categories, big spectra, 2023. [arXiv:2109.11934v4](https://arxiv.org/abs/2109.11934v4).
- [5] Paul Balmer and Giordano Favi. Generalized tensor idempotents and the telescope conjecture. *Proc. Lond. Math. Soc. (3)*, 102(6):1161–1185, 2011.
- [6] Michael Barr. Toposes without points. *J. Pure Appl. Algebra*, 5:265–280, 1974.
- [7] Silvana Bazzoni and Jan Št’ovíček. Smashing localizations of rings of weak global dimension at most one. *Adv. Math.*, 305:351–401, 2017.
- [8] Robert Burklund, Jeremy Hahn, Ishan Levy, and Tomer M. Schlank.  $K$ -theoretic counterexamples to Ravenel’s telescope conjecture, 2023. [arXiv:2310.17459v1](https://arxiv.org/abs/2310.17459v1).
- [9] Dustin Clausen and Akhil Mathew. Hyperdescent and étale  $K$ -theory. *Invent. Math.*, 225(3):981–1076, 2021.



- [10] Nenosuke Funayama. Imbedding infinitely distributive lattices completely isomorphically into Boolean algebras. *Nagoya Math. J.*, 15:71–81, 1959.
- [11] Peter T. Johnstone. *Stone spaces*, volume 3 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1982.
- [12] Jacob Lurie. Tannaka duality for geometric stacks, 2005. [arXiv:math/0412266v2](https://arxiv.org/abs/math/0412266v2).
- [13] Jacob Lurie. *Higher topos theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2009.
- [14] Jacob Lurie. Spectral algebraic geometry (under construction). Available at the author's website, 2018.
- [15] Amnon Neeman. The chromatic tower for  $D(R)$ . *Topology*, 31(3):519–532, 1992. With an appendix by Marcel Bökstedt.
- [16] Douglas C. Ravenel. Localization with respect to certain periodic homology theories. *Amer. J. Math.*, 106(2):351–414, 1984.
- [17] Robert M. Solovay. A model of set-theory in which every set of reals is Lebesgue measurable. *Ann. of Math. (2)*, 92:1–56, 1970.
- [18] John von Neumann. *Continuous geometry*. Princeton Mathematical Series, No. 25. Princeton University Press, Princeton, N.J., 1960. Foreword by Israel Halperin.

MAX PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, 53111 BONN, GERMANY  
Email address: aoki@mpim-bonn.mpg.de