

Generalized symmetry in non-Hermitian systems

Karin Sim,^{1,*} Nicolò Defenu,¹ Paolo Mognini,² and R. Chitra¹

¹*Institute for Theoretical Physics, ETH Zürich, 8093 Zurich, Switzerland*

²*Department of Physics, Stockholm University, AlbaNova University Center, 106 91 Stockholm, Sweden*

(Dated: June 11, 2024)

Despite acute interest in the dynamics of non-Hermitian systems, there is a lack of consensus in the mathematical formulation of non-Hermitian quantum mechanics in the community. Different methodologies are used in the literature to study non-Hermitian dynamics. This ranges from consistent frameworks like biorthogonal quantum mechanics and metric approach characterized by modified inner products, to normalization by time-dependent norms inspired by open quantum systems. In this work, we systematically explore the similarities and differences among these various methods. Utilizing illustrative models with exact solutions, we demonstrate that these methods produce not only quantitatively different results but also distinct physical interpretations. For dissipative systems where non-Hermiticity arises as an approximation, we find that the normalization method in the \mathcal{PT} -broken regime closely aligns with the full master equation solutions. In contrast, for quantum systems where non-Hermiticity can be engineered exactly, incorporating metric dynamics is crucial for the probabilistic interpretation of quantum mechanics, necessitating the generalization of unitary symmetry to non-Hermitian systems. This study lays the groundwork for further exploration of non-Hermitian Hamiltonians, potentially leveraging generalized symmetries for novel physical phenomena.

I. INTRODUCTION

Systems governed by non-Hermitian Hamiltonians [1–3] serve as an ideal playground to explore a wide range of unconventional behaviors. Examples include exceptional points [4], unconventional topology [5–7] and quantum phase transitions without gap closure [8]. Non-Hermiticity arises in many different contexts. For example, it provides an effective description of Lindbladian dynamics under certain conditions in dissipative systems [9] and continuously monitored systems [10, 11]. Non-Hermitian Hamiltonians can also be directly engineered in coupled qubit systems via Naimark dilation [12–14]. Recently, numerous experimental implementations of non-Hermitian systems have emerged, proving that they are far from merely theoretical endeavors. Examples include optical [15] and acoustic [16] systems, ultracold atoms [17] and quantum circuits [12].

Non-Hermitian Hamiltonians which commute with the \mathcal{PT} -operator often possess a real spectrum in the regime where the \mathcal{PT} -symmetry is not spontaneously broken [3, 18, 19]. \mathcal{PT} -symmetry [3, 20] refers to the combined operation of parity and time reversal symmetries. This has prompted the interpretation of this special class of non-Hermitian systems as a natural extension to conventional quantum mechanics [21]. When \mathcal{PT} -symmetry is spontaneously broken, exceptional points (EPs) [22] arise in the spectrum, where the eigenvectors of the Hamiltonian coalesce and the spectrum becomes gapless. Beyond the EP, the spectrum becomes complex. Exceptional points (EPs) have garnered significant recent attention from both theoretical [2, 4, 23] and experimental [24, 25] perspectives.

Dynamics of non-Hermitian systems is a new frontier of exploration which harbors a lot of potential to discover new physical paradigms. A plethora of novel phenomena, such as dynamically induced topological phases, exponentially-enhanced quantum sensing [26] and the divergence of Lieb-Robinson bounds [8, 27] have been reported recently. Despite its increasing popularity, there is a lack of consensus in the mathematical formulation of non-Hermitian quantum mechanics in the community. As non-Hermitian time evolution is intrinsically non-unitary, different methods have been adopted to obtain a normalized physical observable in non-Hermitian systems. Among them is biorthogonal quantum mechanics (BQM) [28], which is a comprehensive formulation of non-Hermitian quantum mechanics consistent with its probabilistic interpretation. BQM is widely adopted within the condensed matter community [29–31] and mainly applied to study systems featuring a time-independent Hamiltonian. The so-called metric framework [32–34] provides a generalization of BQM to the case with time-dependent Hamiltonians. In this framework, one considers a modified Hilbert space endowed by a time-dependent metric which appears as a weight factor in the inner product. The equation of motion of the metric incorporates the dynamics induced by both the Hamiltonian and its Hermitian conjugate. Both of these methods generate an exact time evolution of the non-Hermitian system, and are best suited when the non-Hermiticity of the system is realized in a fundamental manner instead of arising as an approximation. For example, this can be realized in NV centres via Naimark dilation [12].

On the other hand, a lot of the available literature on non-Hermitian dynamics [27, 35–39] skirts the important issue of non-conserved norms during time evolution and simply divides physical quantities by the time-dependent

* simkarin@phys.ethz.ch

norm of the time-evolved state. This approach is ubiquitous in the open quantum systems community and is motivated by the quantum trajectory theory [40–42]. Quantum trajectories are used to find the time evolution of open quantum systems [43], for example when the direct solution of Lindblad master equation is computationally costly. In certain scenarios such as the detection of photons emitted from a cavity, the evolution of the system can be approximated by that with an effective non-Hermitian Hamiltonian interrupted by quantum jumps at random times [40]. By post-selecting quantum trajectories which did not undergo quantum jumps, a non-Hermitian evolution can be realized and the time-evolved state has to be explicitly normalized by hand at every time step [35].

The ostensibly different methodologies lead to very different physics. A prominent example is the quantum Brachistochrone problem [44, 45]. In this problem, the normalization method predicts arbitrarily fast quantum evolutions using \mathcal{PT} -symmetry [44]. However, Ref. [45] shows, using the metric formalism, that it is impossible to achieve faster unitary evolutions using non-Hermitian Hamiltonians than those given by Hermitian Hamiltonians. Another example is our recent finding [46] that the metric formalism predicted non-zero defect density due to \mathcal{PT} -broken modes when crossing exceptional points adiabatically. This violation of adiabaticity was deeply linked to the symmetries of observables, an aspect absent in the by-hand-normalization methods.

These puzzling discrepancies motivate an in-depth comparative study of the different methodologies used to study non-Hermitian systems. In this work, we clarify the differences and similarities between the various methodologies used to study non-Hermitian dynamics and their physical impact. More importantly, we show that the consistent non-Hermitian quantum mechanical formulations, biorthogonal and metric, necessitate a generalization of the notion of unitary symmetries in non-Hermitian settings. Our work highlights the physical origin of the symmetry structure of observables seen in Ref. [46], which arises naturally in the metric formalism.

The paper is structured as follows: we first introduce the mathematical background of the different methodologies in Sec. II, then proceed in Sec. III to show the equivalence of BQM, pseudo-Hermiticity and the metric framework when the Hamiltonian is time-independent and has a real spectrum. In Sec. IV, we study the dynamics of open quantum systems, and compare the results of the metric framework, no-jump approximation and the full master equation. In Sec. V, we present the generalized unitary symmetry in non-Hermitian systems and elaborate on its physical consequence.

II. METHODOLOGIES

Here we briefly describe the different approaches that are used in the study of non-Hermitian systems. We

first present the consistent formulations of non-Hermitian quantum mechanics: biorthogonal and the metric framework, followed by the approaches used in the case of dissipative quantum systems.

A. Biorthogonal quantum mechanics (BQM)

We start with a brief summary of BQM [28] for a general static non-Hermitian Hamiltonian H with a non-degenerate spectrum. We assume a finite-dimensional Hilbert space throughout the paper. The infinite-dimensional case has been discussed in Ref. [28]. The left and right eigenstates of H are defined as

$$\begin{aligned} H|n\rangle_R &= E_n|n\rangle_R \\ H^\dagger|n\rangle_L &= \bar{E}_n|n\rangle_L \end{aligned} \quad (1)$$

where the sets $\{|n\rangle_R\}$ and $\{|n\rangle_L\}$ individually form a complete basis and are linearly independent, though not necessarily orthogonal [28]. The left and right eigenstates satisfy

$${}_L\langle m|n\rangle_R = \delta_{mn}. \quad (2)$$

The identity operator can now be expressed in terms of both left and right eigenstates as,

$$\sum_m |m\rangle_R {}_L\langle m| = \mathbb{1}. \quad (3)$$

Given a general state $|\psi\rangle$, we can decompose it in the right eigenbasis as

$$|\psi\rangle = \sum_n c_n |n\rangle_R, \quad (4)$$

where $c_n = {}_L\langle n|\psi\rangle$. For the non-Hermitian case, we introduce the *associated state*

$$|\tilde{\psi}\rangle := \sum_n c_n |n\rangle_L, \quad (5)$$

as expectation values of observables in the biorthogonal framework are defined as $\langle \hat{F} \rangle = \langle \tilde{\psi} | \hat{F} | \psi \rangle$. In particular, setting $\hat{F} = \mathbb{1}$, we see $\langle \tilde{\psi} | \psi \rangle = \sum_n |c_n|^2$. We note that this does not necessarily equate to one, as $\sum_n |n\rangle_L {}_L\langle n|$ is not necessarily equal to the identity.

The notion of Hermiticity changes in the context of BQM. This can be seen by considering the expansion of an observable \hat{F} in the biorthogonal basis

$$\hat{F} = \sum_{mn} f_{mn} |m\rangle_R {}_L\langle n|, \quad (6)$$

from which it can be seen that the realness of the expectation value $\langle \tilde{\psi} | \hat{F} | \psi \rangle = \langle \psi | \hat{F}^\dagger | \tilde{\psi} \rangle$ is only guaranteed if $f_{mn} = f_{nm}^*$; this defines *biorthogonally Hermitian* operators [28].

B. Metric framework

An alternative to the BQM formalism is the metric approach to non-Hermitian quantum mechanics [33, 34, 47, 48]. Here, we consider a modified Hilbert space endowed with a positive-definite “metric” operator $\rho(t)$, which appears as a weight factor in the inner product as $\langle \cdot, \cdot \rangle_{\rho(t)} := \langle \cdot | \rho(t) | \cdot \rangle$.

The time evolution of $\rho(t)$ follows from the requirement that the norm of the time-evolved state is conserved. Its equation of motion is given by

$$i\dot{\rho}(t) = H^\dagger(t)\rho(t) - \rho(t)H(t), \quad (7)$$

where the overdot denotes time derivative and $H(t)$ is a non-Hermitian Hamiltonian. Here we generalize to include the case of a time-dependent Hamiltonian, since this formalism can be applied to such a case as well. Additionally, dynamics is an integral part of this framework, distinguishing it from the earlier discussion of BQM which is more naturally applied to a stationary Hamiltonian. Using the solution to Eq. (7) and the square-root decomposition $\rho(t) = \eta^\dagger(t)\eta(t)$, we can map the system to an effective Hermitian Hamiltonian

$$h(t) = \eta(t)H(t)\eta^{-1}(t) + i\dot{\eta}(t)\eta^{-1}(t). \quad (8)$$

We assume $\eta(t)$ to be Hermitian, since all non-Hermitian square roots are related to the Hermitian root by similarity transformations; see e.g. Ref. [46] for a discussion on this. The Hamiltonian $h(t)$ acts in a different Hilbert space \mathcal{H} where the non-Hermiticity is encoded in the dynamics of $\eta(t)$.

Time evolution in the Hilbert spaces $\mathcal{H}_{\rho(t)}$ and \mathcal{H} is generated by the respective Hamiltonians, $H(t)$ and $h(t)$, via the time-dependent Schrödinger equation (TDSE)

$$\begin{aligned} i\frac{d}{dt}|\psi(t)\rangle &= H(t)|\psi(t)\rangle \\ i\frac{d}{dt}|\Psi(t)\rangle &= h(t)|\Psi(t)\rangle, \end{aligned} \quad (9)$$

where the states are related by $|\Psi(t)\rangle = \eta(t)|\psi(t)\rangle$. The construction of $\rho(t)$ guarantees that $h(t)$ is Hermitian, such that the unitarity of the time evolution is restored, giving $\langle \psi(t) | \rho(t) | \psi(t) \rangle = \langle \Psi(t) | \Psi(t) \rangle = 1$ at all times t .

The expectation value of an operator $\hat{o} : \mathcal{H} \rightarrow \mathcal{H}$ is given by

$$\begin{aligned} \langle \hat{O}(t) \rangle_{\text{metric}} &= \langle \Psi(t) | \hat{o} | \Psi(t) \rangle \\ &= \langle \psi(t) | \rho(t) \hat{O}(t) | \psi(t) \rangle, \end{aligned} \quad (10)$$

where $\hat{O}(t) : \mathcal{H}_{\rho(t)} \rightarrow \mathcal{H}_{\rho(t)}$ is defined as $\hat{O}(t) = \eta^{-1}(t)\hat{o}\eta(t)$.

C. Open quantum systems

As dissipation is a common source of non-Hermiticity in quantum systems[40–43], we provide a brief recap of

the theory underlying such systems. Within the Born-Markov approximations, the dynamics of an open quantum system is well-approximated by the Born-Markov master equation [49], which can be written in the Lindblad form as follows:

$$\begin{aligned} \dot{\rho}_{\text{me}} &= -i[H, \rho_{\text{me}}] + \sum_j \Gamma_j \rho_{\text{me}} \Gamma_j^\dagger \\ &\quad - \sum_j \frac{1}{2} \{ \rho_{\text{me}}, \Gamma_j^\dagger \Gamma_j \}. \end{aligned} \quad (11)$$

The master equation describes the time evolution of the system density operator ρ_{me} , which represents a probabilistic mixture of pure states. In Eq. (11), H is the system Hamiltonian and each Γ_j is a jump or Lindblad operator, which encodes the interaction between the system and the environment. Here the Hamiltonian H can be time-dependent, but we assume the Lindblad operators to be time-independent. The expectation value of an observable \hat{O} in such a system is defined as $\langle \hat{O} \rangle_{\text{ME}} = \text{Tr}(\rho_{\text{me}} \hat{O})$. For simplicity, we take the case with only one Lindblad operator, i.e. $\Gamma_j = \Gamma$ for $j = 1$ and 0 otherwise. The result is readily generalizable to the case with multiple Lindblad operators.

The direct solution of Eq. (11) is often computationally costly, and the quantum trajectory approach [41] is used as an alternative in such a case. The method entails computing the stochastic evolution of the individual pure states in the ensemble, which correspond to the states conditioned on the measurement outcomes. Numerically, this involves the evolution of a pure state with an effective non-Hermitian Hamiltonian $H_{\text{eff}} = H - \frac{i}{2}\Gamma^\dagger\Gamma$, interrupted by finite evolutions induced by one of the measurement operators at random times. These are known as quantum jumps [41]. The ensemble average of the projectors onto the pure states then give rise to the density operator ρ_{me} , and the expectation values of observables can also be calculated accordingly. The evolution of the individual pure state is discontinuous in time, unlike the continuous evolution of the density matrix which represents the ensemble average.

The so-called no-jump limit [50, 51] corresponds to the evolution of a dissipative system in the absence of quantum jumps. For example, by post-selecting quantum trajectories that did not undergo a jump, we can describe the evolution of the system purely in terms of H_{eff} . This is equivalent to rewriting Eq. (11) in the form

$$\dot{\rho}_{\text{me}} = -i(H_{\text{eff}}\rho_{\text{me}} - \rho_{\text{me}}H_{\text{eff}}^\dagger) + \Gamma\rho_{\text{me}}\Gamma^\dagger, \quad (12)$$

and neglecting the term $\Gamma\rho_{\text{me}}\Gamma^\dagger$, which is known as the recycling or quantum jump term. Curiously, the approximate master equation in the no-jump limit, $\dot{\rho}_{\text{nj}} = -i(H_{\text{eff}}\rho_{\text{nj}} - \rho_{\text{nj}}H_{\text{eff}}^\dagger)$ is identical to the metric equation of motion Eq. (7) with the substitution $H = H_{\text{eff}}^\dagger$. Note however, that the initial conditions are different. For example, a natural choice of initial condition for the metric is $\rho(0) = \mathbb{1}$, as is discussed in Sec.III and Ref. [46]. On

the other hand, a density matrix always take the form $\rho_{\text{nj}}(0) = \frac{1}{2}(\mathbb{1} + \mathbf{r} \cdot \boldsymbol{\sigma})$, $|\mathbf{r}| \leq 1$. Note that $\rho_{\text{nj}}(t)$ does not describe a mixed state and is not trace-preserving [50, 51]. In the no-jump limit, the expectation value of an observable is defined as $\langle \hat{O} \rangle_{\text{nj}} = \text{Tr}(\rho_{\text{nj}} \hat{O}) / \text{Tr}(\rho_{\text{nj}})$. If the initial state is a pure state, the solution to the no-jump master equation is given by $\rho_{\text{nj}}(t) = |\psi(t)\rangle\langle\psi(t)|$ where $i|\dot{\psi}(t)\rangle = H_{\text{eff}}|\psi(t)\rangle$. Consequently, [27, 35–39]:

$$\langle \hat{O}(t) \rangle_{\text{nj}} = \frac{\langle \psi(t) | \hat{O} | \psi(t) \rangle}{\langle \psi(t) | \psi(t) \rangle}, \quad (13)$$

which is the same as tracking the Schrödinger evolution of the pure state $|\psi(t)\rangle$ and normalizing quantities by the time-dependent norm, as was done in many works [27, 35–39]. To summarize, we have presented the different methodologies for handling non-Hermitian Hamiltonians: the fully self-consistent formulation of quantum mechanics encoded by both BQM and the metric, and the norm method used in the case of dissipative systems.

III. METRIC AS A GENERALIZATION OF BQM

In this section, we show that the metric framework is a generalization of the consistent non-Hermitian quantum mechanics encapsulated by BQM. We proceed by first highlighting the equivalence between BQM and a particular similarity transformation, known as pseudo-Hermiticity in the literature [52, 53]. This then motivates the metric framework as a natural generalization of BQM to a dynamical context. For clarity, we consider a general time-independent Hamiltonian H with a real, non-degenerate spectrum and an appropriately normalized biorthogonal basis fulfilling Eqs. (2) and (3).

It has been shown that every Hamiltonian with a real spectrum is pseudo-Hermitian [52]. The definition of pseudo-Hermiticity is the existence of the similarity transformation

$$H^\dagger = SHS^{-1}. \quad (14)$$

In the biorthogonal basis, we can write S and S^{-1} as

$$S = \sum_n |n\rangle_L \langle n|, \quad S^{-1} = \sum_n |n\rangle_R \langle n|, \quad (15)$$

such that $SS^{-1} = \mathbb{1}$ by Eqs. (2) and (3). We have $S|n\rangle_R = |n\rangle_L$ and $S^{-1}|n\rangle_L = |n\rangle_R$, and so $S|\psi\rangle = |\tilde{\psi}\rangle$ according to Eqs. (4) and (5). Furthermore, for any operator F , we have the equivalence

$$\langle \psi | SF | \psi \rangle = \langle \tilde{\psi} | F | \psi \rangle \quad (16)$$

and in particular $\langle \psi | S | \psi \rangle = \langle \tilde{\psi} | \psi \rangle = \sum_n |c_n|^2$.

We now introduce the Hermitian square root of S , which we denote \sqrt{S} , such that $S = \sqrt{S}^2$. As was discussed in Sec. IIB and Ref. [46], we assume this to be

Hermitian. Defining the state $|\Psi\rangle = \sqrt{S}|\psi\rangle$, we see that

$$\langle \Psi | F | \Psi \rangle = \langle \psi | S \tilde{F} | \psi \rangle = \langle \tilde{\psi} | \tilde{F} | \psi \rangle \quad (17)$$

where $\tilde{F} = \sqrt{S}^{-1} F \sqrt{S}$.

Equation (17) closely resembles the equivalence of the Hilbert spaces \mathcal{H} and $\mathcal{H}_{\rho(t)}$ in the metric framework, in the particular case where the metric is time-independent. This is clear when we identify $\sqrt{S} \rightarrow \eta$, $\tilde{F} \rightarrow H$ and $F \rightarrow h$ in Eqs. (7) and (8). In fact, the matrix S , which satisfies Eq. (14), is a time-independent solution to the metric equation of motion Eq. (7) with the initial condition $\rho(0) = S$, such that $\dot{\rho} = 0$. It is then evident that the notions in the metric framework are generalizations of the concepts discussed above, namely BQM and pseudo-Hermiticity, to time-dependent cases.

With respect to dynamics, we emphasize that the initial condition for the metric plays an important role in the time evolution of a non-Hermitian system. The metric equation of motion, Eq. (7), guarantees that $\frac{d}{dt} \langle \psi(t) | \rho(t) | \psi(t) \rangle = 0$, so $\langle \psi(t) | \rho(t) | \psi(t) \rangle = \langle \psi(0) | \rho(0) | \psi(0) \rangle$. This prompts the choice of initial condition $\rho(0) = \mathbb{1}$ [46], which ensures that $\langle \psi(t) | \rho(t) | \psi(t) \rangle = \langle \psi(0) | \psi(0) \rangle = 1$ given a normalized initial state $|\psi(0)\rangle$. Alternatively, the initial condition $\rho(0) = S$ gives rise to a stationary metric, as discussed earlier. With this choice, we would have $\langle \psi(t) | \rho(t) | \psi(t) \rangle = \langle \psi(t) | S | \psi(t) \rangle = \sum_n |c_n(t)|^2$. As discussed in Sec. IIA, this is not necessarily equal to one. Evidently, the dynamical nature of the metric, coupled with an appropriate initial condition, plays an important role in ensuring the subsequent normalization of the states. This is unsurprising, since the metric formalism encompasses a standard Schrödinger-like evolution of a closed quantum system at its core.

IV. DYNAMICS OF NON-HERMITIAN SYSTEMS

With a view to understanding the physical content of the different formalisms discussed earlier, we provide a comprehensive comparison of the dynamics of observables stemming from the metric, the no-jump limit in a dissipative system and the full master equation. We do so through illustrative models of non-Hermiticity engendered by dissipation in two-level systems. We consider two representative cases (i) $[\Gamma^\dagger \Gamma, H] = 0$ and (ii) $[\Gamma^\dagger \Gamma, H] \neq 0$, where H is the system Hamiltonian and Γ is the Lindblad operator present in Eq. (11). Note that H is Hermitian. We uncover behaviors that are sensitive to the underlying models and the initial conditions, but are starkly different in general.

A. $\Gamma^\dagger \Gamma$ commutes with system Hamiltonian

We consider a dissipative system with system Hamiltonian $H = \omega \sigma_z$ and jump operator $\Gamma = 2\sqrt{\gamma} \sigma_- =$

$\sqrt{\gamma}(\sigma_x - i\sigma_y)$, where $\omega, \gamma > 0$. For example, this corresponds physically to the decay process of a two-level system from the excited to the ground state. The effective non-Hermitian Hamiltonian, which appears in the no-jump master equation, is given by $H_{\text{eff}} = (\omega - i\gamma)\sigma_z$ up to a constant term proportional to the identity operator. We note that the energy eigenvalues are always complex in this model.

The metric dynamics can be obtained analytically by solving Eq. (7) with $H = H_{\text{eff}}$. The solution is given by $\rho(t) = \cosh(2\gamma t)\mathbb{1} + \sinh(2\gamma t)\sigma_z$. The Hermitian mapping defined by Eq. (8), which acts in the Hilbert space \mathcal{H} , is given by $h = \omega\sigma_z$. This is a special case in which the Hermitian Hamiltonian h , the system Hamiltonian and the effective non-Hermitian Hamiltonian share an eigenbasis, which is the σ_z basis denoted by $\{|0\rangle, |1\rangle\}$. For this model, the unique steady state solution of the full master equation is given by $\lim_{t \rightarrow \infty} \rho_{\text{me}} = \frac{1}{2}(\mathbb{1} - \sigma_z) = |0\rangle\langle 0|$.

When the system is initially in the ground state of the system Hamiltonian, the $|0\rangle$ state, $\langle \hat{O} \rangle_{\text{metric}} = \langle \hat{O} \rangle_{\text{me}} = \langle \hat{O} \rangle_{\text{nj}} = \langle 0|\hat{O}|0\rangle$ is trivially the steady-state solution of the full master equation for a general observable \hat{O} . If we instead have the initial state $|\psi(0)\rangle = |1\rangle$, we have $\langle \hat{O} \rangle_{\text{metric}} = \langle \hat{O} \rangle_{\text{nj}} = \langle 1|\hat{O}|1\rangle$ which stays in the excited state. On the other hand, the solution of the full master equation,

$$\begin{aligned} \langle \hat{O} \rangle_{\text{me}} &= \langle 0|\hat{O}|0\rangle(1 - e^{-4\gamma t}) + e^{-4\gamma t} \langle \hat{O} \rangle_{\text{metric}} \\ &= \langle 0|\hat{O}|0\rangle(1 - e^{-4\gamma t}) + e^{-4\gamma t} \langle \hat{O} \rangle_{\text{nj}} \end{aligned} \quad (18)$$

approaches the steady state solution in the long-time limit as expected.

However, for initial conditions that are superpositions of both the ground and excited states, we observe different behaviors. With the initial condition $|\psi(0)\rangle = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$, the metric observable oscillates indefinitely as

$$\begin{aligned} \langle \hat{O} \rangle_{\text{metric}} &= \frac{1}{2}\text{Tr}(\hat{O}) - \cos(2\omega t)\text{Re}(\langle 1|\hat{O}|0\rangle) \\ &\quad + \sin(2\omega t)\text{Im}(\langle 1|\hat{O}|0\rangle). \end{aligned} \quad (19)$$

On the other hand, the solution of the full master equation is given by

$$\begin{aligned} \langle \hat{O} \rangle_{\text{me}} &= \frac{1}{2}\text{Tr}(\hat{O}) - \cos(2\omega t)e^{-2\gamma t}\text{Re}(\langle 1|\hat{O}|0\rangle) \\ &\quad + \sin(2\omega t)e^{-2\gamma t}\text{Im}(\langle 1|\hat{O}|0\rangle) \\ &\quad + \frac{e^{-4\gamma t} - 1}{2} \left(\langle 1|\hat{O}|1\rangle - \langle 0|\hat{O}|0\rangle \right), \end{aligned} \quad (20)$$

which can be rewritten as

$$\begin{aligned} \langle \hat{O} \rangle_{\text{me}} - \frac{1}{2}\text{Tr}(\hat{O}) &= \left(\langle \hat{O} \rangle_{\text{metric}} - \frac{1}{2}\text{Tr}(\hat{O}) \right) e^{-2\gamma t} \\ &\quad + \frac{e^{-4\gamma t} - 1}{2} \left(\langle 1|\hat{O}|1\rangle - \langle 0|\hat{O}|0\rangle \right). \end{aligned} \quad (21)$$

Comparing Eqs. (18) and (21), it is evident that while the master equation solution approaches the steady state solution, the metric formalism predicts oscillatory behaviour.

The solution to the no-jump master equation, or equivalently the norm approach, yields

$$\begin{aligned} \langle \hat{O} \rangle_{\text{nj}} &= \frac{1}{2}\text{Tr}(\hat{O}) - \frac{2\cos(2\omega t)e^{-2\gamma t}}{1 + e^{-4\gamma t}}\text{Re}(\langle 1|\hat{O}|0\rangle) \\ &\quad + \frac{2\sin(2\omega t)e^{-2\gamma t}}{1 + e^{-4\gamma t}}\text{Im}(\langle 1|\hat{O}|0\rangle) \\ &\quad + \frac{e^{-4\gamma t} - 1}{2(e^{-4\gamma t} + 1)} \left(\langle 1|\hat{O}|1\rangle - \langle 0|\hat{O}|0\rangle \right). \end{aligned} \quad (22)$$

This is comparable to the full master equation solution, up to some factors proportional to $\frac{1}{1+e^{-4\gamma t}}$ for each term except for the $\frac{1}{2}\text{Tr}(\hat{O})$ term. The factors arise due to renormalization by the trace of the approximate density matrix ρ_{nj} . In contrast to the metric solution, it tends to an asymptotic value that is equal to the steady state solution in the long-time limit: $\lim_{t \rightarrow \infty} \langle \hat{O} \rangle_{\text{nj}} = \langle 0|\hat{O}|0\rangle$.

To summarize, the no-jump expectation value $\langle \hat{O} \rangle_{\text{nj}}$ tends to a constant value in asymptotically long time for general initial conditions away from the eigenbasis of H_{eff} . Even though $\langle \hat{O} \rangle_{\text{nj}}$ approaches the same value as the steady-state solution given by $\langle \hat{O} \rangle_{\text{me}}$, the short-time behaviors of the two expectation values only agree to zeroth order in time in general. On the other hand, the metric expectation value $\langle \hat{O} \rangle_{\text{metric}}$ is governed by the Hermitian mapping $h = \omega\sigma_z$ given by Eq. (8). Note that h is independent of the non-Hermitian coefficient γ . As a result, $\langle \hat{O} \rangle_{\text{metric}}$ does not tend to an asymptotic value in the long-time limit for this model, independent of the initial condition.

B. $\Gamma^\dagger\Gamma$ does not commute with system Hamiltonian

We now turn to a richer model in which the term $\Gamma^\dagger\Gamma$ does not commute with the system Hamiltonian. We set the system Hamiltonian to be $H = \omega\sigma_x$ and take the same jump operator as in the previous subsection, $\Gamma = 2\sqrt{\gamma}\sigma_- = \sqrt{\gamma}(\sigma_x - i\sigma_y)$ where $\omega, \gamma > 0$. For this model, the unique steady state solution of the master equation is given by $\lim_{t \rightarrow \infty} \rho_{\text{me}} = \frac{1}{2}(\mathbb{1} + \mathbf{r}_{ss} \cdot \boldsymbol{\sigma})$, where

$$r_{x,ss} = 0, \quad r_{y,ss} = \frac{2\gamma\omega}{2\gamma^2 + \omega^2}, \quad r_{z,ss} = -\frac{2\gamma^2}{2\gamma^2 + \omega^2}. \quad (23)$$

The effective non-Hermitian Hamiltonian is $H_{\text{eff}} = \omega\sigma_x - i\gamma\sigma_z$ up to a term proportional to the identity operator. The energy eigenvalues are $\pm\sqrt{\omega^2 - \gamma^2}$. There are two parameter regimes: \mathcal{PT} -symmetric regime where $\gamma^2 \leq \omega^2$ and a regime where the \mathcal{PT} -symmetry is spontaneously broken, $\gamma^2 > \omega^2$. In the regime $\gamma^2 > \omega^2$, the energy eigenvalues are purely imaginary and appear as complex conjugate pairs. The significance of the different




FIG. 1. The σ_z expectation value for the model discussed in Sec. IV B, with (a), (c): \mathcal{PT} -symmetric regime $\omega = 1.0, \gamma = 0.5$ and (b), (d): \mathcal{PT} -broken regime $\omega = 1.0, \gamma = 1.5$. The solid lines show the three different approaches discussed in text: metric (blue; labelled $\langle \sigma_z \rangle_{\text{metric}}$), full master equation (green; labelled $\langle \sigma_z \rangle_{\text{me}}$) and master equation in the no-jump limit (red; labelled $\langle \sigma_z \rangle_{\text{nj}}$). The dashed lines with the same colors show the corresponding asymptotic values/steady state solution shown in Eqs. (23), (24) and (25). The initial conditions are (a), (b): the ground state of the σ_z operator, and (c), (d): a random pure state $|\psi\rangle = (0.56 + 0.014i, -0.466 - 0.685i)^T$. The expectation values calculated with the no-jump approximation and the metric framework show parameter-dependent behavior. They are oscillatory and do not tend to a constant value for the parameter regime $\omega^2 > \gamma^2$ ((a), (c)). For $\omega^2 < \gamma^2$ where the energy eigenvalues of H_{eff} are complex ((b), (d)), the solutions approach different asymptotic values. The asymptotic value of the metric solution depends on the initial condition, which leads to a discrepancy in its relative value compared to the steady state solution of the full master equation, as can be seen from (b) and (d). The qualitative behaviors of the no-jump solution and the full master equation solution are similar.

parameter regimes stems from the fact that the effective Hamiltonian H_{eff} now features an exceptional point at $\omega = \gamma$. The solutions obtained by the no-jump approximation and the metric framework display parameter-dependent behaviors: in the \mathcal{PT} -broken regime $\omega^2 < \gamma^2$, the solutions tend to a constant value in the limit $t \rightarrow \infty$, whereas in the \mathcal{PT} -symmetric regime $\omega^2 > \gamma^2$, both methods yield oscillatory solutions. The only qualitative difference is that the metric expectation value displays periodic oscillations, while the oscillations of no-jump expectation value are aperiodic.

This parameter-dependent behavior is observed for various initial states as shown in Fig. 1. For the no-jump approximation, the asymptotic values valid for $\omega^2 < \gamma^2$ are given by

$$\begin{aligned} \lim_{t \rightarrow \infty} \langle \sigma_x \rangle_{\text{nj}} &= 0, & \lim_{t \rightarrow \infty} \langle \sigma_y \rangle_{\text{nj}} &= \frac{\omega}{\gamma} \\ \lim_{t \rightarrow \infty} \langle \sigma_z \rangle_{\text{nj}} &= -\frac{\sqrt{\gamma^2 - \omega^2}}{\gamma}. \end{aligned} \quad (24)$$

They are independent of initial conditions, since division by the time-dependent norm cancels out the contribution of the initial condition. An exception is when the initial state is an eigenbasis of H_{eff} , where the expectation values do not evolve in time. We note that the qualitative behaviors of $\langle \sigma_z \rangle_{\text{nj}}$ and $\langle \sigma_z \rangle_{\text{me}}$ are similar when $\omega^2 < \gamma^2$,

though the asymptotic values are different. This can be seen from Figs. 1(b) and (d).

For the metric solutions, the asymptotic values obtained for the case $\omega^2 < \gamma^2$ are given by

$$\begin{aligned} \lim_{t \rightarrow \infty} \langle \sigma_x \rangle_{\text{metric}} &= x_0, \\ \lim_{t \rightarrow \infty} \langle \sigma_y \rangle_{\text{metric}} &= \frac{-2\omega\sqrt{\gamma^2 - \omega^2}z_0 + (\gamma^2 - 2\omega^2)y_0}{\gamma^2}, \\ \lim_{t \rightarrow \infty} \langle \sigma_z \rangle_{\text{metric}} &= \frac{2\omega\sqrt{\gamma^2 - \omega^2}y_0 + (\gamma^2 - 2\omega^2)z_0}{\gamma^2}. \end{aligned} \quad (25)$$

Unlike in the no-jump case, they depend on the initial conditions: x_0 , y_0 and z_0 are the initial values of the spin expectation in each direction. The sensitivity to initial condition is typical of Schrödinger evolution of states, since the information of the initial state is encoded in the time-evolved state and thus in the expectation value. This emphasizes the fact that the metric framework tracks the Schrödinger dynamics of an intrinsically non-Hermitian system. The steady state solution of the Markovian master equation is independent of the initial condition in general, since the presence of stochastic quantum jumps wipes out the information of the initial state in the long-time limit. Curiously, the no-jump approximation erases information about the initial state,

though it describes the Hamiltonian dynamics of a pure state.

Unsurprisingly, the dependence of the metric solution on initial condition leads to its varying degrees of discrepancy with the full master equation solutions. For example, Fig. 1(b) shows a large discrepancy between $\langle \sigma_z \rangle_{\text{metric}}$ and $\langle \sigma_z \rangle_{\text{me}}$ in the long-time limit. However, this discrepancy is smaller in Fig. 1(d), where the asymptotic value of $\langle \sigma_z \rangle_{\text{metric}}$ is even closer to the steady state solution of the master equation than the no-jump solution.

To summarize, the full master equation, the no-jump approximation and the metric approaches all display different behaviors in general, despite being model and parameter-specific. For the simpler model in Sec. IV A, the metric solution is oscillatory in general and does not tend to an asymptotic value in the long-time limit. On the other hand, the qualitative behavior of the no-jump solution resembles the master equation solution. In the long-time limit, the asymptotic value of the no-jump solution coincides with the steady state solution of the master equation solution. The role of the exceptional points in the dynamics is evident in the more complicated model discussed in Sec. IV B. In the parameter regime where \mathcal{PT} -symmetry is preserved, there are common qualitative features between the metric and no-jump expectation values, which is oscillatory behavior. They only tend to an asymptotic value, resembling the steady state solution, in the parameter regime where \mathcal{PT} -symmetry is spontaneously broken. These asymptotic values generally do not coincide with the steady state solution of the master equation, though the no-jump approximation reproduces the qualitative behavior of the master equation albeit only in the \mathcal{PT} -broken regime. We also found that the asymptotic values of the metric expectation values are sensitive to initial conditions. This sensitivity can be attributed to Schrödinger evolution with an intrinsically non-Hermitian Hamiltonian, in contrast to the dissipative dynamics described by the full master equation. These observations motivates further study of whether there are a class of initial states where the metric formalisms leads to observables which mimic steady state values. Our findings thus highlight the importance of considering the physical origin of non-Hermiticity when studying the dynamics of non-Hermitian systems.

V. GENERALIZED UNITARY SYMMETRY

In this section, we further explore the deep contrast between the consistent formalism of non-Hermitian quantum mechanics, embodied by BQM and metric formalisms, and the oft-used no-jump approximation valid for dissipative systems. Specifically, we focus on aspects of symmetry. We show that the consistent formulation of non-Hermitian quantum mechanics leads to a generalization of the conventional notions of symmetry associated with the Hermitian case. Using a toy model, we

show that the generalization of a discrete symmetry in the model leads to important symmetry constraints for observables. We contrast this with the no-jump formalism where such constraints are absent.

A first point of significance is that the left and right eigenvectors of a non-Hermitian Hamiltonian do not coincide, unlike the Hermitian case. Therefore, the construction of symmetry operators should be generalized to incorporate this aspect. The biorthogonal basis provides a natural path to this generalization. The spectral decomposition for a general non-Hermitian Hamiltonian, parameterized by a parameter k , acting on a finite-dimensional Hilbert space reads as follows:

$$H_k = \sum_n E_{k,n} |k, n\rangle_R \langle k, n|_L. \quad (26)$$

For such a Hamiltonian, we can define the operators

$$\begin{aligned} U_{R,k} &= \sum_n |-k, n\rangle_R \langle k, n|_L \\ U_{L,k} &= \sum_n |-k, n\rangle_L \langle k, n|_R \end{aligned} \quad (27)$$

which satisfy $U_{L,k} U_{R,k}^\dagger = U_{L,k}^\dagger U_{R,k} = 1$. This is the generalization of the notion of unitary transformations to a biorthogonal basis. We note that this is the first time such a generalization in the biorthogonal basis has been made, to the best of our knowledge.

These operators define the mappings between k to $-k$ states, which are given by $U_{R/L,k} |k, n\rangle_{R/L} = |-k, n\rangle_{R/L}$. If we impose the requirement that the spectrum is even with respect to k , such that $E_{k,n} = E_{-k,n}$ in Eq. (26), we have

$$\begin{aligned} U_{R,k} H_k U_{R,k}^{-1} &= U_{R,k} H_k U_{L,k}^\dagger = H_{-k} \\ U_{L,k} H_k^\dagger U_{L,k}^{-1} &= U_{L,k} H_k^\dagger U_{R,k}^\dagger = H_{-k}^\dagger. \end{aligned} \quad (28)$$

This symmetry mapping of the Hamiltonian reflects the even parity of the spectrum of H_k .

We note that a similar notion also arises naturally within the metric framework. To this end, we can define a unitary operator similar to Eq. (27) using the spectrum of the Hermitian mapping h_k , given by Eq. (8). Writing $h_k = \sum_n e_{k,n} |k, n\rangle \langle k, n|$, the symmetry is defined as

$$U_k = \sum_n |-k, n\rangle \langle k, n|. \quad (29)$$

It then follows that the even parity of the spectrum $e_k = e_{-k}$ translates directly to the symmetry $U_k h_k U_k^\dagger = h_{-k}$. This shows that the generalized unitary symmetry is already embedded in the metric framework, and we can access it directly in the observables without the need to consider BQM.

To further explore this notion of a generalized symmetry, we consider the concrete model of two level systems studied in Ref. [46],

$$H_k = k\sigma_x + i\gamma(t)\sigma_y + \Delta(t)\sigma_z, \quad (30)$$

where k denotes the momentum, as was considered in Ref. [46]. This Hamiltonian does not necessarily have to satisfy Eq. (14). In general, the coefficients γ and Δ in Eq. (30) can be time dependent, but we omit the time dependence hereafter for the ease of notation.

Ref. [46] showed that the metric dynamics led to qualitatively different symmetry structures of spin expectation values. This was shown to have important physical consequences compared to the straightforward renormalization of the no-jump formalism. Here we show that this symmetry structure is a direct consequence of the generalized symmetry discussed above. To this end, we consider the left and right instantaneous eigenstates of Eq (30), which are denoted $|k, \pm\rangle_{L/R}$, as in Eq. (1). They are normalized such that they obey Eqs. (2) and (3). The corresponding eigenvalues are $E_k = \pm\sqrt{k^2 - \gamma^2 + \Delta^2}$, such that they satisfy the requirement for Eq. (27). The spectrum is real in the parameter regime $k^2 + \Delta^2 \geq \gamma^2$ where \mathcal{PT} -symmetry is not spontaneously broken, and form complex conjugate pairs otherwise. The exact forms of $U_{L,k}$ and $U_{R,k}$ for our model are given by $U_{R,k} = U_{R,k}^\dagger = \frac{1}{\gamma+k}(\gamma\mathbb{1} - k\sigma_z)$ and $U_{L,k} = U_{L,k}^\dagger = \frac{1}{\gamma-k}(\gamma\mathbb{1} + k\sigma_z)$, such that Eq. (28) is satisfied. For this model, Eq. (29) greatly simplifies, giving $U_k = \sigma_z$. This implies the symmetry $\sigma_z h_k(t) \sigma_z = h_{-k}(t)$.

In the metric framework, the expectation values are calculated using Eq. (10). Note that the normalized time-evolved state $|\Psi(t)\rangle_k$, as defined by Eq. (9), can be represented in the instantaneous eigenstates of $h_k(t)$ as $|\Psi(t)\rangle_k = \sum_{n \in \{\pm\}} c_{n,k}(t) |n(t)\rangle_k$. The aforementioned symmetry $\sigma_z h_k(t) \sigma_z = h_{-k}(t)$ guarantees that the instantaneous eigenstates transform as $\sigma_z |n(t)\rangle_k \propto |n(t)\rangle_{-k}$. Additionally, for the choice of initial condition in Ref. [46] ($|\Psi(0)\rangle_k = (1, 0)^T$ for all k), the time-dependent coefficients of the time-evolved state are even in k , i.e. $c_{n,k}(t) = c_{n,-k}(t)$ for all n at all times. Therefore, with this initial condition, the time-evolved state also transforms as $\sigma_z |\Psi(t)\rangle_k = |\Psi(t)\rangle_{-k}$.

Introducing the pure-state density matrix of the time-evolved state $\hat{\Psi}_k(t) \equiv |\Psi(t)\rangle_k \langle\Psi(t)|_k$, it is easy to verify the symmetry properties of the spin expectation values with respect to $k \rightarrow -k$. The density matrix is given by

$$\hat{\Psi}_k(t) = \frac{1}{2} \left(\mathbb{1} + \sum_{i=x,y,z} \langle\sigma_i(t)\rangle_k \sigma_i \right) \quad (31)$$

where $\langle\sigma_i(t)\rangle_k = \text{Tr}(\hat{\Psi}_k(t) \sigma_i)$. Since the density matrix transforms as $\sigma_z \hat{\Psi}_k \sigma_z = \hat{\Psi}_{-k}$ and $\sigma_z \hat{\Psi}_k \sigma_z = \frac{1}{2} \left(\mathbb{1} + \langle\sigma_z(t)\rangle_k \sigma_z - \sum_{i=x,y} \langle\sigma_i(t)\rangle_k \sigma_i \right)$, we see that $\langle\sigma_i(t)\rangle_k$ is even in k for $i = z$ and odd in k for $i = x, y$.

This symmetry structure of the observables is also visible in the BQM formalism. We expand Eq. (10) for a general observable \hat{F} in the instantaneous biorthogonal

basis

$$\begin{aligned} \langle\hat{F}\rangle_k &= \text{Tr}(\hat{\Psi}_k(t) \hat{F}) \\ &= \sum_{ij \in \{\pm\}} \langle j | \hat{\Psi}_k(t) | i \rangle_R \langle i | \hat{F} | j \rangle_R \end{aligned} \quad (32)$$

where the time dependence in the basis has been omitted for notational ease. Since Eq. (27) implies that $|\pm\rangle_{R/L, -k} = U_{R/L, k} |\pm\rangle_{R/L, k}$, we have

$$\begin{aligned} \langle\hat{F}\rangle_{-k} &= \sum_{m, n \in \{\pm\}} \langle m | U_{L, k} \sigma_z \hat{\Psi}_k \sigma_z U_{R, k} | n \rangle_{R, k} \\ &\quad \times \langle n | U_{L, k} \hat{F} U_{R, k} | m \rangle_{R, k}. \end{aligned} \quad (33)$$

The first factor is independent of the observable \hat{F} , and can be recast as $U_{L, k} \sigma_z \hat{\Psi}_k \sigma_z U_{R, k} = \phi_{x, k} \sigma_x + \phi_{y, k} \sigma_y + \hat{\Psi}_k$. We can also rewrite the second term as $\langle n | U_{L, k} \hat{F} U_{R, k} | m \rangle_{R, k} \equiv c_{nm}^F \langle n | \hat{F} | m \rangle_{R, k}$, where the exact forms of the coefficients $\phi_{x/y, k}$ and c_{nm}^F for $\hat{F} \in \{\sigma_x, \sigma_y, \sigma_z\}$ are summarized in Appendix B. Combining these results, we obtain

$$\begin{aligned} \langle F \rangle_{-k} &= \sum_{m, n \in \{\pm\}} \langle m | \phi_{x, k} \sigma_x + \phi_{y, k} \sigma_y + \hat{\Psi}_k | n \rangle_{R, k} \\ &\quad \times \langle n | F | m \rangle_{R, k} c_{nm}^F. \end{aligned} \quad (34)$$

Evaluating Eq. (34) for $\hat{F} = \sigma_z$ is algebraically simple, since the coefficient $c_{nm}^{\sigma_z} = 1$ for all pairs. We thus have

$$\langle\sigma_z\rangle_{-k} = \text{Tr} \left([\phi_{x, k} \sigma_x + \phi_{y, k} \sigma_y + \hat{\Psi}_k] \sigma_z \right) = \langle\sigma_z\rangle_k. \quad (35)$$

For the cases $F = \sigma_x, \sigma_y$, since the off-diagonal coefficients are non-trivial, a more complex calculation yields $\langle\sigma_i\rangle_{-k} = -\langle\sigma_i\rangle_k$ for $i = x, y$. In summary, we have introduced the generalization of unitary symmetries using the biorthogonal basis, and showed how a generalized parity symmetry in the non-Hermitian Hamiltonian directly manifests in the metric formalism with important consequences for observables.

VI. CONCLUSION

We have demonstrated the similarities and differences among the various methods available in the literature to study non-Hermitian systems. By employing illustrative models with exact solutions, we have shown that these methods yield not only quantitatively different results but also distinct physical interpretations. The physical origin of non-Hermiticity plays a crucial role in these differences.

Our findings indicate that when non-Hermiticity arises from dissipation in the limit where quantum jumps can be neglected, the normalization method is more appropriate, especially in the \mathcal{PT} -broken regime. This method closely reproduces the qualitative behavior of the full

master equation in this regime. Conversely, when non-Hermiticity is directly engineered in the system, such as by Naimark dilation, it is essential to consider the metric dynamics for a consistent quantum treatment. This approach can reveal richer physics, including the manifestation of a generalized symmetry. Notably, our findings align with those of Ref. [54], which highlights starkly different behaviors obtained by BQM and explicit normalization in the context of dynamical phase transitions in non-Hermitian systems.

Our work paves the way for deeper explorations of truly non-Hermitian Hamiltonians, where the physics of the metric and the concept of generalized symmetries could potentially be harnessed for exotic behaviors. An exciting future direction is the extension of this framework to include other discrete and continuous symmetries. This paves the way for the exploration of new paradigms for quantum criticality, spontaneous symmetry breaking and plausibly new classification schemes based on generalized symmetries for topological states.

ACKNOWLEDGMENTS

This research was funded by the Swiss National Science Foundation (SNSF) grant number 200021 207537, by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy EXC2181/1-390900948 (the Heidelberg STRUCTURES Excellence Cluster), the Swiss State Secretariat for Education, Research and Innovation (SERI), and by the Swedish Research Council (2018-00313) and Knut and Alice Wallenberg Foundation (KAW) via the project Dynamic Quantum Matter (2019.0068). The authors would like to thank G. M. Graf for numerous fruitful discussions.

Appendix A: Exact solution for the metric approach

In Sec. IV B, we discuss the case in which the jump operator $\Gamma^\dagger\Gamma$ does not commute with the system Hamiltonian. This leads to a rich dynamics which show qualitative differences using the different methods.

For the non-Hermitian Hamiltonian $H_{\text{eff}} = \omega\sigma_x - i\gamma\sigma_z$, the metric is given by

$$\rho(t) = \frac{\omega^2 - \gamma^2 \cos(2\sqrt{\omega^2 - \gamma^2}t)}{\omega^2 - \gamma^2} \mathbb{1} - \frac{2\gamma\omega \sin^2(\sqrt{\omega^2 - \gamma^2}t)}{\omega^2 - \gamma^2} \sigma_y + \frac{\gamma \sin(2\sqrt{\omega^2 - \gamma^2}t)}{\sqrt{\omega^2 - \gamma^2}} \sigma_z. \quad (\text{A1})$$

This gives rise to the Hermitian mapping

$$h(t) = \left(2\omega + 2\gamma \frac{\rho_y(t)}{1 + \rho_0(t)} \right) \sigma_x \quad (\text{A2})$$

given by Eq. (8).

Appendix B: Generalized unitary symmetry

In Sec. V, we use the result $U_{L,k}\sigma_z\hat{\Psi}_k\sigma_zU_{R,k} = \phi_{x,k}\sigma_x + \phi_{y,k}\sigma_y + \hat{\Psi}_k$ to arrive at Eq. (34). Here we provide the exact form of the coefficients:

$$\begin{aligned} \phi_{x,k} &= \frac{\gamma}{k^2 - \gamma^2} (\gamma\langle\sigma_x\rangle_k - ik\langle\sigma_y\rangle_k), \\ \phi_{y,k} &= \frac{\gamma}{k^2 - \gamma^2} (\gamma\langle\sigma_y\rangle_k + ik\langle\sigma_x\rangle_k). \end{aligned} \quad (\text{B1})$$

We also rewrite the second term in Eq. (33) as ${}_{L,k}\langle n|U_{L,k}FU_{R,k}|m\rangle_{R,k} \equiv c_{nm}^F {}_{L,k}\langle n|F|m\rangle_{R,k}$. The coefficients c_{nm}^F for $F \in \{\sigma_x, \sigma_y, \sigma_z\}$ and the pairs $\{n, m\}$ are summarised in the table below.

F	$\{+, +\}$	$\{-, -\}$	$\{+, -\}$	$\{-, +\}$
σ_z	1	1	1	1
σ_x	-1	-1	$\frac{\gamma E_k + k\Delta}{\gamma E_k - k\Delta}$	$\frac{\gamma E_k - k\Delta}{\gamma E_k + k\Delta}$
σ_y	1	1	$\frac{\gamma\Delta + kE_k}{\gamma\Delta - kE_k}$	$\frac{\gamma\Delta - kE_k}{\gamma\Delta + kE_k}$

-
- [1] Y. Ashida, Z. Gong, and M. Ueda, Non-hermitian physics, *Advances in Physics* **69**, 249 (2020).
[2] E. J. Bergholtz, J. C. Budich, and F. K. Kunst, Exceptional topology of non-hermitian systems, *Rev. Mod. Phys.* **93**, 015005 (2021).
[3] C. M. Bender, Making sense of non-hermitian hamiltonians, *Reports on Progress in Physics* **70**, 947 (2007).
[4] S. Sayyad and F. K. Kunst, Realizing exceptional points of any order in the presence of symmetry, *Phys. Rev. Res.* **4**, 023130 (2022).
[5] Z. Gong, Y. Ashida, K. Kawabata, K. Takasan, S. Higashikawa, and M. Ueda, Topological phases of non-

- hermitian systems, *Phys. Rev. X* **8**, 031079 (2018).
[6] F. K. Kunst, E. Edvardsson, J. C. Budich, and E. J. Bergholtz, Biorthogonal bulk-boundary correspondence in non-hermitian systems, *Phys. Rev. Lett.* **121**, 026808 (2018).
[7] F. Roccati, M. Bello, Z. Gong, M. Ueda, F. Ciccarello, A. Chenu, and A. Carollo, Hermitian and non-hermitian topology from photon-mediated interactions (2023), arXiv:2303.00762 [quant-ph].
[8] N. Matsumoto, K. Kawabata, Y. Ashida, S. Furukawa, and M. Ueda, Continuous phase transition without gap closing in non-hermitian quantum many-body systems,

- Phys. Rev. Lett. **125**, 260601 (2020).
- [9] N. Shibata and H. Katsura, Dissipative spin chain as a non-hermitian kitaev ladder, Phys. Rev. B **99**, 174303 (2019).
- [10] T. Müller, S. Diehl, and M. Buchhold, Measurement-induced dark state phase transitions in long-ranged fermion systems, Phys. Rev. Lett. **128**, 010605 (2022).
- [11] M. Buchhold, Y. Minoguchi, A. Altland, and S. Diehl, Effective theory for the measurement-induced phase transition of dirac fermions, Phys. Rev. X **11**, 041004 (2021).
- [12] Y. Wu, W. Liu, J. Geng, X. Song, X. Ye, C.-K. Duan, X. Rong, and J. Du, Observation of parity-time symmetry breaking in a single-spin system, Science **364**, 878 (2019), <https://www.science.org/doi/pdf/10.1126/science.aaw8205>.
- [13] Y.-C. Chen, M. Gong, P. Xue, H.-D. Yuan, and C.-J. Zhang, Quantum deleting and cloning in a pseudo-unitary system, Frontiers of Physics **16**, 53601 (2021).
- [14] C. Zheng, L. Hao, and G. L. Long, Observation of a fast evolution in a parity-time-symmetric system, Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences **371**, 20120053 (2013), <https://royalsocietypublishing.org/doi/pdf/10.1098/rsta.2012.0053>.
- [15] J. M. Zeuner, M. C. Rechtsman, Y. Plotnik, Y. Lumer, S. Nolte, M. S. Rudner, M. Segev, and A. Szameit, Observation of a topological transition in the bulk of a non-hermitian system, Phys. Rev. Lett. **115**, 040402 (2015).
- [16] X. Zhang, Y. Tian, J.-H. Jiang, M.-H. Lu, and Y.-F. Chen, Observation of higher-order non-hermitian skin effect, Nature Communications **12**, 5377 (2021).
- [17] Q. Liang, D. Xie, Z. Dong, H. Li, H. Li, B. Gadway, W. Yi, and B. Yan, Dynamic signatures of non-hermitian skin effect and topology in ultracold atoms, Phys. Rev. Lett. **129**, 070401 (2022).
- [18] C. M. Bender and S. Boettcher, Real spectra in non-hermitian hamiltonians having \mathcal{PT} symmetry, Phys. Rev. Lett. **80**, 5243 (1998).
- [19] C. M. Bender and D. W. Hook, Pt-symmetric quantum mechanics (2023), arXiv:2312.17386 [quant-ph].
- [20] C. M. Bender, S. Boettcher, and P. N. Meisinger, \mathcal{PT} -symmetric quantum mechanics, Journal of Mathematical Physics **40**, 2201 (1999), https://pubs.aip.org/aip/jmp/article-pdf/40/5/2201/19231175/2201.1_online.pdf.
- [21] J. Gong and Q.-h. Wang, Time-dependent \mathcal{PT} -symmetric quantum mechanics, Journal of Physics A: Mathematical and Theoretical **46**, 485302 (2013).
- [22] W. D. Heiss, The physics of exceptional points, Journal of Physics A: Mathematical and Theoretical **45**, 444016 (2012).
- [23] C. H. Lee, Exceptional bound states and negative entanglement entropy, Phys. Rev. Lett. **128**, 010402 (2022).
- [24] L. Ding, K. Shi, Q. Zhang, D. Shen, X. Zhang, and W. Zhang, Experimental determination of \mathcal{PT} -symmetric exceptional points in a single trapped ion, Phys. Rev. Lett. **126**, 083604 (2021).
- [25] W. Chen, Ş. Kaya Özdemir, G. Zhao, J. Wiersig, and L. Yang, Exceptional points enhance sensing in an optical microcavity, Nature **548**, 192 (2017).
- [26] J. C. Budich and E. J. Bergholtz, Non-hermitian topological sensors, Phys. Rev. Lett. **125**, 180403 (2020).
- [27] B. Dóra and C. P. Moca, Quantum quench in \mathcal{PT} -symmetric luttinger liquid, Phys. Rev. Lett. **124**, 136802 (2020).
- [28] D. C. Brody, Biorthogonal quantum mechanics, Journal of Physics A: Mathematical and Theoretical **47**, 035305 (2013).
- [29] T. Curtright and L. Mezincescu, Biorthogonal quantum systems, Journal of Mathematical Physics **48**, 092106 (2007).
- [30] J. V. Hounguevou, F. A. Dossa, and G. Y. H. Avossevou, Biorthogonal quantum mechanics for non-hermitian multimode and multiphoton jaynes-cummings models, Theoretical and Mathematical Physics **193**, 1464 (2017).
- [31] F. K. Kunst, E. Edvardsson, J. C. Budich, and E. J. Bergholtz, Biorthogonal bulk-boundary correspondence in non-hermitian systems, Phys. Rev. Lett. **121**, 026808 (2018).
- [32] H. B. Geyer, W. D. Heiss, and F. G. Scholtz, The physical interpretation of non-hermitian hamiltonians and other observables, Canadian Journal of Physics **86**, 1195 (2008).
- [33] A. Mostafazadeh, Time-dependent pseudo-hermitian hamiltonians and a hidden geometric aspect of quantum mechanics, Entropy **22**, 10.3390/e22040471 (2020).
- [34] A. Mostafazadeh, Energy observable for a quantum system with a dynamical hilbert space and a global geometric extension of quantum theory, Phys. Rev. D **98**, 046022 (2018).
- [35] X. Turkeshi and M. Schiró, Entanglement and correlation spreading in non-hermitian spin chains, Phys. Rev. B **107**, L020403 (2023).
- [36] B. Dóra, M. Heyl, and R. Moessner, The kibble-zurek mechanism at exceptional points, Nature Communications **10**, 2254 (2019).
- [37] A. Bácsi and B. Dóra, Dynamics of entanglement after exceptional quantum quench, Phys. Rev. B **103**, 085137 (2021).
- [38] B. Dóra, D. Sticlet, and C. P. Moca, Correlations at pt-symmetric quantum critical point, Phys. Rev. Lett. **128**, 146804 (2022).
- [39] B. Longstaff and E.-M. Graefe, Nonadiabatic transitions through exceptional points in the band structure of a pt-symmetric lattice, Phys. Rev. A **100**, 052119 (2019).
- [40] I. Percival, *Quantum State Diffusion* (Cambridge University Press, 1999).
- [41] H. M. Wiseman and G. J. Milburn, Quantum measurement theory, in *Quantum Measurement and Control* (Cambridge University Press, 2009) p. 1–50.
- [42] T. A. Brun, A simple model of quantum trajectories, American Journal of Physics **70**, 719 (2002), https://pubs.aip.org/aapt/ajp/article-pdf/70/7/719/7530993/719.1_online.pdf.
- [43] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, UK).
- [44] C. M. Bender, D. C. Brody, H. F. Jones, and B. K. Meister, Faster than hermitian quantum mechanics, Phys. Rev. Lett. **98**, 040403 (2007).
- [45] A. Mostafazadeh, Quantum brachistochrone problem and the geometry of the state space in pseudo-hermitian quantum mechanics, Phys. Rev. Lett. **99**, 130502 (2007).
- [46] K. Sim, N. Defenu, P. Mognini, and R. Chitra, Quantum metric unveils defect freezing in non-hermitian systems, Phys. Rev. Lett. **131**, 156501 (2023).
- [47] A. Mostafazadeh, Pseudo-hermiticity and generalized pt- and cpt-symmetries, Journal of Mathematical Physics

- 44**, 974 (2003), <https://doi.org/10.1063/1.1539304>.
- [48] A. Mostafazadeh, Time-dependent hilbert spaces, geometric phases, and general covariance in quantum mechanics, *Physics Letters A* **320**, 375 (2004).
- [49] G. Lindblad, On the generators of quantum dynamical semigroups, *Communications in Mathematical Physics* **48**, 119 (1976).
- [50] B. Gulácsi and B. Dóra, Defect production due to time-dependent coupling to environment in the lindblad equation, *Phys. Rev. B* **103**, 205153 (2021).
- [51] M. Coppola, D. Karevski, and G. T. Landi, Conditional no-jump dynamics of non-interacting quantum chains (2023), arXiv:2311.05515 [cond-mat.stat-mech].
- [52] A. Mostafazadeh, Pseudo-Hermiticity versus PT symmetry: The necessary condition for the reality of the spectrum of a non-Hermitian Hamiltonian, *Journal of Mathematical Physics* **43**, 205 (2002), https://pubs.aip.org/aip/jmp/article-pdf/43/1/205/19019524/205_1_online.pdf.
- [53] A. Das, Pseudo-hermitian quantum mechanics, *Journal of Physics: Conference Series* **287**, 012002 (2011).
- [54] Y. Jing, J.-J. Dong, Y.-Y. Zhang, and Z.-X. Hu, Biorthogonal dynamical quantum phase transitions in non-hermitian systems (2023), arXiv:2307.02993 [quant-ph].