The Imaging Database for Epilepsy And Surgery (IDEAS)

Peter N. Taylor^{1,2,3*}, Yujiang Wang^{1,2,3}, Callum Simpson¹, Vytene Janiukstyte¹, Jonathan Horsley¹, Karoline Leiberg¹, Beth Little^{1,2}, Harry Clifford¹, Sophie Adler⁴, Sjoerd B. Vos^{5,6}, Gavin P Winston^{3,6}, Andrew W McEvoy³, Anna Miserocchi³, Jane de Tisi³, John S Duncan³

- 1. CNNP Lab (www.cnnp-lab.com), Interdisciplinary Computing and Complex BioSystems Group, School of Computing, Newcastle University, Newcastle upon Tyne, United Kingdom
- 2. Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, United Kingdom
- 3. UCL Queen Square Institute of Neurology, Queen Square, London, United Kingdom
- 4. UCL Great Ormond Street Institute of Child Health, London, United Kingdom
- 5. Centre for Medical Image Computing, Department of Computer Science, UCL, London, United Kingdom
- 6. Centre for Microscopy, Characterisation, and Analysis, The University of Western Australia, Nedlands, Australia
- 7. Department of Medicine, Division of Neurology, Queen's University, Kingston, Canada

* peter.taylor@newcastle.ac.uk

Abstract

Magnetic resonance imaging (MRI) is a crucial tool to identify brain abnormalities in a wide range of neurological disorders. In focal epilepsy MRI is used to identify structural cerebral abnormalities. For covert lesions, machine learning and artificial intelligence algorithms may improve lesion detection if abnormalities are not evident on visual inspection. The success of this approach depends on the volume and quality of training data.

Herein, we release an open-source dataset of preprocessed MRI scans from 442 individuals with drug-refractory focal epilepsy who had neurosurgical resections, and detailed demographic information. The MRI scan data includes the preoperative 3D T1 and where available 3D FLAIR, as well as a manually inspected complete surface reconstruction and volumetric parcellations. Demographic information includes age, sex, age of onset of epilepsy, location of surgery, histopathology of resected specimen, occurrence and frequency of focal seizures with and without impairment of awareness, focal to bilateral tonic-clonic seizures, number of anti-seizure medications (ASMs) at time of surgery, and a total of 1764 patient years of post-surgical follow up. Crucially, we also include resection masks delineated from post-surgical imaging.

To demonstrate the veracity of our data, we successfully replicated previous studies showing long-term outcomes of seizure freedom in the range of around 50%. Our imaging data replicates findings of group level atrophy in patients compared to controls. Resection locations in the cohort were predominantly in the temporal and frontal lobes.

We envisage our dataset, shared openly with the community, will catalyse the development and application of computational methods in clinical neurology.

Introduction

Large-scale sharing of raw MRI scan data is commonplace in the neurosciences¹, and has led to substantial advances in our understanding of brain function and dysfunction². Such advances are enabled by the association of high quality clinical and demographic metadata with the scan. In neurological conditions such as Alzheimer's³, autism⁴, ADHD⁵, Parkinson's⁶, and traumatic brain injury⁷ there are large MRI data sets available for research. In epilepsy, data sharing was pioneered in the field of seizure prediction, with annotated EEG data available for hundreds of patients^{8–10}. However, given that MRI is crucial in the clinical management of epilepsy, it is surprising that relatively little data are openly available, particularly with high quality clinical and demographic information, but note ^{11–15,45} for related work.

In this study we share anonymised MRI scans from the National Hospital for Neurology and Neurosurgery, UCLH, acquired in 442 individuals with drug-refractory focal epilepsy who proceeded to neurosurgical resection. We also share anonymised demographic and clinical information for all subjects, and masks of subsequently resected tissue. To verify these data we replicate two previous landmark studies^{14,25}.

Methods

Study approval

This study of anonymised data that had been previously acquired was approved by the Health Research Authority, without the necessity to obtain individual subject consent (UCLH epilepsy surgery database: 22/SC/0016), and by the Database Local Data Monitoring Committee. Individuals who declined for their data to be used in anonymised research were not included in the research database.

Patient and data selection

We identified all eligible people who had neurosurgical resections for drug-resistant focal epilepsy at the National Hospital for Neurology & Neurosurgery, London, UK between 01 January 2003 and 30 June 2022 (n=625). From these we excluded individuals with previous neurosurgery (n=38), and those who had not had pre-operative T1 weighted MRI scans of sufficient quality at the Chalfont Centre for Epilepsy (n=145). Sufficient MRI quality was determined based on visual inspection by either PNT, BL, or VJ and considered aspects including motion artefact, ringing, and field of view completeness. Our final dataset contains pre-operative MRI for 442 individuals, and a resection mask for 433 of those individuals, along with most demographic and clinical data.

Clinical & demographic information

We extracted the following information from the UCLH epilepsy surgery database, that is prospectively and regularly updated. Data included: age at MRI scan, sex, age at surgery, type and location of surgery, post-operative pathology, age of epilepsy onset, occurrence and frequency of focal seizures with and without impairment of awareness, focal to bilateral tonic-clonic seizures, number of antiseizure medications (ASM) at time of surgery, yearly outcomes of seizure freedom assessed using the ILAE classification³⁹.

MRI scan information

All pre-operative 3D-T1-weighted scans were acquired using one of two 3T GE scanners. All subjects included in the release have T1-weighted MRI included, whilst 420 subjects also had FLAIR scans acquired in the same session included. Common acquisition protocols used include 3D fast spoiled gradient echo (FSPGR) at a resolution of 0.9375 x 0.9375 x 1.1mm, or magnetisation-prepared rapid acquisition gradient echo (MPRAGE) with a resolution of 1 x 1 x 1mm. Detailed acquisition parameters for each individual scan are shared as part of the data release in accompanying JavaScript object notation (JSON) format using the brain imaging data structure (BIDS).

Quality control

Pre-operative T1 weighted scans were processed with the FreeSurfer 7.3.2 software pipeline 'recon-all'¹⁶. The pipeline performed segmentation and parcellation of cortical tissues. Outputs were then visually inspected following established quality control protocols¹⁴. We paid particular attention to pial and white matter surfaces. Following visual inspection, some reconstructed surfaces were deemed insufficient and manual edits were made using control points and dura edits to improve their accuracy. For some subjects, particularly those with gross pathology, we noted that surface reconstruction may still be imperfect.

Once a scan and segmentation of sufficient quality was identified, we exported region volumes for deep brain areas, and additional thicknesses and surface areas for neocortical regions using the FreeSurfer command 'aparcstats2table'. These volume, thickness and surface area tables are shared as part of our data release.

Data harmonisation and normative modelling

Cortical thickness and volume measurements depend on a subject's age, sex, and the scanning protocol used for data acquisition. It is therefore necessary to account for these covariates to investigate the effects of epilepsy. For this, we used a combination of two methods^{46,47}. First, a data harmonisation technique, ComBat, which removes scanner differences whilst accounting for and preserving covariate and pathology effects. Second, a

normative model, which describes population-level trends in the relationship between variables. Specifically we modelled age, sex, and cortical thickness or volume, and estimated the expected variance in these trends to derive subject-specific deviations⁴⁸.

Both ComBat and normative modelling require an inference of age and sex effects of the healthy population. We therefore used data from two large public healthy control datasets, NKI (n=833) and OASIS (n=542)^{18,19}, for model training. This was in addition to 100 healthy control scans acquired on the same scanners as the patient cohort, which informed on effects of scanning protocols and served as a reference cohort for subsequent abnormality calculations.

For data harmonisation with ComBat, we clustered subjects based on the similarity of their scanning parameters. ComBat then modelled effects of age and sex, as well as the offset and variance of each cluster, so that covariate effects could be retained in the data whilst effects of scanning acquisition were removed¹⁷.

After data harmonisation, we fitted a generalised additive model (GAM) in each region to the healthy controls. For cortical thickness, we used the model formula

 $t \sim 1 + s(age) + sex$,

where *t* is thickness, *s*(*age*) is a smooth function of age, and *sex* is a fixed effect. For subcortical volume, we used the model

 $v \sim 1 + s(age) + sex + ICV$,

where *v* is volume, *s*(*age*) is a smooth function of age, and *sex* and *ICV* (intracranial volume) are fixed effects. To apply this normative model, we predicted values of thickness/volume for each patient and the 100 healthy controls which were acquired on the same scanners. We calculated residuals, i.e. differences between the observed and predicted values, which removed the age and sex effects from the data.

Abnormality calculation

For each subject, we calculated regional abnormalities in cortical thickness and subcortical volume. This was done by z-scoring the residuals, using the mean and standard deviation in each region across the residuals of the 100 reference controls:

$$Z_{ij} = \frac{R_{ij} - \mu_i}{\sigma_i},$$

where Z_{ij} is the z-scored abnormality in region *i* for subject *j*, *R* is the residual, μ is the mean residual across controls and σ is the standard deviation of the residuals across controls. These abnormalities quantify how many standard deviations a subject's regional thickness or volume is away from the control population.

Similar abnormalities were also calculated for healthy controls. In this scenario, each control was held out from the control cohort before the calculation of μ and σ used for its z-scoring.

For patients with mesial temporal lobe epilepsy (mTLE), we quantified the extent of alterations in cortical thickness or subcortical volume across the cohort using Cohen's d. In each region, we calculated the effect size of the difference in abnormality between the cohort of patients with mTLE and healthy controls, and plotted the results using the ENIGMA toolbox²⁰. This approach is broadly similar to that described previously¹⁴.

Generation of resection masks

We used postoperative imaging, in the 433 individuals for whom these were available, to generate masks of the tissue that was subsequently resected. Masks were initially generated automatically using a custom-built software pipeline before manual edits of pipeline outputs were performed where needed. The mask generation pipeline comprised three steps. First, data were processed through FastSurfer²¹ to label brain regions and the lobe where the resection had taken place was identified. Second, the ANTs registration tool²² was used to perform a two-part alignment of the pre-operative and post-operative images to compensate for any distortion of remaining brain tissue into the resection cavity seen in the post-operative image. The registration used the lobe information from step 1 to improve registration accuracy. The third step used a classification algorithm ATROPOS²³ to find differences between the registered images. Step three was then followed by manual checks and edits, when needed, to ensure that the produced resection mask was within and accurately filled anatomical boundaries²⁴.

Data de-identification and anonymisation

MRI scans were converted to NIFTI format to remove any identifying information in scan headers. Individual faces were removed from all MR images by multiplying a dilated binarised brain mask with the original image. All scans were then visually inspected to ensure that the nose, mouth, and ears were not visible.

Date of birth and date of scan or surgery are not shared publicly since these are protected as potentially identifiable information. Instead, data are shared in categories (e.g. 18-22, 23-27, 28-32 years old). Annual ILAE surgery outcomes of up to five years are shared when available. Data beyond five years are not shared to maintain anonymity regarding date of surgery.

Data availability

Data will be shared on the openneuro.org platform and publicly searchable without restriction upon acceptance of the manuscript. For review purposes, links to data can be found in Table S3.

Results

The IDEAS dataset

Demographic information was available for all patients included in our final data release (n=442). Post-operative imaging to delineate a resection mask was also available for most subjects (N=433, 98%). Table 1 summarises the shared data which include: sex, history of FBTCS (12 months pre-surgery), history and frequency of focal seizures with impaired awareness (12 months pre-surgery), history of status epilepticus, histopathological findings, clinical MRI findings, number of anti-seizure medications at time of surgery, age at MRI scan. Figure 1 presents data from a typical case.

	Number of subjects	Notes
T1w MRI	442	
FLAIR MRI	411	
Resection mask	433	
Sex	446	
Age of epilepsy onset	442	
History of FUS	446	N=409=true
Frequency of FUS	406 of 409	
History of FBTCS	442	N=343=true
Frequency of FBTCS	330 of 343	
History of SE	442	N=53=true
Side of resection	442	
Resection type	442	
Pathology	442	
Number of ASMs	442	
Age of epilepsy surgery	442	
12 month ILAE outcome	427	
24 month ILAE outcome	403	
36 month ILAE outcome	356	
48 month ILAE outcome	311	
60 month ILAE outcome	267	

Table 1. Summary of available data



Five-year outcome of adult epilepsy surgery, patterns of seizure remission, and

relapse

For clinical metadata, we replicate a previous study demonstrating long-term outcomes from epilepsy surgery²⁵. Post-operative 12-month outcomes of seizure-freedom were available for 422 patients. Five subjects had multilobar resections. At 12 months, 243 (58%) of the remaining 422 patients were completely seizure-free (figure 2A). After five years of follow up, patients with extra-temporal lobe epilepsy surgery had the lowest estimated proportion of continuous seizure-freedom (44% and 42% for lesionectomy and lobar resection respectively).

Patients with focal aware seizures (FAS) in the first 24 months after surgery were significantly more likely to relapse to have seizures with impaired awareness in the subsequent years (figure 2B, log rank test p<0.001). Sample sizes at each year of follow-up are presented in supplementary table S2. These findings align with those described previously²⁵.

There were large ranges of age of epilepsy onset (median 13, IQR 14 years), age at surgery (median 37, IQR 17 years) and epilepsy duration (mean 21, IQR 20 years). The minimum age at surgery was 16 years old, with the majority over 18, reflecting our clinical practice as an adult epilepsy centre. None of these variables were significantly related to post-operative seizure outcomes (figure 2C,D,E).



Figure 2: Seizure outcome up to 5 years after epilepsy surgery

A Survival plot of proportion of patients remaining seizure-free following epilepsy surgery and with available follow-up at yearly intervals, showing time to first seizure. Each coloured line represents a different surgical procedure (ATLRx=anterior temporal resection. TLesx=temporal lesionectomy. ETLesx=extratemporal lesionectomy. ETLx=extratemporal resection. Hx=hemispherectomy).

B Survival plot of proportion of patients who did (blue), and did not (red), have focal aware seizures (FAS) following surgery in years 1,2, remaining free of seizures with impaired awareness at subsequent years.

C,D,E: 12-month seizure outcome did not differ by age of onset of epilepsy, (*C*), age at surgery (*D*), or duration of epilepsy (*E*). All results broadly replicate those described previously in more extensive cohorts from 1990-2012^{25,40}.

Structural brain abnormalities in focal epilepsies

We investigated brain alterations in mTLE, using similar methods to those described previously¹⁴. Figure 3 shows effect size differences (Cohen's d) in cortical and subcortical brain regions for left, and right, mesial temporal lobe epilepsy (mTLE) (N=122 and N=85 respectively), compared to healthy controls (N=100).

Reduced volumes were noted in several regions including the ipsilateral hippocampus and thalamus. Neocortical thickness was also reduced in several areas. Specifically, in left mTLE the cortical thickness of the caudal middle frontal gyrus was the most reduced ipsilaterally (d=-0.61), followed by the inferior parietal cortex (d=-0.58) and the contralateral precentral gyrus (d=-0.58). In right mTLE, neocortical volumes were reduced in the parietal lobe. Full effect sizes for all regions in all individuals are included in the data release tables.

Other imaging features available in the IDEAS database are available as direct outputs from the FreeSurfer recon-all pipeline, along with fully processed FreeSurfer volumes and surfaces. Outputs include neocortical volumes, surface area and thickness for a further seven parcellations (multi-scale Lausanne⁴¹, Destrieux⁴², HCP-MMP1⁴³, and Whittaker⁴⁴). In addition, volumes of brainstem subregions, hippocampal and amygdala subunits, and hypothalamus segmentation are also available.



other studies.¹⁴

Resection masks & locations

Resection masks were generated for N=433 subjects including N=356 individuals with TLE, and N=58 with frontal lobe resections. Masks are visualised in figure 4, overlaid in the same space. Individual resection masks are available within the IDEAS database.



Discussion

Here we present the Imaging Database for Epilepsy And Surgery (IDEAS). The database contains a large, anonymised sample of neuroimaging, and clinical metadata. The data are organised in an easy-to-use manner, and amenable to future research. Data have been pre-processed, checked for quality and organised using BIDS format. We present demonstrative use-cases for the data, replicating several recent key publications. Data are openly available for the research community.

With the advent of advanced machine learning models and artificial intelligence, we are experiencing a paradigm shift in radiology and beyond. In some task-specific scenarios, AI already exceeds human performance²⁶. In epilepsy, recent work has applied AI techniques for lateralisation³¹⁻³³ and localisation²⁷⁻³⁰ of epileptogenic tissues. Replication of these algorithms on external datasets is crucial to achieve clinical translation. Our data release, including gold standard resections and patient outcome measures represent an opportunity for the field to progress in this area by providing real-world training and test data.

To gain a mechanistic understanding of epilepsy, high quality clinical and demographic data are required. A key component of our data release is the richness of the metadata. Several studies previously reported clinical and demographic factors to be related to post-operative seizure outcomes^{34,35}. For example, Jehi et al (2015)³⁴ reported an association with epilepsy duration, whilst others^{35, 36} report FBTCS, being associated with reduced chance of a seizure free outcome. We also share histopathology classification for the resected tissue, which we envisage may be of use to derive MRI markers of specific pathology³⁷. These collated clinical and demographic features, in a cohort of this size, are rare in epilepsy. Their associations will be vital to investigate underlying mechanisms.

A crucial facet of our data is that the imaging and metadata are linked. The linking of these data is vital to enable research into our understanding of the pathophysiology and functional anatomy of epilepsy, particularly at the level of the individual. Until now such data on a large scale was unavailable in epilepsy. To facilitate scientific endeavours we have shared all

imaging data in a preprocessed format of z-scores, in addition to the raw MRI data. We have also included the metadata in easy-to-read spreadsheets to facilitate future research.

Data quality is crucial. We therefore performed multiple rounds of quality control. For T1weighted MRI data, all scans were visually inspected, and manual edits made to preprocessed data outputs from FreeSurfer when necessary. Clinical and demographic data, were entered prospectively, verified, and regularly updated, since the inception of the UCLH epilepsy surgery database in 1990. All code to process and analyse the data was verified by multiple people. We are sharing the data on the OpenNeuro platform¹, which is version controlled, and we will provide updates when necessary. We welcome any feedback from the community regarding data issues and commit to continue supporting its continued use.

A further key feature of these data is the inclusion of masks delineating the resection zone. In individuals who are seizure-free in the long-term after epilepsy surgery it can be assumed that at least part of the epileptogenic zone (EZ) is located within the resection zone. It cannot be assumed that there is a one-to-one correspondence between the resection and EZ in seizure-free patients. The resection may be larger than the EZ, for example when surgical access was required via non-epileptogenic tissue. Furthermore, the resection may be smaller than the EZ in seizure-free patients, if the patient is still on anti-seizure medication. Thus, the resection masks represent a silver standard approximation of the EZ, and should be regarded as such. Nonetheless, such a silver standard may still be sufficient for training algorithms for prediction³⁸.

Full anonymisation of the data that are shared was an important consideration and we took several steps to protect against identification of individuals. For MRI data, these steps included removal of header information from imaging files using NIFTI format and removal of potentially identifying facial features from MRI. Clinical and demographic data were extracted from the clinical database by the clinical team and these and MRI data were pseudo-anonymised prior to any data processing and linking of MRI and clinical data. All data were irrevocably anonymised prior to preparation for being shareable. Important steps to ensure anonymisation included limiting disclosed follow-up to 5 years from surgery, and reducing specificity of age at surgery, age of onset and duration of epilepsy to 5 year epochs.

The use of previously acquired clinical and investigatory data for research, without the requirement for individual patient consent was, approved by Information Governance of UCL Hospitals, the UCLH/UCL Joint Research Office, the Health Research Authority (UCLH epilepsy surgery database: 22/SC/0016) and the Database Data Monitoring Committee. A key requirement is that all data shared externally to UCL and UCLH are totally and irrevocably anonymised.

Although large and richly characterised, our data have limitations. First, our data reflects resective adult epilepsy surgery clinical practice at a single centre over the last 20 years. As such, no individuals who received laser interstitial thermal therapy or thermocoagulation are included. Second, our surgical practice is confined to adults, and any inferences with regard to paediatric epilepsy are limited. Third, our data release of 3D-T1-weighted and T2-FLAIR MRI represents only part of the information used during pre-surgical evaluation. Many of the individuals having neurosurgical treatment of drug-refractory focal epilepsy also underwent acquisition of data from other modalities, including diffusion MRI, PET, SPECT high density scalp EEG, MEG and intracranial EEG. Neuropsychological and neuropsychiatric data are also typically considered during presurgical evaluation. We hope to be able to include these data in the coming years.

We hope that the IDEAS data will be a valuable resource for the epilepsy research community that will catalyse efforts in data science research in epilepsy. We welcome contributions from other sites, and propose our metadata headers as common data elements to be used for cross site consistency. We will be pleased to concatenate contributions from other centres with our own to establish a larger, multi-centre resource.

Acknowledgements

We thank members of the Computational Neurology, Neuroscience & Psychiatry Lab (www.cnnp-lab.com) for discussions on the analysis and manuscript. We are grateful to the UCLH Epilepsy Surgery Database local Data Monitoring Committee. P.N.T. and Y.W. are both supported by UKRI Future Leaders Fellowships (MR/T04294X/1, MR/V026569/1). Y.W. and B.L. are supported by EPSRC (EP/Y016009/1). G.P.W. and acquisition of control data was supported by the MRC (G0802012, MR/M00841X/1). JSD, JdT are supported by the NIHR UCLH/UCL Biomedical Research Centre. This work was supported by Epilepsy Research UK (grant number P1904).

References

1. Markiewicz CJ, Gorgolewski KJ, Feingold F, et al. The OpenNeuro resource for sharing of neuroscience data. *Elife*. 2021;10:e71774.

2. Biswal BB, Mennes M, Zuo XN, et al. Toward discovery science of human brain function. *Proceedings of the national academy of sciences*. 2010;107(10):4734-4739.

3. Mueller SG, Weiner MW, Thal LJ, et al. Ways toward an early diagnosis in alzheimer's disease: The alzheimer's disease neuroimaging initiative (ADNI). *Alzheimer's & Dementia*. 2005;1(1):55-66.

4. Di Martino A, Yan CG, Li Q, et al. The autism brain imaging data exchange: Towards a large-scale evaluation of the intrinsic brain architecture in autism. *Molecular psychiatry*. 2014;19(6):659-667.

5. ADHD-200Consortium. The ADHD-200 consortium: A model to advance the translational potential of neuroimaging in clinical neuroscience. *Frontiers in systems neuroscience*. 2012;6:62.

6. Marek K, Jennings D, Lasch S, et al. The parkinson progression marker initiative (PPMI). *Progress in neurobiology*. 2011;95(4):629-635.

7. Thompson HJ, Vavilala MS, Rivara FP. Common data elements and federal interagency traumatic brain injury research informatics system for TBI research. *Annual review of nursing research*. 2015;33(1).

8. Lehnertz K, Litt B. The first international collaborative workshop on seizure prediction: Summary and data description. *Clinical neurophysiology*. 2005;116(3):493-505.

9. Wagenaar JB, Brinkmann BH, Ives Z, Worrell GA, Litt B. A multimodal platform for cloud-based collaborative research. In: *2013 6th International IEEE/EMBS Conference on Neural Engineering (NER)*. IEEE; 2013:1386-1389.

10. Ihle M, Feldwisch-Drentrup H, Teixeira CA, et al. EPILEPSIAE–a european epilepsy database. *Computer methods and programs in biomedicine*. 2012;106(3):127-138.

11. Schuch F, Walger L, Schmitz M, et al. An open presurgery MRI dataset of people with epilepsy and focal cortical dysplasia type II. *Scientific Data*. 2023;10(1):475.

12. Sisodiya SM, Whelan CD, Hatton SN, et al. The ENIGMA-epilepsy working group: Mapping disease from large data sets. *Human brain mapping*. 2022;43(1):113-128.

13. Hatton SN, Huynh KH, Bonilha L, et al. White matter abnormalities across different epilepsy syndromes in adults: An ENIGMA-epilepsy study. *Brain*. 2020;143(8):2454-2473.

14. Whelan CD, Altmann A, Botía JA, et al. Structural brain abnormalities in the common epilepsies assessed in a worldwide ENIGMA study. *Brain*. 2018;141(2):391-408.

15. Rodríguez-Cruces R, Camacho-Téllez V, Fajardo A, Concha L. "Temporal lobe epilepsy - UNAM". Published online 2023. doi:doi:10.18112/openneuro.ds004469.v1.1.3

16. Fischl B. FreeSurfer. *Neuroimage*. 2012;62(2):774-781.

17. Fortin JP, Cullen N, Sheline YI, et al. Harmonization of cortical thickness measurements across scanners and sites. *NeuroImage*. 2018;167:104-120. doi:10.1016/j.neuroimage.2017.11.024

18. Nooner KB, Colcombe SJ, Tobe RH, et al. The NKI-Rockland Sample: A Model for Accelerating the Pace of Discovery Science in Psychiatry. *Frontiers in Neuroscience*. 2012;6. doi:10.3389/fnins.2012.00152

19. LaMontagne PJ, Benzinger TLs, Morris JC, et al. *OASIS-3: Longitudinal Neuroimaging, Clinical, and Cognitive Dataset for Normal Aging and Alzheimer Disease*. Radiology; Imaging; 2019. doi:10.1101/2019.12.13.19014902

20. Larivière S, Paquola C, Park B, et al. The ENIGMA toolbox: Multiscale neural contextualization of multisite neuroimaging datasets. *Nature Methods*. 2021;18(7):698-700.

21. Henschel L, Conjeti S, Estrada S, Diers K, Fischl B, Reuter M. Fastsurfer-a fast and accurate deep learning based neuroimaging pipeline. *NeuroImage*. 2020;219:117012.

22. Avants BB, Tustison NJ, Song G, Cook PA, Klein A, Gee JC. A reproducible evaluation of ANTs similarity metric performance in brain image registration. *Neuroimage*. 2011;54(3):2033-2044.

23. Avants BB, Tustison NJ, Stauffer M, Song G, Wu B, Gee JC. The insight ToolKit image registration framework. *Frontiers in neuroinformatics*. 2014;8:44.

24. Taylor PN, Sinha N, Wang Y, et al. The impact of epilepsy surgery on the structural connectome and its relation to outcome. *NeuroImage: Clinical*. 2018;18:202-214.

25. De Tisi J, Bell GS, Peacock JL, et al. The long-term outcome of adult epilepsy surgery, patterns of seizure remission, and relapse: A cohort study. *The Lancet*. 2011;378(9800):1388-1395.

26. Hosny A, Parmar C, Quackenbush J, Schwartz LH, Aerts HJ. Artificial intelligence in radiology. *Nature Reviews Cancer*. 2018;18(8):500-510.

27. Spitzer H, Ripart M, Whitaker K, et al. Interpretable surface-based detection of focal cortical dysplasias: A multi-centre epilepsy lesion detection study. *Brain*. 2022;145(11):3859-3871.

28. David B, Kröll-Seger J, Schuch F, et al. External validation of automated focal cortical dysplasia detection using morphometric analysis. *Epilepsia*. 2021;62(4):1005-1021.

29. Walger L, Adler S, Wagstyl K, et al. Artificial intelligence for the detection of focal cortical dysplasia: Challenges in translating algorithms into clinical practice. *Epilepsia*. Published online 2023.

30. Gill RS, Lee HM, Caldairou B, et al. Multicenter validation of a deep learning detection algorithm for focal cortical dysplasia. *Neurology*. 2021;97(16):e1571-e1582.

31. Gleichgerrcht E, Munsell BC, Alhusaini S, et al. Artificial intelligence for classification of temporal lobe epilepsy with ROI-level MRI data: A worldwide ENIGMA-epilepsy study. *NeuroImage: Clinical*. 2021;31:102765.

32. Johnson GW, Cai LY, Narasimhan S, et al. Temporal lobe epilepsy lateralisation and surgical outcome prediction using diffusion imaging. *Journal of Neurology, Neurosurgery & Psychiatry*. 2022;93(6):599-608.

33. Kaestner E, Rao J, Chang AJ, et al. Convolutional neural network algorithm to determine lateralization of seizure onset in patients with epilepsy: A proof-of-principle study. *Neurology*. Published online 2023.

34. Jehi L, Yardi R, Chagin K, et al. Development and validation of nomograms to provide individualised predictions of seizure outcomes after epilepsy surgery: A retrospective analysis. *The Lancet Neurology*. 2015;14(3):283-290.

35. Bell GS, De Tisi J, Gonzalez-Fraile JC, et al. Factors affecting seizure outcome after epilepsy surgery: An observational series. *Journal of Neurology, Neurosurgery & Psychiatry*. 2017;88(11):933-940.

36. Rowland NC, Englot DJ, Cage TA, Sughrue ME, Barbaro NM, Chang EF. A meta-analysis of predictors of seizure freedom in the surgical management of focal cortical dysplasia. *Journal of neurosurgery*. 2012;116(5):1035-1041.

37. Deleo F, Thom M, Concha L, Bernasconi A, Bernhardt BC, Bernasconi N. Histological and MRI markers of white matter damage in focal epilepsy. *Epilepsy Research*. 2018;140:29-

38. Owen TW, Janiukstyte V, Hall GR, et al. Identifying epileptogenic abnormalities through spatial clustering of MEG interictal band power. *Epilepsia Open*. Published online 2023.

39. Wieser, H.G., Blume, W.T., Fish, D., Goldensohn, E., Hufnagel, A., King, D., Sperling, M.R., Lüders, H. and Pedley, T.A., 2001. Proposal for a new classification of outcome with respect to epileptic seizures following epilepsy surgery. Epilepsia, 42(2), pp.282-286.

40. Bell GS, de Tisi J, Gonzalez-Fraile JC, Peacock JL, McEvoy AW, Harkness WFJ, Foong J, Pope RA, Diehl B, Sander JW, Duncan JS. Factors affecting seizure outcome after epilepsy surgery: an observational series. J Neurol Neurosurg Psychiatry. 2017 Nov;88(11):933-940.

41. Daducci, A., Gerhard, S., Griffa, A., Lemkaddem, A., Cammoun, L., Gigandet, X., Meuli, R., Hagmann, P. and Thiran, J.P., 2012. The connectome mapper: an open-source processing pipeline to map connectomes with MRI. PloS one, 7(12), p.e48121.

42. Destrieux, C., Fischl, B., Dale, A. and Halgren, E., 2010. Automatic parcellation of human cortical gyri and sulci using standard anatomical nomenclature. Neuroimage, 53(1), pp.1-15.

43. Glasser, M.F., Coalson, T.S., Robinson, E.C., Hacker, C.D., Harwell, J., Yacoub, E., Ugurbil, K., Andersson, J., Beckmann, C.F., Jenkinson, M. and Smith, S.M., 2016. A multi-modal parcellation of human cerebral cortex. Nature, 536(7615), pp.171-178.

44. Whitaker, K.J., Vértes, P.E., Romero-Garcia, R., Váša, F., Moutoussis, M., Prabhu, G., Weiskopf, N., Callaghan, M.F., Wagstyl, K., Rittman, T. and Tait, R., 2016. Adolescence is associated with genomically patterned consolidation of the hubs of the human brain connectome. Proceedings of the National Academy of Sciences, 113(32), pp.9105-9110.

45. Spitzer, Hannah, et al. "Interpretable surface-based detection of focal cortical dysplasias: a Multi-centre Epilepsy Lesion Detection study." Brain 145.11 (2022): 3859-3871.

46. Ge, R., Yu, Y., Qi, Y.X., Fan, Y.N., Chen, S., Gao, C., Haas, S.S., New, F., Boomsma, D.I., Brodaty, H. and Brouwer, R.M., 2024. Normative modelling of brain morphometry across the lifespan with CentileBrain: algorithm benchmarking and model optimisation. The Lancet Digital Health, 6(3), pp.e211-e221.

47. Frangou, S., Modabbernia, A., Williams, S.C., Papachristou, E., Doucet, G.E., Agartz, I., Aghajani, M., Akudjedu, T.N., Albajes-Eizagirre, A., Alnæs, D. and Alpert, K.I., 2022. Cortical thickness across the lifespan: Data from 17,075 healthy individuals aged 3–90 years. Human brain mapping, 43(1), pp.431-451.

48. Rutherford, S., Fraza, C., Dinga, R., Kia, S.M., Wolfers, T., Zabihi, M., Berthet, P., Worker, A., Verdi, S., Andrews, D. and Han, L.K., 2022. Charting brain growth and aging at high spatial precision. elife, 11, p.e72904.