

# On Stewart's Perturbation Theorem for SVD

Ren-Cang Li\*

Ninoslav Truhar<sup>†</sup>Lei-Hong Zhang<sup>‡</sup>

June 12, 2024

## Abstract

This paper establishes a variant of Stewart's theorem [Theorem 6.4 of Stewart, *SIAM Rev.*, 15:727–764, 1973] for the singular subspaces associated with the SVD of a matrix subject to perturbations. Stewart's original version uses both the Frobenius and spectral norms, whereas the new variant uses the spectral norm and any unitarily invariant norm that offer choices per convenience of particular applications and lead to sharper bounds than that straightforwardly derived from Stewart's original theorem with the help of the well-known equivalence inequalities between matrix norms. Of interest in their own right, bounds on the solution to two couple Sylvester equations are established for a few different circumstances.

**Keywords:** SVD, perturbation, singular subspaces, coupled Sylvester equations

**Mathematics Subject Classification** 65F15, 15A18

## 1 Introduction

In [15], Stewart established a perturbation theory for the singular subspaces associated with the SVD of a matrix  $G \in \mathbb{C}^{m \times n}$  slightly perturbed. In the same paper he also investigated the eigenspace of a matrix and the deflating subspace of a regular matrix pencil subject to perturbations. But in this paper, we limit our scope to one of his theorems, namely, [15, Theorem 6.4] on SVD, where both the Frobenius and spectral norms are used to measure perturbations. Our goal is to establish a more general and yet sharper version of [15, Theorem 6.4] in the spectral norm and any unitarily invariant norm. Our new version offers flexibility in its applications, for example, the version with the unitarily invariant norm also set to the spectral norm is more convenient to use in our recent work [18] for analyzing the quality of a reduced order model by approximate balanced truncation [1, 2]. Even in the case of using both the Frobenius and spectral

---

\*Department of Mathematics, University of Texas at Arlington, Arlington, TX 76019-0408, USA. Supported in part by NSF DMS-2009689. Email: [rccli@uta.edu](mailto:rccli@uta.edu).

<sup>†</sup>Department of Mathematics, J. J. Strossmayer University of Osijek, Trg Ljudevita Gaja 6, 31000 Osijek, Croatia. Email: [ntruhar@mathos.hr](mailto:ntruhar@mathos.hr).

<sup>‡</sup>School of Mathematical Sciences, Soochow University, Suzhou 215006, Jiangsu, China. Supported in part by the National Natural Science Foundation of China NSFC-11671246 NSFC-12371380, and Jiangsu Shuangchuang Project JSSCTD202209. Email: [longzlh@suda.edu.cn](mailto:longzlh@suda.edu.cn).

norms, our results are slightly better than Stewart's [15, Theorem 6.4] in terms of a less stringent condition and yet a sharper bound.

Given  $G \in \mathbb{C}^{m \times n}$ , let

$$U \equiv \begin{bmatrix} r & m-r \\ U_1 & U_2 \end{bmatrix} \in \mathbb{C}^{m \times m}, \quad V \equiv \begin{bmatrix} r & n-r \\ V_1 & V_2 \end{bmatrix} \in \mathbb{C}^{n \times n} \quad (1.1a)$$

be two unitary matrices such that  $G$  admits the decomposition

$$U^H G V \equiv \begin{bmatrix} U_1^H \\ U_2^H \end{bmatrix} G [V_1, V_2] = \begin{bmatrix} r & n-r \\ m-r & \end{bmatrix} \begin{bmatrix} G_1 & 0 \\ 0 & G_2 \end{bmatrix}, \quad (1.1b)$$

where  $1 \leq r < \min\{m, n\}$ . For example, this can be the SVD of  $G$ , for which  $G_1$  is diagonal with nonnegative diagonal entries as some of the singular values of  $G$  and  $G_2$  is leading diagonal by which we mean that only its entries along the main diagonal line starting at the top-left corner may be nonzero and nonnegative and they are some of the singular values of  $G$ , too. By convention,  $G$  has  $\min\{m, n\}$  singular values  $\sigma_i(G)$  arranged in decreasing order:

$$\sigma_1(G) \geq \sigma_2(G) \geq \cdots \geq \sigma_{\min\{m, n\}}(G). \quad (1.2)$$

Denote the singular value set and its extended set of  $G$  by

$$\text{sv}(G) = \{\sigma_i(G)\}_{i=1}^{\min\{m, n\}}, \quad \text{sv}_{\text{ext}}(G) = \text{sv}(G) \cup \{|m - n| \text{ copies of } 0\text{s}\}, \quad (1.3)$$

where the union is meant to be the multiset union.

Consider now that  $G$  is perturbed to  $\tilde{G} = G + E \in \mathbb{C}^{m \times n}$ , and partition

$$U^H \tilde{G} V \equiv \begin{bmatrix} U_1^H \\ U_2^H \end{bmatrix} (G + E) [V_1, V_2] = \begin{bmatrix} r & n-r \\ m-r & \end{bmatrix} \begin{bmatrix} G_1 + E_{11} & E_{12} \\ E_{21} & G_2 + E_{22} \end{bmatrix}. \quad (1.4)$$

Naturally, one would ask whether  $\tilde{G}$  admits a decomposition that is ‘‘close’’ to the one for  $G$  in (1.1). Stewart [15, Theorem 6.4] provided an answer to that by seeking orthogonal matrices [15, p.760]

$$\check{U} \equiv \begin{bmatrix} r & m-r \\ \check{U}_1 & \check{U}_2 \end{bmatrix} = [U_1, U_2] \begin{bmatrix} I_r & -\Gamma^H \\ \Gamma & I_{m-r} \end{bmatrix} \begin{bmatrix} (I + \Gamma^H \Gamma)^{-1/2} & 0 \\ 0 & (I + \Gamma \Gamma^H)^{-1/2} \end{bmatrix}, \quad (1.5a)$$

$$\check{V} \equiv \begin{bmatrix} r & n-r \\ \check{V}_1 & \check{V}_2 \end{bmatrix} = [V_1, V_2] \begin{bmatrix} I_r & -\Omega^H \\ \Omega & I_{n-r} \end{bmatrix} \begin{bmatrix} (I + \Omega^H \Omega)^{-1/2} & 0 \\ 0 & (I + \Omega \Omega^H)^{-1/2} \end{bmatrix} \quad (1.5b)$$

such that

$$\check{U}^H \tilde{G} \check{V} \equiv \begin{bmatrix} \check{U}_1^H \\ \check{U}_2^H \end{bmatrix} (G + E) [\check{V}_1, \check{V}_2] = \begin{bmatrix} r & n-r \\ m-r & \end{bmatrix} \begin{bmatrix} \check{G}_1 & 0 \\ 0 & \check{G}_2 \end{bmatrix}. \quad (1.6)$$

Theorem 1.1 below is [15, Theorem 6.4] in our notations here.

**Theorem 1.1** ([15, Theorem 6.4]). *Given  $G, \tilde{G} \in \mathbb{C}^{m \times n}$ , let  $G$  be decomposed as in (1.1), and partition  $U^H \tilde{G} V$  according to (1.4). Let*

$$\hat{\varepsilon} = \sqrt{\|E_{12}\|_F^2 + \|E_{21}\|_F^2}, \quad (1.7a)$$

$$\delta = \min_{\mu \in \text{sv}(G_1), \nu \in \text{sv}_{\text{ext}}(G_2)} |\mu - \nu|, \quad (1.7b)$$

$$\underline{\delta} = \delta - \|E_{11}\|_2 - \|E_{22}\|_2, \quad (1.7c)$$

where  $\|\cdot\|_2$  and  $\|\cdot\|_F$  denote the spectral and Frobenius norms, respectively. If

$$\underline{\delta} > 0 \quad \text{and} \quad \frac{\hat{\varepsilon}}{\underline{\delta}} < \frac{1}{2}, \quad (1.8)$$

then there exist  $\Omega \in \mathbb{C}^{(n-r) \times r}$  and  $\Gamma \in \mathbb{C}^{(m-r) \times r}$  satisfying

$$\sqrt{\|\Gamma\|_F^2 + \|\Omega\|_F^2} \leq \frac{1 + \sqrt{1 - 4(\hat{\varepsilon}/\underline{\delta})^2}}{1 - 2(\hat{\varepsilon}/\underline{\delta})^2 + \sqrt{1 - 4(\hat{\varepsilon}/\underline{\delta})^2}} \frac{\hat{\varepsilon}}{\underline{\delta}} < 2 \frac{\hat{\varepsilon}}{\underline{\delta}} \quad (1.9)$$

such that (1.6) with (1.5) holds.

A number of results can be deduced from this informative theorem, e.g., bounds on  $\|U - \check{U}\|_F$  and  $\|V - \check{V}\|_F$ , the explicit expressions of  $\check{G}_1$  and  $\check{G}_2$  in terms of  $\Omega$  and  $\Gamma$ ,  $G_1$ ,  $G_2$ , and  $E_{ij}$ , and also the singular values of  $\check{G}$  is the multiset union of the singular values of  $\check{G}_1$  and  $\check{G}_2$ .

Although the spectral norm  $\|\cdot\|_2$  is used in defining  $\underline{\delta}$ , the Frobenius norm  $\|\cdot\|_F$  is in full display in defining  $\hat{\varepsilon}$  and in bounding  $\Omega$  and  $\Gamma$ . A straightforward version of it, using only the spectral norm can be easily derived in light of the equivalency inequalities

$$\|B\|_2 \leq \|B\|_F \leq \sqrt{\text{rank}(B)} \|B\|_2 \leq \sqrt{\min\{s, t\}} \|B\|_2 \quad \text{for } B \in \mathbb{C}^{s \times t}.$$

For example, we can conclude that if  $\underline{\delta} > 0$  and

$$\tilde{\varepsilon} := \sqrt{\min\{m-r, n-r, r\}} \frac{\sqrt{\|E_{12}\|_2^2 + \|E_{21}\|_2^2}}{\underline{\delta}} < \frac{1}{2}, \quad (1.10)$$

then there exist  $\Omega \in \mathbb{C}^{(n-r) \times r}$  and  $\Gamma \in \mathbb{C}^{(m-r) \times r}$  satisfying

$$\sqrt{\|\Gamma\|_F^2 + \|\Omega\|_F^2} \leq \frac{1 + \sqrt{1 - 4(\tilde{\varepsilon}/\underline{\delta})^2}}{1 - 2(\tilde{\varepsilon}/\underline{\delta})^2 + \sqrt{1 - 4(\tilde{\varepsilon}/\underline{\delta})^2}} \frac{\tilde{\varepsilon}}{\underline{\delta}} < 2 \frac{\tilde{\varepsilon}}{\underline{\delta}} \quad (1.11)$$

such that (1.6) with (1.5) holds. There are two clear drawbacks of such a straightforward version: 1) a much stronger condition in (1.10) than something like

$$\frac{\sqrt{\|E_{12}\|_2^2 + \|E_{21}\|_2^2}}{\underline{\delta}} < \frac{1}{2} \quad (1.12)$$

as one might possibly expect, and 2) a much weaker conclusion in (1.11) than something like

$$\sqrt{\|T\|_2^2 + \|\Omega\|_2^2} \leq \sqrt{\|T\|_F^2 + \|\Omega\|_F^2} < 2 \frac{\sqrt{\|E_{12}\|_2^2 + \|E_{21}\|_2^2}}{\delta} \quad (1.13)$$

also as one might possibly expect. The appearances of  $m$  and  $n$  in (1.10) are particularly unsatisfactory for large (huge)  $m$  and usually modest  $r$ .

Our goal in this paper is to create a version of Theorem 1.1 that uses the spectral norm and any unitarily invariant norm. Our eventual version, Theorem 3.1, when restricted exclusively to the spectral norm, does not exactly coincide with the possible expectations in (1.12) and (1.13), but is very much close to it, namely (1.12) and (1.13) with  $\sqrt{\|E_{12}\|_2^2 + \|E_{21}\|_2^2}$  and  $\sqrt{\|T\|_2^2 + \|\Omega\|_2^2}$  replaced by  $\max\{\|E_{12}\|_2, \|E_{21}\|_2\}$  and  $\max\{\|T\|_2, \|\Omega\|_2\}$ , respectively. In particular,  $m$  and  $n$  disappear altogether.

The rest of this paper is organized as follows. Section 2 states two preliminary results for later use in our proofs. Our main result, a variant of Stewart's, is stated and proved in Section 3. In Section 4, we compare our main result realized for the spectral norm only and for both the spectral and Frobenius norm with Stewart's result in Theorem 1.1 and its potential consequences in (1.10) and (1.11). Finally, we draw our conclusions in Section 5.

**Notation.**  $\mathbb{C}^{n \times m}$  is the set of all  $n \times m$  complex matrices,  $\mathbb{C}^n = \mathbb{C}^{n \times 1}$ , and  $\mathbb{C} = \mathbb{C}^1$ .  $I_n$  (or simply  $I$  if its dimension is clear from the context) is the  $n \times n$  identity matrix. The superscript  $X^H$  is the complex conjugate transpose of a matrix or vector  $X$ . We shall also adopt MATLAB-like convention to access the entries of vectors and matrices. Let  $i : j$  be the set of integers from  $i$  to  $j$  inclusive.  $X_{(k:\ell, i:j)}$ ,  $X_{(k:\ell, :)}$ , and  $X_{(:, i:j)}$  are submatrices of  $X$ , consisting of intersections of row  $k$  to row  $\ell$  and column  $i$  to column  $j$ , row  $k$  to row  $\ell$ , and column  $i$  to column  $j$ , respectively.  $\mathcal{R}(B)$  is the column space of  $B$ , i.e., the subspace spanned by the columns of  $B$ . Finally  $\text{eig}(A)$  is the spectrum of a square matrix  $A$ .

We will continue to adopt the notation introduced so far in this section such as  $\|\cdot\|_2$  and  $\|\cdot\|_F$ . In particular,  $G \in \mathbb{C}^{m \times n}$  has  $\min\{m, n\}$  singular values denoted by  $\sigma_i(G)$  in decreasing order as in (1.2), and accordingly two singular value sets  $\text{sv}(G)$  and  $\text{sv}_{\text{ext}}(G)$  in (1.3), and  $\sigma_{\max}(G) := \sigma_1(G)$  and  $\sigma_{\min}(G) := \sigma_{\min\{m, n\}}(G)$ .

## 2 Preliminaries

In this section, we will state four lemmas that we will need later. The first lemma, Lemma 2.1, is due to Stewart [14, 15], and the second one, Lemma 2.2, summarizes known bounds on the solution to the Sylvester equation with Hermitian coefficient matrices of Davis and Kahan [6] and of Bhatia, Davis, and McIntosh [4]. The third one, Lemma 2.3, establishes bounds on the solution pair to a set of the coupled Sylvester equations and the results are new, except the one for the Frobenius norm. Finally, the fourth lemma, Lemma 2.4, is likely known.

**Lemma 2.1** ([14, Theorem 3.5], [15, Theorem 3.1]). *Let  $\mathbf{T}$  be a bounded linear operator on the Banach space  $(\mathcal{B}, \|\cdot\|)$  that has a bounded inverse. Let  $\phi$  be a continuous function on  $(\mathcal{B}, \|\cdot\|)$  that satisfies, for  $\mathbf{x}, \mathbf{y} \in \mathcal{B}$ ,*

$$(i) \quad \|\phi(\mathbf{x})\| \leq \hat{\varepsilon} \|\mathbf{x}\|^2,$$

$$(ii) \quad \|\phi(\mathbf{x}) - \phi(\mathbf{y})\| \leq 2\hat{\varepsilon} \max\{\|\mathbf{x}\|, \|\mathbf{y}\|\} \|\mathbf{x} - \mathbf{y}\|,$$

for some  $\hat{\varepsilon} \geq 0$ . Let  $0 < \hat{\delta} \leq \|\mathbf{T}^{-1}\|^{-1}$ . Given  $\mathbf{g} \in \mathcal{B}$ , if

$$\kappa_2 := (\hat{\varepsilon}/\hat{\delta}^2)\|\mathbf{g}\| < 1/4,$$

then equation  $\mathbf{T}\mathbf{x} = \mathbf{g} - \phi(\mathbf{x})$  has a solution  $\mathbf{x} \in \mathcal{B}$  that satisfies

$$\|\mathbf{x}\| \leq \frac{1 + \sqrt{1 - 4\kappa_2}}{1 - 2\kappa_2 + \sqrt{1 - 4\kappa_2}} \frac{\|\mathbf{g}\|}{\hat{\delta}} < 2 \frac{\|\mathbf{g}\|}{\hat{\delta}}.$$

We need the notation of unitarily invariant norm to go forward. A matrix norm  $\|\cdot\|_{\text{ui}}$  is called a *unitarily invariant norm* on  $\mathbb{C}^{m \times n}$  if it is a matrix norm and has the following two additional properties [3, 16]:

1.  $\|U^H B V\|_{\text{ui}} = \|B\|_{\text{ui}}$  for all unitary matrices  $U \in \mathbb{C}^{m \times m}$  and  $V \in \mathbb{C}^{n \times n}$  and  $B \in \mathbb{C}^{m \times n}$ ;
2.  $\|B\|_{\text{ui}} = \|B\|_2$ , the spectral norm of  $B$ , if  $\text{rank}(B) = 1$ .

Two commonly used unitarily invariant norms are

$$\begin{aligned} \text{the spectral norm:} \quad & \|B\|_2 = \max_j \sigma_j, \\ \text{the Frobenius norm:} \quad & \|B\|_F = \sqrt{\sum_j \sigma_j^2}, \end{aligned}$$

where  $\sigma_1, \sigma_2, \dots, \sigma_{\min\{m,n\}}$  are the singular values of  $B$ . In what follows,  $\|\cdot\|_{\text{ui}}$  denotes a general unitarily invariant norm.

In this article, for convenience, any  $\|\cdot\|_{\text{ui}}$  we use is generic to matrix sizes in the sense that it applies to matrices of all sizes. Examples include the matrix spectral norm  $\|\cdot\|_2$ , the Frobenius norm  $\|\cdot\|_F$ , and the trace norm. One important property of unitarily invariant norms is

$$\|XYZ\|_{\text{ui}} \leq \|X\|_2 \cdot \|Y\|_{\text{ui}} \cdot \|Z\|_2$$

for any matrices  $X$ ,  $Y$ , and  $Z$  of compatible sizes.

The next lemma summarizes known bounds on the solution of the Sylvester equation  $AX - XB = S$ . These bounds have played important roles in eigenspace variations as demonstrated by Davis and Kahan [6, 1970], and Bhatia, Davis, and McIntosh [4, 1983]. For a brief review, see [10].

**Lemma 2.2.** *Consider matrix equation  $XA - BX = S$ , where  $A \in \mathbb{R}^{r \times r}$  and  $B \in \mathbb{R}^{s \times s}$  are Hermitian, and  $S \in \mathbb{C}^{s \times r}$ . If  $\text{eig}(A) \cap \text{eig}(B) = \emptyset$ , then the equation has a unique solution  $X \in \mathbb{C}^{s \times r}$ . Furthermore, the following statements hold.*

(a) [6] we have

$$\|X\|_F \leq \|S\|_F / \delta, \quad \delta := \min_{\mu \in \text{eig}(A), \nu \in \text{eig}(B)} |\mu - \nu|; \quad (2.1)$$

(b) [6] if there exist  $\alpha < \beta$  and  $\delta > 0$  such that

$$\begin{aligned} \text{either} \quad & \text{eig}(A) \subset [\alpha, \beta] \text{ and } \text{eig}(B) \subset (-\infty, \alpha - \delta] \cup [\beta + \delta, \infty), \\ \text{or} \quad & \text{eig}(A) \subset (-\infty, \alpha - \delta] \cup [\beta + \delta, \infty) \text{ and } \text{eig}(B) \subset [\alpha, \beta], \end{aligned} \quad (2.2)$$

then for any unitarily invariant norm  $\|\cdot\|_{\text{ui}}$ ,

$$\|X\|_{\text{ui}} \leq \|S\|_{\text{ui}}/\delta; \quad (2.3)$$

(c) [4, 5] we have

$$\|X\|_{\text{ui}} \leq (\pi/2)\|S\|_{\text{ui}}/\delta, \quad (2.4)$$

where  $\delta$  is as in (2.1).

The results of Lemma 2.2 have an alternative interpretation through a linear operator

$$\mathbf{L} : X \in \mathbb{C}^{s \times r} \rightarrow \mathbf{L}(X) = XA - BX \in \mathcal{B}, \quad (2.5)$$

endowed with certain unitarily invariant norm  $\|\cdot\|_{\text{ui}}$  on  $\mathbb{C}^{s \times r}$ , including the spectral and Frobenius norms as special ones, where  $A \in \mathbb{R}^{r \times r}$  and  $B \in \mathbb{R}^{s \times s}$  are Hermitian. With any given  $\|\cdot\|_{\text{ui}}$ , there is an induced operator norm  $\|\cdot\|$  on the linear operator from  $\mathbb{C}^{s \times r}$  to itself. Translating the results of Lemma 2.2 yields the following corollary.

**Corollary 2.1.** *Let  $A \in \mathbb{R}^{r \times r}$  and  $B \in \mathbb{R}^{s \times s}$  be Hermitian, and define linear operator  $\mathbf{L}$  on  $(\mathbb{C}^{s \times r}, \|\cdot\|_{\text{ui}})$  as in (2.5). If  $\text{eig}(A) \cap \text{eig}(B) = \emptyset$ , then  $\mathbf{L}$  is invertible, and, furthermore, the following statements hold.*

- (i) *With  $\|\cdot\|_{\text{ui}} = \|\cdot\|_{\text{F}}$  and  $\delta$  as in (2.1), we have  $\|\mathbf{L}^{-1}\|^{-1} = \delta$ ;*
- (ii) *With general  $\|\cdot\|_{\text{ui}}$  and assuming (2.2), then  $\|\mathbf{L}^{-1}\|^{-1} = \delta$ , where  $\delta$  is the largest  $|\mu - \nu|$  for some  $\mu \in \text{eig}(A)$ ,  $\nu \in \text{eig}(B)$  and subject to (2.2);*
- (iii) *With general  $\|\cdot\|_{\text{ui}}$  and  $\delta$  as in (2.1), we have  $\|\mathbf{L}^{-1}\|^{-1} \geq (2/\pi)\delta$ .*

*Proof.* To see these, we note that  $\|\mathbf{L}^{-1}\|^{-1} = \min\{\gamma : \|\mathbf{L}(X)\| \geq \gamma\|(X)\|\}$ . Hence, immediately it follows that  $\|\mathbf{L}^{-1}\|^{-1} \geq \delta$  for items (i) and (ii) and  $\|\mathbf{L}^{-1}\|^{-1} \geq (2/\pi)\delta$  for item (iii). The equality sign in items (i) and (ii) are achieved by letting  $X = \mathbf{xy}^{\text{H}}$ , where  $\mathbf{x}$ ,  $\mathbf{y}$  are unit eigenvectors of  $A$  and  $B$ , respectively, such that  $A\mathbf{x} = \mu\mathbf{x}$  and  $B\mathbf{y} = \nu\mathbf{y}$ , assuming  $\delta = |\mu - \nu|$ , and hence

$$\mathbf{L}(X) = (\mu - \nu)\mathbf{xy}^{\text{H}} = (\mu - \nu)X.$$

Also because  $\text{sv}(X)$  consists of one nonzero singular value 1 and some copies of zeros,

$$\|X\|_{\text{ui}} = \|X\|_2 = 1,$$

implying  $\|\mathbf{L}(X)\|_{\text{ui}} = \delta\|X\|_{\text{ui}}$  for item (i) or item (ii). □

For subspace variation associated with SVD, a set of two coupled Sylvester equations come into play:

$$XA - BY = S, \quad YA^H - B^H X = T, \quad (2.6)$$

where  $A \in \mathbb{C}^{r \times r}$ ,  $B \in \mathbb{C}^{s \times t}$ ,  $S \in \mathbb{C}^{s \times r}$ , and  $T \in \mathbb{C}^{t \times r}$ . Note that  $B$  is possibly a nonsquare matrix, in which case we pad some blocks of zeros to  $B$ ,  $Y$ , and  $S$  or  $T$  to yield an equivalent set of two coupled Sylvester equations with  $B$  being square. Specifically, if  $s > t$ , let

$$B_{\text{ext}} = [B, \mathbf{0}_{s \times (s-t)}], \quad Y_{\text{ext}} = \begin{bmatrix} Y \\ \mathbf{0}_{(s-t) \times r} \end{bmatrix}, \quad S_{\text{ext}} = S, \quad T_{\text{ext}} = \begin{bmatrix} T \\ \mathbf{0}_{(s-t) \times r} \end{bmatrix},$$

whereas if  $s < t$ , let

$$B_{\text{ext}} = \begin{bmatrix} B \\ \mathbf{0}_{(t-s) \times t} \end{bmatrix}, \quad Y_{\text{ext}} = [Y, \mathbf{0}_{(t-s) \times r}], \quad S_{\text{ext}} = \begin{bmatrix} S \\ \mathbf{0}_{(t-s) \times r} \end{bmatrix}, \quad T_{\text{ext}} = T.$$

Then (2.6) is equivalent to

$$XA - B_{\text{ext}} Y_{\text{ext}} = S_{\text{ext}}, \quad Y_{\text{ext}} A^H - B_{\text{ext}}^H X = T_{\text{ext}}, \quad (2.7)$$

which can be merged into one Sylvester equation

$$\begin{bmatrix} 0 & X \\ Y_{\text{ext}} & 0 \end{bmatrix} \begin{bmatrix} 0 & A^H \\ A & 0 \end{bmatrix} - \begin{bmatrix} 0 & B_{\text{ext}} \\ B_{\text{ext}}^H & 0 \end{bmatrix} \begin{bmatrix} 0 & X \\ Y_{\text{ext}} & 0 \end{bmatrix} = \begin{bmatrix} S_{\text{ext}} & 0 \\ 0 & T_{\text{ext}} \end{bmatrix}. \quad (2.8)$$

It can be seen that

$$\text{eig} \left( \begin{bmatrix} 0 & A^H \\ A & 0 \end{bmatrix} \right) = \text{sv}(A) \cup (-\text{sv}(A)), \quad \text{eig} \left( \begin{bmatrix} 0 & B_{\text{ext}} \\ B_{\text{ext}}^H & 0 \end{bmatrix} \right) = \text{sv}_{\text{ext}}(B) \cup (-\text{sv}_{\text{ext}}(B)), \quad (2.9)$$

where negating a set means negating each element of the set, and

$$\left\| \begin{bmatrix} 0 & X \\ Y_{\text{ext}} & 0 \end{bmatrix} \right\|_{\text{ui}} = \left\| \begin{bmatrix} 0 & X \\ Y & 0 \end{bmatrix} \right\|_{\text{ui}}, \quad \left\| \begin{bmatrix} S_{\text{ext}} & 0 \\ 0 & T_{\text{ext}} \end{bmatrix} \right\|_{\text{ui}} = \left\| \begin{bmatrix} S & 0 \\ 0 & T \end{bmatrix} \right\|_{\text{ui}} \quad (2.10)$$

for any unitary invariant norm. Apply Lemma 2.2 to get

**Lemma 2.3.** *Consider a set of two coupled Sylvester equations (2.6), where  $A \in \mathbb{C}^{r \times r}$ ,  $B \in \mathbb{C}^{s \times t}$ ,  $S \in \mathbb{C}^{s \times r}$ , and  $T \in \mathbb{C}^{t \times r}$ . If  $\text{sv}(A) \cap \text{sv}_{\text{ext}}(B) = \emptyset$ , then the set of equations has a unique solution pair  $(X, Y) \in \mathbb{C}^{s \times r} \times \mathbb{C}^{t \times r}$ . Furthermore, the following statements hold:*

(a) *we have [15]*

$$\sqrt{\|X\|_{\mathbb{F}}^2 + \|Y\|_{\mathbb{F}}^2} \leq \sqrt{\|S\|_{\mathbb{F}}^2 + \|T\|_{\mathbb{F}}^2} / \delta, \quad \delta := \min_{\omega \in \text{sv}(A), \gamma \in \text{sv}_{\text{ext}}(B)} |\omega - \gamma|; \quad (2.11)$$

(b) if  $\delta := \sigma_{\min}(A) - \sigma_{\max}(B) > 0$ , then for any unitarily invariant norm  $\|\cdot\|_{\text{ui}}$ ,

$$\left\| \begin{bmatrix} 0 & X \\ Y & 0 \end{bmatrix} \right\|_{\text{ui}} \leq \frac{1}{\delta} \left\| \begin{bmatrix} S & 0 \\ 0 & T \end{bmatrix} \right\|_{\text{ui}}, \quad (2.12a)$$

$$\max\{\|X\|_{\text{ui}}, \|Y\|_{\text{ui}}\} \leq \frac{1}{\delta} \max\{\|S\|_{\text{ui}}, \|T\|_{\text{ui}}\}, \quad (2.12b)$$

and in particular for the spectral norm

$$\max\{\|X\|_2, \|Y\|_2\} \leq \frac{1}{\delta} \max\{\|S\|_2, \|T\|_2\}; \quad (2.13)$$

(c) we have

$$\left\| \begin{bmatrix} 0 & X \\ Y & 0 \end{bmatrix} \right\|_{\text{ui}} \leq \frac{\pi}{2} \frac{1}{\delta} \left\| \begin{bmatrix} S & 0 \\ 0 & T \end{bmatrix} \right\|_{\text{ui}}, \quad (2.14a)$$

$$\max\{\|X\|_{\text{ui}}, \|Y\|_{\text{ui}}\} \leq \pi \frac{1}{\delta} \max\{\|S\|_{\text{ui}}, \|T\|_{\text{ui}}\}, \quad (2.14b)$$

where  $\delta$  is as in (2.11), and in particular for the spectral norm

$$\max\{\|X\|_2, \|Y\|_2\} \leq \frac{\pi}{2} \frac{1}{\delta} \max\{\|S\|_2, \|T\|_2\}. \quad (2.15)$$

*Proof.* Recall (2.9) and (2.10). The inequality in (2.11) is essentially Stewart's [15, Theorem 6.2], but here as a corollary of Lemma 2.2 applied to (2.8). Inequalities (2.12a) and (2.14a) are also corollaries of Lemma 2.2 applied to (2.8). Inequalities (2.13) and (2.15) follow from (2.12) and (2.14a), respectively, due to

$$\left\| \begin{bmatrix} 0 & X \\ Y & 0 \end{bmatrix} \right\|_2 = \max\{\|X\|_2, \|Y\|_2\}, \quad \left\| \begin{bmatrix} S & 0 \\ 0 & T \end{bmatrix} \right\|_2 = \max\{\|S\|_2, \|T\|_2\}. \quad (2.16)$$

It remains to show (2.12b) and (2.14b). Inequality (2.12b) is essentially implied in [17, section 3], but we will present a quick proof anyway. Note that for the case  $\delta := \sigma_{\min}(A) - \sigma_{\max}(B) > 0$ . We have

$$\begin{aligned} \|XA - BY\|_{\text{ui}} &\geq \|XA\|_{\text{ui}} - \|BY\|_{\text{ui}} \\ &\geq \|X\|_{\text{ui}} \|A^{-1}\|_2^{-1} - \|B\|_2 \|Y\|_{\text{ui}} \\ &= \|X\|_{\text{ui}} \sigma_{\min}(A) - \sigma_{\max}(B) \|Y\|_{\text{ui}}. \end{aligned} \quad (2.17a)$$

Similarly, we can get

$$\|YA^{\text{H}} - B^{\text{H}}X\|_{\text{ui}} \geq \sigma_{\min}(A) \|Y\|_{\text{ui}} - \sigma_{\max}(B) \|X\|_{\text{ui}}. \quad (2.17b)$$

There are two cases to consider. If  $\|X\|_{\text{ui}} \geq \|Y\|_{\text{ui}}$ , then by (2.17a) we get

$$\|XA - BY\|_{\text{ui}} \geq \delta \|X\|_{\text{ui}} = \delta \max\{\|X\|_{\text{ui}}, \|Y\|_{\text{ui}}\}; \quad (2.18a)$$



if, on the other hand,  $\|X\|_{\text{ui}} < \|Y\|_{\text{ui}}$ , then by (2.17b) we get

$$\|YA^{\text{H}} - B^{\text{H}}X\|_{\text{ui}} \geq \delta \|Y\|_{\text{ui}} = \delta \max\{\|X\|_{\text{ui}}, \|Y\|_{\text{ui}}\}. \quad (2.18\text{b})$$

Together, (2.18a) and (2.18b) yield

$$\max\{\|XA - BY\|_{\text{ui}}, \|YA^{\text{H}} - B^{\text{H}}X\|_{\text{ui}}\} \geq \delta \max\{\|X\|_{\text{ui}}, \|Y\|_{\text{ui}}\},$$

as expected. Finally, noticing that

$$\begin{aligned} \max\{\|X\|_{\text{ui}}, \|Y\|_{\text{ui}}\} &\leq \left\| \begin{bmatrix} 0 & X \\ Y & 0 \end{bmatrix} \right\|_{\text{ui}}, \\ \left\| \begin{bmatrix} S & 0 \\ 0 & T \end{bmatrix} \right\|_{\text{ui}} &\leq \|S\|_{\text{ui}} + \|T\|_{\text{ui}} \\ &\leq 2 \max\{\|S\|_{\text{ui}}, \|T\|_{\text{ui}}\}, \end{aligned}$$

we see that (2.14a) implies (2.14b).  $\square$

Lemma 2.3 has more than what we need later. In fact, we will only use (2.13) and (2.15) in our later development.

The results of Lemma 2.3 have an alternative interpretation through a linear operator

$$\begin{aligned} \mathbf{T} : \mathcal{B} := \mathbb{C}^{s \times r} \times \mathbb{C}^{t \times r} &\rightarrow \mathbf{T}(X, Y) \in \mathcal{B} \\ (X, Y) &\rightarrow (XA - BY, YA^{\text{H}} - B^{\text{H}}X), \end{aligned} \quad (2.19)$$

endowed with certain norm on  $\mathcal{B}$  to make it a Banach space, including (cf. those used in Lemma 2.3)

$$\|(X, Y)\| = \left\| \begin{bmatrix} 0 & X \\ Y & 0 \end{bmatrix} \right\|_{\text{ui}} \equiv \left\| \begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix} \right\|_{\text{ui}} \quad (2.20\text{a})$$

for any  $(X, Y) \in \mathcal{B}$ , where  $\|\cdot\|_{\text{ui}}$  is any given unitarily invariant norm. Two particular ones are

$$\|(X, Y)\| = \max\{\|X\|_2, \|Y\|_2\} \quad \text{or} \quad \sqrt{\|X\|_{\text{F}}^2 + \|Y\|_{\text{F}}^2}, \quad (2.20\text{b})$$

upon realizing  $\|\cdot\|_{\text{ui}}$  in (2.20a) as the spectral norm or the Frobenius norm. Another possible endowed norm on  $\mathcal{B}$  is

$$\|(X, Y)\| = \max\{\|X\|_{\text{ui}}, \|Y\|_{\text{ui}}\}. \quad (2.20\text{c})$$

With each endowed norm, there is an induced operator norm  $\|\cdot\|$  on the linear operator from  $\mathcal{B}$  to itself. Translating the results of Lemma 2.3 yields the following corollary.

**Corollary 2.2.** *Let  $A \in \mathbb{C}^{r \times r}$ ,  $B \in \mathbb{C}^{s \times t}$ ,  $S \in \mathbb{C}^{s \times r}$ , and  $T \in \mathbb{C}^{t \times r}$ , and define linear operator  $\mathbf{T}$  on  $(\mathcal{B} := \mathbb{C}^{s \times r} \times \mathbb{C}^{t \times r}, \|\cdot\|)$  as in (2.5) where  $\|\cdot\|$  is given by one of those in (2.20). If  $\text{sv}(A) \cap \text{sv}_{\text{ext}}(B) = \emptyset$ , then  $\mathbf{T}$  is invertible, and, furthermore, the following statements hold.*

- (i) *With  $\|(X, Y)\| = \sqrt{\|X\|_{\text{F}}^2 + \|Y\|_{\text{F}}^2}$  and  $\delta$  as in (2.11), we have  $\|\mathbf{T}^{-1}\|^{-1} = \delta$  [15];*

- (ii) With either (2.20a) or (2.20c), if  $\delta := \sigma_{\min}(A) - \sigma_{\max}(B) > 0$ , then  $\|\mathbf{T}^{-1}\|^{-1} = \delta$ ;
- (iii) With (2.20a) and  $\delta$  as in (2.11), we have  $\|\mathbf{T}^{-1}\|^{-1} \geq (2/\pi)\delta$ ;
- (iv) With (2.20c) and  $\delta$  as in (2.11), we have  $\|\mathbf{T}^{-1}\|^{-1} \geq (1/\pi)\delta$ .

*Proof.* To see these, we note that  $\|\mathbf{T}^{-1}\|^{-1} = \min\{\gamma : \|\mathbf{T}(X, Y)\| \geq \gamma\|(X, Y)\|\}$ . Hence, immediately it follows that  $\|\mathbf{T}^{-1}\|^{-1} \geq \delta$  for items (i) and (ii),  $\|\mathbf{T}^{-1}\|^{-1} \geq (2/\pi)\delta$  for item (iii), and  $\|\mathbf{T}^{-1}\|^{-1} \geq (1/\pi)\delta$  for item (iv). The equality sign in items (i) and (ii) are achieved by letting  $X = \mathbf{y}\mathbf{u}^H$  and  $Y = \mathbf{x}\mathbf{v}^H$ , where  $\mathbf{u}, \mathbf{v}, \mathbf{x}, \mathbf{y}$  are unit singular vectors of  $A$  and  $B$ , respectively such that  $A\mathbf{u} = \sigma_{\min}(A)\mathbf{u}$  and  $B\mathbf{x} = \sigma_{\max}(B)\mathbf{x}$ , and hence

$$\mathbf{T}(X, Y) = [\sigma_{\min}(A) - \sigma_{\max}(B)](\mathbf{y}\mathbf{v}^H, \mathbf{x}\mathbf{u}^H).$$

Also because  $\text{sv}(X)$ ,  $\text{sv}(Y)$ ,  $\text{sv}(\mathbf{y}\mathbf{v}^H)$ , and  $\text{sv}(\mathbf{x}\mathbf{u}^H)$  all consist of one nonzero singular value 1 and some copies of zeros,

$$\begin{aligned} \left\| \begin{bmatrix} 0 & X \\ Y & 0 \end{bmatrix} \right\|_{\text{ui}} &\equiv \left\| \begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix} \right\|_{\text{ui}} = \left\| \begin{bmatrix} \mathbf{y}\mathbf{v}^H & 0 \\ 0 & \mathbf{x}\mathbf{u}^H \end{bmatrix} \right\|_{\text{ui}}, \\ \|X\|_{\text{ui}} = \|X\|_2 = 1, \quad \|Y\|_{\text{ui}} = \|Y\|_2 = 1, \\ \|\mathbf{y}\mathbf{v}^H\|_{\text{ui}} = \|\mathbf{y}\mathbf{v}^H\|_2 = 1, \quad \|\mathbf{x}\mathbf{u}^H\|_{\text{ui}} = \|\mathbf{x}\mathbf{u}^H\|_2 = 1, \end{aligned}$$

implying  $\|\mathbf{T}(X, Y)\| = \delta\|(X, Y)\|$  with the respective norm on  $\mathcal{B}$  as specified in item (i) or item (ii).  $\square$

The next lemma is likely known. We state it here with a proof for self-containedness.

**Lemma 2.4.** *Given  $B \in \mathbb{C}^{m \times n}$  and an integer  $1 \leq r < \min\{m, n\}$ , we have*

$$\begin{aligned} \sigma_r(B) &\geq \max\{\sigma_{\min}(B_{(:,1:r)}), \sigma_{\min}(B_{(1:r,:)})\} \geq \sigma_{\min}(B_{(1:r,1:r)}), \\ \sigma_{r+1}(B) &\leq \min\{\sigma_{\max}(B_{(:,r+1:n)}), \sigma_{\max}(B_{(r+1:m,:)})\}. \end{aligned}$$

*Proof.* Partition  $B$  as

$$B = \begin{matrix} & r & n-r \\ \begin{matrix} r \\ m-r \end{matrix} & \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix} \end{matrix},$$

and then we have

$$B^H B = \begin{bmatrix} B_{11}^H B_{11} + B_{21}^H B_{21} & B_{11}^H B_{12} + B_{21}^H B_{22} \\ B_{12}^H B_{11} + B_{22}^H B_{21} & B_{12}^H B_{12} + B_{22}^H B_{22} \end{bmatrix} \in \mathbb{C}^{n \times n}.$$

It follows from Fischer's minimax principle for the symmetric eigenvalue problem (see, e.g., [12, eq. (1.3)], [13, p.206], [16, p.201]) that

$$[\sigma_r(B)]^2 = \lambda_r(B^H B) = \max_{\dim \mathcal{X}=r} \min_{\mathbf{x} \in \mathcal{X} \subset \mathbb{C}^n} \frac{\mathbf{x}^H (B^H B) \mathbf{x}}{\mathbf{x}^H \mathbf{x}}$$

$$\begin{aligned}
&\geq \min_{\mathbf{z} \in \mathbb{C}^r} \frac{\mathbf{z}^H (B_{11}^H B_{11} + B_{21}^H B_{21}) \mathbf{z}}{\mathbf{z}^H \mathbf{z}} \\
&\geq \min_{\mathbf{z} \in \mathbb{C}^r} \frac{\mathbf{z}^H (B_{11}^H B_{11}) \mathbf{z}}{\mathbf{z}^H \mathbf{z}} = [\sigma_{\min}(B_{11})]^2, \tag{2.21}
\end{aligned}$$

where  $\mathcal{X} \subset \mathbb{C}^n$  denotes a subspace of  $\mathbb{C}^n$ ,  $\lambda_r(B^H B)$  is the  $r$ th largest eigenvalue of  $B^H B$ , and the first inequality is due to limiting the subspace to the one composed of vectors with their last  $n - r$  entries being to 0. This proves  $\sigma_r(B) \geq \sigma_{\min}(B_{(:,1:r)}) \geq \sigma_{\min}(B_{(1:r,1:r)})$ . Next, using  $[\sigma_r(B)]^2 = \lambda_r(BB^H)$ , in the same way as in (2.21), we can get  $\sigma_r(B) \geq \sigma_{\min}(B_{(1:r,:)}) \geq \sigma_{\min}(B_{(1:r,1:r)})$ . This completes the proof of the inequalities for  $\sigma_r(B)$ .

Analogously, again by Fischer's minimax principle, we have

$$\begin{aligned}
[\sigma_{r+1}(B)]^2 = \lambda_{r+1}(B^H B) &= \min_{\dim \mathcal{X} = n-r} \max_{\mathbf{x} \in \mathcal{X} \subset \mathbb{C}^n} \frac{\mathbf{x}^H (B^H B) \mathbf{x}}{\mathbf{x}^H \mathbf{x}} \\
&\leq \max_{\mathbf{z} \in \mathbb{C}^{n-r}} \frac{\mathbf{z}^H (B_{12}^H B_{12} + B_{22}^H B_{22}) \mathbf{z}}{\mathbf{z}^H \mathbf{z}} \\
&= \lambda_{\max}(B_{12}^H B_{12} + B_{22}^H B_{22}) = [\sigma_{\max}(B_{(:,r+1:n)})]^2, \tag{2.22}
\end{aligned}$$

where  $\lambda_{r+1}(B^H B)$  is the  $(r + 1)$ st largest eigenvalue of  $B^H B$ , and the first inequality is due to limiting the subspace to the one composed of vectors with their first  $r$  entries being to 0. Next, using  $[\sigma_{r+1}(B)]^2 = \lambda_{r+1}(BB^H)$ , in the same way as in (2.22), we can get  $\sigma_{r+1}(B) \leq \sigma_{\max}(B_{(r+1:m,:)})$ .  $\square$

### 3 Main Result

In order to achieve (1.6), we need some  $\Omega$  and  $\Gamma$  to satisfy

$$\Gamma(G_1 + E_{11}) - (G_2 + E_{22})\Omega = E_{21} - \Gamma E_{12}\Omega, \tag{3.1a}$$

$$\Omega(G_1 + E_{11})^H - (G_2 + E_{22})^H \Gamma = E_{12}^H - \Omega E_{21}^H \Gamma, \tag{3.1b}$$

obtained from setting off-diagonal blocks of  $\check{U}^H \check{G} \check{V}$ , partitioned accordingly, to 0.

In what follows, we will use Lemma 2.1 to prove the existence of a solution pair  $(\Gamma, \Omega) \in \mathbb{C}^{(m-r) \times r} \times \mathbb{C}^{(n-r) \times r}$  to (3.1) with an upper bound under certain conditions.

Keeping in mind that our goal is to create a variant of Theorem 1.1 in a unitarily invariant norm and the spectral norm, and hopefully the variant for the special case of the Frobenius norm is better than Theorem 1.1 in terms of both weaker conditions and stronger results. In the setting of Lemma 2.1, we will use the Banach space

$$\mathcal{B} := \mathbb{C}^{(m-r) \times r} \times \mathbb{C}^{(n-r) \times r}$$

endowed with one of the two norms: for  $(\Gamma, \Omega) \in \mathcal{B}$ ,

$$\|(\Gamma, \Omega)\| := \left\| \begin{bmatrix} 0 & \Gamma \\ \Omega & 0 \end{bmatrix} \right\|_{\text{ui}} \equiv \left\| \begin{bmatrix} \Gamma & 0 \\ 0 & \Omega \end{bmatrix} \right\|_{\text{ui}}, \tag{3.2a}$$

$$\|(\Gamma, \Omega)\| := \max\{\|\Gamma\|_{\text{ui}}, \|\Omega\|_{\text{ui}}\}. \quad (3.2b)$$

The two endowed norms become one for the case  $\|\cdot\|_{\text{ui}} = \|\cdot\|_2$ , the spectral norm, but otherwise are different. The linear operator  $\mathbf{T} : \mathcal{B} \rightarrow \mathcal{B}$  is given by

$$\mathbf{T}(\Gamma, \Omega) = \left( \Gamma(G_1 + E_{11}) - (G_2 + E_{22})\Omega, \Omega(G_1 + E_{11})^{\text{H}} - (G_2 + E_{22})^{\text{H}}\Gamma \right), \quad (3.3)$$

and the continuous function  $\phi : \mathcal{B} \rightarrow \mathcal{B}$  is

$$\phi((\Gamma, \Omega)) = (\Gamma E_{12}\Omega, \Omega E_{21}^{\text{H}}\Gamma). \quad (3.4)$$

It is not hard to verify that  $\|(\Gamma, \Omega)\|$  defined in (3.2) is indeed a norm on  $\mathcal{B}$ .

Compactly, the two equations in (3.1) can be merged into one to take the form

$$\mathbf{T}(\Gamma, \Omega) = (E_{21}, E_{12}^{\text{H}}) - \phi((\Gamma, \Omega)). \quad (3.5)$$

It remains to verify the conditions of Lemma 2.1 for  $\mathbf{T}$  and  $\phi$  we just defined. This is done in the next two lemmas. Let

$$\delta = \min_{\mu \in \text{sv}(G_1), \nu \in \text{sv}_{\text{ext}}(G_2)} |\mu - \nu|, \quad (3.6a)$$

$$\underline{\delta} = \delta - \|E_{11}\|_2 - \|E_{22}\|_2, \quad (3.6b)$$

$$\varepsilon = \max\{\|E_{12}\|_2, \|E_{21}\|_2\}. \quad (3.6c)$$

**Lemma 3.1.** *Suppose  $\underline{\delta} > 0$ . Dependent of different cases, we have for any  $(\Gamma, \Omega) \in \mathcal{B}$*

$$\|\mathbf{T}(\Gamma, \Omega)\| \geq \frac{\underline{\delta}}{c} \|(\Gamma, \Omega)\|, \quad (3.7)$$

which implies  $\|\mathbf{T}^{-1}\|^{-1} \geq \underline{\delta}/c$ , where  $\underline{\delta}$  is as in (3.6b), and

(i) with the endowed norm in (3.2a),

$$c = \begin{cases} 1, & \text{if } \sigma_{\min}(G_1) > \sigma_{\max}(G_2) \text{ or } \|\cdot\|_{\text{ui}} = \|\cdot\|_{\text{F}}, \\ \pi/2, & \text{otherwise;} \end{cases} \quad (3.8)$$

(ii) with the endowed norm in (3.2b),

$$c = \begin{cases} 1, & \text{if } \sigma_{\min}(G_1) > \sigma_{\max}(G_2), \\ \pi, & \text{otherwise.} \end{cases} \quad (3.9)$$

*Proof.* Notice that  $\mathbf{T}(\Gamma, \Omega) = (S, T)$  consists of a set of two coupled Sylvester equations in (2.6) with

$$A = G_1 + E_{11}, \quad B = G_2 + E_{22},$$

and  $\Gamma$  and  $\Omega$  correspond to  $X$  and  $Y$  there, respectively. We claim that

$$\underline{\delta} \leq \min_{\mu \in \text{sv}(G_1 + E_{11}), \nu \in \text{sv}_{\text{ext}}(G_2 + E_{22})} |\mu - \nu| = |\mu' - \nu'|. \quad (3.10)$$

where  $\mu' \in \text{sv}(G_1 + E_{11})$ ,  $\nu' \in \text{sv}_{\text{ext}}(G_2 + E_{22})$  achieve the minimum. By Mirsky's theorem [16, p.204], there are  $\mu \in \text{sv}(G_1)$ ,  $\nu \in \text{sv}_{\text{ext}}(G_2)$  such that  $|\mu' - \mu| \leq \|E_{11}\|_2$  and  $|\nu' - \nu| \leq \|E_{22}\|_2$ . Hence

$$\begin{aligned} |\mu' - \nu'| &= |\mu - \nu + (\mu' - \mu) - (\nu' - \nu)| \\ &\geq |\mu - \nu| - |\mu' - \mu| - |\nu' - \nu| \\ &\geq \delta - \|E_{11}\|_2 - \|E_{22}\|_2 = \underline{\delta}, \end{aligned}$$

yielding (3.10).

If  $\sigma_{\min}(G_1) > \sigma_{\max}(G_2)$ , then  $\delta = \sigma_{\min}(G_1) - \sigma_{\max}(G_2) > 0$ , and hence

$$\begin{aligned} 0 < \underline{\delta} &= \sigma_{\min}(G_1) - \sigma_{\max}(G_2) - \|E_{11}\|_2 - \|E_{22}\|_2 \\ &= [\sigma_{\min}(G_1) - \|E_{11}\|_2] - [\sigma_{\max}(G_2) + \|E_{22}\|_2]. \end{aligned}$$

Keep in mind that

$$\sigma_{\min}(G_1 + E_{11}) \geq \sigma_{\min}(G_1) - \|E_{11}\|_2, \quad \sigma_{\max}(G_2 + E_{22}) \leq \sigma_{\max}(G_2) + \|E_{22}\|_2$$

by Mirsky's theorem [16, p.204], and hence

$$\text{sv}(A) \subset [\sigma_{\min}(G_1) - \|E_{11}\|_2, \infty), \quad \text{sv}_{\text{ext}}(B) \subset [0, \sigma_{\max}(G_2) + \|E_{22}\|_2].$$

This lemma is a consequence of Lemma 2.3.  $\square$

**Remark 3.1.** There are a few comments in order, regarding the assumptions in Lemma 3.1 that ensure (3.7).

- 1) Always  $\text{sv}(G_1) \cap \text{sv}_{\text{ext}}(G_2) = \text{sv}(G_1 + E_{11}) \cap \text{sv}_{\text{ext}}(G_2 + E_{22}) = \emptyset$ , guaranteed by  $\underline{\delta} > 0$ ;
- 2) If  $m \neq n$ , then 0 is an element of both  $\text{sv}_{\text{ext}}(G_2)$  and  $\text{sv}_{\text{ext}}(G_2 + E_{22})$  and hence  $\sigma_{\min}(G_1) > 0$  and  $\sigma_{\min}(G_1 + E_{11}) > 0$ ;
- 3) Two different ways of separation between  $\text{sv}(G_1)$  and  $\text{sv}_{\text{ext}}(G_2)$  are assumed:
  - (i) simply  $\text{sv}(G_1) \cap \text{sv}_{\text{ext}}(G_2) = \emptyset$ ;
  - (ii)  $\sigma_{\min}(G_1) > \sigma_{\max}(G_2)$ , i.e, the interval  $[\sigma_{\max}(G_2), \sigma_{\min}(G_1)]$  separates  $\text{sv}(G_1)$  from  $\text{sv}_{\text{ext}}(G_2)$ .

Assumption (i) of separation is weaker than Assumption (ii) of separation. Each implies the same way of separation between  $\text{sv}(G_1 + E_{11})$  and  $\text{sv}_{\text{ext}}(G_2 + E_{22})$  by  $\underline{\delta} > 0$ ;

- 4) The endowed norm (3.2a) works with both assumptions of separation on the singular values. Correspondingly,  $c = 1$  always for the Frobenius norm, and  $c = 1$  under Assumption (ii) of separation above and  $\pi/2$  otherwise;
- 5) The endowed norm (3.2b) works with both assumptions of separation, too. Correspondingly,  $c = 1$  always for the Frobenius norm, and  $c = 1$  under Assumption (ii) of separation and  $\pi$  otherwise.

Finally, because of (2.16), items (i) and (ii) of Lemma 3.1 overlap at the case  $\|\cdot\|_{\text{ui}} = \|\cdot\|_2$  and  $\sigma_{\min}(G_1) > \sigma_{\max}(G_2)$ .

In what follows, our representation will assume that one of the endowed norm  $\|(\cdot, \cdot)\|$  in (3.2) is selected and fixed, and, along with it, the part of Lemma 3.1, unless explicitly stated otherwise.

**Lemma 3.2.** *For the continuous function  $\phi$  defined in (3.4), we have*

- (i)  $\|\phi((\Gamma, \Omega))\| \leq \varepsilon \|(\Gamma, \Omega)\|^2$ ,
- (ii)  $\|\phi((\Gamma, \Omega)) - \phi((\widehat{\Gamma}, \widehat{\Omega}))\| \leq 2\varepsilon \max\{\|(\Gamma, \Omega)\|, \|(\widehat{\Gamma}, \widehat{\Omega})\|\} \|(\Gamma - \widehat{\Gamma}, \Omega - \widehat{\Omega})\|$ ,

where  $\varepsilon = \max\{\|E_{12}\|_2, \|E_{21}\|_2\}$  is as in (3.6c).

*Proof.* With (3.2a), we have

$$\begin{aligned} \|\phi((\Gamma, \Omega))\| &= \left\| \begin{bmatrix} \Gamma E_{12} \Omega & 0 \\ 0 & \Omega E_{21}^H \Gamma \end{bmatrix} \right\|_{\text{ui}} = \left\| \begin{bmatrix} \Gamma & 0 \\ 0 & \Omega \end{bmatrix} \begin{bmatrix} E_{12} & 0 \\ 0 & E_{21}^H \end{bmatrix} \begin{bmatrix} \Omega & 0 \\ 0 & \Gamma \end{bmatrix} \right\|_{\text{ui}} \\ &\leq \left\| \begin{bmatrix} \Gamma & 0 \\ 0 & \Omega \end{bmatrix} \right\|_{\text{ui}} \left\| \begin{bmatrix} E_{12} & 0 \\ 0 & E_{21}^H \end{bmatrix} \right\|_2 \left\| \begin{bmatrix} \Omega & 0 \\ 0 & \Gamma \end{bmatrix} \right\|_{\text{ui}} \\ &= \max\{\|E_{12}\|_2, \|E_{21}\|_2\} \times \|(\Gamma, \Omega)\|^2; \end{aligned} \quad (3.11)$$

with (3.2b), we have

$$\begin{aligned} \|\phi((\Gamma, \Omega))\| &= \max\{\|\Gamma E_{12} \Omega\|_{\text{ui}}, \|\Omega E_{21}^H \Gamma\|_{\text{ui}}\} \\ &\leq \max\{\|E_{12}\|_2, \|E_{21}\|_2\} \times \|\Gamma\|_{\text{ui}} \|\Omega\|_{\text{ui}} \\ &\leq \max\{\|E_{12}\|_2, \|E_{21}\|_2\} \times \|(\Gamma, \Omega)\|^2. \end{aligned} \quad (3.12)$$

This proves the inequality in item (i). For item (ii), we note

$$\phi((\Gamma, \Omega)) - \phi((\widehat{\Gamma}, \widehat{\Omega})) = \left( \Gamma E_{12} \Omega - \widehat{\Gamma} E_{12} \widehat{\Omega}, \Omega E_{21}^H \Gamma - \widehat{\Omega} E_{21}^H \widehat{\Gamma} \right).$$

For each of the two components, we have

$$\Gamma E_{12} \Omega - \widehat{\Gamma} E_{12} \widehat{\Omega} = (\Gamma - \widehat{\Gamma}) E_{12} \Omega + \widehat{\Gamma} E_{12} (\Omega - \widehat{\Omega}), \quad (3.13a)$$

$$\Omega E_{21}^H \Gamma - \widehat{\Omega} E_{21}^H \widehat{\Gamma} = (\Omega - \widehat{\Omega}) E_{21}^H \Gamma + \widehat{\Omega} E_{21}^H (\Gamma - \widehat{\Gamma}). \quad (3.13b)$$

With (3.2a), we get, similar to the derivation in (3.11),

$$\begin{aligned} &\|\phi((\Gamma, \Omega)) - \phi((\widehat{\Gamma}, \widehat{\Omega}))\| \\ &= \left\| \begin{bmatrix} (\Gamma - \widehat{\Gamma}) E_{12} \Omega + \widehat{\Gamma} E_{12} (\Omega - \widehat{\Omega}) & 0 \\ 0 & (\Omega - \widehat{\Omega}) E_{21}^H \Gamma + \widehat{\Omega} E_{21}^H (\Gamma - \widehat{\Gamma}) \end{bmatrix} \right\|_{\text{ui}} \\ &\leq \left\| \begin{bmatrix} (\Gamma - \widehat{\Gamma}) E_{12} \Omega & 0 \\ 0 & (\Omega - \widehat{\Omega}) E_{21}^H \Gamma \end{bmatrix} \right\|_{\text{ui}} + \left\| \begin{bmatrix} \widehat{\Gamma} E_{12} (\Omega - \widehat{\Omega}) & 0 \\ 0 & \widehat{\Omega} E_{21}^H (\Gamma - \widehat{\Gamma}) \end{bmatrix} \right\|_{\text{ui}} \end{aligned}$$

$$\begin{aligned}
&\leq \varepsilon \|(\Gamma - \widehat{\Gamma}, \Omega - \widehat{\Omega})\| (\|(\Gamma, \Omega)\| + \|(\widehat{\Gamma}, \widehat{\Omega})\|) \\
&\leq 2\varepsilon \max\{\|(\Gamma, \Omega)\|, \|(\widehat{\Gamma}, \widehat{\Omega})\|\} \|(\Gamma - \widehat{\Gamma}, \Omega - \widehat{\Omega})\|;
\end{aligned}$$

with (3.2b), we get, similar to the derivation in (3.12),

$$\begin{aligned}
&\max\{\|\Gamma E_{12}\Omega - \widehat{\Gamma} E_{12}\widehat{\Omega}\|_{\text{ui}}, \|\Omega E_{21}^{\text{H}}\Gamma - \widehat{\Omega} E_{21}^{\text{H}}\widehat{\Gamma}\|_{\text{ui}}\} \\
&\leq \varepsilon \max\{\|\Gamma - \widehat{\Gamma}\|_{\text{ui}}\|\Omega\|_{\text{ui}} + \|\widehat{\Gamma}\|_{\text{ui}}\|\Omega - \widehat{\Omega}\|_{\text{ui}}, \|\Omega - \widehat{\Omega}\|_{\text{ui}}\|\Gamma\|_{\text{ui}} + \|\widehat{\Omega}\|_{\text{ui}}\|\Gamma - \widehat{\Gamma}\|_{\text{ui}}\} \\
&\leq \varepsilon \max\{\|\Gamma - \widehat{\Gamma}\|_{\text{ui}}, \|\Omega - \widehat{\Omega}\|_{\text{ui}}\} \cdot \max\{\|\Omega\|_{\text{ui}} + \|\widehat{\Gamma}\|_{\text{ui}}, \|\Gamma\|_{\text{ui}} + \|\widehat{\Omega}\|_{\text{ui}}\} \\
&\leq \varepsilon \max\{\|\Gamma - \widehat{\Gamma}\|_{\text{ui}}, \|\Omega - \widehat{\Omega}\|_{\text{ui}}\} \cdot 2 \max\{\|(\Gamma, \Omega)\|, \|(\widehat{\Gamma}, \widehat{\Omega})\|\} \\
&= 2\varepsilon \max\{\|(\Gamma, \Omega)\|, \|(\widehat{\Gamma}, \widehat{\Omega})\|\} \|(\Gamma - \widehat{\Gamma}, \Omega - \widehat{\Omega})\|,
\end{aligned}$$

completing the proof of item (ii).  $\square$

Next we apply Lemma 2.1 to ensure a particular solution  $(\Gamma, \Omega) \in \mathcal{B}$  to (3.5).

**Lemma 3.3.** *Let  $\delta$ ,  $\underline{\delta}$ , and  $\varepsilon$  be defined as in (3.6), and suppose the conditions in Lemma 3.1 that ensure (3.7) with constant  $c$  as specified there. If*

$$\underline{\delta} > 0 \quad \text{and} \quad \kappa_2 := \frac{c^2 \varepsilon}{\underline{\delta}^2} \|(E_{21}, E_{12}^{\text{H}})\| < \frac{1}{4}, \quad (3.14)$$

then (3.5) has a solution  $(\Gamma, \Omega)$  that satisfies

$$\|(\Gamma, \Omega)\| \leq \frac{1 + \sqrt{1 - 4\kappa_2}}{1 - 2\kappa_2 + \sqrt{1 - 4\kappa_2}} \frac{c\|(E_{21}, E_{12}^{\text{H}})\|}{\underline{\delta}} < 2 \frac{c\|(E_{21}, E_{12}^{\text{H}})\|}{\underline{\delta}}. \quad (3.15)$$

Here  $\|(E_{21}, E_{12}^{\text{H}})\|$  and  $\|(\Gamma, \Omega)\|$  are understood as the same one of the endowed norms in (3.2) under consideration.

*Proof.* In light of Lemmas 3.1 and 3.2, we find the conclusion is a straightforward consequence of Lemma 2.1 with  $\hat{\varepsilon} = \varepsilon$ ,  $\hat{\delta} = \underline{\delta}/c$ , and  $\|\mathbf{g}\| = \|(E_{21}, E_{12}^{\text{H}})\|$ .  $\square$

Finally, we state the main result of this section.

**Theorem 3.1.** *Given  $G, \widetilde{G} \in \mathbb{C}^{m \times n}$ , let  $G$  be decomposed as in (1.1), and partition  $U^{\text{H}}\widetilde{G}V$  according to (1.4). Let  $\delta$ ,  $\underline{\delta}$ , and  $\varepsilon$  be defined as in (3.6). If (3.14) is satisfied, then the following statements hold:*

- (a) *there exists a solution  $(\Gamma, \Omega) \in \mathbb{C}^{(m-r) \times r} \times \mathbb{C}^{(n-r) \times r}$  to (3.1), satisfying (3.15);*
- (b)  *$\widetilde{G}$  admits the decomposition (1.6), and the singular values of  $\widetilde{G}$  is the multiset union of those of*

$$\begin{aligned} \check{G}_1 &= \check{U}_1^{\text{H}} \widetilde{G} \check{V}_1 \\ &= (I + \Gamma^{\text{H}}\Gamma)^{1/2} (G_1 + E_{11} + E_{12}\Omega) (I + \Omega^{\text{H}}\Omega)^{-1/2} \end{aligned} \quad (3.16a)$$

$$= (I + \Gamma^{\text{H}}\Gamma)^{-1/2} (G_1 + E_{11} + \Gamma^{\text{H}}E_{21}) (I + \Omega^{\text{H}}\Omega)^{1/2}, \quad (3.16b)$$

and

$$\begin{aligned}\check{G}_2 &= \check{U}_2^H \check{G} \check{V}_2 \\ &= (I + \Gamma \Gamma^H)^{1/2} (G_2 + E_{22} - E_{21} \Omega^H) (I + \Omega \Omega^H)^{-1/2}\end{aligned}\quad (3.17a)$$

$$= (I + \Gamma \Gamma^H)^{-1/2} (G_2 + E_{22} - \Gamma E_{12}) (I + \Omega \Omega^H)^{1/2}; \quad (3.17b)$$

(c) We have

$$\sigma_{\min}(\check{G}_1) \geq \sigma_{\min}(G_1) - \|E_{11}\|_2 - 2c \frac{\varepsilon \|(E_{21}, E_{12}^H)\|}{\delta}, \quad (3.18a)$$

$$\sigma_{\max}(\check{G}_2) \leq \sigma_{\max}(G_2) + \|E_{22}\|_2 + 2c \frac{\varepsilon \|(E_{21}, E_{12}^H)\|}{\delta}, \quad (3.18b)$$

where  $\sigma_{\min}(\check{G}_1)$  and  $\sigma_{\max}(\check{G}_2)$  are the smallest singular value of  $\check{G}_1$  and the largest singular value of  $\check{G}_2$ , respectively;

(d) The left and right singular subspaces of  $\check{G}$  associated with the part of its singular values  $\text{sv}(\check{G})$  are spanned by the columns of

$$\begin{aligned}\check{U}_1 &= (U_1 + U_2 \Gamma) (I + \Gamma^H \Gamma)^{-1/2}, \\ \check{V}_1 &= (V_1 + V_2 \Omega) (I + \Omega^H \Omega)^{-1/2},\end{aligned}$$

respectively. In particular,

$$\|\check{U}_1 - U_1\|_{\text{ui}} \leq \|\Gamma\|_{\text{ui}} \leq \|(\Gamma, \Omega)\| \leq 2 \frac{c \|(E_{21}, E_{12}^H)\|}{\delta}, \quad (3.20a)$$

$$\|\check{V}_1 - V_1\|_{\text{ui}} \leq \|\Omega\|_{\text{ui}} \leq \|(\Gamma, \Omega)\| \leq 2 \frac{c \|(E_{21}, E_{12}^H)\|}{\delta}. \quad (3.20b)$$

*Proof.* Only item (c) and the inequalities in (3.20) of item (d) need proofs. For item (c), using (3.16a) for  $\check{G}_1$  and (3.16b) for  $\check{G}_1$  in  $\check{G}_1^H$  below, we get

$$\check{G}_1 \check{G}_1^H = (I + \Gamma^H \Gamma)^{1/2} (G_1 + E_{11} + E_{12} \Omega) (G_1 + E_{11} + \Gamma^H E_{21})^H (I + \Gamma^H \Gamma)^{-1/2}.$$

Hence

$$[\sigma_{\min}(\check{G}_1)]^2 = \lambda_{\min}(\check{G}_1 \check{G}_1^H) = \lambda_{\min}((G_1 + E_{11} + E_{12} \Omega) (G_1 + E_{11} + \Gamma^H E_{21})^H),$$

where  $\lambda_{\min}(\cdot)$  is the smallest eigenvalues of a Hermitian matrix. Therefore, with the help of (3.15), we have

$$\begin{aligned}[\sigma_{\min}(\check{G}_1)]^2 &= \left\| [(G_1 + E_{11} + E_{12} \Omega) (G_1 + E_{11} + \Gamma^H E_{21})^H]^{-1} \right\|_2^{-1} \\ &\geq \|(G_1 + E_{11} + \Gamma^H E_{21})^{-H}\|_2^{-1} \|(G_1 + E_{11} + E_{12} \Omega)^{-1}\|_2^{-1} \\ &\geq (\sigma_{\min}(G_1) - \|E_{11}\|_2 - \|\Gamma\|_2 \|E_{21}\|_2) (\sigma_{\min}(G_1) - \|E_{11}\|_2 - \|E_{12}\|_2 \|\Omega\|_2)\end{aligned}$$



$$\begin{aligned}
&\geq \left( \sigma_{\min}(G_1) - \|E_{11}\|_2 - \|( \Gamma, \Omega )\|_\varepsilon \right)^2 \\
&\geq \left( \sigma_{\min}(G_1) - \|E_{11}\|_2 - 2c \frac{\varepsilon \|(E_{21}, E_{12}^H)\|}{\underline{\delta}} \right)^2,
\end{aligned} \tag{3.21}$$

yielding the first inequality in (3.18). Similarly, we get the second inequality there by using (3.17).

Now we prove the inequalities in (3.20). We have

$$\begin{aligned}
\check{U}_1 - U_1 &= U_1 [(I + \Gamma^H \Gamma)^{-1/2} - I] + U_2 \Gamma (I + \Gamma^H \Gamma)^{-1/2} \\
&= [-U_1, U_2] \begin{bmatrix} I - (I + \Gamma^H \Gamma)^{-1/2} \\ \Gamma (I + \Gamma^H \Gamma)^{-1/2} \end{bmatrix}.
\end{aligned}$$

Let  $\Gamma = Z \Xi W^H$  be the SVD of  $\Gamma$ . We find

$$\begin{bmatrix} I - (I + \Gamma^H \Gamma)^{-1/2} \\ \Gamma (I + \Gamma^H \Gamma)^{-1/2} \end{bmatrix} = \begin{bmatrix} W & \\ & Z \end{bmatrix} \begin{bmatrix} I - (I + \Xi^H \Xi)^{-1/2} \\ \Xi (I + \Xi^H \Xi)^{-1/2} \end{bmatrix} W^H,$$

where for the middle matrix on the right,  $I - (I + \Xi^H \Xi)^{-1/2}$  is diagonal and  $\Xi (I + \Xi^H \Xi)^{-1/2}$  is leading diagonal. Hence the singular values of the middle matrix are given by: for each singular value  $\gamma$  of  $\Gamma$ ,

$$\sqrt{\left(1 - \frac{1}{\sqrt{1 + \gamma^2}}\right)^2 + \left(\frac{\gamma}{\sqrt{1 + \gamma^2}}\right)^2} = \sqrt{2 \left(1 - \frac{1}{\sqrt{1 + \gamma^2}}\right)} \tag{3.22}$$

$$\begin{aligned}
&= \frac{\sqrt{2} \gamma}{[\sqrt{1 + \gamma^2}(\sqrt{1 + \gamma^2} + 1)]^{1/2}} \\
&\leq \gamma.
\end{aligned} \tag{3.23}$$

Therefore, we get<sup>1</sup>

$$\|\check{U}_1 - U_1\|_{\text{ui}} = \left\| \begin{bmatrix} I - (I + \Xi^H \Xi)^{-1/2} \\ \Xi (I + \Xi^H \Xi)^{-1/2} \end{bmatrix} \right\|_{\text{ui}} \leq \|\Gamma\|_{\text{ui}},$$

yielding (3.20a) in light of (3.15). Similarly, we have (3.20b).  $\square$

The lower and upper bound on  $\sigma_{\min}(\check{G}_1)$  and  $\sigma_{\max}(\check{G}_2)$ , respectively, in (3.18), although always true, do not provide useful information, unless also  $\sigma_{\min}(G_1) > \sigma_{\max}(G_2)$ , in which case it can be used to establish a sufficient condition to ensure  $\sigma_{\min}(\check{G}_1) > \sigma_{\max}(\check{G}_2)$ .

<sup>1</sup>By (3.22) and (3.23), we conclude that for the spectral norm

$$\|\check{U}_1 - U_1\|_2 = \frac{\sqrt{2} \|\Gamma\|_2}{[\sqrt{1 + \|\Gamma\|_2^2}(\sqrt{1 + \|\Gamma\|_2^2} + 1)]^{1/2}}, \quad \|\check{V}_1 - V_1\|_2 = \frac{\sqrt{2} \|\Omega\|_2}{[\sqrt{1 + \|\Omega\|_2^2}(\sqrt{1 + \|\Omega\|_2^2} + 1)]^{1/2}}.$$

**Corollary 3.1.** *Given  $G, \tilde{G} \in \mathbb{C}^{m \times n}$ , let  $G$  be decomposed as in (1.1), and partition  $U^H \tilde{G} V$  according to (1.4). Let  $\delta, \underline{\delta}$ , and  $\varepsilon$  be defined as in (3.6), and suppose  $\sigma_{\min}(G_1) > \sigma_{\max}(G_2)$ . If (3.14) with  $c = 1$  is satisfied, then the top  $r$  singular values of  $\tilde{G}$  are exactly the  $r$  singular values of  $\check{G}_1$ , and*

$$\sigma_{\min}(G_1) - \|E_{11}\|_2 \leq \sigma_{\min}(\check{G}_1) \leq \sigma_{\min}(G_1) + \|E_{11}\|_2 + \frac{2\varepsilon^2}{\underline{\delta} + \sqrt{\underline{\delta}^2 + 4\varepsilon^2}}. \quad (3.24)$$

*Proof.* We have all conclusions of Theorem 3.1. By (3.18), we have

$$\sigma_{\min}(\check{G}_1) - \sigma_{\max}(\check{G}_2) \geq \underline{\delta} - 4 \frac{\varepsilon \|(E_{21}, E_{12}^H)\|}{\underline{\delta}} = \underline{\delta} \left[ 1 - 4 \frac{\varepsilon \|(E_{21}, E_{12}^H)\|}{\underline{\delta}^2} \right] > 0,$$

which says that the top  $r$  singular values of  $\tilde{G}$  are exactly the  $r$  singular values of  $\check{G}_1$ . As a result, applying Lemma 2.4 to (1.4), we have

$$\sigma_{\min}(\check{G}_1) = \sigma_r(U^H \tilde{G} V) \geq \sigma_{\min}(G_1 + E_{11}) \geq \sigma_{\min}(G_1) - \|E_{11}\|_2, \quad (3.25)$$

where the last inequality in (3.25) is a consequence of Mirsky's theorem [16, p.204]. This proves the first inequality in (3.24). It can be seen that

$$\sigma_{\min}(G_1 + E_{11}) - \sigma_{\max}(G_2 + E_{22}) \geq \sigma_{\min}(G_1) - \|E_{11}\|_2 - [\sigma_{\max}(G_2) + \|E_{22}\|_2] = \underline{\delta},$$

and hence by [7, Theorem 3] combining with (3.25), we get

$$\sigma_{\min}(\check{G}_1) - \sigma_{\min}(G_1 + E_{11}) \leq \frac{2\varepsilon^2}{\underline{\delta} + \sqrt{\underline{\delta}^2 + 4\varepsilon^2}}, \quad (3.26)$$

yielding the second inequality in (3.24).  $\square$

The sharper lower bound on  $\sigma_{\min}(\check{G}_1)$  in (3.24) than the one by the first inequality in (3.18) is only made possible by first establishing that the top  $r$  singular values of  $\tilde{G}$  are exactly the  $r$  singular values of  $\check{G}_1$ , which relies on the first inequality in (3.18) for a proof, however. More than (3.26) which is just for the smallest singular value only, [7, Theorem 3] yields

$$|\sigma_i(\check{G}_1) - \sigma_i(G_1 + E_{11})| \leq \frac{2\varepsilon^2}{\underline{\delta} + \sqrt{\underline{\delta}^2 + 4\varepsilon^2}} \quad \text{for } 1 \leq i \leq r$$

under the conditions of Corollary 3.1.

## 4 Discussions

Theorem 3.1 serves the same purpose as Stewart's [15, Theorem 6.4], but the former provides a variety of choices of norms for the Banach space  $\mathcal{B}$  and, as a consequence, a number of results dependent of circumstances to use.

Let us begin by realizing Theorem 3.1 and Corollary 3.1 with unitarily invariant norm  $\|\cdot\|_{\text{ui}}$  being set to the Frobenius norm, in order to compare with Theorem 1.1 [15, Theorem 6.4]. We have two endowed norms from (3.2) to choose:

$$\|(\Gamma, \Omega)\| = \sqrt{\|\Gamma\|_{\text{F}}^2 + \|\Omega\|_{\text{F}}^2} \quad \text{or} \quad \|(\Gamma, \Omega)\| = \max\{\|\Gamma\|_{\text{F}}, \|\Omega\|_{\text{F}}\}.$$

We have the following corollary to Theorem 3.1, where we state the conditions and the bounds on  $\|(\Gamma, \Omega)\|$  but omit the others that can be deduced from the bounds on  $\|(\Gamma, \Omega)\|$ .

**Corollary 4.1.** *Given  $G, \tilde{G} \in \mathbb{C}^{m \times n}$ , let  $G$  be decomposed as in (1.1), and partition  $U^{\text{H}}\tilde{G}V$  according to (1.4). Let  $\delta, \underline{\delta}$ , and  $\varepsilon$  be defined as in (3.6).*

(i) *If*

$$\underline{\delta} > 0 \quad \text{and} \quad \kappa_2 := \frac{\varepsilon}{\underline{\delta}^2} \sqrt{\|E_{12}\|_{\text{F}}^2 + \|E_{21}\|_{\text{F}}^2} < \frac{1}{4}, \quad (4.1)$$

*then (3.5) has a solution  $(\Gamma, \Omega)$  that satisfies*

$$\begin{aligned} \sqrt{\|\Gamma\|_{\text{F}}^2 + \|\Omega\|_{\text{F}}^2} &\leq \frac{1 + \sqrt{1 - 4\kappa_2}}{1 - 2\kappa_2 + \sqrt{1 - 4\kappa_2}} \frac{\sqrt{\|E_{12}\|_{\text{F}}^2 + \|E_{21}\|_{\text{F}}^2}}{\underline{\delta}} \\ &< 2 \frac{\sqrt{\|E_{12}\|_{\text{F}}^2 + \|E_{21}\|_{\text{F}}^2}}{\underline{\delta}}. \end{aligned} \quad (4.2)$$

(ii) *If  $\sigma_{\min}(G_1) > \sigma_{\max}(G_2)$  and if*

$$\underline{\delta} > 0 \quad \text{and} \quad \kappa_2 := \frac{\varepsilon}{\underline{\delta}^2} \max\{\|E_{12}\|_{\text{F}}, \|E_{21}\|_{\text{F}}\} < \frac{1}{4}, \quad (4.3)$$

*then (3.5) has a solution  $(\Gamma, \Omega)$  that satisfies*

$$\begin{aligned} \max\{\|\Gamma\|_{\text{F}}, \|\Omega\|_{\text{F}}\} &\leq \frac{1 + \sqrt{1 - 4\kappa_2}}{1 - 2\kappa_2 + \sqrt{1 - 4\kappa_2}} \frac{\max\{\|E_{12}\|_{\text{F}}, \|E_{21}\|_{\text{F}}\}}{\underline{\delta}} \\ &< 2 \frac{\max\{\|E_{12}\|_{\text{F}}, \|E_{21}\|_{\text{F}}\}}{\underline{\delta}}. \end{aligned} \quad (4.4)$$

In comparing Corollary 4.1 with Theorem 1.1 [15, Theorem 6.4], we note Corollary 4.1(i) provides better results than Theorem 1.1 in their conditions: (4.1) *vs.* (1.8), and bounds: (4.2) *vs.* (1.9), because

$$\begin{aligned} \hat{\varepsilon}^2 = \|E_{12}\|_{\text{F}}^2 + \|E_{21}\|_{\text{F}}^2 &\geq \max\{\|E_{12}\|_2, \|E_{21}\|_2\} \sqrt{\|E_{12}\|_{\text{F}}^2 + \|E_{21}\|_{\text{F}}^2} \\ &= \varepsilon \sqrt{\|E_{12}\|_{\text{F}}^2 + \|E_{21}\|_{\text{F}}^2}. \end{aligned}$$

Such an improvement of Corollary 4.1(i) over Theorem 1.1 could be considered marginal because it can be easily recovered by just refining Stewart's relevant estimates [15]. However, the improvement by Corollary 4.1(ii) over Theorem 1.1, under the condition  $\sigma_{\min}(G_1) > \sigma_{\max}(G_2)$ , unlikely can be achieved by simply refining Stewart's arguments there.

Our original motivation to revisit this classical result of Stewart's is the need for a version of Theorem 1.1 in the spectral norm while we are working on an error analysis for model order reduction by balanced truncation [18]. Let us look at what Theorem 3.1 leads to for the spectral norm, for which the two endowed norms from (3.2) collapse to the same one

$$\|(Γ, Ω)\| = \max\{\|Γ\|_2, \|Ω\|_2\}.$$

We have the following corollary to Theorem 3.1, which yields far sharper results than those such as (1.11) that would otherwise have to be derived from Theorem 1.1.

**Corollary 4.2.** *Given  $G, \tilde{G} \in \mathbb{C}^{m \times n}$ , let  $G$  be decomposed as in (1.1), and partition  $U^H \tilde{G} V$  according to (1.4). Let  $\delta, \underline{\delta}$ , and  $\varepsilon$  be defined as in (3.6).*

(i) *If*

$$\underline{\delta} > 0 \quad \text{and} \quad \kappa_2 := \left(\frac{\pi}{2}\right)^2 \frac{\varepsilon^2}{\underline{\delta}^2} < \frac{1}{4}, \quad (4.5)$$

*then (3.5) has a solution  $(Γ, Ω)$  that satisfies*

$$\max\{\|Γ\|_2, \|Ω\|_2\} \leq \frac{1 + \sqrt{1 - 4\kappa_2}}{1 - 2\kappa_2 + \sqrt{1 - 4\kappa_2}} \frac{\pi \varepsilon}{2 \underline{\delta}} < \pi \frac{\varepsilon}{\underline{\delta}}. \quad (4.6)$$

(ii) *If  $\sigma_{\min}(G_1) > \sigma_{\max}(G_2)$  and if*

$$\delta > 0 \quad \text{and} \quad \kappa_2 := \frac{\varepsilon^2}{\delta^2} < \frac{1}{4}, \quad (4.7)$$

*then (3.5) has a solution  $(Γ, Ω)$  that satisfies*

$$\max\{\|Γ\|_2, \|Ω\|_2\} \leq \frac{1 + \sqrt{1 - 4\kappa_2}}{1 - 2\kappa_2 + \sqrt{1 - 4\kappa_2}} \frac{\varepsilon}{\delta} < 2 \frac{\varepsilon}{\delta}. \quad (4.8)$$

During the proof of Lemma 2.3, we commented that inequality (2.12b) had been really implied by Wedin [17, section 3], and inequality (2.11) may also be essentially implied in [17] but with his  $\delta$  defined as

$$\delta := \min \left\{ \min_{\mu \in \text{sv}(A), \nu \in \text{sv}(B)} |\mu - \nu|, \sigma_{\min}(A) \right\} \quad (4.9)$$

which is the same as the one in (2.11) for the case  $s \neq t$  because then  $0 \in \text{sv}_{\text{ext}}(B)$ , but can be different if  $s = t$  for which case  $\text{sv}_{\text{ext}}(B) = \text{sv}(B)$  that may or may not contain 0, however. Both (2.12b) and (2.11) are used to develop *the generalized sin  $\theta$  theorems* for SVD there (see, also, [11, p.21-7]). In the same spirit, we can use (2.12a) and (2.14) to establish more generalized sin  $\theta$  theorems for SVD. In fact, we have the following theorem.

**Theorem 4.1.** *Let  $G \in \mathbb{C}^{m \times n}$  be decomposed as in (1.1), and let  $\tilde{G} \in \mathbb{C}^{m \times n}$  admit a decomposition in the same form as in (1.1), except with tildes on all symbols. Let*

$$R = G \tilde{V}_1 - \tilde{U}_1 \tilde{G}_1, \quad S = G^H \tilde{U}_1 - \tilde{V}_1 \tilde{G}_1^H.$$

If

$$\delta := \min_{\mu \in \text{sv}(\tilde{G}_1), \nu \in \text{sv}_{\text{ext}}(G_2)} |\mu - \nu| > 0,$$

then

$$\left\| \begin{bmatrix} \sin \Theta(\mathcal{R}(U_1), \mathcal{R}(\tilde{U}_1)) & 0 \\ 0 & \sin \Theta(\mathcal{R}(V_1), \mathcal{R}(\tilde{V}_1)) \end{bmatrix} \right\|_{\text{ui}} \leq c \frac{1}{\delta} \left\| \begin{bmatrix} R & 0 \\ 0 & S \end{bmatrix} \right\|_{\text{ui}},$$

where  $c = \pi/2$  in general, but  $c = 1$  if also  $\sigma_{\min}(\tilde{G}_1) > \sigma_{\max}(G_2)$  or for the Frobenius norm. Here  $\Theta(\mathcal{R}(U_1), \mathcal{R}(\tilde{U}_1))$  is the diagonal matrix of the canonical angles between the subspaces  $\mathcal{R}(U_1)$  and  $\mathcal{R}(\tilde{U}_1)$  [11, p.21-2], [16].

The case for  $\|\cdot\|_{\text{ui}} = \|\cdot\|_{\text{F}}$  is not new and Stewart and Sun [16, p.260] credited it to Wedin [17] with a slightly different  $\delta$  (similar to (4.9) we commented moments ago). However, Wedin [17] did not explicitly mention it for the case. Evidently, Wedin could easily had it because of the machinery he had already built in the paper.

*Proof of Theorem 4.1.* It can be seen that

$$U_2^H R = G_2 V_2^H \tilde{V}_1 - U_2^H \tilde{U}_1 \tilde{G}_1, \quad V_2^H S = G_2^H U_2^H \tilde{U}_1 - V_2^H \tilde{V}_1 \tilde{G}_1^H.$$

Or, equivalently,

$$(U_2^H \tilde{U}_1) \tilde{G}_1 - G_2 (V_2^H \tilde{V}_1) = -U_2^H R, \quad (V_2^H \tilde{V}_1) \tilde{G}_1^H - G_2^H (U_2^H \tilde{U}_1) = -V_2^H S,$$

which takes the form of the coupled Sylvester equations (2.6) with  $X = U_2^H \tilde{U}_1$  and  $Y = V_2^H \tilde{V}_1$ . Noticing that the singular values of  $U_2^H \tilde{U}_1$  and those of  $V_2^H \tilde{V}_1$  are the same as the sines of the canonical angles between  $\mathcal{R}(U_1)$  and  $\mathcal{R}(\tilde{U}_1)$  and between  $\mathcal{R}(V_1)$  and  $\mathcal{R}(\tilde{V}_1)$ , respectively, and hence [16, 8, 9]

$$\left\| \begin{bmatrix} U_2^H \tilde{U}_1 & 0 \\ 0 & V_2^H \tilde{V}_1 \end{bmatrix} \right\|_{\text{ui}} = \left\| \begin{bmatrix} \sin \Theta(\mathcal{R}(U_1), \mathcal{R}(\tilde{U}_1)) & 0 \\ 0 & \sin \Theta(\mathcal{R}(V_1), \mathcal{R}(\tilde{V}_1)) \end{bmatrix} \right\|_{\text{ui}},$$

and noticing

$$\left\| \begin{bmatrix} -U_2^H R & 0 \\ 0 & -V_2^H S \end{bmatrix} \right\|_{\text{ui}} = \left\| - \begin{bmatrix} U_2^H & 0 \\ 0 & V_2^H \end{bmatrix} \begin{bmatrix} R & 0 \\ 0 & S \end{bmatrix} \right\|_{\text{ui}} \leq \frac{1}{\delta} \left\| \begin{bmatrix} R & 0 \\ 0 & S \end{bmatrix} \right\|_{\text{ui}},$$

the conclusion in the theorem is a simple consequence of Lemma 2.3.  $\square$

## 5 Concluding Remarks

Stewart's original theorem [15, Theorem 6.4] for the singular subspaces associated with the SVD of a matrix subject to some perturbations uses both the Frobenius and spectral norms. Although it is still versatile to apply in the situations such as only the spectral norm is suitable [18], it may lead to weaker results: stronger conditions and yet less sharp bounds, through equivalency bounds between the spectral norm and the Frobenius norm,

as we argued in Section 1. Consequently it pays to establish variants of Stewart’s original theorem from scratch. Our main contribution in Theorem 3.1 provides perturbation bounds that encompass a variety of circumstances and that are dictated by the conditions as specified by Lemma 3.1, which are further explained in Remark 3.1.

Lemma 2.3 collects bounds, some old and some new, on the solution to the set of coupled Sylvester equations (2.6) under different circumstances. They form a part of the foundation based on which our main results in Theorem 3.1 are derived. Furthermore Lemma 2.3 can be put into good use to establish more generalized  $\sin \theta$  theorems for SVD as exemplified in Theorem 4.1, beyond existing ones due to Wedin [17].

## References

- [1] A. C. Antoulas. *Approximation of Large-Scale Dynamical Systems*. Advances in Design and Control. SIAM, Philadelphia, PA, 2005.
- [2] Peter Benner, Mario Ohlberger, Albert Cohen, and Karen Willcox, editors. *Model Reduction and Approximation: Theory and Algorithms*. Computational Science & Engineering. SIAM, Philadelphia, 2017.
- [3] R. Bhatia. *Matrix Analysis*. Graduate Texts in Mathematics, vol. 169. Springer, New York, 1996.
- [4] R. Bhatia, C. Davis, and A. McIntosh. Perturbation of spectral subspaces and solution of linear operator equations. *Linear Algebra Appl.*, 52-53:45–67, 1983.
- [5] R. Bhatia and P. Rosenthal. How and why to solve the operator equation  $AX - XB = Y$ . *Bull. London Math. Soc.*, 29:1–21, 1997.
- [6] C. Davis and W. Kahan. The rotation of eigenvectors by a perturbation. III. *SIAM J. Numer. Anal.*, 7:1–46, 1970.
- [7] C.-K. Li and R.-C. Li. A note on eigenvalues of perturbed Hermitian matrices. *Linear Algebra Appl.*, 395:183–190, 2005.
- [8] R.-C. Li. Bounds on perturbations of generalized singular values and of associated subspaces. *SIAM J. Matrix Anal. Appl.*, 14:195–234, 1993.
- [9] R.-C. Li. On perturbations of matrix pencils with real spectra. *Math. Comp.*, 62:231–265, 1994.
- [10] R.-C. Li. A bound on the solution to a structured Sylvester equation with an application to relative perturbation theory. *SIAM J. Matrix Anal. Appl.*, 21:440–445, 1999.
- [11] R.-C. Li. Matrix perturbation theory. In L. Hogben, R. Brualdi, and G. W. Stewart, editors, *Handbook of Linear Algebra*, page Chapter 21. CRC Press, Boca Raton, FL, 2nd edition, 2014.

- [12] X. Liang and R.-C. Li. Extensions of Wielandt’s min-max principles for positive semi-definite pencils. *Lin. Multilin. Alg.*, 62(8):1032–1048, 2014.
- [13] B. N. Parlett. *The Symmetric Eigenvalue Problem*. SIAM, Philadelphia, 1998.
- [14] G. W. Stewart. Error bounds for approximate invariant subspaces of closed linear operators. *SIAM J. Numer. Anal.*, 8:796–808, 1971.
- [15] G. W. Stewart. Error and perturbation bounds for subspaces associated with certain eigenvalue problems. *SIAM Rev.*, 15:727–764, 1973.
- [16] G. W. Stewart and J.-G. Sun. *Matrix Perturbation Theory*. Academic Press, Boston, 1990.
- [17] P.-Å. Wedin. Perturbation bounds in connection with singular value decomposition. *BIT*, 12:99–111, 1972.
- [18] L.-H. Zhang and R.-C. Li. Quality of approximate balanced truncation. URL <https://arxiv.org/pdf/2406.05665>, 2024.