Open Packing in Graphs: Bounds and Complexity

Shalu M. A. Kirubakaran V. K.

Indian Institute of Information Technology, Design and Manufacturing, Kancheepuram, Chennai, India.

Abstract

Given a graph G(V, E), a vertex subset S of G is called an open packing in G if no pair of distinct vertices in S have a common neighbour in G. The size of a largest open packing in G is called the open packing number of G and is denoted by $\rho^{o}(G)$. It would be interesting to note that the open packing number is a lower bound for the total domination number in graphs with no isolated vertices [Henning and Slater, 1999]. Given a graph G and a positive integer k, the decision problem OPEN PACKING tests whether G has an open packing of size at least k. The optimization problem MAX-OPEN PACKING takes a graph G as input and finds the open packing number of G.

It is known that OPEN PACKING is NP-complete on split graphs (i.e., the class of $\{2K_2, C_4, C_5\}$ -free graphs) [Ramos et al., 2014]. In this work, we complete the study on the complexity (P vs NPC) of OPEN PACKING on *H*-free graphs for every graph *H* with at least three vertices by proving that OPEN PACKING is (i) NP-complete on $K_{1,3}$ -free graphs and (ii) polynomial time solvable on $(P_4 \cup rK_1)$ -free graphs for every $r \ge 1$. In the course of proving (ii), we show that for every $t \in \{2, 3, 4\}$ and $r \ge 1$, if *G* is a $(P_t \cup rK_1)$ -free graph, then $\rho^{\circ}(G)$ is bounded above by a linear function of *r*. In addition, we find near-optimal upper bounds for the total domination number in the class of $(P_t \cup rK_1)$ -free graphs for every $t \in \{3, 4\}$ and $r \ge 1$. Moreover, we show that OPEN PACKING parameterized by solution size is W[1]-complete on $K_{1,3}$ -free graphs, and MAX-OPEN PACKING is hard to approximate within a factor of $(n^{\frac{1}{2}-\delta})$ for any $\delta > 0$ on $K_{1,3}$ -free graphs unless P = NP. Further, we prove that OPEN PACKING is (a) NP-complete on $K_{1,4}$ -free split graphs and (b) polynomial time solvable on $K_{1,3}$ -free split graphs. We prove a similar dichotomy result on split graphs with degree restrictions on the vertices in the independent set of the clique-independent set partition of the split graphs.

 $Keywords\colon$ Total dominating set, Open packing, $H\mathchar`-free graphs, NP\mathchar`-complete, W[1]\mathchar`-complete, W[1]\mathchar`-complete$

1 Introduction

In a graph G(V, E), a vertex subset D of G is called a *total dominating set* in G if every vertex in V(G) is adjacent to some vertex in D. In other words, $V(G) = \bigcup_{u \in D} N_G(u)$, where $N_G(u)$ denotes the set of vertices in V(G) that are adjacent to u in G. This implies that if D is a total dominating set in G, then $|D \cap N_G(u)| \ge 1$ for every vertex $u \in V(G)$. Note that by the definition of a total dominating set, it is evident that a graph G admits a total dominating set if and only if G has no isolated vertices. The cardinality of a smallest total dominating set in G is called the *total domination number*, $\gamma_t(G)$, of G. The total domination problem is useful in facility location problems, monitoring computer networks, and electronic communications [7]. In this article, we study the total dominating set problem and its dual problem, called open packing in graphs [8]. A vertex subset S of a graph G is called an *open packing* in G if, for every pair of distinct vertices $x, y \in S$, $N_G(x) \cap N_G(y)$ is empty. The *open packing number* of G, denoted by $\rho^o(G)$, is the size of a largest open packing in G. Note that if S is an open packing in G, then $|S \cap N_G(u)| \le 1$ for every $u \in V(G)$. This validates the primal-dual relation between the total dominating set and the open packing, and thus $\rho^o(G) \le \gamma_t(G)$. The decision and the optimisation versions of the problems of total dominating set and open packing are as follows.

Total	Dominating	Set
-------	------------	-----

Instance : A graph G and a positive integer $k \le |V(G)|$. **Question:** Does G has a total dominating set of size at most k?

		1
Instance :	A graph G and a positive integer $k \leq V(G) $.	ĺ
Question:	Is there an open packing of size at least \boldsymbol{k}	
	in G ?	

MIN-TOTAL DOMINATING SET **Instance:** A graph G. **Goal** : Find $\gamma_t(G)$.

Max-Open Packing	
Instance: A graph G . Goal : Find $\rho^o(G)$.	

Total dominating set is one of the well-studied problems in the literature and an extensive list of results can be found in [5, 7, 10]. In the context of our research, it would be interesting to note that TOTAL DOMINATING SET is NP-complete on $K_{1,5}$ -free split graphs [24] and polynomial time solvable on $K_{1,4}$ -free split graphs [20]. Also, TOTAL DOMINATING SET is NP-complete on $K_{1,3}$ free graphs [14] and an optimal total dominating set of a chordal bipartite graph can be found in polynomial time [3]. The study on open packing of graphs was initiated by Henning and Slater [9], and Rall [18] proved that for every non-trivial tree T, $\gamma_t(G) = \rho^o(G)$. In a recent work [21] (yet to be published), we extended this result by proving that the total domination number and the open packing number are equal when the underlying graph is a chordal bipartite graph with no isolated vertices. It is also known that OPEN PACKING is NP-complete (NPC in short) on split graphs (equivalently, the class of $\{2K_2, C_4, C_5\}$ -free graphs) [19] and planar bipartite graphs of maximum degree three (a subclass of K_3 -free graphs) [21]. In this work, we complete the study of OPEN PACKING on *H*-free graphs for every graph *H* with at least three vertices by proving the following theorems.

Theorem 1. Let H be a graph on three vertices. Then, an optimal open packing in H-free graphs can be found in polynomial time if and only if $H \ncong K_3$ unless P = NP.

Theorem 2. For $p \ge 4$, let H be a graph on p vertices. Then, OPEN PACKING is polynomial time solvable on the class of H-free graphs if and only if $H \in \{pK_1, (K_2 \cup (p-2)K_1), (P_3 \cup (p-3)K_1), (P_4 \cup (p-4)K_1)\}$ unless P = NP.

To prove the above theorem, we proved the following results.

- (i) OPEN PACKING is NP-complete on $K_{1,3}$ -free graphs.
- (ii) For $t \in \{2, 3, 4\}$ and $r \ge 1$, if G is a connected $(P_t \cup rK_1)$ -free graph, then
 - (a) $\rho^{o}(G) \leq 2r+1$ and $\gamma_{t}(G) \leq 2r+2$ provided t=4.
 - (b) $\rho^{o}(G) \leq 2r$ and $\gamma_{t}(G) \leq 2r+1$ provided t=3.
 - (c) $\rho^{o}(G) \leq \max\{r+1, 2(r-1)\}$ provided t = 2.

We also show that the bounds on $\rho^o(G)$ given in (ii) are tight. Moreover, we use the bounds on the total domination number in (a) and (b) to show that the results equivalent to those of Theorems 1 and 2 hold for TOTAL DOMINATING SET in *H*-free graphs as well (see Theorem 9). This implies that TOTAL DOMINATING SET and OPEN PACKING are of the same nature in the view of classical complexity (P vs NPC) in *H*-free graphs for every graph *H*. Table 1 provides a comparative study of complexity (P vs NPC) between TOTAL DOMINATING SET and OPEN PACKING in certain classes of graphs. The rest of the results in this article are motivated by Table 1 and the question, 'Does there exist a graph class where these two problems 'differ' from each other in the view of classical complexity (P vs NPC)?'

We answered this question in the affirmative (see Table 2) with the following list of dichotomy results in subclasses of split graphs.

- 1. OPEN PACKING is NP-complete on $K_{1,r}$ -free split graphs for $r \ge 4$, and is polynomial time solvable for $r \le 3$.
- 2. OPEN PACKING is NP-complete on I_r -split graphs (see Section 2 for definition) for $r \ge 3$ and is polynomial time solvable for $r \le 2$.
- 3. TOTAL DOMINATING SET is NP-complete in I_r -split graphs for $r \ge 2$ and is polynomial time solvable when r = 1 (a minor modification in the reduction by Corneil and Perl [1] will also prove that TOTAL DOMINATING SET is NP-complete for I_2 -split graphs).

Graph Classes	Total Dominating Set	Open Packing
Chordal bipartite graphs	P [11, 13]	P [21]
P_4 -free graphs	P [Folklore]	P [Folklore]
Bipartite graphs	NPC [16]	NPC $[21, 22]$
Split Graphs	NPC [1]	NPC [19]
$K_{1,3}$ -free Graphs	NPC [14]	NPC[*]

Table 1: Comparison between TOTAL DOMINATING SET and OPEN PACKING

Table 2: Complexity 'difference' between TOTAL DOMINATING SET and OPEN PACKING in subclasses of split graphs

Graph Classes	Total Dominating Set	Open Packing
I_2 -Split Graphs	NPC [1]	$P[*]^{1}$
$K_{1,4}$ -free Split Graphs	P [20]	NPC[*]

2 Preliminaries

We follow West [23] for terminology and notation. The graphs considered in this work are simple and undirected unless specified otherwise. Given a graph G(V, E), let n and m denote the number of vertices and the number of edges in G, respectively. Given a vertex $x \in V(G)$, the (open) neighbourhood of x in G is defined as $N_G(x) = \{y \in V(G) : xy \in E(G)\}$, and let the degree of a vertex x in G be $deg_G(x) = |N_G(x)|$. The closed neighbourhood of a vertex x in G is defined as $N_G[x] = \{x\} \cup N_G(x)$. We use N(x), N[x] and deg(x) instead of $N_G(x)$, $N_G[x]$ and $deg_G(v)$ respectively, when there is no ambiguity on G. A vertex x in G is called an isolated vertex in G, if $N_G(x) = \emptyset$. Given $U \subseteq V(G)$, the subgraph of G induced by U is denoted as G[U]. Given a graph H, G is said to be H-free if no induced subgraph of G is isomorphic to H. For a vertex $x \in V(G)$, let $E_G(x)$ denote the set of all edges incident on x, and for an edge $e \in E(G)$, let $V_G(e)$ denote the end vertices of e in G. Note that for $u \in V(G)$ and $e \in E(G)$, the edge $e \in E_G(u)$ if and only if $u \in V_G(e)$. For a graph G, the line graph L(G) of G is a graph with V(L(G)) = E(G) and two elements $e, e' \in V(L(G))$ are adjacent in L(G) if and only if $V_G(e) \cap V_G(e') \neq \emptyset$. Given two (not necessarily disjoint) graphs H and H', the graph union $H \cup H'$ is defined as $V(H \cup H') = V(H) \cup V(H')$ and $E(H \cup H') = E(H) \cup E(H')$. If V(H) and V(H')are not disjoint, then we call the graph union $H \cup H'$ as merger of the graphs H and H'. For $p \in \mathbb{N} \cup \{0\}$ and a graph H, the graph pH is defined as the union of p disjoint copies of H. Given two disjoint graphs G and H, we say that (i) a vertex x of the graph G is replaced by the graph H if a graph G' is constructed with $V(G') = (V(G) \setminus \{x\}) \cup V(H)$ and $E(G') = (E(G) \setminus \{xy:$ $y \in V(G)$ and $xy \in E(G)$ $\cup E(H) \cup \{wy : w \in V(H), y \in (V(G) \setminus \{x\}) \text{ and } xy \in E(G)\}$, and (ii) a vertex x of the graph G is *identified* with a vertex y of the graph H, if the vertex y in H is relabelled as x and the graphs G and H are merged.

A vertex subset C of G is called a *clique* in G if every pair of distinct vertices in C is adjacent in G. A vertex subset I of G is called an *independent set* in G if no pair of vertices in I are adjacent in G. The size of a largest independent set in G is called the *independence number* of G, and is denoted by $\alpha(G)$. Let P_n , C_n and K_n denote the path, cycle and complete graph on n vertices, respectively. A set $D \subseteq V(G)$ is called a *dominating set* in G, if every vertex in $V(G) \setminus D$ is adjacent to some vertex in D. An edge subset F of a graph G is called a *matching* in G, if $V_G(e) \cap V_G(e') = \emptyset$ for every pair of distinct $e, e' \in F$. The cardinality of a largest matching in G is called the matching number, $\alpha'(G)$, of G.

A graph G(V, E) is said to be a *split graph*, if there exists a partition $V(G) = C \cup I$ such that C is a clique and I is an independent set in G and is denoted as $G(C \cup I, E)$. Note that for $r \ge 2$, if a split graph G is $K_{1,r}$ -free, then $|N_G(v) \cap I| \le r - 1$ for every $v \in C$. In accordance with this observation, the class of I_r -split graphs is defined for every natural number r as a split graph $G(C \cup I, E)$ with $deg_G(v) = r$ for every $v \in I$.

Given a graph G and a positive integer $k \leq |V(G)|$, the problem INDEPENDENT SET asks whether G has an independent set of size at least k. Given a graph G, the goal of the problem MAX-

 $^{^1 [\}ast]$ denotes the results in this work

INDEPENDENT SET is to find $\alpha(G)$. For $r \in \mathbb{N}$, given a collection of sets X_1, X_2, \ldots, X_r each of cardinality q for some $q \in \mathbb{N}$ and a subset M of $\prod_{i=1}^r X_i$, the r-DIMENSIONAL MATCHING problem asks whether there exists $L \subseteq M$ such that (i) |L| = q and (ii) for every pair of r-tuple $x = (x_1, x_2, \ldots, x_r)$ and $y = (y_1, y_2, \ldots, y_r)$ in $L, x_i \neq y_i$ for $1 \leq i \leq r$. For $r \in \mathbb{N}$, given a non-empty set U, a set \mathcal{W} of r-sized subsets of U and a positive integer k, the r-HITTING SET problem asks whether there exists a subset X of U such that $|X| \leq k$ and $X \cap W \neq \emptyset$ for every $W \in \mathcal{W}$. Also, we refer the reader to [2, 5] for a brief note on parameterized algorithms, intractability and W-hierarchy.

3 H-free Graphs

We dedicate this section to prove the dichotomy result on OPEN PACKING stated in Theorem 2. Observation 1 helps us prove the necessary part of Theorem 2.

Observation 1. For $p \ge 4$, let H be a graph on p vertices such that $H \notin \{P_4 \cup (p-4)K_1, P_3 \cup (p-3)K_1, K_2 \cup (p-2)K_1, pK_1\}$. Then, H contains at least one of $K_3, 2K_2, C_4, K_{1,3}$, or C_5 as an induced subgraph.

Proof. On the contrary, assume that $H \notin \{P_4 \cup (p-4)K_1, P_3 \cup (p-3)K_1, K_2 \cup (p-2)K_1, pK_1\}$ and H is a $\{K_3, 2K_2, C_4, K_{1,3}, C_5\}$ -free graph. Then, the following statements hold.

- (1) *H* has exactly one non-trivial component since *H* is $2K_2$ -free and $H \cong pK_1$.
- (2) For every vertex $v \in V(H)$, $deg_H(v) \leq 2$ since H is $\{K_3, K_{1,3}\}$ -free.
- (3) H is acyclic since H is $\{K_3, 2K_2, C_4, C_5\}$ -free.

Note that by (1), (2), and (3), it is clear that the non-trivial component H' of H is a path and since H is $2K_2$ -free, $H' \in \{K_2, P_3, P_4\}$. This implies that $H \in \{P_4 \cup (p-4)K_1, P_3 \cup (p-3)K_1, K_2 \cup (p-2)K_1\}$, a contradiction.

Remark 1.

OPEN PACKING is NP-complete on (i) $2K_2$ -free graphs, (ii) C_4 -free graphs, (iii) C_5 -free graphs, and (iv) triangle-free graphs because of the facts that OPEN PACKING is NP-complete on (a) split graphs (i.e., the class of $\{2K_2, C_4, C_5\}$ -free graphs) [19] and (b) bipartite graphs (a subclass of K_3 -free graphs) [21, 22] and (c) OPEN PACKING is in NP in every subclass of simple graphs.

Next, we prove the following results on $K_{1,3}$ -free graphs: (i) OPEN PACKING is NP-complete, (ii) MAX-OPEN PACKING is hard to approximate within a factor of $(N^{(\frac{1}{2}-\delta)})$ for any $\delta > 0$ unless P = NP (where N denotes the number of vertices in a $K_{1,3}$ -free graph), and (iii) OPEN PACKING parameterized by solution size is W[1]-complete.

3.1 $K_{1,3}$ -free Graphs

The following construction gives a polynomial time reduction from INDEPENDENT SET on simple graphs to OPEN PACKING on $K_{1,3}$ -free graphs, where the former problem is known to be NP-complete [12].

Construction 1.

Input: A simple graph G with $V(G) = \{u_1, u_2, \dots, u_n\}.$

Output: A $K_{1,3}$ -free graph G'.

Guarantee: G has an independent set of size k if and only if G' has an open packing of size k. Procedure:

- Step 1: Replace each edge e = uu' in G with a three vertex path ueu' in G'.
- Step 2: For every $u \in V(G)$, make $E_G(u)$ a clique in G'.
- Step 3: For every vertex $u_i \in V(G)$ with exactly one edge, say e incident on it in G, introduce a vertex v_i , and two edges $u_i v_i$ and $v_i e$ in G'.

An example of Construction 1 is given in Fig. 1. Also, note that Step 2 of Construction 1 can be viewed as the merger of the line graph of G and the graph obtained in Step 1. Further, $V(G') = V(G) \cup E(G) \cup A$, where $A = \{v_i : 1 \le i \le n \text{ and } deg_G(u_i) = 1\}$ and E(G') =




Figure 1: (a) a simple graph G, (b) $K_{1,3}$ -free graph G' obtained from G using Construction 1.

 $E_1 \cup E_2 \cup E_3 \cup E_4$, where $E_1 = \{ue : u \in V(G), e \in E(G) \text{ and } e \in E_G(u)\}$, $E_2 = E(L(G))$, $E_3 = \{v_i u_i : 1 \le i \le n \text{ and } deg_G(u_i) = 1\}$ and $E_4 = \{v_i e : 1 \le i \le n \text{ and } E_G(u_i) = \{e\}\}$. Hence, $|V(G')| \le 2n + m$ and $|E(G')| = \sum_{i=1}^4 |E_i| \le 2m + \binom{m}{2} + 2n$. Thus, the graph G' can be constructed in quadratic time in the input size.

We can observe that in the graph G', for every vertex $u_i \in V(G)$,

$$N_{G'}(u_i) = \begin{cases} E_G(u_i) \cup \{v_i\} & \text{if } |E_G(u_i)| = \deg_G(u_i) = 1\\ E_G(u_i) & \text{otherwise} \end{cases}$$
(1)

$$N_{G'}(v_i) = \{u_i, e\} \text{ if } E_G(u_i) = \{e\} \text{ i.e., } deg_G(u_i) = 1$$
(2)

The above equations imply that, by Steps 2 and 3 of Construction 1, $N_{G'}(u_i)$ and $N_{G'}(v_i)$ are cliques in G'. Also, by Construction 1, for every $e \in E(G)$ with $V_G(e) = \{u, u'\}$,

$$N_{G'}[e] = N_{G'}[u] \cup N_{G'}[u'] \text{ and so } N_{G'}(e) = (N_{G'}[u] \cup N_{G'}[u']) \setminus \{e\}$$
(3)

This implies that $N_{G'}(e)$ is a union of two cliques in G'. We know that a graph H is $K_{1,3}$ -free if and only if $N_H(x)$ is a union of at most two cliques for all $x \in V(H)$. So, Equations (1), (2), and (3) imply that G' is $K_{1,3}$ -free.

Next, we prove Claims 1-5, which are helpful in completing the proof of guarantee of Construction 1.

Claim 1: A set $S \subseteq V(G)$ is an independent set in the input graph G of Construction 1 if and only if S is an open packing in the output graph G'. Note that by Equation (1), for $i \neq j$, $e = u_i u_j$ in G if and only if $N_{G'}(u_j) \cap N_{G'}(u_j) = \{e\}$, and

Note that by Equation (1), for $i \neq j$, $e = u_i u_j$ in G if and only if $N_{G'}(u_i) \cap N_{G'}(u_j) = \{e\}$, and hence $u_i u_j$ is not an edge in G if and only if $N_{G'}(u_i) \cap N_{G'}(u_j) = \emptyset$. This completes the proof of Claim 1.

Observation 2. Let H be a graph, and let $U = \{x, y, z\}$ be a vertex subset of H that induces a triangle in H. Then, for any open packing S in H, $|S \cap U| \leq 1$.

The above observation is held by the fact that any two vertices in a triangle have a common neighbour (third vertex).

Claim 2: Let $u_i \in V(G)$ such that $deg_G(u_i) = 1$, and let S be an open packing in G' with $v_i \in S$. Then, (i) $u_i \notin S$, (ii) $S' = (S \setminus \{v_i\}) \cup \{u_i\}$ is an open packing in G', and (iii) |S| = |S'|.

Since $deg_G(u_i) = 1$, there exists an edge $e \in E(G)$ such that $E_G(u_i) = \{e\}$. Then, $\{u_i, v_i, e\}$ induces a triangle in G' with $N_{G'}(u_i) = \{e, v_i\}$ and $N_{G'}(v_i) = \{e, u_i\}$. Therefore, by Observation 2, for any open packing S_1 in G', $|S_1 \cap \{u_i, v_i, e\}| \leq 1$. Thus, $v_i \in S$ implies that $u_i, e \notin S$. This proves (i) of Claim 2. Next, we prove (ii). Since $N_{G'}(u_i) = \{e, v_i\}$ and $N_{G'}(v_i) = \{e, u_i\}$, for any vertex $x \in S \setminus \{v_i\}$, x and v_i have no common neighbour in G', which implies that x and u_i have no common neighbour in G'. This, together with the fact that $S \setminus \{v_i\}$ is an open packing in G', implies that $S' = (S \setminus \{v_i\}) \cup \{u_i\}$ is an open packing in G'. Note that (iii) follows from (i).




Figure 2: (a) an edge $e = u_{\ell}u_j$ of the input graph G of Construction 1 and (b) the three vertex path $u_{\ell}eu_j$ produced by Construction 1 along with two vertices y, z of G' such that $y \in N_{G'}(z) \cap N_{G'}(u_{\ell})$ as in the contrary assumption of Claim 4. In figure, dashed lines represent the edges not in G' as observed in the proof of Claim 4.

Claim 3: Let e be an edge incident on a vertex u_i in G. Then, for any open packing S in G', $|S \cap \{e, u_i\}| \leq 1$.

If $deg_G(u_i) = 1$, then $\{u_i, v_i, e\}$ induces a triangle in G' and so $|S \cap \{e, u_i\}| \le 1$ by Observation 2. Also, if $deg_G(u_i) \ge 2$, then $\{u_i, e, f\}$ induces a triangle in G' for some $f \in (E_G(u_i) \setminus \{e\})$. Again by Observation 2, $|S \cap \{u_i, e\}| \le 1$.

Claim 4: Let S be an open packing in G' with $e = u_{\ell}u_j \in S \cap E(G)$ for some $1 \leq \ell < j \leq n$. Then, (i) $u_{\ell}, u_j \notin S$, (ii) $S_{\ell} = (S \setminus \{e\}) \cup \{u_{\ell}\}$ and $S_j = (S \setminus \{e\}) \cup \{u_j\}$ are open packings in G', and (iii) $|S| = |S_{\ell}| = |S_j|$.

Since $e \in S$, $u_{\ell}, u_j \notin S$ because of Claim 3. This proves (i). Also, if $v_{\ell} \in V(G')$, then $v_{\ell} \notin S$ because of Observation 2 and the fact that $\{v_{\ell}, e, u_{\ell}\}$ induces a triangle in G'. Similarly, if $v_j \in V(G')$, then $v_j \notin S$. Next, we prove that $S_{\ell} = (S \setminus \{e\}) \cup \{u_{\ell}\}$ is an open packing in G'. Note that $S \setminus \{e\}$ is an open packing in G'. So, if S_{ℓ} is not an open packing in G, then there exists a vertex $z \in S \setminus \{e\}$ such that $N_{G'}(z) \cap N_{G'}(u_{\ell}) \neq \emptyset$. Let $y \in N_{G'}(z) \cap N_{G'}(u_{\ell})$. Note that $zu_{\ell} \notin E(G')$, else $u_{\ell} \in N_{G'}(z) \cap N_{G'}(e)$, which would be a contradiction to $e, z \in S$ (an open packing in G'). Similarly, $zu_{j} \notin E(G')$. If y = e, then since $e = y \in N_{G'}(z)$, we have $z \in (N_{G'}(e) \setminus (N_{G'}[u_{j}] \cup N_{G'}[u_{\ell}]))$, a contradiction to Equation (3) (see Fig. 2). Therefore, $y \neq e$. This implies that $ye \in E(G')$, since $y, e \in N_{G'}(u_{\ell})$, which is a clique in G'. Then, $y \in N_{G'}(z) \cap N_{G'}(e)$, a contradiction to S being an open packing in G'. Therefore, $S_{\ell} = (S \setminus \{e\}) \cup \{u_{\ell}\}$ is an open packing in G'. A similar argument holds for S_j . Also, note that (iii) follows from (i).

Claim 5: Let S be an open packing of size k in G'. Then, there exists an independent set S^* of size k in G.

Note that $V(G') = V(G) \cup E(G) \cup A$. We construct S^* from S in two phases. First, we construct an open packing S_1 in G' such that $S_1 \subseteq V(G) \cup E(G)$ and $|S_1| = |S|$. If $S \cap A = \emptyset$, then $S_1 := S$. Else $S \cap A \neq \emptyset$ and let $S \cap A = \{v_{i_1}, v_{i_2}, \ldots, v_{i_p}\}$. Then, $S_1 := (S \setminus \{v_{i_1}, v_{i_2}, \ldots, v_{i_p}\}) \cup \{u_{i_1}, u_{i_2}, \ldots, u_{i_p}\}$. By repeated application of Claim 2, S_1 is an open packing in G' such that $|S_1| = |S|$ and $S_1 \subseteq V(G) \cup E(G)$. Next, we construct an open packing S^* of G' from S_1 such that $|S^*| = |S_1|$ and $S^* \subseteq V(G)$. If $S_1 \cap E(G) = \emptyset$, then $S^* := S_1$. If not, let $S_1 \cap E(G) = \{e_1, e_2, \ldots, e_q\}$, and let u_{ℓ_i} be an end vertex of e_i in G for $1 \leq i \leq q$. Note that for $1 \leq i < j \leq q$, $u_{\ell_i} \neq u_{\ell_j}$; otherwise, $\{u_{\ell_i}, e_i, e_j\}$ induces a triangle (by Equation (1)) in G' with $e_i, e_j \in S_1$ (an open packing in G'), a contradiction to Observation 2. Thus, $S^* := (S_1 \setminus \{e_1, e_2, \ldots, e_q\}) \cup \{u_{\ell_1}, u_{\ell_2}, \ldots, u_{\ell_q}\}$. By repeated application of Claim 4, S^* is an open packing in G' with $|S^*| = |S_1| = |S| = k$ and $S^* \subseteq V(G)$. By Claim 1, S^* is an independent set in G. This proves Claim 5.

Claims 1 and 5 prove the guarantee of Construction 1. Thus, we have the following theorem which is an implication of Construction 1 and the fact that INDEPENDENT SET is NP-complete on simple graphs [12].

Theorem 3. OPEN PACKING is NP-complete on $K_{1,3}$ -free graphs.

We use the theorem below to derive an in-approximation result of MAX-OPEN PACKING for $K_{1,3}$ -free graphs.

Theorem 4 ([6]). MAX-INDEPENDENT SET cannot be approximated within a factor of $n^{(1-\epsilon)}$ for any $\epsilon > 0$, in general graphs unless P = NP.

Theorem 5. MAX-OPEN PACKING is hard to approximate within a factor of $N^{\frac{1}{2}-\delta}$ for any $\delta > 0$ in $K_{1,3}$ -free graphs unless P = NP, where N denotes the number of vertices in a $K_{1,3}$ -free graph.

Proof. On the contrary, assume that MAX-OPEN PACKING admits a $N^{\left(\frac{1}{2}-\delta\right)}$ -factor approximation algorithm for some $\delta > 0$ in $K_{1,3}$ -free graphs. Then, an open packing S of the graph G' constructed in Construction 1 with $|S| > \frac{\rho^o(G)}{|V(G')|^{\left(\frac{1}{2}-\delta\right)}}$ can be found in polynomial time. Hence, an independent set U of the input graph G in Construction 1 with |U| = |S| can be found in polynomial time. Note that $N = |V(G')| \le 2n + m \le 2n + \frac{n^2}{2} \le n^2$ for $n \ge 4$. Then,

$$|S| > \frac{\rho^o(G')}{N^{\left(\frac{1}{2} - \delta\right)}}$$
$$|S| > \frac{\rho^o(G')}{(n^2)^{\left(\frac{1}{2} - \delta\right)}} \quad \text{for } n \ge 4$$
$$|S| > \frac{\alpha(G)}{n^{1-2\delta}}$$
$$|U| > \frac{\alpha(G)}{n^{1-2\delta}}$$

Since $\rho^o(G') = \alpha(G)$ by the guarantee of Construction 1. Let $\epsilon = 2\delta$. Then, $\epsilon > 0$ and for $n \ge 4$, we have

$$|U| > \frac{\alpha(G)}{n^{1-\epsilon}}$$

The above inequality implies that MAX-INDEPENDENT SET has a $(n^{1-\epsilon})$ -factor approximation algorithm for some $\epsilon > 0$, which contradicts Theorem 4.

Next, we give a parameterized intractability result for OPEN PACKING on $K_{1,3}$ -free split graphs using Construction 1 and the following result from the literature.

Lemma 1 ([18]). Given a graph G, let the neighbourhood graph $G^{[o]}$ of G be a simple graph with $V(G^{[o]}) = V(G)$ and $E(G^{[o]}) = \{xy : x, y \in V(G), x \neq y \text{ and } N_G(x) \cap N_G(y) \neq \emptyset\}$. Then, a vertex subset S is an open packing in G if and only if S is an independent set in $G^{[o]}$.

Theorem 6. OPEN PACKING parameterized by solution size is W[1]-complete on $K_{1,3}$ -free graphs.

Proof. Note that INDEPENDENT SET parameterized by solution size is W[1]-complete [4]. Lemma 1 implies that every instance (G, k) of OPEN PACKING on $K_{1,3}$ -free graphs can be (FPT) reduced into an instance $(G^{[o]}, k)$ of INDEPENDENT SET. Hence, OPEN PACKING parameterized by solution size on $K_{1,3}$ -free graphs is in W[1]. The guarantee of Construction 1 implies that every instance (G, k) of INDEPENDENT SET can be (FPT) reduced into an instance (G', k) of OPEN PACKING on $K_{1,3}$ -free graphs. Thus, OPEN PACKING parameterized by solution size is W[1]-hard on $K_{1,3}$ -free graphs.

3.2 Subclasses of $(P_4 \cup rK_1)$ -free Graphs

In this section, we prove bounds on the total domination number and the open packing number in the classes of (i) $(P_4 \cup rK_1)$ -free graphs, (ii) $(P_3 \cup rK_1)$ -free graphs, (iii) $(K_2 \cup rK_1)$ -free graphs, and (iv) rK_1 -free graphs. These bounds eventually solve TOTAL DOMINATING SET and OPEN PACKING in polynomial time in the graph classes mentioned above. The following lemmas are the tools used to connect the bounds with polynomial time algorithms.

Lemma 2 (Folklore). Given a graph G and a vertex subset D of G, it can be tested whether D is a total dominating set of G or not in O(n + m) time.

Lemma 3. Given a graph class \mathcal{G} , if there exists $\ell \in \mathbb{N}$ such that $\gamma_t(G) \leq \ell$ for every $G \in \mathcal{G}$, then an optimal total dominating set of G can be found in $O(n^{\ell}(n+m))$ time for every $G \in \mathcal{G}$.

Proof. Let $G \in \mathcal{G}$, and let \mathcal{D}_{ℓ} be the set of all total dominating sets of size at most ℓ in G. Also, let D be an optimal total dominating set in G. Then, since $|D| = \gamma_t(G) \leq \ell$, $D \in \mathcal{D}_{\ell}$. Also, $\mathcal{D}_{\ell} \subseteq \mathcal{V}_{\ell} = \{X \subseteq V(G) : |X| \leq \ell\}$. Note that the collection \mathcal{V}_{ℓ} can be found in $O(n^{\ell})$ time. Also, for every $X \in \mathcal{V}_{\ell}$, it can be tested whether $X \in \mathcal{D}_{\ell}$ in O(n+m) time using Lemma 2. This, together with the fact that $|\mathcal{V}_{\ell}| = O(n^{\ell})$, implies that the collection \mathcal{D}_{ℓ} can be found in $O(n^{\ell}(n+m))$ time. Since every $D' \in \mathcal{D}_{\ell}$ is a total dominating set, $\gamma_t(G) \leq \min\{|D'| : D' \in \mathcal{D}_{\ell}\}$. Further, since $D \in \mathcal{D}_{\ell}, \gamma_t(G) = |D| \geq \min\{|D'| : D' \in \mathcal{D}_{\ell}\}$. So, $\gamma_t(G) = \min\{|D'| : D' \in \mathcal{D}_{\ell}\}$. Since \mathcal{D}_{ℓ} can be generated in $O(n^{\ell}(n+m))$ time, so is $\gamma_t(G)$ and an optimal total dominating set of G.

The following lemma is helpful in proving a result similar to the above lemma for open packing.

Lemma 4 ([21]). Given a graph G and a vertex subset S of G, it can be tested in O(n) time whether S is an open packing in G or not.

Proof of Lemma 4 is given in Appendix A.

Lemma 5. Given a graph class \mathcal{G} , if there exists $k \in \mathbb{N}$ such that $\rho^{o}(G) \leq k$ for every $G \in \mathcal{G}$, then (i) G contains at most $O(n^{k})$ open packings, and (ii) all open packings in G can be computed in $O(n^{k+1})$ time for every $G \in \mathcal{G}$. So, $\rho^{o}(G)$ can be computed in $O(n^{k+1})$ time.

Proof. Let $G \in \mathcal{G}$. Let \mathcal{S} be the set of all open packings in G. We prove that the set \mathcal{S} can be found in $O(n^{k+1})$ time. Let $\mathcal{V}_k = \{X \subseteq V(G) : |X| \leq k\}$. Then, since $\rho^o(G) \leq k, \mathcal{S} \subseteq \mathcal{V}_k$. Note that the set \mathcal{V}_k is of size $O(n^k)$ and can be found in $O(n^k)$ time. Since $\mathcal{S} \subseteq \mathcal{V}_k$, $|\mathcal{S}| = O(n^k)$. Also, for every vertex subset $X \in \mathcal{V}_k$, it can be tested in O(n) time whether $X \in \mathcal{S}$ using Lemma 4. Thus, the set \mathcal{S} can be found in $O(n^{k+1})$ time. Note that $\rho^o(G) = \max\{|S| : S \in \mathcal{S}\}$. Since the set \mathcal{S} can be found in $O(n^{k+1})$ time, $\rho^o(G)$ can be computed in $O(n^{k+1})$ time.

Next, we study the bounds on the total domination number and the open packing number in the class of $(P_t \cup rK_1)$ -free graphs for $t \in \{1, 2, 3, 4\}$ and every non-negative integer $r \ge 3 - t$.

$(P_4 \cup rK_1)$ -free Graphs

In this section, we prove that for $r \geq 1$, $\gamma_t(G) \leq 2r + 2$, and $\rho^o(G) \leq 2r + 1$ in a connected $(P_4 \cup rK_1)$ -free graph G (the bound on the open packing number of $(P_4 \cup rK_1)$ -free graphs along with Lemma 5 will solve the sufficiency part of Theorem 2). Further, we show that the bound on the open packing number in these graph classes is tight (see Remark 3) and that the bound on the total domination number is tight for the class of $(P_4 \cup K_1)$ -free graphs (see Remark 2). The following known lemma solves the problems for P_4 -free graphs.

Lemma 6 (Folklore). Let G be a connected P_4 -free graph. Then, $\rho^o(G) \leq \gamma_t(G) = 2$.

The lemma below shows that for $r \ge 1$, the total domination number of a connected $(P_4 \cup rK_1)$ -free graph is bounded above by a function of r.

Lemma 7. For $r \ge 1$, if G is a connected $(P_4 \cup rK_1)$ -free graph, then $\gamma_t(G) \le 2r+2$.

Proof. We prove this lemma by induction on $r \geq 0$. Note that Lemma 6 shows that the bound holds for the case when r = 0. So, assume that for $r \geq 1$, $\gamma_t(G') \leq 2(r-1)+2$ if G' is a connected $(P_4 \cup (r-1)K_1)$ -free graph. Next, we show that for every connected $(P_4 \cup rK_1)$ -free graph G, $\gamma_t(G) \leq 2r+2$. Let G be a connected $(P_4 \cup rK_1)$ -free graph. Suppose that G is $(P_4 \cup (r-1)K_1)$ free, then by induction assumption $\gamma_t(G) \leq 2(r-1)+2 < 2r+2$. So, assume that G contains a $P_4 \cup (r-1)K_1$ as an induced subgraph. Let $D = \{x_1, x_2, x_3, x_4, y_1, y_2, \ldots, y_{r-1}\}$ be a vertex subset of G such that $x_i x_{i+1} \in E(G)$ for i = 1, 2, 3 and $G[D] \cong P_4 \cup (r-1)K_1$ (see Fig. 3). Since G is $(P_4 \cup rK_1)$ -free, every vertex in $V(G) \setminus D$ is adjacent to some vertex in D. Also, since G is connected, for every $j = 1, 2, \ldots, (r-1)$, there exists a vertex $w_j \in V(G)$ such that $y_j w_j \in E(G)$. Also, the vertices x_1, x_2, x_3 and x_4 are adjacent to a vertex in $\{x_2, x_3\} \subseteq D$. Thus, every vertex in $V(G) = D \cup (V(G) \setminus D)$ is adjacent to some vertex in $\{w_j : 1 \leq j \leq r-1\} \cup D$. So, $\{w_j : 1 \leq j \leq r-1\} \cup D$ is a total dominating set in G, and hence $\gamma_t(G) \leq r-1+|D| \leq 2(r-1)+4 = 2r+2$. \Box




Figure 3: A connected $(P_4 \cup rK_1)$ -free graph G with $D = \{x_1, x_2, x_3, x_4, y_1, y_2, \dots, y_{r-1}\} \subseteq V(G)$ such that $G[D] \cong (P_4 \cup (r-1)K_1)$. It is possible that $w_i = w_j$ for $i \neq j$.

Remark 2.

For r = 1, the bound given in Lemma 7 is tight, and a cycle on six vertices (C_6) is a connected $(P_4 \cup K_1)$ -free graph that satisfies this bound. Also, the graphs obtained by replacing every vertex of C_6 with a complete graph will remain as a $(P_4 \cup K_1)$ -free graph with total domination number four. For every r > 1, there exists a $(P_4 \cup rK_1)$ -free graph G_r with $\gamma_t(G_r) = 2r + 1$ (see Remark 3 and Fig. 4). Therefore, the bound given in Lemma 7 is a near-optimal upper bound.

The following corollary follows from Lemmas 6 and 7, and the fact that for every $r \ge 1$, every $(K_2 \cup rK_1)$ -free graph is a $(P_4 \cup (r-1)K_1)$ -free graph.

Corollary 1. For every $r \ge 1$, the total domination number of a connected $(K_2 \cup rK_1)$ -free graph is bounded above by 2r.

The following observation that follows from the definition of open packing is helpful in proving Lemma 8.

Observation 3 ([17]). Let S be an open packing in a graph G. Then, G[S] is $\{P_3, K_3\}$ -free, i.e., G[S] is a union of K_2 's and K_1 's.

The following lemma proves the bound on the open packing number in the class of $(P_4 \cup rK_1)$ -free graphs.

Lemma 8. For $r \ge 1$, if G is a connected $(P_4 \cup rK_1)$ -free graph, then $\rho^o(G) \le 2r + 1$.

Proof. We use induction on $r \ge 1$ to prove this lemma as well. Firstly, we prove the bound for the case r = 1. Let G be a connected $(P_4 \cup K_1)$ -free graph. Then, $\rho^o(G) \le 2 < 2r + 1 = 3$ if G is P_4 -free by Lemma 6. So, assume that G contains a P_4 . Let $D = \{x_1, x_2, x_3, x_4\}$ be a vertex subset of G that induces a P_4 in G with $x_i x_{i+1} \in E(G)$ for i = 1, 2, 3. Then, since G is $(P_4 \cup K_1)$ free, every vertex in $V(G) \setminus D$ is adjacent to some vertex in $\{x_1, x_2, x_3, x_4\}$. Also, every vertex

in $\{x_1, x_2, x_3, x_4\}$ is adjacent to a vertex in $\{x_2, x_3\}$. So, $V(G) = \bigcup_{i=1}^4 N(x_i)$. Thus, for any open

packing S in G, $S = S \cap V(G) = \bigcup_{i=1}^{4} (S \cap N(x_i))$. Since S is an open packing, $|S \cap N(x_i)| \le 1$ for

i = 1, 2, 3, 4. Thus, $|S| \leq \sum_{i=1}^{4} |S \cap N(x_i)| \leq 4$. So, to prove the claim, it is enough to disprove the

possibility of an open packing of size four in G. On the contrary, assume that there exists an open packing S of size four in G. Then, by the above arguments, we can conclude that $|S \cap N(x_i)| = 1$ for every i = 1, 2, 3, 4. Let $S \cap N(x_i) = \{z_i\}$. So, $S = \{z_1, z_2, z_3, z_4\}$. This, together with the assumption that |S| = 4, implies $z_i \neq z_j$ (and so $x_i z_j \notin E(G)$) for $i, j \in \{1, 2, 3, 4\}$ and $i \neq j$. Note that $z_1 x_1 \in E(G)$ (i.e., $z_1 \neq x_1$) whereas $x_1 x_3, x_1 x_4 \notin E(G)$, so $z_1 \notin \{x_1, x_3, x_4\}$. Further, $z_1 \neq x_2$ since $x_2 x_3 \in E(G)$ and $z_1 x_3 \notin E(G)$. Similarly, it can be proved that $z_4 \notin \{x_1, x_2, x_3, x_4\}$. Therefore, $z_1 z_4 \in E(G)$, else $\{z_1, x_1, x_2, x_3\} \cup \{z_4\}$ induces a $(P_4 \cup K_1)$ in G, a contradiction. Since $z_1 x_4 \in E(G)$, we have $z_1 z_2, z_2 z_4 \notin E(G)$ because of Observation 3. This, together with the fact that $z_1 x_1, z_4 x_4 \in E(G)$, implies that $z_2 \notin \{x_1, x_4\}$. Further, since $x_1 z_2, z_2 x_4 \notin E(G)$, the vertex subset $\{x_1, z_1, z_4, x_4\} \cup \{z_2\}$ induces a $(P_4 \cup K_1)$ in G, a contradiction. So $\rho^o(G) \leq 3$. Next, we prove the case $r \geq 2$.

Induction Assumption: Assume that for every connected $(P_4 \cup (r-1)K_1)$ -free graph G', $\rho^o(G') \le 2(r-1)+1$.

Let G be a $(P_4 \cup rK_1)$ -free graph. If G does not contain any vertex subset that induces a $(P_4 \cup (r-1)K_1)$, then G is $(P_4 \cup (r-1)K_1)$ -free, and so by induction assumption, $\rho^o(G) \leq 2(r-1) + 1 < 2r + 1$. So, assume that G has a vertex subset $D = \{x_1, x_2, x_3, x_4, y_1, y_2, \dots, y_{r-1}\}$ that induces a $(P_4 \cup (r-1)K_1)$ in G with $x_i x_{i+1} \in E(G)$ for i = 1, 2, 3. Since G is $(P_4 \cup rK_1)$ -free, every vertex in $V(G) \setminus D$ is adjacent to some vertex in D, i.e., D is a dominating set in G. So, $V(G) = N[x_1] \cup N[x_2] \cup N[x_3] \cup N[x_4] \cup N[y_1] \cup N[y_2] \cup \ldots \cup N[y_{r-1}]$. Hence, for any open packing S in $G, S = \bigcup_{v \in D} (S \cap N[u])$. Note that $|S \cap N[u]| \leq |S \cap N(u)| + |S \cap \{u\}| \leq 2$. Also, since

$$x_1, x_2, x_3, x_4 \in N(x_2) \cup N(x_3), \ \bigcup_{i=1}^4 N[x_i] = \bigcup_{i=1}^4 N(x_i). \ \text{So}, \ |S \cap (\bigcup_{i=1}^4 N[x_i])| = |S \cap (\bigcup_{i=1}^4 N(x_i))| = |S \cap (\bigcup_{i=1$$

$$|\bigcup_{i=1} (S \cap N(x_i))| \le 4. \text{ Therefore, } |S| = |S \cap V(G)| = |S \cap (N[x_1] \cup N[x_2] \cup N[x_3] \cup N[x_4] \cup N[y_1] \cup N[x_4] \cup N[y_1] \cup N[x_4] \cup N[y_1] \cup N[y_$$

 $N[y_2] \cup \ldots \cup N[y_{r-1}])| = |\bigcup_{i=1}^{n} (S \cap N(x_i))| + |\bigcup_{j=1}^{n} (S \cap N[y_j])| \le 4 + 2(r-1) = 2r+2$. Thus, to prove the claim, it is enough to rule out the possibility of |S| = 2r+2. On the contrary, assume

that there exists an open packing S of size 2r + 2 in G. Then, by the above arguments, we can conclude that $|S \cap N[y_j]| = 2$ for j = 1, 2, ..., r - 1 and $|S \cap (\bigcup_{i=1}^4 N(x_i))| = 4$. Since $|S \cap N(y_i)| \le 1$ and $|S \cap N[y_i]| = 2$, $y_i \in S$ and $S \cap N(y_i) \ne \emptyset$. Let $S \cap N[y_j] = \{w_j, y_j\}$ for j = 1, 2, ..., r - 1, and let $S \cap (\bigcup_{i=1}^4 N(x_i)) = \{z_1, z_2, z_3, z_4\}$ with $x_i z_i \in E(G)$ (then, since $|S \cap N(x_i)| \le 1$, we

have $S \cap N(x_i) = \{z_i\}$ for every i = 1, 2, 3, 4). So, $S = \left(\bigcup_{i=1}^4 \{z_i\}\right) \cup \left(\bigcup_{j=1}^{r-1} \{y_j\}\right) \cup \left(\bigcup_{j=1}^{r-1} \{w_j\}\right)$.

This, together with the fact that |S| = 2r + 2, imply that every vertex in the (2r + 2)-tuple $(z_1, z_2, z_3, z_4, y_1, y_2, \ldots, y_{r-1}, w_1, w_2, \ldots, w_{r-1})$ is distinct. This implies that for $s \in \{1, 2, 3, 4\}$, $z_s u \in E(G)$ for some $u \in D$ if and only if $u = x_s$ because for every $u \in D$, (i) $S \cap N(u) = \{z_i\}$ if $u = x_i$ for $i \in \{1, 2, 3, 4\}$ and (ii) $S \cap N(u) = \{w_j\}$ if $u = y_j$ for $j \in \{1, 2, \ldots, r-1\}$. We complete our claim in two cases based on whether $z_1 z_4 \in E(G)$ or not.

Case 1: $z_1 z_4 \in E(G)$.

Since G[S] is $\{P_3, K_3\}$ -free by Observation 3, $z_1z_2, z_2z_4 \notin E(G)$. So, $z_2 \neq x_1$ since $z_1x_1 \in E(G)$. Then, $\{x_1, z_1, z_4, x_4\} \cup \{z_2\} \cup \{y_1\} \cup \{y_2\} \cup \cdots \cup \{y_{r-1}\}$ induces a $(P_4 \cup rK_1)$ in G, a contradiction. Case 2: $z_1z_4 \notin E(G)$.

Then, $\{z_1, x_1, x_2, x_3\} \cup \{z_4\} \cup \{y_1\} \cup \{y_2\} \cup \cdots \cup \{y_{r-1}\}$ induces a $(P_4 \cup rK_1)$ in G, a contradiction. As we have arrived at a contradiction in both cases, $\rho^o(G) \leq 2r+1$.

Remark 3.

The bound given in Lemma 8 is tight. For example, let G_r be a graph obtained by subdividing exactly r edges of $K_{1,r+1}$ by four vertex paths. Graph G_3 is shown in Fig. 4. The vertex and edge set of G_r can be defined as $V(G_r) = (\bigcup_{i=1}^r \{x_i, y_i, z_i\}) \cup \{u, v\}$ and $E(G_r) = (\bigcup_{i=1}^r \{x_i y_i, y_i z_i, z_i u\}) \cup \{uv\}$. Then, G_r is $(P_4 \cup rK_1)$ -free (proof of this statement is given as Claim 11 in Appendix B), and $S_r = (\bigcup_{i=1}^r \{x_i, y_i\}) \cup \{v\}$ is an open packing in G of size 2r + 1. Also, note that the graph obtained by replacing the vertex u in G_r with a complete graph will remain a $(P_4 \cup rK_1)$ -free graph with open packing number 2r+1. Further, $\gamma_t(G_r) = \rho^o(G) = 2r+1$ with $D = \{u, z_1, z_2, \ldots, z_r, y_1, y_2, \ldots, y_r\}$ as a total dominating set in G_r .

$(P_3 \cup rK_1)$ -free Graphs

In this section, we prove that for $r \ge 1$, $\gamma_t(G) \le 2r + 1$, and $\rho^o(G) \le 2r$ for every connected $(P_3 \cup rK_1)$ -free graph G. Further, we show that the bound on (i) the open packing number in this graph class is tight and (ii) the total domination number is tight for the class of $(P_3 \cup K_1)$ -free graphs.

Remark 4.

Every connected P_3 -free graph is a complete graph, and hence $\rho^o(G) \leq \gamma_t(G) = 2$.

Lemma 9. For $r \ge 1$, if G is a connected $(P_3 \cup rK_1)$ -free graph, then $\gamma_t(G) \le 2r+1$.




Figure 4: A $(P_4 \cup 3K_1)$ -free graph G_3 defined in Remark 3 with an open packing $S_3 = \{x_1, x_2, x_3, y_1, y_2, y_3, v\}$ of size 7 = (2(3) + 1). Also, $\gamma_t(G_3) = 7$ with $D = \{u, z_1, y_1, z_2, y_2, z_3, y_3\}$ as a total dominating set in G_3 .

Proof. Firstly, we prove the case r = 1. Let G be a connected $(P_3 \cup K_1)$ -free graph. If G is P_3 -free, then $\gamma_t(G) = 2$. So, assume that G contains a P_3 . Let $D = \{x_1, x_2, x_3\}$ be a subset of V(G) that induces a P_3 in G with $x_1x_2, x_2x_3 \in E(G)$. Since G is $(P_3 \cup K_1)$ -free, every vertex in $V(G) \setminus D$ is adjacent to some vertex in D. Further, since the vertices in $\{x_1, x_2, x_3\}$ are adjacent to a vertex in $\{x_1, x_2\} \subseteq D$, every vertex in $V(G) = (V(G) \setminus D) \cup D$ is adjacent to some vertex in D. Hence, D is a total dominating set in G, and $\gamma_t(G) \leq |D| = 3 = 2r + 1$. Next, we prove the case $r \geq 2$ by induction on r.

Induction Assumption: For $r \ge 2$, if G' is a connected $(P_3 \cup (r-1)K_1)$ -free graph, then $\gamma_t(G') \le 2(r-1)+1$.

Let G be a $(P_3 \cup rK_1)$ -free graph. Suppose that G does not contain $(P_3 \cup (r-1)K_1)$ as an induced subgraph, then G is $(P_3 \cup (r-1)K_1)$ -free, and so by induction assumption, $\gamma_t(G) \leq 2(r-1)+1 < 2r+1$. So, assume that G contains $(P_3 \cup (r-1)K_1)$ as an induced subgraph. Let $D = \{x_1, x_2, x_3, y_1, y_2, \ldots, y_{r-1}\}$ be a subset of V(G) that induces $(P_3 \cup (r-1)K_1)$ in G with $x_1x_2, x_2x_3 \in E(G)$. Since G is $(P_3 \cup rK_1)$ -free, every vertex in $V(G) \setminus D$ is adjacent to a vertex in D. Also, since D is connected, for every vertex in $\{x_1, x_2, x_3\}$ is adjacent to a vertex in $\{x_1, x_2\} \subseteq D$. Thus, every vertex in $V(G) \setminus D$ is adjacent to some vertex in $D \cup \{w_j : 1 \leq j \leq r-1\}$. Hence, $D \cup \{w_j : 1 \leq j \leq r-1\}$ is a total dominating set in G. So, $\gamma_t(G) \leq |D \cup \{w_j : 1 \leq j \leq r-1\}| = 3 + (r-1) + (r-1) = 2r + 1$.

Next, we prove the bound on the open packing number in the class of $(P_3 \cup rK_1)$ -free graphs.

Lemma 10. For $r \ge 1$, if G is a connected $(P_3 \cup rK_1)$ -free graph, then $\rho^o(G) \le 2r$.

Proof of Lemma 10 is similar to the proof of Lemma 8. However, the complete proof is given in Appendix B. The corollary below follows from Lemmas 9 and 10, and the fact that for $r \ge 3$, every rK_1 -free graph is a $(P_3 \cup (r-2)K_1)$ -free graph as well.

Corollary 2. For $r \ge 3$, let G be a connected rK_1 -free graph. Then, (i) $\gamma_t(G) \le 2r - 3$ and (ii) $\rho^o(G) \le 2(r-2)$.

Remark 5.

For $r \geq 3$, the bound on the open packing number of rK_1 -free graphs given in Corollary 2 (and so the one in Lemma 10) is tight. For example, let H_r be a graph obtained by identifying a leaf vertex of a (distinct) P_3 with every vertex of a complete graph K_{r-2} , then H_r is a rK_1 -free graph (and so $(P_3 \cup (r-2)K_1)$ -free graph). An example of H_r for r=7 is given in Fig. 5. The vertex and edge set of H_r can be defined as $V(H_r) = \bigcup_{i=1}^{r-2} \{x_i, y_i, z_i\}$ and $E(H_r) = E_{r1} \cup E_{r2}$, where $E_{r1} = \bigcup_{i=1}^{r-2} \{x_i y_i, y_i z_i\}$ and $E_{r2} = \{z_i z_j : 1 \leq i < j \leq r-2\}$. Then, $S_r = \{x_1, x_2, \ldots, x_{r-1}, y_1, y_2, \ldots, y_{r-2}\}$ is an open packing (as well as a total dominating set) of size 2(r-2) in H_r . This also implies that the bound given in Lemma 9 is a near-optimal upper bound. Also, for the case r = 1, the bound given in Lemma 9 is tight, and a cycle on five vertices (C_5) is a connected $(P_3 \cup K_1)$ -free graph that satisfies this bound. Further, the graphs obtained by replacing every vertex of C_5 with a complete graph will remain as a $(P_3 \cup K_1)$ -free graph satisfying this bound.




Figure 5: A connected $7K_1$ -free graph H_7 with an open packing (as well as a total dominating set) $S_7 = \{x_1, x_2, x_3, x_4, x_5, y_1, y_2, y_3, y_4, y_5\}$ of size 2(7-2) = 10. Note that H_7 is also a $(K_2 \cup 6K_1)$ -free graph.

$(K_2 \cup rK_1)$ -free Graphs

For $r \ge 1$, it is known by Corollary 1 that $\gamma_t(G) \le 2r$ for every connected $(K_2 \cup rK_1)$ -free graph. In this section, we show that the open packing number of a connected $(K_2 \cup rK_1)$ -free graph is at most max $\{r + 1, 2(r - 1)\}$.

Remark 6.

Let G be a connected $(K_2 \cup K_1)$ -free graph. Then, G is also a connected P_4 -free graph. So, $\rho^o(G) \leq \gamma_t(G) = 2(=r+1)$.

Lemma 11. For $r \ge 2$, the open packing number of a connected $(K_2 \cup rK_1)$ -free graph is at most $\max\{r+1, 2(r-1)\}$.

Proof. We prove the lemma in two cases.

Case 1: r = 2.

Then, $\max\{r+1, 2(r-1)\} = r+1 = 3$. Since every connected $(K_2 \cup 2K_1)$ -free graph G is also a connected $(P_4 \cup K_1)$ -free graph, $\rho^o(G) \le 2(r-1)+1 = 3$ by Lemma 8. Case 2: $r \ge 3$.

Then, $\max\{r+1, 2(r-1)\} = 2(r-1)$. We prove that $\rho^o(G) \leq 2(r-1)$. On the contrary, assume that there exists a connected $(K_2 \cup rK_1)$ -free graph G with $\rho^o(G) > 2(r-1)$. We know that every open packing of size at least 2(r-1) + 1 = 2r - 1 contains an open packing of size 2r - 1. So, to contradict the assumption, it is enough to eliminate the possibility of an open packing of size 2r - 1 in G. Let S be an open packing of size 2r - 1 in G. Then, since G[S] is $\{P_3, K_3\}$ -free, every component of G[S] contains at most two vertices, and so the number of components in G[S] is at least $\lceil \frac{2r-1}{2} \rceil > r - 1$. Also, since 2r - 1 is odd, there exists at least one isolated vertex in G[S] due to the fact that every component of G[S] contains at most two vertex a vertex $v \in V(G)$ such that $uv \in E(G)$. Note that $G[S \setminus \{u\}]$ contains at least (r-1) components. Let G_1, G_2, \ldots, G_p be the components of $G[S \setminus \{u\}]$ for some $p \ge r - 1$. We now contradict our assumption in two cases based on the value of p.

Case 2.1: $p \ge r$.

Let $v_i \in V(G_i)$ for i = 1, 2, ..., r. Since each v_i is from a distinct component of $G[S \setminus \{u\}], v_i \neq v_j$ for $i, j \in \{1, 2, ..., r\}$ and $i \neq j$. Also, since $v_i \in S \setminus \{u\}, v_i \neq u$ and $v_i u \notin E(G[S]) \subseteq E(G)$ by the fact that u is an isolated vertex in G[S]. This implies that $v_i \neq v$ for i = 1, 2, ..., r since $uv \in E(G)$. Further $v_i v \notin E(G)$ since $v \in N(u)$ and $N(u) \cap N(v_i)$ is empty because S is an open packing in G with $u, v_i \in S$ for every i = 1, 2, ..., r. This implies that $\{u, v\} \cup \{v_1, v_2, ..., v_r\}$ induces a $(K_2 \cup rK_1)$ in G, a contradiction.

Case 2.2: p = r - 1.

Then, $G_1, G_2, \ldots, G_{r-1}$ are the components in $G[S \setminus \{u\}]$. Recall that $|V(G_i)| \leq 2$ for every $1 \leq i \leq r-1$. But since $2r-2 = |S \setminus \{u\}| = |\bigcup_{i=1}^{r-1} V(G_i)| \leq 2r-2$, every G_i consists of exactly two vertices. Let $V(G_i) = \{u_i, v_i\}$ for $1 \leq i \leq r-1$. Since G is connected, there exists a shortest path joining u_1 and u. Let $x_1x_2\ldots x_s$ with $u = x_1$ and $u_1 = x_s$ be a shortest path joining u_1 and u in G (see Fig 6). Since S is an open packing in G with $u, v_1 \in S$, $N(u) \cap N(v_1)$ is empty. So,




Figure 6: (a) G[S] along with the path $x_1x_2...x_s$ joining $u = x_1$ and $u_1 = x_s$ in G such that $x_{s-1} \neq v_1$ and (b) G[S] along with the path $x_1x_2...x_s$ joining $u = x_1$ and $u_1 = x_s$ in G such that $x_{s-1} = v_1$. Note that $\{x_2, x_3, ..., x_{s-2}\} \cap \{u_2, v_2, ..., u_{r-1}, v_{r-1}\}$ need not be empty.

 $uu_1 \notin E(G)$ since $u_1v_1 \in E(G)$. Similarly, $N(u) \cap N(u_1)$ is empty. This implies that $s \ge 4$. So, $u \notin \{x_{s-2}, x_{s-1}\}$. We further divide this case into two cases based on whether $x_{s-1} = v_1$ or not. Case 2.2.1: $x_{s-1} \neq v_1$.

Then, since $x_{s-1}x_s = x_{s-1}u_1 \in E(G)$, i.e., $x_{s-1} \in N(u_1)$ and S is an open packing with $u_1, v_i \in S$, $x_{s-1} \notin N(v_i)$ (i.e., $x_{s-1}v_i \notin E(G)$) for every $i = 1, 2, \ldots, r-1$. For similar reasons, $x_{s-1}u, x_{s-1}u_2 \notin E(G)$. The existence of u_2 is guaranteed by the fact that $r-1 \geq 2$. Also, note that $uv_i, u_2v_\ell, v_iv_j \notin E(G[S]) \subseteq E(G)$ for every $i, j, \ell \in \{1, 2, \ldots, r-1\}$ and $i \neq j, \ell \neq 2$ (see Fig. 6). Thus, $\{u_2, v_2\} \cup \{x_{s-1}\} \cup \{v_1\} \cup \{v_3\} \cup \{v_4\} \cup \ldots \cup \{v_{r-1}\} \cup \{u\}$ is a $(K_2 \cup rK_1)$ in G, a contradiction. Case 2.2.2: $x_{s-1} = v_1$.

Then, similar to the above case, it can be proved that $x_{s-2}u, x_{s-2}u_i, x_{s-2}v_2 \notin E(G)$ for every $i = 1, 2, \ldots, r-1$. Also, $uu_i, v_2u_\ell, u_iu_j \notin E(G)$ for every $i, j, \ell \in \{1, 2, \ldots, r-1\}$ and $i \neq j$, $\ell \neq 2$. So, $\{u_2, v_2\} \cup \{x_{s-2}\} \cup \{u_1\} \cup \{u_3\} \cup \{u_4\} \cup \ldots \cup \{u_{r-1}\} \cup \{u\}$ is a $(K_2 \cup rK_1)$ in G, a contradiction.

Remark 7.

The bound given in Lemma 11 is tight. For $r \ge 3$, let H_{r+1} be the graph defined in Remark 5. Then, H_{r+1} is a $(K_2 \cup rK_1)$ -free graph, and the set S_{r+1} in Remark 5 is an open packing of size 2(r-1).

Using the bounds obtained in the subclasses of $(P_4 \cup rK_1)$ -free graphs studied so far, we show that TOTAL DOMINATING SET and OPEN PACKING are in P in the subclasses of $(P_4 \cup rK_1)$ -free graphs.

Theorem 7. Given a graph class \mathcal{G} , TOTAL DOMINATING SET is in P on the class \mathcal{G} if \mathcal{G} is (a) $(P_4 \cup rK_1)$ -free graphs, (b) $(P_3 \cup rK_1)$ -free graphs, (c) $(K_2 \cup rK_1)$ -free graphs, or (d) rK_1 -free graphs.

The above theorem follows from Lemmas 3, 6, 7, and 9, as well as Corollaries 1 and 2. The graph classes $(P_3 \cup rK_1)$ -free graphs, $(K_2 \cup rK_1)$ -free graphs, and rK_1 -free graphs are explicitly stated in Theorem 7 for clarity.

Theorem 8. Given a graph class \mathcal{G} , if \mathcal{G} is (a) $(P_4 \cup rK_1)$ -free graphs $(r \ge 0)$, (b) $(P_3 \cup rK_1)$ -free graphs $(r \ge 0)$, (c) $(K_2 \cup rK_1)$ -free graphs $(r \ge 1)$, or (d) rK_1 -free graphs $(r \ge 3)$, then for every fixed r and for every graph $G \in \mathcal{G}$, (i) there are polynomially many open packings in G, (ii) all open packings in G can be found in polynomial time, and hence $\rho^o(G)$ can be found in polynomial time.

The above theorem follows from Lemmas 5, 6, 8, 10, 11, Corollary 2, Remarks 4, 6, and the fact that for a graph G with components G_1, G_2, \ldots, G_k , $\rho^o(G) = \sum_{i=1}^k \rho^o(G_i)$. Next, we prove Theorem 2.

Theorem 2 (Restated). For $p \ge 4$, let H be a graph on p vertices. Then, OPEN PACKING is polynomial time solvable on the class of H-free graphs if and only if $H \in \{pK_1, (K_2 \cup (p - 2)K_1), (P_3 \cup (p - 3)K_1), (P_4 \cup (p - 4)K_1)\}$ unless P = NP.

Proof. Let H be a graph on p vertices for some $p \ge 4$ such that OPEN PACKING is polynomial time solvable on H-free graphs. Then, we prove that $H \in \{P_4 \cup (p-4)K_1, P_3 \cup (p-3)K_1, K_2 \cup (p-2)K_1, pK_1\}$ unless P = NP. On the contrary, assume that $H \notin \{P_4 \cup (p-4)K_1, P_3 \cup (p-3)K_1, K_2 \cup (p-2)K_1, pK_1\}$. Then, by Observation 1, H contains at least one of (i) K_3 , (ii) $2K_2$, (iii) C_4 , (iv) $K_{1,3}$, or (v) C_5 as its induced subgraph. So, (a) H'-free graphs are a subclass of H-free graphs for some $H' \in \{K_3, 2K_2, C_4, K_{1,3}, C_5\}$ and (b) OPEN PACKING is NP-complete on H'-free graphs for every $H' \in \{K_3, 2K_2, C_4, K_{1,3}, C_5\}$ (by Remark 1 and Theorem 3). Thus, by (a), (b), and the fact that OPEN PACKING is in NP in every subclass of simple graphs, we can conclude that OPEN PACKING is NP-complete on H-free graphs, a contradiction. This proves the necessary part of Theorem 2.

The sufficiency part of Theorem 2 is held by Theorem 8.

Note that Observation 1, along with Theorem 7, and the fact that TOTAL DOMINATING SET is NP-complete for (i) Split graphs ($\{2K_2, C_4, C_5\}$ -free graphs) [1], (ii) Bipartite graphs (a subclass of K_3 -free graphs) [16], and (iii) Line graphs (a subclass of $K_{1,3}$ -free graphs) [14], imply the following theorem, similar to Theorem 2, for TOTAL DOMINATING SET.

Theorem 9. For $p \ge 4$, let H be a graph on p vertices. Then, TOTAL DOMINATING SET is polynomial time solvable on the class of H-free graphs if and only if $H \in \{pK_1, (K_2 \cup (p-2)K_1), (P_3 \cup (p-3)K_1), (P_4 \cup (p-4)K_1)\}$ unless P = NP.

3.3 H on three vertices

In this section, we give the proof of Theorem 1.

Theorem 1 (Restated). Let H be a graph on three vertices. Then, an optimal open packing on H-free graphs can be found in polynomial time if and only if $H \ncong K_3$ unless P = NP.

Proof. Note that $3K_1, (K_2 \cup K_1), P_3$, and K_3 are the only non-isomorphic graph on three vertices. By Theorem 8, an optimal open packing of a *H*-free graph can be found in polynomial time if $H \in \{3K_1, (K_2 \cup K_1), P_3\}$ and OPEN PACKING is NP-complete on K_3 -free graphs by Remark 1. This completes the proof of Theorem 1.

Note that TOTAL DOMINATING SET is NP-complete for K_3 -free graphs [16]. So, by Theorem 7, a result similar to Theorem 1 for TOTAL DOMINATING SETholds. This note completes our study on the complexity (P vs NPC) of TOTAL DOMINATING SET and OPEN PACKING on H-free graphs. Next, we study the complexity of these problems on some subclasses of split graphs.

4 Split Graphs

Ramos et al. [19] proved that OPEN PACKING is NP-complete on split graphs. In this section, we study the complexity of OPEN PACKING in (i) $K_{1,r}$ -free split graphs for $r \ge 2$ and (ii) I_r -split graphs for $r \ge 1$. This study shows the complexity (P vs NPC) difference between TOTAL DOMINATING SET and OPEN PACKING in split graphs (see Table 2).

4.1 $K_{1,r}$ -free Split Graphs

It is known that TOTAL DOMINATING SET is NP-complete on $K_{1,5}$ -free split graphs [24] and is polynomial time solvable on $K_{1,4}$ -free split graphs [20]. In this section, we give a dichotomy result for OPEN PACKING in $K_{1,r}$ -free split graphs by proving that the problem is (i) NP-complete for $r \geq 4$ and (ii) polynomial time solvable for $r \leq 3$.

We begin this section with a construction to prove that OPEN PACKING is NP-complete on $K_{1,4}$ -free split graphs, which is a reduction from INDEPENDENT SET, a well-known NP-complete problem [12].




Figure 7: (a) a graph G and (b) the graph G' constructed from G using Construction 2

Construction 2.

Input: A simple graph G(V, E).

Output: A $K_{1,4}$ -free split graph $G'(C \cup I, E)$.

Guarantee: G has an independent set of size k if and only if G' has an open packing of size k + 1. Procedure:

Step 1: Subdivide each edge e = uu' in G into a three vertex path ueu' in G'.

Step 2: Introduce three new vertices, x, y, and z, and two edges, xy and xz, in G'.

Step 3: Make $E(G) \cup \{y, z\}$ a clique in G'.

An example of Construction 2 is shown in Fig. 7. The vertex set and the edge set of the graph G' are $V(G') = V(G) \cup E(G) \cup \{x, y, z\}$ and $E(G') = \{ue : u \in V(G), e \in E(G) \text{ and } e \in E_G(u)\} \cup \{xy, xz, yz\} \cup \{ye : e \in E(G)\} \cup \{ze : e \in E(G)\} \cup \{ee' : e, e' \in E(G) \text{ and } e \neq e'\}.$ Note that |V(G')| = n + m + 3 and $|E(G')| = 2m + 3 + 2m + \binom{m}{2} = O(m^2)$. Hence, the graph G' can be constructed in polynomial time from G. Also, by Construction 2, $V(G') = C \cup I$ is a clique-independent set partition of G' with $C = E(G) \cup \{y, z\}$ and $I = V(G) \cup \{x\}$. Thus, $G'(C \cup I, E)$ is a split graph.

Claim 6: The graph G' output by Construction 2 is a $K_{1,4}$ -free graph.

On the contrary, assume that $\{w, w_1, w_2, w_3, w_4\}$ induces a $K_{1,4}$ in G' with w as its centre. Then, $\{w_1, w_2, w_3, w_4\}$ is an independent set in G'. So $w \notin I$, since $N_{G'}(v) \subseteq C$ is a clique for every vertex $v \in I$, whereas $N_{G'}(w)$ contains an independent set on four vertices. This implies that $w \in C$. By Construction 2, for every vertex $v \in C$, $|N_{G'}(v) \cap I| \leq 2$, and so w has at most two neighbours in I. Therefore, $|\{w_1, w_2, w_3, w_4\} \cap C| \geq 2$, a contradiction to C being a clique and $\{w_1, w_2, w_3, w_4\}$ being an independent set in G'. Thus, G' is a $K_{1,4}$ -free graph.

The following claims help us prove the guarantee of Construction 2.

Claim 7: A set $S \subseteq V(G)$ is an independent set in the input graph G of Construction 2 if and only if S is an open packing in the output graph G'.

Proof of Claim 7 is similar to the proof of Claim 1, and so it is moved to Appendix C.

Claim 8: For every open packing S in the output graph G' of Construction 2, $|S \cap (C \cup \{x\})| = |S \cap (E(G) \cup \{x, y, z\})| \le 1$.

Observe that in G', the graph induced by $C \cup \{x\}$ consists of a complete graph (G'[C]) on m + 2 vertices together with a vertex x, which is adjacent to exactly two vertices (y and z) in C. So, any two distinct vertices in $C \cup \{x\}$ have a common neighbour. Thus, the claim holds.

Proof of Guarantee of Construction 2. We prove the guarantee of Construction 2 in two cases. Case 1: G has an independent set S of size k.

Then, by Claim 7, $S \subseteq V(G)$ is an open packing in G' of size k. To prove that G' has an open packing of size k + 1, we show that $S \cup \{x\}$ is an open packing in G'. Note that by Construction 2,

 $N_{G'}(u) \cap N_{G'}(x) \subseteq E(G) \cap \{y, z\} = \emptyset$ for every vertex $u \in V(G)$ (in particular, for every vertex $u \in S$). So, $S \cup \{x\}$ is an open packing of size k + 1 in G'.

Case 2: G' has an open packing S of size k + 1.

Then, by Claim 8, $|S \cap (E(G) \cup \{x, y, z\})| \leq 1$. So, $S' = S \setminus (E(G) \cup \{x, y, z\})$ is an open packing of size at least k in G'. Also, by the construction of G', we have $S' \subseteq V(G)$. Hence, by Claim 7, S' is an independent set of size at least k in G.

Thus, by Cases 1 and 2, the guarantee of Construction 2 holds.

The following theorem is held by Construction 2 and the fact that INDEPENDENT SET is NPcomplete on simple graphs [12].

Theorem 10. OPEN PACKING is NP-complete on $K_{1,4}$ -free split graphs.

Similar to that of Theorems 5 and 6, it can be proved that on $K_{1,4}$ -free split graphs (i) OPEN PACKING parameterized by solution size is W[1]-complete and (ii) MAX-OPEN PACKING is hard to approximate within a factor of $N^{(\frac{1}{2}-\epsilon)}$ for any $\epsilon > 0$ unless P = NP, where N denotes the number of vertices in a $K_{1,4}$ -free split graph (proof of (ii) is given as Theorem 16 in Appendix C).

Next, to complete the dichotomy result of OPEN PACKING in $K_{1,r}$ -free split graphs, we prove that a maximum open packing in $K_{1,3}$ -free split graphs can be found in polynomial time, and the following claim and lemma are useful in proving it.

Claim 9: If S is an open packing in a connected split graph $G(C \cup I, E)$ with $S \cap C \neq \emptyset$, then for every vertex $u \in S \cap C$, $S \subseteq N[u]$, and so $|S| \leq 2$.

Let S be an open packing in a connected split graph $G(C \cup I, E)$ with $S \cap C \neq \emptyset$. Let $u \in S \cap C$. Then, we show that $S \subseteq N[u]$. On the contrary, assume that there exists a vertex $v \in S \setminus N[u]$. Since $C \subseteq N[u]$ and $v \notin N[u]$, $v \notin C$, i.e., $v \in I$. Since G is connected, $deg(v) \ge 1$, and so there exists $w \in C$ such that $vw \in E(G)$. Since $v \notin N(u)$, $w \neq u$. Further, since $w, u \in C$, $w \in N(u)$. This implies that $w \in N(u) \cap N(v)$, a contradiction to S is an open packing in G and $u, v \in S$. Therefore, $S \subseteq N[u]$. Since S is an open packing in G, $|S \cap N(u)| \le 1$. So, $|S| \le |S \cap N(u)| + 1 \le 2$.

Lemma 12 ([20]). If $G(C \cup I, E)$ is a $K_{1,3}$ split graph with a vertex $u \in C$ such that $|N(u) \cap I| = 2$, then $|I| \leq 3$.

Theorem 11. OPEN PACKING is polynomial time solvable on $K_{1,3}$ -free split graphs.

Proof. Let $G(C \cup I, E)$ be a $K_{1,3}$ -free split graph. W.L.O.G., assume that $C \neq \emptyset$ and $I \neq \emptyset$. Since G is $K_{1,3}$ -free, $|N(u) \cap I| \leq 2$ for every $u \in C$. We show that $\rho^o(G) \in \{1, 2, |I|\}$ in two cases based on $|N(u) \cap I|$ for $u \in C$.

Case 1: $|N(u) \cap I| \leq 1$ for every $u \in C$.

Let $I = \{x_1, x_2, \ldots, x_k\}$. Let $V_i = N(x_i) \cap C$ for $i = 1, 2, \ldots, k$ and $V_0 = C \setminus (\bigcup_{i=1}^k V_i)$. Then, $V_i \cap V_j = \emptyset$ for every $i, j \in \{1, 2, \ldots, k\}$ and $i \neq j$. If not, let $u' \in V_i \cap V_j$ for some $i, j \in \{1, 2, \ldots, k\}$ with $i \neq j$. Then, $x_i, x_j \in N(u') \cap I$, a contradiction to $|N(u) \cap I| \leq 1$ for every $u \in C$. We prove that in this case $\rho^o(G) \in \{2, |I|\}$, and the proof is divided into two subcases. *Case 1.1:* If k = 1 and $deg(x_1) = 1$, then $\rho^o(G) = 2$.

Let $N(x_1) = \{y_1\}$. Then, $N(x_1) \cap N(y_1) = \{y_1\} \cap N(y_1) = \emptyset$. So, $\{x_1, y_1\}$ is an open packing of size two in G. Thus, $\rho^o(G) \ge 2$. Next, we show that $\rho^o(G) \le 2$. On the contrary, assume that there exists an open packing S of size at least three in G. Then, by Claim 9, $S \cap C = \emptyset$, i.e., $S \subseteq I$. This implies that $|S| \le |I| = 1$, a contradiction. Therefore, $\rho^o(G) \le 2$. As we have already proved that $\rho^o(G) \ge 2$ in this case, we can conclude that $\rho^o(G) = 2$.

Case 1.2: If (i) k > 1 or (ii) k = 1 and $deg(x_1) > 1$, then $\rho^o(G) = |I|$.

Note that the assumption in Case 1.2 is the negation of Case 1.1. As we have already proved that $N(x_i) \cap N(x_j) = V_i \cap V_j = \emptyset$ for every $x_i, x_j \in I$, I is an open packing in G, and hence $\rho^o(G) \ge |I| = k$. Next, we show that $\rho^o(G) \le |I| = k$. On the contrary, assume that $\rho^o(G) > k$. Let S be an open packing in G of size $\rho^o(G)$. Then, |S| > |I| = k, and so $S \cap C \neq \emptyset$. Therefore, by Claim 9, $|S| \le 2$. This implies that $2 \ge |S| > |I| \ge 1$. So, |S| = 2 and |I| = k = 1. Then, $V(G) = C \cup \{x_1\}$, and by the assumption in Case 1.2, $deg(x_1) \ge 2$. Let $S = \{a, b\}$. Since G is constituted of a complete graph (G[C]) together with a vertex x_1 , which is adjacent to at least two vertices in C, for any choice of $a, b \in C \cup \{x_1\}$, a and b will have a common neighbour in G, a contradiction to S is an open packing of size two in G. Therefore, for every open packing S in $G, |S| \le |I| = k$. This implies that $\rho^o(G) \le |I|$. Since we have proved that I is an open packing S in $G, \rho^{o}(G) = |I|.$

Case 2: There exists $u \in C$ such that $|N(u) \cap I| = 2$.

In this case, we prove that $\rho^o(G) \in \{1,2\}$. Note that $|I| \leq 3$ by Lemma 12. Let $u \in C$ such that $|N(u) \cap I| = 2$. Let $v, w \in (N(u) \cap I)$. Then, since $|I| \leq 3$, $|I \setminus \{v, w\}| \leq 1$. This implies that for any open packing S in G with $S \subseteq I$, $|S| \leq |S \cap \{v, w\}| + |S \cap (I \setminus \{v, w\})| \leq 2$ because $N(v) \cap N(w) \neq \emptyset$ with $u \in N(v) \cap N(w)$ (and so $|S \cap \{v, w\}| \leq 1$). Also, by Claim 9, $|S| \leq 2$ if $S \cap C \neq \emptyset$. Since any open packing S in G falls under one of the above two categories (i.e., either $S \subseteq I$ or $S \cap C \neq \emptyset$), $|S| \leq 2$ for every open packing S in G. So, $\rho^o(G) \leq 2$. Next, we complete the proof of Case 2 in two cases.

Case 2.1: If (i) there exist $x, y \in I$ such that $N(x) \cap N(y) = \emptyset$ or (ii) there exists $x \in I$ such that deg(x) = 1, then $\rho^o(G) = 2$.

If there exist $x, y \in I$ such that $N(x) \cap N(y) = \emptyset$, then $\{x, y\}$ is an open packing of size two in G, and so $\rho^o(G) = 2$. Also, if there exists a vertex $x \in I$ such that deg(x) = 1, then it is evident that n[x] is an open packing of size two. Thus, $\rho^o(G) = 2$.

Case 2.2: If (i) $N(x) \cap N(y) \neq \emptyset$ for every $x, y \in I$, and (ii) for every $x \in I$, deg(x) > 1, then $\rho^{o}(G) = 1$.

Note that the assumption in Case 2.2 is the negation of Case 2.1. We prove that $\rho^o(G) = 1$ in this case. As we have already proved that $\rho^o(G) \leq 2$, it is enough to rule out the case $\rho^o(G) = 2$. Since $V(G) \subseteq C \cup I$, where (i) any two vertices in I have a common neighbour, (ii) every vertex $x \in I$ has at least two neighbours in C, and (iii) G[C] is a complete graph, it is evident that any two vertices in V(G) have a common neighbour in G, and so $\rho^o(G) < 2$. Therefore, $\rho^o(G) = 1$.

Hence, we have shown every possible value of $\rho^o(G)$. Note that every $K_{1,3}$ -free split graph G falls under Case 1 or 2. It can be easily verified in polynomial time whether a $K_{1,3}$ -free split graph G falls under Case 1 or Case 2. Also, the conditions given in Cases 1.1, 1.2, 2.1, and 2.2 are polynomial time verifiable. Hence, the open packing number of a $K_{1,3}$ -free split graph can be found in polynomial time.

4.2 I_r -split graphs

We observed that a minor modification in Corneil and Perl's reduction [1] would show that TOTAL DOMINATING SET is NP-complete on I_2 -split graphs. We extended this by proving a dichotomy that TOTAL DOMINATING SET in I_r -split graphs is (i) NP-complete for $r \ge 2$ and (ii) polynomial time solvable for r = 1. In addition, we prove that OPEN PACKING is (i) NP-complete on I_r -split graphs for $r \ge 3$ and (ii) polynomial time solvable on I_r -split graphs for $r \le 2$. We begin the section with Construction 3 to prove that TOTAL DOMINATING SET is NP-complete on I_r -split graphs for $r \ge 2$ through a reduction from r-HITTING SET problem, which is known to be NP-complete for $r \ge 2$ [12].

Construction 3.

Input: A set U and a set W of r-sized subsets of U for some $r \geq 2$.


Output: An I_r -split graph $G(C \cup I, E)$.

Guarantee: For $k \leq |U|$, (U, W) has a hitting set of size k if and only if G has a total dominating set of size k + 1.

Procedure:

- Step 1: For every element $u \in U$, create a vertex u in G. Similarly, for every $W \in W$, create a vertex z_W in G.
- Step 2: Create r + 1 new vertices $\{x_1, x_2, \dots, x_r, y\}$ in G.
- Step 3: Introduce an edge between every pair of distinct vertices of $U \cup \{x_1, x_2, \dots, x_r\}$ in G.
- Step 4: Introduce an edge between a vertex $u \in U$ and z_W for $W \in W$ in G if $u \in W$.
- Step 5: Add an edge between x_i and y for every $1 \le i \le r$.

An example of Construction 3 for the case r = 3 is given in Fig. 8. Note that $V(G) = C \cup I$, where $C = U \cup \{x_1, x_2, \ldots, x_r\}$ and $I = \{z_W : W \in W\} \cup \{y\}$ and $E(G) = \{uv : u, v \in U \cup \{x_1, x_2, \ldots, x_r\}\} \cup \{uz_W : u \in U, W \in W \text{ and } u \in W\} \cup \{x_iy : 1 \leq i \leq r\}$. So, |V(G)| = |U| + |W| + r + 1 and $|E(G)| = \binom{|U|+r}{2} + (r \cdot |W|) + r$. Hence, the graph G can be constructed in polynomial time with respect to |U|, |W|, and r. By the construction of G, it is clear that $V(G) = C \cup I$ is a clique-independent set partition of the vertex set of G. So, G is a split graph. Since the size of every set $W \in W$ is r, $deg_G(z_W) = r$ by the construction of G. Further, since $N_G(y) = \{x_1, x_2, \ldots, x_r\}, deg_G(y) = r$. This implies that $deg_G(u) = r$ for every $u \in I$. Thus,



 $U = \{u_1, u_2, u_3, u_4, u_5, u_6\}$

 $\mathcal{W} = \{W_1, W_2, W_3, W_4\}$

where $W_1 = \{u_1, u_2, u_5\}, W_2 = \{u_3, u_5, u_6\}$ $W_3 = \{u_1, u_3, u_6\}, W_4 = \{u_2, u_4, u_5\}$

(a)



Figure 8: (a) An instance (U, W) of 3-HITTING SET problem and (b) a I_3 -split graph $G(C \cup$ I, E) produced by Construction 3, where $C = U \cup \{x_1, x_2, x_3\}$ is a clique in G and I = $\{z_{W_1}, z_{W_2}, z_{W_3}, z_{W_4}, y\}$ is an independent set in G with $deg_G(z_{W_i}) = deg_G(y) = 3$ for every i = 1, 2, 3, 4 and $V(G) = C \cup I$.

 $G(C \cup I, E)$ is an I_r -split graph.

Proof of Guarantee of Construction 3.

Case 1: The instance (U, W) of r-HITTING SET has a hitting set of size k.

Let $U' = \{u_1, u_2, \ldots, u_k\}$ be a hitting set of size k in (U, \mathcal{W}) . Then, we show that $\{u_1, u_2, \ldots, u_k, x_1\}$ is a total dominating set of size k+1 in G. By the construction of G, every vertex in $C \cup \{y\}$ is adjacent to u_1 or x_1 . So, to prove the claim, it is enough to show that every vertex in $(I \setminus \{y\}) = \{z_W : W \in \mathcal{W}\}$ is adjacent to some vertex in $\{u_1, u_2, \dots, u_k, x_1\}$. Let $W \in \mathcal{W}$. Then, since $\{u_1, u_2, \ldots, u_k\}$ is a hitting set of (U, \mathcal{W}) , there exists $i \in \{1, 2, \ldots, k\}$ such that $u_i \in W$. Hence, by the construction of G, z_W is adjacent to u_i . Since W is an arbitrary set in \mathcal{W} , for every $W \in \mathcal{W}$, the corresponding vertex z_W in G is adjacent to some vertex in $\{u_1, u_2, \ldots, u_k\}$. Thus, every vertex in $V(G) = C \cup I$ is adjacent to some vertex in $\{u_1, u_2, \ldots, u_k, x_1\}$, and hence $\{u_1, u_2, \ldots, u_k, x_1\}$ is a total dominating set of size k + 1 in G. Case 2: G has a total dominating set of size k + 1.

Let D be a total dominating set of size k + 1 in G. Then, $D \cap \{x_1, x_2, \ldots, x_r\} \neq \emptyset$ because $N_G(y) = \{x_1, x_2, \dots, x_r\}$ and $D \cap N_G(y) \neq \emptyset$. Let $D' = D \cap U$. Since $D \cap \{x_1, x_2, \dots, x_r\} \neq \emptyset$ and $U \cap \{x_1, x_2, \dots, x_r\} = \emptyset, |D'| = |D \cap U| \le k$. Next, we show that D' is a hitting set of size at most k of (U, \mathcal{W}) . Let $W \in \mathcal{W}$. Then, since D is a total dominating set, there exists a vertex $u \in D$ such that $uz_W \in E(G)$. Also, since $N_G(z_W) \subseteq U$ by the construction of $G, u \in D \cap U = D'$. So, by the construction of $G, u \in W$. Since W is an arbitrary set in \mathcal{W} , for every $W \in \mathcal{W}$, there exists an element $u \in D'$ such that $u \in W$. Hence, D' is a hitting set of size at most k in (U, W).

The theorem below follows from Construction 3 and the fact that r-HITTING SET is NP-complete for $r \ge 2$ [12].

Theorem 12. TOTAL DOMINATING SET is NP-complete on I_r -split graphs for $r \geq 2$.

The following theorem completes the dichotomy result for TOTAL DOMINATING SET on I_r -split graphs.

Theorem 13. Let $G(C \cup I, E)$ be a connected non-trivial I_1 -split graph. Then, $\gamma_t(G) = \max\{2, |N(I)|\}$, where $N(I) = \bigcup N(u)$. Hence, an optimal total dominating set of G can be found in linear time. $u \in I$

Proof. If $I = \emptyset$, then G is a complete graph, and so $\gamma_t(G) = 2$. Thus, assume that $I \neq \emptyset$. Note that by the definition of I_1 -split graphs, every vertex in I is a pendant vertex in G. Since $N(u) \cap D$ is non-empty for every total dominating set D in G, $N(u) \subseteq D$ for every pendant vertex u in G. This implies that $N(I) \subseteq D$ for every total dominating set D in G, and so $|N(I)| \leq \gamma_t(G)$. We prove the theorem in two cases based on |N(I)|.

Case 1: |N(I)| = 1.

Let $N(I) = \{u\}$. Since I is an independent set in $G, u \in C$. Since C is a clique and $N(I) = \{u\}$, every vertex in $V(G) \setminus \{u\} = (C \cup I) \setminus \{u\}$ is adjacent to u. Let $v \in I$. Then, since $N(I) = \{u\}$, $vu \in E(G)$. This implies that every vertex in $V(G) = (V(G) \setminus \{u\}) \cup \{u\}$ is adjacent to a vertex in $\{u, v\}$. So, $\{u, v\}$ is a total dominating set in G, and hence $\gamma_t(G) \leq 2$. Since $\gamma_t(H) \geq 2$ for every graph H that admits a total dominating set, $\gamma_t(G) = 2$. *Case 2:* $|N(I)| \ge 2$.

Since I is an independent set in G, $N(I) \subseteq C$. Let $u, u' \in N(I)$. Then, since C is a clique and $u, u' \in N(I) \subseteq C$, every vertex in C is adjacent to u or u'. Note that by the definition of N(I), every vertex in I is adjacent to some vertex in N(I). Thus, every vertex in $V(G) = C \cup I$ is adjacent to a vertex in N(I), and so N(I) is a total dominating set. This implies that $\gamma_t(G) \leq |N(I)|$. Recall that $\gamma_t(G) > |N(I)|$. Hence, $\gamma_t(G) = |N(I)|$.

Thus, by Cases 1 and 2, $\gamma_t(G) = \max\{2, |N(I)|\}$. Note that |N(I)| can be found in linear time. So, an optimal total dominating set of G can be found in linear time.

Next, we prove the dichotomy result on OPEN PACKING in I_r -split graphs. Firstly, we show that OPEN PACKING is NP-complete on I_r -split graphs for $r \geq 3$ through a reduction from r-DIMENSIONAL MATCHING problem, which is known to be NP-complete for $r \geq 3$ [12].

Construction 4.

Input: A collection of sets X_1, X_2, \ldots, X_r such that $|X_i| = q$ for $i = 1, 2, \ldots, r$, for some $q \in \mathbb{N}$, and a non-empty set $M \subseteq \prod_{i=1}^{r} X_i$. Output: An I_r -split graph $G(C \cup I, E)$.

Guarantee: $(X_1, X_2, \ldots, X_r, M)$ is a yes-instance of r-DIMENSIONAL MATCHING if and only if G has an open packing of size q.

Procedure:

- For every $1 \leq i \leq r$ and for every $x \in X_i$, create a vertex $z_{(x,i)}$ in G. Similarly, for Step 1: $w \in M$, create a vertex y_w in G.
- Introduce an edge between every pair of distinct vertices of $C = \bigcup_{i=1}^{r} \{z_{(x,i)} : x \in X_i\}$ Step 2: in G.
- Introduce an edge between the vertex $z_{(x,i)} \in C$ and y_w in G if the i^{th} coordinate of Step 3: w is x.

An example of Construction 4 is given in Fig. 9. The vertex and edge sets of the graph G are $V(G) = C \cup I, \text{ where } I = \{y_w \, : \, w \in M\} \text{ and } E(G) = \{xx' \, : \, x, x' \in C\} \cup \{z_{(x,i)}y_w \, : \, x \in X_i, w \in X_i\}$ M, and the *i*th coordinate of w is x. Then, $|V(G)| = (r \cdot q) + |M|$ and $|E(G)| = r \cdot |M| + {r \cdot q \choose 2}$. Hence, the graph G can be constructed in polynomial time with respect to r, q, and |M|. Note that by the construction of G, C is a clique and I is an independent set, and hence $V(G) = C \cup I$ is a clique-independent set partition of V(G). Thus, $G(C \cup I, E)$ is a split graph. Further, by the construction of G, it is clear that for $y_w \in I$, $N_G(y_w) = \{z_{(x_1,1)}, z_{(x_2,2)}, \ldots, z_{(x_r,r)}\}$ if w = (x_1, x_2, \ldots, x_r) . Hence, $deg_G(y_w) = r$ for every $y_w \in I$. Thus, G is an I_r -split graph. Next, we prove the guarantee of Construction 4. The following claim is used in the proof of guarantee of Construction 4.

Claim 10: If S is an open packing in the output graph G of Construction 4 such that $S \cap C$ is non-empty, then |S| = 1.


On the contrary, assume that there exists an open packing S in G such that $S \cap C$ is non-empty and |S| > 1. Let $u \in S \cap C$, and let $z \in S \setminus \{u\}$. Since $r \geq 3$ and $q \geq 1$, $|C| \geq 3$. This implies that C is a clique of size at least three in G, and so any two vertices in C have a common neighbour in G. This says that $z \notin C$. So, $z \in I$, i.e., $z = y_w$ for some $w \in M$. Since $r \geq 3$ and $deg_G(y_w) = r$ with $N_G(y_w) \subseteq C$, $(z =)y_w \in I$ is adjacent to at least two vertices other than u in C. This implies that $N_G(y_w) \cap N_G(u) \neq \emptyset$, a contradiction to $u, z(=y_w) \in S$ and S is an open packing in G.

Proof of Guarantee of Construction 4. We prove the Guarantee of Construction 4 in two cases. Case 1: $(X_1, X_2, \ldots, X_r, M)$ is a yes-instance of r-DIMENSIONAL MATCHING.

$$X_1 = \{x_1, u_1, v_1\} \qquad X_2 = \{x_2, u_2, v_2\} \qquad X_3 = \{x_3, u_3, v_3\} \qquad X_4 = \{x_4, u_4, v_4\}$$
$$M = \{w_1, w_2, w_3, w_4, w_5\}$$

where
$$w_1 = (x_1, u_2, v_3, v_4)$$
 $w_2 = (x_1, x_2, v_3, v_4)$ $w_3 = (u_1, u_2, u_3, u_4)$
 $w_4 = (v_1, v_2, v_3, x_4)$ $w_5 = (v_1, v_2, x_3, x_4)$





(b) $G(C \cup I, E)$

Figure 9: (a) An instance (X_1, X_2, X_3, X_4, M) of 4-DIMENSIONAL MATCHING and (b) An I_r -split graph $G(C \cup I, E)$ corresponding to the instance (X_1, X_2, X_3, X_4, M) produced using Construction 4. The edges between the vertices of C aren't given in the figure for clarity. Note that for every $X_i = \{z_i, u_i, v_i\}$ in the instance of 4-DIMENSIONAL MATCHING, three vertices $z_{(x_i,i)}, z_{(u_i,i)}$ and $z_{(v_i,i)}$ are created in the graph G. W.L.O.G., assume that $\{w_1, w_2, \ldots, w_q\}$ is an *r*-DIMENSIONAL MATCHING of $(X_1, X_2, \ldots, X_r, M)$. Then, for $w_j = (x_1, x_2, \ldots, x_r)$ and $w_p = (u_1, u_2, \ldots, u_r)$ with $p, j \in \{1, 2, \ldots, q\}$ and $p \neq j$, $x_i \neq u_i$ for $i \in \{1, 2, \ldots, r\}$. Therefore, by the construction of G, $z_{(x_t,t)} \neq z_{(u_\ell,\ell)}$ for every $t, \ell \in \{1, 2, \ldots, r\}$. This implies that $N_G(y_{w_j}) \cap N_G(y_{w_p}) = \{z_{(x_1,1)}, z_{(x_2,2)}, \ldots, z_{(x_r,r)}\} \cap \{z_{(u_1,1)}, z_{(u_2,2)}, \ldots, z_{(u_r,r)}\} = \emptyset$. So, $\{y_{w_1}, y_{w_2}, \ldots, y_{w_q}\}$ is an open packing of size q in G. Case 2: G has an open packing S of size q.

We further divide this case into two cases based on $S \cap C$.

Case 2.1: $S \cap C$ is non-empty.

Then, by Claim 10, |S| = q = 1. Therefore, $|X_i| = 1$ for every $i \in \{1, 2, ..., r\}$. Let $X_i = \{x_i\}$ for i = 1, 2, ..., r. Since M is non-empty, there exists exactly one r-tuple $w \in M$ such that $w = (x_1, x_2, ..., x_r)$. Hence, $M = \{w\}$ is an r-DIMENSIONAL MATCHING of $(X_1, X_2, ..., X_r, M)$. Case 2.2: $S \cap C$ is empty.

So, $S \subseteq I$. W.L.O.G., assume that $S = \{y_{w_1}, y_{w_2}, \ldots, y_{w_q}\}$. Then, we show that $L = \{w_1, w_2, \ldots, w_q\}$ is an *r*-DIMENSIONAL MATCHING of $(X_1, X_2, \ldots, X_r, M)$. On the contrary, assume that there exist $j, p \in \{1, 2, \ldots, q\}$ and $i \in \{1, 2, \ldots, r\}$ such that $x_i = u_i$ for $w_j = (x_1, x_2, \ldots, x_r)$ and $w_p = (u_1, u_2, \ldots, u_r)$. Then, by the construction of G, $z_{(x_i,i)} = z_{(u_i,i)} \in N_G(y_{w_j}) \cap N_G(y_{w_p})$, a contradiction to $y_{w_j}, y_{w_p} \in S$ and S is an open packing in G. So, $L = \{w_1, w_2, \ldots, w_q\}$ is an *r*-DIMENSIONAL MATCHING of $(X_1, X_2, \ldots, X_r, M)$.

Theorem 14. For $r \geq 3$, OPEN PACKING is NP-complete on I_r -split graphs.

Proof. Given an I_r -split graph G and a vertex subset S of G, it can be tested in linear time whether S is an open packing in G or not using Lemma 4. Hence, OPEN PACKING is in the class NP on I_r -split graphs. Further, Construction 4 and the fact that r-DIMENSIONAL MATCHING is NP-complete for $r \geq 3$ [12] implies that OPEN PACKING is NP-complete on I_r -split graphs for $r \geq 3$.

The following construction is used to show that OPEN PACKING is polynomial time solvable on a superclass of I_1 -split graphs and I_2 -split graphs. Thus, completing our dichotomy result on I_r -split graphs.

Construction 5.

Input: A split graph $G(C \cup I, E)$ with $1 \le deg_G(v) \le 2$ for every $v \in I$.

Output: A graph G'(V', E').

Guarantee: For $k \in \mathbb{N}$, G has an open packing S such that $S \subseteq I$ and |S| = k if and only if G' has a matching of size k.

Procedure:

- Step 1: For every vertex $u \in C$ of G, create a vertex u in G'.
- Step 2: For every vertex $v \in I$ with $deg_G(v) = 2$ and $N_G(v) = \{u, u'\}$, create an edge e_v with $e_v = uu'$ in G'.
- Step 3: For every vertex $v \in I$ with $deg_G(v) = 1$ and $N_G(v) = \{u\}$, create a vertex v and an edge e_v with $e_v = uv$ in G'.

Let $J = \{v \in I : deg_G(v) = 1\}$. Then, the vertex set and the edge set of G' are $V' = C \cup J$ and $E' = \{e_v : v \in I\}$. Note that $|V'| \leq n$ and $|E'| \leq n$. Since G' is constructed based on $N_G(v)$ for every $v \in I$, it can be constructed in O(n + m) time. Also, note that the graph G' may have parallel edges. Next, we prove the guarantee of the above construction.

Proof of Guarantee of Construction 5. We prove the guarantee of the construction in two cases. Case 1: G has an open packing S such that |S| = k and $S \subseteq I$.

Let $F_S = \{e_v \in E' : v \in S\}$. We show that F_S is a matching in G'. On the contrary, assume that F_S is not a matching in G'. Then, there exist two distinct edges $e_v, e_y \in F_S$ such that $V_{G'}(e_v) \cap V_{G'}(e_y) \neq \emptyset$. Let $u \in V_{G'}(e_v) \cap V_{G'}(e_y)$, i.e., $e_v, e_y \in E_{G'}(u)$. Then, by the construction of G', $u \in C \cup J$ and $v, y \in I$. Firstly, we show that $u \notin J$. If not, $u \in J$. Then, $deg_G(u) = 1$, and so, by Step 3 of Construction 5, $E_{G'}(u) = \{e_u\}$. This implies that $e_v = e_u = e_y$, a contradiction to $v \neq y$. So, $u \notin J$. Hence, $u \in C$, and so by the construction of G', $uv, uy \in E(G)$. This implies that $u \in N_G(v) \cap N_G(y)$, a contradiction to $v, y \in S$ and S is an open packing in G. So, F_S is a matching of size k in G'.

Case 2: F is a matching in G' of size k.

Let $S_F = \{v \in I : y_v \in F\}$. In this case, we prove that S_F is an open packing in G. On the contrary, assume that S_F is not an open packing in G. Then, there exist vertices $v, y \in S_F \subseteq I$




Figure 10: (a) A split graph $G(C \cup I, E)$ with $1 \le deg_G(v) \le 2$ for every $v \in I$ and (b) the graph G'(V', E') corresponding to G constructed by Construction 5.

such that $N_G(v) \cap N_G(y) \neq \emptyset$. Let $u \in (N_G(v) \cap N_G(y)) \subseteq C$. This implies that $u \in V(G') = C \cup J$. Also, by the construction of G', $u \in V_{G'}(e_v) \cap V_{G'}(e_y)$, a contradiction to $e_v, e_y \in F$ and F is a matching in G'. Hence, S_F is an open packing in G such that $S \subseteq I$ and |S| = k.

The following lemma and Construction 5 help us compute a maximum open packing in I_1 -split graphs and I_2 -split graphs.

Lemma 13 ([15]). A maximum matching in a graph with n vertices and m edges can be found in $O(m\sqrt{n})$ time.

Theorem 15. A maximum open packing in a split graph $G(C \cup I, E)$ with $1 \le \deg_G(v) \le 2$ for every $v \in I$ can be found in $O(\max\{n+m, n^{1.5}\})$ time.

Proof. Let $G(C \cup I, E)$ be a split graph with $deg_G(v) \in \{1, 2\}$ for every $v \in I$. Also, let F be a maximum matching of the graph G' constructed in Construction 5, i.e., $\alpha'(G') = |F|$. Then, we find the open packing number of G using F in four parts.

Part 1: If $\alpha'(G') \ge 3$, then $S_F = \{v \in I : e_v \in F\}$ is a maximum open packing in G.

Note that S_F is an open packing in G by Case 2 in the proof of Guarantee of Construction 5. So, $\rho^o(G) \ge |S_F| = \alpha'(G) \ge 3$. Hence, by Claim 9, every maximum open packing in G is a subset of I. Let S be a maximum open packing in G. Then, by Case 1 in the proof of Guarantee of Construction 5, $F_S = \{e_v \in E' : v \in S\}$ is a matching in G'. So, $\alpha'(G') \ge |F_S| = |S| = \rho^o(G) \ge |S_F| = \alpha'(G')$. Hence, $\rho^o(G) = \alpha'(G')$, and S_F is a maximum matching in G'.

Part 2: If $\alpha'(G') = 2$, then $S_F = \{v \in I : e_v \in F\}$ is a maximum open packing in G.

If $\rho^{o}(G) \geq 3$, then by Claim 9 and Construction 5, $\alpha'(G') \geq 3$, a contradiction to the assumption that $\alpha'(G') = 2$. So, $\rho^{o}(G) \leq 2$. By Case 2 in the proof of Guarantee of Construction 5, S_F is an open packing of size two in G. Hence, $\rho^{o}(G) = 2$.

Part 3: If $\alpha'(G') = 1$ and there exists a vertex of degree one in G, then $\rho^o(G) = 2$.

Since $\alpha'(G') = 1$, it is evident, by Claim 9 and Construction 5, that $\rho^o(G) \leq 2$. Let $v \in V(G)$ such that $deg_G(v) = 1$. Then, $N_G[v]$ is an open packing of size two in G. Hence, $\rho^o(G) = 2$.

Part 4: If $\alpha'(G') = 1$ and every vertex in G is of degree at least two, then $\rho^o(G) = 1$.

On the contrary, assume that G has an open packing S of cardinality at least two. Since $\alpha'(G') = 1$, there does not exist an open packing S in G such that $|S| \ge 2$ and $S \subseteq I$ by Construction 5. So, $S \cap C \ne \emptyset$. Let $u \in S \cap C$. Then, by Claim 9, $S \subseteq N_G[u]$. Let $v \in S \setminus \{u\}$. Then, by the assumption on Part 4 that every vertex in G is of degree at least two, for any choice of v in $V(G) \setminus \{u\} = (C \cup I) \setminus \{u\}, u$ and v must have a common neighbour in G, a contradiction to S being an open packing in G with $u, v \in S$. So, $\rho^o(G) = 1$.

Parts 1-4 complete the arguments for finding a maximum open packing in G. Next, we look at the complexity of computing it. Note that

- (a) the graph G' can be constructed in O(n+m) time,
- (b) a maximum matching in G' can be found in $O(|E(G')|\sqrt{V(G')}) = O(n^{1.5})$ time using Lemma 13, and
- (c) degree of every vertex in G can be found in O(n+m) time.

By (a)-(c) and Parts 1-4, a maximum open packing in G can be found in $O(\max\{n+m, n^{1.5}\})$ time.

5 Conclusion

In this article, we have completed the study on the complexity (P vs NPC) of OPEN PACKING on H-free graphs for every graph H with at least three vertices by proving that OPEN PACKING is (i) NP-complete on $K_{1,3}$ -free graphs and (ii) polynomial time solvable on subclasses of $(P_4 \cup rK_1)$ free graphs for every $r \geq 1$. Also, we proved that for a connected $(P_4 \cup rK_1)$ -free graph G, $\gamma_t(G) \leq 2r+2$ and $\rho^o(G) \leq 2r+1$. Since $\rho^o(G) \leq \gamma_t(G)$, it would be interesting to study (i) the bound on the difference between $\gamma_t(G)$ and $\rho^o(G)$ and (ii) the lower bounds for open packing number. Further, we proved that OPEN PACKING is (i) NP-complete on $K_{1,4}$ -free split graphs and (ii) polynomial time solvable on $K_{1,3}$ -free split graphs. We also showed that OPEN PACKING is (i) NP-complete on I_r -split graphs for $r \geq 3$ and (ii) polynomial time solvable on I_r -split graphs for $r \leq 2$. Also, we completed the complexity (P vs NPC) of TOTAL DOMINATING SET on I_r -split graphs by proving that it is NP-complete when $r \geq 2$ and polynomial time solvable when r = 1. Note that I_r -split graphs is a split graph with regularity (equal number of edges incident) on the vertices of the independent set part of the clique-independence partition of split graphs. So, it would be interesting to know the complexity (P vs NPC) of TOTAL DOMINATING SET and OPEN PACKING in split graphs with regularity imposed on (i) the clique part and (ii) both the clique and the independent set parts of the clique-independence partition of split graphs, where the regularity on clique part shall be different from the regularity on independent set part.

References

- Corneil, D.G., Perl, Y.: Clustering and domination in perfect graphs. Discrete Applied Mathematics 9(1), 27–39 (1984). doi:10.1016/0166-218X(84)90088-X
- [2] Cygan, M., Fomin, F.V., Kowalik, L., Lokshtanov, D., Marx, D., Pilipczuk, M., Pilipczuk, M., Saurabh, S.: Parameterized Algorithms. Springer Publishing Company, Incorporated, 1st edn. (2015)
- [3] Damaschke, P., Muller, H., Kratsch, D.: Domination in convex and chordal bipartite graphs. Information Processing Letters 31, 231–236 (1990). doi:10.1016/0020-0190(90)90147-P
- [4] Downey, R.G., Fellows, M.R.: Fixed-parameter tractability and completeness II: On completeness for W[1]. Theoretical Computer Science 141(1), 109–131 (1995). doi:10.1016/0304-3975(94)00097-3
- [5] Downey, R.G., Fellows, M.R.: Fundamentals of Parameterized Complexity. Springer London (2016). doi:10.1007/978-1-4471-5559-1
- [6] Håstard, J.: Clique is hard to approximate within $n^{1-\epsilon}$. Acta Mathematica **182**, 105–142 (1999). doi:10.1007/BF02392825
- [7] Haynes, T.W., Hedetniemi, S., Slater, P.: Fundamentals of Domination in Graphs. CRC Press, 1 edn. (1998). doi:10.1201/9781482246582
- [8] Henning, M.A.: Packing in trees. Discrete Mathematics 186(1), 145–155 (1998). doi:10.1016/S0012-365X(97)00228-8
- [9] Henning, M.A., Slater, P.J.: Open packing in graphs. Journal of Combinatorial Mathematics and Combinatorial Computing 29, 3–16 (1999)

- [10] Henning, M.A., Yeo, A.: Total Domination in Graphs. Springer New York (2015). doi:10.1007/978-1-4614-6525-6
- [11] ISGCI: On relation between 3K₁-free graphs and AT-free graphs. https://www.graphclasses.org/classes/AUTO_399.html
- [12] Karp, R.M.: Reducibility among combinatorial problems. In: Miller, R.E., Thatcher, J.W., Bohlinger, J.D. (eds.) Complexity of Computer Computations. pp. 85–103. Springer US (1972). doi:10.1007/978-1-4684-2001-2_9
- [13] Kratsch, D.: Domination and total domination on asteroidal triple-free graphs. Discrete Applied Mathematics 99(1), 111–123 (2000). doi:10.1016/S0166-218X(99)00128-6
- [14] McRae, A.A.: Generalizing NP-completeness proofs for bipartite graphs and chordal graphs. Ph.D. thesis, Clemson University, USA (1995)
- [15] Micali, S., Vazirani, V.V.: An O(√|V||E|) algorithm for finding maximum matching in general graphs. In: 21st Annual Symposium on Foundations of Computer Science (sfcs 1980). pp. 17– 27 (1980). doi:10.1109/SFCS.1980.12
- [16] Pfaff, J., Laskar, R.C., Hedetniemi, S.T.: NP-completeness of total and connected domination and irredundance for bipartite graphs. Technical Report 428, Department of Mathematical Sciences, Clemson University (1983)
- [17] Raja Chandrasekar, K., Saravanakumar, S.: Open packing number for some classes of perfect graphs. Ural Mathematical Journal 6, 38–43 (2020). doi:10.15826/umj.2020.2.004
- [18] Rall, D.F.: Total domination in categorical products of graphs. Discussiones Mathematicae Graph Theory 25, 35–44 (2005). doi:10.7151/dmgt.1257
- [19] Ramos, I., Santos, V.F., Szwarcfiter, J.L.: Complexity aspects of the computation of the rank of a graph. Discrete Mathematics & Theoretical Computer Science 16 (2014). doi:10.46298/dmtcs.2075
- [20] Renjith, P., Sadagopan, N.: The Steiner tree in $K_{1,r}$ -free split graphs—a dichotomy. Discrete Applied Mathematics **280**, 246–255 (2020). doi:10.1016/j.dam.2018.05.050
- [21] Shalu, M.A., Kirubakaran, V.K.: Total domination number and its lower bound in some subclasses of bipartite graphs. Manuscript-UnderReview (2023)
- [22] Shalu, M.A., Vijayakumar, S., Sandhya, T.P.: A lower bound of the cd-chromatic number and its complexity. In: Gaur, D., Narayanaswamy, N. (eds.) Algorithms and Discrete Applied Mathematics. pp. 344–355. Springer International Publishing (2017). doi:10.1007/978-3-319-53007-9_30
- [23] West, D.B.: Introduction to Graph Theory, Second Edition. Pearson (2018)
- [24] White, K., Farber, M., Pulleyblank, W.: Steiner trees, connected domination and strongly chordal graphs. Networks 15(1), 109–124 (1985). doi:10.1002/net.3230150109

A Open Packing

In this section, we give an algorithm to test whether a subset S of a graph G is an open packing in G or not.

Algorithm 1: Open Packing Testing Algorithm for Graphs
Input : A Graph $G(V, E)$ and a vertex subset S of G. Output: S is/isn't an open packing in G
Initialization: $U = S$, $W = \emptyset$, an empty array Q , $j = 1$ and every vertex in $V(G)$ is unmarked.
while $U \neq \emptyset$ do
Choose a vertex $u \in U$
Set $W = N_G(u)$
while $W \neq \emptyset$ do
Choose a vertex $v \in W$
Set $Q[j] = v$ \\ Q is used only in the analysis of the algorithm
Set $j = j + 1$
if v is marked then
return 'S is not an open packing in G ' and terminate
else
$\operatorname{Mark} v$
Set $W = W \setminus \{v\}$
end
end
Set $U = U \setminus \{u\}$
end
Return 'S is an open packing in G ' and terminate

Observation 4. Given a graph G and a vertex subset S of G, Algorithm 1 determines whether S is an open packing in G or not in O(n) time.

Lemma 4 follows from Observation 4.

Proof of Observation 4. We start by giving some simple clarifications to get a clear understanding of our proof. We use the adjacency list of the graph G as the data structure. We say that a vertex $u \in V(G)$ is visited by the algorithm if the vertex is chosen by at least one of the while loops during some stage of the algorithm. We call the while loop that is performed based on whether U is empty or not the external while loop. We refer to the iterations performed by the external while loop as the iterations of the algorithm. The while loops that are performed based on the set W are called the internal while loops.

We begin with the proof for the termination of the algorithm. Note that if the algorithm visits some vertex that is already marked, then the algorithm will terminate. Otherwise, the algorithm visits no marked vertex in any iteration of the algorithm. However, since W is a finite set for every iteration of the external while loop, the internal while loop iterates only a finite number of times. Hence, every iteration of the external while loop will terminate. Also, since the size of U is finite, the number of iterations performed by the algorithm is finite, and hence, the algorithm would terminate once U becomes empty.

Next, we prove the correctness of Algorithm 1. Let $S = \{u_1, u_2, \ldots, u_k\}$. W.L.O.G., assume that the algorithm chooses the vertices in S in the increasing order of their suffices. We prove the correctness of the algorithm in two cases based on whether the input vertex subset S is an open packing in G or not.

Case 1: The input vertex subset S is an open packing in G. Since every vertex in V(G) is unmarked during the initialization process, no vertex in $N_G(u_1)$ might be marked while examining them. For $2 \leq i \leq k$, the vertices of G that are marked during the previous iterations of the algorithm are in $\bigcup_{s=1}^{i-1} N_G(u_s)$. Since S is an open packing, $N_G(u_i) \cap N_G(u_s) = \emptyset$ for every s < i. These statements imply that no vertex in $N_G(u_i)$ is marked before the i^{th} iteration. So, the algorithm will never encounter a marked vertex, and hence the algorithm will return 'S is an open packing in G'. Thus, the algorithm correctly determines the yes instances of the open packing sets of a graph.

Case 2: The input vertex subset S is not an open packing in G.

Since S is not an open packing in G, there exist i, s such that $1 \leq i < s \leq k$ and $N_G(u_i) \cap N_G(u_s) \neq \emptyset$. W.L.O.G., assume that u_s is the least indexed vertex such that there exists a vertex u_i with i < s and $N_G(u_i) \cap N_G(u_s) \neq \emptyset$. Note that $S' = \{u_1, \ldots, u_{s-1}\}$ is an open packing in G by the choice of s. Hence, by the arguments in Case 1, the algorithm does not end before examining $N_G(u_s)$. Let $N_G(u_i) \cap N_G(u_s) = \{w_1, w_2, \ldots, w_p\}$ for some $p \leq n$. Note that every vertex in $N_G(u_i)$ is marked during i^{th} iteration of the external while loop. So, the algorithm will encounter a marked vertex (w_t for some $1 \leq t \leq p$) while exploring $N_G(u_s)$ during the s^{th} iteration. Hence, the algorithm will return 'S is not an open packing in G'. Thus, the algorithm correctly determines the no instances of the open packing sets of a graph.

Thus, Cases 1 and 2 prove that Algorithm 1 correctly determines whether the input vertex subset S is an open packing in G or not.

Next, we prove that the algorithm runs in O(n) time. To prove this, we show that the size of the list of vertices visited by the algorithm is at most 2n + 1. Let $Q = (v_1, v_2, \ldots, v_r)$ for some $r \in \mathbb{N}$. Then, it is clear that Q is the list of vertices visited by the internal while loop (i.e., every vertex v_i (for $1 \le j \le r$) is visited as a neighbor of u_i for some $i \in \{1, 2, \ldots, k\}$) before the termination of the algorithm. We claim that $v_j \neq v_\ell$ for every $j, \ell \in \{1, 2, \ldots, r-1\}$ and $j \neq \ell$. On the contrary, assume that there exist $j, \ell \in \{1, 2, ..., r-1\}$ such that $j \neq \ell$ but $v_j = v_\ell = u$ for some $u \in V(G)$. W.L.O.G., assume that ℓ is the least integer in $\{1, 2, \ldots, r-1\}$ such that there exists $j < \ell$ with $v_j = v_\ell$. Then, the vertex $u(=v_j)$ is marked by the algorithm when the internal while loop visits the j^{th} vertex in Q. Also, by the assumption on ℓ , every vertex in $(v_1, \ldots, v_{\ell-1})$ is distinct, and hence the algorithm does not terminate without visiting ℓ^{th} vertex of Q. But, when the ℓ^{th} vertex $u = v_{\ell}$ of the array Q is visited by the internal while loop, the algorithm would terminate as the vertex is already marked while visiting the j^{th} vertex (u) in Q, a contradiction to the assumption that $r > \ell$ and the internal while loop visits r vertices in Q before the algorithm terminates. So, $v_j \neq v_\ell$ for every $j, \ell \in \{1, 2, \ldots, r-1\}$ and $j \neq \ell$. This implies that the cardinality of the set $V' = \{v_1, v_2, \dots, v_{r-1}\}$ is r-1. Since $V' \subseteq V(G), r-1 = |V'| \le n$. Hence, $r \le n+1$, i.e., the size of the list of vertices visited by the internal while loop is at most n + 1. Note that for $Q = (v_1, v_2, \ldots, v_r)$ and $S = \{u_1, u_2, \ldots, u_k\}$, the list of vertices visited by the algorithm is a permutation of some subarray of $(v_1, v_2, \ldots, v_r, u_1, u_2, \ldots, u_k)$. We say subarray because the algorithm may terminate before visiting some vertices in S. Hence, the size of the list of vertices visited by the algorithm is at most $|S| + |Q| \le 2n + 1$. Also, note that the edges of G are visited only during the internal while loop (for neighborhood testing). If $E' = (e_1, e_2, \ldots, e_q)$ is the list of edges visited by the algorithm with the suffices denoting the order in which the edges are visited, then by the construction of Q, it is clear that for every e_i $(1 \le i \le q)$ visited by the algorithm, exactly one vertex (v_i) in Q was visited. Since the size of Q is at most $n+1, q \leq n+1$. Hence, the algorithm runs in 3n + 2 = O(n) time.

B Proofs related to Section 3.2

Claim 11: The graph G_r defined in Remark 3 is a $(P_4 \cup rK_1)$ -free graph.

Recall that $V(G_r) = (\bigcup_{i=1}^r \{x_i, y_i, z_i\}) \cup \{u, v\}$ and $E(G_r) = (\bigcup_{i=1}^r \{x_i y_i, y_i z_i, z_i u\}) \cup \{uv\}$. On the contrary to claim, assume that there exists a $(P_4 \cup rK_1)$, say H, in G_r . Let $\{a, b, c, d\} \in V(H)$ such that $G_r[a, b, c, d] \cong P_4$. Note that every P_4 in G_r consists of the vertex u. So, $u \in \{a, b, c, d\}$. Similarly, observe that every P_4 in G consists of exactly one vertex labelled y_i for some $i = 1, 2, \ldots, r$. So, $y_i \in \{a, b, c, d\}$ for some $i = 1, 2, \ldots, r$. W.L.O.G., assume that i = 1. Then, $\{a, b, c, d\} \in \{\{v, u, z_1, y_1\}, \{z_j, u, z_1, y_1\}, \{u, z_1, y_1, x_1\}\}$. Since $H \cong (P_4 \cup rK_1)$ and $\{a, b, c, d\} \subseteq V(H)$ induces the P_4 , $G_r[V(H) \setminus \{a, b, c, d\}] \cong rK_1$ and no vertex in $(V(H) \setminus \{a, b, c, d\}) \subseteq \{x_2, y_2, \ldots, x_r, y_r\}$. But the independence number of $G_r[\{x_2, y_2, \ldots, x_r, y_r\}]$ is (r-1), a contradiction to $(V(H) \setminus \{a, b, c, d\}) \subseteq \{x_2, y_2, \ldots, x_r, y_r\}$ and $(V(H) \setminus \{a, b, c, d\}) \cong rK_1$, i.e., the independence number of an induced subgraph of $G_r[\{x_2, y_2, \ldots, x_r, y_r\}]$ is at least r.

Lemma 10 (Restated). For $r \ge 1$, if G is a connected $(P_3 \cup rK_1)$ -free graph, then $\rho^o(G) \le 2r$.

Proof. We use induction on r to prove this lemma as well. We begin with the proof that $\rho^o(G) \leq 2$

for every connected $(P_3 \cup K_1)$ -free graph G. Note that, by Lemma 9, $\rho^o(G) \leq \gamma_t(G) \leq 3$, for every connected $(P_3 \cup K_1)$ -free graph G. So, to prove the claim, it is enough to disprove the case $\rho^{o}(G) = 3$. On the contrary, assume that there exists a $(P_3 \cup K_1)$ -free graph G with $\rho^{o}(G) = 3$. Since the open packing number of a connected P_3 -free graph is at most two by Remark 4 and $\rho^{o}(G) = 3$, G must contain P_3 as an induced subgraph. Let $D = \{x_1, x_2, x_3\}$ be a subset of V(G) that induces a P_3 in G with $x_1x_2, x_2x_3 \in E(G)$. Then, by the arguments in the proof of Lemma 9,

D is a total dominating set in G, i.e., $V(G) = \bigcup_{i=1}^{3} N(x_i)$. Then, for every open packing S in G,

 $S = \bigcup_{i=1}^{3} (S \cap N(x_i)).$ We know that $|S \cap N(x_i)| \le 1$ for i = 1, 2, 3. This, together with the fact that $S = \bigcup_{i=1}^{3} (S \cap N(x_i)),$ implies that for an open packing S in G of cardinality three, $|S \cap N(x_i)| = 1$

for i = 1, 2, 3. Let $S \cap N(x_i) = \{z_i\}$. Then, $S = \{z_1, z_2, z_3\}$. So, $z_i \neq z_j$ for $i, j \in \{1, 2, 3\}$ and $i \neq j$ (else $|S| \leq 2$). Note that $S \cap N(x_1) = \{z_1\}$. So, $x_1z_2, x_1z_3 \notin E(G)$. Using similar arguments, we can show that $x_i z_j \notin E(G)$ for $i, j \in \{1, 2, 3\}$ and $i \neq j$. Further, since $x_2 x_3 \in E(G)$ and $x_3z_1 \notin E(G), z_1 \neq x_2$. Similarly, $z_3 \neq x_2$. Also, note that if $z_1z_3 \in E(G)$, then $z_1z_2, z_2z_3 \notin E(G)$ since G[S] is $\{P_3, K_3\}$ -free by Observation 3. This, together with the fact that $z_3x_3 \in E(G)$, implies that $z_2 \neq x_3$. Then, $\{z_1, z_3, x_3\} \cup \{z_2\}$ induces a $(P_3 \cup K_1)$ in G, a contradiction. So, $z_1z_3 \notin E(G)$. But, then $\{z_1, x_1, x_2\} \cup \{z_3\}$ induces a $(P_3 \cup K_1)$, a contradiction. So, $\rho^o(G) \leq 2$. Next, for every $r \ge 2$, we show that $\rho^o(G) \le 2r$ for every connected $(P_3 \cup rK_1)$ -free graph G. Induction Assumption: Assume that for $r \geq 2$, $\rho^o(G') \leq 2(r-1)$ for every connected $(P_3 \cup (r-1))$ 1) K_1)-free graph G'.

Let G be a connected $(P_3 \cup rK_1)$ -free graph. If G does not contain any vertex subset that induces a $(P_3 \cup (r-1)K_1)$, then G is $(P_3 \cup (r-1)K_1)$ -free, and so by induction assumption $\rho^{o}(G) \leq 2(r-1) < 2r$. So, assume that G has a vertex subset $D = \{x_1, x_2, x_3, y_1, y_2, \dots, y_{r-1}\}$ that induces a $(P_3 \cup (r-1)K_1)$ in G with $x_i x_{i+1} \in E(G)$ for i = 1, 2. Since G is $(P_3 \cup rK_1)$ -free, every vertex in $V(G) \setminus D$ is adjacent to some vertex in D, i.e., D is a dominating set in G. So, $V(G) = N[x_1] \cup N[x_2] \cup N[x_3] \cup N[y_1] \cup N[y_2] \cup \ldots \cup N[y_{r-1}]$. This implies that, for any open packing S in G, $S = \bigcup_{u \in D} (S \cap N[u])$. Note that $|S \cap N[u]| \le |S \cap N(u)| + |S \cap \{u\}| \le 2$. Also, since

$$x_1, x_2, x_3 \in N(x_1) \cup N(x_2), \bigcup_{i=1}^3 N[x_i] = \bigcup_{i=1}^3 N(x_i). \text{ So, } |S \cap (\bigcup_{i=1}^3 N[x_i])| = |S \cap (\bigcup_{i=1}^3 N(x_i))| \le 3.$$

Therefore, $|S| = |S \cap V(G)| = |S \cap (N[x_1] \cup N[x_2] \cup N[x_3] \cup N[y_1] \cup N[y_2] \cup \ldots \cup N[y_{r-1}])| \le 3 + 2(r-1) = 2r+1.$ Thus, to prove the claim, it is enough to contradict the possibility of $|S| = 2r+1.$ Assume that there exists an open packing S of size $2r+1$ in G. Then, by the above

arguments, we can conclude that $|S \cap N[y_j]| = 2$ for j = 1, 2..., r-1 and $|S \cap (\bigcup_{i=1}^{3} N(x_i))| = 3$. Since $|S \cap N(y_j)| \le 1$ and $|S \cap N[y_j]| = 2$, $y_j \in S$ and $S \cap N(y_j) \ne \emptyset$. Let $S \cap N[y_j] = \{w_j, y_j\}$ for j = 1, 2, ..., r-1 and let $S \cap (\bigcup_{i=1}^{3} N(x_i)) = \{z_1, z_2, z_3\}$ with $x_i z_i \in E(G)$ (then, since $|S \cap N(x_i)| \le 1$, we have $S \cap N(x_i) = \{z_i\}$ for every i = 1, 2, 3). So, $S = (\bigcup_{i=1}^{3} \{z_i\}) \cup (\bigcup_{j=1}^{r-1} \{y_j\}) \cup (\bigcup_{j=1}^{r-1} \{w_j\})$.

This, together with the fact that |S| = 2r + 1, implies that every vertex in the (2r + 1)-tuple $(z_1, z_2, z_3, y_1, y_2, \dots, y_{r-1}, w_1, w_2, \dots, w_{r-1})$ is distinct. Further, since (i) $S \cap N(x_i) = \{z_i\}$ and (ii) $S \cap N(y_i) = \{w_i\}$, we have $z_s u \notin E(G)$ for $s \in \{1, 2, 3\}$ and $u \in D \setminus \{x_s\}$. We complete our claim in two cases based on $z_1 z_3 \in E(G)$ or not.

Case 1: $z_1z_3 \in E(G)$.

Then, by Observation 3, $z_1z_2, z_2z_3 \notin E(G)$. This implies that $z_2 \neq x_3$. So, $\{x_1, z_1, z_3\} \cup$ $\{z_2, y_1, y_2, \ldots, y_{r-1}\}$ induces a $(P_3 \cup rK_1)$ in G, a contradiction. Case 2: $z_1z_3 \notin E(G)$.

Then, $\{z_1, x_1, x_2\} \cup \{z_3, y_1, y_2, \dots, y_{r-1}\}$ induces a $(P_3 \cup rK_1)$ in *G*, a contradiction. In both cases, we arrive at a contradiction. Therefore, $\rho^o(G) \leq 2r$.

\mathbf{C} Proof Related to Section 4

Claim 7 (Restated): A set $S \subseteq V(G)$ is an independent set in the input graph G of Construction 2 if and only if S is an open packing in the output graph G'.

Suppose that $S \subseteq V(G)$ is an independent set in G. On the contrary to Claim 7, assume that S is not an open packing in G'. Then, there exist vertices $u, u' \in S$ and $w \in V(G')$ such that $w \in N_{G'}(u) \cap N_{G'}(u')$. Since $u, u' \in V(G) \subseteq I$, $w \in C = E(G) \cup \{y, z\}$. By the construction of G', $uy, uz \notin E(G')$ for any $u \in V(G)$. Thus, w = e for some $e \in E(G)$. Then, by the construction of G', $e = uu' \in E(G)$, a contradiction to S being an independent set in G.

Conversely, suppose that $S \subseteq V(G)$ is an open packing in G'. If S is not an independent set in G, then there exist $u, u' \in S$ and an $e \in E(G)$ such that uu' = e. Then, by the construction of G', $e \in N_{G'}(u) \cap N_{G'}(u') \neq \emptyset$, a contradiction.

Theorem 16. MAX-OPEN PACKING is hard to approximate within a factor of $N^{(\frac{1}{2}-\epsilon)}$ for any $\epsilon > 0$ in $K_{1,4}$ -free split graphs unless P = NP, where N is the number of vertices in $K_{1,4}$ -free split graphs.

Proof. On the contrary, assume that MAX-OPEN PACKING admits a $N^{\left(\frac{1}{2}-\epsilon\right)}$ -factor approximation algorithm for some $\epsilon > 0$ in $K_{1,4}$ -free split graphs. Then, an open packing S of the graph G'constructed in Construction 2 with $|S| > \frac{\rho^o(G)}{|V(G')|^{\left(\frac{1}{2}-\epsilon\right)}}$ can be found in polynomial time. Hence, an independent set U of the input graph G in Construction 2 with |U| = |S| - 1 can be found in polynomial time. Note that $N = |V(G')| = n + m + 3 \le 2n + \frac{n^2}{2} \le n^2$ for $n \ge 4$. Then,

$$\begin{split} |S| &> \frac{\rho^o(G')}{N^{\left(\frac{1}{2} - \epsilon\right)}} \\ |S| &> \frac{\rho^o(G')}{(n + m + 3)^{\left(\frac{1}{2} - \epsilon\right)}} \\ 2(|S| - 1) &> \frac{\rho^o(G')}{(n^2)^{\left(\frac{1}{2} - \epsilon\right)}} \quad \text{for } |S| \ge 2 \text{ and } n \ge 4 \\ |S| - 1 &> \frac{1}{2} \cdot \frac{\alpha(G) + 1}{n^{1 - 2\epsilon}} \\ |U| &> \frac{1}{2} \cdot \frac{\alpha(G)}{n^{1 - 2\epsilon}} \end{split}$$

Since $\rho^o(G') = \alpha(G) + 1 > \alpha(G)$ by Guarantee of Construction 2. Fix n_0 such that $n_0^{\epsilon} > 2 \implies \frac{1}{n_0^{\epsilon}} < \frac{1}{2}$. Then, for $n \ge \max\{n_0, 4\}$, we have

$$|U| > \frac{\alpha(G)}{n^{\epsilon} \cdot n^{1-2\epsilon}}$$
$$|U| > \frac{\alpha(G)}{n^{1-\epsilon}}$$

The above inequality implies that MAX-INDEPENDENT SET has a $(n^{1-\epsilon})$ -factor approximation algorithm, which contradicts Theorem 4.