

NON-STATIONARY GAUSSIAN RANDOM FIELDS ON HYPERSURFACES: SAMPLING AND STRONG ERROR ANALYSIS

ERIK JANSSON, ANNIKA LANG, AND MIKE PEREIRA

ABSTRACT. A flexible model for non-stationary Gaussian random fields on hypersurfaces is introduced. The class of random fields on curves and surfaces is characterized by a power spectral density of a second order elliptic differential operator. Sampling is done by a Galerkin–Chebyshev approximation based on the surface finite element method and Chebyshev polynomials. Strong error bounds are shown with convergence rates depending on the smoothness of the approximated random field.

1. INTRODUCTION

Random fields are powerful tools for modeling spatially dependent data. They have found uses in a wide range of applications, for instance in geostatistics, cosmological data analysis, climate modeling, and biomedical imaging (Marinucci and Peccati, 2011; Farag, 2014). One challenge in the modeling of spatial data is non-stationary behavior, i.e., different behaviors in different parts of the domain. Another challenge is that the domain may be a non-Euclidean space, for instance, a surface such as the sphere or on the cortical surface of the brain. In this paper, we present a surface finite element-based method to sample a flexible class of non-stationary random fields on curves and surfaces and show its strong convergence. The method, building on the foundational work for stationary fields introduced in Lang and Pereira (2023), is an extension of the stochastic partial differential equation (SPDE) approach pioneered by Whittle (1963) and popularized by Lindgren et al. (2011). The idea behind our method is to color white noise by applying a function of an elliptic differential operator \mathcal{L} . Formally, we study Gaussian random fields on curves and surfaces of the form

$$(1) \quad \mathcal{Z} = \gamma(\mathcal{L})\mathcal{W},$$

where γ is a function, called the *power spectral density*, \mathcal{L} is an elliptic differential operator and \mathcal{W} denotes white noise. By letting the coefficients of the differential operator vary over the domain, we can obtain local, non-stationary behaviors. If $1/\gamma$ is well-defined over \mathbb{R}_+ , one may formally view \mathcal{Z} as the solution to the stochastic partial differential equation

$$(1/\gamma)(\mathcal{L})\mathcal{Z} = \mathcal{W}.$$

2020 *Mathematics Subject Classification.* 60G60, 60H35, 60H15,65C30, 60G15, 58J05, 41A10, 65N30,65M60.

Key words and phrases. Gaussian random fields, non-stationary random fields, stochastic partial differential equations, surface finite element method, Chebyshev approximation, Gaussian processes.

Acknowledgment: The authors thank Julie Rowlett and Iosif Polterovich for helpful discussions. This work was supported in part by the European Union (ERC, StochMan, 101088589), by the Mines Paris - PSL / INRAE “Geolearning” chair, by the Swedish Research Council (VR) through grant no. 2020-04170, by the Wallenberg AI, Autonomous Systems and Software Program (WASP) funded by the Knut and Alice Wallenberg Foundation, and by the Chalmers AI Research Centre (CHAIR) .




FIGURE 1. Examples of random field samples generated with our method.

For instance, consider the three examples depicted in Figure 1. To generate random field samples, there are two components of \mathcal{L} that we can vary: the diffusion matrix and the potential. In Figure 1(a), we use a diffusion matrix to give the field preferred directions, more specifically it elongates field in the northwest-southeast direction. The potential is large over the continents and small over the oceans, effectively “turning off” the random field over land. In Figure 1(b), the potential is small in the front of the brain and large elsewhere, so that the field is only large in the front of the brain. Finally, Figure 1(c) shows the method used to generate a non-stationary random field in the one-dimensional case. To illustrate the value of the field at a point, we move it in the normal direction for a distance proportional to the value of the field. In all three cases, we see that the field behaves locally varying over the domain. With the suggested model, we can achieve preferred directions, local activation, and local deactivation.

The computational method we use to solve Equation (1), i.e., sample the random fields, is based on the surface finite element method (SFEM), a computational method pioneered by Dziuk (1988); Dziuk and Elliott (2013) and that has been used in the context of the generation of random fields in for instance Bonito et al. (2024); Jansson et al. (2022); Lang and Pereira (2023). Our main mathematical contribution is a strong convergence result. Using a functional calculus approach to the finite element discretization error, we obtain a strong rate of convergence of order $\mathcal{O}(K_\alpha(h)h^{\min\{\alpha-d/4;2\}})$, where $d = 2$ for surfaces and $d = 1$ for curves, and $K_\alpha(h)$ is a dimension-dependent logarithmic factor.

The SPDE approach to random fields and their approximation have been studied previously for both surfaces and Euclidean geometries, examples include Bonito et al. (2024); Borovitskiy et al. (2020); Bolin et al. (2020); Cox and Kirchner (2020); Jansson et al. (2022); Lang and Pereira (2023); Lindgren et al. (2011, 2022). However, to the best of our knowledge, we are the first to present a strong error analysis for the non-stationary case on hypersurfaces. Moreover, we achieve this without requiring an explicit approximation of the eigenfunctions of the elliptic operator, and contrary to for instance Borovitskiy et al. (2020), computing the eigenfunctions. In fact, the computation of eigenfunctions, with theoretical guarantees, is a notoriously difficult problem, see e.g., Boffi (2010). Our method circumvents this issue by introducing a Chebyshev approximation.

Our main contribution is a powerful, efficient, and flexible tool for the modeling and sampling of non-stationary random fields on curves and surfaces with proven accuracy. In particular, using tools from complex analysis and operator theory, we derive strong error bounds for approximations of arbitrary sufficiently smooth transformations of elliptic operators, where we do not require assumptions on the approximability of individual eigenfunctions.

The paper is structured as follows: In Section 2, we introduce the relevant deterministic framework. We provide the necessary background on geometry and functional analysis in Section 2.1. This is followed by a description of the main computational tool, surface finite elements in Section 2.2. Finally, Section 2.3 provides the relevant error estimates in the deterministic setting. In Section 3, we collect all material in the stochastic setting. We introduce first the class of considered random fields in Section 3.1 and their Galerkin–Chebyshev approximation in Section 3.2. The proof of its strong convergence is split into the SFEM error in Section 3.3 and the Chebyshev approximation error Section 3.4. For numerical confirmation of our theoretical results, we refer to our earlier work Lang and Pereira (2023) and the results presented in Bonito et al. (2024). The source code used to generate the figures is available at this address: <https://github.com/mike-pereira/SFEMsim>.

2. DETERMINISTIC THEORY: GEOMETRY, FUNCTIONAL ANALYSIS AND FINITE ELEMENTS

Before we are able to approximate random fields on hypersurfaces, we need to introduce and partially extend the existing literature on surface finite element approximations due to so far unconsidered error bounds required in our stochastic setting. We introduce the functional analytic setting in Section 2.1, discuss surface finite element methods in Section 2.2 and show error bounds in the deterministic setting in Section 2.3.

2.1. Geometric and functional analytic setting. Let $\mathcal{M} \subset \mathbb{R}^{d+1}$ be a d -dimensional ($d \leq 2$) compact oriented smooth hypersurface ($k \geq 2$) without boundary, i.e., for any $x_0 \in \mathcal{M}$, there exists an open set $U_{x_0} \subset \mathbb{R}^{d+1}$ containing x_0 and a function $\phi_{x_0} \in C^\infty(U_{x_0})$ such that $\nabla\phi_{x_0} \neq 0$ on $\mathcal{M} \cap U_{x_0}$ and

$$\mathcal{M} \cap U_{x_0} = \{x \in U_{x_0}, \phi_{x_0}(x) = 0\}.$$

The tangent space of \mathcal{M} at $x \in \mathcal{M}$ is the d -dimensional subspace of \mathbb{R}^{d+1} given by $T_x\mathcal{M} = [\nabla\phi_x]^\perp$ (where ∇ denotes the usual gradient of functions of $C^1(\mathbb{R}^{d+1})$ and \perp denotes the orthogonal complement in \mathbb{R}^{d+1} with respect to the standard Euclidean inner product.). Since \mathcal{M} is oriented, there exists a smooth map $\nu : \mathcal{M} \rightarrow \mathbb{R}^{d+1}$ assigning to each point $x \in \mathcal{M}$ a unit vector $\nu(x) = \pm \nabla\phi_x / \|\nabla\phi_x\|$ perpendicular to the tangent space $T_x\mathcal{M}$. Our hypersurface \mathcal{M} is a Riemannian manifold equipped with the metric g that is the pullback of the Euclidean metric on \mathbb{R}^{d+1} . For instance, if $\mathcal{M} = \mathbb{S}^2$, this results in the standard round metric.

Let $\nabla_{\mathcal{M}}$ be the gradient operator acting on differentiable functions of \mathcal{M} , and let $\Delta_{\mathcal{M}}$ denote the Laplace–Beltrami operator on (\mathcal{M}, g) . We denote by dA the surface measure on \mathcal{M} , and by $L^2(\mathcal{M})$ the Hilbert space of dA -measurable square integrable complex-valued functions, equipped with the inner product $(\cdot, \cdot)_{L^2(\mathcal{M})}$ defined by

$$(u, v)_{L^2(\mathcal{M})} = \int_{\mathcal{M}} u\bar{v} \, dA, \quad u, v \in L^2(\mathcal{M}).$$

The *Sobolev spaces* with smoothness index $s \in \mathbb{R}^+$ are then defined via Bessel potentials by

$$H^s(\mathcal{M}) = (I - \Delta_{\mathcal{M}})^{-s/2} L^2(\mathcal{M}),$$

with corresponding norm $\|\cdot\|_{H^s(\mathcal{M})} = \|(I - \Delta_{\mathcal{M}})^{s/2} \cdot\|_{L^2(\mathcal{M})}$. For $s < 0$, $H^s(\mathcal{M})$ is defined as the space of distributions generated by

$$(2) \quad H^s(\mathcal{M}) = \left\{ u = (I - \Delta_{\mathcal{M}})^k v, v \in H^{2k+s}(\mathcal{M}) \right\},$$

where $k \in \mathbb{N}$ is the smallest integer such that $2k + s > 0$. In this case, the corresponding norm is given by $\|u\|_{H^s(\mathcal{M})} = \|v\|_{H^{2k+s}(\mathcal{M})}$. We set $H^0(\mathcal{M}) = L^2(\mathcal{M})$. The reader is referred to Herrmann et al. (2018), Strichartz (1983), Triebel (1985), and references therein for more details on Sobolev spaces defined using Bessel potentials.

In this work, we consider elliptic differential operators associated to bilinear forms $A_{\mathcal{M}}$ given by

$$(3) \quad A_{\mathcal{M}}(u, v) = \int_{\mathcal{M}} (\mathcal{D}\nabla_{\mathcal{M}}u) \cdot (\nabla_{\mathcal{M}}\bar{v}) \, dA + \int_{\mathcal{M}} (Vu)\bar{v} \, dA, \quad u, v \in H^1(\mathcal{M}),$$

where for any $x_0 \in \mathcal{M}$ the diffusion matrix $\mathcal{D}(x_0) = [D_{ij}(x_0)]_{i,j=1}^{d+1}$ is a real-valued, symmetric matrix such that for any $w \in T_{x_0}\mathcal{M}$, $\mathcal{D}(x_0)w \in T_{x_0}\mathcal{M}$ and $(\mathcal{D}(x_0)w) \cdot \bar{w} > 0$ if $w \neq 0$. In particular, since $T_{x_0}\mathcal{M} = [\nu(x_0)^\perp]$, $\mathcal{D}(x_0)$ is simply a matrix admitting $\nu(x_0)$ as an eigenvector with some eigenvalue $\mu_1(x_0) \in \mathbb{R}$, and such that the eigenvalues $\mu_i(x_0)$, $2 \leq i \leq d+1$ associated with its other eigenvectors are positive. Without loss of generality, we may assume that $\mu_1(x_0) = 0$, meaning that $\mathcal{D}\nu = 0$, that the eigenvalues $\mu_i(x_0)$, $2 \leq i \leq d+1$ are uniformly lower-bounded and upper-bounded on \mathcal{M} by positive constants, and that $D_{ij} \in L^\infty(\mathcal{M})$ for any $1 \leq i, j \leq d+1$. Finally, we assume that $V \in L^\infty(\mathcal{M})$ is a real-valued function that satisfies $V_- \leq V \leq V_+$ for some $0 < V_- \leq V_+ < +\infty$.

Throughout this paper, let $A_{\mathcal{M}}$ be coercive and continuous, i.e., there exist positive constants δ and M such that for all $u, v \in H^1(\mathcal{M})$,

$$(4) \quad A_{\mathcal{M}}(u, u) \geq \delta \|u\|_{H^1(\mathcal{M})}^2,$$

$$(5) \quad |A_{\mathcal{M}}(u, v)| \leq M \|u\|_{H^1(\mathcal{M})} \|v\|_{H^1(\mathcal{M})}.$$

Following Yagi (2010, Equation (1.33)), $A_{\mathcal{M}}$ gives rise to an associated elliptic differential operator $\mathcal{L} : H^1(\mathcal{M}) \rightarrow H^{-1}(\mathcal{M})$ defined weakly by

$$A_{\mathcal{M}}(u, v) = \int_{\mathcal{M}} (\mathcal{L}u)\bar{v} \, dA, \quad u, v \in H^1(\mathcal{M}).$$

The spectral properties of this operator are detailed in the next proposition, which is proven in Appendix A.

Proposition 2.1. *Let $\delta > 0$ be the coercivity constant defined in Equation (4). There exists a set of eigenpairs $\{(\lambda_i, e_i)\}_{i \in \mathbb{N}}$ of \mathcal{L} consisting of a sequence of increasing real-valued eigenvalues $0 < \delta \leq \lambda_1 \leq \lambda_2 \leq \dots$ with $\lambda_i \rightarrow +\infty$ as $i \rightarrow +\infty$, and $\{e_i\}_{i \in \mathbb{N}}$ forms an orthonormal basis of $L^2(\mathcal{M})$ where each e_i is real-valued.*

Since the operator \mathcal{L} differs from the Laplace–Beltrami operator only by a zeroth-order potential term and a diffusion function in the second order term, switching between the two operators corresponds to a change of metric on \mathcal{M} . Therefore, the eigenvalue problem for \mathcal{L} is equivalent to that for the Laplace–Beltrami operator on \mathcal{M} equipped with a possibly rough metric if the coefficients of \mathcal{D} are not smooth. The results in Bandara et al. (2021) imply growth rates on the eigenvalues in accordance with Weyl’s law, and more specifically that there exist $c_{\mathcal{M}}, C_{\mathcal{M}} > 0$ such that for any $i \in \mathbb{N}$

$$(6) \quad c_{\mathcal{M}} i^{2/d} \leq \lambda_i \leq C_{\mathcal{M}} i^{2/d}.$$




FIGURE 2. One-dimensional illustration of the lift. The lift is along the normal vector ν to the surface \mathcal{M} .

As a last step in this subsection, we introduce nonlinear functions of \mathcal{L} which allow later in Section 3 for the definition of a variety of Gaussian random fields. For that, we call a function $\gamma : \mathbb{R}_+ \rightarrow \mathbb{R}$ an α -power spectral density if

- 1) γ is extendable to a holomorphic function on $H_{\pi/2} := \{z \in \mathbb{C} : |\arg z| \leq \pi/2\}$.
- 2) There exist constants $C_\gamma > 0$ and $\alpha > 0$ such that for all $z \in H_{\pi/2}$,

$$(7) \quad |\gamma(z)| \leq C_\gamma |z|^{-\alpha}.$$

Applying a power spectral density to \mathcal{L} results in a linear operator $\gamma(\mathcal{L})$ whose action on functions $f \in L^2(\mathcal{M})$ is defined by

$$(8) \quad \gamma(\mathcal{L})f = \sum_{i=1}^{\infty} \gamma(\lambda_i)(f, e_i)_{L^2(\mathcal{M})} e_i,$$

where $\{(\lambda_i, e_i)\}_{i \in \mathbb{N}}$ are the eigenpairs of \mathcal{L} defined in Proposition 2.1. A typical example is the function $\gamma(\lambda) = (\kappa^2 + \lambda)^{-\alpha}$ for $\alpha > d/4$ and $\kappa > 0$, which can be used to obtain Whittle–Matérn random fields (Lang and Pereira, 2023).

The goal of the remainder of this section is to study the approximation of functions of the form $u = \gamma(\mathcal{L})f$, where $f \in L^2(\mathcal{M})$. Formally, if $1/\gamma$ is well-defined on the spectrum of \mathcal{L} , then this is the solution to the partial differential equation $\gamma(\mathcal{L})^{-1}u = f$.

2.2. SFEM–Galerkin approximation. The idea behind the surface finite element method, as introduced by Dziuk (1988), is to work on a polyhedral approximation of the surface that is in some sense close to the true surface \mathcal{M} . More precisely, fix $h > 0$ and let \mathcal{M}_h be a piecewise polygonal surface consisting of non-degenerate simplices (for $d = 2$, triangles and for $d = 1$, line segments) with vertices on \mathcal{M} , and such that h is the size of the largest simplex defined as the in-ball radius. The set of simplices making up the discretized surface is denoted by \mathcal{T}_h , thus meaning that

$$\mathcal{M}_h = \bigcup_{T_j \in \mathcal{T}_h} T_j,$$

and we assume that for any two simplices in \mathcal{T}_h , it holds that their intersection is either empty, or a common edge or vertex.

Following (Dziuk and Elliott, 2013, Section 1.4.1), we assume that the triangulation \mathcal{T}_h is quasi-uniform, shape-regular, and that the number of simplices sharing the same vertex can be upper-bounded by a constant independent of h . In turn, these two assumption imply that $N_h \propto h^{-d}$ where $N_h \in \mathbb{N}$ denotes the number of vertices of \mathcal{M}_h .

The discrete surface \mathcal{M}_h is close to the true surface \mathcal{M} in the sense that \mathcal{M}_h is contained in a small neighborhood around \mathcal{M} defined as follows. First, note that \mathcal{M} can be seen as

the boundary of some bounded open set $G \subset \mathbb{R}^{d+1}$ with exterior normal ν . Then, following Dziuk and Elliott (2013, Section 2.3), we consider that there exists some (small) $\varpi > 0$ such that \mathcal{M}_h is contained in a so-called *tubular neighborhood* U_ϖ of \mathcal{M} defined by

$$U_\varpi = \{x \in \mathbb{R}^{d+1} : |d_s(x)| < \varpi\},$$

where $d_s : \mathbb{R}^{d+1} \rightarrow \mathbb{R}$ denotes the oriented distance function given by

$$d_s(x) = \begin{cases} \inf_{y \in \mathcal{M}} |x - y|, & x \in \mathbb{R}^{d+1} \setminus G, \\ \inf_{y \in \mathcal{M}} -|x - y|, & x \in G. \end{cases}$$

We denote by dA_h the surface measure on \mathcal{M}_h , and by $L^2(\mathcal{M}_h)$ the Hilbert space of dA_h -measurable square integrable functions, equipped with the inner product $(\cdot, \cdot)_{L^2(\mathcal{M}_h)}$ defined by

$$(u_h, v_h)_{L^2(\mathcal{M}_h)} = \int_{\mathcal{M}_h} u_h \bar{v}_h \, dA_h, \quad u_h, v_h \in L^2(\mathcal{M}_h).$$

Following (Bonito et al., 2020, Section 1.2.1), we denote by $\sigma : \mathcal{M}_h \rightarrow \mathbb{R}^+$ the *area element* given by $\sigma = dA/dA_h$, such that for all $v \in L^2(\mathcal{M})$,

$$(9) \quad \int_{\mathcal{M}} v \, dA = \int_{\mathcal{M}_h} \sigma v^{-\ell} \, dA_h,$$

where we next introduce the lift and its inverse denoted by ℓ and $-\ell$, respectively.

A key element of SFEM is that we can move between \mathcal{M} and \mathcal{M}_h using the so-called *lift operator*. To construct the lift operator, we note that $d_s \in C^k(U_\varpi)$ for $k \geq 2$, and that for any $x \in U_\varpi$, there exists a unique $a(x) \in \mathcal{M}$ such that

$$x = a(x) + d_s(x)\nu(a(x)),$$

where ν denotes the normal at $a(x)$ to \mathcal{M} . In particular, this implies that any point $x \in U_\varpi$ can be uniquely described by the pair $(a(x), d_s(x)) \in \mathcal{M} \times \mathbb{R}$, and this procedure defines an isomorphism $p : \mathcal{M}_h \rightarrow \mathcal{M}$ given by

$$p(x) = x - d_s(x)\nu(a(x)), \quad x \in \mathcal{M}_h.$$

Therefore, any function $\eta : \mathcal{M}_h \rightarrow \mathbb{C}$ may be lifted to \mathcal{M} by $\eta^\ell = \eta \circ p^{-1} : \mathcal{M} \rightarrow \mathbb{C}$. Likewise, the inverse lift of any function $\zeta : \mathcal{M} \rightarrow \mathbb{C}$ is given by $\zeta^{-\ell} = \zeta \circ p : \mathcal{M}_h \rightarrow \mathbb{C}$. The procedure is illustrated in Figure 2 in the one-dimensional setting. Note that the points on the discretized surface are *lifted* along the normal ν to the surface \mathcal{M} .

The mapping a is used to define the gradient of functions on \mathcal{M}_h (Dziuk and Elliott, 2013). More specifically, for a differentiable $\eta : \mathcal{M}_h \rightarrow \mathbb{C}$, the gradient is given by

$$(10) \quad \nabla_{\mathcal{M}_h} \eta(x) = \nabla \tilde{\eta}(x) - (\nabla \tilde{\eta}(x) \cdot \nu_h(x))\nu_h(x) \in T_x \mathcal{M}_h,$$

where ν_h is the normal of \mathcal{M}_h and $\tilde{\eta}$ is the continuous extension of η defined by $\tilde{\eta} : x \in U_\varpi \mapsto \tilde{\eta}(x) = \eta^\ell(a(x))$. With this definition, the Laplace–Beltrami operator on \mathcal{M}_h can be defined by $\Delta_{\mathcal{M}_h} = \nabla_{\mathcal{M}_h} \cdot \nabla_{\mathcal{M}_h}$, and Sobolev spaces on \mathcal{M}_h are defined in complete analogy to those on \mathcal{M} .

The analogue of the bilinear form $A_{\mathcal{M}}$ on \mathcal{M}_h is given by

$$(11) \quad A_{\mathcal{M}_h}(u_h, v_h) = \int_{\mathcal{M}_h} (\mathcal{D}^{-\ell} \nabla_{\mathcal{M}_h} u_h) \cdot (\nabla_{\mathcal{M}_h} \bar{v}_h) \, dA_h + \int_{\mathcal{M}_h} (V^{-\ell} u_h) \bar{v}_h \, dA_h, \quad u_h, v_h \in H^1(\mathcal{M}_h),$$

where $\mathcal{D}^{-\ell} = [D_{ij}^{-\ell}]_{i,j=1}^{d+1}$. As in Dziuk and Elliott (2013), we assume that there exists $h_0 \in (0, 1)$ small enough, such that $\mathbf{A}_{\mathcal{M}_h}$ is coercive (and continuous) whenever $h \leq h_0$. Unless stated otherwise, we now assume that this last condition on h is fulfilled.

To conclude this subsection, we introduce the (linear) finite element space S_h on \mathcal{M}_h . The finite element space S_h is defined as the complex span of the standard real-valued *nodal basis* $\psi_1, \dots, \psi_{N_h} : \mathcal{M} \rightarrow \mathbb{R}$, where for any $i \in \{1, \dots, N_h\}$, $\psi_i|_T$ is a polynomial of at most degree one taking the value 1 at the i -th vertex of \mathcal{M}_h and 0 at all the other vertices, i.e.,

$$S_h = \text{span}(\psi_1, \dots, \psi_{N_h}) \subset H^1(\mathcal{M}_h).$$

By construction, S_h is a vector space of dimension N_h . Its counterpart on \mathcal{M} is the lifted finite element space S_h^ℓ given by

$$S_h^\ell = \left\{ \phi_h^\ell, \phi_h \in S_h \right\} \subset H^1(\mathcal{M}).$$

On S_h , we can, as \mathcal{L} on \mathcal{M} , associate to the bilinear form $\mathbf{A}_{\mathcal{M}_h}$ in Equation (11) a linear operator $\mathbf{L}_h : S_h \rightarrow S_h$ which maps any $u_h \in S_h$ to the unique $\mathbf{L}_h u_h \in S_h$ satisfying, for any $v_h \in S_h$, the equality

$$\mathbf{A}_{\mathcal{M}_h}(u_h, v_h) = (\mathbf{L}_h u_h, v_h)_{L^2(\mathcal{M}_h)}.$$

Similarly, if the bilinear form $\mathbf{A}_{\mathcal{M}}$ introduced in Equation (3) is restricted to S_h^ℓ , we can associate it to a linear operator $\mathcal{L}_h : S_h^\ell \rightarrow S_h^\ell$ that maps any $u_h^\ell \in S_h^\ell$ to the unique $\mathcal{L}_h u_h^\ell \in S_h^\ell$ that satisfies, for any $v_h^\ell \in S_h^\ell$, the equality

$$\mathbf{A}_{\mathcal{M}}(u_h^\ell, v_h^\ell) = (\mathcal{L}_h u_h^\ell, v_h^\ell)_{L^2(\mathcal{M})}.$$

Since the bilinear forms $\mathbf{A}_{\mathcal{M}}$ and $\mathbf{A}_{\mathcal{M}_h}$ are coercive, positive definite, Hermitian and have real coefficients, these two operators are diagonalizable in the sense that they each give rise to a set of N_h eigenpairs (Hall, 2013). On the one hand, there exists a sequence $0 \leq \Lambda_1^h \leq \dots \leq \Lambda_{N_h}^h$ and an $L^2(\mathcal{M}_h)$ -orthonormal basis $E_1^h, \dots, E_{N_h}^h$ of S_h such that

$$\mathbf{L}_h E_i^h = \Lambda_i^h E_i^h, \quad 1 \leq i \leq N_h,$$

and similarly there exists a sequence $0 \leq \lambda_1^h \leq \dots \leq \lambda_{N_h}^h$ and an $L^2(\mathcal{M})$ -orthonormal basis $e_1^h, \dots, e_{N_h}^h$ of S_h^ℓ such that

$$\mathcal{L}_h e_i^h = \lambda_i^h e_i^h, \quad 1 \leq i \leq N_h.$$

In particular, using the same approach as in Proposition 2.1, we can assume that the eigenfunctions $\{E_i^h\}_{1 \leq i \leq N_h}$ and $\{e_i^h\}_{1 \leq i \leq N_h}$ are all real-valued. The eigenvalues of the operators \mathcal{L} , \mathcal{L}_h and \mathbf{L}_h are linked to one another through the following lemma, due to Bonito et al. (2018, Lemma 3.1), Bonito et al. (2024, Lemma 4.1) and Strang and Fix (2008, Theorem 6.1).

In the following, $A \lesssim B$ is shorthand for that there is a constant $C > 0$ such that $A \leq CB$.

Lemma 2.2 (Eigenvalue error bounds). *Let $\{\lambda_i\}_{i \in \mathbb{N}}$, $\{\lambda_i^h\}_{1 \leq i \leq N_h}$ and $\{\Lambda_i^h\}_{1 \leq i \leq N_h}$ denote the eigenvalues of the operators \mathcal{L} , \mathcal{L}_h and \mathbf{L}_h , respectively. Then,*

$$(12) \quad \lambda_i \leq \lambda_i^h \lesssim (1 + h^2)\lambda_i, \quad 1 \leq i \leq N_h,$$

and

$$(13) \quad |\lambda_i^h - \Lambda_i^h| \lesssim h^2 \lambda_i^h \lesssim h^2 \lambda_i, \quad 1 \leq i \leq N_h.$$

Finally, we remark that the eigenvalues $\{\Lambda_i^h\}_{1 \leq i \leq N_h}$ of L_h can be linked to the eigenvalues of some classical finite element matrices. Let \mathbf{C} and \mathbf{R} be the so-called *mass matrix* and *stiffness matrix*, respectively, and defined from the nodal basis by

$$(14) \quad \mathbf{C} = [(\psi_k, \psi_l)_{L^2(\mathcal{M}_h)}]_{1 \leq k, l \leq N_h}, \quad \mathbf{R} = [A_{\mathcal{M}_h}(\psi_k, \psi_l)]_{1 \leq k, l \leq N_h}.$$

As defined, \mathbf{C} is a symmetric positive definite matrix and \mathbf{R} is a symmetric positive semi-definite matrix. Let then $\sqrt{\mathbf{C}} \in \mathbb{R}^{N_h \times N_h}$ be an invertible matrix satisfying $\sqrt{\mathbf{C}}(\sqrt{\mathbf{C}})^T = \mathbf{C}$. Then, by Lang and Pereira (2023, Corollary 3.2), the eigenvalues $\{\Lambda_i^h\}_{1 \leq i \leq N_h}$ are also the eigenvalues of the matrix $\mathbf{S} \in \mathbb{R}^{N_h \times N_h}$ defined by

$$(15) \quad \mathbf{S} = (\sqrt{\mathbf{C}})^{-1} \mathbf{R} (\sqrt{\mathbf{C}})^{-T}.$$

Besides, if $\boldsymbol{\psi}$ denotes the vector-valued function given by $\boldsymbol{\psi} = (\psi_1, \dots, \psi_{N_h})^T$, then the mapping $F : \mathbb{R}^{N_h} \rightarrow S_h$, defined by

$$F(\mathbf{v}) = \boldsymbol{\psi}^T (\sqrt{\mathbf{C}})^{-T} \mathbf{v}, \quad \mathbf{v} \in \mathbb{R}^{N_h},$$

is an isomorphism whose inverse maps the eigenfunctions $\{E_i^h\}_{1 \leq i \leq N_h}$ to (orthonormal) eigenvectors of \mathbf{S} . This means in particular that \mathbf{S} can also be written as

$$(16) \quad \mathbf{S} = \mathbf{V} \text{Diag}(\Lambda_1^h, \dots, \Lambda_{N_h}^h) \mathbf{V}^T,$$

where $\mathbf{V} = (F^{-1}(E_1^h) | \dots | F^{-1}(E_{N_h}^h)) \in \mathbb{R}^{N_h \times N_h}$.

2.3. Deterministic error analysis. Based on the introduced framework, we are now in place to quantify the error between functions of the operators \mathcal{L} , \mathcal{L}_h and L_h . Let $P_h : L^2(\mathcal{M}) \rightarrow S_h^\ell$ be the L^2 -projection onto S_h^ℓ and let $P_h : L^2(\mathcal{M}_h) \rightarrow S_h$ the L^2 -projection onto S_h . We note that the operators \mathcal{L} , \mathcal{L}_h , and L_h define norms that are equivalent to the standard Sobolev norms, i.e.,

$$(17) \quad \begin{aligned} \|\mathcal{L}^{1/2} v\|_{L^2(\mathcal{M})} &\sim \|v\|_{H^1(\mathcal{M})}, \quad \|\mathcal{L}_h^{1/2} V_h\|_{L^2(\mathcal{M})} \sim \|V_h\|_{H^1(\mathcal{M}_h)}, \\ \|\mathcal{L}_h^{1/2} v_h\|_{L^2(\mathcal{M})} &\sim \|v_h\|_{H^1(\mathcal{M})}, \end{aligned}$$

for all $v \in H^1(\mathcal{M})$, and all $v_h \in S_h^\ell$ and $V_h \in S_h$.

With that at hand we are ready to state our main result in this section.

Proposition 2.3. *Let γ be an α -power spectral density. Then, there is a constant $C > 0$ such that for all $f \in L^2(\mathcal{M})$*

$$\|\gamma(\mathcal{L}_h) P_h f - \gamma(\mathcal{L}) f\|_{L^2(\mathcal{M})} \leq \begin{cases} Ch^2 \|f\|_{L^2(\mathcal{M})}, & \alpha > 1, \\ Ch^2 |\log h| \|f\|_{L^2(\mathcal{M})}, & \alpha = 1, \\ Ch^{2\alpha} \|f\|_{L^2(\mathcal{M})}, & \alpha < 1. \end{cases}$$

To prove this proposition we rely on a representation of functions of operators based on Cauchy–Stieltjes integrals, which are constructed as follows. Since the sesquilinear form defined by $A_{\mathcal{M}}$ is continuous and coercive, the associated operators \mathcal{L} and \mathcal{L}_h are sectorial with some (common) angle $\theta \in (0, \pi/2)$ (Yagi, 2010, Theorem 2.1). Therefore, the spectra of \mathcal{L} and \mathcal{L}_h are contained in the complement of the set $G_\theta = \{z \in \mathbb{C}, \theta \leq \arg(z) \leq \pi\}$, as




FIGURE 3. Contours used to define the Cauchy–Stieltjes integral representation of operators.

illustrated in Figure 3(a), and the following inequalities are satisfied for any $z \in G_\theta$ (cf. Yagi (2010, Equation (2.2))):

$$(18) \quad \begin{aligned} \|(z - \mathcal{L})^{-1}v\|_{L^2(\mathcal{M})} &\leq C_\theta |z|^{-1} \|v\|_{L^2(\mathcal{M})}, \quad v \in L^2(\mathcal{M}), \\ \|(z - \mathcal{L}_h)^{-1}v_h^\ell\|_{L^2(\mathcal{M})} &\leq C_\theta |z|^{-1} \|v_h^\ell\|_{L^2(\mathcal{M})}, \quad v_h^\ell \in S_h^\ell, \end{aligned}$$

where $C_\theta > 0$ is a generic constant. Note in particular that by definition, G_θ is contained in the resolvent sets of \mathcal{L} and \mathcal{L}_h , and that any power spectral density γ is bounded, holomorphic and satisfies the inequality $|z|^\alpha |\gamma(z)| \lesssim 1$ for any $z \in G_\theta$. Hence, the operators $\gamma(\mathcal{L})$ and $\gamma(\mathcal{L}_h)$ can be defined as functional calculi of the operators \mathcal{L} and \mathcal{L}_h as (Yagi, 2010, Chapter 16, Section 1.2) by

$$(19) \quad \gamma(\mathcal{L}) = \frac{1}{2\pi i} \int_\Gamma \gamma(z)(z - \mathcal{L})^{-1} dz \quad \text{and} \quad \gamma(\mathcal{L}_h) = \frac{1}{2\pi i} \int_\Gamma \gamma(z)(z - \mathcal{L}_h)^{-1} dz,$$

where $\Gamma \subset \mathbb{C}$ is any integral contour surrounding the spectra of \mathcal{L} and \mathcal{L}_h and contained in G_θ . In particular, these new definitions of functions of operators are independent of the choice of Γ , and coincide with the spectral definitions previously introduced in Equation (8) (cf. e.g. (Yagi, 2010, Remark 2.7)).

In the remainder, we split the contour Γ into

$$(20) \quad \Gamma = \Gamma_+ \cup \Gamma_0 \cup \Gamma_-,$$

where Γ_+ is parametrized by $g_+(t) = te^{i\theta}$ for $t \in (\infty, \delta_0]$, Γ_0 by $g_0(t) = \delta_0 e^{it}$ for $t \in [\theta, -\theta]$ and Γ_- by $g_-(t) = te^{i\theta}$ for $t \in [\delta_0, \infty)$ with $\delta_0 < \delta/2$ (see Figure 3(b) for an illustration). We use this contour to prove Proposition 2.3, while relying on the following bounds for the resolvent error along Γ . The proof is included in Appendix A and is an adaption of results from Fujita and Suzuki (1991) and Bonito et al. (2024).

Lemma 2.4. *For any $z \in \Gamma$, $f \in L^2(\mathcal{M})$, and $\beta \in [0, 1]$,*

$$\|(z - \mathcal{L}_h)^{-1}P_h f - (z - \mathcal{L})^{-1}f\|_{L^2(\mathcal{M})} \leq Ch^{2\beta} |z|^{-(1-\beta)}.$$

We now provide a proof for Proposition 2.3.

Proof of Proposition 2.3. Let $f \in L^2(\mathcal{M})$. For any $z \in \Gamma$, set $e_h(z) = (z - \mathcal{L}_h)^{-1}P_h f - (z - \mathcal{L})^{-1}f$, which yields, using the integral representations of $\gamma(\mathcal{L})$ and $\gamma(\mathcal{L}_h)$,

$$\gamma(\mathcal{L}_h)P_h f - \gamma(\mathcal{L})f = \frac{1}{2\pi i} \int_\Gamma \gamma(z)e_h(z) dz.$$

The definition of Γ and its parametrization allow to decompose the integral as

$$\begin{aligned} \gamma(\mathcal{L}_h)P_h f - \gamma(\mathcal{L})f &= \frac{-1}{2\pi i} \int_{\delta_0}^{\infty} \gamma(g_+(t))e_h(g_+(t))g'_+(t) dt - \frac{1}{2\pi i} \int_{-\theta}^{\theta} \gamma(g_0(t))e_h(g_0(t))g'_0(t) dt \\ &\quad + \frac{1}{2\pi i} \int_{\delta_0}^{\infty} \gamma(g_-(t))e_h(g_-(t))g'_-(t) dt, \end{aligned}$$

where we recall that $g'_+(t) = e^{i\theta}$, $g'_-(t) = e^{-i\theta}$ and $g'_0(t) = i\delta_0 e^{it}$. Taking norms on both sides of this equality and using the triangle inequality then gives

$$\begin{aligned} &\|\gamma(\mathcal{L}_h)P_h f - \gamma(\mathcal{L})f\|_{L^2(\mathcal{M})} \\ &\leq \frac{1}{2\pi} \int_{\delta_0}^{\infty} |\gamma(g_+(t))| \|e_h(g_+(t))\|_{L^2(\mathcal{M})} dt + \frac{\delta_0}{2\pi} \int_{-\theta}^{\theta} |\gamma(g_0(t))| \|e_h(g_0(t))\|_{L^2(\mathcal{M})} dt \\ &\quad + \frac{1}{2\pi} \int_{\delta_0}^{\infty} |\gamma(g_-(t))| \|e_h(g_-(t))\|_{L^2(\mathcal{M})} dt \\ &= I_+ + I_0 + I_-. \end{aligned}$$

Let us start by bounding I_0 . We apply Equation (7) and Lemma 2.4 with $\beta = 1$ to obtain

$$I_0 \leq Ch^2 \|f\|_{L^2(\mathcal{M})} \int_{-\theta}^{\theta} |g_0(t)|^{-\alpha} dt = Ch^2 \|f\|_{L^2(\mathcal{M})} \int_{-\theta}^{\theta} \delta_0^{-\alpha} dt \leq Ch^2 \|f\|_{L^2(\mathcal{M})}.$$

To bound I_+ , we distinguish between the three cases $\alpha > 1$, $\alpha < 1$ and $\alpha = 1$. When $\alpha > 1$, we apply Equation (7) and Lemma 2.4 with $\beta = 1$ to get

$$I_+ \leq Ch^2 \|f\|_{L^2(\mathcal{M})} \int_{\delta_0}^{\infty} |g_+(t)|^{-\alpha} dt = Ch^2 \int_{\delta_0}^{\infty} |t|^{-\alpha} dt = C \frac{\delta_0^{-(\alpha-1)}}{\alpha-1} h^2 \|f\|_{L^2(\mathcal{M})}.$$

For $\alpha < 1$, recall that $h < h_0 < 1$ and assume without loss of generality that $h_0^{-2} > \delta_0$. We split $I_+ = I_{+1} + I_{+2}$ with

$$I_{+1} = \int_{\delta_0}^{h^{-2}} |\gamma(g_+(t))| \|e_h(g_+(t))\|_{L^2(\mathcal{M})} dt, \quad I_{+2} = \int_{h^{-2}}^{\infty} |\gamma(g_+(t))| \|e_h(g_+(t))\|_{L^2(\mathcal{M})} dt.$$

Using Equation (7) and Lemma 2.4 with $\beta = 1$ we bound I_{+1} by

$$\begin{aligned} I_{+1} &\leq Ch^2 \|f\|_{L^2(\mathcal{M})} \int_{\delta_0}^{h^{-2}} |g_+(t)|^{-\alpha} dt = Ch^2 \|f\|_{L^2(\mathcal{M})} \int_{\delta_0}^{h^{-2}} |t|^{-\alpha} dt \\ &\leq Ch^2 \|f\|_{L^2(\mathcal{M})} \int_0^{h^{-2}} |t|^{-\alpha} dt = Ch^2 \|f\|_{L^2(\mathcal{M})} \frac{h^{-2(1-\alpha)}}{1-\alpha} \leq Ch^{2\alpha} \|f\|_{L^2(\mathcal{M})}. \end{aligned}$$

We proceed in the same manner to bound I_{+2} , but use Lemma 2.4 with $\beta = 0$,

$$I_{+2} \leq C \|f\|_{L^2(\mathcal{M})} \int_{h^{-2}}^{\infty} |g_+(t)|^{-\alpha} |g_+(t)|^{-1} dt = C \|f\|_{L^2(\mathcal{M})} \int_{h^{-2}}^{\infty} |t|^{-\alpha-1} dt \leq Ch^{2\alpha} \|f\|_{L^2(\mathcal{M})}.$$

For $\alpha = 1$ and the same splitting $I_+ = I_{+1} + I_{+2}$ we obtain for I_{+1} with Lemma 2.4 for $\beta = 1$

$$\begin{aligned} I_{+1} &\leq Ch^2 \|f\|_{L^2(\mathcal{M})} \int_{\delta_0}^{h^{-2}} t^{-1} dt = Ch^2 \|f\|_{L^2(\mathcal{M})} (\log(h^{-2}) - \log \delta_0) \\ &\leq Ch^2 \|f\|_{L^2(\mathcal{M})} (|\log(h^{-2})| + |\log \delta_0|) \leq Ch^2 \|f\|_{L^2(\mathcal{M})} (|\log h| + |\log \delta_0|). \end{aligned}$$

Assuming again $h < h_0$ and $h_0^{-2} > \delta_0$, we finally bound

$$\begin{aligned} I_{+1} &\leq Ch^2 |\log h| \|f\|_{L^2(\mathcal{M})} \left(1 + \left|\frac{\log \delta_0}{\log h}\right|\right) \leq Ch^2 |\log h| \|f\|_{L^2(\mathcal{M})} \left(1 + \left|\frac{\log \delta_0}{\log h_0}\right|\right) \\ &\leq Ch^2 |\log h| \|f\|_{L^2(\mathcal{M})}. \end{aligned}$$

The bound for $I_{+2} \leq Ch^2 \|f\|_{L^2(\mathcal{M})}$ is obtained in the same way as in the case $\alpha > 1$. Hence, $I_+ \leq Ch^2 |\log h| \|f\|_{L^2(\mathcal{M})}$ for $\alpha = 1$.

Finally, note that by symmetry, I_- satisfies the same bounds as I_+ , which finishes the proof adding up all terms. \square

A result similar to Proposition 2.3 can be derived to quantify the error between functions of the operators \mathcal{L}_h and \mathbf{L}_h . To do so, we note that since $\mathbf{A}_{\mathcal{M}_h}$ is continuous and coercive, the associated operator \mathbf{L}_h is also sectorial with some angle in $(0, \pi/2)$. Hence, without loss of generality, the angle $\theta \in (0, \pi/2)$ can be assumed to be large enough to ensure that Equation (18) also holds for \mathbf{L}_h , i.e. that for any $z \in G_\theta$,

$$(21) \quad \|(z - \mathbf{L}_h)^{-1} v_h\|_{L^2(\mathcal{M}_h)} \leq C_\theta |z|^{-1} \|v_h\|_{L^2(\mathcal{M}_h)}, \quad v_h \in S_h,$$

and that an integral representation similar to Equation (19) also holds for $\gamma(\mathbf{L}_h)$, namely:

$$\gamma(\mathbf{L}_h) = \frac{1}{2\pi i} \int_\Gamma \gamma(z) (z - \mathbf{L}_h)^{-1} dz.$$

Proposition 2.5. *Let γ be an α -power spectral density with $\alpha > d/4$. There exists a constant $C > 0$ such that, for any $\tilde{f} \in S_h^\ell$,*

$$(22) \quad \left\| \left(\gamma(\mathcal{L}_h) \tilde{f} \right)^{-\ell} - \gamma(\mathbf{L}_h) \mathbf{P}_h(\sigma \tilde{f}^{-\ell}) \right\|_{L^2(\mathcal{M}_h)} \leq Ch^2 \|\mathcal{L}_h^{-\min\{\alpha+d/4; 1\}/2} \tilde{f}\|_{L^2(\mathcal{M})}.$$

This proposition can be seen as an extension of Bonito et al. (2024, Lemma 4.4) relying on extensions of the estimates proven in Bonito et al. (2024, Lemma A.1). Its proof is similar and therefore postponed to Appendix B.

3. STOCHASTIC THEORY: RANDOM FIELDS ON SURFACES AND CONVERGENCE OF SFEM APPROXIMATION

In this section, we introduce random fields and white noise on surfaces, thus allowing us to make sense of Equation (1) in Section 1. Further, we give approximation methods based on SFEM and prove strong error bounds.

3.1. Random fields on surfaces. Let $(\Omega, \mathcal{S}, \mathbb{P})$ be a complete probability space. We are interested in Gaussian random fields on \mathcal{M} defined as $L^2(\mathcal{M})$ -valued random variables through expansions of the form

$$(23) \quad \mathcal{Z} = \sum_{i \in \mathbb{N}} Z_i e_i,$$

where $\{e_i\}_{i \in \mathbb{N}}$ denotes a real-valued orthonormal basis of $L^2(\mathcal{M})$ composed of eigenfunctions of the operator \mathcal{L} (cf. Proposition 2.1), and $\{Z_i\}_{i \in \mathbb{N}}$ is a sequence of real Gaussian random variables such that $\mathbb{E}[Z_i] = 0$ for any $i \in \mathbb{N}$ and $\sum_{i \in \mathbb{N}} \mathbb{E}[|Z_i|^2] < \infty$. As such, \mathcal{Z} can be seen as

an element of the Hilbert space $L^2(\Omega; L^2(\mathcal{M}))$ of $L^2(\mathcal{M})$ -valued random variables, to which we associate the inner product $(\cdot, \cdot)_{L^2(\Omega; L^2(\mathcal{M}))}$ (and norm $\|\cdot\|_{L^2(\Omega; L^2(\mathcal{M}))}$) defined by

$$(\mathcal{Z}, \mathcal{Z}')_{L^2(\Omega; L^2(\mathcal{M}))} = \mathbb{E} [(\mathcal{Z}, \mathcal{Z}')_{L^2(\mathcal{M})}], \quad \mathcal{Z}, \mathcal{Z}' \in L^2(\Omega; L^2(\mathcal{M})).$$

Finally, in analogy to Equation (23), we formally define the Gaussian white noise on \mathcal{M} by the expansion

$$(24) \quad \mathcal{W} = \sum_{i \in \mathbb{N}} W_i e_i,$$

where $\{W_i\}_{i \in \mathbb{N}}$ is a sequence of independent real standard Gaussian random variables. We observe that even though this expansion does not converge in $L^2(\Omega; L^2(\mathcal{M}))$, it does however converge in $L^2(\mathcal{M}; H^s(\mathcal{M}))$ for $s < -d/2$. Moreover, we have that for any $\phi \in L^2(\mathcal{M})$, the expansion $(\phi, \mathcal{W})_{L^2(\mathcal{M})} = \sum_{i \in \mathbb{N}} W_i (\phi, e_i)_{L^2(\mathcal{M})}$ converges in $L^2(\Omega)$. Further, $(\mathcal{W}, \phi)_{L^2(\mathcal{M})}$ defines a complex Gaussian variable with mean 0, and for any $\phi' \in L^2(\mathcal{M})$, $\text{Cov}((\phi, \mathcal{W})_{L^2(\mathcal{M})}, (\phi', \mathcal{W})_{L^2(\mathcal{M})}) = \mathbb{E}[(\phi, \mathcal{W})_{L^2(\mathcal{M})} \overline{(\phi', \mathcal{W})_{L^2(\mathcal{M})}}] = (\phi, \phi')_{L^2(\mathcal{M})}$. As such, the Gaussian white noise (24) can be interpreted as a *generalized Gaussian random field* over $L^2(\mathcal{M})$.

Circling back to the class of random fields defined in Section 1, we can now make sense of Equation (1) through Equation (8) and Equation (24), thus yielding the definition

$$(25) \quad \mathcal{Z} = \gamma(\mathcal{L})\mathcal{W} = \sum_{i \in \mathbb{N}} \gamma(\lambda_i) W_i e_i,$$

which results in $\mathcal{Z} \in L^2(\Omega; H^s(\mathcal{M}))$ for any $s \geq 0$ such that $4\alpha - d > 2s$. Note in particular that all summands in Equation (25) are real-valued functions, and that therefore \mathcal{Z} is real-valued. In Figure 4 we illustrate the influence of the parameter choices on the resulting field on \mathbb{S}^2 for generalized *non-stationary Whittle–Matérn* fields on \mathbb{S}^2

$$\mathcal{Z} = (\mathcal{L})^{-\alpha} \mathcal{W},$$

where $\alpha > d/2$. By selecting $D_{ij}(x) = \delta_{ij}$ and $V(x) = \kappa^2$ with $\kappa > 0$ one recovers the classical, stationary Whittle–Matérn fields studied in for instance Cox and Kirchner (2020); Bolin et al. (2020); Bonito et al. (2024); Lindgren et al. (2011); Jansson et al. (2022); Whittle (1963). In Figures 4(a) and 4(b) we show this case with $\kappa^2 = 10$ for a rougher field with $\alpha = 0.55$ and a smoother one with $\alpha = 1.5$, respectively.

Two non-stationary fields are shown in Figure 4(c) and Figure 4(d), obtained by varying the coefficient functions D_{ij} and V and setting $\alpha = 0.75$. In Figure 4(c), we keep $D_{ij} = \delta_{ij}$ but use

$$V(x) = \begin{cases} 10^5 & \text{for } x_2^6 + x_1^3 - x_3^2 \in (0.1, 0.5), \\ 10 & \text{else} \end{cases}$$

resulting in the observed localized behavior, where the field is essentially turned off in the region with large V . More specifically, V describes the local correlation length, where a large $V(x)$ corresponds to a small correlation length around x .

Finally, setting $V = 10$ constant again, we show the influence of varying parameters \mathcal{D} in Figure 4(d). To derive suitable coefficients, we select a smooth function f and compute its gradient $\nabla_{\mathbb{S}^2} f$ as well as its skew-gradient X_f given by $x \times \nabla f(x)$ at each point $x \in \mathbb{S}^2$. We set for fixed $\rho_1, \rho_2 > 0$, $\mathcal{D}(x)u = \rho_1(\nabla f(x) \cdot u)\nabla f(x) + \rho_2(X_f(x) \cdot u)X_f(x)$ for any $u \in T_x \mathbb{S}^2$. Here, the inner product refers to the Riemannian inner product associated with the standard




FIGURE 4. Some examples to highlight the influence of the different model parameters.

round metric on \mathbb{S}^2 . Since $\nabla_{\mathbb{S}^2} f(x)$ and $X_f(x)$ both are in $T_x \mathbb{S}^2$, $\mathcal{D}(x)$ is a linear mapping from $T_x \mathbb{S}^2$ into itself. Further, as $\nabla_{\mathbb{S}^2} f$ is perpendicular to $X_f(x)$, by selecting ρ_1 and ρ_2 , we obtain a field that is elongated either orthogonally to the level sets of f (large ρ_1 , small ρ_2) or tangentially to the level sets (small ρ_1 , and large ρ_2). To generate Figure 4(d), we selected $f(x) = x_2$, i.e., the function returning the second coordinate in Cartesian coordinates, $\rho_1 = 1$ and $\rho_2 = 25$.

3.2. Approximation of random fields with surface finite elements. Let γ be an α -power spectral density with $\alpha > d/4$. Following the approach presented by Lang and Pereira (2023), the field \mathcal{Z} is approximated by an expansion similar to that of Equation (25), but involving only quantities defined on the polyhedral surface \mathcal{M}_h . More precisely, we define the SFEM–Galerkin approximation Z_h of the field \mathcal{Z} by the relation

$$(26) \quad Z_h = \sum_{i=1}^{N_h} \gamma(\Lambda_i^h) W_i^h E_i^h,$$

where $\{W_i^h\}_{1 \leq i \leq N_h}$ is a sequence of independent standard Gaussian random variables, whose precise definition is clarified later in this section, and $\{(\Lambda_i^h, E_i^h)\}_{1 \leq i \leq N_h}$ are the eigenpairs of L_h introduced in Section 2.2. Then, as proven in Lang and Pereira (2023, Theorem 3.4), Z_h can be decomposed in the nodal basis $\{\psi_i\}_{1 \leq i \leq N_h}$ of S_h as

$$Z_h = \sum_{i=1}^{N_h} Z_i \psi_i,$$

where the weights (Z_1, \dots, Z_{N_h}) form a centered Gaussian vector which covariance matrix Σ_Z can be expressed using the matrices \mathbf{C} and \mathbf{S} introduced in Equations (14) and (15) as follows:

$$(27) \quad \Sigma_Z = (\sqrt{\mathbf{C}})^{-T} \gamma(\mathbf{S})^2 (\sqrt{\mathbf{C}})^{-1},$$

and, following Equation (16), the function of matrix $\gamma(\mathbf{S})$ is defined as

$$(28) \quad \gamma(\mathbf{S})^2 = \mathbf{V} \text{Diag} \left(\gamma(\Lambda_1^h)^2, \dots, \gamma(\Lambda_{N_h}^h)^2 \right) \mathbf{V}^T.$$

Note that sampling the weights (Z_1, \dots, Z_{N_h}) , and therefore the field Z_h , using directly the expression of their covariance matrix requires in practice to fully diagonalize \mathbf{S} (since Equation (27) involves a function of a matrix). Such an operation would result in a prohibitive computational cost (of order $\mathcal{O}(N_h^3)$ operations). To avoid this cost, we use the Chebyshev trick proposed by Lang and Pereira (2023, Section 4), and approximate Z_h by the field $Z_{h,M}$ defined by

$$(29) \quad Z_{h,M} = \sum_{i=1}^{N_h} P_{\gamma,M}(\Lambda_i^h) W_i^h E_i^h,$$

where $P_{\gamma,M}$ is a Chebyshev polynomial approximation of degree $M \in \mathbb{N}$ of γ over an interval $[\lambda_{\min}, \lambda_{\max}]$ containing all the eigenvalues $\{\Lambda_i^h\}_{1 \leq i \leq N_h}$ of \mathbf{L}_h (which we recall, coincide with the eigenvalues of \mathbf{S}). Such an interval can be obtained as follows. On the one hand, one can take $\lambda_{\min} = V_-$. On the other hand, following Lang and Pereira (2023), a candidate for λ_{\max} is obtained by applying the Gershgorin circle theorem to \mathbf{S} .

Since $Z_{h,M}$ is defined by just replacing the power spectral density γ by the polynomial $P_{\gamma,M}$, its expansion $Z_{h,M}$ into the nodal basis,

$$Z_{h,M} = \sum_{i=1}^{N_h} Z_i^{(M)} \psi_i,$$

is such that the weights $(Z_1^{(M)}, \dots, Z_{N_h}^{(M)})$ now form a centered Gaussian vector with covariance matrix $\Sigma_{Z^{(M)}}$ given by

$$(30) \quad \Sigma_{Z^{(M)}} = (\sqrt{\mathbf{C}})^{-T} P_{\gamma,M}^2(\mathbf{S}) (\sqrt{\mathbf{C}})^{-1}.$$

Hence, the matrix function in Equation (27) is now replaced by a matrix polynomial $P_{\gamma,M}^2(\mathbf{S})$. This eliminates the eigendecomposition need associated with matrix functions and therefore speeds up the computations. Indeed, the weights $(Z_1^{(M)}, \dots, Z_{N_h}^{(M)})$ can be sampled through

$$\begin{pmatrix} Z_1^{(M)} \\ \vdots \\ Z_{N_h}^{(M)} \end{pmatrix} = (\sqrt{\mathbf{C}})^{-T} P_{\gamma,M}(\mathbf{S}) \begin{pmatrix} \varepsilon_1 \\ \vdots \\ \varepsilon_{N_h} \end{pmatrix}$$

where $\varepsilon_1, \dots, \varepsilon_{N_h}$ are independent standard Gaussian random variables, and the matrix-vector product by $P_{\gamma,M}(\mathbf{S})$ can be computed iteratively while just requiring products between \mathbf{S} and vectors. In the next two subsections, we provide error estimates quantifying the error between our target random field Z and its successive SFEM and polynomial approximations by Z_h and $Z_{h,M}$.

3.3. Error analysis of the SFEM discretization. We start with analyzing the error between the random field Z defined on \mathcal{M} and its SFEM approximation Z_h , as stated in the next theorem.

Theorem 3.1. *Let γ be an α -power spectral density with $\alpha > d/4$. Then, there exists $h_0 > 0$, such that for any $h < h_0$ the strong approximation error of the random field \mathcal{Z} by its discretization Z_h^ℓ satisfies the bound*

$$(31) \quad \|\mathcal{Z} - Z_h^\ell\|_{L^2(\Omega; L^2(\mathcal{M}))} \leq K_\alpha(h) h^{\min\{\alpha-d/4; 2\}},$$

where $K_\alpha(h) = 1$ if $\alpha - d/4 < 2$, $K_\alpha(h) = |\log h|$ if $\alpha - d/4 = 2$, and $K_\alpha(h) = |\log h|^{(d-1)/2}$ otherwise.

To prove the strong error estimate, we rely on the deterministic error bounds proven in the previous section, and on several intermediate approximations of \mathcal{Z} defined on the spaces S_h^ℓ (i.e., on \mathcal{M}) and S_h (i.e., on \mathcal{M}_h). These intermediate approximations require in turn to define approximations of the Gaussian white noise \mathcal{W} on the spaces S_h^ℓ and S_h .

We first define on S_h^ℓ (i.e., on \mathcal{M}), the projected white noise $\widetilde{\mathcal{W}}$ as

$$(32) \quad \widetilde{\mathcal{W}} = \sum_{j=1}^{N_h} \xi_j e_j^h,$$

where we recall that $\{e_j^h\}_{1 \leq i \leq N_h}$ denotes an orthonormal basis of eigenfunctions of \mathcal{L}_h , and for any $j \in \{1, \dots, N_h\}$, we take $\xi_j = (e_j^h, \mathcal{W})_{L^2(\mathcal{M})}$. In particular, this last relation implies (by definition of the white noise \mathcal{W}) that ξ_1, \dots, ξ_{N_h} are independent standard Gaussian random variables.

Remark 3.2. By injecting the representation (24) of \mathcal{W} in the definition of ξ_j , we get that $\widetilde{\mathcal{W}}$ can itself be formally represented by $\widetilde{\mathcal{W}} = \sum_{k=1}^{\infty} W_k P_h e_k = P_h \mathcal{W}$. This explains why we refer to it as a projected white noise.

Based on $\widetilde{\mathcal{W}}$, we can then introduce a first approximation, on the space S_h^ℓ , of the field \mathcal{Z} . We denote this approximation by \tilde{Z}_h and define it in analogy to Equation (25) as

$$(33) \quad \tilde{Z}_h = \gamma(\mathcal{L}_h) \widetilde{\mathcal{W}} = \sum_{j=1}^{N_h} \gamma(\lambda_j^h) \xi_j e_j^h.$$

Now, on the space S_h , we define *two* white noise approximations $\widehat{\mathcal{W}}$ and \mathcal{W} which are based on the projected white noise $\widetilde{\mathcal{W}}$:

$$\widehat{\mathcal{W}} = P_h(\sigma \widetilde{\mathcal{W}}^{-\ell}) \quad \text{and} \quad \mathcal{W} = P_h(\sigma^{1/2} \widetilde{\mathcal{W}}^{-\ell}),$$

where σ is the ratio of area measures introduced in Section 2.2. On the one hand, we associate to $\widehat{\mathcal{W}}$ an approximation \widehat{Z}_h of the field \mathcal{Z} on S_h , which we define in analogy to Equation (34) as

$$\widehat{Z}_h = \gamma(L_h) \widehat{\mathcal{W}}.$$

On the other hand, by expanding $\widehat{\mathcal{W}}$ and \mathcal{W} in the orthonormal basis $\{E_j^h\}_{1 \leq i \leq N_h}$ of eigenfunctions of L_h , we obtain alternative representations of these fields, and we can draw a link between \mathcal{W} and the SFEM–Galerkin approximation Z_h , as stated in the next lemma.

Lemma 3.3. *The noises $\widehat{\mathcal{W}}$ and \mathcal{W} can be written as*

$$\widehat{\mathcal{W}} = \sum_{j=1}^{N_h} \alpha_j E_j^h \quad \text{and} \quad \mathcal{W} = \sum_{j=1}^{N_h} \beta_j E_j^h,$$

where $(\alpha_1, \dots, \alpha_{N_h})$ and $(\beta_1, \dots, \beta_{N_h})$ are multivariate normal with mean 0 and respective covariance matrices $\mathbf{A} = [(\sigma E_i^h, E_j^h)_{L^2(\mathcal{M}_h)}]_{1 \leq i, j \leq N_h}$ and $\mathbf{B} = \mathbf{I}$. In particular, it holds that the SFEM–Galerkin approximation \mathbf{Z}_h defined in Equation (26) satisfies

$$(34) \quad \mathbf{Z}_h = \gamma(\mathbf{L}_h)\mathbf{W}$$

where we take for any $j \in \{1, \dots, N_h\}$, $W_j^h = \beta_j$.

Proof. As $\widehat{\mathbf{W}} \in S_h$, we can expand it in the orthonormal basis $\{E_j^h\}_{1 \leq j \leq N_h}$ to get

$$\widehat{\mathbf{W}} = \sum_{j=1}^{N_h} \alpha_j E_j^h$$

where $\alpha_j = (\widehat{\mathbf{W}}, E_j^h)_{L^2(\mathcal{M}_h)} = (\sigma \widetilde{\mathbf{W}}^{-\ell}, E_j^h)_{L^2(\mathcal{M}_h)}$. Then, by definition of σ , of $\widetilde{\mathbf{W}}$ and since $(E_j^h)^\ell \in S_h^\ell$, we further get for any $1 \leq j \leq N_h$,

$$\begin{aligned} \alpha_j &= (\widetilde{\mathcal{W}}, (E_j^h)^\ell)_{L^2(\mathcal{M})} = \sum_{k=1}^{N_h} (\mathcal{W}, e_k^h)_{L^2(\mathcal{M})} (e_k^h, (E_j^h)^\ell)_{L^2(\mathcal{M})} \\ &= \left(\mathcal{W}, \sum_{k=1}^{N_h} (e_k^h, (E_j^h)^\ell)_{L^2(\mathcal{M})} e_k^h \right)_{L^2(\mathcal{M})} = (\mathcal{W}, (E_j^h)^\ell)_{L^2(\mathcal{M})}. \end{aligned}$$

Therefore, by definition of the white noise \mathcal{W} , we can conclude that for any $1 \leq i, j \leq N_h$, α_j is normally distributed with mean 0, and that

$$\mathbb{E}[\alpha_i \alpha_j] = \text{Cov}(\alpha_i, \alpha_j) = \left((E_i^h)^\ell, (E_j^h)^\ell \right)_{L^2(\mathcal{M})} = \left(\sigma E_i^h, E_j^h \right)_{L^2(\mathcal{M}_h)} = A_{ij}.$$

Hence, $(\alpha_1, \dots, \alpha_{N_h})$ is indeed multivariate normal (any linear combination of the α_j being Gaussian by definition of \mathcal{W}) with mean 0 and covariance matrix \mathbf{A} .

Similarly, since $\mathbf{W} \in S_h$, we can write again

$$\mathbf{W} = \sum_{j=1}^{N_h} \beta_j E_j^h,$$

where $\beta_j = (\mathbf{W}, E_j^h)_{L^2(\mathcal{M}_h)} = (\sigma^{1/2} \widetilde{\mathbf{W}}^{-\ell}, E_j^h)_{L^2(\mathcal{M}_h)} = (\sigma \widetilde{\mathbf{W}}^{-\ell}, \sigma^{-1/2} E_j^h)_{L^2(\mathcal{M}_h)}$, and the same computations as before yield that $\beta_j = (\mathcal{W}, (\sigma^{-1/2} E_j^h)^\ell)_{L^2(\mathcal{M})}$. Hence, we can conclude this time that for any $1 \leq i, j \leq N_h$, β_j is also normally distributed with mean 0, and that

$$\mathbb{E}[\beta_i \beta_j] = \left(\sigma(\sigma^{-1/2} E_i^h), (\sigma^{-1/2} E_j^h)^\ell \right)_{L^2(\mathcal{M}_h)} = \left(\sigma(\sigma^{-1/2} E_i^h), (\sigma^{-1/2} E_j^h)^\ell \right)_{L^2(\mathcal{M}_h)} = B_{ij},$$

by orthonormality of $\{E_j^h\}_{1 \leq j \leq N_h}$. In conclusion, $(\beta_1, \dots, \beta_{N_h})$ is indeed multivariate normal with mean 0 and covariance matrix $\mathbf{B} = \mathbf{I}$. In particular, this means that $\beta_1, \dots, \beta_{N_h}$ are independent standard Gaussian variables.

Finally, note that we can write

$$\begin{aligned}\gamma(\mathcal{L}_h)P_h\mathcal{W} &= \sum_{j=1}^{N_h} \gamma(\Lambda_j^h)(P_h\mathcal{W}, E_j^h)_{L^2(\mathcal{M}_h)} E_j^h = \sum_{j=1}^{N_h} \gamma(\Lambda_j^h)(\mathcal{W}, E_j^h)_{L^2(\mathcal{M}_h)} E_j^h \\ &= \sum_{j=1}^{N_h} \gamma(\Lambda_j^h)\beta_j E_j^h = \mathcal{Z}_h\end{aligned}$$

where we take $W_j^h = \beta_j$ in Equation (26). \square

We now circle back to proving Theorem 3.1. Using the intermediate approximations $\tilde{\mathcal{Z}}_h$ and $\hat{\mathcal{Z}}_h$ and the equivalence of the L^2 norms on \mathcal{M} and \mathcal{M}_h , we can upper-bound the error between the field \mathcal{Z} and its SFEM–Galerkin approximation \mathcal{Z}_h by

$$\begin{aligned}\|\mathcal{Z} - \mathcal{Z}_h^\ell\|_{L^2(\Omega; L^2(\mathcal{M}))} &\lesssim \|\mathcal{Z} - \tilde{\mathcal{Z}}_h\|_{L^2(\Omega; L^2(\mathcal{M}))} + \|\tilde{\mathcal{Z}}_h - \hat{\mathcal{Z}}_h^\ell\|_{L^2(\Omega; L^2(\mathcal{M}))} + \|\hat{\mathcal{Z}}_h^\ell - \mathcal{Z}_h^\ell\|_{L^2(\Omega; L^2(\mathcal{M}))} \\ &\lesssim \|\mathcal{Z} - \tilde{\mathcal{Z}}_h\|_{L^2(\Omega; L^2(\mathcal{M}))} + \|(\tilde{\mathcal{Z}}_h)^{-\ell} - \hat{\mathcal{Z}}_h\|_{L^2(\Omega; L^2(\mathcal{M}_h))} + \|\hat{\mathcal{Z}}_h - \mathcal{Z}_h\|_{L^2(\Omega; L^2(\mathcal{M}_h))}.\end{aligned}$$

We derive error estimates for each one of the three terms obtained in the last inequality. We start with the term $\|\mathcal{Z} - \tilde{\mathcal{Z}}_h\|_{L^2(\Omega; L^2(\mathcal{M}))}$.

Lemma 3.4. *It holds that*

$$\|\mathcal{Z} - \tilde{\mathcal{Z}}_h\|_{L^2(\Omega; L^2(\mathcal{M}))} \lesssim C_\alpha(h)h^{\min\{\alpha-d/4; 2\}},$$

where $C_\alpha(h) = |\log h|$ if $\alpha - d/4 = 2$ and $C_\alpha(h) = 1$, otherwise.

Proof. Our aim is to bound the error between $\tilde{\mathcal{Z}}_h$ and \mathcal{Z} , where we remark in particular that $\mathcal{Z} = \gamma(\mathcal{L})\mathcal{W}$ and $\mathcal{Z}_h = \gamma(\mathcal{L}_h)P_h\mathcal{W}$ (cf. Remark 3.2). Let $\eta = \alpha + d/4 > d/2$, and let $\zeta = \alpha - d/4 > 0$, so that $\alpha = (\eta + \zeta)/2$. Further, define the function $\tilde{\gamma}(x) = \gamma(x)x^{\eta/2}$. Note that, since γ is an α -power spectral density and since $\tilde{\gamma}$ decays as

$$|\tilde{\gamma}(x)| \lesssim |x|^{-(\alpha-\eta/2)} = |x|^{-\zeta/2},$$

$\tilde{\gamma}$ is a $(\zeta/2)$ -power spectral density. Now, by definition of $\tilde{\gamma}$,

$$\|\mathcal{Z} - \tilde{\mathcal{Z}}_h\|_{L^2(\Omega; L^2(\mathcal{M}))} = \|\tilde{\gamma}(\mathcal{L})\mathcal{L}^{-\eta/2}\mathcal{W} - \tilde{\gamma}(\mathcal{L}_h)\mathcal{L}_h^{-\eta/2}P_h\mathcal{W}\|_{L^2(\Omega; L^2(\mathcal{M}))},$$

so that the triangle inequality yields

$$\|\mathcal{Z} - \tilde{\mathcal{Z}}_h\|_{L^2(\Omega; L^2(\mathcal{M}))} \leq E_1 + E_2,$$

where we take

$$\begin{aligned}E_1 &= \|(\tilde{\gamma}(\mathcal{L}) - \tilde{\gamma}(\mathcal{L}_h)P_h)\mathcal{L}^{-\eta/2}\mathcal{W}\|_{L^2(\Omega; L^2(\mathcal{M}))}, \\ E_2 &= \|\tilde{\gamma}(\mathcal{L}_h)(P_h\mathcal{L}^{-\eta/2}\mathcal{W} - \mathcal{L}_h^{-\eta/2}P_h\mathcal{W})\|_{L^2(\Omega; L^2(\mathcal{M}))}.\end{aligned}$$

For the term E_1 , note that by Bonito et al. (2024, Lemma 4.2),

$$\|\mathcal{L}^{-\eta/2}\mathcal{W}\|_{L^2(\Omega; L^2(\mathcal{M}))} \leq \frac{\eta}{\eta - d/2}.$$

In particular, $\mathcal{L}^{-\eta/2}\mathcal{W}$ is in $L^2(\Omega; L^2(\mathcal{M}))$ and an application of Proposition 2.3 yields that

$$E_1 \lesssim C_\zeta(h)h^{\min\{\zeta; 2\}},$$

where

$$C_\zeta(h) = \begin{cases} 1, & \zeta \neq 2, \\ |\log h|, & \zeta = 2. \end{cases}$$

For E_2 , we first write $P_h \mathcal{L}^{-\eta/2} \mathcal{W} = \sum_{i=1}^{N_h} \beta_i e_i^h$, where $\beta_i = (P_h \mathcal{L}^{-\eta/2} \mathcal{W}, e_i^h)_{L^2(\mathcal{M})} = (\mathcal{L}^{-\eta/2} \mathcal{W}, e_i^h)_{L^2(\mathcal{M})} = \sum_{k=1}^{\infty} \lambda_k^{-\eta/2} W_k(e_k, e_i^h)_{L^2(\mathcal{M})}$. Further, following the definition of $\tilde{\mathcal{W}}$ in Equation (32), we have $\mathcal{L}_h^{-\eta/2} P_h \mathcal{W} = \sum_{i=1}^{N_h} (\lambda_i^h)^{-\eta/2} \xi_i e_i^h$ where $\xi_i = (e_i^h, \mathcal{W})_{L^2(\mathcal{M})} = \sum_{k=1}^{\infty} W_k(e_i^h, e_k)_{L^2(\mathcal{M})}$. Note in particular that both β_i and ξ_i are real-valued (since the eigenbases $\{e_k\}_{k \in \mathbb{N}}$ and $\{e_i^h\}_{1 \leq i \leq N_h}$ are real-valued). Thus,

$$E_2^2 = \sum_{i=1}^{N_h} |\tilde{\gamma}(\lambda_i^h)|^2 \mathbb{E}[|\beta_i - (\lambda_i^h)^{-\eta/2} \xi_i|^2],$$

where we use the orthonormality of the eigenfunctions. Note that

$$\mathbb{E}[|\beta_i - (\lambda_i^h)^{-\eta/2} \xi_i|^2] = \mathbb{E}[|\beta_i|^2] - 2\mathbb{E}[\beta_i (\lambda_i^h)^{-\eta/2} \xi_i] + \mathbb{E}[(\lambda_i^h)^{-\eta} |\xi_i|^2].$$

First, by the independence of the sequence $\{W_k\}_{k=1}^{\infty}$ and (λ_k, e_k) eigenpairs of \mathcal{L} ,

$$\begin{aligned} \mathbb{E}[|\beta_i|^2] &= \sum_{k=1}^{\infty} \lambda_k^{-\eta} |(e_k, e_i^h)_{L^2(\mathcal{M})}|^2 = \left(\sum_{k=1}^{\infty} \lambda_k^{-\eta} (e_i^h, e_k)_{L^2(\mathcal{M})} e_k, e_i^h \right)_{L^2(\mathcal{M})} \\ &= (\mathcal{L}^{-\eta} e_i^h, e_i^h)_{L^2(\mathcal{M})} = \|\mathcal{L}^{-\eta/2} e_i^h\|_{L^2(\mathcal{M})}^2. \end{aligned}$$

Similarly, and using the fact that (λ_i^h, e_i^h) is an eigenpair of \mathcal{L}_h ,

$$\mathbb{E}[(\lambda_i^h)^{-\eta} |\xi_i|^2] = (\lambda_i^h)^{-\eta} \sum_{k=1}^{\infty} |(e_i^h, e_k)_{L^2(\mathcal{M})}|^2 = (\lambda_i^h)^{-\eta} \|e_i^h\|_{L^2(\mathcal{M})}^2 = \|\mathcal{L}_h^{-\eta/2} e_i^h\|_{L^2(\mathcal{M})}^2$$

and

$$\mathbb{E}[\beta_i (\lambda_i^h)^{-\eta/2} \xi_i] = (\lambda_i^h)^{-\eta/2} \sum_{k=1}^{\infty} \lambda_k^{-\eta/2} |(e_k, e_i^h)_{L^2(\mathcal{M})}|^2 = (\mathcal{L}^{-\eta/2} e_i^h, \mathcal{L}_h^{-\eta/2} e_i^h)_{L^2(\mathcal{M})}.$$

This implies that by Proposition 2.3,

$$\mathbb{E}[|\beta_i - (\lambda_i^h)^{-\eta/2} \xi_i|^2] = \|\mathcal{L}^{-\eta/2} e_i^h - \mathcal{L}_h^{-\eta/2} e_i^h\|_{L^2(\mathcal{M})}^2 \lesssim h^{\min\{2\eta; 4\}} C_\eta(h)^2,$$

where

$$C_\eta(h) = \begin{cases} 1, & \eta \neq 2, \\ |\log h|, & \eta = 2. \end{cases}$$

Hence, it holds that

$$E_2^2 \lesssim h^{\min\{2\eta; 4\}} C_\eta(h)^2 \sum_{i=1}^{N_h} \tilde{\gamma}(\lambda_i^h)^2.$$

It remains to bound $\sum_{i=1}^{N_h} \tilde{\gamma}(\lambda_i^h)^2$. To this end, recall Equation (12) and Equation (6) which yield that

$$\sum_{i=1}^{N_h} \tilde{\gamma}(\lambda_i^h)^2 \lesssim \sum_{i=1}^{N_h} (\lambda_i^h)^{-\zeta} \lesssim \sum_{i=1}^{N_h} i^{-2\zeta/d} \lesssim \begin{cases} N_h^{1-2\zeta/d}, & 2\zeta/d < 1, \\ \log(N_h), & 2\zeta/d = 1, \\ 1, & 2\zeta/d > 1, \end{cases} \lesssim \begin{cases} h^{-2(d/2-\zeta)}, & \zeta < d/2, \\ |\log(h)|, & \zeta = d/2, \\ 1, & \zeta > d/2, \end{cases}$$

where we used in the last inequality that $N_h \propto h^{-d}$. We obtain

$$\sum_{i=1}^{N_h} \tilde{\gamma}(\lambda_i^h)^2 \lesssim h^{2(\zeta - \max\{\zeta; d/2\})} K_\zeta(h)^2$$

with

$$K_\zeta(h) = \begin{cases} 1, & \zeta \neq d/2 \\ |\log(h)|^{1/2}, & \zeta = d/2 \end{cases}$$

We conclude that

$$E_2 \lesssim C_\eta(h) K_\zeta(h) h^{\min\{\eta; 2\} + \zeta - \max\{\zeta; d/2\}}$$

and in particular,

$$h^{-\min\{\zeta; 2\}} E_2 \lesssim C_\eta(h) K_\zeta(h) h^\beta,$$

where we take $\beta = \min\{\eta; 2\} + \zeta - \max\{\zeta; d/2\} - \min\{\zeta; 2\}$. We have two cases. On the one hand, if $\zeta \geq 2$ then, since $\eta > \zeta$, we also have $\eta > 2$ and therefore $\beta = 0$, $C_\eta(h) = 1$ and $K_\zeta(h) = 1$ (given that $d/2 < 2$). On the other hand, if $\zeta < 2$, then $\max\{\zeta; d/2\} < 2$ and $\min\{\zeta; 2\} < \eta$ and therefore $\beta = \min\{\eta; 2\} - \max\{\zeta; d/2\} > 0$. In turn, this means that $C_\eta(h) K_\zeta(h) h^\beta \lesssim |\log(h)|^{3/2} h^\beta \lesssim 1$. Hence, in both cases

$$h^{-\min\{\zeta; 2\}} E_2 \lesssim 1,$$

and we conclude that

$$\|\mathcal{Z} - \tilde{\mathcal{Z}}_h\|_{L^2(\Omega; L^2(\mathcal{M}))} \lesssim h^{\min\{\zeta; 2\}} \left(C_\zeta(h) + h^{-\min\{\zeta; 2\}} E_2 \right) \lesssim C_\zeta(h) h^{\min\{\zeta; 2\}}.$$

The result follows by inserting the definition of ζ . \square

For the error between $(\tilde{\mathcal{Z}}_h)^{-\ell}$ and $\widehat{\mathcal{Z}}_h$ we get the following estimate, inspired by Bonito et al. (2024, Lemma 4.4).

Lemma 3.5. *It holds that*

$$\|(\tilde{\mathcal{Z}}_h)^{-\ell} - \widehat{\mathcal{Z}}_h\|_{L^2(\Omega; L^2(\mathcal{M}_h))} \lesssim |\log(h)|^{(d-1)/2} h^2.$$

Proof. Let $\mathcal{E}_h = \|(\tilde{\mathcal{Z}}_h)^{-\ell} - \widehat{\mathcal{Z}}_h\|_{L^2(\Omega; L^2(\mathcal{M}_h))} = \|(\gamma(\mathcal{L}_h) \widetilde{\mathcal{W}})^{-\ell} - \gamma(\mathcal{L}_h) \mathbb{P}_h(\sigma \widetilde{\mathcal{W}}^{-\ell})\|_{L^2(\Omega; L^2(\mathcal{M}_h))}$. We note that $\widetilde{\mathcal{W}}$ is an S_h^ℓ -valued random variable. Therefore, we can apply Proposition 2.5 to realizations of $\widetilde{\mathcal{W}}$, and take the expectation on both sides to get

$$\mathcal{E}_h \lesssim h^2 \|\mathcal{L}_h^{-\min\{\alpha+d/4; 1\}/2} \widetilde{\mathcal{W}}\|_{L^2(\Omega; L^2(\mathcal{M}))}.$$

We then distinguish two cases. First, if $d = 1$, then we note that $\min\{\alpha + d/4; 1\} \geq \alpha + d/4 > d/2$ since $\alpha > d/4$. Besides, Bonito et al. (2024, Lemma 4.2) yields that for all $r \in (d/2, 2)$, there is a constant $C > 0$ such that

$$(35) \quad \|\mathcal{L}_h^{-r/2} \widetilde{\mathcal{W}}\|_{L^2(\Omega; L^2(\mathcal{M}))}^2 \leq C \frac{r}{r - d/2}.$$

Hence, using estimate (35), we conclude that

$$\mathcal{E}_h \lesssim h^2 \frac{\min\{\alpha + d/4; 1\}}{\min\{\alpha + d/4; 1\} - d/2} \lesssim h^2.$$

If now $d = 2$, since $\alpha + d/4 > d/2 = 1$, we have $\min\{\alpha + d/4; 1\} = 1$. We then note that by the proof of Bonito et al. (2024, Lemma 4.2), for any $\varepsilon > 0$

$$\|\mathcal{L}_h^{-1/2} \widetilde{\mathcal{W}}\|_{L^2(\Omega; L^2(\mathcal{M}))}^2 \lesssim \sum_{j=1}^{N_h} \lambda_j^{-1} \lesssim \lambda_{N_h}^\varepsilon \sum_{j=1}^{N_h} \lambda_j^{-1-\varepsilon}.$$

Apply now Equation (6) to see that

$$\|\mathcal{L}_h^{-1/2} \widetilde{\mathcal{W}}\|_{L^2(\Omega; L^2(\mathcal{M}))}^2 \lesssim N_h^\varepsilon \sum_{j=1}^{N_h} j^{-(1+\varepsilon)} \lesssim N_h^\varepsilon \frac{1+\varepsilon}{\varepsilon} \lesssim h^{-2\varepsilon} \frac{1+\varepsilon}{\varepsilon},$$

meaning that for any $\varepsilon > 0$,

$$\mathcal{E}_h \lesssim \left(\frac{1+\varepsilon}{\varepsilon}\right)^{1/2} h^{2-\varepsilon}.$$

In particular, taking $\varepsilon = |\log(h)|^{-1}$ yields

$$\mathcal{E}_h \lesssim |\log(h)|^{1/2} h^2,$$

which concludes the proof. \square

Finally, for the error between $\widehat{\mathbf{Z}}_h$ and \mathbf{Z}_h we get the following estimate.

Lemma 3.6. *It holds that there is a constant $C > 0$ such that*

$$\|\mathbf{Z}_h - \widehat{\mathbf{Z}}_h\|_{L^2(\Omega; L^2(\mathcal{M}_h))} \leq Ch^2.$$

Proof. To prove this statement, we note that, using the same notations as the ones in the proof of Lemma 3.3,

$$\begin{aligned} \|\mathbf{Z}_h - \widehat{\mathbf{Z}}_h\|_{L^2(\Omega; L^2(\mathcal{M}_h))}^2 &= \left\| \gamma(\mathbf{L}_h) \left(\widehat{\mathbf{W}} - \mathbf{W} \right) \right\|_{L^2(\Omega; L^2(\mathcal{M}_h))}^2 \\ &= \mathbb{E} \left[\left\| \sum_{j=1}^{N_h} \gamma(\Lambda_j^h) (\alpha_j - \beta_j) E_j^h \right\|_{L^2(\mathcal{M}_h)}^2 \right] = \sum_{j=1}^{N_h} \gamma(\Lambda_j^h)^2 \mathbb{E}[|\alpha_j - \beta_j|^2], \end{aligned}$$

where we applied the orthogonality of the eigenfunctions in the last step. Now, following the definition of α_j and β_j given in the proof of Lemma 3.3,

$$\begin{aligned} \mathbb{E}[|\alpha_j - \beta_j|^2] &= \mathbb{E} \left[\left\| (\mathcal{W}, ((1 - \sigma^{-1/2}) E_j^h)^\ell) \right\|_{L^2(\mathcal{M})}^2 \right] = \left\| ((1 - \sigma^{-1/2}) E_j^h)^\ell \right\|_{L^2(\mathcal{M})}^2 \\ &= \left\| (\sigma^{1/2} - 1) E_j^h \right\|_{L^2(\mathcal{M}_h)}^2 \leq \left\| (\sigma^{1/2} - 1) \right\|_{L^\infty(\mathcal{M}_h)}^2. \end{aligned}$$

Besides, for all $x \in \mathcal{M}_h$, it holds that

$$\sigma^{1/2}(x) - 1 = \sqrt{1 + \sigma(x) - 1} - 1 \leq \sqrt{1 + |\sigma(x) - 1|} - 1 \leq 1 + \frac{1}{2}|\sigma(x) - 1| - 1 = \frac{1}{2}|\sigma(x) - 1|.$$

Therefore,

$$\left\| (\sigma^{1/2} - 1) \right\|_{L^\infty(\mathcal{M}_h)}^2 \leq \|(\sigma - 1)\|_{L^\infty(\mathcal{M}_h)}^2 \leq Ch^2,$$

where Dziuk and Elliott (2013, Lemma 4.1) was applied in the final step. We conclude that

$$\|Z_h - \widehat{Z}_h\|_{L^2(\Omega; L^2(\mathcal{M}_h))}^2 \leq Ch^4 \sum_{j=1}^{N_h} \gamma(\Lambda_j^h)^2,$$

and it remains to show that $\sum_{j=1}^{N_h} \gamma(\Lambda_j^h)^2$ is bounded by a constant. To this end, we use Lemma 2.2, which implies that there exists some constant $C_\lambda > 0$ such that

$$|\Lambda_j^h / \lambda_j^h - 1| \leq C_\lambda h^2.$$

Recall then that the mesh size h satisfies $h < h_0$, where $h_0 \in (0, 1)$. Without loss of generality, let us further assume that $C_\lambda h_0 < 1$. Now, by the growth assumption on γ ,

$$\begin{aligned} \sum_{j=1}^{N_h} \gamma(\Lambda_j^h)^2 &\lesssim \sum_{j=1}^{N_h} |\Lambda_j^h|^{-2\alpha} = \sum_{j=1}^{N_h} |\lambda_j^h|^{-2\alpha} \left| 1 + \left(\frac{\Lambda_j^h}{\lambda_j^h} - 1 \right) \right|^{-2\alpha} \leq (1 - Ch_0^2)^{-2\alpha} \sum_{j=1}^{N_h} |\lambda_j^h|^{-2\alpha} \\ &\lesssim \sum_{j=1}^{N_h} |\lambda_j^h|^{-2\alpha}. \end{aligned}$$

Using Equation (12) and Equation (6), we bound

$$\sum_{j=1}^{N_h} |\lambda_j^h|^{-2\alpha} \leq \sum_{j=1}^{N_h} |\lambda_j|^{-2\alpha} \lesssim \sum_{j=1}^{\infty} j^{-4\alpha/d} = \zeta(4\alpha/d),$$

where ζ denotes the Riemann zeta function. Thus, $\sum_{j=1}^{N_h} \gamma(\Lambda_j^h)^2$ is bounded by a constant and

$$\|Z_h - \widehat{Z}_h\|_{L^2(\Omega; L^2(\mathcal{M}_h))}^2 \leq Ch^4,$$

which proves the lemma. \square

Equipped with the estimates derived in Lemmas 3.4 to 3.6 we are ready to prove Theorem 3.1.

Proof of Theorem 3.1. By summing the three estimates derived in Lemmas 3.4 to 3.6, we get

$$\|\mathcal{Z} - Z_h^\ell\|_{L^2(\Omega; L^2(\mathcal{M}))} \lesssim C_\alpha(h) h^{\min\{\alpha-d/4; 2\}} + |\log(h)|^{(d-1)/2} h^2.$$

Let then ϵ_h be the quantity defined by $\epsilon_h = (K_\alpha(h) h^{\min\{\alpha-d/4; 2\}})^{-1} \|\mathcal{Z} - Z_h^\ell\|_{L^2(\Omega; L^2(\mathcal{M}))}$, which satisfies

$$\epsilon_h \lesssim C_\alpha(h) K_\alpha(h)^{-1} + |\log(h)|^{(d-1)/2} K_\alpha(h)^{-1} h^{2-\min\{\alpha-d/4; 2\}}.$$

If $\alpha - d/4 \in (0, 2)$, we have $C_\alpha(h) = K_\alpha(h) = 1$ and $2 - \min\{\alpha - d/4; 2\} > 0$, and therefore $\epsilon_h \lesssim 1+$. Similarly, if $\alpha - d/4 = 2$, it holds $C_\alpha(h) = K_\alpha(h) = |\log h|$ and $2 - \min\{\alpha - d/4; 2\} = 0$, and therefore $\epsilon_h \lesssim 1 + |\log(h)|^{(d-3)/2} \lesssim 1$. Finally, for $\alpha - d/4 > 2$, $C_\alpha(h) = 1$,

$K_\alpha(h) = |\log h|^{(d-1)/2}$ and $2 - \min\{\alpha - d/4; 2\} = 0$, and therefore $\epsilon_h \lesssim |\log h|^{-(d-1)/2} + 1 \lesssim 1$. In conclusion, we obtain for any $\alpha > d/4$, $\epsilon_h \lesssim 1$ which in turn yields

$$\|\mathcal{Z} - \mathbf{Z}_h^\ell\|_{L^2(\Omega; L^2(\mathcal{M}))} \lesssim K_\alpha(h) h^{\min\{\alpha - d/4; 2\}}. \quad \square$$

Having bounded the SFEM error, we are now ready to derive the additional error associated with the Chebyshev approximation which we use to compute SFEM–Galerkin approximations in practice.

3.4. Error analysis of the Chebyshev approximation. To bound the error between $\mathbf{Z}_{h,M}$ and \mathbf{Z}_h , we can directly apply Lang and Pereira (2023, Theorem 5.8). We obtain the following result.

Theorem 3.7. *Let $[\lambda_{\min}, \lambda_{\max}] \subset (0, \infty)$ be the interval on which the Chebyshev polynomial approximation $P_{\gamma, M}$ is computed, and let $\xi = \lambda_{\min}/(\lambda_{\max} - \lambda_{\min})$. Then, there exists a constant $C > 0$ such that the error between the discretized field \mathbf{Z}_h and its approximation $\mathbf{Z}_{h,M}$ is upper-bounded by*

$$(36) \quad \|\mathbf{Z}_{h,M} - \mathbf{Z}_h\|_{L^2(\Omega; L^2(\mathcal{M}_h))} \leq Ch^{-d/2} \epsilon_\xi^{-1} (1 + \epsilon_\xi)^{-M}$$

with $\epsilon_\xi = \xi + \sqrt{\xi(2 + \xi)}$. In particular, setting $\lambda_{\min} = V_-$ and $\lambda_{\max} = \Lambda_{N_h}^h$, the error is bounded by

$$(37) \quad \|\mathbf{Z}_{h,M} - \mathbf{Z}_h\|_{L^2(\Omega; L^2(\mathcal{M}_h))} \leq C_V^{-1} h^{-(d/2+1)} \exp(-C_V h^{-1} M)$$

for some constant $C_V > 0$ proportional to $\sqrt{V_-}$.

Proof. Let $E_\xi \subset \mathbb{C}$ be the ellipse centered at $z = (\lambda_{\min} + \lambda_{\max})/2$, with foci $z_1 = \lambda_{\min}$ and $z_2 = \lambda_{\max}$, and semi-major axis $a_\xi = \lambda_{\max}/2$. In particular, note that $E_\xi \subset H_{\pi/2}$ and for any $z \in E_\xi$, $\operatorname{Re}(z) \geq \lambda_{\min}/2 > 0$. Hence, since γ is a power spectral density, by definition γ is holomorphic and bounded inside E_ξ . We can then adapt the same proof as in Lang and Pereira (2023, Theorem 5.8) to obtain the stated proposition. \square

For a fixed mesh size h , the approximation error converges to 0 as the order of the polynomial approximation M goes to infinity. Choosing M as a function of h that grows fast enough then allows us to ensure the convergence of the approximation error as n goes to infinity (Lang and Pereira, 2023, Section 5.2).

REFERENCES

- L. Bandara, M. Nursultanov, and J. Rowlett, Eigenvalue asymptotics for weighted Laplace equations on rough Riemannian manifolds with boundary, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* (2021), 1843–1878.
- D. Boffi, Finite element approximation of eigenvalue problems, *Acta Numer.* **19** (2010), 1–120.
- D. Bolin, K. Kirchner, and M. Kovács, Numerical solution of fractional elliptic stochastic PDEs with spatial white noise, *IMA J. Numer. Anal.* **40** (2020), 1051–1073.
- A. Bonito, A. Demlow, and R. H. Nochetto, Finite element methods for the Laplace–Beltrami operator, *Handbook of Numerical Analysis*, vol. 21, pp. 1–103, Elsevier, 2020.
- A. Bonito, A. Demlow, and J. Owen, A priori error estimates for finite element approximations to eigenvalues and eigenfunctions of the Laplace–Beltrami operator, *SIAM J. Numer. Anal.* **56** (2018), 2963–2988.
- A. Bonito, D. Guignard, and W. Lei, Numerical approximation of Gaussian random fields on closed surfaces, *Comput. Appl. Math.* **In press** (2024).

- V. Borovitskiy, A. Terenin, P. Mostowsky, and M. P. Deisenroth, Matérn Gaussian processes on Riemannian manifolds, *Advances in Neural Information Processing Systems 33*, 2020.
- S. G. Cox and K. Kirchner, Regularity and convergence analysis in Sobolev and Hölder spaces for generalized whittle–matérn fields, *Numer. Math.* **146** (2020), 819–873.
- G. Dziuk, Finite elements for the Beltrami operator on arbitrary surfaces, *Partial differential equations and calculus of variations*, vol. 1357 of *Lecture Notes in Math.*, pp. 142–155, Springer, Berlin, 1988.
- G. Dziuk and C. M. Elliott, Finite element methods for surface PDEs, *Acta Numer.* **22** (2013), 289–396.
- A. A. Farag, *Biomedical Image Analysis: Statistical and Variational Methods*, Cambridge University Press, Cambridge, 2014.
- H. Fujita and T. Suzuki, Evolution problems, *Handbook of Numerical Analysis*, pp. 789–928, Elsevier, 1991.
- B. C. Hall, *Quantum Theory for Mathematicians*, Springer, New York, 2013.
- L. Herrmann, A. Lang, and C. Schwab, Numerical analysis of lognormal diffusions on the sphere, *Stoch PDE: Anal. Comp.* **6** (2018), 1–44.
- E. Jansson, M. Kovács, and A. Lang, Surface finite element approximation of spherical Whittle–Matérn Gaussian random fields, *SIAM J. on Sci. Comput.* **44** (2022), A825–A842.
- A. Lang and M. Pereira, Galerkin–Chebyshev approximation of Gaussian random fields on compact Riemannian manifolds, *BIT Numer. Math.* **63** (2023), 51.
- F. Lindgren, D. Bolin, and H. Rue, The SPDE approach for Gaussian and non-Gaussian fields: 10 years and still running, *Spat. Stat.* **50** (2022), 100599.
- F. Lindgren, H. Rue, and J. Lindström, An explicit link between Gaussian fields and Gaussian Markov random fields: The stochastic partial differential equation approach, *J. R. Stat. Soc. Ser. B Methodol.* **73** (2011), 423–498.
- D. Marinucci and G. Peccati, *Random Fields on the Sphere. Representation, Limit Theorems and Cosmological Applications*, vol. 389 of *London Mathematical Society Lecture Note Series*, Cambridge University Press, Cambridge, 2011.
- M. A. Shubin, *Pseudodifferential Operators and Spectral Theory*, Springer, Berlin, 2001.
- G. Strang and G. Fix, *An Analysis of the Finite Element Method*, Wellesley-Cambridge Press, Wellesley, MA, 2008.
- R. S. Strichartz, Analysis of the Laplacian on the complete Riemannian manifold, *J. Funct. Anal.* **52** (1983), 48–79.
- M. E. Taylor, *Partial Differential Equations I. Basic Theory*, vol. 115 of *Applied Mathematical Sciences*, Springer, New York, 2011.
- H. Triebel, Spaces of Besov–Hardy–Sobolev type on complete Riemannian manifolds, *Ark. Mat.* **24** (1985), 299–337.
- P. Whittle, Stochastic processes in several dimensions, *Bull. Inst. Int. Stat.* **40** (1963), 974–994.
- A. Yagi, *Abstract Parabolic Evolution Equations and Their Applications*, Springer Monographs in Mathematics, Springer, Berlin, 2010.

APPENDIX A. DETERMINISTIC PROOFS

A.1. Proof of Proposition 2.1.

Proof. A standard result in the spectral theory of elliptic operators on compact Riemannian manifolds (see e.g. (Shubin, 2001, Section 8)) ensures that there exists a set of eigenpairs

$\{(\lambda_i, f_i)\}_{i \in \mathbb{N}}$ of \mathcal{L} such that $0 \leq \lambda_1 \leq \lambda_2 \leq \dots$ with $\lambda_i \rightarrow +\infty$ as $i \rightarrow +\infty$, and $\{f_i\}_{i \in \mathbb{N}}$ is an orthonormal basis of $L^2(\mathcal{M})$ composed of possibly complex-valued functions.

Hence, let us prove that $\lambda_1 \geq \delta$ and that we can build an orthonormal basis $\{e_i\}_{i \in \mathbb{N}}$ of $L^2(\mathcal{M})$ such that each e_i is real-valued and an eigenfunction of \mathcal{L} with eigenvalue λ_i . On the one hand, by definition of \mathcal{L} , λ_1 and f_1 , we obtain

$$\lambda_1 = \lambda_1(f_1, f_1)_{L^2(\mathcal{M})} = \mathbf{A}_{\mathcal{M}}(f_1, f_1) \geq \delta \|f_1\|_{H^1(\mathcal{M})}^2 \geq \delta \|f_1\|_{L^2(\mathcal{M})}^2 = \delta,$$

where for the last two inequalities we used the coercivity of $\mathbf{A}_{\mathcal{M}}$ and the definition of the H^1 -norm.

On the other hand, let $\lambda > 0$ be one of the eigenvalues of \mathcal{L} , and $E_\lambda \subset L^2(\mathcal{M})$ the associated eigenspace. Following again the results from (Shubin, 2001, Section 8), we get that $E_\lambda \subset C^\infty(\mathcal{M})$, $\dim E_\lambda < \infty$ and that if $\lambda' \neq \lambda$ is another eigenvalue of \mathcal{L} , then E_λ and $E_{\lambda'}$ are orthogonal. Besides, E_λ is in fact generated by the set $\{f_j\}_{j \in J_\lambda}$, where $J_\lambda = \{i \in \mathbb{N} : \lambda_i = \lambda\}$ is finite (since $\dim E_\lambda < \infty$).

Take then $u \in E_\lambda$. Hence, for any $v \in H^1(\mathcal{M})$, $\mathbf{A}_{\mathcal{M}}(u, v) = \lambda(u, v)_{L^2(\mathcal{M})}$. But also, by definition of $\mathbf{A}_{\mathcal{M}}$,

$$\mathbf{A}_{\mathcal{M}}(\bar{u}, v) = \mathbf{A}_{\mathcal{M}}(\bar{v}, u) = \overline{\mathbf{A}_{\mathcal{M}}(u, \bar{v})} = \overline{\lambda(u, \bar{v})_{L^2(\mathcal{M})}} = \lambda(\bar{u}, v)_{L^2(\mathcal{M})},$$

where we used the fact that \mathcal{D} is a real symmetric matrix and V is real-valued for the first two equalities. Consequently, we also have $\bar{u} \in E_\lambda$, and so, the real-valued functions $\operatorname{Re}(u) = (u + \bar{u})/2$ and $\operatorname{Im}(u) = (u - \bar{u})/2i$ (corresponding to real and imaginary parts of u) are also in E_λ .

Circling back to the orthonormal basis $\{f_j\}_{j \in J_\lambda}$ of E_λ , we consider the set of real-valued functions $F_\lambda = \{\operatorname{Re}(f_j)\}_{j \in J_\lambda} \cup \{\operatorname{Im}(f_j)\}_{j \in J_\lambda} \subset E_\lambda$, and the subspace $V_\lambda \subset E_\lambda$ generated by F_λ . In particular $\dim V_\lambda \leq \dim E_\lambda$. By applying the Gram–Schmidt orthogonalization process to F_λ , we get an orthonormal basis $\{e_k\}_{1 \leq k \leq \dim V_\lambda}$ of V_λ which by construction is composed of real-valued functions (since F_λ is composed of real-valued functions). Let us show that $\{e_k\}_{1 \leq k \leq \dim V_\lambda}$ is in fact a basis of E_λ , or equivalently that $\dim V_\lambda = \dim E_\lambda$.

We proceed by contradiction. Assume that $\dim V_\lambda < \dim E_\lambda$. This means in particular that the orthogonal complement of V_λ in E_λ , denoted by V_λ^\perp , is not reduced to 0. Let then $0 \neq w \in V_\lambda^\perp$. By linearity, we have, for any $j \in J_\lambda$, $(f_j, w)_{L^2(\mathcal{M})} = (\operatorname{Re}(f_j), w)_{L^2(\mathcal{M})} + i(\operatorname{Im}(f_j), w)_{L^2(\mathcal{M})} = 0$, since $\operatorname{Re}(f_j), \operatorname{Im}(f_j) \in F_\lambda \subset V_\lambda$. Hence, since $w \in E_\lambda$ and $\{f_j\}_{j \in J_\lambda}$ is an orthonormal basis of E_λ , it must hold that $w = 0$, which contradicts our initial claim. Consequently, $\dim V_\lambda = \dim E_\lambda$, and therefore $\{e_k\}_{1 \leq k \leq \dim V_\lambda}$ is an orthonormal basis of E_λ .

Finally, by repeating the construction above to each eigenspace E_λ associated with distinct eigenvalues, and concatenating the obtained bases, we obtain an orthonormal basis $L^2(\mathcal{M})$ (due to the fact that these eigenspaces are orthogonal to one another and span $L^2(\mathcal{M})$). Each element in this basis is an eigenfunction of \mathcal{L} since it is built from a given eigenspace, and is a real-valued function. This concludes our proof. \square

A.2. Proof of Lemma 2.4.

Proof. Let us start by proving the following claims: for any $z \in \Gamma$, $f \in L^2(\mathcal{M})$,

$$(38) \quad \|(z - \mathcal{L}_h)^{-1} P_h f - (z - \mathcal{L})^{-1} f\|_{L^2(\mathcal{M})} \leq Ch^2,$$

$$(39) \quad \|(z - \mathcal{L}_h)^{-1} P_h f - (z - \mathcal{L})^{-1} f\|_{L^2(\mathcal{M})} \leq \frac{C}{|z|}.$$

We omit the proof of Equation (39) and refer the reader to Fujita and Suzuki (1991, Theorem 5.1), where the proof may be adapted verbatim.

To prove Equation (38), we can largely proceed as in Fujita and Suzuki (1991, Theorem 7.1), substituting the Euclidean elliptic regularity estimates and interpolation bounds with their surface counterparts, see for instance Taylor (2011, Chapter 11), Dziuk and Elliott (2013, Lemma 4.3) and Bonito et al. (2024, Equation (4.8)). We must in essence only adapt Fujita and Suzuki (1991, Lemma 7.1) to also hold for $z \in \Gamma_0$, as their estimate only holds for $z \in \partial G_\theta$. In other words, we must show that for any $z = \delta_0 e^{it}$, where $t \in [-\theta, \theta]$, there is a constant $\tilde{\delta} > 0$ such that for all $v \in H^1(\mathcal{M})$,

$$(40) \quad |z| \|v\|_{L^2(\mathcal{M})}^2 + \|v\|_{H^1(\mathcal{M})}^2 \leq \tilde{\delta}^{-1} \left| z \|v\|_{L^2(\mathcal{M})}^2 - \mathbf{A}_{\mathcal{M}}(v, v) \right|.$$

To prove this inequality, note that by coercivity of $\mathbf{A}_{\mathcal{M}}$,

$$\mathbf{A}_{\mathcal{M}}(v, v) - \delta_0 \|v\|_{L^2(\mathcal{M})}^2 \geq \delta \|v\|_{H^1(\mathcal{M})}^2 - \delta_0 \|v\|_{L^2(\mathcal{M})}^2.$$

Let then $\tilde{\delta} = \frac{\delta_0}{1+\delta_0}$. Since the only requirement on δ_0 was that $\delta_0 < \delta/2$, we can then write

$$\begin{aligned} \mathbf{A}_{\mathcal{M}}(v, v) - \delta_0 \|v\|_{L^2(\mathcal{M})}^2 &\geq 2\delta_0 \|v\|_{H^1(\mathcal{M})}^2 - \delta_0 \|v\|_{L^2(\mathcal{M})}^2 = 2\tilde{\delta}(1+\delta_0) \|v\|_{H^1(\mathcal{M})}^2 - \tilde{\delta}(1+\delta_0) \|v\|_{L^2(\mathcal{M})}^2 \\ &= 2\tilde{\delta}(1+\delta_0) \|v\|_{H^1(\mathcal{M})}^2 + \left(\delta_0 \tilde{\delta} - (2\delta_0 + 1)\tilde{\delta} \right) \|v\|_{L^2(\mathcal{M})}^2 \\ &\geq 2\tilde{\delta}(1+\delta_0) \|v\|_{H^1(\mathcal{M})}^2 - (2\delta_0 + 1)\tilde{\delta} \|v\|_{H^1(\mathcal{M})}^2 + \delta_0 \tilde{\delta} \|v\|_{L^2(\mathcal{M})}^2 \\ &= \tilde{\delta} \left(\|v\|_{H^1(\mathcal{M})}^2 + \delta_0 \|v\|_{L^2(\mathcal{M})}^2 \right) = \tilde{\delta} \left(\|v\|_{H^1(\mathcal{M})}^2 + |z| \|v\|_{L^2(\mathcal{M})}^2 \right), \end{aligned}$$

since $|z| = \delta_0$ if $z \in \Gamma_0$. Therefore,

$$\|v\|_{H^1(\mathcal{M})}^2 + |z| \|v\|_{L^2(\mathcal{M})}^2 \leq \tilde{\delta}^{-1} \left(\mathbf{A}_{\mathcal{M}}(v, v) - \delta_0 \|v\|_{L^2(\mathcal{M})}^2 \right).$$

This last inequality implies in particular that $\mathbf{A}_{\mathcal{M}}(v, v) - \delta_0 \|v\|_{L^2(\mathcal{M})}^2 \geq 0$ and therefore that

$$\|v\|_{H^1(\mathcal{M})}^2 + |z| \|v\|_{L^2(\mathcal{M})}^2 \leq \tilde{\delta}^{-1} \left| \mathbf{A}_{\mathcal{M}}(v, v) - \delta_0 \|v\|_{L^2(\mathcal{M})}^2 \right|.$$

Using once again that $|z| = \delta_0$ and that $\mathbf{A}_{\mathcal{M}}(v, v) \geq 0$, we deduce that

$$\|v\|_{H^1(\mathcal{M})}^2 + |z| \|v\|_{L^2(\mathcal{M})}^2 \leq \tilde{\delta}^{-1} \left| \mathbf{A}_{\mathcal{M}}(v, v) \right| - |z| \|v\|_{L^2(\mathcal{M})}^2 \leq \tilde{\delta}^{-1} \left| \mathbf{A}_{\mathcal{M}}(v, v) - z \|v\|_{L^2(\mathcal{M})}^2 \right|$$

by the reverse triangle inequality. To conclude, we have proven that the relation (40) holds and equipped with this estimate, we can proceed as in the proof of Fujita and Suzuki (1991, Theorem 7.1) and obtain Equation (38).

Finally, by interpolating between the bounds obtained in Equations (38) and (39), we obtain that, for any $\beta \in [0, 1]$,

$$\|(z - \mathcal{L}_h)^{-1} P_h f - (z - \mathcal{L})^{-1} f\|_{L^2(\mathcal{M})} \leq C \frac{h^{2\beta}}{|z|^{1-\beta}}. \quad \square$$

APPENDIX B. ERROR BETWEEN FUNCTIONS OF DISCRETE OPERATORS

The aim of this section is to prove Proposition 2.5. To do so, we rely on a geometric consistency estimate, and on extensions of the results of Bonito et al. (2024, Lemma A.1), which we start by stating and proving.

B.1. Geometric consistency estimate. The following geometric consistency estimate quantifies the error between the bilinear forms $\mathbf{A}_{\mathcal{M}}$ and $\mathbf{A}_{\mathcal{M}_h}$. Its proof is a straightforward adaptation of the proof of Dziuk and Elliott (2013, Lemma 4.7) to account for the diffusion matrix \mathcal{D} .

Lemma B.1. *There is a constant $C > 0$ such that for all $h < h_0$ and $u_h, v_h \in S_h^\ell$,*

$$(41) \quad \left| \mathbf{A}_{\mathcal{M}}(u_h^\ell, v_h^\ell) - \mathbf{A}_{\mathcal{M}_h}(u_h, v_h) \right| \leq Ch^2 \|u_h^\ell\|_{H^1(\mathcal{M})} \|v_h^\ell\|_{H^1(\mathcal{M})}.$$

Proof. We first note that, by definition of \mathcal{D} , for any $x_0 \in \mathcal{M}$, and any $w, w' \in T_{x_0}\mathcal{M}$, $(\mathcal{D}(x_0)w) \cdot \overline{w'}$ defines an inner product on $T_{x_0}\mathcal{M}$. We denote by $\|\cdot\|$ the usual Euclidean norm of vectors of $T_{x_0}\mathcal{M} \subset \mathbb{C}^{d+1}$ and by $\|\cdot\|_{\mathcal{D}(x_0)}$ the norm defined by $\|w\|_{\mathcal{D}(x_0)}^2 = (\mathcal{D}(x_0)w) \cdot \overline{w}$, $w \in T_{x_0}\mathcal{M}$.

Let $\Pi = I - \nu\nu^T$ (resp. $\Pi_h = I - \nu_h\nu_h^T$) be the orthogonal projection onto the tangent planes of \mathcal{M} (resp. \mathcal{M}_h), and let $\mathcal{H} : \mathcal{M} \rightarrow \mathbb{R}^{(d+1) \times (d+1)}$ be the extended Weingarten map of \mathcal{M} (cf. Dziuk and Elliott (2013, Definition 2.5)). Recall in particular that $\mathcal{H}(x)\nu(x) = 0$ for any $x \in \mathcal{M}$, meaning in particular that $\mathcal{H}\Pi = \Pi\mathcal{H} = \mathcal{H}$. Finally, we introduce the map $\mathcal{Q}_h : \mathcal{M} \rightarrow \mathbb{R}^{(d+1) \times (d+1)}$ defined as

$$\mathcal{Q}_h = \frac{1}{\sigma^\ell} \Pi(I - d_s^\ell \mathcal{H}) \Pi_h^\ell \mathcal{D} \Pi_h^\ell (I - d_s^\ell \mathcal{H}) \Pi,$$

where d_s is the oriented distance function restricted to \mathcal{M}_h and introduced in Section 2.2. On the one hand, note that for any $u_h, v_h \in S_h$,

$$\begin{aligned} (\mathcal{D}^{-\ell} \nabla_{\mathcal{M}_h} u_h) \cdot \nabla_{\mathcal{M}_h} \overline{v_h} &= (\mathcal{D}^{-\ell} \Pi_h (I - d_s \mathcal{H}^{-\ell}) \Pi^{-\ell} (\nabla_{\mathcal{M}} u_h^\ell)^{-\ell}) \cdot (\Pi_h (I - d_s \mathcal{H}^{-\ell}) \Pi^{-\ell} (\nabla_{\mathcal{M}} \overline{v_h}^\ell)^{-\ell}) \\ &= \sigma (\mathcal{Q}_h^{-\ell} (\nabla_{\mathcal{M}} u_h^\ell)^{-\ell}) \cdot (\nabla_{\mathcal{M}} \overline{v_h}^\ell)^{-\ell}, \end{aligned}$$

which gives, after integrating both sides over \mathcal{M}_h and using Equation (9),

$$(42) \quad \int_{\mathcal{M}_h} (\mathcal{D}^{-\ell} \nabla_{\mathcal{M}_h} u_h) \cdot (\nabla_{\mathcal{M}_h} \overline{v_h}) \, dA_h = \int_{\mathcal{M}} (\mathcal{Q}_h \nabla_{\mathcal{M}} u_h^\ell) \cdot (\nabla_{\mathcal{M}} \overline{v_h}^\ell) \, dA.$$

Let then $\mathbf{A}_{\mathcal{M}_h}^\ell : S_h^\ell \times S_h^\ell \rightarrow \mathbb{R}$ be the Hermitian form defined for any $u_h^\ell, v_h^\ell \in S_h^\ell$ by

$$(43) \quad \mathbf{A}_{\mathcal{M}_h}^\ell(u_h^\ell, v_h^\ell) = \int_{\mathcal{M}} (\mathcal{Q}_h \nabla_{\mathcal{M}} u_h^\ell) \cdot (\nabla_{\mathcal{M}} \overline{v_h}^\ell) \, dA + \int_{\mathcal{M}} (\sigma^\ell)^{-1} V u_h^\ell \overline{v_h}^\ell \, dA.$$

Note that following Equations (9) and (42), $\mathbf{A}_{\mathcal{M}_h}^\ell$ satisfies for any $u_h, v_h \in S_h$ the equality $\mathbf{A}_{\mathcal{M}_h}^\ell(u_h^\ell, v_h^\ell) = \mathbf{A}_{\mathcal{M}_h}(u_h, v_h)$. Therefore, for any $u_h, v_h \in S_h$, we bound

$$(44) \quad \begin{aligned} |\mathbf{A}_{\mathcal{M}}(u_h^\ell, v_h^\ell) - \mathbf{A}_{\mathcal{M}_h}(u_h, v_h)| &= |\mathbf{A}_{\mathcal{M}}(u_h^\ell, v_h^\ell) - \mathbf{A}_{\mathcal{M}_h}^\ell(u_h^\ell, v_h^\ell)| \\ &\leq \left| \int_{\mathcal{M}} ((\mathcal{Q}_h - \mathcal{D}) \nabla_{\mathcal{M}} u_h^\ell) \cdot (\nabla_{\mathcal{M}} \overline{v_h}^\ell) \, dA \right| + \left| \int_{\mathcal{M}} (1 - (\sigma^\ell)^{-1}) V u_h^\ell \overline{v_h}^\ell \, dA \right|. \end{aligned}$$

We now bound these two terms. Recall that Dziuk and Elliott (2013, Lemma 4.1) shows

$$(45) \quad \|\sigma\|_{L^\infty(\mathcal{M}_h)} \lesssim 1, \quad \|\sigma^{-1}\|_{L^\infty(\mathcal{M}_h)} \lesssim 1, \quad \|\sigma - 1\|_{L^\infty(\mathcal{M}_h)} \lesssim h^2, \quad \|\sigma^{-1} - 1\|_{L^\infty(\mathcal{M}_h)} \lesssim h^2.$$

Hence, since V takes positive values,

$$\begin{aligned} \left| \int_{\mathcal{M}} (1 - (\sigma^\ell)^{-1}) V u_h^\ell \overline{v_h}^\ell \, dA \right| &\leq \int_{\mathcal{M}} |1 - (\sigma^\ell)^{-1}| V |u_h^\ell| |v_h^\ell| \, dA \\ &\leq \|1 - \sigma^{-1}\|_{L^\infty(\mathcal{M}_h)} \int_{\mathcal{M}} V |u_h^\ell| |v_h^\ell| \, dA, \end{aligned}$$

which in turn gives (using the Cauchy–Schwartz inequality and Equation (45)),

$$(46) \quad \left| \int_{\mathcal{M}} (1 - (\sigma^\ell)^{-1}) V u_h^\ell \bar{v}_h^\ell \, dA \right| \lesssim h^2 \left(\int_{\mathcal{M}} V |u_h^\ell|^2 \, dA \right)^{1/2} \left(\int_{\mathcal{M}} V |v_h^\ell|^2 \, dA \right)^{1/2}.$$

To bound the other term, we first introduce for any $B \in \mathbb{R}^{(d+1) \times (d+1)}$ the notation $\|B\| = \sup_{\|x\|=1} \|Bx\|$. Then we have

$$\begin{aligned} \|\mathcal{Q}_h - \mathcal{D}\| &= \|(\sigma^\ell)^{-1} (\Pi(I - d_s^\ell \mathcal{H}) \Pi_h^\ell \mathcal{D} \Pi_h^\ell (I - d_s^\ell \mathcal{H}) \Pi - \mathcal{D}) + ((\sigma^\ell)^{-1} - 1) \mathcal{D}\| \\ &\leq \|(\sigma^\ell)^{-1}\|_{L^\infty(\mathcal{M})} \|\Pi(I - d_s^\ell \mathcal{H}) \Pi_h^\ell \mathcal{D} \Pi_h^\ell (I - d_s^\ell \mathcal{H}) \Pi - \mathcal{D}\| \\ &\quad + \|(\sigma^\ell)^{-1} - 1\|_{L^\infty(\mathcal{M})} \|\mathcal{D}\|. \end{aligned}$$

By Equation (45) and since \mathcal{D} has bounded eigenvalues over \mathcal{M} , we obtain

$$(47) \quad \|\mathcal{Q}_h - \mathcal{D}\| \lesssim \|\Pi(I - d_s^\ell \mathcal{H}) \Pi_h^\ell \mathcal{D} \Pi_h^\ell (I - d_s^\ell \mathcal{H}) \Pi - \mathcal{D}\| + h^2,$$

where the constant in the inequality is independent of the location on \mathcal{M} . We split the first term on the right into

$$\begin{aligned} &\|\Pi(I - d_s^\ell \mathcal{H}) \Pi_h^\ell \mathcal{D} \Pi_h^\ell (I - d_s^\ell \mathcal{H}) \Pi - \mathcal{D}\| \\ &= \|\Pi \Pi_h^\ell \mathcal{D} \Pi_h^\ell \Pi - \mathcal{D} - \Pi \Pi_h^\ell \mathcal{D} \Pi_h^\ell d_s^\ell \mathcal{H} \Pi - d_s^\ell \mathcal{H} \Pi_h^\ell \mathcal{D} \Pi_h^\ell (I - d_s^\ell \mathcal{H}) \Pi\| \\ &\leq \|\Pi \Pi_h^\ell \mathcal{D} \Pi_h^\ell \Pi - \mathcal{D}\| + \|\Pi \Pi_h^\ell \mathcal{D} \Pi_h^\ell d_s^\ell \mathcal{H} \Pi\| + \|d_s^\ell \mathcal{H} \Pi_h^\ell \mathcal{D} \Pi_h^\ell (I - d_s^\ell \mathcal{H}) \Pi\|. \end{aligned}$$

Since $\|d_s\|_{L^\infty(\mathcal{M}_h)} \lesssim h^2$ by Dziuk and Elliott (2013, Lemma 4.1) and \mathcal{H} is defined independently of h , we conclude that

$$(48) \quad \|\Pi(I - d_s^\ell \mathcal{H}) \Pi_h^\ell \mathcal{D} \Pi_h^\ell (I - d_s^\ell \mathcal{H}) \Pi - \mathcal{D}\| \lesssim \|\Pi \Pi_h^\ell \mathcal{D} \Pi_h^\ell \Pi - \mathcal{D}\| + h^2.$$

We notice that $\mathcal{D} = \Pi \mathcal{D} \Pi$, since by definition of \mathcal{D} , $\mathcal{D}v = 0$, which implies

$$\begin{aligned} \|\Pi \Pi_h^\ell \mathcal{D} \Pi_h^\ell \Pi - \mathcal{D}\| &= \|\Pi \Pi_h^\ell \Pi \mathcal{D} \Pi_h^\ell \Pi - \Pi \mathcal{D} \Pi\| = \|(\Pi \Pi_h^\ell \Pi - \Pi) \mathcal{D} \Pi_h^\ell \Pi + \Pi \mathcal{D} (\Pi \Pi_h^\ell \Pi - \Pi)\| \\ &\leq \|\Pi \Pi_h^\ell \Pi - \Pi\| \|\mathcal{D} \Pi_h^\ell \Pi\| + \|\Pi \mathcal{D}\| \|\Pi \Pi_h^\ell \Pi - \Pi\|. \end{aligned}$$

Using that $\|\Pi \Pi_h^\ell \Pi - \Pi\| \lesssim h^2$ by the proof of Dziuk and Elliott (2013, Lemma 4.1), we deduce that $\|\Pi \Pi_h^\ell \mathcal{D} \Pi_h^\ell \Pi - \mathcal{D}\| \lesssim h^2$. Injecting this inequality into Equation (48), and the resulting inequality into Equation (47), we conclude that

$$\|\mathcal{Q}_h - \mathcal{D}\| \lesssim h^2.$$

This allows us to write

$$\begin{aligned} \left| \int_{\mathcal{M}} ((\mathcal{Q}_h - \mathcal{D}) \nabla_{\mathcal{M}} u_h^\ell) \cdot (\nabla_{\mathcal{M}} \bar{v}_h^\ell) \, dA \right| &\leq \int_{\mathcal{M}} \|(\mathcal{Q}_h - \mathcal{D}) \nabla_{\mathcal{M}} u_h^\ell\| \|\nabla_{\mathcal{M}} v_h^\ell\| \, dA \\ &\lesssim \int_{\mathcal{M}} h^2 \|\nabla_{\mathcal{M}} u_h^\ell\| \|\nabla_{\mathcal{M}} v_h^\ell\| \, dA \leq h^2 \int_{\mathcal{M}} (\mu_{\min})^{-1} \|\nabla_{\mathcal{M}} u_h^\ell\|_{\mathcal{D}} \|\nabla_{\mathcal{M}} v_h^\ell\|_{\mathcal{D}} \, dA, \end{aligned}$$

where $\mu_{\min} : \mathcal{M} \rightarrow \mathbb{R}_+$ maps any $x \in \mathcal{M}$ to the smallest eigenvalue of $\mathcal{D}(x)$ associated with an eigenvector in ν^\perp . This last inequality is a consequence of the fact that by construction $\nabla_{\mathcal{M}} u_h^\ell, \nabla_{\mathcal{M}} v_h^\ell \in \nu^\perp$ and using the characterization of eigenvalues through Rayleigh quotients. Since the non-zero eigenvalues of \mathcal{D} are uniformly bounded above and below by positive constants, we conclude that

$$\left| \int_{\mathcal{M}} ((\mathcal{Q}_h - \mathcal{D}) \nabla_{\mathcal{M}} u_h^\ell) \cdot (\nabla_{\mathcal{M}} \bar{v}_h^\ell) \, dA \right| \lesssim h^2 \int_{\mathcal{M}} \|\nabla_{\mathcal{M}} u_h^\ell\|_{\mathcal{D}} \|\nabla_{\mathcal{M}} v_h^\ell\|_{\mathcal{D}} \, dA.$$

Then, using the Cauchy–Schwartz inequality yields

(49)

$$\left| \int_{\mathcal{M}} ((\mathcal{Q}_h - \mathcal{D})\nabla_{\mathcal{M}} u_h^\ell) \cdot (\nabla_{\mathcal{M}} \bar{v}_h^\ell) \, dA \right| \lesssim h^2 \left(\int_{\mathcal{M}} \|\nabla_{\mathcal{M}} u_h^\ell\|_{\mathcal{D}}^2 \, dA \right)^{1/2} \left(\int_{\mathcal{M}} \|\nabla_{\mathcal{M}} v_h^\ell\|_{\mathcal{D}}^2 \, dA \right)^{1/2}.$$

Inserting the derived bounds Equation (46) and Equation (49) into Equation (44), we derive

$$\begin{aligned} |\mathbf{A}_{\mathcal{M}}(u_h^\ell, v_h^\ell) - \mathbf{A}_{\mathcal{M}_h}(u_h, v_h)| &\lesssim h^2 \left(\int_{\mathcal{M}} \|\nabla_{\mathcal{M}} u_h^\ell\|_{\mathcal{D}}^2 \, dA \right)^{1/2} \left(\int_{\mathcal{M}} \|\nabla_{\mathcal{M}} v_h^\ell\|_{\mathcal{D}}^2 \, dA \right)^{1/2} \\ &\quad + h^2 \left(\int_{\mathcal{M}} V |u_h^\ell|^2 \, dA \right)^{1/2} \left(\int_{\mathcal{M}} V |v_h^\ell|^2 \, dA \right)^{1/2}. \end{aligned}$$

Note that for any $u \in H^1(\mathcal{M})$,

$$\mathbf{A}_{\mathcal{M}}(u, u) \geq \int_{\mathcal{M}} V |u|^2 \, dA, \quad \text{and} \quad \mathbf{A}_{\mathcal{M}}(u, u) \geq \int_{\mathcal{M}} \mathcal{D} \nabla_{\mathcal{M}} u \cdot \nabla_{\mathcal{M}} \bar{u} \, dA = \int_{\mathcal{M}} \|\nabla_{\mathcal{M}} u\|_{\mathcal{D}}^2 \, dA,$$

so we obtain

$$\left| \mathbf{A}_{\mathcal{M}}(u_h^\ell, v_h^\ell) - \mathbf{A}_{\mathcal{M}_h}^\ell(u_h^\ell, v_h^\ell) \right| \lesssim h^2 \sqrt{\mathbf{A}_{\mathcal{M}}(u_h^\ell, u_h^\ell)} \sqrt{\mathbf{A}_{\mathcal{M}}(v_h^\ell, v_h^\ell)}.$$

Finally, due to Equation (5),

$$\sqrt{\mathbf{A}_{\mathcal{M}}(u_h^\ell, u_h^\ell)} \sqrt{\mathbf{A}_{\mathcal{M}}(v_h^\ell, v_h^\ell)} \lesssim \|u_h^\ell\|_{H^1(\mathcal{M})} \|v_h^\ell\|_{H^1(\mathcal{M})},$$

and the result follows. \square

B.2. Norm and error estimates of discrete operators. We now prove some estimates for the norm of shifted inverses of the operators \mathcal{L}_h and \mathbf{L}_h , and for the error between inverses of these two operators. These results can be seen as extensions of the ones stated in Bonito et al. (2024, Lemma A.1).

Lemma B.2. *Let $v_h \in S_h^\ell$ and $V_h \in S_h$ be arbitrary. Then, for all $z \in \Gamma$, for any $q \in [-1, 1]$, and any $r \in (0, 2)$,*

$$(50) \quad \|\mathcal{L}_h^{1/2}(z - \mathcal{L}_h)^{-1} v_h\|_{L^2(\mathcal{M})} \lesssim |z|^{-(1+q)/2} \|\mathcal{L}_h^{q/2} v_h\|_{L^2(\mathcal{M})},$$

$$(51) \quad \|\mathbf{L}_h(z - \mathbf{L}_h)^{-1} V_h\|_{L^2(\mathcal{M}_h)} \lesssim |z|^{-1/2} \|V_h\|_{H^1(\mathcal{M}_h)},$$

$$(52) \quad \|(z - \mathcal{L}_h)^{-1} v_h\|_{L^2(\mathcal{M})} \lesssim |z|^{r/2-1} \|\mathcal{L}_h^{-r/2} v_h\|_{L^2(\mathcal{M})},$$

$$(53) \quad \left\| (\mathcal{L}_h^{-1} v_h)^{-\ell} - \mathbf{L}_h^{-1} \mathbf{P}_h(\sigma v_h^{-\ell}) \right\|_{H^1(\mathcal{M}_h)} \lesssim h^2 \|\mathcal{L}_h^{-1/2} v_h\|_{L^2(\mathcal{M})}.$$

Proof. We start with the proof of Equation (50). To this end, let $z \in \Gamma$ and $q \in [-1, 1]$. For any $s \in [0, 1]$ and $v_h \in S_h^\ell$, we expand

$$\mathcal{L}_h^s v_h = \sum_{i=1}^{N_h} (\lambda_i^h)^s (v_h, e_i^h)_{L^2(\mathcal{M})} e_i^h,$$

implying that

$$\|\mathcal{L}_h^s v_h\|_{L^2(\mathcal{M})}^2 = \sum_{i=1}^{N_h} |\lambda_i^h|^{2s} |(v_h, e_i^h)_{L^2(\mathcal{M})}|^2 = \sum_{i=1}^{N_h} \left(|\lambda_i^h| |(v_h, e_i^h)_{L^2(\mathcal{M})}| \right)^{2s} |(v_h, e_i^h)_{L^2(\mathcal{M})}|^{2(1-s)}.$$

Hölder's inequality yields

$$(54) \quad \begin{aligned} \|\mathcal{L}_h^s v_h\|_{L^2(\mathcal{M})}^2 &\leq \left(\sum_{i=1}^{N_h} |\lambda_i^h|^2 |(v_h, e_i^h)_{L^2(\mathcal{M})}|^2 \right)^s \left(\sum_{i=1}^{N_h} |(v_h, e_i^h)_{L^2(\mathcal{M})}|^2 \right)^{1-s} \\ &= \|\mathcal{L}_h v_h\|_{L^2(\mathcal{M})}^{2s} \|v_h\|_{L^2(\mathcal{M})}^{2(1-s)}. \end{aligned}$$

Let $w_h = (z - \mathcal{L}_h)^{-1} v_h$, then we obtain by Equation (18) that

$$\|w_h\|_{L^2(\mathcal{M})} \lesssim |z|^{-1} \|v_h\|_{L^2(\mathcal{M})}.$$

Hence, adding and subtracting $z(z - \mathcal{L}_h)^{-1}$ yields

$$\begin{aligned} \|\mathcal{L}_h w_h\|_{L^2(\mathcal{M})} &= \|z(z - \mathcal{L}_h)^{-1} v_h - v_h\|_{L^2(\mathcal{M})} \leq |z| \|(z - \mathcal{L}_h)^{-1} v_h\|_{L^2(\mathcal{M})} + \|v_h\|_{L^2(\mathcal{M})} \\ &\lesssim 2\|v_h\|_{L^2(\mathcal{M})}. \end{aligned}$$

Combining this estimate with Equation (54), we get for any $s \in [0, 1]$

$$(55) \quad \|\mathcal{L}_h^s w_h\|_{L^2(\mathcal{M})} \leq \|\mathcal{L}_h w_h\|_{L^2(\mathcal{M})}^s \|w_h\|_{L^2(\mathcal{M})}^{(1-s)} \leq |z|^{-(1-s)} \|v_h\|_{L^2(\mathcal{M})}^s \|v_h\|_{L^2(\mathcal{M})}^{1-s} = |z|^{s-1} \|v_h\|_{L^2(\mathcal{M})},$$

yielding for any $q \in [-1, 1]$ that

$$(56) \quad \begin{aligned} \|\mathcal{L}_h^{1/2} (z - \mathcal{L}_h)^{-1} v_h\|_{L^2(\mathcal{M})} &= \|\mathcal{L}_h^{1-(1+q)/2} (z - \mathcal{L}_h)^{-1} \mathcal{L}_h^{q/2} v_h\|_{L^2(\mathcal{M})} \\ &\lesssim |z|^{-(1+q)/2} \|\mathcal{L}_h^{q/2} v_h\|_{L^2(\mathcal{M})}, \end{aligned}$$

where we applied Equation (55) with $s = 1 - (1 + q)/2$ in the last step. Hence, we retrieve Equation (50).

Next, to prove Equation (51), we first observe that the estimates in Equation (55) and Equation (56) carry over to the case when \mathbf{L}_h is used instead of \mathcal{L}_h , as the proof in essence is a standard manipulation of a finite eigenexpansion. In particular, we obtain (by taking $q = 1$ and $s = 0$)

$$(57) \quad \|\mathbf{L}_h (z - \mathbf{L}_h)^{-1} V_h\|_{L^2(\mathcal{M}_h)} \lesssim |z|^{-1/2} \|\mathbf{L}_h^{1/2} V_h\|_{L^2(\mathcal{M}_h)}.$$

Now, Equation (57) combined with the equivalence of norms in Equation (17), yields that

$$\|\mathbf{L}_h (z - \mathbf{L}_h)^{-1} V_h\|_{L^2(\mathcal{M}_h)} \lesssim |z|^{-1/2} \|V_h\|_{H^1(\mathcal{M}_h)},$$

which shows Equation (51).

To bound Equation (52), we apply Equation (55) with $s = r/2$ to obtain

$$\|(z - \mathcal{L}_h)^{-1} v_h\|_{L^2(\mathcal{M})} = \|\mathcal{L}_h^{r/2} (z - \mathcal{L}_h)^{-1} \mathcal{L}_h^{-r/2} v_h\|_{L^2(\mathcal{M})} \lesssim |z|^{r/2-1} \|\mathcal{L}_h^{-r/2} v_h\|_{L^2(\mathcal{M})}.$$

Finally, to prove the bound in Equation (53), we rely on the geometric consistency estimate of Lemma B.1. Let $u_h = \mathcal{L}_h^{-1} v_h$ and let $U_h = \mathbf{L}_h^{-1} \mathbf{P}_h(\sigma v_h^{-\ell})$. Note that by definition of \mathcal{L}_h^{-1} ,

$$(\mathcal{L}_h u_h, w_h)_{L^2(\mathcal{M})} = \mathbf{A}_{\mathcal{M}}(u_h, w_h) = (v_h, w_h)_{L^2(\mathcal{M})},$$

for all $w_h \in S_h^\ell$. Likewise, for \mathbf{L}_h^{-1} we obtain

$$(\mathbf{L}_h U_h, W_h)_{L^2(\mathcal{M}_h)} = \mathbf{A}_{\mathcal{M}_h}(U_h, W_h) = (\mathbf{P}_h(\sigma v_h^{-\ell}), W_h)_{L^2(\mathcal{M}_h)} = (\sigma v_h^{-\ell}, W_h)_{L^2(\mathcal{M}_h)} = (v_h, W_h^\ell)_{L^2(\mathcal{M})},$$

for all $W_h \in S_h$, where we used the definition of σ in the last step. Let us now select a fixed, but arbitrary, $\Xi_h \in S_h$. Then, by combining the last two equations,

$$\begin{aligned} |\mathbf{A}_{\mathcal{M}_h}(u_h^{-\ell} - U_h, \Xi_h)| &= |\mathbf{A}_{\mathcal{M}_h}(u_h^{-\ell}, \Xi_h) - \mathbf{A}_{\mathcal{M}_h}(U_h, \Xi_h)| = |\mathbf{A}_{\mathcal{M}_h}(u_h^{-\ell}, \Xi_h) - (v_h, \Xi_h^{\ell})_{L^2(\mathcal{M})}| \\ &= |\mathbf{A}_{\mathcal{M}_h}(u_h^{-\ell}, \Xi_h) - \mathbf{A}_{\mathcal{M}}(u_h, \Xi_h^{\ell})|, \end{aligned}$$

meaning that an application of Lemma B.1 results in the bound

$$(58) \quad |\mathbf{A}_{\mathcal{M}_h}(u_h^{-\ell} - U_h, \Xi_h)| \lesssim h^2 \|u_h\|_{H^1(\mathcal{M})} \|\Xi_h^{\ell}\|_{H^1(\mathcal{M})}.$$

Further, note that for any $\Xi_h^{\ell} \in S_h^{\ell}$, the equivalence of norms (17) gives

$$(59) \quad \|\Xi_h^{\ell}\|_{H^1(\mathcal{M})}^2 \sim \|\mathcal{L}_h^{1/2} \Xi_h^{\ell}\|_{L^2(\mathcal{M})}^2 = (\mathcal{L}_h \Xi_h^{\ell}, \Xi_h^{\ell})_{L^2(\mathcal{M})} = \mathbf{A}_{\mathcal{M}}(\Xi_h^{\ell}, \Xi_h^{\ell}),$$

where the last equality comes from the definition of \mathcal{L}_h . And similarly, for any $\Xi_h \in S_h$, we have

$$(60) \quad \|\Xi_h\|_{H^1(\mathcal{M}_h)}^2 \sim \mathbf{A}_{\mathcal{M}_h}(\Xi_h, \Xi_h).$$

Then, applying the triangle inequality to the (last) right-hand side of Equation (59) gives

$$\|\Xi_h^{\ell}\|_{H^1(\mathcal{M})}^2 \lesssim \mathbf{A}_{\mathcal{M}_h}(\Xi_h, \Xi_h) + |\mathbf{A}_{\mathcal{M}}(\Xi_h^{\ell}, \Xi_h^{\ell}) - \mathbf{A}_{\mathcal{M}_h}(\Xi_h, \Xi_h)| \lesssim \|\Xi_h\|_{H^1(\mathcal{M}_h)}^2 + h^2 \|\Xi_h^{\ell}\|_{H^1(\mathcal{M})}^2,$$

where we used Equation (60) and Lemma B.1 to derive the second inequality. This means in particular that there exists $C > 0$ independent of h such that $\|\Xi_h^{\ell}\|_{H^1(\mathcal{M})}^2 \leq C(\|\Xi_h\|_{H^1(\mathcal{M}_h)}^2 + h^2 \|\Xi_h^{\ell}\|_{H^1(\mathcal{M})}^2)$. Recall that $h \in (0, h_0)$ for some $h_0 \in (0, 1)$ small enough. Assuming that especially $1 - Ch_0^2 > 0$ yields $\|\Xi_h^{\ell}\|_{H^1(\mathcal{M})}^2 \leq C(1 - Ch_0^2)^{-1} \|\Xi_h\|_{H^1(\mathcal{M}_h)}^2 \leq C(1 - Ch_0^2)^{-1} \|\Xi_h\|_{H^1(\mathcal{M}_h)}^2$, which allows us to conclude that

$$(61) \quad \|\Xi_h^{\ell}\|_{H^1(\mathcal{M})}^2 \lesssim \|\Xi_h\|_{H^1(\mathcal{M}_h)}^2.$$

Now, applying successively Equation (60) and Equation (58) with $\Xi_h = u_h^{-\ell} - U_h$, we obtain

$$\|u_h^{-\ell} - U_h\|_{H^1(\mathcal{M}_h)}^2 \lesssim h^2 \|u_h\|_{H^1(\mathcal{M})} \|u_h - U_h^{\ell}\|_{H^1(\mathcal{M})} \lesssim h^2 \|u_h\|_{H^1(\mathcal{M})} \|u_h^{-\ell} - U_h\|_{H^1(\mathcal{M}_h)},$$

where the last inequality is derived from applying Equation (61). Therefore, we end up with

$$\|u_h - U_h^{\ell}\|_{H^1(\mathcal{M})} \lesssim h^2 \|u_h\|_{H^1(\mathcal{M})} \lesssim h^2 \|\mathcal{L}_h^{-1/2} v_h\|_{L^2(\mathcal{M})},$$

where the equivalence of norms (17) together with the definition of u_h are used in the final step. This concludes the proof of Equation (53). \square

B.3. Proof of Proposition 2.5. Based on the results in the previous subsections, we can now move on to the proof of Proposition 2.5.

Proof. Let $\tilde{f} \in S_h^{\ell}$. We introduce the inverse lift operator $\mathcal{P}_{\ell} : L^2(\mathcal{M}) \rightarrow L^2(\mathcal{M}_h)$ which maps any $F \in L^2(\mathcal{M})$ to $\mathcal{P}_{\ell} F = F^{-\ell}$. Let then $\mathcal{E}_h = \|(\gamma(\mathcal{L}_h) \tilde{f})^{-\ell} - \gamma(\mathbf{L}_h) \mathbf{P}_h(\sigma \tilde{f}^{-\ell})\|_{L^2(\mathcal{M}_h)} = \|\mathcal{P}_{\ell} \gamma(\mathcal{L}_h) \tilde{f} - \gamma(\mathbf{L}_h) \mathbf{P}_h(\sigma \mathcal{P}_{\ell} \tilde{f})\|_{L^2(\mathcal{M}_h)}$. Note that by the integral representations of the operators (19)

$$\mathcal{E}_h = \left\| \frac{1}{2\pi i} \int_{\Gamma} \gamma(z) \mathcal{F}(z) \, dz \right\|_{L^2(\mathcal{M}_h)},$$

where we take for any $z \in \Gamma$, $\mathcal{F}(z) = \mathcal{P}_\ell(z - \mathcal{L}_h)^{-1} - (z - \mathbf{L}_h)^{-1} \mathbf{P}_h \sigma \mathcal{P}_\ell$. Similarly, as in the proof of Proposition 2.3, we use the splitting (20) of Γ and the triangle inequality to deduce that

$$(62) \quad \begin{aligned} \mathcal{E}_h &\leq \frac{1}{2\pi} \int_{\Omega_+} |\gamma(g_+(t))| \|\mathcal{F}(g_+(t))\|_{L^2(\mathcal{M}_h)} dt + \frac{\delta_0}{2\pi} \int_{\Omega_0} |\gamma(g_0(t))| \|\mathcal{F}(g_0(t))\|_{L^2(\mathcal{M}_h)} dt \\ &\quad + \frac{1}{2\pi} \int_{\Omega_-} |\gamma(g_-(t))| \|\mathcal{F}(g_-(t))\|_{L^2(\mathcal{M}_h)} dt, \end{aligned}$$

where we take $\Omega_+ = \Omega_- = [\delta_0, \infty)$ and $\Omega_0 = [-\theta, \theta]$. For $* \in \{+, 0, -\}$, let us then introduce the quantity

$$\mathcal{E}_h^* = \int_{\Omega_*} |\gamma(g_*(t))| \|\mathcal{F}(g_*(t))\|_{L^2(\mathcal{M}_h)} dt,$$

so that Equation (62) may be rewritten as $\mathcal{E}_h \lesssim \mathcal{E}_h^+ + \mathcal{E}_h^0 + \mathcal{E}_h^-$ and in particular, $g_*(t) \in \Gamma$ for any $t \in \Omega_*$.

We now fix $* \in \{+, 0, -\}$ and bound the term \mathcal{E}_h^* . First, for any $z \in \Gamma$, we rewrite $\mathcal{F}(z)$ and split

$$\begin{aligned} \mathcal{F}(z) &= (z - \mathbf{L}_h)^{-1} \mathbf{L}_h \left((z \mathbf{L}_h^{-1} - I) \mathcal{P}_\ell \mathcal{L}_h^{-1} - \mathbf{L}_h^{-1} \mathbf{P}_h \sigma \mathcal{P}_\ell (z \mathcal{L}_h^{-1} - I) \right) \mathcal{L}_h (z - \mathcal{L}_h)^{-1} \\ &= (z - \mathbf{L}_h)^{-1} \mathbf{L}_h \left(z \mathbf{L}_h^{-1} (I - \mathbf{P}_h \sigma) \mathcal{P}_\ell \mathcal{L}_h^{-1} + \mathbf{L}_h^{-1} \mathbf{P}_h \sigma \mathcal{P}_\ell - \mathcal{P}_\ell \mathcal{L}_h^{-1} \right) \mathcal{L}_h (z - \mathcal{L}_h)^{-1} \\ &= \mathcal{F}_1(z) + \mathcal{F}_2(z), \end{aligned}$$

where we take $\mathcal{F}_1(z) = (z - \mathbf{L}_h)^{-1} \mathbf{L}_h \left(z \mathbf{L}_h^{-1} (I - \mathbf{P}_h \sigma) \mathcal{P}_\ell \mathcal{L}_h^{-1} \right) \mathcal{L}_h (z - \mathcal{L}_h)^{-1} = z(z - \mathbf{L}_h)^{-1} (I - \mathbf{P}_h \sigma) \mathcal{P}_\ell (z - \mathcal{L}_h)^{-1}$ and $\mathcal{F}_2(z) = (z - \mathbf{L}_h)^{-1} \mathbf{L}_h \left(\mathbf{L}_h^{-1} \mathbf{P}_h \sigma \mathcal{P}_\ell - \mathcal{P}_\ell \mathcal{L}_h^{-1} \right) \mathcal{L}_h (z - \mathcal{L}_h)^{-1}$. Hence, by the triangle inequality,

$$(63) \quad \mathcal{E}_h^* \lesssim \int_{\Gamma_t} |\gamma(g_*(t))| \left(\|\mathcal{F}_1(g_*(t)) \tilde{f}\|_{L^2(\mathcal{M}_h)} + \|\mathcal{F}_2(g_*(t)) \tilde{f}\|_{L^2(\mathcal{M}_h)} \right) dt.$$

We first bound $\|\mathcal{F}_1(z) \tilde{f}\|_{L^2(\mathcal{M}_h)}$. Using successively Equation (21) and the geometric estimates in Bonito et al. (2018, Corollary 2.2) results in

$$\begin{aligned} \|\mathcal{F}_1(z) \tilde{f}\|_{L^2(\mathcal{M}_h)} &= |z| \left\| (z - \mathbf{L}_h)^{-1} (I - \mathbf{P}_h \sigma) \mathcal{P}_\ell (z - \mathcal{L}_h)^{-1} \tilde{f} \right\|_{L^2(\mathcal{M}_h)} \\ &\lesssim \left\| (I - \mathbf{P}_h \sigma) \mathcal{P}_\ell (z - \mathcal{L}_h)^{-1} \tilde{f} \right\|_{L^2(\mathcal{M}_h)} \lesssim h^2 \left\| (z - \mathcal{L}_h)^{-1} \tilde{f} \right\|_{L^2(\mathcal{M})}. \end{aligned}$$

Using then Equation (52), we conclude that, for any $p \in (0, 2)$,

$$(64) \quad \left\| \mathcal{F}_1(z) \tilde{f} \right\|_{L^2(\mathcal{M}_h)} \lesssim h^2 |z|^{-(1-p/2)} \left\| \mathcal{L}_h^{-p/2} \tilde{f} \right\|_{L^2(\mathcal{M})}^2.$$

To bound $\|\mathcal{F}_2(z) \tilde{f}\|_{L^2(\mathcal{M}_h)}$, we apply Equations (51) and (53) to obtain

$$\begin{aligned} \|\mathcal{F}_2(z) \tilde{f}\|_{L^2(\mathcal{M}_h)} &\lesssim |z|^{-1/2} \left\| (\mathbf{L}_h^{-1} \mathbf{P}_h \sigma \mathcal{P}_\ell - \mathcal{P}_\ell \mathcal{L}_h^{-1}) \mathcal{L}_h (z - \mathcal{L}_h)^{-1} \tilde{f} \right\|_{H^1(\mathcal{M}_h)} \\ &\lesssim |z|^{-1/2} h^2 \left\| \mathcal{L}_h^{1/2} (z - \mathcal{L}_h)^{-1} \tilde{f} \right\|_{L^2(\mathcal{M})} \end{aligned}$$

and with Equation (50) for any $p \in [-1, 1]$ ($p = -q$)

$$(65) \quad \left\| \mathcal{F}_2(z) \tilde{f} \right\|_{L^2(\mathcal{M}_h)} \lesssim h^2 |z|^{-(1-p/2)} \left\| \mathcal{L}_h^{-p/2} \tilde{f} \right\|_{L^2(\mathcal{M})}.$$

Using Equations (64) and (65) with $p = \min\{\alpha + d/4; 1\}$ together with Equation (63) gives

$$\mathcal{E}_h^* \lesssim \int_{\Omega_*} |\gamma(g_*(t))| h^2 |g_*(t)|^{-(1-p/2)} \|\mathcal{L}_h^{-p/2} \tilde{f}\|_{L^2(\mathcal{M})} dt,$$

which yields in turn (since γ is an α -power spectral density)

$$\mathcal{E}_h^* \lesssim h^2 \|\mathcal{L}_h^{-p/2} \tilde{f}\|_{L^2(\mathcal{M})} \int_{\Omega_*} |g_*(t)|^{-(1+\alpha-p/2)} dt \lesssim h^2 \|\mathcal{L}_h^{-\min\{\alpha+d/4; 1\}/2} \tilde{f}\|_{L^2(\mathcal{M})},$$

since $\alpha - p/2 = \max\{\alpha - (\alpha + d/4)/2; \alpha - 1/2\} = \max\{(\alpha - d/4)/2; \alpha - 1/2\} \geq (\alpha - d/4)/2 > 0$. Finally, since this inequality holds for any $*$ $\in \{+, 0, -\}$, we retrieve the claim (22) using Equation (62). □

(Erik Jansson)

DEPARTMENT OF MATHEMATICAL SCIENCES
CHALMERS UNIVERSITY OF TECHNOLOGY & UNIVERSITY OF GOTHENBURG
S-412 96 GÖTEBORG, SWEDEN.

Email address: erikjans@chalmers.se

(Annika Lang)

DEPARTMENT OF MATHEMATICAL SCIENCES
CHALMERS UNIVERSITY OF TECHNOLOGY & UNIVERSITY OF GOTHENBURG
S-412 96 GÖTEBORG, SWEDEN.

Email address: annika.lang@chalmers.se

(Mike Pereira)

DEPARTMENT OF GEOSCIENCES AND GEOENGINEERING
MINES PARIS
PSL UNIVERSITY
F-77 305 FONTAINEBLEAU, FRANCE.

Email address: mike.pereira@minesparis.psl.eu