

# The Behavior of Tree-Width and Path-Width under Graph Operations and Graph Transformations

Frank Gurski<sup>1,\*</sup> and Robin Weishaupt<sup>1</sup>

<sup>1</sup>University of Düsseldorf, Institute of Computer Science, 40225 Düsseldorf, Germany

\*Corresponding Author: frank.gurski@hhu.de

Tree-width and path-width are well-known graph parameters. Many NP-hard graph problems allow polynomial-time solutions, when restricted to graphs of bounded tree-width or bounded path-width. In this work, we study the behavior of tree-width and path-width under various unary and binary graph transformations. Doing so, for considered transformations we provide upper and lower bounds for the tree-width and path-width of the resulting graph in terms of the tree-width and path-width of the initial graphs or argue why such bounds are impossible to specify. Among the studied, unary transformations are vertex addition, vertex deletion, edge addition, edge deletion, subgraphs, vertex identification, edge contraction, edge subdivision, minors, powers of graphs, line graphs, edge complements, local complements, Seidel switching, and Seidel complementation. Among the studied, binary transformations we consider the disjoint union, join, union, substitution, graph product, 1-sum, and corona of two graphs.

**Keywords:** tree-width, path-width, graph operations, graph transformations

## 1 Introduction

A *graph parameter* is a function that associates every graph with a non-negative integer. One of the most famous graph parameters is tree-width, which was defined by Robertson and Seymour [RS86a]. Graphs of bounded tree-width are interesting from an algorithmic point of view, as several NP-complete graph problems can be solved in polynomial time for graph classes of bounded tree-width. For example, tree-decompositions allow for many efficient algorithms in dynamic programming [Arn85, AP89, Hag00, KZN00]. The same holds for the similar graph parameter path-width. That is, because every path-decomposition can be interpreted as a special case of a tree-decomposition. Both parameters play a crucial role in the field of structural graph theory, especially in the graph minor theory of Robertson and Seymour [RS85].

Trees and forests have tree-width at most one. Series parallel graphs have tree-width at most two [Bod98]. Outerplanar graphs (and subclasses, such as cactus graphs and maximal outerplanar graphs) have tree-width at most two and  $k$ -outerplanar graphs have tree-width at most  $3k - 1$  [Bod98]. Halin graphs have tree-width at most three [Bod98]. For further classes of graphs with bounded tree-width, we refer to the works by Bodlaender [Bod86, Bod88, Bod98].

Determining whether the tree-width or path-width of a graph is at most a given value  $w$  is NP-complete [ACP87]. However, for every fixed integer  $k$ , one can decide in linear time whether a given graph  $G$  has tree-width or path-width  $k$ , see Bodlaender [Bod96]. For an in-depth overview of tree-width and path-width, we refer again to the work by Bodlaender [Bod98].

A *graph transformation*  $f$  is a function that creates a new graph  $f(G_1, \dots, G_n)$  from a number of  $n \geq 1$  input graphs  $G_1, \dots, G_n$ . Examples of graph transformations are taking an induced subgraph of a graph, adding an edge to a graph, or generating the join of two graphs. A *graph operation* is a graph transformation which is deterministic and invariant under isomorphism. Examples of graph operations are the edge complementation of a graph or generating the join of two graphs.<sup>1</sup> The graph theory books by Bondy and Murty [BM76] and Harary [Har69] provide a large number of graph transformations.

The impact of graph operations, which can be defined by monadic second order formulas (so-called MS transductions), on graph parameters can often be shown in a very short way. Unfortunately, the resulting bounds are most of the time rather imprecise [Cou06, CE12].

Transformations that reduce graphs can be used to characterize classes of graphs by forbidden subgraphs. For example, the property that a graph has tree-width at most  $k$  is preserved under the graph transformation “taking minors”. This fact is used to show that the set of graphs of tree-width at most  $k$  can be characterized by a finite set of forbidden minors [RS85].

The effect of graph transformations on graph parameters is well studied, e.g., for bandwidth [CO86], for tree-width [Bod98], for clique-width [Cou14, HOSG08, Gur17], and for rank-width [HOSG08].

In this paper we study the behavior of tree-width and path-width under various graph transformations and graph operations. Thereby, we consolidate known results from various works and prove novel results. Doing so, this work tries to provide a comprehensive overview of the effects of unary and binary graph transformations on tree-width and path-width. Therefore, the paper is organized as follows. In Section 2, we recall the definitions of tree-width and path-width. In Section 3, we consider the effects of the unary graph transformations vertex deletion, vertex addition, edge deletion, edge addition, subgraphs, vertex identification, edge contraction, edge subdivision, minors, powers of graphs, line graphs, edge complements, local complements, Seidel switching, and Seidel complementation on tree-width and path-width. Whenever it is possible to bound the tree-width or path-width of the resulting graph  $f(G)$ , we show how to compute the corresponding decomposition in time linear in the size of the corresponding decomposition for  $G$ . In Section 4, we give an overview of the effects of the binary graph operations disjoint union, join, union, substitution, graph products, 1-sum, and corona on tree-width and path-width. If it is possible to bound the tree-width or path-width of the combined graph  $f(G_1, G_2)$  in the tree-width or path-width of graphs  $G_1$  and  $G_2$ , we show how to compute the corresponding decomposition in time linear in the size of the corresponding decompositions for  $G_1$  and  $G_2$ . Finally, we summarize our results and provide some conclusions as well as an outlook for future work in Section 5.

---

<sup>1</sup>Please note that by our definition the two graph transformations taking an induced subgraph of a graph and adding an edge to a graph are no graph operations.

## 2 Preliminaries

### 2.1 Graphs

We work with finite, undirected *graphs*  $G$ , where  $V(G)$  is a finite set of *vertices* and  $E(G) \subseteq \{\{u, v\} \mid u, v \in V(G), u \neq v\}$  is a finite set of *edges* without loops or multiple edges. For a vertex  $v \in V(G)$ , we denote by  $N_G(v)$  the set of all vertices adjacent to  $v$  in  $G$ , i.e.,  $N_G(v) = \{w \in V(G) \mid \{v, w\} \in E(G)\}$ . The vertex set  $N_G(v)$  is called the set of all *neighbors* of  $v$  in  $G$  or the *neighborhood* of  $v$  in  $G$ . Note that  $v$  itself does not belong to  $N_G(v)$ . The *degree* of a vertex  $v \in V(G)$ , denoted by  $\deg_G(v)$ , is the number of neighbors of  $v$  in  $G$ , i.e.,  $\deg_G(v) = |N_G(v)|$ , and the *maximum degree* of  $G$ , denoted by  $\Delta(G)$ , is the maximum over the vertices' degrees, i.e.,  $\Delta(G) = \max_{v \in V(G)} \deg_G(v)$ . In this work, we discuss graphs only up to isomorphism. This allows us to define the path with  $n$  vertices as  $P_n = (\{v_1, \dots, v_n\}, \{\{v_1, v_2\}, \dots, \{v_{n-1}, v_n\}\})$ , which is useful in several examples. For definitions of further special graphs, we refer to the book by Brandstädt et al. [BLS99].

### 2.2 Tree-width

One of the most famous tree structured graph classes are graphs of bounded tree-width. The first notion equivalent to tree-width, the dimension of a graph, was introduced by Bertelè and Brioschi [BB73] in 1973. The subsequent definition of tree-width was given by Robertson and Seymour [RS86a] in 1986.

**Definition 1 (Tree-width)** A tree-decomposition of a graph  $G$  is a pair  $(\mathcal{X}, T)$ , where  $T$  is a tree and  $\mathcal{X} = \{X_u \subseteq V(G) \mid u \in V(T)\}$  is a family of subsets of  $V(G)$ , called bags, such that the following three conditions hold:

$$(tw-1) \bigcup_{u \in V(T)} X_u = V(G),^2$$

(tw-2) for every edge  $\{v_1, v_2\} \in E(G)$ , there is a node  $u \in V(T)$  with  $v_1, v_2 \in X_u$ , and

(tw-3) for every vertex  $v \in V(G)$  the subgraph of  $T$  induced by the nodes  $u \in V(T)$  with  $v \in X_u$  is connected.

The width of a tree-decomposition  $(\mathcal{X} = \{X_u \mid u \in V(T)\}, T)$  is defined as

$$\max_{u \in V(T)} |X_u| - 1.$$

Finally, the tree-width of a graph  $G$ , denoted by  $tw(G)$ , is the smallest integer  $k$ , such that there is a tree-decomposition  $(\mathcal{X}, T)$  for  $G$  of width  $k$ .

Obviously, the tree-width of every graph  $G$  is bounded by  $|V(G)| - 1$ , as one can always place all its vertices in a single bag. Contrarily, determining whether the tree-width of some given graph is at most some given integer  $w$  is NP-complete, even for bipartite graphs and complements of bipartite graphs [ACP87].

*Example 1 (Tree-decomposition)* In Figure 1 we depict a graph together with a corresponding tree-decomposition of width 1.

---

<sup>2</sup>Please note that (tw-1) has no influence on the width of a tree-decomposition. It merely ensures that isolated vertices are covered by a tree-decomposition, which is useful for the design of algorithms along tree-decompositions.




Figure 1: A graph (left) and an associated tree-decomposition (right).

Next to the previous definition of tree-width, tree-width can also be defined by bramble numbers [Ree97] (also known as the size of screens [ST93]) or partial  $k$ -trees [Ros74]. We refer to the work by Bodlaender [Bod98] for an overview of equivalent definitions.

Throughout the course of our work, we make use of the following lemma by Scheffler [Sch89] several times.

**Lemma 1** *Every graph of tree-width at most  $k$  has a vertex of degree at most  $k$ .*

**Proof** Let  $G$  be a graph with  $\text{tw}(G) = k$  and  $(\mathcal{X}, T)$  a tree-decomposition of width  $k$  for  $G$ , such that  $|V(T)|$  is as small as possible. Let  $\ell$  be a leaf of  $T$  and denote its parent in  $T$  as  $v$ . Since  $|V(T)|$  is as small as possible, there must exist a vertex  $u \in X_\ell$  with  $u \notin X_v$ . Otherwise,  $X_\ell \subset X_v$  holds and we can delete  $X_\ell$ , contradicting the minimality of  $V(T)$ . Thus, due to (tw-2) and (tw-3), all neighbors of  $u$  must be in  $X_\ell$  and, as  $X_\ell$  contains at most  $k + 1$  vertices, it follows that  $u$  has degree at most  $k$ .  $\square$

The subsequent proposition is stated by Bodlaender [Bod98], who in turn credits it to Rose [Ros74]. As Rose's proof makes use of  $k$ -trees, subsequently, for the sake of self-containment, we provide a proof only making use of Lemma 1 and tree-width.

**Proposition 2** *Let  $G$  be a graph. Then, it holds that*

$$|E(G)| \leq \text{tw}(G)|V(G)| - \frac{1}{2}\text{tw}(G)(\text{tw}(G) + 1).$$

**Proof** By Lemma 1 we know that there exists at least one vertex  $v$  in  $G$  of degree at most  $\text{tw}(G)$ , such that  $v$  possesses at most  $\text{tw}(G)$  incident edges in  $G$ . Let us denote the graph we obtain by removing  $v$  from  $G$  as  $G_1$ . Then, we know that from  $G$  to  $G_1$  we removed at most  $\text{tw}(G)$  edges from  $G$ . It is easy to see that removing  $v$  and all its incident edges from  $G$  can reduce the tree-width of  $G$  but not increase it. Hence,  $\text{tw}(G_1) \leq \text{tw}(G)$  holds and we can repeat the previous argument for  $G_1$ , i.e., remove a vertex as well as at most  $\text{tw}(G)$  edges from  $G_1$ . Repeating this argument for all vertices of  $G$  results in  $|E(G)| \leq \text{tw}(G)|V(G)|$ . However, at one point, there are at most  $\text{tw}(G)$  vertices left. This remaining graph can possess a tree-width of at most  $\text{tw}(G) - 1$ , such that by Lemma 1 it follows that in this graph there is a vertex with degree at most  $\text{tw}(G) - 1$ . Removing this vertex from the remaining graph, we remove at most  $\text{tw}(G) - 1$  edges. Repeating this argument for all of the  $\text{tw}(G)$  remaining vertices in sequence yields

$$\begin{aligned} |E(G)| &\leq \text{tw}(G)(|V(G)| - \text{tw}(G)) + \sum_{i=1}^{\text{tw}(G)} (\text{tw}(G) - i) \\ &= \text{tw}(G)(|V(G)| - \text{tw}(G)) + \text{tw}(G)\text{tw}(G) - \sum_{i=1}^{\text{tw}(G)} i \\ &= \text{tw}(G)|V(G)| - \frac{1}{2}\text{tw}(G)(\text{tw}(G) + 1). \end{aligned}$$

□

While the previous proposition provides an upper bound for the number of edges of a graph in terms of the number of its vertices as well as its tree-width, the ensuing proposition by Kneis et al. [KMRR09] provides an upper bound for the tree-width of a graph in terms of the number of its edges and vertices.

**Proposition 3** *Let  $G$  be a graph. Then, it holds that*

$$tw(G) \leq \frac{|E(G)|}{5.769} + \mathcal{O}(\log(|V(G)|)).$$

Recall that a graph is said to be  $k$ -connected if and only if every pair of its vertices is connected by at least  $k$  vertex-disjoint paths [Whi32]. Making use of Lemma 1, we obtain the following lower bound for the tree-width of  $k$ -connected graphs.

**Lemma 4** *Every  $k$ -connected graph has tree-width at least  $k$ .*

**Proof** Let  $G$  be a  $k$ -connected graph. By definition, it holds that for all vertices  $u, v \in V(G)$  there exists  $k$  vertex-disjoint paths from  $u$  to  $v$  in  $G$ . Consequently,  $\deg_G(u) \geq k$  as well as  $\deg_G(v) \geq k$  must be true. In other words, every vertex of  $G$  has degree at least  $k$ . By the contraposition of Lemma 1 we immediately obtain  $tw(G) \geq k$ . □

To conclude the preliminary subsection on tree-width, we cite two lemmas by Bodlaender and Möhring [BM93] of which we make use in future sections.

**Lemma 5 (Clique containment)** *Let  $(\mathcal{X}, T)$  be a tree-decomposition for a graph  $G$  and  $C \subseteq V(G)$  a clique in  $G$ . Then, there exists some  $u \in V(T)$  with  $C \subseteq X_u$ .*

**Lemma 6 (Complete bipartite subgraph containment)** *Let  $(\mathcal{X}, T)$  be a tree-decomposition for a graph  $G$  and  $A, B \subseteq V(G)$  with  $A \cap B = \emptyset$  and  $\{\{u, v\} \mid u \in A, v \in B\} \subseteq E(G)$ . Then, there exists a vertex  $u \in V(T)$  with  $A \subseteq X_u$  or  $B \subseteq X_u$ .*

## 2.3 Path-width

In 1983, three years ahead of their definition for tree-width, Robertson and Seymour introduced the notion of path-width [RS83].

**Definition 2 (Path-width)** *A path-decomposition of a graph  $G$  is a sequence  $(X_1, \dots, X_r)$  of subsets of  $V(G)$ , called bags, such that the subsequent three conditions hold:*

(pw-1)  $\bigcup_{1 \leq i \leq r} X_i = V(G)$ ,<sup>3</sup>

(pw-2) *for every edge  $\{u, v\} \in E(G)$  there is a set  $X_i$ ,  $1 \leq i \leq r$ , with  $u, v \in X_i$ , and*

(pw-3) *for all  $i, j, \ell$  with  $1 \leq i < j < \ell \leq r$  it holds that  $X_i \cap X_\ell \subseteq X_j$ .*

*The width of a path-decomposition  $(X_1, \dots, X_r)$  is defined as*

$$\max_{1 \leq i \leq r} |X_i| - 1.$$

*Finally, the path-width of a graph  $G$ , denoted by  $pw(G)$ , is the smallest integer  $k$ , such that there is a path-decomposition  $(X_1, \dots, X_r)$  for  $G$  of width  $k$ .*

---

<sup>3</sup>Please note that (pw-1) has no influence on the width of a path-decomposition, see footnote 2.

Similar to the tree-width of a graph, the path-width of a graph  $G$  is limited by  $|V(G)|-1$ , as all vertices of  $G$  can be stored in a single bag. However, deciding whether the path-width of a graph is at most some given integer  $w$  is NP-complete [KF79]. Even for special graph classes, such as bipartite graphs, complements of bipartite graphs [ACP87], chordal graphs [Gus93], bipartite distance hereditary graphs [KBMK93], and planar graphs with maximum vertex degree 3 [MS88], it is NP-complete to decide if the path-width is at most a given integer  $w$ . On the other hand, there are also special graph classes, such as permutation graphs [BKK93], circular arc graphs [ST07], and co-graphs [BM93], for which one can decide in polynomial time whether the path-width is at most a given integer  $w$ .

*Example 2 (Path-decomposition)*  $\mathcal{X} = (\{a, b, c\}, \{c, f, g\}, \{c, d, e\})$  is a path-decomposition of width 2 for the graph shown in Figure 1.

Besides the definition of path-width by Robertson and Seymour, there are further definitions of path-width, for example by vertex separation number [Kin92]. Again, we refer to the work by Bodlaender [Bod98] for an overview of equivalent definitions.

With a small adaption to the proof for Lemma 1, we obtain a similar result for path-width.

**Lemma 7** *Every graph of path-width at most  $k$  has a vertex of degree at most  $k$ .*

Having this lemma, we obtain the subsequent proposition for path-width with the same proof as used for Proposition 2, adapted to path-width.

**Proposition 8** *Let  $G$  be a graph. Then, it holds that*

$$|E(G)| \leq pw(G)|V(G)| - \frac{1}{2}pw(G)(pw(G) + 1).$$

The same work by Kneis et al. [KMRR09] that provides an upper bound for the tree-width of a graph in terms of its edges and vertices also provides an upper bound for the path-width of a graph in terms of its edges and vertices.

**Proposition 9** *Let  $G$  be a graph. Then, it holds that*

$$pw(G) \leq \frac{|E(G)|}{5.769} + \mathcal{O}(\log(|V(G)|)).$$

Following the same argumentation as for Lemma 1, we obtain the subsequent result for the path-width of  $k$ -connected graphs.

**Lemma 10** *Every  $k$ -connected graph has path-width at least  $k$ .*

As every path is a tree, every path-decomposition  $\mathcal{X} = (X_1, \dots, X_r)$  can be interpreted as a tree-decomposition, such that the results from Lemma 5 and 6 also hold for path-decompositions. Subsequently, we provide reformulations of the results by Bodlaender and Möhring [BM93], adjusted for path-width.

**Lemma 11 (Clique containment)** *Let  $\mathcal{X} = (X_1, \dots, X_r)$  be a path-decomposition for a graph  $G$  and  $C \subseteq V(G)$  a clique in  $G$ . Then, there exists some  $1 \leq i \leq r$  with  $C \subseteq X_i$ .*

**Lemma 12 (Complete bipartite subgraph containment)** *Let  $\mathcal{X} = (X_1, \dots, X_r)$  be a path-decomposition for a graph  $G$  and  $A, B \subseteq V(G)$  with  $A \cap B = \emptyset$  and  $\{\{u, v\} \mid u \in A, v \in B\} \subseteq E(G)$ . Then, there exists some  $1 \leq i \leq r$  with  $A \subseteq X_i$  or  $B \subseteq X_i$ .*

## 2.4 Graph Parameters and Relations

In this subsection, we provide results setting the parameters tree-width and path-width into relation with each other as well as with other, well-known graph parameters, such as

1. the *independence number*  $\alpha(G)$ , specifying the size of a largest independent set in  $G$ ,
2. the *clique number*  $\omega(G)$ , specifying the size of a largest clique in  $G$ , and
3. the *chromatic number*  $\chi(G)$ , specifying the least number of distinct colors required to color all vertices of  $G$ , such that no two adjacent vertices obtain the same color.

With the subsequent theorem, we prove that the tree-width of a graph is always bounded by the path-width of the graph, whereas the path-width of a graph cannot be bounded only in the graph's tree-width, but in the tree-width and the number of vertices.

**Theorem 13** *For a graph  $G$  it holds that*

$$tw(G) \leq pw(G) \quad \text{and} \quad pw(G) \leq (tw(G) + 1)(\log_3(2|V(G)| + 1) + 1) - 1.$$

**Proof** The first relation follows by the definitions of tree-width and path-width, as every path-decomposition can also be interpreted as a tree-decomposition.

The second relation, making use of the idea of Theorem 6 in the work by Korach and Solel [KS93], can be shown as follows. Let  $(\mathcal{X}, T)$  be a tree-decomposition for  $G$  and denote its bags by  $X_u \in \mathcal{X}$  for  $u \in V(T)$ . According to the result (10.16) in the book by Kleinberg and Tardos [KT04], we can assume  $(\mathcal{X}, T)$  to be a non-redundant tree-decomposition for  $G$ , i.e.,  $|V(T)| \leq |V(G)|$  holds. Following Scheffler [Sch89], the path-width of any tree  $S$  can be limited from above by  $pw(S) \leq \log_3(2|V(S)| + 1)$ . Thus, since  $(\mathcal{X}, T)$  is a non-redundant tree-decomposition for  $G$ , it follows that there exists a path-decomposition  $\mathcal{Y}$  for  $T$  of path-width  $pw(T) \leq \log_3(2|V(G)| + 1)$ . We denote the bags of  $\mathcal{Y}$  by  $Y_i \in \mathcal{Y}$  for  $1 \leq i \leq r$ . Given this path-decomposition  $\mathcal{Y}$  for  $T$ , we construct  $\mathcal{Z} = (Z_1, \dots, Z_r)$  with  $Z_i = \bigcup_{u \in Y_i} X_u$ . By Theorem 6 in the work by Korach and Solel [KS93], it follows that  $\mathcal{Z}$  is a path-decomposition for  $G$ . We know that  $|X_u| \leq tw(G) + 1$  holds for all  $u \in V(T)$  and  $|Y_i| \leq \log_3(2|V(G)| + 1) + 1$  for  $1 \leq i \leq r$ . Consequently, it follows that  $|Z_i| \leq (tw(G) + 1)(\log_3(2|V(G)| + 1) + 1)$  is true for  $1 \leq i \leq r$ , such that we obtain  $pw(G) \leq (tw(G) + 1)(\log_3(2|V(G)| + 1) + 1) - 1$ .  $\square$

By Lemma 5 and Lemma 11 we know that for a complete graph with  $n \in \mathbb{N}$  vertices,  $K_n$ , it holds that  $tw(K_n) = n - 1 = pw(K_n)$ . Consequently, the first inequality stated in Theorem 13 is tight. The second inequality is asymptotically optimal, as trees have tree-width one while their path-width grows asymptotically in their height, which is at least logarithmic in the number of vertices.

The same two lemmas imply that for a graph  $G$  every clique of  $G$  is subset of at least one bag of every tree-decomposition and path-decomposition for  $G$ . With this fact at hand, the subsequent proposition follows immediately.

**Proposition 14** *For a graph  $G$  it holds that*

$$\omega(G) - 1 \leq tw(G) \quad \text{and} \quad \omega(G) - 1 \leq pw(G).$$

Chlebíkóv [Chl02] proved the subsequent proposition's first bound, i.e., the one for tree-width. To do so, he argued that by splitting a graph  $G$ 's vertices into an independent

set  $I$  of size  $\alpha(G)$  and  $|V(G)| - \alpha(G)$  subsets  $V_i \subseteq V(G) \setminus I$  of size one, it follows that  $G$  is a subgraph of a complete  $(|V(G)| - \alpha(G) + 1)$ -partite graph. Then, it is easy to see that the bags  $X_v = (V(G) \setminus I) \cup \{v\}$  for  $v \in I$  allow a tree-decomposition of width  $|V(G)| - \alpha(G)$  for the  $(|V(G)| - \alpha(G) + 1)$ -partite graph. As the same bags  $X_v$ ,  $v \in I$ , allow a path-decomposition as well, the same bound follows for path-width.

**Proposition 15** *For a graph  $G$  it holds that*

$$\alpha(G) + tw(G) \leq |V(G)| \quad \text{and} \quad \alpha(G) + pw(G) \leq |V(G)|.$$

Finally, Chlebíková [Chl02] also proved the subsequent proposition's first bound. Making use of the first bound in Theorem 13, we extended this result to path-width, too.

**Proposition 16** *For a graph  $G$  it holds that*

$$\chi(G) \leq tw(G) + 1 \quad \text{and} \quad \chi(G) \leq pw(G) + 1.$$

### 3 Unary Graph Operations and Graph Transformations

Let  $G$  be a non-empty graph and  $f$  a unary graph transformation that creates a new graph  $f(G)$  from  $G$ . In this section of our work, we consider the tree-width and path-width of graph  $f(G)$  with respect to the tree-width and path-width of  $G$ . Especially, we consider the graph transformations vertex deletion, vertex addition, edge deletion edge addition, taking a subgraph, edge subdivision, vertex identification, edge contraction, taking a minor, powers of graphs, line graphs, edge complements, local complements, Seidel switching, and Seidel complementation.

#### 3.1 Vertex Deletion and Vertex Addition

**Vertex Deletion** Let  $G$  be a graph and  $v \in V(G)$  a vertex of  $G$ . By  $G - v$  we denote the graph we obtain from  $G$  when removing  $v$  and all its incident edges, i.e.,

$$G - v = (V(G) \setminus \{v\}, E(G) \setminus \{\{v, u\} \mid u \in N_G(v)\}).$$

Having the graph operation of *vertex deletion* defined, subsequently we consider the tree-width and path-width of  $G - v$ .

**Theorem 17** *For a graph  $G$  and vertex  $v \in V(G)$ , it holds that*

$$tw(G) - 1 \leq tw(G - v) \leq tw(G) \quad \text{and} \\ pw(G) - 1 \leq pw(G - v) \leq pw(G).$$

**Proof** Removing  $v$  from every bag of a tree-decomposition (path-decomposition) for  $G$  and deleting all resulting, empty bags, we obtain a tree-decomposition (path-decomposition) for  $G - v$ . Consequently, we obtain  $tw(G - v) \leq tw(G)$  and  $pw(G - v) \leq pw(G)$ . Adding  $v$  and its incident edges to  $G - v$  results in  $G$ , such that the lower bounds follow from the upper bounds of Theorem 18.  $\square$



**Vertex Addition** Let  $G$  be a graph,  $N \subseteq V(G)$  a subset of vertices from  $G$ , and  $v \notin V(G)$  a newly introduced vertex. By  $G +_N v$  we denote the graph we obtain from  $G$  when inserting  $v$  with neighborhood  $N_G(v) = N$ , i.e.,

$$G +_N v = (V(G) \cup \{v\}, E(G) \cup \{\{v, u\} \mid u \in N\}).$$

In the special case, when  $N_G(v) = \{u\}$  holds for a vertex  $u \in V(G)$ , we call  $v$  a *pendant vertex* of  $G$ . If  $N_G(v) = V(G) \setminus \{v\}$  is true, we call  $v$  a *dominating vertex* of  $G$ . With *vertex addition* defined, we consider the tree-width and path-width of graph  $G +_N v$  in the ensuing theorem.

**Theorem 18** *For a graph  $G$ , a subset of vertices  $N \subseteq V(G)$ , and a vertex  $v \notin V(G)$  it holds that*

$$\begin{aligned} tw(G) &\leq tw(G +_N v) \leq tw(G) + 1 \quad \text{and} \\ pw(G) &\leq pw(G +_N v) \leq pw(G) + 1. \end{aligned}$$

**Proof** Introducing  $v$  to all bags of a tree-decomposition (path-decomposition) of  $G$ , we obtain a tree-decomposition (path-decomposition) of  $G +_N v$ . Thereby, it follows that  $tw(G +_N v) \leq tw(G) + 1$  and  $pw(G +_N v) \leq pw(G) + 1$  is true. Removing  $v$  from  $G +_N v$ , we obtain  $G$ . Consequently, the lower bounds follow from the upper bounds of Theorem 17.  $\square$

We can always add a pendant vertex to a graph without increasing the graph's tree-width. To do so, we introduce a new bag that contains the new vertex and its solely neighbor. Afterwards, the new bag is connected to any bag of a tree-decomposition for the graph, which already contains the neighbor.

**Corollary 19** *For a graph  $G$ , a vertex  $u \in V(G)$ , and a newly introduced vertex  $v \notin V(G)$ , it holds that*

$$tw(G +_{\{u\}} v) = \max(tw(G), 1).$$

Contrarily, the subsequent example shows that the previous statement does not hold with respect to path-width. Introducing a pendant vertex to a graph can increase its path-width.

*Example 3* The path-decomposition  $\mathcal{X} = (\{a, b\}, \{b, c\}, \{c, f\}, \{c, d\}, \{d, e\})$  shows that the path-width of graph  $C$  in Figure 2 is one. Introducing pendant vertex  $g$  to  $C$ , as depicted in graph  $I(K_{1,3})$  of the same figure, the path-width increases to two,  $\mathcal{X}' = (\{a, b\}, \{b, c\}, \{c, f, g\}, \{c, d\}, \{d, e\})$ .

### 3.2 Edge Addition and Edge Deletion

**Edge Deletion** Let  $G$  be a graph and  $v, w \in V(G)$  two vertices. For  $\{v, w\} \in E(G)$ , we define by  $G - \{v, w\}$  the graph we obtain from  $G$  by deleting edge  $\{v, w\}$ , i.e.,

$$G - \{v, w\} = (V(G), E(G) \setminus \{\{v, w\}\}).$$

With *edge deletion* defined, the subsequent theorem shows that removing an edge from a graph decreases the width of the graph at most by one.

**Theorem 20** *For a graph  $G$  and two different vertices  $v, w \in V(G)$  it holds that*

$$\begin{aligned} tw(G) - 1 &\leq tw(G - \{v, w\}) \leq tw(G) \quad \text{and} \\ pw(G) - 1 &\leq pw(G - \{v, w\}) \leq pw(G). \end{aligned}$$

**Proof** The upper bound follows immediately, because a tree-decomposition (path-decomposition) for  $G$  is also a tree-decomposition (path-decomposition) for  $G - \{v, w\}$ .

As  $G$  can be obtained from  $G - \{v, w\}$  by adding edge  $\{v, w\}$ , the lower bound follows from the upper bound of Theorem 21.  $\square$

**Edge Addition** Let  $G$  be a graph and  $v, w \in V(G)$  two vertices. For  $\{v, w\} \notin E(G)$ , we define by  $G + \{v, w\}$  the graph we obtain from  $G$  when adding edge  $\{v, w\}$ , i.e.,

$$G + \{v, w\} = (V(G), E(G) \cup \{\{v, w\}\}).$$

Having *edge addition* defined, our next theorem shows that inserting an edge into a graph increases the graph's width at most by one.

**Theorem 21** *For a graph  $G$  and two different vertices  $v, w \in V(G)$  it holds that*

$$\begin{aligned} tw(G) &\leq tw(G + \{v, w\}) \leq tw(G) + 1 \quad \text{and} \\ pw(G) &\leq pw(G + \{v, w\}) \leq pw(G) + 1. \end{aligned}$$

**Proof** Given a tree-decomposition (path-decomposition) for  $G$ , we obtain a tree-decomposition (path-decomposition) for  $G + \{v, w\}$  by adding one of the two vertices,  $v$  or  $w$ , to all its bags. Consequently,  $tw(G + \{v, w\}) \leq tw(G) + 1$  and  $pw(G + \{v, w\}) \leq pw(G) + 1$  hold.

The lower bounds follow by the fact that a tree-decomposition (path-decomposition) for  $G + \{v, w\}$  is also a tree-decomposition (path-decomposition) for  $G$ .  $\square$

### 3.3 Subgraph

So far we have only studied unary graph operations within this section of our work. However, in this subsection, we deviate from this path and study the unary graph transformation of *taking a subgraph*. This act of modifying a graph is not deterministic, since there are various subgraphs one can take from any given graph and it is not explicitly defined which one should be taken. Consequently, taking a subgraph of a graph multiple times can result in different subgraphs, making the studied modification a graph transformation but no graph operation.

Note, that taking a subgraph of a graph can be interpreted as a sequence of vertex deletion and edge deletion operations. Hence, every subgraph of a graph can be obtained by deleting selected vertices and edges from the original graph, such that the subsequent corollary follows immediately by Theorem 17 and Theorem 20.

**Corollary 22** *For a graph  $G$  and any subgraph  $H$  of  $G$  it holds that  $tw(H) \leq tw(G)$  and  $pw(H) \leq pw(G)$ .*

### 3.4 Vertex Identification

For a graph  $G$  and two different vertices  $v, w \in V(G)$ , the *identification* of  $v$  and  $w$  in  $G$ , denoted by  $\text{Ident}(G, v, w)$ , consists of vertex set  $(V(G) \setminus \{v, w\}) \cup \{u\}$  and edge set

$$\begin{aligned} E(G) \setminus \{\{x, y\} \mid x \in V(G), y \in \{v, w\}\} \\ \cup \{\{x, u\} \mid x \in (N_G(v) \cup N_G(w)) \setminus \{v, w\}\}. \end{aligned}$$

Given this definition of vertex identification, the ensuing result summarizes the graph operation's effect on the tree-width and path-width of the involved graph.

**Theorem 23** For a graph  $G$  and two different vertices  $v, w \in V(G)$  it holds that

$$\begin{aligned} tw(G) - 1 &\leq tw(\text{Ident}(G, v, w)) \leq tw(G) + 1 \quad \text{and} \\ pw(G) - 1 &\leq pw(\text{Ident}(G, v, w)) \leq pw(G) + 1. \end{aligned}$$

**Proof** For the upper bounds, let  $(\mathcal{X}, T)$  be a tree-decomposition ( $\mathcal{X}$  be a path-decomposition) for  $G$  of width  $tw(G)$  ( $pw(G)$ ). To obtain a tree-decomposition (path-decomposition) for  $\text{Ident}(G, v, w)$ , we proceed as follows. In a first step, replace all occurrences of  $v$  and  $w$  in all bags of  $\mathcal{X}$  by  $u$  and denote the result by  $\mathcal{X}'$ . Since  $\{v, w\}$  is not necessarily an edge of  $G$ ,  $\mathcal{X}'$  could violate (tw-3) ((pw-3)), i.e., the bags of  $\mathcal{X}'$  containing  $u$  might not be connected. In this case, we add  $u$  to all bags between the disconnected components. With that,  $(\mathcal{X}', T)$  is a valid tree-decomposition ( $\mathcal{X}'$  is a valid path-decomposition) for  $\text{Ident}(G, v, w)$ . To obtain  $\mathcal{X}'$  from  $\mathcal{X}$ , we increased the width at most by one, such that  $tw(\text{Ident}(G, v, w)) \leq tw(G) + 1$  ( $pw(\text{Ident}(G, v, w)) \leq pw(G) + 1$ ) follows.

For the lower bounds, we proceed as follows. First, rename vertex  $u$  of  $\text{Ident}(G, v, w)$  to  $v$  and denote the resulting graph by  $G'$ . Next, add a new vertex  $w$  with neighborhood  $N_G(w)$  to  $G'$ . By Theorem 18 it follows that  $tw(G' +_{N_G(w)} w) \leq tw(G') + 1$  and  $pw(G' +_{N_G(w)} w) \leq pw(G') + 1$  are true. Since  $G$  is a subgraph of  $G' +_{N_G(w)} w$ , with Corollary 22 we obtain that

$$\begin{aligned} tw(G) &\leq tw(G' +_{N_G(w)} w) \leq tw(\text{Ident}(G, v, w)) + 1 \quad \text{and} \\ pw(G) &\leq pw(G' +_{N_G(w)} w) \leq pw(\text{Ident}(G, v, w)) + 1 \end{aligned}$$

hold, which yields the lower bounds  $tw(G) - 1 \leq tw(\text{Ident}(G, v, w))$  and  $pw(G) - 1 \leq pw(\text{Ident}(G, v, w))$ .  $\square$

Note that the upper bounds specified in the previous theorem are tight, since a vertex identification on the end vertices of a path results in a cycle with increased tree-width and path-width.

### 3.5 Edge Contraction

In the case that the two vertices  $v, w \in V(G)$  of a vertex identification  $\text{Ident}(G, v, w)$  are adjacent, i.e.,  $\{v, w\} \in E(G)$ , we call the operation an *edge contraction*.

**Theorem 24** For a graph  $G$  and two different vertices  $v, w \in V(G)$  with  $\{v, w\} \in E(G)$  it holds that

$$\begin{aligned} tw(G) - 1 &\leq tw(\text{Ident}(G, v, w)) \leq tw(G) \quad \text{and} \\ pw(G) - 1 &\leq pw(\text{Ident}(G, v, w)) \leq pw(G). \end{aligned}$$

**Proof** Let  $(\mathcal{X}, T)$  be a tree-decomposition ( $\mathcal{X}$  be a path-decomposition) for  $G$  of width  $tw(G)$  ( $pw(G)$ ). We replace all occurrences of  $v$  and  $w$  in all bags of  $\mathcal{X}$  by  $u$  and denote the resulting decomposition by  $\mathcal{X}'$ . Since  $v$  and  $w$  are adjacent in  $G$ , we know by (tw-2) ((pw-2)) that there is at least one bag in  $\mathcal{X}$  which contains  $v$  and  $w$ . Consequently, (tw-3) ((pw-3)) must hold for  $\mathcal{X}'$  and it follows that  $(\mathcal{X}', T)$  is a valid tree-decomposition ( $\mathcal{X}'$  is a valid path-decomposition) for  $\text{Ident}(G, v, w)$  of width at most  $tw(G)$  ( $pw(G)$ ).<sup>4</sup>

---

<sup>4</sup>Because  $\{v, w\} \in E(G)$  is true, we can argue that (tw-3) ((pw-3)) holds for  $\mathcal{X}'$ . This argument is not valid for an arbitrary vertex identification, as  $v$  and  $w$  are not guaranteed to be adjacent, see Theorem 23.

The lower bounds follow by the same argumentation as for the lower bounds in the proof of Theorem 23.  $\square$

Contracting any edge of a clique  $K_n$  of size  $n$  results in a clique  $K_{n-1}$  of size  $n - 1$ . From Lemma 5 (Lemma 11) we know that  $\text{tw}(K_n) = n - 1$  ( $\text{pw}(K_n) = n - 1$ ) holds. Consequently, the lower bounds specified in the previous theorem are tight.

### 3.6 Edge Subdivision

Let  $G$  be a graph,  $u \notin V(G)$  a newly introduced vertex, and  $\{v, w\} \in E(G)$  an edge of  $G$ . The *edge subdivision* of  $\{v, w\}$  in  $G$ , denoted by  $\text{Subdiv}(G, v, w)$ , consists of vertex set  $V(G) \cup \{u\}$  and edge set  $(E(G) \setminus \{\{v, w\}\}) \cup \{\{v, u\}, \{u, w\}\}$ .

With this definition at hand, the subsequent theorem states the effect of an edge subdivision on the tree-width and path-width of a given graph.

**Theorem 25** *For a graph  $G$  and an edge  $\{v, w\} \in E(G)$  it holds that*

$$\begin{aligned} \text{tw}(\text{Subdiv}(G, v, w)) &= \text{tw}(G) \quad \text{and} \\ \text{pw}(G) &\leq \text{pw}(\text{Subdiv}(G, v, w)) \leq \text{pw}(G) + 1. \end{aligned}$$

**Proof**  $G$  is isomorphic to  $\text{Ident}(\text{Subdiv}(G, v, w), u, w)$ , such that the upper bounds of Theorem 24 yield

$$\begin{aligned} \text{tw}(G) &= \text{tw}(\text{Ident}(\text{Subdiv}(G, v, w), u, w)) \leq \text{tw}(\text{Subdiv}(G, v, w)) \quad \text{and} \\ \text{pw}(G) &= \text{pw}(\text{Ident}(\text{Subdiv}(G, v, w), u, w)) \leq \text{pw}(\text{Subdiv}(G, v, w)), \end{aligned}$$

resulting in this theorem's lower bounds.

For the upper bound of tree-width, let us distinguish the subsequent two cases.

**Case 1:**  $\text{tw}(G) = 1$

In this case,  $G$  is a forest. Since an edge subdivision does not alter this fact,  $\text{Subdiv}(G, v, w)$  is still a forest and  $\text{tw}(\text{Subdiv}(G, v, w)) = 1$  must hold.

**Case 2:**  $\text{tw}(G) > 1$

In this case, the biggest bag of every tree-decomposition of  $G$  contains at least three vertices. Furthermore, by (tw-2) it follows that in every tree-decomposition of  $G$  there is at least one bag which contains  $v$  and  $w$ . Let us denote this bag by  $X$ . Adding a new bag  $X'$  with vertices  $\{u, v, w\}$  and making it adjacent to  $X$  results in a tree-decomposition of  $\text{Subdiv}(G, v, w)$  with unaltered size, such that  $\text{tw}(\text{Subdiv}(G, v, w)) \leq \text{tw}(G)$  follows.

For the upper bound of path-width, we proceed as follows. By (pw-2), we know that in every path-decomposition of  $G$  there exists a bag containing  $v$  and  $w$ . Adding  $u$  to this bag, we obtain a valid path-decomposition for  $\text{Subdiv}(G, v, w)$ . Consequently,  $\text{pw}(\text{Subdiv}(G, v, w)) \leq \text{pw}(G) + 1$  follows.  $\square$

The upper bound for path-width given in the previous theorem is tight. The path-width of the caterpillar<sup>5</sup> graph  $C$  in Figure 2 equals one. Contrarily, the path-width of the graph obtained by subdividing edge  $\{c, f\}$  of  $C$ , depicted as  $I(K_{1,3})$  in Figure 2, is two.

Latest after subdividing all edges of a graph, the resulting graph must be bipartite. The resulting graph, one obtains after subdividing all edges of a graph  $G$ , is called the *incidence graph* of  $G$ , denoted by  $I(G)$ . The subsequent corollary provides bounds for a graph's incident graph and follows from Theorem 25.

<sup>5</sup>A *caterpillar* graph is a tree for which the removal of all pendant vertices results in a chordless path.




Figure 2: Special graphs.

**Corollary 26** *For a graph  $G$  it holds that*

$$tw(I(G)) = tw(G) \quad \text{and} \\ pw(G) \leq pw(I(G)) \leq pw(G) + 1.$$

**Proof** The result for tree-width follows directly by Theorem 25.

Contracting all subdivided edges of  $I(G)$ , we obtain  $G$ . By Theorem 24 we know that edge contractions do not increase the resulting graph's path-width, such that  $pw(G) \leq pw(I(G))$  follows.

Denote by  $\mathcal{X}$  a path-decomposition for  $G$  of width  $pw(G)$  and let  $\{v, w\} \in E(G)$  be an edge of  $G$  as well as  $x_{\{v,w\}}$  the node introduced to subdivide the edge. By (pw-2) we know that there is a bag  $X$  in  $\mathcal{X}$  with  $v, w \in X$ . We introduce a new bag  $X' = X \cup \{x_{\{v,w\}}\}$  to  $\mathcal{X}$  as successor of  $X$  and denote the resulting decomposition  $\mathcal{X}'$ . Then,  $\mathcal{X}'$  is a path-decomposition for  $\text{Subdiv}(G, v, w)$  of width  $pw(G) + 1$ . We can repeat this step for every edge of  $G$  and finally, obtain a path-decomposition for  $I(G)$  of width  $pw(G) + 1$ , such that  $pw(I(G)) \leq pw(G) + 1$  follows.  $\square$

### 3.7 Minor

Every graph  $H$  one may obtain from a given graph  $G$ , by applying a finite sequence of arbitrary edge deletion and edge contraction operations, is called a *minor* of  $G$ . Obviously, taking a minor is a graph transformation but not a graph operation, as the applied sequence of operations is not deterministic.

Given this definition, the subsequent theorem follows immediately from Theorem 20 and Theorem 24.

**Theorem 27** *For a graph  $G$  and a minor  $H$  of  $G$ , it holds that  $tw(H) \leq tw(G)$  and  $pw(H) \leq pw(G)$ .*

In other words, the fact that a graph has tree-width (path-width) at most  $k \in \mathbb{N}$  is preserved under the graph transformation of taking a minor.

Subsequently, we cite one of the probably most important theorems in graph theory, the *minor theorem*, shown by Robertson and Seymour [RS04].

**Theorem 28 (Minor theorem)** *In every infinite set of graphs there are two graphs, such that one of them is a minor of the other.*

Before the minor theorem was proven by Robertson and Seymour, it was known as Wagner's conjecture [Wag70]. The following theorem is an important corollary of the minor theorem [RS04].

**Theorem 29** *Every set of graphs that is closed under minors can be defined by a finite set of forbidden minors.*

For  $k \in \mathbb{N}$ , Theorem 27 implies that the set of graphs with tree-width (path-width) at most  $k$  is closed under minors. Consequently, Theorem 29 then implies that the set of graphs with tree-width (path-width) at most  $k$  can be characterized by a finite set of forbidden minors. For small values of  $k$  these sets of forbidden minors are known. For example, Kinnersley and Langston [KL94] characterized the set of all graphs with path-width at most one by the forbidden minors  $K_3$  and  $I(K_{1,3})$ , c.f. Figure 2. Furthermore, they characterized the set of graphs with path-width at most two by 110 forbidden minors [KL94]. The set of all graphs with tree-width at most one is characterized by the forbidden minor  $K_3$  and the set of all graphs with tree-width at most two is characterized by the forbidden minor  $K_4$ , see the work by Bodlaender and van Antwerpen-de Fluiter [BvAdF01]. For the four forbidden minors that characterize the set of all graphs with tree-width at most three, see the work by Arnborg, Proskurowski, and Corneil [APC90].

The main algorithmic consequence of the minor theorem is stated in the subsequent theorem, the *minor test*, also shown by Robertson and Seymour [RS95].

**Theorem 30 (Minor test)** *For a fixed graph  $H$  and a given graph  $G$ , one can decide in time in  $\mathcal{O}(|V(G)|^3)$  whether  $H$  is a minor of  $G$ .<sup>6</sup>*

From a parameterized point of view, the minor test is fixed-parameter tractable with respect to the parameter  $\text{size}(H)$ . However, the constant behind the  $\mathcal{O}$ -notation in Theorem 30 depends on the parameter  $\text{size}(H)$  and is huge. Kawarabayashi, Kobayashi, and Reed [KKR12] showed an improved version of the previous result, allowing to decide the minor test in time in  $\mathcal{O}(|V(G)|^2)$ .

The ensuing corollary follows immediately by Theorem 29 and the application of Theorem 30 for every forbidden minor.

**Corollary 31** *For every set of graphs  $\mathcal{G}$  that is closed under minors and every graph  $G$  one can decide in time polynomial in the size of  $G$  whether  $G \in \mathcal{G}$  holds.*

Nevertheless, an implicit precondition of the corollary is that the set of all forbidden minors characterizing  $\mathcal{G}$  is known.

Another result by Robertson and Seymour [RS86b] is the so-called *grind minor theorem*, also known as the *excluded grid theorem*.

**Theorem 32 (Grid minor theorem)** *There is a function  $f: \mathbb{N} \rightarrow \mathbb{N}$ , such that every graph with tree-width at least  $f(k)$  has a  $k \times k$ -grid graph as minor.*

Even if the grid minor theorem seems very technical at first, it has a direct effect for the tree-width of a graph, as the subsequent theorem from the same work by Robertson and Seymour [RS86b] shows.

**Theorem 33** *For every planar graph  $H$  there is an integer  $k \in \mathbb{N}$ , such that every graph without  $H$  as minor has tree-width at most  $k$ .*

Ahead of this result for tree-width, Robertson and Seymour [RS83] proved an analogue result with respect to path-width.

**Theorem 34** *For every forest  $H$  there is an integer  $k \in \mathbb{N}$ , such that every graph without  $H$  as minor has path-width at most  $k$ .*

---

<sup>6</sup>A more precise formulation of the runtime would be  $\mathcal{O}(f(k)|V(G)|^3)$  for some function  $f$  and  $k \leq \text{size}(H)$ .

### 3.8 Power of a Graph

For  $d \in \mathbb{N}$  and a graph  $G$ , we denote the  $d$ -th power of  $G$  by  $G^d$ . Thereby,  $G^d$  possesses the same set of vertices as  $G$ , i.e.,  $V(G^d) = V(G)$ , and two vertices of  $G^d$  are adjacent if and only if there exists a path of length at most  $d$  in  $G$  between these vertices.

To begin, the subsequent lemma provides an upper bound on the number of neighbors each vertex might possess in the  $d$ -th power of a graph  $G$ .

**Lemma 35** *Given  $d \in \mathbb{N}$  and a graph  $G$ , for every vertex  $v \in V(G)$  it holds that*

$$\deg_{G^d}(v) \leq \Delta(G) \sum_{i=0}^{d-1} (\Delta(G) - 1)^i.$$

**Proof** Let  $v \in V(G)$  be a vertex of  $G$ . To obtain an upper bound on the number of neighbors  $v$  might possess in  $G^d$ , we derive an upper bound for the number of vertices reachable from  $v$  within a distance of at most  $d$  in  $G$ . By definition, we know that  $\deg_G(v) \leq \Delta(G)$  holds, such that  $v$  has at most  $\Delta(G)$  neighbors in  $G$ . By the same argument, every neighbor  $u$  of  $v$  in  $G$  has at most  $\Delta(G)$  neighbors. However, one of those neighbors of  $u$  is  $v$ , such that  $u$  has at most  $\Delta(G) - 1$  not yet considered neighbors. Repeating the previous argument  $d - 1$  times, we obtain an upper bound on the number of vertices with exact distance  $d$  to  $v$ ,

$$\Delta(G)(\Delta(G) - 1)^{d-1}.$$

Since we are interested in an upper bound on the number of vertices with a distance of at most  $d$  to  $v$ , we need to sum up the upper bounds for all intermediate distances from one up to  $d$ . Doing so, we obtain

$$\Delta(G) + \Delta(G)(\Delta(G) - 1) + \dots + \Delta(G)(\Delta(G) - 1)^{d-1} = \Delta(G) \sum_{i=0}^{d-1} (\Delta(G) - 1)^i.$$

As every vertex, which is reachable from  $v$  in  $G$  within a distance of at most  $d$ , is adjacent to  $v$  in  $G^d$ , it follows that

$$\deg_{G^d}(v) \leq \Delta(G) \sum_{i=0}^{d-1} (\Delta(G) - 1)^i$$

holds. □

Note that the previously stated upper bound is tight. To see this, consider a tree  $T$  with root  $v \in V(T)$  and three binary subtrees of equal, but arbitrary depth adjacent to  $v$ .

Having this upper bound for a vertex's number neighbors in the  $d$ -th power of a graph at hand, the subsequent theorem makes use of it to study the effect of raising a graph to the power of  $d$  with respect to tree-width and path-width.

**Theorem 36** *For  $d \in \mathbb{N}$  and a graph  $G$  it holds that*

$$tw(G) \leq tw(G^d) \leq (tw(G) + 1) \left( 1 + \Delta(G) \sum_{i=0}^{d-1} (\Delta(G) - 1)^i \right) - 1 \quad \text{and}$$

$$pw(G) \leq pw(G^d) \leq (pw(G) + 1) \left( 1 + \Delta(G) \sum_{i=0}^{d-1} (\Delta(G) - 1)^i \right) - 1.$$

**Proof** The lower bounds follow by Corollary 22, as  $G$  is a subgraph of  $G^d$ .

For the upper bounds, denote by  $(\mathcal{X}, T)$  a tree-decomposition (by  $\mathcal{X}$  a path-decomposition) of  $G$ . Constructing  $G^d$ , we know all neighbors of every vertex  $v \in V(G)$  in  $G^d$ . To obtain a tree-decomposition (path-decomposition) for  $G^d$ , for every vertex  $v \in V(G)$ , we add all neighbors of  $v$  in  $G^d$ , i.e.,  $N_{G^d}(v)$ , to every bag  $X \in \mathcal{X}$  containing  $v$ . We denote the resulting bag by  $X'$  and the resulting set of bags by  $\mathcal{X}'$ . Afterwards, by Lemma 35, every bag of  $\mathcal{X}'$  contains at most

$$\begin{aligned} & (\text{tw}(G) + 1) \left( 1 + \Delta(G) \sum_{i=0}^{d-1} (\Delta(G) - 1)^i \right) \\ & \left( (\text{pw}(G) + 1) \left( 1 + \Delta(G) \sum_{i=0}^{d-1} (\Delta(G) - 1)^i \right) \right) \end{aligned}$$

vertices. Taking  $G$  to the power of  $d$  does not alter the set of vertices, i.e.,  $V(G) = V(G^d)$ , so that  $(\mathcal{X}', T)$  ( $\mathcal{X}'$ ) still satisfies (tw-1) ((pw-1)). Adding, for every vertex  $v \in V(G)$ , all neighbors of  $v$  in  $G^d$  to every bag containing  $v$ , all new edges introduced to  $G^d$  are covered, such that  $(\mathcal{X}', T)$  ( $\mathcal{X}'$ ) satisfies (tw-2) ((pw-2)). To see that  $(\mathcal{X}', T)$  ( $\mathcal{X}'$ ) satisfies (tw-3) ((pw-3)), let  $v \in V(G)$  be any vertex of  $G$  and denote by  $X'_s, X'_t \in \mathcal{X}'$  two bags with  $v \in X'_s, X'_t$ . We separate three cases: (i) If it holds that  $v \in X_s, X_t$  for  $X_s, X_t \in \mathcal{X}$ , it immediately follows that  $v$  is also in all bags connecting  $X'_s$  and  $X'_t$ , since  $(\mathcal{X}, T)$  ( $\mathcal{X}$ ) satisfies (tw-3) ((pw-3)). (ii) If it holds that  $v \in X_s$  but  $v \notin X_t$ , there must be a vertex  $u \in X_t$  with  $\text{dist}_G(u, v) \leq d$ , as otherwise  $v \in X'_t$  would not hold. Since  $\text{dist}_G(u, v) \leq d$ , there is a path  $p_G$  between  $u$  and  $v$  in  $G$  of length at most  $d$ . Consequently, with  $(\mathcal{X}, T)$  ( $\mathcal{X}$ ) satisfying (tw-2) ((pw-2)) and (tw-3) ((pw-3)), there must also be a path  $p_T$  between  $s$  and  $t$ , such that for every  $t'$  in  $p_T$  there is a vertex  $v'$  in  $p_G$  with  $v' \in X_{t'}$ . For every such vertex  $v'$  it holds that  $\text{dist}_G(v', v) \leq \text{dist}_G(u, v) \leq d$ , so that by the construction of  $\mathcal{X}'$  it follows that  $v \in X'_{t'}$  holds. Therefore, also in this case all bags between  $X'_s$  and  $X'_t$  must contain  $v$ . (iii) Finally, if  $v \notin X_s$  and  $v \notin X_t$ , there must be at least one bag  $X_r \in \mathcal{X}$  with  $v \in X_r$  and we can repeat the previous argument for  $X'_r, X'_s$  and  $X'_r, X'_t$ . Consequently,  $(\mathcal{X}', T)$  ( $\mathcal{X}'$ ) is a tree-decomposition (path-decomposition) for  $G^d$  of width

$$\begin{aligned} & (\text{tw}(G) + 1) \left( 1 + \Delta(G) \sum_{i=0}^{d-1} (\Delta(G) - 1)^i \right) - 1 \\ & \left( (\text{pw}(G) + 1) \left( 1 + \Delta(G) \sum_{i=0}^{d-1} (\Delta(G) - 1)^i \right) - 1 \right) \end{aligned}$$

and the upper bound follows.  $\square$

### 3.9 Line Graph

In this subsection, we study the graph operation of creating a graph's line graph, using the notation as formulated by Harary and Norman [HZ60]. For a graph  $G$ , its *line graph*,  $L(G)$ , is defined by

$$\begin{aligned} V(L(G)) &= \{x_{\{u,v\}} \mid \{u,v\} \in E(G)\} \quad \text{and} \\ E(L(G)) &= \{\{x_e, x_f\} \mid |e \cap f| = 1\}. \end{aligned}$$

In other words, the line graph  $L(G)$  of a graph  $G$  has a vertex for every edge of  $G$  and two vertices of  $L(G)$  are adjacent, if the corresponding edges in  $G$  are adjacent. The concept of



a line graph, although not explicitly called line graph, was first used by Whitney [Whi32] in 1932.

Furthermore, the underlying graph  $G$  of a given line graph  $L(G)$  is called the *root graph* of  $L(G)$ .

Given a graph  $G$ , it is possible to bound the tree-width (path-width) of its line graph in the tree-width (path-width) of  $G$  and  $G$ 's maximum vertex degree.

**Theorem 37** *For a graph  $G$  it holds that*

$$tw(G) - 1 \leq tw(L(G)) \leq (tw(G) + 1)\Delta(G) - 1 \quad \text{and}$$

$$\frac{1}{2}(pw(G) + 1) - 1 \leq pw(L(G)) \leq (pw(G) + 1)\Delta(G) - 1.$$

**Proof** The stated lower bound for the tree-width of  $L(G)$  was shown by Harvey and Wood [HW18] in Proposition 2.3.

A slightly weaker bound can be obtained as follows. Let  $(\mathcal{X}, T)$  be a tree-decomposition for  $L(G)$  of width  $tw(L(G))$ . In every bag of  $(\mathcal{X}, T)$  replace each edge of  $G$  by both of its endpoints. Then, we can obtain a tree-decomposition for  $G$  of width at most  $2(tw(L(G)) + 1) - 1$ , such that  $\frac{1}{2}(tw(G) + 1) - 1 \leq tw(L(G))$  follows. A similar argument results in the stated lower bound for path-width.

The upper bounds are known from several works [Ats08, Bod93, CFR03] and can be obtained as follows. Let  $(\mathcal{X}, T)$  be a tree-decomposition ( $\mathcal{X}$  a path-decomposition) for  $G$  of width  $tw(G)$  ( $pw(G)$ ). If we replace every bag  $X_u \in \mathcal{X}$  by the set of all edges incident to at least one vertex in  $X_u$ , we obtain a tree-decomposition (path-decomposition) for  $L(G)$  of width at most  $(tw(G) + 1)\Delta(G) - 1$  ( $(pw(G) + 1)\Delta(G) - 1$ ).  $\square$

Stricter upper bounds than the ones shown in Theorem 37 can be found in Theorem 1.3 of the work by Harvey and Wood [HW18].

Furthermore, it is easy to confirm that for every graph  $G$ , the edges incident to a vertex of  $G$  with degree  $\Delta(G)$  form a clique of size  $\Delta(G)$  in  $L(G)$ . With Lemma 5 and Lemma 11, the subsequent corollary follows immediately.

**Corollary 38** *For a graph  $G$  it holds that  $tw(L(G)) \geq \Delta(G) - 1$  and  $pw(L(G)) \geq \Delta(G) - 1$ .*

For special graphs the inequality turns into an equality, as the subsequent result shows.

**Proposition 39** *Let  $G$  be a graph. If  $tw(G) = 1$  is true, it holds that  $tw(L(G)) = \Delta(G) - 1$ . If  $pw(G) = 1$  is true, it holds that  $pw(L(G)) = \Delta(G) - 1$ .*

The statement for tree-width was shown by Harvey and Wood [HW18]. The statement for path-width follows by two arguments. For the lower bound we refer to Corollary 38. For the upper bound, we know from Section 3.7 that a graph  $G$  of path-width one can be identified as disjoint union of caterpillar graphs, allowing to construct a path-decomposition of width at most  $\Delta(G) - 1$  for  $L(G)$ .

Beside providing a direct equation how to obtain a line graph's tree-width (path-width) from its root graph's tree-width (path-width), if the root graph is a forest (caterpillar graph), the previous proposition shows that a line graph's tree-width (path-width) can not be bounded in its root graph's tree-width (path-width).

We conclude this subsection with a theorem by Harvey and Wood [HW15], showing how to derive a line graph's tree-width (path-width) from its root graph, if the root graph is a complete graph.

**Theorem 40** For  $n \geq 2$  it holds that

$$tw(L(K_n)) = pw(L(K_n)) = \begin{cases} \binom{\frac{n-1}{2}}{\frac{n-1}{2}} + n - 2, & \text{if } n \text{ is odd,} \\ \binom{\frac{n-2}{2}}{\frac{n}{2}} + n - 2, & \text{if } n \text{ is even.} \end{cases}$$

### 3.10 Edge Complement

In this subsection of our work, we study the graph operation of creating a graph's edge complement graph. Thereby, the *edge complement graph* of a graph  $G$ , denoted by  $\text{co-}G$ , has the same vertex set as  $G$  and two vertices are adjacent in  $\text{co-}G$  if and only if they are not adjacent in  $G$ , i.e.,

$$\begin{aligned} V(\text{co-}G) &= V(G) \quad \text{and} \\ E(\text{co-}G) &= \{\{u, v\} \mid u, v \in V(G), u \neq v, \{u, v\} \notin E(G)\}. \end{aligned}$$

Let  $K_{1,\ell}$  be a star graph with one dominating vertex  $v$  in the center and  $\ell$  vertices  $u_i$ ,  $1 \leq i \leq \ell$ , as satellites, all only adjacent to  $v$ . Then, the edge complement graph of  $K_{1,\ell}$ ,  $\text{co-}K_{1,\ell}$ , consists of an isolated vertex  $v$  and a clique of size  $\ell$  formed by all satellites  $u_i$ ,  $1 \leq i \leq \ell$ . Since neither  $K_3$  nor  $I(K_{1,3})$  is a minor of  $K_{1,\ell}$ , we know from Subsection 3.7 that  $\text{pw}(K_{1,\ell}) = \text{tw}(K_{1,\ell}) = 1$  holds. However, by Lemma 5 and Lemma 11 it follows that  $\text{tw}(\text{co-}K_{1,\ell}) = \text{pw}(\text{co-}K_{1,\ell}) = \ell - 1$  is true. Therefore, it is generally impossible to bound the tree-width (path-width) of an edge complement graph  $\text{co-}G$  in the tree-width (path-width) of the original graph  $G$ .

Nevertheless, Joret and Wood [JW12] proved the subsequent theorem, providing a lower bound for the tree-width of a graph's edge complement graph.<sup>7</sup>

**Theorem 41** For a graph  $G$  it holds that

$$tw(G) + tw(\text{co-}G) \geq |V(G)| - 2.$$

In their work, Joret and Wood also showed that the specified bound is tight. As  $\text{pw}(G) \geq \text{tw}(G)$  is true for every graph  $G$ , the subsequent corollary follows immediately.

**Corollary 42** For a graph  $G$  it holds that

$$pw(G) + pw(\text{co-}G) \geq |V(G)| - 2.$$

For the path with four vertices,  $P_4$ , it holds that  $\text{co-}P_4 \cong P_4$  as well as  $\text{pw}(P_4) = 1$ . Consequently, we obtain

$$\text{pw}(P_4) + \text{pw}(\text{co-}P_4) = \text{pw}(P_4) + \text{pw}(P_4) = 2 = |V(P_4)| - 2,$$

such that the bound specified in Corollary 42 is tight, too.

### 3.11 Local Complementation

In his work, Bouchet [Bou94] introduced the graph operation local complementation. Given a graph  $G$  and a vertex  $v \in V(G)$ , the *local complementation* of  $G$ , denoted by  $LC(G, v)$ , is defined by

$$\begin{aligned} V(LC(G, v)) &= V(G) \quad \text{and} \\ E(LC(G, v)) &= E(G) \setminus \{\{u, w\} \mid u, w \in N_G(v)\} \\ &\quad \cup \{\{u, w\} \mid u, w \in N_G(v), u \neq w, \{u, w\} \notin E(G)\}. \end{aligned}$$

<sup>7</sup>Formulating bounds of the form  $f(G) + f(\text{co-}G)$  for a graph parameter  $f$  is known as *Nordhaus-Gaddum problem*.

In other words,  $LC(G, v)$  is obtained from graph  $G$  by replacing the subgraph of  $G$  induced by  $N_G(v)$  with its edge complement. Recall that  $v \notin N_G(v)$  holds, such that the neighborhood of  $v$  in  $LC(G, v)$  is the same as in  $G$ .

Denote by  $K_{1,\ell}$  the star graph we made already use of in Subsection 3.10. Applying a local complementation to the dominating vertex  $v$ , it is easy to see that  $LC(K_{1,\ell}, v)$  equals a clique of size  $\ell + 1$ . The star  $K_{1,\ell}$  has tree-width and path-width one, while  $LC(K_{1,\ell}, v)$  has tree-width and path-width  $\ell$ . Therefore, in general the tree-width (path-width) of a graph  $G$ 's local complement  $LC(G, v)$  can not be bounded in the tree-width (path-width) of  $G$ .

### 3.12 Seidel Switching

The Seidel switching operation was introduced by the Dutch mathematician J. J. Seidel in connection with regular structures, such as systems of equiangular lines, strongly regular graphs, or so-called two-graphs [Sei74, Sei76, ST81]. Several examples for applications of Seidel switching can be found in algorithms, e.g., in a polynomial-time algorithm for the  $P_3$ -structure recognition problem [Hay96] or in an algorithm for the construction of bi-join decompositions of graphs [dMR05].

For a graph  $G$  and a vertex  $v \in V(G)$ , the graph resulting from a *Seidel switching* operation, denoted by  $S(G, v)$ , is defined as follows. The vertex set of  $S(G, v)$  is the same as the vertex set of  $G$ , i.e.,  $V(S(G, v)) = V(G)$ , and the edge set of  $S(G, v)$  is defined as

$$E(S(G, v)) = E(G) \setminus \{\{v, w\} \mid w \in N_G(v)\} \\ \cup \{\{v, w\} \mid w \in V(G) \setminus (N_G(v) \cup \{v\})\}.$$

In other words, every neighbor of  $v$  in  $G$  is a non-neighbor of  $v$  in  $S(G, v)$  and every non-neighbor of  $v$  in  $G$  is a neighbor of  $v$  in  $S(G, v)$ .

Given this definition of Seidel switching, subsequently, extending a result by Bodlaender and Hage [BH12], we show that a single Seidel switching operation increases or decreases a graph's tree-width and path-width at most by one.

**Theorem 43** *For a graph  $G$  and a vertex  $v \in V(G)$  it holds that*

$$tw(G) - 1 \leq tw(S(G, v)) \leq tw(G) + 1 \quad \text{and} \\ pw(G) - 1 \leq pw(S(G, v)) \leq pw(G) + 1.$$

**Proof** For the upper bounds, let  $(\mathcal{X}, T)$  be a tree-decomposition ( $\mathcal{X}$  a path-decomposition) for  $G$  of width  $tw(G)$  ( $pw(G)$ ). When we add  $v$  to all bags of  $\mathcal{X}$ , denoting the resulting set of bags by  $\mathcal{X}'$ , we obtain a tree-decomposition  $(\mathcal{X}', T)$  (a path-decomposition  $\mathcal{X}'$ ) for  $S(G, v)$  of width at most  $tw(G) + 1$  ( $pw(G) + 1$ ). Consequently,  $tw(S(G, v)) \leq tw(G) + 1$  ( $pw(S(G, v)) \leq pw(G) + 1$ ) follows.

Since  $S(S(G, v), v) = G$  holds, we can derive the lower bound from the upper bound via  $tw(G) = tw(S(S(G, v), v)) \leq tw(S(G, v)) + 1$  ( $pw(G) = pw(S(S(G, v), v)) \leq pw(S(G, v)) + 1$ ).  $\square$

Note that the bounds shown in Theorem 43 are tight. To convince oneself of this fact, consider the path with five vertices,  $P_5$ . Its tree-width and path-width is one. Denote by  $v$  one of the two vertices in  $P_5$  with degree one. Then,  $S(P_5, v)$  contains  $K_3$  as minor, such that a tree-width and path-width of at least two follows. Following the example in the opposite direction, i.e., applying the Seidel switching operation to  $S(P_5, v)$  for the same vertex  $v$ , provides an example that the lower bound is tight, too.

In 1980, Colbourn and Corneil [CC80] studied the complexity of the decision problem whether two graphs are switching equivalent. In their work, they proved that this decision

problem is polynomial time equivalent to the decision problem of graph isomorphism. Thereby, two graphs  $G$  and  $G'$  with the same vertex set  $V$  are called *switching equivalent*, if there exists a sequence of vertices  $(v_1, \dots, v_\ell)$  in  $V$ , such that for  $G^0 = G$  and  $G^i = S(G^{i-1}, v_i)$ ,  $1 \leq i \leq \ell$ , it holds that  $G^\ell = G'$ . In 2012, Bodlaender and Hage [BH12] considered in their work the tree-width of switching classes. By the definition of switching equivalence and via Theorem 43, we can formulate the subsequent corollary, contributing to the research on the tree-width of switching classes, initiated by Bodlaender and Hage.

**Corollary 44** *Let  $G$  and  $G'$  be two switching equivalent graph and denote by  $(v_1, \dots, v_\ell)$  a sequence of vertices, such that  $G^\ell = G'$  is true. Then, it holds that*

$$tw(G') \leq tw(G) + \ell \quad \text{and} \quad pw(G') \leq pw(G) + \ell.$$

### 3.13 Seidel Complementation

Limouzy [Lim10] defined the Seidel complementation operation in order to give a characterization for permutation graphs. For a graph  $G$  and a vertex  $v \in V(G)$ , the graph resulting from the *Seidel complementation* operation, denoted by  $SC(G, v)$ , has the same vertices as  $G$ , i.e.,  $V(SC(G, v)) = V(G)$ , and edge set

$$E(SC(G, v)) = E(G) \Delta \{\{x, y\} \mid \{v, x\} \in E(G), \{v, y\} \notin E(G)\}.$$

In other words, the edge set of  $SC(G, v)$  equals the edge set of  $G$  with edges and non-edges between the neighborhood and non-neighborhood of  $v$  complemented.

Let  $G$  be a graph that consists of two parts. The first part is a star with vertex  $v$  as dominating vertex and  $\ell \in \mathbb{N}$  satellites  $u_1, \dots, u_\ell$ , adjacent to  $v$ . The second part is a set of  $\ell$  isolated vertices,  $w_1, \dots, w_\ell$ . Since  $G$  does neither possess a  $K_3$  nor an  $I(K_{1,3})$  as minor, we know that  $tw(G) = pw(G) = 1$  holds. Applying the Seidel complementation operation to vertex  $v$ ,  $SC(G, v)$  contains a complete, bipartite subgraph formed by the vertices  $u_1, \dots, u_\ell, w_1, \dots, w_\ell$ . By Lemma 6 (Lemma 12) it follows that  $tw(SC(G, v)) \geq \ell - 1$  ( $pw(SC(G, v)) \geq \ell - 1$ ) is true. Consequently, we conclude that, given a graph  $G$  and a vertex  $v \in V(G)$ , the tree-width (path-width) of  $SC(G, v)$  cannot be bounded in the tree-width (path-width) of  $G$ .

## 4 Binary Graph Operations

Let  $G_1, G_2$  be two non-empty graphs and  $f$  a binary graph operation that creates a new graph  $f(G_1, G_2)$  from  $G_1$  and  $G_2$ . In this section, we consider the tree-width and path-width of  $f(G_1, G_2)$  with respect to the tree-widths and path-widths of the initial graphs  $G_1$  and  $G_2$ . In particular, we study the binary graph operations disjoint union, join, union, substitution, various types of graph products, 1-sum, and corona.

### 4.1 Disjoint Union

The *disjoint union* of two vertex-disjoint graphs  $G_1$  and  $G_2$ , denoted by  $G_1 \oplus G_2$ , is defined as the graph with vertex set  $V(G_1) \cup V(G_2)$  and edge set  $E(G_1) \cup E(G_2)$ .

Bodlaender and Möhring [BM90, BM93] proved the subsequent theorem with respect to the tree-width and path-width of a graph which is the disjoint union of two vertex-disjoint graphs.

**Theorem 45** *Let  $G_1$  and  $G_2$  be two vertex-disjoint graphs, then it holds that  $tw(G_1 \oplus G_2) = \max(tw(G_1), tw(G_2))$  and  $pw(G_1 \oplus G_2) = \max(pw(G_1), pw(G_2))$ .*




Figure 3:  $G_1$ ,  $G_2$ , and  $G_1 \cup G_2$  for  $m = n = 3$  in Example 4.

These bounds imply that the tree-width and path-width of a graph can be derived from the tree-width and path-width of its connected components.

**Corollary 46** *Let  $G$  be a graph. It holds that*

1. *the tree-width of  $G$  is the maximum tree-width and*
2. *the path-width of  $G$  is the maximum path-width*

*of its connected components.*

## 4.2 Join

The *join* of two vertex-disjoint graphs  $G_1$  and  $G_2$ , denoted by  $G_1 \otimes G_2$ , is defined as the graph with vertex set  $V(G_1) \cup V(G_2)$  and edge set

$$E(G_1) \cup E(G_2) \cup \{\{v_1, v_2\} \mid v_1 \in V(G_1), v_2 \in V(G_2)\}.$$

As for the disjoint union of two graphs, Bodlaender and Möhring [BM90, BM93] proved the subsequent theorem with respect to the tree-width and path-width of a graph that is the join of two vertex-disjoint graphs.

**Theorem 47** *Let  $G_1$  and  $G_2$  be two vertex-disjoint graphs. Then, for the join of  $G_1$  and  $G_2$  it holds that  $tw(G_1 \otimes G_2) = \min(tw(G_1) + |V(G_2)|, tw(G_2) + |V(G_1)|)$  and  $pw(G_1 \otimes G_2) = \min(pw(G_1) + |V(G_2)|, pw(G_2) + |V(G_1)|)$ .*

The combination of Theorem 45 and 47 implies that for every co-graph  $G$ , it holds that  $tw(G) = pw(G)$  and both widths can be computed in linear time [BM90, BM93].

## 4.3 Union

The *union* of two graphs  $G_1$  and  $G_2$  with  $V(G_1) = V(G_2)$ , denoted by  $G_1 \cup G_2$ , is defined as the graph with vertices  $V(G_1)$  and edge set  $E(G_1) \cup E(G_2)$ . Thus, two vertices are adjacent in  $G_1 \cup G_2$  if and only if they are adjacent in  $G_1$  or in  $G_2$ .

In general, it is not possible to bound the tree-width (path-width) of the union of two graphs in terms of the tree-widths (path-widths) of the individual graphs. To see why that is the case, consider the subsequent example.

*Example 4* For  $m, n \in \mathbb{N}$ , define the set of vertices  $V = \{v_{i,j} \mid 1 \leq i \leq m, 1 \leq j \leq n\}$ . Next, denote by  $G_1$  the disjoint union of  $m$  paths with  $n$  vertices from  $V$ ,  $P_n^i = (\{v_{i,j} \mid 1 \leq j \leq n\}, \{\{v_{i,j}, v_{i,j+1}\} \mid 1 \leq j < n\})$ ,  $1 \leq i \leq m$ , and by  $G_2$  the disjoint union of  $n$  paths with  $m$  vertices from  $V$ ,  $P_m^j = (\{v_{i,j} \mid 1 \leq i \leq m\}, \{\{v_{i,j}, v_{i+1,j}\} \mid 1 \leq i < m\})$ ,  $1 \leq j \leq n$ . Since paths have a tree-width (path-width) of 1, it follows by Theorem 45 that  $tw(G_1) = tw(G_2) = 1$  ( $pw(G_1) = pw(G_2) = 1$ ) holds.

The union,  $G_1 \cup G_2$ , of  $G_1$  and  $G_2$  is an  $n \times m$ -grid graph, and Bodlaender [Bod98] proved that the tree-width (path-width) of an  $n \times m$ -grid graph equals  $\min(n, m)$ . Consequently, it is not possible to bound the tree-width (path-width) of  $G_1 \cup G_2$  in the tree-widths (path-widths) of  $G_1$  and  $G_2$ .

See Figure 3 for an explicit example of  $G_1, G_2$ , and the resulting union  $G_1 \cup G_2$ .

#### 4.4 Substitution

Let  $G_1$  and  $G_2$  be two vertex-disjoint graphs and  $v \in V(G_1)$  a vertex. The *substitution* of  $v$  by  $G_2$  in  $G_1$ , denoted by  $G_1[v/G_2]$ , is defined as the graph with vertex set  $(V(G_1) \setminus \{v\}) \cup V(G_2)$  and edge set

$$(E(G_1) \setminus \{\{v, w\} \mid w \in N_{G_1}(v)\}) \cup \{\{u, w\} \mid u \in V(G_2), w \in N_{G_1}(v)\} \cup E(G_2).$$

The vertex set  $V(G_2)$  is called a *module* of graph  $G_1[v/G_2]$ , since all vertices of  $V(G_2)$  are adjacent to the same vertices of  $V(G_1[v/G_2]) \setminus V(G_2)$  in  $G_1[v/G_2]$ .

Given this definition of substitution, the subsequent theorem considers the tree-width (path-width) of  $G_1[v/G_2]$  in terms of the tree-width (path-width) of  $G_1$  and  $G_2$ .

**Theorem 48** *For two vertex-disjoint graphs  $G_1, G_2$  and vertex  $v \in V(G_1)$  it holds that*

$$\max(\text{tw}(G_1), \text{tw}(G_2)) \leq \text{tw}(G_1[v/G_2]) \leq \min(\text{tw}(G_1) + |V(G_2)|, \text{tw}(G_2) + |V(G_1)|) - 1$$

and

$$\max(\text{pw}(G_1), \text{pw}(G_2)) \leq \text{pw}(G_1[v/G_2]) \leq \min(\text{pw}(G_1) + |V(G_2)|, \text{pw}(G_2) + |V(G_1)|) - 1.$$

**Proof** The lower bounds follow by Corollary 22, as  $G_1$  and  $G_2$  are subgraphs of  $G_1[v/G_2]$ . For the upper bounds, we can

1. replace  $v$  in every bag of a tree-decomposition (path-decomposition) of width  $\text{tw}(G_1)$  ( $\text{pw}(G_1)$ ) for  $G_1$  by  $V(G_2)$  or
2. add to every bag of a tree-decomposition (path-decomposition) of width  $\text{tw}(G_2)$  ( $\text{pw}(G_2)$ ) for  $G_2$  the set  $V(G_1) \setminus \{v\}$ .

As both alternatives result in a valid tree-decomposition (path-decomposition) for  $G_1[v/G_2]$ , the upper bound follows.  $\square$

Note that the upper bounds described in the previous theorem are strict, as for two cliques  $K_n, K_m$  and a vertex  $v \in V(K_n)$ , we have  $K_n[v/K_m] = K_{n+m-1}$  with

$$\begin{aligned} \text{tw}(K_{n+m-1}) &= n + m - 2 = \min(\text{tw}(K_n) + m, \text{tw}(K_m) + n) - 1 \quad \text{and} \\ \text{pw}(K_{n+m-1}) &= n + m - 2 = \min(\text{pw}(K_n) + m, \text{pw}(K_m) + n) - 1. \end{aligned}$$

Besides this general upper bound for tree-width, the ensuing proposition can provide an even stricter upper bound in specific situations.

**Proposition 49** *Let  $G_1, G_2$  be two vertex-disjoint graphs and  $v \in V(G_1)$  a non-isolated vertex. Then, it holds that*

$$\text{tw}(G_1[v/G_2]) \leq \max(\text{tw}(G_1) - 1, \text{tw}(G_2)) + |N_{G_1}(v)|.$$

**Proof** Let  $(\mathcal{X}_1, T_1)$  be a tree-decomposition for  $G_1$  of width  $\text{tw}(G_1)$  and  $(\mathcal{X}_2, T_2)$  a tree-decomposition for  $G_2$  of width  $\text{tw}(G_2)$ . Replace  $v$  by  $N_{G_1}(v)$  in every bag of  $\mathcal{X}_1$  and denote the modified set of bags by  $\mathcal{X}'_1$ . Then,  $(\mathcal{X}'_1, T_1)$  is a tree-decomposition for  $G_1 \setminus \{v\}$  of width at most  $\text{tw}(G_1) - 1 + |N_{G_1}(v)|$ . Subsequently, add  $N_{G_1}(v)$  to all bags of  $\mathcal{X}_2$  and denote the modified set of bags by  $\mathcal{X}'_2$ . Let  $v_1 \in V(T_1)$  be a vertex of  $T_1$  with  $N_{G_1}(v) \subseteq X_{v_1}$  for  $X_{v_1} \in \mathcal{X}'_1$  and  $v_2$  any vertex of  $T_2$ . Then,  $(\mathcal{X}, T)$  with  $\mathcal{X} = \mathcal{X}'_1 \cup \mathcal{X}'_2$ ,  $V(T) = V(T_1) \cup V(T_2)$ , and  $E(T) = E(T_1) \cup E(T_2) \cup \{v_1, v_2\}$  is a tree-decomposition for  $G_1[v/G_2]$  of width at most  $\max(\text{tw}(G_1) - 1, \text{tw}(G_2)) + |N_{G_1}(v)|$ .  $\square$

## 4.5 Graph Product

The graph product of two vertex-disjoint graphs  $G_1$  and  $G_2$  is a new graph with vertex set  $V(G_1) \times V(G_2)$  and an edge set derived from the adjacency, equality, or non-adjacency of vertices in the original graphs  $G_1$  and  $G_2$ . In this work, we consider the cartesian [Sab59], categorical [Wei62], co-normal [HW67], lexicographic [Har58], and normal [Sab59] graph product, as well as the symmetric difference [HW67] and the rejection [HW67].<sup>8</sup> All graph products, their respective notations, and the edge sets of the resulting graphs are listed in Table 1. For more exhaustive definitions and in-depths results on these graph products, we refer to the works by Imrich and Klavzar [IK00] and Jensen and Toft [JT94].

Graph product	Notation	Edge set $\{(u_1, u_2), (v_1, v_2)\}  $
Cartesian	$G_1 \times G_2$	$(u_1 = v_1 \wedge \{u_2, v_2\} \in E(G_2)) \vee (u_2 = v_2 \wedge \{u_1, v_1\} \in E(G_1))$
Categorical	$G_1 \star_{\text{cat}} G_2$	$\{u_1, v_1\} \in E(G_1) \wedge \{u_2, v_2\} \in E(G_2)$
Co-Normal	$G_1 \vee G_2$	$\{u_1, v_1\} \in E(G_1) \vee \{u_2, v_2\} \in E(G_2)$
Lexicographic	$G_1[G_2]$	$(\{u_1, v_1\} \in E(G_1)) \vee (u_1 = v_1 \wedge \{u_2, v_2\} \in E(G_2))$
Normal (Strong)	$G_1 \star G_2$	$(u_1 = v_1 \wedge \{u_2, v_2\} \in E(G_2)) \vee (\{u_1, v_1\} \in E(G_1) \wedge u_2 = v_2) \vee (\{u_1, v_1\} \in E(G_1) \wedge \{u_2, v_2\} \in E(G_2))$
Symmetric difference	$G_1 \Delta G_2$	$\{u_1, v_1\} \in E(G_1) \vee \{u_2, v_2\} \in E(G_2)$
Rejection	$G_1   G_2$	$\{u_1, v_1\} \notin E(G_1) \wedge \{u_2, v_2\} \notin E(G_2)$

Table 1: Graph products

Besides these graph products, Teh and Yap defined the  $\gamma$ -product of two graphs  $G_1$  and  $G_2$  as  $\overline{G_1} \vee \overline{G_2}$  [TY64]. Transforming the edge set of the  $\gamma$ -product of two graphs  $G_1$  and  $G_2$ , it follows that  $\overline{G_1} \vee \overline{G_2} = G_1 \star G_2$  holds. In other words, the  $\gamma$ -product is merely a different formulation for the normal product of two graphs. Furthermore, transforming the edge set of  $G_1[G_2]$ , we obtain  $G_1[G_2] = \overline{G_1} \star \overline{G_2}$  while transforming the edge set of  $G_1 \Delta G_2$  results in  $G_1 \Delta G_2 = (G_1 \times G_2) \cup (G_1 \star_{\text{cat}} \overline{G_2}) \cup (\overline{G_1} \star_{\text{cat}} G_2)$ .

For two paths  $P_n, P_m$  with  $n, m \in \mathbb{N}$  it holds that  $P_n \times P_m$  is an  $n \times m$ -grid graph. Consequently, for two graphs  $G_1$  and  $G_2$ , it is generally not possible to bound the tree-width (path-width) of  $G_1 \times G_2$  from above by the tree-widths (path-widths) of  $G_1$  and  $G_2$ , c.f. Example 4. With  $P_n \times P_m$  being a subgraph of  $P_n \vee P_m, P_n \star P_m$ , and  $P_n \Delta P_m$ , the same result follows by Corollary 22 for the co-normal and normal graph product as well as the symmetric difference. Next,  $P_n \star_{\text{cat}} P_m$  has an  $a \times a$ -grid graph with  $a$

<sup>8</sup>Weichsel introduced the categorical graph product as the “Kronecker product”, while Harary and Wilcox referred to it as “conjunction”. The normal graph product was introduced by Sabidussi as the “strong product”. The co-normal graph product was introduced by Harary and Wilcox as “disjunction” and the lexicographic graph product was initially defined by Harary as “composition”.

proportional to  $\min(n, m)$  as subgraph, so that by the same argument as before, the tree-width (path-width) of  $G_1 \star_{\text{cat}} G_2$  can generally not be bound from above by the tree-widths (path-widths) of  $G_1$  and  $G_2$ . For  $n, m \in \mathbb{N}$ , let  $I_n$  and  $I_m$  denote the graphs that contain  $n$  and  $m$  isolated vertices. By our earlier observation, we know that  $I_n | I_m = K_n \star K_m$  holds. Consequently, the tree-width (path-width) of  $I_n | I_m$  cannot be bound from above by the tree-widths (path-widths) of  $I_n$  and  $I_m$ , as  $\text{tw}(I_n) = \text{pw}(I_n) = 1$  holds, while  $K_n$  and  $K_m$  are subgraphs of  $I_n | I_m$ , such that by Lemma 5 (Lemma 11)  $\text{tw}(I_n | I_m) \geq \max(n, m) - 1$  ( $\text{pw}(I_n | I_m) \geq \max(n, m) - 1$ ) follows. Therefore, for two graphs  $G_1$  and  $G_2$ , the tree-width (path-width) of the rejection  $G_1 | G_2$  can in general not be bound from above by the tree-widths (path-widths) of  $G_1$  and  $G_2$ . The subsequent corollary summarizes these observations.

**Corollary 50** *Let  $G_1$  and  $G_2$  be two graphs. It is not possible to provide an upper bound for the tree-width (path-width) of the two graph's cartesian, categorical, co-normal, and normal graph product, their rejection or their symmetric difference in terms of the tree-widths (path-widths) of  $G_1$  and  $G_2$ .*

Lower bounds for the tree-width (using the notation of bramble number [Ree97]) of the cartesian and the normal product of two graphs are given in terms of Hadwiger, PI, and bramble number in the work by by Kozawa, Otachi, and Yamazaki [KOY14].

Having discussed all previously defined graph products but the lexicographic graph product, subsequently we provide bounds for this operation. First, for two graphs  $G_1$  and  $G_2$ , it holds that  $G_1$  and  $G_2$  are subgraphs of  $G_1[G_2]$ . Consequently, by Corollary 22 it follows that the tree-width (path-width) of  $G_1[G_2]$  is at least as big as the maximum over the tree-widths (path-widths) of  $G_1$  and  $G_2$ . Second, to obtain a tree-decomposition (path-decomposition) for  $G_1[G_2]$ , we begin with a tree-decomposition (path-decomposition) for  $G_1$  and replace every vertex  $v_i \in V(G_1)$  in every bag by  $(v_i, v_j)$  for all  $v_j \in V(G_2)$ . This results in a tree-decomposition (path-decomposition) for  $G_1[G_2]$  of width  $(\text{tw}(G_1) + 1)|V(G_2)| - 1$  ( $(\text{pw}(G_1) + 1)|V(G_2)| - 1$ ).

**Corollary 51** *Let  $G_1$  and  $G_2$  be two vertex-disjoint graphs. It holds that*

$$\begin{aligned} \max(\text{tw}(G_1), \text{tw}(G_2)) &\leq \text{tw}(G_1[G_2]) \leq (\text{tw}(G_1) + 1)|V(G_2)| - 1 \quad \text{and,} \\ \max(\text{pw}(G_1), \text{pw}(G_2)) &\leq \text{pw}(G_1[G_2]) \leq (\text{pw}(G_1) + 1)|V(G_2)| - 1. \end{aligned}$$

Bodlaender et al. [BGHK95] have shown that if  $G_2$  is a clique, the upper bounds for  $\text{tw}(G_1[G_2])$  and  $\text{pw}(G_1[G_2])$  as stated above are tight.

**Theorem 52** *Let  $G$  be a graph and  $q \in \mathbb{N}$ . It holds that  $\text{tw}(G[K_q]) = (\text{tw}(G) + 1)q - 1$  and  $\text{pw}(G[K_q]) = (\text{pw}(G) + 1)q - 1$ .*

## 4.6 1-Sum

Let  $G_1$  and  $G_2$  be two vertex-disjoint graphs and  $v \in V(G_1)$  and  $w \in V(G_2)$  two vertices. The 1-sum of  $G_1$  and  $G_2$ , denoted by  $G_1 \oplus_{v,w} G_2$ , consists of the disjoint union of  $G_1$  and  $G_2$  with vertices  $v$  and  $w$  identified. More specifically, graph  $G_1 \oplus_{v,w} G_2$  has vertex set  $(V(G_1) \cup V(G_2) \cup \{z\}) \setminus \{v, w\}$  for a newly introduced vertex  $z$  and edge set

$$\begin{aligned} E(G_1) \cup E(G_2) \setminus \{ \{v, v_1\} \mid v_1 \in N_{G_1}(v) \} \\ \setminus \{ \{w, w_1\} \mid w_1 \in N_{G_2}(w) \} \\ \cup \{ \{z, z_1\} \mid z_1 \in (N_{G_1}(v) \cup N_{G_2}(w)) \}. \end{aligned}$$

Having the 1-sum of two graphs  $G_1$  and  $G_2$  formally defined, in the subsequent theorem we consider the tree-width and path-width of  $G_1 \oplus_{v,w} G_2$ .



**Theorem 53** *Let  $G_1$  and  $G_2$  be two vertex-disjoint graphs and  $v \in V(G_1)$  as well as  $w \in V(G_2)$  two vertices. Then, it holds that*

$$\begin{aligned} tw(G_1 \oplus_{v,w} G_2) &= \max(tw(G_1), tw(G_2)) \quad \text{and} \\ \max(pw(G_1), pw(G_2)) &\leq pw(G_1 \oplus_{v,w} G_2) \leq \max(pw(G_1), pw(G_2)) + 1. \end{aligned}$$

**Proof** The lower bounds follow by Corollary 22, as  $G_1$  and  $G_2$  are subgraphs of  $G_1 \oplus_{v,w} G_2$ .

Let  $(\mathcal{X}_1, T_1)$  be a tree-decomposition for  $G_1$  of width  $tw(G_1)$  and  $(\mathcal{X}_2, T_2)$  be a tree-decomposition for  $G_2$  of width  $tw(G_2)$ . To define a tree-decomposition  $(\mathcal{X}, T)$  for  $G_1 \oplus_{v,w} G_2$ , we replace every occurrence of  $v$  in  $\mathcal{X}_1$  and every occurrence of  $w$  in  $\mathcal{X}_2$  by  $z$ . Then, we choose a vertex  $u_1$  in  $V(T_1)$ , such that  $z$  belongs to  $X_{u_1} \in \mathcal{X}_1$ , and a vertex  $u_2$  in  $V(T_2)$ , such that  $z$  belongs to  $X_{u_2} \in \mathcal{X}_2$ . We define  $T$  as the disjoint union of  $T_1$  and  $T_2$  with the additional edge  $\{u_1, u_2\}$  and  $\mathcal{X}$  as the union of  $\mathcal{X}_1$  and  $\mathcal{X}_2$ . This results in a tree-decomposition  $(\mathcal{X}, T)$  for  $G_1 \oplus_{v,w} G_2$  of width  $\max(tw(G_1), tw(G_2))$ .

In order to define a path-decomposition for  $G_1 \oplus_{v,w} G_2$ , let  $\mathcal{X}_1$  be a path-decomposition for  $G_1$  of width  $pw(G_1)$  and  $\mathcal{X}_2$  be a path-decomposition for  $G_2$  of width  $pw(G_2)$ . Then, we can either proceed as for tree-width, replacing  $v$  in all bags of  $\mathcal{X}_1$  and  $w$  in all bags of  $\mathcal{X}_2$  by  $z$ , and concatenate both paths of bags to a new path  $\mathcal{X}$ , resulting in a path-decomposition for  $G_1 \oplus_{v,w} G_2$  of width  $\max(pw(G_1), pw(G_2))$ , or, if the resulting concatenation violates (pw-3), additionally add  $z$  to all remaining bags of  $\mathcal{X}$ , resulting in a path-decomposition of width at most  $\max(pw(G_1), pw(G_2)) + 1$ .  $\square$

If  $v \in V(G_1)$  and  $w \in V(G_2)$  have degree at least one in  $G_1$  and  $G_2$ , i.e., are no isolated vertices, the new vertex  $z$  in  $G_1 \oplus_{v,w} G_2$  is called an *articulation vertex* of  $G_1 \oplus_{v,w} G_2$ , since  $(G_1 \oplus_{v,w} G_2) - z$  has more connected components than  $G_1 \oplus_{v,w} G_2$ . For a graph  $G$ , a maximal, biconnected subgraph without any articulation vertex is called a *block* or *biconnected component* of  $G$ . The bounds of Theorem 53 for tree-width imply that the tree-width of a graph equals the maximum tree-width of its biconnected components.

**Corollary 54** *Let  $G$  be a graph. It holds that the tree-width of  $G$  equals the maximum tree-width of its biconnected components.*

Contrarily, the subsequent example shows that Corollary 54 does not hold for path-width.

*Example 5* Denote the vertices of  $P_3$  by  $V(P_3) = \{v_1, v_2, v_3\}$  and the vertices of  $P_5$  by  $V(P_5) = \{u_1, u_2, u_3, u_4, u_5\}$ . Then, the incidence graph  $I(K_{1,3})$  from Figure 2 can be created as  $P_3 \oplus_{v_3, u_3} P_5$ . We know that  $pw(P_3) = pw(P_5) = 1$  holds and showed in Example 2 that  $pw(I(K_{1,3})) = 2$  is true. As all biconnected components of  $I(K_{1,3})$  are subgraphs of  $P_3$  or  $P_5$ , it follows that all biconnected components have a path-width of one. Consequently, the path-width of  $I(K_{1,3})$  cannot equal the maximum path-width of any of its biconnected components.

However, the bounds of Theorem 53 for path-width imply that the path-width of a graph can be bounded by its number of biconnected components and their maximum path-widths.

## 4.7 Corona

Frucht and Harary [FH70] introduced the corona of two graphs when they constructed a graph whose automorphism group is the wreath product of the two graphs' automorphism

groups.<sup>9</sup> The *corona* of two vertex-disjoint graphs  $G_1$  and  $G_2$ , denoted by  $G_1 \wedge G_2$ , consists of the disjoint union of one copy of  $G_1$  and  $|V(G_1)|$  copies of  $G_2$ , where each vertex of the copy of  $G_1$  is connected to all vertices of one copy of  $G_2$ . In other words,  $|V(G_1)||V(G_2)|$  edges are inserted in the disjoint union of the  $|V(G_1)| + 1$  graphs.

Alternatively, the corona of  $G_1$  and  $G_2$  can also be obtained by applying 1-sum and dominating vertex operations as follows. Let  $V(G_1) = \{v_1, \dots, v_n\}$  be the vertex set of  $G_1$ . For  $i = 1, \dots, n$ , we take a copy of  $G_2$ , insert a dominating vertex  $w_i$  (cf. Section 3.1) in that copy, and obtain the resulting graph  $G_{2,i}$ . Then, the subsequent sequence of 1-sums,

$$(\dots((G_1 \oplus_{v_1, w_1} G_{2,1}) \oplus_{v_2, w_2} G_{2,2}) \dots) \oplus_{v_n, w_n} G_{2,n}, \quad (1)$$

results in the corona  $G_1 \wedge G_2$  of  $G_1$  and  $G_2$ .

With this observation at hand, we can bound the tree-width (path-width) of  $G_1 \wedge G_2$  in the tree-width (path-width) of its combined graphs as follows.

**Theorem 55** *Let  $G_1$  and  $G_2$  be two vertex-disjoint graphs. Then, it holds that*

$$\begin{aligned} \max(\text{tw}(G_1), \text{tw}(G_2)) &\leq \text{tw}(G_1 \wedge G_2) \leq \max(\text{tw}(G_1), \text{tw}(G_2)) + 1 \quad \text{and} \\ \max(\text{pw}(G_1), \text{pw}(G_2)) &\leq \text{pw}(G_1 \wedge G_2) \leq \max(\text{pw}(G_1), \text{pw}(G_2)) + |V(G_1)|. \end{aligned}$$

**Proof** The lower bounds follow by Corollary 22 since  $G_1$  and  $G_2$  are subgraphs of  $G_1 \wedge G_2$ .

For the upper bounds we make use of our earlier observation that we can obtain the corona of  $G_1$  and  $G_2$  by applying 1-sum and dominating vertex operations as described in Equation (1). By Theorem 18 it follows that  $\text{tw}(G_{2,i}) \leq \text{tw}(G_2) + 1$  and  $\text{pw}(G_{2,i}) \leq \text{pw}(G_2) + 1$ . By Theorem 53 and Equation (1) it follows that  $\text{tw}(G_1 \wedge G_2) \leq \max(\text{tw}(G_1), \text{tw}(G_2)) + 1$  and  $\text{pw}(G_1 \wedge G_2) \leq \max(\text{pw}(G_1), \text{pw}(G_2)) + |V(G_1)|$ .  $\square$

With the previous theorem proved, we asked ourselves whether there exists a constant integer  $c \in \mathbb{N}$ , such that for all graphs  $G_1, G_2$  it holds that  $\text{pw}(G_1 \wedge G_2) \leq \max(\text{pw}(G_1), \text{pw}(G_2)) + c$ , similar to the upper bound for tree-width. The subsequent proposition provides a negative answer to this question.

**Proposition 56** *For  $n \geq 2$  and  $m \geq 1$  it holds that*

$$\text{pw}(K_n \wedge K_m) = n + \max\left(0, m - \left\lfloor \frac{n}{2} \right\rfloor\right) - 1.$$

**Proof** We write  $K_n = (\{v_1, \dots, v_n\}, \{\{v_i, v_j\} \mid 1 \leq i < j \leq n\})$  as well as  $K_m = (\{u_1, \dots, u_m\}, \{\{u_i, u_j\} \mid 1 \leq i < j \leq m\})$ . For  $1 \leq i \leq n$  we denote the  $n$  copies of  $K_m$  in  $K_n \wedge K_m$  as  $K_m^i$  with  $V(K_m^i) = \{u_1^i, \dots, u_m^i\}$ . By construction of  $K_n \wedge K_m$ , every vertex  $v_i$  of  $K_n$  gets connected to all vertices of  $K_m^i$ , resulting in  $n$  cliques of size  $m + 1$  which we denote by  $K_{m+1}^i$  with  $V(K_{m+1}^i) = V(K_m^i) \cup \{v_i\}$  for  $1 \leq i \leq n$ .

Since  $K_{m+1}^i$ ,  $1 \leq i \leq n$ , and  $K_n$  are subgraphs of  $K_n \wedge K_m$ , it follows by Corollary 22 that  $\max(n - 1, m) \leq \text{pw}(K_n \wedge K_m)$  must hold.

Next, let us construct the ensuing path-decomposition  $\mathcal{X} = (X_1, \dots, X_{n+1})$  for  $K_n \wedge K_m$ . We define

1.  $X_i = V(K_m^i) \cup \{v_1, \dots, v_i\}$  for  $1 \leq i \leq \lfloor \frac{n}{2} \rfloor$ ,
2.  $X_{\lfloor \frac{n}{2} \rfloor + 1} = V(K_n)$ , and

---

<sup>9</sup>Please be unconcerned, the corona of graphs has nothing to do with the global pandemic of coronavirus disease 2019 (COVID-19).

3.  $X_{i+1} = V(K_m^i) \cup \{v_i, \dots, v_n\}$  for  $\lfloor \frac{n}{2} \rfloor < i \leq n$ .

It is easy to check that  $\mathcal{X}$  satisfies all three requirements of a path-decomposition for  $K_n \wedge K_m$ . Furthermore, we note that (a)  $\max_{1 \leq i \leq \lfloor \frac{n}{2} \rfloor} |X_i| = m + \lfloor \frac{n}{2} \rfloor$ , (b)  $|X_{\lfloor \frac{n}{2} \rfloor + 1}| = n$ , and (c)  $\max_{\lfloor \frac{n}{2} \rfloor < i \leq n} |X_{i+1}| = m + n - \lfloor \frac{n}{2} \rfloor$  holds.

Next, let us differentiate two cases,  $m \leq \lfloor \frac{n}{2} \rfloor$  and  $m > \lfloor \frac{n}{2} \rfloor$ :

**Case 1:**  $m \leq \lfloor \frac{n}{2} \rfloor$

In this case we have for (a)  $m + \lfloor \frac{n}{2} \rfloor \leq \lfloor \frac{n}{2} \rfloor + \lfloor \frac{n}{2} \rfloor \leq n$  and for (c)  $m + (n - \lfloor \frac{n}{2} \rfloor) \leq \lfloor \frac{n}{2} \rfloor + n - \lfloor \frac{n}{2} \rfloor = n$ , such that  $\max_{1 \leq i \leq n+1} |X_i| = n$  holds. Thus, we know that the path-decomposition  $\mathcal{X}$  has a width of  $n - 1$ , yielding

$$\max(n - 1, m) = n - 1 \leq \text{pw}(K_n \wedge K_m) \leq n - 1.$$

Consequently, in this case we obtain

$$\text{pw}(K_n \wedge K_m) = n - 1 = n + \max\left(0, m - \left\lfloor \frac{n}{2} \right\rfloor\right) - 1.$$

**Case 2:**  $m > \lfloor \frac{n}{2} \rfloor$

In this case we have for (a)  $m + n - \lfloor \frac{n}{2} \rfloor = m + \lceil \frac{n}{2} \rceil \geq m + \lfloor \frac{n}{2} \rfloor$  and for (c)  $m + n - \lfloor \frac{n}{2} \rfloor = m + \lceil \frac{n}{2} \rceil \geq \lfloor \frac{n}{2} \rfloor + 1 + \lceil \frac{n}{2} \rceil = n + 1 \geq n$ , such that  $\max_{1 \leq i \leq n+1} |X_i| = m + n - \lfloor \frac{n}{2} \rfloor$  holds. Thus, the width of path-decomposition  $\mathcal{X}$  is  $m + n - \lfloor \frac{n}{2} \rfloor - 1 = n + \max(0, m - \lfloor \frac{n}{2} \rfloor) - 1$  and it follows that

$$\text{pw}(K_n \wedge K_m) \leq n + \max(0, m - \left\lfloor \frac{n}{2} \right\rfloor) - 1$$

holds. Lastly, we show that there cannot exist any path-decomposition  $\mathcal{X}'$  for  $K_n \wedge K_m$  of width smaller than  $n + \max(0, m - \lfloor \frac{n}{2} \rfloor) - 1$ . To do so, let us assume that  $\mathcal{X}'$  is a minimum path-decomposition for  $K_n \wedge K_m$  of smallest possible width.

As  $K_n$  is a subgraph of  $K_n \wedge K_m$ , we know by Lemma 11 that there must exist at least one bag  $X \in \mathcal{X}'$  with  $V(K_n) \subseteq X$ . Furthermore, as  $K_{m+1}^i$  is a subgraph of  $K_n \wedge K_m$  for  $1 \leq i \leq n$ , we know by the same argument that there must exist at least one bag  $Y_i \in \mathcal{X}'$  with  $V(K_{m+1}^i) \subseteq Y_i$  for every  $1 \leq i \leq n$ .

Assuming that  $X$  and  $Y_i$ ,  $1 \leq i \leq n$ , contain no more than the previously mentioned sets of vertices,  $V(K_n \wedge K_m) = X \cup \bigcup_{i=1}^n Y_i$  is true and for every edge  $\{u, v\} \in E(K_n \wedge K_m)$  it holds that  $u, v \in X$  if  $u, v \in V(K_n)$  or  $u, v \in Y_i$  if  $u, v \in V(K_{m+1}^i)$ . Consequently  $\mathcal{X}'$  satisfies (pw-1) and (pw-2) already.

Thus, it is left to argue that (i) for every  $1 \leq i \leq n$  the only bag containing  $V(K_m^i)$  is  $Y_i$  and (ii) the least bag-size increasing way to satisfy (pw-3) for  $X, Y_1, \dots, Y_n$  is by introducing additional copies of subsets of  $V(K_n)$  to  $Y_1, \dots, Y_n$ .

For every  $u_j^i \in V(K_m^i)$ ,  $1 \leq j \leq m$ ,  $1 \leq i \leq n$ , it holds that  $N_{K_n \wedge K_m}(u_j^i) \subseteq Y_i$ . Consequently, if  $u_j^i$  is present in any other bag of  $\mathcal{X}'$  except  $Y_i$ , this contradicts our assumption of  $\mathcal{X}'$  being minimal, as we could remove  $u_j^i$  from all bags in  $\mathcal{X}'$  except  $Y_i$  and still have a valid path-decomposition. Hence, (i) holds.

Since  $\mathcal{X}'$  must form a path, there can exist at most one bag preceding  $X$  and one bag succeeding  $X$  in the path. The most efficient way to ensure that (pw-3) is satisfied is by allocating the first  $\lfloor \frac{n}{2} \rfloor$  vertices from  $X$  to its predecessor and the remaining  $\lceil \frac{n}{2} \rceil$  vertices to its successor. Thus, (ii) holds and  $\mathcal{X}' = \mathcal{X}$  follows.

Consequently,  $\mathcal{X}'$  has a width of at least  $n + \max(0, m - \lfloor \frac{n}{2} \rfloor) - 1$ , as well, such that

$$\text{pw}(K_n \wedge K_m) = n + \max(0, m - \lfloor \frac{n}{2} \rfloor) - 1$$

follows.

This shows the statement of the proposition.  $\square$

Note that for  $n = 1$  one can verify that  $\text{pw}(K_1 \wedge K_m) = \text{pw}(K_{m+1}) = m$  holds.

## 5 Conclusions and Outlook

In Section 3, we have shown how the tree-width or path-width of a given graph changes, if we apply a certain, unary graph transformation  $f$  to this graph. In all cases, in which it is possible to bound the tree-width or path-width of the resulting graph  $f(G)$ , we have also shown how to compute the corresponding decomposition in time linear in the size of the corresponding decomposition for  $G$ . Table 2 summarizes the results. It is noteworthy that the behavior of tree-width and path-width under the considered transformations is almost identical.

transformation $f$	$\text{tw}(f(G))$	$\text{pw}(f(G))$
vertex deletion	$k$	$k$
vertex addition	$k + 1$	$k + 1$
edge deletion	$k$	$k$
edge addition	$k + 1$	$k + 1$
subgraph	$k$	$k$
vertex identification	$k + 1$	$k + 1$
edge contraction	$k$	$k$
edge subdivision	$k$	$k + 1$
minor	$k$	$k$
switching	$k + 1$	$k + 1$

Table 2: Let  $G$  be a graph of tree-width (path-width)  $k$  and  $f$  the unary graph transformation in the first column. The second column lists the upper bound for the tree-width and the third column the upper bound for the path-width of  $f(G)$ .

Furthermore, in Section 4, we have considered various binary graph operations  $f$ , which create a new graph  $f(G_1, G_2)$  out of two graphs  $G_1$  and  $G_2$ . In all cases, in which it is possible to bound the tree-width or path-width of the combined graph  $f(G_1, G_2)$  in the tree-width or path-width of  $G_1$  and  $G_2$ , we have shown how to compute the corresponding decomposition in time linear in the size of the corresponding decompositions for  $G_1$  and  $G_2$ , such that our results are constructive. In Table 3, we summarize these results, which show that, with an exception for the corona of two graphs, the behavior of tree-width and path-width under the considered operations is nearly identical.

The results in Sections 3.2 and 3.3 allow to generalize known results on the stability of trees and forests [WR21a, WR21b] to the stability of graph classes of bounded tree-width [GRW24].

Most of our results provide tight upper and lower bounds for the tree-width and path-width of the resulting graph in terms of the tree-width and path-width of the initial graphs or argue why such bounds are impossible.

operation $f$	$\text{tw}(f(G_1, G_2))$	$\text{pw}(f(G_1, G_2))$
disjoint union	$\max(k_1, k_2)$	$\max(k_1, k_2)$
join	$\min(k_1 + n_2, k_2 + n_1)$	$\min(k_1 + n_2, k_2 + n_1)$
substitution	$\min(k_1 + n_2, k_2 + n_1) - 1$	$\min(k_1 + n_2, k_2 + n_1) - 1$
lexicographic product	$(k_1 + 1)n_2 - 1$	$(k_1 + 1)n_2 - 1$
1-sum	$\max(k_1, k_2)$	$\max(k_1, k_2) + 1$
corona	$\max(k_1, k_2) + 1$	$\max(k_1, k_2) + n_1$

Table 3: Let  $G_1$  and  $G_2$  be two graphs of tree-width (path-width)  $k_1$  and  $k_2$ , respectively, and  $f$  the binary graph operation from the first column. By  $n_1$  we denote the number of vertices of  $G_1$  and by  $n_2$  the number of vertices of  $G_2$ . The second column lists the upper bound of the tree-width and the third column lists the upper bound of the path-width for graph  $f(G_1, G_2)$ .

For the following case it remains to show that our bounds are the best possible ones or to provide stricter bounds. The bounds for the tree-width and path-width of the power of a graph, as shown in Section 3.8, are very rough as all vertex degrees are approximated by the maximum degree of the graph. Furthermore, neither did we provide lower bounds for the tree-width (path-width) of the categorical or co-normal graph product of two graphs, nor for the symmetric difference or rejection of two graphs.

## References

- [ACP87] S. Arnborg, D.G. Corneil, and A. Proskurowski. Complexity of finding embeddings in a  $k$ -tree. *SIAM Journal of Algebraic and Discrete Methods*, 8(2):277–284, 1987.
- [AP89] S. Arnborg and A. Proskurowski. Linear time algorithms for NP-hard problems restricted to partial  $k$ -trees. *Discrete Applied Mathematics*, 23:11–24, 1989.
- [APC90] S. Arnborg, A. Proskurowski, and D.G. Corneil. Forbidden minors characterization of partial 3-trees. *Discrete Mathematics*, 80:1–19, 1990.
- [Arn85] S. Arnborg. Efficient algorithms for combinatorial problems on graphs with bounded decomposability – A survey. *BIT*, 25:2–23, 1985.
- [Ats08] A. Atserias. On digraph coloring problems and treewidth duality. *European Journal of Combinatorics*, 29(4):796–820, 2008.
- [BB73] U. Bertelè and F. Brioschi. On Non-serial Dynamic Programming. *Journal of Combinatorial Theory*, 14(2):137–148, 1973.
- [BGHK95] H.L. Bodlaender, J.R. Gilbert, H. Hafsteinsson, and T. Kloks. Approximating treewidth, pathwidth, frontsize, and shortest elimination tree. *Journal of Algorithms*, 18(2):238–255, 1995.
- [BH12] H.L. Bodlaender and J. Hage. On switching classes, NLC-width, cliquewidth and treewidth. *Theoretical Computer Science*, 429:30–35, 2012.
- [BKK93] H.L. Bodlaender, T. Kloks, and D. Kratsch. Treewidth and pathwidth of permutation graphs. In *Proceedings of International Colloquium on Automata*,

- Languages and Programming (ICALP)*, volume 700 of *LNCS*, pages 114–125, Heidelberg and Berlin, Germany, 1993. Springer-Verlag.
- [BLS99] A. Brandstädt, V.B. Le, and J.P. Spinrad. *Graph Classes: A Survey*. SIAM Monographs on Discrete Mathematics and Applications. SIAM, Philadelphia, 1999.
- [BM76] J. Bondy and U. Murty. *Graph Theory with Applications*. North-Holland, New York, Amsterdam and Oxford, 1976.
- [BM90] H.L. Bodlaender and R.H. Möhring. The pathwidth and treewidth of co-graphs. In *Proceedings of Scandinavian Workshop on Algorithm Theory (SWAT)*, volume 447 of *LNCS*, pages 301–309, Heidelberg and Berlin, Germany, 1990. Springer-Verlag.
- [BM93] H.L. Bodlaender and R.H. Möhring. The pathwidth and treewidth of co-graphs. *SIAM J. Disc. Math.*, 6(2):181–188, 1993.
- [Bod86] H.L. Bodlaender. Classes of graphs with bounded treewidth. Technical Report RUU-CS-86-22, Universiteit Utrecht, 1986.
- [Bod88] H. Bodlaender. Classes of graphs with bounded tree-width. *Bulletin of the European Association for Theoretical Computer Science*, 36:116–125, 1988.
- [Bod93] H.L. Bodlaender. Complexity of path-forming games. *Theoretical Computer Science*, 110:215–245, 1993.
- [Bod96] H.L. Bodlaender. A linear-time algorithm for finding tree-decompositions of small treewidth. *SIAM Journal on Computing*, 25(6):1305–1317, 1996.
- [Bod98] H.L. Bodlaender. A partial  $k$ -arboretum of graphs with bounded treewidth. *Theoretical Computer Science*, 209:1–45, 1998.
- [Bou94] A. Bouchet. Circle graph obstructions. *Journal of Combinatorial Theory, Series B*, 60:107–144, 1994.
- [BvAdF01] H.L. Bodlaender and B. van Antwerpen-de Fluiter. Parallel algorithms for series parallel graphs and graphs with tree-width two. *Algorithmica*, 29:543–559, 2001.
- [CC80] C.J. Colbourn and D.G. Corneil. On deciding switching equivalence of graphs. *Discrete Applied Mathematics*, 2:181–184, 1980.
- [CE12] B. Courcelle and J. Engelfriet. *Graph Structure and Monadic Second-Order Logic. A Language-Theoretic Approach*. Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge, 2012.
- [CFR03] G. Calinescu, C.G. Fernandes, and B. Reed. Multicuts in unweighted graphs and digraphs with bounded degree and bounded tree-width. *Journal of Algorithms*, 48:333–359, 2003.
- [Chl02] J. Chlebíková. Partial  $k$ -trees with maximum chromatic number. *Discrete Mathematics*, 259(1–3):269–276, 2002.
- [CO86] J. Chvatalova and J. Opatrny. The bandwidth problem and operations on graphs. *Discrete Mathematics*, 61(2-3):141–150, 1986.

- [Cou06] B. Courcelle. The Monadic Second-Order Logic of Graphs XV: A conjecture by D. Seese. *Journal of Applied Logic*, 4:79–114, 2006.
- [Cou14] B. Courcelle. Clique-width and edge contraction. *Information Processing Letters*, 114:42–44, 2014.
- [dMR05] F. de Montgolfier and M. Rao. The bi-join decomposition. *Electronic Notes in Discrete Mathematics*, 22:173–177, 2005.
- [FH70] R. Frucht and F. Haray. On the coronas of two graphs. *Aequationes Math.*, 4:322–324, 1970.
- [GRW24] F. Gurski, J. Rothe, and R. Weishaupt. Stability, vertex stability, and unfrozenness for special graph classes. *Theory of Computing Systems*, 68:75–102, 2024.
- [Gur17] F. Gurski. The behavior of clique-width under graph operations and graph transformations. *Theory of Computing Systems*, 60(2):346–376, 2017.
- [Gus93] J. Gusted. On the pathwidth of chordal graphs. *Discrete Applied Mathematics*, 45(3):233–248, 1993.
- [Hag00] T. Hagerup. Dynamic algorithms for graphs of bounded treewidth. *Algorithmica*, 27(3):292–315, 2000.
- [Har58] F. Harary. On the number of bicolored graphs. *Pacific Journal of Mathematics*, 8(4):743–755, 1958.
- [Har69] F. Harary. *Graph Theory*. Addison-Wesley Publishing Company, Massachusetts, 1969.
- [Hay96] R.B. Hayward. Recognizing  $P_3$ -structure: A switching approach. *Journal of Combinatorial Theory, Series B*, 66(2):247–262, 1996.
- [HOSG08] P. Hliněný, S. Oum, D. Seese, and G. Gottlob. Width parameters beyond tree-width and their applications. *Computer Journal*, 51(3):326–362, 2008.
- [HW67] F. Harary and G.W. Wilcox. Boolean operations on graphs. *Mathematica Scandinavica*, 20:41–51, 1967.
- [HW15] D.J. Harvey and D.R. Wood. Treewidth of the line graph of a complete graph. *Journal of Graph Theory*, 79(1):48–54, 2015.
- [HW18] D.J. Harvey and D.R. Wood. The treewidth of line graphs. *Journal of Combinatorial Theory, Series B*, 132:157–179, 2018.
- [HZ60] F. Harary and R. Z.Norman. Some properties of line digraphs. *Rendiconti del Circolo Matematico di Palermo*, pages 161–168, 1960.
- [IK00] W. Imrich and S. Klavzar. *Product Graphs: Structure and Recognition*. Series in Discrete Mathematics and Optimization. Wiley-Interscience, New York, USA, 2000.
- [JT94] T.R. Jensen and B. Toft. *Graph Coloring Problems*. John Wiley & Sons, New York, 1994.

- [JW12] G. Joret and D.R. Wood. Nordhaus-Gaddum for treewidth. *European Journal of Combinatorics*, 33:488–490, 2012.
- [KBMK93] T. Kloks, H. Bodlaender, H. Müller, and D. Kratsch. Computing treewidth and minimum fill-in: All you need are the minimal separators. In *Proceedings of the Annual European Symposium on Algorithms (ESA)*, volume 726 of *LNCS*, pages 260–271, Heidelberg and Berlin, Germany, 1993. Springer-Verlag.
- [KF79] T. Kashiwabara and T. Fujisawa. NP-completeness of the problem of finding a minimum-clique-number interval graph containing a given graph as a subgraph. In *Proceedings of the International Symposium on Circuits and Systems*, pages 657–660, 1979.
- [Kin92] N.G. Kinnersley. The vertex separation number of a graph equals its path width. *Information Processing Letters*, 42:345–350, 1992.
- [KKR12] K. Kawarabayashi, Y. Kobayashi, and B. Reed. The disjoint paths problem in quadratic time. *Journal of Combinatorial Theory, Series B*, pages 424–435, 2012.
- [KL94] N.G. Kinnersley and M.A. Langston. Obstruction set isolation for the gate matrix layout problem. *Discrete Applied Mathematics*, 54:169–213, 1994.
- [KMRR09] J. Kneis, D. Mölle, S. Richter, and P. Rossmanith. A bound on the pathwidth of sparse graphs with applications to exact algorithms. *SIAM Journal on Discrete Mathematics*, 23(1):407–427, 2009.
- [KOY14] K. Kozawa, Y. Otachi, and K. Yamazaki. Lower bounds for treewidth of product graphs. *Discrete Applied Mathematics*, 162:251–258, 2014.
- [KS93] E. Korach and N. Solel. Tree-width, path-width, and cutwidth. *Discrete Applied Mathematics*, 43:97–101, 1993.
- [KT04] J. Kleinberg and E. Tardos. *Algorithm Design*. Addison-Wesley Publishing Company, Massachusetts, 2004.
- [KZN00] M.A. Kashem, X. Zhou, and T. Nishizeki. Algorithms for generalized vertex-rankings of partial  $k$ -trees. *Theoretical Computer Science*, 240(2):407–427, 2000.
- [Lim10] V. Limouzy. Seidel minor, permutation graphs and combinatorial properties. In *Proceedings of the International Symposium on Algorithms and Computation*, volume 6506 of *LNCS*, pages 194–205, Heidelberg and Berlin, Germany, 2010. Springer-Verlag.
- [MS88] B. Monien and I.H. Sudborough. Min cut is NP-complete for edge weighted trees. *Theoretical Computer Science*, 58:209–229, 1988.
- [Ree97] B.A. Reed. *Tree Width and Tangles: A New Connectivity Measure and Some Applications*, pages 87–162. London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge, 1997.
- [Ros74] D.J. Rose. On simple characterizations of  $k$ -trees. *Discrete Mathematics*, 7:317–322, 1974.



- [RS83] N. Robertson and P.D. Seymour. Graph minors I. Excluding a forest. *Journal of Combinatorial Theory, Series B*, 35:39–61, 1983.
- [RS85] N. Robertson and P.D. Seymour. *Graph Minors – A Survey*, pages 153–171. Cambridge Univ. Press, Cambridge, 1985.
- [RS86a] N. Robertson and P.D. Seymour. Graph minors II. Algorithmic aspects of tree width. *Journal of Algorithms*, 7:309–322, 1986.
- [RS86b] N. Robertson and P.D. Seymour. Graph minors V. Excluding a planar graph. *Journal of Combinatorial Theory, Series B*, 41:92–114, 1986.
- [RS95] N. Robertson and P.D. Seymour. Graph minors. XIII. The Disjoint Paths Problem. *Journal of Combinatorial Theory, Series B*, 63:65–110, 1995.
- [RS04] N. Robertson and P.D. Seymour. Graph minors XX. Wagners conjecture. *Journal of Combinatorial Theory, Series B*, 92:325–357, 2004.
- [Sab59] G. Sabidussi. Graph multiplication. *Mathematische Zeitschrift*, 72:446–457, 1959.
- [Sch89] P. Scheffler. *Die Baumweite von Graphen als ein Mass für die Kompliziertheit algorithmischer Probleme*. PhD thesis, Akademie der Wissenschaften der DDR, 1989.
- [Sei74] J.J. Seidel. Graphs and two-graphs. In *Proceedings of the 5th Southeastern Conf. on Combinatorics, Graph Theory, and Computing*, Winnipeg, Canada, 1974. Utilitas Mathematica Publishing.
- [Sei76] J.J. Seidel. A survey of two-graphs. In *Proceedings of Colloquio Internazionale sulle Teorie Combinatorie*, volume 17, pages 481–511, Rome, 1976. Accademia Nazionale dei Lincei.
- [ST81] J.J. Seidel and D.E. Taylor. Two-graphs, a second survey. In *Algebraic Methods in Graph Theory*, volume II, pages 689–711, 1981.
- [ST93] P.D. Seymour and R. Thomas. Graph Searching and a Min-Max Theorem for Tree-Width. *Journal of Combinatorial Theory, Series B*, 58:22–33, 1993.
- [ST07] K. Suchan and I. Todinca. Pathwidth of circular-arc graphs. In *Proceedings of Graph-Theoretical Concepts in Computer Science (WG)*, volume 4769 of *LNCS*, pages 258–269, Heidelberg and Berlin, Germany, 2007. Springer-Verlag.
- [TY64] H. H. Teh and H. D. Yap. Some construction problems of homogeneous graphs. *Bulletin of the Mathematical Society of Nanyang University*, pages 164–196, 1964.
- [Wag70] K. Wagner. *Graphentheorie*, volume 248/248a. B. J. Hochschultaschenbücher, Mannheim, 1970.
- [Wei62] P. Weichsel. The kronecker product of graphs. *Proceedings of the American Mathematical Society*, 3:47–52, 1962.
- [Whi32] H. Whitney. Congruent graphs and the connectivity of graphs. *American Journal of Mathematics*, 54:150–168, 1932.

- [WR21a] R. Weishaupt and J. Rothe. Stability of special graph classes. In Claudio Sacerdoti Coen and Ivano Salvo, editors, *Proceedings of the 22nd Italian Conference on Theoretical Computer Science*, volume 3072 of *CEUR Workshop Proceedings*, pages 234–248, RWTH Aachen, Germany, 2021. CEUR-WS.org.
- [WR21b] R. Weishaupt and J. Rothe. Stability of special graph classes. Technical Report arXiv:2106.01496 [cs.CC], arXiv.org, June 2021.