

COMPUTING CONGRUENCES OF FINITE INVERSE SEMIGROUPS

LUNA ELLIOTT, ALEX LEVINE, AND JAMES D. MITCHELL

ABSTRACT. In this paper we present an algorithm for computing a congruence on an inverse semigroup from a collection of generating pairs. This algorithm uses a myriad of techniques from computational group theory, automata, and the theory of inverse semigroups. An initial implementation of this algorithm outperforms existing implementations by several orders of magnitude.

1. INTRODUCTION

In this paper we are concerned with the question of computing two-sided congruences of a finite inverse semigroup. The class of inverse semigroups lies somewhere between the classes of groups, and semigroups, with more useful structure than semigroups in general, and possibly less structure than groups. A *semigroup* is a set, usually denoted S , with an associative binary operation, usually indicated by juxtaposing elements of S . An *inverse semigroup* is a semigroup S such that for every element $s \in S$ there exists a unique $s' \in S$ with $ss's = s$ and $s'ss' = s'$. The element s' is usually denoted s^{-1} , a choice which is at least partially justified by the fact that if S is a group and $s \in S$, then s^{-1} is just the usual group theoretic inverse of s . On the other hand, if S is not a group and $s \in S$, then neither ss^{-1} nor $s^{-1}s$ is necessarily equal to the identity of S , not least because S need not have an identity.

Two-sided congruences are to semigroups what normal subgroups are to groups. However, a *two-sided congruence* on a semigroup S is not a subsemigroup of S but rather an equivalence relation $\rho \subseteq S \times S$ with the property that if $(x, y) \in \rho$ and $s \in \rho$, then $(xs, ys), (sx, sy) \in \rho$. Although there is a definition of one-sided congruences also, akin to the notion of subgroups of groups, we will be solely concerned with two-sided congruences, and so we will drop the “two-sided” and henceforth refer to “congruences” to exclusively mean “two-sided congruences”. Equivalently, ρ is a congruence on S if $(xs, yt) \in \rho$ whenever $(x, y), (s, t) \in \rho$, making ρ a subsemigroup of $S \times S$ rather than S . Although congruences on semigroups and normal subgroups on group are analogous notions, the definition for semigroups is a special case of the notion of congruences in universal algebra; see, for example, [3, Section 5]. For a congruence ρ , it will be convenient for us to write $x =_{\rho} y$ instead of $(x, y) \in \rho$.

Like the normal subgroups of a group, the congruences of a semigroup form a lattice with the meet $\rho \wedge \sigma$ of congruences ρ and σ being $\rho \cap \sigma$, and join $\rho \vee \sigma$ being the least congruence containing ρ and σ .

If G is a group and $A \subseteq G$, then algorithms for determining the least normal subgroup $\langle\langle A \rangle\rangle$ of G containing A are one of the core components of computational group theory. Following the nomenclature of GAP [7] we refer to such algorithms as *normal closure* algorithms. For example, normal closure algorithms for permutation groups are considered in [21, Section 5.4.1]; for groups in general, in [9, Section 3.3.2]; or for computing all normal subgroups in [11].

Congruences of inverse semigroups have also been studied extensively in the literature. For example, the lattices of congruences of various semigroups, including many inverse semigroups, have been completely described, for example in [13, 14, 15, 24]; and from the perspective of computation in [1, 2, 4, 23]. With the single exception of [23], the existing algorithms [1, 2, 4], and their implementations in [16, 17, 18], for computing individual congruences on an inverse semigroup do not use any of the specific structure of inverse semigroups. We will say more about the exception below.

The following notions have been, and will be here, indispensable for the study of congruences on inverse semigroups. Let S be an inverse semigroup. We denote the set of idempotents of S by $E(S)$. The *kernel* of a congruence ρ on an inverse semigroup S is the inverse subsemigroup

$$\text{Ker}(\rho) = \{s \in S \mid \text{there exists } e \in E(S), s =_{\rho} e\} \leq S$$

2020 *Mathematics Subject Classification.* 20M18, 20-08, 20M30.

Key words and phrases. inverse semigroup, congruence, computational algebra.

and the *trace* of ρ is the restriction of ρ to the idempotents of S :

$$\text{Tr}(\rho) = \rho \cap (E(S) \times E(S)).$$

If S is an inverse semigroup and ρ is a congruence on $E(S)$, then ρ is said to be *normal in S* if $s^{-1}xs =_{\rho} s^{-1}ys$ whenever $s \in S$ and $x =_{\rho} y$. Similarly, if T is a subsemigroup of S , then T is *normal in S* if $s^{-1}ts \in T$ for all $s \in S$ and all $t \in T$. If T is a normal subsemigroup of S and τ is a normal congruence on $E(S)$, then (T, τ) is (rather unimaginatively) called a *congruence pair* if the following conditions hold:

(CP1) $ae \in T$ and $e =_{\tau} a^{-1}a$ implies that $a \in T$;

(CP2) $a \in T$ implies that $aa^{-1} =_{\tau} a^{-1}a$;

for all $a \in S$ and all $e \in E(S)$. It is well-known that the congruences of an inverse semigroup S are in one-to-one correspondence with the congruence pairs on S ; see, for example, [10, Theorem 5.3.3]. The kernel-trace description originates in [20], and is described in almost books about semigroup theory; in addition to [10, Theorem 5.3.3], see [8, Proposition 1.3], [12, Section 5.1], [19, Chapter III].

In this paper, we present various mathematical results that can be combined into an algorithm for computing a congruence on a finite inverse semigroup. The aim of these results is to allow the efficient computation of the least congruence R^{\sharp} on an inverse semigroup S from a collection of generating pairs $R \subseteq S \times S$. By “compute” a congruence ρ we mean that we have a representation of the congruence that is amenable to computation (i.e. that is not larger than necessary, and can be computed relatively quickly), and that can be used to answer questions about ρ such as whether or not $(x, y) \in S \times S$ belongs to ρ ; what is the number of classes in ρ ; and what are the elements of x/ρ ?

A preliminary implementation of this algorithm in [16] and [7], indicates that there is, for some examples, a 1000 times speedup when compared to the existing implementation of [23] in [17] (which uses the kernel and trace); a 1500 times speed up in comparison to the implementation in [17] and [16] (for semigroups in general).

Although significantly faster than existing implementations, it is worth mentioning that neither the time nor space complexity of the algorithms we present is polynomial in the size of the input. For instance, one key step in the algorithm we present is computing the trace of a congruence on an inverse semigroup S . If S is the symmetric inverse monoid on the set $\{1, \dots, n\}$, then S can be represented using $O(n)$ space. However, $|E(S)| = 2^n$, and computing the idempotents in this case has complexity $O(2^n)$. The complexity of the other steps in the algorithm are somewhat harder to describe; but they also depend on $|E(S)|$. It seems unlikely to the authors that there is a sub-exponential algorithm for computing a congruence on an inverse semigroup.

The paper is organized as follows. In [Section 2](#) we provide some details of the prerequisite notions from semigroup theory that we require. In [Section 3](#) we describe data structures for inverse semigroups, and their quotients, that uses the theory of Green’s relations, the action of an inverse semigroup on its idempotents by conjugation, and an analogue of Schreier’s Lemma. The data structure consists of a generating set X for the inverse semigroup S , a certain automata-like graph Γ_X encoding the action (of the previous sentence) and its strongly connected components, and a finite sequence G_1, \dots, G_m of groups. For a quotient of S , the data structure consists of the generating set X for S , a quotient of the graph Γ_X , and a sequence of normal subgroups N_1, \dots, N_n of the groups in the data structure for S . In [Section 4](#) we describe how to compute the trace of a congruence using Γ_X and a (guaranteed to terminate) variant of the Todd-Coxeter Algorithm from [4]. In [Section 5](#), we show how to obtain relatively small collections of elements Y_i of each group G_i such that the normal closure $\langle\langle Y_i \rangle\rangle$ is the required normal subgroup N_i . In [Section 6](#) we show how to obtain the elements of the kernel of a congruence as a translate of the preimage of a coset of a normal subgroup under a homomorphism of groups. In the final section [Section 7](#), we describe a completely separate algorithm for computing the maximum idempotent separating congruence on an inverse subsemigroup of a finite symmetric inverse monoid.

2. PRELIMINARIES

Let S be an inverse semigroup. We denote the set of idempotents of S by $E(S)$. If $s, t \in S$, then we write $s \leq t$ if there exists $e \in E(S)$ such that $s = te$. The relation \leq is a partial order on S (see, for example, [12, Proposition 1.4.7]), usually referred to as the *natural partial order* on S . This definition may appear to be inherently “right-handed”, but it is not, since $s \leq t$ if and only if there exists $f \in E(S)$ such that $s = ft$ [12, Lemma 1.4.6]. Similarly, if $s \leq t$ and $u \leq v$, then $su \leq tv$ [12, Proposition 1.4.7].

We define a **word graph** $\Gamma = (N, E)$ over the alphabet A to be a directed graph with nodes N and edges $E \subseteq N \times A \times N$. Word graphs are just finite state automata without initial or terminal states.

If $(\alpha, a, \beta) \in E$ is an edge in a word graph $\Gamma = (N, E)$, then α is the **source**, a is the **label**, and β is the **target** of (α, a, β) . A word graph Γ is **complete** if for every node α and every letter $a \in A$ there is at least one edge with source α labelled by a . A word graph $\Gamma = (N, E)$ is **finite** if the sets of nodes N and edges E are finite. A word graph is **deterministic** if for every node $\alpha \in N$ and every $a \in A$ there is at most one edge with source α and label a . Complete deterministic word graphs are just unary algebras with universe N and operations $f_a : N \rightarrow N$ defined by $(\alpha)f_a = \beta$ whenever (α, a, β) is an edge in Γ ; see [3] for more details. The perspective of unary algebras maybe helpful, for those familiar with this notion, when we define word graph quotients and homomorphisms, these are identical to the notions of quotients and homomorphisms of the associated unary algebras. If $\alpha, \beta \in N$, then an (α, β) -**path** is a sequence of edges $(\alpha_0, a_0, \alpha_1), \dots, (\alpha_{n-1}, a_{n-1}, \alpha_n) \in E$ where $\alpha_0 = \alpha$ and $\alpha_n = \beta$ and $a_0, \dots, a_{n-1} \in A$. If $\alpha, \beta \in V$ and there is an (α, β) -path in Γ , then we say that β is **reachable** from α . If α is a node in a word graph Γ , then the **strongly connected component of α** is the set of all nodes β such that β is reachable from α and α is reachable from β . If $\Gamma_1 = (N_1, E_1)$ and $\Gamma_2 = (N_2, E_2)$ are word graphs over the same alphabet A , then $\phi : N_1 \rightarrow N_2$ is a **homomorphism** if $(\alpha, a, \beta) \in E_1$ implies $((\alpha)\phi, a, (\beta)\phi) \in E_2$. If κ is an equivalence relation on the nodes of a word graph $\Gamma = (N, E)$, then we define the **quotient Γ/κ of Γ by κ** to be the word graph with nodes $\{\alpha/\kappa \mid \alpha \in N\}$ and edges $\{(\alpha/\kappa, a, \beta/\kappa) \mid (\alpha, a, \beta) \in E\}$. Of course, even if Γ is deterministic, the quotient Γ/κ is not necessarily deterministic. If Γ is deterministic, then Γ/κ is deterministic if and only if κ is a congruence on the unary algebra associated to Γ .

The **kernel** of a congruence ρ on an inverse semigroup S is the inverse subsemigroup

$$\text{Ker}(\rho) = \{s \in S \mid \text{there exists } e \in E(S), s =_\rho e\} \leq S$$

and the **trace** of ρ is the restriction of ρ to the idempotents of S :

$$\text{Tr}(\rho) = \rho \cap (E(S) \times E(S)).$$

If ρ is a congruence on an inverse semigroup S , then ρ is said to be **normal** if $s^{-1}xs =_\rho s^{-1}ys$ whenever $s \in S$ and $x =_\rho y$. Similarly, if T is a subsemigroup of S , then T is **normal** if $sts^{-1} \in T$ for all $s \in S$ and all $t \in T$. If T is a normal subsemigroup of S and τ is a normal congruence on $E(S)$, then (T, τ) is (rather unimaginatively) called a **congruence pair** if the following conditions hold:

(CP1) $ae \in T$ and $e =_\tau a^{-1}a$ implies that $a \in T$;

(CP2) $a \in T$ implies that $aa^{-1} =_\tau a^{-1}a$

for all $a \in S$ and all $e \in E(S)$. It is well-known that the congruences of an inverse semigroup S are in one-to-one correspondence with the congruence pairs on S ; see, for example, [10, Theorem 5.3.3].

If S is a semigroup, then we denote by S^1 either: $S \cup \{1_S\}$ with an identity $1_S \notin S$ adjoined; or just S in the case that S already has an identity.

The final ingredient that we require in this paper is that of Green's relations. If $s, t \in S$, then **Green's \mathcal{R} -relation** is the equivalence relation on S defined by $(s, t) \in \mathcal{R}$ if and only if $sS^1 = \{sx \mid x \in S^1\} = tS^1$. Green's \mathcal{L} -relation is defined analogously; Green's \mathcal{H} -relation is just $\mathcal{L} \cap \mathcal{R}$; and Green's \mathcal{D} -relations is defined to be $\mathcal{L} \circ \mathcal{R}$. If S is finite, then $(s, t) \in \mathcal{D}$ if and only if $S^1sS^1 = S^1tS^1$. Green's relations are fundamental to the study of semigroups; we refer the reader to any of [10, 12, 8, 19] for further details. If T is a subsemigroup of S (denoted $T \leq S$), then we may write \mathcal{K}^S and \mathcal{K}^T to distinguish the Green's relations on S and T when $\mathcal{K} \in \{\mathcal{L}, \mathcal{R}, \mathcal{H}, \mathcal{D}\}$. If $s \in S$, then we denote the equivalence class of Green's \mathcal{K} -relation containing s by K_s or K_s^S if we want to indicate the semigroup containing the class.

Theorem 2.1 (Location Theorem). *Let S be a finite semigroup and let $a, b \in S$ be such that $(a, b) \in \mathcal{D}$. Then the following are equivalent*

- (a) $(ab, a), (ab, b) \in \mathcal{D}$;
- (b) $(a, ab) \in \mathcal{R}$ and $(ab, b) \in \mathcal{L}$;
- (c) *there exists an idempotent $(e, a) \in \mathcal{L}$ and $(e, b) \in \mathcal{R}$.*

We will make repeated use of the following straightforward result also.

Lemma 2.2. *If S is a finite inverse semigroup, $e, f \in E(S)$ are such that $e \leq f$, and $(e, f) \in \mathcal{D}^S$, then $e = f$.*

3. A DATA STRUCTURE FOR INVERSE SEMIGROUPS AND THEIR QUOTIENTS

In this section, we describe the data structure for inverse semigroups given in [5, Section 5.6]. If S is an inverse semigroup, then the data structure for S consists of the following:

- (A1) a generating set X for S ;
- (A2) the word graph Γ_X with nodes $E(S)$ and edges $\{(e, x, x^{-1}ex) \mid e \in E(S), x \in X\}$;
- (A3) the strongly connected components of Γ_X ;
- (A4) a generating set for one group \mathcal{H} -class per strongly connected component of Γ_X .

Given X , the word graph Γ_X can be found in $O(|E(S)||X|)$ time and space (assuming that products in S can be found in constant time). The strongly connected components of Γ_X can be found from Γ_X using algorithms from graph theory (such as those of Gabow [6] or Tarjan [22]). Given the strongly connected components of Γ_X , the groups from (A4) can be determined using the analogue of Schreier's Lemma given in [5, Proposition 2.3(c) and Algorithm 3]. This data structure can be used to answer many of the fundamental questions about S that arise in a computational setting, such as membership testing in S , determining the Green's structure, and the size of S ; see [5] for more details.

If S is an inverse semigroup S , $R \subseteq S \times S$, and $R^\sharp = \rho$, we will show how to compute a data structure for the quotient S/ρ from the data structure for S . This data structure consists of:

- (Q1) the generating set X for S ;
- (Q2) the quotient word graph $\Gamma_X/\text{Tr}(\rho)$ with nodes $E(S)/\text{Tr}(\rho)$ and edges $\{(e/\text{Tr}(\rho), x, (x^{-1}ex)/\text{Tr}(\rho)) \mid e \in E(S), x \in X\}$;
- (Q3) the strongly connected components of $\Gamma_X/\text{Tr}(\rho)$;
- (Q4) a generating sets for one group \mathcal{H} -class per strongly connected component of $\Gamma_X/\text{Tr}(\rho)$.

Clearly for (Q2) we must compute $\text{Tr}(\rho)$; and given (Q2) we can compute the strongly connected components as we did for S itself. Without a representation of ρ (beyond R) we have no means of representing X/ρ , and hence we cannot determine the generating sets for the group \mathcal{H} -classes required in (Q4). We show how to compute $\text{Tr}(\rho)$ from R in Section 4; and show how to compute the required group \mathcal{H} -classes in Section 5.

The quotient data structure is sufficient for representing the inverse semigroup S/ρ , and can be used to compute various aspects of ρ , such as the number of classes. But it does not suffice for other purposes, such as checking membership in ρ , or computing the elements of s/ρ . For the latter, we require a means of computing the kernel $\text{Ker}(\rho)$ of ρ from the data structure. We do this in Section 6.

Throughout this paper we will use the notation from this section for S , the congruence ρ , and the associated data structures.

4. COMPUTING THE TRACE

In this section we show how to compute the trace of a congruence on the inverse semigroup S from the set of generating pairs $R \subseteq S \times S$.

Lemma 4.1 (Generating pairs for the trace). *If S is an inverse semigroup and $\rho = R^\sharp$ is a congruence of S , then the trace $\text{Tr}(\rho)$ of ρ is the least normal congruence of $E(S)$ in S containing*

$$\{(aea^{-1}, beb^{-1}) \mid e \in E(S), (a, b) \in R\}.$$

Proof. Let N denote the set of pairs in the statement, and let ν be the least normal congruence on $E(S)$ containing N . We must show that $\nu = \text{Tr}(\rho)$.

If $(a, b) \in R$ and $e \in E(S)$ are arbitrary, then certainly $a =_\rho b$ and so $ae =_\rho be$ and $a^{-1} =_\rho b^{-1}$. Hence $aea^{-1} =_\rho beb^{-1}$ and so $aea^{-1} =_{\text{Tr}(\rho)} beb^{-1}$. Therefore $\nu \subseteq \text{Tr}(\rho)$.

For the converse containment, suppose that $e =_{\text{Tr}(\rho)} f$. Then $e =_\rho f$, and hence $e =_{R^\sharp} f$. So there exist $s_0 = e, s_1, \dots, s_n = f$ where $s_i = p_i u_i q_i$ and $s_{i+1} = p_i v_i q_i$ for some $p_i, q_i \in S^1$ and $(u_i, v_i) \in R$ for all i . We set $e_i = s_i s_i^{-1}$ for every i . Then $e_0 = e$ and $e_n = f$. For every i , $e_i = s_i s_i^{-1} = p_i u_i q_i q_i^{-1} u_i^{-1} p_i^{-1}$ and $e_{i+1} = s_{i+1} s_{i+1}^{-1} = p_i v_i q_i q_i^{-1} v_i^{-1} p_i^{-1}$. Since $q_i q_i^{-1} \in E(S)$ and $(u_i, v_i) \in R$, it follows that $(u_i q_i q_i^{-1} u_i^{-1}, v_i q_i q_i^{-1} v_i^{-1}) \in N \subseteq \nu$ by definition. Hence, since ν is normal, $e_i = p_i u_i q_i q_i^{-1} u_i^{-1} p_i^{-1} =_\nu p_i v_i q_i q_i^{-1} v_i^{-1} p_i^{-1} = e_{i+1}$ for all i . Thus $e = e_0 =_\nu e_n = f$, as required. \square

For the remainder of this section we require S to be a monoid, by adjoining an identity 1_S if necessary. If σ is any equivalence relation on $E(S)$, then we define Γ_X/σ to be the word graph with nodes $E(S)/\sigma$ and




FIGURE 1. Diagram of the word graph Γ_X from Example 4.3, where x_1 is represented in magenta, x_2 in blue and x_3 in orange. Each node is the idempotent that is the identity on the set in its label. Shaded nodes correspond to the idempotents belonging to the only non-singleton class of $\text{Tr}(\rho)$.

edges $(e/\sigma, x, (x^{-1}ex)/\sigma)$ for all $e \in E(S)$ and all $x \in X$. It is routine to verify that σ is a normal congruence on $E(S)$ with respect to S if and only if Γ_X/σ is deterministic. In this case, $\text{Tr}(\sigma)$ is completely determined by Γ_X/σ . If $e = x_1 \dots x_n \in E(S)$ where $x_i \in X$ labels a path from $1_S/\sigma$ to f/σ in Γ_X/σ , then $e =_{\text{Tr}(\sigma)} f$. Conversely, if $e =_{\text{Tr}(\sigma)} f$ and $e = x_1 \dots x_n$ and $f = y_1 \dots y_m$ where $x_i, y_j \in X$, then $x_1 \dots x_n$ and $y_1 \dots y_m$ both label $(1_S, e/\sigma)$ -paths in Γ_X/σ .

We have established the following result.

Lemma 4.2 (Normal congruences as quotients of word graphs). *There is a one-to-one correspondence between the normal congruences of $E(S)$ and the deterministic quotients of Γ_X .*

The trace $\text{Tr}(\rho)$ of a congruence $\rho = R^\sharp$ on an inverse semigroup S can therefore be computed by:

- (T1) computing the set from Lemma 4.1;
- (T2) find the greatest quotient of Γ_X containing R using the variant of the Todd-Coxeter Algorithm described in Section 5 of [4].

Next we consider an example to illustrate the steps (T1) and (T2). Each element of a finite symmetric inverse monoid is expressible as a product of chains and disjoint cycles. So we write (i_1, \dots, i_n) for a cycle and $[i_1, \dots, i_n]$ for a chain. When points are fixed we write (i) to denote that i is fixed as omitted points are not in the domain of the described partial permutation.

Example 4.3. In this example we show how to compute the trace of the least congruence ρ on the symmetric inverse monoid I_4 (consisting of the partial permutations on the set $\{1, 2, 3, 4\}$) containing the pair:

$$(a, b) := ((1)(2)(3), (1\ 2\ 3)).$$

We use the following generating set for I_4 :

$$X := \{x_1 := (1\ 2\ 3\ 4), \quad x_2 := (1\ 2)(3)(4), \quad x_3 := [4\ 3\ 2\ 1]\}.$$

If N is the set of generating pairs for $\text{Tr}(\rho)$ from Lemma 4.1, then a maximal subset M of N such that $M \cap M^{-1} = \emptyset$ is:

$$M := \{((1), (2)), ((1), (3)), ((2), (3)), ((1)(2), (2)(3)), ((1)(2), (1)(3)), ((2)(3), (1)(3))\}.$$

Obviously, M also generates $\text{Tr}(\rho)$. A diagram of the word graph Γ_X in this example can be seen in Figure 1. A diagram of the greatest quotient of Γ_X containing (a, b) is shown in Figure 2.

5. COMPUTING THE GROUP \mathcal{H} -CLASSES OF THE QUOTIENT

In this section we show how to compute the group \mathcal{H} -class component (Q4) of the quotient data structure.

We will repeatedly make use of the following simple lemma, which we record for the sake of completeness.

Lemma 5.1. *If S is finite, $e, f \in E(S)$, $x, y \in S$, and $fxey \mathcal{D} f$, then $fxey = fxy$.*




FIGURE 2. Diagram of the maximum quotient of the word graph Γ_X from Example 4.3 by the generating pairs of $\text{Tr}(\rho)$ from Lemma 4.1, where x_1 is represented in magenta, x_2 in blue and x_3 in orange.

Proof. We may assume without loss of generality that S is an inverse subsemigroup of the symmetric inverse monoid I_X for some finite set X . Since $f\mathcal{D}fxy$, it follows that $\text{rank}(f) = \text{rank}(fxy) \leq \text{rank}(fxy) \leq \text{rank}(f)$, yielding equality throughout. In particular, $\text{rank}(fxy) = \text{rank}(fxy)$, and since e is an idempotent and S is finite, it follows that $fxy = fxy$, as required. \square

If s/ρ is an idempotent in S/ρ , then by Lallement's Lemma there exists $e \in E(S)$ such that $e/\rho = s/\rho$, and so $E(S) \cap s/\rho = e/\text{Tr}(\rho)$. We define $f \in E(S)$ to be the meet of $e/\text{Tr}(\rho)$, that is,

$$f = \bigwedge e/\text{Tr}(\rho),$$

and we denote the group \mathcal{H} -class of f in S by G .

The following lemma describes the group \mathcal{H} -classes in the quotient S/ρ in terms of the group \mathcal{H} -classes in S and a normal subgroup.

Lemma 5.2. *If $N = H_f^S \cap (f/\rho)$, then the following hold:*

- (a) N is a normal subgroup of $G = H_f^S$;
- (b) f/ρ is an inverse subsemigroup of S and N is the minimum non-empty ideal of f/ρ ;
- (c) the group \mathcal{H} -class $H_{e/\rho}^{S/\rho}$ is isomorphic to G/N .

Proof. (a) Let $g, h \in N$. Since f is an idempotent $h^{-1} \in f/\rho$ and so $gh^{-1} \in f^2/\rho = f/\rho$. Thus N is a subgroup of $G = H_f^S$. If $n \in N$ and $g \in G$, then $g^{-1}ng = {}_\rho g^{-1}fg = f$ and so $g^{-1}ng \in f/\rho$ and N is normal.

(b) We first show that f/ρ is an inverse subsemigroup of S . Let $s, t \in f/\rho$. Then $st = {}_\rho f^2 = f$ and $s^{-1} = {}_\rho f^{-1} = f$ and so $st, s^{-1} \in f/\rho$. Thus f/ρ is an inverse subsemigroup of S . Next we show that N is the minimum non-empty ideal of f/ρ . Clearly, since $f \in N$, N is non-empty. Let $n \in N$ and $s \in f/\rho$. Then $sn \in N$ if and only if $(sn, f) \in \mathcal{H}^S$ if and only if $sn(sn)^{-1} = (sn)^{-1}sn = f$.

It follows that $sn(sn)^{-1}, (sn)^{-1}sn \in f/\rho$, so $sn(sn)^{-1} \geq f$ and $(sn)^{-1}sn \geq f$. On the other hand, $nf = f$ so $(sn)^{-1}sn = (sn)^{-1}snf \leq f$. Thus $(sn)^{-1}sn = f$. As $sn(sn)^{-1} \geq (sn)^{-1}sn$ and $(sn(sn)^{-1}, (sn)^{-1}sn) \in \mathcal{D}$, it follows from Lemma 2.2 that $(sn)^{-1}sn = f$ also.

We have shown that N is a left ideal, by symmetry it is a right ideal also. Every non-empty ideal of f/ρ contains fs for some $s \in f/\rho$. Thus $f = fs(fs)^{-1} \in I$ since this is the unique idempotent in N . Therefore the ideal I contains all of N and so N is the minimum ideal.

(c) We define

$$\psi: G/N \longrightarrow H_{e/\rho}^{S/\rho} \quad \text{by} \quad Ng \mapsto g/\rho.$$

To show that ψ is well-defined, we must show that ψ maps into $H_{e/\rho}^{S/\rho}$ and that ψ does not depend on the choice of coset representative. Let $Ng \in G/N$. Then $(Ng)\psi((Ng)\psi)^{-1} = (g/\rho) \cdot (g^{-1}/\rho) = f/\rho = e/\rho$ and by symmetry $((Ng)\psi)^{-1}(Ng)\psi = e/\rho$ and so $(Ng)\psi \in H_{e/\rho}^{S/\rho}$. If $h \in Ng$, then $h = ng$ for some $n \in N$ and so $h = ng = {}_\rho fg = g$. So $(Nh)\psi = h/\rho = g/\rho = (Ng)\psi$ and ψ is well-defined. We will next show that ψ is a homomorphism. Let $Ng, Nh \in G/N$. Then

$$(Ng \cdot Nh)\psi = (Ngh)\psi = gh/\rho = (g/\rho) \cdot (h/\rho) = (Ng)\psi \cdot (Nh)\psi,$$

and ψ is a homomorphism. To show ψ is injective, let $Ng, Nh \in G/N$ be such that $(Ng)\psi = (Nh)\psi$. Then $g/\rho = h/\rho$ and so $gh^{-1} \in f/\rho$. Thus $gh^{-1} \in N$, and it follows that $Ng = Nh$, as required. It remains to show that ψ is surjective. Let $h/\rho \in H_{e/\rho}^{S/\rho}$. Then $h \in G$ and so $h/\rho = (Nh)\psi$ and ψ is surjective. \square

The next lemma is the key result in this section, permitting us to express N in terms of R and the word graph Γ_X of S , and allowing for the efficient computation of N .

Lemma 5.3 (Generating the normal subgroups). *If the strongly connected component of f in Γ_X is $\{e_1 = f, e_2, \dots, e_r\}$ for some r , and for every i , we choose $s_i \in S$ to be the label of an (e_1, e_i) -path in Γ_X , then $N = H_f^S \cap (f/\rho)$ is the normal closure of*

$$\{fs_iab^{-1}s_i^{-1} \mid (a, b) \in R, i \in \{1, \dots, r\}\} \cap H_f^S$$

in H_f^S .

Proof. Let N' denote the normal closure of the set in the statement.

To show that $N' \subseteq N$, suppose that $i \in \{1, \dots, k\}$ and $(a, b) \in R$ are such that $fs_iab^{-1}s_i^{-1} \in H_f^S$. We must show that $fs_iab^{-1}s_i^{-1} \in N$; that is, $fs_iab^{-1}s_i^{-1} =_{\rho} f$ (this is sufficient because, by Lemma 5.2(a), N is a normal subgroup of G). We begin by showing that $f = fs_i aa^{-1}s_i^{-1}$. Also $s_i aa^{-1}s_i^{-1} \in E(S)$ and so $fs_i aa^{-1}s_i^{-1} \leq f$. On the other hand, since $fs_iab^{-1}s_i^{-1} \in H_f^S$, which is a group, it follows that

$$\begin{aligned} f &= (fs_iab^{-1}s_i^{-1})(fs_iab^{-1}s_i^{-1})^{-1} && f \text{ is the identity of } H_f^S \\ &= fs_iab^{-1}s_i^{-1}s_i b a^{-1}s_i^{-1}f \\ &\leq fs_i aa^{-1}s_i^{-1}f && b^{-1}s_i^{-1}s_i b \in E(S) \\ &= fs_i aa^{-1}s_i^{-1} && \text{idempotents commute in } S. \end{aligned}$$

It follows that $f = fs_i aa^{-1}s_i^{-1}$, and so, in particular, $f =_{\rho} fs_i aa^{-1}s_i^{-1}$. Since $(a, b) \in R$ we have $a =_{\rho} b$, it follows that $a^{-1} =_{\rho} b^{-1}$, and so $fs_iab^{-1}s_i^{-1} =_{\rho} fs_i aa^{-1}s_i^{-1}$. Therefore by the transitivity of ρ , $fs_iab^{-1}s_i^{-1} =_{\rho} f$, as required.

For the converse containment ($N \subseteq N'$), suppose that $g \in N = H_f^S \cap (f/\rho)$. Since $f =_{\rho} g$, there exists an elementary sequence

$$a_0 = f, \dots, a_i = p_i b_i q_i, a_{i+1} = p_i c_i q_i, \dots, a_n = g$$

where $p_i, q_i \in S$ and (b_i, c_i) or $(c_i, b_i) \in R$ for all i . By assumption, $a_i =_{\rho} a_0 = f$, or equivalently, $a_i \in f/\rho$ for every i . Thus, since f/ρ is a subsemigroup of S and N is the minimum non-empty ideal of f/ρ (Lemma 5.2(b)), $fa_i f \in N$ for all i .

We will show that $fa_k f \in N'$ for every k by induction. Certainly, $a_0 = f \in N'$ since f is the identity of G and $N' \leq G$. Assume that $fa_k f \in N'$ for all $k \leq i$. To prove that $fa_{i+1} f \in N'$, it suffices to show that

$$(fa_i f)(fa_{i+1} f)^{-1} \in N'.$$

But $(fa_i f)(fa_{i+1} f)^{-1} \in N$, and is thus \mathcal{H} -related to f , hence

$$\begin{aligned} (fa_i f)(fa_{i+1} f)^{-1} &= (fp_i b_i q_i f)(fq_i^{-1} c_i^{-1} p_i^{-1} f) \\ &= fp_i b_i c_i^{-1} p_i^{-1} f && \text{by Lemma 5.1.} \end{aligned}$$

If $t = fp_i b_i c_i^{-1} p_i^{-1} f$, then $t = (fa_i f)(fa_{i+1} f)^{-1} \in N \leq G$ since $fa_i f, fa_{i+1} f \in N$.

By assumption $fa_i f \in N' \leq G$, and f is the identity of the group G . Hence $(fa_i f)^{-1}(fa_i f) = f$ and so

$$f = (fa_i f)^{-1}(fa_i f) = (b_i q_i f)^{-1}(p_i^{-1} f p_i)(b_i q_i f).$$

This shows that f and $p_i^{-1} f p_i$ belong to the same strongly connected component of Γ_X , and so there exists $j \in \{1, \dots, k\}$ such that $s_j^{-1} f s_j = p_i^{-1} f p_i$. Clearly,

$$s_j s_j^{-1} f = f s_j s_j^{-1} f = f s_j s_j^{-1} s_j s_j^{-1} f = s_j s_j^{-1} f s_j s_j^{-1} f = f^2 = f,$$

and $fp_i p_i^{-1} = f^1$.

¹We may assume without loss of generality that S is an inverse subsemigroup of I_n . Since $fp_i p_i^{-1} = f|_{\text{dom}(p_i)}$. But $\text{rank}(f) = \text{rank}(s_j s_j^{-1} f s_j s_j^{-1}) = \text{rank}(s_j^{-1} f s_j) = \text{rank}(p_i^{-1} f p_i)$, and so $\text{dom}(p_i) \cap \text{dom}(f) = \text{im}(p_i^{-1}) \cap \text{dom}(f) = \text{dom}(f)$ (otherwise $\text{rank}(p_i^{-1} f p_i) < \text{rank}(f)$). In other words $\text{dom}(f) \subseteq \text{dom}(p_i)$ and so $fp_i p_i^{-1} = f|_{\text{dom}(p_i)} = f$.

If $u = fp_i s_j^{-1} f$, then

$$s_j p_i^{-1} \cdot f p_i s_j^{-1} f = s_j s_j^{-1} f s_j s_j^{-1} f = s_j s_j^{-1} f = f \text{ and } f p_i s_j^{-1} f \cdot s_j p_i^{-1} = f p_i p_i^{-1} f p_i p_i^{-1} = f p_i p_i^{-1} = f.$$

In particular, $(u, f) \in \mathcal{H}$, and so $u \in G$. Since $t \in G$ also, $u^{-1} t u \in N'$ if and only if $t \in N'$ since N' is a normal subgroup of G . But

$$\begin{aligned} u^{-1} t u &= (f p_i s_j^{-1} f)^{-1} \cdot (f p_i b_i c_i^{-1} p_i^{-1} f) \cdot (f p_i s_j^{-1} f) \\ &= f s_j \cdot p_i^{-1} f p_i \cdot b_i c_i^{-1} \cdot p_i^{-1} f p_i \cdot s_j^{-1} f \\ &= f s_j \cdot s_j^{-1} f s_j \cdot b_i c_i^{-1} \cdot s_j^{-1} f s_j \cdot s_j^{-1} f & s_j^{-1} f s_j = p_i^{-1} f p_i \\ &= f s_j b_i c_i^{-1} s_j^{-1} f \in N'. \end{aligned}$$

Hence $t \in N'$, and so $(f a_i f)(f a_{i+1} f)^{-1} \in N'$, and so $f a_{i+1} f \in N'$, as required. We have shown that $f a_i f \in N'$ for all i , and so, in particular, $f a_n f = f g f = g \in N'$. \square

The algorithm for computing the normal subgroups component (Q4) of the quotient data structure is to

- (N1) find one $e \in E(S)$ for every strongly connected component of $\Gamma_X / \text{Tr}(\rho)$;
- (N2) for each representative $e \in E(S)$ from (N1), set N to be the trivial group, and iterate through the generating set given in Lemma 5.3 for $H_f^S \cap (f/\rho)$ where f is the meet of $e/\text{Tr}(\rho)$, taking the normal closure of N and each generator.

Next, we continue the example started in Example 4.3, and compute the generating sets for the normal subgroups in the quotient using Lemma 5.3.

Example 5.4. We compute the normal subgroup component (Q4) of the quotient data structure for the least congruence ρ on the symmetric inverse monoid I_4 containing the pair:

$$(a, b) := ((1)(2)(3), (1\ 2\ 3)).$$

Firstly the graph $\Gamma_X / \text{Tr}(\rho)$ given in Figure 2 clearly has 3 strongly connected components, and so we are attempting to compute 3 normal subgroups. If the representatives chosen in (N1) are $1_{I_4} = (1)(2)(3)(4)$, $(1)(2)(3)$, and \emptyset , then the normal subgroups found in (N2) are the trivial group, $\langle (1\ 3\ 2) \rangle$, and the trivial group, respectively. Thus the quotient groups are (up to isomorphism) the symmetric group on $\{1, 2, 3, 4\}$, the cyclic group of order 2, and the trivial group, respectively.

At this point, the number of classes of the congruence ρ is just the size of the quotient semigroup which equals $4! \cdot 1^2 + 2 \cdot 4^2 + 1 \cdot 1^2 = 57$ where each summand equals $|G/N|$ multiplied by the size of the strongly connected component of e to the power of 2; see [5, Section 5.6] for more details about how to use the quotient data structure to compute with the quotient inverse semigroup S/ρ .

6. COMPUTING A CLASS OF A CONGRUENCE

In this section we consider the question of how to enumerate the elements of x/ρ for an arbitrary $x \in S$. One consequence of this description will be a means of enumerating the elements of the kernel $\text{Ker}(\rho)$ of the congruence ρ .

Throughout this section we fix $x \in S$ with the aim of enumerating x/ρ . We require the following definitions, which are the key ingredients in this section:

$$U_x = \bigcup D_{x/\rho}^{S/\rho} = \{y \in S \mid (y/\rho, x/\rho) \in \mathcal{D}^{S/\rho}\},$$

$\mu, \nu: S \rightarrow E(S)$ defined by

$$(y)\mu = \min((yy^{-1})/\text{Tr}(\rho)) \quad \text{and} \quad (y)\nu = \min((y^{-1}y)/\text{Tr}(\rho)),$$

and, finally, $\phi_x: U_x \rightarrow S$ defined by

$$(y)\phi_x = (y)\mu \cdot y \cdot (y)\nu.$$

The following lemma collects various properties of U_x and ϕ_x that are used repeatedly throughout this section.

- Lemma 6.1.**
- (a) If $y =_\rho z$, then $y =_\rho (y)\phi_x =_\rho (z)\phi_x$;
 - (b) $\text{im}(\phi_x) \subseteq U_x$ and $(y)\phi_x = y$ for all $y \in \text{im}(\phi_x)$;
 - (c) If $y, z \in U_x$, then $(y/\rho, z/\rho) \in \mathcal{D}^{S/\rho}$;

- (d) If $y \in U_x$ and $(y, z) \in \mathcal{D}^S$, then $z \in U_x$;
- (e) If $y \in U_x$, then $(y)\phi_x \leq y$;
- (f) If $y, z \in U_x$ and $y \leq z$, then $y =_\rho z$;
- (g) If $y \in U_x$, then $(y)\phi_x \cdot ((y)\phi_x)^{-1} = (y)\mu$ and $((y)\phi_x)^{-1} \cdot (y)\phi_x = (y)\nu$.
- (h) If $x, y \in U_x$ are such that $(x)\mu =_\rho (y)\mu$, then $(x)\mu = (y)\mu$. Similarly, if $(x)\nu =_\rho (y)\nu$, then $(x)\nu = (y)\nu$.

Proof. (a) Suppose that $y =_\rho z$. Then by definition $(y)\phi_x = (y)\mu \cdot y \cdot (y)\nu$. Since $(y)\mu =_\rho yy^{-1}$ and $(y)\nu =_\rho y^{-1}y$, it follows that $(y)\phi_x = (y)\mu \cdot y \cdot (y)\nu =_\rho yy^{-1} \cdot y \cdot y^{-1}y = y$

Similarly $(z)\phi_x =_\rho z$, hence $(z)\phi_x =_\rho z =_\rho y =_\rho (y)\phi_x$ as required.

(b) Let $y \in \text{im}(\phi_x)$. Then there exists $z \in U_x$ such that $(z)\phi_x = y$. By part (a), $(z)\phi_x =_\rho z$ and so $y = (z)\phi_x =_\rho z$. In particular, $y/\rho = z/\rho$ and so $(y/\rho, x/\rho) = (z/\rho, x/\rho) \in \mathcal{D}^{S/\rho}$, since $z \in U_x$. This shows that $y \in U_x$.

Next, $y =_\rho z$ implies that $(y)\mu = (z)\mu$ ([Lemma 6.1\(h\)](#)). A similar argument shows that $(y)\nu = (z)\nu$. In particular, since $(y)\mu = (z)\mu$ and $(y)\nu = (z)\nu$ are idempotents, $(z)\mu = (y)\mu \cdot (z)\mu$ and $(z)\nu = (z)\nu \cdot (y)\nu$. Hence

$$\begin{aligned}
y &= (z)\phi_x = (z)\mu \cdot z \cdot (z)\nu \\
&= (y)\mu \cdot (z)\mu \cdot z \cdot (z)\nu \cdot (y)\nu \\
&= (y)\mu \cdot (z)\phi_x \cdot (y)\nu \\
&= (y)\mu \cdot y \cdot (y)\nu \\
&= (y)\phi_x.
\end{aligned}$$

(c) If $y, z \in U_x$, then $(y/\rho, x/\rho), (z/\rho, x/\rho) \in \mathcal{D}^{S/\rho}$ by definition. Thus, since ρ is transitive, $(y/\rho, z/\rho) \in \mathcal{D}^{S/\rho}$.

(d) If $(y, z) \in \mathcal{D}^S$, then $(y/\rho, z/\rho) \in \mathcal{D}^{S/\rho}$. Since $y \in U_x$, $(y/\rho, x/\rho)$, and so, again by the transitivity of ρ , $(z/\rho, x/\rho) \in \mathcal{D}^{S/\rho}$. Therefore $z \in U_x$.

(e) This follows immediately from the definition of ϕ_x , since $(z)\phi_x = (z)\mu \cdot z \cdot (z)\nu \leq z$.

(f) Since $y \leq z$, it follows that $x/\rho \leq y/\rho$. But by assumption $y, z \in U_x$ and so $(y/\rho, z/\rho) \in \mathcal{D}^{S/\rho}$, by part (c). Therefore $y/\rho = z/\rho$, as required.

(g) By part (a), $(y)\phi_x((y)\phi_x)^{-1} =_\rho yy^{-1}$ and so $(y)\phi_x((y)\phi_x)^{-1} \geq (y)\mu$, since $(y)\mu$ is the minimum of its trace class. On the other hand, $(y)\mu \cdot (y)\phi_x = (y)\phi_x$ and so $(y)\mu(y)\phi_x((y)\phi_x)^{-1} = (y)\phi_x((y)\phi_x)^{-1}$. Therefore $(y)\phi_x((y)\phi_x)^{-1} \leq (y)\mu$.

The proof for ν follows by symmetry.

(h) By definition both $(x)\mu$ and $(y)\mu$ are the minimum elements in their trace classes. Since $(x)\mu =_\rho (y)\mu$, these trace classes coincide, and so $(x)\mu = (y)\mu$. The proof for ν is the same. \square

For the next lemma it will be convenient to use the languages of groupoids. The set U_x naturally forms a groupoid with $*$: $U_x \times U_x \rightarrow U_x$ defined by

$$y * z = yz \quad \text{whenever } y, z, yz \in U_x \text{ and } yz \mathcal{D}^S y \mathcal{D}^S z$$

and where the inverse operation coincides with that on S . The connected components of U_x are just the \mathcal{D} -classes of S ; for further details see [[12](#), Section 3.1].

Lemma 6.2. *If $u \in U_x$, then $\phi_x|_{D_u^S}$ is a functor (or equivalently a groupoid morphism).*

Proof. Suppose that $y, z \in D_u^S$ are such that $yz \in D_u^S$. It suffices to show that $(y)\phi_x \cdot (z)\phi_x = (yz)\phi_x$. Since $yz \mathcal{D}^S u$, $y^{-1}y = zz^{-1}$ (by the [Location Theorem 2.1](#)), and so $(z)\mu = (y)\nu$. It follows that

$$\begin{aligned}
(y)\phi_x \cdot (z)\phi_x &= ((y)\mu \cdot y \cdot (y)\nu)((z)\mu \cdot z \cdot (z)\nu) \\
&= (y)\mu \cdot y \cdot (y)\nu \cdot z \cdot (z)\nu && \text{since } (y)\nu = (z)\mu \in E(S) \\
&= (y)\mu \cdot yz \cdot (z)\nu && \text{by Lemma 5.1.}
\end{aligned}$$

Since $(yz)(yz)^{-1} = yzz^{-1}y^{-1} = yy^{-1}yy^{-1} = yy^{-1}$, it follows that $(y)\mu = (yz)\mu$ and similarly, $(z)\nu = (yz)\nu$. Therefore

$$(y)\phi_x \cdot (z)\phi_x = (y)\mu \cdot yz \cdot (z)\nu = (yz)\mu \cdot yz \cdot (yz)\nu = (yz)\phi_x. \quad \square$$

Lemma 6.3. (a) *If $y \in \text{im}(\phi_x)$ and $z \in U_x$ is such that $z \leq y$, then $y = z$.*

(b) *If $e =_{\text{Tr}(\rho)} f$, $e \in \text{im}(\phi_x)$ is an idempotent, and $f \in U_x$ is also an idempotent, then $e \leq f$.*

Proof. Suppose that $y \in \text{im}(\phi_x)$ and $z \in U_x$ are such that $z \leq y$. If $z < y$, then $zz^{-1} < yy^{-1}$. Hence it suffices to show that yy^{-1} is minimal in U_x .

Since $y \in \text{im}(\phi_x)$, **Lemma 6.1(a)** shows that $(y)\phi_x = y \in U_x$. We begin by showing that yy^{-1} is the minimum in its trace class; this will establish part (b). By the definition of μ , it suffices to show that $yy^{-1} = (u)\mu$. This follows from **Lemma 6.1(g)**.

By the definition of ϕ_x , $y = (u)\mu \cdot u \cdot (u)\nu$ and so

$$yy^{-1} = ((u)\mu \cdot u \cdot (u)\nu)((u)\mu \cdot u \cdot (u)\nu)^{-1} = (u)\mu \cdot u \cdot (u)\nu \cdot ((u)\nu)^{-1} \cdot u^{-1} \cdot ((u)\mu)^{-1} \leq (u)\mu \cdot (u)\mu^{-1} = (u)\mu,$$

the last equality holds because $(u)\mu$ is an idempotent.

For the converse inequality, by **Lemma 6.1(a)**, $y = (u)\phi_x =_{\rho} u$, and so $yy^{-1} =_{\rho} uu^{-1} =_{\rho} (u)\mu$. So, $yy^{-1} =_{\rho} (u)\mu$, and since $(u)\mu$ is the minimum in its trace class, $yy^{-1} \geq (u)\mu$. We have shown that $yy^{-1} = (u)\mu$, meaning that yy^{-1} is the minimum in its trace class.

If $z \in U_x$ and $z \leq yy^{-1}$, then z is an idempotent, and $z/\rho \leq yy^{-1}/\rho$ (homomorphisms preserve the natural partial order). Since $y, z \in U_x$, it follows that $(y/\rho, z/\rho) \in \mathcal{D}^{S/\rho}$, by **Lemma 6.1(c)**. Since $(y, yy^{-1}) \in \mathcal{D}^S$, and homomorphisms preserve Green's \mathcal{D} -relation, $(y/\rho, yy^{-1}/\rho) \in \mathcal{D}^{S/\rho}$. Thus $(yy^{-1}/\rho, z/\rho) \in \mathcal{D}^{S/\rho}$ and $z/\rho \leq yy^{-1}/\rho$, which implies that $z/\rho = yy^{-1}/\rho$. Hence $z = yy^{-1}$ by **Lemma 6.1(h)** and so yy^{-1} is minimal in U_x , as required. \square

Lemma 6.4. *The set $\text{im}(\phi_x)$ is a \mathcal{D}^S -class.*

Proof. Suppose that $y \in \text{im}(\phi_x)$. We will show that $D_y^S = \text{im}(\phi_x)$.

(\supseteq) Suppose that $z \in \text{im}(\phi_x)$. Then $y, z \in U_x$ (**Lemma 6.1(b)**) and so $(y/\rho, z/\rho) \in \mathcal{D}^{S/\rho}$ (**Lemma 6.1(c)**). Hence there exists $s \in S$ such that $(s/\rho, y/\rho) \in \mathcal{D}^{S/\rho}$, $zz^{-1} =_{\rho} ss^{-1}$, and $yy^{-1} =_{\rho} s^{-1}s$. This implies that $s \in U_x$ and so $(s)\phi_x \in \text{im}(\phi_x)$. We will show that $(s)\phi_x((s)\phi_x)^{-1} = zz^{-1}$ and $((s)\phi_x)^{-1}(s)\phi_x = yy^{-1}$. It will follow from this that $(z, (s)\phi_x), ((s)\phi_x, y) \in \mathcal{D}^S$ implying that $(z, y) \in \mathcal{D}^S$ which will conclude the proof.

By **Lemma 6.1(a)**, $s =_{\rho} (s)\phi_x$ and so $(s)\phi_x((s)\phi_x)^{-1} =_{\rho} ss^{-1} =_{\rho} zz^{-1}$. Since $z \in \text{im}(\phi_x) \subseteq U_x$, it follows from the definition of U_x that $zz^{-1} \in U_x$ also. **Lemma 6.1(b)** implies that $z = (z)\phi_x$ and since ϕ_x is a functor (**Lemma 6.2**), $(zz^{-1})\phi_x = (z)\phi_x \cdot (z^{-1})\phi_x = (z)\phi_x \cdot ((z)\phi_x)^{-1} = zz^{-1}$ (the second to last equality holds because functors preserve inverses). Hence $zz^{-1} \in \text{im}(\phi_x)$, and similarly, $(s)\phi_x((s)\phi_x)^{-1} \in \text{im}(\phi_x)$. But $(s)\phi_x((s)\phi_x)^{-1} =_{\rho} zz^{-1}$, and so **Lemma 6.1(g,h)** implies that both $zz^{-1} = (s)\phi_x((s)\phi_x)^{-1}$. By symmetry $yy^{-1} = ((s)\phi_x)^{-1}(s)\phi_x$, as required.

(\subseteq) If $z \in D_y^S$, then $z \in U_x = \text{dom}(\phi_x)$, by **Lemma 6.1(d)**. It follows that $(z)\phi_x \leq z$ (by **Lemma 6.1(e)**) and so (assuming without loss of generality that S is an inverse semigroup of partial perms)

$$\text{rank}((z)\phi_x) \leq \text{rank}(z) = \text{rank}(y) = \text{rank}((y)\phi_x) = \text{rank}((z)\phi_x),$$

(the last equality holds since $\text{im}(\phi_x) \subseteq D_y^S$). Hence $(z)\phi_x = z$ and so $z \in \text{im}(\phi_x)$. \square


If $y, z \in S$ and $(y, z) \in \mathcal{D}^S$, then in the next theorem we will denote the intersection of the \mathcal{R} -class R_y^S of y and the \mathcal{L} -class L_z^S of z by $H_{y,z}$.

We can finally state and prove the main result in this section which will allow us to compute the elements in the congruence class $x/\rho \cap H_{e,f}$ where $e, f \in E(S)$, as a translate of the preimage of a coset of a normal subgroup under the functor ϕ_x .

Theorem 6.5. *If $e, f \in E(S)$ are such that $(e, f) \in \mathcal{D}^S$ and $x \in S$, then there exists $s \in S$ such that*

$$H_{e,f} \cap x/\rho = ((H_{(e)\phi_x, (e)\phi_x} \cap e/\rho) xs^{-1}) \phi_x|_{H_{e,e}}^{-1} \cdot s,$$

where $e =_{\rho} xx^{-1}$, $f =_{\rho} x^{-1}x$, and $s^{-1}es = f$.


 FIGURE 3. A diagram of the proof of [Theorem 6.5](#).

Before giving the proof of [Theorem 6.5](#) note that $H_{(e)\phi_x, (e)\phi_x} \cap e/\rho$ is a normal subgroup of $H_{(e)\phi_x, (e)\phi_x}$ by [Lemma 5.2\(a\)](#). Hence $(H_{(e)\phi_x, (e)\phi_x} \cap e/\rho)xs^{-1}$ is a coset of a normal subgroup. Since $H_{e,e}$ is a group, and ϕ_x is a functor ([Lemma 6.2](#)), it follows that $\phi_x|_{H_{e,e}} : H_{e,e} \rightarrow H_{(e)\phi_x, (e)\phi_x}$ is a group homomorphism. The ideas of the proof are visualised in [Figure 3](#).

Proof of [Theorem 6.5](#). We start by noting that:

$$fs^{-1}s = s^{-1}ess^{-1}s = s^{-1}es = f,$$

which will be useful in both parts of the proof below.

(\subseteq) Let $t \in H_{e,f} \cap x/\rho$. Since $t \in H_{e,f}$, $ts^{-1}s = t$ (Green's Lemma [[10](#), Lemma 2.2.1]) and so $ts^{-1} \in H_{e,e}$. In particular, $(ts^{-1}, e) \in \mathcal{D}^S$ and $e = \rho xx^{-1} \in U_x$ and so $e \in U_x$. Thus $ts^{-1} \in U_x$ ([Lemma 6.1\(d\)](#)) and so it suffices to show that $(ts^{-1})\phi_x \in (H_{(e)\phi_x, (e)\phi_x} \cap e/\rho)xs^{-1}$.

We start by showing that $(ts^{-1})\phi_x \cdot sx^{-1} =_{\rho} e$:

$$\begin{aligned}
(ts^{-1})\phi_x \cdot sx^{-1} &=_{\rho} (ts^{-1})sx^{-1} && ts^{-1} =_{\rho} (ts^{-1})\phi_x \text{ by Lemma 6.1(a)} \\
&=_{\rho} (xs^{-1})sx^{-1} && t =_{\rho} x \text{ by assumption} \\
&= (xx^{-1}xs^{-1})sx^{-1} \\
&=_{\rho} (xfs^{-1}ss^{-1})sx^{-1} && fs^{-1}s =_{\rho} x^{-1}x \\
&= xfs^{-1}sx^{-1} \\
&= xfx^{-1} && fs^{-1}s = f \\
&=_{\rho} xx^{-1}xx^{-1} && f =_{\rho} x^{-1}x \\
&= xx^{-1} \\
&=_{\rho} e.
\end{aligned}$$

It remains to show that $(ts^{-1})\phi_x \in H_{(e)\phi_x, (e)\phi_x} xs^{-1}$, or, by Green's Lemma, equivalently that $(ts^{-1})\phi_x sx^{-1} \in H_{(e)\phi_x, (e)\phi_x}$.

In order to do this, we start by showing that $sx^{-1} \in U_x$. Since $(sx^{-1}, sx^{-1}xs^{-1}) \in \mathcal{D}^S$, it follows that $(sx^{-1}/\rho, sx^{-1}xs^{-1}/\rho) \in \mathcal{D}^{S/\rho}$. But $sx^{-1}xs^{-1} =_{\rho} sfs^{-1} = ss^{-1}ess^{-1} = ess^{-1} = e =_{\rho} xx^{-1}$ and so $(sx^{-1}xs^{-1}/\rho, x/\rho) \in \mathcal{D}^{S/\rho}$. By transitivity, $(sx^{-1}/\rho, x/\rho) \in \mathcal{D}^{S/\rho}$ and so $sx^{-1} \in U_x$ (Lemma 6.1(d)). By the definitions of μ and ν :

$$(1) \quad (ts^{-1})\nu =_{\rho} (ts^{-1})^{-1}ts^{-1} = st^{-1}ts =_{\rho} sx^{-1}xs^{-1} = sx^{-1}(sx^{-1})^{-1} =_{\rho} (sx^{-1})\mu$$

so $(ts^{-1})\nu = (sx^{-1})\mu$ (Lemma 6.1(h)). On the other hand, $ts^{-1}st^{-1} \leq tt^{-1}$ and $tt^{-1} \in U_x$ since $t =_{\rho} x$. If $ts^{-1}st^{-1} \in U_x$, then $(ts^{-1})\mu = ts^{-1}st^{-1} =_{\text{Tr}(\rho)} tt^{-1} = (t)\mu$ (Lemma 6.1(f)). To show that $ts^{-1}st^{-1} \in U_x$ it suffices to show that $ts^{-1}st^{-1}$ is ρ -related to an element of U_x :

$$\begin{aligned}
ts^{-1}st^{-1} &= ts^{-1}st^{-1}tt^{-1} \\
&=_{\rho} ts^{-1}sx^{-1}xt^{-1} && t^{-1}t =_{\rho} x^{-1}x \\
&=_{\rho} ts^{-1}sft^{-1} && x^{-1}x =_{\rho} f \\
&= ts^{-1}ss^{-1}est^{-1} && s^{-1}es = f \\
&= ts^{-1}est^{-1} \\
&=_{\rho} tft^{-1} && s^{-1}es = f \\
&=_{\rho} tx^{-1}xt^{-1} && x^{-1}x =_{\rho} f \\
&=_{\rho} xx^{-1}xx^{-1} && t =_{\rho} x \\
&=_{\rho} xx^{-1} \in U_x.
\end{aligned}$$

Hence $(ts^{-1})\mu = (t)\mu$ (Lemma 6.1(h)). By the assumption at the start of the proof, $t =_{\rho} x$ and so $tt^{-1} =_{\rho} xx^{-1}$, and so $(t)\mu = (xx^{-1})\mu$. Since $e \in E(S)$, $(e)\phi_x = (e)\mu =_{\rho} (xx^{-1})\mu$, and again by Lemma 6.1(h), $(e)\mu = (xx^{-1})\mu$. We have shown that

$$(2) \quad (ts^{-1})\mu = (e)\phi_x.$$

By a similar argument, $(sx^{-1})\nu = (x^{-1})\nu = (x)\mu = (e)\phi_x$.

It follows that

$$\begin{aligned}
(ts^{-1})\phi_x \cdot sx^{-1} &= (ts^{-1})\phi_x \cdot (ts^{-1})\nu \cdot sx^{-1} && \text{by the definition of } \phi_x \\
&= (ts^{-1})\phi_x \cdot (sx^{-1})\mu \cdot sx^{-1} && \text{by (1)} \\
&= (ts^{-1})\phi_x \cdot (sx^{-1})\mu \cdot sx^{-1} \cdot (sx^{-1})\nu && \text{by Lemma 5.1} \\
&= (ts^{-1})\phi_x \cdot (sx^{-1})\phi_x && sx^{-1} \in U_x.
\end{aligned}$$

We set $a = (ts^{-1})\phi_x$ and $b = (sx^{-1})\phi_x$. By (1), $(ts^{-1})\nu = (sx^{-1})\mu$ and so by Lemma 6.1(g), $a^{-1}a = bb^{-1}$ and so $(ab, a) \in \mathcal{D}^S$. The Location Theorem 2.1 then implies that $ab \in H_{aa^{-1}, b^{-1}b}$. But

$$\begin{aligned} aa^{-1} &= (ts^{-1})\phi_x((ts^{-1})\phi_x)^{-1} \\ &= (ts^{-1})\mu && \text{by Lemma 6.1(g)} \\ &= (e)\phi_x && \text{by (2).} \end{aligned}$$

Similarly, $b^{-1}b = (sx^{-1})\nu = (e)\phi_x$. Whence $(ts^{-1})\phi_x \cdot sx^{-1} = (ts^{-1})\phi_x \cdot (sx^{-1})\phi_x = ab \in H_{aa^{-1}, b^{-1}b} = H_{(e)\phi_x, (e)\phi_x}$, as required.

(\supseteq) Let $t \in ((H_{(e)\phi_x, (e)\phi_x} \cap e/\rho)xs^{-1})\phi_x|_{H_{e,e}}^{-1} \cdot s$ be arbitrary. We must show that $t \in H_{e,f}$ and $t \in x/\rho$. Since $t \in \text{dom}(\phi_x|_{H_{e,e}})s$, there exists $h \in H_{e,e}$ such that $t = hs \in H_{e,e}s = H_{e,f}$.

It remains to prove that $t =_{\rho} x$:

$$\begin{aligned} t &= hs && \text{where } h \in ((H_{(e)\phi_x, (e)\phi_x} \cap e/\rho)xs^{-1})\phi_x|_{H_{e,e}}^{-1} \\ &=_{\rho} (h)\phi_x s && \text{by Lemma 6.1(a)} \\ &=_{\rho} exs^{-1}s && \text{by the choice of } h \\ &= exx^{-1}xs^{-1}s \\ &=_{\rho} exfs^{-1}s && f =_{\rho} x^{-1}x \\ &= exf && fs^{-1}s = f \\ &=_{\rho} xx^{-1}xx^{-1}x \\ &= x, \end{aligned}$$

as required. \square

The algorithm for iterating through the elements of the set x/ρ is then as follows:

- (X1) determine the data structure for the semigroup S consisting of: the generating set X , the word graph Γ_X , the strongly connected components of Γ_X ; and one group \mathcal{H} -class per strongly connected component of Γ_X using the algorithms described in [5, Section 5.6];
- (X2) determine the data structure for the quotient S/ρ consisting of: the generating set X ; the quotient word graph $\Gamma_X/\text{Tr}(\rho)$; the strongly connected components of $\Gamma_X/\text{Tr}(\rho)$; and the quotient groups G/N using the algorithms described in Section 4 and Section 5;
- (X3) for every pair $\{e, f\}$ of idempotents where $(e, xx^{-1}), (f, x^{-1}x) \in \text{Tr}(\rho)$ and e and f belong to the same strongly connected component of Γ_X determine the set $H_{e,f} \cap x/\rho$ using Theorem 6.5.

Example 6.6. We continue Example 5.4 by computing x/ρ where $x = [1\ 2\ 4](3) \in I_4$. Steps (X1) and (X2) were covered in Example 4.3 and Example 5.4, respectively. For step (X3), we iterate though all the pairs of idempotents e, f such that

$$e =_{\rho} xx^{-1} = (1)(2)(3) \quad \text{and} \quad f =_{\rho} x^{-1}x = (2)(3)(4).$$

In this case, there is only one such pair when $e = (1)(2)(3)$ and $f = (2)(3)(4)$. We then compute $((H_{(e)\phi_x, (e)\phi_x} \cap e/\rho)xs^{-1})\phi_x|_{H_{e,e}}^{-1} \cdot s$ where $s \in I_4$ is any fixed element such that $s^{-1}es = f$; such as $s = [1\ 2\ 3\ 4]$. Since $\text{Tr}(\rho)$ equals $\Delta_{D_x \cap E(S)} = \{(d, d) \mid d \in D_x \cap E(S)\}$ when restricted to the idempotents of the \mathcal{D} -class of x , it follows that ϕ_x is the identity function. Thus $((H_{e,e} \cap e/\rho)xs^{-1}) \cdot s = (H_{e,e} \cap e/\rho)x$. We calculated in the previous example that $H_{e,e} \cap e/\rho$ is the alternating group on $\{1, 2, 3\}$, that is $\{(1)(2)(3), (1\ 2\ 3), (1\ 3\ 2)\}$. Translating this by $x = [1\ 2\ 4](3)$ gives

$$x/\rho = \{(1)(2)(3) \cdot [1\ 2\ 4](3), (1\ 2\ 3) \cdot [1\ 2\ 4](3), (1\ 3\ 2) \cdot [1\ 2\ 4](3)\} = \{[1\ 2\ 4](3), [1\ 4](2\ 3), [1\ 3\ 4](2)\}.$$

7. THE MAXIMUM IDEMPOTENT-SEPARATING CONGRUENCE

In this section we give a method for computing the maximum idempotent-separating congruence on a finite inverse subsemigroup of a symmetric inverse monoid. We achieve this using a description of the maximum idempotent-separating congruence via centralisers. We begin with the definition of a centraliser.

If S is a semigroup and A is a subset of S , then the **centraliser** of A in S is the set

$$C_S(A) = \{s \in S \mid sa = as \text{ for all } a \in A\}.$$

Let μ be the congruence defined by $a =_\mu b$ if and only if $aea^{-1} = beb^{-1}$ for all $e \in E(S)$.

Lemma 7.1 (cf. Section 5.2 in [12]). *The congruence μ is the maximum idempotent-separating congruence on S .*

We have that $\text{Ker}(\mu) = C_S(E(S))$ and since μ is the maximum idempotent-separating congruence on an inverse semigroup S , $\text{Tr}(\mu) = \Delta_{E(S)}$. For the remainder of this section, we discuss how to compute $C_S(E(S))$ when $S \leq I_n$.

If S is an inverse semigroup of partial permutations of degree n , and $X \subseteq \{1, \dots, n\}$, then the (**setwise**) **stabiliser** of X with respect to S is

$$\text{Stab}_S(X) = \{g \in S \mid (X)g = X\} \leq S.$$

Proposition 7.2. *If $S \leq I_n$ is an inverse semigroup, then*

$$C(E(S)) = \bigcup_{e \in E(S)} \bigcap_{f \leq e} \text{Stab}_{S \cap \text{Sym}(\text{dom}(e))}(\text{dom}(f)).$$

Where $\text{Sym}(\text{dom}(e))$ denotes the group of permutations of $\text{dom}(e)$, and f is taken to be in S .

Proof. (\subseteq) Let $s \in C(E(S))$ and $e = ss^{-1}$. As $se = es = s$ and $\text{dom}(e) = \text{dom}(s)$ it follows that

$$\text{dom}(e) = \text{dom}(s) = \text{dom}(se) = (\text{dom}(e))s^{-1}.$$

Thus s^{-1} bijectively maps $\text{dom}(e) = \text{dom}(s)$ to itself. So s does too, and so $s \in S \cap \text{Sym}(\text{dom}(e))$. Let $f \leq e$. To conclude that s belongs to the right side of the equality in the statement, it suffices to show that $(\text{dom}(f))s = \text{dom}(f)$. By assumption $fs = sf$, so

$$\text{dom}(f)s = \text{im}(fs) = \text{im}(sf) = \text{im}(s) \cap \text{dom}(f) = \text{dom}(e) \cap \text{dom}(f) = \text{dom}(f).$$

(\supseteq) Let s be an element of the right hand side of the equality in the statement of the proposition. Then there exists $e \in E(S)$ such that for all $f \leq e$, we have $s \in \text{Stab}_{S \cap \text{Sym}(\text{dom}(e))}(\text{dom}(f))$. In particular, this holds when $f = e$, and so $s \in S \cap \text{Sym}(\text{dom}(e))$. Let $g \in E(S)$ be arbitrary. We need to show that $sg = gs$. Since s is an element of a subgroup with identity e , it follows that $ss^{-1} = s^{-1}s = e$. If we define $f = eg$, then as $f \leq e$, $(\text{dom}(f))s = \text{dom}(f)$. This implies that $(\text{dom}(f))s^{-1} = \text{dom}(f)$ and so

$$gs = ges = fs = s|_{\text{dom}(f)} = s|_{\text{dom}(f)s^{-1}} = sf = seg = sg. \quad \square$$

If $A \subseteq \mathcal{P}(X)$ for some set X , then we say that A is a **boolean algebra** (on X) if A is closed under taking finite (possibly empty) unions, and is also closed under taking complements in X . Each boolean algebra is partially ordered by \subseteq and contains the empty set, which is called the 0 of the algebra. The complement of 0 (the universal set) is similarly called the 1. Note that this is consistent with standard meet semilattice notation. If $Y \subseteq X$ we write Y^c to denote the complement of Y in X . We say that an element of a boolean algebra is an **atom** if it is a minimal non-zero element. If B is a boolean algebra, then we define $A(B)$ to be the set of atoms of B . For any finite boolean algebra B , $B = \{\cup X \mid X \subseteq A(B)\}$. If $S \leq I_n$ is an inverse semigroup, then we define $B(S) \leq \mathcal{P}(\{1, 2, \dots, n\})$ to be the least boolean algebra containing the set of domains (or equivalently images) of the elements of S , noting that such a boolean algebra exists as the intersection of two boolean algebras is always a boolean algebra.

Theorem 7.3. *If $S \leq I_n$ is an inverse semigroup, then*

$$C(E(S)) = \{s \in S \mid (b)s = b \text{ for all } b \in A(B(S)) \text{ such that } b \subseteq \text{dom}(s)\}.$$

Proof. (\subseteq) Let $s \in C(E(S))$. We must show that $(b)s = b$ for all $b \in A(B(S))$ such that $b \subseteq \text{dom}(s)$. Let $b \in A(B(S))$ be such that $b \subseteq \text{dom}(s)$ and let

$$\begin{aligned} X &= \{Y \subseteq \{1, \dots, n\} \mid (Y)s \subseteq Y \text{ and } (Y^c)s \subseteq Y^c\} \\ X' &= \{Y \subseteq \{1, \dots, n\} \mid (Y)s \subseteq Y \text{ and } (Y)s^{-1} \subseteq Y\}. \end{aligned}$$

We show that $X = X'$. Let $Y \in X$. Then $(Y)s \subseteq Y$ and $(Y^c)s \subseteq Y^c$. So s moves nothing from Y to Y^c and nothing from Y^c to Y , and thus the same must hold for s^{-1} . In particular, $(Y)s^{-1} \subseteq Y$ and so $Y \in X'$ and

$X \subseteq X'$. Now suppose $Y \in X'$. Then $(Y)s \subseteq Y$ and $(Y)s^{-1} \subseteq Y$. The later implies that s cannot move anything from Y^c into Y and so $(Y^c)s \subseteq Y^c$ and $Y \in X$. Thus $X' \subseteq X$ and so $X' = X$.

Note that X is a boolean algebra, as from the definition of X it is closed under complements, and from the definition of X' it is closed under unions. Let $D = \{\text{dom}(t) \mid t \in S\}$. By definition, the least boolean algebra containing D is $B(S)$. We will show that $D \subseteq X$. This will be sufficient because, together with fact that X is a boolean algebra, this implies that $B(S) \subseteq X$, which in turn implies that $(b)s \subseteq b$. Since b is an atom $(b)s$ cannot be a proper subset of b .

So let $d \in D$ be arbitrary, and let $f_d \in S$ be an idempotent with $\text{dom}(f_d) = d$. As $s \in C(E(S))$, we have $sf_d = f_d s$. The image of sf_d is $d \cap \text{im}(s)$ and the image of $f_d s$ is $(d)s$. Thus $(d)s = d \cap \text{im}(s)$ and so $(d)s \subseteq d$. Since $s^{-1} \in C(E(S))$, we similarly get that $(d)s^{-1} \subseteq d$. It follows that $d \in X$, as required.

(\supseteq) Let $s \in S$ be such that for all $b \in A(B(S))$ with $b \subseteq \text{dom}(s)$, we have $(b)s = b$. Let $e \in S$ be an idempotent. We will show that $se = es$. Let $b_1, \dots, b_k \in A(B(S))$ be distinct such that $\text{dom}(e) \cap \text{dom}(s) = b_1 \cup \dots \cup b_k$. Note that, from the assumption on s , $(b_i)s = b_i$ for all $1 \leq i \leq k$ so

$$\text{dom}(e) \cap \text{dom}(s) = b_1 \cup \dots \cup b_k = \text{dom}(e) \cap \text{im}(s).$$

For all $x \in \{1, \dots, n\}$ we have that

$$\begin{aligned} (\{x\})es &= \begin{cases} \emptyset & \text{if } x \notin b_1 \cup \dots \cup b_k \\ \{x\}s & \text{if } x \in b_1 \cup \dots \cup b_k \end{cases} \\ &= (\{x\})se. \end{aligned}$$

Therefore, $se = es$ and so $s \in C(E(S))$, as required. \square

Example 7.4. We compute the maximum idempotent-separating congruence μ of the semigroup I_4 . The first step is to construct $C(E(I_4))$ using [Theorem 7.3](#). The set of domains of elements of I_4 is just $\mathcal{P}(I_4)$, and so $B(I_4) = \mathcal{P}(\{1, 2, 3, 4\})$. It follows that $A(B(I_4))$ is the set of singleton subsets of $\{1, 2, 3, 4\}$. From [Theorem 7.3](#), it follows that

$$C(E(I_4)) = \{s \in I_4 \mid (i)s = i \text{ for all } i \in \{1, 2, 3, 4\} \text{ such that } i \in \text{dom}(s)\}.$$

This implies that $C(E(I_4))$ is precisely the set of elements of I_4 which act as the identity on their domains, which is just $E(I_4)$ and so $\text{Ker}(\mu) = C(E(I_4)) = E(I_4)$. Since μ is idempotent-separating, we already know that $\text{Tr}(\mu) = \Delta_{E(S)}$, and so we have computed the kernel and trace for μ , which fully describes the congruence. In this case, the kernel and trace equal those of the trivial congruence Δ_S , and so $\mu = \Delta_S$.

ACKNOWLEDGEMENTS

The authors were supported by a Heilbronn Institute for Mathematical Research Small Grant during part of this work. The second named author was supported the Heilbronn Institute for Mathematical Research during this work. The authors would also like to thank the University of Manchester for hosting them during part of the work on this paper.

REFERENCES

- [1] M. Anagnostopoulou-Merkouri, R. Cirpons, J. D. Mitchell, and M. Tsalakou. *Computing finite index congruences of finitely presented semigroups and monoids*. 2023. eprint: [arXiv:2302.06295](#).
- [2] J. Araújo, R. B. Pereira, W. Bentz, C. Chow, J. Ramires, L. Sequeira, and C. Sousa. *CREAM: a Package to Compute [Auto, Endo, Iso, Mono, Epi]-morphisms, Congruences, Divisors and More for Algebras of Type $(2^n, 1^n)$* . 2022. eprint: [arXiv:2202.00613](#).
- [3] S. N. Burris and H. P. Sankappanavar. *A Course in Universal Algebra*. en. 1981st ed. Graduate texts in mathematics. New York, NY: Springer, Oct. 2011.
- [4] T. D. H. Coleman, J. D. Mitchell, F. L. Smith, and M. Tsalakou. “The Todd–Coxeter algorithm for semigroups and monoids”. In: *Semigroup Forum* (May 2024). ISSN: 1432-2137. DOI: [10.1007/s00233-024-10431-z](#). URL: [http://dx.doi.org/10.1007/s00233-024-10431-z](#).
- [5] J. East, A. Egri-Nagy, J. D. Mitchell, and Y. Péresse. “Computing finite semigroups”. In: *Journal of Symbolic Computation* 92 (May 2019), pp. 110–155. ISSN: 0747-7171. DOI: [10.1016/j.jsc.2018.01.002](#). URL: [http://dx.doi.org/10.1016/j.jsc.2018.01.002](#).

- [6] H. N. Gabow. “Path-based depth-first search for strong and biconnected components”. In: *Information Processing Letters* 74.34 (2000), pp. 107–114. ISSN: 0020-0190. DOI: [http://dx.doi.org/10.1016/S0020-0190\(00\)00051-X](http://dx.doi.org/10.1016/S0020-0190(00)00051-X). URL: <http://www.sciencedirect.com/science/article/pii/S002001900000051X>.
- [7] *GAP – Groups, Algorithms, and Programming, Version 4.13.0*. The GAP Group. 2024. URL: <https://www.gap-system.org>.
- [8] P. A. Grillet. *Semigroups: An Introduction to the Structure Theory*. Nov. 2017. DOI: [10.4324/9780203739938](https://doi.org/10.4324/9780203739938). URL: <http://dx.doi.org/10.4324/9780203739938>.
- [9] D. F. Holt, B. Eick, and E. A. O’Brien. *Handbook of Computational Group Theory*. Chapman and Hall/CRC, Jan. 2005. ISBN: 9780429147944. DOI: [10.1201/9781420035216](https://doi.org/10.1201/9781420035216). URL: <http://dx.doi.org/10.1201/9781420035216>.
- [10] J. M. Howie. *Fundamentals of semigroup theory*. Vol. 12. London Mathematical Society Monographs. New Series. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1995, pp. x+351. ISBN: 0-19-851194-9.
- [11] A. Hulpke. “Computing normal subgroups”. In: *Proceedings of the 1998 international symposium on Symbolic and algebraic computation*. ISSAC98. ACM, Aug. 1998. DOI: [10.1145/281508.281612](https://doi.org/10.1145/281508.281612). URL: <http://dx.doi.org/10.1145/281508.281612>.
- [12] M. V. Lawson. *Inverse Semigroups: The Theory of Partial Symmetries*. World Scientific, Nov. 1998. ISBN: 9789812816689. DOI: [10.1142/3645](https://doi.org/10.1142/3645). URL: <http://dx.doi.org/10.1142/3645>.
- [13] A. E. Liber. “On symmetric generalized groups”. In: *Mat. Sbornik N.S.* 33/75 (1953), pp. 531–544.
- [14] A. I. Mal’cev. “Multiplicative congruences of matrices”. In: *Doklady Akad. Nauk SSSR (N.S.)* 90 (1953), pp. 333–335.
- [15] A. I. Mal’cev. “Symmetric groupoids”. In: *Mat. Sbornik N.S.* 31/73 (1952), pp. 136–151.
- [16] J. D. Mitchell et al. *libsemigroups – C++ library - version 2.7.3*. Jan. 2024. URL: <https://libsemigroups.rtf.d.io>.
- [17] J. D. Mitchell et al. *Semigroups – GAP package - version 5.3.7*. Mar. 2024. URL: <https://semigroups.github.io/Semigroups/>.
- [18] R. B. Pereira et al. *The CREAM GAP Package - Algebra CongRuences, Endomorphisms and Automorphisms. version 2.0.14*. June 2022. URL: <https://gitlab.com/rmbper/cream>.
- [19] M. Petrich. *Inverse semigroups*. Pure and Applied Mathematics (New York). A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York, 1984, pp. x+674. ISBN: 0-471-87545-7.
- [20] H. E. Scheiblich. “Kernels of inverse semigroup homomorphisms”. In: *J. Austral. Math. Soc.* 18 (1974), pp. 289–292.
- [21] Á. Seress. *Permutation Group Algorithms*. Cambridge University Press, Mar. 2003. ISBN: 9780511546549. DOI: [10.1017/cbo9780511546549](https://doi.org/10.1017/cbo9780511546549). URL: <http://dx.doi.org/10.1017/CB09780511546549>.
- [22] R. Tarjan. “Depth-first search and linear graph algorithms”. In: *SIAM J. Comput.* 1.2 (1972), pp. 146–160. ISSN: 0097-5397.
- [23] M. Torpey. “Semigroup congruences : computational techniques and theoretical applications”. PhD thesis. University of St Andrews, 2019. DOI: [10.17630/10023-17350](https://doi.org/10.17630/10023-17350).
- [24] Z.-P. Wang. “Congruences on graph inverse semigroups”. In: *J. Algebra* 534 (2019), pp. 51–64. ISSN: 0021-8693,1090-266X. DOI: [10.1016/j.jalgebra.2019.06.020](https://doi.org/10.1016/j.jalgebra.2019.06.020). URL: <https://doi.org/10.1016/j.jalgebra.2019.06.020>.