

Coralai: Intrinsic Evolution of Embodied Neural Cellular Automata Ecosystems

Aidan Barbieux¹ and Rodrigo Canaan¹

¹California Polytechnic State University, San Luis Obispo
aidanbx@email.com

Abstract

This paper presents Coralai, a framework for exploring diverse ecosystems of Neural Cellular Automata (NCA). Organisms in Coralai utilize modular, GPU-accelerated Taichi kernels to interact, enact environmental changes, and evolve through local survival, merging, and mutation operations implemented with HyperNEAT and PyTorch. We provide an exploratory experiment implementing physics inspired by slime mold behavior showcasing the emergence of competition between sessile and mobile organisms, cycles of resource depletion and recovery, and symbiosis between diverse organisms. We conclude by outlining future work to discover simulation parameters through measures of multi-scale complexity and diversity. Code for Coralai is available at <https://github.com/aidanbx/coralai>, video demos are available at <https://www.youtube.com/watch?v=NL8IZQY02-8>.

Introduction

Most, if not all, of life exists within diverse networks of organisms that interact with each other, physics, and their environment to collectively develop resilience. Simulating this complexity manually is impractical and limited by our understanding of the system. Conversely, the emergence of complex behavior from simple rules, as demonstrated by many Cellular Automata or Neural Soups, can be difficult to interpret, control, and compute Plantec et al. (2023); Gregor and Besse (2021). Neural Cellular Automata (NCA) enable efficient training of low-level rules to create systems which self-organize to a defined state and have shown success in a wide variety of tasks (Mordvintsev et al., 2022; Pande and Grattarola, 2023; Stovold, 2023). By leaving trained NCA running, minimal evolutionary properties emerge Sinapayen (2023). By explicitly evolving rules to survive and interact with ‘physics,’ they display more diverse, lifelike behaviors Randazzo and Mordvintsev (2023). This shows significant potential to model the emergence of complex ecosystems, but current work is limited to specific behavior around plant-like growth.

Coralai is designed for the exploration of a wide variety of organism, environment, and physics configurations in




Figure 1: A single update in Coralai. (A): The sensing of substrate and production of actions. (B): Applying the actions of the entire system to the substrate via ‘physics’. (C): Visualization and radiation of the substrate resulting in mutation of CPPN genomes and generation of dense network parameters.

evolved NCA ecosystems. Organisms in Coralai produce abstract output which can then be arbitrarily transformed by physics to enact change in the environment, interact with other organisms, and evolve. Experiments can be defined at a high level and arbitrary physics can be implemented using GPU-accelerated Taichi kernels (Hu et al., 2019). Interactive simulations run with simple visualizations in real-time and could be scaled to larger systems. These features provide the basis of a powerful experimental platform for studying the emergence of complex, resilient ecosystems.

System Design

Each experiment in Coralai is defined by Substrate, Sensors, Actuators, Kernel, and Physics. The substrate is a tensor with width, height, and named channels/sub-channels. One such channel is *GenomeIndex*, which locates the neural

network parameters governing the behavior of the cell. In discrete time steps, each cell is updated by sending a neighborhood (the Kernel) of channels (the Sensors) to the cell’s network to receive values for the Actuator channels, which are then transformed via Physics to produce changes in the substrate.

The networks of each cell are generated via HyperNEAT (Hypercube Based Neuroevolution of Augmented Topologies) (Stanley et al., 2009) implemented in PyTorch which uses small Compositional Pattern Producing Networks (CPPN) to generate the weights and biases of a larger network. Coralai uses only the mutation and crossover operators of NEAT, which are applied via the Physics.

Experiment

To test Coralai, we conducted exploratory experiments resembling the physical affordances exploited by slime molds. This was chosen for the well-studied behaviors slime molds can exhibit with simple physiologies, such as network optimization, transfer learning, and decision making Ray et al. (2019); Vogel and Dussutour (2016); Vallverdu et al. (2017), developing the work of Barbieux and Canaan (2023).

The main channels of the experiment are Infrastructure, Energy, and Communication (shown with 3 sub-channels in figure 1). Energy is added and removed from the substrate in a day/night cycle. Organisms can transform Energy into Infrastructure via Invest and Liquidate actuators. Communication channels are simply normalized and activated via a sigmoid function. The kernel is a Moore’s neighborhood with offsets ordered by increasing angular distance to a Rotation channel of each cell.

Organisms spread via Explore actuators matched to each offset in the kernel. Exploration spreads infrastructure to a neighboring cell determined by the maximal actuator. If the quantity of the exploratory infrastructure exceeds that of the explored cell, the explored cell adopts the genome of the original cell and the rotation of the exploration. Otherwise, the infrastructure is taken while the current genome and rotation are maintained. If multiple cells explore a given cell, the one with maximal exploratory investment wins.

We ran exploratory simulations modulating substrate size, initial conditions, and hyper-parameters. We observed a wide variety of behaviors and patterns requiring further quantitative analysis. Figure 2 provides noteworthy examples showing mobile and sessile organisms, dominance overcome by diversity, and an ecosystem at equilibrium. We additionally observed cycles of greedy organisms consuming excess energy and causing mass-extinction followed by diversity and resilience and networked organisms spreading via spore-like cells upon depletion of an energy pocket.

Simulations of 1024x1024 cells with 512 unique organisms ran at 20 FPS on an M1 chip. Runs were limited to manual set up and observation with various initial conditions and physical parameters.




Figure 2: Notable experiment runs. Visual observation shows competition between mobile and sessile organisms in (A), extinction and replacement in (B), a variety of organisms stabilized in an ecosystem in (C), and competing slime-mold-like organisms in (D). Energy is displayed as red, infrastructure by green, and genome/species by blue.

Discussion and Future Work

Evolving diverse Neural Cellular Automata to collectively survive physics provides an easy-to-use and powerful basis to study artificial life. However, due to the large parameter spaces and possible physics in Coralai, intelligent search strategies should be employed. To move beyond manual observation and implementation of physics, we suggest meta-evolution of both parameters and physical rules to optimize known indicators of life.

One promising metric is multi-scale complexity (MSC) to detect self-organizing criticality, a crucial component of living systems (Bagrov et al., 2020; Bak et al., 1989). MSC calculation via renormalization-groups (but not meta-evolution) is implemented in Coralai and described in more detail by Barbieux (2024). We also suggest ecological measures of diversity (Eysenbach et al., 2018). Coralai can use NEAT’s measure of genetic similarity (Stanley and Miikkulainen, 2002) and tracks phylogeny to enable this.

Finally, we suggest future experiments that cater to niche formation, promoted through physics for obstacles, local weather patterns, ‘cross-pollination’ between simulations, among other techniques.

References

- Bagrov, A. A., Iakovlev, I. A., Iliasov, A. A., Katsnelson, M. I., and Mazurenko, V. V. (2020). Multiscale structural complexity of natural patterns. *Proceedings of the National Academy of Sciences*, 117(48):30241–30251. Publisher: Proceedings of the National Academy of Sciences.
- Bak, P., Chen, K., and Creutz, M. (1989). Self-organized criticality in the 'Game of Life'. *Nature*, 342(6251):780–782. Publisher: Nature Publishing Group.
- Barbieux, A. (2024). Coralai: Emergent Ecosystems of Neural Cellular Automata. *Digital Commons*.
- Barbieux, A. and Canaan, R. (2023). EINCASM: Emergent Intelligence in Neural Cellular Automaton Slime Molds. In *Artificial Life 2023*. MIT Press.
- Eysenbach, B., Gupta, A., Ibarz, J., and Levine, S. (2018). Diversity is All You Need: Learning Skills without a Reward Function. arXiv:1802.06070 [cs].
- Gregor, K. and Besse, F. (2021). Self-Organizing Intelligent Matter: A blueprint for an AI generating algorithm. arXiv:2101.07627 [cs].
- Hu, Y., Li, T.-M., Anderson, L., Ragan-Kelley, J., and Durand, F. (2019). Taichi: a language for high-performance computation on spatially sparse data structures. *ACM Transactions on Graphics*, 38(6):1–16.
- Mordvintsev, A., Randazzo, E., and Fouts, C. (2022). Growing Isotropic Neural Cellular Automata. In *The 2022 Conference on Artificial Life*, Online. MIT Press.
- Pande, R. and Grattarola, D. (2023). Hierarchical Neural Cellular Automata. In *Artificial Life 2023*. MIT Press.
- Plantec, E., Hamon, G., Etcheverry, M., Oudeyer, P.-Y., Moulin-Frier, C., and Chan, B. W.-C. (2023). Flow-Lenia: Towards open-ended evolution in cellular automata through mass conservation and parameter localization. arXiv:2212.07906 [nlin].
- Randazzo, E. and Mordvintsev, A. (2023). Biomaker CA: a Biome Maker project using Cellular Automata. arXiv:2307.09320 [cs].
- Ray, S. K., Valentini, G., Shah, P., Haque, A., Reid, C. R., Weber, G. F., and Garnier, S. (2019). Information Transfer During Food Choice in the Slime Mold *Physarum polycephalum*. *Frontiers in Ecology and Evolution*, 7:67.
- Sinapayen, L. (2023). Self-replication, spontaneous mutations, and exponential genetic drift in neural cellular automata. In *Artificial Life 2023*. MIT Press.
- Stanley, K. O., D'Ambrosio, D. B., and Gauci, J. (2009). A Hypercube-Based Encoding for Evolving Large-Scale Neural Networks. *Artificial Life*, 15(2):185–212. Conference Name: Artificial Life.
- Stanley, K. O. and Miikkulainen, R. (2002). Evolving Neural Networks through Augmenting Topologies. *Evolutionary Computation*, 10(2):99–127.
- Stovold, J. (2023). Neural Cellular Automata Can Respond to Signals. In *The 2023 Conference on Artificial Life*. arXiv:2305.12971 [cs].
- Vallverdu, J., Castro, O., Mayne, R., Talanov, M., Levin, M., Baluska, F., Gunji, Y., Dussutour, A., Zenil, H., and Adamatzky, A. (2017). Slime mould: the fundamental mechanisms of cognition. arXiv:1712.00414 [cs].
- Vogel, D. and Dussutour, A. (2016). Direct transfer of learned behaviour via cell fusion in non-neural organisms. *Proceedings of the Royal Society B: Biological Sciences*, 283(1845):20162382. Publisher: Royal Society.