

# FINDING PURE NASH EQUILIBRIA IN LARGE RANDOM GAMES

ANDREA COLLEVECCHIO, TUAN-MINH NGUYEN, AND ZIWEN ZHONG

**ABSTRACT.** Best Response Dynamics (BRD) is a class of strategy updating rules to find Pure Nash Equilibria (PNE) in a game. At each step, a player is randomly picked and they switches to a “best response” strategy based on the strategies chosen by others, so that the new strategy profile maximises their payoff. If no such strategy exists, a different player will be chosen randomly. When no player wants to change their strategy anymore, the process reaches a PNE and will not deviate from it. On the other hand, either PNE could not exist, or BRD could be “trapped” within a subgame that has no PNE. We prove that BRD typically converges to PNE when the game has  $N$  players, each having two actions. Our results are more general and are described as follows. We study a class of random walks in a random medium on the  $N$ -dimensional hypercube. The medium is determined by a random game with  $N$  players, each with two actions available, and i.i.d. payoffs. The medium contains obstacles that can be of two types. The first type is composed of the PNE of the game, while the other obstacles are known in the literature as traps or sink equilibria. The class of processes we analyze includes BRD, simple random walks (SRW) on the hypercube, and many other nearest neighbour processes. We prove that, with high probability, these processes reach a PNE before hitting any trap.

## CONTENTS

1. Introduction.	2
1.1. Motivation	2
1.2. Further background	4
1.3. Basic concepts from Game Theory	5
1.4. Random games	6
1.5. Main result	7
1.6. Strategies of the proof	8
2. Randomly oriented hypercube and bond percolation	8
3. Proof of Theorem 1.3	11
3.1. Strong construction of $\mathbf{X}^{(N)}$	11
3.2. Absorbing states clustering together	11
Appendix.	17
Acknowledgement	17
References	18

---

2020 *Mathematics Subject Classification.* 91A10, 91A06, 60K35, 60K37.

*Key words and phrases.* random games, pure Nash equilibria, best-response dynamics, percolation, random walks on an oriented hypercube.

## 1. INTRODUCTION.

1.1. **Motivation.** Understanding the “typical” structure of games is paramount to designing algorithms that solve games efficiently, i.e. find Mixed Nash Equilibria (MNE) in a reasonable time. In this context, an NE is a profile in the set of randomized strategies such that no player, considered individually, has an incentive to change strategy. A class of MNE is formed by the Pure Nash Equilibria (PNE), which is on the space of strategies rather than probability measures of strategies. Whereas a finite game always exhibits MNE, they might contain no PNE. The reader can think of the familiar Rock-Paper-Scissors game. Although the definitions and properties of mixed strategies and mixed equilibria are clear, the concept of PNE is “more natural”. It is worth mentioning that Osborne and Rubinstein discussed the interpretation of mixed equilibria in Section 3.2 of [17] with two different paragraphs individually signed by the two authors as they could not reach an agreement. Generally, PNE exhibit a stronger epistemic foundation than mixed equilibria. As mentioned earlier, the main issue of PNE is their existence. In order to understand the typical behaviour of a game, it is natural to consider what happens in the stochastic case, where the payoffs are random. We consider the simplest model in which players’ payoffs are identically independent distributed. It is natural to address the following questions.

- *How many PNE does a random game with  $N$  players, each having two actions, typically have?* Answers to this question were given in the literature as follows.


- (1) [20, 14, 24, 21, 22] focused on the case when the payoffs are atomless, i.e. the probability that two payoffs are equal, which we denote by  $\alpha$ , is equal to zero. In this case the number of PNE, which we denote by  $\#PNEs$  converges to a Poisson with mean one.
- (2) [3] studied the case where ties are allowed, i.e.  $\alpha \in (0, 1)$ . In this case

$$\frac{(\#PNEs) - (1 + \alpha)^N}{(1 + \alpha)^{N/2}}$$


converges in distribution to a standard normal. This is used to prove that  $\#PNEs$  is concentrated around  $(1 + \alpha)^N$ . For example, consider a win-loose game, where the payoffs are i.i.d. Bernoulli(1/2), with 100 players. In this case  $\alpha = 1/2$ , and the number of PNE is “very close” to  $(1.5)^{100}$ .

- *How to find PNE?* A natural way is to devise iterative procedures that converge to a PNE. For instance, some adaptive procedures start from a strategy profile and allow a single player (picked *at random*) to choose a different strategy. In the case of Best Response Dynamics (BRD), introduced by Gilboa and Matsui [6, 12], the selected player myopically switches to a “best response” action to the actions chosen by other players. The new strategy profile corresponding to this best response gives the highest possible payoff to the player. If no such strategy exists, the player will not move, and a different player will be chosen at random. The process is either absorbed by PNE, if they exist, or could get stuck on some subgraphs called ‘traps’ or ‘sink equilibria’ (see e.g. [7]). The following results for random games with  $N$  players, each having two actions, were proved in [3].

- i. If  $\alpha = 0$ , then BRD converges to a PNE conditionally on the existence of the latter, which holds with probability  $1 - e^{-1} + o(1)$ .



(A) When  $\alpha = 0$  (no ties),  
 $\#\text{PNEs} \xrightarrow{d} \text{Poisson}(1)$ .



(B) When  $\alpha \in (0, 1)$ ,  
 $(1 + \alpha)^{-N/2} (\#\text{PNEs} - (1 + \alpha)^N) \xrightarrow{d} \mathcal{N}(0, 1)$ .

FIGURE 1.

- ii. If  $0 < \alpha < 2^{3/4} - 1 \approx 0.6818$ , then the BRD converges to a PNE with high probability, as shown in [3], as there are many PNE, as mentioned above, and there are no traps, i.e. subgraphs that prevent BRD from reaching PNE, with very high probability.

For a formal definition of PNE, traps and BRD see Section 1.3.




FIGURE 2. Iterations needed for Best Response Dynamics (red) and Simple Random Walk (blue) to reach a PNE for  $N = 15$  and  $\alpha \in \{0.5, 0.6, 0.7, 0.8, 0.9\}$ , with 500 trials. The boxes and the whiskers are corresponding to  $(0.25, 0.75)$  and  $(0.05, 0.95)$  quantile intervals respectively.

The goal of this paper to fill the gap, and study in detail the case where there is a competition between PNE and traps. We prove that the process still converges to PNE, for any  $\alpha \in (0, 1)$ . Our result (see Theorem 1.3 below) holds for a large class of nearest neighbor processes. In the proof, we use a coupling between random games and percolation on hypercubes established in [3] to describe the geometrical structure of the model.

The study of random subgraphs of a given graph has been an intensively active field for over half a century, lying at the intersection of probability theory, statistical physics, combinatorics and computer science. In the simplest setting, each edge of the graph is considered independently, retained with probability  $p$ , and otherwise deleted. One classical example is the Erdős-Rényi random graph, and another is bond percolation on  $\mathbb{Z}^d$  (see [10]). The term percolation originates from a physical interpretation, asking whether the resulting subgraph is sufficiently connected that water can percolate through it. For this reason, regardless of the form of the original graph, random subgraph models are often referred to as percolation models. A key feature of such models is the existence of a critical threshold  $p = p_c$ , such that for  $p > p_c$  a giant connected component (or cluster) emerges.

**1.2. Further background.** The probability that a random two-person game contains a PNE was first studied for zero-sum games in [9] and for games with i.i.d. payoffs in [8]. The result in [8] was later extended to  $N$ -person games with i.i.d. payoffs in [5] and [18]. Moreover, it was shown that if the distribution of the payoffs does not admit atoms, then the number of PNEs is asymptotically Poisson(1) when the dimension of the game grows (see, for example, [19, 21, 20]). Much more difficult to study is the case where the distribution admits atoms. This case was studied in [3], where the authors obtained a central limit theorem for the number of PNEs, with a description of the rate of convergence. Other models with a large number of PNEs related to vector payoffs and common payoffs were investigated in [23] and [24]. It is worth mentioning that the framework of games with random payoffs was extended to graphical games in [4], in which a graph is defined whose vertices represent the players of the game and edges correspond to the strategic interaction between players. The payoff of vertex  $v$  depends only on the strategies of  $v$  and its neighbours. The authors in [4] investigated how the structure of the graph affects the existence of PNEs and studied the convergences to Poisson(1) of the number of PNEs. The relation between the one-shot deviation property and the stochastic stability of random games having a PNE was established in [16]. Recently, [1] studied the connection between the dominance solvability of random games and the iterated elimination of pure-strategy strictly dominated actions required for the convergence to PNE.

As for the adaptive processes on random games, the behaviour of BRD and better response dynamics (bRD) on games with i.i.d. atomless payoffs, two players, and a large number of actions was studied in [2]. In particular, the authors in [2] established that BRD does not converge to PNE, whereas bRD does whenever at least one PNE exists. More precisely, BRD is absorbed in a trap of length  $\sqrt{N}$ , where  $N$  is the number of actions, with high probability, for  $N$  large enough. Additionally, the behaviour of BRD in random games with correlated payoffs was studied in [15]. The authors in [15] described a phase transition in the convergence of BRD, in terms of the correlation parameter of the payoffs.

The case of  $N$  players each with two actions available was treated in [3], where the authors proved that BRD converges whenever  $\alpha \in (0, 2^{3/4} - 1)$  where  $\alpha$  is the probability that two payoffs coincide. In the present paper we show that BRD reaches a PNE before hitting any trap for all  $\alpha \in (0, 1)$  and for all large enough games, i.e., when  $N \rightarrow \infty$ . Our result holds not only for BRD but also for a large class of processes.

**1.3. Basic concepts from Game Theory.** A game  $\Gamma$  is a triplet  $(\mathcal{P}, (S_i)_{i \in \mathcal{P}}, (g_i)_{i \in \mathcal{P}})$ , where  $\mathcal{P}$  is the set of players,  $S_i$  is the set of strategies of each player  $i \in \mathcal{P}$  and  $g_i: S_i \rightarrow \mathbb{R}$  be the payoff function of player  $i$ .

Let  $N$  be a natural number which is not smaller than 2. Set  $[N] := \{1, 2, \dots, N\}$ . We assume throughout this paper that  $\Gamma = \Gamma_N$  is a game with  $N$  players and each player has exactly two strategies. More specifically, we set  $\mathcal{P} = [N]$  and  $S_i = \{0, 1\}$  for each  $i \in [N]$ .

Set  $\mathcal{V}_N := S_1 \times S_2 \times \dots \times S_N = \{0, 1\}^N$ . An element  $u \in \mathcal{V}_N$  is called strategy profile. For each  $i \in [N]$ , denote by  $u^{(i)}$  the strategy profile that differs from  $u$  exactly in the action of the  $i$ -th player. A pure Nash equilibrium in a normal form game is a profile of strategies, one for each player, such that, given the choice of the other players, no player has an incentive to make a different choice. In other words, deviations from equilibrium are not profitable for any player.

**Definition 1.1.** A strategy profile  $u \in \mathcal{V}_N$  is called Pure Nash Equilibrium (PNE) of the game  $\Gamma_N$  if

$$g_i(u) \geq g_i(u^{(i)}) \text{ for all } i \in [N].$$

An example with two players is given in Table 1. In this example, the strategy profile  $(0, 0)$  is a PNE. We can identify the game's strategies as the vertices of a hypercube  $\mathcal{H}_N = (\mathcal{V}_N, \mathcal{E}_N)$  where an edge connects two vertices if they differ in exactly one coordinate. Define the *Hamming distance*  $d: \mathcal{V}_N \times \mathcal{V}_N \mapsto \{0, 1, \dots, N\}$  as

$$d(u, v) := \text{card}\{i \in [N]: u_i \neq v_i\},$$

in which for a set  $A$ , we denote by  $\text{card}(A)$  the cardinality of  $A$ . We write  $u \sim v$  to denote that  $u$  and  $v$  are *neighbours* in  $\mathcal{H}_N$ , i.e., are connected by an edge which is equivalent to having Hamming distance one. A subset of vertices  $W \subset \mathcal{V}_N$  is *connected* if for any distinct vertices  $u, v \in W$  there is a path  $(x_0, x_1, \dots, x_k)$  such that  $x_0 = u$ ,  $x_k = v$  and  $\{x_{i-1}, x_i\} \in \mathcal{E}_N$  for all  $i \in [k]$ .

	P2		
		0	1
P1			
	0	(0.322, 0.412)	(0.469, 0.233)
	1	(0.214, 0.878)	(0.202, 0.311)

TABLE 1. Representation of a game  $\Gamma_2$  on  $\{0, 1\}^2$ .




FIGURE 3. Representation of  $\Gamma_3$  on  $\{0, 1\}^3$ .

The vertex  $(0, 0, \dots, 0) \in \mathcal{V}_N$  is denoted by  $\mathbf{0}$ . In order to identify PNEs, it is sufficient to know the ranking among the payoffs and not their specific values. This suggests that it is enough to describe the game using a partially oriented hypercube  $\vec{H}(\Gamma_N)$ , as follows: to each pair of neighbors  $u, u^{(i)}$  is associated either an oriented edge pointing from  $u$  to  $u^{(i)}$  if  $g_i(u) < g_i(u^{(i)})$ , denoted by  $(u, u^{(i)})$ , or an unoriented edge if  $g_i(u) = g_i(u^{(i)})$ . In this context, a PNE is a vertex  $u$  of  $\vec{H}(\Gamma_N)$  which is incident only to edges that are either unoriented or pointing towards  $u$ .

For an oriented subgraph  $\vec{G}$  of  $\vec{H}(\Gamma_N)$ , we denote by  $\mathcal{V}_N(\vec{G})$  and  $\vec{\mathcal{E}}_N(\vec{G})$  the sets of vertices and oriented edges of  $\vec{G}$  respectively. For a pair of distinct vertices  $u, v \in \mathcal{V}_N(\vec{G})$ , we say  $v$  is *accessible* from  $u$  in  $\vec{G}$  if there is an oriented path  $(x_0, x_1, \dots, x_k)$  such that  $x_0 = u$ ,  $x_k = v$  and  $(x_{i-1}, x_i) \in \vec{\mathcal{E}}_N(\vec{G})$  for all  $i \in [k]$ . We say  $\vec{G}$  is *strongly connected* if every vertex is accessible from every other vertex in that graph.

A *trap* is a strongly connected induced oriented subgraph  $\vec{T}$  of  $\vec{H}(\Gamma_N)$  with four or more vertices, such that, for all  $u \in \mathcal{V}_N(\vec{T})$  and all  $v \notin \mathcal{V}_N(\vec{T})$ , we have that  $v$  is not accessible from  $u$  in  $\vec{H}(\Gamma_N)$ . As  $\mathcal{H}_N$  is a bipartite graph, the smallest cycle of the graph are squares, i.e., cycles of length four. Fig. 4 shows an example of a game with four players. The subgraph displayed in Fig. 4(A) is a trap of size six, i.e., the cardinality of its vertex set is six.




FIGURE 4.

We define a *Best Response Dynamic* (BRD) process as follows. It is a Markov chain on the oriented hypercube  $\vec{H}(\Gamma_N)$ . It starts from  $\mathbf{0}$  and jumps to the nearest neighbours until reaching a PNE. At each step, the process chooses an edge uniformly at random among the ones incident to the present position and pointing out from it, then the process moves towards that direction. The process stops when it reaches a PNE. To see an example, the reader can refer to Fig. 3. The red dots are PNEs of the game. The process starts at  $\mathbf{0}$ , and it follows one of the two blue edges, which share the same probability of being chosen. This is equivalent to picking a player randomly and asking if they want to change their strategy, upon which they agree only if they would receive a better payoff.

**1.4. Random games.** We aim to study random games with  $N$  players, each having two actions, an i.i.d. payoffs. As mentioned above, our goal is to prove that a large class of random walks in random media on the hypercube  $\mathcal{H}_N$ . These processes, which include BRD, converge to a PNE with a probability quickly converging to 1. Then we appeal to the first Borel-Cantelli lemma to show that this convergence holds for all large  $N$ .

Consider a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ , on which the following sequence of random games is defined. Let  $\Xi_N$  be a game with  $N$  players, each with two strategies available, and random i.i.d. payoffs. Formally, for each  $u \in \mathcal{V}_N$ , the payoff  $g_i(u)$  is the realization of a random variable  $Z_i^u$ , and the random variables  $(Z_i^u)_{i \in [N], u \in \mathcal{V}_N}$  are i.i.d.. We use  $Z$  to denote a generic independent copy of  $Z_i^u$  and let  $\alpha := \mathbb{P}(Z_1 = Z_2)$  and  $\beta := (1 - \alpha)/2$ , where  $Z_1$  and  $Z_2$  are i.i.d. copies of  $Z$ .

We associate to the random game  $\Xi_N$  a random partially oriented graph  $\vec{\mathcal{H}}_N^\beta = (\mathcal{V}_N, \vec{\mathcal{E}}_N)$  in the way described in the previous section. In particular,  $(u, u^{(i)}) \in \vec{\mathcal{E}}_N$  if and only if  $Z_i^u < Z_i^{u^{(i)}}$ . The choices of orientation for each edge in  $\vec{\mathcal{E}}_N$ , uniquely determined by a distinct pair of random variables in  $(Z_i^u)_{i \in [N], u \in \mathcal{V}_N}$ , are independent to each other. We provide a proof to the edge-independent property of  $\vec{\mathcal{H}}_N^\beta$  in the Appendix, [Lemma 3.7](#). The set of oriented edges  $\vec{\mathcal{E}}_N$  can be seen as a random medium induced by  $\Xi_N$  on the unoriented hypercube  $\mathcal{H}_N$ .

We denote by  $\mathbb{Z}_+$  and  $\mathbb{N} = \mathbb{Z}_+ \setminus \{0\}$  respectively the set of non-negative integers and the set of natural numbers. Let  $\mathbf{X}^{(N)} = (X_t^{(N)})_{t \in \mathbb{Z}_+}$  be a discrete-time random walk in the random medium induced by  $\Xi_N$ , in which its distribution is defined as follows. The process takes values in  $\mathcal{V}_N$ . For each fixed outcome  $\vec{H}$  of  $\vec{\mathcal{H}}_N^\beta$ , we denote by  $P_{\vec{H}}$  the conditional probability measure corresponding to  $\mathbb{P}$  on the event  $\{\vec{\mathcal{H}}_N^\beta = \vec{H}\}$ . The distribution of  $\mathbf{X}^{(N)}$  under  $P_{\vec{H}}$  is called the *quenched law*. Let  $\mathfrak{H}_N$  be the set of all possible outcomes of  $\vec{\mathcal{H}}_N^\beta$ . The distribution of  $\mathbf{X}^{(N)}$ , which is called the *annealed law*, is given by

$$(1.1) \quad \mathbb{P}(\mathbf{X}^{(N)} \in A) = \sum_{\vec{H} \in \mathfrak{H}_N} P_{\vec{H}}(\mathbf{X}^{(N)} \in A) \mathbb{P}(\vec{\mathcal{H}}_N^\beta = \vec{H})$$

for each Borel measurable subset  $A$  of  $\mathcal{V}_N^{\mathbb{N}}$ .

**1.5. Main result.** From now on, we assume the quenched law of  $\mathbf{X}^{(N)}$  satisfies the following conditions.

**Assumption 1.2.** For any fixed outcome  $\vec{H}$  of  $\vec{\mathcal{H}}_N^\beta$ , we assume that under  $P_{\vec{H}}$ , the process  $\mathbf{X}^{(N)}$  is a time-homogeneous Markov chain taking values on  $\mathcal{V}_N$  such that  $P_{\vec{H}}(X_0^{(N)} = \mathbf{0}) = 1$  and its transition probabilities satisfy the following conditions. If  $x$  is a PNE in  $\vec{H}$ , then  $P_{\vec{H}}(X_t^{(N)} = x \mid X_{t-1}^{(N)} = x) = 1$ . Otherwise,  $P_{\vec{H}}(X_t^{(N)} \sim X_{t-1}^{(N)} \mid X_{t-1}^{(N)} = x) = 1$  and for each  $y \sim x$ ,

$$P_{\vec{H}}(X_t^{(N)} = y \mid X_{t-1}^{(N)} = x) \geq \kappa_1 N^{-\kappa_2} \delta_{(x,y) \in \vec{H}},$$

where  $\delta$  is the usual Dirac mass,  $\kappa_1$  and  $\kappa_2$  are positive constants which do not depend on  $N$ ,  $x, y$  and  $\vec{H}$ .

The class of processes described in [Assumption 1.2](#) contains BRD, and both simple random walk (SRW) on  $\mathcal{H}_N$  and SRW on the connected component of  $\vec{\mathcal{H}}_N^\beta$  containing  $\mathbf{0}$ .

Define  $\text{PNEs}_N$  to be the set of PNEs in  $\vec{\mathcal{H}}_N^\beta$ . Let

$$\tau^{(N)} := \inf\{t: X_t^{(N)} \in \text{PNEs}_N\}$$

be the absorbing time of the process  $\mathbf{X}^{(N)}$  to a PNE. Let

$$\xi^{(N)} := \inf\{t: X_t^{(N)} \in \mathcal{V}_N(\vec{\mathcal{F}}) \text{ for some trap } \vec{\mathcal{F}} \text{ of } \vec{\mathcal{H}}_N^\beta\}.$$



be the first time the process  $\mathbf{X}^{(N)}$  hits a trap. We note that these random times are not necessarily stopping times with respect to the natural filtration of the process.

We say that a sequence of events  $(A_N)_N$  holds *with small probability* (w.s.p.) if  $\sum_N \mathbb{P}(A_N) < \infty$ . We say that  $(A_N)_N$  holds *with high probability* (w.h.p.) if  $(A_N^c)_N$  holds w.s.p..

**Theorem 1.3.** *Assume that  $\alpha \in (0, 1)$  i.e.  $\beta \in (0, 1/2)$ . Let  $(\mathbf{X}^{(N)})_N$  be a sequence of processes such that for each  $N$ , the distribution of  $\mathbf{X}^{(N)}$  is given by (1.1) and its quenched law satisfies [Assumption 1.2](#). Then the sequence of events  $(\{\tau^{(N)} < \xi^{(N)}\})_{N \geq 2}$  holds w.h.p., yielding that for large enough  $N$ , the process  $\mathbf{X}^{(N)}$  eventually hits a PNE, almost surely.*

*Remark 1.1.* [Theorem 1.3](#) generalises the result in [\[3\]](#), where the authors proved that BRD converges to a PNE for  $0 < \alpha < 2^{3/4} - 1 \approx 0.68$ , by using different tools. With such a parameter  $\alpha$ , it is very likely to have at least one PNE and no traps, which in turn implies that with high probability, the best response dynamics converges to a PNE. In contrast, it was shown in [\[11\]](#) that if at each step the players are not chosen at random, but they follow a fixed deterministic cyclic order, then the process, called ‘clockwork’ BRD, does not converge to a PNE for either a large number of players or a large number of strategies. Moreover, [\[2\]](#) studies a two-player game with  $N$  actions, and it was there proved that it is unlikely for BRD to reach a PNE.

**1.6. Strategies of the proof.** In [Section 2](#), we couple the random oriented hypercube  $\vec{\mathcal{H}}_N^\beta$  with an i.i.d. bond percolation on  $\mathcal{H}_N$ , in which the geometry of the latter is well-described in [\[13\]](#). The coupling hints that even though there are many traps, the PNEs outnumber them and they are all, roughly, uniformly distributed on the hypercube. Hence, the process  $\mathbf{X}^{(N)}$  is very likely to hit a PNE rather than a trap.

The proof of [Theorem 1.3](#) is presented in [Section 3](#). Our strategy can be roughly described as follows. We cover the set of absorbing states such that each group of traps and PNEs which are close to each other within a certain distance will be included in a cover set called cluster. We show in [Lemma 3.3](#) that the process  $\mathbf{X}^{(N)}$  can visit a large number of clusters satisfying a ‘regular’ property. We next demonstrate in [Lemma 3.5](#) that with high probability these clusters contain PNEs only. We then establish in [Lemma 3.6](#) a lower bound for the probability that upon a visit to the boundary of a regular cluster, the process will reach a PNE inside this cluster. The reciprocal of this lower bound is polynomial in  $N$ , but much smaller than the number of regular clusters. Combining the above-mentioned results, we deduce that with high probability, the process eventually hits a PNE before hitting any trap.

## 2. RANDOMLY ORIENTED HYPERCUBE AND BOND PERCOLATION

Denote by  $\overleftarrow{\mathcal{H}}_N^\beta$  the oriented hypercube which has a dual orientation with respect to  $\vec{\mathcal{H}}_N^\beta$ . More precisely,  $\overleftarrow{\mathcal{H}}_N^\beta$  is obtained from  $\vec{\mathcal{H}}_N^\beta$  by inverting the directions of the oriented edges in  $\vec{\mathcal{H}}_N^\beta$  and keeping the unoriented ones still unoriented. The graph  $\overleftarrow{\mathcal{H}}_N^\beta$  is called the *dual graph* of  $\vec{\mathcal{H}}_N^\beta$ . Let  $\overleftarrow{\mathcal{L}}_N^{\beta, \mathbf{0}}$  be the collection of vertices  $u$  such that there exists an oriented path in  $\overleftarrow{\mathcal{H}}_N^\beta$  connecting  $\mathbf{0}$  to  $u$ , i.e. the set of vertices accessible from  $\mathbf{0}$  in  $\overleftarrow{\mathcal{H}}_N^\beta$ . Define  $\overleftarrow{\mathcal{M}}_N^{\beta, \mathbf{0}} := \mathcal{V}_N \setminus \overleftarrow{\mathcal{L}}_N^{\beta, \mathbf{0}}$ . In words,  $\overleftarrow{\mathcal{M}}_N^{\beta, \mathbf{0}}$  contains all the vertices that are not reachable from  $\mathbf{0}$  in  $\overleftarrow{\mathcal{H}}_N^\beta$ .



We next define a coupling between the randomly oriented hypercube describing the game and an *i.i.d.* bond percolation which was previously mentioned in [3].

**Definition 2.1.** Fix a parameter  $\beta \in (0, 1)$ . Each edge in  $\mathcal{H}_N$  is *open* with probability  $\beta$  independently of other edges, and is *closed* otherwise. The random subgraph with vertex set  $\mathcal{V}_N$ , obtained by deleting the closed edges, is called (bond) percolation with parameter  $\beta$ .

Recall that we assume  $\beta = (1 - \alpha)/2 \in (0, 1/2]$  as  $\alpha \in [0, 1)$ . Define the process

$$\mathcal{Q}_1 = \{\mathbf{0}\} \text{ and, for each } k \in \mathbb{N}, \mathcal{Q}_{k+1} = \mathcal{Q}_k \cup \{v \in \mathcal{Q}_k^c : \exists u \in \mathcal{Q}_k \text{ such that } (v, u) \in \vec{\mathcal{E}}_N\}.$$

Start with a bond percolation on  $\mathcal{H}_N$  with parameter  $\beta$  that is independent of the random variables  $(Z_i^u)_{i \in [N], u \in \mathcal{V}_N}$ . Call the resulting graph  $\mathcal{B}_1$ . For every  $k \in \mathbb{N}$  we will update  $\mathcal{B}_k$  by changing the status of some edges at each stage, in such a way that  $\mathcal{B}_{k+1}$  is still a bond percolation with parameter  $\beta$ . For each edge  $e \in \mathcal{E}_N$ , we define  $\mathcal{B}_k[e]$  the status (open or closed) of edge  $e$  in  $\mathcal{B}_k$ . We obtain  $\mathcal{B}_{k+1}$  from  $\mathcal{B}_k$ , by updating *all and only* the edges in  $\mathcal{E}_N$  that connect an element of  $\mathcal{Q}_k$  to an element of  $\partial\mathcal{Q}_k$ . More precisely, for any  $u \in \mathcal{Q}_k$  and any  $w \in \partial\mathcal{Q}_k$ , with  $u \sim w$ , set

$$\mathcal{B}_{k+1}[u, w] = \begin{cases} \text{open,} & \text{if } (w, u) \in \vec{\mathcal{E}}_N, \\ \text{closed,} & \text{otherwise;} \end{cases}$$

for all other edges  $e \in \mathcal{E}_N$ , we have  $\mathcal{B}_{k+1}[e] = \mathcal{B}_k[e]$ .

Set  $\mathcal{B}_N^\beta := \mathcal{B}_{2N}$ . We note that  $\mathcal{B}_N^\beta$  is not necessarily connected. Let  $\mathcal{L}_N^{\beta, \mathbf{0}}$  be the set of vertices in the connected component of the percolation graph  $\mathcal{B}_N^\beta$  that contains  $\mathbf{0}$ . By construction, we notice that  $\mathcal{Q}_{2N}$  is exactly equal to  $\mathcal{L}_N^{\beta, \mathbf{0}}$  and that

$$(2.1) \quad \mathcal{L}_N^{\beta, \mathbf{0}} = \overleftarrow{\mathcal{L}}_N^{\beta, \mathbf{0}}.$$

The coupling with percolation mentioned above allows us to give a detailed description of the geometry of PNEs. Denote by  $\mathcal{L}_N^\beta$  the set of vertices in the largest connected component of the percolation  $\mathcal{B}_N^\beta$ . The following result strengthens the statement of Proposition 2 in [3].

**Lemma 2.2.** *Assume that  $\alpha \in [0, 1)$ , i.e.  $\beta \in (0, 1/2]$ . There exists a constant  $\theta \in (0, 1)$  such that*

$$\mathbb{P}(\mathcal{L}_N^\beta \neq \overleftarrow{\mathcal{L}}_N^{\beta, \mathbf{0}}) \leq \theta^N.$$

*Proof.* Recall from (2.1) that  $\mathcal{L}_N^{\beta, \mathbf{0}} = \overleftarrow{\mathcal{L}}_N^{\beta, \mathbf{0}}$ . Hence, it is sufficient to show that there exists a constant  $\theta \in (0, 1)$  such that

$$(2.2) \quad \mathbb{P}(\mathcal{L}_N^\beta \neq \mathcal{L}_N^{\beta, \mathbf{0}}) \leq \theta^N.$$

We first consider the case  $\beta \in (0, 1/2)$ . Let  $\mathcal{M}_N^\beta := \mathcal{V}_N \setminus \mathcal{L}_N^\beta$ , the *fragment of the percolation*  $\mathcal{B}_N^\beta$ . Fix  $\epsilon > 0$  and set

$$\zeta_N := \mathbb{E} \left[ \text{card}(\mathcal{M}_N^\beta) \right] + \epsilon \sqrt{N(2(1 - \beta))^N}.$$

From the proof of Lemma 1 in [3], we have

$$(2.3) \quad \mathbb{P}(\mathcal{L}_N^\beta \neq \mathcal{L}_N^{\beta, \mathbf{0}}) \leq \frac{\zeta_N}{2^N} + \mathbb{P}\left(\text{card}\left(\mathcal{M}_N^\beta\right) \geq \zeta_N\right).$$

Furthermore, in virtue of Theorem 3 and inequality (2) in [13], we have that for  $\beta \in (0, 1/2)$ ,

$$(2.4) \quad \mathbb{E}\left[\text{card}\left(\mathcal{M}_N^\beta\right)\right] = (2(1-\beta))^N(1 + O(N(1-\beta)^N)) \text{ and } \mathbb{P}\left(\text{card}\left(\mathcal{M}_N^\beta\right) \geq \zeta_N\right) \leq \theta^N,$$

for some  $\theta \in (0, 1)$  which depends only on  $\epsilon$  and  $\beta$ . Combining (2.3) and (2.4), we infer that (2.2) holds for all  $\beta \in (0, 1/2)$ . Similarly as in the proof of Proposition 2 in [3], the case  $\beta = 1/2$  is proved by using the fact that  $\mathcal{L}_N^\beta$  is monotone in  $\beta$ .  $\square$

For  $r \in \mathbb{N}$  and  $v \in \mathcal{V}_N$ , set  $\text{Ball}_r(v) = \{u \in \mathcal{V}_N : d(u, v) = r\}$ , where we recall that  $d$  is the Hamming distant on  $\mathcal{H}_N$ . Let

$$m_\beta := \lfloor (-\log_2(1-\beta))^{-1} \rfloor.$$

The following result follows from [13, Proof of Theorem 2(a)] (see also Remark 5 in [3]), with a slight adjustment which is a direct consequence of Lemma 2.2.

**Lemma 2.3.** *Fix  $\beta \in (0, 1/2]$ . There exist  $\delta > 0$  and  $\theta \in (0, 1)$  such that*

$$\mathbb{P}\left(\exists v : \text{card}\left(\text{Ball}_{\lceil \delta N \rceil}(v) \setminus \overleftarrow{\mathcal{L}}_N^{\beta, \mathbf{0}}\right) > m_\beta\right) \leq \theta^N.$$

Define

$$\text{Abso}_N := \{\mathcal{U} \subseteq \mathcal{V}_N : \text{either } \mathcal{U} = \{v\} \text{ for some } v \in \text{PNEs}_N \text{ or } \mathcal{U} = \mathcal{V}_N(\vec{\mathcal{H}})\}$$

$$\text{for some trap } \vec{\mathcal{H}} \text{ of } \vec{\mathcal{H}}_N^\beta\},$$

$$\text{Abso}_N^* := \{\mathcal{U} \subseteq \mathcal{V}_N : \mathcal{U} \in \text{Abso}_N \text{ and } \text{card}(\mathcal{U}) \leq m_\beta\} \quad \text{and} \quad \text{Trap}_N^* := \text{Abso}_N^* \setminus \text{PNEs}_N.$$

**Lemma 2.4.** *Assume that  $\alpha \in (0, 1)$ , there exists a constant  $\theta \in (0, 1)$  such that*

$$\mathbb{P}\left(\exists \mathcal{U} \in \text{Abso}_N : \mathcal{U} \cap \overleftarrow{\mathcal{L}}_N^{\beta, \mathbf{0}} \neq \emptyset\right) \leq \theta^N.$$

*Proof.* Assume that there exists  $\mathcal{U} \in \text{Abso}_N$  such that  $\mathcal{U} \cap \overleftarrow{\mathcal{L}}_N^{\beta, \mathbf{0}} \neq \emptyset$ . Each edge in  $\vec{\mathcal{H}}_N^\beta$  connecting a vertex of  $\mathcal{U}$  to one in  $\mathcal{U}^c$  is either unoriented or pointing towards  $\mathcal{U}$ . As  $\mathcal{U} \cap \overleftarrow{\mathcal{L}}_N^{\beta, \mathbf{0}} \neq \emptyset$ , the vertex  $\mathbf{0}$  is accessible in  $\vec{\mathcal{H}}_N^\beta$  from a vertex in  $\mathcal{U}$ . Hence we must have that  $\mathbf{0} \in \mathcal{U}$ .

To complete the proof, it is sufficient to show that there exists a constant  $\theta \in (0, 1)$  such that

$$(2.5) \quad \mathbb{P}(\exists \mathcal{U} \in \text{Abso}_N : \mathbf{0} \in \mathcal{U}) \leq \theta^N.$$

Fix a small  $\epsilon \in (0, 1)$ . In virtue of [3, Theorem 2], we have

$$\mathbb{P}\left(\text{card}(\text{PNEs}_N) \geq (1-\epsilon)(1+\alpha)^N\right) \geq 1 - \theta^N$$

for some constant  $\theta \in (0, 1)$  which depends only on  $\epsilon$  and  $\beta$ . A PNE is said to be ‘good’ if it is incident to at least  $(\beta - \epsilon)N$  oriented edges. Using a Cramer-Chernoff bound, we notice that, with probability at least  $1 - \theta^N$ , there are at least  $(1 - 2\epsilon)(1 + \alpha)^N$  good PNEs. On the event that there exists an element  $\mathcal{U} \in \text{Abso}_N$  containing  $\mathbf{0}$ , all PNEs in  $\text{PNEs}_N \setminus \{\mathcal{U}\}$  belong to  $\mathcal{V}_N \setminus \overleftarrow{\mathcal{L}}_N^{\beta, \mathbf{0}}$ .

Hence, with probability at least  $1 - \theta^N$ , there exists a good PNE  $v$  that belongs to  $\mathcal{V}_N \setminus \overleftarrow{\mathcal{L}}_N^{\beta, \mathbf{0}}$ . Hence,  $v$  is incident at least  $(\beta - \epsilon)N$  vertices that also belong to  $\mathcal{V}_N \setminus \overleftarrow{\mathcal{L}}_N^{\beta, \mathbf{0}}$ , which in turns implies  $\text{card}(\text{Ball}_{\lceil \delta N \rceil}(v) \setminus \overleftarrow{\mathcal{L}}_N^{\beta, \mathbf{0}}) > m_\beta$ . Combining this fact with [Lemma 2.3](#), we obtain [\(2.5\)](#).  $\square$

Combining [Lemma 2.3](#) and [Lemma 2.4](#), we obtain the following result.

**Corollary 2.5.** *Fix  $\alpha \in (0, 1)$ . There exists a constant  $\theta \in (0, 1)$  such that*

$$\mathbb{P}\left(\text{Abso}_N \neq \text{Abso}_N^*\right) \leq \theta^N.$$

### 3. PROOF OF [THEOREM 1.3](#)

**3.1. Strong construction of  $\mathbf{X}^{(N)}$ .** Without loss of generality, we use the following strong construction of the process  $\mathbf{X}^{(N)}$  throughout this section.

Let  $(\chi_k^v)_{k \in \mathbb{N}, v \in \mathcal{V}_N}$  be independent exponential random variables with parameter 1, which are independent of  $(Z_i^u)_{i \in [N], u \in \mathcal{V}_N}$ . For a fixed outcome  $\vec{H}$  of  $\vec{\mathcal{H}}_N^\beta$ , let  $(p_{\vec{H}}(x, y))_{x, y \in \mathcal{V}_N}$  be a transition probability matrix such that if  $x$  is not a PNE in  $\vec{H}$  then  $\sum_{y: y \sim x} p_{\vec{H}}(x, y) = 1$  and

$$p_{\vec{H}}(x, y) \geq \kappa_1 N^{-\kappa_2} \delta_{(x, y) \in \vec{H}},$$

otherwise  $p_{\vec{H}}(x, x) = 1$ . Let  $X_0^{(N)} = \mathbf{0}$ . Assume that for some  $t \in \mathbb{N}$ ,  $X_{t-1}^{(N)} = u$  and  $\vec{\mathcal{H}}_N^\beta = \vec{H}$ . The next position of the process is given by

$$X_t^{(N)} = \text{argmin}_{v \in \mathcal{V}_N} \frac{\chi_t^v}{p_{\vec{H}}(u, v)}.$$

It is immediate that conditional on the event  $\{\vec{\mathcal{H}}_N^\beta = \vec{H}\}$ , the process  $\mathbf{X}^{(N)} = (X_t^{(N)})_{t \in \mathbb{Z}_+}$  above-constructed is a Markov chain with transition probability matrix  $(p_{\vec{H}}(x, y))_{x, y \in \mathcal{V}_N}$ . Furthermore, the annealed law of  $\mathbf{X}^{(N)}$  is given by [\(1.1\)](#) and the quenched law satisfies [Assumption 1.2](#).

**3.2. Absorbing states clustering together.** Let  $(\eta^v)_{v \in \mathcal{V}_N}$  be independent exponential random variables with parameter 1, which are also independent of  $(\xi_k^v)_{k \in \mathbb{N}, v \in \mathcal{V}_N}$  and  $(Z_i^v)_{i \in [N], v \in \mathcal{V}_N}$ .

For each element  $\mathcal{U} \in \text{Abso}_N^*$ , denote by  $\text{des}(\mathcal{U})$  one of the vertices in  $\mathcal{U}$  such that

$$\text{des}(\mathcal{U}) = \text{argmin}_{v \in \mathcal{U}} \eta^v,$$

i.e.  $\text{des}(\mathcal{U})$  is uniformly chosen among vertices in  $\mathcal{U}$ . We call  $\text{des}(\mathcal{U})$  the *designated vertex* of  $\mathcal{U}$ . Let  $\mathcal{D}$  be the set of designated vertices

$$\mathcal{D} := \{u \in \mathcal{V}_N : u = \text{des}(\mathcal{U}) \text{ for some } \mathcal{U} \in \text{Abso}_N^*\}.$$

For two distinct vertices  $u = (u_1, u_2, \dots, u_N)$  and  $v = (v_1, v_2, \dots, v_N)$  we say  $u$  is smaller than  $v$  in the *lexicographic order* if  $u_j < v_j$  for the first  $j$  where  $u_j$  and  $v_j$  differ. For each vertex  $u \in \mathcal{V}_N$ , let  $\ll_u$  be a order on  $\mathcal{V}_N$  which is defined as follows. For two distinct vertices  $x, y \in \mathcal{V}_N$ , we denote  $x \ll_u y$  if either  $d(x, u) < d(y, u)$  or  $d(x, u) = d(y, u)$  and  $x$  is smaller than  $y$  in the lexicographic order. Note that for any pair of distinct vertices  $x$  and  $y$ , we have either  $x \ll_u y$  or  $y \ll_u x$ .

Set

$$\Pi_N := \{u \in \mathcal{V}_N : d(u, \mathcal{D}) \leq m_\beta - 1\}.$$

Notice that  $\Pi_N$  is not necessarily connected but it is partitioned into connected component(s). For each vertex  $u \in \Pi_N$ , let  $V(u)$  be the connected component of  $\Pi_N$  containing  $u$ . Let  $\mathfrak{d}_1, \mathfrak{d}_2, \dots, \mathfrak{d}_j$  be the designated vertices in  $V(u)$  which are arranged in the ascending order corresponding to  $\llcorner_u$ . Set

$$\mathcal{D}^{(u)} := \{\mathfrak{d}_k : 1 \leq k \leq \min\{j, m_\beta\}\} \quad \text{and} \quad \mathcal{C}(u) := \{v \in \mathcal{V}_N : d(v, \mathcal{D}^{(u)}) \leq m_\beta - 1\}.$$

We call  $\mathcal{C}(u)$  the *cluster* of  $u$ .

**Definition 3.1.** Let  $\sigma^{(N)}$  be the first time the process  $\mathbf{X}^{(N)}$  hits a designated vertex, i.e.

$$\sigma^{(N)} := \inf\{t: X_t^{(N)} \in \mathcal{D}\}.$$

We define a new process  $\tilde{\mathbf{X}}^{(N)} = (\tilde{X}_t^{(N)})_{t \in \mathbb{Z}_+}$  as follows. We set  $\tilde{X}_t^{(N)} := X_t^{(N)}$  for  $0 \leq t \leq \sigma^{(N)}$  and at time  $t > \sigma^{(N)}$ , we set

$$\tilde{X}_t^{(N)} = \operatorname{argmin}_{u \sim \tilde{X}_{t-1}^{(N)}} \chi_t^u$$

In other words, the process moves like a simple symmetric random walk on  $\mathcal{H}$  after hitting  $\mathcal{D}$ . We call the process  $\tilde{\mathbf{X}}^{(N)}$  an *extension* of  $\mathbf{X}^{(N)}$ .

Note that  $\sigma^{(N)}$  is a stopping time w.r.t. the filtration  $(\mathcal{F}_n)_{n \in \mathbb{Z}_+}$  where  $\mathcal{F}_n$  is generated by  $(X_t)_{0 \leq t \leq n}$  and  $\mathcal{D}$ . For the sake of simplicity, we use from now on the notations  $\mathbf{X}$  and  $\tilde{\mathbf{X}}$  instead of  $\mathbf{X}^{(N)}$  and  $\tilde{\mathbf{X}}^{(N)}$ .




FIGURE 5. Every pair of designated vertices (in red) is connected with a dash line if the Hamming distance between them is less than or equal to  $2m_\beta - 2$ .

We define recursively a sequence of random times  $(S_k)_{k \in \mathbb{Z}_+}$  as follows. Set  $S_0 = 0$ , and let

$$S_k := \inf \left\{ t > S_{k-1} : \tilde{X}_t \in \Pi_N \setminus \bigcup_{j=1}^{k-1} \mathcal{C}(\tilde{X}_{S_j}) \right\}.$$

Note that  $(S_k)_{k \in \mathbb{N}}$  are stopping times w.r.t. the filtration  $(\tilde{\mathcal{F}}_n)_{n \in \mathbb{Z}_+}$  where  $\tilde{\mathcal{F}}_n$  is generated by  $(\tilde{X}_t)_{1 \leq t \leq n}$ ,  $\vec{\mathcal{H}}_N^\beta$  and  $\mathcal{D}$ .

We notice that  $\mathcal{C}(\tilde{X}_{S_k}) \neq \mathcal{C}(\tilde{X}_{S_j})$  for all  $1 \leq j \leq k-1$ . Roughly speaking, a new cluster  $\mathcal{C}(\tilde{X}_{S_k})$  is revealed by the process  $\tilde{\mathbf{X}}$  at each time  $S_k$  for  $k \in \mathbb{N}$ .

**Definition 3.2.** We say a cluster  $\mathcal{C}(u)$  is *good* if  $d(\mathcal{D} \cap \mathcal{C}(u), \mathcal{D} \setminus \mathcal{C}(u)) \geq 2m_\beta - 1$ . We define

$$\mathcal{J}_k := \{S_k < \infty\} \cap \{\mathcal{C}(\tilde{X}_{S_k}) \text{ is good}\}.$$

Note that on  $\mathcal{J}_k$ , the cluster  $\mathcal{C}(\tilde{X}_{S_k})$  is a connected component of  $\Pi_N$  containing no more than  $m_\beta$  designated vertices. Let

$$q_N := \lfloor e^{(\log N)^2} \rfloor.$$

**Lemma 3.3.** *There exists a constant  $\theta \in (0, 1)$  such that*

$$\mathbb{P}\left(\bigcap_{k=1}^{q_N} \mathcal{J}_k\right) \geq 1 - q_N \theta^N.$$

*Proof.* It sufficient to prove that there exists a constant  $\theta \in (0, 1)$  such that for  $1 \leq k \leq q_N$ , we have

$$(3.1) \quad \mathbb{P}\left(\bigcap_{i=1}^{k-1} \mathcal{J}_i \cap \mathcal{J}_k^c\right) \leq \theta^N.$$

By [Definition 3.1](#), we notice that the process  $(\tilde{X}_t)$  can not be absorbed by any element in  $\text{Abso}_N^*$ . On the event  $\bigcap_{i=1}^{k-1} \mathcal{J}_i \cap \{S_k = \infty\}$  we have either the process is absorbed by an element  $\mathcal{U} \in \text{Abso}_N \setminus \text{Abso}_N^*$  or  $\mathcal{D} \subseteq \bigcup_{i=1}^{k-1} \mathcal{C}(\tilde{X}_{S_i})$ . By reason of [Corollary 2.5](#), the former case occurs with probability not larger than  $\theta^N$  for some constant  $\theta \in (0, 1)$ . In the latter case, the number of PNEs is not larger than  $4q_N m_\beta^2$ . By reason of [\[3, Theorem 2\]](#), this case occurs with probability not larger than  $\theta^N$ . Hence

$$(3.2) \quad \mathbb{P}\left(\bigcap_{i=1}^{k-1} \mathcal{J}_i \cap \{S_k = \infty\}\right) \leq \theta^N.$$

On the other hand, on the event  $\{S_k < \infty\} \cap \{\mathcal{C}(\tilde{X}_{S_k}) \text{ is not good}\}$ , there are more than  $m_\beta$  designated vertices in  $\text{Ball}_{4m_\beta^2}(\tilde{X}_{S_k})$ . Hence, by reason of [Lemma 2.3](#) and [Lemma 2.4](#), we infer that

$$(3.3) \quad \mathbb{P}(S_k < \infty, \mathcal{C}(\tilde{X}_{S_k}) \text{ is not good}) \leq \theta^N.$$

Combining [\(3.2\)](#) and [\(3.3\)](#), we obtain [\(3.1\)](#). □

We define

$$(3.4) \quad \mathcal{B}_k := \{\mathcal{D} \cap \mathcal{C}(\tilde{X}_{S_k}) \subseteq \text{PNEs}_N\}.$$

In the remaining part of the paper, we use  $\text{Const}$  to denote a generic **positive** constant that can vary in different expressions and is allowed to depend on  $\alpha$ , but does not depend on  $N$ .

**Lemma 3.4.** *Let  $\mathfrak{d}^* \subseteq \mathcal{V}_N$  be a deterministic subset of vertices such that  $1 \leq \text{card}(\mathfrak{d}^*) \leq m_\beta$ . We have*

$$\mathbb{P}(\mathfrak{d}^* \subseteq \text{PNEs}_N \mid \mathfrak{d}^* \subseteq \mathcal{D}) \geq 1 - \text{Const} \times \left(\frac{1 + \alpha}{2}\right)^{3N}.$$

*Proof.* Denote by  $\{\mathfrak{d}_1, \mathfrak{d}_2, \dots, \mathfrak{d}_j\}$  the distinct elements of  $\mathfrak{d}^*$ . We have

$$(3.5) \quad \mathbb{P}(\mathfrak{d}^* \subseteq \text{PNEs}_N) \geq \alpha^{j^2} \left(\frac{1+\alpha}{2}\right)^{jN} \geq \alpha^{m_\beta^2} \left(\frac{1+\alpha}{2}\right)^{jN},$$

where in the first inequality of (3.5), we use the following facts:

- The probability that a vertex is a PNE is equal to  $\left(\frac{1+\alpha}{2}\right)^N$ .
- Two PNEs could be adjacent, where the edge connecting them must be unoriented. The quantity  $\alpha^{j^2}$  is the lower bound for the probability that the edges connecting any  $j$  vertices are unoriented.

The last inequality of (3.5) follows from the assumption that  $1 \leq j = \text{card}(\mathfrak{d}^*) \leq m_\beta$ .

Let  $\mathbf{G}(\mathfrak{d}^*)$  be the subgraph induced by  $\vec{\mathcal{H}}_N^\beta$  on the vertex set

$$\bigcup_{\mathcal{T} \in \text{Abso}_N^* : \mathcal{T} \cap \mathfrak{d}^* \neq \emptyset} \mathcal{T}.$$

Note that an element  $\mathcal{T} \in \text{Abso}_N^*$  has no more than  $m_\beta$  vertices. As  $j = \text{card}(\mathfrak{d}^*) \leq m_\beta$ , the cardinality of all possible outcomes of  $\mathbf{G}(\mathfrak{d}^*)$  is bounded by  $N^{m_\beta^3}$ .

To complete the proof, we show that

$$(3.6) \quad \mathbb{P}(\mathfrak{d}^* \subseteq \mathcal{D}) \leq \left(\frac{1+\alpha}{2}\right)^{jN} + \text{Const} \times N^{m_\beta^3} \left(\frac{1+\alpha}{2}\right)^{(j+3)N}.$$

It is immediate that the first quantity appearing in the right-hand side of (3.6) is an upper bound for the probability that  $\mathfrak{d}^* \subseteq \text{PNEs}_N$ . Let  $\mathcal{R}$  be a possible outcome of  $\mathbf{G}(\mathfrak{d}^*)$  that contains at least one trap in  $\text{Trap}_N^*$ . We finish the proof by proving that the quantity  $\left(\frac{1+\alpha}{2}\right)^{(j+3)N}$  appearing in the right-hand side of (3.6) is an upper bound for the probability that  $\mathfrak{d}^* \subseteq \mathcal{D}$  and  $\mathbf{G}(\mathfrak{d}^*) = \mathcal{R}$ . For  $1 \leq k \leq j$ , let  $R_k$  be the PNE/Trap in  $\mathcal{R}$  containing  $\mathfrak{d}_k$ . We have

$$(3.7) \quad \mathbb{P}(\mathbf{G}(\mathfrak{d}^*) = \mathcal{R}, \mathfrak{d}^* \subseteq \mathcal{D}) \leq \text{Const} \times \prod_{k=1}^j \left(\frac{1+\alpha}{2}\right)^{\text{card}(R_k) \cdot N} \leq \text{Const} \times \left(\frac{1+\alpha}{2}\right)^{(j+3)N}.$$

where in the first inequality we use the following facts:

- Each pair  $R_i$  and  $R_k$  can be connected by at most  $|R_i| \cdot |R_k|$  unoriented edges. Thus each  $R_k$  can have at most  $\binom{j}{2} m_\beta^2$  unoriented edges connecting one of its vertices to the vertices of the remaining subgraphs  $R_i$ , with  $i \neq k$ , which gives a contribution of constant order in (3.7).
- The probability of a fixed trap of size  $r$  is bounded by  $\text{Const} \times \left(\frac{1+\alpha}{2}\right)^{rN}$ .

In the second inequality, we use the fact that  $\mathcal{R}$  contains at least one trap and thus  $\sum_{k=1}^j \text{card}(R_k) \geq j+3$ .  $\square$

Recall that  $q_N = \lfloor e^{(\log N)^2} \rfloor$ .

**Lemma 3.5.** *There exists  $\theta \in (0, 1)$  such that for all  $N$  large enough we have that*

$$\mathbb{P} \left( \bigcap_{k=1}^{q_N} \mathcal{B}_k \cap \mathcal{J}_k \right) \geq 1 - q_N \theta^N.$$

*Proof.* Fix  $k \in \{1, 2, \dots, q_N\}$ . Let  $\mathcal{Q}_k := \bigcap_{i=1}^k \mathcal{B}_i \cap \mathcal{J}_i$ . Assume that  $\mathbb{P}(\mathcal{J}_k, \mathcal{Q}_{k-1}) > 0$ . We first prove that

$$(3.8) \quad \mathbb{P}\left(\mathcal{B}_k \mid \mathcal{J}_k, \mathcal{Q}_{k-1}\right) \geq 1 - \text{Const} \times \left(\frac{1+\alpha}{2}\right)^{3N}.$$

There exists a deterministic subset  $\mathfrak{d}^* \subseteq \mathcal{V}_N$  such that

$$(3.9) \quad \mathbb{P}(\mathcal{C}(\tilde{X}_{S_k}) \cap \mathcal{D} = \mathfrak{d}^*, \mathcal{J}_k, \mathcal{Q}_{k-1}) > 0.$$

Let

$$W(\mathfrak{d}^*) := \{u \in \mathcal{V}_N : d(u, \mathfrak{d}^*) \leq m_\beta - 1\} \quad \text{and} \quad \mathcal{B}(\mathfrak{d}^*) := \{\mathfrak{d}^* \subseteq \text{PNEs}_N\}.$$

The assumption (3.9) implies that  $\text{card}(\mathfrak{d}^*) \leq m_\beta$  and  $W(\mathfrak{d}^*)$  is connected. Note that the event  $\mathcal{B}(\mathfrak{d}^*)$  adapts the definition  $\mathcal{B}_k$  to the case  $\mathcal{C}(\tilde{X}_{S_k}) \cap \mathcal{D} = \mathfrak{d}^*$ . Set

$$(3.10) \quad \mathcal{R}_k(\mathfrak{d}^*) = \left\{ d(\mathfrak{d}^*, \mathcal{D} \setminus \mathfrak{d}^*) \geq 2m_\beta - 1 \right\} \cap \left\{ S_k = \inf\{t \in \mathbb{Z}_+ : \tilde{X}_t \in W(\mathfrak{d}^*)\} < \infty \right\}.$$

Notice that

$$(3.11) \quad \{\mathcal{C}(\tilde{X}_{S_k}) \cap \mathcal{D} = \mathfrak{d}^*\} \cap \mathcal{J}_k = \{\mathfrak{d}^* \subseteq \mathcal{D}\} \cap \mathcal{R}_k(\mathfrak{d}^*).$$

We thus have

$$(3.12) \quad \begin{aligned} \mathbb{P}\left(\mathcal{B}_k \mid \mathcal{C}(\tilde{X}_{S_k}) \cap \mathcal{D} = \mathfrak{d}^*, \mathcal{J}_k, \mathcal{Q}_{k-1}\right) &= \mathbb{P}\left(\mathcal{B}(\mathfrak{d}^*) \mid \mathfrak{d}^* \subseteq \mathcal{D}, \mathcal{R}_k(\mathfrak{d}^*), \mathcal{Q}_{k-1}\right) \\ &= \mathbb{P}(\mathcal{B}(\mathfrak{d}^*) \mid \mathfrak{d}^* \subseteq \mathcal{D}). \end{aligned}$$

where in the last identity we use the independence which is justified as follows. For a subset  $W \subset \mathcal{V}_N$ , let  $W^\circ = \{x \in W : \text{if } y \sim x \text{ then } y \in W\}$  be the set of interior vertices of  $W$ .

- The event  $\mathcal{B}(\mathfrak{d}^*)$  is determined by the orientation of edges connecting vertices in  $W(\mathfrak{d}^*)^\circ$ .
- The event  $\mathcal{R}_k(\mathfrak{d}^*) \cap \mathcal{Q}_{k-1}$  is determined by  $(\eta^v)_{v \in \mathcal{V}_N \setminus W(\mathfrak{d}^*)^\circ}$ ,  $(\chi_k^v)_{k \in \mathbb{N}, v \in \mathcal{V}_N \setminus W(\mathfrak{d}^*)^\circ}$  and the orientation of edges connecting vertices in  $\mathcal{V}_N \setminus W(\mathfrak{d}^*)^\circ$ .

In virtue of [Lemma 3.4](#), we have

$$(3.13) \quad \mathbb{P}(\mathcal{B}(\mathfrak{d}^*) \mid \mathfrak{d}^* \subseteq \mathcal{D}) \geq 1 - \text{Const} \times \left(\frac{1+\alpha}{2}\right)^{3N}.$$

Combining (3.13) with (3.12), and using the law of total probabilities over  $\mathfrak{d}^*$ , we obtain (3.8).

On the other hand, using independence properties similarly as above, we notice that

$$(3.14) \quad \mathbb{P}(\mathcal{J}_k \mid \mathcal{Q}_{k-1}) = \mathbb{P}\left(\mathcal{J}_k \mid \bigcap_{i=1}^{k-1} \mathcal{J}_i\right).$$

Combining (3.8), (3.14) and [Lemma 3.3](#), we have

$$\begin{aligned} \mathbb{P}\left(\bigcap_{k=1}^{q_N} \mathcal{B}_k \cap \mathcal{J}_k\right) &= \prod_{k=1}^{q_N} \mathbb{P}(\mathcal{B}_k \mid \mathcal{J}_k, \mathcal{Q}_{k-1}) \mathbb{P}(\mathcal{J}_k \mid \mathcal{Q}_{k-1}) \\ &= \mathbb{P}\left(\bigcap_{k=1}^{q_N} \mathcal{J}_k\right) \prod_{k=1}^{q_N} \mathbb{P}(\mathcal{B}_k \mid \mathcal{J}_k, \mathcal{Q}_{k-1}) \\ &\geq 1 - q_N \theta^N \end{aligned}$$



for some constant  $\theta \in (0, 1)$ . This ends the proof of the lemma.  $\square$

Let

$$\mathcal{A}_k = \{ \tilde{X}_{S_k+m_\beta-1} \in \mathcal{C}(\tilde{X}_{S_k}) \cap \mathcal{D} \} \quad \text{and} \quad \mathcal{G}_k = \bigcap_{i=1}^k \mathcal{A}_i^c \cap \mathcal{B}_i \cap \mathcal{J}_i.$$

**Lemma 3.6.** *There exist positive constants  $K_1$  and  $K_2$  which does not depends on  $N$  such that for  $1 \leq k \leq q_N$ , we have*

$$\mathbb{P}(\mathcal{A}_k \mid \mathcal{B}_k \cap \mathcal{J}_k \cap \mathcal{G}_{k-1}) \geq K_1 N^{-K_2}.$$

*Proof.* Consider a deterministic subset  $\mathfrak{d}^* \subseteq \mathcal{V}_N$  and a deterministic vertex  $x_1$  such that  $d(x_1, \mathfrak{d}^*) = m_\beta - 1$  such that

$$\mathbb{P}(\tilde{X}_{S_k} = x_1, \mathcal{C}(\tilde{X}_{S_k}) \cap \mathcal{D} = \mathfrak{d}^*, \mathcal{B}_k, \mathcal{J}_k, \mathcal{G}_{k-1}) > 0.$$

Recall from (3.10)-(3.11) that  $\{\mathcal{C}(\tilde{X}_{S_k}) \cap \mathcal{D} = \mathfrak{d}^*\} \cap \mathcal{J}_k = \{\mathfrak{d}^* \subset \mathcal{D}\} \cap \mathcal{R}_k(\mathfrak{d}^*)$ , where

$$\mathcal{R}_k(\mathfrak{d}^*) = \left\{ d(\mathfrak{d}^*, \mathcal{D} \setminus \mathfrak{d}^*) \geq 2m_\beta - 1 \right\} \cap \left\{ S_k = \inf\{t \in \mathbb{Z}_+ : \tilde{X}_t \in W(\mathfrak{d}^*)\} < \infty \right\}.$$

Let  $x_1, x_2, \dots, x_{m_\beta-1}$  be deterministic vertices in a shortest path in  $\mathcal{H}_N$  which connects  $x_1$  and  $\mathfrak{d}^*$  with  $x_{m_\beta-1} \in \mathfrak{d}^*$ . We have

$$\begin{aligned} & \mathbb{P}\left( (x_j, x_{j+1}) \in \vec{\mathcal{E}}_N \text{ for all } 1 \leq j \leq m_\beta - 2 \mid \tilde{X}_{S_k} = x_1, \mathcal{C}(\tilde{X}_{S_k}) \cap \mathcal{D} = \mathfrak{d}^*, \mathcal{B}_k, \mathcal{J}_k, \mathcal{G}_{k-1} \right) \\ &= \mathbb{P}\left( (x_j, x_{j+1}) \in \vec{\mathcal{E}}_N \text{ for all } 1 \leq j \leq m_\beta - 2 \mid \tilde{X}_{S_k} = x_1, \mathfrak{d}^* \subseteq \text{PNEs}_N, \mathcal{R}_k(\mathfrak{d}^*), \mathcal{G}_{k-1} \right) \\ (3.15) \quad &= \mathbb{P}\left( (x_j, x_{j+1}) \in \vec{\mathcal{E}}_N \text{ for all } 1 \leq j \leq m_\beta - 2 \mid \mathfrak{d}^* \subseteq \text{PNEs}_N \right) \\ &= \beta^{m_\beta-2} \mathbb{P}\left( (x_{m_\beta-2}, x_{m_\beta-1}) \in \vec{\mathcal{E}}_N \mid x_{m_\beta-1} \in \text{PNEs}_N \right) = \frac{2}{1+\alpha} \beta^{m_\beta-1}, \end{aligned}$$

where in the second and the third identities we use the follows facts:

- The orientation of edges on the path  $(x_1, x_2, \dots, x_{m_\beta-1})$  is independent of the event  $\{\tilde{X}_{S_k} = x_1\} \cap \mathcal{R}_k(\mathfrak{d}^*) \cap \mathcal{G}_{k-1}$ .
- The orientation of edges on the path  $(x_1, x_2, \dots, x_{m_\beta-2})$  is independent of the event  $\{\mathfrak{d}^* \setminus \{x_{m_\beta-1}\} \subseteq \text{PNEs}_N\}$ .

Using (3.15) and Assumption 1.2, we obtain

$$\begin{aligned} & \mathbb{P}\left( \mathcal{A}_k \mid \tilde{X}_{S_k} = x_1, \mathcal{C}(\tilde{X}_{S_k}) \cap \mathcal{D} = \mathfrak{d}^*, \mathcal{B}_k, \mathcal{J}_k, \mathcal{G}_{k-1} \right) \\ & \geq \mathbb{P}\left( \tilde{X}_{S_k+j} = x_{j+1} \text{ for all } 1 \leq j \leq m_\beta - 2 \mid \tilde{X}_{S_k} = x_1, \mathcal{C}(\tilde{X}_{S_k}) \cap \mathcal{D} = \mathfrak{d}^*, \mathcal{B}_k, \mathcal{J}_k, \mathcal{G}_{k-1} \right) \\ & \geq \text{Const} \times N^{-\kappa_2 m_\beta}. \end{aligned}$$

The proof ends by using the law of total probabilities over  $\mathfrak{d}^*$  and  $x_1$ .  $\square$

*Proof of Theorem 1.3.* Recall that  $q_N = \lfloor e^{(\log N)^2} \rfloor$ . In virtue of Lemma 3.5, there exists  $\theta \in (0, 1)$  such that

$$(3.16) \quad \mathbb{P} \left( \bigcap_{k=1}^{q_N} \mathcal{B}_k \cap \mathcal{J}_k \right) \geq 1 - q_N \theta^N.$$

Using Lemma 3.6, we obtain that

$$(3.17) \quad \begin{aligned} \mathbb{P} \left( \bigcap_{k=1}^{q_N} \mathcal{A}_k^c \cap \mathcal{B}_k \cap \mathcal{J}_k \right) &\leq (1 - K_1 N^{-K_2})^{q_N} \\ &\leq e^{-K_1 q_N N^{-K_2}}. \end{aligned}$$

The right-hand side of (3.17) is summable. Hence  $\bigcup_{k=1}^{q_N} \mathcal{A}_k \cap \bigcap_{k=1}^{q_N} \mathcal{B}_k \cap \mathcal{J}_k$  occurs with high probability. We notice that

$$\bigcup_{k=1}^{q_N} \mathcal{A}_k \cap \bigcap_{k=1}^{q_N} \mathcal{B}_k \cap \mathcal{J}_k \subseteq \bigcup_{k=1}^{q_N} \mathcal{A}_k \cap \bigcap_i^{k-1} \mathcal{A}_i^c \cap \bigcap_i^k \mathcal{B}_i \cap \mathcal{J}_i = \bigcup_{k=1}^{q_N} \mathcal{A}_k \cap \mathcal{B}_k \cap \mathcal{J}_k \cap \mathcal{G}_{k-1},$$

where we recall that  $\mathcal{G}_k = \bigcap_{i=1}^k \mathcal{A}_i^c \cap \mathcal{B}_i \cap \mathcal{J}_i$ . It follows that with high probability there exists  $k \in [q_N]$  such that  $\mathcal{A}_k \cap \mathcal{B}_k \cap \mathcal{J}_k \cap \mathcal{G}_{k-1}$  occurs. Recall that  $\sigma^{(N)} = \inf\{t: \tilde{X}_t^{(N)} \in \mathcal{D}\}$ . Notice that on the event  $\mathcal{A}_k \cap \mathcal{B}_k \cap \mathcal{J}_k \cap \mathcal{G}_{k-1} \cap \{\text{Abso}_N = \text{Abso}_N^*\}$ , we have  $\tilde{X}_t = X_t$  for all  $0 \leq t \leq S_k + m_\beta - 1$  and  $S_k + m_\beta - 1 = \sigma^{(N)} = \tau^{(N)} < \xi^{(N)}$ . In virtue of Corollary 2.5, we note that,  $\text{Abso}_N = \text{Abso}_N^*$  with high probability. Hence  $\tau^{(N)} < \xi^{(N)}$  with high probability.  $\square$

## APPENDIX.

**Lemma 3.7.** *Each edge in  $\vec{\mathcal{H}}_N^\beta$  is either unoriented, with probability  $\alpha$ , or follows a given orientation, with probability  $\beta$ . The state of each edge is independent of the others.*

*Proof of Lemma 3.7.* Fix a subset of edges  $e_1, e_2, \dots, e_n \in \vec{\mathcal{E}}_N$ . Denote by  $u_i, v_i \in \mathcal{V}_N$  the endpoints of edge  $e_i$ , where  $v_i = u_i^{(j)}$  for some  $j \in [N]$ . The orientation of  $e_i$  is determined uniquely by the two random variables  $Z_j^{u_i}$  and  $Z_j^{v_i}$ , which are disjoint from other edges' depending random variables. It suffices to prove that the orientation of  $e_n$  is independent of the orientation of the edges  $\{e_1, e_2, \dots, e_{n-1}\}$ , and the result trivially holds as the class of random variables  $(Z_i^u)_{i \in [N], u \in \mathcal{V}_N}$  are i.i.d..  $\square$

## ACKNOWLEDGEMENT

We would like to thank Alex Scott and Marco Scarsini for very helpful discussion. The work of AC and TN was partially supported by ARC DP230102209. ZZ was supported by the Australian Government Research Training Program (RTP) Scholarship.

## REFERENCES

- [1] N. ALON, K. RUDOV, AND L. YARIV, *Dominance solvability in random games*, arXiv e-prints, (2021).
- [2] B. AMIET, A. COLLEVECCHIO, AND K. HAMZA, *When “better” is better than “best”*, *Oper. Res. Lett.*, 49 (2021), pp. 260–264.
- [3] B. AMIET, A. COLLEVECCHIO, M. SCARSINI, AND Z. ZHONG, *Pure Nash equilibria and best-response dynamics in random games*, *Math. Oper. Res.*, 46 (2021), pp. 1552–1572.
- [4] C. DASKALAKIS, A. G. DIMAKIS, AND E. MOSSEL, *Connectivity and equilibrium in random games*, *Ann. Appl. Probab.*, 21 (2011), pp. 987–1016.
- [5] M. DRESHER, *Probability of a pure equilibrium point in  $n$ -person games*, *J. Combinatorial Theory*, 8 (1970), pp. 134–145.
- [6] I. GILBOA AND A. MATSUI, *Social stability and equilibrium*, *Econometrica*, 59 (1991), pp. 859–867.
- [7] M. GOEMANS, V. MIRROKNI, AND A. VETTA, *Sink equilibria and convergence*, in 46th Annual IEEE Symposium on Foundations of Computer Science (FOCS’05), Oct 2005, pp. 142–151.
- [8] K. GOLDBERG, A. J. GOLDMAN, AND M. NEWMAN, *The probability of an equilibrium point*, *J. Res. Nat. Bur. Standards Sect. B*, 72B (1968), pp. 93–101.
- [9] A. J. GOLDMAN, *The probability of a saddlepoint*, *Amer. Math. Monthly*, 64 (1957), pp. 729–730.
- [10] G. R. GRIMMETT, H. KESTEN, AND Y. ZHANG, *Random walk on the infinite cluster of the percolation model*, *Probab. Theory Related Fields*, 96 (1993), pp. 33–44.
- [11] T. HEINRICH, Y. JANG, L. MUNGO, M. PANGALLO, A. SCOTT, B. TARBUSH, AND S. WIESE, *Best-response dynamics, playing sequences, and convergence to equilibrium in random games*, *Internat. J. Game Theory*, 52 (2023), pp. 703–735.
- [12] A. MATSUI, *Best response dynamics and socially stable strategies*, *J. Econom. Theory*, 57 (1992), pp. 343–362.
- [13] C. MCDIARMID, A. SCOTT, AND P. WITHERS, *The component structure of dense random subgraphs of the hypercube*, *Random Structures Algorithms*, 59 (2021), pp. 3–24.
- [14] A. MCLENNAN, *The expected number of Nash equilibria of a normal form game*, *Econometrica*, 73 (2005), pp. 141–174.
- [15] H. A. MIMUN, M. QUATTROPANI, AND M. SCARSINI, *Best-response dynamics in two-person random games with correlated payoffs*, *Games Econom. Behav.*, 145 (2024), pp. 239–262.
- [16] J. NEWTON AND R. SAWA, *Conventions and social choice in large games*, SSRN e-prints, (2024).
- [17] M. J. OSBORNE AND A. RUBINSTEIN, *A course in game theory*, MIT Press, Cambridge, MA, 1994.
- [18] G. P. PAPAVALOPOULOS, *On the probability of existence of pure equilibria in matrix games*, *J. Optim. Theory Appl.*, 87 (1995), pp. 419–439.
- [19] I. Y. POWERS, *Limiting distributions of the number of pure strategy Nash equilibria in  $N$ -person games*, *Internat. J. Game Theory*, 19 (1990), pp. 277–286.
- [20] Y. RINOTT AND M. SCARSINI, *On the number of pure strategy Nash equilibria in random games*, *Games Econom. Behav.*, 33 (2000), pp. 274–293.
- [21] W. STANFORD, *A note on the probability of  $k$  pure Nash equilibria in matrix games*, *Games Econom. Behav.*, 9 (1995), pp. 238–246.
- [22] ———, *The limit distribution of pure strategy Nash equilibria in symmetric bimatrix games*, *Math. Oper. Res.*, 21 (1996), pp. 726–733.
- [23] ———, *On the distribution of pure strategy equilibria in finite games with vector payoffs*, *Math. Social Sci.*, 33 (1997), pp. 115–127.
- [24] ———, *On the number of pure strategy Nash equilibria in finite common payoffs games*, *Econom. Lett.*, 62 (1999), pp. 29–34.

SCHOOL OF MATHEMATICS AND MONASH DATA FUTURES INSTITUTE, MONASH UNIVERSITY, MELBOURNE, AUSTRALIA

*Email address:* `Andrea.Collevecchio@monash.edu`

SCHOOL OF MATHEMATICS, MONASH UNIVERSITY, MELBOURNE, AUSTRALIA

*Email address:* `tuanminh.nguyen@monash.edu`

SCHOOL OF MATHEMATICS, MONASH UNIVERSITY, MELBOURNE, AUSTRALIA

*Email address:* `ziwen.zhong@monash.edu`