Deep neural networks with ReLU, leaky ReLU, and softplus activation provably overcome the curse of dimensionality for space-time solutions of semilinear partial differential equations

Julia Ackermann¹, Arnulf Jentzen^{2,3}, Benno Kuckuck⁴, and Joshua Lee Padgett⁵

¹ Department of Mathematics & Informatics, University of Wuppertal, Germany, e-mail: jackermann@uni-wuppertal.de

² School of Data Science and Shenzhen Research Institute of Big Data, The Chinese University of Hong Kong, Shenzhen (CUHK-Shenzhen), China, e-mail: ajentzen@cuhk.edu.cn

³ Applied Mathematics: Institute for Analysis and Numerics, University of Münster, Germany, e-mail: ajentzen@uni-muenster.de

⁴ Applied Mathematics: Institute for Analysis and Numerics, University of Münster, Germany, e-mail: bkuckuck@uni-muenster.de

> ⁵ Data & Analytics, Toyota Financial Services, Texas, USA, e-mail: josh.padgett@toyota.com

> > June 18, 2024

Abstract

It is a very challenging topic in applied mathematics to solve high-dimensional nonlinear partial differential equations (PDEs). Standard approximation methods for nonlinear PDEs such as finite difference and finite element methods suffer under the so-called curse of dimensionality (COD) in the sense that the number of computational operations of the numerical approximation method grows at least exponentially in the PDE dimension and with such methods it is essentially impossible to approximately solve high-dimensional PDEs even when the fastest currently available computers are used. However, in the last years great progress has been made in this area of research through suitable deep learning (DL) based methods for PDEs in which deep neural networks (DNNs) are used to approximate solutions of PDEs. Despite the remarkable success of such DL methods in numerical simulations, it remains a fundamental open problem of research to prove (or disprove) that such methods can overcome the COD in the approximation of PDEs. However, there are nowadays several partial error analysis results for DL methods for high-dimensional nonlinear PDEs in the literature which prove that DNNs can overcome the COD in the sense that the number of parameters of the approximating DNN grows at most polynomially in both the reciprocal $1/\varepsilon$ of the prescribed approximation accuracy $\varepsilon > 0$ and the PDE dimension $d \in \mathbb{N} = \{1, 2, 3, ...\}$. In the main result of this article we prove that for all $T, p \in (0, \infty)$ it holds that solutions $u_d : [0, T] \times \mathbb{R}^d \to \mathbb{R}, d \in \mathbb{N}$, of semilinear heat equations with Lipschitz continuous nonlinearities can be approximated in the L^p -sense on space-time regions without the COD by DNNs with the rectified linear unit (ReLU), the leaky ReLU, or the softplus activation function. In previous articles similar results have been established not for space-time regions but for the solutions $u_d(T, \cdot), d \in \mathbb{N}$, at the terminal time T.

Contents

| T | Intr | oduction | 2 |
|-----|---|--|--|
| 2 | Proj 2.1 2.2 | perties of solutions of partial differential equations (PDEs)Perturbation estimates for solutions of PDEsTemporal regularity estimates for solutions of PDEs | 7 7 10 |
| 3 | Arti | ficial neural network (ANN) calculus | 15 |
| | 3.1 | ANNs | 16 |
| | 3.2 | Realizations of ANNs | 16 |
| | 3.3 | Activation ANNs | 17 |
| | 3.4 | Compositions, powers, and extensions of ANNs | 18 |
| | 3.5 | Parallelizations of ANNs | 18 |
| | 3.6 | Scalar multiplications and sums of ANNs | 21 |
| 4 | ANN approximations for linear interpolations of multilevel Picard (MLP) | | |
| 4 | ANI | N approximations for linear interpolations of multilevel Picard (MLP) | |
| 4 | ANI appi | N approximations for linear interpolations of multilevel Picard (MLP) roximations | 22 |
| 4 | ANI appr 4.1 | N approximations for linear interpolations of multilevel Picard (MLP) roximations Properties of linear interpolation functions | 22 24 |
| 4 | ANI app 4.1 4.2 | N approximations for linear interpolations of multilevel Picard (MLP) roximations Properties of linear interpolation functions | 22 24 25 |
| 4 | ANI appr 4.1 4.2 4.3 | N approximations for linear interpolations of multilevel Picard (MLP) roximations Properties of linear interpolation functions | 22 24 25 27 |
| 4 | ANI appr 4.1 4.2 4.3 4.4 | N approximations for linear interpolations of multilevel Picard (MLP) roximations Properties of linear interpolation functions | 22 24 25 27 29 |
| 4 | ANI appr 4.1 4.2 4.3 4.4 4.5 | N approximations for linear interpolations of multilevel Picard (MLP) roximations Properties of linear interpolation functions | 22 24 25 27 29 32 |
| 4 | ANI app 4.1 4.2 4.3 4.4 4.5 4.6 | N approximations for linear interpolations of multilevel Picard (MLP) roximations Properties of linear interpolation functions | 22 24 25 27 29 32 35 |
| 4 5 | ANI appr 4.1 4.2 4.3 4.4 4.5 4.6 ANI | N approximations for linear interpolations of multilevel Picard (MLP) roximations Properties of linear interpolation functions | 22 24 25 27 29 32 35 37 |
| 4 5 | ANI appr 4.1 4.2 4.3 4.4 4.5 4.6 ANI 5.1 | N approximations for linear interpolations of multilevel Picard (MLP) roximations Properties of linear interpolation functions | 22 24 25 27 29 32 35 37 38 |

1 Introduction

It is a very challenging topic in applied mathematics to solve high-dimensional *partial differential equations* (PDEs). The dimensionality corresponds here to the number of dimensions/degrees of freedom of the domain set on which solutions of the PDE are defined. Classical deterministic numerical approximation methods for PDEs such as finite difference methods (see, e.g., Jovanović & Süli [57]) typically suffer from the so-called *curse of* dimensionality (COD) (cf., e.g., Bellman [13], Novak & Ritter [67], and Novak & Woźniakowski [68, Chapter 1]) in the sense that the number of computational operations of the numerical method grows at least exponentially in the PDE dimension $d \in \mathbb{N} = \{1, 2, 3, ...\}$ and with such numerical methods it is basically impossible to approximately compute solutions of even moderate high-dimensional PDEs, say, 30-dimensional PDE solutions (corresponding to d = 30).

Great progress has been made in this field of research using suitable *deep learning* (DL) based approximation methods for high-dimensional PDEs. More specifically, in about the last 7 years there have arisen a large number of articles in which suitable DL based approximation methods – involving deep artificial neural networks (ANNs) trained by stochastic gradient descent optimization methods – have been proposed and used to approximately solve high-dimensional PDEs. For example, we refer to [4, 5, 7, 17, 23, 40, 43, 59, 69, 70, 71]for DL methods which are based on stochastic representations (involving forward stochastic differential equations (SDEs) or forward backward stochastic differential equations (BSDEs)) of the PDE under consideration such as deep BSDE and deep Kolmogorov methods, we refer to [14,39,42,60,72,76] for DL methods which are based on the classical or strong formulation of the PDE under consideration such as physics-informed neural network (PINN) and deep Galerkin (DG) methods, and we refer to [3, 18, 26, 77, 80] for DL methods which are based on weak or variational formulations of the PDE under consideration. We also refer, for instance, to the survey articles Beck et al. [11], Blechschmidt & Ernst [16], Cuomo et al. [22], E et al. [24], Germain et al. [30], and Karniadakis et al. [58] and the monograph Jentzen et al. [54, Chapters 16–18] for further references and details.

Despite the remarkable success of such DL methods in numerical simulations, it remains a fundamental open problem of research to prove (or disprove) that such methods can indeed overcome the COD in the approximation of PDEs. Actually, even in the situation of onedimensional PDEs and one-dimensional abstract target functions it remains a challenging open research problem to prove (or disprove) that such methods do indeed converge (cf., e.g., [19, 28, 29, 32, 53, 55, 61, 74, 78]).

However, there are nowadays several partial error analyses for DL methods for highdimensional PDEs in the scientific literature which prove that ANNs have the fundamental capacity to overcome the COD in the sense that the number of parameters of the approximating ANN grows at most polynomially in both the reciprocal $1/\varepsilon$ of the prescribed approximation accuracy $\varepsilon > 0$ and the dimension $d \in \mathbb{N}$ of the PDE¹. Such ANN approximation results for high-dimensional PDEs have first been obtained for linear PDEs of the Kolmogorov type (see, e.g., [2, 15, 20, 27, 33, 34, 35, 36, 38, 41, 56, 73, 79]) and, thereafter, have been extended to certain classes of nonlinear PDEs (see, e.g., [1, 21, 46, 62, 63, 64]). We also refer to the survey articles Beck et al. [11, Section 6] and E et al. [24, Section 7] and the monograph Jentzen et al. [54, Section 18.4] for further reading on such ANN approximation results.

In this work we prove in Theorem 1.1 in this introductory section that for every arbitrarily large moment $p \in (0, \infty)$ and every arbitrarily large time horizon $T \in (0, \infty)$ it holds that deep ANNs with the *rectified linear unit* (ReLU), the leaky ReLU, or the softplus activation

¹This polynomial growth property in both the inverse $1/\varepsilon$ of the prescribed approximation accuracy $\varepsilon > 0$ and the PDE dimension $d \in \mathbb{N}$ is sometimes referred to as polynomial tractability in the literature (cf., e.g., Novak & Woźniakowski [68, Section 4.4.1]).

overcome the COD in the $L^p([0,T] \times [0,1]^d; \mathbb{R})$ -approximation of solutions of a class of semilinear heat PDEs with Lipschitz continuous nonlinearities (see (1.6) in Theorem 1.1 for details). Theorem 1.1 follows from the more general results in Theorem 5.3, Corollary 5.4, and Corollary 5.5 in Section 5 and in Theorem 1.1 in this introductory section we restrict ourselves to measuring the error with respect to the Lebesgue integral on the simple spacetime region $[0,T] \times [0,1]^d$ while in our more general results in Theorem 5.3, Corollary 5.4, and Corollary 5.5 we consider more general measures on more general space-time regions to measure the error between the exact solution of the PDE and its deep ANN approximation. In our preliminary article [1] we also showed such an ANN approximation result for semilinear heat PDEs but we restricted ourselves to deep ANN approximations for the PDE solution on some spatial regions (subsets of \mathbb{R}^d) evaluated at the terminal time T instead of on space-time regions as in this work. We now present the precise statement of Theorem 1.1 in a self-contained fashion in full mathematical details and, thereafter, we provide further explanatory sentences regarding the statement of Theorem 1.1.

Theorem 1.1. Let $T, \kappa, p \in (0, \infty)$, let $f : \mathbb{R} \to \mathbb{R}$ be Lipschitz continuous, for every $d \in \mathbb{N}$ let $u_d \in C^{1,2}([0,T] \times \mathbb{R}^d, \mathbb{R})$ satisfy for all $t \in [0,T]$, $x \in \mathbb{R}^d$ that

$$\frac{\partial}{\partial t}u_d(t,x) = \Delta_x u_d(t,x) + f(u_d(t,x)), \tag{1.1}$$

let $\nu \in \{0, 1\}$, $\alpha \in \mathbb{R} \setminus \{-1, 1\}$, let $\alpha \colon \mathbb{R} \to \mathbb{R}$ satisfy for all $x \in \mathbb{R}$ that $a(x) = \nu \max\{x, \alpha x\} + (1 - \nu) \ln(1 + \exp(x))$, for every $d \in \mathbb{N}$, $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ let $\mathbf{A}(x) \in \mathbb{R}^d$ satisfy

$$\mathbf{A}(x) = (a(x_1), \dots, a(x_d)), \tag{1.2}$$

for every $L \in \mathbb{N}$, $l_0, l_1, \ldots, l_L \in \mathbb{N}$, $\Phi = ((W_1, B_1), \ldots, (W_L, B_L)) \in (\times_{k=1}^L (\mathbb{R}^{l_k \times l_{k-1}} \times \mathbb{R}^{l_k}))$ let $\mathcal{R}(\Phi) \colon \mathbb{R}^{l_0} \to \mathbb{R}^{l_L}$ and $\mathcal{P}(\Phi) \in \mathbb{N}$ satisfy for all $v_0 \in \mathbb{R}^{l_0}$, $v_1 \in \mathbb{R}^{l_1}$, \ldots , $v_L \in \mathbb{R}^{l_L}$ with $\forall k \in \{1, 2, \ldots, L-1\} \colon v_k = \mathbf{A}(W_k v_{k-1} + B_k)$ that

$$(\mathcal{R}(\Phi))(v_0) = W_L v_{L-1} + B_L$$
 and $\mathcal{P}(\Phi) = \sum_{k=1}^L l_k (l_{k-1} + 1),$ (1.3)

for every $d \in \mathbb{N}$ let

$$\mathbf{N}_{d} = \bigcup_{H \in \mathbb{N}} \bigcup_{(l_{0}, l_{1}, \dots, l_{H+1}) \in \{d\} \times \mathbb{N}^{H} \times \{1\}} (\times_{k=1}^{H+1} (\mathbb{R}^{l_{k} \times l_{k-1}} \times \mathbb{R}^{l_{k}})),$$
(1.4)

and assume for all $d \in \mathbb{N}$, $\varepsilon \in (0,1]$ that there exists $\mathbf{G} \in \mathbf{N}_d$ such that for all $t \in [0,T]$, $x = (x_1, \ldots, x_d) \in \mathbb{R}^d$ it holds that $\mathcal{P}(\mathbf{G}) \leq \kappa d^{\kappa} \varepsilon^{-\kappa}$ and

$$\varepsilon \left(|u_d(t,x)| + \sum_{k=1}^d \left| \frac{\partial}{\partial x_k} u_d(0,x) \right| \right) + |u_d(0,x) - (\mathcal{R}(\mathbf{G}))(x)| \le \varepsilon \kappa d^{\kappa} (1 + \sum_{k=1}^d |x_k|^{\kappa}).$$
(1.5)

Then there exists $c \in \mathbb{R}$ such that for all $d \in \mathbb{N}$, $\varepsilon \in (0,1]$ there exists $\mathbf{U} \in \mathbf{N}_{d+1}$ such that

$$\left[\int_{[0,T]\times[0,1]^d} |u_d(y) - (\mathcal{R}(\mathbf{U}))(y)|^p \,\mathrm{d}y\right]^{1/p} \le \varepsilon \qquad and \qquad \mathcal{P}(\mathbf{U}) \le cd^c \varepsilon^{-c}. \tag{1.6}$$

Theorem 1.1 is an immediate consequence of Corollary 5.5 in Subsection 5.2 below. Corollary 5.5, in turn, follows from Theorem 5.3, which is the main theorem of this article. In the following we add a few explanatory comments on the conclusion of Theorem 1.1 and the mathematical objects appearing in Theorem 1.1. The real number T > 0 in Theorem 1.1 describes the time horizon of the PDEs whose solutions we intend to approximate by ANNs in Theorem 1.1. The real number $\kappa > 0$ in Theorem 1.1 is a constant which we use to formulate the regularity and approximation assumption in (1.5) in Theorem 1.1. The real number p > 0 in Theorem 1.1 is a constant which determines the way how we measure the error between the PDE solution and its ANN approximation, that is, we measure the error between the PDE solution and its ANN approximation in the L^p -distance; see (1.6) in Theorem 1.1.

The function $f : \mathbb{R} \to \mathbb{R}$ in Theorem 1.1 is the nonlinearity in the PDEs whose solutions we intend to approximate by ANNs in Theorem 1.1. It is assumed to be Lipschitz continuous in the sense that there exists $c \in \mathbb{R}$ such that for all $x, y \in \mathbb{R}$ we have that

$$|f(x) - f(y)| \le c|x - y|.$$
(1.7)

In (1.1) in Theorem 1.1 we present the semilinear heat PDEs whose solutions we intend to approximate by ANNs in Theorem 1.1 and the functions $u_d: [0,T] \times \mathbb{R}^d \to \mathbb{R}, d \in \mathbb{N}$, in Theorem 1.1 are the solutions of the PDEs in (1.1).

The function $a: \mathbb{R} \to \mathbb{R}$ in Theorem 1.1 is the activation function for the approximating ANNs in Theorem 1.1. The real numbers $\nu, \alpha \in \mathbb{R}$ are two parameters that determine the concrete choice of the activation function $a: \mathbb{R} \to \mathbb{R}$. In particular, in the case $\nu = 0$ we have that a is nothing else but the *softplus activation* (see, e.g., [54, Section 1.2.5]), in the case $\nu = \alpha = 1$ we have that a is nothing else but the *ReLU activation* (see, e.g., [54, Section 1.2.3]), and in the case $\nu = 1, \alpha \in (0, 1)$ we have that a is nothing else but the *leaky ReLU activation* with leaky factor α (see, e.g., [54, Section 1.2.11]). We also note that in Theorem 1.1 we have for every $d \in \mathbb{N}$ that the function $\mathbb{R}^d \ni x \mapsto \mathbf{A}(x) \in \mathbb{R}^d$ is the *d*-dimensional version of the one-dimensional activation function $a: \mathbb{R} \to \mathbb{R}$.

The sets \mathbf{N}_d , $d \in \mathbb{N}$, in (1.4) describe the sets of the approximating ANNs in Theorem 1.1. Moreover, we note that for all $d \in \mathbb{N}$ and every $\Phi \in \mathbf{N}_d$ we have that the function $\mathcal{R}(\Phi) \colon \mathbb{R}^d \to \mathbb{R}$ in Theorem 1.1 is the realization function associated to the ANN Φ . Furthermore, we observe that for all $d \in \mathbb{N}$ and every $\Phi \in \mathbf{N}_d$ we have that the natural number $\mathcal{P}(\Phi)$ specifies the number of scalar real parameters used to describe the ANN Φ . In particular, we note that for all $d \in \mathbb{N}$ and every $\Phi \in \mathbf{N}_d$ we have that $\mathcal{P}(\Phi)$ is connected to the amount of memory (the amount of bits) needed to store Φ on a computer.

In Theorem 1.1 we also impose the assumption that the solutions $u_d: [0, T] \times \mathbb{R}^d \to \mathbb{R}$, $d \in \mathbb{N}$, of the PDEs in (1.1) grow at most polynomially in the PDE dimension and the spatial variable. This growth assumption is the subject of the regularity and approximation assumption in (1.5) in Theorem 1.1. More formally, observe that (1.5) implies that for all $d \in \mathbb{N}, t \in [0, T], x = (x_1, \ldots, x_d) \in \mathbb{R}^d$ we have that

$$|u_d(t,x)| \le \kappa d^{\kappa} (1 + \sum_{k=1}^d |x_k|^{\kappa}).$$
(1.8)

The assumption in (1.5) also ensures that the gradients of the initial values of the PDE solutions $u_d: [0,T] \times \mathbb{R}^d \to \mathbb{R}, d \in \mathbb{N}$, grow at most polynomially in the PDE dimension and the spatial variable. More formally, we note that (1.5) implies that for all $d \in \mathbb{N}$, $x = (x_1, \ldots, x_d) \in \mathbb{R}^d$ we have that

$$\sum_{k=1}^{d} \left| \frac{\partial}{\partial x_k} u_d(0, x) \right| \le \kappa d^{\kappa} \left(1 + \sum_{k=1}^{d} |x_k|^{\kappa} \right).$$
(1.9)

In addition, in Theorem 1.1 we also assume that the initial value functions $\mathbb{R}^d \ni x \mapsto u_d(0,x) \in \mathbb{R}, d \in \mathbb{N}$, of the PDE solutions $u_d: [0,T] \times \mathbb{R}^d \to \mathbb{R}, d \in \mathbb{N}$, can be approximated by ANNs without the COD in the sense of (1.5). More specifically, we observe that (1.5) ensures that for every arbitrarily large PDE dimension $d \in \mathbb{N}$ and every arbitrarily small prescribed approximation accuracy $\varepsilon \in (0,1]$ we have that there exists an ANN $\mathbf{G} \in \mathbf{N}_d$ such that for all $x = (x_1, \ldots, x_d) \in \mathbb{R}^d$ we have that the approximation error

$$|u_d(0,x) - (\mathcal{R}(\mathbf{G}))(x)| \tag{1.10}$$

between the initial value function $u_d(0, \cdot)$ evaluated at x and the realization $\mathcal{R}(\mathbf{G})$ of the ANN \mathbf{G} evaluated at x is bounded by $\varepsilon \kappa d^{\kappa}(1 + \sum_{k=1}^{d} |x_k|^{\kappa})$ and such that the number of parameters $\mathcal{P}(\mathbf{G})$ of the approximating ANN \mathbf{G} is bounded by $\kappa d^{\kappa} \varepsilon^{-\kappa}$.

In the above described setup Theorem 1.1 concludes in (1.6) that there exists a constant $c \in \mathbb{R}$ which is independent of the PDE dimension and the approximation accuracy such that for every arbitrarily large PDE dimension $d \in \mathbb{N}$ and every arbitrarily small prescribed approximation accuracy $\varepsilon \in (0, 1]$ we have that there must exist an ANN $\mathbf{U} \in \mathbf{N}_{d+1}$ such that the L^p -approximation error

$$\left[\int_{[0,T]\times[0,1]^d} |u_d(y) - (\mathcal{R}(\mathbf{U}))(y)|^p \,\mathrm{d}y\right]^{1/p} \tag{1.11}$$

is smaller than or equal to the prescribed approximation accuracy ε and such that the number of parameters $\mathcal{P}(\mathbf{U})$ of the approximating ANN \mathbf{U} (connected to the amount of memory to store \mathbf{U}) is bounded by $cd^c\varepsilon^{-c}$ and thus grows at most polynomially, in both, the reciprocal $1/\varepsilon$ of the prescribed approximation accuracy $\varepsilon > 0$ and the PDE dimension $d \in \mathbb{N}$.

The arguments in our proof of Theorem 1.1 are based on so-called *multilevel Picard* (MLP) approximation methods (see Hutzenthaler et al. [47]) and on the ANN representations for MLP methods in our preliminary article [1]. MLP methods are certain nonlinear Monte Carlo methods (see Hutzenthaler et al. [47] and E et al. [25]) that have been shown to overcome the COD for certain classes of semilinear PDEs (see, e.g., [8,9,25,31,44,45,47,48, 49,50,52,65,66]) and related problems (see, e.g., [12,51]). In our proof of Theorem 1.1 we employ that suitable MLP methods provably overcome the COD in the L^p -approximation of PDEs of the form (1.1) (see our preliminary work Hutzenthaler et al. [48]) and we design suitable deep ANNs that appropriately approximate temporal linear interpolations of such MLP approximations (see Corollary 4.13 in Subsection 4.6 below).

The remainder of this work is structured in the following way. In Section 2 we establish appropriate elementary perturbation and regularity estimates for solutions of PDEs. In our proofs of the ANN approximation results for PDEs in this work (such as Theorem 1.1 above) we employ certain ingredients of a suitable calculus for ANNs from the literature and in Section 3 we recall such ingredients of this ANN calculus. One of our main goals in Section 4 is to construct and study suitable ANNs (with general/abstract activations) which approximate linear interpolations of appropriate MLP approximations (see Corollary 4.13 in Section 4 for details). In Section 5 we employ some of the findings of Sections 2, 3, and 4 to prove the space-time ANN approximation results for semilinear heat PDEs in Theorem 5.3, Corollary 5.4, and Corollary 5.5. Theorem 1.1 in this introductory section is an immediate consequence of Corollary 5.5.

2 Properties of solutions of partial differential equations (PDEs)

The PDEs in the ANN approximation results for PDEs in this work (see Section 5 and Theorem 1.1 in the introduction) can be reformulated as suitable *stochastic fixed-point equations* (SFPEs) and the resulting SFPEs can then be solved approximately without the COD by means of certain nonlinear Monte Carlo methods, specifically, by means of MLP methods. Our proofs of the ANN approximation results for PDEs in this work (see Section 5 and Theorem 1.1 in the introduction) exploit this reformulation of the PDEs as SFPEs. In this section we establish certain elementary perturbation and regularity estimates for solutions of such SFPEs. We employ those perturbation and regularity estimates for solutions of SFPEs and PDEs, respectively, in the proofs of our ANN approximation results in Section 5 and Theorem 1.1.

In particular, in Corollary 2.3 we provide an elementary upper bound for the absolute value of the difference of two solutions u_1 and u_2 of SFPEs at the same space-time evaluation point but with different (perturbed) nonlinearities f_1 and f_2 and different (perturbed) terminal/initial value functions g_1 and g_2 . In this aspect we note that SFPEs are usually formulated as terminal value problems and in this aspect we also note the elementary fact that initial value PDE problems can be reformulated as terminal value PDE problems and vice versa (see, for example, [6, Remark 3.3]). Furthermore, in Corollary 2.7 we establish elementary temporal 1/2-Hölder continuity properties for solutions of SFPEs. In particular, Corollary 2.7 provides an upper bound for the absolute value of the difference of the solution u of an SFPE evaluated at the same spatial point but at different time points. We employ Corollary 2.3 and Corollary 2.7 in our proof of the ANN approximation result in Theorem 5.3 in Section 5.

2.1 Perturbation estimates for solutions of PDEs

Definition 2.1 (Standard and maximum norms). We denote by $\|\cdot\|: (\cup_{d\in\mathbb{N}}\mathbb{R}^d) \to \mathbb{R}$ and $\|\|\cdot\|: (\cup_{d\in\mathbb{N}}\mathbb{R}^d) \to \mathbb{R}$ the functions which satisfy for all $d \in \mathbb{N}$, $x = (x_1, \ldots, x_d) \in \mathbb{R}^d$ that $\|x\| = [\sum_{i=1}^d |x_i|^2]^{1/2}$ and $\|\|x\|\| = \max_{i\in\{1,2,\ldots,d\}} |x_i|.$

Lemma 2.2. Let $d \in \mathbb{N}$, $T, L, \mathfrak{L} \in (0, \infty)$, $p \in [1, \infty)$, let $f_1, f_2 \in C([0, T] \times \mathbb{R}^d \times \mathbb{R}, \mathbb{R})$ satisfy for all $i \in \{1, 2\}$, $s, t \in [0, T]$, $x \in \mathbb{R}^d$, $v, w \in \mathbb{R}$ that

$$|f_i(t,x,0)| \le \mathfrak{L}(1+||x||)^p, \qquad |f_1(t,x,v) - f_1(t,x,w)| \le L|v-w|, \tag{2.1}$$

and
$$|f_2(s, x, v) - f_2(t, x, w)| \le L(|s - t| + |v - w|),$$
 (2.2)

let $F_i: C([0,T] \times \mathbb{R}^d, \mathbb{R}) \to C([0,T] \times \mathbb{R}^d, \mathbb{R}), i \in \{1,2\}, \text{ satisfy for all } i \in \{1,2\}, t \in [0,T], x \in \mathbb{R}^d, v \in C([0,T] \times \mathbb{R}^d, \mathbb{R}) \text{ that}$

$$(F_i(v))(t,x) = f_i(t,x,v(t,x)),$$
(2.3)

let $g \in C(\mathbb{R}^d, \mathbb{R})$, let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $W : [0, T] \times \Omega \to \mathbb{R}^d$ be a standard Brownian motion, let $u_1, u_2 \in C([0, T] \times \mathbb{R}^d, \mathbb{R})$ satisfy for all $i \in \{1, 2\}$, $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$\mathbb{E}\left[|g(x+W_{T-t})| + \int_{t}^{T} |(F_{i}(u_{i}))(s, x+W_{s-t})| \,\mathrm{d}s\right] < \infty$$
(2.4)

and
$$u_i(t,x) = \mathbb{E}[g(x+W_{T-t})] + \int_t^T \mathbb{E}[(F_i(u_i))(s,x+W_{s-t})] \,\mathrm{d}s,$$
 (2.5)

and let $\mathfrak{u}_{i,\mathfrak{t}}: [0, T - \mathfrak{t}] \times \mathbb{R}^d \to \mathbb{R}$, $\mathfrak{t} \in [0, T]$, $i \in \{1, 2\}$, and $\mathfrak{f}_{i,\mathfrak{t}}: [0, T - \mathfrak{t}] \times \mathbb{R}^d \times \mathbb{R} \to \mathbb{R}$, $\mathfrak{t} \in [0, T]$, $i \in \{1, 2\}$, be the functions which satisfy for all $i \in \{1, 2\}$, $\mathfrak{t} \in [0, T]$, $t \in [0, T - \mathfrak{t}]$, $x \in \mathbb{R}^d$, $v \in \mathbb{R}$ that

$$\mathfrak{u}_{i,\mathfrak{t}}(t,x) = u_i(t+\mathfrak{t},x) \quad and \quad \mathfrak{f}_{i,\mathfrak{t}}(t,x,v) = f_i(t+\mathfrak{t},x,v) \tag{2.6}$$

(cf. Definition 2.1). Then

- (i) it holds for all $i \in \{1, 2\}$, $\mathfrak{t} \in [0, T]$ that $\mathfrak{f}_{i, \mathfrak{t}} \in C([0, T \mathfrak{t}] \times \mathbb{R}^d \times \mathbb{R}, \mathbb{R})$ and $\mathfrak{u}_{i, \mathfrak{t}} \in C([0, T \mathfrak{t}] \times \mathbb{R}^d, \mathbb{R})$,
- (ii) it holds for all $i \in \{1, 2\}$, $\mathfrak{t} \in [0, T]$, $t \in [0, T \mathfrak{t}]$, $x \in \mathbb{R}^d$ that

$$\mathbb{E}\left[\left|g(x+W_{(T-\mathfrak{t})-t})\right| + \int_{t}^{T-\mathfrak{t}} \left|\mathfrak{f}_{i,\mathfrak{t}}\left(s, x+W_{s-t}, \mathfrak{u}_{i,\mathfrak{t}}(s, x+W_{s-t})\right)\right| \,\mathrm{d}s\right] < \infty, \qquad (2.7)$$

(iii) it holds for all $i \in \{1, 2\}$, $\mathfrak{t} \in [0, T]$, $t \in [0, T - \mathfrak{t}]$, $x \in \mathbb{R}^d$ that

$$\mathbf{u}_{i,\mathfrak{t}}(t,x) = \mathbb{E}[g(x+W_{(T-\mathfrak{t})-t})] + \int_{t}^{T-\mathfrak{t}} \mathbb{E}\big[\mathfrak{f}_{i,\mathfrak{t}}\big(s,x+W_{s-t},\mathfrak{u}_{i,\mathfrak{t}}(s,x+W_{s-t})\big)\big] \,\mathrm{d}s, \quad (2.8)$$

and

(iv) it holds for all $i \in \{1, 2\}$, $\mathfrak{t} \in [0, T]$, $s, t \in [0, T - \mathfrak{t}]$, $x \in \mathbb{R}^d$, $v, w \in \mathbb{R}$ that

$$|\mathfrak{f}_{i,\mathfrak{t}}(t,x,0)| \le \mathfrak{L}(1+||x||)^p, \qquad |\mathfrak{f}_{1,\mathfrak{t}}(t,x,v) - \mathfrak{f}_{1,\mathfrak{t}}(t,x,w)| \le L|v-w|, \qquad (2.9)$$

and
$$|\mathfrak{f}_{2,\mathfrak{t}}(s,x,v) - \mathfrak{f}_{2,\mathfrak{t}}(t,x,w)| \le L(|s-t|+|v-w|).$$
 (2.10)

Proof of Lemma 2.2. Observe that the fact that $u_1, u_2 \in C([0,T] \times \mathbb{R}^d, \mathbb{R})$, the fact that $f_1, f_2 \in C([0,T] \times \mathbb{R}^d \times \mathbb{R}, \mathbb{R})$, and (2.6) establish item (i). Note that (2.3), (2.4), (2.5), (2.6), Fubini's theorem, and a change of variables ensure that for all $\mathfrak{t} \in [0,T]$, $t \in [0,T-\mathfrak{t}]$, $x \in \mathbb{R}^d$, $i \in \{1,2\}$ it holds that

$$\infty > \mathbb{E} \left[|g(x + W_{T-(t+\mathfrak{t})})| + \int_{(t+\mathfrak{t})}^{T} |(F_i(u_i))(s, x + W_{s-(t+\mathfrak{t})})| \, \mathrm{d}s \right]$$

= $\mathbb{E} \left[|g(x + W_{T-t-\mathfrak{t}})| + \int_{t}^{(T-\mathfrak{t})} |(F_i(u_i))(s + \mathfrak{t}, x + W_{s-t})| \, \mathrm{d}s \right]$
= $\mathbb{E} \left[|g(x + W_{(T-\mathfrak{t})-t})| + \int_{t}^{(T-\mathfrak{t})} |\mathfrak{f}_{i,\mathfrak{t}}(s, x + W_{s-t}, \mathfrak{u}_{i,\mathfrak{t}}(s, x + W_{s-t}))| \, \mathrm{d}s \right]$ (2.11)

and

$$\begin{aligned} \mathbf{u}_{i,t}(t,x) &= u_i(t+\mathbf{t},x) \\ &= \mathbb{E}[g(x+W_{T-(t+\mathbf{t})})] + \int_{(t+\mathbf{t})}^T \mathbb{E}[(F_i(u_i))(s,x+W_{s-(t+\mathbf{t})})] \, \mathrm{d}s \\ &= \mathbb{E}[g(x+W_{T-(t+\mathbf{t})})] + \mathbb{E}\left[\int_{(t+\mathbf{t})}^T (F_i(u_i))(s,x+W_{s-(t+\mathbf{t})}) \, \mathrm{d}s\right] \\ &= \mathbb{E}[g(x+W_{(T-\mathbf{t})-t})] + \mathbb{E}\left[\int_t^{(T-\mathbf{t})} \mathbf{f}_{i,\mathbf{t}}(s,x+W_{s-t},\mathbf{u}_{i,\mathbf{t}}(s,x+W_{s-t}))] \, \mathrm{d}s\right] \\ &= \mathbb{E}[g(x+W_{(T-\mathbf{t})-t})] + \int_t^{(T-\mathbf{t})} \mathbb{E}[\mathbf{f}_{i,\mathbf{t}}(s,x+W_{s-t},\mathbf{u}_{i,\mathbf{t}}(s,x+W_{s-t}))] \, \mathrm{d}s. \end{aligned}$$
(2.12)

This proves items (ii) and (iii). Combining (2.1), (2.2), and (2.6) establishes item (iv). The proof of Lemma 2.2 is thus complete.

Corollary 2.3. Let $d \in \mathbb{N}$, $T, L, \mathfrak{L}, B \in (0, \infty)$, $p, q \in [1, \infty)$, $f_1, f_2 \in C([0, T] \times \mathbb{R}^d \times \mathbb{R}, \mathbb{R})$, $g_1, g_2 \in C(\mathbb{R}^d, \mathbb{R})$ satisfy for all $i \in \{1, 2\}$, $t \in [0, T]$, $x \in \mathbb{R}^d$, $v, w \in \mathbb{R}$ that

$$|f_i(t, x, v) - f_i(t, x, w)| \le L|v - w|, \qquad \max\{|f_i(t, x, 0)|, |g_i(x)|\} \le \mathfrak{L}(1 + ||x||)^p, \quad (2.13)$$

and
$$\max\{|f_1(t,x,v) - f_2(t,x,v)|, |g_1(x) - g_2(x)|\} \le B((1+||x||)^{pq} + |v|^q), \quad (2.14)$$

let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $W: [0, T] \times \Omega \to \mathbb{R}^d$ be a standard Brownian motion, and let $u_1, u_2 \in C([0, T] \times \mathbb{R}^d, \mathbb{R})$ satisfy for all $i \in \{1, 2\}, t \in [0, T], x \in \mathbb{R}^d$ that

$$\mathbb{E}\left[|g_i(x+W_{T-t})| + \int_t^T |f_i(s, x+W_{s-t}, u_i(s, x+W_{s-t}))| \,\mathrm{d}s\right] < \infty$$
(2.15)

and
$$u_i(t,x) = \mathbb{E}[g_i(x+W_{T-t})] + \int_t^T \mathbb{E}[f_i(s,x+W_{s-t},u_i(s,x+W_{s-t}))] \,\mathrm{d}s$$
 (2.16)

(cf. Definition 2.1). Then it holds for all $t \in [0,T]$, $x \in \mathbb{R}^d$ that

$$|u_1(t,x) - u_2(t,x)| \le B \left(e^{LT} (T+1) \right)^{q+1} \left(\mathfrak{L}^q + 1 \right) 3^{pq-1} \left(1 + \|x\|^{pq} + \sup_{s \in [0,T]} \mathbb{E}[\|W_s\|^{pq}] \right).$$
(2.17)

Proof of Corollary 2.3. Throughout this proof let $\mathfrak{u}_{i,\mathfrak{t}}: [0, T - \mathfrak{t}] \times \mathbb{R}^d \to \mathbb{R}, \mathfrak{t} \in [0, T], i \in \{1, 2\}$, and $\mathfrak{f}_{i,\mathfrak{t}}: [0, T - \mathfrak{t}] \times \mathbb{R}^d \times \mathbb{R} \to \mathbb{R}, \mathfrak{t} \in [0, T], i \in \{1, 2\}$, be the functions which satisfy for all $\mathfrak{t} \in [0, T], t \in [0, T - \mathfrak{t}], x \in \mathbb{R}^d, v \in \mathbb{R}, i \in \{1, 2\}$ that $\mathfrak{u}_{i,\mathfrak{t}}(t, x) = u_i(t + \mathfrak{t}, x)$ and $\mathfrak{f}_{i,\mathfrak{t}}(t, x, v) = f_i(t + \mathfrak{t}, x, v)$. Observe that (2.14) implies that for all $\mathfrak{t} \in [0, T], t \in [0, T - \mathfrak{t}], x \in \mathbb{R}^d, v \in \mathbb{R}$ it holds that

$$|\mathfrak{f}_{1,\mathfrak{t}}(t,x,v) - \mathfrak{f}_{2,\mathfrak{t}}(t,x,v)| = |f_1(t+\mathfrak{t},x,v) - f_2(t+\mathfrak{t},x,v)| \le B\big((1+||x||)^{pq} + |v|^q\big) \quad (2.18)$$

(cf. Definition 2.1). This, Lemma 2.2, and Hutzenthaler et al. [46, Lemma 2.3] (applied for every $\mathfrak{t} \in [0,T]$ with $T \curvearrowleft T - \mathfrak{t}$, $L \curvearrowleft L$, $B \backsim \mathfrak{L}$, $\delta \backsim B$, $u_1 \curvearrowleft \mathfrak{u}_{1,\mathfrak{t}}$, $u_2 \backsim \mathfrak{u}_{2,\mathfrak{t}}$, $f_1 \curvearrowleft \mathfrak{f}_{1,\mathfrak{t}}, f_2 \curvearrowleft \mathfrak{f}_{2,\mathfrak{t}}, g_1 \curvearrowleft g_1, g_2 \curvearrowleft g_2, p \curvearrowleft p, q \backsim q$ in the notation of Hutzenthaler et al. [46, Lemma 2.3]) show that for all $\mathfrak{t} \in [0,T], t \in [0,T-\mathfrak{t}], x \in \mathbb{R}^d$ it holds that

$$\mathbb{E}\Big[\Big|\mathfrak{u}_{1,\mathfrak{t}}(t,x+W_t) - \mathfrak{u}_{2,\mathfrak{t}}(t,x+W_t)\Big|\Big] \le B\Big(e^{LT}(T+1)\Big)^{q+1}\Big(\mathfrak{L}^q+1\Big)\Big(1 + \|x\| + \big(\mathbb{E}[\|W_{T-\mathfrak{t}}\|^{pq}]\big)^{\frac{1}{pq}}\Big)^{pq}.$$
(2.19)

This and Jensen's inequality prove that for all $\mathfrak{t} \in [0,T], x \in \mathbb{R}^d$ it holds that

$$|u_{1}(\mathfrak{t},x) - u_{2}(\mathfrak{t},x)| = |\mathfrak{u}_{1,\mathfrak{t}}(0,x) - \mathfrak{u}_{2,\mathfrak{t}}(0,x)| = \mathbb{E}\left[|\mathfrak{u}_{1,\mathfrak{t}}(0,x+W_{0}) - \mathfrak{u}_{2,\mathfrak{t}}(0,x+W_{0})|\right] \\ \leq B\left(e^{LT}(T+1)\right)^{q+1} (\mathfrak{L}^{q}+1)3^{pq-1}\left(1+\|x\|^{pq} + \sup_{s\in[0,T]}\mathbb{E}[\|W_{s}\|^{pq}]\right).$$

$$(2.20)$$

The proof of Corollary 2.3 is thus complete.

2.2 Temporal regularity estimates for solutions of PDEs

Lemma 2.4. Let $d \in \mathbb{N}$, $T, \mathfrak{L} \in (0, \infty)$, $p \in [1, \infty)$, let $g \in C^1(\mathbb{R}^d, \mathbb{R})$ satisfy for all $x \in \mathbb{R}^d$ that $\|\nabla g(x)\| \leq \mathfrak{L}(1+\|x\|)^p$, let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, and let $W: [0, T] \times \Omega \to \mathbb{R}^d$ be a standard Brownian motion (cf. Definition 2.1). Then it holds for all $t, \mathfrak{t} \in [0, T]$, $x \in \mathbb{R}^d$ that

$$\mathbb{E}\left[|g(x+W_{\mathfrak{t}}) - g(x+W_{t})|\right] \le 8^{p+1} \mathfrak{L}\left(1 + \|x\|^{p} + \sup_{s \in [0,T]} \left(\mathbb{E}[\|W_{s}\|^{2p}]\right)^{\frac{1}{2}}\right) \sqrt{|\mathfrak{t} - t|} \sqrt{d}.$$
(2.21)

Proof of Lemma 2.4. Note that the fundamental theorem of calculus, the multivariate chain rule, and the Cauchy–Schwarz inequality assure that for all $t, t \in [0, T], x \in \mathbb{R}^d$ it holds that

$$|g(x+W_{\mathfrak{t}}) - g(x+W_{t})| \leq \int_{0}^{1} \left\| \nabla g \left((W_{\mathfrak{t}} - W_{t})z + x + W_{t} \right) \right\| \|W_{\mathfrak{t}} - W_{t}\| \, \mathrm{d}z.$$
(2.22)

Next we combine the fact that for all $x \in \mathbb{R}^d$ it holds that $\|\nabla g(x)\| \leq \mathfrak{L}(1+\|x\|)^p$, the triangle inequality, and Jensen's inequality to obtain that for all $t, \mathfrak{t} \in [0, T]$, $x \in \mathbb{R}^d$, $z \in [0, 1]$ it holds that

$$\begin{aligned} \left\| \nabla g \big((W_{\mathfrak{t}} - W_{t}) z + x + W_{t} \big) \right\| &\leq \mathfrak{L} \big(1 + \left\| (W_{\mathfrak{t}} - W_{t}) z + x + W_{t} \right\| \big)^{p} \\ &\leq \mathfrak{L} \big(1 + \left\| W_{\mathfrak{t}} \right\| + \left\| x \right\| + 2 \left\| W_{t} \right\| \big)^{p} \\ &\leq 4^{p-1} \mathfrak{L} \big(1 + \left\| W_{\mathfrak{t}} \right\|^{p} + \left\| x \right\|^{p} + 2^{p} \left\| W_{t} \right\|^{p} \big). \end{aligned}$$

$$(2.23)$$

This and (2.22) demonstrate that for all $t, t \in [0, T], x \in \mathbb{R}^d$ it holds that

$$|g(x+W_{\mathfrak{t}}) - g(x+W_{\mathfrak{t}})| \le 4^{p-1}\mathfrak{L}\left(1 + ||W_{\mathfrak{t}}||^{p} + ||x||^{p} + 2^{p}||W_{\mathfrak{t}}||^{p}\right)||W_{\mathfrak{t}} - W_{\mathfrak{t}}||.$$
(2.24)

Combining this, the Cauchy–Schwarz inequality, and Jensen's inequality establishes that for all $t, t \in [0, T], x \in \mathbb{R}^d$ it holds that

$$\mathbb{E}\left[\left|g(x+W_{\mathfrak{t}})-g(x+W_{t})\right|\right] \leq 4^{p-1}\mathfrak{L}\left(\mathbb{E}\left[\left(1+\|W_{\mathfrak{t}}\|^{p}+\|x\|^{p}+2^{p}\|W_{t}\|^{p}\right)^{2}\right]\right)^{\frac{1}{2}}\left(\mathbb{E}\left[\|W_{\mathfrak{t}}-W_{t}\|^{2}\right]\right)^{\frac{1}{2}} \leq 4^{p}\mathfrak{L}\left(1+\|x\|^{2p}+\mathbb{E}\left[\|W_{\mathfrak{t}}\|^{2p}\right]+2^{2p}\mathbb{E}\left[\|W_{t}\|^{2p}\right]\right)^{\frac{1}{2}}\left(\mathbb{E}\left[\|W_{|\mathfrak{t}-t|}\|^{2}\right]\right)^{\frac{1}{2}} \leq 4^{p}\mathfrak{L}\left(1+\|x\|^{2p}+(1+2^{2p})\sup_{s\in[0,T]}\mathbb{E}\left[\|W_{s}\|^{2p}\right]\right)^{\frac{1}{2}}\left(|\mathfrak{t}-t|d\right)^{\frac{1}{2}} \leq 4^{p}(1+2^{p})\mathfrak{L}\left(1+\|x\|^{p}+\sup_{s\in[0,T]}\left(\mathbb{E}\left[\|W_{s}\|^{2p}\right]\right)^{\frac{1}{2}}\right)\left(|\mathfrak{t}-t|d\right)^{\frac{1}{2}}.$$
(2.25)

The proof of Lemma 2.4 is thus complete.

Lemma 2.5. Let $T, \kappa \in (0, \infty)$, $p, r, q, q \in [1, \infty)$, let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, for every $d \in \mathbb{N}$ let $W^d \colon [0, T] \times \Omega \to \mathbb{R}^d$ be a standard Brownian motion, and for every $d \in \mathbb{N}$ let $\nu_d \colon \mathcal{B}(\mathbb{R}^{d+1}) \to [0, \infty)$ be a measure with

$$\int_{\mathbb{R}^{d+1}} (1 + \|y\|^{p^2 q \mathfrak{q}}) \nu_d(\mathrm{d}y) \le \kappa d^{r p^2 q \mathfrak{q}}$$

$$(2.26)$$

(cf. Definition 2.1). Then

(i) it holds for all $d \in \mathbb{N}$, $\mathfrak{p} \in [1, \infty)$, $s \in [0, T]$ that

$$\mathbb{E}[\|W_s^d\|^{\mathfrak{p}}] \le 1 + (1+2T)^{\mathfrak{p}} \left(\frac{d}{2} + \mathfrak{p}\right)^{\mathfrak{p}}$$
(2.27)

and

(ii) there exists $\mathfrak{C} \in [1,\infty)$ such that for all $d \in \mathbb{N}$ it holds that

$$\int_{[0,T]\times\mathbb{R}^d} \left(1 + \|x\|^{p^2q\mathfrak{q}} + \sup_{s\in[0,T]} \mathbb{E}\left[\|W_s^d\|^{2p^2q\mathfrak{q}}\right]\right) \nu_d(\mathrm{d}t,\mathrm{d}x) \le \mathfrak{C}d^{(r+2)p^2q\mathfrak{q}}.$$
 (2.28)

Proof of Lemma 2.5. Observe that the fact that for all $d \in \mathbb{N}$, $s \in (0, T]$ the random variable $||W_s^d/\sqrt{s}||^2$ is chi-square distributed with d degrees of freedom, Jensen's inequality, and, for instance, (2.35) in [75], show that for all $d \in \mathbb{N}$, $k \in \mathbb{N}$, $s \in [0, T]$ it holds that

$$\left(\mathbb{E}[\|W_s^d\|^k]\right)^2 \le \mathbb{E}\left[\|W_s^d\|^{2k}\right] \le \frac{(2s)^k \Gamma(\frac{d}{2}+k)}{\Gamma(\frac{d}{2})} = (2s)^k \prod_{j=0}^{k-1} \left(\frac{d}{2}+j\right) \le (2s)^k \left(\frac{d}{2}+k-1\right)^k.$$
(2.29)

This ensures that for all $d \in \mathbb{N}$, $\mathfrak{p} \in [1, \infty)$, $s \in [0, T]$ it holds that

$$\mathbb{E}[\|W_s^d\|^{\mathfrak{p}}] \le 1 + \mathbb{E}[\|W_s^d\|^{\lceil p\rceil}] \le 1 + (2s)^{\frac{\lceil \mathfrak{p}\rceil}{2}} \left(\frac{d}{2} + \lceil \mathfrak{p}\rceil - 1\right)^{\frac{\lceil \mathfrak{p}\rceil}{2}} \le 1 + (1 + 2T)^{\mathfrak{p}} \left(\frac{d}{2} + \mathfrak{p}\right)^{\mathfrak{p}}.$$
(2.30)

This and (2.26) ensure that for all $d \in \mathbb{N}$ it holds that

$$\int_{[0,T]\times\mathbb{R}^{d}} \left(1 + \|x\|^{p^{2}qq} + \sup_{s\in[0,T]} \mathbb{E}\left[\|W_{s}^{d}\|^{2p^{2}qq}\right]\right) \nu_{d}(\mathrm{d}t,\mathrm{d}x) \\
\leq \kappa d^{rp^{2}qq} + \kappa d^{rp^{2}qq} \left(1 + (1+2T)^{2p^{2}qq} \left(\frac{d}{2} + 2p^{2}qq\right)^{2p^{2}qq}\right) \\
\leq 2\kappa d^{rp^{2}qq} + \kappa (1+2T)^{2p^{2}qq} d^{(r+2)p^{2}qq} \left(\frac{1}{2} + 2p^{2}qq\right)^{2p^{2}qq} \\
\leq \kappa \left(2 + (1+2T)^{2p^{2}qq} \left(\frac{1}{2} + 2p^{2}qq\right)^{2p^{2}qq}\right) d^{(r+2)p^{2}qq} \qquad (2.31)$$

(cf. Definition 2.1). The proof of Lemma 2.5 is thus complete.

Lemma 2.6. Let $d \in \mathbb{N}$, $T, L, \mathfrak{L} \in (0, \infty)$, $p \in [1, \infty)$, let $f \in C([0, T] \times \mathbb{R}^d \times \mathbb{R}, \mathbb{R})$ and $g \in C^1(\mathbb{R}^d, \mathbb{R})$ satisfy for all $s, t \in [0, T]$, $x \in \mathbb{R}^d$, $v, w \in \mathbb{R}$ that

$$|f(s, x, v) - f(t, x, w)| \le L(|s - t| + |v - w|)$$
(2.32)

and
$$\max\{|f(t,x,0)|, |g(x)|, \|\nabla g(x)\|\} \le \mathfrak{L}(1+\|x\|)^p,$$
 (2.33)

let $F: C([0,T] \times \mathbb{R}^d, \mathbb{R}) \to C([0,T] \times \mathbb{R}^d, \mathbb{R})$ satisfy for all $t \in [0,T]$, $x \in \mathbb{R}^d$, $v \in C([0,T] \times \mathbb{R}^d, \mathbb{R})$ that

$$(F(v))(t,x) = f(t,x,v(t,x)),$$
(2.34)

let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $W : [0, T] \times \Omega \to \mathbb{R}^d$ be a standard Brownian motion, and let $u \in C([0, T] \times \mathbb{R}^d, \mathbb{R})$ satisfy for all $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$\mathbb{E}\left[|g(x+W_{T-t})| + \int_{t}^{T} |(F(u))(s, x+W_{s-t})| \,\mathrm{d}s\right] < \infty$$
(2.35)

and
$$u(t,x) = \mathbb{E}[g(x+W_{T-t})] + \int_{t}^{T} \mathbb{E}[(F(u))(s,x+W_{s-t})] \,\mathrm{d}s$$
 (2.36)

(cf. Definition 2.1). Then it holds for all $\mathfrak{t} \in [0,T]$, $t \in [0,\mathfrak{t}]$, $x \in \mathbb{R}^d$ that

$$\mathbb{E}\left[\left|u(\mathfrak{t}, x+W_{t})-u(t, x+W_{t})\right|\right] \le e^{2LT}(T+1)^{2}(L+1)(\mathfrak{L}+1)8^{p+2}\left(1+\|x\|^{p}+\sup_{s\in[0,T]}\mathbb{E}\left[\|W_{s}\|^{2p}\right]\right)\sqrt{|\mathfrak{t}-t|}\sqrt{d}.$$
(2.37)

Proof of Lemma 2.6. Note that (2.36) and the triangle inequality show that for all $\delta \in (0, T]$, $x \in \mathbb{R}^d$, $t \in [0, T - \delta]$ it holds that

$$|u(t + \delta, x) - u(t, x)| \le \mathbb{E}[|g(x + W_{T-(t+\delta)}) - g(x + W_{T-t})|] + \left| \int_{t+\delta}^{T} \mathbb{E}[(F(u))(s, x + W_{s-(t+\delta)})] \, \mathrm{d}s - \int_{t}^{T} \mathbb{E}[(F(u))(s, x + W_{s-t})] \, \mathrm{d}s \right|.$$
(2.38)

In the next step we combine (2.35), Fubini's theorem, and a change of variables to obtain that for all $\delta \in (0, T]$, $x \in \mathbb{R}^d$, $t \in [0, T - \delta]$ it holds that

$$\int_{t+\delta}^{T} \mathbb{E}\left[(F(u))(s, x + W_{s-(t+\delta)}) \right] ds = \mathbb{E}\left[\int_{t+\delta}^{T} (F(u))(s, x + W_{s-(t+\delta)}) ds \right]$$
$$= \mathbb{E}\left[\int_{t}^{T-\delta} (F(u))(s + \delta, x + W_{s-t}) ds \right]$$
$$= \int_{t}^{T-\delta} \mathbb{E}\left[(F(u))(s + \delta, x + W_{s-t}) \right] ds.$$
(2.39)

This, (2.34), (2.35), and the triangle inequality imply that for all $\delta \in (0,T], x \in \mathbb{R}^d$,

 $t \in [0, T - \delta]$ it holds that

$$\begin{aligned} \left| \int_{t+\delta}^{T} \mathbb{E} \left[(F(u))(s, x + W_{s-(t+\delta)}) \right] \mathrm{d}s - \int_{t}^{T} \mathbb{E} \left[(F(u))(s, x + W_{s-t}) \right] \mathrm{d}s \right| \\ &= \left| \int_{t}^{T-\delta} \mathbb{E} \left[(F(u))(s + \delta, x + W_{s-t}) \right] \mathrm{d}s - \int_{t}^{T-\delta} \mathbb{E} \left[(F(u))(s, x + W_{s-t}) \right] \mathrm{d}s \right| \\ &- \int_{T-\delta}^{T} \mathbb{E} \left[(F(u))(s, x + W_{s-t}) \right] \mathrm{d}s \right| \\ &\leq \int_{t}^{T-\delta} \mathbb{E} \left[\left| f \left(s + \delta, x + W_{s-t}, u(s + \delta, x + W_{s-t}) \right) - f \left(s, x + W_{s-t}, u(s, x + W_{s-t}) \right) \right| \right] \mathrm{d}s \\ &+ \int_{T-\delta}^{T} \mathbb{E} \left[\left| f \left(s, x + W_{s-t}, u(s, x + W_{s-t}) \right) \right| \right] \mathrm{d}s. \end{aligned}$$

$$(2.40)$$

In addition, observe that the fact that for all $s, t \in [0, T]$, $x \in \mathbb{R}^d$, $v, w \in \mathbb{R}$ it holds that $|f(s, x, v) - f(t, x, w)| \le L(|s-t|+|v-w|)$ shows that for all $\delta \in (0, T]$, $x \in \mathbb{R}^d$, $t \in [0, T-\delta]$ it holds that

$$\int_{t}^{T-\delta} \mathbb{E}\left[\left|f\left(s+\delta, x+W_{s-t}, u(s+\delta, x+W_{s-t})\right) - f\left(s, x+W_{s-t}, u(s, x+W_{s-t})\right)\right|\right] \mathrm{d}s$$

$$\leq L\delta T + L \int_{t}^{T-\delta} \mathbb{E}\left[\left|u(s+\delta, x+W_{s-t}) - u(s, x+W_{s-t})\right|\right] \mathrm{d}s.$$
(2.41)

Furthermore, note that the fact that for all $s, t \in [0, T]$, $x \in \mathbb{R}^d$, $v, w \in \mathbb{R}$ it holds that $|f(s, x, v) - f(t, x, w)| \leq L(|s - t| + |v - w|)$, the fact that for all $t \in [0, T]$, $x \in \mathbb{R}^d$ it holds that $|f(t, x, 0)| \leq \mathfrak{L}(1 + ||x||)^p$, and the triangle inequality prove that for all $\delta \in (0, T]$, $x \in \mathbb{R}^d$, $t \in [0, T - \delta]$, $s \in [T - \delta, T]$ it holds that

$$\mathbb{E}\left[\left|f\left(s, x + W_{s-t}, u(s, x + W_{s-t})\right)\right|\right] \\
\leq \mathbb{E}\left[\left|f\left(s, x + W_{s-t}, u(s, x + W_{s-t})\right) - f\left(s, x + W_{s-t}, 0\right)\right|\right] + \mathbb{E}\left[\left|f\left(s, x + W_{s-t}, 0\right)\right|\right] \quad (2.42) \\
\leq L\mathbb{E}\left[\left||u(s, x + W_{s-t})|\right] + \mathfrak{LE}\left[(1 + ||x + W_{s-t}||)^{p}\right].$$

Combining this, (2.38), (2.40), and (2.41) demonstrates that for all $\delta \in (0, T]$, $x \in \mathbb{R}^d$, $t \in [0, T - \delta]$ it holds that

$$|u(t + \delta, x) - u(t, x)| \leq \mathbb{E} \left[|g(x + W_{T-(t+\delta)}) - g(x + W_{T-t})| \right] + L\delta T + L \int_{t}^{T-\delta} \mathbb{E} \left[|u(s + \delta, x + W_{s-t}) - u(s, x + W_{s-t})| \right] ds + L \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s, x + W_{s-t})|] ds + \mathcal{L} \int_{T-\delta}^{T} \mathbb{E} [|u(s$$

Fubini's theorem and the fact that W has independent and stationary increments therefore establish that for all $\delta \in (0, T]$, $x \in \mathbb{R}^d$, $t \in [0, T - \delta]$ it holds that

$$\mathbb{E}\left[\left|u(t+\delta, x+W_{t})-u(t, x+W_{t})\right|\right] \\
\leq \mathbb{E}\left[\left|g(x+W_{T-\delta})-g(x+W_{T})\right|\right] + L\delta T \\
+ L \int_{t}^{T-\delta} \mathbb{E}\left[\left|u(s+\delta, x+W_{s})-u(s, x+W_{s})\right|\right] \mathrm{d}s \\
+ L \int_{T-\delta}^{T} \mathbb{E}\left[\left|u(s, x+W_{s})\right|\right] \mathrm{d}s + \mathfrak{L} \int_{T-\delta}^{T} \mathbb{E}\left[(1+\|x+W_{s}\|)^{p}\right] \mathrm{d}s.$$
(2.44)

In addition, observe that Hutzenthaler et al. [46, Lemma 2.2] (applied with $d \curvearrowleft d, T \backsim T$, $L \curvearrowleft L, B \curvearrowleft \mathfrak{L}, p \curvearrowleft p, q \backsim 1, f_1 \curvearrowleft f, g_1 \curvearrowleft g, u_1 \curvearrowleft u$ in the notation of Hutzenthaler et al. [46, Lemma 2.2]), the triangle inequality, and Jensen's inequality ensure that for all $x \in \mathbb{R}^d, s \in [0, T]$ it holds that

$$\mathbb{E}\left[|u(s, x+W_s)|\right] \le e^{LT}(T+1)\mathfrak{L}3^{p-1}\left(1+\|x\|^p+\sup_{r\in[0,T]}\mathbb{E}[\|W_r\|^p]\right).$$
(2.45)

The triangle inequality and Jensen's inequality hence imply that for all $\delta \in (0, T]$, $x \in \mathbb{R}^d$, $t \in [0, T - \delta]$ it holds that

$$L\delta T + L \int_{T-\delta}^{T} \mathbb{E}[|u(s, x + W_{s})|] \, ds + \mathfrak{L} \int_{T-\delta}^{T} \mathbb{E}[(1 + ||x + W_{s}||)^{p}] \, ds$$

$$\leq L\delta T + L\delta e^{LT} (T+1)\mathfrak{L} 3^{p-1} (1 + ||x||^{p} + \sup_{r \in [0,T]} \mathbb{E}[||W_{r}||^{p}])$$

$$+ \mathfrak{L} \delta 3^{p-1} (1 + ||x||^{p} + \sup_{s \in [0,T]} \mathbb{E}[||W_{s}||^{p}])$$

$$\leq \delta e^{LT} (T+1) (L+1) (\mathfrak{L} + 1) 3^{p} (1 + ||x||^{p} + \sup_{s \in [0,T]} \mathbb{E}[||W_{s}||^{p}])$$

$$\leq \sqrt{\delta} e^{LT} (T+1)^{2} (L+1) (\mathfrak{L} + 1) 3^{p+1} (1 + ||x||^{p} + \sup_{s \in [0,T]} \mathbb{E}[||W_{s}||^{2p}]).$$
(2.46)

Combining this, (2.44), and Lemma 2.4 shows that for all $\delta \in (0, T]$, $x \in \mathbb{R}^d$, $t \in [0, T - \delta]$ it holds that

$$\mathbb{E}\left[\left|u(t+\delta,x+W_{t})-u(t,x+W_{t})\right|\right] \leq 8^{p+1}\mathfrak{L}\left(1+\|x\|^{p}+1+\sup_{s\in[0,T]}\mathbb{E}\left[\|W_{s}\|^{2p}\right]\right)\sqrt{\delta}\sqrt{d} + L\int_{t}^{T-\delta}\mathbb{E}\left[\left|u(s+\delta,x+W_{s})-u(s,x+W_{s})\right|\right]ds + \sqrt{\delta}e^{LT}(T+1)^{2}(L+1)(\mathfrak{L}+1)3^{p+1}\left(1+\|x\|^{p}+\sup_{s\in[0,T]}\mathbb{E}\left[\|W_{s}\|^{2p}\right]\right) \leq e^{LT}(T+1)^{2}(L+1)(\mathfrak{L}+1)8^{p+2}\left(1+\|x\|^{p}+\sup_{s\in[0,T]}\mathbb{E}\left[\|W_{s}\|^{2p}\right]\right)\sqrt{\delta}\sqrt{d} + L\int_{t}^{T-\delta}\mathbb{E}\left[\left|u(s+\delta,x+W_{s})-u(s,x+W_{s})\right|\right]ds$$
(2.47)

(cf. Definition 2.1). This, (2.45), Lemma 2.5, and Gronwall's integral inequality (see, for example, [48, Corollary 2.2]) prove that for all $\delta \in (0, T]$, $x \in \mathbb{R}^d$, $t \in [0, T - \delta]$ it holds that

$$\mathbb{E}\left[|u(t+\delta, x+W_t) - u(t, x+W_t)|\right] \\
\leq e^{2LT}(T+1)^2(L+1)(\mathfrak{L}+1)8^{p+2}\left(1+\|x\|^p + \sup_{s\in[0,T]}\mathbb{E}\left[\|W_s\|^{2p}\right]\right)\sqrt{\delta}\sqrt{d}.$$
(2.48)

The proof of Lemma 2.6 is thus complete.

Corollary 2.7. Let $d \in \mathbb{N}$, $T, L, \mathfrak{L} \in (0, \infty)$, $p \in [1, \infty)$, let $f \in C([0, T] \times \mathbb{R}^d \times \mathbb{R}, \mathbb{R})$ and $g \in C^1(\mathbb{R}^d, \mathbb{R})$ satisfy for all $s, t \in [0, T]$, $x \in \mathbb{R}^d$, $v, w \in \mathbb{R}$ that

$$|f(s, x, v) - f(t, x, w)| \le L(|s - t| + |v - w|)$$
(2.49)

and
$$\max\{|f(t,x,0)|, |g(x)|, \|\nabla g(x)\|\} \le \mathfrak{L}(1+\|x\|)^p,$$
 (2.50)

let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let $W: [0, T] \times \Omega \to \mathbb{R}^d$ be a standard Brownian motion, and let $u \in C([0, T] \times \mathbb{R}^d, \mathbb{R})$ satisfy for all $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$\mathbb{E}\left[\left|g(x+W_{T-t})\right| + \int_{t}^{T} \left|f\left(s, x+W_{s-t}, u(s, x+W_{s-t})\right)\right| \,\mathrm{d}s\right] < \infty$$
(2.51)

and
$$u(t,x) = \mathbb{E}[g(x+W_{T-t})] + \int_t^T \mathbb{E}[f(s,x+W_{s-t},u(s,x+W_{s-t}))] \,\mathrm{d}s$$
 (2.52)

(cf. Definition 2.1). Then it holds for all $\mathfrak{t}, t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$|u(\mathbf{t}, x) - u(t, x)| \le e^{2LT} (T+1)^2 (L+1) (\mathfrak{L}+1) 8^{p+2} (1+||x||^p + \sup_{s \in [0,T]} \mathbb{E} [||W_s||^{2p}]) \sqrt{|\mathbf{t}-t|} \sqrt{d}.$$
(2.53)

Proof of Corollary 2.7. Let $\mathfrak{u}_{\mathfrak{t}}: [0, T-\mathfrak{t}] \times \mathbb{R}^{d} \to \mathbb{R}, \mathfrak{t} \in [0, T], \text{ and } \mathfrak{f}_{\mathfrak{t}}: [0, T-\mathfrak{t}] \times \mathbb{R}^{d} \times \mathbb{R} \to \mathbb{R}, \mathfrak{t} \in [0, T], be the functions which satisfy for all <math>\mathfrak{t} \in [0, T], t \in [0, T-\mathfrak{t}], x \in \mathbb{R}^{d}, v \in \mathbb{R}$ that $\mathfrak{u}_{\mathfrak{t}}(t, x) = u(t + \mathfrak{t}, x)$ and $\mathfrak{f}_{\mathfrak{t}}(t, x, v) = f(t + \mathfrak{t}, x, v)$. Note that Lemma 2.2 and Lemma 2.6 (applied for every $\mathfrak{t} \in [0, T]$ with $d \frown d, T \frown T - \mathfrak{t}, L \frown L, \mathfrak{L} \frown \mathfrak{L}, p \frown p, f \frown \mathfrak{f}_{\mathfrak{t}}, u \frown \mathfrak{u}_{\mathfrak{t}}, g \frown g$ in the notation of Lemma 2.6) demonstrate that for all $\mathfrak{t} \in [0, T], t \in [0, T-\mathfrak{t}], x \in \mathbb{R}^{d}$ it holds that

$$|u(t + \mathfrak{t}, x) - u(\mathfrak{t}, x)| = |\mathfrak{u}_{\mathfrak{t}}(t, x) - \mathfrak{u}_{\mathfrak{t}}(0, x)| = \mathbb{E}[|\mathfrak{u}_{\mathfrak{t}}(t, x + W_{0}) - \mathfrak{u}_{\mathfrak{t}}(0, x + W_{0})|] \\ \leq e^{2LT}(T+1)^{2}(L+1)(\mathfrak{L}+1)8^{p+2}(1+||x||^{p} + \sup_{s \in [0,T]} \mathbb{E}[||W_{s}||^{2p}])\sqrt{t}\sqrt{d}.$$

$$(2.54)$$

This establishes that for all $\mathfrak{t} \in [0, T]$, $t \in [\mathfrak{t}, T]$, $x \in \mathbb{R}^d$ it holds that

$$|u(t,x) - u(\mathfrak{t},x)| \le e^{2LT}(T+1)^2(L+1)(\mathfrak{L}+1)8^{p+2}(1+||x||^p + \sup_{s\in[0,T]}\mathbb{E}[||W_s||^{2p}])\sqrt{|t-\mathfrak{t}|}\sqrt{d}.$$
(2.55)

The proof of Corollary 2.7 is thus complete.

3 Artificial neural network (ANN) calculus

In our proofs of the ANN approximation results in Sections 4 and 5 and Theorem 1.1 in the introduction we make use of a suitable calculus for ANNs from the literature (cf., for example, [37, Section 2], [54, Section 1.3 and Chapter 2], and the references therein). In this section we recall the ingredients of this ANN calculus that we need in the later ANN approximation results of this work. The notions in this section can – often in a slightly modified form – be found in [54, Section 1.3 and Chapter 2] and [1, Section 2], for instance.

3.1 ANNs

Definition 3.1 (ANNs). We denote by \mathbf{N} the set given by

$$\mathbf{N} = \bigcup_{L \in \mathbb{N}} \bigcup_{l_0, l_1, \dots, l_L \in \mathbb{N}} \left(\times_{k=1}^L (\mathbb{R}^{l_k \times l_{k-1}} \times \mathbb{R}^{l_k}) \right), \tag{3.1}$$

for every $L \in \mathbb{N}$, $l_0, l_1, \ldots, l_L \in \mathbb{N}$, $\Phi \in (\times_{k=1}^L (\mathbb{R}^{l_k \times l_{k-1}} \times \mathbb{R}^{l_k})) \subseteq \mathbf{N}$ we denote by $\mathcal{P}(\Phi), \mathcal{L}(\Phi)$, $\mathcal{I}(\Phi), \mathcal{O}(\Phi), \mathcal{H}(\Phi) \in \mathbb{R}$ the numbers given by

$$\mathcal{P}(\Phi) = \sum_{k=1}^{L} l_k (l_{k-1} + 1), \qquad \mathcal{L}(\Phi) = L, \qquad \mathcal{I}(\Phi) = l_0, \qquad \mathcal{O}(\Phi) = l_L, \qquad (3.2)$$

and $\mathcal{H}(\Phi) = L - 1$, for every $n \in \mathbb{N}_0, L \in \mathbb{N}, l_0, l_1, \dots, l_L \in \mathbb{N}, \Phi \in (\times_{k=1}^L (\mathbb{R}^{l_k \times l_{k-1}} \times \mathbb{R}^{l_k})) \subseteq \mathbb{N}$ we denote by $\mathbb{D}_n(\Phi) \in \mathbb{R}$ the number given by

$$\mathbb{D}_n(\Phi) = \begin{cases} l_n & : n \le L\\ 0 & : n > L, \end{cases}$$
(3.3)

for every $\Phi \in \mathbf{N}$ we denote by $\mathcal{D}(\Phi) \in \mathbb{R}^{\mathcal{L}(\Phi)+1}$ the vector given by

$$\mathcal{D}(\Phi) = (\mathbb{D}_0(\Phi), \mathbb{D}_1(\Phi), \dots, \mathbb{D}_{\mathcal{L}(\Phi)}(\Phi)), \tag{3.4}$$

and for every $L \in \mathbb{N}$, $l_0, l_1, \ldots, l_L \in \mathbb{N}$, $\Phi = ((W_1, B_1), \ldots, (W_L, B_L)) \in (\times_{k=1}^L (\mathbb{R}^{l_k \times l_{k-1}} \times \mathbb{R}^{l_k})) \subseteq \mathbb{N}$, $n \in \{1, 2, \ldots, L\}$ we denote by $\mathcal{W}_{n,\Phi} \in \mathbb{R}^{l_n \times l_{n-1}}$, $\mathcal{B}_{n,\Phi} \in \mathbb{R}^{l_n}$ the matrix and the vector given by

$$\mathcal{W}_{n,\Phi} = W_n \quad \text{and} \quad \mathcal{B}_{n,\Phi} = B_n.$$
 (3.5)

Definition 3.2 (ANN). We say that Φ is an ANN if and only if it holds that $\Phi \in \mathbf{N}$ (cf. Definition 3.1).

3.2 Realizations of ANNs

Definition 3.3 (Multidimensional version). Let $a \colon \mathbb{R} \to \mathbb{R}$ be a function and let $d \in \mathbb{N}$. Then we denote by $\mathfrak{M}_{a,d} \colon \mathbb{R}^d \to \mathbb{R}^d$ the function which satisfies for all $x = (x_1, \ldots, x_d) \in \mathbb{R}^d$ that

$$\mathfrak{M}_{a,d}(x) = (a(x_1), \dots, a(x_d)). \tag{3.6}$$

Definition 3.4 (Realization associated to an ANN). Let $a \colon \mathbb{R} \to \mathbb{R}$ be a function and let $\Phi \in \mathbb{N}$ (cf. Definition 3.1). Then we denote by $\mathcal{R}_a(\Phi) \colon \mathbb{R}^{\mathcal{I}(\Phi)} \to \mathbb{R}^{\mathcal{O}(\Phi)}$ the function which satisfies for all $x_0 \in \mathbb{R}^{\mathbb{D}_0(\Phi)}$, $x_1 \in \mathbb{R}^{\mathbb{D}_1(\Phi)}$, ..., $x_{\mathcal{L}(\Phi)} \in \mathbb{R}^{\mathbb{D}_{\mathcal{L}(\Phi)}(\Phi)}$ with $\forall k \in \{1, 2, ..., \mathcal{L}(\Phi)\} \colon x_k = \mathfrak{M}_{a_{\mathbb{I}_{(0,\mathcal{L}(\Phi))}(k)+\mathrm{id}_{\mathbb{R}}\mathbb{I}_{\{\mathcal{L}(\Phi)\}}(k), \mathbb{D}_k(\Phi)}(\mathcal{W}_{k,\Phi}x_{k-1} + \mathcal{B}_{k,\Phi})$ that

$$(\mathcal{R}_a(\Phi))(x_0) = x_{\mathcal{L}(\Phi)} \tag{3.7}$$

(cf. Definition 3.3).

3.3 Activation ANNs

Definition 3.5 (Identity matrices). Let $d \in \mathbb{N}$. Then we denote by $I_d \in \mathbb{R}^{d \times d}$ the identity matrix in $\mathbb{R}^{d \times d}$.

Definition 3.6 (Activation ANNs). Let $d \in \mathbb{N}$. Then we denote by $\mathbf{i}_d \in ((\mathbb{R}^{d \times d} \times \mathbb{R}^d) \times (\mathbb{R}^{d \times d} \times \mathbb{R}^d)) \subseteq \mathbf{N}$ the ANN given by $\mathbf{i}_d = ((\mathbf{I}_d, 0), (\mathbf{I}_d, 0))$ (cf. Definitions 3.1 and 3.5).

In Lemma 4.5 in Section 4 below we establish an elementary representation result for hat functions in terms of ANNs with the leaky ReLU activation functions. We employ Lemma 4.5 in our proofs of the ANN approximation results for PDEs in Section 5 and Theorem 1.1 in the introduction, respectively. Our proof of Lemma 4.5, in turn, is based on the elementary fact that hat functions (and other piecewise linear functions) can be exactly represented by ANNs with the ReLU activation function (cf., for example, (4.10) and [1, Lemma 4.10]) and the elementary fact that multidimensional versions of the ReLU activation function can be exactly represented by leaky ReLU ANNs. These representations of multidimensional versions of the ReLU activation through leaky ReLU ANNs are precisely the subject of the following elementary result, Lemma 3.7 below. Our statement of Lemma 3.7 employs the notion of the activation ANNs in Definition 3.6 above. In this context we note that the realization function function under consideration (cf., for example, [54, Lemma 3.2.2]).

Lemma 3.7. Let $d \in \mathbb{N}$, $a \in \mathbb{R} \setminus \{-1, 1\}$ and let $a \colon \mathbb{R} \to \mathbb{R}$ and $\mathfrak{r} \colon \mathbb{R} \to \mathbb{R}$ satisfy for all $x \in \mathbb{R}$ that $a(x) = \max\{x, ax\}$ and $\mathfrak{r}(x) = \max\{x, 0\}$. Then it holds for all $x \in \mathbb{R}^d$ that

$$\left(\mathcal{R}_{\mathfrak{r}}(\mathfrak{i}_d)\right)(x) = \frac{|1-a|}{(1-a)(1-a^2)} \left[a\left(\mathcal{R}_a(\mathfrak{i}_d)\right)\left(\frac{-|1-a|x}{1-a}\right) + \left(\mathcal{R}_a(\mathfrak{i}_d)\right)\left(\frac{|1-a|x}{1-a}\right)\right] \quad (3.8)$$

(cf. Definitions 3.4 and 3.6).

Proof of Lemma 3.7. Observe that, for instance, item (iii) in Lemma 3.2 in [1] establishes that $\mathcal{R}_{\mathfrak{r}}(\mathfrak{i}_d) = \mathfrak{M}_{\mathfrak{r},d}$ and $\mathcal{R}_a(\mathfrak{i}_d) = \mathfrak{M}_{a,d}$ (cf. Definitions 3.3, 3.4, and 3.6). In the next step we note that the fact that for all $x, y \in \mathbb{R}$ it holds that $\max\{x, y\} = \frac{1}{2}(x + y + |x - y|)$ ensures that for all $x \in \mathbb{R}$ it holds that

$$2a\left(\frac{|1-a|x}{1-a}\right) = \frac{|1-a|x}{1-a} + \frac{|1-a|ax}{1-a} + \left|\frac{|1-a|x}{1-a} - \frac{|1-a|ax}{1-a}\right|$$

$$= \frac{|1-a|(1+a)x}{1-a} + |1-a||x|.$$
(3.9)

Therefore, we obtain for all $x \in \mathbb{R}$ that

$$aa\left(\frac{-|1-a|x}{1-a}\right) + a\left(\frac{|1-a|x}{1-a}\right)$$

= $\frac{1}{2}\left(\frac{-a|1-a|(1+a)x}{1-a} + a|1-a||-x| + \frac{|1-a|(1+a)x}{1-a} + |1-a||x|\right)$
= $\frac{1}{2}\left(\frac{(1-a)|1-a|(1+a)x}{1-a} + (1+a)|1-a||x|\right)$
= $\frac{|1-a|(1+a)(x+|x|)}{2}.$ (3.10)

Hence, we obtain for all $x \in \mathbb{R}$ that

$$\frac{|1-a|}{(1-a)(1-a^2)} \left[aa \left(\frac{-|1-a|x}{1-a} \right) + a \left(\frac{|1-a|x}{1-a} \right) \right] = \frac{|1-a|}{(1-a)(1-a)(1+a)} \frac{|1-a|(1+a)(x+|x|)}{2} = \frac{x+|x|}{2} = \max\{x,0\} = \mathfrak{r}(x).$$
(3.11)

Combining this with the fact that $\mathcal{R}_{\mathfrak{r}}(\mathfrak{i}_d) = \mathfrak{M}_{\mathfrak{r},d}$ and the fact that $\mathcal{R}_a(\mathfrak{i}_d) = \mathfrak{M}_{a,d}$ establishes (3.8). The proof of Lemma 3.7 is thus complete.

3.4 Compositions, powers, and extensions of ANNs

Definition 3.8 (Composition of ANNs). Let $\Phi, \Psi \in \mathbf{N}$ satisfy $\mathcal{I}(\Phi) = \mathcal{O}(\Psi)$ (cf. Definition 3.1). Then we denote by $\Phi \bullet \Psi \in \mathbf{N}$ the ANN which satisfies for all $k \in \mathbb{N} \cap (0, \mathcal{L}(\Phi) + \mathcal{L}(\Psi))$ that $\mathcal{L}(\Phi \bullet \Psi) = \mathcal{L}(\Phi) + \mathcal{L}(\Psi) - 1$ and

$$(\mathcal{W}_{k,\Phi\bullet\Psi},\mathcal{B}_{k,\Phi\bullet\Psi}) = \begin{cases} (\mathcal{W}_{k,\Psi},\mathcal{B}_{k,\Psi}) & : k < \mathcal{L}(\Psi) \\ (\mathcal{W}_{1,\Phi}\mathcal{W}_{\mathcal{L}(\Psi),\Psi},\mathcal{W}_{1,\Phi}\mathcal{B}_{\mathcal{L}(\Psi),\Psi} + \mathcal{B}_{1,\Phi}) & : k = \mathcal{L}(\Psi) \\ (\mathcal{W}_{k-\mathcal{L}(\Psi)+1,\Phi},\mathcal{B}_{k-\mathcal{L}(\Psi)+1,\Phi}) & : k > \mathcal{L}(\Psi). \end{cases}$$
(3.12)

Definition 3.9 (Affine transformation ANNs). Let $m, n \in \mathbb{N}, W \in \mathbb{R}^{m \times n}, B \in \mathbb{R}^m$. Then we denote by $\mathbf{A}_{W,B} \in (\mathbb{R}^{m \times n} \times \mathbb{R}^m) \subseteq \mathbf{N}$ the ANN given by $\mathbf{A}_{W,B} = (W, B)$ (cf. Definition 3.1).

Definition 3.10 (Powers of ANNs). Let $\Phi \in \mathbf{N}$ satisfy $\mathcal{I}(\Phi) = \mathcal{O}(\Phi)$ (cf. Definition 3.1). Then we denote by $\Phi^{\bullet n} \in \{\Psi \in \mathbf{N} : \mathcal{I}(\Psi) = \mathcal{O}(\Psi) = \mathcal{I}(\Phi)\}, n \in \mathbb{N}_0$, the ANNs which satisfy for all $n \in \mathbb{N}_0$ that

$$\Phi^{\bullet n} = \begin{cases} \mathbf{A}_{\mathbf{I}_{\mathcal{O}(\Phi)},0} & : n = 0\\ \Phi \bullet (\Phi^{\bullet(n-1)}) & : n > 0 \end{cases}$$
(3.13)

(cf. Definitions 3.5, 3.8, and 3.9).

Definition 3.11 (Extensions of ANNs). Let $L \in \mathbb{N}$, $\Phi, \Psi \in \mathbb{N}$ satisfy $\mathcal{L}(\Phi) \leq L$ and $\mathcal{O}(\Phi) = \mathcal{I}(\Psi) = \mathcal{O}(\Psi)$ (cf. Definition 3.1). Then we denote by $\mathcal{E}_{L,\Psi}(\Phi) \in \mathbb{N}$ the ANN given by

$$\mathcal{E}_{L,\Psi}(\Phi) = (\Psi^{\bullet(L-\mathcal{L}(\Phi))}) \bullet \Phi$$
(3.14)

(cf. Definitions 3.8 and 3.10).

3.5 Parallelizations of ANNs

Definition 3.12 (Parallelization of ANNs with the same length). Let $n \in \mathbb{N}$, $\Phi = (\Phi_1, \ldots, \Phi_n) \in \mathbb{N}^n$ satisfy $\mathcal{L}(\Phi_1) = \mathcal{L}(\Phi_2) = \ldots = \mathcal{L}(\Phi_n)$ (cf. Definition 3.1). Then we denote by $\mathbf{P}_n(\Phi) \in \mathbb{N}$ the ANN which satisfies that $\mathcal{L}(\mathbf{P}_n(\Phi)) = \mathcal{L}(\Phi_1)$ and that for all $k \in \{1, 2, \ldots, \mathcal{L}(\Phi_1)\}$ it

holds that

$$\mathcal{W}_{k,\mathbf{P}_{n}(\Phi)} = \begin{pmatrix}
\mathcal{W}_{k,\Phi_{1}} & 0 & 0 & \cdots & 0 \\
0 & \mathcal{W}_{k,\Phi_{2}} & 0 & \cdots & 0 \\
0 & 0 & \mathcal{W}_{k,\Phi_{3}} & \cdots & 0 \\
\vdots & \vdots & \vdots & \ddots & \vdots \\
0 & 0 & 0 & \cdots & \mathcal{W}_{k,\Phi_{n}}
\end{pmatrix} \quad \text{and} \quad \mathcal{B}_{k,\mathbf{P}_{n}(\Phi)} = \begin{pmatrix}
\mathcal{B}_{k,\Phi_{1}} \\
\mathcal{B}_{k,\Phi_{2}} \\
\vdots \\
\mathcal{B}_{k,\Phi_{n}}
\end{pmatrix}. \quad (3.15)$$

Definition 3.13 (Parallelization of ANNs with different lengths). Let $n \in \mathbb{N}$, $\Psi \in \mathbb{N}$, $\Phi = (\Phi_1, \ldots, \Phi_n) \in \mathbb{N}^n$ satisfy $\mathcal{H}(\Psi) = 1$ and $\mathcal{O}(\Phi_1) = \ldots = \mathcal{O}(\Phi_n) = \mathcal{I}(\Psi) = \mathcal{O}(\Psi)$ (cf. Definition 3.1). Then we denote by $P_{n,\Psi}(\Phi) \in \mathbb{N}$ the ANN given by

$$P_{n,\Psi}(\Phi) = \mathbf{P}_n \left(\mathcal{E}_{\max_{k \in \{1,2,\dots,n\}} \mathcal{L}(\Phi_k),\Psi}(\Phi_1), \dots, \mathcal{E}_{\max_{k \in \{1,2,\dots,n\}} \mathcal{L}(\Phi_k),\Psi}(\Phi_n) \right)$$
(3.16)

(cf. Definitions 3.11 and 3.12).

In our ANN approximation results in Section 5 we reformulate the PDE under consideration from a terminal value PDE problem (as PDEs are often formulated in the finance/stochastic analysis literature) to an initial value PDE problem (as PDEs are often presented in the physics literature). In particular, in Corollary 5.4 in Section 5 the PDE approximation problem is formulated as a terminal value PDE problem in which the terminal value function of the PDE is considered to be (explicitly representable or) approximable by ANNs without the COD (see (5.75) and (5.78) in Corollary 5.4 for details) and in Corollary 5.5 in Section 5 the PDE approximation problem is formulated as an initial value PDE problem in which the initial value function of the PDE is considered to be (explicitly representable or) approximable by ANNs without the COD (see (5.102) and (5.104)in Corollary 5.5 for details). In the next elementary result, Lemma 3.14 below, we provide a suitable elementary transformation result for ANN approximations that allows us to suitably shift/transform the temporal variable of the realization functions of space-time ANN approximations. In our proof of Corollary 5.5 we apply Lemma 3.14 in conjunction with the ANN approximation result in Corollary 5.4 to establish the ANN approximation result in Corollary 5.5. Our proof of Lemma 3.14, in turn, is based on applications of appropriate ANN calculus results in the literature (cf., for instance, [1, Section 2.1], [37, Sections 2.2] and 2.3, and 54, Section 2.2).

Lemma 3.14. Let $T, c \in \mathbb{R}$, $d, \mathfrak{d} \in \mathbb{N}$, $a \in C(\mathbb{R}, \mathbb{R})$, $\mathbf{F}, \mathfrak{J}, \mathbf{G} \in \mathbf{N}$ satisfy $\mathcal{D}(\mathfrak{J}) = (1, \mathfrak{d}, 1)$, $\mathcal{R}_a(\mathfrak{J}) = \mathrm{id}_{\mathbb{R}}, \mathcal{R}_a(\mathbf{F}) \in C(\mathbb{R}^{d+1}, \mathbb{R})$, and

$$\mathbf{G} = \mathbf{F} \bullet \mathbf{P}_{d+1,\mathfrak{J}} \big(\mathbf{A}_{c,T}, \mathfrak{J}, \mathfrak{J}, \dots, \mathfrak{J} \big)$$
(3.17)

(cf. Definitions 3.1, 3.4, 3.8, 3.9, and 3.13). Then

- (i) it holds that $\mathcal{R}_a(\mathbf{G}) \in C(\mathbb{R}^{d+1}, \mathbb{R})$,
- (ii) it holds for all $s \in \mathbb{R}$, $x \in \mathbb{R}^d$ that $(\mathcal{R}_a(\mathbf{G}))(s, x) = (\mathcal{R}_a(\mathbf{F}))(T + cs, x)$, and
- (iii) it holds that $\mathcal{P}(\mathbf{G}) \leq \mathcal{P}(\mathbf{F}) (1 + 32\mathfrak{d}^2 d^2 + 2\mathfrak{d} d) \leq 96\mathfrak{d}^2 d^2 \mathcal{P}(\mathbf{F}).$

Proof of Lemma 3.14. Throughout this proof let $\Phi \in \mathbf{N}$ satisfy

$$\Phi = \mathcal{P}_{d+1,\mathfrak{J}} \big(\mathbf{A}_{c,T}, \mathfrak{J}, \mathfrak{J}, \dots, \mathfrak{J} \big)$$
(3.18)

(cf. Definitions 3.1, 3.9, and 3.13). Observe that (3.12), (3.13), (3.14), (3.16), and (3.18) show that

$$\Phi = \mathbf{P}_{d+1}(\mathcal{E}_{2,\mathfrak{J}}(\mathbf{A}_{c,T}),\mathfrak{J},\mathfrak{J},\ldots,\mathfrak{J})$$
(3.19)

(cf. Definitions 3.11 and 3.12). Next note that, for example, item (i) in Corollary 2.23 in [37] and, for instance, item (vi) in Proposition 2.6 in [37] establish item (i). Observe that, for example, item (vi) in Proposition 2.6 in [37] implies that for all $s \in \mathbb{R}$, $x \in \mathbb{R}^d$ it holds that

$$\left(\mathcal{R}_{a}(\mathbf{G})\right)(s,x) = \left(\left(\mathcal{R}_{a}(\mathbf{F})\right) \circ \left(\mathcal{R}_{a}(\Phi)\right)\right)(s,x)$$
(3.20)

(cf. Definition 3.4). Furthermore, note that the fact that $\mathcal{R}_a(\mathfrak{J}) = \mathrm{id}_{\mathbb{R}}$ and, for instance, item (ii) in Corollary 2.23 in [37] show that for all $s \in \mathbb{R}$, $x \in \mathbb{R}^d$ it holds that

$$\left(\mathcal{R}_{a}(\Phi)\right)(s,x) = \left(\left(\mathcal{R}_{a}\left(\mathbf{A}_{c,T}\right)\right)(s),x\right) = (cs+T,x).$$
(3.21)

This and (3.20) prove item (ii). Observe that, for example, item (v) in Proposition 2.6 in [37] shows that

$$\mathcal{P}(\mathbf{G}) \le \mathcal{P}(\mathbf{F}) + \mathcal{P}(\Phi) + \mathbb{D}_{1}(\mathbf{F}) \cdot \mathbb{D}_{\mathcal{L}(\Phi)-1}(\Phi).$$
(3.22)

Moreover, note that (3.19) and, for instance, [37, Proposition 2.20] prove that

$$\mathcal{L}(\Phi) - 1 = 1, \qquad \mathbb{D}_1(\Phi) = \mathbb{D}_1\left(\mathcal{E}_{2,\mathfrak{J}}(\mathbf{A}_{c,T})\right) + \sum_{k=1}^d \mathbb{D}_1(\mathfrak{J}), \qquad (3.23)$$

and
$$\mathcal{P}(\Phi) \leq \frac{1}{2} \left[\mathcal{P}(\mathcal{E}_{2,\mathfrak{J}}(\mathbf{A}_{c,T})) + \sum_{k=1}^{d} \mathcal{P}(\mathfrak{J}) \right]^{2}.$$
 (3.24)

Furthermore, observe that, for example, [54, Lemma 2.2.11] demonstrates that

$$\mathbb{D}_1\big(\mathcal{E}_{2,\mathfrak{J}}\big(\mathbf{A}_{c,T}\big)\big) = \mathbb{D}_1(\mathfrak{J}). \tag{3.25}$$

Combining this, (3.23), and the fact that $\mathcal{D}(\mathfrak{J}) = (1, \mathfrak{d}, 1)$ establishes that

$$\mathbb{D}_1(\Phi) = (d+1)\mathfrak{d} \le 2\mathfrak{d}d. \tag{3.26}$$

Moreover, note that, for instance, item (ii) in Lemma 2.13 in [37] ensures that

$$\mathcal{P}(\mathcal{E}_{2,\mathfrak{J}}(\mathbf{A}_{c,T})) \leq \mathfrak{d}\mathcal{P}(\mathbf{A}_{c,T}) + \mathfrak{d} + 1 = 2\mathfrak{d} + \mathfrak{d} + 1 \leq 4\mathfrak{d}.$$
(3.27)

Next, observe that the fact that $\mathcal{D}(\mathfrak{J}) = (1, \mathfrak{d}, 1)$ implies that

$$\mathcal{P}(\mathfrak{J}) = 2\mathfrak{d} + \mathfrak{d} + 1 \le 4\mathfrak{d}. \tag{3.28}$$

This, (3.24), and (3.27) show that

$$\mathcal{P}(\Phi) \le \frac{1}{2} \left(4\mathfrak{d} + 4\mathfrak{d} d \right)^2 \le 32\mathfrak{d}^2 d^2.$$
(3.29)

Combining this, (3.22), (3.26), and, for example, [1, Lemma 2.4] proves that

$$\mathcal{P}(\mathbf{G}) \le \mathcal{P}(\mathbf{F}) + 32\mathfrak{d}^2 d^2 + \mathbb{D}_1(\mathbf{F}) \cdot 2\mathfrak{d}d \le \mathcal{P}(\mathbf{F}) (1 + 32\mathfrak{d}^2 d^2 + 2\mathfrak{d}d).$$
(3.30)

This demonstrates item (iii). The proof of Lemma 3.14 is thus complete.

3.6 Scalar multiplications and sums of ANNs

Definition 3.15 (Scalar multiplications of ANNs). Let $\lambda \in \mathbb{R}$, $\Phi \in \mathbb{N}$ (cf. Definition 3.1). Then we denote by $\lambda \circledast \Phi \in \mathbb{N}$ the ANN given by

$$\lambda \circledast \Phi = \mathbf{A}_{\lambda \mathbf{I}_{\mathcal{O}(\Phi)}, 0} \bullet \Phi \tag{3.31}$$

(cf. Definitions 3.5, 3.8, and 3.9).

Definition 3.16 (Summation ANNs). Let $m, n \in \mathbb{N}$. Then we denote by $\mathfrak{S}_{m,n} \in (\mathbb{R}^{m \times (nm)} \times \mathbb{R}^m)$ the ANN given by $\mathfrak{S}_{m,n} = \mathbf{A}_{(\mathbf{I}_m \ \mathbf{I}_m \ \dots \ \mathbf{I}_m),0}$ (cf. Definitions 3.5 and 3.9).

Definition 3.17 (Transpose of matrices). Let $m, n \in \mathbb{N}$, $A \in \mathbb{R}^{m \times n}$. Then we denote by $A^* \in \mathbb{R}^{n \times m}$ the transpose of A.

Definition 3.18 (Vectorization ANNs). Let $m, n \in \mathbb{N}$. Then we denote by $\mathfrak{T}_{m,n} \in (\mathbb{R}^{(nm) \times m} \times \mathbb{R}^{nm})$ the ANN given by $\mathfrak{T}_{m,n} = \mathbf{A}_{(\mathbf{I}_m \ \mathbf{I}_m \ \dots \ \mathbf{I}_m)^*,0}$ (cf. Definitions 3.5, 3.9, and 3.17).

Definition 3.19 (Sums of ANNs with the same length). Let $u \in \mathbb{Z}$, $v \in \mathbb{Z} \cap [u, \infty)$, $\Phi_u, \Phi_{u+1}, \ldots, \Phi_v \in \mathbb{N}$ satisfy for all $k \in \mathbb{Z} \cap [u, v]$ that $\mathcal{L}(\Phi_k) = \mathcal{L}(\Phi_u)$, $\mathcal{I}(\Phi_k) = \mathcal{I}(\Phi_u)$, and $\mathcal{O}(\Phi_k) = \mathcal{O}(\Phi_u)$ (cf. Definition 3.1). Then we denote by $\bigoplus_{k=u}^v \Phi_k$ (we denote by $\Phi_u \oplus \Phi_{u+1} \oplus \ldots \oplus \Phi_v$) the ANN given by

$$\overset{v}{\underset{k=u}{\oplus}} \Phi_k = \left(\mathfrak{S}_{\mathcal{O}(\Phi_u), v-u+1} \bullet \left[\mathbf{P}_{v-u+1}(\Phi_u, \Phi_{u+1}, \dots, \Phi_v) \right] \bullet \mathfrak{T}_{\mathcal{I}(\Phi_u), v-u+1} \right) \in \mathbf{N}$$
(3.32)

(cf. Definitions 3.8, 3.12, 3.16, and 3.18).

Definition 3.20 (Sums of ANNs with different lengths). Let $u \in \mathbb{Z}$, $v \in \mathbb{Z} \cap [u, \infty)$, Φ_u , $\Phi_{u+1}, \ldots, \Phi_v, \Psi \in \mathbb{N}$ satisfy for all $k \in \mathbb{Z} \cap [u, v]$ that $\mathcal{I}(\Phi_k) = \mathcal{I}(\Phi_u)$, $\mathcal{O}(\Phi_k) = \mathcal{I}(\Psi) = \mathcal{O}(\Psi)$, and $\mathcal{H}(\Psi) = 1$ (cf. Definition 3.1). Then we denote by $\boxplus_{k=u,\Psi}^v \Phi_k$ (we denote by $\Phi_u \boxplus_\Psi \Phi_{u+1} \boxplus_\Psi \ldots \boxplus_\Psi \Phi_v$) the ANN given by

$$\underset{k=u,\Psi}{\overset{v}{\boxplus}} \Phi_k = \left[\underset{k=u}{\overset{v}{\oplus}} \mathcal{E}_{\max_{j \in \{u,u+1,\dots,v\}} \mathcal{L}(\Phi_j),\Psi}(\Phi_k) \right] \in \mathbf{N}$$
(3.33)

(cf. Definitions 3.11 and 3.19).

In the following elementary result, Lemma 3.21 below, we collect a few basic properties for sums of ANNs with different lengths (see Definition 3.20 above). Lemma 3.21 is a direct consequence of [1, Lemma 2.20] and [54, Lemma 2.2.11], for example.

Lemma 3.21 (Elementary properties of sums of ANNs with different lengths). Let $a \in C(\mathbb{R}, \mathbb{R})$, $L \in \mathbb{N}$, $u \in \mathbb{Z}$, $v \in \mathbb{Z} \cap [u, \infty)$, $\Phi_u, \Phi_{u+1}, \ldots, \Phi_v, \mathfrak{J}, \mathbf{G} \in \mathbf{N}$ satisfy for all $k \in \mathbb{Z} \cap [u, v]$ that $L = \max_{k \in \mathbb{Z} \cap [u, v]} \mathcal{L}(\Phi_k)$, $\mathcal{I}(\Phi_k) = \mathcal{I}(\Phi_u)$, $\mathcal{O}(\Phi_k) = \mathcal{I}(\mathfrak{J}) = \mathcal{O}(\mathfrak{J})$, $\mathcal{H}(\mathfrak{J}) = 1$, $\mathcal{R}_a(\mathfrak{J}) = \mathrm{id}_{\mathbb{R}}$, and $\mathbf{G} = \bigoplus_{k=u,\mathfrak{J}}^v \Phi_k$ (cf. Definitions 3.1, 3.4, and 3.20). Then

(i) it holds that $\mathcal{L}(\mathbf{G}) = L$,

(*ii*) it holds that

$$\mathcal{D}(\mathbf{G})$$

$$= \left(\mathcal{I}(\Phi_u), \sum_{k=u}^{v} \mathbb{D}_1 \left(\mathcal{E}_{L,\mathfrak{J}}(\Phi_k) \right), \sum_{k=u}^{v} \mathbb{D}_2 \left(\mathcal{E}_{L,\mathfrak{J}}(\Phi_k) \right), \dots, \sum_{k=u}^{v} \mathbb{D}_{L-1} \left(\mathcal{E}_{L,\mathfrak{J}}(\Phi_k) \right), \mathcal{O}(\Phi_u) \right),$$
(3.34)

(iii) it holds that

$$\||\mathcal{D}(\mathbf{G})|\| \le (v-u+1) \max\Big\{\mathbb{D}_1(\mathfrak{J}), \max_{k \in \mathbb{Z} \cap [u,v]} |||\mathcal{D}(\Phi_k)|||\Big\},$$
(3.35)

- (iv) it holds that $\mathcal{R}_a(\mathbf{G}) \in C(\mathbb{R}^{\mathcal{I}(\Phi_u)}, \mathbb{R}^{\mathcal{O}(\Phi_u)})$, and
- (v) it holds for all $x \in \mathbb{R}^{\mathcal{I}(\Phi_u)}$ that

$$(\mathcal{R}_a(\mathbf{G}))(x) = \sum_{k=u}^{v} (\mathcal{R}_a(\Phi_k))(x)$$
(3.36)

(cf. Definitions 2.1 and 3.11).

Proof of Lemma 3.14. Note that, for instance, [1, Lemma 2.20] (applied with $\psi \curvearrowleft \mathbf{G}$, $(h_k)_{k \in \mathbb{Z} \cap [u,v]} \curvearrowleft (1)_{k \in \mathbb{Z} \cap [u,v]}, (B_k)_{k \in \mathbb{Z} \cap [u,v]} \backsim (0)_{k \in \mathbb{Z} \cap [u,v]}$ in the notation of [1, Lemma 2.20]), (3.12), and (3.31) establish items (ii), (iv), and (v). Observe that item (ii) implies item (i). Note that, for example, [54, Lemma 2.2.11] establishes that for all $k \in \mathbb{Z} \cap [u, v]$ it holds that

$$\||\mathcal{D}(\mathcal{E}_{L,\mathfrak{J}}(\Phi_k))|\| \le \max\{\mathbb{D}_1(\mathfrak{J}), \||\mathcal{D}(\Phi_k)\||\}$$
(3.37)

(cf. Definitions 2.1, 3.1, and 3.11). Furthermore, observe that item (ii) ensures that

$$\||\mathcal{D}(\mathbf{G})|\| \le \max\Big\{\mathcal{I}(\Phi_u), \mathcal{O}(\Phi_u), (v-u+1)\max_{k\in\mathbb{Z}\cap[u,v]}\||\mathcal{D}(\mathcal{E}_{L,\mathfrak{J}}(\Phi_k))\||\Big\}.$$
(3.38)

This and (3.38) imply item (iii). The proof of Lemma 3.14 is thus complete.

4 ANN approximations for linear interpolations of multilevel Picard (MLP) approximations

One of the main goals of this section is to construct and study in Corollary 4.13 below suitable ANNs (with general/abstract activations) which approximate linear interpolations of appropriate MLP approximations (see (4.84) in Corollary 4.13 in Subsection 4.6 below for details). We employ Corollary 4.13 in our proofs of the ANN approximation results for PDEs in Section 5 and Theorem 1.1 in the introduction, respectively. In our proof of Corollary 4.13 we employ the abstract ANN approximation result for interpolation functions in Proposition 4.12 and the ANN approximation result for MLP approximations at fixed time points in [1, Proposition 3.9]. Moreover, the statements and our proofs of Proposition 4.12 and Corollary 4.13 build up on the concepts and results of the ANN calculus from Section 3, [37, Sections 2.2 and 2.3], [1, Section 2.1], and [54, Section 2.2], respectively.

In Proposition 4.12 and Corollary 4.13 the activation function of the considered ANNs is a general continuous function $a: \mathbb{R} \to \mathbb{R}$ which fulfills, among other assumptions, the condition that it makes the class of ANNs with this activation function flexible enough to exactly represent the one-dimensional identity function $\mathrm{id}_{\mathbb{R}} = (\mathbb{R} \ni x \mapsto x \in \mathbb{R})$ and to approximately represent the product function $\mathbb{R}^2 \ni (v, w) \mapsto vw \in \mathbb{R}$ (see (4.56) in Proposition 4.12 and (4.79) in Corollary 4.13 for details) in a suitable way. In the ANN approximation results in Lemma 4.7, Lemma 4.8, Corollary 4.9, Corollary 4.10 (ReLU and leaky ReLU activations), and Corollary 4.11 (softplus activation) in Subsections 4.3 and 4.4 below we verify that the assumption in Corollary 4.13 that the considered ANNs can approximately represent the product function in a suitable way is satisfied in the situation of the ReLU activation function $\mathfrak{r} = (\mathbb{R} \ni x \mapsto \max\{x, 0\} \in \mathbb{R})$, in the situation of the leaky ReLU activation functions $\mathbb{R} \ni x \mapsto \max\{x, ax\} \in \mathbb{R}$ for $a \in (0, 1)$, and in the situation of the softplus activation function $\mathbb{R} \ni x \mapsto \ln(1 + \exp(x)) \in \mathbb{R}$. Lemma 4.7 (an appropriate ANN approximation result for the square function $\mathbb{R} \ni x \mapsto x^2 \in \mathbb{R}$) is an extension of Grohs et al. [37, Proposition 3.4] in which a result similar to Lemma 4.7 has been established in the special situation of the ReLU activation function and Lemma 4.8 (a suitable ANN approximation result for the product function $\mathbb{R}^2 \ni (v, w) \mapsto vw \in \mathbb{R}$) is an extension of Grohs et al. [37, Proposition 3.5] in which a result similar to Lemma 4.8 has been established in the special situation of the ReLU activation function. Our proofs of Lemma 4.7 and Lemma 4.8 are strongly based on the proofs in Grohs et al. [37, Proposition 3.4 and Proposition 3.5].

In Subsection 4.1 we recall in Definition 4.1 (which coincides with Definition 4.5 in [1]), roughly speaking, the concept of a continuous piecewise linear (more accurately, piecewise affine) function which interpolates certain given values at certain given points/arguments/ positions while being affine on the intervals between two neighboring position points and in Lemma 4.2, Lemma 4.3, and Lemma 4.4 we collect a few elementary and well-known properties of such linear interpolation functions. In (4.5) in Lemma 4.4 we recall that such a linear interpolation function can be written as a linear combination of hat functions: the left hand side of (4.5) is the considered linear interpolation function from Definition 4.1 and the right hand side of (4.5) is the linear combination of the hat functions. The linear combination on the right hand side of (4.5) consists of a sum of (K + 1) summands (where $K \in \mathbb{N}$ is an arbitrary natural number) consisting of coefficients (real numbers) multiplied with appropriate hat functions. Roughly speaking, we apply (4.5) in Lemma 4.4 in the situation where the left hand side of (4.5) are linear interpolations of MLP approximations and where we then want to approximate the right hand side of (4.5) through ANN approximations to thereby obtain ANN approximations for MLP approximations.

Taking this into account, in Lemma 4.5 (ReLU and leaky ReLU activations) and Lemma 4.6 (softplus activation) in Subsection 4.2 we study ANN approximations for hat functions (which appear on the right hand side of (4.5)) and in Lemma 4.7, Lemma 4.8, Corollary 4.9, Corollary 4.10 (ReLU and leaky ReLU activations), and Corollary 4.11 (softplus activation) in Subsections 4.3 and 4.4 we study ANN approximations for the production function $\mathbb{R}^2 \ni (v, w) \mapsto vw \in \mathbb{R}$ (which appears on the right hand side of (4.5) to present the products (the multiplications) of the coefficients with the hat functions on the right hand side of (4.5)). In the situation of the ReLU activation a result similar to Lemma 4.5 has been established in Grohs et al. [37, Lemma 3.9]. Our overall approach in this section to employ (4.5) in Lemma 4.4 to approximate suitable linear interpolations through ANN approximations is

strongly inspired by the approach in Grohs et al. [37, Section 3].

4.1 Properties of linear interpolation functions

Definition 4.1 (Linear interpolation function). Let $K \in \mathbb{N}$, $\mathfrak{t}_0, \mathfrak{t}_1, \ldots, \mathfrak{t}_K, f_0, f_1, \ldots, f_K \in \mathbb{R}$ satisfy $\mathfrak{t}_0 < \mathfrak{t}_1 < \ldots < \mathfrak{t}_K$. Then we denote by $\mathscr{L}^{f_0, f_1, \ldots, f_K}_{\mathfrak{t}_0, \mathfrak{t}_1, \ldots, \mathfrak{t}_K} : \mathbb{R} \to \mathbb{R}$ the function which satisfies for all $k \in \{1, 2, \ldots, K\}$, $x \in (-\infty, \mathfrak{t}_0)$, $y \in [\mathfrak{t}_{k-1}, \mathfrak{t}_k)$, $z \in [\mathfrak{t}_K, \infty)$ that $\mathscr{L}^{f_0, f_1, \ldots, f_K}_{\mathfrak{t}_0, \mathfrak{t}_1, \ldots, \mathfrak{t}_K}(x) = f_0$, $\mathscr{L}^{f_0, f_1, \ldots, f_K}_{\mathfrak{t}_0, \mathfrak{t}_1, \ldots, \mathfrak{t}_K}(z) = f_K$, and

$$\mathscr{L}_{\mathfrak{t}_{0},\mathfrak{t}_{1},\ldots,\mathfrak{t}_{K}}^{f_{0},f_{1},\ldots,f_{K}}(y) = f_{k-1} + \left(\frac{y-\mathfrak{t}_{k-1}}{\mathfrak{t}_{k}-\mathfrak{t}_{k-1}}\right)(f_{k}-f_{k-1}).$$

$$(4.1)$$

Lemma 4.2. Let $K \in \mathbb{N}$, $\mathfrak{t}_0, \mathfrak{t}_1, \ldots, \mathfrak{t}_K, f_0, f_1, \ldots, f_K \in \mathbb{R}$ satisfy $\mathfrak{t}_0 < \mathfrak{t}_1 < \ldots < \mathfrak{t}_K$. Then it holds for all $k \in \{1, 2, \ldots, K\}$, $t \in [\mathfrak{t}_{k-1}, \mathfrak{t}_k]$ that

$$\left|\mathscr{L}_{\mathfrak{t}_{0},\mathfrak{t}_{1},\ldots,\mathfrak{t}_{K}}^{f_{0},f_{1},\ldots,f_{K}}(t) - f_{k}\right| \leq |f_{k} - f_{k-1}|$$
(4.2)

(cf. Definition 4.1).

Proof of Lemma 4.2. Note that (4.1) implies that for all $k \in \{1, 2, ..., K\}$, $t \in [\mathfrak{t}_{k-1}, \mathfrak{t}_k]$ it holds that

$$\mathscr{L}_{\mathfrak{t}_{0},\mathfrak{t}_{1},\ldots,\mathfrak{t}_{K}}^{f_{0},f_{1},\ldots,f_{K}}(t) - f_{k} = f_{k-1} + \left(\frac{t - \mathfrak{t}_{k-1}}{\mathfrak{t}_{k} - \mathfrak{t}_{k-1}}\right) \left(f_{k} - f_{k-1}\right) - f_{k} = \left(\frac{t - \mathfrak{t}_{k}}{\mathfrak{t}_{k} - \mathfrak{t}_{k-1}}\right) \left(f_{k} - f_{k-1}\right) \quad (4.3)$$

(cf. Definition 4.1). Combining this and the fact that for all $k \in \{1, 2, ..., K\}, t \in [\mathfrak{t}_{k-1}, \mathfrak{t}_k]$ it holds that $\left|\frac{t-\mathfrak{t}_k}{\mathfrak{t}_k-\mathfrak{t}_{k-1}}\right| \leq 1$ shows (4.2). The proof of Lemma 4.2 is thus complete.

Lemma 4.3. Let $\mathfrak{t}_0, \mathfrak{t}_1, \mathfrak{t}_2 \in \mathbb{R}$ satisfy $\mathfrak{t}_0 < \mathfrak{t}_1 < \mathfrak{t}_2$. Then it holds for all $t \in \mathbb{R}$ that

$$\mathscr{D}_{\mathfrak{t}_{0},\mathfrak{t}_{1},\mathfrak{t}_{2}}^{0,1,0}(t) = \frac{t-\mathfrak{t}_{0}}{\mathfrak{t}_{1}-\mathfrak{t}_{0}}\mathbb{1}_{(\mathfrak{t}_{0},\mathfrak{t}_{1}]}(t) + \frac{\mathfrak{t}_{2}-t}{\mathfrak{t}_{2}-\mathfrak{t}_{1}}\mathbb{1}_{(\mathfrak{t}_{1},\mathfrak{t}_{2})}(t)$$

$$(4.4)$$

(cf. Definition 4.1).

Proof of Lemma 4.3. Observe that (4.1) establishes (4.4). The proof of Lemma 4.3 is thus complete.

Lemma 4.4. Let $K \in \mathbb{N}$, $T \in (0, \infty)$, $\mathfrak{t}_{-1}, \mathfrak{t}_0, \mathfrak{t}_1, \ldots, \mathfrak{t}_K, \mathfrak{t}_{K+1}, f_0, f_1, \ldots, f_K \in \mathbb{R}$ satisfy $\mathfrak{t}_{-1} < 0 = \mathfrak{t}_0 < \mathfrak{t}_1 < \ldots < \mathfrak{t}_K = T < \mathfrak{t}_{K+1}$. Then it holds for all $t \in [0, T]$ that

$$\mathscr{L}^{f_0, f_1, \dots, f_K}_{\mathfrak{t}_0, \mathfrak{t}_1, \dots, \mathfrak{t}_K}(t) = \sum_{k=0}^K \mathscr{L}^{0, 1, 0}_{\mathfrak{t}_{k-1}, \mathfrak{t}_k, \mathfrak{t}_{k+1}}(t) f_k \tag{4.5}$$

(cf. Definition 4.1).

Proof of Lemma 4.4. Note that Lemma 4.3 and, for instance, items (i) and (iii) in Lemma 4.6 in [1] show that for all $t \in [0, T]$ it holds that

$$\begin{split} \sum_{k=0}^{K} \mathscr{L}_{t_{k-1}, t_k, t_{k+1}}^{0,1,0}(t) f_k &= \mathbb{1}_{\{t_0\}}(t) f_0 + \sum_{k=1}^{K} \frac{t - t_{k-1}}{t_k - t_{k-1}} \mathbb{1}_{(t_{k-1}, t_k]}(t) f_k + \sum_{k=0}^{K-1} \frac{t_{k+1} - t}{t_{k+1} - t_k} \mathbb{1}_{(t_k, t_{k+1})}(t) f_k \\ &= \mathbb{1}_{\{t_0\}}(t) f_0 + \sum_{k=1}^{K} \frac{t - t_{k-1}}{t_k - t_{k-1}} \mathbb{1}_{(t_{k-1}, t_k]}(t) f_k + \sum_{k=1}^{K} \frac{t_k - t}{t_k - t_{k-1}} \mathbb{1}_{(t_{k-1}, t_k)}(t) f_{k-1} \\ &= \mathbb{1}_{\{t_0\}}(t) f_0 + \sum_{k=1}^{K} \left(\frac{t - t_{k-1}}{t_k - t_{k-1}} f_k + \frac{t_k - t}{t_k - t_{k-1}} f_{k-1} \right) \mathbb{1}_{(t_{k-1}, t_k]}(t) \\ &= \mathbb{1}_{\{t_0\}}(t) \mathscr{L}_{t_0, t_1, \dots, t_K}^{f_0, f_1, \dots, f_K}(t) + \sum_{k=1}^{K} \mathscr{L}_{t_0, t_1, \dots, t_K}^{f_0, f_1, \dots, f_K}(t) \mathbb{1}_{(t_{k-1}, t_k]}(t) \\ &= \mathscr{L}_{t_0, t_1, \dots, t_K}^{f_0, f_1, \dots, f_K}(t) \end{split}$$

(cf. Definition 4.1). The proof of Lemma 4.4 is thus complete.

4.2 ANN representations and approximations for hat functions

Lemma 4.5. Let $\mathfrak{t}_0, \mathfrak{t}_1, \mathfrak{t}_2 \in \mathbb{R}$ satisfy $\mathfrak{t}_0 < \mathfrak{t}_1 < \mathfrak{t}_2$, let $c_0, c_1, c_2 \in \mathbb{R}$ satisfy

$$c_0 = \frac{1}{\mathfrak{t}_1 - \mathfrak{t}_0}, \qquad c_1 = -\frac{1}{\mathfrak{t}_2 - \mathfrak{t}_1} - \frac{1}{\mathfrak{t}_1 - \mathfrak{t}_0}, \qquad and \qquad c_2 = \frac{1}{\mathfrak{t}_2 - \mathfrak{t}_1}, \tag{4.7}$$

let $a \in \mathbb{R} \setminus \{-1, 1\}$, let $a \colon \mathbb{R} \to \mathbb{R}$ satisfy for all $x \in \mathbb{R}$ that $a(x) = \max\{x, ax\}$, and let $\mathbf{H} \in \mathbf{N}$ be given by

$$\mathbf{H} = \begin{bmatrix} 2\\ \bigoplus \\ j=0 \end{bmatrix} \left(\left(\frac{|1-a|ac_j}{(1-a)(1-a^2)} \right) \otimes \left(\mathbf{i}_1 \bullet \mathbf{A}_{\frac{-|1-a|}{1-a},\frac{|1-a|\mathbf{i}_j}{1-a}} \right) \right) \end{bmatrix} \\ \oplus \begin{bmatrix} 2\\ \bigoplus \\ j=0 \end{bmatrix} \left(\left(\frac{|1-a|c_j}{(1-a)(1-a^2)} \right) \otimes \left(\mathbf{i}_1 \bullet \mathbf{A}_{\frac{|1-a|}{1-a},\frac{-|1-a|\mathbf{i}_j}{1-a}} \right) \right) \end{bmatrix}$$
(4.8)

(cf. Definitions 3.1, 3.6, 3.8, 3.9, 3.15, and 3.19). Then

- (i) it holds that $\mathcal{R}_a(\mathbf{H}) \in C(\mathbb{R}, \mathbb{R})$,
- (ii) it holds for all $t \in \mathbb{R}$ that $(\mathcal{R}_a(\mathbf{H}))(t) = \mathscr{L}^{0,1,0}_{\mathfrak{t}_0,\mathfrak{t}_1,\mathfrak{t}_2}(t)$,
- (iii) it holds that $\mathcal{D}(\mathbf{H}) = (1, 6, 1)$, and
- (iv) it holds that $\mathcal{P}(\mathbf{H}) = 19$
- (cf. Definitions 3.4 and 4.1).

Proof of Lemma 4.5. Let $\mathfrak{r} \in C(\mathbb{R}, \mathbb{R})$ satisfy for all $x \in \mathbb{R}$ that $\mathfrak{r}(x) = \max\{x, 0\}$ and let $\mathbf{F} \in \mathbf{N}$ be given by

$$\mathbf{F} = \mathbf{A}_{1,0} \bullet \bigoplus_{j=0}^{2} \left(c_j \circledast \left(\mathbf{i}_1 \bullet \mathbf{A}_{1,-\mathbf{i}_j} \right) \right)$$
(4.9)

(cf. Definitions 3.1, 3.6, 3.8, 3.9, 3.15, and 3.19). Observe that [1, Lemma 4.10] proves that for all $t \in \mathbb{R}$ it holds that

$$(\mathcal{R}_{\mathfrak{r}}(\mathbf{F}))(t) = \mathscr{L}^{0,1,0}_{\mathfrak{t}_0,\mathfrak{t}_1,\mathfrak{t}_2}(t)$$
(4.10)

(cf. Definitions 3.4 and 4.1). In addition, note that, for example, item (iii) in Lemma 4.9 in [1] and, for instance, [1, Lemma 3.2] demonstrate that for all $t \in \mathbb{R}$ it holds that

$$(\mathcal{R}_{\mathfrak{r}}(\mathbf{F}))(t) = \sum_{j=0}^{2} c_j \mathfrak{r}(t-\mathfrak{t}_j) = \sum_{j=0}^{2} c_j (\mathcal{R}_{\mathfrak{r}}(\mathfrak{i}_1))(t-\mathfrak{t}_j).$$
(4.11)

Moreover, observe that, for example, [1, Lemma 2.19] establishes that for all $t \in \mathbb{R}$ it holds that $\mathcal{R}_a(\mathbf{H}) \in C(\mathbb{R}, \mathbb{R})$ and

$$\begin{aligned} (\mathcal{R}_{a}(\mathbf{H}))(t) \\ &= \left[\sum_{j=0}^{2} \left(\frac{|1-a|ac_{j}|}{(1-a)(1-a^{2})} \right) \left(\mathcal{R}_{a}(\mathfrak{i}_{1}) \right) \left(\frac{-|1-a|t}{1-a} + \frac{|1-a|\mathfrak{t}_{j}|}{1-a} \right) \right] \\ &+ \left[\sum_{j=0}^{2} \left(\frac{|1-a|c_{j}|}{(1-a)(1-a^{2})} \right) \left(\mathcal{R}_{a}(\mathfrak{i}_{1}) \right) \left(\frac{|1-a|t}{1-a} + \frac{-|1-a|\mathfrak{t}_{j}|}{1-a} \right) \right] \\ &= \sum_{j=0}^{2} \frac{|1-a|c_{j}|}{(1-a)(1-a^{2})} \left[a \left(\mathcal{R}_{a}(\mathfrak{i}_{1}) \right) \left(\frac{-|1-a|(t-\mathfrak{t}_{j})|}{1-a} \right) + \left(\mathcal{R}_{a}(\mathfrak{i}_{1}) \right) \left(\frac{|1-a|(t-\mathfrak{t}_{j})|}{1-a} \right) \right]. \end{aligned}$$

$$(4.12)$$

This, Lemma 3.7, (4.10), and (4.11) ensure that for all $t \in \mathbb{R}$ it holds that

$$(\mathcal{R}_{a}(\mathbf{H}))(t) = \sum_{j=0}^{2} c_{j}(\mathcal{R}_{\mathfrak{r}}(\mathfrak{i}_{1}))(t-\mathfrak{t}_{j}) = (\mathcal{R}_{\mathfrak{r}}(\mathbf{F}))(t) = \mathscr{L}_{\mathfrak{t}_{0},\mathfrak{t}_{1},\mathfrak{t}_{2}}^{0,1,0}(t).$$
(4.13)

This implies items (i) and (ii). Note that, for instance, item (i) in Lemma 2.19 in [1] and, for example, item (i) in Lemma 3.2 in [1] show that

$$\mathcal{D}(\mathbf{H}) = \left(\mathcal{I}(\mathfrak{i}_1), 6\mathbb{D}_1(\mathfrak{i}_1), \mathcal{O}(\mathfrak{i}_1)\right) = (1, 6, 1).$$
(4.14)

This proves item (iii). Observe that item (iii) demonstrates that

$$\mathcal{P}(\mathbf{H}) = 6(1+1) + (6+1) = 19. \tag{4.15}$$

This establishes item (iv). The proof of Lemma 4.5 is thus complete. \Box

Lemma 4.6. Let $\mathfrak{t}_0, \mathfrak{t}_1, \mathfrak{t}_2 \in \mathbb{R}$ satisfy $\mathfrak{t}_0 < \mathfrak{t}_1 < \mathfrak{t}_2$, let $\varepsilon \in (0, 1]$, $q \in (1, \infty)$, $\mathfrak{L} = \max\{|\mathfrak{t}_1 - \mathfrak{t}_0|^{-1}, |\mathfrak{t}_2 - \mathfrak{t}_1|^{-1}\}$ and let $a \colon \mathbb{R} \to \mathbb{R}$ satisfy for all $x \in \mathbb{R}$ that $a(x) = \ln(1 + \exp(x))$. Then there exists $\mathbf{H} \in \mathbf{N}$ such that for all $t \in \mathbb{R}$ it holds that

$$\begin{aligned}
\mathcal{R}_{a}(\mathbf{H}) &\in C(\mathbb{R}, \mathbb{R}), \qquad \mathcal{P}(\mathbf{H}) \leq 12 \left(\max\{1, 4\mathfrak{L}\} \right)^{q/(q-1)} 2^{q/(q-1)} \varepsilon^{-q/(q-1)}, \\
and \qquad \left| (\mathcal{R}_{a}(\mathbf{H}))(t) - \mathscr{L}^{0,1,0}_{t_{0}, t_{1}, t_{2}}(t) \right| \leq \varepsilon \max\{1, |t|^{q}\}
\end{aligned} \tag{4.16}$$

(cf. Definitions 3.1, 3.4, and 4.1).

Proof of Lemma 4.6. Let $f: [\mathfrak{t}_0, \mathfrak{t}_2] \to \mathbb{R}$ satisfy for all $t \in [\mathfrak{t}_0, \mathfrak{t}_2]$ that

$$f(t) = \mathscr{L}^{0,1,0}_{\mathfrak{t}_0,\mathfrak{t}_1,\mathfrak{t}_2}(t) \tag{4.17}$$

(cf. Definition 4.1). This and, for instance, [1, item (i) in Lemma 4.6] ensure that $f(\mathfrak{t}_0) = 0$, $f(\mathfrak{t}_1) = 1$, and $f(\mathfrak{t}_2) = 0$. In addition, note that the triangle inequality and (4.4) imply that for all $t, s \in [\mathfrak{t}_0, \mathfrak{t}_2]$ it holds that

$$|f(t) - f(s)| \le |f(t)| + |f(s)| \le 1 + 1 = 2.$$
(4.18)

This and, for example, item (i) in Lemma 4.7 in [1] show that for all $t, s \in \mathbb{R}$ it holds that

$$\begin{aligned} \left| \mathscr{L}_{\mathbf{t}_{0},\mathbf{t}_{1},\mathbf{t}_{2}}^{0,1,0}(t) - \mathscr{L}_{\mathbf{t}_{0},\mathbf{t}_{1},\mathbf{t}_{2}}^{0,1,0}(s) \right| &= \left| \mathscr{L}_{\mathbf{t}_{0},\mathbf{t}_{1},\mathbf{t}_{2}}^{f(\mathbf{t}_{0}),f(\mathbf{t}_{1}),f(\mathbf{t}_{2})}(t) - \mathscr{L}_{\mathbf{t}_{0},\mathbf{t}_{1},\mathbf{t}_{2}}^{f(\mathbf{t}_{0}),f(\mathbf{t}_{1}),f(\mathbf{t}_{2})}(s) \right| \\ &\leq \max\left\{ \frac{2}{|\mathbf{t}_{1} - \mathbf{t}_{0}|}, \frac{2}{|\mathbf{t}_{2} - \mathbf{t}_{1}|} \right\} |t - s| = 2\mathfrak{L}|t - s|. \end{aligned}$$
(4.19)

This and [1, Corollary 4.14] (applied with $\varepsilon \cap 2^{-1}\varepsilon$, $L \cap 2\mathfrak{L}$, $q \cap q$, $f \cap \mathscr{L}^{0,1,0}_{\mathfrak{t}_0,\mathfrak{t}_1,\mathfrak{t}_2}$, $a \cap a$ in the notation of [1, Corollary 4.14]) prove that there exists $\mathbf{H} \in \mathbf{N}$ such that (4.16) is satisfied (cf. Definition 3.1). The proof of Lemma 4.6 is thus complete.

4.3 ANN approximations for the square function

Lemma 4.7. Let $f: \mathbb{R} \to \mathbb{R}$ satisfy for all $x \in [0,1]$, $y \in \mathbb{R} \setminus [0,1]$ that $f(x) = x^2$ and $f(y) = \max\{y, 0\}$, let $\varepsilon \in (0,1]$, $q \in (2,\infty)$, $\delta \in \mathbb{R}$ satisfy $\delta = 2^{-1}4^{-2/(q-2)}\varepsilon^{q/(q-2)}$, let $a \in C(\mathbb{R}, \mathbb{R})$, $\mathbf{G} \in \mathbf{N}$, $r, c \in [1,\infty)$ satisfy for all $x \in \mathbb{R}$ that $\mathcal{R}_a(\mathbf{G}) \in C(\mathbb{R}, \mathbb{R})$, $\mathcal{P}(\mathbf{G}) \leq c\delta^{-r}$, and

$$|(\mathcal{R}_a(\mathbf{G}))(x) - f(x)| \le \delta \max\{1, |x|^q\},\tag{4.20}$$

let $W_1 \in \mathbb{R}^{2 \times 1}$, $W_2 \in \mathbb{R}^{1 \times 2}$ be given by

$$W_1 = \begin{pmatrix} \left(\frac{\varepsilon}{4}\right)^{1/(q-2)} \\ -\left(\frac{\varepsilon}{4}\right)^{1/(q-2)} \end{pmatrix} \quad and \quad W_2 = \left(\left(\frac{\varepsilon}{4}\right)^{-2/(q-2)} \quad \left(\frac{\varepsilon}{4}\right)^{-2/(q-2)} \right), \quad (4.21)$$

and let $\Phi \in \mathbf{N}$ be given by

$$\Phi = \mathbf{A}_{W_{2},0} \bullet \mathbf{P}_{2}(\mathbf{G},\mathbf{G}) \bullet \mathbf{A}_{W_{1},0}$$
(4.22)

(cf. Definitions 3.1, 3.4, 3.8, 3.9, and 3.12). Then

(i) it holds that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}, \mathbb{R})$,

(ii) it holds for all $x \in \mathbb{R}$ that $|(\mathcal{R}_a(\Phi))(x) - x^2| \leq \varepsilon \max\{1, |x|^q\}$, and

(*iii*) it holds that $\mathcal{P}(\Phi) \leq 2^{r+2} 4^{2r/(q-2)} c \varepsilon^{-rq/(q-2)}$.

Proof of Lemma 4.7. Throughout this proof let $\tilde{\varepsilon}, \tilde{q} \in \mathbb{R}$ satisfy $\tilde{\varepsilon} = \varepsilon/4$ and $\tilde{q} = 1/(q-2)$. Observe that, for instance, [37, Proposition 2.19] and, for example, item (vi) in Proposition 2.6 in [37] demonstrate that for all $x \in \mathbb{R}$ it holds that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}, \mathbb{R})$ and

$$(\mathcal{R}_{a}(\Phi))(x) = ((\mathcal{R}_{a}(\mathbf{A}_{W_{2},0})) \circ (\mathcal{R}_{a}(\mathbf{P}_{2}(\mathbf{G},\mathbf{G}))) \circ (\mathcal{R}_{a}(\mathbf{A}_{W_{1},0})))(x) = (\mathcal{R}_{a}(\mathbf{A}_{W_{2},0}))((\mathcal{R}_{a}(\mathbf{P}_{2}(\mathbf{G},\mathbf{G})))(\tilde{\varepsilon}^{\tilde{q}}x, -\tilde{\varepsilon}^{\tilde{q}}x)) = (\mathcal{R}_{a}(\mathbf{A}_{W_{2},0}))((\mathcal{R}_{a}(\mathbf{G}))(\tilde{\varepsilon}^{\tilde{q}}x), (\mathcal{R}_{a}(\mathbf{G}))(-\tilde{\varepsilon}^{\tilde{q}}x)) = \tilde{\varepsilon}^{-2\tilde{q}}((\mathcal{R}_{a}(\mathbf{G}))(\tilde{\varepsilon}^{\tilde{q}}x) + (\mathcal{R}_{a}(\mathbf{G}))(-\tilde{\varepsilon}^{\tilde{q}}x))$$

$$(4.23)$$

(cf. Definitions 3.4, 3.9, and 3.12). This establishes item (i). Note that the fact that for all $x \in \mathbb{R} \setminus [0, 1]$ it holds that $f(x) = \max\{x, 0\}$ ensures that for all $x \in \mathbb{R} \setminus [-1, 1]$ it holds that

$$|x| = \max\{x, 0\} - \min\{x, 0\} = \max\{x, 0\} + \max\{-x, 0\} = f(x) + f(-x).$$
(4.24)

The triangle inequality and (4.20) therefore imply that for all $x \in \mathbb{R} \setminus [-1, 1]$ it holds that

$$\begin{aligned} \left| (\mathcal{R}_{a}(\mathbf{G}))(x) + (\mathcal{R}_{a}(\mathbf{G}))(-x) - |x| \right| &\leq \left| (\mathcal{R}_{a}(\mathbf{G}))(x) - f(x) \right| + \left| (\mathcal{R}_{a}(\mathbf{G}))(-x) - f(-x) \right| \\ &\leq \delta \max\{1, |x|^{q}\} + \delta \max\{1, |-x|^{q}\} = 2\delta |x|^{q}. \end{aligned}$$

$$(4.25)$$

This and (4.23) show that for all $x \in \mathbb{R} \setminus [-\tilde{\varepsilon}^{-\tilde{q}}, \tilde{\varepsilon}^{-\tilde{q}}]$ it holds that

$$\begin{aligned} \left| (\mathcal{R}_{a}(\Phi))(x) - \tilde{\varepsilon}^{-\tilde{q}} |x| \right| &= \left| \tilde{\varepsilon}^{-2\tilde{q}} \left(\left((\mathcal{R}_{a}(\mathbf{G})) \left(\tilde{\varepsilon}^{\tilde{q}} x \right) + (\mathcal{R}_{a}(\mathbf{G})) \left(-\tilde{\varepsilon}^{\tilde{q}} x \right) \right) - |\tilde{\varepsilon}^{\tilde{q}} x| \right) \right| \\ &\leq \tilde{\varepsilon}^{-2\tilde{q}} 2\delta |\tilde{\varepsilon}^{\tilde{q}} x|^{q} = 2\delta \tilde{\varepsilon} |x|^{q}. \end{aligned}$$

$$(4.26)$$

Next, observe that the triangle inequality proves that for all $x \in \mathbb{R} \setminus [-\tilde{\varepsilon}^{-\tilde{q}}, \tilde{\varepsilon}^{-\tilde{q}}]$ it holds that

$$\begin{aligned} \left| \tilde{\varepsilon}^{-\tilde{q}} |x| - x^2 \right| &\leq \tilde{\varepsilon}^{-\tilde{q}} |x| + |x|^2 = |x|^q \left(\tilde{\varepsilon}^{-\tilde{q}} |x|^{-(q-1)} + |x|^{-(q-2)} \right) \\ &\leq |x|^q \left(\tilde{\varepsilon}^{-\tilde{q}} \tilde{\varepsilon}^{(q-1)\tilde{q}} + \tilde{\varepsilon}^{(q-2)\tilde{q}} \right) = 2|x|^q \tilde{\varepsilon}. \end{aligned}$$
(4.27)

Combining this, (4.26), the triangle inequality, and the fact that $\delta \in (0, 1]$ demonstrates that for all $x \in \mathbb{R} \setminus [-\tilde{\varepsilon}^{-\tilde{q}}, \tilde{\varepsilon}^{-\tilde{q}}]$ it holds that

$$\begin{aligned} \left| (\mathcal{R}_{a}(\Phi))(x) - x^{2} \right| &\leq \left| (\mathcal{R}_{a}(\Phi))(x) - \tilde{\varepsilon}^{-\tilde{q}} |x| \right| + \left| \tilde{\varepsilon}^{-\tilde{q}} |x| - x^{2} \right| \\ &\leq 2\delta \tilde{\varepsilon} |x|^{q} + 2|x|^{q} \tilde{\varepsilon} \leq 4\tilde{\varepsilon} |x|^{q} \leq \varepsilon \max\{1, |x|^{q}\}. \end{aligned}$$

$$(4.28)$$

Furthermore, note that the triangle inequality and (4.20) establish that for all $x \in [0, \infty)$ it holds that

$$|(\mathcal{R}_{a}(\mathbf{G}))(-x)| \leq |(\mathcal{R}_{a}(\mathbf{G}))(-x) - f(-x)| + |f(-x)| \\ \leq \delta \max\{1, |-x|^{q}\} + 0 = \delta \max\{1, |x|^{q}\}.$$
(4.29)

Combining this, the triangle inequality, and (4.20) ensures that for all $x \in [0, 1]$ it holds that

$$\begin{aligned} \left| (\mathcal{R}_{a}(\mathbf{G}))(x) + (\mathcal{R}_{a}(\mathbf{G}))(-x) - x^{2} \right| &= \left| (\mathcal{R}_{a}(\mathbf{G}))(x) + (\mathcal{R}_{a}(\mathbf{G}))(-x) - f(x) \right| \\ &\leq \left| (\mathcal{R}_{a}(\mathbf{G}))(x) - f(x) \right| + \left| (\mathcal{R}_{a}(\mathbf{G}))(-x) \right| \\ &\leq \delta \max\{1, |x|^{q}\} + \delta \max\{1, |x|^{q}\} = 2\delta. \end{aligned}$$

$$(4.30)$$

This and (4.23) imply that for all $x \in [-\tilde{\varepsilon}^{-\tilde{q}}, \tilde{\varepsilon}^{-\tilde{q}}]$ it holds that

$$\begin{aligned} |(\mathcal{R}_{a}(\Phi))(x) - x^{2}| &\leq \tilde{\varepsilon}^{-2\tilde{q}} |(\mathcal{R}_{a}(\mathbf{G}))(\tilde{\varepsilon}^{\tilde{q}}x) + (\mathcal{R}_{a}(\mathbf{G}))(-\tilde{\varepsilon}^{\tilde{q}}x) - (\tilde{\varepsilon}^{\tilde{q}}x)^{2}| \\ &\leq \tilde{\varepsilon}^{-2\tilde{q}}2\delta = \varepsilon^{-2\tilde{q}} \cdot 4^{2\tilde{q}} \cdot 2 \cdot 2^{-1} \cdot 4^{-2\tilde{q}} \cdot \varepsilon^{q\tilde{q}} = \varepsilon \\ &\leq \varepsilon \max\{1, |x|^{q}\}. \end{aligned}$$
(4.31)

This and (4.28) show item (ii). Observe that, for instance, [37, Corollary 2.9] proves that

$$\mathcal{P}(\mathbf{A}_{W_{2},0} \bullet \mathbf{P}_{2}(\mathbf{G},\mathbf{G}) \bullet \mathbf{A}_{W_{1},0}) \leq \mathcal{P}(\mathbf{A}_{W_{2},0} \bullet \mathbf{P}_{2}(\mathbf{G},\mathbf{G})) \leq \mathcal{P}(\mathbf{P}_{2}(\mathbf{G},\mathbf{G}))$$
(4.32)

(cf. Definitions 3.1 and 3.8). This, the fact that $\mathcal{P}(\mathbf{G}) \leq c\delta^{-r}$, and, for example, [37, Corollary 2.21] demonstrate that

$$\mathcal{P}(\Phi) \le \mathcal{P}(\mathbf{P}_2(\mathbf{G}, \mathbf{G})) \le 4\mathcal{P}(\mathbf{G}) \le 4c\delta^{-r} = 4c2^r 4^{2r/(q-2)}\varepsilon^{-rq/(q-2)}.$$
(4.33)

This establishes item (iii). The proof of Lemma 4.7 is thus complete.

4.4 ANN approximations for the product function

Lemma 4.8. Let $\varepsilon \in (0,1]$, $q \in (2,\infty)$, $\delta \in \mathbb{R}$ satisfy $\delta = (2^{q-1}+1)^{-1}\varepsilon$, let $a \in C(\mathbb{R},\mathbb{R})$, $\Phi \in \mathbb{N}$, $r, c \in [1,\infty)$ satisfy for all $x \in \mathbb{R}$ that $\mathcal{R}_a(\Phi) \in C(\mathbb{R},\mathbb{R})$, $\mathcal{P}(\Phi) \leq c\delta^{-r}$, and

$$|(\mathcal{R}_a(\Phi))(x) - x^2| \le \delta \max\{1, |x|^q\},$$
(4.34)

let $W_1 \in \mathbb{R}^{3 \times 2}$, $W_2 \in \mathbb{R}^{1 \times 3}$ be given by

$$W_1 = \begin{pmatrix} 1 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix} \quad and \quad W_2 = \begin{pmatrix} \frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} \end{pmatrix}, \quad (4.35)$$

and let $\Gamma \in \mathbf{N}$ be given by

$$\Gamma = \mathbf{A}_{W_{2},0} \bullet \mathbf{P}_{3}(\Phi, \Phi, \Phi) \bullet \mathbf{A}_{W_{1},0}$$
(4.36)

(cf. Definitions 3.1, 3.4, 3.8, 3.9, and 3.12). Then

- (i) it holds that $\mathcal{R}_a(\Gamma) \in C(\mathbb{R}^2, \mathbb{R})$,
- (ii) it holds for all $x, y \in \mathbb{R}$ that $|(\mathcal{R}_a(\Gamma))(x, y) xy| \leq \varepsilon \max\{1, |x|^q, |y|^q\}$, and
- (iii) it holds that $\mathcal{P}(\Gamma) \leq 9c(2^{q-1}+1)^r \varepsilon^{-r}$.

Proof of Lemma 4.8. Note that, for instance, [37, Proposition 2.19] and, for example, item (vi) in Proposition 2.6 in [37] show that for all $x, y \in \mathbb{R}$ it holds that $\mathcal{R}_a(\Gamma) \in C(\mathbb{R}^2, \mathbb{R})$ and

$$(\mathcal{R}_{a}(\Gamma))(x,y) = \left(\left(\mathcal{R}_{a}(\mathbf{A}_{W_{2},0}) \right) \circ \left(\mathcal{R}_{a}\left(\mathbf{P}_{3}(\Phi,\Phi,\Phi) \right) \right) \circ \left(\mathcal{R}_{a}(\mathbf{A}_{W_{1},0}) \right) \right)(x,y) \\ = \left(\mathcal{R}_{a}(\mathbf{A}_{W_{2},0}) \right) \left(\left(\mathcal{R}_{a}\left(\mathbf{P}_{3}(\Phi,\Phi,\Phi) \right) \right)(x+y,x,y) \right) \\ = \left(\mathcal{R}_{a}(\mathbf{A}_{W_{2},0}) \right) \left(\left(\mathcal{R}_{a}(\Phi) \right)(x+y), \left(\mathcal{R}_{a}(\Phi) \right)(x), \left(\mathcal{R}_{a}(\Phi) \right)(y) \right) \\ = \frac{1}{2} \left(\mathcal{R}_{a}(\Phi) \right)(x+y) - \frac{1}{2} \left(\mathcal{R}_{a}(\Phi) \right)(x) - \frac{1}{2} \left(\mathcal{R}_{a}(\Phi) \right)(y) \right)$$
(4.37)

(cf. Definitions 3.4, 3.9, and 3.12). This proves item (i). Observe that (4.37), the fact that for all $x, y \in \mathbb{R}$ it holds that $(x + y)^2 - x^2 - y^2 = 2xy$, the triangle inequality, and (4.34) ensure that for all $x, y \in \mathbb{R}$ it holds that

$$2|(\mathcal{R}_{a}(\Gamma))(x,y) - xy| = |(\mathcal{R}_{a}(\Phi))(x+y) - (x+y)^{2} - ((\mathcal{R}_{a}(\Phi))(x) - x^{2}) - ((\mathcal{R}_{a}(\Phi))(y) - y^{2})| \le |(\mathcal{R}_{a}(\Phi))(x+y) - (x+y)^{2}| + |(\mathcal{R}_{a}(\Phi))(x) - x^{2}| + |(\mathcal{R}_{a}(\Phi))(y) - y^{2}| \le \delta \max\{1, |x+y|^{q}\} + \delta \max\{1, |x|^{q}\} + \delta \max\{1, |y|^{q}\}.$$
(4.38)

Furthermore, note that Jensen's inequality implies that for all $x, y \in \mathbb{R}$ it holds that

$$\max\{1, |x+y|^{q}\} + \max\{1, |x|^{q}\} + \max\{1, |y|^{q}\}
\leq 2^{q-1} \max\{1, |x|^{q} + |y|^{q}\} + \max\{1, |x|^{q}\} + \max\{1, |y|^{q}\}
\leq (2^{q-1} + 1) (\max\{1, |x|^{q}\} + \max\{1, |y|^{q}\})
\leq 2(2^{q-1} + 1) \max\{1, |x|^{q}, |y|^{q}\}.$$
(4.39)

This and (4.38) show that for all $x, y \in \mathbb{R}$ it holds that

$$\left| \left(\mathcal{R}_{a}(\Gamma) \right)(x,y) - xy \right| \leq \delta(2^{q-1} + 1) \max\{1, |x|^{q}, |y|^{q}\} = \varepsilon \max\{1, |x|^{q}, |y|^{q}\}.$$
(4.40)

This proves item (ii). Observe that, for instance, [37, Corollary 2.9], [37, Corollary 2.21], and the fact that $\mathcal{P}(\Phi) \leq c\delta^{-r}$ demonstrate that

$$\mathcal{P}(\Gamma) = \mathcal{P}(\mathbf{A}_{W_{2},0} \bullet \mathbf{P}_{3}(\Phi, \Phi, \Phi) \bullet \mathbf{A}_{W_{1},0}) \leq \mathcal{P}(\mathbf{A}_{W_{2},0} \bullet \mathbf{P}_{3}(\Phi, \Phi, \Phi))$$

$$\leq \mathcal{P}(\mathbf{P}_{3}(\Phi, \Phi, \Phi)) \leq 9\mathcal{P}(\Phi) \leq 9c\delta^{-r} = 9c(2^{q-1}+1)^{r}\varepsilon^{-r}$$
(4.41)

(cf. Definitions 3.1 and 3.8). This establishes item (iii). The proof of Lemma 4.8 is thus complete. $\hfill \Box$

Corollary 4.9. Let $f: \mathbb{R} \to \mathbb{R}$ satisfy for all $x \in [0,1]$, $y \in \mathbb{R} \setminus [0,1]$ that $f(x) = x^2$ and $f(y) = \max\{y, 0\}$, let $\varepsilon \in (0,1]$, $q \in (2,\infty)$, $r, c \in [1,\infty)$, let $\delta \in \mathbb{R}$ satisfy $\delta = 2^{-1}4^{-2/(q-2)}(2^{q-1}+1)^{-q/(q-2)}\varepsilon^{q/(q-2)}$, let $a \in C(\mathbb{R},\mathbb{R})$, and let $\mathbf{G} \in \mathbf{N}$ satisfy for all $x \in \mathbb{R}$ that $\mathcal{R}_a(\mathbf{G}) \in C(\mathbb{R},\mathbb{R})$, $\mathcal{P}(\mathbf{G}) \leq c\delta^{-r}$, and

$$|(\mathcal{R}_a(\mathbf{G}))(x) - f(x)| \le \delta \max\{1, |x|^q\}$$

$$(4.42)$$

(cf. Definitions 3.1 and 3.4). Then there exists $\Gamma \in \mathbf{N}$ such that for all $x, y \in \mathbb{R}$ it holds that

$$\mathcal{R}_{a}(\Gamma) \in C(\mathbb{R}^{2}, \mathbb{R}), \qquad \mathcal{P}(\Gamma) \leq 36 \cdot 2^{\frac{(q^{2}+q+2)r}{q-2}} c\varepsilon^{-\frac{rq}{q-2}},$$

and $|(\mathcal{R}_{a}(\Gamma))(x, y) - xy| \leq \varepsilon \max\{1, |x|^{q}, |y|^{q}\}.$ (4.43)

Proof of Corollary 4.9. Note that Lemma 4.7 (applied with $f \curvearrowleft f$, $\mathbf{G} \curvearrowleft \mathbf{G}$, $a \curvearrowleft a$, $q \curvearrowleft q$, $\varepsilon \curvearrowleft (2^{q-1}+1)^{-1}\varepsilon$, $r \curvearrowleft r$, $c \backsim c$ in the notation of Lemma 4.7) ensures that there exists $\Phi \in \mathbf{N}$ such that for all $x \in \mathbb{R}$ it holds that

$$\mathcal{R}_{a}(\Phi) \in C(\mathbb{R}, \mathbb{R}), \qquad \mathcal{P}(\Phi) \leq 2^{r+2} 4^{\frac{2r}{q-2}} c \left((2^{q-1}+1)^{-1} \varepsilon \right)^{-\frac{rq}{q-2}}
\text{and} \qquad |(\mathcal{R}_{a}(\Phi))(x) - x^{2}| \leq (2^{q-1}+1)^{-1} \varepsilon \max\{1, |x|^{q}\}$$
(4.44)

(cf. Definitions 3.1 and 3.4). Therefore, Lemma 4.8 (applied with $\Phi \curvearrowleft \Phi$, $a \curvearrowleft a$, $q \backsim q$, $\varepsilon \backsim \varepsilon$, $r \backsim \frac{rq}{q-2}$, $c \backsim 2^{r+2}4^{2r/(q-2)}c$ in the notation of Lemma 4.8) demonstrates that there exists $\Gamma \in \mathbf{N}$ such that for all $x, y \in \mathbb{R}$ it holds that

$$\mathcal{R}_{a}(\Gamma) \in C(\mathbb{R}^{2}, \mathbb{R}), \qquad \mathcal{P}(\Gamma) \leq 9 \cdot 2^{r+2} 4^{\frac{2r}{q-2}} c(2^{q-1}+1)^{\frac{rq}{q-2}} \varepsilon^{-\frac{rq}{q-2}},$$

and
$$|(\mathcal{R}_{a}(\Gamma))(x, y) - xy| \leq \varepsilon \max\{1, |x|^{q}, |y|^{q}\}.$$

$$(4.45)$$

Moreover, observe that the fact that $2^{q-1} + 1 \leq 2^q$ ensures that

$$9 \cdot 2^{r+2} 4^{\frac{2r}{q-2}} c (2^{q-1}+1)^{\frac{rq}{q-2}} = 36 \cdot 2^{r+\frac{4r}{q-2}} c (2^{q-1}+1)^{\frac{rq}{q-2}} \leq 36 \cdot 2^{r+\frac{4r}{q-2}} c 2^{\frac{rq^2}{q-2}} = 36 \cdot 2^{\frac{(q^2+q+2)r}{q-2}} c.$$

$$(4.46)$$

The proof of Corollary 4.9 is thus complete.

Corollary 4.10. Let $\varepsilon \in (0, 1]$, $q \in (2, \infty)$, $\alpha \in \mathbb{R} \setminus \{-1, 1\}$ and let $a \colon \mathbb{R} \to \mathbb{R}$ satisfy for all $x \in \mathbb{R}$ that $a(x) = \max\{x, \alpha x\}$. Then there exists $\Gamma \in \mathbb{N}$ such that for all $x, y \in \mathbb{R}$ it holds that

$$\mathcal{R}_{a}(\Gamma) \in C(\mathbb{R}^{2}, \mathbb{R}), \qquad \mathcal{P}(\Gamma) \leq 864 \cdot 2^{\frac{q^{3}+3q^{2}-2q}{(q-2)(q-1)}} \varepsilon^{-\frac{q^{2}}{(q-2)(q-1)}}, \qquad (4.47)$$
and
$$|(\mathcal{R}_{a}(\Gamma))(x, y) - xy| \leq \varepsilon \max\{1, |x|^{q}, |y|^{q}\}$$

(cf. Definitions 3.1 and 3.4).

Proof of Corollary 4.10. Let $\delta = 2^{-1}4^{-2/(q-2)}(2^{q-1}+1)^{-q/(q-2)}\varepsilon^{q/(q-2)}$ and let $f: \mathbb{R} \to \mathbb{R}$ satisfy for all $x \in [0,1]$, $y \in \mathbb{R} \setminus [0,1]$ that $f(x) = x^2$ and $f(y) = \max\{y,0\}$. Note that for all $x, y \in \mathbb{R}$ it holds that $|f(x) - f(y)| \leq 2|x - y|$. Therefore, [1, Corollary 4.13] (applied with $\varepsilon \frown \delta$, $L \frown 2$, $q \frown q$, $\alpha \frown \alpha$, $f \frown f$, $a \frown a$ in the notation of [1, Corollary 4.13]) establishes that there exists $\mathbf{G} \in \mathbf{N}$ such that for all $x \in \mathbb{R}$ it holds that

$$\mathcal{R}_{a}(\mathbf{G}) \in C(\mathbb{R}, \mathbb{R}), \qquad \mathcal{P}(\mathbf{G}) \leq 24 \cdot 4^{\frac{q}{q-1}} \cdot \delta^{-\frac{q}{q-1}},$$

and
$$|(\mathcal{R}_{a}(\mathbf{G}))(x) - f(x)| \leq \delta \max\{1, |x|^{q}\}$$
(4.48)

(cf. Definitions 3.1 and 3.4). This and Corollary 4.9 (applied with $f \curvearrowleft f$, $\mathbf{G} \curvearrowleft \mathbf{G}$, $a \backsim a$, $q \backsim q$, $\varepsilon \backsim \varepsilon$, $r \backsim \frac{q}{q-1}$, $c \backsim 24 \cdot 4^{q/(q-1)}$ in the notation of Corollary 4.9) imply that there exists $\Gamma \in \mathbf{N}$ such that for all $x, y \in \mathbb{R}$ it holds that

$$\mathcal{R}_{a}(\Gamma) \in C(\mathbb{R}^{2}, \mathbb{R}), \qquad \mathcal{P}(\Gamma) \leq 36 \cdot 2^{\frac{(q^{2}+q+2)q}{(q-2)(q-1)}} \cdot 24 \cdot 4^{\frac{q}{q-1}} \varepsilon^{-\frac{q^{2}}{(q-2)(q-1)}}, \qquad (4.49)$$

and $|(\mathcal{R}_{a}(\Gamma))(x, y) - xy| \leq \varepsilon \max\{1, |x|^{q}, |y|^{q}\}.$

Moreover, observe that the fact that $\frac{(q^2+q+2)q}{(q-2)(q-1)} + \frac{2q}{q-1} = \frac{(q^2+3q-2)q}{(q-2)(q-1)}$ shows that

$$36 \cdot 2^{\frac{(q^2+q+2)q}{(q-2)(q-1)}} \cdot 24 \cdot 4^{\frac{q}{q-1}} = 864 \cdot 2^{\frac{(q^2+3q-2)q}{(q-2)(q-1)}}.$$
(4.50)

The proof of Corollary 4.10 is thus complete.

Corollary 4.11. Let $\varepsilon \in (0,1]$, $q \in (2,\infty)$ and let $a \colon \mathbb{R} \to \mathbb{R}$ satisfy for all $x \in \mathbb{R}$ that $a(x) = \ln(1 + \exp(x))$. Then there exists $\Gamma \in \mathbf{N}$ such that for all $x, y \in \mathbb{R}$ it holds that

$$\mathcal{R}_{a}(\Gamma) \in C(\mathbb{R}^{2}, \mathbb{R}), \qquad \mathcal{P}(\Gamma) \leq 1728 \cdot 2^{\frac{q^{3}+3q^{2}-2q}{(q-2)(q-1)}} \varepsilon^{-\frac{q^{2}}{(q-2)(q-1)}},$$

$$and \qquad |(\mathcal{R}_{a}(\Gamma))(x, y) - xy| \leq \varepsilon \max\{1, |x|^{q}, |y|^{q}\}$$

$$(4.51)$$

(cf. Definitions 3.1 and 3.4).

Proof of Corollary 4.11. Let $\delta = 2^{-1}4^{-2/(q-2)}(2^{q-1}+1)^{-q/(q-2)}\varepsilon^{q/(q-2)}$ and let $f: \mathbb{R} \to \mathbb{R}$ satisfy for all $x \in [0,1]$, $y \in \mathbb{R} \setminus [0,1]$ that $f(x) = x^2$ and $f(y) = \max\{y,0\}$. Note that for all $x, y \in \mathbb{R}$ it holds that $|f(x) - f(y)| \leq 2|x - y|$. Hence, [1, Corollary 4.14] (applied with $\varepsilon \frown \delta/2$, $L \frown 2$, $q \frown q$, $f \frown f$, $a \frown a$ in the notation of [1, Corollary 4.14]) demonstrates that there exists $\mathbf{G} \in \mathbf{N}$ such that for all $x \in \mathbb{R}$ it holds that

$$\mathcal{R}_{a}(\mathbf{G}) \in C(\mathbb{R}, \mathbb{R}), \qquad \mathcal{P}(\mathbf{G}) \leq 12 \cdot 4^{\frac{q}{q-1}} \cdot 2^{\frac{q}{q-1}} \delta^{-\frac{q}{q-1}},$$

and
$$|(\mathcal{R}_{a}(\mathbf{G}))(x) - f(x)| \leq \delta \max\{1, |x|^{q}\}$$
(4.52)

(cf. Definitions 3.1 and 3.4). This and Corollary 4.9 (applied with $f \curvearrowleft f$, $\mathbf{G} \curvearrowleft \mathbf{G}$, $a \backsim a$, $q \backsim q$, $\varepsilon \backsim \varepsilon$, $r \backsim \frac{q}{q-1}$, $c \backsim 12 \cdot 4^{q/(q-1)} \cdot 2^{q/(q-1)}$ in the notation of Corollary 4.9) prove that there exists $\Gamma \in \mathbf{N}$ such that for all $x, y \in \mathbb{R}$ it holds that

$$\mathcal{R}_{a}(\Gamma) \in C(\mathbb{R}^{2}, \mathbb{R}), \qquad \mathcal{P}(\Gamma) \leq 36 \cdot 2^{\frac{(q^{2}+q+2)q}{(q-2)(q-1)}} \cdot 12 \cdot 4^{\frac{q}{q-1}} \cdot 2^{\frac{q}{q-1}} \varepsilon^{-\frac{q^{2}}{(q-2)(q-1)}}, \qquad (4.53)$$
and
$$|(\mathcal{R}_{a}(\Gamma))(x, y) - xy| \leq \varepsilon \max\{1, |x|^{q}, |y|^{q}\}.$$

In addition, observe that the fact that $\frac{q}{q-1} \leq 2$ demonstrates that

$$36 \cdot 2^{\frac{(q^2+q+2)q}{(q-2)(q-1)}} \cdot 12 \cdot 4^{\frac{q}{q-1}} \cdot 2^{\frac{q}{q-1}} = 432 \cdot 2^{\frac{(q^2+3q-2)q}{(q-2)(q-1)}} \cdot 2^{\frac{q}{q-1}} \le 1728 \cdot 2^{\frac{(q^2+3q-2)q}{(q-2)(q-1)}}.$$
(4.54)

The proof of Corollary 4.11 is thus complete.

4.5 ANN approximations for interpolation functions

Proposition 4.12. Let $K, d, \mathfrak{d} \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $q \in (2, \infty)$, $a, h_0, h_1, \ldots, h_K \in C(\mathbb{R}, \mathbb{R})$, $f_0, f_1, \ldots, f_K \in C(\mathbb{R}^d, \mathbb{R})$, $\mathbf{H}_0, \mathbf{H}_1, \ldots, \mathbf{H}_K, \mathbf{F}_0, \mathbf{F}_1, \ldots, \mathbf{F}_K \in \mathbf{N}$ satisfy for all $k \in \{0, 1, \ldots, K\}$, $t \in \mathbb{R}$ that $\mathcal{R}_a(\mathbf{F}_k) = f_k, \mathcal{R}_a(\mathbf{H}_k) \in C(\mathbb{R}, \mathbb{R})$, and

$$|h_k(t) - (\mathcal{R}_a(\mathbf{H}_k))(t)| \le \varepsilon \max\{1, |t|^q\},$$
(4.55)

let $\mathfrak{J}, \Gamma \in \mathbf{N}$ satisfy for all $v, w \in \mathbb{R}$ that $\mathcal{D}(\mathfrak{J}) = (1, \mathfrak{d}, 1), \mathcal{R}_a(\mathfrak{J}) = \mathrm{id}_{\mathbb{R}}, \mathcal{R}_a(\Gamma) \in C(\mathbb{R}^2, \mathbb{R}),$ and

$$|vw - (\mathcal{R}_a(\Gamma))(v, w)| \le \varepsilon \max\{1, |v|^q, |w|^q\},$$

$$(4.56)$$

and let $\Phi \in \mathbf{N}$ be given by

$$\Phi = \bigoplus_{k=0,\mathfrak{J}}^{K} \left(\Gamma \bullet \mathcal{P}_{2,\mathfrak{J}}(\mathbf{H}_{k}, \mathbf{F}_{k}) \right)$$
(4.57)

(cf. Definitions 3.1, 3.4, 3.8, 3.13, and 3.20). Then

- (i) it holds that $\mathcal{R}_a(\Phi) \in C(\mathbb{R}^{d+1}, \mathbb{R})$,
- (ii) it holds for all $t \in \mathbb{R}$, $x \in \mathbb{R}^d$ that

$$\left(\mathcal{R}_{a}(\Phi)\right)(t,x) = \sum_{k=0}^{K} \left(\mathcal{R}_{a}(\Gamma)\right) \left(\left(\mathcal{R}_{a}(\mathbf{H}_{k})\right)(t), f_{k}(x)\right),$$
(4.58)

(iii) it holds for all $t \in \mathbb{R}$, $x \in \mathbb{R}^d$ that

$$\left| \left(\mathcal{R}_{a}(\Phi) \right)(t,x) - \sum_{k=0}^{K} h_{k}(t) f_{k}(x) \right| \leq 2\varepsilon \sum_{k=0}^{K} \left(1 + |f_{k}(x)|^{q} \right) \left(\max\{1, |t|^{q}\} + |h_{k}(t)| \right)^{q}, \quad (4.59)$$

(iv) it holds that

$$\mathcal{L}(\Phi) \le 2\mathcal{L}(\Gamma) \left[\max_{k \in \{0,1,\dots,K\}} \mathcal{L}(\mathbf{H}_k) \right] \left[\max_{k \in \{0,1,\dots,K\}} \mathcal{L}(\mathbf{F}_k) \right],$$
(4.60)

(v) it holds that

$$\left\|\left|\mathcal{D}(\Phi)\right\|\right\| \le 2\mathfrak{d}(K+1) \left\|\left|\mathcal{D}(\Gamma)\right\|\right\| \left[\max_{k \in \{0,1,\dots,K\}} \left\|\left|\mathcal{D}(\mathbf{H}_k)\right\|\right\|\right] \left[\max_{k \in \{0,1,\dots,K\}} \left\|\left|\mathcal{D}(\mathbf{F}_k)\right\|\right\|\right], \quad (4.61)$$

and

(vi) it holds that

$$\mathcal{P}(\Phi) \leq 16\mathfrak{d}^{2}(K+1)^{2} \left[\mathcal{P}(\Gamma)\right]^{3} \left[\max_{k \in \{0,1,\dots,K\}} \mathcal{P}(\mathbf{H}_{k})\right]^{3} \\ \cdot \left[\max_{k \in \{0,1,\dots,K\}} \mathcal{L}(\mathbf{F}_{k})\right] \left[\max_{k \in \{0,1,\dots,K\}} \left\|\left|\mathcal{D}(\mathbf{F}_{k})\right|\right\|\right]^{2}$$
(4.62)

(cf. Definition 2.1).

Proof of Proposition 4.12. Note that, for example, item (i) in Corollary 2.23 in [37], item (vi) in Proposition 2.6 in [37], and item (iv) in Lemma 3.21 establish item (i). Observe that item (v) in Lemma 3.21 establishes that for all $t \in \mathbb{R}$, $x \in \mathbb{R}^d$ it holds that

$$\left(\mathcal{R}_{a}(\Phi)\right)(t,x) = \sum_{k=0}^{K} \left(\mathcal{R}_{a}\left(\Gamma \bullet \mathsf{P}_{2,\mathfrak{J}}(\mathbf{H}_{k},\mathbf{F}_{k})\right)\right)(t,x)$$
(4.63)

(cf. Definitions 3.4, 3.8, and 3.13). In the next step we note that, for instance, item (vi) in Proposition 2.6 in [37], item (ii) in Corollary 2.23 in [37], and the fact that for all $k \in$

 $\{0, 1, \ldots, K\}$ it holds that $\mathcal{R}_a(\mathbf{F}_k) = f_k$ ensure that for all $t \in \mathbb{R}, x \in \mathbb{R}^d, k \in \{0, 1, \ldots, K\}$ it holds that

$$(\mathcal{R}_{a}(\Gamma \bullet P_{2,\mathfrak{J}}(\mathbf{H}_{k}, \mathbf{F}_{k})))(t, x) = (\mathcal{R}_{a}(\Gamma))(\mathcal{R}_{a}(P_{2,\mathfrak{J}}(\mathbf{H}_{k}, \mathbf{F}_{k}))(t, x)) = (\mathcal{R}_{a}(\Gamma))((\mathcal{R}_{a}(\mathbf{H}_{k}))(t), f_{k}(x)).$$

$$(4.64)$$

This and (4.63) imply item (ii). Observe that item (ii) and the triangle inequality show that for all $t \in \mathbb{R}$, $x \in \mathbb{R}^d$ it holds that

$$\left| \left(\mathcal{R}_{a}(\Phi) \right)(t,x) - \sum_{k=0}^{K} h_{k}(t) f_{k}(x) \right| \leq \sum_{k=0}^{K} \left| \left(\mathcal{R}_{a}(\Gamma) \right) \left(\left(\mathcal{R}_{a}(\mathbf{H}_{k}) \right)(t), f_{k}(x) \right) - h_{k}(t) f_{k}(x) \right|.$$
(4.65)

In the next step we combine the triangle inequality, the fact that for all $v, w \in \mathbb{R}$ it holds that $|vw - (\mathcal{R}_a(\Gamma))(v, w)| \leq \varepsilon \max\{1, |v|^q, |w|^q\}$, and the fact that for all $t \in \mathbb{R}, k \in \{0, 1, \ldots, K\}$ it holds that $|h_k(t) - (\mathcal{R}_a(\mathbf{H}_k))(t)| \leq \varepsilon \max\{1, |t|^q\}$ to obtain that for all $t \in \mathbb{R}, x \in \mathbb{R}^d$, $k \in \{0, 1, \ldots, K\}$ it holds that

$$\begin{aligned} \left| \left(\mathcal{R}_{a}(\Gamma) \right) \left(\left(\mathcal{R}_{a}(\mathbf{H}_{k}) \right)(t), f_{k}(x) \right) - h_{k}(t) f_{k}(x) \right| \\ &\leq \left| \left(\mathcal{R}_{a}(\Gamma) \right) \left(\left(\mathcal{R}_{a}(\mathbf{H}_{k}) \right)(t), f_{k}(x) \right) - \left(\mathcal{R}_{a}(\mathbf{H}_{k}) \right)(t) f_{k}(x) \right| + \left| \left(\mathcal{R}_{a}(\mathbf{H}_{k}) \right)(t) - h_{k}(t) \right| |f_{k}(x)| \\ &\leq \varepsilon \max \left\{ 1, \left| \left(\mathcal{R}_{a}(\mathbf{H}_{k}) \right)(t) \right|^{q}, |f_{k}(x)|^{q} \right\} + \varepsilon \max \left\{ 1, |t|^{q} \right\} |f_{k}(x)| \\ &\leq \varepsilon \max \left\{ 1, \left| \left(\mathcal{R}_{a}(\mathbf{H}_{k}) \right)(t) \right|^{q} \right\} \left(1 + |f_{k}(x)|^{q} \right) + \varepsilon \max \left\{ 1, |t|^{q} \right\} \left(1 + |f_{k}(x)|^{q} \right). \end{aligned}$$

$$(4.66)$$

In the next step we note that the triangle inequality and the fact that for all $t \in \mathbb{R}$, $k \in \{0, 1, \ldots, K\}$ it holds that $|h_k(t) - (\mathcal{R}_a(\mathbf{H}_k))(t)| \leq \varepsilon \max\{1, |t|^q\}$ prove that for all $t \in \mathbb{R}$, $k \in \{0, 1, \ldots, K\}$ it holds that

$$\left| \left(\mathcal{R}_a(\mathbf{H}_k) \right)(t) \right| \le \left| \left(\mathcal{R}_a(\mathbf{H}_k) \right)(t) - h_k(t) \right| + |h_k(t)| \le \max\{1, |t|^q\} + |h_k(t)|.$$

$$(4.67)$$

This and (4.66) demonstrate that for all $t \in \mathbb{R}$, $x \in \mathbb{R}^d$, $k \in \{0, 1, \dots, K\}$ it holds that

$$\left| \left(\mathcal{R}_a(\Gamma) \right) \left(\left(\mathcal{R}_a(\mathbf{H}_k) \right)(t), f_k(x) \right) - h_k(t) f_k(x) \right| \le 2\varepsilon \left(\max\{1, |t|^q\} + |h_k(t)| \right)^q \left(1 + |f_k(x)|^q \right).$$

$$\tag{4.68}$$

This and (4.65) establish item (iii). Observe that item (i) in Lemma 3.21 and, for example, item (ii) in Proposition 2.6 in [37] ensure that

$$\mathcal{L}(\Phi) = \max_{k \in \{0,1,\dots,K\}} \mathcal{L}\left(\Gamma \bullet \left(\mathcal{P}_{2,\mathfrak{J}}(\mathbf{H}_k, \mathbf{F}_k)\right)\right) = \max_{k \in \{0,1,\dots,K\}} \left(\mathcal{L}(\Gamma) + \mathcal{L}\left(\mathcal{P}_{2,\mathfrak{J}}(\mathbf{H}_k, \mathbf{F}_k)\right) - 1\right)$$
(4.69)

(cf. Definition 3.1). In addition, note that, for instance, item (ii) in Lemma 2.13 in [37] implies that for all $k \in \{0, 1, ..., K\}$ it holds that

$$\mathcal{L}(\mathbf{P}_{2,\mathfrak{J}}(\mathbf{H}_{k},\mathbf{F}_{k})) = \mathcal{L}(\mathbf{P}_{2}(\mathcal{E}_{\max\{\mathcal{L}(\mathbf{H}_{k}),\mathcal{L}(\mathbf{F}_{k})\},\mathfrak{J}}(\mathbf{H}_{k}),\mathcal{E}_{\max\{\mathcal{L}(\mathbf{H}_{k}),\mathcal{L}(\mathbf{F}_{k})\},\mathfrak{J}}(\mathbf{F}_{k}))))$$

$$= \max\{\mathcal{L}(\mathbf{H}_{k}),\mathcal{L}(\mathbf{F}_{k})\}$$

$$\leq \left[\max_{j\in\{0,1,\dots,K\}}\mathcal{L}(\mathbf{H}_{j})\right]\left[\max_{j\in\{0,1,\dots,K\}}\mathcal{L}(\mathbf{F}_{j})\right]$$
(4.70)

(cf. Definitions 3.11 and 3.12). This and (4.69) show item (iv). Observe that item (iii) in Lemma 3.21 proves that

$$\left\|\left|\mathcal{D}(\Phi)\right\|\right| \le (K+1) \max\left\{\mathfrak{d}, \max_{k \in \{0,1,\dots,K\}} \left\|\left|\mathcal{D}\left(\Gamma \bullet \mathcal{P}_{2,\mathfrak{J}}(\mathbf{H}_{k}, \mathbf{F}_{k})\right)\right|\right\|\right\}$$
(4.71)

(cf. Definition 2.1). In addition, note that, for example, item (i) in Proposition 2.6 in [37] demonstrates that for all $k \in \{0, 1, ..., K\}$ it holds that

$$\left\| \left| \mathcal{D} \left(\Gamma \bullet \mathcal{P}_{2,\mathfrak{J}}(\mathbf{H}_{k}, \mathbf{F}_{k}) \right) \right\| \right\| \leq \max \left\{ \left\| \left| \mathcal{D} \left(\mathcal{P}_{2,\mathfrak{J}}(\mathbf{H}_{k}, \mathbf{F}_{k}) \right) \right\| \right\|, \left\| \mathcal{D} (\Gamma) \right\| \right\}.$$
(4.72)

Next we combine, for instance, item (i) in Proposition 2.20 in [37] and, for example, [54, Lemma 2.2.11] to obtain that for all $k \in \{0, 1, ..., K\}$ it holds that

$$\begin{aligned} \left\| \left\| \mathcal{D} \left(\mathbf{P}_{2, \mathfrak{J}}(\mathbf{H}_{k}, \mathbf{F}_{k}) \right) \right\| &= \left\| \left\| \mathcal{D} \left(\mathbf{P}_{2} \left(\mathcal{E}_{\max\{\mathcal{L}(\mathbf{H}_{k}), \mathcal{L}(\mathbf{F}_{k})\}, \mathfrak{J}}(\mathbf{H}_{k}), \mathcal{E}_{\max\{\mathcal{L}(\mathbf{H}_{k}), \mathcal{L}(\mathbf{F}_{k})\}, \mathfrak{J}}(\mathbf{F}_{k}) \right) \right) \right\| \\ &\leq \left\| \left\| \mathcal{D} \left(\mathcal{E}_{\max\{\mathcal{L}(\mathbf{H}_{k}), \mathcal{L}(\mathbf{F}_{k})\}, \mathfrak{J}}(\mathbf{H}_{k}) \right) \right\| \right\| + \left\| \left\| \mathcal{D} \left(\mathcal{E}_{\max\{\mathcal{L}(\mathbf{H}_{k}), \mathcal{L}(\Xi_{k})\}, \mathfrak{J}}(\mathbf{F}_{k}) \right) \right\| \right\| \\ &\leq \max\left\{ \mathfrak{d}, \left\| \left\| \mathcal{D}(\mathbf{H}_{k}) \right\| \right\} + \max\left\{ \mathfrak{d}, \left\| \left\| \mathcal{D}(\mathbf{F}_{k}) \right\| \right\} \\ &\leq \mathfrak{d} \left\| \left\| \mathcal{D}(\mathbf{H}_{k}) \right\| + \mathfrak{d} \left\| \left\| \mathcal{D}(\mathbf{F}_{k}) \right\| \right\| \leq 2\mathfrak{d} \left\| \left| \mathcal{D}(\mathbf{H}_{k}) \right\| \right\| \right\| \mathcal{D}(\mathbf{F}_{k}) \right\|. \end{aligned}$$

$$(4.73)$$

This and (4.72) establish that for all $k \in \{0, 1, \dots, K\}$ it holds that

$$\left\| \left| \mathcal{D} \left(\Gamma \bullet \mathcal{P}_{2,\mathfrak{J}}(\mathbf{H}_k, \mathbf{F}_k) \right) \right\| \le 2\mathfrak{d} \left\| \mathcal{D}(\mathbf{H}_k) \right\| \left\| \mathcal{D}(\mathbf{F}_k) \right\| \left\| \mathcal{D}(\Gamma) \right\|.$$

$$(4.74)$$

This and (4.71) ensure item (v). Observe that (3.2), (3.3), and (3.4) imply that

$$\mathcal{P}(\Phi) = \sum_{m=1}^{\mathcal{L}(\Phi)} \mathbb{D}_{m}(\Phi) \left(\mathbb{D}_{m-1}(\Phi) + 1 \right) \le \sum_{m=1}^{\mathcal{L}(\Phi)} \||\mathcal{D}(\Phi)||| \left(|||\mathcal{D}(\Phi)||| + |||\mathcal{D}(\Phi)||| \right) = 2\mathcal{L}(\Phi) |||\mathcal{D}(\Phi)|||^{2}.$$
(4.75)

Next, note that item (iv) and, for instance, [1, Lemma 2.4] show that

$$\mathcal{L}(\Phi) \le 2\mathcal{P}(\Gamma) \left[\max_{k \in \{0,1,\dots,K\}} \mathcal{P}(\mathbf{H}_k) \right] \left[\max_{k \in \{0,1,\dots,K\}} \mathcal{L}(\mathbf{F}_k) \right].$$
(4.76)

In the next step we combine item (v) and, for example, [1, Lemma 2.4] to obtain that

$$\||\mathcal{D}(\Phi)|\| \le 2\mathfrak{d}(K+1)\mathcal{P}(\Gamma)\left[\max_{k\in\{0,1,\dots,K\}}\mathcal{P}(\mathbf{H}_k)\right]\left[\max_{k\in\{0,1,\dots,K\}}\||\mathcal{D}(\mathbf{F}_k)\||\right].$$
(4.77)

Combining this, (4.75), and (4.76) proves item (vi). The proof of Proposition 4.12 is thus complete.

4.6 ANN approximations for linear interpolations of MLP approximations

Corollary 4.13. Let $K, d, \mathfrak{d}, M \in \mathbb{N}$, $T \in (0, \infty)$, $q \in (2, \infty)$, $\mathfrak{t}_{-1}, \mathfrak{t}_0, \mathfrak{t}_1, \ldots, \mathfrak{t}_K, \mathfrak{t}_{K+1} \in \mathbb{R}$ satisfy $\mathfrak{t}_{-1} < 0 = \mathfrak{t}_0 < \mathfrak{t}_1 < \ldots < \mathfrak{t}_K = T < \mathfrak{t}_{K+1}$, let $a \in C(\mathbb{R}, \mathbb{R})$, $\mathfrak{J}, \mathbf{F}, \mathbf{G} \in \mathbf{N}$ satisfy $\mathcal{D}(\mathfrak{J}) = (1, \mathfrak{d}, 1)$, $\mathcal{R}_a(\mathfrak{J}) = \mathrm{id}_{\mathbb{R}}$, $\mathcal{R}_a(\mathbf{F}) \in C(\mathbb{R}, \mathbb{R})$, and $\mathcal{R}_a(\mathbf{G}) \in C(\mathbb{R}^d, \mathbb{R})$, for every $k \in \{0, 1, \ldots, K\}$, $\gamma \in (0, 1]$ let $\mathbf{H}_{k, \gamma} \in \mathbf{N}$ satisfy for all $t \in \mathbb{R}$ that $\mathcal{R}_a(\mathbf{H}_{k, \gamma}) \in C(\mathbb{R}, \mathbb{R})$ and

$$\left|\mathscr{L}^{0,1,0}_{\mathfrak{t}_{k-1},\mathfrak{t}_k,\mathfrak{t}_{k+1}}(t) - \left(\mathcal{R}_a(\mathbf{H}_{k,\gamma})\right)(t)\right| \le \gamma \max\{1, |t|^q\},\tag{4.78}$$

for every $\gamma \in (0,1]$ let $\Gamma_{\gamma} \in \mathbf{N}$ satisfy for all $v, w \in \mathbb{R}$ that $\mathcal{R}_{a}(\Gamma_{\gamma}) \in C(\mathbb{R}^{2},\mathbb{R})$ and

$$\left|vw - \left(\mathcal{R}_{a}(\Gamma_{\gamma})\right)(v,w)\right| \leq \gamma \max\left\{1, |v|^{q}, |w|^{q}\right\},\tag{4.79}$$

let $\Theta = \bigcup_{n \in \mathbb{N}} \mathbb{Z}^n$, for every $\theta \in \Theta$ let $\mathcal{U}^{\theta} \colon [0,T] \to [0,T]$ and $W^{\theta} \colon [0,T] \to \mathbb{R}^d$ be functions, let $U_n^{\theta} \colon [0,T] \times \mathbb{R}^d \to \mathbb{R}$, $n \in \mathbb{N}_0$, $\theta \in \Theta$, satisfy for all $\theta \in \Theta$, $n \in \mathbb{N}$, $t \in [0,T]$, $x \in \mathbb{R}^d$ that $U_0^{\theta}(t,x) = 0$ and

$$U_{n}^{\theta}(t,x) = \frac{1}{M^{n}} \left[\sum_{k=1}^{M^{n}} (\mathcal{R}_{a}(\mathbf{G})) (x + W_{T-t}^{(\theta,0,-k)}) \right]$$

$$+ \sum_{i=0}^{n-1} \frac{(T-t)}{M^{n-i}} \left[\sum_{k=1}^{M^{n-i}} ((\mathcal{R}_{a}(\mathbf{F}) \circ U_{i}^{(\theta,i,k)}) - \mathbb{1}_{\mathbb{N}}(i) (\mathcal{R}_{a}(\mathbf{F}) \circ U_{\max\{i-1,0\}}^{(\theta,-i,k)})) (\mathcal{U}_{t}^{(\theta,i,k)}, x + W_{\mathcal{U}_{t}^{(\theta,i,k)}-t}^{(\theta,i,k)}) \right],$$
(4.80)

let $\mathbf{U}_{n,t}^{\theta} \in \{\Phi \in \mathbf{N} : \mathcal{R}_{a}(\Phi) \in C(\mathbb{R}^{d}, \mathbb{R})\}, t \in [0, T], n \in \mathbb{N}_{0}, \theta \in \Theta, \text{ satisfy for all } \theta \in \Theta, n \in \mathbb{N}, t \in [0, T] \text{ that } \mathbf{U}_{0,t}^{\theta} = ((0 \ 0 \ \dots \ 0), 0) \in \mathbb{R}^{1 \times d} \times \mathbb{R}^{1} \text{ and}$

$$\mathbf{U}_{n,t}^{\theta} = \begin{bmatrix} M^{n} \oplus \left(\frac{1}{M^{n}} \circledast \left(\mathbf{G} \bullet \mathbf{A}_{\mathbf{I}_{d}, W_{T-t}^{(\theta,0,-k)}} \right) \right) \end{bmatrix} \\
\equiv \mathfrak{J} \begin{bmatrix} n-1 \oplus \left(\frac{(T-t)}{M^{n-i}} \right) \circledast \left(M^{n-i} \oplus \left(\left(\mathbf{F} \bullet \mathbf{U}_{i,\mathcal{U}_{t}^{(\theta,i,k)}} \right) \bullet \mathbf{A}_{\mathbf{I}_{d}, W_{\mathcal{U}_{t}^{(\theta,i,k)}-t}} \right) \right) \end{bmatrix} \end{bmatrix} \\
\equiv \mathfrak{J} \begin{bmatrix} n-1 \oplus \left(\frac{(t-T) \mathbb{I}_{\mathbb{N}}(i)}{M^{n-i}} \right) \circledast \left(M^{n-i} \oplus \left(\left(\mathbf{F} \bullet \mathbf{U}_{i,\mathcal{U}_{t}^{(\theta,i,k)}} \right) \bullet \mathbf{A}_{\mathbf{I}_{d}, W_{\mathcal{U}_{t}^{(\theta,i,k)}-t}} \right) \right) \end{bmatrix} \end{bmatrix} \end{bmatrix}$$

$$(4.81)$$

and for every $\theta \in \Theta$, $\gamma \in (0, 1]$, $n \in \mathbb{N}$ let $\Phi_{\gamma,n}^{\theta} \in \mathbf{N}$ satisfy

$$\Phi_{\gamma,n}^{\theta} = \bigoplus_{k=0,\mathfrak{J}}^{K} \left(\Gamma_{\gamma} \bullet \mathcal{P}_{2,\mathfrak{J}} \left(\mathbf{H}_{k,\gamma}, \mathbf{U}_{n,\mathfrak{t}_{k}}^{\theta} \right) \right)$$
(4.82)

(cf. Definitions 3.1, 3.4, 3.5, 3.8, 3.9, 3.13, 3.15, 3.19, 3.20, and 4.1). Then

(i) it holds for all $\theta \in \Theta$, $\gamma \in (0,1]$, $n \in \mathbb{N}$, $t \in \mathbb{R}$, $x \in \mathbb{R}^d$ that $\mathcal{R}_a(\Phi_{\gamma,n}^\theta) \in C(\mathbb{R}^{d+1},\mathbb{R})$ and

$$\left(\mathcal{R}_{a}(\Phi_{\gamma,n}^{\theta})\right)(t,x) = \sum_{k=0}^{K} \left(\mathcal{R}_{a}(\Gamma_{\gamma})\right) \left(\left(\mathcal{R}_{a}(\mathbf{H}_{k,\gamma})\right)(t), U_{n}^{\theta}(\mathbf{t}_{k},x)\right),$$
(4.83)

(ii) it holds for all $\theta \in \Theta$, $\gamma \in (0, 1]$, $n \in \mathbb{N}$, $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$\left| \mathscr{L}_{\mathfrak{t}_{0},\mathfrak{t}_{1},\ldots,\mathfrak{t}_{K}}^{U_{n}^{\theta}(\mathfrak{t}_{0},x),U_{n}^{\theta}(\mathfrak{t}_{1},x),\ldots,U_{n}^{\theta}(\mathfrak{t}_{K},x)}(t) - \left(\mathcal{R}_{a}(\Phi_{\gamma,n}^{\theta}) \right)(t,x) \right| \\ \leq 2\gamma \left(1 + (T+1)^{q} \right)^{q} \left(\sum_{k=0}^{K} \left(1 + |U_{n}^{\theta}(\mathfrak{t}_{k},x)|^{q} \right) \right),$$

$$(4.84)$$

and

(iii) it holds for all
$$\theta \in \Theta$$
, $\gamma \in (0, 1]$, $n \in \mathbb{N}$ that

$$\mathcal{P}\left(\Phi_{\gamma,n}^{\theta}\right) \leq 16\left(\max\{\mathfrak{d},\mathcal{L}(\mathbf{G})\} + \mathcal{L}(\mathbf{F})\right) \left[\max\{\mathfrak{d}, \||\mathcal{D}(\mathbf{F})\||, \||\mathcal{D}(\mathbf{G})\||\}\right]^{2} \left[n^{\frac{1}{2}}(3M)^{n}\right]^{2} \\ \cdot \left[\mathcal{P}(\Gamma_{\gamma})\right]^{3} \left[\max_{k\in\{0,1,\dots,K\}}\mathcal{P}(\mathbf{H}_{k,\gamma})\right]^{3} (K+1)^{2} \mathfrak{d}^{2}$$

$$(4.85)$$

(cf. Definition 2.1).

Proof of Corollary 4.13. Observe that items (ii) and (v) in Proposition 3.9 in [1] and items (i) and (ii) in Proposition 4.12 (applied for every $\gamma \in (0, 1]$, $n \in \mathbb{N}$, $\theta \in \Theta$ with $K \curvearrowleft K$, $d \curvearrowleft d$, $\mathfrak{d} \frown \mathfrak{d}$, $\varepsilon \frown \gamma$, $q \frown q$, $a \frown a$, $(h_k)_{k \in \{0,1,\ldots,K\}} \frown (\mathscr{L}^{0,1,0}_{\mathfrak{t}_{k-1},\mathfrak{t}_k,\mathfrak{t}_{k+1}})_{k \in \{0,1,\ldots,K\}}$, $(f_k)_{k \in \{0,1,\ldots,K\}} \frown (\mathbb{R}^d \ni x \mapsto U^{\theta}_n(\mathfrak{t}_k, x) \in \mathbb{R})_{k \in \{0,1,\ldots,K\}}$, $(\mathbf{H}_k)_{k \in \{0,1,\ldots,K\}} \frown (\mathbf{H}_{k,\gamma})_{k \in \{0,1,\ldots,K\}}$, $(\mathbf{F}_k)_{k \in \{0,1,\ldots,K\}} \frown (\mathbf{U}^{\theta}_{n,\mathfrak{t}_k})_{k \in \{0,1,\ldots,K\}}, \mathfrak{J} \frown \mathfrak{J}, \Gamma \frown \Gamma_{\gamma}, \Phi \frown \Phi^{\theta}_{\gamma,n}$ in the notation of Proposition 4.12) demonstrate item (i). Note that item (iii) in Proposition 4.12 and Lemma 4.4 establish that for all $t \in [0,T], x \in \mathbb{R}^d, \gamma \in (0,1], n \in \mathbb{N}, \theta \in \Theta$ it holds that

$$\left| \mathscr{L}_{\mathfrak{t}_{0},\mathfrak{t}_{1},\ldots,\mathfrak{t}_{K}}^{U_{n}^{\theta}(\mathfrak{t}_{0},x),U_{n}^{\theta}(\mathfrak{t}_{1},x),\ldots,U_{n}^{\theta}(\mathfrak{t}_{K},x)}(t) - \left(\mathcal{R}_{a}(\Phi_{\gamma,n}^{\theta}) \right)(t,x) \right| \\ \leq 2\gamma \sum_{k=0}^{K} \left(1 + |U_{n}^{\theta}(\mathfrak{t}_{k},x)|^{q} \right) \left(\max\{1,|t|^{q}\} + \left| \mathscr{L}_{\mathfrak{t}_{k-1},\mathfrak{t}_{k},\mathfrak{t}_{k+1}}^{0,1,0}(t) \right| \right)^{q}.$$

$$(4.86)$$

The fact that for all $t \in [0, T]$ it holds that $\max\{1, |t|^q\} \leq (1+T)^q$ and the fact that for all $t \in \mathbb{R}, k \in \{0, 1, \ldots, K\}$ it holds that $\mathscr{L}^{0,1,0}_{\mathfrak{t}_{k-1},\mathfrak{t}_k,\mathfrak{t}_{k+1}}(t) \in [0, 1]$ hence ensure item (ii). Observe that item (iii) in Proposition 3.9 in [1] implies that for all $n \in \mathbb{N}, \theta \in \Theta, t \in [0, T]$ it holds that

$$\mathcal{L}(\mathbf{U}_{n,t}^{\theta}) \le \max\{\mathfrak{d}, \mathcal{L}(\mathbf{G})\} + n\mathcal{H}(\mathbf{F}) \le n\big(\max\{\mathfrak{d}, \mathcal{L}(\mathbf{G})\} + \mathcal{L}(\mathbf{F})\big).$$
(4.87)

Furthermore, note that item (iv) in Proposition 3.9 in [1] shows that for all $n \in \mathbb{N}$, $\theta \in \Theta$, $t \in [0, T]$ it holds that

$$\left\|\left|\mathcal{D}(\mathbf{U}_{n,t}^{\theta})\right\|\right\|^{2} \leq \left(\max\{\mathfrak{d}, \left\|\left|\mathcal{D}(\mathbf{F})\right|\right\|, \left\|\left|\mathcal{D}(\mathbf{G})\right|\right\|\}\right)^{2} \left((3M)^{n}\right)^{2}$$
(4.88)

(cf. Definition 2.1). This, (4.87), and item (vi) in Proposition 4.12 prove item (iii). The proof of Corollary 4.13 is thus complete.

5 ANN approximations for solutions of semilinear heat PDEs

In this section we establish the main ANN approximation results of this work. In particular, in Theorem 5.3 in Subsection 5.1 we show that for every arbitrarily large absolute moment $\mathbf{q} \in [2, \infty)$ and every arbitrarily large time horizon $T \in (0, \infty)$ we have that the solutions $u_d: [0, T] \times \mathbb{R}^d \to \mathbb{R}, d \in \mathbb{N}$, of the semilinear heat PDEs in (5.18) below can be approximated on $[0, T] \times \mathbb{R}^d, d \in \mathbb{N}$, in the L^q -sense with respect to the measures $\nu_d: \mathcal{B}(\mathbb{R}^{d+1}) \to [0, \infty),$ $d \in \mathbb{N}$, in (5.10) (see (5.19)) without the COD (see (5.20)) through realizations of ANNs with a general activation function provided that there exist

- (i) appropriate ANN approximations $\Gamma_{\varepsilon} \in \mathbf{N}, \ \varepsilon \in (0, 1]$, for the product function (see (5.11) and (5.12) below),
- (ii) a shallow ANN representation $\mathfrak{J} \in \mathbf{N}$ for the one-dimensional identity function $\mathrm{id}_{\mathbb{R}} = (\mathbb{R} \ni x \mapsto x \in \mathbb{R}),$
- (iii) appropriate ANN approximations $\mathbf{H}_{K,k,\varepsilon} \in \mathbf{N}, \varepsilon \in (0,1], k \in \{0,1,\ldots,K\}, K \in \mathbb{N}$, for the hat functions (see (5.13) and (5.14) below),
- (iv) appropriate ANN approximations $\mathbf{F}_{0,\varepsilon} \in \mathbf{N}$, $\varepsilon \in (0, 1]$, for the nonlinearity in the PDE (see (5.15), (5.16), and (5.17) below), and
- (v) appropriate ANN approximations $\mathbf{F}_{d,\varepsilon} \in \mathbf{N}, \varepsilon \in (0,1], d \in \mathbb{N}$, for the terminal value functions (see (5.15) and (5.16) below).

In our proof of Theorem 5.3 we employ, among other things, Corollary 4.13 from Section 4, Corollary 2.3, Lemma 2.5, and Corollary 2.7 from Section 2, the strong L^p -error estimates for the employed MLP approximations from Hutzenthaler et al. [48], the elementary complexity estimate in Lemma 5.1, and the elementary and well-known measurability property in Lemma 5.2.

In Corollary 5.4 and Corollary 5.5 in Subsection 5.2 we then specialize Theorem 5.3 to the situation of ANNs with the ReLU, the leaky ReLU, and the softplus activation function. Our proofs of Corollary 5.4 and Corollary 5.5, respectively, employ the general ANN approximation result for PDEs in Theorem 5.3, the ANN representation and approximation results for hat functions in Subsection 4.2, the ANN approximation results for the product function from Subsection 4.4, the ANN representation results for the one-dimensional identity function in [1, Section 3.2], and the ANN approximation results for Lipschitz continuous nonlinearities in [1, Section 4.2].

5.1 ANN approximations for PDEs with general activation functions

Lemma 5.1. Let $L, T \in (0, \infty)$, $p \in [2, \infty)$, $\mathfrak{B} \in [1, \infty)$, $(m_k)_{k \in \mathbb{N}} \subseteq \mathbb{N}$ satisfy for all $k \in \mathbb{N}$ that $\liminf_{j \to \infty} m_j = \infty$, $\limsup_{j \to \infty} \frac{(m_j)^{p/2}}{j} < \infty$, and $m_{k+1} \leq \mathfrak{B}m_k$, and let $(N_{\varepsilon})_{\varepsilon \in (0,1]}$ satisfy for all $\varepsilon \in (0, 1]$ that

$$N_{\varepsilon} = \inf\left(\left\{n \in \mathbb{N} \colon \left[(1 + LT)(m_n)^{-\frac{1}{2}} \exp\left(\frac{(m_n)^{\frac{p}{2}}}{n}\right)\right]^n \le \varepsilon\right\} \cup \{\infty\}\right).$$
(5.1)

Then

- (i) it holds for all $\varepsilon \in (0, 1]$ that $N_{\varepsilon} < \infty$ and
- (ii) it holds for all $\delta \in (0, \infty)$ that

$$\sup_{\varepsilon \in (0,1]} \left(\varepsilon^{2+\delta} (N_{\varepsilon})^{1/2} (3m_{N_{\varepsilon}})^{N_{\varepsilon}} \right) < \infty.$$
(5.2)

Proof of Lemma 5.1. Throughout this proof let $(\mathfrak{m}_n)_{n\in\mathbb{N}}$ satisfy for all $n\in\mathbb{N}$ that

$$\mathfrak{m}_{n} = \left[(1+2LT)(m_{n})^{-\frac{1}{2}} \exp\left(\frac{(m_{n})^{\frac{p}{2}}}{n}\right) \right]^{n}.$$
(5.3)

Observe that the fact that $\limsup_{j\to\infty} {(m_j)^{p/2}/j} < \infty$ and the fact that $\liminf_{j\to\infty} m_j = \infty$ show that $\limsup_{n\to\infty} \mathfrak{m}_n = 0$. Combining this and (5.1) proves item (i). Note that (5.1) and (5.3) demonstrate that for all $\varepsilon \in (0, 1]$ with $N_{\varepsilon} \in \mathbb{N} \cap [2, \infty)$ it holds that $\mathfrak{m}_{N_{\varepsilon}-1} > \varepsilon$. Hence, (5.3) shows that for all $\delta \in (0, \infty)$, $\varepsilon \in (0, 1]$ with $N_{\varepsilon} \in \mathbb{N} \cap [2, \infty)$ it holds that

$$(N_{\varepsilon})^{1/2} (3m_{N_{\varepsilon}})^{N_{\varepsilon}} \leq (N_{\varepsilon})^{1/2} (3m_{N_{\varepsilon}})^{N_{\varepsilon}} (\varepsilon^{-1} \mathfrak{m}_{N_{\varepsilon}-1})^{2+\delta} \leq \varepsilon^{-(2+\delta)} \sup_{n \in \mathbb{N}} \left((n+1)^{\frac{1}{2}} (3m_{n+1})^{n+1} (\mathfrak{m}_{n})^{2+\delta} \right)$$

$$\leq \varepsilon^{-(2+\delta)} \sup_{n \in \mathbb{N}} \left((n+1)^{\frac{1}{2}} (m_{n+1})^{n+1} \left[3(1+2LT) \exp\left(\frac{(m_{n})^{\frac{p}{2}}}{n}\right) (m_{n})^{-\frac{1}{2}} \right]^{n(2+\delta)} \right).$$
(5.4)

Next, observe that the fact that for all $n \in \mathbb{N}$ it holds that $m_{n+1} \leq \mathfrak{B}m_n$ establishes that for all $n \in \mathbb{N}$ it holds that $(m_{n+1})^{n+1} \leq m_{n+1}\mathfrak{B}^n(m_n)^n \leq m_1\mathfrak{B}^{2n}(m_n)^n$. Combining this and (5.4) establishes that for all $\delta \in (0, \infty)$, $\varepsilon \in (0, 1]$ with $N_{\varepsilon} \in \mathbb{N} \cap [2, \infty)$ it holds that

$$(N_{\varepsilon})^{1/2} (3m_{N_{\varepsilon}})^{N_{\varepsilon}} \le \varepsilon^{-(2+\delta)} \sup_{n \in \mathbb{N}} \left((n+1)^{\frac{1}{2}} m_1 (m_n)^{-\frac{n\delta}{2}} \left[3(1+2LT) \mathfrak{B} \exp\left(\frac{(m_n)^{\frac{p}{2}}}{n}\right) \right]^{n(2+\delta)} \right).$$
(5.5)

This, the fact that $\mathfrak{B} \in [1,\infty)$, and the fact that for all $n \in \mathbb{N}$, $\delta \in (0,\infty)$ it holds that

$$(m_1)^{-\frac{\delta}{2}} \left(\exp\left((m_1)^{\frac{p}{2}}\right) \right)^{2+\delta} \ge (m_1)^{-\frac{\delta}{2}} \left(1 + (m_1)^{\frac{p}{2}}\right)^{2+\delta} \ge (m_1)^{\frac{(p-1)\delta}{2}+p} \ge 1$$
(5.6)

ensure that for all $\delta \in (0, \infty)$, $\varepsilon \in (0, 1]$ it holds that

$$(N_{\varepsilon})^{1/2} (3m_{N_{\varepsilon}})^{N_{\varepsilon}} \leq \varepsilon^{-(2+\delta)} m_1 \sup_{n \in \mathbb{N}} \left[(n+1)^{\frac{1}{2n}} (m_n)^{-\frac{\delta}{2}} \left[3(1+2LT)\mathfrak{B} \exp\left(\frac{(m_n)^{\frac{p}{2}}}{n}\right) \right]^{2+\delta} \right]^n.$$

$$(5.7)$$

Next we combine the fact that $\limsup_{n\to\infty} (m_n)^{p/2}/n < \infty$, the fact that $\liminf_{n\to\infty} m_n = \infty$, and the fact that $\lim_{n\to\infty} (n+1)^{1/(2n)} = 1$ to obtain that for all $\delta \in (0,\infty)$ it holds that

$$\sup_{n \in \mathbb{N}} \left[(n+1)^{\frac{1}{2n}} (m_n)^{-\frac{\delta}{2}} \left[3(1+2LT) \mathfrak{B} \exp\left(\frac{(m_n)^{\frac{p}{2}}}{n}\right) \right]^{2+\delta} \right]^n < \infty.$$
(5.8)

This and (5.7) imply item (ii). The proof of Lemma 5.1 is thus complete.

Lemma 5.2. Let (Ω, \mathcal{F}) be a measurable space, let $T \in (0, \infty)$, $d, K \in \mathbb{N}$, for every $k \in \{0, 1, \ldots, K\}$ let $h_k \colon [0, T] \to \mathbb{R}$ and $f_k \colon \mathbb{R}^d \times \Omega \to \mathbb{R}$ be measurable, and let $\psi \colon \mathbb{R}^2 \to \mathbb{R}$ be measurable. Then

$$[0,T] \times \mathbb{R}^d \times \Omega \ni (t,x,\omega) \mapsto \sum_{k=0}^K \psi(h_k(t), f_k(x,\omega)) \in \mathbb{R}$$
(5.9)

is measurable.

Proof of Lemma 5.2. Note that the fact that for all $k \in \{0, 1, ..., K\}$ it holds that h_k , f_k , and ψ are measurable shows that for all $k \in \{0, 1, ..., K\}$ it holds that $[0, T] \times \mathbb{R}^d \times \Omega \ni (t, x, \omega) = \psi(h_k(t), f_k(x, \omega))$ is measurable. This establishes (5.9). The proof of Lemma 5.2 is thus complete.

Theorem 5.3. Let $L, \kappa, \alpha_0, \alpha_1, \beta_0, \beta_1, T \in (0, \infty), r, p \in [1, \infty), q \in [2, \infty), q \in (2, \infty), a \in C(\mathbb{R}, \mathbb{R})$, for every $d \in \mathbb{N}_0$ let $f_d \in C^{\min\{d,1\}}(\mathbb{R}^{\max\{d,1\}}, \mathbb{R})$, for every $d \in \mathbb{N}$ let $\nu_d \colon \mathcal{B}(\mathbb{R}^{d+1}) \to [0, \infty)$ be a measure with

$$\int_{\mathbb{R}^{d+1}} (1 + \|y\|^{p^2 q \mathfrak{q}}) \nu_d(\mathrm{d}y) \le \kappa d^{r p^2 q \mathfrak{q}}, \tag{5.10}$$

let $\mathfrak{J} \in \mathbf{N}$ satisfy $\mathcal{H}(\mathfrak{J}) = 1$ and $\mathcal{R}_a(\mathfrak{J}) = \mathrm{id}_{\mathbb{R}}$, for every $\varepsilon \in (0, 1]$ let $\Gamma_{\varepsilon} \in \mathbf{N}$ satisfy for all $v, w \in \mathbb{R}$ that

$$\mathcal{R}_a(\Gamma_\varepsilon) \in C(\mathbb{R}^2, \mathbb{R}), \qquad \mathcal{P}(\Gamma_\varepsilon) \le \kappa \varepsilon^{-r}, \qquad and$$

$$(5.11)$$

$$|vw - (\mathcal{R}_a(\Gamma_{\varepsilon}))(v, w)| \le \varepsilon \max\{1, |v|^q, |w|^q\},$$
(5.12)

for every $K \in \mathbb{N}$, $k \in \{0, 1, \dots, K\}$, $\varepsilon \in (0, 1]$ let $\mathbf{H}_{K,k,\varepsilon} \in \mathbf{N}$ satisfy for all $t \in \mathbb{R}$ that

$$\mathcal{R}_{a}(\mathbf{H}_{K,k,\varepsilon}) \in C(\mathbb{R},\mathbb{R}), \qquad \mathcal{P}(\mathbf{H}_{K,k,\varepsilon}) \leq \kappa K^{r} \varepsilon^{-r}, \qquad and \qquad (5.13)$$

$$\left|\mathscr{L}^{0,1,0}_{\frac{(k-1)T}{K},\frac{kT}{K},\frac{(k+1)T}{K}}(t) - \left(\mathcal{R}_a(\mathbf{H}_{K,k,\varepsilon})\right)(t)\right| \le \varepsilon \max\{1,|t|^q\},\tag{5.14}$$

for every $d \in \mathbb{N}_0$, $\varepsilon \in (0, 1]$ let $\mathbf{F}_{d,\varepsilon} \in \mathbf{N}$ satisfy for all $x \in \mathbb{R}^{\max\{d,1\}}$, that

$$\mathcal{R}_{a}(\mathbf{F}_{d,\varepsilon}) \in C(\mathbb{R}^{\max\{d,1\}},\mathbb{R}), \quad \varepsilon^{\alpha_{\min\{d,1\}}} \mathcal{L}(\mathbf{F}_{d,\varepsilon}) + \varepsilon^{\beta_{\min\{d,1\}}} |||\mathcal{D}(\mathbf{F}_{d,\varepsilon})||| \le \kappa (\max\{d,1\})^{p},$$
(5.15)

and

$$\varepsilon \|\nabla f_{\max\{d,1\}}(x)\| + \varepsilon |(\mathcal{R}_a(\mathbf{F}_{d,\varepsilon}))(x)| + |f_d(x) - (\mathcal{R}_a(\mathbf{F}_{d,\varepsilon}))(x)| \le \varepsilon \kappa (\max\{d,1\})^p (1 + \|x\|)^p,$$
(5.16)

assume for all $v, w \in \mathbb{R}$, $\varepsilon \in (0, 1]$ that

$$\max\{|f_0(v) - f_0(w)|, |(\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(v) - (\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(w)|\} \le L|v - w|,$$
(5.17)

and let $\delta, \eta \in (0, \infty)$ satisfy $\eta = \max\{\alpha_0, \alpha_1\} + 2\max\{\beta_0, \beta_1\} + 2\delta + 24r + 8$ (cf. Definitions 2.1, 3.1, 3.4, and 4.1). Then

(i) for every $d \in \mathbb{N}$ there exists a unique at most polynomially growing viscosity solution $u_d \in C([0,T] \times \mathbb{R}^d, \mathbb{R})$ of

$$\frac{\partial}{\partial t}u_d(t,x) + \frac{1}{2}\Delta_x u_d(t,x) + f_0(u_d(t,x)) = 0$$
(5.18)

with $u_d(T, x) = f_d(x)$ for $(t, x) \in (0, T) \times \mathbb{R}^d$ and

(ii) there exists $c \in \mathbb{R}$ such that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ there exists $\mathbf{U} \in \{\Phi \in \mathbf{N} : \mathcal{R}_a(\Phi) \in C(\mathbb{R}^{d+1}, \mathbb{R})\}$ such that

$$\left[\int_{[0,T]\times\mathbb{R}^d} |u_d(y) - (\mathcal{R}_a(\mathbf{U}))(y)|^{\mathfrak{q}} \nu_d(\mathrm{d}y)\right]^{1/\mathfrak{q}} \le \varepsilon$$
(5.19)

and
$$\mathcal{P}(\mathbf{U}) \le cd^{(2(r+1)p^2 + (p+1)^2)(6r+\eta)q+3p} \varepsilon^{-\eta}.$$
 (5.20)

Proof of Theorem 5.3. Throughout this proof let $\mathfrak{B} \in [1,\infty)$, $(m_k)_{k\in\mathbb{N}} \subseteq \mathbb{N}$ satisfy for all $k \in \mathbb{N}$ that $\liminf_{j\to\infty} m_j = \infty$, $\limsup_{j\to\infty} (m_j)^{qq/2}/j < \infty$, and $m_{k+1} \leq \mathfrak{B}m_k$, let $\mathfrak{d} \in \mathbb{N}$ satisfy $\mathcal{D}(\mathfrak{J}) = (1,\mathfrak{d},1)$, let $\Theta = \bigcup_{n\in\mathbb{N}}\mathbb{Z}^n$, let $(\Omega,\mathcal{F},\mathbb{P})$ be a probability space, let $\mathfrak{u}^{\theta} \colon \Omega \to [0,1]$, $\theta \in \Theta$, be i.i.d. random variables, assume for all $t \in (0,1)$ that $\mathbb{P}(\mathfrak{u}^0 \leq t) = t$, let $\mathcal{U}^{\theta} \colon [0,T] \times \Omega \to [0,T]$, $\theta \in \Theta$, satisfy for all $t \in [0,T]$, $\theta \in \Theta$ that $\mathcal{U}^{\theta}_t = t + (T-t)\mathfrak{u}^{\theta}$, let $W^{d,\theta} \colon [0,T] \times \Omega \to \mathbb{R}^d$, $d \in \mathbb{N}$, $\theta \in \Theta$, be independent standard Brownian motions, assume for every $d \in \mathbb{N}$ that $(\mathcal{U}^{\theta})_{\theta\in\Theta}$ and $(W^{d,\theta})_{\theta\in\Theta}$ are independent, for every $d, j \in \mathbb{N}, \varepsilon \in (0,1]$ let $U^{d,\theta}_{n,j,\varepsilon} \colon [0,T] \times \mathbb{R}^d \times \Omega \to \mathbb{R}, n \in \mathbb{N}_0, \theta \in \Theta$, satisfy for all $\theta \in \Theta, n \in \mathbb{N}, t \in [0,T], x \in \mathbb{R}^d$, $\omega \in \Omega$ that $U^{d,\theta}_{0,j,\varepsilon}(t,x,\omega) = 0$ and

$$U_{n,j,\varepsilon}^{d,\theta}(t,x) = \frac{1}{(m_j)^n} \left[\sum_{k=1}^{(m_j)^n} (\mathcal{R}_a(\mathbf{F}_{d,\varepsilon})) (x + W_{T-t}^{d,(\theta,0,-k)}) \right] \\ + \sum_{i=0}^{n-1} \frac{(T-t)}{(m_j)^{n-i}} \left[\sum_{k=1}^{(m_j)^{n-i}} \left[((\mathcal{R}_a(\mathbf{F}_{0,\varepsilon})) (U_{i,j,\varepsilon}^{d,(\theta,i,k)} (\mathcal{U}_t^{(\theta,i,k)}, x + W_{\mathcal{U}_t^{(\theta,i,k)}-t}^{d,(\theta,i,k)})) - \mathbb{1}_{\mathbb{N}}(i) ((\mathcal{R}_a(\mathbf{F}_{0,\varepsilon})) (U_{\max\{i-1,0\},j,\varepsilon}^{d,(\theta,i,k)} (\mathcal{U}_t^{(\theta,i,k)}, x + W_{\mathcal{U}_t^{(\theta,i,k)}-t}^{d,(\theta,i,k)})) \right] \right],$$
(5.21)

for every $d, j \in \mathbb{N}, \varepsilon \in (0, 1]$ let $\mathbf{U}_{n, j, t}^{d, \theta, \varepsilon} \colon \Omega \to \{\Phi \in \mathbf{N} \colon \mathcal{R}_{a}(\Phi) \in C(\mathbb{R}^{d}, \mathbb{R})\}, t \in [0, T], n \in \mathbb{N}_{0}, \theta \in \Theta$, satisfy for all $\theta \in \Theta, n \in \mathbb{N}, t \in [0, T], \omega \in \Omega$ that $\mathbf{U}_{0, j, t}^{d, \theta, \varepsilon}(\omega) = ((0 \ 0 \ \dots \ 0), 0) \in \mathbb{R}^{1 \times d} \times \mathbb{R}^{1}$ and

for every $K \in \mathbb{N}, k \in \mathbb{Z}$ let $\mathfrak{t}_{K,k} \in \mathbb{R}$ satisfy $\mathfrak{t}_{K,k} = \frac{kT}{K}$, for every $d, n, j, K \in \mathbb{N}, \theta \in \Theta$, $\varepsilon, \gamma \in (0, 1]$ let $\Phi_{\gamma, n, j, K}^{d, \theta, \varepsilon} \colon \Omega \to \mathbf{N}$ satisfy for all $\omega \in \Omega$ that

$$\Phi_{\gamma,n,j,K}^{d,\theta,\varepsilon}(\omega) = \underset{k=0,\mathfrak{J}}{\overset{K}{\boxplus}} \Big[\Gamma_{\gamma} \bullet P_{2,\mathfrak{J}} \Big(\Big(\mathbf{H}_{K,k,\gamma}, \mathbf{U}_{n,j,\mathfrak{t}_{K,k}}^{d,\theta,\varepsilon}(\omega) \Big) \Big) \Big],$$
(5.23)

assume without loss of generality that $\max\{|f_0(0)|, \mathfrak{d}, 1\} \leq \kappa$, let $(\mathfrak{a}_d)_{d \in \mathbb{N}}$ satisfy for all $d \in \mathbb{N}$ that

$$\mathfrak{a}_d = e^{2LT} (T+1)^2 (L+1) (\kappa d^p + 1) 8^{p+2} d^{\frac{1}{2}}, \qquad (5.24)$$

let $(\mathfrak{b}_d)_{d\in\mathbb{N}}$ satisfy for all $d\in\mathbb{N}$ that

$$\mathbf{b}_{d} = 2^{p-1} \kappa d^{p} \left(e^{LT} (T+1) \right)^{p+1} \left((\kappa d^{p})^{p} + 1 \right) 3^{p^{2}-1},$$
(5.25)

let $(\mathfrak{c}_d)_{d\in\mathbb{N}}$ satisfy for all $d\in\mathbb{N}$ that

$$\mathbf{c}_d = 2^p \kappa d^p (T+1) e^{LT} \sqrt{q\mathbf{q} - 1},\tag{5.26}$$

let $(\mathfrak{m}_{j,n})_{(j,n)\in\mathbb{N}^2}$ satisfy for all $j,n\in\mathbb{N}$ that

$$\mathbf{m}_{j,n} = \left[(1+2LT)(m_j)^{-\frac{1}{2}} \exp\left(\frac{(m_j)^{\frac{qq}{2}}}{n}\right) \right]^n,$$
(5.27)

let $(c_d)_{d\in\mathbb{N}}$ satisfy for all $d\in\mathbb{N}$ that

$$c_d = \left[\int_{[0,T] \times \mathbb{R}^d} \left(1 + \|x\|^{p^2 q \mathfrak{q}} + \sup_{s \in [0,T]} \mathbb{E} \left[\|W_s^{d,0}\|^{2p^2 q \mathfrak{q}} \right] \right) \nu_d(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}},\tag{5.28}$$

let $(N_{\varepsilon})_{\varepsilon \in (0,1]}$ satisfy for all $\varepsilon \in (0,1]$ that

$$N_{\varepsilon} = \inf \left\{ n \in \mathbb{N} \colon \mathfrak{m}_{n,n} \le \varepsilon \right\},\tag{5.29}$$

let $(K_{\varepsilon})_{\varepsilon \in (0,1]}$ satisfy for all $\varepsilon \in (0,1]$ that

$$K_{\varepsilon} = \inf \{ n \in \mathbb{N} \colon n \ge \varepsilon^{-2} \}, \tag{5.30}$$

let $(\delta_{d,\varepsilon})_{(d,\varepsilon)\in\mathbb{N}\times(0,1]}$ satisfy for all $d\in\mathbb{N}, \varepsilon\in(0,1]$ that

$$\delta_{d,\varepsilon} = \frac{\varepsilon}{1 + 180(T+1)2^p(\mathbf{a}_d + \mathbf{b}_d + \mathbf{c}_d)c_d},\tag{5.31}$$

and let $(\gamma_{d,\varepsilon})_{(d,\varepsilon)\in\mathbb{N}\times(0,1]}$ satisfy for all $d\in\mathbb{N}, \varepsilon\in(0,1]$ that

$$\gamma_{d,\varepsilon} = \frac{\varepsilon}{2\kappa d^{rp^2q} (K_{\varepsilon} + 1) \left(1 + (T+1)^q\right)^q \left(1 + 3^{3p^2q} c_d \left((\mathfrak{c}_d)^q + (T+1)^q ((\mathfrak{a}_d)^q + (\mathfrak{b}_d)^q) + \kappa^q d^{pq}\right)\right)}$$
(5.32)

(cf. Definitions 2.1, 3.1, 3.4, 3.5, 3.8, 3.9, 3.13, 3.15, 3.19, and 3.20). Observe that the triangle inequality and the assumption that for all $d \in \mathbb{N}$, $x \in \mathbb{R}^d$, $\varepsilon \in (0, 1]$ it holds that

 $\varepsilon |(\mathcal{R}_a(\mathbf{F}_{d,\varepsilon}))(x)| + |f_d(x) - (\mathcal{R}_a(\mathbf{F}_{d,\varepsilon}))(x)| \le \varepsilon \kappa d^p (1 + ||x||)^p$ prove that for all $d \in \mathbb{N}, x \in \mathbb{R}^d$ it holds that

$$|f_d(x)| \le |f_d(x) - (\mathcal{R}_a(\mathbf{F}_{d,1}))(x)| + |(\mathcal{R}_a(\mathbf{F}_{d,1}))(x)| \le \kappa d^p (1 + ||x||)^p.$$
(5.33)

This, the assumption that for all $w, z \in \mathbb{R}$ it holds that $|f_0(w) - f_0(z)| \leq L|w-z|$, Beck et al. [10, Theorem 1.1] (applied for every $d \in \mathbb{N}$ with $d \frown d, m \frown d, L \frown L+1, T \frown T, \mu \frown (\mathbb{R}^d \ni x \mapsto (0, 0, \dots, 0) \in \mathbb{R}^d), \sigma \frown (\mathbb{R}^d \ni x \mapsto I_d \in \mathbb{R}^{d \times d}), f \frown (\mathbb{R}^d \times \mathbb{R} \ni (x, w) \mapsto f_0(w) \in \mathbb{R}), g \frown f_d, W \frown W^{d,0}$ in the notation of Beck et al. [10, Theorem 1.1]), and Fubini's theorem demonstrate that for every $d \in \mathbb{N}$ there exists a unique at most polynomially growing viscosity solution $u_d \in C([0,T] \times \mathbb{R}^d, \mathbb{R})$ of (5.18) with $u_d(T,x) = f_d(x), x \in \mathbb{R}^d$, and that it moreover holds for all $d \in \mathbb{N}, t \in [0,T], x \in \mathbb{R}^d$ that $\mathbb{E}[|f_d(x+W^{d,0}_{T-t})| + \int_t^T |f_0(u_d(s,x+W^{d,0}_{s-t}))| \, ds] < \infty$ and

$$u_d(t,x) = \mathbb{E}\left[f_d(x+W_{T-t}^{d,0})\right] + \int_t^T \mathbb{E}\left[f_0\left(u_d(s,x+W_{s-t}^{d,0})\right)\right] \mathrm{d}s.$$
(5.34)

This proves item (i). Note that the assumption that for all $w, z \in \mathbb{R}, \varepsilon \in (0, 1]$ it holds that $|(\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(z) - (\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(w)| \leq L|z-w|$, the assumption that for all $d \in \mathbb{N}_0, x \in \mathbb{R}^{\max\{d,1\}}, \varepsilon \in (0, 1]$ it holds that $|(\mathcal{R}_a(\mathbf{F}_{d,\varepsilon}))(x)| \leq \kappa (\max\{d,1\})^p (1+||x||)^p$, and Beck et al. [10, Theorem 1.1] (applied for every $d \in \mathbb{N}, \varepsilon \in (0, 1]$ with $d \curvearrowleft d, m \curvearrowleft d, L \curvearrowleft L+1, T \backsim T, \mu \curvearrowleft (\mathbb{R}^d \ni x \mapsto (0, 0, \ldots, 0) \in \mathbb{R}^d), \sigma \curvearrowleft (\mathbb{R}^d \ni x \mapsto I_d \in \mathbb{R}^{d \times d}), f \backsim (\mathbb{R}^d \times \mathbb{R} \ni (x, w) \mapsto (\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(w) \in \mathbb{R}), g \backsim \mathcal{R}_a(\mathbf{F}_{d,\varepsilon}), W \backsim W^{d,0}$ in the notation of Beck et al. [10, Theorem 1.1]) establish that for every $d \in \mathbb{N}, \varepsilon \in (0, 1]$ there exists a unique at most polynomially growing $v_{d,\varepsilon} \in C([0,T] \times \mathbb{R}^d, \mathbb{R})$ such that for all $t \in [0,T], x \in \mathbb{R}^d$ it holds that $\mathbb{E}[|(\mathcal{R}_a(\mathbf{F}_{d,\varepsilon}))(x+W^{d,0}_{T-t})| + \int_t^T |(\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(v_{d,\varepsilon}(s,x+W^{d,0}_{s-t}))| \, ds] < \infty$ and

$$v_{d,\varepsilon}(t,x) = \mathbb{E}\left[\left(\mathcal{R}_a(\mathbf{F}_{d,\varepsilon})\right)(x+W_{T-t}^{d,0})\right] + \int_t^T \mathbb{E}\left[\left(\mathcal{R}_a(\mathbf{F}_{0,\varepsilon})\right)\left(v_{d,\varepsilon}(s,x+W_{s-t}^{d,0})\right)\right] \mathrm{d}s.$$
(5.35)

Observe that item (i) in Lemma 3.3 in Hutzenthaler et al. [48] ensures that for all $\varepsilon \in (0, 1]$, $n \in \mathbb{N}_0, d, j \in \mathbb{N}, \theta \in \Theta$ it holds that $U_{n,j,\varepsilon}^{d,\theta} \colon [0,T] \times \mathbb{R}^d \times \Omega \to \mathbb{R}$ is measurable. Therefore, we obtain that for all $\varepsilon \in (0,1], n \in \mathbb{N}_0, d, j \in \mathbb{N}, \theta \in \Theta, t \in [0,T] \mathbb{R}^d \times \Omega \ni (x,\omega) \mapsto U_{n,j,\varepsilon}^{d,\theta}(t,x,\omega) \in \mathbb{R}$ is measurable. Combining this, the fact that for all $\gamma \in (0,1]$ it holds that $\mathcal{R}_a(\Gamma_\gamma) \in C(\mathbb{R}^2,\mathbb{R})$, the fact that for all $K \in \mathbb{N}, k \in \{0,1,\ldots,K\}, \gamma \in (0,1]$ it holds that $\mathcal{R}_a(\mathbf{H}_{K,k,\gamma}) \in C(\mathbb{R},\mathbb{R})$, item (i) in Corollary 4.13, and Lemma 5.2 proves that for all $\gamma, \varepsilon \in (0,1], n, j, K, d \in \mathbb{N}, \theta \in \Theta$ it holds that $[0,T] \times \mathbb{R}^d \times \Omega \ni (t,x,\omega) \mapsto (\mathcal{R}_a(\Phi_{\gamma,n,j,K}^{d,\theta,\varepsilon}(\omega)))(t,x) \in \mathbb{R}$ is measurable. Next we combine the fact that for all $\varepsilon \in (0,1], n \in \mathbb{N}_0, d, j \in \mathbb{N}, \theta \in \Theta$, $t \in [0,T]$ it holds that $\mathbb{R}^d \times \Omega \ni (x,\omega) \mapsto U_{n,j,\varepsilon}^{d,\theta}(t,x,\omega) \in \mathbb{R}$ is measurable, the fact that for all $\varepsilon \in (0,1], n, j, K, d \in \mathbb{N}, \theta \in \Theta$ it holds that $[0,T] \ni t \mapsto \mathcal{L}_{t,k-1}^{d,\theta,t_{K,k+1}}(t) \in \mathbb{R}$ is continuous, Lemma 4.4, and Lemma 5.2 to obtain that for all $\varepsilon \in (0,1], n, j, K, d \in \mathbb{N}, \theta \in \Theta$ it holds that for all $\varepsilon \in (0,1], n, j, K, d \in \mathbb{N}, \theta \in \Theta$ it holds that $[0,T] \ni t \mapsto \mathcal{L}_{t,k-1}^{0,1,0}(t,x,k+1)$.

$$[0,T] \times \mathbb{R}^d \times \Omega \ni (t,x,\omega) \mapsto \mathscr{L}^{U^{d,\theta}_{n,j,\varepsilon}(\mathfrak{t}_{K,0},x,\omega),U^{d,\theta}_{n,j,\varepsilon}(\mathfrak{t}_{K,1},x,\omega),\dots,U^{d,\theta}_{n,j,\varepsilon}(\mathfrak{t}_{K,K},x,\omega)}(t) \in \mathbb{R} \qquad (5.36)$$

is measurable (cf. Definition 4.1). Note that the triangle inequality implies that for all

 $d, K, j, n \in \mathbb{N}, \varepsilon, \gamma \in (0, 1]$ it holds that

$$\begin{split} & \left[\int_{[0,T]\times\mathbb{R}^{d}} \mathbb{E} \Big[\left| u_{d}(t,x) - \left(\mathcal{R}_{a} \left(\Phi_{\gamma,n,j,K}^{d,0,\varepsilon} \right) \right)(t,x) \right|^{\mathfrak{q}} \right] \nu_{d}(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}} \\ & \leq \left[\int_{[0,T]\times\mathbb{R}^{d}} \mathbb{E} \Big[\left| u_{d}(t,x) - \mathscr{L}_{\mathfrak{t}_{K,0},\ldots,\mathfrak{t}_{K,K}}^{U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,0},x),\ldots,U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,K},x)}(t) \right|^{\mathfrak{q}} \Big] \nu_{d}(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}} \\ & + \left[\int_{[0,T]\times\mathbb{R}^{d}} \mathbb{E} \Big[\left| \mathscr{L}_{\mathfrak{t}_{K,0},\ldots,\mathfrak{t}_{K,K}}^{U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,0},x),\ldots,U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,K},x)}(t) - \left(\mathcal{R}_{a} \left(\Phi_{\gamma,n,j,K}^{d,0,\varepsilon} \right) \right)(t,x) \right|^{\mathfrak{q}} \Big] \nu_{d}(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}}. \end{split}$$

$$(5.37)$$

Moreover, observe that the triangle inequality and Jensen's inequality show that for all $d, K, j, n \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $k \in \{1, 2, ..., K\}$, $t \in [0, T]$, $x \in \mathbb{R}^d$ it holds that

$$\mathbb{E}\Big[|u_{d}(t,x) - \mathscr{L}_{\mathfrak{t}_{K,0},\dots,\mathfrak{t}_{K,K}}^{U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,0},x),\dots,U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,K},x)}(t)|^{\mathfrak{q}} \Big] \\
\leq 4^{\mathfrak{q}-1} \Big(|u_{d}(t,x) - u_{d}(\mathfrak{t}_{K,k},x)|^{\mathfrak{q}} + |u_{d}(\mathfrak{t}_{K,k},x) - v_{d,\varepsilon}(\mathfrak{t}_{K,k},x)|^{\mathfrak{q}} \\
+ \mathbb{E}\Big[|v_{d,\varepsilon}(\mathfrak{t}_{K,k},x) - U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,k},x)|^{\mathfrak{q}} \Big] \\
+ \mathbb{E}\Big[|U_{d,\varepsilon}^{d,0}(\mathfrak{t}_{K,k},x) - \mathscr{L}_{\mathfrak{t}_{K,0},\dots,\mathfrak{t}_{K,K}}^{U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,K},x)}(t)|^{\mathfrak{q}} \Big] \Big).$$
(5.38)

Furthermore, note that Lemma 4.2, the triangle inequality, and Jensen's inequality prove that for all $d, K, j, n \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $k \in \{1, 2, ..., K\}$, $x \in \mathbb{R}^d$, $t \in [\mathfrak{t}_{K,k-1}, \mathfrak{t}_{K,k}]$ it holds that

$$\mathbb{E}\Big[|U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,k},x) - \mathscr{L}_{\mathfrak{t}_{K,0},\ldots,\mathfrak{t}_{K,K}}^{d,0}(\mathfrak{t}_{K,0,x}),\ldots,U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,K},x)}(t)|^{\mathfrak{q}} \Big] \\
\leq \mathbb{E}\Big[|U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,k},x) - U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,k-1},x)|^{\mathfrak{q}} \Big] \\
\leq 5^{\mathfrak{q}-1} \bigg(\mathbb{E}\Big[|U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,k},x) - v_{d,\varepsilon}(\mathfrak{t}_{K,k},x)|^{\mathfrak{q}} \Big] + |v_{d,\varepsilon}(\mathfrak{t}_{K,k},x) - u_{d}(\mathfrak{t}_{K,k},x)|^{\mathfrak{q}} \\
+ |u_{d}(\mathfrak{t}_{K,k},x) - u_{d}(\mathfrak{t}_{K,k-1},x)|^{\mathfrak{q}} + |u_{d}(\mathfrak{t}_{K,k-1},x) - v_{d,\varepsilon}(\mathfrak{t}_{K,k-1},x)|^{\mathfrak{q}} \\
+ \mathbb{E}\Big[|v_{d,\varepsilon}(\mathfrak{t}_{K,k-1},x) - U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,k-1},x)|^{\mathfrak{q}} \Big] \Big).$$
(5.39)

This and (5.38) demonstrate that for all $d, K, j, n \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $k \in \{1, 2, \dots, K\}$, $x \in \mathbb{R}^d$,

 $t \in [\mathfrak{t}_{K,k-1},\mathfrak{t}_{K,k}]$ it holds that

$$\mathbb{E}\Big[\left|u_{d}(t,x) - \mathscr{D}_{\mathsf{t}_{K,0},\dots,\mathsf{t}_{K,K}}^{U_{n,j,\varepsilon}^{d,0}(\mathsf{t}_{K,0},x),\dots,\mathsf{U}_{n,j,\varepsilon}^{d,0}(\mathsf{t}_{K,K},x)}(t)\right|^{\mathfrak{q}}\Big] \\
\leq 20^{\mathfrak{q}-1} \Big(\left|u_{d}(t,x) - u_{d}(\mathsf{t}_{K,k},x)\right|^{\mathfrak{q}} + 2\left|u_{d}(\mathsf{t}_{K,k},x) - v_{d,\varepsilon}(\mathsf{t}_{K,k},x)\right|^{\mathfrak{q}} \\
+ \left|u_{d}(\mathsf{t}_{K,k},x) - u_{d}(\mathsf{t}_{K,k-1},x)\right|^{\mathfrak{q}} + \left|u_{d}(\mathsf{t}_{K,k-1},x) - v_{d,\varepsilon}(\mathsf{t}_{K,k-1},x)\right|^{\mathfrak{q}} \\
+ 2\mathbb{E}\Big[\left|v_{d,\varepsilon}(\mathsf{t}_{K,k},x) - U_{n,j,\varepsilon}^{d,0}(\mathsf{t}_{K,k},x)\right|^{\mathfrak{q}}\Big] + \mathbb{E}\Big[\left|v_{d,\varepsilon}(\mathsf{t}_{K,k-1},x) - U_{n,j,\varepsilon}^{d,0}(\mathsf{t}_{K,k-1},x)\right|^{\mathfrak{q}}\Big]\Big) \\
\leq 20^{\mathfrak{q}-1} \Big(2\Big(\sup_{s\in[\mathsf{t}_{K,k-1},\mathsf{t}_{K,k}]}\left|u_{d}(s,x) - u_{d}(\mathsf{t}_{K,k},x)\right|^{\mathfrak{q}}\Big) + 3\Big(\sup_{s\in[0,T]}\left|u_{d}(s,x) - v_{d,\varepsilon}(s,x)\right|^{\mathfrak{q}}\Big) \\
+ 3\Big(\sup_{s\in[0,T]}\mathbb{E}\Big[\left|v_{d,\varepsilon}(s,x) - U_{n,j,\varepsilon}^{d,0}(s,x)\right|^{\mathfrak{q}}\Big]\Big)\Big).$$
(5.40)

In addition, observe that (5.33), (5.34), the fact that for all $w, z \in \mathbb{R}$ it holds that $|f_0(z) - f_0(w)| \leq L|z-w|$, the fact that for all $d \in \mathbb{N}$, $x \in \mathbb{R}^d$ it holds that $\|\nabla f_d(x)\| \leq \kappa d^p (1+\|x\|)^p$, Corollary 2.7 (applied for every $d \in \mathbb{N}$ with $u \curvearrowleft u_d$, $T \curvearrowleft T$, $d \backsim d$, $W \curvearrowleft W^{d,0}$, $g \backsim f_d$, $f \backsim ([0,T] \times \mathbb{R}^d \times \mathbb{R} \ni (t, x, z) \mapsto f_0(z) \in \mathbb{R})$, $L \backsim L$, $\mathfrak{L} \curvearrowleft \kappa d^p$, $p \backsim p$ in the notation of Corollary 2.7), (5.24), and Jensen's inequality establish that for all $d, K \in \mathbb{N}$, $k \in \{1, 2, \ldots, K\}, x \in \mathbb{R}^d$ it holds that

$$\begin{split} \sup_{s \in [\mathfrak{t}_{K,k-1},\mathfrak{t}_{K,k}]} &|u_d(s,x) - u_d(\mathfrak{t}_{K,k},x)|^{\mathfrak{q}} \leq \left(\mathfrak{t}_{K,k} - \mathfrak{t}_{K,k-1}\right)^{\frac{q}{2}} d^{\frac{q}{2}} \left(e^{2LT} (T+1)^2 (L+1) (\kappa d^p + 1)\right)^{\mathfrak{q}} \\ &\cdot 8^{(p+2)\mathfrak{q}} \left(1 + \|x\|^p + \sup_{s \in [0,T]} \mathbb{E} \left[\|W_s^{d,0}\|^{2p}\right]\right)^{\mathfrak{q}} \\ &\leq \left[\frac{T}{K}\right]^{\frac{q}{2}} (\mathfrak{a}_d)^{\mathfrak{q}} 3^{\mathfrak{q}-1} \left(1 + \|x\|^{p\mathfrak{q}} + \sup_{s \in [0,T]} \mathbb{E} \left[\|W_s^{d,0}\|^{2p\mathfrak{q}}\right]\right) \\ &\leq \left[\frac{T}{K}\right]^{\frac{q}{2}} (\mathfrak{a}_d)^{\mathfrak{q}} 3^{\mathfrak{q}} \left(1 + \|x\|^{p^2q\mathfrak{q}} + \sup_{s \in [0,T]} \mathbb{E} \left[\|W_s^{d,0}\|^{2p^2q\mathfrak{q}}\right]\right). \end{split}$$

$$(5.41)$$

In addition, note that (5.16) and Jensen's inequality ensure that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $z \in \mathbb{R}$, $x \in \mathbb{R}^d$ it holds that

$$|f_0(z) - (\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(z)| \le \varepsilon \kappa d^p (1+|z|)^p \le \varepsilon 2^{p-1} \kappa d^p \big((1+||x||)^{p^2} + |z|^p \big).$$
(5.42)

In addition, observe that (5.16) implies that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $z \in \mathbb{R}$, $x \in \mathbb{R}^d$ it holds that

$$|f_d(x) - (\mathcal{R}_a(\mathbf{F}_{d,\varepsilon}))(x)| \le \varepsilon \kappa d^p (1 + ||x||)^p \le \varepsilon 2^{p-1} \kappa d^p ((1 + ||x||)^{p^2} + |z|^p).$$
(5.43)

This, (5.42), (5.17), (5.33), (5.34), (5.35), the assumption that for all $d \in \mathbb{N}_0$, $x \in \mathbb{R}^{\max\{d,1\}}$, $\varepsilon \in (0, 1]$ it holds that $|(\mathcal{R}_a(\mathbf{F}_{d,\varepsilon}))(x)| \leq \kappa (\max\{d,1\})^p (1+||x||)^p$, Corollary 2.3 (applied for every $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ with $u_1 \curvearrowleft u_d$, $u_2 \curvearrowleft v_{d,\varepsilon}$, $T \curvearrowleft T$, $d \backsim d$, $W \curvearrowleft W^{d,0}$, $L \backsim L$,

 $\mathfrak{L} \curvearrowright \kappa d^p, \ B \curvearrowleft \varepsilon 2^{p-1} \kappa d^p, \ p \backsim p, \ q \backsim p, \ f_1 \curvearrowleft ([0,T] \times \mathbb{R}^d \times \mathbb{R} \ni (t,x,z) \mapsto f_0(z) \in \mathbb{R}), \\ f_2 \curvearrowleft ([0,T] \times \mathbb{R}^d \times \mathbb{R} \ni (t,x,z) \mapsto (\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(z) \in \mathbb{R}), \ g_1 \curvearrowleft f_d, \ g_2 \curvearrowleft \mathcal{R}_a(\mathbf{F}_{d,\varepsilon}) \text{ in the notation of Corollary 2.3}, \text{ and } (5.25) \text{ show that for all } d \in \mathbb{N}, \ \varepsilon \in (0,1], \ x \in \mathbb{R}^d \text{ it holds that}$

$$\sup_{s \in [0,T]} |u_d(s,x) - v_{d,\varepsilon}(s,x)| \le \varepsilon \mathfrak{b}_d \left(1 + ||x||^{p^2} + \sup_{s \in [0,T]} \mathbb{E} \left[||W_s^{d,0}||^{p^2} \right] \right).$$
(5.44)

This and Jensen's inequality prove that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $x \in \mathbb{R}^d$ it holds that

$$\sup_{s\in[0,T]} |u_d(s,x) - v_{d,\varepsilon}(s,x)|^{\mathfrak{q}} \le \varepsilon^{\mathfrak{q}}(\mathfrak{b}_d)^{\mathfrak{q}} 3^{\mathfrak{q}} \left(1 + ||x||^{p^2q\mathfrak{q}} + \sup_{s\in[0,T]} \mathbb{E}\left[||W_s^{d,0}||^{2p^2q\mathfrak{q}}\right]\right).$$
(5.45)

Next we combine (5.35), the fact that for all $\varepsilon \in (0, 1]$, $w, z \in \mathbb{R}$ it holds that $|(\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(w) - (\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(z)| \leq L|w-z|$, the fact that for all $\varepsilon \in (0, 1]$, $d \in \mathbb{N}_0$, $x \in \mathbb{R}^{\max\{d,1\}}$ it holds that $|(\mathcal{R}_a(\mathbf{F}_{d,\varepsilon}))(x)| \leq \kappa (\max\{d,1\})^p (1+||x||)^p$, Hutzenthaler et al. [48, Corollary 3.15] (applied for every $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $j \in \mathbb{N}$ with $\mathfrak{p} \curvearrowleft \mathfrak{q}$, $d \curvearrowleft d$, $m \curvearrowleft m_j$, $T \curvearrowleft T$, $L \backsim L$, $\mathfrak{L} \hookrightarrow 2^{p-1}\kappa d^p$, $p \backsim p$, $u \backsim v_{d,\varepsilon}$, $f \backsim ([0,T] \times \mathbb{R}^d \times \mathbb{R} \ni (t,x,z) \mapsto (\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(z) \in \mathbb{R})$, $g \backsim \mathcal{R}_a(\mathbf{F}_{d,\varepsilon})$, $\Theta \curvearrowleft \Theta$, $(W^\theta)_{\theta \in \Theta} \curvearrowleft (W^{d,\theta})_{\theta \in \Theta}$, $(U_n^\theta)_{(n,\theta) \in \mathbb{Z} \times \Theta} \curvearrowleft (U_{n,j,\varepsilon}^{d,\theta})_{(n,\theta) \in \mathbb{Z} \times \Theta}$, $(\mathfrak{u}^\theta)_{\theta \in \Theta}$, $(\mathfrak{U}^\theta)_{\theta \in \Theta} \curvearrowleft (\mathfrak{U}^\theta)_{\theta \in \Theta}$ in the notation of Hutzenthaler et al. [48, Corollary 3.15]), and (5.26) to obtain that for all $d, j, n \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $x \in \mathbb{R}^d$ it holds that

$$\sup_{s\in[0,T]} \mathbb{E}\left[|v_{d,\varepsilon}(s,x) - U_{n,j,\varepsilon}^{d,0}(s,x)|^{\mathfrak{q}}\right]$$

$$\leq (\mathfrak{c}_d)^{\mathfrak{q}}\left[(1+2LT)^n(m_j)^{-\frac{n}{2}}\exp\left(\frac{(m_j)^{\frac{\mathfrak{q}}{2}}}{\mathfrak{q}}\right)\right]^{\mathfrak{q}}\left(\sup_{s\in[0,T]} \mathbb{E}\left[(1+\|x+W_s^{d,0}\|^p)^{\mathfrak{q}}\right]\right).$$
(5.46)

In the next step we note that Jensen's inequality and the triangle inequality demonstrate that for all $d \in \mathbb{N}$, $x \in \mathbb{R}^d$, $s \in [0, T]$ it holds that

$$\mathbb{E}\left[\left(1 + \|x + W_{s}^{d,0}\|^{p}\right)^{\mathfrak{q}}\right] \leq 2^{\mathfrak{q}-1} \left(1 + 2^{p\mathfrak{q}-1} \left(\|x\|^{p\mathfrak{q}} + \mathbb{E}\left[\|W_{s}^{d,0}\|^{2p^{2}q\mathfrak{q}}\right]\right)\right) \\ \leq 2^{\mathfrak{q}-1} \left(1 + 2^{p\mathfrak{q}-1} \left(\|x\|^{p^{2}q\mathfrak{q}} + \mathbb{E}\left[\|W_{s}^{d,0}\|^{2p^{2}q\mathfrak{q}}\right]\right) + 2^{p\mathfrak{q}}\right) \qquad (5.47) \\ \leq 2^{(p+1)\mathfrak{q}} \left(1 + \|x\|^{p^{2}q\mathfrak{q}} + \mathbb{E}\left[\|W_{s}^{d,0}\|^{2p^{2}q\mathfrak{q}}\right]\right).$$

In the next step we observe that (5.27) establishes that for all $j, n \in \mathbb{N}$ it holds that

$$(1+2LT)^{n}(m_{j})^{-\frac{n}{2}}\exp\left(\frac{(m_{j})^{\frac{q}{2}}}{\mathfrak{q}}\right) \leq \left[(1+2LT)(m_{j})^{-\frac{1}{2}}\exp\left(\frac{(m_{j})^{\frac{qq}{2}}}{n}\right)\right]^{n} = \mathfrak{m}_{j,n}.$$
 (5.48)

This, (5.46), and (5.47) ensure that for all $d, j, n \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $x \in \mathbb{R}^d$ it holds that

$$\sup_{s \in [0,T]} \mathbb{E} \Big[|v_{d,\varepsilon}(s,x) - U_{n,j,\varepsilon}^{d,0}(s,x)|^{\mathfrak{q}} \Big] \le (\mathfrak{c}_d)^{\mathfrak{q}} (\mathfrak{m}_{j,n})^{\mathfrak{q}} 2^{(p+1)\mathfrak{q}} \bigg(1 + \|x\|^{p^2q\mathfrak{q}} + \sup_{s \in [0,T]} \mathbb{E} \Big[\|W_s^{d,0}\|^{2p^2q\mathfrak{q}} \Big] \bigg).$$
(5.49)

Combining this, (5.40), (5.41), and (5.45) implies that for all $d, K, j, n \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $x \in \mathbb{R}^d$, $t \in [0, T]$ it holds that

$$\mathbb{E}\Big[\Big|u_{d}(t,x) - \mathscr{D}_{\mathfrak{t}_{K,0},\dots,\mathfrak{t}_{K,K}}^{U^{d,0}_{n,j,\varepsilon}(\mathfrak{t}_{K,0},x),\dots,U^{d,0}_{n,j,\varepsilon}(\mathfrak{t}_{K,K},x)}(t)\Big|^{\mathfrak{q}}\Big] \\
\leq 20^{\mathfrak{q}-1}\left(2\Big[\frac{T}{K}\Big]^{\frac{\mathfrak{q}}{2}}(\mathfrak{a}_{d})^{\mathfrak{q}}3^{\mathfrak{q}} + 3\varepsilon^{\mathfrak{q}}(\mathfrak{b}_{d})^{\mathfrak{q}}3^{\mathfrak{q}} + 3(\mathfrak{c}_{d})^{\mathfrak{q}}(\mathfrak{m}_{j,n})^{\mathfrak{q}}2^{(p+1)\mathfrak{q}}\right) \\
\cdot \left(1 + \|x\|^{p^{2}q\mathfrak{q}} + \sup_{s\in[0,T]}\mathbb{E}\big[\|W^{d,0}_{s}\|^{2p^{2}q\mathfrak{q}}\big]\big).$$
(5.50)

The triangle inequality and (5.28) hence show that for all $d, K, j, n \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$\begin{bmatrix}
\int_{[0,T]\times\mathbb{R}^d} \mathbb{E}\Big[\left| u_d(t,x) - \mathscr{L}^{U^{d,0}_{n,j,\varepsilon}(\mathfrak{t}_{K,0},x),\dots,U^{d,0}_{n,j,\varepsilon}(\mathfrak{t}_{K,K},x)}(t) \right|^{\mathfrak{q}} \right] \nu_d(\,\mathrm{d}t,\,\mathrm{d}x) \end{bmatrix}^{1/\mathfrak{q}} \leq 20\Big(6(T+1)\mathfrak{a}_d K^{-\frac{1}{2}} + 9\varepsilon\mathfrak{b}_d + 2^p 6\mathfrak{c}_d\mathfrak{m}_{j,n}\Big) c_d.$$
(5.51)

Combining this, (5.27), (5.29), (5.30), and item (i) in Lemma 5.1 proves that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$\begin{bmatrix}
\int_{[0,T]\times\mathbb{R}^d} \mathbb{E}\Big[|u_d(t,x) - \mathscr{L}^{U^{d,0}_{N_{\varepsilon},N_{\varepsilon},\varepsilon}(\mathfrak{t}_{K_{\varepsilon},0,x}),\dots,U^{d,0}_{N_{\varepsilon},N_{\varepsilon},\varepsilon}(\mathfrak{t}_{K_{\varepsilon},K_{\varepsilon},x})}(t)|^{\mathfrak{q}}]\nu_d(\,\mathrm{d}t,\,\mathrm{d}x) \Big]^{1/\mathfrak{q}} \\
\leq 20\big(6(T+1)\mathfrak{a}_d(K_{\varepsilon})^{-\frac{1}{2}} + 9\varepsilon\mathfrak{b}_d + 2^p6\mathfrak{c}_d\mathfrak{m}_{N_{\varepsilon},N_{\varepsilon}}\big)c_d \\
\leq 180(T+1)2^p\big(\mathfrak{a}_d + \mathfrak{b}_d + \mathfrak{c}_d\big)c_d\varepsilon.
\end{cases}$$
(5.52)

Note that (5.12), (5.14), item (ii) in Corollary 4.13, the triangle inequality, and (5.10) demonstrate that for all $d, K, j, n \in \mathbb{N}$, $\varepsilon, \gamma \in (0, 1]$ it holds that

$$\begin{bmatrix}
\int_{[0,T]\times\mathbb{R}^{d}} \mathbb{E}\left[\left|\mathscr{L}_{\mathfrak{t}_{K,0},\ldots,\mathfrak{t}_{K,K}}^{U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,0},x),\ldots,U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,K},x)}(t) - \left(\mathcal{R}_{a}\left(\Phi_{\gamma,n,j,K}^{d,0,\varepsilon}\right)\right)(t,x)\right|^{\mathfrak{q}}\right]\nu_{d}(dt, dx)\right]^{1/\mathfrak{q}} \\
\leq 2\gamma\left(1 + (T+1)^{q}\right)^{q}\left[(K+1)\kappa d^{rp^{2}q} + \sum_{k=0}^{K}\left[\int_{[0,T]\times\mathbb{R}^{d}} \mathbb{E}\left[\left|U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,k},x)\right|^{q}\right]\nu_{d}(dt, dx)\right]^{1/\mathfrak{q}}\right].$$
(5.53)

Next, observe that Jensen's inequality and the triangle inequality establish that for all $d, K, j, n \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $k \in \{0, 1, \dots, K\}$ it holds that

$$\begin{bmatrix}
\int_{[0,T]\times\mathbb{R}^{d}} \mathbb{E}\left[\left|U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,k},x)\right|^{q\mathfrak{q}}\right]\nu_{d}(\,\mathrm{d}t,\,\mathrm{d}x)\right]^{1/\mathfrak{q}} \\
\leq 3^{q-1} \left(\left[\int_{[0,T]\times\mathbb{R}^{d}} \mathbb{E}\left[\left|U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,k},x)-v_{d,\varepsilon}(\mathfrak{t}_{K,k},x)\right|^{q\mathfrak{q}}\right]\nu_{d}(\,\mathrm{d}t,\,\mathrm{d}x)\right]^{1/\mathfrak{q}} \\
+ \left[\int_{[0,T]\times\mathbb{R}^{d}}\left|v_{d,\varepsilon}(\mathfrak{t}_{K,k},x)-v_{d,\varepsilon}(T,x)\right|^{q\mathfrak{q}}\nu_{d}(\,\mathrm{d}t,\,\mathrm{d}x)\right]^{1/\mathfrak{q}} \\
+ \left[\int_{[0,T]\times\mathbb{R}^{d}}\left|v_{d,\varepsilon}(T,x)\right|^{q\mathfrak{q}}\nu_{d}(\,\mathrm{d}t,\,\mathrm{d}x)\right]^{1/\mathfrak{q}}\right).$$
(5.54)

Next, note that (5.35), the fact that for all $\varepsilon \in (0, 1]$, $w, z \in \mathbb{R}$ it holds that $|(\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(w) - (\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(z)| \leq L|w-z|$, the fact that for all $\varepsilon \in (0, 1]$, $d \in \mathbb{N}_0$, $x \in \mathbb{R}^{\max\{d,1\}}$ it holds that $|(\mathcal{R}_a(\mathbf{F}_{d,\varepsilon}))(x)| \leq \kappa(\max\{d,1\})^p (1+||x||)^p$, Hutzenthaler et al. [48, Corollary 3.15] (applied for every $d \in \mathbb{N}$, $\varepsilon \in (0,1]$, $j \in \mathbb{N}$ with $\mathfrak{p} \curvearrowleft q\mathfrak{q}$, $d \curvearrowleft d$, $m \curvearrowleft m_j$, $T \curvearrowleft T$, $L \backsim L$, $\mathfrak{L} \curvearrowleft 2^{p-1}\kappa d^p$, $p \backsim p$, $u \backsim v_{d,\varepsilon}$, $f \backsim ([0,T] \times \mathbb{R}^d \times \mathbb{R} \ni (t,x,z) \mapsto (\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(z) \in \mathbb{R})$, $g \curvearrowleft \mathcal{R}_a(\mathbf{F}_{d,\varepsilon})$, $\Theta \curvearrowleft \Theta$, $(W^\theta)_{\theta \in \Theta} \curvearrowleft (W^{d,\theta})_{\theta \in \Theta}$, $(U^\theta_n)_{(n,\theta) \in \mathbb{Z} \times \Theta} \curvearrowleft (U^{d,\theta}_{n,j,\varepsilon})_{(n,\theta) \in \mathbb{Z} \times \Theta}$, $(\mathfrak{u}^\theta)_{\theta \in \Theta} \curvearrowleft (\mathfrak{U}^\theta)_{\theta \in \Theta}$ in the notation of Hutzenthaler et al. [48, Corollary 3.15]), (5.26), (5.27), Jensen's inequality, and the triangle inequality ensure that for all $d, K, j, n \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $k \in \{0, 1, \ldots, K\}$, $x \in \mathbb{R}^d$ it holds that

$$\mathbb{E}\Big[|U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,k},x) - v_{d,\varepsilon}(\mathfrak{t}_{K,k},x)|^{q\mathfrak{q}} \Big] \\
\leq (\mathfrak{c}_{d})^{q\mathfrak{q}} \Big[(1 + 2LT)^{n} (m_{j})^{-\frac{n}{2}} \exp\left(\frac{(m_{j})^{\frac{q\mathfrak{q}}{2}}}{q\mathfrak{q}}\right) \Big]^{q\mathfrak{q}} (\sup_{s \in [0,T]} \mathbb{E}\big[(1 + ||x + W_{s}^{d,0}||^{p})^{q\mathfrak{q}} \big]) \quad (5.55) \\
\leq (\mathfrak{c}_{d})^{q\mathfrak{q}} (\mathfrak{m}_{j,n})^{q\mathfrak{q}} 2^{(p+1)q\mathfrak{q}} (1 + ||x||^{p^{2}q\mathfrak{q}} + \sup_{s \in [0,T]} \mathbb{E}\big[||W_{s}^{d,0}||^{2p^{2}q\mathfrak{q}} \big]).$$

In the next step we observe that Jensen's inequality and the triangle inequality imply that for all $d, K \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $k \in \{0, 1, \dots, K\}$ it holds that

$$\begin{bmatrix}
\int_{[0,T]\times\mathbb{R}^{d}} |v_{d,\varepsilon}(\mathfrak{t}_{K,k},x) - v_{d,\varepsilon}(T,x)|^{q\mathfrak{q}} \nu_{d}(\,\mathrm{d}t,\,\mathrm{d}x) \end{bmatrix}^{1/\mathfrak{q}} \\
\leq 3^{q-1} \left(\left[\int_{[0,T]\times\mathbb{R}^{d}} |v_{d,\varepsilon}(\mathfrak{t}_{K,k},x) - u_{d}(\mathfrak{t}_{K,k},x)|^{q\mathfrak{q}} \nu_{d}(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}} \\
+ \left[\int_{[0,T]\times\mathbb{R}^{d}} |u_{d}(\mathfrak{t}_{K,k},x) - u_{d}(T,x)|^{q\mathfrak{q}} \nu_{d}(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}} \\
+ \left[\int_{[0,T]\times\mathbb{R}^{d}} |u_{d}(T,x) - v_{d,\varepsilon}(T,x)|^{q\mathfrak{q}} \nu_{d}(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}} \right).$$
(5.56)

In the next step we note that (5.28), (5.44), and Jensen's inequality show that for all $d, K \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $k \in \{0, 1, \dots, K\}$, $\mathfrak{t} \in \{\mathfrak{t}_{K,k}, T\}$ it holds that

$$\begin{split} & \left[\int_{[0,T]\times\mathbb{R}^{d}} |u_{d}(\mathfrak{t},x) - v_{d,\varepsilon}(\mathfrak{t},x)|^{q\mathfrak{q}} \nu_{d}(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}} \\ & \leq \varepsilon^{q}(\mathfrak{b}_{d})^{q} \left[\int_{[0,T]\times\mathbb{R}^{d}} (1 + \|x\|^{p^{2}} + \sup_{s\in[0,T]} \mathbb{E}\left[\|W_{s}^{d,0}\|^{p^{2}} \right] \right)^{q\mathfrak{q}} \nu_{d}(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}} \\ & \leq 3^{q} \varepsilon^{q}(\mathfrak{b}_{d})^{q} \left[\int_{[0,T]\times\mathbb{R}^{d}} (1 + \|x\|^{p^{2}q\mathfrak{q}} + \sup_{s\in[0,T]} \mathbb{E}\left[\|W_{s}^{d,0}\|^{2p^{2}q\mathfrak{q}} \right] \right) \nu_{d}(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}} \\ & = 3^{q} \varepsilon^{q}(\mathfrak{b}_{d})^{q} c_{d}. \end{split}$$
(5.57)

Furthermore, observe that (5.33), (5.34), the fact that for all $w, z \in \mathbb{R}$ it holds that $|f_0(z) - f_0(w)| \leq L|z-w|$, the fact that for all $d \in \mathbb{N}$, $x \in \mathbb{R}^d$ it holds that $\|\nabla f_d(x)\| \leq \kappa d^p (1+\|x\|)^p$, Corollary 2.7 (applied for every $d \in \mathbb{N}$ with $u \curvearrowleft u_d$, $T \curvearrowleft T$, $d \curvearrowleft d$, $W \curvearrowleft W^{d,0}$, $g \curvearrowleft f_d$, $f \curvearrowleft ([0,T] \times \mathbb{R}^d \times \mathbb{R} \ni (t,x,z) \mapsto f_0(z) \in \mathbb{R}), L \curvearrowleft L, \mathfrak{L} \curvearrowleft \kappa d^p, p \curvearrowleft p$ in the notation of Corollary 2.7), (5.24), Jensen's inequality, and (5.28) prove that for all $d, K \in \mathbb{N}, k \in \{0, 1, \ldots, K\}$ it holds that

$$\begin{split} & \left[\int_{[0,T]\times\mathbb{R}^d} |u_d(\mathfrak{t}_{K,k},x) - u_d(T,x)|^{q\mathfrak{q}} \,\nu_d(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}} \\ & \leq \left(T - \mathfrak{t}_{K,k}\right)^{\frac{q}{2}} d^{\frac{q}{2}} \left(e^{2LT} (T+1)^2 (L+1) (\kappa d^p + 1) 8^{p+2} \right)^q \\ & \cdot \left[\int_{[0,T]\times\mathbb{R}^d} \left(1 + \|x\|^p + \sup_{s\in[0,T]} \mathbb{E} \left[\|W_s^{d,0}\|^{2p} \right] \right)^{q\mathfrak{q}} \,\nu_d(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}} \\ & \leq T^{\frac{q}{2}} (\mathfrak{a}_d)^q 3^q \left[\int_{[0,T]\times\mathbb{R}^d} \left(1 + \|x\|^{p^2q\mathfrak{q}} + \sup_{s\in[0,T]} \mathbb{E} \left[\|W_s^{d,0}\|^{2p^2q\mathfrak{q}} \right] \right) \nu_d(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}} \\ & = T^{\frac{q}{2}} (\mathfrak{a}_d)^q 3^q c_d. \end{split}$$
(5.58)

Combining this, (5.56), and (5.57) demonstrates that for all $d, K \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $k \in \{0, 1, \ldots, K\}$ it holds that

$$\left[\int_{[0,T]\times\mathbb{R}^d} |v_{d,\varepsilon}(\mathfrak{t}_{K,k},x) - v_{d,\varepsilon}(T,x)|^{q\mathfrak{q}} \nu_d(\,\mathrm{d}t,\,\mathrm{d}x)\right]^{1/\mathfrak{q}} \leq 3^{q-1} \left(3^q 2\varepsilon^q(\mathfrak{b}_d)^q c_d + T^{\frac{q}{2}}(\mathfrak{a}_d)^q 3^q c_d\right) \leq 3^{2q} (T+1)^{\frac{q}{2}} \left((\mathfrak{a}_d)^q + (\mathfrak{b}_d)^q\right) c_d.$$
(5.59)

In addition, note that (5.35), the fact that for all $d \in \mathbb{N}$, $x \in \mathbb{R}^d$, $\varepsilon \in (0, 1]$ it holds that $|(\mathcal{R}_a(\mathbf{F}_{d,\varepsilon}))(x)| \leq \kappa d^p (1 + ||x||)^p$, Jensen's inequality, and (5.28) establish that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$\begin{split} \left[\int_{[0,T]\times\mathbb{R}^d} |v_{d,\varepsilon}(T,x)|^{q\mathfrak{q}} \,\nu_d(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}} &= \left[\int_{[0,T]\times\mathbb{R}^d} \left| \left(\mathcal{R}_a(\mathbf{F}_{d,\varepsilon}) \right)(x) \right|^{q\mathfrak{q}} \,\nu_d(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}} \\ &\leq \kappa^q d^{pq} \left[\int_{[0,T]\times\mathbb{R}^d} (1+\|x\|)^{pq\mathfrak{q}} \,\nu_d(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}} \\ &\leq \kappa^q d^{pq} 2^{\frac{p^2q\mathfrak{q}-1}{\mathfrak{q}}} \left[\int_{[0,T]\times\mathbb{R}^d} (1+\|x\|)^{p^2q\mathfrak{q}} \,\nu_d(\,\mathrm{d}t,\,\mathrm{d}x) \right]^{1/\mathfrak{q}} \\ &\leq \kappa^q d^{pq} 2^{p^2q} c_d. \end{split}$$
(5.60)

Combining this, (5.28), (5.54), (5.55), and (5.59) ensures that for all $d, K, j, n \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $k \in \{0, 1, \ldots, K\}$ it holds that

$$\begin{bmatrix}
\int_{[0,T]\times\mathbb{R}^{d}} \mathbb{E}\Big[|U_{n,j,\varepsilon}^{d,0}(\mathfrak{t}_{K,k},x)|^{q\mathfrak{q}} \Big] \nu_{d}(dt, dx) \end{bmatrix}^{1/\mathfrak{q}} \\
\leq 3^{q-1} c_{d} \big((\mathfrak{c}_{d})^{q}(\mathfrak{m}_{j,n})^{q} 2^{(p+1)q} + 3^{2q} (T+1)^{\frac{q}{2}} ((\mathfrak{a}_{d})^{q} + (\mathfrak{b}_{d})^{q}) + \kappa^{q} d^{pq} 2^{p^{2}q} \big) \\
\leq 3^{3p^{2}q} c_{d} \big((\mathfrak{c}_{d})^{q}(\mathfrak{m}_{j,n})^{q} + (T+1)^{\frac{q}{2}} ((\mathfrak{a}_{d})^{q} + (\mathfrak{b}_{d})^{q}) + \kappa^{q} d^{pq} \big).$$
(5.61)

This and (5.53) imply that for all $d, K, j, n \in \mathbb{N}, \varepsilon, \gamma \in (0, 1]$ it holds that

$$\begin{bmatrix}
\int_{[0,T]\times\mathbb{R}^{d}} \mathbb{E}\Big[|\mathscr{L}_{\mathfrak{t}_{K,0},\dots,\mathfrak{t}_{K,K}}^{U^{d,0}_{n,j,\varepsilon}(\mathfrak{t}_{K,0},x),\dots,U^{d,0}_{n,j,\varepsilon}(\mathfrak{t}_{K,K},x)}(t) - \big(\mathcal{R}_{a}\big(\Phi^{d,0,\varepsilon}_{\gamma,n,j,K}\big)\big)(t,x)\big|^{\mathfrak{q}}\Big]\nu_{d}(\,\mathrm{d}t,\,\mathrm{d}x) \end{bmatrix}^{1/\mathfrak{q}} \\
\leq 2\gamma\big(1 + (T+1)^{q}\big)^{q}(K+1)\kappa d^{rp^{2}q} \\
\cdot \big[1 + 3^{3p^{2}q}c_{d}\big((\mathfrak{c}_{d})^{q}(\mathfrak{m}_{j,n})^{q} + (T+1)^{\frac{q}{2}}((\mathfrak{a}_{d})^{q} + (\mathfrak{b}_{d})^{q}) + \kappa^{q}d^{pq}\big)\Big].$$
(5.62)

Combining this, (5.27), (5.29), item (i) in Lemma 5.1, (5.30), and (5.32) shows that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$\begin{split} & \left[\int_{[0,T]\times\mathbb{R}^d} \mathbb{E} \Big[\left| \mathscr{L}_{\mathfrak{t}_{\kappa_{\varepsilon},0},n}^{U_{N_{\varepsilon},N_{\varepsilon},\varepsilon}^{d}}(\mathfrak{t}_{\kappa_{\varepsilon},0,x}), \dots, \mathcal{U}_{N_{\varepsilon},N_{\varepsilon},\varepsilon}^{d,0}(\mathfrak{t}_{\kappa_{\varepsilon},K_{\varepsilon},x})}(t) - \left(\mathcal{R}_{a} \left(\Phi_{\gamma_{d,\varepsilon},N_{\varepsilon},N_{\varepsilon},K_{\varepsilon}}^{d,0,\varepsilon} \right) \right)(t,x) \right|^{\mathfrak{q}} \Big] \nu_{d}(\,\mathrm{d}t,\,\mathrm{d}x) \Big]^{1/\mathfrak{q}} \\ & \leq 2\gamma_{d,\varepsilon} \Big(K_{\varepsilon} + 1 \Big) \kappa d^{rp^{2}q} \Big(1 + (T+1)^{q} \Big)^{q} \\ & \cdot \Big(1 + 3^{3p^{2}q} c_{d} \Big((\mathfrak{c}_{d})^{q} (\mathfrak{m}_{N_{\varepsilon},N_{\varepsilon}})^{q} + (T+1)^{\frac{q}{2}} ((\mathfrak{a}_{d})^{q} + (\mathfrak{b}_{d})^{q}) + \kappa^{q} d^{pq} \Big) \Big) \\ & \leq 2\gamma_{d,\varepsilon} \Big(K_{\varepsilon} + 1 \Big) \kappa d^{rp^{2}q} \Big(1 + (T+1)^{q} \Big)^{q} \Big(1 + 3^{3p^{2}q} c_{d} \Big((\mathfrak{c}_{d})^{q} + (T+1)^{\frac{q}{2}} ((\mathfrak{a}_{d})^{q} + (\mathfrak{b}_{d})^{q}) + \kappa^{q} d^{pq} \Big) \Big) \\ &= \varepsilon. \end{split}$$

$$\tag{5.63}$$

Combining this, (5.31), (5.37), and (5.52) proves that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$\begin{bmatrix}
\int_{[0,T]\times\mathbb{R}^d} \mathbb{E}\Big[|u_d(t,x) - \big(\mathcal{R}_a\big(\Phi^{d,0,\delta_{d,\varepsilon}}_{\gamma_d,\delta_{d,\varepsilon}}, N_{\delta_{d,\varepsilon}}, N_{\delta_{d,\varepsilon}}, K_{\delta_{d,\varepsilon}}\big)\big)(t,x)|^{\mathfrak{q}} \Big]\nu_d(\,\mathrm{d}t,\,\mathrm{d}x) \end{bmatrix}^{1/\mathfrak{q}} \leq \big(180(T+1)2^p\big(\mathfrak{a}_d + \mathfrak{b}_d + \mathfrak{c}_d\big)c_d + 1\big)\delta_{d,\varepsilon} = \varepsilon.$$
(5.64)

Combining this and Fubini's theorem demonstrates that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$\mathbb{E}\bigg[\int_{[0,T]\times\mathbb{R}^d} |u_d(t,x) - \big(\mathcal{R}_a\big(\Phi^{d,0,\delta_{d,\varepsilon}}_{\gamma_{d,\delta_{d,\varepsilon}},N_{\delta_{d,\varepsilon}},N_{\delta_{d,\varepsilon}},K_{\delta_{d,\varepsilon}}}\big)\big)(t,x)\big|^{\mathfrak{q}}\,\nu_d(\,\mathrm{d} t,\,\mathrm{d} x)\bigg] \le \varepsilon^{\mathfrak{q}}.\tag{5.65}$$

This establishes that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ there exists $\omega_{d,\varepsilon} \in \Omega$ such that

$$\int_{[0,T]\times\mathbb{R}^d} \left| u_d(t,x) - \left(\mathcal{R}_a \left(\Phi^{d,0,\delta_{d,\varepsilon}}_{\gamma_{d,\delta_{d,\varepsilon}},N_{\delta_{d,\varepsilon}},K_{\delta_{d,\varepsilon}}}(\omega_{d,\varepsilon}) \right) \right)(t,x) \right|^{\mathfrak{q}} \nu_d(\,\mathrm{d}t,\,\mathrm{d}x) \le \varepsilon^{\mathfrak{q}}.$$
(5.66)

Observe that item (iii) in Corollary 4.13 ensures that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$\mathcal{P}\left(\Phi_{\gamma_{d,\delta_{d,\varepsilon}},N_{\delta_{d,\varepsilon}},N_{\delta_{d,\varepsilon}},K_{\delta_{d,\varepsilon}}}^{d,0,\delta_{d,\varepsilon}}(\omega_{d,\varepsilon})\right) \leq 16\left(\max\{\mathfrak{d},\mathcal{L}(\mathbf{F}_{d,\delta_{d,\varepsilon}})\} + \mathcal{L}(\mathbf{F}_{0,\delta_{d,\varepsilon}})\right) \left[\max\{\mathfrak{d},\left\|\left|\mathcal{D}(\mathbf{F}_{0,\delta_{d,\varepsilon}})\right\|\right|,\left\|\left|\mathcal{D}(\mathbf{F}_{d,\delta_{d,\varepsilon}})\right\|\right|\right\}\right]^{2} \\ \cdot \left[\left(N_{\delta_{d,\varepsilon}}\right)^{1/2} \left(3m_{N_{\delta_{d,\varepsilon}}}\right)^{N_{\delta_{d,\varepsilon}}}\right]^{2} \left[\mathcal{P}(\Gamma_{\gamma_{d,\delta_{d,\varepsilon}}})\right]^{3} \left[\max_{k\in\{0,1,\dots,K_{\delta_{d,\varepsilon}}\}} \mathcal{P}\left(\mathbf{H}_{K_{\delta_{d,\varepsilon}},k,\gamma_{d,\delta_{d,\varepsilon}}}\right)\right]^{3} \\ \cdot (K_{\delta_{d,\varepsilon}} + 1)^{2} \mathfrak{d}^{2}.$$
(5.67)

Moreover, note that the fact that for all $d \in \mathbb{N}$, $\varepsilon \in (0,1]$ it holds that $\mathcal{L}(\mathbf{F}_{0,\varepsilon}) \leq \kappa \varepsilon^{-\alpha_0}$, $\||\mathcal{D}(\mathbf{F}_{0,\varepsilon})|\| \leq \kappa \varepsilon^{-\beta_0}$, $\mathcal{L}(\mathbf{F}_{d,\varepsilon}) \leq \kappa d^p \varepsilon^{-\alpha_1}$, and $\||\mathcal{D}(\mathbf{F}_{d,\varepsilon})\|| \leq \kappa d^p \varepsilon^{-\beta_1}$ implies that for all $d \in \mathbb{N}$, $\varepsilon \in (0,1]$ it holds that

$$\left(\max\{\boldsymbol{\mathfrak{d}}, \mathcal{L}(\mathbf{F}_{d,\delta_{d,\varepsilon}})\} + \mathcal{L}(\mathbf{F}_{0,\delta_{d,\varepsilon}}) \right) \left[\max\{\boldsymbol{\mathfrak{d}}, \left\| \left| \mathcal{D}(\mathbf{F}_{0,\delta_{d,\varepsilon}}) \right\| \right|, \left\| \left| \mathcal{D}(\mathbf{F}_{d,\delta_{d,\varepsilon}}) \right\| \right| \right\} \right]^{2}$$

$$\leq \left(\kappa d^{p}(\delta_{d,\varepsilon})^{-\alpha_{1}} + \kappa(\delta_{d,\varepsilon})^{-\alpha_{0}} \right) \left(\max\{\kappa, \kappa(\delta_{d,\varepsilon})^{-\beta_{0}}, \kappa d^{p}(\delta_{d,\varepsilon})^{-\beta_{1}} \} \right)^{2}$$

$$\leq 2\kappa^{3} d^{3p} \max\{ (\delta_{d,\varepsilon})^{-\alpha_{1}}, (\delta_{d,\varepsilon})^{-\alpha_{0}} \} \left(\max\{ (\delta_{d,\varepsilon})^{-\beta_{0}}, (\delta_{d,\varepsilon})^{-\beta_{1}} \} \right)^{2}$$

$$\leq 2\kappa^{3} d^{3p}(\delta_{d,\varepsilon})^{-(\max\{\alpha_{0},\alpha_{1}\}+2\max\{\beta_{0},\beta_{1}\})}.$$

$$(5.68)$$

In addition, observe that item (ii) in Lemma 5.1 shows that there exists $\bar{c} \in (0, \infty)$ such that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$\left[\left(N_{\delta_{d,\varepsilon}} \right)^{1/2} \left(3m_{N_{\delta_{d,\varepsilon}}} \right)^{N_{\delta_{d,\varepsilon}}} \right]^2 \le (\delta_{d,\varepsilon})^{-2(2+\delta)} \bar{c}.$$
(5.69)

In the next step we note that (5.24), (5.25), (5.26), and Lemma 2.5 prove that there exists $\mathfrak{C} \in [1, \infty)$ such that for all $d \in \mathbb{N}$ it holds that $\mathfrak{a}_d \leq \mathfrak{C} d^{3p+1}$, $\mathfrak{b}_d \leq \mathfrak{C} d^{(p+1)p}$, $\mathfrak{c}_d \leq \mathfrak{C} d^p$, and $c_d \leq \mathfrak{C} d^{(r+2)p^2q}$. Combining (5.31) and (5.32) therefore shows that there exists $\overline{\mathfrak{C}} \in [1, \infty)$ such that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$(\gamma_{d,\delta_{d,\varepsilon}})^{-1} \le \bar{\mathfrak{C}} d^{2(r+1)p^2q + \max\{(p+1)p,3p+1\}q} (\delta_{d,\varepsilon})^{-1} (K_{\delta_{d,\varepsilon}} + 1)$$
(5.70)

and

$$(\delta_{d,\varepsilon})^{-1} \le \bar{\mathfrak{C}}\varepsilon^{-1}d^{(r+2)p^2q+\max\{(p+1)p,3p+1\}}.$$
(5.71)

Next we combine the fact that for all $\gamma \in (0, 1]$ it holds that $\mathcal{P}(\Gamma_{\gamma}) \leq \kappa \gamma^{-r}$, the fact that for all $K \in \mathbb{N}$, $k \in \{0, 1, \ldots, K\}$, $\gamma \in (0, 1]$ it holds that $\mathcal{P}(\mathbf{H}_{K,k,\gamma}) \leq \kappa K^r \gamma^{-r}$, (5.70), and the fact that for all $\varepsilon \in (0, 1]$ it holds that $K_{\varepsilon} + 1 = (K_{\varepsilon} - 1) + 2 \leq \varepsilon^{-2} + 2 \leq 3\varepsilon^{-2}$ to obtain that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$\left[\mathcal{P}\left(\Gamma_{\gamma_{d,\delta_{d,\varepsilon}}}\right)\right]^{3} \left[\max_{k\in\{0,1,\dots,K_{\delta_{d,\varepsilon}}\}} \mathcal{P}\left(\mathbf{H}_{K_{\delta_{d,\varepsilon}},k,\gamma_{d,\delta_{d,\varepsilon}}}\right)\right]^{3} (K_{\delta_{d,\varepsilon}}+1)^{2} \\
\leq \kappa^{6} (\gamma_{d,\delta_{d,\varepsilon}})^{-6r} (K_{\delta_{d,\varepsilon}})^{3r} (K_{\delta_{d,\varepsilon}}+1)^{2} \\
\leq \kappa^{6} \bar{\mathfrak{C}}^{6r} d^{6r(2(r+1)p^{2}q+\max\{(p+1)p,3p+1\}q)} (\delta_{d,\varepsilon})^{-6r} (K_{\delta_{d,\varepsilon}}+1)^{9r+2} \\
\leq 3^{9r+2} \kappa^{6} \bar{\mathfrak{C}}^{6r} d^{6rq(2(r+1)p^{2}+(p+1)^{2})} (\delta_{d,\varepsilon})^{-6r-2(9r+2)}.$$
(5.72)

Combining this, (5.67), (5.68), (5.69), and the fact that $\eta = \max\{\alpha_0, \alpha_1\} + 2\max\{\beta_0, \beta_1\} + 2\delta + 24r + 8$ demonstrates that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$\mathcal{P}\left(\Phi_{\gamma_{d,\delta_{d,\varepsilon}},N_{\delta_{d,\varepsilon}},N_{\delta_{d,\varepsilon}},K_{\delta_{d,\varepsilon}}}^{d,0,\delta_{d,\varepsilon}}(\omega_{d,\varepsilon})\right) \leq 32\kappa^{3}d^{3p}(\delta_{d,\varepsilon})^{-(\max\{\alpha_{0},\alpha_{1}\}+2\max\{\beta_{0},\beta_{1}\})}(\delta_{d,\varepsilon})^{-2(2+\delta)}\bar{c} \\
\cdot 3^{9r+2}\kappa^{6}\bar{\mathfrak{C}}^{6r}d^{6rq(2(r+1)p^{2}+(p+1)^{2})}(\delta_{d,\varepsilon})^{-6r-2(9r+2)}\mathfrak{d}^{2} \qquad (5.73) \\
\leq 32\kappa^{9}\bar{\mathfrak{C}}^{6r}\mathfrak{d}^{2}3^{9r+2}\bar{c}d^{6rq(2(r+1)p^{2}+(p+1)^{2})+3p}(\delta_{d,\varepsilon})^{-\eta}.$$

This and (5.71) establish that there exists $c \in (0, \infty)$ such that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$\mathcal{P}\Big(\Phi_{\gamma_{d,\delta_{d,\varepsilon}},N_{\delta_{d,\varepsilon}},N_{\delta_{d,\varepsilon}},K_{\delta_{d,\varepsilon}}}^{d,0,\delta_{d,\varepsilon}}(\omega_{d,\varepsilon})\Big) \leq cd^{6rq(2(r+1)p^{2}+(p+1)^{2})+3p}d^{\eta((r+2)p^{2}q+\max\{(p+1)p,3p+1\})}\varepsilon^{-\eta} \leq cd^{(2(r+1)p^{2}+(p+1)^{2})(6r+\eta)q+3p}\varepsilon^{-\eta}.$$
(5.74)

This and (5.66) prove item (ii). The proof of Theorem 5.3 is thus complete.

5.2 ANN approximations for PDEs with specific activation functions

Corollary 5.4. Let $\kappa, T, \mathfrak{c} \in (0, \infty)$, $r \in [1, \infty)$, $p \in (1, \infty)$, $\mathfrak{q} \in [2, \infty)$, $q \in (2, \infty)$, $\mathfrak{a} \in \mathbb{R} \setminus \{-1, 1\}, \nu \in \{0, 1\}, let f : \mathbb{R} \to \mathbb{R}$ be Lipschitz continuous, for every $d \in \mathbb{N}$ let $u_d \in C^{1,2}([0, T] \times \mathbb{R}^d, \mathbb{R})$ satisfy for all $t \in [0, T], x \in \mathbb{R}^d$ that

$$\frac{\partial}{\partial t}u_d(t,x) + \mathfrak{c}\Delta_x u_d(t,x) + f(u_d(t,x)) = 0, \qquad (5.75)$$

for every $d \in \mathbb{N}$ let $\mu_d \colon \mathcal{B}(\mathbb{R}^d) \to [0,\infty)$ be a measure with

$$\int_{\mathbb{R}^d} (1 + \|x\|^{p^2 q \mathfrak{q}}) \,\mu_d(\mathrm{d}x) \le \kappa d^{r p^2 q \mathfrak{q}},\tag{5.76}$$

let $a: \mathbb{R} \to \mathbb{R}$ satisfy for all $x \in \mathbb{R}$ that $a(x) = \nu \max\{x, ax\} + (1 - \nu) \ln(1 + \exp(x))$, and assume for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ that there exists $\mathbf{G} \in \mathbf{N}$ such that for all $t \in [0, T]$, $x \in \mathbb{R}^d$ it holds that

$$\mathcal{R}_a(\mathbf{G}) \in C(\mathbb{R}^d, \mathbb{R}), \qquad \mathcal{P}(\mathbf{G}) \le \kappa d^p \varepsilon^{-\kappa}, \qquad and$$
(5.77)

$$\varepsilon \|\nabla_x u_d(T, x)\| + \varepsilon |u_d(t, x)| + |u_d(T, x) - (\mathcal{R}_a(\mathbf{G}))(x)| \le \varepsilon \kappa d^p (1 + \|x\|)^p$$
(5.78)

(cf. Definitions 2.1, 3.1, and 3.4). Then there exists $c \in \mathbb{R}$ such that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ there exists $\mathbf{U} \in \mathbf{N}$ such that

$$\mathcal{R}_{a}(\mathbf{U}) \in C(\mathbb{R}^{d+1}, \mathbb{R}), \qquad \mathcal{P}(\mathbf{U}) \leq cd^{c}\varepsilon^{-c}, \qquad and$$

$$(5.79)$$

$$\left[\int_0^T \int_{\mathbb{R}^d} |u_d(t,x) - (\mathcal{R}_a(\mathbf{U}))(t,x)|^{\mathfrak{q}} \,\mu_d(\mathrm{d}x) \,\mathrm{d}t\right]^{1/\mathfrak{q}} \le \varepsilon.$$
(5.80)

Proof of Corollary 5.4. Throughout this proof let $\not : \mathbb{R} \to \mathbb{R}$ satisfy for all $w \in \mathbb{R}$ that

$$f(w) = (2\mathbf{c})^{-1} f(w), \tag{5.81}$$

for every $d \in \mathbb{N}$ let $u_d \colon [0, 2\mathfrak{c}T] \times \mathbb{R}^d \to \mathbb{R}$ satisfy for all $t \in [0, 2\mathfrak{c}T], x \in \mathbb{R}^d$ that

$$u_d(t,x) = u_d((2\mathfrak{c})^{-1}t,x), \tag{5.82}$$

let $L \in [1, \infty)$ satisfy for all $v, w \in \mathbb{R}$ that

$$|f(v) - f(w)| \le L|v - w|, \tag{5.83}$$

for every $d \in \mathbb{N}$, $\varepsilon \in (0,1]$ let $\mathbf{F}_{d,\varepsilon} \in \mathbf{N}$ satisfy for all $t \in [0,T]$, $x \in \mathbb{R}^d$ that $\mathcal{R}_a(\mathbf{F}_{d,\varepsilon}) \in C(\mathbb{R}^d,\mathbb{R})$, $\mathcal{P}(\mathbf{F}_{d,\varepsilon}) \leq \kappa d^p \varepsilon^{-\kappa}$, and

$$\varepsilon \|\nabla_x u_d(T, x)\| + \varepsilon |u_d(t, x)| + |u_d(T, x) - (\mathcal{R}_a(\mathbf{F}_{d,\varepsilon}))(x)| \le \varepsilon \kappa d^p (1 + \|x\|)^p, \tag{5.84}$$

for every $d \in \mathbb{N}$ let $\nu_d \colon \mathcal{B}(\mathbb{R}^{d+1}) \to \mathbb{R}$ satisfy for all $B \in \mathcal{B}(\mathbb{R}^{d+1})$ that

$$\nu_d(B) = \frac{1}{2\mathfrak{c}} \int_0^{2\mathfrak{c}T} \int_{\mathbb{R}^d} \mathbb{1}_B(t, x) \,\mu_d(\mathrm{d}x) \,\mathrm{d}t,$$
(5.85)

let

$$\tilde{r} = \max\left\{r, \frac{q^2}{(q-2)(q-1)}, \frac{q}{q-1}\right\},$$
(5.86)

and let

$$\tilde{\kappa} = \max\left\{2^{\frac{p^2q\mathfrak{q}}{2}}\kappa T(2\mathfrak{c}T+1)^{p^2q\mathfrak{q}}, 2\kappa+4+(2\mathfrak{c})^{-1}(L+|f(0)|), 1728\cdot 2^{\frac{q^3+3q^2-2q}{(q-2)(q-1)}}, 19\cdot\left(2+\frac{8}{T}\right)^{\frac{q}{q-1}}\right\}$$
(5.87)

(cf. Definitions 2.1, 3.1, and 3.4). Observe that the fact that for all $d \in \mathbb{N}$ it holds that $u_d \in C^{1,2}([0,T] \times \mathbb{R}^d, \mathbb{R})$, the fact that for all $d \in \mathbb{N}$, $t \in [0,T]$, $x \in \mathbb{R}^d$ it holds that $|u_d(t,x)| \leq \kappa d^p (1+||x||)^p$, and (5.75) ensure that for all $d \in \mathbb{N}$, $t \in [0, 2\mathfrak{c}T]$, $x \in \mathbb{R}^d$ it holds that that $u_d \in C^{1,2}([0,2\mathfrak{c}T] \times \mathbb{R}^d, \mathbb{R})$, that u_d is at most polynomially growing, and that

$$\frac{\partial}{\partial t}u_d(t,x) + \frac{1}{2}\Delta_x u_d(t,x) + \not f(u_d(t,x)) = 0.$$
(5.88)

In the next step we note that the triangle inequality and (5.84) imply that for all $d \in \mathbb{N}$, $x \in \mathbb{R}^d$, $\varepsilon \in (0, 1]$ it holds that

$$\varepsilon \|\nabla_{x} u_{d}(2\mathfrak{c}T, x)\| + \varepsilon |(\mathcal{R}_{a}(\mathbf{F}_{d,\varepsilon}))(x)| + |u_{d}(2\mathfrak{c}T, x) - (\mathcal{R}_{a}(\mathbf{F}_{d,\varepsilon}))(x)|$$

$$\leq \varepsilon \|\nabla_{x} u_{d}(T, x)\| + \varepsilon (|u_{d}(T, x)| + |(\mathcal{R}_{a}(\mathbf{F}_{d,\varepsilon}))(x) - u_{d}(T, x)|) + |u_{d}(T, x) - (\mathcal{R}_{a}(\mathbf{F}_{d,\varepsilon}))(x)|$$

$$\leq \varepsilon \|\nabla_{x} u_{d}(T, x)\| + \varepsilon |u_{d}(T, x)| + 2|u_{d}(T, x) - (\mathcal{R}_{a}(\mathbf{F}_{d,\varepsilon}))(x)|$$

$$\leq 2\varepsilon \kappa d^{p} (1 + ||x||)^{p}.$$
(5.89)

Moreover, observe that the fact that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that $\mathcal{P}(\mathbf{F}_{d,\varepsilon}) \leq \kappa d^p \varepsilon^{-\kappa}$ and, for instance, [1, Lemma 2.4] show that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$\varepsilon^{\kappa} \mathcal{L}(\mathbf{F}_{d,\varepsilon}) + \varepsilon^{\kappa} ||| \mathcal{D}(\mathbf{F}_{d,\varepsilon}) ||| \le \varepsilon^{\kappa} \kappa d^{p} \varepsilon^{-\kappa} = \kappa d^{p}.$$
(5.90)

In addition, note that (5.85), the fact that for all $d \in \mathbb{N}$ it holds that μ_d is a finite measure, Jensen's inequality, Fubini's theorem, and (5.76) prove that for all $d \in \mathbb{N}$ it holds that ν_d is a finite measure on $(\mathbb{R}^{d+1}, \mathcal{B}(\mathbb{R}^{d+1}))$ with

$$\int_{\mathbb{R}^{d+1}} (1 + \|y\|^{p^{2}qq}) \nu_{d}(\mathrm{d}y) \\
\leq 2^{\frac{p^{2}qq}{2} - 1} \left(\frac{1}{2\mathfrak{c}} \int_{0}^{2\mathfrak{c}T} \int_{\mathbb{R}^{d}} |t|^{p^{2}qq} \mu_{d}(\mathrm{d}x) \,\mathrm{d}t + \frac{1}{2\mathfrak{c}} \int_{0}^{2\mathfrak{c}T} \int_{\mathbb{R}^{d}} (1 + \|x\|^{p^{2}qq}) \mu_{d}(\mathrm{d}x) \,\mathrm{d}t \right) \\
\leq 2^{\frac{p^{2}qq}{2} - 1} \left(\frac{\kappa d^{rp^{2}qq}}{2\mathfrak{c}} \int_{0}^{2\mathfrak{c}T} t^{p^{2}qq} \,\mathrm{d}t + T\kappa d^{rp^{2}qq} \right) \\
\leq 2^{\frac{p^{2}qq}{2}} \kappa T (2\mathfrak{c}T + 1)^{p^{2}qq} d^{rp^{2}qq}.$$
(5.91)

In addition, observe that the fact that for all $v, w \in \mathbb{R}$ it holds that $|f(v) - f(w)| \leq L|v - w|$ demonstrates that for all $v, w \in \mathbb{R}$ it holds that $|f(v) - f(w)| \leq (2\mathfrak{c})^{-1}L|v - w|$. Hence, [1, Corollary 4.13] and [1, Corollary 4.14] ensure that for every $\varepsilon \in (0, 1]$ there exists $\mathbf{F}_{0,\varepsilon} \in \mathbf{N}$ such that

(I) it holds that $\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}) \in C(\mathbb{R},\mathbb{R})$,

(II) it holds for all $v, w \in \mathbb{R}$ that $|(\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(v) - (\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(w)| \le (2\mathfrak{c})^{-1}L|v-w|,$

(III) it holds for all $w \in \mathbb{R}$ that $|(\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(w) - \not e(w)| \le 2\varepsilon \max\{1, |w|^p\}$, and

(IV) it holds that $\mathcal{P}(\mathbf{F}_{0,\varepsilon}) \leq 24 \left(\max\{1, \mathfrak{c}^{-1}L\} \right)^{\frac{p}{p-1}} \varepsilon^{-\frac{p}{p-1}}$.

Note that, for example, [1, Lemma 2.4] and item (IV) establish that for all $\varepsilon \in (0, 1]$ it holds that

$$\varepsilon^{\frac{p}{p-1}} \mathcal{L}(\mathbf{F}_{0,\varepsilon}) + \varepsilon^{\frac{p}{p-1}} \||\mathcal{D}(\mathbf{F}_{0,\varepsilon})||| \le 24 (\max\{1, \mathfrak{c}^{-1}L\})^{\frac{p}{p-1}}.$$
(5.92)

In addition, observe that the triangle inequality, the fact that for all $v, w \in \mathbb{R}$ it holds that $|\mathcal{F}(v) - \mathcal{F}(w)| \leq (2\mathfrak{c})^{-1}L|v - w|$, item (III), and the fact that for all $w \in \mathbb{R}$ it holds that $1 + |w|^p \leq (1 + |w|)^p$ ensure that for all $\varepsilon \in (0, 1]$, $w \in \mathbb{R}$ it holds that

$$\varepsilon |(\mathcal{R}_{a}(\mathbf{F}_{0,\varepsilon}))(w)| + |\mathbf{f}(w) - (\mathcal{R}_{a}(\mathbf{F}_{0,\varepsilon}))(w)|$$

$$\leq \varepsilon (|(\mathcal{R}_{a}(\mathbf{F}_{0,\varepsilon}))(w) - \mathbf{f}(w)| + |\mathbf{f}(w) - \mathbf{f}(0)| + |\mathbf{f}(0)|) + |\mathbf{f}(w) - (\mathcal{R}_{a}(\mathbf{F}_{0,\varepsilon}))(w)|$$

$$\leq 2|(\mathcal{R}_{a}(\mathbf{F}_{0,\varepsilon}))(w) - \mathbf{f}(w)| + \varepsilon ((2\mathbf{c})^{-1}L|w| + (2\mathbf{c})^{-1}|f(0)|)$$

$$\leq 4\varepsilon \max\{1, |w|^{p}\} + \varepsilon(2\mathbf{c})^{-1}(L + |f(0)|)(1 + |w|)$$

$$\leq \varepsilon (4 + (2\mathbf{c})^{-1}(L + |f(0)|))(1 + |w|)^{p}.$$
(5.93)

This and (5.89) imply that for all $\varepsilon \in (0, 1]$, $x \in \mathbb{R}$ it holds that

$$\varepsilon \|\nabla_x u_1(2\mathfrak{c}T, x)\| + \varepsilon |(\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(x)| + |\not(x) - (\mathcal{R}_a(\mathbf{F}_{0,\varepsilon}))(x)| \\ \le \varepsilon (2\kappa + 4 + (2\mathfrak{c})^{-1}(L + |f(0)|))(1 + |x|)^p.$$
(5.94)

In the next step we combine items (iii) and (v) in Lemma 3.5 in [1], item (i) in Lemma 3.5 in [1], and item (ii) in Lemma 3.8 in [1] to obtain that there exists $\mathfrak{J} \in \mathbf{N}$ such that

$$\mathcal{H}(\mathfrak{J}) = 1, \qquad \mathcal{D}(\mathfrak{J}) = (1, 2, 1), \qquad \text{and} \qquad \mathcal{R}_a(\mathfrak{J}) = \mathrm{id}_{\mathbb{R}}.$$
 (5.95)

Next we combine Corollary 4.10 and Corollary 4.11 to obtain that for every $\varepsilon \in (0, 1]$ there exists $\Gamma_{\varepsilon} \in \mathbf{N}$ such that for all $v, w \in \mathbb{R}$ it holds that

$$\mathcal{R}_{a}(\Gamma_{\varepsilon}) \in C(\mathbb{R}^{2}, \mathbb{R}), \qquad \mathcal{P}(\Gamma_{\varepsilon}) \leq 1728 \cdot 2^{\frac{q^{3}+3q^{2}-2q}{(q-2)(q-1)}} \varepsilon^{-\frac{q^{2}}{(q-2)(q-1)}},$$

and $|vw - (\mathcal{R}_{a}(\Gamma_{\varepsilon}))(v, w)| \leq \varepsilon \max\{1, |v|^{q}, |w|^{q}\}.$ (5.96)

In the next step we note that Lemma 4.5 and Lemma 4.6 show that for every $K \in \mathbb{N}$, $k \in \{0, 1, \ldots, K\}$, $\varepsilon \in (0, 1]$ there exists $\mathbf{H}_{K,k,\varepsilon} \in \mathbf{N}$ such that for all $t \in \mathbb{R}$ it holds that

$$\mathcal{R}_{a}(\mathbf{H}_{K,k,\varepsilon}) \in C(\mathbb{R},\mathbb{R}), \qquad \left| \left(\mathcal{R}_{a}(\mathbf{H}_{K,k,\varepsilon}) \right)(t) - \mathscr{L}^{0,1,0}_{\frac{(k-1)T}{K},\frac{kT}{K},\frac{(k+1)T}{K}}(t) \right| \leq \varepsilon \max\{1,|t|^{q}\},$$

and
$$\mathcal{P}(\mathbf{H}_{K,k,\varepsilon}) \leq 19 \cdot \left(\max\left\{1,\frac{4K}{T}\right\} \right)^{\frac{q}{q-1}} 2^{\frac{q}{q-1}} \varepsilon^{-\frac{q}{q-1}} \leq 19 \cdot \left(2 + \frac{8}{T}\right)^{\frac{q}{q-1}} K^{\frac{q}{q-1}} \varepsilon^{-\frac{q}{q-1}}$$
(5.97)

(cf. Definition 4.1). Moreover, observe that Theorem 5.3 (applied with $L \curvearrowleft (2\mathfrak{c})^{-1}L$, $\kappa \curvearrowleft \tilde{\kappa}, \alpha_0 \curvearrowleft \frac{p}{p-1}, \alpha_1 \curvearrowleft \kappa, \beta_0 \curvearrowleft \frac{p}{p-1}, \beta_1 \curvearrowleft \kappa, T \curvearrowleft 2\mathfrak{c}T, r \curvearrowleft \tilde{r}, p \curvearrowleft p, \mathfrak{q} \backsim$ $\mathfrak{q}, q \curvearrowleft q, a \curvearrowleft a, f_0 \curvearrowleft \mathfrak{f}, (f_d)_{d\in\mathbb{N}} \curvearrowleft (\mathbb{R}^d \ni x \mapsto u_d(2\mathfrak{c}T, x) \in \mathbb{R})_{d\in\mathbb{N}}, (\nu_d)_{d\in\mathbb{N}} \curvearrowleft$ $(\nu_d)_{d\in\mathbb{N}}, \mathfrak{J} \curvearrowleft \mathfrak{J}, (\mathbf{F}_{d,\varepsilon})_{(d,\varepsilon)\in\mathbb{N}_0\times(0,1]} \curvearrowleft (\mathbf{F}_{d,\varepsilon})_{(d,\varepsilon)\in\mathbb{N}_0\times(0,1]}, (\Gamma_{\varepsilon})_{\varepsilon\in(0,1]} \curvearrowleft (\Gamma_{\varepsilon})_{\varepsilon\in(0,1]}, \delta \backsim 1,$ $((\mathbf{H}_{K,k,\varepsilon})_{k\in\{0,1,\dots,K\}})_{(K,\varepsilon)\in\mathbb{N}\times(0,1]} \curvearrowleft ((\mathbf{H}_{K,k,\varepsilon})_{k\in\{0,1,\dots,K\}})_{(K,\varepsilon)\in\mathbb{N}\times(0,1]}, \eta \backsim 3\max\{\frac{p}{p-1},\kappa\}+2+$ $24\tilde{r}+8$ in the notation of Theorem 5.3) proves that there exist $(\mathbf{U}_{d,\varepsilon})_{(d,\varepsilon)\in\mathbb{N}\times(0,1]} \subseteq \mathbf{N}$ and $c \in (0,\infty)$ which satisfy for all $d \in \mathbb{N}, \varepsilon \in (0,1]$ that $\mathcal{R}_a(\mathbf{U}_{d,\varepsilon}) \in C(\mathbb{R}^{d+1},\mathbb{R}),$ $\mathcal{P}(\mathbf{U}_{d,\varepsilon}) \leq cd^c\varepsilon^{-c},$ and

$$\left[\int_{[0,2\epsilon T]\times\mathbb{R}^d} |\boldsymbol{u}_d(t,x) - (\mathcal{R}_a(\mathbf{U}_{d,\varepsilon}))(t,x)|^{\mathfrak{q}} \,\nu_d(\mathrm{d} t,\mathrm{d} x)\right]^{1/\mathfrak{q}} \leq \varepsilon.$$
(5.98)

Furthermore, note that Lemma 3.14 (applied for every $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ with $T \frown 0$, $c \frown 2\mathfrak{c}$, $\mathbf{F} \frown \mathbf{U}_{d,\varepsilon}$, $a \frown a$, $d \frown d$, $\mathfrak{d} \frown 2$, $\mathfrak{J} \frown \mathfrak{J}$ in the notation of Lemma 3.14) demonstrates that for every $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ there exists $\tilde{\mathbf{U}}_{d,\varepsilon} \in \mathbf{N}$ such that for all $s \in [0, T]$, $x \in \mathbb{R}^d$ it holds that

$$\mathcal{R}_{a}(\tilde{\mathbf{U}}_{d,\varepsilon}) \in C(\mathbb{R}^{d+1},\mathbb{R}), \qquad (\mathcal{R}_{a}(\tilde{\mathbf{U}}_{d,\varepsilon}))(s,x) = (\mathcal{R}_{a}(\mathbf{U}_{d,\varepsilon}))(2\mathfrak{c}s,x), \\
\text{and} \qquad \mathcal{P}(\tilde{\mathbf{U}}_{d,\varepsilon}) \leq \mathcal{P}(\mathbf{U}_{d,\varepsilon})384d^{2} \leq 384cd^{c+2}\varepsilon^{-c} \leq (384c+2)d^{(384c+2)}\varepsilon^{-(384c+2)}.$$
(5.99)

Combining this, (5.82), (5.85), Fubini's theorem, and a change of variables establishes that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$\int_{[0,2\mathfrak{c}T]\times\mathbb{R}^d} |\boldsymbol{u}_d(t,x) - (\mathcal{R}_a(\mathbf{U}_{d,\varepsilon}))(t,x)|^{\mathfrak{q}} \,\boldsymbol{\nu}_d(\mathrm{d}t,\mathrm{d}x) \\
= \frac{1}{2\mathfrak{c}} \int_{\mathbb{R}^d} \int_0^{2\mathfrak{c}T} |\boldsymbol{u}_d((2\mathfrak{c})^{-1}t,x) - (\mathcal{R}_a(\tilde{\mathbf{U}}_{d,\varepsilon}))((2\mathfrak{c})^{-1}t,x)|^{\mathfrak{q}} \,\mathrm{d}t \,\boldsymbol{\mu}_d(\mathrm{d}x) \qquad (5.100) \\
= \int_{\mathbb{R}^d} \int_0^T |\boldsymbol{u}_d(s,x) - (\mathcal{R}_a(\tilde{\mathbf{U}}_{d,\varepsilon}))(s,x)|^{\mathfrak{q}} \,\mathrm{d}s \,\boldsymbol{\mu}_d(\mathrm{d}x).$$

Combining this, Fubini's theorem, and (5.98) ensures that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$\left[\int_0^T \int_{\mathbb{R}^d} |u_d(t,x) - (\mathcal{R}_a(\tilde{\mathbf{U}}_{d,\varepsilon}))(t,x)|^{\mathfrak{q}} \, \mu_d(\mathrm{d}x) \, \mathrm{d}t\right]^{1/\mathfrak{q}} \le \varepsilon.$$
(5.101)

The proof of Corollary 5.4 is thus complete.

Corollary 5.5. Let $\kappa, T, \mathfrak{c}, p \in (0, \infty)$, $b_1 \in \mathbb{R}$, $b_2 \in (b_1, \infty)$, $\alpha \in \mathbb{R} \setminus \{-1, 1\}$, $\nu \in \{0, 1\}$, let $f : \mathbb{R} \to \mathbb{R}$ be Lipschitz continuous, for every $d \in \mathbb{N}$ let $u_d \in C^{1,2}([0, T] \times \mathbb{R}^d, \mathbb{R})$ satisfy for all $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$\frac{\partial}{\partial t}u_d(t,x) = \mathfrak{c}\Delta_x u_d(t,x) + f(u_d(t,x)), \qquad (5.102)$$

let $a: \mathbb{R} \to \mathbb{R}$ satisfy for all $x \in \mathbb{R}$ that $a(x) = \nu \max\{x, ax\} + (1 - \nu) \ln(1 + \exp(x))$, and assume for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ that there exists $\mathbf{G} \in \mathbf{N}$ such that for all $t \in [0, T]$, $x \in \mathbb{R}^d$ it holds that

$$\mathcal{R}_a(\mathbf{G}) \in C(\mathbb{R}^d, \mathbb{R}), \qquad \mathcal{P}(\mathbf{G}) \le \kappa d^{\kappa} \varepsilon^{-\kappa}, \qquad and \qquad (5.103)$$

$$\varepsilon \|\nabla_x u_d(0,x)\| + \varepsilon |u_d(t,x)| + |u_d(0,x) - (\mathcal{R}_a(\mathbf{G}))(x)| \le \varepsilon \kappa d^{\kappa} (1+\|x\|)^{\kappa}$$
(5.104)

(cf. Definitions 2.1, 3.1, and 3.4). Then there exists $c \in \mathbb{R}$ such that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ there exists $\mathbf{U} \in \mathbf{N}$ such that

$$\mathcal{R}_{a}(\mathbf{U}) \in C(\mathbb{R}^{d+1}, \mathbb{R}), \qquad \mathcal{P}(\mathbf{U}) \leq cd^{c}\varepsilon^{-c}, \qquad and \qquad (5.105)$$

$$\left[\int_{[0,T]\times[b_1,b_2]^d} \frac{|u_d(y) - (\mathcal{R}_a(\mathbf{U}))(y)|^p}{(b_2 - b_1)^d} \,\mathrm{d}y\right]^{1/p} \le \varepsilon.$$
(5.106)

Proof of Corollary 5.5. Throughout this proof let $(\delta_{\varepsilon,\mathfrak{p}})_{(\varepsilon,\mathfrak{p})\in(0,1]\times(0,2)}$ satisfy for all $\varepsilon \in (0,1]$, $\mathfrak{p} \in (0,2)$ that

$$\delta_{\varepsilon,\mathfrak{p}} = \varepsilon \left(\max\{1, T\} \right)^{\frac{1}{2} - \frac{1}{\mathfrak{p}}},\tag{5.107}$$

for every $d \in \mathbb{N}$ let $\mu_d \colon \mathcal{B}(\mathbb{R}^d) \to [0, 1]$ be the uniform distribution on $[b_1, b_2]^d$, and for every $d \in \mathbb{N}$ let $u_d \colon [0, T] \times \mathbb{R}^d \to \mathbb{R}$ satisfy for all $t \in [0, T]$, $x \in \mathbb{R}^d$ that

$$u_d(t,x) = u_d(T-t,x).$$
 (5.108)

Observe that the fact that for all $d \in \mathbb{N}$ it holds that $u_d \in C^{1,2}([0,T] \times \mathbb{R}^d, \mathbb{R})$ and (5.102) imply that for all $d \in \mathbb{N}$, $t \in [0,T]$, $x \in \mathbb{R}^d$ it holds that $u_d \in C^{1,2}([0,T] \times \mathbb{R}^d, \mathbb{R})$ and

$$\frac{\partial}{\partial t}\boldsymbol{u}_d(t,x) = -\boldsymbol{\mathfrak{c}}\Delta_x\boldsymbol{u}_d(t,x) - f(\boldsymbol{u}_d(t,x)).$$
(5.109)

Next, note that (5.104) shows that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $t \in [0, T]$, $x \in \mathbb{R}^d$ it holds that

$$\varepsilon \|\nabla_x u_d(T, x)\| + \varepsilon |u_d(t, x)| + |u_d(T, x) - (\mathcal{R}_a(\mathbf{G}))(x)| \le \varepsilon \kappa d^{\kappa} (1 + \|x\|)^{\kappa}$$
(5.110)

(cf. Definitions 2.1 and 3.4). Next, observe that the fact that for all $d \in \mathbb{N}$ it holds that μ_d is the uniform distribution on $[b_1, b_2]^d$ and, for instance, [36, Lemma 3.15] prove that for all $d \in \mathbb{N}$, $\mathfrak{q} \in [2, \infty)$, $q \in (2, \infty)$ it holds that

$$\int_{\mathbb{R}^{d}} (1 + \|x\|^{(\kappa+1)^{2}q\mathfrak{q}}) \mu_{d}(\mathrm{d}x) = 1 + \frac{1}{|b_{1} - b_{2}|^{d}} \int_{[b_{1}, b_{2}]^{d}} \|x\|^{(\kappa+1)^{2}q\mathfrak{q}} \,\mathrm{d}x \\
\leq 1 + d^{(\kappa+1)^{2}q\mathfrak{q}} \max\{|b_{1}|^{(\kappa+1)^{2}q\mathfrak{q}}, |b_{2}|^{(\kappa+1)^{2}q\mathfrak{q}}\} \\
\leq 2d^{(\kappa+1)^{2}q\mathfrak{q}} \max\{1, |b_{1}|^{3\mathfrak{q}(\kappa+1)^{2}}, |b_{2}|^{3\mathfrak{q}(\kappa+1)^{2}}\}.$$
(5.111)

In addition, note that Corollary 5.4 (applied for every $\mathbf{q} \in [2, \infty)$ with $T \frown T$, $\mathbf{c} \frown \mathbf{c}$, $r \frown 1$, $p \frown \kappa + 1$, $\mathbf{q} \frown \mathbf{q}$, $q \frown 3$, $\alpha \frown \alpha$, $\nu \frown \nu$, $a \frown a$, $f \frown f$, $(u_d)_{d \in \mathbb{N}} \frown (u_d)_{d \in \mathbb{N}}$, $(\mu_d)_{d \in \mathbb{N}} \frown (\mu_d)_{d \in \mathbb{N}}$, $(\mu_d)_{d \in \mathbb{N}} \frown (\mu_d)_{d \in \mathbb{N}}$, $\mathbf{G} \frown \mathbf{G}$, $\kappa \frown 2 \max\{1, \kappa, |b_1|^{3\mathfrak{q}(\kappa+1)^2}, |b_2|^{3\mathfrak{q}(\kappa+1)^2}\}$ in the notation of Corollary 5.4) demonstrates that there exist $(\mathbf{U}_{d,\varepsilon,\mathfrak{q}})_{(d,\varepsilon,\mathfrak{q})\in\mathbb{N}\times(0,1]\times[2,\infty)} \subseteq \mathbf{N}$ and $(c_{\mathfrak{q}})_{\mathfrak{q}\in[2,\infty)} \subseteq \mathbb{R}$ which satisfy for all $d \in \mathbb{N}$, $\varepsilon \in (0,1]$, $\mathbf{q} \in [2,\infty)$ that $\mathcal{R}_a(\mathbf{U}_{d,\varepsilon,\mathfrak{q}}) \in C(\mathbb{R}^{d+1},\mathbb{R})$, $\mathcal{P}(\mathbf{U}_{d,\varepsilon,\mathfrak{q}}) \leq c_{\mathfrak{q}} d^{c_{\mathfrak{q}}} \varepsilon^{-c_{\mathfrak{q}}}$, and

$$\left[\int_{0}^{T}\int_{\mathbb{R}^{d}}|\boldsymbol{u}_{d}(t,x)-(\mathcal{R}_{a}(\mathbf{U}_{d,\varepsilon,\mathfrak{q}}))(t,x)|^{\mathfrak{q}}\,\boldsymbol{\mu}_{d}(\mathrm{d}x)\,\mathrm{d}t\right]^{1/\mathfrak{q}}\leq\varepsilon\tag{5.112}$$

(cf. Definition 3.1). Next we combine items (iii) and (v) in Lemma 3.5 in [1], item (i) in Lemma 3.5 in [1], and item (ii) in Lemma 3.8 in [1] to obtain that there exists $\mathfrak{J} \in \mathbf{N}$ such

that $\mathcal{D}(\mathfrak{J}) = (1, 2, 1)$ and $\mathcal{R}_a(\mathfrak{J}) = \mathrm{id}_{\mathbb{R}}$. Next, observe that Lemma 3.14 (applied for every $d \in \mathbb{N}, \varepsilon \in (0, 1], \mathfrak{q} \in [2, \infty)$ with $T \frown T, c \frown -1, \mathbf{F} \frown \mathbf{U}_{d,\varepsilon,\mathfrak{q}}, a \frown a, d \frown d, \mathfrak{d} \frown 2, \mathfrak{J} \frown \mathfrak{J}$ in the notation of Lemma 3.14) establishes that for every $d \in \mathbb{N}, \varepsilon \in (0, 1], \mathfrak{q} \in [2, \infty)$ there exists $\tilde{\mathbf{U}}_{d,\varepsilon,\mathfrak{q}} \in \mathbf{N}$ such that for all $s \in [0, T], x \in \mathbb{R}^d$ it holds that

$$\mathcal{R}_{a}(\tilde{\mathbf{U}}_{d,\varepsilon,\mathfrak{q}}) \in C(\mathbb{R}^{d+1},\mathbb{R}), \qquad (\mathcal{R}_{a}(\tilde{\mathbf{U}}_{d,\varepsilon,\mathfrak{q}}))(s,x) = (\mathcal{R}_{a}(\mathbf{U}_{d,\varepsilon,\mathfrak{q}}))(T-s,x),$$

and
$$\mathcal{P}(\tilde{\mathbf{U}}_{d,\varepsilon,\mathfrak{q}}) \leq \mathcal{P}(\mathbf{U}_{d,\varepsilon,\mathfrak{q}}) 384d^{2} \leq 384c_{\mathfrak{q}}d^{c_{\mathfrak{q}}+2}\varepsilon^{-c_{\mathfrak{q}}} \leq (384c_{\mathfrak{q}}+2)d^{(384c_{\mathfrak{q}}+2)}\varepsilon^{-(384c_{\mathfrak{q}}+2)}.$$

(5.113)

This ensures that for every $\mathbf{q} \in [2, \infty)$ there exists $\tilde{c}_{\mathbf{q}} \in \mathbb{R}$ such that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$ it holds that

$$\mathcal{P}(\tilde{\mathbf{U}}_{d,\varepsilon,\mathfrak{q}}) \le \tilde{c}_{\mathfrak{q}} d^{\tilde{c}_{\mathfrak{q}}} \varepsilon^{-\tilde{c}_{\mathfrak{q}}}.$$
(5.114)

Furthermore, note that (5.108), (5.112), (5.113), Fubini's theorem, a change of variables, and the fact that for all $d \in \mathbb{N}$ it holds that μ_d is the uniform distribution on $[b_1, b_2]^d$ imply that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $\mathfrak{q} \in [2, \infty)$ it holds that

$$\varepsilon \ge \left[\int_0^T \int_{\mathbb{R}^d} |u_d(T-t,x) - (\mathcal{R}_a(\tilde{\mathbf{U}}_{d,\varepsilon,\mathfrak{q}}))(T-t,x)|^{\mathfrak{q}} \mu_d(\mathrm{d}x) \,\mathrm{d}t \right]^{1/\mathfrak{q}}$$
$$= \left[\int_0^T \int_{\mathbb{R}^d} |u_d(s,x) - (\mathcal{R}_a(\tilde{\mathbf{U}}_{d,\varepsilon,\mathfrak{q}}))(s,x)|^{\mathfrak{q}} \mu_d(\mathrm{d}x) \,\mathrm{d}s \right]^{1/\mathfrak{q}}$$
$$= \left[\int_{[0,T] \times [b_1, b_2]^d} \frac{|u_d(y) - (\mathcal{R}_a(\tilde{\mathbf{U}}_{d,\varepsilon,\mathfrak{q}}))(y)|^{\mathfrak{q}}}{(b_2 - b_1)^d} \,\mathrm{d}y \right]^{1/\mathfrak{q}}.$$
(5.115)

Combining the fact that for all $\varepsilon \in (0, 1]$, $\mathfrak{p} \in (0, 2)$ it holds that $\delta_{\varepsilon, \mathfrak{p}} \in (0, 1]$ and Jensen's inequality hence shows that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $\mathfrak{p} \in (0, 2)$ it holds that

$$\begin{bmatrix}
\int_{[0,T]\times[b_{1},b_{2}]^{d}} \frac{|u_{d}(y) - (\mathcal{R}_{a}(\tilde{\mathbf{U}}_{d,\delta_{\varepsilon,\mathfrak{p}},2}))(y)|^{\mathfrak{p}}}{(b_{2} - b_{1})^{d}} \,\mathrm{d}y \end{bmatrix}^{2/\mathfrak{p}} \\
\leq T^{\frac{2}{\mathfrak{p}}-1} \left[\int_{[0,T]\times[b_{1},b_{2}]^{d}} \frac{|u_{d}(y) - (\mathcal{R}_{a}(\tilde{\mathbf{U}}_{d,\delta_{\varepsilon,\mathfrak{p}},2}))(y)|^{2}}{(b_{2} - b_{1})^{d}} \,\mathrm{d}y \right] \leq T^{\frac{2}{\mathfrak{p}}-1} (\delta_{\varepsilon,\mathfrak{p}})^{2}.$$
(5.116)

This and (5.107) prove that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $\mathfrak{p} \in (0, 2)$ it holds that

$$\left[\int_{[0,T]\times[b_1,b_2]^d} \frac{|u_d(y) - (\mathcal{R}_a(\tilde{\mathbf{U}}_{d,\delta_{\varepsilon,\mathfrak{p}},2}))(y)|^{\mathfrak{p}}}{(b_2 - b_1)^d} \,\mathrm{d}y\right]^{1/\mathfrak{p}} \le T^{\frac{1}{\mathfrak{p}} - \frac{1}{2}} \delta_{\varepsilon,\mathfrak{p}} \le \varepsilon.$$
(5.117)

Furthermore, observe that (5.107) and (5.114) demonstrate that for all $d \in \mathbb{N}$, $\varepsilon \in (0, 1]$, $\mathfrak{p} \in (0, 2)$, and $\hat{c}_{\mathfrak{p}} = \tilde{c}_2(\max\{1, T\})^{(\frac{1}{\mathfrak{p}} - \frac{1}{2})\tilde{c}_2}$ it holds that

$$\mathcal{P}(\tilde{\mathbf{U}}_{d,\delta_{\varepsilon,\mathfrak{p}},2}) \leq \tilde{c}_2 d^{\tilde{c}_2} \varepsilon^{-\tilde{c}_2} \left(\max\{1,T\} \right)^{\left(\frac{1}{\mathfrak{p}}-\frac{1}{2}\right)\tilde{c}_2} \leq \hat{c}_{\mathfrak{p}} d^{\hat{c}_{\mathfrak{p}}} \varepsilon^{-\hat{c}_{\mathfrak{p}}}.$$
(5.118)

Combining this with (5.114), (5.115), and (5.117) establishes (5.105) and (5.106). The proof of Corollary 5.5 is thus complete.

Acknowledgments

This project has been partially funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy EXC 2044-390685587, Mathematics Münster: Dynamics-Geometry-Structure (third author). This project has also been partially funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) in the frame of the priority programme SPP 2298 'Theoretical Foundations of Deep Learning' – Project no. 464123384 (third author). This work has also been partially supported by the Internal Project Fund from Shenzhen Research Institute of Big Data under Grant T00120220001 (second author).

References

- ACKERMANN, J., JENTZEN, A., KRUSE, T., KUCKUCK, B., AND PADGETT, J. L. Deep neural networks with ReLU, leaky ReLU, and softplus activation provably overcome the curse of dimensionality for Kolmogorov partial differential equations with Lipschitz nonlinearities in the L^p-sense. arXiv:2309.13722 (2023), 52 pages.
- [2] BAGGENSTOS, J., AND SALIMOVA, D. Approximation properties of residual neural networks for Kolmogorov PDEs. Discrete Contin. Dyn. Syst. Ser. B 28, 5 (2023), 3193– 3215.
- [3] BAO, G., YE, X., ZANG, Y., AND ZHOU, H. Numerical solution of inverse problems by weak adversarial networks. *Inverse Problems* 36, 11 (2020), 115003, 31.
- [4] BECK, C., BECKER, S., CHERIDITO, P., JENTZEN, A., AND NEUFELD, A. Deep splitting method for parabolic PDEs. SIAM J. Sci. Comput. 43, 5 (2021), A3135– A3154.
- [5] BECK, C., BECKER, S., GROHS, P., JAAFARI, N., AND JENTZEN, A. Solving the Kolmogorov PDE by means of deep learning. J. Sci. Comput. 88, 3 (2021), Paper No. 73, 28.
- [6] BECK, C., E, W., AND JENTZEN, A. Machine learning approximation algorithms for high-dimensional fully nonlinear partial differential equations and second-order backward stochastic differential equations. arXiv:1709.05963 (2017), 56 pages.
- [7] BECK, C., E, W., AND JENTZEN, A. Machine learning approximation algorithms for high-dimensional fully nonlinear partial differential equations and second-order backward stochastic differential equations. J. Nonlinear Sci. 29, 4 (2019), 1563–1619.
- [8] BECK, C., GONON, L., AND JENTZEN, A. Overcoming the curse of dimensionality in the numerical approximation of high-dimensional semilinear elliptic partial differential equations. *arXiv:2003.00596* (2020), 50 pages. Accepted in Partial Differ. Equ. Appl.
- [9] BECK, C., HORNUNG, F., HUTZENTHALER, M., JENTZEN, A., AND KRUSE, T. Overcoming the curse of dimensionality in the numerical approximation of Allen-Cahn

partial differential equations via truncated full-history recursive multilevel Picard approximations. J. Numer. Math. 28, 4 (2020), 197–222.

- [10] BECK, C., HUTZENTHALER, M., AND JENTZEN, A. On nonlinear Feynman–Kac formulas for viscosity solutions of semilinear parabolic partial differential equations. *Stochastics and Dynamics* 21, 08 (2021), 2150048.
- [11] BECK, C., HUTZENTHALER, M., JENTZEN, A., AND KUCKUCK, B. An overview on deep learning-based approximation methods for partial differential equations. *Discrete Contin. Dyn. Syst. Ser. B* 28, 6 (2023), 3697–3746.
- [12] BECK, C., JENTZEN, A., KLEINBERG, K., AND KRUSE, T. Nonlinear Monte Carlo methods with polynomial runtime for Bellman equations of discrete time highdimensional stochastic optimal control problems. arXiv:2303.03390 (2023), 33 pages.
- [13] BELLMAN, R. Dynamic programming. Princeton Landmarks in Mathematics. Princeton University Press, Princeton, NJ, 2010. Reprint of the 1957 edition, With a new introduction by Stuart Dreyfus.
- [14] BERG, J., AND NYSTRÖM, K. A unified deep artificial neural network approach to partial differential equations in complex geometries. *Neurocomputing* 317 (2018), 28– 41.
- [15] BERNER, J., GROHS, P., AND JENTZEN, A. Analysis of the generalization error: empirical risk minimization over deep artificial neural networks overcomes the curse of dimensionality in the numerical approximation of Black-Scholes partial differential equations. SIAM J. Math. Data Sci. 2, 3 (2020), 631–657.
- [16] BLECHSCHMIDT, J., AND ERNST, O. G. Three ways to solve partial differential equations with neural network—a review. *GAMM-Mitt.* 44, 2 (2021), Paper No. e202100006, 29.
- [17] CHAN-WAI-NAM, Q., MIKAEL, J., AND WARIN, X. Machine learning for semi linear PDEs. J. Sci. Comput. 79, 3 (2019), 1667–1712.
- [18] CHEN, F., HUANG, J., WANG, C., AND YANG, H. Friedrichs learning: weak solutions of partial differential equations via deep learning. *SIAM J. Sci. Comput.* 45, 3 (2023), A1271–A1299.
- [19] CHERIDITO, P., JENTZEN, A., AND ROSSMANNEK, F. Non-convergence of stochastic gradient descent in the training of deep neural networks. J. Complexity 64 (2021), Paper No. 101540, 10.
- [20] CHERIDITO, P., AND ROSSMANNEK, F. Efficient Sobolev approximation of linear parabolic PDEs in high dimensions. arXiv:2306.16811 (2023), 28 pages.
- [21] CIOICA-LICHT, P. A., HUTZENTHALER, M., AND WERNER, P. T. Deep neural networks overcome the curse of dimensionality in the numerical approximation of semilinear partial differential equations. arXiv:2205.14398 (2022), 34 pages.

- [22] CUOMO, S., SCHIANO DI COLA, V., GIAMPAOLO, F., ROZZA, G., RAISSI, M., AND PICCIALLI, F. Scientific machine learning through physics-informed neural networks: where we are and what's next. J. Sci. Comput. 92, 3 (2022), Paper No. 88, 62.
- [23] E, W., HAN, J., AND JENTZEN, A. Deep learning-based numerical methods for highdimensional parabolic partial differential equations and backward stochastic differential equations. *Commun. Math. Stat.* 5, 4 (2017), 349–380.
- [24] E, W., HAN, J., AND JENTZEN, A. Algorithms for solving high dimensional PDEs: from nonlinear Monte Carlo to machine learning. *Nonlinearity* 35, 1 (2022), 278–310.
- [25] E, W., HUTZENTHALER, M., JENTZEN, A., AND KRUSE, T. Multilevel Picard iterations for solving smooth semilinear parabolic heat equations. *Partial Differ. Equ. Appl.* 2, 6 (2021), Paper No. 80, 31.
- [26] E, W., AND YU, B. The deep Ritz method: A deep learning-based numerical algorithm for solving variational problems. *Commun. Math. Stat.* 6, 1 (2018), 1–12.
- [27] ELBRÄCHTER, D., GROHS, P., JENTZEN, A., AND SCHWAB, C. DNN expression rate analysis of high-dimensional PDEs: Application to option pricing. *Constr. Approx.* (2021), 1–69.
- [28] GENTILE, R. Function approximation guarantees for a shallow neural network trained by gradient flow. Master's thesis, University of Central Florida (2022). https://stars.library.ucf.edu/etd2020/1203.
- [29] GENTILE, R., AND WELPER, G. Approximation results for gradient descent trained shallow neural networks in 1d. arXiv:2209.08399 (2022), 49 pages.
- [30] GERMAIN, M., PHAM, H., AND WARIN, X. Neural Networks-Based Algorithms for Stochastic Control and PDEs in Finance. Cambridge University Press, 2023, pp. 426– -452.
- [31] GILES, M. B., JENTZEN, A., AND WELTI, T. Generalised multilevel Picard approximations. arXiv:1911.03188 (2019), 61 pages.
- [32] GONON, L. Random feature neural networks learn Black-Scholes type PDEs without curse of dimensionality. J. Mach. Learn. Res. 24 (2023), Paper No. [189], 51.
- [33] GONON, L., GROHS, P., JENTZEN, A., KOFLER, D., AND ŠIŠKA, D. Uniform error estimates for artificial neural network approximations for heat equations. *IMA J. Numer. Anal.* 42, 3 (2022), 1991–2054.
- [34] GONON, L., AND SCHWAB, C. Deep ReLU network expression rates for option prices in high-dimensional, exponential Lévy models. *Finance Stoch.* 25, 4 (2021), 615–657.
- [35] GONON, L., AND SCHWAB, C. Deep ReLU neural networks overcome the curse of dimensionality for partial integrodifferential equations. Anal. Appl. (Singap.) 21, 1 (2023), 1–47.

- [36] GROHS, P., HORNUNG, F., JENTZEN, A., AND VON WURSTEMBERGER, P. A proof that artificial neural networks overcome the curse of dimensionality in the numerical approximation of Black-Scholes partial differential equations. *Mem. Amer. Math. Soc.* 284, 1410 (2023), v+93.
- [37] GROHS, P., HORNUNG, F., JENTZEN, A., AND ZIMMERMANN, P. Space-time error estimates for deep neural network approximations for differential equations. Adv. Comput. Math. 49, 1 (2023), Paper No. 4, 78.
- [38] GROHS, P., JENTZEN, A., AND SALIMOVA, D. Deep neural network approximations for solutions of PDEs based on Monte Carlo algorithms. *Partial Differ. Equ. Appl. 3*, 4 (2022), Paper No. 45, 41.
- [39] GU, Y., YANG, H., AND ZHOU, C. SelectNet: self-paced learning for high-dimensional partial differential equations. J. Comput. Phys. 441 (2021), Paper No. 110444, 18.
- [40] HAN, J., JENTZEN, A., AND E, W. Solving high-dimensional partial differential equations using deep learning. Proc. Natl. Acad. Sci. USA 115, 34 (2018), 8505–8510.
- [41] HORNUNG, F., JENTZEN, A., AND SALIMOVA, D. Space-time deep neural network approximations for high-dimensional partial differential equations. arXiv:2006.02199 (2020), 52 pages. Accepted in J. Comput. Math.
- [42] HU, Z., SHUKLA, K., KARNIADAKIS, G. E., AND KAWAGUCHI, K. Tackling the curse of dimensionality with physics-informed neural networks. arXiv:2307.12306 (2023), 37 pages.
- [43] HURÉ, C., PHAM, H., AND WARIN, X. Deep backward schemes for high-dimensional nonlinear PDEs. *Math. Comp.* 89, 324 (2020), 1547–1579.
- [44] HUTZENTHALER, M., JENTZEN, A., AND KRUSE, T. Overcoming the curse of dimensionality in the numerical approximation of parabolic partial differential equations with gradient-dependent nonlinearities. *Found. Comput. Math.* 22, 4 (2022), 905–966.
- [45] HUTZENTHALER, M., JENTZEN, A., KRUSE, T., AND NGUYEN, T. A. Multilevel Picard approximations for high-dimensional semilinear second-order PDEs with Lipschitz nonlinearities. arXiv:2009.02484 (2020), 37 pages.
- [46] HUTZENTHALER, M., JENTZEN, A., KRUSE, T., AND NGUYEN, T. A. A proof that rectified deep neural networks overcome the curse of dimensionality in the numerical approximation of semilinear heat equations. *Partial Differ. Equ. Appl.* 1, 2 (2020), Paper No. 10, 34.
- [47] HUTZENTHALER, M., JENTZEN, A., KRUSE, T., NGUYEN, T. A., AND VON WURSTEMBERGER, P. Overcoming the curse of dimensionality in the numerical approximation of semilinear parabolic partial differential equations. *Proc. A.* 476, 2244 (2020), 20190630, 25.

- [48] HUTZENTHALER, M., JENTZEN, A., KUCKUCK, B., AND PADGETT, J. L. Strong L^perror analysis of nonlinear Monte Carlo approximations for high-dimensional semilinear partial differential equations. arXiv:2110.08297 (2021), 42 pages. Revision requested from Numer. Algorithms.
- [49] HUTZENTHALER, M., JENTZEN, A., AND VON WURSTEMBERGER, P. Overcoming the curse of dimensionality in the approximative pricing of financial derivatives with default risks. *Electron. J. Probab.* 25 (2020), Paper No. 101, 73.
- [50] HUTZENTHALER, M., AND KRUSE, T. Multilevel Picard approximations of highdimensional semilinear parabolic differential equations with gradient-dependent nonlinearities. SIAM J. Numer. Anal. 58, 2 (2020), 929–961.
- [51] HUTZENTHALER, M., KRUSE, T., AND NGUYEN, T. A. Multilevel Picard approximations for McKean-Vlasov stochastic differential equations. J. Math. Anal. Appl. 507, 1 (2022), Paper No. 125761, 14.
- [52] HUTZENTHALER, M., AND NGUYEN, T. A. Multilevel Picard approximations of highdimensional semilinear partial differential equations with locally monotone coefficient functions. *Appl. Numer. Math.* 181 (2022), 151–175.
- [53] IBRAGIMOV, S., JENTZEN, A., AND RIEKERT, A. Convergence to good non-optimal critical points in the training of neural networks: Gradient descent optimization with one random initialization overcomes all bad non-global local minima with high probability. arXiv:2212.13111 (2022), 98 pages.
- [54] JENTZEN, A., KUCKUCK, B., AND VON WURSTEMBERGER, P. Mathematical Introduction to Deep Learning: Methods, Implementations, and Theory. arXiv:2310.20360, 2023.
- [55] JENTZEN, A., AND RIEKERT, A. Non-convergence to global minimizers for Adam and stochastic gradient descent optimization and constructions of local minimizers in the training of artificial neural networks. arXiv:2402.05155 (2024), 36 pages.
- [56] JENTZEN, A., SALIMOVA, D., AND WELTI, T. A proof that deep artificial neural networks overcome the curse of dimensionality in the numerical approximation of Kolmogorov partial differential equations with constant diffusion and nonlinear drift coefficients. *Commun. Math. Sci.* 19, 5 (2021), 1167–1205.
- [57] JOVANOVIĆ, B. S., AND SÜLI, E. Analysis of finite difference schemes, vol. 46 of Springer Series in Computational Mathematics. Springer, London, 2014. For linear partial differential equations with generalized solutions.
- [58] KARNIADAKIS, G. E., KEVREKIDIS, I. G., LU, L., PERDIKARIS, P., WANG, S., AND YANG, L. Physics-informed machine learning. *Nat. Rev. Phys.* 3 (2021), 422–440.
- [59] KREMSNER, S., STEINICKE, A., AND SZÖLGYENYI, M. A deep neural network algorithm for semilinear elliptic PDEs with applications in insurance mathematics. *Risks* 8, 4 (2020).

- [60] LU, L., MENG, X., MAO, Z., AND KARNIADAKIS, G. E. DeepXDE: a deep learning library for solving differential equations. SIAM Rev. 63, 1 (2021), 208–228.
- [61] LU, L., SHIN, Y., SU, Y., AND KARNIADAKIS, G. E. Dying ReLU and initialization: theory and numerical examples. *Commun. Comput. Phys.* 28, 5 (2020), 1671–1706.
- [62] NEUFELD, A., AND NGUYEN, T. A. Rectified deep neural networks overcome the curse of dimensionality when approximating solutions of McKean–Vlasov stochastic differential equations. arXiv:2312.07042 (2023), 21 pages.
- [63] NEUFELD, A., AND NGUYEN, T. A. Rectified deep neural networks overcome the curse of dimensionality in the numerical approximation of gradient-dependent semilinear heat equations. arXiv:2403.09200 (2024), 22 pages.
- [64] NEUFELD, A., NGUYEN, T. A., AND WU, S. Deep ReLU neural networks overcome the curse of dimensionality when approximating semilinear partial integro-differential equations. arXiv:2310.15581 (2023), 38 pages.
- [65] NEUFELD, A., NGUYEN, T. A., AND WU, S. Multilevel Picard approximations overcome the curse of dimensionality in the numerical approximation of general semilinear PDEs with gradient-dependent nonlinearities. arXiv:2311.11579 (2023), 62 pages.
- [66] NEUFELD, A., AND WU, S. Multilevel Picard approximation algorithm for semilinear partial integro-differential equations and its complexity analysis. arXiv:2205.09639 (2023), 55 pages.
- [67] NOVAK, E., AND RITTER, K. The curse of dimension and a universal method for numerical integration. In *Multivariate approximation and splines (Mannheim, 1996)*, vol. 125 of *Internat. Ser. Numer. Math. Birkhäuser*, Basel, 1997, pp. 177–187.
- [68] NOVAK, E., AND WOŹNIAKOWSKI, H. Tractability of multivariate problems. Vol. 1: Linear information, vol. 6 of EMS Tracts in Mathematics. European Mathematical Society (EMS), Zürich, 2008.
- [69] NÜSKEN, N., AND RICHTER, L. Solving high-dimensional Hamilton-Jacobi-Bellman PDEs using neural networks: perspectives from the theory of controlled diffusions and measures on path space. *Partial Differ. Equ. Appl. 2*, 4 (2021), Paper No. 48, 48.
- [70] PHAM, H., WARIN, X., AND GERMAIN, M. Neural networks-based backward scheme for fully nonlinear PDEs. *Partial Differ. Equ. Appl.* 2, 1 (2021), Paper No. 16, 24.
- [71] RAISSI, M. Forward-Backward Stochastic Neural Networks: Deep Learning of Highdimensional Partial Differential Equations. arXiv:1804.07010 (2018), 17 pages.
- [72] RAISSI, M., PERDIKARIS, P., AND KARNIADAKIS, G. E. Physics-informed neural networks: a deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. J. Comput. Phys. 378 (2019), 686–707.

- [73] REISINGER, C., AND ZHANG, Y. Rectified deep neural networks overcome the curse of dimensionality for nonsmooth value functions in zero-sum games of nonlinear stiff systems. Anal. Appl. (Singap.) 18, 6 (2020), 951–999.
- [74] SHIN, Y., AND KARNIADAKIS, G. E. Trainability of ReLU networks and datadependent initialization. Journal of Machine Learning for Modeling and Computing 1, 1 (2020), 39–74.
- [75] SIMON, M. K. Probability distributions involving Gaussian random variables: A handbook for engineers and scientists. Springer Science & Business Media, 2007.
- [76] SIRIGNANO, J., AND SPILIOPOULOS, K. DGM: a deep learning algorithm for solving partial differential equations. J. Comput. Phys. 375 (2018), 1339–1364.
- [77] VALSECCHI OLIVA, P., WU, Y., HE, C., AND NI, H. Towards fast weak adversarial training to solve high dimensional parabolic partial differential equations using XNODE-WAN. J. Comput. Phys. 463 (2022), Paper No. 111233, 17.
- [78] WELPER, G. Approximation results for gradient descent trained neural networks. arXiv:2309.04860 (2023), 69 pages.
- [79] XIAO, J., AND WANG, X. Empirical risk minimization over artificial neural networks overcomes the curse of dimensionality in the numerical approximation of linear Kolmogorov partial differential equations with unbounded initial functions. arXiv:2310.12582 (2023), 23 pages.
- [80] ZANG, Y., BAO, G., YE, X., AND ZHOU, H. Weak adversarial networks for highdimensional partial differential equations. J. Comput. Phys. 411 (2020), 109409, 14.