

PROJECTION METHODS FOR OPERATOR LEARNING AND UNIVERSAL APPROXIMATION

EMANUELE ZAPPALA

ABSTRACT. We obtain a new universal approximation theorem for continuous operators on arbitrary Banach spaces using the Leray-Schauder mapping. Moreover, we introduce and study a method for operator learning in Banach spaces L^p of functions with multiple variables, based on orthogonal projections on polynomial bases. We derive a universal approximation result for operators where we learn a linear projection and a finite dimensional mapping under some additional assumptions. For the case of $p = 2$, we give some sufficient conditions for the approximation results to hold. This article serves as the theoretical framework for a deep learning methodology whose implementation will be provided in subsequent work.

CONTENTS

1. Introduction	1
2. Nonlinear Projections for Operator Learning in Banach spaces	2
3. Learning Linear Projections on Banach Spaces of functions	4
4. Learning Linear Projections on the Hilbert space	6
5. Approximations for Fixed Points	7
6. Future Perspectives	9
References	9

1. INTRODUCTION

Operator learning is a branch of deep learning involved with approximating (potentially highly nonlinear) continuous operators between Banach spaces. The interest of operator learning lies in the fact that it allows to model complex phenomena, e.g. dynamical systems, whose underlying governing equations are not known [12, 17, 18]. The study of operator learning was initiated by the theoretical work [2], whose implementation was given in [12].

Projection methods, e.g. Galerkin methods, are approaches for finding solutions of an operator equation by approximating this on prescribed subspaces through a projection [3, 9]. After projecting the operator equation on a subspace, it is not necessarily true that this equation has a solution. When projected solutions exist, upon increasing the dimension of the subspaces one would want the solutions to converge to a solution of the original equation. This is not necessarily the case. The main question of projection methods is whether projected solutions exist, and converge to a solution of the original non-projected equation.

We can formulate the problem of operator learning in relation to projection methods as follows. We want to learn projections on (finite dimensional) subspaces, and a map between subspaces such that we can approximate a target operator between Banach spaces. This operator is assumed to satisfy an operator equation whose solutions model the data, as formulated in [17, 18] for the case

of integral and integro-differential operators. Our problem is therefore two-fold. First, we want to learn an operator whose solutions of a corresponding operator equation model data. Second, we want to approximate the operator on a projected space, and solve the projected operator equation. A similar approach for a class of integral operators was employed in [19].

In the present article we theoretically address these problems, and derive a methodology for operator learning based on learning projections and mappings between projected spaces. The concrete implementation of the present results will be deferred to subsequent work. Here, we obtain a universal approximation theorem for operators between Banach spaces using the Leray-Schauder mappings. This result is more general than the main result in [2] since the latter refers to Banach spaces of functions.

The main issue with the approach based on Leray-Schauder mappings is that we need to find elements of the Banach spaces that approximate given compact subsets. Doing this in practice is not simple. An algorithm that concretely implements this approach should also obtain the points whose linear subspace will be used for the Leray-Schauder (nonlinear) projection. Additionally, it is unclear how to determine whether the projected equations produce solutions that approximate solutions to a given operator equation, even when we approximate the operator appropriately.

To address the issues raised above, we consider the more specific (and extremely important) case of L^p spaces. In this case, we show that given a set of orthogonal polynomials with respect to a quasi-inner product with some mild additional assumptions, we can find a triple of neural networks defining two projections and a map between the projected spaces that approximate the given operator with arbitrary precision. When $p = 2$, and we are working with the Hilbert space L^2 , we give some simple sufficient conditions for the aforementioned results to be applicable. We then turn to the problem of approximating solutions of the projected operator equation. We provide in this case some sufficient conditions for the projection to admit solutions for each n , and such that the solutions converge to a solution of the operator equation when $n \rightarrow \infty$.

This article is organized as follows. In Section 2 we provide the universal approximation result for general Banach spaces by means of Leray-Schauder mappings. In Section 3 we consider L^p spaces, and obtain a universal approximation result with linear projections on finite spaces of polynomials. In Section 4 we consider the case $p = 2$, and give some simple sufficient conditions for the universal approximation results to be applicable, along with examples. In Section 5 we consider operator equations for operator learning problems. We determine some conditions under which our framework produces solutions to the projected equations that converge to solutions of the operator equation, reformulated as a fixed point problem. We conclude with some remarks, in Section 6, that describe future work on the algorithmic implementation of the deep learning methodology for operator learning based on this theoretical work.

2. NONLINEAR PROJECTIONS FOR OPERATOR LEARNING IN BANACH SPACES

Theorem 2.1. *Let X and Y be Banach spaces, let $T : X \rightarrow Y$ be a continuous (possibly nonlinear) map, and let $K \subset X$ be a compact subset. Then, for any choice of $\epsilon > 0$ there exist natural numbers $n, m \in \mathbb{N}$, finite dimensional subspaces $E_n \subset X$ and $E_m \subset Y$, continuous maps $P_n : X \rightarrow E_n$ and $P_m : Y \rightarrow E_m$, and a neural network $f_{n,m} : \mathbb{R}^n \rightarrow \mathbb{R}^m$ such that for every $x \in K$*

$$(1) \quad \|T(x) - \phi_m^{-1} f_{n,m} \phi_n P_n(x)\| < \epsilon,$$

where $\phi_k : E_k \rightarrow \mathbb{R}^k$ indicates an isomorphism between the finite dimensional space E_k and \mathbb{R}^k .

Proof. Since T is continuous, by the Heine-Cantor Theorem it follows that it is uniformly continuous over the compact K . Therefore, we can choose $\delta > 0$ such that $\|T(x_1) - T(x_2)\| < \frac{\epsilon}{3}$ whenever $\|x_1 - x_2\| < \delta$, with $x_1, x_2 \in K$. Corresponding to such choice of $\delta > 0$ we can find a δ -net in K , which we denote by $\{x_i\}_{i=1}^n$. Let E_n be the subspace of X spanned by x_1, \dots, x_n . Using known arguments of Leray-Schauder, we can find a continuous map $P_n : X \rightarrow E_n$ such that for every $x \in K$, we have

$$(2) \quad \|x - P_n(x)\| < \delta.$$

Let us now define the compact $L = TP_n(K)$. We can then find an $\frac{\epsilon}{3}$ -net for L , which we indicate by $\{y_j\}_{j=1}^m$, and set E_m to be the linear span of y_1, \dots, y_m . Applying again the arguments of Leray-Schauder, there exists a continuous map $P_m : X \rightarrow E_m$ satisfying the property that

$$(3) \quad \|x - P_m(x)\| < \frac{\epsilon}{3},$$

for any choice of x in L .

We define an isomorphism $\phi_n : E_n \rightarrow \mathbb{R}^n$ obtained through the identification $\phi_n(x_i) = e_i$, where e_i is the canonical basis vector of \mathbb{R}^n . Since it is defined over finite dimensional spaces, and it is linear, ϕ_n is continuous and has continuous inverse. Similarly, we can identify E_m with \mathbb{R}^m through a map ϕ_m which is defined analogously to ϕ_n .

The map T induces a continuous map $T_{n,m} : E_n \rightarrow E_m$ which is obtained as $T_{n,m} = P_m T|_{E_n}$. Corresponding to such $T_{n,m}$, we introduce $F_{n,m} : \mathbb{R}^n \rightarrow \mathbb{R}^m$ by filling the commutative diagram

$$\begin{array}{ccc} E_n & \xrightarrow{T_{n,m}} & E_m \\ \phi_n^{-1} \uparrow & & \downarrow \phi_m \\ \mathbb{R}^n & \xrightarrow{F_{n,m}} & \mathbb{R}^m \end{array}.$$

The function $F_{n,m}$ is continuous, since it is a composition of continuous functions. As such, using the universal approximation properties of neural networks [4, 5, 14], we can find a neural network $f_{n,m} : \mathbb{R}^n \rightarrow \mathbb{R}^m$ such that

$$(4) \quad \|F_{n,m}(x) - f_{n,m}(x)\| < \frac{\epsilon}{3\|\phi_m^{-1}\|},$$

for all x in the compact $\phi_n P_n(K)$.

Finally, for all $x \in K$ we have

$$\begin{aligned} \|T(x) - (\phi_m^{-1} f_{n,m} \phi_n) P_n(x)\| &\leq \|T(x) - TP_n(x)\| + \|TP_n(x) - P_m TP_n(x)\| \\ &\quad + \|P_m TP_n(x) - (\phi_m^{-1} f_{n,m} \phi_n) P_n(x)\| \\ &< \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} \\ &= \epsilon, \end{aligned}$$

where $\|T(x) - TP_n(x)\| < \frac{\epsilon}{3}$ because of (2), the definition of uniform continuity, and the choice of δ , $\|TP_n(x) - P_m TP_n(x)\| < \frac{\epsilon}{3}$ because of (3), and $\|P_m TP_n(x) - (\phi_m^{-1} f_{n,m} \phi_n) P_n(x)\| < \frac{\epsilon}{3}$ because of the choice of the neural network $f_{n,m}$, i.e. (4), and the fact that $P_m T|_{E_n} = T_{n,m} = (\phi_m^{-1} F_{n,m} \phi_n)$ by construction. This completes the proof. \square

Remark 2.2. The operators P_n and P_m have an explicit and very simple construction, that was used by Leray and Schauder in the proof of their celebrated fixed point theorem. We recall it here, to further elucidate the functioning of the approximation given in Theorem 2.1. As in the proof,

let K be a compact, E_n an n -dimensional space spanned by x_1, \dots, x_n , and $\epsilon > 0$ chosen and fixed. We define $P_n : K \rightarrow E_n$ by the assignment

$$P_n x = \frac{\sum_{i=1}^n \mu_i(x) x_i}{\sum_{i=1}^n \mu_i(x)},$$

where

$$(5) \quad \mu_i(x) = \begin{cases} \epsilon - \|x - x_i\|, & \|x - x_i\| \leq \epsilon \\ 0, & \|x - x_i\| > \epsilon \end{cases}$$

for all $i = 1, \dots, n$.

We will refer to these operators as *Leray-Schauder projections*, following the same convention as [8], although they are not linear.

Corollary 2.3. *With the assumption of Theorem 2.1, the neural network $f_{n,m}$ can be chosen to have a single hidden layer.*

Proof. This fact follows from the proof of Theorem 2.1, with the observation that $f_{n,m}$ needs to approximate the function $F_{n,m}$ on a compact. Then, from the theory of neural network approximation, see [5], we can choose $f_{n,m}$ with a single hidden layer with the necessary properties. \square

The simplicity of Leray-Schauder projections hides a fundamental implementational problem of the theoretical framework of Theorem 2.1. Namely, we do not know how to choose the points used to obtain the spaces E_n for the (nonlinear) projections. In a general Banach space, this might not be a trivial issue. In the rest of this article, we will develop a framework based on orthogonal multivariate polynomials to address this issue in the case of Banach spaces of functions L_μ^p , and the Hilbert space L_μ^2 , with some measure μ .

3. LEARNING LINEAR PROJECTIONS ON BANACH SPACES OF FUNCTIONS

In this section we assume to work on the Banach space $L_\mu^p(S)$ where μ is some fixed measure, and S is a μ -measurable subset of \mathbb{R}^d , which we will assume to be compact throughout. We will also assume that the measure is normalized, i.e. $\mu(S) = 1$, for simplicity since the reasoning can be adjusted to the case where $\mu(S) \neq 1$. A typical example would be the L^p space on $[0, 1]^{\times d}$ with Lebesgue measure μ .

Let $\rho : S \rightarrow \mathbb{R}$ be a μ -integrable function in $L_\mu^q(S)$, where $\frac{1}{p} + \frac{1}{q} = 1$. Then, we say that ρ is a weight function in $L_\mu^p(S)$. Let $\{p_k\}_{k=0}^r$ be a class of polynomials, then we say that they are orthogonal with respect to ρ if

$$\int_S p_k p_\ell \rho d\mu = \begin{cases} 0 & k \neq \ell \\ \neq 0 & k = \ell \end{cases}.$$

If the polynomials p_k are normalized to 1, i.e. $\int_S p_k^2 \rho d\mu = 1$, we will say that they are orthonormal. The weight function ρ , along with a class of orthogonal polynomials $\{p_k\}_{k=0}^r$ defines a projection on the subspace E_r spanned by the polynomials through the functional defined according to the assignment

$$\mathcal{L}(f) = \int_S f(\mathbf{x}) \rho(\mathbf{x}) d\mu.$$

The functional \mathcal{L} defines a quasi-inner product. The projection on E_r is then explicitly given by $P_r(f) = \sum_k \mathcal{L}(f p_k) \frac{p_k}{\mathcal{L}(p_k^2)}$. If the functional is positive (i.e. if ρ has non-negative values), then we

can normalize the polynomials by $\sqrt{\mathcal{L}(p_k^2)}$, and simply define $P_r(f) = \sum_k \mathcal{L}(fp_k)p_k$, where each p_k is normalized to 1. The maps P_n are directly seen to be continuous. More generally, when \mathcal{L} is an arbitrary functional such that

$$\mathcal{L}(p_k p_\ell) = \begin{cases} 0 & k \neq \ell \\ \neq 0 & k = \ell \end{cases},$$

we will still say that p_k are orthogonal (or orthonormal) polynomials with respect to \mathcal{L} , and in this case we can define a projection P_n as before for all n .

Definition 3.1. A *neural projection operator* $\mathfrak{S}_{n,m,r}$ is a quadruple $(F_{n,m}, \rho_1, \rho_2, \{p_k^1\}_{k=0}^{r_1}, \{p_k^2\}_{k=0}^{r_2})$, where $F_{n,m} : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a neural network, $\rho_i : \mathbb{R}^d \rightarrow \mathbb{R}$ are neural network weight functions, and $\{p_k^i\}_{k=0}^{r_i}$ is a set of orthogonal polynomials with respect to ρ_i .

This class of deep learning models consist of a learnable projection over a multivariate orthogonal polynomial basis, along with a neural network that models a projected operator. We want to show now that neural projection operators are universal approximators of continuous (possibly nonlinear) operators in L^p spaces.

The work of Kowalski [6, 7] and Xu [15, 16] has characterized the algebraic properties that the polynomials $\{p_k\}_{k=0}^\infty$ need to satisfy to be an orthogonal basis for the space of polynomials with respect to some functional \mathcal{L} . They have also shown that there are bases of such polynomials. In [6], Kowalski has also given a condition for which the functional \mathcal{L} is continuous on the space of polynomials in $\|\cdot\|_2$ norm. We assume that $\{p_k\}_{k=0}^\infty$ is orthogonal with respect to a functional \mathcal{L} is continuous in $\|\cdot\|_p$ norm. We want to show that given a basis and a continuous functional, we can approximate any continuous operator between L^p spaces with a neural projection operator.

Theorem 3.2. *Let $T : L_\mu^{p_1}(S) \rightarrow L_\mu^{p_2}(S)$ be a continuous operator, and let X be a compact subset of $L_\mu^{p_1}(S)$. Let $\{p_k^i\}_{k=0}^\infty$ and \mathcal{L}^i be as above, for $i = 1, 2$. Then, for any choice of $\epsilon > 0$, we can find a neural projection operator $\mathfrak{S}_{n,m,r}$ such that*

$$(6) \quad \|T(f) - \phi_m^{-1} f_{n,m} \phi_n \hat{P}_n(f)\|_{p_2} < \epsilon,$$

for all $f \in X$, where ϕ_i are isomorphisms as before, and \hat{P}_n is a learned continuous linear map.

Proof. Given the fact that \mathcal{L} is continuous by assumption on the space of polynomials in the $\|\cdot\|_p$ norm, where p is either p_1 or p_2 , using the density of polynomials in $L_\mu^p(S)$, we can extend \mathcal{L} to the whole $L_\mu^p(S)$ continuously. Using the Riesz representation theorem for L^p spaces, we can find a function ρ in $L_\mu^q(S)$, where $\frac{1}{p} + \frac{1}{q} = 1$, such that $\mathcal{L}(f) = \int f \rho d\mu$ for all $f \in L_\mu^p(S)$. Moreover, $\|\mathcal{L}\| = \|\rho\|_q$. For any fixed number of polynomials $p_{i_1}, p_{i_2}, \dots, p_{i_n}$ we have a projection $P_n : L_\mu^p(S) \rightarrow E_n$ defined as

$$P_n(f) = \sum_{k=1}^n \mathcal{L}(fp_{i_k}) \frac{p_{i_k}}{\mathcal{L}(p_{i_k}^2)},$$

where E_n indicates the span of the polynomials. We will call the projections P_n^1 and P_n^2 , corresponding to the polynomials $\{p_i^1\}$ and $\{p_j^2\}$, respectively. Similarly, ρ_i ($i = 1, 2$) will denote the corresponding weight functions. For notational simplicity, $\|\rho_i\|$ will indicate $\|\rho_i\|_{q_i}$, $i = 1, 2$.

Let $\epsilon > 0$ be fixed. For simplicity, we just write $\|\cdot\|$ for the p_1 -norms. We now set $Y = T(X)$, which is compact. We choose an ϵ' -net in Y , where $\epsilon' := \min\{\frac{\epsilon}{9\|\rho_1\|}, \frac{\epsilon}{9}\}$, consisting of polynomials f_1, \dots, f_d , due to compactness of X and density of polynomials in $L_\mu^p(S)$. We can then find ℓ

polynomials $p_{i_1}, \dots, p_{i_\ell}$ of $\{p_k\}_{k=0}^\infty$ such that f_1, \dots, f_d are in the linear span of $p_{i_1}, \dots, p_{i_\ell}$. Let E_n be the span of p_0, \dots, p_n where $n > \max\{i_1, \dots, i_\ell\}$. We let $P_n^1 : L_\mu^{p_1}(S) \rightarrow E_n$ denote the projection discussed above. Let now $M > 0$ be larger than $\|f\|_p$ for all $f \in X$ (since X is bounded such M exists), and define $N = \sum_{k=1}^n \frac{\|p_k\|_\infty \|p_k\|_p}{|\mathcal{L}(p_k^2)|}$. Using the density of polynomials in $L_\mu^{q_1}(S)$, we can find a polynomial ζ_1 such that $\|\zeta_1 - \rho_1\| < \frac{\epsilon}{18MN}$. Using the results on universal approximation of continuous functions by means of neural networks [4, 5, 14], we can find a neural network $\hat{\rho}_1$ such that $\|\hat{\rho}_1 - \rho_1\|_\infty < \frac{\epsilon}{9MN}$ on the compact S . Therefore, we also have that $\|\hat{\rho}_1 - \rho_1\|_{q_1} < \frac{\epsilon}{9MN}$. Let us define the linear map $\hat{P}_n : L_\mu^{p_1}(S) \rightarrow E_n$, defined by $\hat{P}_n(f) = \sum_{k=1}^n \int f p_k \hat{\rho}_1 d\mu \cdot \frac{p_k}{\mathcal{L}(p_k^2)}$.

We want to show that for all f in X , we have $\|\hat{P}_n(f) - f\| < \frac{\epsilon}{3}$. By construction, we can find a polynomial f_r , for some r in $\{1, \dots, n\}$, such that $\|f - f_r\| < \epsilon'$. Then, we have

$$\begin{aligned} \|\hat{P}_n(f) - f\| &\leq \|\hat{P}_n(f) - P_n(f)\| + \|P_n(f) - P_n(f_r)\| + \|P_n(f_r) - f\| \\ &= \|\hat{P}_n(f) - P_n(f)\| + \|P_n(f - f_r)\| + \|f_r - f\| \\ &\leq \sum_{k=1}^n \|f\| \|\hat{\rho}_1 - \rho_1\|_{q_1} \|p_k\|_\infty \frac{\|p_k\|_p}{|\mathcal{L}(p_k^2)|} + \|\rho_1\|_{q_1} \|f - f_r\| + \epsilon' \\ &< \frac{\epsilon}{9} + \frac{\epsilon}{9} + \frac{\epsilon}{9} \\ &= \frac{\epsilon}{3}. \end{aligned}$$

Similarly, one can construct \hat{P}_m^2 which maps onto an m -dimensional space E_m such that $\|\hat{P}_m^2(f) - f\| < \frac{\epsilon}{3}$ whenever f is in Y , where the norm is p_2 -norm. We can therefore now repeat the same construction of the proof of Theorem 2.1 where we use \hat{P}_n^1 and \hat{P}_m^2 instead of the Leray-Schauder (nonlinear) projections. We will not repeat the details, as they are virtually identical to the other proof. This will give us a neural network $f_{n,m}$ such that (6) is satisfied, therefore completing the proof. \square

Remark 3.3. Theorem 3.2 has the fundamental assumption that the functionals \mathcal{L}^i determined by the polynomials $\{p_k^i\}_{k=0}^\infty$ are continuous in the p -norm. We will consider the important case $p = 2$ where proceeding as Kowalski in [6] we can have sufficient conditions for this to happen. In particular, ensuring such conditions in a deep learning algorithm would allow the approximation result of Theorem 3.2 to be applicable. We will introduce such a deep learning methodology in a subsequent work.

4. LEARNING LINEAR PROJECTIONS ON THE HILBERT SPACE

We now consider the particularly important example of the Hilbert space $L^2([-1, 1]^n)$, which is of great importance in applications, including deep learning. We recall the following condition considered by Kowalski in [6].

Hypothesis 4.1 (Kowalski). *For each $k = 0, 1, \dots$ there exist matrices A_k, B_k, C_k such that*

$$\text{rank} A_k = r_n^{k+1},$$

where r_n^{k+1} is the number of degree $k+1$ polynomials in $\{p_i\}$. The recursion formula

$$x\vec{p}_i = A_i\vec{p}_i + B_i\vec{p}_i + C_i\vec{p}_{i-1}$$

holds, where the symbol \vec{p}_i represents vectors consisting of all polynomials in the family $\{p_k\}$ of degree i . For any arbitrary sequence of matrices such that $D_k A_k = \mathbb{1}$ the recursion

$$I_0, \quad I_{j+1} = D_j \text{bp}(I_j C_{j+1}^T),$$

gives positive definite matrices, where bp is the operation defined in [6] that performs block permutation.

We have the following useful result, which is proved using the same approach of Theorem 2 in [6].

Lemma 4.2. *Let $\{p_k\}_{k=0}^\infty$ be a family of polynomials satisfying Hypothesis 4.1. Let $\{q_k\}$ be an orthogonal polynomial complete sequence, and suppose that*

$$(7) \quad q_k = \sum_{j=0}^{\infty} C_{kj} p_j,$$

where $\sum_{k=0}^{\infty} C_{k1}^2 < \infty$. Then, there exists a function $\rho \in L^2([-1, 1]^n)$, and numbers $m_k \neq 0$, such that the functional $\mathcal{L}(f) := \int f \rho d\mu$ is continuous over $L^2([-1, 1]^n)$ and it satisfies $\mathcal{L}(p_i p_k) = \delta_{ik} m_k$.

The following is a special case of Theorem 3.2 when $p = 2$, by applying Lemma 4.2.

Theorem 4.3. *Let $\{p_k\}_{k=0}^\infty$ be a family of polynomials as in Lemma 4.2, let $T : L^2([-1, 1]^n) \rightarrow L^2([-1, 1]^n)$ be a continuous operator, and let $X \subset L^2([-1, 1]^n)$ be compact. Then, for any choice of $\epsilon > 0$, we can find a neural projection operator $\mathfrak{S}_{n,m,r}$ such that*

$$(8) \quad \|T(x) - \phi_m^{-1} f_{n,m} \phi_n \hat{P}_n(x)\|_2 < \epsilon,$$

for all $x \in X$, where ϕ_i are isomorphisms as before, and \hat{P}_n is a learned continuous linear map.

Proof. The main observation is that Lemma 4.2 allows us to use the same proof as in Theorem 3.2 since the functional \mathcal{L} is continuous on $L^2([-1, 1]^n)$. \square

As a particular case of the previous result one can take any complete sequence of orthogonal polynomials, since in this case $C_{kj} = \delta_{kj}$ in Equation (7), and $\sum C_{k1} = C_{11} < \infty$, and where $\rho = 1$, so that $\mathcal{L}(fg) = \int fg d\mu = \langle f, g \rangle$.

5. APPROXIMATIONS FOR FIXED POINTS

We now consider the problem of solving an equation of type

$$(9) \quad T(x) + f = x,$$

where f is a fixed element of the Banach space X , and $T : X \rightarrow X$ is an operator which is possibly nonlinear. The element $x \in X$ satisfying Equation (9) is a fixed point for the operator. Our interest in such a problem stems from the fact that it is possible to frame operator learning tasks in terms of fixed point problems as in [17–19]. Our fundamental question in this section is whether it is possible to project Equation (9) to a finite dimensional space, and if taking the limit $n \rightarrow \infty$ one recovers a solution to Equation (9). If this is the case, then we can define an algorithm based on the framework of this article where we learn an operator in projected form (we also learn the projection) and solve the corresponding equation in the projected space. One would have the guarantee that upon taking a high enough projection dimension n the learned operator would approximate T and the solution would give an approximation to the real fixed point of T in Equation (9).

We use the same framework as in Section 3, and we assume that the function $\rho \in L_\mu^q(S)$ (with $\frac{1}{p} + \frac{1}{q} = 1$) is given with the property that the projection on E_n it induces from $L_\mu^p(S)$ is continuous for all n . We define

$$(10) \quad G_k(f) = \int f p_k \rho d\mu.$$

So, the projection P_n of Section 3 can be written as

$$(11) \quad P_n(g) = \sum_{k=0}^n \int g p_k \rho d\mu \cdot \frac{p_k}{G_k(p_k)} = \sum_{k=0}^n G_k(g) \cdot \frac{p_k}{G_k(p_k)}.$$

Hypothesis 5.1. *We make the following assumptions.*

- (1) *The operator $T : L_\mu^p(S) \rightarrow L_\mu^p(S)$ is completely continuous.*
- (2) *The operator T is Frechet differentiable.*
- (3) *The value 1 is not an eigenvalue of the Frechet derivative of T at 0.*
- (4) *The topological index of T is nonzero.*
- (5) *We have $\sum_{k=0}^{\infty} \frac{\|p_k\|_\infty \|p_k\|_p}{G_k(p_k)} < \infty$.*

Theorem 5.2. *Under the assumptions of Hypothesis 5.1, for any choice of n , the projected equation*

$$(12) \quad T_n(x_n) + f_n = x_n,$$

where $T_n(y) = P_n T(y)$, and $f_n = P_n f$, admits a unique solution x_n^* . Moreover, $x_n^* \rightarrow x^*$ where x^* is a solution to Equation (9).

Proof. Let us consider the function $g \in L_\mu^p(S)$. We have

$$\begin{aligned} \|P_n(g)\|_p &= \left\| \sum_{k=0}^n \int g p_k \rho d\mu \cdot \frac{p_k}{G_k(p_k)} \right\|_p \\ &\leq \sum_{k=0}^n \left| \int g p_k \rho d\mu \right| \cdot \left\| \frac{p_k}{G_k(p_k)} \right\|_p \\ &\leq \sum_{k=0}^n \int |g p_k \rho| d\mu \cdot \frac{\|p_k\|_p}{|G_k(p_k)|} \\ &\leq \sum_{k=0}^n \|g p_k\|_p \|\rho\|_q \frac{\|p_k\|_p}{|G_k(p_k)|} \\ &\leq \sum_{k=0}^n \|g\|_p \|p_k\|_\infty \|\rho\|_q \frac{\|p_k\|_p}{|G_k(p_k)|} \\ &= \|g\|_p \|\rho\|_q \sum_{k=0}^n \|p_k\|_\infty \frac{\|p_k\|_p}{|G_k(p_k)|}. \end{aligned}$$

Therefore, from Hypothesis 5.1 (5) it follows that the projections P_n are uniformly bounded by the finite number $\|\rho\|_q \sum_{k=0}^{\infty} \frac{\|p_k\|_\infty \|p_k\|_p}{G_k(p_k)}$. The assumptions in Hypothesis 5.1 now guarantee that the framework of [8] on Galerkin's Method is applicable, and as a result we obtain that the projected equation has a unique solution for all n , and these solutions converge to a solution of Equation (9). \square

The rate of convergence of the approximated solutions to the solution of the original equation can be studied using the methods of [1]. There is an important case where we can apply Theorem 5.2.

This is the case also considered in Section 4 where $p = 2$ (i.e. we have the Hilbert space), and we use an orthonormal basis p_k . Here $\rho = 1$ is just the identity function, G_k is also the identity since the projection maps over normalized elements. In this case it is known that $\{P_n\}$ is uniformly bounded.

6. FUTURE PERSPECTIVES

We now conclude with a few remarks on future work based on the results of this paper. More specifically, we describe the algorithmic perspectives that follow from our present results. The approach concerns learning operators between L^p spaces in a suitable projected space. We want to both learn projections P_n and P_m (possibly coinciding) via learning a basis of polynomials in order to apply Theorem 3.2. To do so, we need the polynomials to be orthogonal in the sense of Section 4. This can be done using the algebraic characterization of Kowalski [6, 7], and Xu [15, 16]. These approaches generalize the well known one-dimensional case of recursion formulas of the Favard's Theorem. Therefore, one can recursively construct a family of orthogonal polynomials.

In addition, we need to learn a mapping between the projected spaces that approximates a given operator. Our objective is to learn this operator in such a way that its solutions approximate the solutions of a projected operator equation as in Equation 9, which is a fixed point problem. A similar type of approach for specific projections on Chebyshev polynomials (spectral methods) and integral nonlinear operators has been pursued in [19]. The approach described in this article is much more general, and we expect that it is much more widely applicable in practice.

The results obtained in this article show that the methodology described produces a universal approximator under mild additional assumptions on the continuity of the corresponding functional. Moreover, as discussed in Section 4, in the fundamental case of $p = 2$ there are some direct conditions that can be imposed guaranteeing that the continuity of the functional holds, therefore giving a direct method for obtaining universal approximators. Additionally, Section 5 shows that upon increasing the dimension of the projections the solutions obtained in the projected spaces converge to the solutions of the operator equation that is being modeled. Therefore, the model has some good convergence properties under the framework of Hypothesis 5.1.

Guaranteeing that the needed assumptions are satisfied during the learning process is an interesting computational problem, and leveraging the theoretical framework described in this article in practice is of its own interest in machine learning. As such, we will pursue the algorithmic implementation of this work in a separate article, where we also showcase concrete examples of these methods in data science applications.

REFERENCES

- [1] Kendall E Atkinson and Florian A Potra, *Projection and iterated projection methods for nonlinear integral equations*, SIAM journal on numerical analysis **24** (1987), no. 6, 1352–1373.
- [2] Tianping Chen and Hong Chen, *Universal approximation to nonlinear operators by neural networks with arbitrary activation functions and its application to dynamical systems*, IEEE transactions on neural networks **6** (1995), no. 4, 911–917.
- [3] Clive AJ Fletcher, *Computational galerkin methods*, Springer, 1984.
- [4] Ken-Ichi Funahashi, *On the approximate realization of continuous mappings by neural networks*, Neural networks **2** (1989), no. 3, 183–192.
- [5] Kurt Hornik, Maxwell Stinchcombe, and Halbert White, *Multilayer feedforward networks are universal approximators*, Neural networks **2** (1989), no. 5, 359–366.
- [6] MA Kowalski, *Orthogonality and recursion formulas for polynomials in n variables*, SIAM Journal on Mathematical Analysis **13** (1982), no. 2, 316–323.

- [7] ———, *The recursion formulas for orthogonal polynomials in n variables*, SIAM Journal on Mathematical Analysis **13** (1982), no. 2, 309–315.
- [8] Yu P Krasnosel'skii, *Topological methods in the theory of nonlinear integral equations*, Pergamon Press (1964).
- [9] Mark Aleksandrovich Krasnosel'skii, Gennadi M Vainikko, RP Zabreyko, Ya B Ruticki, and V Va Stet'senko, *Approximate solution of operator equations*, Springer Science & Business Media, 2012.
- [10] MA Krasnosel'skii and PP Zabreiko, *Geometrical Methods of Nonlinear Analysis*, Springer-Verlag, Berlin, 1984, MR 85b **47057**.
- [11] Moshe Leshno and Allan Pinkus, *Multilayer feedforward networks with a nonpolynomial activation function can approximate any function*, Neural networks **6** (1993), no. 6, 861–867.
- [12] Lu Lu, Pengzhan Jin, Guofei Pang, Zhongqiang Zhang, and George Em Karniadakis, *Learning nonlinear operators via DeepONet based on the universal approximation theorem of operators*, Nature machine intelligence **3** (2021), no. 3, 218–229.
- [13] Zhou Lu, Hongming Pu, Feicheng Wang, Zhiqiang Hu, and Liwei Wang, *The expressive power of neural networks: A view from the width*, Advances in neural information processing systems **30** (2017).
- [14] Allan Pinkus, *Approximation theory of the MLP model in neural networks*, Acta numerica **8** (1999), 143–195.
- [15] Yuan Xu, *On multivariate orthogonal polynomials*, SIAM journal on mathematical analysis **24** (1993), no. 3, 783–794.
- [16] ———, *Multivariate orthogonal polynomials and operator theory*, Transactions of the American Mathematical Society **343** (1994), no. 1, 193–202.
- [17] Emanuele Zappala, Antonio Henrique de Oliveira Fonseca, Josue Ortega Caro, and David van Dijk, *Neural Integral Equations*, arXiv:2209.15190 (2022).
- [18] Emanuele Zappala, Antonio H de O Fonseca, Andrew H Moberly, Michael J Higley, Chadi Abdallah, Jessica A Cardin, and David van Dijk, *Neural integro-differential equations*, Proceedings of the AAAI Conference on Artificial Intelligence, 2023, pp. 11104–11112.
- [19] Emanuele Zappala, *Spectral methods for Neural Integral Equations*, arXiv preprint arXiv:2312.05654 (2023).

DEPARTMENT OF MATHEMATICS AND STATISTICS, IDAHO STATE UNIVERSITY, PHYSICAL SCIENCE COMPLEX —
921 S. 8TH AVE., STOP 8085 — POCA TELLO, ID 83209

Email address: emanuelezappala@isu.edu