

# Harnessing spontaneous emission of correlated photon pairs from ladder-type giant atoms

Zhao-Min Gao,<sup>1</sup> Jia-Qi Li,<sup>1</sup> Ying-Huan Wu,<sup>1</sup> Wen-Xiao Liu,<sup>1,2</sup> and Xin Wang<sup>1,\*</sup>

<sup>1</sup>*Institute of Theoretical Physics, School of Physics,  
Xi'an Jiaotong University, Xi'an 710049, People's Republic of China*

<sup>2</sup>*Department of Electronic Engineering, North China University of Water  
Resources and Electric Power, Zhengzhou 450046, People's Republic of China*

(Dated: June 19, 2024)

The realization of correlated multi-photon processes usually depends on the interaction between nonlinear media and atoms. However, the nonlinearity of optical materials is generally weak, making it still very challenging to achieve correlated multi-photon dynamics at the few-photon level. Meanwhile, giant atoms, with their capability for multi-point coupling, which is a novel paradigm in quantum optics, mostly focus on the single photon field. In this work, using the method described in *Phys. Rev. Res.* **6**, 013279 (2024), we reveal that the ladder-type three-level giant atom spontaneously emits strongly correlated photon pairs with high efficiency by designing and optimizing the target function. In addition, by encoding local phases into the optimal coupling sequence, directional two-photon correlated transfer can be achieved. This method does not require a nonlinear waveguide and can be realized in the conventional environment. We show that the photon pairs emitted in both the bidirectional and the chiral case exhibit strong correlation properties in both time and space. Such correlated photon pairs have great potential applications for quantum information processing. For example, numerical results show that our proposal can realize the two-photon mediated cascaded quantum system.

## I. INTRODUCTION

Giant atoms, which are different from point-like small atoms, can be comparable to or much larger than the wavelength of the interacting field in size [1–7]. Therefore, the dipole approximation becomes invalid, enabling giant atoms to achieve multi-point coupling with waveguides [8–14]. At present, most studies on giant atoms focus on the field of single-photon processes, such as frequency-dependent couplings [15–20] and chiral quantum optics [20–28]. These studies have led to significant advancements in controlling light-matter interactions at the single-photon level. However, only a few studies have delved into the multi-photon dynamics of giant atoms [29, 30], for which nonlinear media are required.

Meanwhile, multi-photon dynamics are crucial for building long-range, high-fidelity quantum information and quantum communication networks [31–38], typically implemented in nonlinear media [39–48]. Multi-photon processes have made significant progress in the fields of quantum entanglement dynamics, quantum interferometry, dissipative quantum dynamics and optical parameter processes *et al.* [49–59]. However, there are still great challenges in the field of multi-photon problems, such as photon loss, nonlinear effects and quantum noise, which limit the stable transmission of multi-photon states [60–67]. Moreover, the weak nonlinearity of the optical materials imposes great limitations on the multi-photon dynamics.

In recent years, the exploration of quantum optics within the structured environment of giant atoms has been proposed [68]. Unlike previous studies of giant atoms which only considered a few coupling points, this study proposes to use the coupling of multiple points  $x_1, \dots, x_N$  with interaction strength  $g(x_i)$ , where the distances between different coupling points are unequal. The optimal coupling sequence is obtained by using the sequential quadratic programming algorithm to optimize the objective function of the interaction between the giant atom and the conventional waveguide in momentum space, and by using an inverse Fourier transform to obtain the coupling sequence in real space. With this method, exotic quantum phenomena such as simulated band-gap environment and broadband chiral emission are realized. Here, we extend this optimization method to multi-level giant atom. We find that the multi-photon processes which are correlated in both time and space can be realized in conventional linear waveguides without requiring any nonlinear media. Moreover, we can achieve high-fidelity quantum state transfer between two three-level giant atoms under this scheme.

In this paper, we propose to use a ladder-type three-level giant atomic platform to achieve correlated two-photon processes in a linear waveguide. Firstly, we consider the bidirectional propagation of photons. By calculating the spontaneous decay process numerically and analytically, we prove that the three-level giant atom emits two correlated photons simultaneously, rather than in cascades. In addition, we obtain the optimal coupling sequence by designing the objective function and optimizing it. Using this coupling sequence, we explore the spatial correlation properties of the two photons. Secondly, we also focus on the case where the

---

\* wangxin.phy@xjtu.edu.cn

correlated two photons can only propagate in the same direction. Finally, the two-photon mediated cascaded quantum system based on our method is proposed.

## II. BIDIRECTIONAL TWO-PHOTON EMISSION

### A. Model and Equations

As shown in Fig. 1(a), we consider a ladder-type three-level giant atom with ground state  $|g\rangle$  and first (second) excited states  $|e\rangle$  ( $|f\rangle$ ), coupled to a one-dimensional conventional waveguide. The three-level giant artificial atom can be easily implemented with superconducting qubit circuits [12, 69, 70]. As shown in Fig. 1(b) the energy of the first (second) excited state is denoted as  $\omega_{eg}$  ( $\omega_{fg}$ ), respectively. The Hamiltonian of the system is (setting  $\hbar=1$ )

$$H = \omega_{fg}\sigma_{fg}^+\sigma_{fe}^- + \omega_{eg}\sigma_{eg}^+\sigma_{eg}^- + \sum_k \omega_k a_k^\dagger a_k + \sum_k g_k a_k^\dagger \sigma_{fe}^- + \sum_{k,k'} g_{k'} a_{k'}^\dagger \sigma_{eg}^- + \text{H.c.}, \quad (1)$$

where  $a_k$  ( $a_k^\dagger$ ) is photon annihilation (creation) operator, with  $k$  being the wave vector in momentum space, and  $\sigma_{fe}^+ = |f\rangle\langle e|$  ( $\sigma_{eg}^+ = |e\rangle\langle g|$ ) denotes the atomic transition operator. We consider the giant atom coupling to the waveguide at multiple points  $x_1, \dots, x_N$  with distinct interaction strength  $g(x_i)$ . The interaction strength in momentum space is expressed as  $g_k = \sum_{x_i} g(x_i) e^{-ikx_i}$  [69–71]. In our model, the waveguide is considered to be conventional without periodical structure, with linear dispersion  $\omega_k = c|k|$ , and  $c$  being the group velocity.

In the rotating-wave approximation, the total Hamiltonian of the system is

$$H_I = \sum_k (\omega_k - \omega_{eg}) a_k^\dagger a_k + \sum_k g_k a_k^\dagger e^{2i\Delta t} \sigma_{fe}^- + \sum_{k,k'} g_{k'} a_{k'}^\dagger \sigma_{eg}^- + \text{H.c.}, \quad (2)$$

where  $\Delta = \omega_{eg} - \omega_0 = \omega_0 - \omega_{fe}$  [see Fig. 1(b)]. In the two-excitation subspace, the state of the system can be expressed as

$$|\psi(t)\rangle = c_f(t)|f,0\rangle + c_{e,k}(t)|e,k\rangle + c_{g,k,k'}(t)|g,k,k'\rangle. \quad (3)$$

Here,  $c_{g,k,k'}(t)$  denotes the probability amplitude of the giant atom being in the ground state, while simultaneously exciting two photons with wave vector  $k$  and  $k'$  in the waveguide. Additionally,  $c_{e,k}(t)$  ( $c_f(t)$ ) is the probability amplitude of the intermediate state (the second excited state).

For small atoms, the interaction between the emitter and waveguide is point-like coupling, the coupling

strength can be considered as a constant. Under these conditions, it becomes challenging to observe peculiar phenomena in quantum optics when small emitters interact with conventional linear waveguide. Therefore, we propose employing a three-level giant atom to realize two-photon processes.

Initially, the three-level atom is considered to be in the second excited state, while the waveguide is in its vacuum state, i.e.,  $|\psi(t=0)\rangle = |f,0\rangle$ . According to Schrödinger equation, we derive the following coupling equations

$$\begin{cases} i\dot{c}_f(t) = \sum_k g_k^* e^{-2i\Delta t} c_{e,k}(t), & (4a) \\ i\dot{c}_{e,k}(t) = (\omega_k - \omega_{eg}) c_{e,k}(t) \\ + g_k e^{2i\Delta t} c_f(t) + \sum_{k'} g_{k'}^* c_{g,k,k'}(t), & (4b) \\ i\dot{c}_{g,k,k'}(t) = (\omega_k + \omega_{k'} - 2\omega_{eg}) c_{g,k,k'}(t) \\ + g_{k'} c_{e,k}(t). & (4c) \end{cases}$$

Assuming the objective coupling function in momentum space following the form of window function, specifically confined to the range  $[\pm k_0 - \delta_w/c, \pm k_0 + \delta_w/c]$ , with  $c|k_0| = \omega_{fg}/2$ , and a window function width of  $2\delta_w$ . In this work, we take  $k_0$  and  $\omega_0 = c|k_0|$  as units of wave vector and frequency, respectively. To avoid the emission of a single photon, the coupling strength is zero at frequencies  $\omega_{fe}$  and  $\omega_{eg}$ , and is identical near  $\omega_0$ , resulting in no mode in the waveguide resonating with the two transition frequencies of the giant atom. On the contrary, the photons will be significantly excited via a second-order process around  $\omega_0$ . The process of releasing two photons from the three-level giant atom is no longer cascaded, but it is possible to result in the simultaneous production of two correlated photons. Therefore, we calculate the evolution by using adiabatic elimination, i.e.,  $\dot{c}_{e,k}(t) = 0$ . We obtain

$$c_{e,k}(t) = -\frac{g_k e^{2i\Delta t} c_f(t) + \sum_{k'} g_{k'}^* c_{g,k,k'}(t)}{\omega_k - \omega_{eg}}. \quad (5)$$

After substituting Eq. (5) into Eq. (4b,4c), we can calculate  $c_{g,k,k'}(t)$  as

$$c_{g,k,k'}(t) = i \frac{g_{k'} g_k}{\omega_k - \omega_{eg}} \int_0^t e^{-i\delta'(t-t')} e^{2i\Delta t} c_f(t') dt', \quad (6)$$

with

$$\delta' = (\omega_k + \omega_{k'} - 2\omega_{eg}) - \frac{g_{k'}^2}{\omega_k - \omega_{eg}}. \quad (7)$$

The Stark shift of the state  $|f,0\rangle$  is derived as

$$\delta\omega_f = \sum_k \frac{g_k^2}{\omega_k - \omega_{eg}}. \quad (8)$$

Therefore, the probability amplitude  $c_f(t)$  is rewritten as

$$\begin{aligned} \dot{c}_f(t) &= i\delta\omega_f c_f(t) \\ &- \sum_{k,k'} \frac{g_k^2 g_{k'}^2}{(\omega_k - \omega_{eg})^2} \int_0^t e^{-i(\delta'+2\Delta)t} c_f(t') dt'. \end{aligned} \quad (9)$$




FIG. 1. (a) Sketch of the bidirectional model: A three-level giant atom couples to the linear conventional waveguide at positions  $x_1, \dots, x_N$ , which photons can travel in both directions. (b) Energy level diagram of the ladder-type three-level giant atom. (c) The dispersion relation of the linear waveguide,  $\omega_k = c|k|$ .




FIG. 2. (a), (b) The numerical and analytical results of the spontaneous emission when the objective function takes the form of window function, setting  $\delta_w/\omega_0 = 0.033$  with  $\Delta/\omega_0 =$  (a) 0.067, (b) 0.20. (c) The spontaneous decay rate changes with the detuning between the intermediate state frequency  $\omega_{eg}$  and  $\omega_0$ , with  $\delta_w/\omega_0 = 0.033$ . (d) The spontaneous decay rate changes with the width of the window function, with  $\Delta/\omega_0 = 0.33$ .

By defining

$$\tilde{c}_f(t) \approx c_f(t) e^{-i \sum_k \frac{g_k^2}{\omega_k - \omega_{eg}} t}, \quad (10)$$

we can renormalize energy level of  $|f\rangle$  with dynamical Stark shift as

$$\dot{\tilde{c}}_f(t) = - \sum_{k,k'} \frac{g_k^2 g_{k'}^2}{(\omega_k - \omega_{eg})^2} \int_0^t e^{-i \delta_{k,k'}(t-t')} \tilde{c}_f(t') dt', \quad (11)$$

where  $\delta_{k,k'} = \omega_k + \omega_{k'} - \omega_{fg}$  is the two-photon detuning. The modes satisfying  $\omega_k + \omega_{k'} = \omega_{fg}$  have great contribution to the dynamics of the two-photon process (cf. Eq. (11)). By employing the Wigner-Weisskopf theory [72], the summation over  $k$  is transformed into an integral, i.e.,  $\sum_k \rightarrow \frac{L}{2\pi} \int_{-\pi}^{\pi} dk$ , with  $L \rightarrow \infty$  being the waveguide length, and Eq. (11) is written as

$$\dot{\tilde{c}}_f(t) = - \left( \frac{L}{2\pi} \right)^2 \int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{g_k^2 g_{k'}^2}{(\omega_k - \omega_{eg})^2} e^{-i \delta_{k,k'}(t-t')} \tilde{c}_f(t') dk dk' dt'. \quad (12)$$

Given that we consider a linear waveguide, where  $\omega_k =$

$c|k|$ , and obtain

$$\dot{\tilde{c}}_f(t) = - \left( \frac{L}{2\pi} \right)^2 \frac{1}{c^2} \int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{g_k^2 g_{k'}^2}{(\omega_k - \omega_{eg})^2} e^{-i \delta_{k,k'}(t-t')} \tilde{c}_f(t') d\omega_k d\omega_{k'} dt'. \quad (13)$$

To calculate the spontaneous decay rate, we perform a variable substitution for  $\omega_k$  and  $\omega_{k'}$ , i.e.,

$$\begin{cases} \omega_+ = \omega_k + \omega_{k'}, \\ \omega_- = \omega_k - \omega_{k'}. \end{cases} \quad (14)$$

We calculate the frequency-dependent component in Eq. (13) and obtain

$$\begin{aligned} & \int_0^{\infty} \int_0^{\infty} \frac{1}{(\omega_k - \omega_{eg})^2} e^{-i \delta_{k,k'}(t-t')} d\omega_k d\omega_{k'} \\ &= 2 \int_0^{\infty} \frac{1}{\omega_+ + \omega_- - 2\omega_{eg}} \Big|_{-2\delta_w}^{2\delta_w} e^{-i(\omega_+ - \omega_f)(t-t')} d\omega_+. \end{aligned} \quad (15)$$

We assume that the coupling strength is much smaller than the bandwidth around  $\omega_0$ . Moreover, since the atomic spontaneous radiation is centered at the

transition frequency, we can employ the integral

$$\int_0^\infty e^{-i(\omega_+ - \omega_f)(t-t')} d\omega_+ \simeq 2\pi\delta(t-t'),$$

and obtain

$$\dot{c}_f(t) = \frac{L^2 |g_{k_0}|^4 \delta_w}{2\pi c^2} \frac{\tilde{c}_f(t)}{\Delta^2 - \delta_w^2}. \quad (16)$$

Therefore, the exponential decay is calculated as

$$c_f(t) = e^{-\frac{\Gamma}{2}t}, \quad \Gamma = \frac{L^2 |g_{k_0}|^4}{\pi c^2} \frac{\delta_w}{\Delta^2 - \delta_w^2}, \quad (17)$$

with  $\Gamma$  representing the spontaneous decay rate of emitting two correlated photons, note that this rate depends on the width of the window function and  $\Delta$ . As depicted in Fig. 2(a,b), we numerically and analytically show the evolution of the three-level giant atom for different values  $\delta_w$  and  $\Delta$ , observing discrepancies between numerical and analytical results. To determine appropriate values, we explore how the agreement between numerical and analytical results varies with changes in  $\delta_w$  and  $\Delta$  [see Fig. 2(c,d)]. We find that the agreement improves as the window function narrows or as  $\omega_e$  moves further from  $\omega_0$ .

### B. Optimization for bidirectional case




FIG. 3. The optimizing coupling function in momentum space for bidirectional correlated two-photon decay. The dashed red line represents the objective function, and the solid blue line is the optimized coupling function. The real-space coupling sequence  $\{x_i, g(x_i)\}$  is listed in Table. I [see Appendix. B]. We set  $|k_2|/k_0 = 1.067$  ( $|k_1|/k_0 = 0.933$ ) and  $N = 50$ .

In this section, we present the method to realize the optimization of target coupling for giant atoms. We assume that the coupling strength of the right ( $k > 0$ ) and left directions ( $k < 0$ ) are identical, i.e.,




FIG. 4. (a) The evolution of probability amplitudes of the three energy states  $|f\rangle$ ,  $|e\rangle$  and  $|g\rangle$  of the giant atom with time. We set the parameter as  $\omega_e/\omega_0 = 1.167$  and  $g_0 = 0.015$ . (b) The probability of the giant atom's transition to the intermediate state varies with  $\Delta$ , with  $\delta_w/\omega_0 = 0.067$ .

$g_{-k} = g_k^*$ , indicating that the two-photon field propagates bidirectionally in the waveguide [68]. By designing an objective function in the form of a window function in momentum space, the coupling strength is required to be exactly zero at the atomic transition frequencies between the three energy levels, as shown in Fig. 1(b), and maximum at  $\omega_0 = \omega_f/2$ , i.e.,

$$g_k^T/g_0 = \begin{cases} 0, & \{\omega_{fe} - c\delta_w < c|k| < \omega_{fe} + c\delta_w\}, \\ 1, & \{\omega_0 - c\delta_w < c|k| < \omega_0 + c\delta_w\}, \\ 0, & \{\omega_{eg} - c\delta_w < c|k| < \omega_{eg} + c\delta_w\}. \end{cases} \quad (18)$$

To obtain the coupling sequence in real space, we define the constraint condition as

$$C_m = \int_{-k_{\max}}^{k_{\max}} dk ||g_k| - |g_k^T|| w(k), \quad (19)$$

which represents the difference between the optimized coupling function  $g_k$  and the objective coupling function  $g_k^T$ , with  $w(k)$  being the weight function which describes the similarity between the optimized and objective functions in different regimes. In our work, to achieve correlated two-photon emission, we focus on the modes around  $c|k| = \omega_0$  and  $c|k| = \omega_{fe}$  ( $\omega_{eg}$ ). Due to




FIG. 5. (a) The two-photon field distributions in real-space at  $\Delta t = 100$ . (b) The two-photon field distribution in momentum space. (c) The second-order correlation function of two photons in the case of bidirectional propagation. We set  $g_0 = 0.02$ , other parameters are the same with those in Fig. 4.

the large detuning effect, the modes far away from these frequencies contribute little to the dynamics of the system. Therefore, we assign much greater weights to these three regions mentioned in Eq. (18) compared to others.

The coupling mechanism of linear quantum electrodynamics (QED) elements requires that  $g(x_i)$  are of the same sign. Employing the method described in Ref. [68], we optimize the objective function with a sequential quadratic programming algorithm, obtaining the optical coupling sequence  $\{x_i/x_0, A(x_i)\}$  in real space, as listed in Table I [see Appendix. B]. When  $\delta_w$  is large, the numerical results agree well with the analytical results. However, when the width of the window function is narrow, the required number of coupling points  $N$  in the optimization process will be large, which is very demanding experimentally. Therefore, we set  $|k_2|/k_0 = 1.067$ , and the coupling point number is  $N = 50$  in this work.

Furthermore, the optimal coupling function in momentum space is obtained by Fourier transformation  $g_k = \sum_{x_i} g(x_i) e^{-ikx_i}$ , which is illustrated in Fig. 3. For the modes far away from the atomic transition frequency  $\omega_0$ , the coupling strength  $g_k$  remains almost constant with respect to  $k$ . Due to  $g(x_i)$  having the same sign, the coupling around  $k \approx 0$  (the dc component) is strong. To achieve correlated two-photon emission, we set the atomic transition frequency  $\omega_0$  far away from  $c|k| = 0$ , making the influence of these low-frequency modes on the system dynamics negligible due to the large detuning.

### C. Super-correlated photon pairs

Based on the theoretical analysis in Sec. II A, when adopting the coupling function depicted in Fig. 3, the correlated radiation of two photons can be achieved. To explore this two-photon process, we plot the evolution of the three-level giant atom shown in Fig. 4(a). We

find that the probability amplitudes of the giant atom being in the state  $|e\rangle$  is extremely low, indicating that the  $|e\rangle$  is only an intermediate state during the correlated decay process. In Fig. 4(b), the population in the intermediate state decreases as  $\omega_e$  moves further from  $\omega_0$ . This reduction in state  $|e\rangle$  population contributes to the improved agreement between numerical and analytical results with increasing  $\omega_e$ . The evolution of the state  $|f\rangle$  approximately follows the form of exponential decay, demonstrating that the three-level atom decays from the state  $|f\rangle$  to the ground state  $|g\rangle$  by simultaneously releasing two photons into the waveguide. Therefore, the two photons are strongly correlated in space, and their second-order correlation function exhibits strong bunching.

We explore the wavefunction of two photons in both real and momentum space, focusing on the direction of their propagation. The atomic wavefunction, which describes the momentum distribution of two photons in  $k$  space, is written as

$$\psi(k, k') = a_k^\dagger a_{k'}^\dagger |\psi(t=0)\rangle, \quad (20)$$

with the initial state being  $|f, 0\rangle$ . According to the numerical method in the Appendix. A, we derive the wavefunction in  $k$ -space.

The real space wavefunction for two correlated photons,  $|\psi(x, y)\rangle$ , is obtained via Fourier transform, which represents the probability of detecting two photons simultaneously at position  $(x, y)$ , that is

$$|\psi(x, y)\rangle = a_k^\dagger a_{k'}^\dagger e^{ikx} e^{ik'y} |f, 0\rangle. \quad (21)$$

where  $a_{k(k')}^\dagger$  is the operator that creates a photon within momentum in the waveguide.

The distributions of the two photons in both momentum and real space are shown in Fig. 5(a,b). In momentum space, the energy of the two emitted photons is conserved, matching the energy of the three-level atom. For bidirectional propagation, the condition






FIG. 6. (a) Sketch of the chiral model: A three-level giant atom couples to the linear conventional waveguide, which photons can only travel in the right direction. (b) Energy level diagram of the ladder-type three-level giant atom.

$c|k_x| + c|k_y| = \omega_{fg}$  is satisfied in both right and left  $k$  directions, which indicates that there is no energy loss during the propagation of two photons. In real space, we can clearly observe the bidirectional propagation of the two photons in the waveguide. We take  $x_0$  ( $y_0$ ) =  $2\pi/k_0$  as the units of the space length.

The second-order correlation function is defined as

$$g^{(2)}(x, y) = \frac{\langle a_x^\dagger a_y^\dagger a_y a_x \rangle}{\langle a_x^\dagger a_x \rangle \langle a_y^\dagger a_y \rangle}, \quad (22)$$

with  $r = x - y$  signifies the spatial separation between the two photons, which are plotted in Fig. 5(c) for the field evolution at  $\Delta t = 100$ . Notably, the strongest correlation peaks occur both at  $r = 2c\Delta t$  and  $r = 0$ , which shows that the two photons can propagate either in the same direction or in opposite directions. In conclusion, in terms of spatial correlation, our scheme can realize bidirectional propagation of correlated two-photons.




FIG. 7. The optimizing coupling sequence in  $k$ -space for chiral case. The dashed blue line represents the objective function, and the solid orange line is the optimized coupling function. The real-space coupling sequence  $\{x_i, A(x_i), \theta(x_i)\}$  is listed in Table II [see Appendix B]. The blue region represents the discrepancy between the objective function and the optimized result.

### III. DIRECTIONAL TWO-PHOTON CHIRAL EMISSION

In our bidirectional propagation case study, we assume that the giant atom interacts with the waveguide via linear conventional elements. Therefore, there is no phase difference between the coupling strength of each coupling point. However, if we consider coupling with nonlinear QED elements, a local phase  $\theta(x_i)$  will be encoded at  $x_i$ , then the coupling strength is  $g(x_i) = A(x_i) e^{i\theta(x_i)}$  [28, 68]. The coupling strength in  $k$ -space becomes asymmetric, i.e.,  $|g_k| \neq |g_{-k}|$ .

The objective function in the form of window function is designed as

$$g_k^T = \begin{cases} 1, & \{\omega_0 - \delta_w < c|k| < \omega_0 + \delta_w\}, \\ 0, & \text{others.} \end{cases} \quad (23)$$

According to the same coupling method discussed in Sec. II B, we obtain the optimized coupling sequence  $\{x_i/x_0, A(x_i), \theta(x_i)\}$  in real space with  $N = 50$ . The corresponding data of the real space coupling sequence under the chiral case are presented in Table II [see Appendix B].

By employing the Fourier transform, we illustrate the optimized coupling function in momentum space in Fig. 7. For modes around  $\omega_0/c$ , the coupling strength  $g_k$  varies slightly with  $k$ , but it approaches zero at  $\omega_{fe}/c$  ( $\omega_{eg}/c$ ) and  $\omega_0/c$  in the left direction. Similarly, the effect of the strong coupling between the emitter and the waveguide at  $k \approx 0$  on our dynamics can be considered negligible.

Under these conditions, the giant atom only interacts




FIG. 8. (a) The second-order correlation function of two photons in real space at  $\Delta t = 100$ . (b) The two-photon field distribution in real space. (c) The two-photon field distribution in momentum space. Other parameters are the same with those in Fig. 5.

with the modes which propagates to the right direction, enabling the observation of correlated two-photon emission, as shown in Fig. 6(a). Initially, a virtual process occurs where the giant emitter transitions from the  $|f\rangle$  state to the  $|e\rangle$  state, releasing a photon with momentum around  $+k_0$  which propagates to the right direction. Due to the non-conservation of energy, the atom keeps on rapidly decaying from the intermediate state  $|e\rangle$  to the ground state  $|g\rangle$  by emitting an identical photon  $+k_0$ , as shown in Fig. 6(b). We find that these two photons are released simultaneously and propagate to the same direction, leading to a strong spatial correlation in one direction.

As depicted in Fig. 8(a), the two-photon correlation peaks exclusively at  $r = 0$ , indicating that the two photons only propagate in the same direction. To further investigate the propagation direction of the photons, we focus on the two-photon distributions in real space in Fig. 8(b). The results reveals a distribution predominantly in the right direction, thus confirming the right emission. Furthermore, in momentum space, we observe the satisfaction of energy conservation  $ck_x + ck_y = \omega_{fg}$ , which is depicted in Fig. 8(c). Therefore, we can achieve directional correlated two-photon emission without energy loss. The same approach can be taken for photon pairs that achieve propagation in the left direction.

#### IV. TWO-PHOTON MEDIATED CASCADED QUANTUM SYSTEM

In this section, based on the directional two-photon chiral emission discussed above, we consider the two-photon mediated cascade quantum network for three-level giant atoms. We take the simplest setups where two giant atoms are coupled to a conventional linear waveguide for example (cf. Fig. 9). The coupling centers of two giant atoms are  $x_{A,B}$ . Now we derive the cascaded master equation for this two-photon mediated system.




FIG. 9. Diagram of a two-photon mediated cascaded quantum system, illustrating the emission and rightward propagation of two photons between two distant three-level atoms.

Given that two giant atoms are identical and two photons are emitted simultaneously, the interaction Hamiltonian for Eq. (6, 9) is given by

$$H_{\text{int}} = \sum_{i=A,B} \sum_{k,k'} g_{i,kk'} e^{i\delta_{k,k'}t} \sigma_i^+ a_k a_{k'} + \text{H.c.}, \quad (24)$$

$$g_{i,kk'} = \frac{g_{ik} g_{ik'}}{\omega_k - \omega_{eg}}, \quad g_{Bk} = g_{Ak} e^{ikd_s}, \quad (25)$$

where  $g_{k,k'}$  represents the two-photon coupling strength, and  $\sigma^+ = |f\rangle\langle g|$  denotes the atom transition operator. Here,  $d_s$  is the separation distance between two giant atoms. Due to the state  $|e\rangle$  being adiabatically eliminated, the giant atoms are treated as two-level qubits which decay to their ground state by emitting two correlated photons. In this scenario, two giant atoms are unidirectionally mediated by correlated photon pairs. The cascaded master equation governing the system's evolution can be derived as [72, 73]

$$\begin{aligned} \dot{\rho}_s(t) = & -i\text{Tr}_R [H_{\text{int}}, \rho(t)] \\ & - \text{Tr}_R \int_{t_0}^t [H_{\text{int}}(t), [H_{\text{int}}(t'), \rho(t')]] dt', \end{aligned} \quad (26)$$

with  $\rho$  ( $\rho_s$ ) being the density matrix operator of the system (two emitters). By substitution Eq. (25) into

Eq. (26), we obtain [74]

$$\dot{\rho}_s(t) = - \sum_{i,j} \int_{t_0}^t dt' A(x_i, x_j; t, t') (\sigma_i^+ \sigma_j^- \rho_s(t') - \sigma_j^- \rho_s(t') \sigma_i^+) + \text{H.c.}, \quad (27)$$

with  $A(x_i, x_j; t, t')$  being the spatiotemporal correlation function for the propagating field. According to Eq. (13), it is defined as

$$A(x_i, x_j; t, t') = \sum_{k,k'} g_{i,kk'} g_{j,kk'}^* e^{i\delta_{k,k'}(t-t')} \approx \frac{L^2 |g_{k_0}|^4}{4(2\pi)^2 c^2} \frac{\delta_w}{\Delta^2 - \delta_w^2} 2\pi \delta\left(t - t' - \frac{x_{ij}}{c}\right), \quad (28)$$

where  $x_{ij} = x_i - x_j$ . In our proposal, we consider the waveguide to be a linear medium  $\omega_k = c|k|$ . When the separation distance  $d_s$  is significantly smaller than the size of wavepacket, the time-retardation effect is negligible. Given that two atoms chirally decay/absorb two correlated photons, the master equation is simplified as [73]

$$\dot{\rho}_s(t) = \frac{\Gamma_R}{2} \sum_{i=A,B} [\sigma_i^+ \sigma_i^- \rho_s(t) - \sigma_i^- \rho_s(t) \sigma_i^+] - \Gamma_R [\sigma_B^+ \sigma_A^- \rho_s(t) - \sigma_A^- \rho_s(t) \sigma_B^+] + \text{H.c.}, \quad (29)$$

where

$$\Gamma_R = \frac{L^2 |g_{k_0}|^4}{8\pi c^2} \frac{\delta_w}{\Delta^2 - \delta_w^2}$$

is the rate of the spontaneous decay to the right direction. The evolution of the system can be described by

$$\begin{aligned} \dot{\rho}_s(t) &= -iH_{\text{eff}}\rho_s + i\rho_s H_{\text{eff}}^\dagger + \mathcal{L}\rho_s \mathcal{L}^\dagger, \\ H_{\text{eff}} &= -i \sum_{i=A,B} \frac{\Gamma_R}{2} \sigma_i^+ \sigma_i^- - i\Gamma_R \sigma_B^+ \sigma_A^-, \end{aligned} \quad (30)$$

with  $\mathcal{L}$  being the collective jump operator,  $\mathcal{L} = \sqrt{\Gamma_R}(\sigma_A^- + \sigma_B^-)$ . From the last term of the Eq. (30), we derive that the two-photon state emitted by emitter A is reabsorbed by the emitter B, but does not flow back into the emitter A. Note that the cascaded master equation ignores the time decay effects by tracing over the waveguide's degrees of freedom. To overcome these constraints, we employ a simulation based on the unitary dynamics governed by the Hamiltonian Eq. (24). As discussed in Appendix. A, this numerical method can effectively capture the time delay effects.

Assuming that the emitter A is initially in  $|f\rangle$ , Fig. 10(a) illustrates the temporal evolution of the population on the two emitters being in  $|f\rangle$  (denoted as  $|c_{fA(B)}|^2$ ). It is clear that emitter B reabsorbs the correlated photon pair from emitter A. In parallel, as depicted in Fig. 10(b), we show the evolution of the field distribution in real space along the  $x = y$




FIG. 10. (a) The evolution of probability amplitudes between two giant emitters. Initially, emitter A is excited and spontaneously decay. After a retardation time, emitter B is excited at  $t = d_s/c$ . (b) The distribution of the field along the  $x = y$  direction varies with time. We set  $d_s/x_0 = 50$ .

direction over time. We find that at the initial of the evolution, the emitter A decays spontaneously, emitting a two-photon field that propagates in the right direction. After a propagating time  $t = (x_B - x_A)/c$ , the emitted field reaches emitter B, and be absorbed without being scattered back to emitter A, indicating that the two-photon mediated cascaded system is realized. Compared with chiral setups mediated by single photon, our proposal can mediate three-level giant atoms, and two correlated photons serve as correlated flying qubits, which might underpin unique application in many-body physics and quantum information processing [29, 35].

## V. CONCLUSION

In this paper, we explore how to realize two-photon correlated emission by considering a three-level giant atom interacting with a conventional linear waveguide. In contrast to previous studies, this proposal does



not need nonlinear medium. By optimizing the objective coupling function using sequential quadratic programming algorithm, the emission of a single photon is prevented, while the process of two correlated photons being emitted from the giant atom is significantly enhanced. Meanwhile, the optimal coupling function in momentum space are illustrated by Fourier transform. Considering the emitter coupling to the waveguide via linear elements, the coupling strength in right and left directions are symmetric. We show the spatial field distribution and the second-order correlation function, revealing the bidirectional propagation of photons in the waveguide and the strong spatial correlations between two identical itinerant photons. Meanwhile, this scheme ensures energy conservation without photon loss, thereby addressing the issue of photon loss in multi-photon dynamics.

Similarly, when the coupling element are adopted as nonlinear element, local phase are encoded at each coupling point. When the giant atom interacting with the waveguide in either the right or left direction, we achieve directional chiral two-photon correlated emission, which both photons propagate in the same direction. Since our proposal does not need nonlinear medium, the efficiency of emitting correlated photons can be much higher. Moreover, we show that the emitted two photons can mediate remote three-level atoms,

indicating our work can enable the cascaded process of quantum information in qutrit systems, which can be configured for quantum information and many-body quantum physics.

## VI. ACKNOWLEDGMENTS

The quantum simulations are based on QuTiP [75, 76]. X.W. is supported by the National Natural Science Foundation of China (NSFC) ( No. 12174303).

### Appendix A: The numerical simulation method

We briefly introduce the numerical method in this work. The specific steps are as follows:

(a) We discretize a set of  $M = 2000$  modes within the range of  $k \in (-2k_0, 2k_0]$ , corresponding to the discretization of a finite waveguide with a length  $L = 2000\lambda_0$  in real space, with  $\lambda_0 = 2\pi/k_0$ . The considerable length of  $L$  is crucial in preventing the propagating wavepacket from reaching the simulation boundary of the waveguide.

(b) In the two-excitation subspace, the state of the system is denoted as  $|\psi(t)\rangle = c_f(t)|f, 0\rangle + c_{e,k}(t)|e, k\rangle + c_{g,k,k'}(t)|g, k, k'\rangle$ . The dimensions of the Hamiltonian matrix is  $M^2 + M + 1$ , which is expressed as

$$\begin{pmatrix} \omega_f & g_{k1} & g_{k2} & \cdots & g_{kM} & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 & \cdots & 0 & \cdots & 0 \\ g_{k1}^* & \omega_{k1} + \omega_e & 0 & \cdots & 0 & g_{k1} & g_{k2} & \cdots & g_{kM} & 0 & \cdots & 0 & \cdots & 0 & \cdots & 0 \\ g_{k2}^* & 0 & \omega_{k2} + \omega_e & \ddots & 0 & 0 & 0 & \cdots & 0 & g_{k1} & \cdots & g_{kM} & \cdots & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & 0 & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 & \cdots & 0 & \cdots & 0 \\ g_{kM}^* & 0 & 0 & 0 & \omega_{kM} + \omega_e & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 & \cdots & g_{k1} & \cdots & g_{kM} \\ 0 & g_{k1}^* & 0 & 0 & 0 & \omega_{k1} + \omega_{k1} & 0 & \cdots & 0 & 0 & \cdots & 0 & \cdots & 0 & \cdots & 0 \\ 0 & g_{k2}^* & 0 & 0 & 0 & 0 & \omega_{k1} + \omega_{k2} & \ddots & 0 & 0 & \cdots & 0 & \cdots & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & 0 & \ddots & \cdots & 0 & \cdots & 0 & \cdots & 0 \\ 0 & g_{kM}^* & 0 & 0 & 0 & 0 & 0 & 0 & \omega_{k1} + \omega_{kM} & 0 & \cdots & 0 & \cdots & 0 & \cdots & 0 \\ 0 & 0 & g_{k1}^* & 0 & 0 & 0 & 0 & 0 & 0 & \omega_{k2} + \omega_{k1} & 0 & 0 & \cdots & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \ddots & 0 & \cdots & 0 & \cdots & 0 \\ 0 & 0 & g_{kM}^* & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \omega_{k2} + \omega_{kM} & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \ddots & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & g_{k1}^* & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \omega_{kM} + \omega_{k1} & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & 0 & 0 & 0 & g_{kM}^* & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \omega_{kM} + \omega_{kM} \end{pmatrix}$$

where  $\omega_{k_i}$  is the frequency at mode  $k_i$  and  $g_{k_i}$  is the coupling strength between the giant atom with mode  $k_i$ .

(c) We assume the giant atom initially in the second excited state and there is no photon in the waveguide, i.e.,  $|\psi(t=0)\rangle = |f, 0\rangle$ , we employ Qutip package [75, 76] to numerically solve the Schrödinger equation. We can

obtain the probability of the three-level giant emitter and the field distribution in both real space and momentum space.

### Appendix B: Optimized coupling sequence

position $x_i/x_0$	-6.97974	-6.1012	-5.85101	-5.13147	-4.87762	-4.62711	-4.37708	-4.12705	-3.87654	-3.6265
amplitude $g(x_i)$	0.01244	0.024889	0.019228	0.035245	0.036471	0	0	0.04685	0.047951	0
position $x_i/x_0$	-3.37631	-3.12612	-2.87609	-2.62542	-2.37539	-2.12536	-1.87532	-1.62497	-1.37492	-1.12481
amplitude $g(x_i)$	0	0.056732	0.057731	0	0	0.064517	0.064959	0	0	0.069555
position $x_i/x_0$	-0.87483	-0.62465	-0.37443	-0.12434	0.125658	0.376449	0.626466	0.876546	1.126562	1.376579
amplitude $g(x_i)$	0.069423	0	0	0.0717	0.070923	0	0	0.070929	0.06927	0
position $x_i/x_0$	1.626723	1.877073	2.127106	2.377138	2.62733	2.878317	3.12835	3.381033	3.630324	3.880516
amplitude $g(x_i)$	0	0.067023	0.064766	0	0	0.060415	0.057325	0	0	0.051466
position $x_i/x_0$	4.130548	4.381695	4.885579	5.140068	5.628992	5.882367	6.141789	6.40344	6.881541	7.13237
amplitude $g(x_i)$	0.047489	0	0.041289	0.035751	0	0.027585	0.02458	0	0.012783	0.011244

TABLE I. The optimized coupling sequence in the case of bidirectional photon propagation in real space.

position $x_i/x_0$	-7.14176	-6.86085	-6.57867	-6.27723	-5.99664	-5.71971	-5.44167	-5.16728	-4.88574	-4.59289
amplitude $A(x_i)$	0.005645	0.009507	0.00452	0.0083	0.018594	0.015474	0.000308	0.021212	0.030514	0.013119
phase $\theta(x_i)$	-0.30258	0.70513	1.5708	-0.65417	0.27106	1.1329	1.5708	-0.34866	0.62847	1.4027
position $x_i/x_0$	-4.22891	-3.94545	-3.63653	-3.35117	-3.06548	-2.76914	-2.48218	-2.19666	-1.90779	-1.61892
amplitude $A(x_i)$	0.022308	0.040937	0.025162	0.014754	0.045659	0.040736	0.002907	0.040073	0.05328	0.024942
phase $\theta(x_i)$	-0.5722	0.38919	1.2498	-1.0289	-0.00959	0.847	1.5708	-0.46378	0.42104	1.3122
position $x_i/x_0$	-1.33496	-1.05367	-0.77144	-0.48584	-0.19703	0.077228	0.35786	0.642509	0.92504	1.21284
amplitude $A(x_i)$	0.023473	0.055365	0.045693	0.000698	0.044941	0.05631	0.027011	0.022645	0.056513	0.046022
phase $\theta(x_i)$	-0.97905	-0.07208	0.82751	1.5708	-0.59173	0.32023	1.1815	-1.1586	-0.21906	0.70155
position $x_i/x_0$	1.801634	2.081588	2.370613	2.67635	2.956144	3.259493	3.558705	3.850913	4.158241	4.446152
amplitude $A(x_i)$	0.044359	0.051272	0.019786	0.025138	0.049259	0.032616	0.009067	0.039377	0.034449	0.003942
phase $\theta(x_i)$	-0.67889	0.27091	1.0651	-1.1264	-0.16069	0.72162	-1.5708	-0.51485	0.39107	1.5708
position $x_i/x_0$	4.744091	5.034389	5.316889	5.595251	5.893508	6.17362	6.455643	6.732413	7.009979	7.322878
amplitude $A(x_i)$	0.024155	0.029487	0.013626	0.008667	0.020083	0.014884	0.001918	0.007753	0.009316	0.004281
phase $\theta(x_i)$	-0.9162	-0.01658	0.95508	-1.4322	-0.51606	0.44446	1.4502	-1.0517	-0.08926	0.76751

TABLE II. The optimized coupling sequence in the case of directional photon propagation in real space.

- [1] Y.-P. Peng and W.-Z. Jia, Single-photon scattering from a chain of giant atoms coupled to a one-dimensional waveguide, *Phys. Rev. A* **108**, 043709 (2023).
- [2] L.-T. Xu and L.-Z. Guo, Catch and release of propagating bosonic field with non-Markovian giant atom, *New J. Phys.* **26**, 013025 (2024).
- [3] X.-L. Yin, W.-B. Luo, and J.-Q. Liao, Non-markovian disentanglement dynamics in double-giant-atom waveguide-QED systems, *Phys. Rev. A* **106**, 063703 (2022).
- [4] A. Soro, C. S. Muñoz, and A. F. Kockum, Interaction between giant atoms in a one-dimensional structured environment, *Phys. Rev. A* **107**, 013710 (2023).
- [5] X.-L. Yin and J.-Q. Liao, Generation of two-giant-atom entanglement in waveguide-QED systems, *Phys. Rev. A* **108**, 023728 (2023).
- [6] X.-Y. Li, W. Zhao, and Z.-H. Wang, Controlling photons by phonons via giant atom in a waveguide QED setup, *Opt. Lett.* **48**, 3595 (2023).
- [7] W.-J. Gu, H. Huang, Z. Yi, L. Chen, L.-H. Sun, and H.-T. Tan, Correlated two-photon scattering in a one-dimensional waveguide coupled to two- or three-level giant atoms, *Phys. Rev. A* **108**, 053718 (2023).
- [8] L. Du, Y. Zhang, J.-H. Wu, A. F. Kockum, and Y. Li, Giant atoms in a synthetic frequency dimension, *Phys. Rev. Lett.* **128**, 223602 (2022).
- [9] S. Longhi, Photonic simulation of giant atom decay, *Opt. Lett.* **45**, 3017 (2020).
- [10] A. M. Vadiraj, A. Ask, T. G. McConkey, I. Nsanzineza, C. W. S. Chang, A. F. Kockum, and C. M. Wilson,

- Engineering the level structure of a giant artificial atom in waveguide quantum electrodynamics, *Phys. Rev. A* **103**, 023710 (2021).
- [11] L. Du, Y.-T. Chen, Y. Zhang, and Y. Li, Giant atoms with time-dependent couplings, *Phys. Rev. Res.* **4**, 023198 (2022).
- [12] A. Frisk Kockum, P. Delsing, and G. Johansson, Designing frequency-dependent relaxation rates and lamb shifts for a giant artificial atom, *Phys. Rev. A* **90**, 013837 (2014).
- [13] W. Zhao and Z.-H. Wang, Single-photon scattering and bound states in an atom-waveguide system with two or multiple coupling points, *Phys. Rev. A* **101**, 053855 (2020).
- [14] X.-J. Zhang, C.-G. Liu, Z.-R. Gong, and Z.-H. Wang, Quantum interference and controllable magic cavity QED via a giant atom in a coupled resonator waveguide, *Phys. Rev. A* **108**, 013704 (2023).
- [15] L. Du, Y. Zhang, and Y. Li, A giant atom with modulated transition frequency, *Front. Phys.* **18**, 12301 (2023).
- [16] J. Zhou, X.-L. Yin, and J.-Q. Liao, Chiral and nonreciprocal single-photon scattering in a chiral-giant-molecule waveguide-QED system, *Phys. Rev. A* **107**, 063703 (2023).
- [17] A. F. Kockum, G. Johansson, and F. Nori, Decoherence-Free Interaction between Giant Atoms in Waveguide Quantum Electrodynamics, *Phys. Rev. Lett.* **120**, 140404 (2018).
- [18] K. H. Lim, W.-K. Mok, and L.-C. Kwek, Oscillating bound states in non-markovian photonic lattices, *Phys. Rev. A* **107**, 023716 (2023).
- [19] W.-Z. Jia and M.-T. Yu, Atom-photon dressed states in a waveguide-QED system with multiple giant atoms, *Opt. Express* **32**, 9495 (2024).
- [20] A. Soro and A. F. Kockum, Chiral quantum optics with giant atoms, *Phys. Rev. A* **105**, 023712 (2022).
- [21] L. Du and Y. Li, Single-photon frequency conversion via a giant  $\Lambda$ -type atom, *Phys. Rev. A* **104**, 023712 (2021).
- [22] L. Du, L.-Z. Guo, and Y. Li, Complex decoherence-free interactions between giant atoms, *Phys. Rev. A* **107**, 023705 (2023).
- [23] A. Carollo, D. Cilluffo, and F. Ciccarello, Mechanism of decoherence-free coupling between giant atoms, *Phys. Rev. Res.* **2**, 043184 (2020).
- [24] H. Xiao, L.-J. Wang, Z.-H. Li, X. Chen, and L. Yuan, Bound state in a giant atom-modulated resonators system, *npj Quantum Inf.* **8**, 80 (2022).
- [25] X. Wang, T. Liu, A. F. Kockum, H.-R. Li, and F. Nori, Tunable Chiral Bound States with Giant Atoms, *Phys. Rev. Lett.* **126**, 043602 (2021).
- [26] L.-Z. Guo, A. F. Kockum, F. Marquardt, and G. Johansson, Oscillating bound states for a giant atom, *Phys. Rev. Res.* **2**, 043014 (2020).
- [27] S.-J. Guo, Y.-D. Wang, T. Purdy, and J. Taylor, Beyond spontaneous emission: Giant atom bounded in the continuum, *Phys. Rev. A* **102**, 033706 (2020).
- [28] Y.-T. Chen, L. Du, L.-Z. Guo, Z.-H. Wang, Y. Zhang, Y. Li, and J.-H. Wu, Nonreciprocal and chiral single-photon scattering for giant atoms, *Commun. Phys.* **5**, 215 (2022).
- [29] X. Wang, J.-Q. L. Z.-H. Wang, A. F. Kockum, L. Du, T. Liu, and F. Nori, Nonlinear chiral quantum optics with giant-emitter pairs (2024), [arXiv:2404.09829 \[quant-ph\]](https://arxiv.org/abs/2404.09829).
- [30] W.-J. Gu, L. Chen, Z. Yi, S.-J. Liu, and G.-X. Li, Tunable photon-photon correlations in waveguide QED systems with giant atoms, *Phys. Rev. A* **109**, 023720 (2024).
- [31] J. Grebel, H.-X. Yan, M.-H. Chou, G. Andersson, C. R. Conner, Y. J. Joshi, J. M. Miller, R. G. Povey, H. Qiao, X. Wu, and A. N. Cleland, Bidirectional multiphoton communication between remote superconducting nodes, *Phys. Rev. Lett.* **132**, 047001 (2024).
- [32] F. Dell'Anno, S. De Siena, and F. Illuminati, Multiphoton quantum optics and quantum state engineering, *Phys. Rep.* **428**, 53 (2006).
- [33] M. Zhang, L.-T. Feng, Z. yuan Zhou, Y. Chen, H. Wu, M. Li, S.-M. Gao, G.-P. Guo, G. can Guo, D. Dai, and X. Ren, Generation of multiphoton quantum states on silicon, *Light Sci. Appl.* **8**, 41 (2019).
- [34] C. Liu, J.-F. Huang, and L. Tian, Deterministic generation of multi-photon bundles in a quantum rabi model, *Sci. China Phys. Mech. Astron.* **66**, 220311 (2022).
- [35] X. Wang, J.-Q. Li, T. Liu, A. Miranowicz, and F. Nori, Long-range four-body interactions in structured nonlinear photonic waveguides (2024), [arXiv:2401.02714 \[quant-ph\]](https://arxiv.org/abs/2401.02714).
- [36] B. J. Metcalf, N. Thomas-Peter, J. B. Spring, D. Kundys, M. A. Broome, P. C. Humphreys, X.-M. Jin, M. Barbieri, W. Steven Kolthammer, J. C. Gates, B. J. Smith, N. K. Langford, P. G. Smith, and I. A. Walmsley, Multiphoton quantum interference in a multiport integrated photonic device, *Nat. Commun.* **4**, 1356 (2013).
- [37] S. Mahmoodian, M. Čepulkovskis, S. Das, P. Lodahl, K. Hammerer, and A. S. Sørensen, Strongly correlated photon transport in waveguide quantum electrodynamics with weakly coupled emitters, *Phys. Rev. Lett.* **121**, 143601 (2018).
- [38] Y.-G. Ke, A. V. Poshakinskiy, C. Lee, Y. S. Kivshar, and A. N. Poddubny, Inelastic scattering of photon pairs in qubit arrays with subradiant states, *Phys. Rev. Lett.* **123**, 253601 (2019).
- [39] P.-C. Li and S.-I. Chu, Multiphoton quantum dynamics of many-electron atomic and molecular systems in intense laser fields, *Chinese Phys. B.* **29**, 083202 (2020).
- [40] A. S. Prasad, J. Hinney, S. Mahmoodian, K. Hammerer, S. Rind, P. Schneeweiss, A. S. Sørensen, J. Volz, and A. Rauschenbeutel, Correlating photons using the collective nonlinear response of atoms weakly coupled to an optical mode, *Nat. Photonics* **14**, 719 (2020).
- [41] G. Calajó and D. E. Chang, Emergence of solitons from many-body photon bound states in quantum nonlinear media, *Phys. Rev. Res.* **4**, 023026 (2022).
- [42] S. H. Cantu, A. V. Venkatramani, W.-C. Xu, L. Zhou, B. Jelenković, M. D. Lukin, and V. Vuletić, Repulsive photons in a quantum nonlinear medium, *Nat. Phys.* **16**, 921 (2020).
- [43] G. S. He, Multi-Photon Excitation Based Nonlinear Optical Effects and Applications, *Prog. Optics*, **64**, 155 (2019).
- [44] G. S. He, Multi-photon nonlinear optical effects, in *Nonlinear Optics and Photonics* (Oxford University Press, 2014).
- [45] Y. V. G. S. Murti and C. Vijayan, Basics of multi-photon microscopy, in *Physics of Nonlinear Optics* (Springer International Publishing, Cham, 2021) pp. 157–168.
- [46] T. Peyronel, O. Firstenberg, Q. Y. Liang, S. Hofferberth, A. V. Gorshkov, T. Pohl, M. D. Lukin, and V. Vuletić,

- Quantum nonlinear optics with single photons enabled by strongly interacting atoms, *Nature* **488**, 57 (2012).
- [47] O. A. Iversen and T. Pohl, Strongly correlated states of light and repulsive photons in chiral chains of three-level quantum emitters, *Phys. Rev. Lett.* **126**, 083605 (2021).
- [48] Z.-H. Wang, T. Jaako, P. Kirton, and P. Rabl, Supercorrelated radiance in nonlinear photonic waveguides, *Phys. Rev. Lett.* **124**, 213601 (2020).
- [49] B. Y. Nikitchuk, E. V. Anikin, and N. S. Maslova, Multi-photon Rabi oscillations in the presence of the classical noise in a quantum nonlinear oscillator (2023), [arXiv:2312.01070 \[quant-ph\]](https://arxiv.org/abs/2312.01070).
- [50] I. García-Elcano, A. González-Tudela, and J. Bravo-Abad, Tunable and robust long-range coherent interactions between quantum emitters mediated by weyl bound states, *Phys. Rev. Lett.* **125**, 163602 (2020).
- [51] J.-W. Pan, Z.-B. Chen, C.-Y. Lu, H. Weinfurter, A. Zeilinger, and M. Żukowski, Multiphoton entanglement and interferometry, *Rev. Mod. Phys.* **84**, 777 (2012).
- [52] M. Barbieri, T. Weinhold, B. Lanyon, A. Gilchrist, K. Resch, M. Almeida, and A. White, Parametric downconversion and optical quantum gates: two's company, four's a crowd, *J. Mod. Opt.* **56**, 209 (2009).
- [53] D. Llewellyn, Y.-H. Ding, I. I. Faruque, S. Paesani, D. Bacco, R. Santagati, Y.-J. Qian, Y. Li, Y.-F. Xiao, M. Huber, M. Malik, G. F. Sinclair, X.-Q. Zhou, K. Rottwitt, J. L. O'Brien, J. G. Rarity, Q.-H. Gong, L. K. Oxenlowe, J.-W. Wang, and M. G. Thompson, Chip-to-chip quantum teleportation and multi-photon entanglement in silicon, *Nat. Phys.* **16**, 148–153 (2019).
- [54] Z. Hou, J.-F. Tang, C.-J. Huang, Y.-F. Huang, G.-Y. Xiang, C.-F. Li, and G.-C. Guo, Entangled-state time multiplexing for multiphoton entanglement generation, *Phys. Rev. Appl.* **19**, L011002 (2023).
- [55] H.-x. Zheng and H. U. Baranger, Persistent quantum beats and long-distance entanglement from waveguide-mediated interactions, *Phys. Rev. Lett.* **110**, 113601 (2013).
- [56] T. Seidelmann, T. K. Bracht, B. U. Lehner, C. Schimpf, M. Cosacchi, M. Cygorek, A. Vagov, A. Rastelli, D. E. Reiter, and V. M. Axt, Two-photon excitation with finite pulses unlocks pure dephasing-induced degradation of entangled photons emitted by quantum dots, *Phys. Rev. B* **107**, 235304 (2023).
- [57] Z.-G. Xia, X. Xiang, R. Quan, R.-F. Dong, T. Liu, and S.-G. Zhang, Dispersion broadening of the two-photon quantum interference width due to temporal filtering, *Phys. Rev. Appl.* **20**, 044080 (2023).
- [58] M. E. Mycroft, T. McDermott, A. Buraczewski, and M. Stobińska, Proposal for the distribution of multiphoton entanglement with optimal rate-distance scaling, *Phys. Rev. A* **107**, 012607 (2023).
- [59] J. M. Raimond, M. Brune, and S. Haroche, Manipulating quantum entanglement with atoms and photons in a cavity, *Rev. Mod. Phys.* **73**, 565 (2001).
- [60] K.-Y. Qian, K. Wang, L.-Z. Chen, Z.-H. Hou, M. Krenn, S.-N. Zhu, and X.-S. Ma, Multiphoton non-local quantum interference controlled by an undetected photon, *Nat. Commun.* **14** (2023).
- [61] S. Massar, F. Devaux, and E. Lantz, Multiphoton correlations between quantum images, *Phys. Rev. A* **108**, 013705 (2023).
- [62] K. Barkemeyer, A. Knorr, and A. Carmele, Heisenberg treatment of multiphoton pulses in waveguide QED with time-delayed feedback, *Phys. Rev. A* **106**, 023708 (2022).
- [63] H. Le Jeannic, A. Tiranov, J. Carolan, T. Ramos, Y. Wang, M. H. Appel, S. Scholz, A. D. Wieck, A. Ludwig, N. Rotenberg, L. Midolo, J. J. García-Ripoll, A. S. Sørensen, and P. Lodahl, Dynamical photon-photon interaction mediated by a quantum emitter, *Nat. Phys.* **18**, 1191–1195 (2022).
- [64] L. Wang, Z. Yi, L.-H. Sun, and W.-J. Gu, Nonreciprocal two-photon transmission and statistics in a chiral waveguide QED system, *Chinese Phys. B* **31**, 054206 (2022).
- [65] T. Shi, D. E. Chang, and J. I. Cirac, Multiphoton-scattering theory and generalized master equations, *Phys. Rev. A* **92**, 053834 (2015).
- [66] S. Li, Z.-C. Ni, L.-B. Zhang, Y.-Y. Cai, J.-S. Mai, S.-C. Wen, P. Zheng, X.-W. Deng, S. Liu, Y. Xu, and D.-P. Yu, Autonomous stabilization of fock states in an oscillator against multiphoton losses, *Phys. Rev. Lett.* **132**, 203602 (2024).
- [67] P. Fraser, The impact of distinct loss mechanisms on multi-photon states, *Eur. J. Phys* **42**, 035303 (2021).
- [68] X. Wang, H.-B. Zhu, T. Liu, and F. Nori, Realizing quantum optics in structured environments with giant atoms, *Phys. Rev. Res.* **6**, 013279 (2024).
- [69] A. Blais, A. L. Grimsmo, S. M. Girvin, and A. Wallraff, Circuit quantum electrodynamics, *Rev. Mod. Phys.* **93**, 025005 (2021).
- [70] J. Koch, T. M. Yu, J. Gambetta, A. A. Houck, D. I. Schuster, J. Majer, A. Blais, M. H. Devoret, S. M. Girvin, and R. J. Schoelkopf, Charge-insensitive qubit design derived from the cooper pair box, *Phys. Rev. A* **76**, 042319 (2007).
- [71] X. Wang, Y.-F. Lin, J.-Q. Li, W.-X. Liu, and H.-R. Li, Chiral SQUID-metamaterial waveguide for circuit-QED, *New J. Phys.* **24**, 123010 (2022).
- [72] M. O. Scully and M. S. Zubairy, *Quantum Optics* (Cambridge University Press, 1997).
- [73] X. Wang, Y.-F. Lin, J.-Q. Li, W.-X. Liu, and H.-R. Li, Chiral SQUID-metamaterial waveguide for circuit-qed, *New Journal of Physics* **24**, 123010 (2022).
- [74] G. Calajó, M. J. A. Schuetz, H. Pichler, M. D. Lukin, P. Schneeweiss, J. Volz, and P. Rabl, Quantum acousto-optic control of light-matter interactions in nanophotonic networks, *Phys. Rev. A* **99**, 053852 (2019).
- [75] J. R. Johansson, P. D. Nation, and F. Nori, Qutip: An open-source Python framework for the dynamics of open quantum systems, *Comput. Phys. Commun.* **183**, 1760 (2012).
- [76] J. R. Johansson, P. D. Nation, and F. Nori, Qutip 2: A Python framework for the dynamics of open quantum systems, *Comput. Phys. Commun.* **184**, 1234 (2013).