

Tutorial on one-dimensional q -Fourier transforms

Hartmut Wachter*
 An der Schafscheuer 56
 D-91781 Weißenburg, Federal Republic of Germany

June 21, 2024

Abstract

This paper is an introductory text to the theory of q -deformed Fourier transforms, as first discussed by Rogov and Olshanetsky. We derive the well-known results in detail, present them in a format that suits our needs, and include some new findings on specific aspects of the theory.

Contents

1	Introduction	1
2	Exponential functions	2
3	Trigonometric functions	7
4	Integrals	9
5	Distributions	13
6	Fourier transforms	20
7	Fourier transforms on $\mathbb{G}_{q^{1/2}, x_0}$	34

1 Introduction

The first three chapters review some basics of the so-called q -calculus. These include the Jackson derivatives and the Jackson integrals as q -analogs of ordinary derivatives or integrals. The eigenfunctions of the Jackson derivative lead us to q -analogs of the exponential function. Using the q -exponential, we introduce q -analogs of the trigonometric functions and derive q -versions of the Taylor expansion. In the subsequent chapters, we present a q -version of the Hilbert space

*E-Mail: Hartmut.Wachter@gmx.de

of square-integrable functions and study the spaces of test functions and q -distributions. Finally, we discuss q -analogs of the classical Fourier transforms of position or momentum space.

Our approach uses the considerations from Ref. [1], adapting them to our specific requirements. We have also added some new results relevant to our further investigations. Remember, there are other q -versions of Fourier transforms apart from the ones discussed in this paper [3,4]. The q -deformed Fourier transforms used in this paper are the best option for our needs because they closely resemble the classical Fourier transform.

2 Exponential functions

The **Jackson derivative** of a real function f is defined as follows [7]:

$$D_q f(x) = \frac{f(x) - f(qx)}{x - qx}. \quad (1)$$

In the limit $q \rightarrow 1$, the Jackson derivative becomes the ordinary derivative of a real function [8]:

$$\lim_{q \rightarrow 1} D_q f(x) = \frac{df(x)}{dx}. \quad (2)$$

The definition in Eq. (1) implies the following rule for the Jackson derivative of a power function ($\alpha \in \mathbb{R}$)

$$D_q x^\alpha = [[\alpha]]_q x^{\alpha-1}, \quad (3)$$

The **antisymmetric q -numbers** are defined as follows:

$$[[\alpha]]_q = \frac{1 - q^\alpha}{1 - q}. \quad (4)$$

For $q \rightarrow 1$, the antisymmetric q -number becomes the corresponding real number:

$$\lim_{q \rightarrow 1} [[\alpha]]_q = \alpha. \quad (5)$$

The **q -factorial** is introduced in analogy to the undeformed case ($n \in \mathbb{N}$):

$$[[n]]_q! = [[1]]_q [[2]]_q \dots [[n-1]]_q [[n]]_q, \quad [[0]]_q! = 1. \quad (6)$$

Accordingly, we define the **q -binomial coefficients** as follows ($k \in \mathbb{N}$):

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{[[n]]_q!}{[[n-k]]_q! [[k]]_q!}. \quad (7)$$

By mathematical induction, we can proof the following formula for the Jackson derivative [8]:

$$D_q^n f(x) = (-1)^n \frac{q^{-n(n-1)/2}}{(1-q)^n x^n} \sum_{m=0}^n \begin{bmatrix} n \\ m \end{bmatrix}_q (-1)^m q^{m(m-1)/2} f(q^{n-m}x). \quad (8)$$

Moreover, the Jackson derivative satisfies a **product rule**:

$$\begin{aligned} D_q(f(x)g(x)) &= f(qx)D_qg(x) + g(x)D_qf(x) \\ &= f(x)D_qg(x) + g(qx)D_qf(x). \end{aligned} \quad (9)$$

We introduce the q -**exponential** as eigenfunction of the Jackson derivative [9], i. e.

$$D_q \exp_q(x) = \exp_q(x) \quad (10)$$

with the normalization condition

$$\exp_q(0) = 1. \quad (11)$$

To get a power series expansion of the q -exponential, we make the following ansatz:

$$\exp_q(x) = \sum_{k=0}^{\infty} (C_q)_k x^k. \quad (12)$$

Due to Eqs. (10) and (11), the coefficients $(C_q)_k$ satisfy the following recurrence relations:

$$(C_q)_{k+1} = \frac{1}{[[k+1]]_q} (C_q)_k, \quad (C_q)_0 = 1. \quad (13)$$

By iteration, we find:

$$(C_q)_k = \frac{1}{[[k]]_q!}. \quad (14)$$

Thus, the power series expansion for the q -exponential takes the following form:

$$\exp_q(x) = \sum_{k=0}^{\infty} \frac{1}{[[k]]_q!} x^k. \quad (15)$$

For $q \rightarrow 1$, the q -exponential becomes the ordinary exponential function [cf. Eq. (5)]:

$$\lim_{q \rightarrow 1} \exp_q(x) = e^x. \quad (16)$$

The q -exponential in Eq. (15) is related to the following functions ($0 < q < 1$) [8, 10]:

$$\begin{aligned} E_q(z) &= \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2} z^k}{(q; q)_k} = (-z; q)_{\infty}, \\ e_q(z) &= \sum_{k=0}^{\infty} \frac{z^k}{(q; q)_k} = \frac{1}{(z; q)_{\infty}}. \end{aligned} \quad (17)$$

In the formulae above, we have used the q -**Pochhammer symbol** ($a \in \mathbb{C}$, $n \in \mathbb{N}_0$):

$$(a; q)_n = \begin{cases} 1 & \text{für } n = 0, \\ (1-a)(1-aq) \dots (1-aq^{n-1}) & \text{für } n \geq 1. \end{cases} \quad (18)$$

Accordingly, we have:

$$(a; q)_\infty = \lim_{n \rightarrow \infty} (a; q)_n = \prod_{j=0}^{\infty} (1 - qa^j). \quad (19)$$

The power series of $e_q(z)$ converges for $|z| < 1$ and that of $E_q(z)$ for $z \in \mathbb{C}$. Due to the identities

$$[[n]]_q! = \frac{(q; q)_n}{(1 - q)^n} \quad (20)$$

and [10]

$$(a; q)_n = (a^{-1}; q^{-1})_n (-a)^n q^{n(n-1)/2}, \quad (n \in \mathbb{N}_0), \quad (21)$$

it holds:

$$e_q((1 - q)z) = \exp_q(z), \quad E_q((1 - q)z) = \exp_{q^{-1}}(z). \quad (22)$$

Comparing the two identities in Eq. (22), we find:

$$e_q((1 - q)z) = E_{q^{-1}}((1 - q^{-1})z). \quad (23)$$

Due to Eq. (21) and the two power series expansions in Eq. (17), we have:

$$e_{q^{-1}}(z) = E_q(-qz). \quad (24)$$

The representations of $E_q(z)$ and $e_q(z)$ as products imply the following identity [cf. Eq. (17)]:

$$e_q(z) E_q(-z) = 1. \quad (25)$$

As we can see from its representation in Eq. (17), the q -exponential function $e_q(z)$ is a meromorphic function with poles at $z = q^{-k}$, $k \in \mathbb{N}_0$. Hence, there is a partial fraction decomposition for $e_q(z)$ [1], i. e.

$$e_q(z) = \sum_{k=0}^{\infty} \frac{c_{k,n}}{1 - zq^k} \quad (26)$$

with

$$c_{k,n} = \lim_{z \rightarrow q^{-k}} (1 - zq^k) e_q(z) = \frac{(-1)^k q^{k(k+1)/2}}{(q; q)_\infty (q; q)_k}. \quad (27)$$

The series in Eq. (26) converges for $0 < q < 1$ and $z \neq q^{-2k}$.

We use the q -exponential in Eq. (15) to calculate q -translations [9, 11, 12]. To this end, we introduce the operator:

$$\exp_q(a|D_q) = \sum_{k=0}^{\infty} \frac{a^k}{[[k]]_q!} D_q^k. \quad (28)$$

Applying this operator to a function $f(x)$, we get the following q -version of **Taylor's formula**:

$$f(a \oplus x) = \exp_q(a|D_q) \triangleright f(x) = \sum_{k=0}^{\infty} \frac{a^k}{[[k]]_q!} D_q^k f(x). \quad (29)$$

Similarly, it holds:

$$f(x \bar{\oplus} a) = \sum_{k=0}^{\infty} \frac{1}{[[k]]_q!} (D_q^k f(x)) a^k. \quad (30)$$

Applying the operator in Eq. (28) to a power function with a natural number as the exponent, we obtain a **q -version of the binomial theorem**:

$$\begin{aligned} (a \bar{\oplus} x)^n &= \exp_q(a|D_q) \triangleright x^n = \sum_{k=0}^{\infty} \frac{a^k}{[[k]]_q!} D_q^k \triangleright x^n \\ &= \sum_{k=0}^{\infty} \frac{a^k}{[[k]]_q!} [[n]]_q \cdots [[n-k+1]]_q x^{n-k} = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q a^{n-k} x^k. \end{aligned} \quad (31)$$

The q -binomial theorem implies the identities

$$\begin{aligned} (a \bar{\oplus} 0)^n &= (a \oplus x)|_{x=0} = a^n \\ (0 \bar{\oplus} a)^n &= (x \bar{\oplus} a)|_{x=0} = a^n, \end{aligned} \quad (32)$$

which can be generalized as follows:

$$\begin{aligned} f(a \bar{\oplus} 0) &= f(a \bar{\oplus} x)|_{x=0} = f(a), \\ f(0 \bar{\oplus} a) &= f(x \bar{\oplus} y)|_{x=0} = f(a). \end{aligned} \quad (33)$$

There is a second operator for calculating q -translations:

$$\begin{aligned} \exp_{q^{-1}}(-a|D_q) &= \sum_{k=0}^{\infty} \frac{(-a)^k}{[[k]]_{q^{-1}}!} D_q^k \\ &= \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{[[k]]_q!} (-a)^k D_q^k. \end{aligned} \quad (34)$$

Applying this operator to a function $f(x)$, we get:

$$\begin{aligned} f((\bar{\ominus} a) \bar{\oplus} x) &= \exp_{q^{-1}}(-a|D_q) \triangleright f(x) \\ &= \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{[[k]]_q!} (-a)^k D_q^k f(x). \end{aligned} \quad (35)$$

Accordingly, we have:

$$f(x \bar{\oplus} (\bar{\ominus} a)) = \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{[[k]]_q!} (D_q^k f(x)) (-a)^k. \quad (36)$$

The operators in Eqs. (28) and (34) are inverses of each other. We show this

by the following calculation:

$$\begin{aligned}
& \exp_{q^{-1}}(-a|D_q) \exp_q(a|D_q) \triangleright f(x) = \exp_q(a|D_q) \exp_{q^{-1}}(-a|D_q) \triangleright f(x) \\
&= \sum_{k,j=0}^{\infty} \frac{(-1)^k q^{k(k-1)/2}}{[[k]]_q! [[j]]_q!} a^{k+j} D_q^{k+j} f(x) \\
&= \sum_{k,j=0}^{\infty} \frac{(-1)^k q^{k(k-1)/2}}{[[k+j]]_q!} \frac{[[k+j]]_q!}{[[k]]_q! [[j]]_q!} a^{k+j} D_q^{k+j} f(x) \\
&= \sum_{n=0}^{\infty} \frac{1}{[[n]]_q!} \sum_{k=0}^n (-1)^k q^{k(k-1)/2} \frac{[[n]]_q!}{[[k]]_q! [[n-k]]_q!} a^n D_q^n f(x) \\
&= \sum_{n=0}^{\infty} \frac{1}{[[n]]_q!} \sum_{k=0}^n (-1)^k q^{k(k-1)/2} \begin{bmatrix} n \\ k \end{bmatrix}_q a^n D_q^n f(x) = f(x). \tag{37}
\end{aligned}$$

In the last step of the above calculation, we used the following identity [8]:

$$\sum_{k=0}^n (-1)^k q^{k(k-1)/2} \begin{bmatrix} n \\ k \end{bmatrix}_q = (1; q)_n = \begin{cases} 1 & \text{if } n = 0, \\ 0 & \text{if } n \geq 1. \end{cases} \tag{38}$$

If we shift the q -exponential by a q -translation, the following **addition theorem** holds [11]:

$$\exp_q(x \bar{\oplus} a) = \exp_q(x) \exp_q(a). \tag{39}$$

We prove this rule as follows [cf. Eq. (31)]:

$$\begin{aligned}
\exp_q(x) \exp_q(a) &= \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \frac{1}{[[k]]_q!} \frac{1}{[[l]]_q!} x^k a^l \\
&= \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \frac{1}{[[k+l]]_q!} \frac{[[k+l]]_q!}{[[k]]_q! [[l]]_q!} x^k a^l \\
&= \sum_{n=0}^{\infty} \frac{1}{[[n]]_q!} \sum_{l=0}^n \begin{bmatrix} n \\ l \end{bmatrix}_q x^{n-l} a^l \\
&= \sum_{n=0}^{\infty} \frac{1}{[[n]]_q!} (x \bar{\oplus} a)^n = \exp_q(x \bar{\oplus} a). \tag{40}
\end{aligned}$$

Using Eq. (35), we can calculate so-called **q -inversions**:¹

$$f((\bar{\ominus} x)) = \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{[[k]]_q!} (-x)^k (D_q^k f)(0). \tag{41}$$

For powers of x , we find:

$$(\bar{\ominus} x)^n = q^{n(n-1)/2} (-x)^n. \tag{42}$$

¹In Chap. 6, we use the following notation for the q -inversion: $\bar{\underline{x}}_x(f(x)) = f(\bar{\ominus} x)$.

Taking Eqs. (10) and (11) into account, we also get [cf. Eq. (15)]:

$$\begin{aligned}\exp_q(\bar{\ominus} x) &= \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{[[k]]_q!} (-x)^k \exp_q(0) \\ &= \sum_{k=0}^{\infty} \frac{1}{[[k]]_{q^{-1}}!} (-x)^k = \exp_{q^{-1}}(-x).\end{aligned}\quad (43)$$

By similar reasoning as in Eq. (37), we can prove the following identities:

$$\exp_q(\bar{\ominus} x) \exp_q(x) = \exp_q(x) \exp_q(\bar{\ominus} x) = 1.$$

3 Trigonometric functions

We use the q -exponential function $e_q(z)$ to introduce q -versions of trigonometric functions [8], i. e.

$$\begin{aligned}\widetilde{\sin}_q(z) &= \frac{1}{2i} (e_q(iz) - e_q(-iz)) = \sum_{n=0}^{\infty} \frac{(-1)^n z^{2n+1}}{(q; q)_{2n+1}}, \\ \widetilde{\cos}_q(z) &= \frac{1}{2} (e_q(iz) + e_q(-iz)) = \sum_{n=0}^{\infty} \frac{(-1)^n z^{2n}}{(q; q)_{2n}},\end{aligned}\quad (44)$$

and

$$\begin{aligned}\widetilde{\text{Sin}}_q(z) &= \frac{1}{2i} (E_q(iz) - E_q(-iz)), \\ \widetilde{\text{Cos}}_q(z) &= \frac{1}{2} (E_q(iz) + E_q(-iz)).\end{aligned}\quad (45)$$

Using the partial fraction decomposition in Eq. (26) of the previous chapter, we get the following expressions for the q -trigonometric functions in Eq. (44):

$$\begin{aligned}\widetilde{\sin}_q(z) &= \frac{z}{(q; q)_{\infty}} \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+3)/2}}{(q; q)_k (1 + z^2 q^{2k})}, \\ \widetilde{\cos}_q(z) &= \frac{1}{(q; q)_{\infty}} \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+1)/2}}{(q; q)_k (1 + z^2 q^{2k})}.\end{aligned}\quad (46)$$

For $x \in \mathbb{R}$, we can estimate both series in Eq. (46) by the following inequalities:

$$\begin{aligned}\left| \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+3)/2}}{(q; q)_k (1 + x^2 q^{2k})} \right| &\leq \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{(q; q)_k (1 + x^2)} = \frac{E_q(1)}{1 + x^2}, \\ \left| \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+1)/2}}{(q; q)_k (1 + x^2 q^{2k})} \right| &\leq \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2} q^{-k}}{(q; q)_k (1 + x^2)} = \frac{E_q(q^{-1})}{1 + x^2}.\end{aligned}\quad (47)$$

For $0 < q < 1$ and $k \in \mathbb{N}_0$, the inequalities above result from

$$\frac{1}{1 + x^2 q^{2k}} \leq \frac{q^{-2k}}{1 + x^2}. \quad (48)$$

Using the results in Eq. (47), we finally get [also cf. Eq. (17) of the previous chapter]:

$$|\widetilde{\sin}_q(x)| \leq \frac{(-1; q)_\infty |x|}{(q; q)_\infty (1 + x^2)}, \quad |\widetilde{\cos}_q(x)| \leq \frac{(-q; q)_\infty}{(q; q)_\infty (1 + x^2)}. \quad (49)$$

There are also inequalities for the q -trigonometric functions in Eq. (45) (cf. Ref. [1]):

$$|\widetilde{\text{Sin}}_q(x)| \leq |x|, \quad |\widetilde{\text{Cos}}_q(x)| \leq 1. \quad (50)$$

The q -trigonometric functions in Eqs. (44) and (45) do not become the ordinary trigonometric functions if we take $q \rightarrow 1$. For this reason, we introduce the new q -trigonometric functions [cf. Eq. (22) of the previous chapter]

$$\begin{aligned} \sin_q(x) &= \widetilde{\sin}_q((1 - q)x) = \frac{1}{2i} (\exp_q(ix) - \exp_q(-ix)), \\ \cos_q(x) &= \widetilde{\cos}_q((1 - q)x) = \frac{1}{2} (\exp_q(ix) + \exp_q(-ix)), \end{aligned} \quad (51)$$

and

$$\begin{aligned} \text{Sin}_q(x) &= \widetilde{\text{Sin}}_q((1 - q)x) = \frac{1}{2i} (\exp_{q^{-1}}(ix) - \exp_{q^{-1}}(-ix)), \\ \text{Cos}_q(x) &= \widetilde{\text{Cos}}_q((1 - q)x) = \frac{1}{2} (\exp_{q^{-1}}(ix) + \exp_{q^{-1}}(-ix)). \end{aligned} \quad (52)$$

Due to Eq. (16) of the previous chapter, we regain the ordinary trigonometric functions if $q \rightarrow 1$:

$$\lim_{q \rightarrow 1^-} \sin_q(x) = \sin(x), \quad \lim_{q \rightarrow 1^-} \cos_q(x) = \cos(x). \quad (53)$$

From the expressions in Eqs. (51) and (52), we get the following identities:

$$\text{Sin}_q(x) = \sin_{q^{-1}}(x), \quad \text{Cos}_q(x) = \cos_{q^{-1}}(x). \quad (54)$$

Furthermore, there is a q -version of the identity $\sin^2(x) + \cos^2(x) = 1$ (also cf. Ref. [13]):

$$\sin_q(x) \sin_{q^{-1}}(x) + \cos_q(x) \cos_{q^{-1}}(x) = 1. \quad (55)$$

By applying the Jackson derivatives to the q -trigonometric functions in Eqs. (51) and (52), we find (also cf. Ref. [13]):

$$\begin{aligned} D_q \sin_q(x) &= \cos_q(x), & D_q \text{Sin}_q(x) &= \text{Cos}_q(qx), \\ D_q \cos_q(x) &= -\sin_q(x), & D_q \text{Cos}_q(x) &= -\text{Sin}_q(qx). \end{aligned} \quad (56)$$

Accordingly, $\sin_q(\omega x)$ and $\cos_q(\omega x)$ are linearly independent solutions of the difference equation (also cf. Ref. [8])

$$D_q^2 f(x) + \omega^2 f(x) = 0, \quad (57)$$

and the q -trigonometric functions $\text{Sin}_q(\omega x)$ and $\text{Cos}_q(\omega x)$ satisfy the difference equation

$$D_q^2 f(x) + \omega^2 f(q^2 x) = 0. \quad (58)$$

4 Integrals

For $z > 0$ and $0 < q < 1$, the one-dimensional **Jackson integral** is defined as follows [14]:

$$\begin{aligned} \int_z^{z.\infty} d_q x f(x) &= (1-q)z \sum_{j=1}^{\infty} q^{-j} f(q^{-j}z), \\ \int_0^z d_q x f(x) &= (1-q)z \sum_{j=0}^{\infty} q^j f(q^j z). \end{aligned} \quad (59)$$

Accordingly, we have:

$$\begin{aligned} \int_0^{z.\infty} d_q x f(x) &= \int_0^z d_q x f(x) + \int_z^{z.\infty} d_q x f(x) \\ &= (1-q)z \sum_{j=-\infty}^{\infty} q^{-j} f(q^{-j}z). \end{aligned} \quad (60)$$

So far, we have only considered q -integrals with a non-negative integration range. We can define q -integrals with a negative integration range as well ($z < 0$ and $0 < q < 1$):

$$\begin{aligned} \int_{-\infty}^z d_q x f(x) &= (q-1)z \sum_{j=1}^{\infty} q^{-j} f(q^{-j}z), \\ \int_z^0 d_q x f(x) &= (q-1)z \sum_{j=0}^{\infty} q^j f(q^j z). \end{aligned} \quad (61)$$

Accordingly, we have:

$$\int_{z.\infty}^0 d_q x f(x) = (q-1)z \sum_{j=-\infty}^{\infty} q^j f(q^j z). \quad (62)$$

The q -integrals in Eqs. (60) and (62) can be combined to form a single

q -integral over the interval $] -\infty, \infty[$:

$$\begin{aligned} \int_{-z.\infty}^{z.\infty} d_q x f(x) &= \int_0^{z.\infty} d_q x f(x) + \int_{-\infty.z}^0 d_q x f(x) \\ &= (1-q)z \sum_{j=-\infty}^{\infty} q^j [f(q^j z) + f(-q^j z)]. \end{aligned} \quad (63)$$

Eq. (63) implies:

$$\int_{-x.\infty}^{x.\infty} d_q z f(z) = \int_{-x.\infty}^{x.\infty} d_q z f(-z). \quad (64)$$

A look at Eq. (60) or Eq. (62) shows that only the lattice points $\{q^k x \mid k \in \mathbb{Z}\}$ contribute to the q -integral over $]0, \infty[$ or $] -\infty, 0[$. Thus, the integrand f has to satisfy the following boundary condition for the q -integral over $]0, \infty[$ or $] -\infty, 0[$ to converge ($q < 1$):

$$\lim_{k \rightarrow -\infty} f(\pm q^k x) = 0. \quad (65)$$

If f is continuous at 0, the boundary condition above implies:

$$\begin{aligned} \int_0^{z.\infty} d_q x D_q f(x) &= (1-q) \sum_{j=-\infty}^{\infty} z q^{-j} \frac{f(q^{-j} z) - f(q^{-j+1} z)}{(1-q)q^{-j} z} \\ &= \lim_{k \rightarrow \infty} \sum_{j=-k}^k [f(q^{-j} z) - f(q^{-j+1} z)] \\ &= \lim_{k \rightarrow \infty} [f(q^{-k} z) - f(q^{k+1} z)] = -f(0). \end{aligned} \quad (66)$$

A similar reasoning shows:

$$\int_{-\infty.z}^0 d_q x D_q f(x) = f(0). \quad (67)$$

We can combine Eqs. (66) and (67) to give the following result ($m \in \mathbb{N}$):

$$\begin{aligned} \int_{-z.\infty}^{z.\infty} d_q x D_q^m f(x) &= \int_0^{z.\infty} d_q x D_q^m f(x) + \int_{-\infty.z}^0 d_q x D_q^m f(x) \\ &= -D_q^{m-1} f(0) + D_q^{m-1} f(0) = 0. \end{aligned} \quad (68)$$

If the two functions f and g vanish at infinity,² Eq. (68) and Eq. (9) of Chap. 2 imply the following **q -deformed rule for integration by parts**:

$$\int_{-z.\infty}^{z.\infty} d_q x f(x) D_q g(x) = - \int_{-z.\infty}^{z.\infty} d_q x [D_q f(x)] g(qx). \quad (69)$$

Applying this rule several times, we get the following identity [1] ($k \in \mathbb{N}$):

$$\int_{-z.\infty}^{z.\infty} d_q x f(x) D_q^k g(x) = (-1)^k q^{-k(k-1)/2} \int_{-z.\infty}^{z.\infty} d_q x [D_q^k f(x)] g(q^k x). \quad (70)$$

²The weaker condition $\lim_{k \rightarrow -\infty} f(\pm q^k a) g(\pm q^k a) = 0$ is also sufficient.

Using Eq. (68), we can show that the q -integral over $]-\infty, \infty[$ is invariant under translation [4, 11, 15]:

$$\begin{aligned} \int_{-z.\infty}^{z.\infty} d_q x f(x \oplus a) &= \int_{-z.\infty}^{z.\infty} d_q x \sum_{k=0}^{\infty} \frac{1}{[[k]]_q!} [D_q^k f(x)] a^k \\ &= \sum_{k=0}^{\infty} \frac{1}{[[k]]_q!} \int_{-z.\infty}^{z.\infty} d_q x [D_q^k f(x)] a^k = \int_{-z.\infty}^{z.\infty} d_q x f(x). \end{aligned} \quad (71)$$

In the first step, we used Eq. (30) of Chap. 2. In the last step, the summands to $k > 0$ vanish due to Eq. (68). Similarly, it holds:

$$\int_{-z.\infty}^{z.\infty} d_q x f(a \oplus x) = \int_{-z.\infty}^{z.\infty} d_q x f(x). \quad (72)$$

Using the Jackson integral in Eq. (60), we can introduce the function $\Theta_q(z)$ ($0 < q < 1$):

$$\Theta_q(z) = \int_0^{z.\infty} d_q x x^{-1} \sin_q(x) = (1-q) \sum_{m=-\infty}^{\infty} \sin_q(q^m z). \quad (73)$$

Due to the definition of the improper q -integral in Eq. (60), it holds ($k \in \mathbb{Z}$):

$$\Theta_q(q^k z) = \Theta_q(z). \quad (74)$$

Next, we show that $\Theta_q(z)$ is related to the function

$$\mathbf{Q}(z; q) = (1-q) \sum_{m=-\infty}^{\infty} \frac{1}{zq^m + z^{-1}q^{-m}}. \quad (75)$$

To this end, we rewrite the series in Eq. (73) using the partial fraction decompositions from Eq. (46) of the previous chapter [1]:

$$\begin{aligned} \Theta_q(z) &= \frac{1-q}{(q; q)_{\infty}} \sum_{m=-\infty}^{\infty} q^m \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+3)/2} (1-q)z}{(q; q)_k [1 + (1-q)^2 z^2 q^{2(m+k)}]} \\ &= \frac{1-q}{(q; q)_{\infty}} \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+3)/2}}{(q; q)_k} \sum_{m=-\infty}^{\infty} \frac{(1-q)z q^m}{1 + (1-q)^2 z^2 q^{2(m+k)}} \\ &= \frac{1-q}{(q; q)_{\infty}} \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+1)/2}}{(q; q)_k} \sum_{m=-\infty}^{\infty} \frac{(1-q)z q^m}{1 + (1-q)^2 z^2 q^{2m}} \\ &= (1-q) \frac{E_q(-q)}{(q; q)_{\infty}} \sum_{m=-\infty}^{\infty} \frac{(1-q)z q^m}{1 + (1-q)^2 z^2 q^{2m}} = \mathbf{Q}((1-q)z; q). \end{aligned} \quad (76)$$

In the second step, we changed the order of summation, and in the next step, we replaced $m+k$ by m . The two final steps are a consequence of Eq. (17) in Chap. 2 and Eq. (75).

Next, we show that the following limit holds ($z \in \mathbb{R}^+$):

$$\lim_{q \rightarrow 1^-} \mathbf{Q}(z; q) = \frac{\pi}{2}. \quad (77)$$

In Eq. (75), we replace the deformation parameter q by e^{-h} ($h > 0$):

$$\begin{aligned} \mathbf{Q}(z; e^{-h}) &= (1 - e^{-h}) \sum_{m=-\infty}^{\infty} \frac{1}{z e^{-hm} + z^{-1} e^{hm}} \\ &= h \sum_{m=-\infty}^{\infty} \frac{z^{-1} e^{hm}}{1 + (z^{-1} e^{hm})^2} + O(h^2). \end{aligned} \quad (78)$$

We take the limit $h \rightarrow 0^+$ that corresponds to the limit $q \rightarrow 1$:

$$\begin{aligned} \lim_{h \rightarrow 0^+} \mathbf{Q}(z; e^{-h}) &= \lim_{h \rightarrow 0^+} \sum_{m=-\infty}^{\infty} \frac{z^{-1} e^{hm}}{1 + (z^{-1} e^{hm})^2} h = \int_{-\infty}^{\infty} \frac{z^{-1} e^x}{1 + (z^{-1} e^x)^2} dx \\ &= \int_{-\infty}^{\infty} \frac{d(z^{-1} e^x)}{1 + (z^{-1} e^x)^2} = \int_0^{\infty} \frac{dy}{1 + y^2} = \frac{1}{2} \int_{-\infty}^{\infty} \frac{dy}{1 + y^2} = \frac{\pi}{2}. \end{aligned} \quad (79)$$

In the above calculation, the series becomes an improper integral for $h \rightarrow 0^+$. We solved this improper integral using the substitution rule. Since the limit in Eq. (77) is independent of z , we also have:

$$\lim_{q \rightarrow 1^-} \Theta_q(z) = \lim_{q \rightarrow 1^-} \mathbf{Q}((1-q)z; q) = \frac{\pi}{2}. \quad (80)$$

Accordingly, Eq. (73) implies ($z \in \mathbb{R}^+$):

$$\lim_{q \rightarrow 1} \Theta_q(z) = \lim_{q \rightarrow 1} \int_0^{zq^{-M}} d_q x x^{-1} \sin_q(x) = \int_0^{\infty} dx x^{-1} \sin(x) = \frac{\pi}{2}. \quad (81)$$

Using the power series expansions in Eq. (44) of the previous chapter, we can compute **q -integrals of q -trigonometric functions**. For example, we find [1] ($0 < q < 1$, $M \in \mathbb{N}$):

$$\begin{aligned} \int_0^{zq^{-M}} d_q x \cos_q(x) &= (1-q)z \sum_{m=-M}^{\infty} q^m \cos_q(q^m z) \\ &= \sum_{m=-M}^{\infty} (1-q)z q^m \sum_{k=0}^{\infty} \frac{(-1)^k (1-q)^{2k} q^{2mk} z^{2k}}{(q; q)_{2k}} \\ &= \sum_{k=0}^{\infty} \frac{(-1)^k (1-q)^{2k+1} z^{2k+1}}{(q; q)_{2k}} \sum_{m=-M}^{\infty} q^{m(2k+1)} \\ &= \sum_{k=0}^{\infty} \frac{(-1)^k (1-q)^{2k+1} z^{2k+1}}{(q; q)_{2k}} \frac{(q^{-M})^{2k+1}}{1 - q^{2k+1}} \\ &= \sum_{k=0}^{\infty} \frac{(-1)^k (1-q)^{2k+1} (q^{-M} z)^{2k+1}}{(q; q)_{2k+1}} = \sin_q(q^{-M} z). \end{aligned} \quad (82)$$

In the third step of the above calculation, we used the following identities:

$$\begin{aligned} \sum_{m=-M}^{\infty} q^{m(2k+1)} &= \sum_{m=0}^{\infty} q^{(m-M)(2k+1)} = q^{-M(2k+1)} \sum_{m=0}^{\infty} q^{m(2k+1)} \\ &= \frac{q^{-M(2k+1)}}{1 - q^{2k+1}}. \end{aligned} \quad (83)$$

By similar reasoning, we can prove the following identity [1]:

$$\int_0^{zq^{-M}} d_q x \sin_q(x) = (1-q)z \sum_{m=-M}^{\infty} q^m \sin_q(q^m z) = 1 - \cos_q(q^{-M} z). \quad (84)$$

If M becomes infinite, Eqs. (82) and (84) imply

$$\begin{aligned} \int_0^{\infty} d_q z \cos_q(z) &= (1-q)z \lim_{M \rightarrow \infty} \sum_{m=-M}^{\infty} q^m \cos_q(q^m z) \\ &= \lim_{M \rightarrow \infty} \sin_q(q^{-M} z) = 0 \end{aligned} \quad (85)$$

and

$$\begin{aligned} \int_0^{\infty} d_q z \sin_q(z) &= (1-q)z \lim_{M \rightarrow \infty} \sum_{m=-M}^{\infty} q^m \sin_q(q^m z) \\ &= 1 - \lim_{M \rightarrow \infty} \cos_q(q^{-M} z) = 1. \end{aligned} \quad (86)$$

In the above calculations, the final identity holds since the q -trigonometric functions \sin_q and \cos_q vanish at infinity [cf. the inequalities in (49) of the previous chapter].

5 Distributions

$\mathcal{M}(x)$ denotes the set of functions with the domain being the following q -lattice:

$$\mathbb{G}_{q,x_0} = \{\pm x_0 q^m \mid m \in \mathbb{Z}\} \cup \{0\}. \quad (87)$$

Using the q -integral over $]-\infty, \infty[$ [cf. Eq. (63) of Chap. 4] we can introduce a q -scalar product on $\mathcal{M}(x)$ [16] [$f, g \in \mathcal{M}(x)$]:

$$\langle f, g \rangle_q = \int_{-x_0}^{x_0} d_q z \overline{f(z)} g(z) = \sum_{\varepsilon=\pm} \sum_{j=-\infty}^{\infty} (1-q) q^j x \overline{f(\varepsilon q^j x)} g(\varepsilon q^j x). \quad (88)$$

Here, \bar{f} denotes the complex conjugate of f . The q -scalar product induces a q -norm on $\mathcal{M}(x)$ [17, 18]:

$$\|f\|_q^2 = \langle f, f \rangle_q = \sum_{\varepsilon=\pm} \sum_{j=-\infty}^{\infty} (1-q) q^j x |f(\varepsilon q^j x)|^2. \quad (89)$$

The sum in the above equation takes the value zero if and only if $f(\pm q^m x) = 0$ for all $m \in \mathbb{Z}$. Any function that vanishes on the q -lattice can be identified with the zero function. This observations ensure that the q -scalar product is positive definite. The set of functions with finite q -norm forms a **q -version of the Hilbert space of square-integrable functions** [6, 19, 20]:

$$L_q^2 = \{f \mid \langle f, f \rangle_q < \infty\}. \quad (90)$$

Next, we consider all functions of $\mathcal{M}(x)$ that are continuous at the origin of the lattice \mathbb{G}_{q, x_0} ($0 < q < 1$):

$$\lim_{m \rightarrow \infty} f(\pm x_0 q^m) = f(0) \quad (91)$$

In addition, there is a set of positive constants $C_{k,l}(q)$ such that the following conditions hold ($k, l \in \mathbb{N}_0$):

$$|x^k D_q^l f(x)| \leq C_{k,l}(q) \quad \text{for } x \in \mathbb{G}_{q, x_0}. \quad (92)$$

These conditions imply:

$$\lim_{m \rightarrow \infty} (D_q^l f)(\pm x q^{-m}) = 0. \quad (93)$$

The functions of $\mathcal{M}(x)$ with the above properties make up the new set $\mathcal{S}(x)$, which is a **q -version of the space of test functions for tempered distributions**. For these q -deformed test functions, we introduce a family of norms:

$$\|f\|_{k,l} = \max_{x \in \mathbb{G}_{q, x_0}} |x^k D_q^l f(x)|. \quad (94)$$

This family of norms induces a topology on the set $\mathcal{S}(x)$ [1].

A continuous linear functional $l : \mathcal{S}(x) \rightarrow \mathbb{C}$ is called a **q -distribution**. A q -distribution l is called regular if there is a function $f \in \mathcal{M}(x)$ with the following property:

$$l(g) = \langle f, g \rangle_q = \int_{-\infty}^{\infty} d_q x \overline{f(x)} g(x). \quad (95)$$

We can use regular q -distributions to obtain new q -distributions. Let $f_n(x)$ be a sequence of functions that determine a sequence of regular q -distributions. If the limit

$$l(g) = \lim_{n \rightarrow \infty} \langle f_n, g \rangle_q \quad (96)$$

exists for all $g \in \mathcal{S}(x)$, the mapping $l : g \mapsto l(g)$ will be a q -distribution [1].

Let l be a regular q -distribution represented by a function f . We can calculate the Jackson derivatives of l as follows ($k \in \mathbb{N}$, $g \in \mathcal{S}(x)$):

$$\begin{aligned} (D_q^k l)(g) &= \int_{-\infty}^{\infty} d_q x \overline{D_q^k f(x)} g(x) = \int_{-\infty}^{\infty} d_q x \left[D_q^k \overline{f(x)} \right] g(x) \\ &= (-1)^k q^{-k(k-1)/2} \int_{-\infty}^{\infty} d_q x \overline{f(q^k x)} D_q^k g(x) \\ &= (-1)^k q^{-k(k+1)/2} \int_{-\infty}^{\infty} d_q x \overline{f(x)} (D_q^k g)(q^{-k} x). \end{aligned} \quad (97)$$

In the second last step, we used Eq. (70) of Chap. 4, and in the final step, we rescaled the integration variable. We can simplify the result of Eq. (97) by applying the following identity:

$$(D_q^k g)(q^{-k}x) = q^{k(k-1)/2} D_{q^{-1}}^k g(x). \quad (98)$$

This way, we obtain:

$$(D_q^k l)(g) = \int_{-\infty}^{\infty} d_q x \overline{f(x)} ((-q^{-1} D_{q^{-1}})^k g)(x) = l((-q^{-1} D_{q^{-1}})^k g). \quad (99)$$

By introducing the functions [1]

$$\phi_m^\varepsilon(x) = \begin{cases} 1 & \text{for } x = \varepsilon q^m x_0, \\ 0 & \text{for } x \neq \varepsilon q^m x_0, \end{cases} \quad (100)$$

we can define **step functions on the q -lattice**:

$$\theta_q^\pm(z) = \sum_{n=-\infty}^{\infty} \phi_n^\pm(z) \quad (101)$$

The functions θ_q^ε determine regular q -distributions ($g(x) \in \mathcal{S}(x)$, $0 < q < 1$):

$$\theta_q^\varepsilon(g) = \langle \theta_q^\varepsilon, g \rangle_q = \int_0^\infty d_q z g(\varepsilon z) = (1-q)x_0 \sum_{m=-\infty}^{\infty} q^m g(\varepsilon x_0 q^m). \quad (102)$$

Next, we define a **q -version of the delta distribution**:

$$\delta_q(g) = \langle \delta_q, g \rangle_q = \lim_{m \rightarrow \infty} \frac{g(x_0 q^m) + g(-x_0 q^m)}{2} = g(0). \quad (103)$$

As the function g is continuous at the origin, it holds

$$2\delta_q(g) = \delta_q^+(g) + \delta_q^-(g) \quad (104)$$

where ($\varepsilon \in \{+, -\}$)

$$\delta_q^\varepsilon(g) = \langle \delta_q^\varepsilon, g \rangle_q = \lim_{m \rightarrow \infty} g(\varepsilon x_0 q^m) = g(0^\varepsilon). \quad (105)$$

Using Eq. (99), we can calculate the **Jackson derivative for the q -distribution** θ_q^\pm ($0 < q < 1$, $g \in \mathcal{S}(x)$, $\varepsilon \in \{+, -\}$):

$$\begin{aligned} (D_q \theta_q^\varepsilon)(g) &= -q^{-1} \langle \theta_q^\varepsilon, D_{q^{-1}} g \rangle_q \\ &= -(q^{-1} - 1)x_0 \sum_{m=-\infty}^{\infty} q^m \frac{g(\varepsilon x_0 q^m) - g(\varepsilon x_0 q^{m-1})}{(1-q^{-1})\varepsilon x_0 q^m} \\ &= \varepsilon \sum_{m=-\infty}^{\infty} [g(\varepsilon x_0 q^m) - g(\varepsilon x_0 q^{m-1})] \\ &= \varepsilon \lim_{M \rightarrow \infty} \sum_{m=-M}^M [g(\varepsilon x_0 q^m) - g(\varepsilon x_0 q^{m-1})] \\ &= \varepsilon \lim_{M \rightarrow \infty} [g(\varepsilon x_0 q^M) - g(\varepsilon x_0 q^{-M-1})] = \varepsilon g(0^\varepsilon). \end{aligned} \quad (106)$$

In the last step of the calculation above, we used the boundary conditions in Eqs. (91) and (93). Due to $g(0^\varepsilon) = \delta_q^\varepsilon(g)$, the result in Eq. (106) gives us the following derivative rule:

$$(D_q \theta_q^\varepsilon)(g) = \varepsilon \delta_q^\varepsilon(g). \quad (107)$$

With this relation, we can represent both parts of the q -delta distribution as the limit of regular q -distributions. To this end, we proceed as follows:

$$\delta_q^\varepsilon(g) = \varepsilon (D_q \theta_q^\varepsilon)(g) = \langle \varepsilon D_q \theta_q^\varepsilon, g \rangle_q = \int_{-\infty}^{\infty} d_q x \overline{\varepsilon D_q \theta_q^\varepsilon(x)} g(x). \quad (108)$$

It remains to calculate the Jackson derivative for $\varepsilon \theta_q^\varepsilon(z)$:

$$\begin{aligned} \varepsilon D_q \theta_q^\varepsilon(x) &= \varepsilon \sum_{n=-\infty}^{\infty} D_q \phi_n^\varepsilon(x) = \varepsilon \sum_{n=-\infty}^{\infty} \frac{\phi_n^\varepsilon(x) - \phi_n^\varepsilon(qx)}{(1-q)x} \\ &= \varepsilon (1-q)^{-1} x^{-1} \sum_{n=-\infty}^{\infty} (\phi_n^\varepsilon(x) - \phi_{n-1}^\varepsilon(x)) \\ &= \varepsilon (1-q)^{-1} x^{-1} \lim_{N \rightarrow \infty} \sum_{n=-N}^N (\phi_n^\varepsilon(x) - \phi_{n-1}^\varepsilon(x)) \\ &= \varepsilon (1-q)^{-1} x^{-1} \lim_{N \rightarrow \infty} (\phi_N^\varepsilon(x) - \phi_{-N-1}^\varepsilon(x)). \end{aligned} \quad (109)$$

Plugging this result into Eq. (108), we finally get:

$$\begin{aligned} \delta_q^\varepsilon(g) &= \lim_{N \rightarrow \infty} \int_{-\infty}^{\infty} d_q x \frac{\varepsilon (\phi_N^\varepsilon(x) - \phi_{-N-1}^\varepsilon(x))}{(1-q)x} g(x) \\ &= \lim_{N \rightarrow \infty} \int_{-\infty}^{\infty} d_q x \frac{\varepsilon \phi_N^\varepsilon(x)}{(1-q)x} g(x) - \lim_{N \rightarrow \infty} \int_{-\infty}^{\infty} d_q x \frac{\varepsilon \phi_{-N-1}^\varepsilon(x)}{(1-q)x} g(x) \\ &= \lim_{N \rightarrow \infty} \int_{-\infty}^{\infty} d_q x \frac{\varepsilon \phi_N^\varepsilon(x)}{(1-q)x} g(x). \end{aligned} \quad (110)$$

In the final step, we have omitted the second q -integral since it vanishes due to Eq. (93).

Next, we want to find the **Jackson derivatives of the q -delta distribution**. To this end, we perform the following calculation ($k \in \mathbb{N}$):

$$\begin{aligned} (D_q^k \delta_q^\varepsilon)(g) &= \varepsilon D_q^{k+1} \theta_q^\varepsilon(g) = \varepsilon (-q^{-1})^k (D_q \theta_q^\varepsilon)(D_{q^{-1}}^k g) \\ &= (-q^{-1})^k \delta_q^\varepsilon(D_{q^{-1}}^k g) = (-q^{-1})^k (D_{q^{-1}}^k g)(0^\varepsilon). \end{aligned} \quad (111)$$

The first two identities follow from Eq. (108) or Eq. (99). Due to Eq. (104), the above result yields the following formulas:

$$(D_q^k \delta_q)(g) = (-q^{-1})^k \delta_q(D_{q^{-1}}^k g) = (-q^{-1})^k (D_{q^{-1}}^k g)(0). \quad (112)$$

We want to study how the q -delta distribution behaves under q -translations. We assume that we have a function $\delta_q(x)$ with the following property:

$$\delta_q(g) = \int_{-\infty}^{\infty} d_q x \delta_q(x) g(x) = g(0). \quad (113)$$

Using the q -Taylor formula in Eqs. (35) and (36) of Chap. 2, we get:

$$\begin{aligned} \delta_q(x \bar{\oplus} (\bar{\ominus} a)) &= \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{[[k]]_q!} (D_q^k \delta_q(x)) (-a)^k, \\ \delta_q((\bar{\ominus} a) \bar{\oplus} x) &= \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{[[k]]_q!} (-a)^k D_q^k \delta_q(x). \end{aligned} \quad (114)$$

Using the expressions above, we do the following calculation :

$$\begin{aligned} \int_{-\infty}^{\infty} d_q x \delta_q((\bar{\ominus} qa) \bar{\oplus} x) f(x) &= \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{[[k]]_q!} (-qa)^k \int_{-\infty}^{\infty} d_q x (D_q^k \delta_q(x)) f(x) \\ &= \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{[[k]]_q!} (-qa)^k (-1)^k q^{-k(k-1)/2} \int_{-\infty}^{\infty} d_q x \delta_q(q^k x) D_q^k f(x) \\ &= \sum_{k=0}^{\infty} \frac{1}{[[k]]_q!} a^k \int_{-\infty}^{\infty} d_q x \delta_q(x) (D_q^k f)(q^{-k} x) \\ &= \sum_{k=0}^{\infty} \frac{1}{[[k]]_q!} a^k (D_q^k f)(0) = f(a). \end{aligned} \quad (115)$$

In the second step, we employed the rules for integration by parts [cf. Eq. (70) of the previous chapter]. In the third step, we changed the scaling of the integration variable. The second last step follows from Eq. (113), and the final step results from Eqs. (29) and (33) of Chap. 2. In the same way, we can derive the following identity:

$$\int_{-\infty}^{\infty} d_q x f(x) \delta_q(x \bar{\oplus} (\bar{\ominus} qa)) = f(a). \quad (116)$$

The q -delta functions are related to the functions $\phi_m^{\pm}(x)$ in Eq. (100). Concretely, the identities in Eqs. (115) and (116) are satisfied for $a \in \mathbb{G}_{q, x_0}$ if it holds ($0 < q < 1$):

$$\delta_q((\bar{\ominus} qa) \bar{\oplus} x)|_{a=\pm x_0 q^m} = \delta_q(x \bar{\oplus} (\bar{\ominus} qa))|_{a=\pm x_0 q^m} = \frac{\phi_m^{\pm}(x)}{(1-q)x_0 q^m}. \quad (117)$$

The functions $\phi_m^{\pm}(x)$ can be represented by infinitely differentiable functions. To this end, we introduce the following function:

$$\psi(x, a) = \begin{cases} 0 & \text{if } x < -a, \\ e^{-e^{x/(x^2-a^2)}} a^2 e & \text{if } -a \leq x \leq a, \\ a^2 e & \text{if } a < x. \end{cases} \quad (118)$$

We calculate the partial derivative of $\psi(x, a)$ with respect to x :

$$\frac{\partial}{\partial x}\psi(x, a) = \begin{cases} 0 & \text{if } x < -a, \\ e^{x/(x^2-a^2)-e^{x/(x^2-a^2)}} \frac{x^2+a^2}{(x^2-a^2)^2} a^2 e & \text{if } -a \leq x \leq a, \\ 0 & \text{if } a < x. \end{cases} \quad (119)$$

This function is infinitely differentiable with respect to x and has a bell-shaped graph that is symmetrical to the axis $x = 0$ [21]. Accordingly, $\partial_x \psi(x, a)$ takes its maximum value at $x = 0$, namely $\partial_x \psi(0, a) = 1$. Furthermore, we define a function that assigns to y half the distance from y to the next lattice point qy or $q^{-1}y$:

$$b_q(y) = \frac{1}{2} \min \{ |(1 - q^{-1})y|, |(1 - q)y| \}. \quad (120)$$

Using $\partial_x \psi(x, a)$ and $b_q(y)$, we can introduce the following function:

$$\phi_q(x, y) = \frac{\partial}{\partial x} \psi(x - y, b_q(y)). \quad (121)$$

This function has the properties

$$\phi_q(x, x) = 1 \quad \text{for } x \in \mathbb{R} \quad (122)$$

and

$$\phi_q(x, y) = 0 \quad \text{for } x \in \mathbb{R} \setminus [y - b_q(y), y + b_q(y)]. \quad (123)$$

This enables us to identify the function $\phi_m^\varepsilon(x)$ on the one-dimensional q -lattice \mathbb{G}_{q, x_0} with an infinitely differentiable function:

$$\phi_m^\varepsilon(x) = \phi_q(x, \varepsilon x_0 q^m). \quad (124)$$

Finally, the result above leads to the following representations [cf. Eq. (117)]:

$$\delta_q((\bar{\ominus} qa) \bar{\oplus} x) = \delta_q(x \bar{\oplus} (\bar{\ominus} qa)) = \frac{\phi_q(x, a)}{(1 - q)|a|} = \frac{\phi_q(x, a)}{(1 - q)|x|}. \quad (125)$$

The step functions θ_q^ε given in Eq. (101) are q -periodic, as we can see from the following calculation ($\varepsilon \in \{+, -\}$):

$$\theta_q^\varepsilon(qz) = \sum_{n=-\infty}^{\infty} \phi_n^\varepsilon(qz) = \sum_{n=-\infty}^{\infty} \phi_{n-1}^\varepsilon(z) = \sum_{n=-\infty}^{\infty} \phi_n^\varepsilon(z) = \theta_q^\varepsilon(z). \quad (126)$$

If we introduce a scaling operator Λ with

$$\Lambda f(x) = f(qx), \quad (127)$$

Eq. (126) implies that the step functions θ_q^ε are invariant under the action of Λ ($n \in \mathbb{N}$, $\varepsilon \in \{+, -\}$):

$$\Lambda^n \theta_q^\varepsilon(x) = \theta_q^\varepsilon(q^n x) = \theta_q^\varepsilon(x). \quad (128)$$

Under the action of the scaling operator, the q -delta function behaves as follows:

$$\Lambda^n \delta_q(x) = \delta_q(q^n x) = q^{-n} \delta_q(x). \quad (129)$$

We prove these identities by the following calculation:

$$\begin{aligned} & \int_{-\infty}^{\infty} d_q x [\Lambda^n \delta_q(x)] g(x) = \int_{-\infty}^{\infty} d_q x \delta_q(q^n x) g(x) \\ & = q^{-n} \int_{-\infty}^{\infty} d_q x \delta_q(x) g(q^{-n} x) = q^{-n} g(0) = \int_{-\infty}^{\infty} d_q p q^{-n} \delta_q(x) g(x). \end{aligned} \quad (130)$$

The result above, which holds for any function $g \in \mathcal{S}(x)$, confirms the identity in Eq. (129).

We regain the shifted q -delta functions given in Eq. (125) by applying a Jackson derivative to a step function on the q -lattice. To show this, we perform the following calculation [$g(x) \in \mathcal{S}(x)$, $\varepsilon \in \{+, -\}$, $m \in \mathbb{Z}$]:

$$\begin{aligned} & \int_{-\infty}^{\infty} d_q x g(x) D_q \sum_{n=-\infty}^m \phi_n^\varepsilon(x) = \int_{-\infty}^{\infty} d_q x g(x) \sum_{n=-\infty}^m \frac{\phi_n^\varepsilon(x) - \phi_n^\varepsilon(qx)}{(1-q)x} \\ & = \int_{-\infty}^{\infty} d_q x \frac{g(x)}{(1-q)x} \sum_{n=-\infty}^m [\phi_n^\varepsilon(x) - \phi_{n-1}^\varepsilon(x)] \\ & = \int_{-\infty}^{\infty} d_q x g(x) \frac{\phi_m^\varepsilon(x)}{(1-q)x} - \lim_{k \rightarrow -\infty} \int_{-\infty}^{\infty} d_q x g(x) \frac{\phi_k^\varepsilon(x)}{(1-q)x} \\ & = \varepsilon g(\varepsilon x_0 q^m) - \varepsilon \lim_{k \rightarrow -\infty} g(\varepsilon x_0 q^k) = \varepsilon g(\varepsilon x_0 q^m). \end{aligned} \quad (131)$$

Moreover, Eq. (116) implies:

$$\int_{-\infty}^{\infty} d_q x g(x) \delta_q(x \bar{\oplus} (\bar{\ominus} qa))|_{a=\varepsilon x_0 q^m} = g(\varepsilon x_0 q^m). \quad (132)$$

Thus, the following identity holds in the set of regular q -distributions:

$$D_q \sum_{n=-\infty}^m \phi_n^\varepsilon(x) = \varepsilon \delta_q(x \bar{\oplus} (\bar{\ominus} qa))|_{a=\varepsilon x_0 q^m}. \quad (133)$$

We can identify the sum on the left-hand side of the above equation with a shifted step function on the q -lattice:

$$\sum_{n=-\infty}^m \phi_n^\varepsilon(x) = \theta_q^\varepsilon(x \bar{\oplus} (\bar{\ominus} qa))|_{a=\varepsilon x_0 q^m}. \quad (134)$$

Combining Eq. (133) and Eq. (134), we finally obtain:

$$D_{q,x} \theta_q^\varepsilon(x \bar{\oplus} (\bar{\ominus} qa))|_{a=\varepsilon x_0 q^m} = \varepsilon \delta_q(x \bar{\oplus} (\bar{\ominus} qa))|_{a=\varepsilon x_0 q^m}. \quad (135)$$

To verify the identity in Eq. (134), we do the following calculation:

$$\begin{aligned}
\int_{-x_0.∞}^{x_0.∞} d_q x \sum_{n=-∞}^m \phi_n^\varepsilon(x) g(x) &= (1-q) \sum_{j=-∞}^∞ x_0 q^j \sum_{n=-∞}^m \phi_n^\varepsilon(\varepsilon x_0 q^j) g(\varepsilon x_0 q^j) \\
&= (1-q) \sum_{n=-∞}^m x_0 q^n g(\varepsilon x_0 q^n) = (1-q) \sum_{n=-∞}^0 x_0 q^{m+n} g(\varepsilon x_0 q^{m+n}) \\
&= (1-q) \sum_{n=1}^∞ x_0 q^{m+1-n} g(\varepsilon x_0 q^{m+1-n}) = \varepsilon \int_{\varepsilon x_0 q^{m+1}}^{\varepsilon x_0.∞} d_q x g(x). \quad (136)
\end{aligned}$$

Similarly to Eq. (115), we have:

$$\begin{aligned}
\int_{-x_0.∞}^{x_0.∞} d_q x g(x) \theta_q^\varepsilon(x \oplus (\ominus qa)) &= \\
&= \sum_{k=0}^∞ \frac{q^{k(k-1)/2}}{[[k]]_q!} (-qa)^k \int_{-x_0.∞}^{x_0.∞} d_q x g(x) D_q^k \theta_q^\varepsilon(x) \\
&= \sum_{k=0}^∞ \frac{(qa)^k}{[[k]]_q!} \int_{-x_0.∞}^{x_0.∞} d_q x \theta_q^\varepsilon(q^k x) D_q^k g(x) \\
&= \int_{-x_0.∞}^{x_0.∞} d_q x \theta_q^\varepsilon(x) g(x \oplus qa) = \varepsilon \int_0^{\varepsilon x_0.∞} d_q x g(x \oplus qa). \quad (137)
\end{aligned}$$

We rewrite the last expression of Eq. (137) as follows [4]:

$$\varepsilon \int_0^{\varepsilon x_0.∞} d_q x g(x \oplus qa) = \varepsilon \int_{\ominus qa}^{\varepsilon x_0.∞} d_q x g(x) = \varepsilon \int_{qa}^{\varepsilon x_0.∞} d_q x g(x). \quad (138)$$

Putting the results in Eqs. (136)-(138) together, we get the identity from which we can read off Eq. (134):

$$\begin{aligned}
\int_{-x_0.∞}^{x_0.∞} d_q x \sum_{n=-∞}^m \phi_n^\varepsilon(x) g(x) &= \varepsilon \int_{\varepsilon x_0 q^{m+1}}^{\varepsilon x_0.∞} d_q x g(x) \\
&= \int_{-x_0.∞}^{x_0.∞} d_q x g(x) \theta_q^\varepsilon(x \oplus (\ominus qa)) \Big|_{a=\varepsilon x_0 q^m}. \quad (139)
\end{aligned}$$

6 Fourier transforms

In this chapter, we consider q -versions of one-dimensional Fourier transforms. The q -deformed Fourier transforms are mappings between the function spaces $\mathcal{M}_R(x)$ and $\mathcal{M}(p)$. In this respect, x and p play the role of position and momentum variables.

We use the q -deformed exponential in Eq. (15) of Chap. 2 to introduce q -versions of one-dimensional plane waves [15]:

$$\exp_q(x|ip) = \sum_{k=0}^∞ \frac{x^k (ip)^k}{[[k]]_q!}. \quad (140)$$

With the help of the Jackson derivative D_q , we introduce the momentum operator

$$\hat{P} = i^{-1}D_q. \quad (141)$$

The q -deformed plane waves in Eq. (140) are eigenfunctions of \hat{P} :

$$i^{-1}D_{q,x} \triangleright \exp_q(x|ip) = \sum_{k=1}^{\infty} \frac{[[k]]}{[[k]]_q!} x^{k-1} (ip)^{k-1} p \quad (142)$$

$$= \exp_q(x|ip)p. \quad (143)$$

Additionally, we need the following function:

$$\begin{aligned} \exp_{q^{-1}}(-ip|x) &= \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2} (-ip)^k x^k}{[[k]]_q!} \\ &= \sum_{k=0}^{\infty} \frac{(-ip)^k x^k}{[[k]]_{q^{-1}}!}. \end{aligned} \quad (144)$$

The momentum operator acts on this function as follows:

$$\begin{aligned} i^{-1}D_{q,x} \triangleright \exp_{q^{-1}}(-ip|x) &= i^{-1} \sum_{k=1}^{\infty} \frac{(-i)^k q^{k(k-1)/2}}{[[k-1]]_q!} p^k x^{k-1} \\ &= -\exp_{q^{-1}}(-ip|qx)p. \end{aligned} \quad (145)$$

As was shown in Ref. [1], the q -deformed plane waves satisfy **completeness relations and orthogonality conditions**, given by

$$\int_{-x_0.\infty}^{x_0.\infty} d_q p \exp_q(x_0|ip) \exp_{q^{-1}}(-ip|qx) = \begin{cases} \frac{4\Theta_0}{(1-q)x_0} & \text{für } x = x_0, \\ 0 & \text{für } x \neq x_0, \end{cases} \quad (146)$$

$$\int_{-p_0.\infty}^{p_0.\infty} d_q x \exp_{q^{-1}}(-ip_0|qx) \exp_q(x|ip) = \begin{cases} \frac{4\Theta_0}{(1-q)p_0} & \text{für } p = p_0, \\ 0 & \text{für } p \neq p_0, \end{cases} \quad (147)$$

with [cf. Eq. (73) in Chap. 4]

$$\Theta_0 = \Theta_q(x_0 p_0). \quad (148)$$

Next, we introduce q -Fourier transforms between $\mathcal{M}(x)$ and $\mathcal{M}(p)$ [1, 15]:

$$\begin{aligned} \mathcal{F}_q(\phi)(p) &= \frac{1}{2\sqrt{\Theta_0}} \int_{-x_0.\infty}^{x_0.\infty} d_q x \phi(x) \exp_q(x|ip), \\ \mathcal{F}_q^{-1}(\psi)(x) &= \frac{1}{2\sqrt{\Theta_0}} \int_{-p_0.\infty}^{p_0.\infty} d_q p \psi(p) \exp_{q^{-1}}(-ip|qx). \end{aligned} \quad (149)$$

We show that the mappings \mathcal{F}_q and \mathcal{F}_q^{-1} are inverse to each other. Let ϕ_n^\pm and ψ_n^\pm with $n \in \mathbb{Z}$ denote the basis functions of $\mathcal{M}(x)$ and $\mathcal{M}(p)$, respectively [cf. Eq. (100) of the previous chapter]. It suffices to prove the following identities:

$$\begin{aligned} (\mathcal{F}_q^{-1} \circ \mathcal{F}_q)(\phi_n^\pm) &= \phi_n^\pm, \\ (\mathcal{F}_q \circ \mathcal{F}_q^{-1})(\psi_n^\pm) &= \psi_n^\pm. \end{aligned} \quad (150)$$

To prove the first of the two identities above, we do the following calculation:

$$\begin{aligned} (\mathcal{F}_q^{-1} \circ \mathcal{F}_q)(\phi_n^\pm(x)) &= \\ &= \frac{1}{4\Theta_0} \int_{-p_0.\infty}^{p_0.\infty} d_q p \int_{-x_0.\infty}^{x_0.\infty} d_q \xi \phi_n^\pm(\xi) \exp_q(\xi|ip) \exp_{q^{-1}}(-ip|qx) \\ &= \frac{(1-q)x_0 q^n}{4\Theta_0} \int_{-p_0.\infty}^{p_0.\infty} d_q p \exp_q(\pm x_0 q^n | ip) \exp_{q^{-1}}(-ip|qx) \\ &= \frac{(1-q)x_0}{4\Theta_0} \int_{-p_0.\infty}^{p_0.\infty} d_q p \exp_q(x_0|ip) \exp_{q^{-1}}(-ip|q(\pm q^{-n})x). \end{aligned} \quad (151)$$

This result, together with Eq. (146), implies:

$$(\mathcal{F}_q^{-1} \circ \mathcal{F}_q)(\phi_n^\pm(x)) = \begin{cases} 1 & \text{for } x = \pm x_0 q^n, \\ 0 & \text{for } x \neq \pm x_0 q^n. \end{cases} \quad (152)$$

We can verify the second identity of Eq. (150) in the same manner:

$$\begin{aligned} (\mathcal{F}_q \circ \mathcal{F}_q^{-1})(\psi_n^\pm(p)) &= \\ &= \frac{1}{4\Theta_0} \int_{-x_0.\infty}^{x_0.\infty} d_q x \int_{-p_0.\infty}^{p_0.\infty} d_q p' \psi_n^\pm(p') \exp_{q^{-1}}(-ip'|qx) \exp_q(x|ip) \\ &= \frac{(1-q)p_0 q^n}{4\Theta_0} \int_{-x_0.\infty}^{x_0.\infty} d_q x \exp_{q^{-1}}(-i(\pm q^n)p_0|qx) \exp_q(x|ip) \\ &= \frac{(1-q)p_0}{4\Theta_0} \int_{-x_0.\infty}^{x_0.\infty} d_q x \exp_{q^{-1}}(-ip_0|qx) \exp_q(x|\pm iq^{-n}p). \end{aligned} \quad (153)$$

Taking into account Eq. (147), we end up with:

$$(\mathcal{F}_q \circ \mathcal{F}_q^{-1})(\psi_n^\pm(p)) = \begin{cases} 1 & \text{for } p = \pm p_0 q^n, \\ 0 & \text{for } p \neq \pm p_0 q^n. \end{cases} \quad (154)$$

For $x \in \mathbb{G}_{q,x_0}$ and $p \in \mathbb{G}_{q,p_0}$, Eqs. (146) and (147) become ($n, m \in \mathbb{Z}$ and $\varepsilon, \varepsilon' = \pm 1$)

$$\begin{aligned} \int_{-p_0.\infty}^{p_0.\infty} d_q p \exp_q(\varepsilon x_0 q^n | ip) \exp_{q^{-1}}(-ip|q\varepsilon' x_0 q^m) &= \\ &= q^{-n} \int_{-p_0.\infty}^{p_0.\infty} d_q p \exp_q(x_0|ip) \exp_{q^{-1}}(-ip|q\varepsilon\varepsilon' x_0 q^{m-n}) \\ &= \frac{4\Theta_0}{(1-q)x_0 q^n} \delta_{nm} \delta_{\varepsilon\varepsilon'} \end{aligned} \quad (155)$$

or

$$\begin{aligned}
& \int_{-x_0.\infty}^{x_0.\infty} d_q x \exp_{q^{-1}}(-i\varepsilon p_0 q^n | qx) \exp_q(x | i\varepsilon' p_0 q^m) = \\
& = q^{-n} \int_{-x_0.\infty}^{x_0.\infty} d_q x \exp_{q^{-1}}(-ip_0 | qx) \exp_q(x | i\varepsilon' p_0 q^{m-n}) \\
& = \frac{4\Theta_0}{(1-q)p_0 q^n} \delta_{nm} \delta_{\varepsilon\varepsilon'}. \tag{156}
\end{aligned}$$

We get the second integral expression in the equations above by rescaling the integration variable. Next, we rewrite the above identities using the following functions:

$$u_p(x) = \frac{1}{2\sqrt{\Theta_0}} \exp_q(x | ip), \quad u_p^*(x) = \frac{1}{2\sqrt{\Theta_0}} \exp_{q^{-1}}(-ip | qx). \tag{157}$$

The *completeness relation* in Eq. (155) yields

$$\begin{aligned}
& \int_{-p_0.\infty}^{p_0.\infty} d_q p u_p(\varepsilon x_0 q^n) u_p^*(\varepsilon' x_0 q^m) = \frac{\delta_{nm} \delta_{\varepsilon\varepsilon'}}{(1-q)x_0 q^n} \\
& = \delta_q(x \oplus (\bar{\ominus} qy))|_{x=\varepsilon x_0 q^n; y=\varepsilon' x_0 q^m}. \tag{158}
\end{aligned}$$

Similarly, the *orthonormality relation* in Eq. (156) takes on the following form:

$$\begin{aligned}
& \int_{-x_0.\infty}^{x_0.\infty} d_q x u_{\varepsilon p_0 q^n}^*(x) u_{\varepsilon' p_0 q^m}(x) = \frac{\delta_{nm} \delta_{\varepsilon\varepsilon'}}{(1-q)p_0 q^n} \\
& = \delta_q((\bar{\ominus} qp) \oplus p')|_{p=\varepsilon p_0 q^n; p'=\varepsilon' p_0 q^m}. \tag{159}
\end{aligned}$$

In writing down the above relations, we used the expressions for the q -delta functions in Eq. (117) of the previous chapter and took into account the following identities [cf. Eq. (100) of the previous chapter]:

$$\phi_m^\varepsilon(\varepsilon' x_0 q^n) = \delta_{mn} \delta_{\varepsilon\varepsilon'}, \quad \psi_m^\varepsilon(\varepsilon' p_0 q^n) = \delta_{mn} \delta_{\varepsilon\varepsilon'}. \tag{160}$$

Due to the completeness relation in Eq. (158), we can expand a function $f(x)$ in position space $\mathcal{M}(x)$ in terms of the functions $u_p^*(x)$:

$$f(x) = \int_{-p_0.\infty}^{p_0.\infty} d_q p a_p u_p^*(x) \quad \text{with} \quad a_p = \mathcal{F}_q(f)(p). \tag{161}$$

Similarly, we can expand a function $g(p)$ in position space $\mathcal{M}(p)$ in terms of the functions $u_p(x)$:

$$g(p) = \int_{-x_0.\infty}^{x_0.\infty} d_q x b_x u_p(x) \quad \text{with} \quad b_x = \mathcal{F}_q^{-1}(g)(x). \tag{162}$$

Next, we consider a scaling operator that acts on functions in position or momentum space as follows:

$$\Lambda f(x) = f(qx), \quad \Lambda g(p) = g(q^{-1}p), \tag{163}$$

We show that the scaling operator Λ commutes with q -Fourier transforms as follows [1]:

$$\mathcal{F}_q \Lambda = q^{-1} \Lambda \mathcal{F}_q, \quad \mathcal{F}_q^{-1} \Lambda = q \Lambda \mathcal{F}_q^{-1}. \quad (164)$$

We prove the first relation by the following calculation:

$$\begin{aligned} \mathcal{F}_q(\Lambda\phi(x))(p) &= \mathcal{F}_q(\phi(qx))(p) \\ &= \frac{1}{2\sqrt{\Theta_0}} \int_{-x_0.\infty}^{x_0.\infty} d_q x \phi(qx) \exp_q(x|ip) \\ &= \frac{q^{-1}}{2\sqrt{\Theta_0}} \int_{-x_0.\infty}^{x_0.\infty} d_q x \phi(x) \exp_q(x|iq^{-1}p) \\ &= q^{-1}(\mathcal{F}_q \phi)(q^{-1}p) = q^{-1} \Lambda(\mathcal{F}_q \phi)(p). \end{aligned} \quad (165)$$

Similar considerations lead to the second relation in Eq. (164):

$$\begin{aligned} \mathcal{F}_q^{-1}(\Lambda\psi(p))(x) &= \mathcal{F}_q^{-1}(\psi(q^{-1}p))(x) \\ &= \frac{1}{2\sqrt{\Theta_0}} \int_{-p_0.\infty}^{p_0.\infty} d_q p \psi(q^{-1}p) \exp_{q^{-1}}(-ip|qx) \\ &= \frac{q}{2\sqrt{\Theta_0}} \int_{-p_0.\infty}^{p_0.\infty} d_q p \psi(p) \exp_{q^{-1}}(-ip|qqx) \\ &= q \mathcal{F}_q^{-1}(\psi)(qx) = q \Lambda \mathcal{F}_q^{-1}(\psi)(x). \end{aligned} \quad (166)$$

As is well known, ordinary Fourier transforms turn derivative operators and multiplication operators into each other. The q -deformed Fourier transforms show this property, as well [1], since we have

$$\mathcal{F}_q(D_{q,x}\phi) = -i\Lambda p \mathcal{F}_q(\phi), \quad \mathcal{F}_q(x\phi) = -iD_{q,p} \mathcal{F}_q(\phi) \quad (167)$$

and

$$\mathcal{F}_q^{-1}(D_{q,p}\psi) = i\mathcal{F}_q^{-1}(\psi)x, \quad \mathcal{F}_q^{-1}(p\psi) = iq^{-1}\Lambda^{-1}D_{q,x}\mathcal{F}_q^{-1}(\psi). \quad (168)$$

We only prove Eq. (168) since Eq. (167) can be proven by similar reasoning. The first identity in Eq. (168) can be verified as follows:

$$\begin{aligned} \mathcal{F}_q^{-1}(D_{q,p}\psi) &= \frac{1}{2\sqrt{\Theta_0}} \int_{-p_0.\infty}^{p_0.\infty} d_q p [D_{q,p}\psi(p)] \exp_{q^{-1}}(-ip|qx) \\ &= -\frac{1}{2\sqrt{\Theta_0}} \int_{-p_0.\infty}^{p_0.\infty} d_q p \psi(qp) D_{q,p} \exp_{q^{-1}}(-ip|qx) \\ &= \frac{iq}{2\sqrt{\Theta_0}} \int_{-p_0.\infty}^{p_0.\infty} d_q p \psi(qp) \exp_{q^{-1}}(-iqp|qx) x \\ &= \frac{i}{2\sqrt{\Theta_0}} \int_{-p_0.\infty}^{p_0.\infty} d_q p \psi(p) \exp_{q^{-1}}(-ip|qx) x \\ &= i\mathcal{F}_q^{-1}(\psi)x. \end{aligned} \quad (169)$$

To check the second identity in Eq. (168), we proceed as follows:

$$\begin{aligned}
\mathcal{F}_q^{-1}(p\psi) &= \frac{1}{2\sqrt{\Theta_0}} \int_{-p_0.\infty}^{p_0.\infty} d_q p p \psi(p) \exp_{q^{-1}}(-ip|qx) \\
&= \frac{i}{2\sqrt{\Theta_0}} \int_{-p_0.\infty}^{p_0.\infty} d_q p \psi(p) D_{q,x} \exp_{q^{-1}}(-ip|x) \\
&= iD_{q,x} \mathcal{F}_q^{-1}(\psi)(q^{-1}x) = iq^{-1}\Lambda^{-1}D_{q,x} \mathcal{F}_q^{-1}(\psi)(x). \tag{170}
\end{aligned}$$

In the two computations above, we have applied Eq. (145).

The ordinary Fourier transform is a unitary operator. Thus, the scalar product of two given functions has the same value as the scalar product of their Fourier transforms. In this respect, the q -deformed Fourier transforms in Eq. (149) are unitary maps on the Hilbert space L_q^2 with the scalar product in Eq. (88) of the previous chapter. However, the q -deformed Fourier transforms of the two factors of the q -scalar product slightly differ from each other. Concretely, we have

$$\langle g, f \rangle_{q,x} = \langle \psi, \phi \rangle_{q,p} \tag{171}$$

with

$$\begin{aligned}
\phi(p) &= \mathcal{F}_q(f)(p) = \frac{1}{2\sqrt{\Theta_0}} \int_{-x_0.\infty}^{x_0.\infty} d_q x \exp_q(ip|x) f(x), \\
\psi(p) &= \tilde{\mathcal{F}}_q(g)(p) = \frac{1}{2\sqrt{\Theta_0}} \int_{-x_0.\infty}^{x_0.\infty} d_q x \exp_{q^{-1}}(ip|qx) g(x), \tag{172}
\end{aligned}$$

and

$$\begin{aligned}
f(x) &= \mathcal{F}_q^{-1}(\phi)(x) = \frac{1}{2\sqrt{\Theta_0}} \int_{-p_0.\infty}^{p_0.\infty} d_q p \phi(p) \exp_{q^{-1}}(-ip|qx), \\
g(x) &= \tilde{\mathcal{F}}_q^{-1}(\psi)(x) = \frac{1}{2\sqrt{\Theta_0}} \int_{-p_0.\infty}^{p_0.\infty} d_q p \exp_q(x|-ip) \psi(p). \tag{173}
\end{aligned}$$

To prove Eq. (171), we insert the expressions for the q -Fourier transforms into one of the two scalar products and apply Eq. (155) or Eq. (156).

Our aim is to consider the **q -Fourier transforms of elementary functions and q -distributions**. First, we calculate the **q -Fourier transform of x^{-1}** :

$$\begin{aligned}
\mathcal{F}_q(x^{-1})(p) &= \frac{1}{2\sqrt{\Theta_0}} \int_{-x_0.\infty}^{x_0.\infty} d_q x x^{-1} \exp_q(x|ip) = \\
&= \frac{1-q}{2\sqrt{\Theta_0}} \sum_{\varepsilon=\pm} \sum_{m=-\infty}^{\infty} \varepsilon x_0 q^m x_0^{-1} q^{-m} \exp_q(\varepsilon x_0 q^m | ip) \\
&= \frac{1-q}{2\sqrt{\Theta_0}} \sum_{m=-\infty}^{\infty} [\exp_q(ix_0 q^m p) - \exp_q(-ix_0 q^m p)] \\
&= \frac{i(1-q)}{\sqrt{\Theta_0}} \sum_{m=-\infty}^{\infty} \sin_q(x_0 q^m p) = \frac{i}{\sqrt{\Theta_0}} \Theta_q(x_0 p). \tag{174}
\end{aligned}$$

For the final step in the above computation, we applied Eq. (73) of Chap. 4. We can simplify the final expression in Eq. (174) since p can only take the values $\pm p_0 q^m$ with $m \in \mathbb{Z}$:

$$\begin{aligned} \frac{i}{\sqrt{\Theta_0}} \Theta_q(\pm x_0 p_0 q^m) &= \pm \frac{i}{\sqrt{\Theta_0}} \Theta_q(x_0 p_0 q^m) = \pm \frac{i}{\sqrt{\Theta_0}} \Theta_q(x_0 p_0) \\ &= \pm \frac{i}{\sqrt{\Theta_0}} \Theta_0 = \pm i\sqrt{\Theta_0}. \end{aligned} \quad (175)$$

In the first step, we took into account that Θ_q is an odd function due to its definition [cf. Eq. (73) of Chap. 4]:

$$\Theta_q(-z) = -\Theta_q(z). \quad (176)$$

In the second step, we used the fact that Θ_q is also a q -periodic function [cf. Eq. (74) of Chap. 4]. The penultimate step follows from Eq. (148). Combining the results of Eqs. (174) and (175), we finally obtain [1]:

$$\mathcal{F}_q(x^{-1})(p) = i\sqrt{\Theta_0} \operatorname{sgn}(p) = i\sqrt{\Theta_0} [\theta_q^+(p) - \theta_q^-(p)]. \quad (177)$$

Next, we calculate the q -**Fourier transform of 1**. Eq. (177) together with Eq. (107) of the previous chapter and Eq. (167) implies [1, 15]:

$$\begin{aligned} \mathcal{F}_q(1)(p) &= \mathcal{F}_q(xx^{-1})(p) = -iD_{q,p} \mathcal{F}_q(x^{-1})(p) \\ &= \sqrt{\Theta_0} D_{q,p} [\theta_q^+(p) - \theta_q^-(p)] = 2\sqrt{\Theta_0} \delta_q(p). \end{aligned} \quad (178)$$

Using this result and Eq. (64) of Chap. 4, we can show that $\delta_q(p)$ is even:

$$\begin{aligned} \delta_q(-p) &= \frac{1}{2\sqrt{\Theta_0}} \mathcal{F}_q(1)(-p) = \frac{1}{4\Theta_0} \int_{-x_0.\infty}^{x_0.\infty} d_q x \exp_q(x|-ip) \\ &= \frac{1}{4\Theta_0} \int_{-x_0.\infty}^{x_0.\infty} d_q x \exp_q(ip|-x) = \\ &= \frac{1}{4\Theta_0} \int_{-x_0.\infty}^{x_0.\infty} d_q x \exp_q(ip|x) = \delta_q(p). \end{aligned} \quad (179)$$

Eq. (178) implies that the shifted q -delta function can be written as the q -**Fourier transform of a q -exponential function**:

$$\begin{aligned} \delta_q(a \bar{\oplus} x) &= \frac{1}{2\sqrt{\Theta_0}} \mathcal{F}_q(1)(a \bar{\oplus} x) = \frac{1}{4\Theta_0} \int_{-p_0.\infty}^{p_0.\infty} d_q p \exp_q(ip|a \bar{\oplus} x) \\ &= \frac{1}{4\Theta_0} \int_{-p_0.\infty}^{p_0.\infty} d_q p \exp_q(ip|a) \exp_q(ip|x) = \frac{1}{2\sqrt{\Theta_0}} \mathcal{F}_q(\exp_q(ip|a))(x). \end{aligned} \quad (180)$$

In the second last step of the above computation, we applied the addition theorem for q -exponentials [cf. Eq. (39) of Chap. 2].

Next, we compute the q -**Fourier transform of $\theta_q^+ - \theta_q^-$** . Using Eq. (101) of the previous chapter and Eq. (51) of Chap. 3, we get [1]:

$$\begin{aligned}
\mathcal{F}_q(\theta_q^+ - \theta_q^-)(p) &= \frac{1}{2\sqrt{\Theta_0}} \int_0^{x_0 \cdot \infty} d_q x \exp_q(ix|p) - \frac{1}{2\sqrt{\Theta_0}} \int_{-x_0 \cdot \infty}^0 d_q x \exp_q(ix|p) \\
&= \frac{(1-q)x_0}{2\sqrt{\Theta_0}} \sum_{m=-\infty}^{\infty} q^m [\exp_q(ix_0 q^m p) - \exp_q(-ix_0 q^m p)] \\
&= \frac{i(1-q)x_0}{\sqrt{\Theta_0}} \sum_{m=-\infty}^{\infty} q^m \sin_q(x_0 q^m p) \\
&= \frac{i}{\sqrt{\Theta_0}} p^{-1} \lim_{M \rightarrow \infty} (1-q)x_0 p \sum_{m=-M}^{\infty} q^m \sin_q(x_0 q^m p) \\
&= \frac{i}{\sqrt{\Theta_0}} p^{-1} \lim_{M \rightarrow \infty} [1 - \cos_q(x_0 q^{-M} p)] = \frac{i}{\sqrt{\Theta_0}} p^{-1}. \quad (181)
\end{aligned}$$

The above computation follows the same logic as the one in Eq. (86) of Chap. 3.

From the q -Fourier transform of $\theta_q^+ - \theta_q^-$, we derive the q -**Fourier transform of the q -delta function**. Using Eq. (107) of the previous chapter and Eq. (167), we get [1]:

$$\begin{aligned}
\mathcal{F}_q(\delta_q)(p) &= \frac{1}{2} \mathcal{F}_q(D_q(\theta_q^+ - \theta_q^-))(p) \\
&= -\frac{i}{2} \Lambda p \mathcal{F}_q(\theta_q^+ - \theta_q^-)(p) = \frac{1}{2\sqrt{\Theta_0}}. \quad (182)
\end{aligned}$$

A direct computation using Eq. (103) of the previous chapter and Eq. (11) of Chap. 2 gives the same result:

$$\begin{aligned}
\mathcal{F}_q(\delta_q)(p) &= \frac{1}{2\sqrt{\Theta_0}} \int_{-x_0 \cdot \infty}^{x_0 \cdot \infty} d_q x \delta_q(x) \exp_q(x|ip) \\
&= \frac{1}{2\sqrt{\Theta_0}} \exp_q(x|ip)|_{x=0} = \frac{1}{2\sqrt{\Theta_0}}. \quad (183)
\end{aligned}$$

We can use the results from Eqs. (178) and (181) to find the q -**Fourier transforms of θ_q^+ and θ_q^-** . Due to $1 = \theta_q^+ + \theta_q^-$, we have [1]:

$$\begin{aligned}
\mathcal{F}_q(\theta_q^+) &= \frac{1}{2} \mathcal{F}_q(\theta_q^+ - \theta_q^- + 1) = \frac{i}{2\sqrt{\Theta_0}} p^{-1} + \sqrt{\Theta_0} \delta_q(p), \\
\mathcal{F}_q(\theta_q^-) &= \frac{1}{2} \mathcal{F}_q(\theta_q^- - \theta_q^+ + 1) = \frac{-i}{2\sqrt{\Theta_0}} p^{-1} + \sqrt{\Theta_0} \delta_q(p). \quad (184)
\end{aligned}$$

Furthermore, we can calculate the q -**Fourier transform of a power function with an integer exponent**. As we show below, it holds [1] ($n \in \mathbb{N}_0$)

$$\mathcal{F}_q(x^n) = i^n 2\sqrt{\Theta_0} q^{-n(n+1)/2} [[n]]_q! p^{-n} \delta_q(p) \quad (185)$$

and

$$\mathcal{F}_q(x^{-n-1}) = i^{n+1} \sqrt{\Theta_0} ([[n]]_q!)^{-1} p^n \operatorname{sgn}(p). \quad (186)$$

From Eq. (185), we obtain a formula for calculating the q -Fourier transform of a function that can be represented as a power series:

$$\mathcal{F}_q(g) = 2\sqrt{\Theta_0} \sum_{n=0}^{\infty} i^n q^{-n(n+1)/2} p^{-n} \delta_q(p) (D_q^n g)(0). \quad (187)$$

To prove Eq. (185), we do the following calculation:

$$\begin{aligned} \delta_q(p) &= \frac{1}{2\sqrt{\Theta_0}} \mathcal{F}_q(1)(p) = \frac{1}{2\sqrt{\Theta_0}} \frac{\mathcal{F}_q(D_q^n x^n)(p)}{[[n]]_q!} \\ &= \frac{1}{2\sqrt{\Theta_0}} \frac{(-i)^n}{[[n]]_q!} (\Lambda p)^n \mathcal{F}_q(x^n)(p). \end{aligned} \quad (188)$$

The first identity follows from Eq. (178), the second one from the identity

$$1 = ([[n]]_q!)^{-1} D_q^n x^n, \quad (189)$$

and the last one from Eq. (167). Solving Eq. (188) for $\mathcal{F}_q(x^n)$ finally gives:

$$\begin{aligned} \mathcal{F}_q(x^n)(p) &= 2\sqrt{\Theta_0} i^n [[n]]_q! (p^{-1} \Lambda^{-1})^n \delta_q(p) \\ &= 2\sqrt{\Theta_0} i^n q^{-n(n+1)/2} [[n]]_q! p^{-n} \delta_q(p). \end{aligned} \quad (190)$$

In the last step, we applied Eq. (129) of the previous chapter and the following identities:

$$(p^{-1} \Lambda^{-1})^n = q^{-1-2-\dots-(n-1)} p^{-n} \Lambda^{-n} = q^{-n(n-1)/2} p^{-n} \Lambda^{-n}. \quad (191)$$

Next, we prove the formula in Eq. (186):

$$\begin{aligned} \mathcal{F}_q(x^{-n-1})(p) &= (-1)^n \frac{q^{n(n+1)/2}}{[[n]]_q!} \mathcal{F}_q(D_q^n x^{-1})(p) \\ &= i^n \frac{q^{n(n+1)/2}}{[[n]]_q!} (\Lambda p)^n \mathcal{F}_q(x^{-1})(p) \\ &= i^{n+1} \frac{\sqrt{\Theta_0}}{[[n]]_q!} p^n \operatorname{sgn}(p). \end{aligned} \quad (192)$$

In the first step, we applied the result of the following calculation:

$$\begin{aligned} D_q^k z^{-1} &= [[-1]]_q [[-2]]_q \dots [[-k]]_q z^{-k-1} \\ &= (-1)^k q^{-k(k+1)/2} [[k]]_q! z^{-k-1}. \end{aligned} \quad (193)$$

The second step in Eq. (192) follows from Eq. (167) and the last step from Eq. (177) if we take into account

$$(\Lambda p)^n = q^{-n(n+1)/2} p^n \Lambda^n \quad (194)$$

and

$$\Lambda \operatorname{sgn}(p) = \operatorname{sgn}(p). \quad (195)$$

Finally, we calculate the q -**Fourier transform of the function** $\Theta_q(z)$ [1]:

$$\begin{aligned} \mathcal{F}_q(\Theta_q) &= \frac{1}{2\sqrt{\Theta_0}} \int_{-x_0.\infty}^{x_0.\infty} d_q x \Theta_q(x) \exp_q(x|ip) \\ &= \frac{(1-q)x_0}{2\sqrt{\Theta_0}} \sum_{\varepsilon=\pm} \sum_{m=-\infty}^{\infty} q^m \Theta_q(\varepsilon x_0 q^m) \exp_q(\varepsilon i x_0 q^m p) \\ &= \frac{(1-q)x_0 \Theta_q(x_0)}{2\sqrt{\Theta_0}} \sum_{m=-\infty}^{\infty} q^m [\exp_q(i x_0 q^m p) - \exp_q(-i x_0 q^m p)] \\ &= \frac{i \Theta_q(x_0)}{\sqrt{\Theta_0}} p^{-1}. \end{aligned} \quad (196)$$

For the third step, we took into account that the function $\Theta_q(z)$ is even and q -periodic [cf. Eq. (74) of Chap. 4]. The last step results from the same considerations we used in Eq. (181).

We take another look at Eq. (113) of the previous chapter. This identity holds for functions being continuous at the origin. In particular, we can apply it to functions having a power series expansion around the origin. We can remove this restriction if we establish the following identities for $n \in \mathbb{Z}$:

$$\int_{-\infty}^{\infty} d_q x \delta_q(x) x^n = \begin{cases} 1, & \text{für } n = 0, \\ 0, & \text{für } n \neq 0. \end{cases} \quad (197)$$

This way, we can apply Eq. (113) of the previous chapter to a function for which we have a Laurent series centered at the origin:

$$\int_{-\infty}^{\infty} d_q x \delta_q(x) g(x) = g_0 \quad \text{with} \quad g(z) = \sum_{k=-\infty}^{\infty} g_k z^k. \quad (198)$$

We aim to prove Eq. (197). To this end, we compute the n -th **Jackson derivative of** $\delta_q(x)$. Using Eq. (8) of Chap. 2 and Eq. (129) of the previous chapter, we obtain ($n \in \mathbb{N}$):

$$\begin{aligned} D_q^n \delta_q(x) &= (-1)^n \frac{q^{-n(n+1)/2}}{(1-q)^n x^n} \sum_{m=0}^n \begin{bmatrix} n \\ m \end{bmatrix}_q (-q)^m q^{m(m-1)/2} \delta_q(x) \\ &= (-1)^n \frac{q^{n(n+1)/2}}{(1-q)^n x^n} (q; q)_n \delta_q(x) \\ &= (-1)^n q^{-n(n+1)/2} [[n]]_q! x^{-n} \delta_q(x). \end{aligned} \quad (199)$$

In the second step, we used the Gaussian binomial formula [13]:

$$(z; q)_n = \sum_{m=0}^n \begin{bmatrix} n \\ m \end{bmatrix}_q (-z)^m q^{m(m-1)/2}. \quad (200)$$

The final step in Eq. (199) follows from Eq. (20) of Chap. 2. Using the result in Eq. (199) and the rule in Eq. (70) of Chap. 4, we can perform the following computation:

$$\begin{aligned} \int_{-x_0.\infty}^{x_0.\infty} d_q x x^{-n} \delta_q(x) &= \frac{(-1)^n q^{n(n+1)/2}}{[[n]]_q!} \int_{-x_0.\infty}^{x_0.\infty} d_q x D_q^n \delta_q(x) \\ &= \frac{1}{[[n]]_q!} \int_{-x_0.\infty}^{x_0.\infty} d_q x \delta_q(q^n x) \cdot D_q^n 1 = 0. \end{aligned} \quad (201)$$

The results above are consistent with our earlier considerations. For example, we can derive the expression for $\mathcal{F}_q(x^n)$ in Eq. (185) using the derivative rule in Eq. (199) ($n \in \mathbb{N}_0$):

$$\begin{aligned} \mathcal{F}_q(x^n) &= (-i)^n D_{q,p}^n \mathcal{F}_q(1)(p) = 2\sqrt{\Theta_0} (-i)^n D_{q,p}^n \delta_q(p) \\ &= 2\sqrt{\Theta_0} i^n q^{-n(n+1)/2} [[n]]_q! p^{-n} \delta_q(p). \end{aligned} \quad (202)$$

With Eq. (197) and the series expansions in Eq. (144), we obtain:

$$\begin{aligned} \mathcal{F}_q^{-1}(p^{-n} \delta_q(p)) &= \frac{1}{2\sqrt{\Theta_0}} \int_{-p_0.\infty}^{p_0.\infty} d_q p \delta_q(p) p^{-n} \exp_{q^{-1}}(-ip|qx) \\ &= \frac{1}{2\sqrt{\Theta_0}} \frac{(-i)^n q^{n(n+1)/2} x^n}{[[n]]_q!}. \end{aligned} \quad (203)$$

Using the results of Eqs. (202) and (203), we immediately check the following identity:

$$(\mathcal{F}_q^{-1} \circ \mathcal{F}_q)(x^n) = x^n. \quad (204)$$

Furthermore, we can rewrite the expression in Eq. (114) of the previous chapter by applying the derivative rule in Eq. (199):

$$\delta_q((\bar{\ominus} qa) \bar{\oplus} x) = \sum_{k=0}^{\infty} \frac{q^{k(k+1)/2}}{[[k]]_q!} (-a)^k (D_q)^k \delta_q(x) = \sum_{k=0}^{\infty} a^k \delta_q(x) x^{-k}. \quad (205)$$

In the same manner, we obtain:

$$\delta_q(x \bar{\oplus} (\bar{\ominus} qa)) = \sum_{k=0}^{\infty} x^{-k} \delta_q(x) a^k. \quad (206)$$

The right-hand side of Eq. (205) or Eq. (206) satisfies Eq. (116) of the previous chapter if the function f has a power series expansion around the origin. If should Eq. (116) of the last chapter should hold when f has a pole at the origin, we need to extend the summation on the right-hand side of Eq. (205) or Eq. (206) to negative integers [cf. Eq. (197)]:

$$\begin{aligned} \delta_q((\bar{\ominus} qa) \bar{\oplus} x) &= \sum_{k=-\infty}^{\infty} a^k \delta_q(x) x^{-k}, \\ \delta_q(x \bar{\oplus} (\bar{\ominus} qa)) &= \sum_{k=-\infty}^{\infty} x^{-k} \delta_q(x) a^k. \end{aligned} \quad (207)$$

Using these expressions and taking into account Eq. (197), we can perform the following calculation ($n \in \mathbb{N}$):

$$\begin{aligned} \int_{-x_0.\infty}^{x_0.\infty} d_q x \delta_q((\bar{\ominus} qa) \bar{\oplus} x) x^{-n} &= \int_{-x_0.\infty}^{x_0.\infty} d_q x \sum_{k=-\infty}^{\infty} a^k \delta_q(x) x^{-k} x^{-n} \\ &= \sum_{k=-\infty}^{\infty} a^k \int_{-x_0.\infty}^{x_0.\infty} d_q x \delta_q(x) x^{-(k+n)} = \sum_{k=-\infty}^{\infty} a^k \delta_{k,-n} = a^{-n}. \end{aligned} \quad (208)$$

For completeness, we calculate the q -Fourier transform of the function $u_p^*(x)$ [cf. Eq. (157)], using Eq. (187) and Eq. (207):

$$\begin{aligned} \mathcal{F}_q(u_p^*(x)) &= \frac{1}{2\sqrt{\Theta_0}} \mathcal{F}_q(\exp_{q^{-1}}(-ip'|qx))(p) \\ &= \sum_{n=0}^{\infty} i^n \frac{q^{-n(n+1)/2}}{p^n} \delta_q(p) (D_{q,x}^n \exp_{q^{-1}}(-ip'|qx))|_{x=0} \\ &= \sum_{n=0}^{\infty} p^{-n} \delta_q(p) p'^n = \delta_q(p \bar{\oplus} (\bar{\ominus} qp')). \end{aligned} \quad (209)$$

In the third step of the calculation above, we used the result of the following calculation:

$$\begin{aligned} (D_{q,x}^n \exp_{q^{-1}}(-ip'|qx))|_{x=0} &= \sum_{k=0}^{\infty} \frac{q^{k(k+1)/2}}{[[k]]_q!} (-ip')^k (D_{q,x}^n x^k)|_{x=0} \\ &= \sum_{k=0}^{\infty} \frac{q^{k(k+1)/2}}{[[k]]_q!} (-ip')^k [[n]]_q! \begin{bmatrix} k \\ n \end{bmatrix}_q x^{k-n}|_{x=0} \\ &= q^{n(n+1)/2} (-ip')^n. \end{aligned} \quad (210)$$

We can apply the reasoning above to the **inverse q -Fourier transform** \mathcal{F}_q^{-1} . This way, we find:

$$\mathcal{F}_q^{-1}(\delta_q(p)) = \frac{1}{2\sqrt{\Theta_0}} \int_{-p_0.\infty}^{p_0.\infty} d_q p \delta_q(p) \exp_{q^{-1}}(-ip|qx) = \frac{1}{2\sqrt{\Theta_0}}. \quad (211)$$

We rewrite the inverse q -Fourier transform by using q -inversions [cf. Eq. (43) of Chap. 2]:

$$\begin{aligned} \mathcal{F}_q^{-1}(f) &= \frac{1}{2\sqrt{\Theta_0}} \int_{-p_0.\infty}^{p_0.\infty} d_q p \exp_q(ip|\bar{\ominus} qx) = \\ &= \frac{1}{2\sqrt{\Theta_0}} \bar{\mathcal{S}}_x \left(\int_{-p_0.\infty}^{p_0.\infty} d_q p \exp_q(ip|qx) \right). \end{aligned} \quad (212)$$

With the help of Eq. (42) of Chap. 2 and a procedure similar to that in Eq. (181),

we get:

$$\begin{aligned}
\mathcal{F}_q^{-1}(\theta_q^+ - \theta_q^-) &= \frac{1}{2\sqrt{\Theta_0}} \bar{\mathcal{S}}_x \left(\int_0^{p_0 \cdot \infty} d_q p \exp_q(ip|qx) - \int_{-p_0 \cdot \infty}^0 d_q p \exp_q(ip|qx) \right) \\
&= \frac{i(1-q)p_0}{q\sqrt{\Theta_0}} \bar{\mathcal{S}}_x \left(\sum_{m=-\infty}^{\infty} q^m \sin_q(p_0 q^m x) \right) \\
&= \bar{\mathcal{S}}_x \left(\frac{i}{q\sqrt{\Theta_0} x} \right) = -\frac{i}{\sqrt{\Theta_0}} x^{-1}.
\end{aligned} \tag{213}$$

In the same manner, we obtain [also cf. Eq. (174)]:

$$\begin{aligned}
\mathcal{F}_q^{-1}(p^{-1}) &= \frac{1}{2\sqrt{\Theta_0}} \bar{\mathcal{S}}_x \left(\int_{-p_0 \cdot \infty}^{p_0 \cdot \infty} d_q p p^{-1} \exp_q(ip|qx) \right) \\
&= \frac{i(1-q)}{\sqrt{\Theta_0}} \bar{\mathcal{S}}_x \left(\sum_{m=-\infty}^{\infty} \sin_q(p_0 q^{m+1} x) \right) \\
&= \frac{i}{\sqrt{\Theta_0}} \bar{\mathcal{S}}_x(\Theta_q(p_0 x)) = -i\sqrt{\Theta_0} \operatorname{sgn}(x).
\end{aligned} \tag{214}$$

We apply this result to calculate $\mathcal{F}_q^{-1}(1)$ [also cf. Eq. (178)]:

$$\begin{aligned}
\mathcal{F}_q^{-1}(1)(x) &= \mathcal{F}_q^{-1}(pp^{-1})(x) = iq^{-1}\Lambda^{-1}D_{q,x}\mathcal{F}_q^{-1}(p^{-1})(x) \\
&= q^{-1}\sqrt{\Theta_0}\Lambda^{-1}D_{q,x}\operatorname{sgn}(x) = 2\sqrt{\Theta_0}\delta_q(x).
\end{aligned} \tag{215}$$

In the last step of the above calculation, we used Eqs. (107) and (129) from Chap. 5. From Eq. (215), we derive an expression for $\mathcal{F}_q^{-1}(p^n)$ with $n \in \mathbb{N}$, taking into account Eq. (168):

$$\begin{aligned}
\delta_q(x) &= \frac{1}{2\sqrt{\Theta_0}} \mathcal{F}_q^{-1}(1)(x) = \frac{1}{2\sqrt{\Theta_0}} \frac{\mathcal{F}_q^{-1}(D_{q,p}^n p^n)(x)}{[[n]]_q!} \\
&= \frac{1}{2\sqrt{\Theta_0}} \frac{i^n \mathcal{F}_q^{-1}(p^n)(x)}{[[n]]_q!} x^n.
\end{aligned} \tag{216}$$

The result above leads to the following identity:

$$\mathcal{F}_q^{-1}(p^n)(x) = i^{-n} 2\sqrt{\Theta_0} [[n]]_q! x^{-n} \delta_q(x). \tag{217}$$

For a function f that has a power series expansion around the origin, the formula above implies the following identity [also cf. Eq. (187)]:

$$\mathcal{F}_q^{-1}(f(p))(x) = 2\sqrt{\Theta_0} \sum_{n=0}^{\infty} i^{-n} (D_{q,p}^n f)(0) \delta_q(x) x^{-n}. \tag{218}$$

Due to this identity, we can calculate the inverse q -Fourier transform of the function $u_p(x)$:

$$\begin{aligned}\mathcal{F}_q^{-1}(u_p(x'))(x) &= \sum_{n=0}^{\infty} i^{-n} (D_{q,p}^n \exp_q(x' | ip)) \Big|_{p=0} \delta_q(x) x^{-n} \\ &= \sum_{n=0}^{\infty} x'^n \delta_q(x) x^{-n} = \delta_q((\ominus qx') \oplus x).\end{aligned}\quad (219)$$

Finally, we derive an expression for $\mathcal{F}_q^{-1}(p^{-n-1})$ with $n \in \mathbb{N}$. The following calculation uses considerations similar to that in Eq. (192):

$$\begin{aligned}\mathcal{F}_q^{-1}(p^{-n-1})(x) &= \frac{(-1)^n q^{n(n+1)/2}}{[[n]]_q!} \mathcal{F}_q^{-1}(D_{q,p}^n p^{-1})(x) \\ &= \frac{(-i)^n q^{n(n+1)/2}}{[[n]]_q!} x^n \mathcal{F}_q^{-1}(p^{-1})(x) \\ &= \sqrt{\Theta_0} \frac{(-i)^{n+1} q^{n(n+1)/2}}{[[n]]_q!} x^n \operatorname{sgn}(x).\end{aligned}\quad (220)$$

With our findings, we can directly confirm the identities ($n \in \mathbb{N}$)

$$\begin{aligned}(\mathcal{F}_q \circ \mathcal{F}_q^{-1})(\delta_q) &= \delta_q, \\ (\mathcal{F}_q \circ \mathcal{F}_q^{-1})(1) &= 1, \\ (\mathcal{F}_q \circ \mathcal{F}_q^{-1})(p^n) &= p^n, \\ (\mathcal{F}_q \circ \mathcal{F}_q^{-1})(p^{-n-1}) &= p^{-n-1},\end{aligned}\quad (221)$$

and

$$\begin{aligned}(\mathcal{F}_q^{-1} \circ \mathcal{F}_q)(\delta_q) &= \delta_q, \\ (\mathcal{F}_q^{-1} \circ \mathcal{F}_q)(1) &= 1, \\ (\mathcal{F}_q^{-1} \circ \mathcal{F}_q)(x^n) &= x^n, \\ (\mathcal{F}_q^{-1} \circ \mathcal{F}_q)(x^{-n-1}) &= x^{-n-1}.\end{aligned}\quad (222)$$

To prove the last identity in Eq. (221) or Eq. (222), we additionally need [cf. Eq. (181)]

$$\begin{aligned}\mathcal{F}_q(x^n \operatorname{sgn}(x))(p) &= (-i)^n D_{q,p}^n \mathcal{F}_q(\operatorname{sgn}(x))(p) = \\ &= \frac{i^{n+1} [[n]]_q!}{\sqrt{\Theta_0} q^{n(n+1)/2}} p^{-n-1}\end{aligned}\quad (223)$$

or [cf. Eq. (213)]

$$\begin{aligned}\mathcal{F}_q^{-1}(p^n \operatorname{sgn}(p))(x) &= i^n q^{-n(n+1)/2} \Lambda^{-n} D_{q,x}^n \mathcal{F}_q^{-1}(\operatorname{sgn}(p))(x) \\ &= \frac{(-i)^{n+1} [[n]]_q!}{\sqrt{\Theta_0}} x^{-n-1}.\end{aligned}\quad (224)$$

Note that we have confirmed $(\mathcal{F}_q^{-1} \circ \mathcal{F}_q)(x^n) = x^n$ using Eqs. (202) and (203). To prove $(\mathcal{F}_q \circ \mathcal{F}_q^{-1})(p^n) = p^n$ in the same way, we additionally need:

$$\begin{aligned} \mathcal{F}_q(x^{-n} \delta_q(x))(p) &= \frac{1}{2\sqrt{\Theta_0}} \int_{-x_0, \infty}^{x_0, \infty} d_q x x^{-n} \delta_q(x) \exp_q(x|ip) \\ &= \frac{1}{2\sqrt{\Theta_0}} \frac{(ip)^n}{[[n]]_q!}. \end{aligned} \quad (225)$$

7 Fourier transforms on $\mathbb{G}_{q^{1/2}, x_0}$

To obtain an improper q -deformed integral over the lattice $\mathbb{G}_{q^{1/2}, x_0}$, we can add an improper q -integral over the lattice \mathbb{G}_{q, x_0} to an improper q -integral over the lattice $\mathbb{G}_{q, q^{1/2}x_0}$ ($0 < q < 1$) [6, 19, 20]:

$$\begin{aligned} \frac{1}{1+q^{1/2}} \left(\int_{-x_0, \infty}^{x_0, \infty} d_q x f(x) + \int_{-x_0 q^{1/2}, \infty}^{x_0 q^{1/2}, \infty} d_q x f(x) \right) &= \\ &= \frac{1-q}{1+q^{1/2}} \sum_{\varepsilon=\pm} \sum_{m=-\infty}^{\infty} \left[x_0 q^m f(\varepsilon x_0 q^m) + x_0 q^{m+1/2} f(\varepsilon x_0 q^{m+1/2}) \right] \\ &= (1-q^{1/2}) \sum_{m=-\infty}^{\infty} x_0 q^{m/2} \left[f(x_0 q^{m/2}) + f(-x_0 q^{m/2}) \right] \\ &= \int_{-x_0, \infty}^{x_0, \infty} d_{q^{1/2}} x f(x). \end{aligned} \quad (226)$$

Note that we have multiplied the sum of the two q -integrals by the factor $(1+q^{1/2})^{-1}$ to regain the ordinary integral for $q \rightarrow 1$.

The $q^{1/2}$ -integral in Eq. (226) consists of q -integrals invariant under q -translations. Thus, under q -translations the $q^{1/2}$ -integral is also invariant. By analogy with Eq. (68) from Chap. 4, the following holds ($m \in \mathbb{N}_0$):

$$\begin{aligned} \int_{-x, \infty}^{x, \infty} d_{q^{1/2}} z D_q^m f(z) &= \\ &= \frac{1}{1+q^{1/2}} \left(\int_{-x_0, \infty}^{x_0, \infty} d_q x D_q^m f(x) + \int_{-x_0 q^{1/2}, \infty}^{x_0 q^{1/2}, \infty} d_q x D_q^m f(x) \right) \\ &= 0. \end{aligned} \quad (227)$$

This identity, together with Eq. (30) of Chap. 2, implies:

$$\begin{aligned} \int_{-x, \infty}^{x, \infty} d_{q^{1/2}} z f(z \oplus a) &= \int_{-x, \infty}^{x, \infty} d_{q^{1/2}} z \sum_{k=0}^{\infty} \frac{1}{[[k]]_q!} [D_q^k f(z)] a^k \\ &= \sum_{k=0}^{\infty} \frac{1}{[[k]]_q!} \int_{-x, \infty}^{x, \infty} d_{q^{1/2}} z [D_q^k f(z)] a^k = \int_{-x, \infty}^{x, \infty} d_{q^{1/2}} z f(z). \end{aligned} \quad (228)$$

By similar reasoning, we obtain

$$\int_{-x.\infty}^{x.\infty} d_{q^{1/2}} z f(a \bar{\oplus} z) = \int_{-x.\infty}^{x.\infty} d_{q^{1/2}} z f(z). \quad (229)$$

The **rule for integration by parts** also holds for the improper $q^{1/2}$ -integral. It gets this property from the q -integrals of which it is composed [cf. Eq. (70) in Chap. 4]:

$$\int d_{q^{1/2}} x f(x) D_q^k g(x) = (-1)^k q^{-k(k-1)/2} \int d_{q^{1/2}} x [D_q^k f(x)] g(q^k x). \quad (230)$$

Next, we show how to formulate q -Fourier transforms using the $q^{1/2}$ -integrals from Eq. (226). Instead of Eqs. (146) and (147) of the last chapter, it holds

$$\int_{-p_0.\infty}^{p_0.\infty} d_{q^{1/2}} p \exp_q(x_0 | ip) \exp_{q^{-1}}(-ip | qx) = \begin{cases} \frac{\text{vol}_q}{(1-q^{1/2})x_0} & \text{for } x = x_0, \\ 0 & \text{for } x \neq x_0, \end{cases} \quad (231)$$

and

$$\int_{-x_0.\infty}^{x_0.\infty} d_{q^{1/2}} x \exp_{q^{-1}}(-ip_0 | qx) \exp_q(x | ip) = \begin{cases} \frac{\text{vol}_q}{(1-q^{1/2})p_0} & \text{for } p = p_0, \\ 0 & \text{for } p \neq p_0, \end{cases} \quad (232)$$

with

$$\text{vol}_q = \frac{8 [\Theta_q(x_0 p_0) + \Theta_q(q^{1/2} x_0 p_0)]}{(1 + q^{1/2})^2}. \quad (233)$$

Remarkably, we obtain the identities in Eqs. (231) and (232) from the identities in Eqs. (146) and (147) of the previous chapter by the following substitutions:

$$\begin{aligned} \int_{-x_0.\infty}^{x_0.\infty} d_q x &\rightarrow \int_{-x_0.\infty}^{x_0.\infty} d_{q^{1/2}} x, & \int_{-p_0.\infty}^{p_0.\infty} d_q p &\rightarrow \int_{-p_0.\infty}^{p_0.\infty} d_{q^{1/2}} p, \\ 4\Theta_0 &\rightarrow \text{vol}_q, & 1 - q &\rightarrow 1 - q^{1/2}. \end{aligned} \quad (234)$$

Accordingly, the defining expression for the q -deformed Fourier transform also changes [cf. Eq. (149) of the previous chapter]:

$$\begin{aligned} \mathcal{F}_{q^{1/2}}(\phi)(p) &= \text{vol}_q^{-1/2} \int_{-x_0.\infty}^{x_0.\infty} d_{q^{1/2}} x \phi(x) \exp_q(x | ip), \\ \mathcal{F}_{q^{1/2}}^{-1}(\psi)(x) &= \text{vol}_q^{-1/2} \int_{-p_0.\infty}^{p_0.\infty} d_{q^{1/2}} p \psi(p) \exp_{q^{-1}}(-ip | qx). \end{aligned} \quad (235)$$

For all $n \in \mathbb{Z}$, it holds [cf. Eq. (100) of Chap. 5]

$$(\mathcal{F}_{q^{1/2}}^{-1} \circ \mathcal{F}_{q^{1/2}})(\phi_{n/2}^\pm)(x) = \begin{cases} 1 & \text{for } x = \pm x_0 q^{n/2}, \\ 0 & \text{for } x \neq \pm x_0 q^{n/2}, \end{cases} \quad (236)$$

and

$$(\mathcal{F}_{q^{1/2}} \circ \mathcal{F}_{q^{1/2}}^{-1})(\psi_{n/2}^\pm)(p) = \begin{cases} 1 & \text{for } p = \pm p_0 q^{n/2}, \\ 0 & \text{for } p \neq \pm p_0 q^{n/2}. \end{cases} \quad (237)$$

We can prove Eq. (236) by the following calculation:

$$\begin{aligned} & \frac{1}{\text{vol}_q} \int_{-p_0 \cdot \infty}^{p_0 \cdot \infty} d_{q^{1/2}} p \int_{-x_0 \cdot \infty}^{x_0 \cdot \infty} d_{q^{1/2}} \xi \phi_{n/2}^\pm(\xi) \exp_q(\xi |ip) \exp_{q^{-1}}(-ip|qx) \\ &= \frac{1 - q^{1/2}}{\text{vol}_q} x_0 q^{n/2} \int_{-p_0 \cdot \infty}^{p_0 \cdot \infty} d_{q^{1/2}} p \exp_q(\pm x_0 q^{n/2} |ip) \exp_{q^{-1}}(-ip|qx) \\ &= \frac{1 - q^{1/2}}{\text{vol}_q} x_0 \int_{-p_0 \cdot \infty}^{p_0 \cdot \infty} d_{q^{1/2}} p \exp_q(x_0 |ip) \exp_{q^{-1}}(-ip|\pm q^{-n/2} qx). \end{aligned} \quad (238)$$

Due to Eq. (231), the result above implies Eq. (236). Eq. (237) can be proved in a similar way.

The substitutions in Eq. (234) also change the normalization of the **momentum eigenfunctions**:

$$u_p(x) = \text{vol}_q^{-1/2} \exp_q(ix|p), \quad u_p^*(x) = \text{vol}_q^{-1/2} \exp_{q^{-1}}(-ip|qx). \quad (239)$$

These momentum eigenfunctions again satisfy **completeness relations and orthogonality conditions**. Due to the identities in Eqs. (231) and (232), it holds ($n, m \in \mathbb{Z}$; $\varepsilon, \varepsilon' = \pm 1$)

$$\begin{aligned} & \int_{-p_0 \cdot \infty}^{p_0 \cdot \infty} d_{q^{1/2}} p u_p(\varepsilon q^{n/2}) u_p^*(\varepsilon' q^{m/2}) = \frac{\delta_{\varepsilon \varepsilon'} \delta_{nm}}{(1 - q^{1/2}) x_0 q^{n/2}} \\ &= \delta_{q^{1/2}}((\bar{\ominus} qx) \bar{\oplus} y)|_{x=\varepsilon x_0 q^{n/2}; y=\varepsilon' x_0 q^{m/2}} \end{aligned} \quad (240)$$

and

$$\begin{aligned} & \int_{-x_0 \cdot \infty}^{x_0 \cdot \infty} d_{q^{1/2}} x u_{\varepsilon q^{n/2}}^*(x) u_{\varepsilon' q^{m/2}}(x) = \frac{\delta_{\varepsilon \varepsilon'} \delta_{nm}}{(1 - q^{1/2}) p_0 q^{n/2}} \\ &= \delta_{q^{1/2}}((\bar{\ominus} qp) \bar{\oplus} p')|_{p=\varepsilon p_0 q^{n/2}; p'=\varepsilon' p_0 q^{m/2}}, \end{aligned} \quad (241)$$

where $\delta_{q^{1/2}}$ denotes the delta distribution for the $q^{1/2}$ -integral:

$$\int_{-x_0 \cdot \infty}^{x_0 \cdot \infty} d_{q^{1/2}} x \delta_{q^{1/2}}(x) f(x) = f(0). \quad (242)$$

Note that the last identity in Eq. (240) or Eq. (241) holds due to Eq. (255).

We can apply the substitutions in Eq. (234) to Eqs. (161)-(173) of the previous chapter. The derivations of these identities remain valid if we change the improper Jackson integrals and the normalization factors according to Eq. (234).

For completeness, we calculate the **classical limit of the q -deformed volume element** introduced in Eq. (233). Using Eq. (80) of Chap. 4, we find:

$$\begin{aligned} \lim_{q \rightarrow 1^-} \text{vol}_q &= \lim_{q \rightarrow 1^-} \frac{8 [\Theta_q(x_0 p_0) + \Theta_q(q^{1/2} x_0 p_0)]}{(1 + q^{1/2})^2} \\ &= \frac{8 \left(\frac{\pi}{2} + \frac{\pi}{2} \right)}{4} = 2\pi. \end{aligned} \quad (243)$$

We summarize some more **properties of the $q^{1/2}$ -delta function**. First, we calculate the q -derivatives of $\delta_{q^{1/2}}(z)$ by using Eq. (230):

$$\begin{aligned}
(D_q^k \delta_{q^{1/2}})(g) &= \int_{-x_0.\infty}^{x_0.\infty} d_{q^{1/2}} x [D_q^k \delta_{q^{1/2}}(x)] g(x) \\
&= (-1)^k q^{-k(k+1)/2} \int_{-x_0.\infty}^{x_0.\infty} d_{q^{1/2}} x \delta_{q^{1/2}}(x) (D_q^k g)(q^{-k}x) \\
&= (-1)^k q^{-k(k+1)/2} (D_q^k g)(0). \tag{244}
\end{aligned}$$

Furthermore, $\delta_{q^{1/2}}$ is again related to the q -derivative of a step function on the $q^{1/2}$ -lattice:

$$D_q \theta_{q^{1/2}}^\pm(x) = \frac{\pm 2}{1+q^{1/2}} \delta_{q^{1/2}}(x) \quad \text{with} \quad \theta_{q^{1/2}}^\pm(x) = \sum_{n=-\infty}^{\infty} \phi_{n/2}^\pm(x). \tag{245}$$

Note that Eq. (245) is a distributional identity. If g is a test function that is continuous at the origin of the lattice $\mathbb{G}_{q^{1/2}, x_0}$, it holds:

$$(D_q \theta_{q^{1/2}}^\pm)(g) = \frac{\pm 2}{1+q^{1/2}} g(0^\pm) = \frac{\pm 2}{1+q^{1/2}} \delta_{q^{1/2}}(g). \tag{246}$$

We prove this relation by the following calculation (we restrict ourselves to $\theta_{q^{1/2}}^+$; there is a similar calculation for $\theta_{q^{1/2}}^-$):

$$\begin{aligned}
(D_q \theta_{q^{1/2}}^+)(g) &= \int_{-x_0.\infty}^{x_0.\infty} d_{q^{1/2}} x [D_q \theta_{q^{1/2}}^+(x)] g(x) \\
&= \frac{-q^{-1}}{1+q^{1/2}} \left(\int_0^{x_0.\infty} d_q x (D_q g)(q^{-1}x) + \int_0^{x_0 q^{1/2}.\infty} d_q x (D_q g)(q^{-1}x) \right) \\
&= \frac{-1}{1+q^{1/2}} (-g(0^+) - g(0^+)) = \frac{2}{1+q^{1/2}} g(0^+). \tag{247}
\end{aligned}$$

In the second step of the calculation above, we used Eq. (97) from Chap. 5 and Eq. (226). The penultimate step follows from Eq. (66) of Chap. 4 and the fact that g vanishes at infinity.

From Eq. (245), we obtain a representation of $\delta_{q^{1/2}}^\pm(z)$ as the limit of regular distributions [also see the calculation in Eq. (110) of Chap. 5] ($0 < q < 1$):

$$\begin{aligned}
\delta_{q^{1/2}}(z) &= \pm \frac{1+q^{1/2}}{2} D_q \theta_{q^{1/2}}^\pm(z) = \pm \frac{1+q^{1/2}}{2} \sum_{n=-\infty}^{\infty} D_q \phi_{n/2}^\pm(z) \\
&= \pm \frac{1+q^{1/2}}{2} \sum_{n=-\infty}^{\infty} \frac{\phi_{n/2}^\pm(z) - \phi_{n/2}^\pm(qz)}{(1-q)z} \\
&= \pm \sum_{n=-\infty}^{\infty} \frac{\phi_{n/2}^\pm(z) - \phi_{(n-1)/2}^\pm(z) + \phi_{(n-1)/2}^\pm(z) - \phi_{(n-2)/2}^\pm(z)}{2(1-q^{1/2})z} \\
&= \pm (1-q^{1/2})^{-1} z^{-1} \lim_{n \rightarrow \infty} \phi_{n/2}^\pm(z). \tag{248}
\end{aligned}$$

Note that the identities above refer to distributions. Hence, for any test function g on the lattice $\mathbb{G}_{q^{1/2}, x_0}$, we have:

$$\int_{-x_0. \infty}^{x_0. \infty} d_{q^{1/2}} x \delta_{q^{1/2}}(x) g(x) = \pm \lim_{n \rightarrow \infty} \int_{-x_0. \infty}^{x_0. \infty} d_{q^{1/2}} x \frac{\phi_{n/2}^{\pm}(x)}{(1 - q^{1/2})x} g(x). \quad (249)$$

We examine the **scaling properties of the $q^{1/2}$ -step functions and the $q^{1/2}$ -delta distribution**. From the definition of the scaling operator, it follows ($k, n \in \mathbb{Z}$):

$$\Lambda^{k/2} \phi_{n/2}^{\pm}(z) = \phi_{n/2}^{\pm}(q^{k/2} z) = \phi_{(n-k)/2}^{\pm}(z). \quad (250)$$

Moreover, we have:

$$\begin{aligned} \Lambda^{k/2} \theta_{q^{1/2}}^{\pm}(z) &= \sum_{n=-\infty}^{\infty} \Lambda^k \phi_{n/2}^{\pm}(z) = \sum_{n=-\infty}^{\infty} \phi_{(n-k)/2}^{\pm}(z) \\ &= \sum_{n=-\infty}^{\infty} \phi_{n/2}^{\pm}(z) = \theta_{q^{1/2}}^{\pm}(z). \end{aligned} \quad (251)$$

Using the results above, we can derive the scaling property of $\delta_{q^{1/2}}$. With the help of Eq. (245), we do the following calculation:

$$\begin{aligned} \delta_{q^{1/2}}(q^{k/2} z) &= \Lambda^{k/2} \delta_{q^{1/2}}(z) = \pm \frac{1 + q^{1/2}}{2} \Lambda^{k/2} D_q \theta_{q^{1/2}}^{\pm}(z) \\ &= \pm \frac{1 + q^{1/2}}{2} q^{-k/2} D_q \Lambda^{k/2} \theta_{q^{1/2}}^{\pm}(z) \\ &= \pm \frac{1 + q^{1/2}}{2} q^{-k/2} D_q \theta_{q^{1/2}}^{\pm}(z) = q^{-k/2} \delta_{q^{1/2}}(z). \end{aligned} \quad (252)$$

We modify the identities of Eqs. (114)-(116) of Chap. 5 so that they apply to the lattice $\mathbb{G}_{q^{1/2}, x_0}$. To this end, we apply the substitutions in Eq. (234) to these identities and write $\delta_{q^{1/2}}$ instead of δ_q . For example, we get

$$\begin{aligned} \int_{-x_0. \infty}^{x_0. \infty} d_{q^{1/2}} x \delta_{q^{1/2}}((\bar{\ominus} qa) \bar{\oplus} x) f(x) &= f(a), \\ \int_{-x_0. \infty}^{x_0. \infty} d_{q^{1/2}} x f(x) \delta_{q^{1/2}}(x \bar{\oplus} (\bar{\ominus} qa)) &= f(a) \end{aligned} \quad (253)$$

where

$$\begin{aligned} \delta_{q^{1/2}}(x \bar{\oplus} (\bar{\ominus} qa)) &= \delta_{q^{1/2}}(x \bar{\oplus} (\bar{\ominus} qa)) \\ &= \sum_{k=0}^{\infty} \frac{q^{k(k+1)/2} (-a)^k}{[[k]]_q!} D_q^k \delta_{q^{1/2}}(x). \end{aligned} \quad (254)$$

Moreover, we have ($m \in \mathbb{Z}$, $0 < q < 1$):

$$\begin{aligned} \delta_{q^{1/2}}((\bar{\ominus} qa) \bar{\oplus} x) \Big|_{a=\pm x_0 q^{m/2}} &= \delta_{q^{1/2}}(x \bar{\oplus} (\bar{\ominus} qa)) \Big|_{a=\pm x_0 q^{m/2}} \\ &= \frac{\phi_{m/2}^{\pm}(x)}{(1 - q^{1/2}) x_0 q^{m/2}}. \end{aligned} \quad (255)$$

In the following, we give **expressions for q -deformed Fourier transforms on the lattice $\mathbb{G}_{q^{1/2}, x_0}$** . In complete analogy to Eq. (174) of the previous chapter, we can calculate $\mathcal{F}_{q^{1/2}}(x^{-1})$ with the following result:

$$\begin{aligned}\mathcal{F}_{q^{1/2}}(x^{-1})(p) &= \text{vol}_q^{-1/2} \int_{-x_0.\infty}^{x_0.\infty} d_{q^{1/2}} x x^{-1} \exp_q(x|ip) \\ &= \frac{2i [\Theta_q(x_0 p) + \Theta_q(q^{1/2} x_0 p)]}{(1 + q^{1/2}) \text{vol}_q^{1/2}}.\end{aligned}\quad (256)$$

For the lattice $\mathbb{G}_{q^{1/2}, x_0}$, the variable p can only take the values $\pm x_0 p_0 q^{m/2}$ with $m \in \mathbb{Z}$. Thus, it holds [cf. Eq. (233)]:

$$\begin{aligned}\frac{2i [\Theta_q(\pm x_0 p_0 q^{m/2}) + \Theta_q(\pm x_0 p_0 q^{(m+1)/2})]}{(1 + q^{1/2}) \text{vol}_q^{1/2}} &= \pm \frac{2i [\Theta_q(x_0 p_0) + \Theta_q(q^{1/2} x_0 p_0)]}{(1 + q^{1/2}) \text{vol}_q^{1/2}} \\ &= \pm \frac{i}{\sqrt{2}} [\Theta_q(x_0 p_0) + \Theta_q(q^{1/2} x_0 p_0)]^{1/2} \\ &= \pm \frac{i}{4} (1 + q^{1/2}) \text{vol}_q^{1/2}.\end{aligned}\quad (257)$$

Combining our results so far, we finally obtain:

$$\mathcal{F}_{q^{1/2}}(x^{-1})(p) = \frac{i}{4} (1 + q^{1/2}) \text{vol}_q^{1/2} \text{sgn}(p).\quad (258)$$

The expression for $\mathcal{F}_{q^{1/2}}(x^{-1})$ enables us to calculate an expression for $\mathcal{F}_{q^{1/2}}(1)$. Analogous to Eq. (178) of the last chapter and considering Eq. (245), we find:

$$\begin{aligned}\mathcal{F}_{q^{1/2}}(1)(p) &= \mathcal{F}_{q^{1/2}}(x x^{-1})(p) = -i D_{q,p} \mathcal{F}_{q^{1/2}}(x^{-1})(p) \\ &= \frac{1}{4} (1 + q^{1/2}) \text{vol}_q^{1/2} D_{q,p} [\theta_{q^{1/2}}^+(p) - \theta_{q^{1/2}}^-(p)] \\ &= \text{vol}_q^{1/2} \delta_{q^{1/2}}(p).\end{aligned}\quad (259)$$

This result implies the following identities for the shifted $q^{1/2}$ -delta function:

$$\begin{aligned}\delta_{q^{1/2}}(a \bar{\oplus} x) &= \frac{1}{\text{vol}_q^{1/2}} \mathcal{F}_{q^{1/2}}(1)(a \bar{\oplus} x) \\ &= \frac{1}{\text{vol}_q} \int_{-p_0.\infty}^{p_0.\infty} d_{q^{1/2}} p \exp_q(ip|a \bar{\oplus} x) \\ &= \frac{1}{\text{vol}_q} \int_{-p_0.\infty}^{p_0.\infty} d_{q^{1/2}} p \exp_q(ip|a) \exp_q(ip|x) \\ &= \frac{1}{\text{vol}_q^{1/2}} \mathcal{F}_{q^{1/2}}(\exp_q(ip|a))(x).\end{aligned}\quad (260)$$

Next, we come to the $q^{1/2}$ -Fourier transform of $\theta_{q^{1/2}}^+ - \theta_{q^{1/2}}^-$. Analogous to Eq. (181) of the previous chapter, we do the following calculation:

$$\begin{aligned}
\mathcal{F}_{q^{1/2}}(\theta_{q^{1/2}}^+ - \theta_{q^{1/2}}^-)(p) &= \\
&= \text{vol}_q^{-1/2} \left(\int_0^{x_0 \cdot \infty} d_{q^{1/2}} x \exp_q(x|ip) - \int_{-x_0 \cdot \infty}^0 d_{q^{1/2}} x \exp_q(-x|ip) \right) \\
&= \frac{(1 - q^{1/2})x_0}{\text{vol}_q^{1/2}} \sum_{m=-\infty}^{\infty} q^{m/2} \left[\exp_q(ix_0 q^{m/2} p) - \exp_q(-ix_0 q^{m/2} p) \right] \\
&= \frac{2i(1 - q^{1/2})x_0}{\text{vol}_q^{1/2}} \sum_{m=-\infty}^{\infty} q^{m/2} \sin_q(x_0 q^{m/2} p). \tag{261}
\end{aligned}$$

We rewrite the series in the last expression in the following way [cf. Eq. (181) of Chap. 6]:

$$\begin{aligned}
\sum_{m=-\infty}^{\infty} q^{m/2} \sin_q(x_0 q^{m/2} p) &= \sum_{m=-\infty}^{\infty} q^m \left[\sin_q(x_0 q^m p) + q^{1/2} \sin_q(x_0 q^{m+1/2} p) \right] \\
&= \frac{(1 - q)x_0 p}{(1 - q)x_0 p} \lim_{M \rightarrow \infty} \sum_{m=-M}^{\infty} \left[q^m \sin_q(x_0 q^m p) + q^{m+1/2} \sin_q(x_0 q^{m+1/2} p) \right] \\
&= \frac{1}{(1 - q)x_0 p} \lim_{M \rightarrow \infty} \left[1 - \cos_q(x_0 q^{-M} p) + 1 - \cos_q(x_0 q^{-M+1/2} p) \right] \\
&= \frac{2}{(1 - q)x_0 p}. \tag{262}
\end{aligned}$$

Plugging this result into Eq. (261), we finally obtain:

$$\mathcal{F}_{q^{1/2}}(\theta_{q^{1/2}}^+ - \theta_{q^{1/2}}^-)(p) = \frac{4i}{(1 + q^{1/2}) \text{vol}_q^{1/2}} p^{-1}. \tag{263}$$

Using the formula above and Eq. (245), we can immediately calculate the $q^{1/2}$ -Fourier transform of $\delta_{q^{1/2}}$:

$$\begin{aligned}
\mathcal{F}_{q^{1/2}}(\delta_{q^{1/2}})(p) &= \frac{1 + q^{1/2}}{4} \mathcal{F}_{q^{1/2}}(D_q(\theta_{q^{1/2}}^+ - \theta_{q^{1/2}}^-)) \\
&= -\frac{i(1 + q^{1/2})}{4} \Lambda p \mathcal{F}_{q^{1/2}}(\theta_{q^{1/2}}^+ - \theta_{q^{1/2}}^-) = \text{vol}_q^{-1/2}. \tag{264}
\end{aligned}$$

Furthermore, we can give the $q^{1/2}$ -Fourier transform of θ_q^+ and θ_q^- . Taking into account

$$2\theta_{q^{1/2}}^{\pm} = \pm(\theta_{q^{1/2}}^+ - \theta_{q^{1/2}}^-) + 1 \tag{265}$$

and Eqs. (259) and (263), it follows:

$$\mathcal{F}_{q^{1/2}}(\theta_{q^{1/2}}^{\pm}) = \frac{\pm 2i}{(1 + q^{1/2}) \text{vol}_q^{1/2}} p^{-1} + \frac{1}{2} \text{vol}_q^{1/2} \delta_{q^{1/2}}(p). \tag{266}$$

We also determine an expression for $\mathcal{F}_{q^{1/2}}(x^k)$ with $k \in \mathbb{Z}$. We first consider the case of the positive exponents. Using Eq. (189) from the previous chapter and Eq. (259), we perform the following calculation ($n \in \mathbb{N}_0$):

$$\begin{aligned}\delta_{q^{1/2}}(p) &= \text{vol}_q^{-1/2} \mathcal{F}_{q^{1/2}}(1)(p) = \text{vol}_q^{-1/2} \frac{\mathcal{F}_{q^{1/2}}(D_q^n x^n)(p)}{[[n]]_q!} \\ &= \text{vol}_q^{-1/2} \frac{(-i)^n}{[[n]]_q!} (\Lambda p)^n \mathcal{F}_{q^{1/2}}(x^n)(p).\end{aligned}\quad (267)$$

Solving for $\mathcal{F}_{q^{1/2}}(x^n)$ gives:

$$\begin{aligned}\mathcal{F}_{q^{1/2}}(x^n)(p) &= i^n \text{vol}_q^{1/2} [[n]]_q! (\Lambda p)^{-n} \delta_{q^{1/2}}(p) \\ &= i^n \text{vol}_q^{1/2} q^{-n(n+1)/2} [[n]]_q! p^{-n} \delta_{q^{1/2}}(p).\end{aligned}\quad (268)$$

From this result, we obtain a formula to calculate the $q^{1/2}$ -Fourier transform of a function that has a power series about the origin:

$$\mathcal{F}_{q^{1/2}}(f)(p) = \text{vol}_q^{1/2} \delta_q(p) \sum_{n=0}^{\infty} i^n q^{-n(n+1)/2} p^{-n} (D_q^n f)(0). \quad (269)$$

Finally, we treat the case of the negative exponents. Using Eq. (193) of the previous chapter and Eq. (258), we obtain:

$$\begin{aligned}\mathcal{F}_{q^{1/2}}(x^{-n-1})(p) &= \frac{(-1)^n q^{n(n+1)/2}}{[[n]]_q!} \mathcal{F}_{q^{1/2}}(D_q^n x^{-1})(p) \\ &= \frac{i^n q^{n(n+1)/2}}{[[n]]_q!} (\Lambda p)^n \mathcal{F}_q(x^{-1})(p) \\ &= \frac{i^{n+1} (1 + q^{1/2})}{4 [[n]]_q!} \text{vol}_q^{1/2} p^n \text{sgn}(p).\end{aligned}\quad (270)$$

Analogous to the previous chapter, we derive **expressions for the inverse $q^{1/2}$ -Fourier transforms**. This way, we have [cf. Eq. (213) of Chap. 6 and Eq. (226)]

$$\begin{aligned}\mathcal{F}_{q^{1/2}}^{-1}(\theta_{q^{1/2}}^+ - \theta_{q^{1/2}}^-)(x) &= \\ &= \text{vol}_q^{-1/2} \underline{\mathcal{S}}_x \left(\int_0^{p_0 \cdot \infty} d_{q^{1/2}} p \exp_q(ip|qx) - \int_{-p_0 \cdot \infty}^0 d_{q^{1/2}} p \exp_q(ip|qx) \right) \\ &= -\frac{4i}{(1 + q^{1/2}) \text{vol}_q^{1/2} x}\end{aligned}\quad (271)$$

and³ [cf. Eq. (224) of the previous chapter]

$$\begin{aligned}\mathcal{F}_{q^{1/2}}^{-1}(p^n \text{sgn}(p))(x) &= i^n q^{-n(n+1)/2} \Lambda^{-n} D_{q,x}^n \mathcal{F}_{q^{1/2}}^{-1}(\text{sgn}(p))(x) \\ &= (-i)^{n+1} [[n]]_q! \frac{4}{(1 + q^{1/2}) \text{vol}_q^{1/2}} x^{-n-1}.\end{aligned}\quad (272)$$

³Note that $\text{sgn} = \theta_{q^{1/2}}^+ - \theta_{q^{1/2}}^-$.

Moreover, it holds [cf. Eq. (214) of the previous chapter]

$$\begin{aligned}\mathcal{F}_{q^{1/2}}^{-1}(p^{-1})(x) &= \text{vol}_q^{-1/2} \bar{\mathcal{S}}_x \left(\int_{-p_0 \cdot \infty}^{p_0 \cdot \infty} d_{q^{1/2}} p p^{-1} \exp_q(ip|qx) \right) \\ &= -\frac{i}{4} (1 + q^{1/2}) \text{vol}_q^{1/2} \text{sgn}(x)\end{aligned}\quad (273)$$

and [cf. Eq. (215) of the previous chapter]

$$\begin{aligned}\mathcal{F}_{q^{1/2}}^{-1}(1)(x) &= \mathcal{F}_{q^{1/2}}^{-1}(pp^{-1})(x) = iq^{-1} \Lambda^{-1} D_{q,x} \mathcal{F}_{q^{1/2}}^{-1}(p^{-1})(x) \\ &= \text{vol}_q^{1/2} \delta_{q^{1/2}}(x).\end{aligned}\quad (274)$$

We can use Eq. (273) to derive an expression for $\mathcal{F}_{q^{1/2}}^{-1}(p^{-n-1})$ with $n \in \mathbb{N}$ [cf. Eq. (220) of the previous chapter]. This way, we get:

$$\mathcal{F}_{q^{1/2}}^{-1}(p^{-n-1})(x) = \frac{(1 + q^{1/2}) \text{vol}_q^{1/2}}{4} \frac{(-i)^{n+1} q^{n(n+1)/2}}{[[n]]_q!} x^n \text{sgn}(x).\quad (275)$$

Similarly, we obtain an expression for $\mathcal{F}_{q^{1/2}}^{-1}(p^n)$ with $n \in \mathbb{N}$ by using Eq. (274) [cf. Eq. (217) of the previous chapter]:

$$\mathcal{F}_{q^{1/2}}^{-1}(p^n)(x) = i^{-n} \text{vol}_q^{1/2} [[n]]_q! x^{-n} \delta_{q^{1/2}}(x).\quad (276)$$

Consequently, we have the following formula for a function f being analytic at the origin:

$$\mathcal{F}_{q^{1/2}}^{-1}(f(p))(x) = \text{vol}_q^{1/2} \delta_{q^{1/2}}(x) \sum_{n=0}^{\infty} i^{-n} (D_{q,p}^n f)(0) x^{-n}.\quad (277)$$

In Eqs. (221) and (222) of the previous chapter, we can replace q by $q^{1/2}$ due to the identities

$$\begin{aligned}\mathcal{F}_{q^{1/2}}^{-1}(\delta_{q^{1/2}}(p))(x) &= \text{vol}_q^{-1/2}, \\ \mathcal{F}_{q^{1/2}}^{-1}(p^{-n} \delta_{q^{1/2}}(p))(x) &= \text{vol}_q^{-1/2} q^{n(n+1)/2} \frac{(-ix)^n}{[[n]]_q!},\end{aligned}\quad (278)$$

and ($n \in \mathbb{N}$)

$$\begin{aligned}\mathcal{F}_{q^{1/2}}(x^{-n} \delta_{q^{1/2}}(x))(p) &= \text{vol}_q^{-1/2} \frac{(ip)^n}{[[n]]_q!}, \\ \mathcal{F}_{q^{1/2}}(x^n \text{sgn}(x))(p) &= \frac{i^{n+1} 4 [[n]]_q!}{(1 + q^{1/2}) \text{vol}_q^{1/2} q^{n(n+1)/2}} p^{-n-1}.\end{aligned}\quad (279)$$

References

- [1] M. A. Olshanetsky and V.-B. K. Rogov. The q -Fourier transform of q -distributions. arXiv:q-alg/9712055v2, 1998.
- [2] H. Wachter. Momentum and position representations for the q -deformed euclidean space. arXiv:math-ph/1910.02283, 2019.
- [3] R. Askey, N. M. Atakishiyev, and S. K. Suslov. *An Analog of the Fourier Transformation for A q -Harmonic Oscillator*. Springer US, Boston, MA, 1993. arXiv:hep-th/0306134.
- [4] T. H. Koornwinder. Special functions and q -commuting variables. arXiv:q-alg/9608008, 1996.
- [5] J. Seifert. *q -Deformierte Ein-Teilchen-Quantenmechanik*. Dissertation, Fak. f. Phys., LMU München, 1996.
- [6] B. L. Cerchiai, R. Hinterding, J. Madore, and J. Wess. A calculus based on a q -deformed Heisenberg algebra. *Eur. Phys. J. C*, 8:547–558, 1999.
- [7] F. N. Jackson. q -Difference equations. *Amer. J. Math.*, 32:305–314, 1910.
- [8] A. U. Klimyk and K. Schmüdgen. *Quantum groups and their representations*. Springer, Berlin - Heidelberg - New York, 1997.
- [9] S. Majid. Free braided differential calculus, braided binomial theorem and the braided exponential map. *J. Math. Phys.*, 34:4843–4856, 1993.
- [10] G. Gasper and M. Rahman. *Basic Hypergeometric Series*. Cambridge University Press, Cambridge/UK, 1990.
- [11] C. Chryssomalakos and B. Zumino. Translations, integrals and Fourier transforms in the quantum plan. In A. Ali, J. Ellis, and S. Randjbar-Daemi, editors, *Salamfestschrift, Proceedings of the Conference on Highlights of Particle and Condensed Matter Physics*, ICTP, Trieste/Italy, 1993.
- [12] V.-B. K. Rogov. q -Convolution and its q -Fourier transform. arXiv:q-alg/0010094, 2000.
- [13] V. Kac and P. Cheung. *Quantum Calculus*. Universitext. Springer, New York - Berlin - Heidelberg, 2002.
- [14] F. N. Jackson. On q -definite integrals. *Quart. J. Pure and Appl. Math.*, 41:193–203, 1910.
- [15] A. Kempf and S. Majid. Algebraic q -integration and Fourier theory on quantum and braided spaces. *J. Math. Phys.*, 35:6802–6837, 1994.
- [16] A. L. Ruffing. *Quantensymmetrische Quantentheorie und Gittermodelle für Oszillatorwechselwirkungen*. Dissertation, Fak. f. Phys., LMU München, 1996.

- [17] C. Jambor. *Non-Commutative Analysis on Quantum Spaces*. Dissertation, Fak. f. Phys., LMU München, 2004.
- [18] A. Lavagno. Basic-deformed quantum mechanics. *Rept. Math. Phys.*, 64:79–91, 2009.
- [19] R. Hinterding. *Eindimensionale q -deformierte Quantenmechanik und Streuprobleme*. Dissertation, Fak. f. Phys., LMU München, 2000.
- [20] J. Wess. q -Deformed Heisenberg algebras. In H. Gausterer, H. Grosse, and L. Pittner, editors, *Proceedings of the 38. Internationale Universitätswochen für Kern- und Teilchenphysik*, volume 543 of *Lec. Notes in Phys.* Springer, 2000.
- [21] S. Grossmann. *Funktionalanalysis im Hinblick auf Anwendungen in der Physik*. Studententext Physik. Aula-Verlag, Wiesbaden, 1988.