

An Algorithm for the Assignment Game Beyond Additive Valuations

Eric Balkanski*, Christopher En† and Yuri Faenza‡

Columbia University, IEOR Department

June 21, 2024

Abstract

The assignment game, introduced by Shapley and Shubik (1971), is a classic model for two-sided matching markets between buyers and sellers. In the original assignment game, it is assumed that payments lead to transferable utility and that buyers have unit-demand valuations for the items being sold. Two important and mostly independent lines of work have studied more general settings with imperfectly transferable utility and gross substitutes valuations. Multiple efficient algorithms have been proposed for computing a competitive equilibrium, the standard solution concept in assignment games, in these two settings. Our main result is an efficient algorithm for computing competitive equilibria in a setting with both imperfectly transferable utility and gross substitutes valuations. Our algorithm combines augmenting path techniques from maximum matching and algorithms for matroid intersection. We also show that, in a mild generalization of our model, computing a competitive equilibrium is NP-hard.

1 Introduction

The assignment game, introduced over half a century ago by [Shapley and Shubik \(1971\)](#), is a standard model to study two-sided markets between buyers and sellers. One of the main solution concepts in this game, the competitive equilibrium, is an efficient outcome composed of allocations and payments where the buyers are envy-free. Analyzing and computing competitive equilibria in the assignment game has applications in housing, hiring, and internet advertising markets. In the original model, each seller sells a single copy of an item (unit supply), and each buyer is interested in acquiring a single item (unit demand). Buyers and sellers are assumed to have *transferable utility* (TU), meaning that payments are used to share surplus between agents without losing utility to frictions. [Shapley and Shubik \(1971\)](#) showed that competitive equilibria always exist by casting the problem as a linear program. This existential result thus also serves as an efficient algorithm for computing the entire set of competitive equilibria. There has since been substantial work studying various extensions of the assignment game. The first main area of extension is to *imperfectly transferable utility* (ITU), which is when frictions, taxes, or fees impede the transfer of money between agents. The second is with more complex valuation functions, in particular *gross substitutes* valuations, which describe substitutable goods. Since its introduction by [Kelso Jr and Crawford \(1982\)](#), gross substitutes has proven to be a very general yet powerful condition on valuations that, with a simple no-complementarities requirement, captures many desirable properties.

A first body of work has relaxed the transferable utility assumption and instead considers settings with *imperfectly transferable utility* (ITU). This relaxation is motivated by the various impediments to the transfer of utility between agents, such as nonlinear taxes or fees a buyer may face when making a trade. [Kaneko \(1982\)](#) and [Quinzii \(1984\)](#) gave nonconstructive proofs that competitive equilibria exist when buyers have unit demands, even under ITU, and [Demange and Gale \(1985\)](#) described various structural properties of the set of competitive equilibria under unit demands and ITU. Unlike in the TU setting, the initial existential and structural results under ITU do not imply efficient algorithms for computing competitive equilibria. A subsequent line of work has thus focused on the computational

*eb3224@columbia.edu

†Corresponding Author: ce2456@columbia.edu

‡yf2414@columbia.edu

side of competitive equilibria under ITU. In particular, [Alkan and Gale \(1990\)](#) provided an algorithm for computing a competitive equilibrium in this setting, though they did not provide a polynomial bound on the running time. [Alkan \(1992\)](#) later improved the algorithm and showed that the new version runs in polynomial time.

Another rich line of work that generalizes the initial model of [Shapley and Shubik \(1971\)](#) has relaxed the unit demand assumption for buyers in the market by considering richer classes of valuation functions. [Crawford and Knoer \(1981\)](#) extend the assignment game to a many-to-one model where each buyer may match with multiple sellers, and prove the existence of competitive equilibria. In their model, buyers are assumed to have *separable* (also known as *additive*) valuations, and to demand a fixed number of matches. [Kelso Jr and Crawford \(1982\)](#) consider an even broader class of valuation functions by introducing *gross substitutes* valuations as a sufficient condition for the existence of competitive equilibria when buyers have valuations over substitutable items. Since this introduction, gross substitutability has proven to be a very general yet powerful condition on valuations that, with a simple no-complementarities requirement, captures many desirable properties. On the computational side, [Kelso Jr and Crawford \(1982\)](#) show that under gross substitutes valuations, a natural ascending auction procedure called *Walrasian tâtonnement* is guaranteed to find a competitive equilibrium. While this process is continuous and primarily used as a proof device in [Kelso Jr and Crawford \(1982\)](#), it was later adapted to run in polynomial time (see, e.g., [\(Paes Leme, 2017\)](#)). [Murota \(1996b\)](#) and [Murota \(1996c\)](#) present the first purely combinatorial cycle-cancelling algorithm for the same problem that runs in strongly polynomial time.

Even though gross substitutes and imperfectly transferable utilities have been studied mostly independently, there are many application areas where markets exhibit both item substitutabilities and nonlinear frictions, taxes, or fees. In hiring markets, there is some amount of substitutability between the skillsets of workers, and scaling taxes on salaries causes the utility of firms and workers to not be perfectly transferable. In online ad marketplaces, advertisers wish to buy clicks or impressions in various online mediums, and different mediums allow for varying levels of user engagement. These scenarios have motivated the study of generalized assignment games that feature both ITU and gross substitutes valuations. As we discuss in more detail in [Section 1.2](#), even though some existential and structural results have been shown in similar or more general settings, there is, to the best of our knowledge, no existing computational result in such a setting. The combination of the nonlinearity in the buyers' valuations (gross substitutes) and the nonlinearity of their loss in utility due to frictions (ITU) introduces novel algorithmic challenges that are further discussed in [Section 1.1](#).

Our main contribution is a polynomial-time algorithm for computing a competitive equilibrium under imperfectly transferable utilities and gross substitutes valuations. Our model for ITU, which we call the QITU model (short for Quasilinear and Imperfectly Transferable Utility model, and described formally in [Section 2.1](#)) with gross substitutes valuations generalizes the gross substitutes TU models of [Kelso Jr and Crawford \(1982\)](#) and the unit-demand ITU models of [Demange and Gale \(1985\)](#), [Alkan and Gale \(1990\)](#), and [Alkan \(1992\)](#).

Theorem 1. *In the QITU model with gross substitutes valuations, there is a polynomial time algorithm that computes a competitive equilibrium.*

As in previous models, we take as input oracles for the valuation functions of each buyer, plus the full description of functions defining the rate at which utility can be transferred between agents. In addition to achieving the first efficient algorithm under ITU and gross substitutes valuations, this result also unifies many of the existing algorithmic results in the previously listed models that are special cases of the QITU model with gross substitutes valuations.

We next show that the assumption that buyers have quasilinear utilities is required for our result by considering a relaxation of our setting, called the NQ model, where a single buyer's utility is a minor generalization of a quasilinear utility function. In particular, this buyer's utility function is piecewise linear in money, with a single breakpoint. See [Section 6](#) for a formal definition. This model is of interest as a way to model budgets or risk aversion in buyers, which are common properties in many real-world markets. However, we prove a negative result showing that computing a competitive equilibrium in this setting is NP-hard.

Theorem 2. *Computing a competitive equilibrium in the NQ model is NP-hard.*

The paper is organized as follows. We conclude the introductory section with an overview of the techniques used to prove [Theorem 1](#) ([Section 1.1](#)) and a discussion of additional related work ([Section 1.2](#)). In [Section 2](#), we formally introduce the QITU model and state the problem of interest. In [Section 3](#), we introduce the basic structures the algorithm from [Theorem 1](#) operates on and formally describe the

algorithm. In Section 4, we introduce and analyze the main subroutine of the algorithm, used to increase prices in the spirit of Walrasian tâtonnement. This is the most technical section and the main technical contribution of the paper. In Section 5, we analyze the remaining subroutines and combine our results into the proof for Theorem 1. In Section 6, we present Theorem 2. Appendix A contains helper results on gross substitutes valuations used in the other sections. Missing proofs from Sections 3, 4, and 6 appear in Appendix B, C, and D, respectively.

1.1 Technical overview

Similar to auction mechanisms for the classical max weight bipartite matching problem (e.g., (Demange et al., 1986)) and for certain generalizations of the assignment game including Alkan (1992); Dütting et al. (2011), our algorithm alternates the search for an alternating path with a procedure that increases the prices of certain items. We introduce the concept of *partial stability*, a relaxed version of the equilibrium property, which we show is maintained as invariant throughout the algorithm. This invariant allows us to extend previous techniques to the setting with additive utilities where each buyer may match multiple sellers, and each seller has multiple copies of its item that can match multiple buyers, i.e., the many-to-many setting. The alternating path subroutine is employed to increase the size of the current matching while preserving partial stability. The non-uniform price increase kicks in when, with the current prices, no larger partially stable matching exists. To select the correct price increase, we employ a modified version of the Walrasian tâtonnement procedure on a *marginal demand graph*, which indicates which buyers demand which items at the current set of prices. We further show that this procedure converges to an equilibrium.

The main differences from Alkan (1992) and Dütting et al. (2011), and the main obstacles for designing our algorithm, arise from the nonlinearity of gross substitutes valuations (as opposed to the many-to-many case with additive utilities) that causes the marginal value that a buyer has for an item to change when changes to the current matching are made. Due to these changes in marginal value, the *indifference sets* of other items that a buyer can swap for without losing utility also change. As a result, instead of finding augmenting paths on a fixed graph, the algorithm operates on a marginal demand graph that evolves with each change to the current partial matching. Thus, finding an augmenting path becomes more complex because we must preserve partial stability. We also note that an augmenting path in the marginal demand graph may not preserve buyer utilities under gross substitutes valuations. Instead, we take advantage of matroidal properties of gross substitutes demand sets to find augmenting paths that do not reduce buyer utility.

Furthermore, when calculating price increases, Dütting et al. (2011) use a linear programming duality trick seen in Alkan (1989) to compute the price increase via a maximum weight matching problem. The nonadditive nature of our setting means this trick no longer applies directly, as the entire family of demanded sets for any given buyer is not easily expressible within a static marginal demand graph or a single linear program. Instead, we can leverage properties of gross substitutes valuations to solve a related weighted common matroid basis problem and then use the solution to define an appropriate linear program. From there, we can reapply the duality trick to compute the desired price increase.

1.2 Further related work

A large body of work extends the models of Shapley and Shubik (1971) and Demange and Gale (1985). Rochford (1984) presents an alternative solution concept for the unit demand TU model based on “threat utility” and shows that these solutions also always exist within the set of competitive equilibria. Roth and Sotomayor (1988) show that the same results hold in the setting with unit demand and ITU. Hatfield and Milgrom (2005) present a very general setting encompassing gross substitutes valuations and ITU, though prices are assumed to be drawn from a finite set. They show that the set of competitive equilibria is nonempty and forms a bounded lattice, and present a generalization of the Gale-Shapley algorithm for computing competitive equilibria. However, because of the discretization of prices, their model is incomparable to ours. Hatfield et al. (2013) present a more complex model that removes the two-sided assumption, instead allowing all agents to simultaneously make multiple trades as both a buyer and a seller. They identify *full substitutability*, a generalization of gross substitutability that accounts for both purchases and sales, as a sufficient condition for the existence of competitive equilibria. Fleiner et al. (2019) then develop a similar one-sided matching model that incorporates ITU, encompassing our model. They show that competitive equilibria are guaranteed to exist under full substitutability, though their results rely on fixed-point theorems and are thus intrinsically non-constructive.

There has also been substantial work studying gross substitutes, as this class of functions naturally appears in many other contexts. Nisan and Segal (2006) provide another competitive equilibrium algorithm for the gross substitutes TU model of Kelso Jr and Crawford (1982) based on linear programming, which requires a separation-based algorithm such as the ellipsoid method. Dress and Wenzel (1990) define *valuated matroids* as an extension of matroid rank functions to settings with weights. Dress and Terhalle (1995a) introduce *matroidal maps* as the exact class of functions for which a greedy function implements a demand oracle at any set of prices. Murota (1996a) introduces M^{\natural} -concavity as a natural extension of convexity to the integral lattice. Fujishige and Yang (2003) then show that these concepts are all equivalent to gross substitutes.

Algorithmically, there has been less work beyond the basic ITU model of Demange and Gale (1985) and the gross substitutes model of Kelso Jr and Crawford (1982). Colini-Baldeschi et al. (2015) develop a polynomial algorithm for a setting inspired by online ad auctions where buyers demand multiple items, but valuations for items are correlated across buyers, and utility is perfectly transferable except for a single discontinuity, used to model a budget constraint. Goel et al. (2015) consider a related setting where items have public valuations and allocations must satisfy certain polyhedral or polymatroidal constraints. Dütting et al. (2015) provide a polynomial algorithm for a setting with private budgets, where values are additive and correlated across buyers.

2 Preliminaries

2.1 Problem statement

In the QITU model, there is a set B of n buyers and a set S of m items. Each item $j \in S$ has a capacity $cap(j)$. Each buyer $i \in B$ has a valuation function $v_i : 2^S \rightarrow \mathbb{R}$ over bundles of items. We assume that the algorithm is given oracle access to each v_i and that evaluating $v_i(T)$ for any $T \subseteq S$ takes $O(1)$ runtime. For each buyer-item pair (i, j) , there is an *effective price* function $q_{ij} : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ where $q_{ij}(p_j)$ represents the utility lost by agent i for paying price p_j for item j . *Imperfectly transferable utility* is the case where the effective price functions need not be the identity. In particular, we assume that q_{ij} is piecewise linear, continuous, strictly increasing in p_j , and that $q_{ij}(0) = 0$. Also, there is some price $p_j < \infty$ such that $q_{ij}(p_j) = v_i(\{j\})$. The right derivative of the effective price function is denoted q'_{ij} . The utility function $u_i : 2^S \times \mathbb{R}_+^S \rightarrow \mathbb{R}$ of agent i is defined as

$$u_i(T, p) := v_i(T) - \sum_{j \in T} q_{ij}(p_j). \quad (1)$$

In this paper, on top of the assumptions of the QITU model, we assume that buyers have valuation functions satisfying *gross substitutes*. The gross substitutes property states that for an item j that is demanded by buyer i at prices p , if the prices of some subset of items that does not include j increase, then j must still be demanded at the new prices.

Definition 3 (Gross Substitutes Kelso Jr and Crawford (1982)). The *demand correspondence* $F_p(i)$ of a buyer is the family of sets of items that maximize utility for buyer i , under prices $p = (p_1, \dots, p_m)$:

$$F_p(i) := \{T \subseteq S \mid u_i(T, p) \geq u_i(T', p), \forall T' \subseteq S\} \quad (2)$$

Any $T \in F_p(i)$ is called a *demand set at price p* . A valuation function v_i satisfies *gross substitutes* if for any price vectors $p, p' \in \mathbb{R}^S$ such that $p \leq p'$, and $T \in F_p(i)$ there is a set $T' \in F_{p'}(i)$ such that $T \cap \{j \mid p_j = p'_j\} \subseteq T'$.

Gross substitutes valuation functions capture a “no complementarities” condition between items, so that all items are substitutable goods. While the definition given here is from classical demand theory, and appears to be a quality of the utility function, in Appendix A we present alternative definitions that explicitly describe the valuation function without relying on prices.

A *many-to-many matching* $\mu \subseteq B \times S$ is a subset of buyer-item pairs such that each item j is matched to at most $cap(j)$ buyers, and each buyer is matched to each item at most once (no multi-edges). We will also write $\mu(i)$ to denote the set of items i is matched to, and $\mu(j)$ as the set of buyers to which j is matched. An *outcome* is a pair (μ, p) with a matching μ and a vector of prices $p = (p_1, \dots, p_m)$ for each item. An item j is *fully matched* if $|\mu(j)| = cap(j)$, and is *undermatched* otherwise. An outcome is *feasible* if all items that are not fully matched have price 0.

The primary solution concept we will be concerned with is that of *competitive equilibrium*, which is a feasible outcome such that each buyer is matched to a set of items that maximizes their utility at the prices specified by the outcome.

Definition 4 (Competitive equilibrium). An outcome (μ, p) is a *competitive equilibrium* if the outcome is feasible, and for every buyer i , we have $\mu(i) \in F_p(i)$.

Our goal is to design an algorithm that, given an input to the QITU model with gross substitute valuations, returns a competitive equilibrium in time polynomial in the input size.

Remark 5. Note that previous one-to-one models such as those presented by [Shapley and Shubik \(1971\)](#) and [Demange and Gale \(1985\)](#) have “sellers” instead of “items,” where a seller j has some nonnegative reserve value s_j for their endowed item. There is a reduction from the problem of finding a competitive equilibrium in the setting with sellers to the problem of finding a competitive equilibrium in the setting with items. To see this, given an instance of a problem with sellers, consider the following edited instance. Set each seller’s reserve price to 0, and let each buyer i with value v_{ij} for item j have a new value $v_{ij} - s_j$. Then, given an equilibrium (μ, p) in the original instance, we can obtain an equilibrium in the edited instance given by (μ, p') where $p'_j = p_j - s_j$. Similarly, given any equilibrium (μ, p') in the edited instance, we can obtain an equilibrium in the original instance given by (μ, p) , where $p_j = p'_j + s_j$.

Throughout the remainder of the paper until Section 6, we work exclusively within the QITU model. In particular, in Section 2.2 and Appendix A, we introduce several equivalent definitions of gross substitutes that are only valid under the assumptions of the QITU model.

2.2 Gross substitutes and stability

The *marginal value* of a set of items T' given T is defined as

$$v_i(T' \mid T) := v_i(T \cup T') - v_i(T) \quad (3)$$

Similarly, the *marginal utility* of T' given T at prices p is defined as

$$u_i(T' \mid T, p) := u_i(T \cup T', p) - u_i(T, p) \quad (4)$$

To simplify definitions and proofs, we assume the existence of $|S|$ dummy items with infinite capacity, with the property that for any buyer b_i , any dummy item j_0 , and any bundle of items T , the marginal value of the dummy item is $v_i(j_0 \mid T) = 0$. Notice that this assumption is without loss of generality since, at any competitive equilibrium, each dummy item will not be fully matched, which means it must have price $p_{j_0} = 0$. Thus, $u_i(j_0 \mid T, p) = 0$ for all buyers i and all bundles T . Essentially, dummy items are always worth nothing and cost nothing.

[Gul and Stacchetti \(2000\)](#) showed a helpful equivalent definition of gross substitutes, known as the *single improvement property*.

Lemma 6 (Single improvement property (SI) [Gul and Stacchetti \(2000\)](#)). *A valuation function v_i satisfies gross substitutes if and only if given any price vector p and any $T, T' \subseteq S$ such that $u_i(T, p) < \max_{T' \subseteq S} u_i(T', p)$, then there exist X, Y with $|X|, |Y| \leq 1$ such that $u_i(T \cup X \setminus Y, p) > u_i(T, p)$.*

Using Lemma 6, we can show that, under gross substitutes valuations, competitive equilibria coincide with the notion of (pairwise) stability from matching theory.

Definition 7 (Stability). An outcome (μ, p) is *stable* if every buyer i has no profitable *additions*, *swaps*, or *drops*:

- **Additions:** For every item $j \notin \mu(i)$, we have $u_i(j \mid \mu(i), p) \leq 0$.
- **Swaps:** For every $j_1 \in \mu(i)$ and $j_2 \notin \mu(i)$, we have $u_i(\mu(i), p) \geq u_i(\mu(i) \cup \{j_2\} \setminus \{j_1\}, p)$.
- **Drops:** For every $j \in \mu(i)$, we have $u_i(\mu(i) \setminus \{j\}, p) \leq u_i(\mu(i), p)$.

Intuitively, stability ensures an outcome is “fair” since each buyer gets their favorite bundle of items at the current prices.

Lemma 8 (Competitive equilibrium coincides with feasible and stable). *An outcome is a competitive equilibrium if and only if it is feasible and stable.*

Proof. First, let (μ, p) be a competitive equilibrium. By definition, (μ, p) is feasible. If there was a profitable addition, swap, or drop for buyer i from the initial bundle $\mu(i)$, then the resulting bundle would have greater utility, contradicting the definition of a competitive equilibrium. Thus, (μ, p) is stable.

Let (μ, p) be feasible and stable. We wish to show $\mu(i) \in F_p(i)$. This is equivalent to showing that $u_i(\mu(i), p) \geq u_i(T, p)$ for all $T \subseteq S$. Suppose for contradiction that there existed T such that $u_i(\mu(i), p) < u_i(T, p)$. Then, by Lemma 6, there exists $X, Y \subseteq S$ with $|X|, |Y| \leq 1$ such that $u_i(\mu(i) \cup X \setminus Y, p) > u_i(\mu(i), p)$. If $|X| = |Y| = 1$, this corresponds to a profitable swap. If $|X| = 1$ and $|Y| = 0$, this corresponds to a profitable addition. If $|X| = 0$ and $|Y| = 1$, this corresponds to a profitable drop, contradicting stability. We conclude that such T cannot exist, so (μ, p) is a competitive equilibrium. \square

2.3 Matroids

A key component of the algorithm and the analysis is matroids.

Definition 9 (Matroid). A matroid M is a pair (E, \mathcal{I}) , where E is a finite *ground set* and \mathcal{I} is a family of subsets of E known as *independent sets*, that satisfies:

1. $\emptyset \in \mathcal{I}$.
2. If $A_1 \in \mathcal{I}$ and $A_2 \subseteq A_1$, then $A_2 \in \mathcal{I}$.
3. If $A_1, A_2 \in \mathcal{I}$ and $|A_1| > |A_2|$, then there exists $x \in A_1 \setminus A_2$ such that $A_2 \cup \{x\} \in \mathcal{I}$.

A *basis* $T \in \mathcal{I}$ of a matroid is a maximal independent set.

We will primarily make use of the well known basis exchange property (see, e.g., Oxley (2006)).

Lemma 10 (Basis exchange property). *Let T_1, T_2 be two distinct bases of a matroid, and let $j_1 \in T_1 \setminus T_2$. Then, there exists $j_2 \in T_2 \setminus T_1$ such that $T_1 \cup \{j_2\} \setminus \{j_1\}$ is also a basis.*

We will take advantage of existing algorithms for solving minimum weight matroid intersection problems. These algorithms require access to independence oracles, defined below.

Definition 11 (Independence oracle). Given a matroid $M = (E, \mathcal{I})$, an independence oracle for M is a subroutine which indicates whether a set $F \subseteq E$ is a member of \mathcal{I} .

3 The Algorithmic Framework

In this section, we first introduce the main structures necessary to present the algorithm. In Section 3.1, we introduce the *marginal demand graph*, which is the basic graph the algorithm operates on. In Section 3.2, we introduce *maximal alternating trees*, which are the fundamental structure used in the algorithm. In Section 3.3, we describe the main algorithm in detail.

To describe and analyze our algorithm, it is helpful to think of a buyer i as a collection of $|S|$ identical unit-demand buyers, which we refer to as *unit-buyers*. Each unit-buyer can be matched to at most one item. Thus, we define $\text{ubuy}(i)$ to be this set of unit-buyers corresponding to i , with $|\text{ubuy}(i)| = |S|$, and we define the set U of all the unit-buyers as $U = \bigcup_{i \in B} \text{ubuy}(i)$. Given a unit-buyer k , we let $\text{buy}(k)$ be the buyer such that $k \in \text{ubuy}(\text{buy}(k))$. Let $\text{copy}(k) = \text{ubuy}(\text{buy}(k))$ be the set of all unit-buyers who share the same original buyer with k . When it is clear from context, given a unit-buyer k , we often write $u_k := u_{\text{buy}(k)}$ to refer to the utility function of the buyer associated with k . Similarly, we often write $q_{kj} = q_{\text{buy}(k)j}$ to refer to the effective price function of the buyer associated with k . We refer to this setting as the *many-to-one market* and to the original one as the *many-to-many market*.

3.1 Marginal demand graphs

A *many-to-one outcome* (ν, p) is a matching $\nu \subseteq U \times S$ and a set of prices p such that each unit-buyer is matched to at most one item, each item j is matched to at most $\text{cap}(j)$ unit-buyers, and no two unit-buyers corresponding to the same buyer are matched to the same item. An outcome (μ, p) of the many-to-many market is said to be the *many-to-many projection of the many-to-one outcome* (ν, p) if $(i, j) \in \mu$ if and only if $(k, j) \in \nu$ for some $k \in \text{ubuy}(i)$. We define a many-to-one outcome (ν, p) to be stable and feasible if its many-to-many projection is stable and feasible, respectively. Similarly, we define a many-to-one outcome to be a competitive equilibrium if its many-to-many projection is a competitive equilibrium.




Figure 1: An example of a marginal demand graph and MAT. Note that $\text{ubuy}(1) = \{1, 2\}$, $\text{ubuy}(2) = \{3, 4\}$, and $\text{ubuy}(3) = \{5, 6\}$. Solid lines represent the current matching, and dashed lines represent unmatched edges in each (unit-)buyer’s demanded set.

Definition 12 (Partial stability). An outcome (μ, p) in the many-to-many market is *partially stable* if for all buyers i and all sets of items T with $|T| \leq |\mu(i)|$, we have $u_i(T, p) \leq u_i(\mu(i), p)$. A many-to-one outcome (ν, p) is defined to be partially stable if its many-to-many projection is partially stable.

When the price vector p is clear from context, we may also refer to the matching μ as partially stable. Partial stability means that a buyer’s current matched bundle is their most preferred bundle of the current size (or smaller). Using the (SI) property (Lemma 6), partial stability can also be interpreted as “no profitable swaps or drops.”

Given a partially stable many-to-one outcome (ν, p) , define the *marginal demand correspondence* of a unit-demand buyer k as the set of items $F_{\nu, p}(k) \subseteq S$ where $j_1 \in F_{\nu, p}(k)$ if

- **Matched:** $j_1 = \nu(k)$, or
- **Potential alternative:** $j_1 \notin \nu(\text{copy}(k))$, k is matched to an item other than j_1 , and $u_k(j_1 \mid \nu(\text{copy}(k)) \setminus \nu(j_1), p) = 0$, or
- **Favorite item:** $j_1 \notin \nu(\text{copy}(k))$, k is unmatched in ν , and

$$u_k(j_1 \mid \nu(\text{copy}(k)), p) \geq \max \left\{ 0, \max_{j_2 \notin \nu(\text{copy}(k))} u_k(j_2 \mid \nu(\text{copy}(k)), p) \right\}.$$

Essentially, $F_{\nu, p}(k)$ contains the item j_1 matched to k , as well as any other item $\text{buy}(k)$ would swap with j_1 without decreasing their utility, as long as one of their copies isn’t already matched to it. If k is unmatched, then $F_{\nu, p}(k)$ contains each unmatched item whose addition to $\nu(\text{copy}(k))$ would lead to the largest utility increase, as long as the increase is nonnegative. The marginal demand correspondence of an item j is given by $F_{\nu, p}^{-1}(j) := \{k \mid j \in F_{\nu, p}(k)\}$. The marginal demand correspondence for a set of unit-buyers $U' \subseteq U$ is given by $F_{\nu, p}(U') := \bigcup_{k \in U'} F_{\nu, p}(k)$. Similarly, for a set of items $S' \subseteq S$, we define $F_{\nu, p}^{-1}(S') := \bigcup_{j \in S'} F_{\nu, p}^{-1}(j)$. Next, we define the main graph upon which the algorithm will operate.

Definition 13 (Marginal demand graph). Given a partially stable outcome (ν, p) , the *many-to-one marginal demand graph* $D(\nu, p)$ is the graph $(U \cup S, E)$ where $(k, j) \in E$ if and only if $j \in F_{\nu, p}(k)$.

Our first important result describes the conditions necessary to verify if an outcome is a competitive equilibrium. This result is based on the concept of maximal alternating trees, described next.

3.2 Maximal alternating trees

Definition 14 (Maximal alternating tree (MAT)). Let $H = (U \times S, E)$ be a graph, let $\nu \subseteq E$ be a many-to-one matching, and let $k_0 \in U$ be an unmatched unit-buyer. An *alternating tree* \mathcal{T} in H rooted at k_0 is a tree subgraph of H such that all paths from the root k_0 are alternating between edges in the matching ν and edges not in ν , beginning with an unmatched edge. An alternating tree \mathcal{T} is *maximal* if

- for every unit-buyer $k \in \mathcal{T}$, all items $j \in S$ with $(k, j) \in H$ are in \mathcal{T} , and
- for every item $j \in \mathcal{T}$, the unit-buyers matched to j are in \mathcal{T} .

When $H = D(\nu, p)$ is a marginal demand graph, these requirements can be rewritten as

- for every unit-buyer $k \in \mathcal{T}$, we have $F_{\nu,p}(k) \subseteq \mathcal{T}$, and
- for every item $j \in \mathcal{T}$, we have $\nu(j) \subseteq \mathcal{T}$.

We give an example of a marginal demand graph and of a MAT in Figure 1.

Remark 15. MATs are a common structure seen in bipartite matching problems. It is a standard result that whenever a vertex (i.e. a unit-buyer) is not matched, a MAT exists and can be found efficiently using a process similar to a breadth-first search, which we leverage in Algorithm 4 and prove in Lemma 34.

Note that given the existence of dummy items, every competitive equilibrium can be expanded into a competitive equilibrium such that all unit-buyers are matched, by matching any unmatched unit-buyers to dummy items. This expansion does not change the utility of any buyer, so each buyer retains a demanded set at the current prices. We can show that the absence of a MAT serves as a certificate of a competitive equilibrium.

Lemma 16 (MATs and competitive equilibrium). *Let (ν, p) be a feasible and partially stable outcome. Then, (ν, p) is a competitive equilibrium if and only if there does not exist a MAT in $D(\nu, p)$.*

We defer the proof of this result to Appendix B. Since we know that a MAT exists if and only if an unmatched unit-buyer exists, this result shows that if we can match every unit-buyer while maintaining partial stability, we will arrive at a competitive equilibrium.

3.3 Description of the algorithm

MATs are the main structure that the algorithm will operate on. At each iteration, the algorithm identifies a MAT which we call the *current MAT*, and divides it into one of two categories: either it contains an undermatched item, or all of its items are all fully matched. In the first case, the algorithm finds an augmenting path from the unmatched root buyer k_0 to the undermatched item, and increases the size of the matching ν . In the latter case, it carefully increases the prices of items in the MAT until buyers in the MAT start demanding other items, thus increasing the size of the current MAT.

The algorithm is formally defined in Algorithm 1. Starting with prices $p = 0$ and an empty matching ν , it iterates until there is no unmatched unit-buyer k_0 . At each outer-iteration (lines 2-9), the algorithm constructs the marginal demand graph $D(\nu, p)$, and identifies a MAT \mathcal{T} rooted at an unmatched unit-buyer k_0 using subroutine FINDMAT (line 3) that is described in Section 5.1. If the MAT \mathcal{T} has an undermatched item, the algorithm uses the subroutine AUGMENTINGPATH, which is described in Section 5.2, to augment matching ν along an alternating path from the unmatched root buyer to that item (lines 8-9). Otherwise, the algorithm uses the subroutine FINDPRICEINCREASE, which is described in Section 4, to conduct a version of Walrasian tâtonnement by finding a price increase that increases the size of the MAT \mathcal{T} (lines 4-7). By repeating this process, the algorithm can iteratively augment the current matching, until no more buyers demand additional items.

ALGORITHM 1: Competitive equilibrium algorithm for the many-to-many assignment game

Input: Buyers B , sellers S , capacities cap , valuation functions v_i , effective price functions q_{ij}

Output: Competitive equilibrium (μ, p)

```

1 Initialize prices  $p_j \leftarrow 0$  for all  $j \in S$  and matching  $\nu \leftarrow \emptyset$ 
2 while there is a unit-buyer  $k_0 \in [|B||S|]$  s.t.  $\nu(k_0) = \emptyset$  do
3    $\mathcal{T} \leftarrow \text{FINDMAT}(k_0, \nu, D(\nu, p))$  /* Compute current MAT rooted at  $k_0$  */
4   while  $\exists j \in S(\mathcal{T})$  s.t.  $|\nu(j)| < cap(j)$  do /* loop: Walrasian tâtonnement */
5      $\nu, p' \leftarrow \text{FINDPRICEINCREASE}(\mathcal{T}, p, \nu)$ 
6      $p \leftarrow p + p'$ 
7      $\mathcal{T} \leftarrow \text{FINDMAT}(k_0, \nu, D(\nu, p))$  /* Update current MAT rooted at  $k_0$  */
8      $P \leftarrow \text{AUGMENTINGPATH}(\mathcal{T}, \nu)$  /* Once MAT has undermatched item, augment */
9      $\nu \leftarrow \nu \Delta P$ 
10  $\mu \leftarrow \{(\text{buy}(k), j) \mid (k, j) \in \nu\}$ 
11 return  $(\mu, p)$ 

```

4 The FindPriceIncrease Subroutine

The primary subroutine of Algorithm 1 and the main technical contribution of this paper is FINDPRICEINCREASE, described in Algorithm 3. Given a MAT \mathcal{T} where all items are fully matched, the

subroutine finds a direction vector d and step size $\lambda^* > 0$ such that after increasing the current set of prices by $p \leftarrow p + \lambda^*d$, the size of the MAT \mathcal{T} increases. This process is conceptually similar to the classical Walrasian tâtonnement process, although price increases are not uniform and thus much more difficult to compute. Proofs omitted from Section 4 are presented in Appendix C.

4.1 MAT-preserving price increases

In this section, we introduce MAT-preserving price increases, which are similar to structures from Alkan (1992) and Dütting et al. (2011). To do this, we need some additional notation.

Given a subgraph H of the marginal demand graph $D(\nu, p)$, we define $U(H) = U \cap V(H)$ and $S(H) = S \cap V(H)$, where $V(G)$ is the vertices of G . Also, define $B(H) = \{\text{buy}(k) \mid k \in U(H)\}$. Similarly, if E' is a set of edges in $D(\nu, p)$, then we define $V(E') = \{i \mid (i, j) \in E'\} \cup \{j \mid (i, j) \in E'\}$, as well as $U(E') = U \cap V(E')$, and $S(E') = S \cap V(E')$.

Definition 17 (MAT-preserving price increase). Given a partially stable and feasible many-to-one outcome (ν, p) , a *price increase* at (ν, p) is a vector $d \geq 0$ over the items S . A MAT-preserving price increase d with respect to the marginal demand graph $D(\nu, p)$ and a MAT \mathcal{T} rooted at k_0 where all items are fully matched, satisfies

- (a) $d_j > 0$ if and only if $j \in S(\mathcal{T})$.
- (b) There is some partially stable matching ν' at p and MAT \mathcal{T}' rooted at k_0 in $D(\nu', p)$ such that
 - i. ν' coincides with ν on $U \setminus U(\mathcal{T}) \times S \setminus S(\mathcal{T})$.
 - ii. $V(\mathcal{T}) = V(\mathcal{T}')$.
 - iii. For sufficiently small $\lambda > 0$, we have that $(\nu', p + \lambda d)$ is partially stable and $\mathcal{T}' \subseteq D(\nu', p + \lambda d)$.

Note that the current MAT rooted at k_0 changes from \mathcal{T} to \mathcal{T}' after the MAT-preserving price increase. The reason is that given a specific MAT \mathcal{T} , there may not exist a price increase that preserves exactly the edges of \mathcal{T} (as shown in Example 18 below). However, as we will later see in Algorithm 3, the matching ν can be adjusted to find a price increase that preserves the edges of another MAT \mathcal{T}' that contains the same unit-buyers and items. In this way, the size of the MAT rooted at k_0 remains the same after a sufficiently small price increase along d .

Example 18. Consider the MAT rooted at $k_0 = 3$ in Figure 2(a). The three unit-buyers correspond to distinct buyers. For each unit buyer k , we have $v_k(\mathcal{T}) = |\mathcal{T}|$. We have $p_1 = p_2 = 0$, and $\text{cap}(1) = \text{cap}(2) = 1$. Then, define the effective price functions as follows:

$$\begin{array}{lll} q_{11}(p_1) = 2p_1 & q_{21}(p_1) = p_1 & q_{31}(p_1) = p_1 \\ q_{12}(p_2) = p_1 & q_{22}(p_2) = 2p_2 & q_{32}(p_2) = p_2 \end{array}$$

We can verify that both $\nu = \{(1, 1), (2, 2)\}$ and $\nu' = \{(1, 2), (2, 1)\}$ are partially stable, and that the marginal demand graph $D(\nu, p)$ is a complete bipartite graph. However, there is no price increase d such that for small $\lambda > 0$, we have $u_1(2 \mid 1, p + \lambda d) \leq 0$ and $u_2(1 \mid 2, p + \lambda d) \leq 0$. Instead, we can change the matching to ν' and the MAT to \mathcal{T}' as given in Figure 2(b). Setting $d_1 = d_2 = 1$, we can verify that for small $\lambda > 0$, ν' is partially stable at $p + \lambda d$, and that $\mathcal{T}' \subseteq D(\nu', p + \lambda d)$.

The following lemma gives a sufficient condition for a price increase d to be a MAT-preserving price increase. This condition depends on the slopes $q'_{kj}(p)$ of the effective price functions $q_{kj}(p)$.

Lemma 19 (MAT-preserving price increase and slopes of effective price functions). *Given a partially stable outcome (ν, p) and MAT \mathcal{T} , let d be a price increase that satisfies condition (a) from Definition 17. Let ν' and \mathcal{T}' satisfy conditions (bi.) and (bii.). If $q'_{kj_1}(p_{j_1})d_{j_1} \leq q'_{kj_2}(p_{j_2})d_{j_2}$ for all $(k, j_1) \in \mathcal{T}'$ and $j_2 \in F_{\nu', p}(k)$, then d is a MAT-preserving price increase with respect to the marginal demand graph $D(\nu, p)$ and MAT \mathcal{T} .*

The proof of this result is deferred to Appendix C.1.




Figure 2: An example of a MAT to be preserved in a MAT-preserving price increase. Solid lines represent the current matching, and dashed lines represent the other edges in the marginal demand graph given the current matching.

Overview of the remainder of Section 4. To find a MAT-preserving price increase with respect to a marginal demand graph $D(\nu, p)$ and a MAT \mathcal{T} , it remains to find a price increase d , matching ν' , and a MAT \mathcal{T}' that satisfy the conditions of Lemma 19. The first step is to find a minimum weight perfect matching ν^* over $U(\mathcal{T}) \setminus \{k_0\} \times S(\mathcal{T})$ such that $\nu \setminus E(\mathcal{T}) \cup \nu^*$ is partially stable, where the weights depend on the slopes of the effective price functions. Section 4.2 shows such a matching can be found by solving a minimum weight common basis problem over two matroids. Section 4.3 then shows that this minimum weight perfect matching ν^* can be used to construct a linear program whose optimal solution is then transformed into a price increase that satisfies the conditions of Lemma 19. Then, in Section 4.4 we give the global description of the subroutine.

4.2 Finding a minimum weight partially stable perfect matching via the weighted common basis problem

For the remainder of Section 4, fix an outcome (ν, p) , with many-to-many projection μ . Also, fix a MAT \mathcal{T} with root k_0 , where all of the items in \mathcal{T} are fully matched (recall that subroutine FINDPRICEINCREASE is only called over MATs \mathcal{T} whose items are fully matched). We define $B'(\mathcal{T}) := \{\text{buy}(k) \mid k \in U(\mathcal{T}) \setminus \{k_0\}\}$ to be the buyers corresponding to unit-buyers in \mathcal{T} , other than the root k_0 , and $W(\mathcal{T}) := B'(\mathcal{T}) \times S(\mathcal{T})$ to be the complete set of edges between these buyers and the items in \mathcal{T} . The first step of the FINDPRICEINCREASE subroutine is to assign each edge $(i, j) \in W(\mathcal{T})$ a weight $\log q'_{ij}(p_j)$, which is the log of the slope of the effective price function of buyer i for item j at price p_j . We wish to compute a minimum weight set of edges $\mu^* \subseteq W(\mathcal{T})$ such that μ^* perfectly matches the buyers and items in $W(\mathcal{T})$, and $\mu \setminus E(\mathcal{T}) \cup \mu^*$ is partially stable. We call such a matching μ^* a *minimum weight partially stable perfect matching*. The weights are designed to eventually apply Lemma 19 after an exponentiation operation.

The partial stability constraint can be formulated as a matroid constraint over the edges. This constraint also ensures that all unit-buyers are matched. A second matroid constraint is used to ensure that the items are perfectly matched as well. Restricting to bases of the matroids imposes the maximum cardinality constraint. Putting these results together, we can cast the minimum weight partially stable perfect matching problem as a minimum weight common basis problem. The proofs in this section are deferred to Appendix C.2.

We start by defining M_B , the first of the two matroids of interest. Essentially, a basis in M_B is a set of edges that maximizes the sum of the utilities to the buyers, subject to the bundle sizes originally allocated to each buyer. Thus, each buyer must be assigned a bundle that it demands at the current prices under the size constraint.

Lemma 20 (Buyer demand matroid). *Let (ν, p) be a partially stable outcome with many-to-many projection μ and \mathcal{T} a MAT in $D(\nu, p)$. For $i \in B'(\mathcal{T})$, let $\mathcal{E}_i = \{E \subseteq \{i\} \times S(\mathcal{T}) : |E| = |U'(\mathcal{T}) \cap \text{ubuy}(i)|\}$. Define the family of sets*

$$\mathcal{J}_B = \left\{ \bigcup_{i \in B'(\mathcal{T})} S_i \mid S_i \in \arg \max_{F \in \mathcal{E}_i} u_i(S(F) \mid \mu(i) \setminus S(\mathcal{T}), p) \right\}$$

and \mathcal{I}_B to be the family of all subsets of \mathcal{J}_B . Then, $M_B = (W(\mathcal{T}), \mathcal{I}_B)$ is a matroid and \mathcal{J}_B are its bases.

We can efficiently implement an independence oracle for M_B .

Lemma 21 (Independence oracle for M_B). *There exists an independence oracle for M_B that runs in $O(|B|^2|S|^2)$ time.*

Next, we can show that a matching is \mathcal{T} -perfect (i.e., matches each item fully) if and only if it is a basis in the *item capacity matroid*.

Definition 22 (Item capacity matroid M_S). Let (ν, p) be a partially stable outcome and \mathcal{T} a MAT at (ν, p) . Let $\mathcal{I}_S = \{E \subseteq W(\mathcal{T}) : |E(j)| \leq \text{cap}(j) \text{ for all } j \in S(\mathcal{T})\}$ and $M_S = (W(\mathcal{T}), \mathcal{I}_S)$, where $E(j) = \{(i, j) \in E\}$.

M_S is a partition matroid (see, e.g., Oxley (2006)). The next two lemmas characterize the bases of M_S and give an independence oracle for M_S . They are well known facts from bipartite matching (see, e.g., Schrijver (2003)), so the proofs are omitted.

Lemma 23 (Basis of M_S if and only if saturates item capacities). *The family \mathcal{J}_S of bases of M_S is given by all and only the set of edges $E \subseteq W(\mathcal{T})$ such that $|E(j)| = \text{cap}(j)$ for all $j \in S(\mathcal{T})$.*

We can also efficiently compute an independence oracle for M_S .

Lemma 24 (Independence oracle for M_S). *There exists an algorithm that identifies whether a set E is independent in M_S in $O(|B|^2|S|)$ time.*

Definition 25 (Lift). Given a many-to-many matching μ , a *lift* of μ is a many-to-one matching ν that satisfies $\{(\text{buy}(k), j) \mid (k, j) \in \nu\} = \mu$. We define the function $\text{LIFT}(\mu)$ to output a lift of μ , selecting one arbitrarily if more than one exist.

We can combine the characterization of M_B and M_S to show that a set of edges in $W(\mathcal{T})$ that is a minimum weight common basis of M_B and M_S has a lift that is a partially stable perfect matching on $V(\mathcal{T}) \setminus \{k_0\}$.

Lemma 26 (Min weight common basis finds min weight partially stable perfect matching). *A lift $\nu^* = \text{LIFT}(\mu^*)$ of the optimal solution μ^* to the weighted minimum common basis problem given by*

$$\min_{E \in \mathcal{J}_B \cap \mathcal{J}_S} \sum_{(i,j) \in W(\mathcal{T})} \log q'_{ij}(p_j) \quad (5)$$

is a minimum weight partially stable perfect matching on $V(\mathcal{T}) \setminus \{k_0\}$, where the weight of an edge (k, j) is also given by $\log q'_{kj}(p_j)$.

We also show in the proof of Lemma 26 that the original partially stable matching μ is a basis of both M_B and M_S , though it may not have minimum weight. We also know that an efficient algorithm exists for finding a common basis of minimum weight.

Lemma 27 (Common matroid basis algorithm). *There exists an algorithm COMMONBASIS that finds the minimum weight common matroid basis in $O(|B|^3|S|^3)$ oracle calls.*

The algorithm repurposes an existing weighted matroid intersection algorithm, such as those by Lawler (1975) and Frank (1981), and applies it to a transformed set of weights.

4.3 The duality trick

In this section, we describe the linear programming duality trick that is used to find a MAT-preserving price increase. In particular, we show that an optimal solution to a linear program defined as a function of $\nu^* = \text{LIFT}(\mu^*)$, the lift of the optimal solution to the minimum weight common basis problem defined in Section 4.2, satisfies complementary slackness constraints with ν^* itself. We begin with the definition of this linear program.

Definition 28 (Linear program for MAT-preserving price increase). Given a partially stable outcome (ν, p) , with unmatched unit-buyer k_0 and corresponding MAT \mathcal{T} where all items in $S(\mathcal{T})$ are fully matched, define the MAT-preserving price increase linear program $LP(\nu, p, \mathcal{T})$ over variables ω, ρ as:

$$\begin{aligned} \max \quad & \sum_{k \in U(\mathcal{T})} \omega_k + \sum_{j \in S(\mathcal{T})} \text{cap}(j) \rho_j \\ \text{s.t.} \quad & \omega_k + \rho_j \leq \log q'_{kj}(p_j) \quad \forall (k, j) \in D(\nu, p) \cap V(\mathcal{T}) \setminus \{k_0\} \end{aligned}$$

We can now state the main technical contribution of this section, Lemma 29, showing that the constraints in $LP(\nu^*, p, \mathcal{T})$ corresponding to edges $(k, j) \in \nu^*$ are tight. These constraints being tight for (ω^*, ρ^*) are crucial to then be able to use ρ^* to compute the desired price increase via complementary slackness.

Lemma 29 (Min-weight partially stable perfect matching satisfies equality constraints). *Let μ^* be an optimal solution to the weighted minimum common basis problem given by*

$$\min_{E \in \mathcal{J}_B \cap \mathcal{J}_S} \sum_{(i,j) \in W(\mathcal{T})} \log q'_{ij}(p_j) \quad (6)$$

and let $\nu^* = \text{LIFT}(\mu^*)$. Let (ω^*, ρ^*) be an optimal solution to $LP(\nu^*, p, \mathcal{T})$. Then, for all $(k, j) \in \nu^*$, we have $\omega_k^* + \rho_j^* = \log q'_{kj}(p_j)$.

Before delving into the proof of Lemma 29, let us provide some intuition on why it holds and how the result differentiates itself from the literature.

Remark 30. Similar to Alkan (1989) and Dütting et al. (2011), the dual of $LP(\nu, p, \mathcal{T})$ is a minimum weight perfect matching linear program on $D(\nu, p) \cap W(\mathcal{T})$, with the weight of an edge (k, j) given by $w((k, j)) = \log q'_{kj}(p_j)$. Lemma 29 is essentially referring to the complementary slackness conditions between the two linear programs. However, unlike the one-to-one matching version of the problem studied in Alkan (1989) and Dütting et al. (2011), we cannot simply define LP and the corresponding dual min-weight matching linear program from an arbitrary initial matching ν , in order to obtain an optimal matching ν^* and corresponding optimal dual solution (ω^*, ρ^*) . This failure is because the optimal matching for the dual minimum weight matching program may not be partially stable. As a result, it is crucial that we use the minimum weight partially stable perfect matching ν^* as computed by the common matroid basis problem as the input for LP .

Once (ω^*, ρ^*) have been obtained, Alkan (1989) and Dütting et al. (2011) massage it to a new solution and then apply a “duality trick” that sets $d_j \leftarrow e^{-\rho_j^*}$, as to obtain the required price increase. We also apply a similar procedure via Algorithm 2 and Lemma 32.

As a first step towards the proof of Lemma 29, we need a helper result on the structure of the marginal demand graph. Essentially, if a buyer i has many unit-buyer copies that all have matched and unmatched edges in the marginal demand graph, we can either swap all of the edges simultaneously while preserving partial stability, or we can find additional unmatched edges in the marginal demand graph.

Lemma 31 (Structure of marginal demand graph). *Let (ν, p) be a feasible and partially stable outcome. Assume there exist, for some $z \geq 2$, $(k_1, j_1), \dots, (k_z, j_z) \in \nu$, and $(k_1, j'_1), \dots, (k_z, j'_z) \in D(\nu, p) \setminus \nu$, where $k_1, \dots, k_z \in \text{ubuy}(i)$ for some i and are distinct and all of the j'_ℓ are distinct. Then, at least one of the following is true:*

- (a) *For all subsets $X \subseteq [z]$, the bundle $\nu(\text{ubuy}(i)) \cup \{j'_x \mid x \in X\} \setminus \{j_x \mid x \in X\}$ maximizes $u_i(\cdot, p)$, among bundles of size $|\nu(\text{ubuy}(i))|$.*
- (b) *There exists a sequence of distinct indices $x_1, \dots, x_s \in [z]$ s.t. $(k_{x_1}, j'_{x_2}, k_{x_2}, j'_{x_3}, k_{x_3}, \dots, k_{x_s}, j'_{x_1}, k_{x_1})$ is a cycle in $D(\nu, p) \setminus \nu$.*

The proof of this result is deferred to Appendix C.3. See Figure 3 for an example of the lemma. This intermediate result will help us show that if a negative weight alternating cycle exists in the marginal demand graph, then a negative weight alternating cycle exists that also preserves partial stability when augmented.

We can now prove Lemma 29.

Proof of Lemma 29. Consider the linear program $DP(\nu, p, \mathcal{T})$:

$$\begin{aligned} \min \quad & \sum_{k \in U(\mathcal{T}), j \in S(\mathcal{T})} \lambda_{kj} \log q'_{kj}(p_j) \\ \text{s.t.} \quad & \sum_{j \in F_{\nu, p}(k)} \lambda_{kj} = 1 && \forall k \in U(\mathcal{T}) \setminus k_0 \quad (\text{unit-buyer constraints}) \\ & \sum_{k \in F_{\nu, p}^{-1}(j)} \lambda_{kj} = \text{cap}(j) && \forall j \in S(\mathcal{T}) \quad (\text{item capacity constraint}) \\ & \lambda_{kj} \geq 0 && \forall (k, j) \in D(\nu, p) \cap V(\mathcal{T}) \setminus \{k_0\} \end{aligned}$$




Figure 3: An example of the structure of the marginal demand graph as shown in Lemma 31, where $k_1, k_2, k_3 \in \text{ubuy}(i)$. Consider the portion of a marginal demand graph shown in Figure 3(a). The matching ν as shown by the solid lines is partially stable. Then, by Lemma 31, either the matching shown in Figure 3(b) also provides i with optimal utility $u_i(\cdot, p)$ among bundles of size $|\nu(\text{ubuy}(i))|$, or we can find additional unmatched edges in the marginal demand graph. If part (b) of Lemma 31 holds for the sequence $x_1 = 1, x_2 = 2$, then the marginal demand graph would also contain the edges in Figure 3(c).

$DP(\nu^*, p, \mathcal{T})$ is a minimum weight perfect matching linear program, and is the dual of $LP(\nu^*, p, \mathcal{T})$. We wish to show that ν^* is an optimal solution for $DP(\nu^*, p, \mathcal{T})$. Let $y \in \{0, 1\}^{U(\mathcal{T}) \setminus \{k_0\} \times S(\mathcal{T})}$ be indicators for the edges in the residual network at ν^* , i.e., $y_{kj} = 1$ if and only if $\nu_{kj}^* = 0$ and $y_{jk} = 1$ if and only if $\nu_{kj}^* = 1$, with y_{kj} and y_{jk} equal to 0 otherwise, for each $(k, j) \in D(\nu, p) \cap V(\mathcal{T}) \setminus \{k_0\}$. It is well known that ν^* is a minimum weight perfect matching if and only if there does not exist a negative-weight cycle in the residual network at ν^* (see, e.g., Schrijver (2003)). Suppose for contradiction that there exists a negative weight cycle

$$C = (j_1, k_1, j_2, k_2, \dots, j_z, k_z) \quad (7)$$

with fewest edges, where $\nu_{k_\ell j_\ell}^* = 1$ and $\nu_{k_\ell j_{\ell+1}}^* = 0$ for all ℓ (and $j_{z+1} := j_1$). Suppose first $\nu^* \Delta C$ is a partially stable perfect matching. Then by construction its weight is strictly smaller than the weight of ν^* , a contradiction to Lemma 26. Thus, $\nu^* \Delta C$ is not partially stable. In particular, some buyer must appear in multiple unit-buyer copies in C . Indeed, if this was not the case, then by definition of the marginal demand graph, we have

$$u_{k_\ell}(\nu^*(\text{copy}(k_\ell)), p) = u_{k_\ell}(\nu^*(\text{copy}(k_\ell)) \cup \{j_{\ell+1}\} \setminus \{j_\ell\}, p) \quad (8)$$

That is, at the augmented matching $\nu^* \Delta C$, every buyer has the same utility as in ν^* , which means that $\nu^* \Delta C$ is also partially stable, a contradiction. There are now two possibilities:

- (1) $\nu^* \Delta C$ matches multiple unit-buyer copies of the same buyer to the same item.
- (2) $\nu^* \Delta C$ does not match multiple unit-buyer copies of the same buyer to the same item.

Suppose (1) is true, and let i be a buyer with multiple unit-buyer copies matched to the same item in $\nu^* \Delta C$. Then, i must have multiple unit-buyers in $V(C)$. Each such unit-buyer $k_x \in \text{ubuy}(i) \cap V(C)$ then has an edge $(k_x, j'_x) \in D(\nu, p) \setminus \nu^*$, given by one of its edges in C . We can now apply Lemma 31 to these unit-buyers $k_x \in \text{ubuy}(i) \cap V(C)$, with the matched edges in C given by (k_x, j_x) , and the unmatched edges in C given by $(k_x, j'_x) = (k_x, j_{x+1})$. Suppose for contradiction that Lemma 31 (a) is true. Then, all of the j'_x are distinct, and so augmenting along C would not match multiple unit-buyers in $\text{ubuy}(i) \cap V(C)$ to the same item, a contradiction. We see that Lemma 31 (b) must be true for i .

Now suppose (2) is true. It must be the case that at least one buyer i has multiple unit-buyer copies in $V(C)$. Then, each such unit-buyer $k_x \in \text{ubuy}(i) \cap V(C)$ has an edge $(k_x, j'_x) \in D(\nu, p) \setminus \nu^*$, given by one of its edges in C . We can now apply Lemma 31. Suppose for contradiction that Lemma 31 (a) is true for every buyer i with multiple unit-buyer copies in $V(C)$. Fixing any such i , we know by Lemma 31 (a) that by setting $X = \{x \mid k_x \in \text{ubuy}(i) \cap V(C)\}$, the bundle

$$\nu^*(\text{ubuy}(i)) \cup \{j'_x \mid x \in X\} \setminus \{j_x \mid x \in X\} \quad (9)$$

maximizes $u_i(\cdot, p)$ among bundles of size $|\nu^*(\text{ubuy}(i))|$, where $(k_x, j_x) \in C \cap \nu^*$ and $(k_x, j'_x) \in C \setminus \nu^*$. However, expression (9) is exactly equal to $(\nu^* \Delta C)(\text{ubuy}(i))$. Thus, $\nu^* \Delta C$ matches this buyer i with a

bundle maximizing $u_i(\cdot, p)$ among sets of size $|\nu^*(\text{ubuy}(i))|$. Since i was an arbitrary buyer with multiple copies in C and clearly $\nu^* \triangle C$ matches every buyer i' with exactly one unit-buyer copy in C to a bundle of size $|\nu^*(\text{ubuy}(i'))|$ maximizing $u_{i'}(\cdot, p)$, $\nu^* \triangle C$ is partially stable, a contradiction. It follows that Lemma 31 (b) must be true for at least one buyer i with multiple unit-buyer copies in $V(C)$.

Thus, if either (1) or (2) is true, there must exist a buyer i with multiple unit-buyer copies in $V(C)$ for which Lemma 31 (b) applies with j_x (resp. j'_x) being the edges of $C \cap \nu^*$ (resp. $C \setminus \nu^*$) incident to his unit-buyer copies. Fix one such buyer i , with unit-buyers $k_1, \dots, k_z \in \text{ubuy}(i) \cap V(C)$, matched edges $(k_x, j_x) \in C$, and unmatched edges $(k_x, j'_x) = (k_x, j_{x+1}) \in C$ for each $1 \leq x \leq z$. Let x_1, \dots, x_s be the sequence of indices described in Lemma 31 (b). Then, $(k_{x_1}, j'_{x_2}, k_{x_2}, j'_{x_3}, k_{x_3}, \dots, k_{x_s}, j'_{x_1})$ is a cycle in $D(\nu, p) \setminus \nu$. Call each of the edges of the form $(k_{x_\ell}, j'_{x_\ell})$ in this cycle a *chord*. Now, consider the set of cycles C_1, \dots, C_s , where C_ℓ is given by

$$C_\ell = (k_{x_\ell}, j'_{x_{\ell+1}} = j_{x_{\ell+1}+1}, k_{x_{\ell+1}+1}, j_{x_{\ell+1}+2}, k_{x_{\ell+1}+2}, \dots, k_{x_{\ell-1}}, j_{x_\ell}, k_{x_\ell}) \quad (10)$$

That is, C_ℓ is the ‘‘shortcutted’’ cycle obtained by first traversing the chord $(k_{x_\ell}, j'_{x_{\ell+1}})$, then following C until we return back to k_{x_ℓ} . Note that

$$w(k_{x_\ell}, j'_{x_{\ell+1}}) = w(k_{x_{\ell+1}}, j'_{x_{\ell+1}}) \quad (11)$$

where w is the objective weight function from DP , as the unit-buyers k_{x_ℓ} and $k_{x_{\ell+1}}$ are copies of the same buyer. It follows that the weight of each C_ℓ is given by

$$\begin{aligned} w(C_\ell) &= w(k_{x_\ell}, j_{x_{\ell+1}+1}) + w(j_{x_{\ell+1}+1}, k_{x_{\ell+1}+1}) + w(k_{x_{\ell+1}+1}, j_{x_{\ell+1}+2}) \\ &\quad + \dots + w(k_{x_{\ell-1}}, j_{x_\ell-1}) + w(j_{x_\ell-1}, k_{x_\ell}) \\ &= w(k_{x_{\ell+1}}, j_{x_{\ell+1}+1}) + w(j_{x_{\ell+1}+1}, k_{x_{\ell+1}+1}) + w(k_{x_{\ell+1}+1}, j_{x_{\ell+1}+2}) \\ &\quad + \dots + w(k_{x_{\ell-1}}, j_{x_\ell-1}) + w(j_{x_\ell-1}, k_{x_\ell}) \end{aligned}$$

That is, the weight of C_ℓ is equal to the sum of the weights of the arcs of C from $k_{x_{\ell+1}}$ to k_{x_ℓ} . Adding the weights of C_1, \dots, C_s gives us the weight of the arcs of C from k_{x_2} to k_{x_1} , then k_{x_3} to k_{x_2} , and so on, completing with the arc from $k_{x_{s+1}} = k_{x_1}$ to k_{x_s} . In other words, we have started a grand cycle from k_{x_1} , followed C an integer m times around, and finishing back at k_{x_1} . It follows that

$$\sum_{\ell=1}^s w(C_\ell) = m \cdot w(C) < 0. \quad (12)$$

There must then be at least one cycle C_ℓ with $w(C_\ell) < 0$. Furthermore, this cycle has strictly fewer edges than C since two copies of the same buyer cannot be at distance 2 in C . This contradicts the choice of C , concluding the proof. \square

Finally, once we have the optimal solution (ω^*, ρ^*) to $LP(\nu^*, p, \mathcal{T})$, we apply a transformation to them, described in Algorithm 2, that guarantees there exists a MAT \mathcal{T}' containing ν^* such that, for the transformed solution, the constraints of $LP(\nu^*, p, \mathcal{T})$ corresponding to edges in \mathcal{T}' are all tight. We note that, before the transformation, complementary slackness guarantees that, for the optimal solution (ω^*, ρ^*) , the constraints $\omega_k^* + \rho_j^* \leq \log q'_{kj}(p_j)$ corresponding to edges $(k, j) \in \nu^*$ are tight. Also note that given two subgraphs H, H' , we may abuse notation and write $H \cap V(H')$ to denote the subgraph of H contained in $U(H') \times S(H')$.

In Algorithm 2, H_ρ is the graph of edges $(k, j) \in D(\nu^*, p) \cap V(\mathcal{T})$ such that the constraint $\omega_k + \rho_j \leq \log q'_{kj}(p_j)$ is tight. The goal is for H_ρ to contain a MAT \mathcal{T}' containing ν^* . Algorithm 2 computes the initial, incomplete version of \mathcal{T}' in line 5. Then, in lines 6 to 8, it uniformly increases the entries of ρ and decreases the entries of ω corresponding to nodes that are not in \mathcal{T}' . This translation of ω and ρ preserves tightness for all edges for which the dual constraint is tight, and allows at least one more constraint to tighten, and so the corresponding edge between $U(\mathcal{T}')$ and $S(\mathcal{T}) \setminus S(\mathcal{T}')$ can enter H_ρ . The new edge is then naturally added to \mathcal{T}' as well, as every edge from a unit-buyer $k \in U(\mathcal{T}')$ must be contained in any MAT containing k by definition of a MAT. This process is then repeated on the transformed solution (ω, ρ) . The loop terminates once \mathcal{T}' covers the same unit-buyers and items as \mathcal{T} , at which point Algorithm 2 returns the final modified solution (ω, ρ) .

ALGORITHM 2: CONNECTMAT

Input: Matching ν^* , prices p , original MAT \mathcal{T} rooted at k_0 , optimal solution (ω^*, ρ^*) to $LP(\nu^*, p, \mathcal{T})$

Output: Solution (ω, ρ) to $LP(\nu^*, p, \mathcal{T})$ such that there exists a MAT \mathcal{T}' such that primal constraints corresponding to edges in \mathcal{T}' are tight over (ω, ρ)

```
1  $\omega, \rho \leftarrow \omega^*, \rho^*$  ;
2  $\omega_{k_0}^* \leftarrow \min_{j \in S(\mathcal{T})} \log q'_{k_0 j}(p_j) - \rho_j$  ;
3 do
4    $H_\rho \leftarrow \{(k, j) \in D(\nu^*, p) \cap V(\mathcal{T}) \mid \omega_k + \rho_j = \log q'_{k j}(p_j)\}$ 
5    $\mathcal{T}' \leftarrow \text{FINDMAT}(H_\rho, \nu^*, k_0)$ 
6    $\epsilon \leftarrow \min_{(k, j) \in D(\nu^*, p): k \in U(\mathcal{T}'), j \in S(\mathcal{T}) \setminus S(\mathcal{T}')} \log q'_{k j}(p_j) - \omega_k - \rho_j$  /*  $\epsilon$  tightens a constr. */
7   Set  $\rho_j \leftarrow \rho_j + \epsilon$  for all  $j \in S(\mathcal{T}) \setminus S(\mathcal{T}')$ 
8   Set  $\omega_k \leftarrow \omega_k - \epsilon$  for all  $k \in U(\mathcal{T}) \setminus U(\mathcal{T}')$  /* by choice of  $\epsilon$ , one constraint tightens */
9 while  $|\mathcal{T}'| < |\mathcal{T}|$  /* so,  $H_\rho$  increases by one edge each iteration */
10 return  $\omega, \rho$ 
```

Lemma 32 (Correctness and runtime of Algorithm 2). *Algorithm 2 outputs ω, ρ such that H_ρ contains a MAT \mathcal{T}' containing ν^* . Furthermore, it terminates in $O(|B||S|^3)$ time.*

The proof of this result is deferred to Appendix C.3.

4.4 The MAT-preserving price increase algorithm

We can now describe Algorithm 3, FINDPRICEINCREASE, for computing the MAT-preserving price increase d and step size λ^* . Let COMMONBASIS($\mathcal{I}_B, \mathcal{I}_S, w$) be an algorithm that takes in two matroid independence oracles and a weight function and outputs the minimum weight common basis. Finally, let LINSEG $_{i,j}(p_j)$ be the highest index ℓ such that p_j belongs to the ℓ^{th} linear segment for the effective price function q_{ij} .

Algorithm 3 first solves the minimum weight common basis problem on M_B and M_S . Then, it uses the solution ν^* to define an appropriate version of LP , where by Lemma 29, the optimal solution satisfies complementary slackness with ν^* . The algorithm calls CONNECTMAT in line 4 to modify the dual variables ω^*, ρ^* , and also extends ω to include ω_{k_0} while satisfying the corresponding constraints involving ω_{k_0} . Next, it uses the duality trick to compute the price increase d (line 5). Finally, it calculates the maximum step size λ^* the price can be increased along d (line 8), until a new edge enters the marginal demand graph $D(p + \lambda^*d)$ (line 6) or the price $p + \lambda^*d$ leaves the current linear domain (line 7).

ALGORITHM 3: FINDPRICEINCREASE

Input: MAT \mathcal{T} with respect to ν with root k_0 in $D(\nu, p)$, where all items in $S(\mathcal{T})$ are fully matched

Output: MAT-preserving price increase and step size λ^*d for \mathcal{T} and corresponding matching ν'

```
1  $\mu^* \leftarrow \text{COMMONBASIS}(\mathcal{I}_B, \mathcal{I}_S, w)$ 
2  $\nu^* \leftarrow \text{LIFT}(\mu^*) \cup (\nu \setminus E(\mathcal{T}))$  /*  $\nu^*$  is an optimal solution to  $LP$  */
3  $\omega^*, \rho^* \leftarrow LP(\nu^*, p, \mathcal{T})$  /* define  $LP$  using max weight common basis */
4  $\omega^*, \rho^* \leftarrow \text{CONNECTMAT}(\nu^*, p, \mathcal{T}, \omega, \rho)$  /* modifies dual variables to preserve full MAT */
5 Set  $d_j \leftarrow e^{-\rho_j^*}$  for all  $j \in S(\mathcal{T})$  and  $d_j \leftarrow 0$  otherwise /* duality trick */
6  $\lambda_1^* \leftarrow \min\{\lambda \geq 0 : (k, j) \in D(\nu^*, p + \lambda d) \text{ for some } k \in U(\mathcal{T}) \text{ and } j \notin F_{\nu^*, p}(k)\}$ 
7  $\lambda_2^* \leftarrow \min\{\lambda \geq 0 : \text{LINSEG}_{\text{buy}(k), j}(p_j) \neq \text{LINSEG}_{\text{buy}(k), j}(p_j + \lambda d_j) \text{ for some } k \in U \text{ and } j \in S\}$ 
8  $\lambda^* \leftarrow \min\{\lambda_1^*, \lambda_2^*\}$  /* max step size to preserve MAT and stay within linear domain */
9 return  $\nu^*, \lambda^*d$ 
```

We prove in Appendix C.4 the correctness and runtime of Algorithm 3.

Lemma 33 (Correctness and runtime of Algorithm 3). *Algorithm 3 finds a MAT-preserving price increase d and maximal step size λ^* such that $(\nu^*, p + \lambda^*d)$ is partially stable, $p + \lambda^*d$ remains within the same linear domain, and an alternating tree \mathcal{T}' with the same unit-buyers, items, and root k_0 as \mathcal{T} is preserved in $D(\nu^*, p + \lambda^*d)$. Furthermore, if $p + \lambda^*d$ does not reach the border of the current linear domain, the number of items in the MAT rooted at k_0 increases by 1. The algorithm runs in time $O(|B|^5|S|^5)$.*

5 The Main Result

In this section, we complete the proof of the main result. In Section 5.1, we analyze FINDMAT, the subroutine for computing a MAT. In Section 5.2, we analyze AUGMENTINGPATH, the subroutine for increasing the size of the matching. Finally, in Section 5.3, we prove Theorem 1.

5.1 The FindMAT subroutine

Given an unmatched unit-buyer k_0 that demands at least one item, FINDMAT, formally described as Algorithm 4 below, finds a MAT \mathcal{T} rooted at k_0 . The process is essentially a breadth-first search on the edges of the marginal demand graph. Initially, \mathcal{T} is a tree with a single vertex, k_0 . For each item $j \in F_{\nu,p}(k_0)$ demanded by k_0 , the edge (k_0, j) is added to \mathcal{T} and, for all buyers $k \in \nu(j)$ matched to j , the edges (k, j) are also added to \mathcal{T} (line 5). The algorithm then repeats this process with the newly added buyers k to \mathcal{T} , but without connecting to a vertex that is already in \mathcal{T} .

ALGORITHM 4: FINDMAT

Input: Graph H , matching $\nu \subseteq H$, and unmatched unit-buyer k_0 with an edge in H

Output: MAT \mathcal{T} in H rooted at k_0

```

1 Initialize  $\mathcal{T} \leftarrow \emptyset$ ,  $U' \leftarrow \{k_0\}$ 
2 while  $\exists k \in U'$  do
3   for  $j \notin V(\mathcal{T})$  s.t.  $(k, j) \in H$  do
4      $U'' \leftarrow \nu(j) \setminus V(\mathcal{T})$ 
5      $\mathcal{T} \leftarrow \mathcal{T} \cup \{(k, j)\} \cup \{(k', j) \mid k' \in U''\}$ 
6      $U' \leftarrow U' \cup U'' \setminus \{k\}$ 
7 return  $\mathcal{T}$ 

```

Lemma 34. *For any many-to-one outcome (ν, p) and unmatched unit-buyer k_0 that demands at least one item, Algorithm 4 finds a MAT rooted at k_0 in $O(|B||S|^2)$ running time.*

Proof of Lemma 34. First, we will show that the output \mathcal{T} of Algorithm 4 is an alternating tree. Edges are only added to \mathcal{T} in line 5. We also know that $j \notin V(\mathcal{T})$ chosen in line 3 and $k' \in U''$ chosen in lines 4-5 must not already be in $V(\mathcal{T})$. We see that \mathcal{T} is acyclic, and so it is a tree. A unit-buyer k can only be added to \mathcal{T} if it is k_0 or on an edge (k', j) in line 5. We also know by line 4 that (k', j) is a matched edge. Then, consider the edge (k, j) added in line 5. When k was added to \mathcal{T} , it was also added via a matched edge, and since unit-buyers have at most one match, we see that (k, j) is not matched. We see that \mathcal{T} is alternating as well.

Now we will show that \mathcal{T} is maximal. Consider any unit-buyer $k \in U(\mathcal{T})$. We know that $k = k_0$, or $k = k'$ in some iteration of line 5. In the first case, k is added to U' in line 1. In the second case, k is added to U' in line 6. In both cases, k is not removed from U' until after it is selected in line 2. When this happens, all items $j \notin V(\mathcal{T})$ with $(k, j) \in H$ are added to \mathcal{T} via the edge (k, j) , unless they were already in $V(\mathcal{T})$. Thus, for every unit-buyer $k \in V(\mathcal{T})$, all items j with an edge to k are also in $V(\mathcal{T})$. Next, consider any item $j \in V(\mathcal{T})$. We know that j must be added to $V(\mathcal{T})$ in line 5. At the same time, all of its matched edges (k', j) are also added to \mathcal{T} unless k' is already in the tree. We see that for every item $j \in V(\mathcal{T})$, all of its matched unit-buyers are in $V(\mathcal{T})$ as well, and conclude that \mathcal{T} is maximal.

Now, we consider the runtime of Algorithm 4. In each iteration of the for loop, an item $j \notin V(\mathcal{T})$ is selected and added to $V(\mathcal{T})$ permanently. As a result, the for loop can iterate at most $O(|S|)$ times throughout the entire algorithm. In each iteration, lines 4-6 require at most $O(|B||S|)$ time. We conclude that the total runtime is at most $O(|B||S|^2)$. \square

5.2 The AugmentingPath subroutine

The final subroutine is AUGMENTINGPATH, called in line 8 of Algorithm 1. The subroutine finds the shortest augmenting path from the root k_0 of a MAT \mathcal{T} to a leaf j that is undermatched.

Note that if an augmenting path P contains two unit-buyer copies k_1, k_2 of the same buyer i , then it is possible that after augmenting the current matching ν along P , then buyer i is no longer matched to a bundle maximizing $u_i(\cdot, p)$ among bundles of size $|\nu(\text{ubuy}(i))|$. This possibility is due to the non-linear nature of gross substitutes valuations, so that by swapping one item, the marginal value of another item may change unpredictably. We can show that by selecting the shortest such path to augment, we can preserve partial stability while increasing the size of the matching. The main idea of the proof is to use Lemma 31 to find shortcuts in the augmenting path; if no shortcuts exist, then we show the augmenting path must preserve partial stability.

Lemma 35 (Augmenting preserves partial stability). *Let (ν, p) be a partially stable outcome. Let \mathcal{T} be a MAT rooted at unmatched unit-buyer k_0 . Let P be the shortest alternating path from k_0 to any item j_0 in \mathcal{T} that is not fully matched, and let $\nu \Delta P$ be the matching obtained by augmenting ν along P . Then, $\nu \Delta P$ is partially stable at p .*

Proof of Lemma 35. We wish to show that after augmentation, each buyer i is matched to a bundle that optimizes $u_i(\cdot, p)$ among bundles of size $|\nu \Delta P(i)|$. Let (k_0, j_1) be the first edge in P . Add an auxiliary item j_a with price $p_a = 0$, and edit the valuation function of $i_0 = \text{buy}(k_0)$ to be v'_i , defined by

$$v'_i(T) = \begin{cases} v_i(T) & j_a \notin T \\ v_i(T \setminus \{j_a\}) + u_i(j_1 \mid \nu(\text{ubuy}(i)), p) & j_a \in T. \end{cases} \quad (13)$$

Also define the corresponding utility function

$$u'_i(T, p) = v'_i(T) - \sum_{j \in T} q_{ij}(p_j) \quad (14)$$

For all $T \subseteq S$ such that $j_a \notin T$, we have $v'_i(j_a \mid T) = u'_i(j_a \mid T, p) = u_i(j_1 \mid \nu(\text{ubuy}(i)), p)$. In other words, the marginal value and marginal utility of j_a under v'_i and u'_i is always equal to $u_i(j_1 \mid \nu(\text{ubuy}(i)), p)$. To see that v'_i still satisfies gross substitutes, notice that $v'_i(T)$ is the sum of a gross substitutes and an additive function, for which it is well-known to satisfy gross substitutes as well (see, e.g., [de Vries et al. \(2020\)](#)).

Next, edit the current matching to become $\nu' = \nu \cup \{(k_0, j_a)\}$ and the augmenting path P to become the alternating path $P' = P \cup \{(k_0, j_a)\}$. This step is necessary to ensure that every buyer with a unit-buyer in P has the same number of matches before and after augmentation. Before augmenting along P' , every buyer $i \neq i_0$ still has the bundle $\nu'(i) = \nu(i)$, which maximizes $u_i(\cdot, p)$ among bundles of size $|\nu'(i)|$. For buyer i_0 , the bundle $\nu'(i_0) = \nu(i_0) \cup \{j_a\}$ also maximizes $u_{i_0}(\cdot, p)$ among bundles of size $|\nu'(i_0)|$. To see this, we can write

$$\begin{aligned} u'_i(\nu'(i_0), p) &= u'_i(\nu(i_0) \cup \{j_a\}, p) \\ &= u'_i(\nu(i_0), p) + u'_i(j_a \mid \nu(i_0), p) \\ &= u'_i(\nu(i_0), p) + u'_i(j_1 \mid \nu(i_0), p) \\ &= u'_i(\nu(i_0) \cup \{j_1\}, p) \end{aligned}$$

The last term is known to maximize $u_{i_0}(\cdot, p)$ among bundles of size $|\nu'(i_0)|$, by definition of the marginal demand graph. It follows that ν' is partially stable.

Consider any buyer i with a single unit-buyer copy k in P' . Then, by definition of the marginal demand graph, $u_i(\nu(i), p) = u_i((\nu \Delta P)(i), p)$, and i retains optimal utility among bundles of size $|\nu(i)|$.

Now, consider any buyer i with multiple unit-buyer copies k_1, \dots, k_z in P' , appearing from first to last in that order. We can then apply Lemma 31 to this sequence, with matched edges $(k_\ell, j_\ell) \in P' \cap \nu'$ and unmatched edges $(k_\ell, j'_\ell) \in P' \setminus \nu'$. Note that if $i = i_0$, our use of j_a, P' , and ν' is what allows these edges to exist. Assume for contradiction that i has less utility $u_i(\cdot, p)$ after augmentation along P' , so that $\nu'(i) \cup \{j'_1, \dots, j'_z\} \setminus \{j_1, \dots, j_z\}$ does not maximize $u_i(\cdot, p)$ among bundles of size $|\nu'(i)|$. Then Lemma 31 (a) is not true, and so Lemma 31 (b) must be true instead. Thus, there is a sequence of distinct indices x_1, \dots, x_s such that $(k_{x_1}, j'_{x_2}, k_{x_2}, j'_{x_3}, k_{x_3}, \dots, k_{x_s}, j'_{x_1})$ is a cycle in $D(\nu, p) \setminus \nu$. Without loss of generality, let x_1 be the smallest index in x_1, \dots, x_s . Then there is an edge (k_{x_1}, j'_{x_2}) in $D(\nu, p)$. Thus, there is an alternating path from k_0 to j_0 that is shorter than P , given by following P from k_0 to k_{x_1} , then going to j'_{x_2} , then following P to j_0 . This alternating path contradicts our selection of P as the shortest such path. We conclude that each buyer i has the same utility after augmentation, which means that $\nu' \Delta P' = \nu \Delta P$ is also partially stable. \square

We can use any existing algorithm for computing a shortest augmenting path, such as that described in [Schrijver \(2003\)](#). This result is well-known, so the proof is omitted.

Lemma 36 (Shortest augmenting path algorithm). *There exists an algorithm AUGMENTINGPATH which, given a root node k_0 , a matching ν , and a graph $D(\nu, p)$, finds a shortest augmenting path in $O(|B||S|^2)$ time.*

5.3 Main result

We can now put everything together to establish the correctness and runtime of Algorithm 1.

Theorem 1. *In the QITU model with gross substitutes valuations, there is a polynomial time algorithm that computes a competitive equilibrium.*

Proof. First, we will show that the outcome (ν, p) maintained by Algorithm 1 is always partially stable. We begin with $(\nu, p) = (\emptyset, 0)$, which is partially stable. In each iteration of the outer while loop from lines 2-9, we know by Lemma 34 that line 3 correctly computes a MAT \mathcal{T} . As long as all items in the MAT are fully matched, the algorithm executes the inner while loop from lines 4-7. In each iteration of the inner while loop, we know by Lemma 33 that lines 5-7 updates ν, p while preserving partial stability and weakly increasing the size of \mathcal{T} . Then, we know by Lemma 35 that augmenting ν in lines 8-9 also preserves partial stability. We see that ν, p remains partially stable throughout the algorithm.

Then, the algorithm only terminates if all unit-buyers k_0 are matched, which means that no MATs exist in $D(\nu, p)$. By Lemma 16, we know that when combined with partial stability, this implies that the output (μ, p) is a competitive equilibrium. As a result, if the algorithm terminates, it returns a competitive equilibrium.

Now we analyze the runtime of Algorithm 1, and show that it terminates. Consider each iteration of the inner while loop. By Lemma 33, either the number of items in \mathcal{T} increases, or we reach the edge of the current linear domain, and $\text{LINSEG}_{i,j}(p_j)$ increases by one for some (i, j) . \mathcal{T} can increase at most $O(|S|)$ times, and we can increment linear domains at most $O(K)$ times, where K is the total number of linear segments in the effective price functions. Each iteration is dominated by the runtime of FINDPRICEINCREASE , which requires $O(|B|^5|S|^5)$ time by Lemma 33. The total runtime of the inner while loop is thus $O((K + |S|)|B|^5|S|^5)$. Finally, in each iteration of the outer while loop, an unmatched unit-buyer k_0 is selected in line 2, then matched along an augmenting path in line 9. Thus, the number of unmatched unit-buyers decreases by one, for at most $O(|B||S|)$ iterations. Each iteration is dominated by the runtime of the inner while loop, for a total runtime of $O((K + |S|)|B|^6|S|^6)$. \square

6 Hardness of Non-Quasilinear Utility Setting

A natural variation of our model, which we call the NQ model, allows a single buyer to have non-quasilinear utility. More precisely, similar to the QITU model, given a bundle T and a set of prices p , the utility of each buyer i is given by

$$u_i(T, p) = v_i(T) - r_i \left(\sum_{j \in T} q_{ij}(p_j) \right). \quad (15)$$

where $r_i : \mathbb{R} \rightarrow \mathbb{R}$ is a *value-for-money* function. For all buyers except one, r_i is the identity function, exactly as in the QITU model. However, for one buyer, the value-for-money function r_i is a piecewise linear continuous function with one breakpoint. We assume, moreover, that valuations v_i are additive and the effective price functions q_{ij} are the identity (i.e., utility is transferable). Note that assumptions on v_i and q_{ij} are more restrictive than in the QITU model.

Theorem 2. *Computing a competitive equilibrium in the NQ model is NP-hard.*

We postpone the full proof to Appendix D. The approach is to reduce an instance of the integer knapsack problem to an instance of the NQ model. Each knapsack item becomes an item in the assignment game, and we have a main buyer who values each item equal to the knapsack value of the item. The main buyer also has a non-quasilinear utility function that loses utility sharply after paying more than the knapsack budget. We add auxiliary buyers that force each item to have price equal to its knapsack cost in any competitive equilibrium, and so the main buyer's optimal bundle is equivalent to the optimal knapsack bundle.

Acknowledgements

Christopher En and Yuri Faenza acknowledge the support of the NSF Grant 2046146 *CAREER: An Algorithmic Theory of Matching Markets*. Eric Balkanski acknowledges the support of NSF grants CCF-2210501 and IIS-2147361.

References

- Alkan, A. (1989). Existence and computation of matching equilibria. *European Journal of Political Economy*, 5(2-3):285–296.
- Alkan, A. (1992). Equilibrium in a matching market with general preferences. In *Equilibrium and Dynamics: Essays in Honour of David Gale*, pages 1–16. Springer.
- Alkan, A. and Gale, D. (1990). The core of the matching game. *Games and economic behavior*, 2(3):203–212.
- Ausubel, L. M. (2004). An efficient ascending-bid auction for multiple objects. *American Economic Review*, 94(5):1452–1475.
- Chade, H., Eeckhout, J., and Smith, L. (2017). Sorting through search and matching models in economics. *Journal of Economic Literature*, 55(2):493–544.
- Colini-Baldeschi, R., Leonardi, S., Henzinger, M., and Starnberger, M. (2015). On multiple keyword sponsored search auctions with budgets. *ACM Transactions on Economics and Computation (TEAC)*, 4(1):1–34.
- Crawford, V. P. and Knoer, E. M. (1981). Job matching with heterogeneous firms and workers. *Econometrica: Journal of the Econometric Society*, pages 437–450.
- de Vries, S., Friedrich, U., and Raach, S. (2020). Geometry of gross substitutes valuations. *Discrete Applied Mathematics*, 272:16–23.
- Demange, G. and Gale, D. (1985). The strategy structure of two-sided matching markets. *Econometrica: Journal of the Econometric Society*, pages 873–888.
- Demange, G., Gale, D., and Sotomayor, M. (1986). Multi-item auctions. *Journal of political economy*, 94(4):863–872.
- Dress, A. W. and Terhalle, W. (1995a). Rewarding maps: On greedy optimization of set functions. *Advances in Applied Mathematics*, 16(4):464–483.
- Dress, A. W. and Terhalle, W. (1995b). Well-layered maps—a class of greedily optimizable set functions. *Applied Mathematics Letters*, 8(5):77–80.
- Dress, A. W. and Wenzel, W. (1990). Valuated matroids: a new look at the greedy algorithm. *Applied Mathematics Letters*, 3(2):33–35.
- Dupuy, A., Galichon, A., Jaffe, S., and Kominers, S. D. (2020). Taxation in matching markets. *International Economic Review*, 61(4):1591–1634.
- Dütting, P. and Henzinger, M. (2010). Mechanisms for the marriage and the assignment game. In *Algorithms and Complexity: 7th International Conference, CIAC 2010, Rome, Italy, May 26-28, 2010. Proceedings 7*, pages 6–12. Springer.
- Dütting, P., Henzinger, M., and Starnberger, M. (2015). Auctions for heterogeneous items and budget limits. *ACM Transactions on Economics and Computation (TEAC)*, 4(1):1–17.
- Dütting, P., Henzinger, M., and Weber, I. (2009). Bidder optimal assignments for general utilities. In *Internet and Network Economics: 5th International Workshop, WINE 2009, Rome, Italy, December 14-18, 2009. Proceedings 5*, pages 575–582. Springer.
- Dütting, P., Henzinger, M., and Weber, I. (2011). An expressive mechanism for auctions on the web. In *Proceedings of the 20th international conference on World wide web*, pages 127–136.
- Dütting, P., Henzinger, M., and Weber, I. (2013). Sponsored search, market equilibria, and the hungarian method. *Information Processing Letters*, 113(3):67–73.
- Edmonds, J. and Karp, R. M. (1972). Theoretical improvements in algorithmic efficiency for network flow problems. *Journal of the ACM (JACM)*, 19(2):248–264.

- Fleiner, T., Jagadeesan, R., Jankó, Z., and Teytelboym, A. (2019). Trading networks with frictions. *Econometrica*, 87(5):1633–1661.
- Frank, A. (1981). A weighted matroid intersection algorithm. *Journal of Algorithms*, 2(4):328–336.
- Fujishige, S. and Yang, Z. (2003). A note on kelso and crawford’s gross substitutes condition. *Mathematics of Operations Research*, 28(3):463–469.
- Galichon, A., Kominers, S. D., and Weber, S. (2019). Costly concessions: An empirical framework for matching with imperfectly transferable utility. *Journal of Political Economy*, 127(6):2875–2925.
- Goel, G., Mirrokni, V., and Leme, R. P. (2015). Polyhedral clinching auctions and the adwords polytope. *Journal of the ACM (JACM)*, 62(3):1–27.
- Google (2024). Adsense revenue share.
- Gul, F. and Stacchetti, E. (2000). The english auction with differentiated commodities. *Journal of Economic theory*, 92(1):66–95.
- Hatfield, J. W., Kominers, S. D., Nichifor, A., Ostrovsky, M., and Westkamp, A. (2013). Stability and competitive equilibrium in trading networks. *Journal of Political Economy*, 121(5):966–1005.
- Hatfield, J. W., Kominers, S. D., Nichifor, A., Ostrovsky, M., and Westkamp, A. (2019). Full substitutability. *Theoretical Economics*, 14(4):1535–1590.
- Hatfield, J. W. and Milgrom, P. R. (2005). Matching with contracts. *American Economic Review*, 95(4):913–935.
- Kaneko, M. (1982). The central assignment game and the assignment markets. *Journal of Mathematical Economics*, 10(2-3):205–232.
- Kelso Jr, A. S. and Crawford, V. P. (1982). Job matching, coalition formation, and gross substitutes. *Econometrica: Journal of the Econometric Society*, pages 1483–1504.
- Lawler, E. L. (1975). Matroid intersection algorithms. *Mathematical programming*, 9(1):31–56.
- Lehmann, B., Lehmann, D., and Nisan, N. (2001). Combinatorial auctions with decreasing marginal utilities. In *Proceedings of the 3rd ACM conference on Electronic Commerce*, pages 18–28.
- Murota, K. (1996a). Convexity and steinitz’s exchange property. *Advances in Mathematics*, 124(2):272–310.
- Murota, K. (1996b). Valuated matroid intersection i: Optimality criteria. *SIAM Journal on Discrete Mathematics*, 9(4):545–561.
- Murota, K. (1996c). Valuated matroid intersection ii: Algorithms. *SIAM Journal on Discrete Mathematics*, 9(4):562–576.
- Murota, K. and Shioura, A. (1999). M-convex function on generalized polymatroid. *Mathematics of operations research*, 24(1):95–105.
- Nisan, N. and Segal, I. (2006). The communication requirements of efficient allocations and supporting prices. *Journal of Economic Theory*, 129(1):192–224.
- Oxley, J. G. (2006). *Matroid Theory (Oxford Graduate Texts in Mathematics)*. Oxford University Press, Inc., USA.
- Paes Leme, R. (2017). Gross substitutability: An algorithmic survey. *Games and Economic Behavior*, 106:294–316.
- Quinzii, M. (1984). Core and competitive equilibria with indivisibilities. *International Journal of Game Theory*, 13:41–60.
- Reijnierse, H., van Gellekom, A., and Potters, J. A. (2002). Verifying gross substitutability. *Economic Theory*, 20:767–776.

- Rochford, S. C. (1984). Symmetrically pairwise-bargained allocations in an assignment market. *Journal of Economic Theory*, 34(2):262–281.
- Roth, A. E. and Sotomayor, M. (1988). Interior points in the core of two-sided matching markets. *Journal of Economic Theory*, 45(1):85–101.
- Schrijver, A. (2003). *Combinatorial optimization: polyhedra and efficiency*, volume 24. Springer.
- Segal-Halevi, E., Hassidim, A., and Aumann, Y. (2016). Demand-flow of agents with gross-substitute valuations. *Operations Research Letters*, 44(6):757–760.
- Shapley, L. S. and Shubik, M. (1971). The assignment game i: The core. *International Journal of game theory*, 1:111–130.

A Properties of Gross Substitutes

We now take an aside to present some helper results we will need for the next two sections. First, several equivalent definitions of gross substitutes will be useful.

Proposition 37. *The following are equivalent:*

1. (GS) v_i satisfies gross substitutes.
2. (SI) Given any price vector p and any $T \subseteq S$, if $u_i(T, p) < \max_{T' \subseteq S} u_i(T', p)$, then there exist X, Y with $|X|, |Y| \leq 1$ such that $u_i(T \cup X \setminus Y, p) > u_i(T, p)$ [Gul and Stacchetti \(2000\)](#).
3. (GR) Given any price vector p , a greedy procedure that iteratively selects the item j with greatest positive marginal utility $u_i(j | T, p)$ finds the optimal solution to $\max_{T \subseteq S} u_i(T, p)$ [Dress and Terhalle \(1995b\)](#).
4. (WL) v_i is submodular, and for any p and any positive integer ℓ , a greedy function optimizes $\max_{T \subseteq S, |T|=\ell} u_i(T, p)$. This is known as the well-layered property [Dress and Terhalle \(1995b\)](#).
5. (ISO) v_i is submodular, and for all sets T and distinct items $j_1, j_2, j_3 \notin T$,

$$v_i(j_1, j_2 | T) + v_i(j_3 | T) \leq \max \left[v_i(j_1, j_3 | T) + v_i(j_2 | T), v_i(j_2, j_3 | T) + v_i(j_1 | T) \right]$$

[Reijnierse et al. \(2002\)](#)

It is well known (see, e.g., [Paes Leme \(2017\)](#)) that gross substitutes is preserved with any initial endowment and prices.

Lemma 38. *Let $v : 2^S \rightarrow \mathbb{R}$ satisfy gross substitutes. Then for any $T \subseteq S$ and any price vector p over the items, the function $u(\cdot | T, p) : 2^{S \setminus T} \rightarrow \mathbb{R}$ also satisfies gross substitutes.*

It is also well-known that gross substitutes is preserved when an additional ‘‘capacity constraint’’ is added. The following Lemma is folklore, but we include a proof for completeness.

Lemma 39 (Capacity constraint preserves gross substitutes). *Let $v : 2^S \rightarrow \mathbb{R}$ satisfy gross substitutes. Then, for any fixed constant ℓ , the function $\tilde{v} : 2^S \rightarrow \mathbb{R}$ given by*

$$\tilde{v}(T) = \begin{cases} v(T) & |T| \leq \ell \\ v(T) - M(|T| - \ell) & o.w. \end{cases} \quad (16)$$

where M is a constant also satisfies gross substitutes. Similarly, $\bar{v} : 2^S \rightarrow \mathbb{R}$ given by

$$\bar{v}(T) = \begin{cases} v(T) + M|T| & |T| \leq \ell \\ v(T) + M\ell - M(|T| - \ell) & o.w. \end{cases} \quad (17)$$

satisfies gross substitutes.

Proof of Lemma 39. We will show that (WL) holds for \tilde{v} . For submodularity, consider a set $T \subseteq S$ and items $j_1, j_2 \notin T$. If $|T| \leq \ell - 1$ we have by submodularity of v that

$$\begin{aligned} \tilde{v}(T \cup \{j_1\}) + \tilde{v}(T \cup \{j_2\}) &= v(T \cup \{j_1\}) + v(T \cup \{j_2\}) \\ &\geq v(T \cup \{j_1, j_2\}) + v(T) \geq \tilde{v}(T \cup \{j_1, j_2\}) + \tilde{v}(T). \end{aligned}$$

If $|T| \geq \ell$ then

$$\begin{aligned} \tilde{v}(T \cup \{j_1\}) + \tilde{v}(T \cup \{j_2\}) &= v(T \cup \{j_1\}) + v(T \cup \{j_2\}) - 2M(|T| + 1 - \ell) \\ &\geq v(T \cup \{j_1, j_2\}) + v(T) - 2M(|T| + 1 - \ell) = \tilde{v}(T \cup \{j_1, j_2\}) + \tilde{v}(T). \end{aligned}$$

Thus, \tilde{v} is submodular. Next, we will show that \tilde{v} is well-layered. Because v is well-layered, we know that for $\ell' \leq \ell$ a greedy algorithm optimizes

$$\max_{T \subseteq S, |T|=\ell'} u_i(T, p) = \max_{T \subseteq S, |T|=\ell'} \tilde{u}_i(T, p) \quad (18)$$

For all $|T| > \ell$, we know that

$$\tilde{u}_i(T, p) = u_i(T, p) - M(|T| - \ell) \quad (19)$$

As a result, if $u_i(T, p) \geq u_i(T', p)$ for sets T, T' of the same size, then $\tilde{u}_i(T, p) \geq \tilde{u}_i(T', p)$ as well. It follows that when $\ell' > \ell$ a greedy function optimizes

$$\max_{T \subseteq S, |T|=\ell'} \tilde{u}_i(T, p) \quad (20)$$

as well. We conclude that (WL) holds for \tilde{v} , and so \tilde{v} satisfies gross substitutes.

Next, we will show that (WL) holds for \bar{v} . For submodularity, consider a set $T \subseteq S$ and items $j_1, j_2 \notin T$. If $|T| \leq \ell - 1$ we have by submodularity of v that

$$\begin{aligned} \bar{v}(T \cup \{j_1\}) + \bar{v}(T \cup \{j_2\}) &= v(T \cup \{j_1\}) + v(T \cup \{j_2\}) + 2M(|T| + 1) \\ &\geq v(T \cup \{j_1, j_2\}) + v(T) + 2M(|T| + 1) \geq \bar{v}(T \cup \{j_1, j_2\}) + \bar{v}(T). \end{aligned}$$

If $|T| \geq \ell$ then

$$\begin{aligned} \bar{v}(T \cup \{j_1\}) + \bar{v}(T \cup \{j_2\}) &= v(T \cup \{j_1\}) + v(T \cup \{j_2\}) + 2M\ell - 2M(|T| + 1 - \ell) \\ &\geq v(T \cup \{j_1, j_2\}) + v(T) + 2M\ell - 2M(|T| + 1 - \ell) = \bar{v}(T \cup \{j_1, j_2\}) + \bar{v}(T). \end{aligned}$$

Thus, \bar{v} is submodular. Next, we will show that \bar{v} is well-layered. Because v is well-layered, we know that for $\ell' \leq \ell$ a greedy algorithm optimizes

$$\max_{T \subseteq S, |T|=\ell'} u_i(T, p) = \max_{T \subseteq S, |T|=\ell'} \bar{u}_i(T, p) \quad (21)$$

For all $|T| \leq \ell$, we know that

$$\bar{u}_i(T, p) = u_i(T, p) + M|T| \quad (22)$$

For all $|T| > \ell$, we know that

$$\bar{u}_i(T, p) = u_i(T, p) + M\ell - M(|T| - \ell) \quad (23)$$

Thus, if $u_i(T, p) \geq u_i(T', p)$ for sets T, T' of the same size, then $\bar{u}_i(T, p) \geq \bar{u}_i(T', p)$ as well. It follows that when $\ell' > \ell$ a greedy function optimizes

$$\max_{T \subseteq S, |T|=\ell'} \bar{u}_i(T, p) \quad (24)$$

as well. We conclude that (WL) holds for \bar{v} , and so \bar{v} satisfies gross substitutes. \square

It has also been shown that the gross substitutes is preserved under convolution.

Lemma 40 (Convolution property [Lehmann et al. \(2001\)](#) [Murota \(1996a\)](#)). *Let $v_1, \dots, v_z : S \rightarrow \mathbb{R}$ satisfy gross substitutes. Given a set $T \subseteq S$, let $\Pi(T)$ be the set of partitions of T into z groups. Thus, if $\pi \in \Pi(T)$, then $\bigcup_{x=1}^z \pi_x = T$, and $\pi_x \cap \pi_y = \emptyset$ for $x \neq y$. Then, the function $v^* : S \rightarrow \mathbb{R}$ given by*

$$v^*(T) = \max_{\pi \in \Pi(T)} \sum_{x=1}^z v_x(\pi_x(T)) \quad (25)$$

also satisfies gross substitutes.

Another very useful property was shown by [Fujishige and Yang \(2003\)](#).

Lemma 41 ([Fujishige and Yang \(2003\)](#)). *Let v satisfy gross substitutes. Let T_1, T_2, S be disjoint sets with $|T_1| = |T_2|$. Let $j \in T_1$. Then, we have*

$$v(T_1 | S) + v(T_2 | S) \leq \max_{j' \in T_2 \setminus T_1} v(T_1 \cup \{j'\} \setminus \{j\} | S) + v(T_2 \cup \{j\} \setminus \{j'\} | S) \quad (26)$$

In particular, when $|T_1| = |T_2| = 2$, this becomes

$$v(j_1, j_2 | S) + v(j_3, j_4 | S) \leq \max \left[v(j_1, j_3 | S) + v(j_2, j_4 | S), \right. \quad (27)$$

$$\left. v(j_1, j_4 | S) + v(j_2, j_3 | S) \right] \quad (28)$$

Next, we have the matroid property of gross substitutes valuations, which shows that optimal bundles that also have a fixed size form the bases of a matroid. This proof is similar to a proof of [Gul and Stacchetti \(2000\)](#) that the optimal bundles of minimum size form the bases of a matroid.

Lemma 42 (Matroid property). *Let $v : 2^S \rightarrow \mathbb{R}$ satisfy gross substitutes, $\ell \in \mathbb{N}$ with $\ell \leq |S|$ and p any price vector. Then, the set of bundles maximizing $u(\cdot, p)$ among bundles of size ℓ form the bases of a matroid.*

Proof. We show that the exchange property holds. Let T_1, T_2 be two bundles maximizing $u(\cdot, p)$ among bundles of size ℓ , such that $u(T_1, p) = u(T_2, p) = u^*$. Fix $j_1 \in T_1 \setminus T_2$. Let \bar{v} be defined as in [Lemma 39](#), for a large constant M . Thus, bundles maximizing $u(\cdot, p)$ among bundles of size ℓ also maximize $\bar{u}(\cdot, p)$ among bundles of size ℓ . Consider the price vector p' given by $p'_j = p_j$ if $j \in T_1 \cup T_2 \setminus \{j_1\}$, and $p'_j = p_j + \epsilon$ otherwise. T_1 then no longer maximizes $\bar{u}(\cdot, p)$ among bundles of size ℓ , as it gives utility $\bar{u}(T_1, p') = u^* - \epsilon < u^* = \bar{u}(T_2, p)$. By (SI), there is a utility-improving local move. It is clear by construction of \bar{u} that we cannot profitably add or drop an item, so we must be able to profitably swap an item. There is then some $j_2 \notin T_1$ such that

$$\bar{u}(T_1, p') < \bar{u}(T_1 \cup \{j_2\} \setminus \{j_1\}, p') \quad (29)$$

Suppose $j_2 \notin T_2$. Then $p'_{j_2} = p_{j_2} + \epsilon$, which means that

$$\bar{u}(T_1, p) - \epsilon = \bar{u}(T_1, p') < \bar{u}(T_1 \cup \{j_2\} \setminus \{j_1\}, p') = \bar{u}(T_1 \cup \{j_2\} \setminus \{j_1\}, p) - \epsilon \quad (30)$$

By construction of \bar{v} and \bar{u} , we know that $\bar{u}(T, p) < \bar{u}(T', p)$ if and only if $u(T, p) < u(T', p)$. It follows that

$$u(T_1, p) < u(T_1 \cup \{j_2\} \setminus \{j_1\}, p) \quad (31)$$

contradicting optimality of T_1 . We see that $j_2 \in T_2 \setminus T_1$. Furthermore, we have that

$$u(T_1, p) - \epsilon = u(T_1, p') < u(T_1 \cup \{j_2\} \setminus \{j_1\}, p') = u(T_1 \cup \{j_2\} \setminus \{j_1\}, p) \quad (32)$$

for arbitrarily small $\epsilon > 0$. We see that it must be the case that $u(T_1 \cup \{j_2\} \setminus \{j_1\}, p) = u(T_1, p) = u^*$. Thus, the exchange property holds, and we conclude that the bundles which maximize $u(\cdot, p)$ among bundles of size ℓ form the bases of a matroid. \square

B Proof of Lemma 16

In this section, we prove [Lemma 16](#).

Proof of Lemma 16. First, assume (ν, p) is a competitive equilibrium such that all unit-buyers are matched. Because a MAT requires an unmatched unit-buyer as a root, a MAT cannot exist in $D(\nu, p)$.

Now, assume that a MAT does not exist in $D(\nu, p)$. We know by [Remark 15](#) that if an unmatched unit-buyer exists, a MAT must exist. Since there do not exist any MATs, all unit-buyers must be matched by ν . By definition, we know that stability is implied by partial stability plus unit-buyer perfectness. Since we are given partial stability, we have stability, and since we are given feasibility, we have that (ν, p) is a competitive equilibrium. \square

C Proofs for Section 4

In this section, we provide proofs omitted from [Section 4](#).

C.1 Proofs for Section 4.1

We prove [Lemma 19](#), which shows how to verify a MAT-preserving price increase using the slopes of the effective price functions.

Proof. First, we will show that for all $(k, j_1) \in \mathcal{T}'$ and all $j_2 \in F_{\nu', p}(k)$, we have $u_k(j_1 \mid T, p + \lambda d) \geq u_k(j_2 \mid T, p + \lambda d)$. Define $T = \nu'(\text{copy}(k) \setminus \{k\})$. We can write

$$u_k(j_1 \mid T, p + \lambda d) = v_k(j_1 \mid T) - q_{kj_1}(p_{j_1} + \lambda d_{j_1}) = v_k(j_1 \mid T) - q_{kj_1}(p_{j_1}) - \lambda q'_{kj_1}(p_{j_1})d_{j_1} \quad (33)$$

where the first equality is by definition and the second is by piecewise linearity. Note that since λ is arbitrarily small, all prices are within the same linear domain. Similarly, we have

$$u_k(j_2 | T, p + \lambda d) = v_k(j_2 | T) - q_{k j_2}(p_{j_2} + \lambda d_{j_2}) = v_k(j_2 | T) - q_{k j_2}(p_{j_2}) - \lambda q'_{k j_2}(p_{j_2}) d_{j_2} \quad (34)$$

Then, since $j_1, j_2 \in F_{\nu', p}(k)$, we know that

$$v_k(j_1 | T) - q_{k j_1}(p_{j_1}) = u_k(j_1 | T, p) = u_k(j_2 | T, p) = v_k(j_2 | T) - q_{k j_2}(p_{j_2}). \quad (35)$$

Since $q'_{k j_1}(p_{j_1}) d_{j_1} \leq q'_{k j_2}(p_{j_2}) d_{j_2}$, we know that

$$v_k(j_1 | T) - q_{k j_1}(p_{j_1}) - \lambda q'_{k j_1}(p_{j_1}) d_{j_1} \geq v_k(j_2 | T) - q_{k j_2}(p_{j_2}) - \lambda q'_{k j_2}(p_{j_2}) d_{j_2} \quad (36)$$

$$\implies u_k(j_1 | T, p + \lambda d) \geq u_k(j_2 | T, p + \lambda d). \quad (37)$$

Next, we will show that $(\nu', p + \lambda d)$ is partially stable, and $\mathcal{T}' \subseteq D(\nu', p + \lambda d)$. Let $(k, j_1) \in \nu'$. First, assume $d_{j_1} = 0$. Then, for any other item j_2 not matched to a copy of k , we have

$$\begin{aligned} u_k(j_1 | T, p + \lambda d) &= v_k(j_1 | T) - q_{k j_1}(p_{j_1} + \lambda d_{j_1}) \\ &= v_k(j_1 | T) - q_{k j_1}(p_{j_1}) \\ &= u_k(j_1 | T, p) \\ &\geq u_k(j_2 | T, p) \\ &= v_k(j_2 | T) - q_{k j_2}(p_{j_2}) \\ &\geq v_k(j_2 | T) - q_{k j_2}(p_{j_2} + \lambda d_{j_2}) \\ &= u_k(j_2 | T, p + \lambda d). \end{aligned}$$

Now assume $d_{j_1} > 0$. Then $(k, j_1) \in \mathcal{T}'$ by definition of a MAT-preserving price increase, and so

$$u_k(j_1 | T, p + \lambda d) \geq u_k(j_2 | T, p + \lambda d). \quad (38)$$

For $j_2 \notin F_{\nu, p}(k)$ and not matched to a copy of k , we have

$$u_k(j_1 | T, p) > u_k(j_2 | T, p). \quad (39)$$

Thus, for sufficiently small $\lambda > 0$, it is also true that

$$u_k(j_1 | T, p + \lambda d) > u_k(j_2 | T, p + \lambda d). \quad (40)$$

We conclude that $(\nu', p + \lambda d)$ is partially stable. Now, we will show that $\mathcal{T}' \subseteq D(\nu', p + \lambda d)$. Fix any $(k, j_1) \in \mathcal{T}'$ and any j_2 not matched to a copy of k . For $j_2 \in F_{\nu', p}(k)$, including any item potentially matched to k , we have

$$u_k(j_1 | T, p + \lambda d) \geq u_k(j_2 | T, p + \lambda d). \quad (41)$$

Then, for $j_2 \notin F_{\nu', p}(k)$ and not matched to a copy of k , we have

$$u_k(j_1 | T, p) > u_k(j_2 | T, p). \quad (42)$$

and as before for sufficiently small $\lambda > 0$ it is also true that

$$u_k(j_1 | T, p + \lambda d) > u_k(j_2 | T, p + \lambda d). \quad (43)$$

Thus, $j_1 \in F_{\nu', p + \lambda d}(k)$. We conclude that $\mathcal{T}' \subseteq D(\nu', p + \lambda d)$, and so \mathcal{T}' is indeed contained in the marginal demand graph $D(\nu', p + \lambda d)$. \square

C.2 Proofs for Section 4.2

Recall that we are given a price vector p , a partially stable outcome (ν, p) and a MAT \mathcal{T} in $D(\nu, p)$.

Proof of Lemma 20. Let \mathcal{J}_B be as in the definition of the lemma. Then clearly \mathcal{J}_B can be alternatively defined as follows:

$$\mathcal{J}_B = \arg \max_{E \in \mathcal{E}} \sum_{i \in B'(\mathcal{T})} u_i(S(E(i)) | \mu(i) \setminus S(\mathcal{T}), p), \quad (44)$$

where $\mathcal{E} = \prod_{i \in B'(\mathcal{T})} \mathcal{E}_i$ and $E(i) = \{(i, j) \in E\}$.

Let μ be the many-to-many projection of ν , and let $\hat{\mu} := \mu \setminus B(\mathcal{T}) \times S(\mathcal{T})$ be the edges of μ that do not include buyers and items in \mathcal{T} . For $T \subseteq S(\mathcal{T})$ and $i \in B'(\mathcal{T})$, define

$$\tilde{v}_i(T) = \begin{cases} v_i(T \mid \hat{\mu}(i)) & |T| \leq |U'(\mathcal{T}) \cap \text{ubuy}(i)| \\ v_i(T \mid \hat{\mu}(i)) - M(|T| - |U'(\mathcal{T}) \cap \text{ubuy}(i)|) & \text{o.w.}, \end{cases} \quad (45)$$

where M is a large constant. Then, for a set of edges $E \subseteq W(\mathcal{T})$, define

$$\hat{v}_i(E) := \tilde{v}_i(E(i)) \quad (46)$$

$$\hat{u}_i(E) := \hat{v}_i(E) - \sum_{(i,j) \in E} q_{ij}(p_j). \quad (47)$$

Similarly, for $E \subseteq W(\mathcal{T})$ define

$$v^*(E) := \max_{\pi \in \Pi(E)} \sum_{i \in B'(\mathcal{T})} \hat{v}_i(\pi_i) \quad (48)$$

$$u^*(E) := v^*(E) - \sum_{(i,j) \in E} q_{ij}(p_j). \quad (49)$$

We claim that

$$\mathcal{J}_B = \arg \max_{E \subseteq W(\mathcal{T}) : |E| = |\nu \cap \mathcal{T}|} u^*(E). \quad (50)$$

First, assume E is an element of the right-hand side of (50). Then, by construction of \hat{v}_i , we know that for an optimal partition $\pi \in \Pi(E)$ maximizing (48), we have

$$|\pi_i| = |U'(\mathcal{T}) \cap \text{ubuy}(i)|. \quad (51)$$

This equality is because all $\sum_{j \in S(\mathcal{T})} \text{cap}(j) = \sum_{i \in B'(\mathcal{T})} |U'(\mathcal{T}) \cap \text{ubuy}(i)|$ item slots in E must be allocated, and by construction of \hat{v}_i , any buyer i allocated more than $|U'(\mathcal{T}) \cap \text{ubuy}(i)|$ elements of E would lose astronomically large utility M . Now, consider the set of edges $\mu \cap W(\mathcal{T})$ and its partition $\pi^* \in \Pi(\mu \cap W(\mathcal{T}))$ given by $\pi_i^* = \mu \cap (\{i\} \times S(\mathcal{T}))$ for $i \in B'(\mathcal{T})$. By partial stability of ν , we know that

$$\hat{u}_i(\pi_i^*) \geq \hat{u}_i(E_i) \quad (52)$$

for all $E_i \subseteq W(\mathcal{T})$ with $|E_i| \leq |U'(\mathcal{T}) \cap \text{ubuy}(i)|$. Thus, we also know that

$$u^*(\mu \cap W(\mathcal{T})) = \sum_{i \in B'(\mathcal{T})} \hat{u}_i(\pi_i^*) \geq \sum_{i \in B'(\mathcal{T})} \hat{u}_i(\pi_i) \quad (53)$$

for all $E \subseteq W(\mathcal{T})$ with $|E| = |\nu \cap \mathcal{T}|$ and all partitions $\pi \in \Pi(E)$. It follows that for any E that maximizes $u^*(E)$ with optimal partition π , we cannot have $(i', j) \in \pi_i$, where $i' \neq i$. Thus, E also maximizes (44), and is an element of \mathcal{J}_B .

Now assume E is an element of \mathcal{J}_B . Then, $E(i)$ achieves maximum utility for buyer i 's utility function $u_i(\cdot \mid \mu(i) \setminus S(\mathcal{T}), p)$, among all bundles of size $|U'(\mathcal{T}) \cap \text{ubuy}(i)|$. Consider the partition π^* over E given by $\pi_i^* = E(i)$. Then, we know that

$$\begin{aligned} \hat{u}_i(\pi_i^*) &= u_i(E(i) \mid \mu(i) \setminus S(\mathcal{T}), p) \\ &\geq u_i(E'(i) \mid \mu(i) \setminus S(\mathcal{T}), p) \\ &\geq \hat{u}_i(\pi_i') \end{aligned}$$

for any set of edges $E' \subseteq W(\mathcal{T})$ and any partition π' of E' . We see that $E(i)$ maximizes $\hat{u}_i(\cdot)$ among sets of size $|U'(\mathcal{T}) \cap \text{ubuy}(i)|$. It follows that E , via the partition π^* , maximizes $u^*(\cdot)$ among sets of size $|\nu \cap \mathcal{T}|$. We conclude that (50) holds.

It then suffices to show that the right-hand side of (50), that is, $\arg \max_{E \subseteq W(\mathcal{T}) : |E| = |\nu \cap \mathcal{T}|} u^*(E)$, is the set of bases of a matroid. Suppose for the moment that \hat{v}_i satisfies gross substitutes for $i \in B'(\mathcal{T})$. Then v^* satisfies gross substitutes by Lemma 40. Hence, we can apply Lemma 42 to conclude that $\arg \max_{E \subseteq W(\mathcal{T}) : |E| = |\nu \cap \mathcal{T}|} u^*(E)$ form the bases of a matroid, as required.

Thus, let us show that, for $i \in B'(\mathcal{T})$, the valuation function \hat{v}_i satisfies gross substitutes. To do this, we will show that the property (ISO) from Proposition 37 holds. Fix any $E \in W(\mathcal{T})$, and distinct edges $(i_1, j_1), (i_2, j_2), (i_3, j_3) \in W(\mathcal{T}) \setminus E$. Let $\bar{E} = \{j \mid (i, j) \in E\}$. We consider several cases.

Case 1: $i_1 = i_2 = i_3 = i$. Then, we can write

$$\begin{aligned} \hat{v}_i((i_1, j_1), (i_2, j_2) \mid E) + \hat{v}_i((i_3, j_3) \mid E) &= \tilde{v}_i(j_1, j_2 \mid \bar{E}) + \tilde{v}_i(j_3 \mid \bar{E}) \\ &\leq \max \left[\tilde{v}_i(j_1, j_3 \mid \bar{E}) + \tilde{v}_i(j_2 \mid \bar{E}), \tilde{v}_i(j_2, j_3 \mid \bar{E}) + \tilde{v}_i(j_1 \mid \bar{E}) \right]. \end{aligned}$$

The first equality is by definition of \hat{v} , and the last inequality is by (ISO) on the function \tilde{v}_i . Then, we can rewrite this as

$$= \max \left[\hat{v}_i((i_1, j_1), (i_3, j_3) \mid E) + \tilde{v}_i((i_2, j_2) \mid E), \hat{v}_i((i_2, j_2), (i_3, j_3) \mid E) + \tilde{v}_i((i_1, j_1) \mid E) \right] \quad (54)$$

as desired.

Case 2: $i_1 \neq i$. Then, we have

$$\begin{aligned} \hat{v}_i((i_1, j_1), (i_2, j_2) \mid E) + \hat{v}_i((i_3, j_3) \mid E) &= \tilde{v}_i(j_2 \mid \bar{E}) + \tilde{v}_i(j_3 \mid \bar{E}) \\ &\leq \max \left[\tilde{v}_i(j_3 \mid \bar{E}) + \tilde{v}_i(j_2 \mid \bar{E}), \tilde{v}_i(j_2, j_3 \mid \bar{E}) + \tilde{v}_i(j_1 \mid \bar{E}) \right] \\ &= \max \left[\hat{v}_i((i_1, j_1), (i_3, j_3) \mid E) + \hat{v}_i((i_2, j_2) \mid E), \hat{v}_i((i_2, j_2), (i_3, j_3) \mid E) + \hat{v}_i((i_1, j_1) \mid E) \right] \end{aligned}$$

The inequality follows since the first element in the max is equal to the LHS of the inequality.

Case 3: $i_2 \neq i$. This case follows from a symmetric argument to the previous case.

Case 4: $i_3 \neq i$. Then, we have

$$\begin{aligned} \hat{v}_i((i_1, j_1), (i_2, j_2) \mid E) + \hat{v}_i((i_3, j_3) \mid E) &= \tilde{v}_i(j_1, j_2 \mid \bar{E}) \\ &\leq \tilde{v}_i(j_1 \mid \bar{E}) + \tilde{v}_i(j_2 \mid \bar{E}) \\ &\leq \max \left[\hat{v}_i((i_1, j_1), (i_3, j_3) \mid E) + \hat{v}_i((i_2, j_2) \mid E), \hat{v}_i((i_2, j_2), (i_3, j_3) \mid E) + \hat{v}_i((i_1, j_1) \mid E) \right] \end{aligned}$$

The middle inequality follows from submodularity of v_i . We see that in all cases, (ISO) holds. We conclude that \hat{v}_i satisfies gross substitutes. Thus, the elements of (50) form the bases of a matroid. \square

We next prove Lemma 21, and show that an independence oracle for M_B can be implemented efficiently. The oracle, described formally in Algorithm 5, proceeds as follows. Let $E \subseteq W(\mathcal{T})$ be the input set and μ' be obtained by restricting the many-to-many projection μ of ν to edges in \mathcal{T} , where (ν, p) is the current partially stable outcome. Using Lemma 20, we show that μ' is a basis of M_B . The main idea is to use the exchange property (Lemma 10) to iteratively replace in μ' a buyer-item pair $(i, j) \in E \setminus \mu'$ with a buyer-item pair $(i', j') \in \mu' \setminus E$. If this exchange is possible, the algorithm updates μ' and repeats until E is contained in the basis μ' (line 5), thus showing that E is independent. If there is no such exchange, the algorithm concludes that E is not independent, and terminates (line 8).

ALGORITHM 5: M_B -INDEPENDENCE

Input: Partially stable outcome (ν, p) , MAT \mathcal{T} where all items are fully matched, set of edges $E \subseteq B'(\mathcal{T}) \times S(\mathcal{T})$.

Output: *True* if $E \in \mathcal{I}_B$ or *False* otherwise.

```

1  $\mu' \leftarrow \{(buy(k), j) \mid (k, j) \in \nu \cap \mathcal{T}\}$ 
2 while  $E \not\subseteq \mu'$  do
3    $(i, j) \leftarrow$  any edge in  $E \setminus \mu'$ 
4   if  $\exists (i', j') \in \mu' \setminus E$  s.t.  $u^*(\mu') = u^*(\mu' \cup \{(i', j')\} \setminus \{(i, j)\})$  then           /* check exchange */
5      $\mu' \leftarrow \mu' \cup \{(i', j')\} \setminus \{(i, j)\}$                                        /* make exchange */
6   else
7     return False
8 return True

```

Proof of Lemma 21. We first claim that $\mu' = \{(\text{buy}(k), j) \mid (k, j) \in \nu \cap \mathcal{T}\}$ (as defined in line 1 of Algorithm 5) is a basis of M_B . To see this, we can write using the definitions of u^* , v^* , and \tilde{v} that

$$\begin{aligned} u^*(\mu') &= v^*(\mu') - \sum_{(i,j) \in \mu'} q_{ij}(p_j) \\ &= \sum_{i \in B'(\mathcal{T})} \left(\tilde{v}_i(\mu'(i)) - \sum_{j \in \mu'(i)} q_{ij}(p_j) \right) \\ &= \sum_{i \in B'(\mathcal{T})} \left(v_i(\mu'(i) \mid \hat{\mu}(i)) - \sum_{j \in \mu'(i)} q_{ij}(p_j) \right) \\ &= \sum_{i \in B'(\mathcal{T})} u_i(\mu'(i) \mid \hat{\mu}(i), p). \end{aligned}$$

Then, for any $T'_i \subseteq S(\mathcal{T})$ with $|T'_i| = |U'(\mathcal{T}) \cap \text{ubuy}(i)|$, we have $|T'_i \cup \hat{\mu}(i)| = |\mu(i)|$. By partial stability of (ν, p) , we have

$$u_i(\mu'(i) \mid \hat{\mu}(i), p) = u_i(\mu(i), p) - u_i(\hat{\mu}(i), p) \geq u_i(T'_i \cup \hat{\mu}(i), p) - u_i(\hat{\mu}(i), p) = u_i(T'_i \mid \hat{\mu}(i), p).$$

Let E' be a basis of M_B . By (51), E' is the disjoint union of a collection of sets $\{T'_i \mid i \in B'(\mathcal{T})\}$ with $|T'_i| = |U'(\mathcal{T}) \cap \text{ubuy}(i)|$ for all i . Thus, we have

$$\begin{aligned} u^*(\mu') &= \sum_{i \in B'(\mathcal{T})} u_i(\mu'(i) \mid \hat{\mu}(i), p) \geq \sum_{i \in B'(\mathcal{T})} u_i(T'_i \mid \hat{\mu}(i), p) \\ &= \sum_{i \in B'(\mathcal{T})} \left(v_i(T'_i \mid \hat{\mu}(i)) - \sum_{j \in T'_i} q_{ij}(p_j) \right) \\ &= \sum_{i \in B'(\mathcal{T})} \left(\hat{v}_i(E'_i) - \sum_{(i,j) \in E'_i} q_{ij}(p_j) \right) \\ &= v^*(E') - \sum_{(i,j) \in E'} q_{ij}(p_j) \\ &= u^*(E'), \end{aligned}$$

where $E'_i = \{(i, j) \mid j \in T'_i\}$ and $E' = \bigcup_{i \in B'(\mathcal{T})} E'_i$. It follows that that μ' is a basis of M_B .

Suppose first E is independent. Then $E \subseteq \mu'$ for some basis E' . By the basis exchange property (Lemma 10) for each $(i', j') \in E \setminus \mu'$ there is some $(i, j) \in \mu' \setminus E'$ such that $\mu' \cup \{(i', j')\} \setminus \{(i, j)\}$ is also a basis. To find such a pair $(i, j), (i', j')$, following (50), it suffices to verify if $\mu' \cup \{(i', j')\} \setminus \{(i, j)\}$ is an optimal set of edges for u^* . If this is true, then the if statement in line 4 activates, and setting $\mu' \leftarrow \mu' \cup \{(i', j')\} \setminus \{(i, j)\}$ in line 5 preserves the fact that μ' is a basis of M_B and strictly decreases $|E \setminus \mu'|$. Thus, the while loop repeats $O(|E|) = O(|B||S|)$ times until $E \subseteq \mu'$, at which point the algorithm completes the while loop and correctly returns *True*.

Now suppose E is not independent. Since, as argued above, every repetition of line 4 decreases $|E \setminus \mu'|$ while preserving the fact that μ' is a basis of \mathcal{J}_B , at some iteration there is no selection of (i, j) that makes line 4 true. Thus, the algorithm moves to line 7 and correctly returns *False*.

Now we will analyze the runtime of the algorithm. Line 1 requires $O(|B||S|)$. The while loop from lines 2-8 requires at most $O(|B||S|)$ iterations, since in each iteration it either terminates in line 8 or increases $|E \cap \mu'|$ by 1. The if statement in line 4 requires $O(|B||S|)$ time to verify, by enumeration. We conclude that the total runtime is $O(|B|^2|S|^2)$. \square

Now, we can prove Lemma 26, showing that the minimum weight common basis problem correctly solves the minimum weight partially stable perfect matching problem.

Proof of Lemma 26. Notice that the original partially stable matching ν , when restricted to $\nu \cap \mathcal{T}$, has its many-to-one projection μ' in \mathcal{J}_B . This membership is because by definition of partial stability, $\mu'(i)$ gives bidder i the maximum value for $u_i(\cdot \mid \mu(i) \setminus S(\mathcal{T}), p)$ among sets of size $|U'(\mathcal{T}) \cap \text{ubuy}(i)|$. It follows

that any element $\mu'' \in \mathcal{J}_B$ must achieve the same value of $u_i(\cdot \mid \mu(i) \setminus S(\mathcal{T}), p)$ for each i , and thus $\mu'' \cup (\mu \cap \mathcal{T})$ is a partially stable matching, aside from the fact that items may be allocated beyond their capacity. It follows from Lemma 23 and the definition of partial stability that μ^* is exactly a minimum weight partially stable matching on $W(\mathcal{T})$. Then, by definition, ν^* is also partially stable on $V(\mathcal{T}) \setminus \{k_0\}$. Furthermore, for any many-to-one matching $\nu \subseteq U'(\mathcal{T}) \times S(\mathcal{T})$, we know that ν has the same weight as its many-to-many projection $\{(\text{buy}(k), j) \mid (k, j) \in \nu\}$. It follows that ν^* is also of minimum weight. \square

To solve the minimum weight common basis problem, we can use any existing algorithm for the maximum weight matroid intersection problem, such as those presented by Lawler (1975) and Frank (1981).

Lemma 43 (Max weight matroid intersection algorithm Lawler (1975), Frank (1981)). *There exists an algorithm MATROIDINTERSECTION which, given matroid independence oracles for M_B and M_S , can find a maximum weight matroid intersection in $O(|B|^3|S|^3)$ oracle calls.*

Then, we can transform the weights to change a minimum weight common basis problem into a maximum weight matroid intersection problem, to prove Lemma 27.

Proof of Lemma 27. Define weights w over the edges in $W(\mathcal{T})$ given by

$$w'((i, j)) = \log q'_{ij}(p_j) - \min_{(i', j') \in W(\mathcal{T})} \log q'_{i'j'}(p_{j'}), \quad (55)$$

$$w((i, j)) = \left(\sum_{j \in S(\mathcal{T})} \text{cap}(j) \right) \left(\max_{(i', j') \in W(\mathcal{T})} w'((i', j')) \right) + 1 - w'((i, j)). \quad (56)$$

These weights are the typical transformation used to turn a minimum weight maximum cardinality matching problem into a maximum weight matching problem, by first translating and reflecting the weight of each edge to be negative then adding a large constant to the weight of each edge. We see that if $|E| > |E'|$, then $w(E) > w(E')$. Thus, any maximum weight intersection must be of maximum cardinality. Since we know there exists a common basis of M_B and M_S , i.e. the original matching ν , any maximum weight intersection must also have size at least $|\nu|$, and thus is a basis of both M_B and M_S . We see that by applying MATROIDINTERSECTION to the weights given by w , we obtain a minimum weight common basis. \square

C.3 Proofs for Section 4.3

First, we use properties of gross substitutes to prove Lemma 31, describing the structure of the marginal demand graph.

Proof of Lemma 31. By hypothesis and definition of demand graph, the set $\{j_1, \dots, j_z, j'_1, \dots, j'_z\}$ is composed of $2z$ distinct elements. For each $X \subseteq [z]$, define $T(X) := \nu(\text{ubuy}(i)) \cup \{j'_x \mid x \in X\} \setminus \{j_x \mid x \in X\}$.

Assume (a) is not true. Then, let y be the largest number such that for all sets X with $|X| \leq y$, the bundle $T(X)$ optimizes $u_i(\cdot, p)$ among bundles of size $|\nu(\text{ubuy}(i))|$. Note that by definition of the marginal demand graph, this is always true for $y = 1$. Relabel the pairs so that $T([y+1])$ does not optimize $u_i(\cdot, p)$ among bundles of size $|\nu(\text{ubuy}(i))|$. Define $T = \nu(\text{ubuy}(i)) \cup \{j'_3, \dots, j'_{y+1}\} \setminus \{j_1, \dots, j_{y+1}\}$. By Lemma 41, we know that

$$u_i(j_1, j'_2 \mid T, p) + u_i(j'_1, j_2 \mid T, p) \leq \max \left[u_i(j_1, j_2 \mid T, p) + u_i(j'_1, j'_2 \mid T, p), \right. \\ \left. u_i(j_1, j'_1 \mid T, p) + u_i(j_2, j'_2 \mid T, p) \right].$$

By selection of T , we know that $T \cup \{j'_1, j_2\}$, $T \cup \{j_1, j'_2\}$, and $T \cup \{j_1, j_2\}$ optimize $u_i(\cdot, p)$ among bundles of size $|\nu(\text{ubuy}(i))|$. Then, we have

$$u_i(j_1, j'_2 \mid T, p) = u_i(j'_1, j_2 \mid T, p) = u_i(j_1, j_2 \mid T, p) = \max_{j, j'} u_i(j, j' \mid T, p) := u^*. \quad (57)$$

As a result, we have

$$u^* + u^* \leq \max \left[u^* + u_i(j'_1, j'_2 \mid T, p), u_i(j_1, j'_1 \mid T, p) + u_i(j_2, j'_2 \mid T, p) \right]. \quad (58)$$

Furthermore, each of the four terms on the RHS of (58) are bounded from above by u^* . Since $T \cup \{j'_1, j'_2\} = T([y+1])$, we know this bundle does not maximize $u_i(\cdot, p)$ among bundles of size $|\nu(\text{ubuy}(i))|$, so we have $u_i(j'_1, j'_2 | T, p) < u^*$. This strict inequality means that the first argument of the max is less than $2u^*$, so the second argument of the max must be equal to $2u^*$. It follows that

$$u_i(j_1, j'_1 | T, p) = u_i(j_2, j'_2 | T, p) = u^*. \quad (59)$$

That is, $T \cup \{j_1, j'_1\}$ and $T \cup \{j_2, j'_2\}$ optimize $u_i(\cdot, p)$ among bundles of size $|\nu(\text{ubuy}(i))|$. By Lemma 42, the set of bundles which optimize $u_i(\cdot, p)$ among bundles of size $|\nu(\text{ubuy}(i))|$ form the bases of a matroid. Consider two such bundles, $\nu(\text{ubuy}(i))$ and $T \cup \{j_1, j'_1\}$. By the bijective basis exchange property, there is a bijection $f : \{j_2, \dots, j_{y+1}\} \rightarrow \{j'_1, j'_3, \dots, j'_{y+1}\}$ such that for every $j \in \{j_2, \dots, j_{y+1}\} = \nu(\text{ubuy}(i)) \setminus (T \cup \{j_1, j'_1\})$, we have $\nu(\text{ubuy}(i)) \cup \{f(j)\} \setminus \{j\}$ optimizes $u_i(\cdot, p)$ among bundles of size $|\nu(\text{ubuy}(i))|$ as well. Let $x_1 = 1$, and let x_2 so that $j'_{x_1} = f(j_{x_2})$. We see that $\nu(\text{ubuy}(i)) \cup \{j'_1\} \setminus \{j_{x_2}\}$ is an optimal size $|\nu(\text{ubuy}(i))|$ bundle, which means by definition of the marginal demand graph that $(k_{x_2}, j'_1) \in D(\nu, p)$. Setting x_3 so that $j'_{x_2} = f(j_{x_3})$, we then see by the same reasoning that $(k_{x_3}, j'_{x_2}) \in D(\nu, p)$. This pattern continues until $j'_{x_{t-1}} = f(j_{x_t})$, where $x_t = 2$. Thus, we have a sequence x_1, \dots, x_t such that $(k_{x_{\ell+1}}, j'_{x_\ell})$ for $1 \leq \ell \leq t-1$, with $x_1 = 1$ and $x_t = 2$. Overall, this gives the path

$$(k_1 = k_{x_1}, j'_{x_1}, k_{x_2}, j'_{x_2}, \dots, j'_{x_{t-1}}, k_{x_t} = k_2) \quad (60)$$

Now consider bundles $\nu(\text{ubuy}(i))$ and $T \cup \{j_2, j'_2\}$, which optimize $u_i(\cdot, p)$ among bundles of size $|\nu(\text{ubuy}(i))|$. By the bijective basis exchange property, there is a bijection $g : \{j_1, j_3, \dots, j_{y+1}\} \rightarrow \{j'_2, \dots, j'_{y+1}\}$ such that for every $j \in \{j_1, j_3, \dots, j_{y+1}\}$, we have $\nu(\text{ubuy}(i)) \cup \{g(j)\} \setminus \{j\}$ is an optimal size $|\nu(\text{ubuy}(i))|$ bundle as well. Keeping $x_t = 2$, let $j'_{x_t} = g(j_{x_{t+1}})$. By the same reasoning as before, we see that $(k_{x_{t+1}}, j_{x_t}) \in D(\nu, p)$. Again, this creates a sequence until $j'_{x_s} = g(j_{x_{s+1}})$, where $x_{s+1} = 1$. We thus have another sequence x_t, \dots, x_{s+1} such that $(k_{x_{\ell+1}}, j'_{x_\ell})$ for $t \leq \ell \leq s+1$, with $x_t = 2$ and $x_{s+1} = 1$. This gives us the path

$$(k_2 = k_{x_t}, j'_{x_t}, k_{x_{t+1}}, j'_{x_{t+1}}, \dots, k_{x_s}, j'_{x_s}, k_{x_{s+1}} = k_1) \quad (61)$$

Combining this with the previous path gives us a closed walk

$$(k_1 = k_{x_1}, j'_{x_1}, k_{x_2}, j'_{x_2}, \dots, j'_{x_{t-1}}, k_{x_t} = k_2, j'_{x_t}, k_{x_{t+1}}, j'_{x_{t+1}}, \dots, k_{x_s}, j'_{x_s}, k_{x_{s+1}} = k_1) \quad (62)$$

in $D(\nu, p) \setminus \nu$. However, some nodes and edges may be repeated in this walk. If there are indices $1 \leq r < r' \leq s$ such that $x_r = x_{r'}$, then we can shortcut the cycle to become

$$(k_1 = k_{x_1}, j'_{x_1}, k_{x_2}, \dots, j'_{x_{r-1}}, k_{x_r} = k_{x_{r'}}, j'_{x_{r'}}, k_{x_{r'+1}}, \dots, k_{x_s}, j'_{x_s}, k_{x_{s+1}} = k_1)$$

This gives us a new, shorter sequence of indices and corresponding cycle. Furthermore, we can see that the cycle still alternates between the original edges $(k_{x_\ell}, j'_{x_\ell})$ and our newly found edges $(k_{x_\ell}, j'_{x_{\ell-1}})$. If there continue to be duplicate indices in this sequence, then we can repeat this shortcutting process, until the final sequence does not have any duplicate indices, satisfying (b). \square

Next, it has also been shown by Edmonds and Karp (1972) that LP and DP can be solved efficiently.

Lemma 44 (Complexity of LP Edmonds and Karp (1972)). *LP can be solved in $O(|B|^3|S|^3)$ time.*

Finally, we can prove Lemma 32 on the correctness and runtime of Algorithm 2.

Proof of Lemma 32. First, note that $\omega_{k_0}^*$ is chosen in line 2 so that the constraint $\omega_{k_0} + \rho_j \leq \log q'_{k_0 j}(p_j)$ is satisfied for all $j \in S(\mathcal{T})$, and furthermore at least one of these constraints is tight.

Consider the main loop of Algorithm 2. We can verify that in each iteration, if $k \in U(\mathcal{T}')$ and $j \in S(\mathcal{T}')$, then $\omega_k + \rho_j < (=) \log q'_{k j}(p_j)$ before the iteration implies $\omega_k + \rho_j < (=) \log q'_{k j}(p_j)$ after the iteration as well. Similarly, if $k \in U(\mathcal{T}) \setminus U(\mathcal{T}')$ and $j \in S(\mathcal{T}) \setminus S(\mathcal{T}')$, then the relationship $\omega_k + \rho_j < (=) \log q'_{k j}(p_j)$ is preserved. As a result, once an agent enters \mathcal{T}' , they remain there throughout the loop.

Next, immediately after ending each iteration of the loop, there is at least one $k \in U(\mathcal{T}')$ and $j \in S(\mathcal{T}) \setminus S(\mathcal{T}')$ such that $\omega_k + \rho_j = \log q'_{k j}(p_j)$. It follows that after the next execution of lines 4-5, (k, j) enters \mathcal{T}' and j enters $S(\mathcal{T}')$. We also know that j was matched to at least one other unit-buyer $k_1 \in U(\mathcal{T}) \setminus U(\mathcal{T}')$, which enters $U(\mathcal{T}')$ after line 5 as well. Thus, the size of \mathcal{T}' increases each iteration, for at most $|S(\mathcal{T})|$ iterations, at which point all items in $S(\mathcal{T})$ and all of their matched unit-buyers, which is all of $U(\mathcal{T})$, are in \mathcal{T}' .

For the runtime, note that the loop takes $O(|S|)$ iterations, since each iteration adds at least one unit-buyer and item to \mathcal{T}' . Each iteration of lines 4-8 take $O(|B||S|^2)$, by Lemma 34. We conclude that the entire algorithm takes $O(|B||S|^3)$ time. \square

C.4 Proofs for Section 4.4

In this section, we prove Lemma 33 on the correctness and runtime of Algorithm 3.

Proof of Lemma 33. First, by Lemma 29, ν^* is an optimal solution for $DP(\nu^*, p, \mathcal{T})$. By duality, we see that for ω^*, ρ^* as computed in line 3, we have $\omega_k^* + \rho_j^* = \log q'_{\text{buy}(k)j}(p_j)$ if $x_{kj} = 1$, and $\omega_k^* + \rho_j^* \leq \log q'_{\text{buy}(k)j}(p_j)$ for all $(k, j) \in D(\nu, p) \times U(\mathcal{T}) \setminus \{k_0\} \times S(\mathcal{T})$.

After line 4 in Algorithm 3, we know by Lemma 32 that there exists a MAT \mathcal{T}' that contains all of $U(\mathcal{T}) \cup S(\mathcal{T})$, such that for each edge $(k, j) \in \mathcal{T}'$, we have $\omega_k^* + \rho_j^* = \log q'_{\text{buy}(k)j}(p_j)$. For all $(k, j) \in D(\nu^*, p)$, we also still have $\omega_k^* + \rho_j^* \leq \log q'_{\text{buy}(k)j}(p_j)$. As a result, by taking e to the power of these (in)equalities, it follows that for $(k, j_1) \in \mathcal{T}'$ and $(k, j_2) \in D(\nu^*, p)$,

$$q'_{k j_1}(p_{j_1})d_{j_1} = q'_{k j_1}(p_{j_1})e^{-\rho_{j_1}^*} = e^{\omega_k^*} \leq q'_{k j_2}(p_{j_2})e^{-\rho_{j_2}^*} = q'_{k j_2}(p_{j_2})d_{j_2}. \quad (63)$$

By Lemma 19, d is a MAT-preserving price increase, and so ν^* is partially stable at $p + \lambda^*d$, and $\mathcal{T}' \subseteq D(\nu^*, p + \lambda^*d)$.

Now we will verify that the algorithm also finds the correct step size λ^* . In line 6, we ensure that λ_1^* is the minimum step size such that a new edge in $U(\mathcal{T}) \times S \setminus S(\mathcal{T})$ enters the demand graph. In line 7, we ensure that λ_2^* is the minimum step size required to reach the edge of the current linear domain. It follows that $p + \lambda^*d$ either has a new edge enter the demand graph, which is also adds an edge and an item to the MAT rooted at k_0 , or $p + \lambda^*d$ reaches the edge of the current linear domain.

Now we will consider the runtime of Algorithm 3. Line 1 takes $O(|B|^5|S|^5)$, by Lemmas 21, 24, and 27. Line 3 can be solved in $O(|B|^3|S|^3)$ time by Lemma 44. Line 4 takes $O(|B||S|^3)$ times by Lemma 32. Lines 5-8 require $O(|B||S|^2)$. In total, the entire algorithm takes $O(|B|^5|S|^5)$ time. \square

D Proofs for Section 6

In this section, we provide the proof for Theorem 2.

Proof of Theorem 2. We will reduce the integer knapsack problem to an instance of our problem. Consider any knapsack instance with items S , where each item $j \in S$ has value w_j and cost c_j , and we have a maximum budget C . We assume that w_j, c_j , and C are all integers. The goal is to maximize

$$\begin{aligned} & \max \sum_{j \in S} w_j x_j \\ & \text{s.t. } \sum_{j \in S} c_j x_j \leq C \\ & \quad x_j \in \{0, 1\} \quad \forall j \in S \end{aligned}$$

Now consider the following instance of the assignment game. The item set S is the same, and each item j has capacity $\text{cap}(j) = 2$. For each item j , define three buyers i_j^1, i_j^2, i_j^3 that have additive, quasilinear utilities given by

$$\begin{aligned} v_{i_j^\ell}(j) &= c_j \\ v_{i_j^\ell}(j') &= 0 \quad \forall j' \neq j \\ v_{i_j^\ell}(T) &= \sum_{j' \in T} v_{i_j^\ell}(j') \\ r_{i_j^\ell}(p) &= p \end{aligned}$$

Thus, i_j^1, i_j^2 , and i_j^3 only value item j , at c_j . Also define a buyer i_0 that has utility given by

$$\begin{aligned} v_{i_0}(T) &= \sum_{j \in T} w_j \\ r_{i_0}(p) &= \begin{cases} \frac{1}{2C}p & p \leq C \\ \frac{1}{2} + (1 + \sum_{j \in S} w_j)(p - C) & p > C \end{cases} \end{aligned}$$

Now consider any competitive equilibrium (μ, p) . If an item j has $p_j < c_j$, then all three buyers i_j^1, i_j^2, i_j^3 demand j , which would not be a competitive equilibrium. If $p_j > c_j$, then none of the buyers i_j^1, i_j^2, i_j^3 demand j , so at most one buyer (i_0) can match to j . Thus, j is undermatched but has a nonzero price, which is impossible. It follows that $p_j = c_j$ for all items $j \in S$. Then, consider the set of items $\mu(i_0)$ matched to i_0 . For any bundle T where the sum of the costs c_j greater than C , we have

$$\begin{aligned} u_{i_0}(T, p) &= v_{i_0}(T) - r_{i_0} \left(\sum_{j \in T} p_j \right) \\ &= \sum_{j \in T} w_j - r_{i_0} \left(\sum_{j \in T} c_j \right) \\ &\leq \sum_{j \in T} w_j - \left(\frac{1}{2} + 1 + \sum_{j \in S} w_j \right) \\ &\leq -\frac{3}{2}, \end{aligned}$$

where the first inequality follows from the fact that $\sum_{j \in T} c_j > C$ implies $\sum_{j \in T} c_j \geq C + 1$ with integer data. We see that i_0 cannot be matched to any bundle T whose costs sum to greater than C . Now consider $\mu(i_0)$, and any bundle T where the sum of the costs is at most C . Thus,

$$\begin{aligned} u_{i_0}(\mu(i_0), p) &\geq u_{i_0}(T, p) \\ \sum_{j \in \mu(i_0)} w_j - r_{i_0} \left(\sum_{j \in \mu(i_0)} p_j \right) &\geq \sum_{j \in T} w_j - r_{i_0} \left(\sum_{j \in T} p_j \right) \\ \sum_{j \in \mu(i_0)} w_j - r_{i_0} \left(\sum_{j \in \mu(i_0)} c_j \right) &\geq \sum_{j \in T} w_j - r_{i_0} \left(\sum_{j \in T} c_j \right). \end{aligned}$$

Since the costs sum to at most C , we know by the definition of r_{i_0} that

$$r_{i_0} \left(\sum_{j \in \mu(i_0)} c_j \right), r_{i_0} \left(\sum_{j \in T} c_j \right) \leq \frac{1}{2}. \quad (64)$$

As a result, since the w_j are integer, we know that

$$\sum_{j \in \mu(i_0)} w_j \geq \sum_{j \in T} w_j. \quad (65)$$

Thus, i_0 must be matched to a bundle $\mu(i_0)$ with total cost at most C that has the maximum total weight, which is exactly the optimal solution to the original integer knapsack problem. We conclude that computing a competitive equilibrium is at least as hard as the integer knapsack problem, which is NP-hard \square