

A NOTE ON THE THRESHOLD NUMBERS OF CYCLES

RUNZE WANG

ABSTRACT. A graph $G = (V, E)$ is said to be a k -threshold graph with thresholds $\theta_1 < \theta_2 < \dots < \theta_k$ if there is a map $r : V \rightarrow \mathbb{R}$ such that $uv \in E$ if and only if $\theta_i \leq r(u) + r(v)$ holds for an odd number of $i \in [k]$. The *threshold number* of G , denoted by $\Theta(G)$, is the smallest positive integer k such that G is a k -threshold graph. In this paper, we determine the exact threshold numbers of cycles by proving

$$\Theta(C_n) = \begin{cases} 1 & \text{if } n = 3, \\ 2 & \text{if } n = 4, \\ 4 & \text{if } n \geq 5, \end{cases}$$

where C_n is the cycle with n vertices.

1. INTRODUCTION

In this paper, for $a, b \in \mathbb{Z}$ with $a < b$, we denote $\{i \in \mathbb{Z} : a \leq i \leq b\}$ by $[a, b]$. For $c \in \mathbb{N}$, we denote $\{j \in \mathbb{N} : 1 \leq j \leq c\}$ by $[c]$.

A graph $G = (V, E)$ is said to be a *threshold graph* if there is a map $f : V \rightarrow \mathbb{R}$ such that $uv \in E$ if and only if $f(u) + f(v) \geq 0$. Threshold graphs were introduced by Chvátal and Hammer [2] in 1977, and have been extensively studied [3, 4, 5, 8] since then.

Jamison and Sprague [6] introduced multithreshold graphs as a generalization of threshold graphs. A graph $G = (V, E)$ is said to be a k -threshold graph with thresholds $\theta_1 < \theta_2 < \dots < \theta_k$ if there is a map $r : V \rightarrow \mathbb{R}$ such that $uv \in E$ if and only if $\theta_i \leq r(u) + r(v)$ holds for an odd number of $i \in [k]$. We say $r(v)$ is the *rank* of v . We say such a rank assignment $r : V \rightarrow \mathbb{R}$ is a $(\theta_1, \theta_2, \dots, \theta_k)$ -*representation* of G .

If G is a k -threshold graph, then the k thresholds divide \mathbb{R} into $k + 1$ parts in the form

$$\text{NO} \mid \text{YES} \mid \text{NO} \mid \text{YES} \mid \dots,$$

and every edge rank sum lands in a YES part, every nonedge rank sum lands in a NO part.

Jamison and Sprague [6] proved that if a graph has n vertices, then it is a k -threshold graph for some $k \leq \binom{n}{2}$. So every finite graph G has a smallest positive integer $\Theta(G)$, called the *threshold number* of G , such that G is a $\Theta(G)$ -threshold graph.

Date: June 2024.

For specific graphs: Jamison and Sprague [6] proved that the threshold number of any path is at most 2, and the threshold number of any caterpillar (obtained by attaching leaves to vertices in a path) is also at most 2; Chen and Hao [1] determined the exact threshold numbers of k -partite graphs with every part having at least $k + 1$ vertices; Kittipassorn and Sumalroj [7] determined the exact threshold numbers of multipartite graphs with every part having 3 vertices or every part having 4 vertices; Wang [9] determined the exact threshold numbers of multipartite graphs with every part having at most 3 vertices.

In this paper, we determine the exact threshold numbers of cycles.

Theorem 1.1. *Let $n \geq 3$ be an integer, let C_n be the cycle with n vertices, then*

$$\Theta(C_n) = \begin{cases} 1 & \text{if } n = 3, \\ 2 & \text{if } n = 4, \\ 4 & \text{if } n \geq 5. \end{cases}$$

In Section 2, we prove Theorem 1.1; In Section 3, we give some remarks and propose a problem.

2. PROOF OF THEOREM 1.1

For C_n , the cycle with n vertices, we randomly pick a vertex and label it v_1 , then counterclockwise label the other vertices v_2, v_3, \dots, v_n . Note that we can continue this labeling process, so each vertex v_i can also be represented as v_{i+kn} for any $k \in \mathbb{N}$.

Let us do the proof of Theorem 1.1.

Proof of Theorem 1.1. It is easy to check that $\Theta(C_3) = 1$.

For C_n with $n \geq 4$, we first show that $\Theta(C_n) \geq 2$.

In C_n with $n \geq 4$, v_1v_2 and v_3v_4 are edges, v_1v_3 and v_2v_4 are nonedges. Assume by contradiction that $\Theta(C_n) = 1$ and $r : V \rightarrow \mathbb{R}$ is a (θ) -representation of C_n for some θ . Then by the definition of multithreshold graphs, we have $r(v_1) + r(v_2) \geq \theta$, $r(v_3) + r(v_4) \geq \theta$, $r(v_1) + r(v_3) < \theta$, and $r(v_2) + r(v_4) < \theta$. But then we will have

$$(r(v_1) + r(v_2)) + (r(v_3) + r(v_4)) \geq 2\theta > (r(v_1) + r(v_3)) + (r(v_2) + r(v_4)),$$

contradiction. So $\Theta(C_n) \geq 2$.

For C_4 , if we let $\theta_1 = -1.1$ and $\theta_2 = 1.1$, and let $r(v_1) = 1$, $r(v_2) = -2$, $r(v_3) = 3$, and $r(v_4) = -4$, then every edge rank sum is -1 or 1 , and every nonedge rank sum is either smaller than -1.1 , or greater than 1.1 , so we have a $(-1.1, 1.1)$ -representation of C_4 . So $\Theta(C_4) \leq 2$, and hence $\Theta(C_4) = 2$.

Now let us assume $n \geq 5$.

Firstly, we show $\Theta(C_n) \leq 4$ by construction. Let $r(v_i) = (-1)^{i-1}i$ for $i \in [n-1]$, and let $r(v_n) = (-1)^{n-1}(n-0.5)$.

Case I. n is odd.

In this case $r(v_n) = n-0.5$. We let $\theta_1 = -1.1$, $\theta_2 = 1.1$, $\theta_3 = r(v_1) + r(v_n) - 0.1 = n+0.4$, and $\theta_4 = r(v_1) + r(v_n) + 0.1 = n+0.6$. We have $\theta_1 < \theta_2 < \theta_3 < \theta_4$. Denote $\{\theta_1, \theta_2, \theta_3, \theta_4\}$ by \mathcal{S} .

Then for an edge not involving v_n , its rank sum is -1 or 1 , which is between θ_1 and θ_2 . For edge $v_{n-1}v_n$, its rank sum is 0.5 , which is between θ_1 and θ_2 . For edge v_1v_n , its rank sum is $1 + (n-0.5) = n+0.5$, which is between θ_3 and θ_4 . We have checked every edge rank sum is greater than or equal to an odd number of elements in \mathcal{S} .

For a nonedge not involving v_n , assume its two vertices are v_i and v_j with $i, j \in [n-1]$ and $|i-j| \neq 1$, then:

- If $r(v_i) + r(v_j) < 0$, then by $|i-j| \neq 1$, we know that $r(v_i) + r(v_j) \leq -2 < \theta_1$, so $|\{s \in \mathcal{S} : s \leq r(v_i) + r(v_j)\}| = 0$, an even number.
- If $r(v_i) + r(v_j) > 0$, then by $|i-j| \neq 1$, we know that $r(v_i) + r(v_j) \geq 2 > \theta_2$. And $r(v_i) + r(v_j)$ is an integer, so it is either smaller than θ_3 or greater than θ_4 . So $|\{s \in \mathcal{S} : s \leq r(v_i) + r(v_j)\}|$ is 2 or 4, an even number.

And for a nonedge involving v_n , assume its the other vertex is v_i with $i \in [2, n-2]$, then:

- If $r(v_i) > 0$, then it is at least 3, so $r(v_i) + r(v_n) \geq 3 + (n-0.5) = n+2.5 > \theta_4$, so $|\{s \in \mathcal{S} : s \leq r(v_i) + r(v_j)\}| = 4$, an even number.
- If $r(v_i) < 0$, then $-(n-3) \leq r(v_i) \leq -2$, and $\theta_2 < 2.5 \leq r(v_i) + r(v_n) \leq n-2.5 < \theta_3$, so $|\{s \in \mathcal{S} : s \leq r(v_i) + r(v_j)\}| = 2$, an even number.

We have also checked every nonedge rank sum is greater than or equal to an even number of elements in \mathcal{S} . So we have a $(\theta_1, \theta_2, \theta_3, \theta_4)$ -representation of C_n , and hence $\Theta(C_n) \leq 4$.

Case II. n is even.

In this case, $r(v_n) = -n+0.5$. The only difference from Case I is we swap the positions of θ_1, θ_2 and θ_3, θ_4 , so we let $\theta_1 = r(v_1) + r(v_n) - 0.1 = -n+1.4$, $\theta_2 = r(v_1) + r(v_n) + 0.1 = -n+1.6$, $\theta_3 = -1.1$, and $\theta_4 = 1.1$. We have $\theta_1 < \theta_2 < \theta_3 < \theta_4$. Denote $\{\theta_1, \theta_2, \theta_3, \theta_4\}$ by \mathcal{S}' . The same as in Case I, we can check every edge rank sum is greater than or equal to an odd number of elements in \mathcal{S}' , and every nonedge rank sum is greater than or equal to an even number of elements in \mathcal{S}' . So we also have $\Theta(C_n) \leq 4$ in this case.

Now we have proved the upper bound $\Theta(C_n) \leq 4$.

And we already know $\Theta(C_n) \geq 2$. So by eliminating the possibility of $\Theta(C_n) = 2$ or $\Theta(C_n) = 3$, we can conclude that $\Theta(C_n) = 4$.

Firstly assume $\Theta(C_n) = 2$. So we have two thresholds θ_1, θ_2 , and a (θ_1, θ_2) -representation $r : V \rightarrow \mathbb{R}$. Two thresholds divide \mathbb{R} into three parts in the form

$$\text{NO} \mid \text{YES} \mid \text{NO}.$$

The two NO parts will be called smaller NO and larger NO.

We have three cases. The proofs of the first two follow the same pattern, the proof of the third one is slightly different.

Case i. $2 \nmid n$.

For vertices v_1, v_2, v_3, v_4 : We know v_1v_2 and v_3v_4 are edges, so both $r(v_1) + r(v_2)$ and $r(v_3) + r(v_4)$ are in YES. We know v_1v_3 and v_2v_4 are nonedges, so both $r(v_1) + r(v_3)$ and $r(v_2) + r(v_4)$ are in NO. And by the fact that $(r(v_1) + r(v_2)) + (r(v_3) + r(v_4)) = (r(v_1) + r(v_3)) + (r(v_2) + r(v_4))$, we know that $r(v_1) + r(v_3)$ and $r(v_2) + r(v_4)$ are in different NO's. Without loss of generality, we assume $r(v_1) + r(v_3)$ is in smaller NO and $r(v_2) + r(v_4)$ is in larger NO.

Then for vertices v_2, v_3, v_4, v_5 : Similarly we know that $r(v_2) + r(v_4)$ and $r(v_3) + r(v_5)$ are in different NO's, we already assumed $r(v_2) + r(v_4)$ is in larger NO, so $r(v_3) + r(v_5)$ is in smaller NO.

Repeatedly, we have that if i is odd then $r(v_i) + r(v_{i+2})$ is in smaller NO, if i is even then $r(v_i) + r(v_{i+2})$ is in larger NO. We take $i = n + 1$, because in this case $n + 1$ is even, we have $r(v_1) + r(v_3) = r(v_{n+1}) + r(v_{n+3})$ is in larger NO, which contradicts our assumption $r(v_1) + r(v_3)$ is in smaller NO.

Case ii. $2 \mid n$, but $4 \nmid n$.

For vertices $v_1, v_2, v_{\frac{n}{2}+1}, v_{\frac{n}{2}+2}$: We know v_1v_2 and $v_{\frac{n}{2}+1}v_{\frac{n}{2}+2}$ are edges, so both $r(v_1) + r(v_2)$ and $r(v_{\frac{n}{2}+1}) + r(v_{\frac{n}{2}+2})$ are in YES. We know $v_1v_{\frac{n}{2}+1}$ and $v_2v_{\frac{n}{2}+2}$ are nonedges, so both $r(v_1) + r(v_{\frac{n}{2}+1})$ and $r(v_2) + r(v_{\frac{n}{2}+2})$ are in NO. And by the fact that $(r(v_1) + r(v_2)) + (r(v_{\frac{n}{2}+1}) + r(v_{\frac{n}{2}+2})) = (r(v_1) + r(v_{\frac{n}{2}+1})) + (r(v_2) + r(v_{\frac{n}{2}+2}))$, we know that $r(v_1) + r(v_{\frac{n}{2}+1})$ and $r(v_2) + r(v_{\frac{n}{2}+2})$ are in different NO's. Without loss of generality, we assume $r(v_1) + r(v_{\frac{n}{2}+1})$ is in smaller NO and $r(v_2) + r(v_{\frac{n}{2}+2})$ is in larger NO.

Then for vertices $v_2, v_3, v_{\frac{n}{2}+2}, v_{\frac{n}{2}+3}$: Similarly we know that $r(v_2) + r(v_{\frac{n}{2}+2})$ and $r(v_3) + r(v_{\frac{n}{2}+3})$ are in different NO's, we already assumed $r(v_2) + r(v_{\frac{n}{2}+2})$ is in larger NO, so $r(v_3) + r(v_{\frac{n}{2}+3})$ is in smaller NO.

Repeatedly, we have that if i is odd then $r(v_i) + r(v_{\frac{n}{2}+i})$ is in smaller NO, if i is even then $r(v_i) + r(v_{\frac{n}{2}+i})$ is in larger NO. We take $i = \frac{n}{2} + 1$, because in this case $\frac{n}{2} + 1$ is

even, we have $r(v_{\frac{n}{2}+1}) + r(v_1) = r(v_{\frac{n}{2}+1}) + r(v_{n+1})$ is in larger NO, which contradicts our assumption $r(v_1) + r(v_{\frac{n}{2}+1})$ is in smaller NO.

Case iii. $4 \mid n$.

For vertices $v_1, v_2, v_{\frac{n}{2}}, v_{\frac{n}{2}+1}$: We know v_1v_2 and $v_{\frac{n}{2}}v_{\frac{n}{2}+1}$ are edges, so both $r(v_1) + r(v_2)$ and $r(v_{\frac{n}{2}}) + r(v_{\frac{n}{2}+1})$ are in YES. We know $v_1v_{\frac{n}{2}}$ and $v_2v_{\frac{n}{2}+1}$ are nonedges, so both $r(v_1) + r(v_{\frac{n}{2}})$ and $r(v_2) + r(v_{\frac{n}{2}+1})$ are in NO. And by the fact that $(r(v_1) + r(v_2)) + (r(v_{\frac{n}{2}}) + r(v_{\frac{n}{2}+1})) = (r(v_1) + r(v_{\frac{n}{2}})) + (r(v_2) + r(v_{\frac{n}{2}+1}))$, we know that $r(v_1) + r(v_{\frac{n}{2}})$ and $r(v_2) + r(v_{\frac{n}{2}+1})$ are in different NO's. Without loss of generality, we assume $r(v_1) + r(v_{\frac{n}{2}})$ is in smaller NO and $r(v_2) + r(v_{\frac{n}{2}+1})$ is in larger NO.

Then for vertices $v_2, v_3, v_{\frac{n}{2}+1}, v_{\frac{n}{2}+2}$: Similarly we know that $r(v_2) + r(v_{\frac{n}{2}+1})$ and $r(v_3) + r(v_{\frac{n}{2}+2})$ are in different NO's, we already assumed $r(v_2) + r(v_{\frac{n}{2}+1})$ is in larger NO, so $r(v_3) + r(v_{\frac{n}{2}+2})$ is in smaller NO.

Repeatedly, we have that if i is odd then $r(v_i) + r(v_{\frac{n}{2}+i-1})$ is in smaller NO, if i is even then $r(v_i) + r(v_{\frac{n}{2}+i-1})$ is in larger NO. So $r(v_{\frac{n}{2}+1}) + r(v_n)$ is in smaller NO, as $\frac{n}{2} + 1$ is odd. But then for vertices $v_{\frac{n}{2}}, v_{\frac{n}{2}+1}, v_n, v_1$: We know $v_{\frac{n}{2}}v_{\frac{n}{2}+1}$ and v_nv_1 are edges, so both $r(v_{\frac{n}{2}}) + r(v_{\frac{n}{2}+1})$ and $r(v_n) + r(v_1)$ are in YES. We know $v_{\frac{n}{2}}v_1$ and $v_{\frac{n}{2}+1}v_n$ are nonedges, and both $r(v_{\frac{n}{2}}) + r(v_1)$ and $r(v_{\frac{n}{2}+1}) + r(v_n)$ are in smaller NO. But then we will have $(r(v_{\frac{n}{2}}) + r(v_{\frac{n}{2}+1})) + (r(v_n) + r(v_1)) > (r(v_{\frac{n}{2}}) + r(v_1)) + (r(v_{\frac{n}{2}+1}) + r(v_n))$, contradiction.

In each of the three cases, we get a contradiction, so $\Theta(C_n)$ cannot be 2.

Then assume $\Theta(C_n) = 3$. So we have three thresholds $\theta_1, \theta_2, \theta_3$, and a $(\theta_1, \theta_2, \theta_3)$ -representation $r : V \rightarrow \mathbb{R}$. Three thresholds divide \mathbb{R} into four parts in the form

$$\text{NO} \mid \text{YES} \mid \text{NO} \mid \text{YES}.$$

The two NO parts will be called smaller NO and larger NO, the two YES parts will be called smaller YES and larger YES.

Firstly we note that there must be an edge rank sum in larger YES, as otherwise two thresholds would be enough, $\Theta(C_n) = 3$ would not hold. We may assume v_1v_2 is in larger YES.

We also make the following claim.

Claim. Assume v_iv_{i+1} and v_jv_{j+1} are disjoint edges in C_n , then $r(v_i) + r(v_{i+1})$ and $r(v_j) + r(v_{j+1})$ cannot both be in larger YES.

Proof of Claim. Assume by contradiction that both $r(v_i) + r(v_{i+1})$ and $r(v_j) + r(v_{j+1})$ are in larger YES. Then by the equation $(r(v_i) + r(v_{i+1})) + (r(v_j) + r(v_{j+1})) = (r(v_i) + r(v_j)) + (r(v_{i+1}) + r(v_{j+1}))$, we know that $\max\{r(v_i) + r(v_j), r(v_{i+1}) + r(v_{j+1})\} \geq \min\{r(v_i) + r(v_{i+1}), r(v_j) + r(v_{j+1})\}$, so at least one of $r(v_i) + r(v_j)$ and $r(v_{i+1}) + r(v_{j+1})$ is in larger YES. However we can see that both v_iv_j and $v_{i+1}v_{j+1}$ are nonedges, contradiction. \square

By this claim, we know that edge v_2v_3 and edge v_nv_1 cannot both have rank sums in larger YES, because they are disjoint edges. Moreover, any edge v_iv_{i+1} with $i \in [3, n-1]$ cannot have rank sum in larger YES, because v_1v_2 and v_iv_{i+1} are disjoint edges.

We have two cases:

Case 1. None of v_2v_3 and v_nv_1 has rank sum in larger YES.

We have that both $r(v_2) + r(v_3)$ and $r(v_n) + r(v_1)$ are in smaller YES, so by a similar argument as in the proof of $\Theta(C_n) \neq 2$, we know $r(v_1) + r(v_3)$ and $r(v_2) + r(v_n)$ are in different NO's. Without loss of generality, we may assume $r(v_1) + r(v_3)$ is in smaller NO.

Because v_2v_4 is a nonedge and $r(v_1) + r(v_2)$ is in larger YES, we have $r(v_2) + r(v_4) < r(v_1) + r(v_2)$, so $r(v_4) < r(v_1)$. Now, v_3v_4 is an edge, however $r(v_3) + r(v_4) < r(v_3) + r(v_1)$, which means $r(v_3) + r(v_4)$ is in smaller NO, contradiction.

Case 2. Exactly one of v_2v_3 and v_nv_1 has rank sum in larger YES.

Without loss of generality, we assume $r(v_2) + r(v_3)$ is in larger YES, $r(v_n) + r(v_1)$ is in smaller YES. We have that both $r(v_3) + r(v_4)$ and $r(v_n) + r(v_1)$ are in smaller YES, so by a similar argument as in the proof of $\Theta(C_n) \neq 2$, we know $r(v_3) + r(v_n)$ and $r(v_4) + r(v_1)$ are in different NO's. Without loss of generality, we may assume $r(v_3) + r(v_n)$ is in smaller NO.

Now we look at vertices v_2, v_3, v_{n-1}, v_n : We know v_2v_3 is an edge with $r(v_2) + r(v_3)$ in larger YES, $v_{n-1}v_n$ is an edge with $r(v_{n-1}) + r(v_n)$ in smaller YES. Now, because $r(v_3) + r(v_n)$ is in smaller NO, we have $r(v_3) + r(v_n) < r(v_{n-1}) + r(v_n)$. Because v_2v_{n-1} is a nonedge, we have $r(v_2) + r(v_{n-1})$ is in NO (larger or smaller), and we know $r(v_2) + r(v_3)$ is in larger YES, so $r(v_2) + r(v_{n-1}) < r(v_2) + r(v_3)$. But then we have $(r(v_3) + r(v_n)) + (r(v_2) + r(v_{n-1})) < (r(v_{n-1}) + r(v_n)) + (r(v_2) + r(v_3))$, contradiction.

Note that if $n = 5$, then v_4 and v_{n-1} are the same vertex, but this does not affect our argument.

In either case, we get a contradiction, so $\Theta(C_n)$ cannot be 3.

In summary, for $n \geq 5$, We have showed $\Theta(C_n) \geq 2$, $\Theta(C_n) \leq 4$, $\Theta(C_n) \neq 2$, and $\Theta(C_n) \neq 3$, thus $\Theta(C_n) = 4$. \square

3. REMARKS

Actually there is another way to see that $\Theta(C_n) \leq 4$, which is given by combining the following two propositions and the fact that we can get C_n by adding an edge to P_n .

Proposition 3.1 (Jamison and Sprague [6]). *Let n be a positive integer, let P_n be the path with n vertices, then $\Theta(P_n) \leq 2$.*

Proposition 3.2 (Jamison and Sprague [6]). *Let $G = (V, E)$ be a graph with $\Theta(G) = k$, and let $u, v \in V$ be distinct vertices.*

(a) *If uv is an edge, then the graph $G - uv$ obtained by deleting edge uv from E has threshold number at most $k + 2$.*

(b) *If uv is a nonedge, then the graph $G + uv$ obtained by adding edge uv to E has threshold number at most $k + 2$.*

The author adopted the constructive proof, as it helps readers understand how things work.

For other specific graphs, we can make the following observations.

Proposition 3.3. *The threshold number of a linear forest is at most 2, where a linear forest is a disjoint union of paths.*

Proposition 3.4. *The threshold number of a ladder is at most 2, where the ladder of length n , denoted by L_n , is defined to be $P_2 \square P_n$, the box product of the path of length 2 and the path of length n .*

Inspired by ladders, we propose the following problem.

Problem 3.5. *Determine the exact threshold numbers of grids, where the grid with m rows and n columns, denoted by $G_{m \times n}$, is defined to be $P_m \square P_n$, the box product of the path of length m and the path of length n .*

REFERENCES

- [1] CHEN, G., AND HAO, Y. Multithreshold multipartite graphs. *J. Graph Theory* 100, 4 (2022), 727–732.
- [2] CHVÁTAL, V., AND HAMMER, P. L. Aggregation of inequalities in integer programming. In *Studies in integer programming (Proc. Workshop, Bonn, 1975)*, vol. Vol. 1 of *Ann. Discrete Math.* North-Holland, Amsterdam-New York-Oxford, 1977, pp. 145–162.
- [3] GOLUBIC, M. C. *Algorithmic graph theory and perfect graphs*, second ed., vol. 57 of *Annals of Discrete Mathematics*. Elsevier Science B.V., Amsterdam, 2004. With a foreword by Claude Berge.
- [4] GOLUBIC, M. C., AND JAMISON, R. E. Rank-tolerance graph classes. *J. Graph Theory* 52, 4 (2006), 317–340.
- [5] GOLUBIC, M. C., AND TRENK, A. N. *Tolerance graphs*, vol. 89 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2004.
- [6] JAMISON, R. E., AND SPRAGUE, A. P. Multithreshold graphs. *J. Graph Theory* 94, 4 (2020), 518–530.
- [7] KITTIPASSORN, T., AND SUMALROJ, T. Multithreshold multipartite graphs with small parts. *Discrete Math.* 347, 7 (2024), Paper No. 113979, 15.
- [8] MAHADEV, N. V. R., AND PELED, U. N. *Threshold graphs and related topics*, vol. 56 of *Annals of Discrete Mathematics*. North-Holland Publishing Co., Amsterdam, 1995.

- [9] WANG, R. Threshold numbers of multipartite graphs with small parts. *arXiv e-prints* (2024), arXiv:2406.12063.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF MEMPHIS, MEMPHIS, TN 38152,
USA

Email address: `runze.w@hotmail.com`