

TRANSVERSAL HAMILTON PATHS AND CYCLES

YANGYANG CHENG, WANTING SUN, GUANGHUI WANG, AND LAN WEI

ABSTRACT. Given a collection $\mathcal{G} = \{G_1, G_2, \dots, G_m\}$ of graphs on the common vertex set V of size n , an m -edge graph H on the same vertex set V is *transversal* in \mathcal{G} if there exists a bijection $\varphi : E(H) \rightarrow [m]$ such that $e \in E(G_{\varphi(e)})$ for all $e \in E(H)$. Denote $\delta(\mathcal{G}) := \min\{\delta(G_i) : i \in [m]\}$. In this paper, we first establish a minimum degree condition for the existence of transversal Hamilton paths in \mathcal{G} : if $n = m+1$ and $\delta(\mathcal{G}) \geq \frac{n-1}{2}$, then \mathcal{G} contains a transversal Hamilton path. This solves a problem proposed by [Li, Li and Li, J. Graph Theory, 2023]. As a continuation of the transversal version of Dirac's theorem [Joos and Kim, Bull. Lond. Math. Soc., 2020] and the stability result for transversal Hamilton cycles [Cheng and Staden, arXiv:2403.09913v1], our second result characterizes all graph collections with minimum degree at least $\frac{n}{2} - 1$ and without transversal Hamilton cycles. We obtain an analogous result for transversal Hamilton paths. The proof is a combination of the stability result for transversal Hamilton paths or cycles, transversal blow-up lemma, along with some structural analysis.

1. INTRODUCTION

Let $G = (V(G), E(G))$ be a graph with vertex set $V(G)$ and edge set $E(G)$. For a vertex v of G , denote by $N_G(v)$ the neighborhood of v and let $d_G(v) = |N_G(v)|$ be the degree of v . The subscripts are omitted when G is clear from the context. Let $\delta(G)$ be the minimum degree of G . Denote by P_n , C_n , K_n and $K_{t,n-t}$ the path, cycle, complete graph and complete bipartite graph (with parts of size t and $n-t$) on n vertices, respectively.

Let $\mathcal{G} = \{G_1, G_2, \dots, G_m\}$ be a collection of not necessarily distinct graphs with common vertex set V . We often think of each G_i having the color i . Let H be a graph on the vertex set V . We say H is *rainbow* in \mathcal{G} if there exists an injection $\varphi : E(H) \rightarrow [m]$ such that $e \in E(G_{\varphi(e)})$ for all $e \in E(H)$. In addition, if $|E(H)| = m$, then φ is a bijection and we say H is *transversal* in \mathcal{G} . Let $\delta(\mathcal{G}) = \min\{\delta(G_i) : i \in [m]\}$ be the minimum degree of \mathcal{G} . Transversal often appears in infinitary combinatorics under several similar definitions (see, [1, 24]), and it is also extensively studied in the context of Latin squares (see [45] for a survey).

The following general question was proposed by Joos and Kim in [32].

Question 1.1. *Let H be a graph with m edges, \mathbf{G} be a family of graphs, and $\mathcal{G} = \{G_1, G_2, \dots, G_m\}$ be a collection of graphs with common vertex set V such that $G_i \in \mathbf{G}$ for all $i \in [m]$. Which properties imposed on \mathbf{G} guarantee a transversal copy of H ?*

By taking $G_1 = G_2 = \dots = G_m$, we need to study properties for \mathbf{G} such that each graph in \mathbf{G} contains H as a subgraph. However, this alone is not always sufficient. For example, Aharoni, DeVos, de la Maza, Montejano and Šámal [4] proved that if $\mathcal{G} = \{G_1, G_2, G_3\}$ is a collection of graphs on a common vertex set of size n and $|E(G_i)| > (\frac{26-2\sqrt{7}}{81})n^2$ for $i \in [3]$, then \mathcal{G} contains a rainbow triangle. Moreover, the constant $\frac{26-2\sqrt{7}}{81}$ is optimal. However, Mantel's theorem states that any n -vertex graph with more than $\lfloor \frac{n^2}{4} \rfloor$ edges must contain a triangle.

Y. Cheng is supported by the PhD studentship of ERC Advanced Grant (883810). W. Sun is supported by the China Postdoctoral Science Foundation (2023M742092). G. Wang and L. Wei are supported by the Natural Science Foundation of China (12231018) and Young Taishan Scholars Program of Shandong Province.

In this paper, we investigate the minimum degree condition for a graph collection to guarantee the existence of transversal Hamilton paths or cycles. Hamiltonicity of graphs is one of the fundamental problems in extremal graph theory and structural graph theory. In 1952, Dirac [23] proved that an n -vertex graph contains a Hamilton cycle if its minimum degree is no less than $\frac{n}{2}$. Ore [41] relaxed this condition by considering the sum of degrees of two non-adjacent vertices. In general, there are many other sufficient conditions that guarantee the Hamiltonicity of a graph, such as Pósa's condition [42], Bondy's condition [7] and so on. For more problems and results about Hamiltonicity of graphs, we refer the reader to [9, 28, 29, 39, 43].

In 2020, Aharoni [4] conjectured that Dirac's theorem can be extended to a transversal version.

Conjecture 1.2 ([4]). *Suppose $\mathcal{G} = \{G_1, \dots, G_n\}$ is a collection of graphs with the same vertex set V of size n . If $\delta(\mathcal{G}) \geq \frac{n}{2}$, then \mathcal{G} contains a transversal Hamilton cycle.*

Cheng, Wang and Zhao [21] solved this conjecture asymptotically, and it was completely confirmed by Joos and Kim [32]. Bradshaw, Halasz and Stacho [13] extended the result of Conjecture 1.2 by showing any such graph collection has at least $(\frac{cn}{e})^{cn}$ transversal Hamilton cycles for some constant $c > \frac{1}{68}$. Anastos and Chakraborti [6] improved $(\frac{cn}{e})^{cn}$ to $(Cn)^{2n}$ for some constant $C > 0$. Bradshaw [12] studied the Hamiltonicity in bipartite graph collections. Bowtell, Morris, Pehova and Staden [11] showed that \mathcal{G} contains every Hamilton cycle pattern if $\delta(\mathcal{G}) \geq (1 + o(1))\frac{n}{2}$. The research of graph collections has also been generalized to random graph collections. Ferber, Han and Mao [25] provided a transversal version of the Dirac's theorem in random graph collections.

In addition to Dirac's theorem, many other classical results of extremal graph theory have been generalized. Cheng, Han, Wang and Wang [17] studied the minimum degree condition for the existence of transversal K_t -factors in (hyper)graph collections, which is an asymptotical version of the rainbow Hajnal-Szemerdi theorem [31]. Montgomery, Myesser and Pehova [40] gave asymptotically tight transversal versions of Hajnal-Szemerdi theorem [31] and Kühn-Osthus theorem [34] on factors, and a transversal generalisation of the theorem of Komlós, Sárközy and Szemerédi [33] on spanning trees. Gupta, Hamann, Müyesser, Parczyk and Sgueglia [30] gave a general approach to transversal versions of several classical Dirac-type results. Cheng and Staden [19] established the transversal blow-up lemma, which is an effective tool for transversal embedding problems. Chakraborti, Im, Kim and Liu [15] extended the bandwidth theorem [10] to graph collections. More relevant developments on this topic, we refer the reader to [2, 3, 5, 18, 35, 37, 38, 44].

Li, Li and Li [36] established a sufficient condition for the existence of a rainbow Hamilton path in $\mathcal{G} = \{G_1, \dots, G_n\}$. However, the edges of a rainbow Hamilton path only come from $n - 1$ graphs in \mathcal{G} . Hence, they proposed the following problem: whether a collection of $n - 1$ graphs is sufficient to guarantee the existence of transversal Hamilton paths. Our first result solves the above problem and prove the following result.

Theorem 1.3. *Let $\mathcal{G} = \{G_1, \dots, G_{n-1}\}$ be a collection of graphs on a common vertex set V of size n . If $\delta(\mathcal{G}) \geq \frac{n-1}{2}$, then \mathcal{G} has a transversal Hamilton path.*

The proof of Theorem 1.3 is inspired by Joos and Kim [32], which proved the transversal version of Dirac's theorem by constructing auxiliary digraphs. The difficulty of our proof is that we can only find a rainbow cycle of length at least $n - 2$ when $\delta(\mathcal{G}) \geq \frac{n-1}{2}$. If the longest rainbow

cycle inside \mathcal{G} has length $n-2$, then we need to analyze the relation between two vertices outside the cycle by using a series of rotations and more structural analysis.

For graphs G and H , denote by $G \cup H$ the disjoint union of G and H . We use $G \vee H$ to denote the graph obtained from $G \cup H$ by adding an edge between each vertex of G and each vertex of H . A k -*matching* in a graph is a set consisting of k edges with no shared vertices. For any vertex subsets $X, Y \subseteq V(G)$, let $G[X]$ be the induced subgraph of G on X . Let $G[X, Y]$ be the subgraph of G with vertex set $X \cup Y$ and edge set $\{xy : x \in X, y \in Y \text{ and } xy \in E(G)\}$. Denote by $G - xy$ the graph obtained from G by deleting the edge xy , where $xy \in E(G)$. Let $\mathcal{G} = \{G_1, G_2, \dots, G_m\}$ be collection of graphs on a common vertex set. Then define $\mathcal{G}[X] := \{G_i[X] : i \in [m]\}$ and $\mathcal{G}[X, Y] := \{G_i[X, Y] : i \in [m]\}$. If $E(G) = \emptyset$, then we simply write $G = \emptyset$. Similarly, if there is no edge in G_i for all $i \in [m]$, then we write $\mathcal{G} = \emptyset$.

Let $0 < \kappa < 1$ be a positive constant. Given two graph collections $\mathcal{G} = \{G_1, \dots, G_m\}$ and $\mathcal{H} = \{H_1, \dots, H_m\}$ defined on the vertex sets V_1 and V_2 respectively, where $|V_1| = |V_2| = n$ and $(1 - \kappa)n \leq m \leq (1 + \kappa)n$, we say \mathcal{G} is κ -close to \mathcal{H} if by adding or deleting at most κn^3 edges of \mathcal{G} we can obtain a copy of \mathcal{H} . A graph collection $\mathcal{G}' = \{G'_1, \dots, G'_m\}$ is a *spanning collection* of \mathcal{G} if G'_i is a spanning subgraph of G_i for each $i \in [m]$. For any vertex set V of size n , we say $A \cup B$ is an *equitable partition* of V if $|A| + |B| = n$ and $||A| - |B|| \leq 1$.

Definition 1.4 (\mathcal{H}_s^t , half-split graph collection). *Given integers $s, t \geq 0$, let \mathcal{H}_s^t be the graph collection on a common vertex set of size n obtained by taking s copies of $K_{\lfloor \frac{n}{2} \rfloor} \cup K_{\lfloor \frac{n}{2} \rfloor}$ and t copies of $K_{\lfloor \frac{n}{2} \rfloor, \lfloor \frac{n}{2} \rfloor}$, where they are defined on the same equitable partition.*

We say that a graph collection $\mathcal{G} = \{G_1, \dots, G_m\}$ on a common vertex set V of size n is half-split if there is a subset $A \subseteq V$ with $|A| = \lfloor \frac{n}{2} \rfloor$ such that $\mathcal{G}[A] = \emptyset$ and all graphs in $\mathcal{G}[A, V \setminus A]$ are complete bipartite graphs.

Cheng and Staden [20] characterized the stability result for transversal Hamilton paths or cycles when the minimum degree of the graph collection is slightly lower than $\frac{n}{2}$.

Theorem 1.5 ([20]). *For all $\kappa > 0$, there exist $\mu > 0$ and n_0 such that the following holds for all integers $n \geq n_0$. Let $\mathcal{G} = \{G_1, \dots, G_m\}$ be a collection of graphs with the common vertex set V of size n and $\delta(\mathcal{G}) \geq (\frac{1}{2} - \mu)n$.*

- (i) *If $m = n$ and \mathcal{G} contains no transversal Hamilton cycles, then either \mathcal{G} is κ -close to \mathcal{H}_s^t for some $s \in [n]$ with $t = n - s$ is odd, or \mathcal{G} is κ -close to a half-split graph collection.*
- (ii) *If $m = n - 1$ and \mathcal{G} contains no transversal Hamilton paths, then either \mathcal{G} is κ -close to \mathcal{H}_{n-1}^0 , or \mathcal{G} is κ -close to a half-split graph collection.*

As a continuation of the transversal version of Dirac's theorem [32] and the stability result for transversal Hamilton cycles (i.e., Theorem 1.5 (i)), we characterize all graph collections with minimum degree at least $\frac{n}{2} - 1$ and without transversal Hamilton cycles. Note that in a graph collection \mathcal{G} , the order of colors does not affect the overall structure of \mathcal{G} . Thus, when we say G_i (for example, G_1) is a graph in \mathcal{G} , it is usually arbitrary.

Theorem 1.6. *For sufficiently large n , let \mathcal{C} be a set of n colors, and let $\mathcal{G} = \{G_i : i \in \mathcal{C}\}$ be a collection of graphs with common vertex set V of size n and $\delta(\mathcal{G}) \geq \frac{n}{2} - 1$. Assume that \mathcal{G} contains no transversal Hamilton cycles.*

- (i) *If n is odd, then one of the following holds:*
 - (a) *\mathcal{G} is the half-split graph collection,*

- (b) there exists a fixed vertex u with $d_{G_i}(u) = n - 1$ and an equitable partition $A \cup B$ of $V \setminus \{u\}$ such that $G_i[V \setminus \{u\}] = G_i[A] \cup G_i[B] = 2K_{\frac{n-1}{2}}$ for all $i \in \mathcal{C}$.
- (ii) If n is even, then one of the following holds:
- (a) \mathcal{G} is a spanning collection of \mathcal{H}_{n-t}^t for some odd integer $t \in [n]$,
- (b) there exists a partition $A \cup B$ of V with $|A| = \frac{n}{2} - 1$ such that either $G_i[B] = \emptyset$ for all but at most one $i \in \mathcal{C}$, or $E(G_i[B]) \subseteq \{uv\}$ for fixed $u, v \in B$ and all $i \in \mathcal{C}$,
- (c) there exists an equitable partition $A \cup B$ of V such that $\mathcal{G}[A, B]$ contains no rainbow 2-matching, that is, one of the following holds:
- $G_i = G_i[A] \cup G_i[B] = 2K_{\frac{n}{2}}$ for all but at most one $i \in \mathcal{C}$,
 - there is a vertex $u \in A$ such that $E(G_i[A, B]) = E(G_i[\{u\}, B])$ for all $i \in \mathcal{C}$,
 - there are vertices $u \in A$ and $v \in B$ such that $E(G_1[A, B]) \subseteq \{uw : w \in B\} \cup \{wv : w \in A\}$ and $E(G_i[A, B]) \subseteq \{uv\}$ for all $i \in \mathcal{C} \setminus \{1\}$ (see Figure 1 (a)),
 - there are vertices $u, u' \in A$ and $v, v' \in B$ such that $E(G_1[A, B]) = \{uv, u'v'\}$, $E(G_2[A, B]) = \{uv', u'v\}$ and $G_j = G_j[A] \cup G_j[B] = 2K_{\frac{n}{2}}$ for all $j \in \mathcal{C} \setminus \{2\}$ (see Figure 1 (b)).




FIGURE 1. Extremal graph collections in Theorem 1.6. Here the dotted line means that the edge may not exist, and the triangle stands for a complete graph.

Remark 1.7. In fact, in Theorem 1.6 (ii) (b)-(c), based on the local structure of \mathcal{G} and the minimum degree condition, the global structure of \mathcal{G} can be deduced immediately. For example, $|A| = \frac{n}{2} - 1$ and $G_i[B] = \emptyset$ imply that $G_i[A, B] = K_{\frac{n}{2}-1, \frac{n}{2}+1}$ and $G_i[A]$ can be any graph.

For an n -vertex graph G with $\delta(G) \geq \frac{n}{2} - 1$, Büyükçolak-Gözüpek-Özkan-Sibel-Shalom [14] and Fu-Gao-Wang-Yang [26] characterized the structure of G without Hamilton cycles if $n (\geq 3)$ is odd and $n (\geq 22)$ is even, respectively. In fact, by applying Theorem 1.6 with $G_1 = G_2 = \dots = G_n = G$, we can deduce their results when n is sufficiently large.

Corollary 1.8 ([14, 26]). *For sufficiently large n , let G be an n -vertex graph with vertex set V . Assume that G contains no Hamilton cycles and $\delta(G) \geq \frac{n}{2} - 1$.*

- (i) *If n is odd, then either $G = K_1 \vee (K_{\frac{n-1}{2}} \cup K_{\frac{n-1}{2}})$ or G is a spanning subgraph of $\frac{n+1}{2}K_1 \vee K_{\frac{n-1}{2}}$.*
- (ii) *If n is even, then G is a spanning subgraph of $K_1 \vee (K_{\frac{n}{2}-1} \cup K_{\frac{n}{2}})$ or $(K_2 \cup (\frac{n}{2} - 1)K_1) \vee K_{\frac{n}{2}-1}$.*

Recall that Theorem 1.3 gives a sufficient condition for the existence of transversal Hamilton paths in a collection of $n - 1$ graphs. In contrast to the Hamilton cycle case, our next result

characterizes all collections of $n - 1$ graphs with minimum degree at least $\frac{n-3}{2}$ and containing no transversal Hamilton paths.

Theorem 1.9. *For sufficiently large n , let \mathcal{C} be a set of $n - 1$ colors, and let $\mathcal{G} = \{G_i : i \in \mathcal{C}\}$ be a collection of graphs with the common vertex set V of size n . Assume that $\delta(\mathcal{G}) \geq \frac{n-3}{2}$ and \mathcal{G} contains no transversal Hamilton paths.*

(i) *If n is even, then one of the following holds:*

- \mathcal{G} is \mathcal{H}_{n-1}^0 ,
- there exists a partition $A \cup B$ of V with $|A| = \frac{n}{2} + 1$ such that $G_i[A] = \emptyset$ for all $i \in \mathcal{C}$.

(ii) *If n is odd, then one of the following holds:*

- \mathcal{G} is a spanning collection of \mathcal{H}_{n-1}^0 ,
- there exists a partition $A \cup B$ of V with $|A| = \frac{n+3}{2}$ such that either $G_i[A] = \emptyset$ for all $i \in \mathcal{C}$, or $E(G_i[A]) \subseteq \{uv\}$ for fixed vertices $u, v \in A$ and all $i \in \mathcal{C}$.

Notice that given a collection \mathcal{G} of $n - 1$ graphs on a common vertex set of size n , the graph collection obtained from adding a complete graph to \mathcal{G} has a transversal Hamilton cycle if and only if \mathcal{G} has a transversal Hamilton path. Thus, Theorem 1.9 (i) is an immediate consequence of Theorem 1.6 (ii). For Theorem 1.9 (ii), it can be proved by using similar arguments as Theorem 1.6, so we omit its proof. Note that we just need to find a transversal Hamilton path inside \mathcal{G} , that is, we don't need to connect its two endpoints, hence the proof of Theorem 1.9 (ii) is much simpler.

2. NOTATION AND ORGANIZATION

2.1. Notation. First of all, we state that all terminology and notation on graph theory not defined in this paper are the same as those used in the textbook [22].

For a vertex subset $U, W \subseteq V(G)$ and a vertex $v \in V(G)$, let $N_G(v, U) = N_G(v) \cap U$ and $d_G(v, U) = |N_G(v, U)|$. Let $N_G(U, W) = \cup_{u \in U} N_G(u, W)$. Denote by S_r the star with r vertices, where the vertex with degree $r - 1$ is the center of S_r . Let $\mathcal{G} = \{G_i : i \in \mathcal{C}\}$ be a graph collection and H be a rainbow subgraph inside \mathcal{G} . Denote by $col(H)$ the set of colors appearing in H . Let $P_t = v_1 v_2 \dots v_t$ be a rainbow path inside \mathcal{G} . For $1 \leq i < j \leq t$, denote by $v_i P v_j := v_i v_{i+1} \dots v_j$ a rainbow subpath of P , and the color of each edge inherits its color on P . We say P is *maximal* in \mathcal{G} if it is not a proper rainbow subpath of any other rainbow paths inside \mathcal{G} . Two rainbow paths P and Q are said to be *disjoint* if $V(P) \cap V(Q) = \emptyset$ and $col(P) \cap col(Q) = \emptyset$.

Let D be a digraph. We denote its vertex set by $V(D)$ and its arc set by $A(D)$. For vertices u and v in D , the arc from u to v is denoted by \vec{uv} . Given $v \in V(D)$, let $N_D^+(v)$ and $N_D^-(v)$ be the out-neighborhood and in-neighborhood of v in D , respectively. Denote by $d_D^+(v)$ and $d_D^-(v)$ the out-degree and in-degree of v , respectively.

For a positive integer n , we write $[n] := \{1, 2, \dots, n\}$ and $[a, b] := \{a, a + 1, \dots, b\}$ for two positive integers $a < b$. For any two constants $\alpha, \beta \in (0, 1)$, we write $\alpha \ll \beta$ if there exists a function $f = f(\beta)$ such that the subsequent arguments hold for all $0 < \alpha \leq f$.

2.2. Organization. In the remainder of this section, we give a sketch of the proof of Theorem 1.6. In Section 3, we prove Theorem 1.3, which establishes a minimum degree condition to guarantee the existence of transversal Hamilton paths. In Section 4, we give the proof of Theorem 1.6. The proof is split into three steps: constructing short disjoint rainbow paths to cover bad

vertices and bad colors; using the connecting tool (Claim 4.5) to connect those rainbow paths; applying transversal blow-up lemma to construct long rainbow paths to cover all left vertices and colors. In order to use the transversal blow-up lemma, we introduce Theorem 4.3 to balance the number of vertices in two parts, whose proof is presented in Section 5. Finally, some concluding remarks are given in Section 6.

2.3. Sketch of the proof of Theorem 1.6. Assume $0 < \frac{1}{n} \ll \delta \ll \eta \ll 1$. Let $\mathcal{G} = \{G_1, G_2, \dots, G_n\}$ be a collection of graphs with common vertex set V of size n . Assume that $\delta(\mathcal{G}) \geq \frac{n}{2} - 1$ and \mathcal{G} contains no transversal Hamilton cycles. Denote $\mathcal{C}_1 := \{i \in [n] : G_i \text{ is close to } K_{\lceil \frac{n}{2} \rceil} \cup K_{\lfloor \frac{n}{2} \rfloor}\}$, $\mathcal{C}_2 := \{i \in [n] : G_i \text{ is close to } K_{\lceil \frac{n}{2} \rceil, \lfloor \frac{n}{2} \rfloor}\}$ and $\mathcal{C}_{bad} := [n] \setminus (\mathcal{C}_1 \cup \mathcal{C}_2)$. By [20, Lemma 5.3] (see also Lemma 4.2 in Section 4), we know that $|\mathcal{C}_1| + |\mathcal{C}_2| \geq (1 - 3\delta)n$. In view of Definition 4.1, we can fix a characteristic partition (A_i, B_i, C_i) for each graph G_i with $i \in \mathcal{C}_1 \cup \mathcal{C}_2$. By swapping the labels of A_i and B_i , we have $|A_1 \Delta A_i|, |B_1 \Delta B_i| < \delta n$ for every $i \in (\mathcal{C}_1 \cup \mathcal{C}_2) \setminus \{1\}$.

Based on the sizes of $|\mathcal{C}_1|$ and $|\mathcal{C}_2|$, we distinguish our proof into three cases: $|\mathcal{C}_1| < \eta n$, $|\mathcal{C}_2| < \eta n$, $|\mathcal{C}_1| \geq \eta n$ and $|\mathcal{C}_2| \geq \eta n$. We will take the first case as an example to illustrate the general idea of our proof.

Firstly, expand $A_1 \cup B_1$ to an equitable partition $A \cup B$ of V . For a vertex $v \in A$, there are three bad cases: v lies in almost all B_i for $i \in \mathcal{C}_1 \cup \mathcal{C}_2$; v lies in almost all C_i for $i \in \mathcal{C}_1 \cup \mathcal{C}_2$; and v lies in both A_i and B_i for many $i \in \mathcal{C}_1 \cup \mathcal{C}_2$. Then move vertices in the first case from A to B , and do the same operation for such vertices in B . Denote the set of vertices in the other two bad cases by V_{bad} . Now, the number of vertices in A and B may be unbalanced, but their difference is still small (no more than $9\delta n$). In order to use the transversal blow-up lemma (see [19] or Claim 4.4 in Section 4) to find long rainbow paths inside \mathcal{G} , the following four steps are needed.

Step 1. Balance the number of vertices in A and B . If there exists a set of disjoint rainbow paths in $\mathcal{G}[B]$ such that after deleting their vertices, the number of vertices in A and B are balanced, then we are done. Otherwise, we will show that \mathcal{G} must be a graph collection described in Theorem 1.6.

Step 2. Deal with vertices in V_{bad} . By using colors in \mathcal{C}_2 , we are to find a series of disjoint rainbow P_3 such that all centers of them are all vertices in V_{bad} and the endpoints of them are unused vertices in $(A \cup B) \setminus V_{bad}$.

Step 3. Deal with colors in \mathcal{C}_{bad} . Choose a maximal rainbow matching M by using colors in \mathcal{C}_{bad} and avoiding vertices used in Steps 1-2. It is routine to check that for each $j \in \mathcal{C}_{bad} \setminus \text{col}(M)$, the graph G_j is close to $K_{\lfloor \frac{n}{2} \rfloor} \cup K_{\lceil \frac{n}{2} \rceil}$.

Step 4. Deal with colors in $\mathcal{C}_1 \cup (\mathcal{C}_{bad} \setminus \text{col}(M))$. We greedily select two rainbow paths with colors $\mathcal{C}_1 \cup (\mathcal{C}_{bad} \setminus \text{col}(M))$ and avoiding the used vertices in the above three steps. The lengths of those two rainbow paths are determined by the parity of $|\mathcal{C}_1 \cup (\mathcal{C}_{bad} \setminus \text{col}(M))|$.

Based on the minimum degree condition and the characteristic partition of extremal graphs, one may use colors in \mathcal{C}_2 to connect all rainbow paths obtained in the above four steps into a single short rainbow path, say P , such that $V_{bad} \subseteq V(P)$, $\mathcal{C}_1 \cup \mathcal{C}_{bad} \subseteq \text{col}(P)$, $|A \setminus V(P)|$ and $|B \setminus V(P)|$ are almost equal. By applying the transversal blow-up lemma and some structural analysis, the exact structure of graphs in \mathcal{G} can be deduced.

3. PROOF OF THEOREM 1.3

In this section, we prove Theorem 1.3. The following lemma allows us to find a rainbow path of length at least $n - 3$ in \mathcal{G} .

Lemma 3.1 ([36]). *Suppose $\mathcal{G} = \{G_1, \dots, G_n\}$ is a collection of graphs with common vertex set V of size n , and $\delta(\mathcal{G}) \geq \frac{n-1}{2}$. Then one of the following holds:*

- \mathcal{G} has a rainbow cycle of length at least $n - 1$,
- n is odd and \mathcal{G} consists of n copies of $K_1 \vee 2K_{\frac{n-1}{2}}$.

Proof of Theorem 1.3. We prove it by contradiction. Suppose that \mathcal{G} does not contain transversal Hamilton paths. Let $V = \{x_1, \dots, x_n\}$. Firstly, we give the following claim.

Claim 3.2. \mathcal{G} has a rainbow cycle of length at least $n - 2$.

Proof of Claim 3.2. Suppose that \mathcal{G} contains no rainbow cycle of length at least $n - 2$. By Lemma 3.1, we know $\{G_1, \dots, G_{n-1}, K_n\}$ contains a rainbow cycle of length at least $n - 1$. Thus, \mathcal{G} contains a rainbow path of length at least $n - 3$. Let $P = x_1x_2 \dots x_{n-2}$ be such a rainbow path, where $x_i x_{i+1} \in E(G_i)$ for $i \in [n - 3]$. It is obvious that $x_1x_{n-2} \notin E(G_{n-1}) \cup E(G_{n-2})$.

We claim that $|N_{G_{n-2}}(x_1) \cap \{x_{n-1}, x_n\}| + |N_{G_{n-1}}(x_{n-2}) \cap \{x_{n-1}, x_n\}| \leq 2$. Otherwise, there exists $x_j \in N_{G_{n-2}}(x_1) \cap N_{G_{n-1}}(x_{n-2})$ for some $j \in \{n - 1, n\}$. It follows that $x_j x_1 \dots x_{n-2} x_j$ forms a rainbow cycle of length $n - 1$ in \mathcal{G} , a contradiction. Let

$$A = \{j \in [n - 4] : x_1 x_{j+1} \in E(G_{n-2})\}, \quad B = \{j \in [2, n - 3] : x_j x_{n-2} \in E(G_{n-1})\}.$$

If $A \cap B \neq \emptyset$, then choose $j_1 \in A \cap B$ and $x_1 x_{j_1+1} \dots x_{n-2} x_{j_1} x_{j_1-1} \dots x_1$ is a rainbow cycle of length $n - 2$ in \mathcal{G} , a contradiction. Thus, $A \cap B = \emptyset$. Notice that $A \cup B \subseteq [n - 3]$. Hence $|A| + |B| \leq n - 3$. Recall that $x_1 x_{n-2} \notin E(G_{n-1}) \cup E(G_{n-2})$. Then, $|A| \geq \frac{n-1}{2} - |N_{G_{n-2}}(x_1) \cap \{x_{n-1}, x_n\}|$ and $|B| \geq \frac{n-1}{2} - |N_{G_{n-1}}(x_{n-2}) \cap \{x_{n-1}, x_n\}|$. Therefore, $|N_{G_{n-2}}(x_1) \cap \{x_{n-1}, x_n\}| + |N_{G_{n-1}}(x_{n-2}) \cap \{x_{n-1}, x_n\}| = 2$ and $|A| + |B| = n - 3$. By symmetry, one of the following holds:

- (A1) $N_{G_{n-2}}(x_1) \cap \{x_{n-1}, x_n\} = \{x_{n-1}\}$ and $N_{G_{n-1}}(x_{n-2}) \cap \{x_{n-1}, x_n\} = \{x_n\}$,
- (A2) $N_{G_{n-2}}(x_1) \cap \{x_{n-1}, x_n\} = \{x_{n-1}\}$ and $N_{G_{n-1}}(x_{n-2}) \cap \{x_{n-1}, x_n\} = \{x_{n-1}\}$,
- (A3) $N_{G_{n-2}}(x_1) \cap \{x_{n-1}, x_n\} = \{x_{n-1}, x_n\}$ and $N_{G_{n-1}}(x_{n-2}) \cap \{x_{n-1}, x_n\} = \emptyset$.

If (A1) holds, then \mathcal{G} contains a transversal Hamilton path $x_{n-1}x_1 \dots x_{n-2}x_n$, a contradiction. If (A2) holds, then \mathcal{G} contains a rainbow $(n - 1)$ -cycle $x_{n-1}x_1 \dots x_{n-2}x_{n-1}$, which is also a contradiction. Hence (A3) holds.

It is easy to see that $x_{n-3}x_n \notin E(G_{n-3}) \cup E(G_{n-1})$. Otherwise, $x_n x_1 \dots x_{n-3} x_n$ is a rainbow cycle of length $n - 2$ in \mathcal{G} , a contradiction. For convenience, denote $x^1 := x_n$, $x^i := x_{i-1}$ for $i \in [2, n - 2]$, $x^{n-1} := x_{n-1}$ and $x^n := x_{n-2}$. Then $P' := x^1 x^2 \dots x^{n-2}$ is a rainbow path inside \mathcal{G} with $n - 3, n - 1 \notin \text{col}(P')$. Let

$$A' = \{j \in [n - 4] : x^1 x^{j+1} \in E(G_{n-1})\}, \quad B' = \{j \in [2, n - 3] : x^j x^{n-2} \in E(G_{n-3})\}.$$

Notice that $x^1 x^{n-1} \notin E(G_{n-1})$, otherwise $x^{n-1} x^1 P' x^{n-2} x^n$ forms a transversal Hamilton path, a contradiction. It follows from (A3) that $x^1 x^n \notin E(G_{n-1})$ and $x^2 x^{n-1} \in E(G_{n-2})$. Then $x^{n-1} x^{n-2} \notin E(G_{n-3})$. Together with $x_{n-3}x_n \notin E(G_{n-3}) \cup E(G_{n-1})$, we have $|A'| + |B'| \geq \frac{n-1}{2} + \frac{n-1}{2} - 1 = n - 2$. Since $A' \cup B' \subseteq [n - 3]$, one has $A' \cap B' \neq \emptyset$ and let $j_2 \in A' \cap B'$. Hence, $x^1 x^{j_2+1} P' x^{n-2} x^{j_2} P' x^1$ is a rainbow cycle of length $n - 2$ in \mathcal{G} , a contradiction.

This completes the proof of Claim 3.2. □

Based on Claim 3.2, we distinguish our proof into the following two cases.

Case 1. \mathcal{G} has a rainbow cycle of length $n - 1$.

Let $C = x_1x_2 \dots x_{n-1}x_1$ be a rainbow cycle inside \mathcal{G} with $x_ix_{i+1} \in E(G_i)$ for $i \in [n-1]$, where we identify x_n with x_1 . Let y be the vertex in $V \setminus V(C)$. Since \mathcal{G} has no transversal Hamilton path, one has $yx_i, yx_{i+1} \notin E(G_i)$ for $i \in [n-1]$.

We construct the following auxiliary digraph D with vertex set V and arc set

$$A(D) = \bigcup_{i \in [n-1]} \{\overrightarrow{x_i z} : x_i z \in E(G_i) \text{ and } z \neq x_{i+1}\}.$$

By the definition of D , we have $d_D^+(y) = 0$. Furthermore, $d_D^-(y) = 0$. Otherwise, there exists $x_i \in N_D^-(y)$ for some $i \in [n-1]$. Then, $yx_i \dots x_1x_{n-1} \dots x_{i+1}$ is a transversal Hamilton path in \mathcal{G} , a contradiction. It follows from $\delta(\mathcal{G}) \geq \frac{n-1}{2}$ that $d_D^+(x_i) \geq \frac{n-1}{2} - 1$ for all $i \in [n-1]$. Therefore,

$$|A(D)| \geq (n-1)\left(\frac{n-1}{2} - 1\right) = (n-1)\frac{n-3}{2}. \quad (1)$$

Without loss of generality, assume that $d_D^-(x_1) = \max\{d_D^-(x_i) : i \in [n-1]\}$. Then we claim that $d_D^-(x_1) \geq \frac{n-3}{2}$. Otherwise, $d_D^-(x_1) \leq \frac{n-5}{2}$. This implies that $|A(D)| = \sum_{i \in [n-1]} d_D^-(x_i) \leq (n-1)\frac{n-5}{2} < (n-1)\frac{n-3}{2}$, which contradicts (1). Let

$$I_1^1 = \{i \in [2, n-2] : x_i \in N_D^-(x_1)\}, \quad I_2^1 = \{i \in [2, n-2] : x_{i+1}y \in E(G_1)\}.$$

Recall that $x_1y, x_2y \notin E(G_1)$. Then $|I_1^1| + |I_2^1| \geq \frac{n-3}{2} + \frac{n-1}{2} = n-2$. But $I_1^1 \cup I_2^1 \subseteq [2, n-2]$. Hence $I_1^1 \cap I_2^1 \neq \emptyset$. Choose $i_1 \in I_1^1 \cap I_2^1$. Therefore, $x_2 \dots x_{i_1}x_1x_{n-1} \dots x_{i_1+1}y$ is a transversal Hamilton path in \mathcal{G} , a contradiction.

Case 2. \mathcal{G} has no rainbow cycle of length $n - 1$.

By Claim 3.2, there exists a rainbow cycle of length $n - 2$ in \mathcal{G} , say $C = x_1x_2 \dots x_{n-2}x_1$ with $x_ix_{i+1} \in E(G_i)$ for $i \in [n-2]$, where we identify x_{n-1} with x_1 . Let $\{y, y'\} := V \setminus V(C)$. Now, we utilize the following auxiliary digraph D with vertex set V and arc set

$$A(D) = \bigcup_{i \in [n-2]} \{\overrightarrow{x_i z} : x_i z \in E(G_i) \text{ and } z \neq x_{i+1}\}.$$

Clearly, $d_D^+(y) = d_D^+(y') = 0$ and $d_D^+(x_i) \geq \frac{n-1}{2} - 1$ for each $i \in [n-3]$. Then, $|A(D)| \geq (n-2)\frac{n-3}{2}$. Without loss of generality, assume that $d_D^-(x_1) = \max\{d_D^-(x_i) : i \in [n-2]\}$. In what follows, we are going to prove that $d_D^-(x_1) \leq \frac{n-5}{2}$. Suppose that $d_D^-(x_1) \geq \frac{n-3}{2}$. We distinguish the proof into two subcases according to yy' is an edge in G_{n-1} or not.

Subcase 2.1. $yy' \in E(G_{n-1})$.

Let

$$I_1^2 = \{i \in [2, n-3] : x_{i+1}y \in E(G_1)\}, \quad I_2^2 = \{i \in [2, n-3] : x_i \in N_D^-(x_1)\}.$$

Notice that $x_1y, x_2y \notin E(G_1)$, otherwise either $y'yx_1x_{n-2} \dots x_2$ or $y'yx_2 \dots x_{n-2}x_1$ is a transversal Hamilton path in \mathcal{G} , a contradiction. Recall that $d_D^-(x_1) \geq \frac{n-3}{2}$. Hence, $|I_1^2| + |I_2^2| \geq \frac{n-3}{2} + \frac{n-3}{2} = n-3$. Together with $I_1^2 \cap I_2^2 \subseteq [2, n-3]$, there exists an i_2 in $I_1^2 \cap I_2^2$. Hence, $y'yx_{i_2+1} \dots x_{n-2}x_1x_{i_2} \dots x_2$ is a transversal Hamilton path in \mathcal{G} (see Figure 2 (a)), which is a contradiction.

Subcase 2.2. $yy' \notin E(G_{n-1})$.

Let

$$I_1^3 = \{i \in [n-2] : x_i \in N_D^-(y)\}, \quad I_2^3 = \{i \in [n-2] : x_{i+1}y \in E(G_{n-1})\}.$$




FIGURE 2.

It is routine to check that $I_1^3 \cap I_2^3 = \emptyset$. Otherwise, let $i_3 \in I_1^3 \cap I_2^3$. Then $x_1 \dots x_{i_3} y x_{i_3+1} \dots x_{n-2} x_1$ is a rainbow cycle of length $n - 1$ in \mathcal{G} , which contradicts the assumption of Case 2. Note that $I_1^3 \cup I_2^3 \subseteq [n - 2]$ and $|I_2^3| \geq \frac{n-1}{2}$. Then $|I_1^3| \leq \frac{n-3}{2}$, i.e., $d_D^-(y) \leq \frac{n-3}{2}$. Similarly, we have $d_D^-(y') \leq \frac{n-3}{2}$. Now, we give the following claim.

Claim 3.3. $yy' \notin E(G_1)$.

Proof of Claim 3.3. Suppose to the contrary that $yy' \in E(G_1)$. In view of Subcase 2.1, we can assume that $\{G_i[V \setminus \{y, y'\}] : i \in [2, n - 1]\}$ contains no transversal Hamilton cycles. Hence $x_1 x_2 \notin E(G_{n-1})$. Let

$$I_1^4 = \{i \in [2, n - 3] : x_i \in N_D^-(x_1)\}, \quad I_2^4 = \{i \in [2, n - 3] : x_{i+1} x_2 \in E(G_{n-1})\}.$$

Then $|I_1^4| + |I_2^4| \geq \frac{n-3}{2} + \frac{n-1}{2} = n - 2$. Since $I_1^4 \cup I_2^4 \subseteq [2, n - 2]$, one has $I_1^4 \cap I_2^4 \neq \emptyset$. Let $i_4 \in I_1^4 \cap I_2^4$. Then $x_1 x_{i_4} x_{i_4-1} \dots x_2 x_{i_4+1} \dots x_{n-2} x_1$ is a transversal Hamilton cycle inside $\{G_i[V \setminus \{y, y'\}] : i \in [2, n - 1]\}$, a contradiction. \square

Define

$$I_1^5 = \{i \in [n - 2] : x_i y \in E(G_1)\}, \quad I_2^5 = \{i \in [n - 2] : x_{i+1} y \in E(G_{n-1})\}.$$

By Claim 3.3 and the fact that $yy' \notin E(G_{n-1})$, we have $|I_1^5| + |I_2^5| \geq \frac{n-1}{2} + \frac{n-1}{2} = n - 1$. Then $I_1^5 \cap I_2^5 \neq \emptyset$ and choose $i_5 \in I_1^5 \cap I_2^5$. Hence, $P'' := x_2 \dots x_{i_5} y x_{i_5+1} \dots x_{n-2} x_1$ is a rainbow path inside \mathcal{G} . For convenience, we rewrite $P'' = x^1 \dots x^{n-1}$ such that $x^1 := x_2$ and $x^{n-1} := x_1$. Let

$$I_1^6 = \{i \in [n - 3] \setminus \{i_5 - 1\} : y' x^{i+1} \in E(G_{i_5})\}, \quad I_2^6 = \{i \in [n - 3] \setminus \{i_5\} : x^i \in N_D^-(x^{n-1})\}.$$

Clearly, $yy' \notin E(G_{i_5})$. Otherwise, $y' y x_{i_5+1} \dots x_{n-2} x_1 x_2 \dots x_{i_5}$ is a transversal Hamilton path in \mathcal{G} , a contradiction. Furthermore, $y' x^{n-1}, y' x^1 \notin E(G_{i_5})$. Notice that $y = x^{i_5}$ and $d_D^+(y) = d_D^+(y') = 0$. Hence $|I_1^6| + |I_2^6| \geq \frac{n-1}{2} + \frac{n-3}{2} = n - 2$. Therefore, $I_1^6 \cap I_2^6 \neq \emptyset$ and choose $i_6 \in I_1^6 \cap I_2^6$. Then $i_6 \notin \{i_5 - 1, i_5\}$ and so $x^{i_6}, x^{i_6+1} \neq y$. By the definition of I_2^6 and D , we have $x^{i_6} x^{i_6+1}, x^{i_6} x^{n-1} \in E(G_{i_6+1})$ if $i_6 \leq i_5 - 1$, and $x^{i_6} x^{i_6+1}, x^{i_6} x^{n-1} \in E(G_{i_6})$ if $i_6 \geq i_5 + 1$. Hence, $P'' - \{x^{i_6} x^{i_6+1}\} + \{y' x^{i_6+1}, x^{i_6} x^{n-1}\}$ forms a transversal Hamilton path in \mathcal{G} (see Figure 2 (b)), which is also a contradiction.

Therefore, we obtain $d_D^-(x_i) \leq d_D^-(x_1) \leq \frac{n-5}{2}$ for all $i \in [n - 2]$. Recall that $d_D^-(y), d_D^-(y') \leq \frac{n-3}{2}$. Thus $|A(D)| \leq (n - 2) \frac{n-5}{2} + (n - 3) < (n - 2) \frac{n-3}{2}$, a contradiction. This implies that there exists a transversal Hamilton path in \mathcal{G} .

This completes the proof of Theorem 1.3. \square

4. PROOF OF THEOREM 1.6

4.1. Preliminaries. In this subsection, we give some preliminaries. Firstly, we define the extremality for a single graph.

Definition 4.1 ([20]). *Let $\epsilon > 0$ and G be a graph on vertex set V of size n . We call G is ϵ -extremal if there exists a partition $A \cup B \cup C$ of V such that $|A| = |B| = (\frac{1}{2} - \epsilon)n$, and one of the following holds:*

- (i) $d_G(a, A) \geq (\frac{1}{2} - 2\epsilon)n$ for all $a \in A$ and $d_G(b, B) \geq (\frac{1}{2} - 2\epsilon)n$ for all $b \in B$;
- (ii) $d_G(a, B) \geq (\frac{1}{2} - 2\epsilon)n$ for all $a \in A$ and $d_G(b, A) \geq (\frac{1}{2} - 2\epsilon)n$ for all $b \in B$.

For convenience, we say G is $(\epsilon, K_{\lceil \frac{n}{2} \rceil} \cup K_{\lfloor \frac{n}{2} \rfloor})$ -extremal if (i) holds; and G is $(\epsilon, K_{\lceil \frac{n}{2} \rceil, \lfloor \frac{n}{2} \rfloor})$ -extremal if (ii) holds. We always call (A, B, C) a characteristic partition of G .

Let $\mathcal{G} = \{G_1, \dots, G_n\}$ be a graph collection on a common vertex set V of size n and $\delta(\mathcal{G}) \geq \frac{n}{2} - 1$. Let $0 < \alpha, \beta, \delta < 1$. We say \mathcal{G} is (α, β) -strongly stable if G_i is α -extremal for at most $(1 - \beta)n$ colors $i \in [n]$. Assume that G_1, \dots, G_m are all α -extremal graphs inside \mathcal{G} . By Definition 4.1, we can fix a characteristic partition (A_i, B_i, C_i) of G_i for each $i \in [m]$. We say G_i and G_j are δ -crossing if G_i and G_j are α -extremal and $|A_i \Delta A_j| \geq \delta n$, $|A_i \Delta B_j| \geq \delta n$. For $i \in [m]$, define I_i to be the set of $j \in [m]$ such that G_i and G_j are δ -crossing. Under the above characteristic partitions, we say \mathcal{G} is (α, δ) -weakly stable if $\sum_{i \in [m]} |I_i| \geq 2\delta m^2$.

Cheng and Staden [20, Lemma 5.3] proved the following result.

Lemma 4.2 ([20]). *Assume $0 < \frac{1}{n} \ll \mu \ll \alpha \ll \beta, \epsilon \ll \delta \ll 1$. Let $\mathcal{G} = \{G_1, \dots, G_n\}$ be a graph collection on a common vertex set V of size n and $\delta(\mathcal{G}) \geq (\frac{1}{2} - \mu)n$. If \mathcal{G} is either (α, β) -strongly stable or (ϵ, δ) -weakly stable, then \mathcal{G} contains a transversal Hamilton cycle.*

The following theorem characterizes the structure of almost balanced graph collections (i.e., almost all graphs in the graph collection have an almost balanced common vertex bipartition such that their subgraphs induced by the bipartition are close to complete bipartite graphs) that do not contain transversal Hamilton cycles. For the sake of readability, we postpone its proof in Section 5.

Theorem 4.3. *Let $0 < \frac{1}{n} \ll \delta \ll 1$ and $0 \leq \gamma \leq 3\delta$. Assume \mathcal{C} is a set of n colors and $\mathcal{G} = \{G_i : i \in \mathcal{C}\}$ is collection of graphs on a common vertex set V of size n with $\delta(\mathcal{G}) \geq \frac{n}{2} - 1$. Let $\mathcal{C}' \cup \mathcal{C}''$ be a partition of \mathcal{C} with $|\mathcal{C}''| \leq \delta n$, and let $A \cup B$ be a partition of V with $|A| = \lceil \frac{n}{2} - 1 \rceil + \gamma n$. Suppose that $G_i[B] = \emptyset$ for all $i \in \mathcal{C}'$ and \mathcal{G} does not contain a transversal Hamilton cycle. Then \mathcal{G} must be one of the graph collections described in Theorem 1.6.*

4.2. Proof of Theorem 1.6. In this subsection, we give the proof of Theorem 1.6.

Proof of Theorem 1.6. Suppose that \mathcal{G} does not contain transversal Hamilton cycles. We are to prove that \mathcal{G} must be a graph collection described in Theorem 1.6. Let $0 < \frac{1}{n} \ll \alpha \ll \beta, \epsilon \ll \delta \ll \eta \ll 1$. By Lemma 4.2, we know \mathcal{G} is not (α, β) -strongly stable. Thus, there are at least $(1 - \beta)n$ graphs in \mathcal{G} such that each of them is α -extremal, and hence ϵ -extremal. Therefore, there exists a constant $\beta' (< \beta)$ such that the number of ϵ -extremal graphs in \mathcal{G} is exactly $(1 - \beta')n$. Without loss of generality, assume that for each $i \in [(1 - \beta')n]$, G_i is ϵ -extremal. By Definition 4.1, we can fix a characteristic partition $(\tilde{A}_i, \tilde{B}_i, \tilde{C}_i)$ for each G_i with $i \in [(1 - \beta')n]$.

Under the above characteristic partition, applying Lemma 4.2 again yields that \mathcal{G} is not (ϵ, δ) -weakly stable. Therefore, $\sum_{i \in [(1 - \beta')n]} |I_i| < 2\delta(1 - \beta')^2 n^2$. Without loss of generality, assume

that $|I_1| = \min\{|I_i| : i \in [(1 - \beta')n]\}$. Then $|I_1| \leq 2\delta(1 - \beta')n$. For every $i \in [(1 - \beta')n] \setminus I_1$, based on the definition of I_1 , we have either $|\tilde{A}_1 \Delta \tilde{A}_i| < \delta n$ or $|\tilde{A}_1 \Delta \tilde{B}_i| < \delta n$. By swapping the labels of \tilde{A}_i and \tilde{B}_i , we have $|\tilde{A}_1 \Delta \tilde{A}_i|, |\tilde{B}_1 \Delta \tilde{B}_i| < \delta n$ for every $i \in [(1 - \beta')n] \setminus I_1$. Denote $[m] := [(1 - \beta')n] \setminus I_1$ and $\mathcal{C}_{bad} := [m + 1, n]$. Then $m \geq (1 - \beta' - 2\delta)n \geq (1 - 3\delta)n$. By adding colors to \mathcal{C}_{bad} if necessary we may assume $m = (1 - 3\delta)n$. Recall that $\alpha \ll \epsilon$. Hence for each $i \in [2, m]$, there exists a new partition (A_i, B_i, C_i) of G_i such that

- (B1) if G_i is $(\epsilon, K_{\lceil \frac{n}{2} \rceil} \cup K_{\lfloor \frac{n}{2} \rfloor})$ -extremal, then for $Y \in \{A, B\}$ we have $\tilde{Y}_i \subseteq Y_i$ and $d_{G_i}(v, Y_i) \geq (\frac{1}{2} - 3\sqrt{\delta})n$ for each vertex $v \in Y_i$,
- (B2) if G_i is $(\epsilon, K_{\lceil \frac{n}{2} \rceil, \lfloor \frac{n}{2} \rfloor})$ -extremal, then for $Y \in \{A, B\}$ we have $\tilde{Y}_i \subseteq Y_i$ and $d_{G_i}(v, Z_i) \geq (\frac{1}{2} - 3\sqrt{\delta})n$ for each vertex $v \in Y_i$,
- (B3) subject to (B1)-(B2), each vertex in V lies in as many as possible $A_i \cup B_i$ for $i \in [2, m]$.

Denote $(A_1, B_1, C_1) := (\tilde{A}_1, \tilde{B}_1, \tilde{C}_1)$. Expand $A_1 \cup B_1$ to be an equitable partition $A \cup B$ of V . Let $\mathcal{C}_1 \cup \mathcal{C}_2$ be a partition of $[m]$ with

$$\begin{aligned} \mathcal{C}_1 &= \{i \in [m] : G_i \text{ is } (\epsilon, K_{\lceil \frac{n}{2} \rceil} \cup K_{\lfloor \frac{n}{2} \rfloor})\text{-extremal}\}, \\ \mathcal{C}_2 &= \{i \in [m] : G_i \text{ is } (\epsilon, K_{\lceil \frac{n}{2} \rceil, \lfloor \frac{n}{2} \rfloor})\text{-extremal}\}. \end{aligned}$$

Hence for $i \in [2, m]$ and $Y \in \{A, B\}$, we have $|Y_1 \Delta Y_i| \leq |\tilde{Y}_1 \Delta \tilde{Y}_i| + |\tilde{C}_i| \leq \delta n + 2\epsilon n < 2\delta n$.

In the following of our proof, we shall emphasize that if we say a set of vertices $V' \subseteq B$ is moved from B to A , then the partition of V becomes to $(A \cup V') \cup (B \setminus V')$. In order not to introduce too much notation, we still denote the resulted partition by $A \cup B$. Moreover, unless otherwise specified, we assume $Y \in \{A, B\}$ and $\{Y, Z\} = \{A, B\}$.

Now, we define the set of bad vertices. Let $\hat{\mathcal{C}} := \cup_{k \in [2]} \psi(\mathcal{C}_k)$, where $\psi(\mathcal{C}_k) = \mathcal{C}_k$ if $|\mathcal{C}_k| \geq \eta n$ and $\psi(\mathcal{C}_k) = \emptyset$ otherwise. Clearly, $|\hat{\mathcal{C}}| \geq (1 - 3\delta - \eta)n$. Define

$$\begin{aligned} X &= \{x \in V : x \notin A_i \cup B_i \text{ for at least } 3\sqrt{\delta}|\hat{\mathcal{C}}| \text{ colors } i \in \hat{\mathcal{C}}\}, \\ X_Y &= \{x \in Y \setminus X : x \notin Y_i \text{ for at least } 10\sqrt{\delta}|\hat{\mathcal{C}}| \text{ colors } i \in \hat{\mathcal{C}}\}. \end{aligned}$$

Recall that for all $i \in \hat{\mathcal{C}}$, we have $|C_i| \leq 2\epsilon n$ and $|Y_1 \Delta Y_i| < 2\delta n$. Hence $3\sqrt{\delta}|\hat{\mathcal{C}}||X| \leq 2\epsilon n|\hat{\mathcal{C}}|$ and $10\sqrt{\delta}|\hat{\mathcal{C}}||X_Y| \leq \sum_{i \in \hat{\mathcal{C}}} |Y \setminus Y_i| \leq (2\delta + 2\epsilon)n|\hat{\mathcal{C}}|$. It follows that $|X| \leq \sqrt{\epsilon}n$ and $|X_Y| < \frac{1}{4}\sqrt{\delta}n$.

For $k \in [2]$, define

$$X_Y^k = \begin{cases} \{x \in X_Y : x \in Y_i \text{ for at least } 3\sqrt{\delta}n \text{ colors } i \in \mathcal{C}_k\}, & \text{if } |\mathcal{C}_k| \geq \eta n, \\ \emptyset, & \text{otherwise.} \end{cases}$$

Recall that $\{Y, Z\} = \{A, B\}$. For a vertex $x \in X_Y \setminus (\cup_{k \in [2]} X_Y^k)$, we know $x \in Y_i$ for at most $6\sqrt{\delta}n$ colors $i \in \hat{\mathcal{C}}$ and $x \in A_i \cup B_i$ for at least $(1 - 3\sqrt{\delta})|\hat{\mathcal{C}}|$ colors $i \in \hat{\mathcal{C}}$. Thus, $x \in Z_i$ for at least $(1 - 3\sqrt{\delta})|\hat{\mathcal{C}}| - 6\sqrt{\delta}n \geq (1 - 10\sqrt{\delta})|\hat{\mathcal{C}}|$ colors $i \in \hat{\mathcal{C}}$. This implies that $x \in Y \setminus Y_i$ for at least $(1 - 10\sqrt{\delta})|\hat{\mathcal{C}}|$ colors $i \in \hat{\mathcal{C}}$. Hence

$$|X_Y \setminus (\cup_{k \in [2]} X_Y^k)| (1 - 10\sqrt{\delta})|\hat{\mathcal{C}}| \leq \sum_{i \in \hat{\mathcal{C}}} |Y \setminus Y_i| \leq \sum_{i \in \hat{\mathcal{C}}} (|Y \setminus Y_1| + |Y_1 \Delta Y_i|) \leq (2\delta + 2\epsilon)n|\hat{\mathcal{C}}|.$$

It follows that $|X_Y \setminus (\cup_{k \in [2]} X_Y^k)| \leq 4\delta n$. Then move vertices in $X_A \setminus (\cup_{k \in [2]} X_A^k)$ to B and vertices in $X_B \setminus (\cup_{k \in [2]} X_B^k)$ to A . Without loss of generality, assume that $|B| - |A| = r$, where $0 \leq r \leq 8\delta n + \sigma$, where $\sigma = 0$ if n is even and $\sigma = 1$ if n is odd. Define $V_{bad} := \cup_{k \in [2]} (X_A^k \cup X_B^k) \cup X$. Obviously, $|V_{bad}| < (\sqrt{\epsilon} + \frac{1}{2}\sqrt{\delta})n$. Moreover, each vertex in $Y \setminus V_{bad}$ lies in Y_i for at least $(1 - 10\sqrt{\delta})|\hat{\mathcal{C}}|$ colors $i \in \hat{\mathcal{C}}$.

Next, we give an application of the transversal blow-up lemma (see [19]) for embedding transversal Hamilton paths inside very dense bipartite graph collections, which can be proved by minor modifications to the proof of [20, Lemma 6.1], and we omit the proof here.

Claim 4.4. *Assume $W, Z \in \{A, B\}$. Let $W^* \subseteq W \setminus V_{bad}$ and $Z^* \subseteq Z \setminus V_{bad}$, where $|W^*|, |Z^*| \geq \eta n$, $W^* \cap Z^* = \emptyset$ and $|W^*| - |Z^*| = t \in \{0, 1\}$. Let $T^* = Z^*$ if $t = 0$ and $T^* = W^*$ if $t = 1$. Let $\mathcal{C}^* \subseteq \mathcal{C}$ satisfy $|\mathcal{C}^*| = |W^*| + |Z^*| - 1$, where $\mathcal{C}^* \subseteq \mathcal{C}_1$ if $W = Z$ and $\mathcal{C}^* \subseteq \mathcal{C}_2$ if $W \neq Z$. Let $W^- \subseteq W^*$ and $T^+ \subseteq T^*$ with $|W^-|, |T^+| \geq \frac{\eta n}{8}$. Then there is a transversal Hamilton path in $\{G_i[W^*, Z^*] : i \in \mathcal{C}^*\}$ starting in W^- and ending in T^+ .*

The subsequent result can be used to connect two disjoint short rainbow paths into a single short rainbow path.

Claim 4.5 (Connecting tool). *Assume $P = u_1 u_2 \dots u_s$ and $Q = v_1 v_2 \dots v_t$ are two disjoint rainbow paths inside \mathcal{G} and $s + t \leq 5\eta n$.*

- (i) *If $u_s \in A \setminus V_{bad}$, $v_1 \in B \setminus V_{bad}$ and $|\mathcal{C}_2 \setminus \text{col}(P \cup Q)| \geq 11\sqrt{\delta}n$, then there are three colors $c_1, c_2, c_3 \in \mathcal{C}_2 \setminus \text{col}(P \cup Q)$ and two vertices $w_1 \in B \setminus (V(P \cup Q) \cup V_{bad})$, $w'_1 \in A \setminus (V(P \cup Q) \cup V_{bad})$ such that $u_1 P u_s w_1 w'_1 v_1 Q v_t$ is a rainbow path with colors $\text{col}(P \cup Q) \cup \{c_1, c_2, c_3\}$.*
- (ii) *If $u_s, v_1 \in Y \setminus V_{bad}$ and $|\mathcal{C}_2 \setminus \text{col}(P \cup Q)| \geq 11\sqrt{\delta}n$, then there are two colors $c_1, c_2 \in \mathcal{C}_2 \setminus \text{col}(P \cup Q)$ and one vertex $w_1 \in Z \setminus (V(P \cup Q) \cup V_{bad})$ such that $u_1 P u_s w_1 v_1 Q v_t$ is a rainbow path with colors $\text{col}(P \cup Q) \cup \{c_1, c_2\}$.*
- (iii) *If $u_s, v_1 \in Y \setminus V_{bad}$ and $|\mathcal{C}_1 \setminus \text{col}(P \cup Q)| \geq 11\sqrt{\delta}n$, then there are two colors $c_1, c_2 \in \mathcal{C}_1 \setminus \text{col}(P \cup Q)$ and one vertex $w_1 \in Y \setminus (V(P \cup Q) \cup V_{bad})$ such that $u_1 P u_s w_1 v_1 Q v_t$ is a rainbow path with colors $\text{col}(P \cup Q) \cup \{c_1, c_2\}$. (See Figure 3.)*




FIGURE 3. Connecting tool.

Proof of Claim 4.5. We only give the proof of (i), the other two statements can be proved by a similar discussion, whose procedures are omitted.

Note that $u_s, v_1 \notin V_{bad}$. Then u_s (resp. v_1) lies in A_i (resp. B_i) for at least $(1 - 10\sqrt{\delta})|\hat{\mathcal{C}}|$ colors $i \in \hat{\mathcal{C}}$. Since $|\mathcal{C}_2 \setminus \text{col}(P \cup Q)| \geq 11\sqrt{\delta}n$, we obtain that u_s (resp. v_1) lies in A_i (resp. B_i) for at least $11\sqrt{\delta}n - 10\sqrt{\delta}|\hat{\mathcal{C}}| \geq \sqrt{\delta}n$ colors $i \in \mathcal{C}_2$. Hence there are two colors $c_1, c_2 \in \mathcal{C}_2 \setminus \text{col}(P \cup Q)$ such that $u_s \in A_{c_1}$ and $v_1 \in B_{c_2}$. It is routine to check that

$$\begin{aligned}
& |N_{G_{c_1}}(u_s) \cap (B \setminus V_{bad})| \\
& \geq |N_{G_{c_1}}(u_s) \cap B_{c_1}| - |B_1 \Delta B_{c_1}| - |X \cup X_A \cup X_B| \\
& \geq \left(\frac{1}{2} - 3\sqrt{\delta}\right)n - (2\delta + \sqrt{\epsilon} + \frac{1}{2}\sqrt{\delta})n \geq \left(\frac{1}{2} - 5\sqrt{\delta}\right)n.
\end{aligned}$$

Hence there exists a vertex $w_1 \in N_{G_{c_1}}(u_s) \cap (B \setminus V_{bad})$ that avoids $V(P \cup Q)$. Similarly, since $w_1 \notin V_{bad}$, there is a color $c_3 \in \mathcal{C}_2 \setminus (\text{col}(P \cup Q) \cup \{c_1, c_2\})$ such that $w_1 \in B_{c_3}$. On the other hand,

$$\begin{aligned} & |N_{G_{c_2}}(v_1) \cap N_{G_{c_3}}(w_1) \cap (A \setminus V_{bad})| \\ & \geq |N_{G_{c_2}}(v_1) \cap (A \setminus V_{bad})| + |N_{G_{c_3}}(w_1) \cap (A \setminus V_{bad})| - \frac{n}{2} \\ & \geq \left(\frac{1}{2} - 10\sqrt{\delta}\right)n. \end{aligned}$$

Hence, there is a vertex $w'_1 \in N_{G_{c_2}}(v_1) \cap N_{G_{c_3}}(w_1) \cap (A \setminus V_{bad})$ that avoids $V(P \cup Q)$. Therefore, $u_1 P u_s w_1 w'_1 v_1 Q v_t$ is a rainbow path inside \mathcal{G} with colors $\text{col}(P \cup Q) \cup \{c_1, c_2, c_3\}$. \square

Based on the sizes of $|\mathcal{C}_1|$ and $|\mathcal{C}_2|$, our proof is distinguished into the following three cases.

Case 1. $|\mathcal{C}_1| < \eta n$.

In this case, each vertex in $Y \setminus (X \cup X_Y^2)$ belongs to Y_i for at least $(1 - 10\sqrt{\delta})|\mathcal{C}_2|$ colors $i \in \mathcal{C}_2$. Recall that $|B| - |A| = r$ with $0 \leq r \leq 8\delta n + \sigma$, where $\sigma = 0$ if n is even and $\sigma = 1$ if n is odd. The proof is divided into four steps, as shown in the Figure 4.




FIGURE 4. Step 1 - Step 3. The vertex with red (resp. green) lies in $A \setminus V_{bad}$ (resp. $B \setminus V_{bad}$), and the symbol “ \blacklozenge ” denotes a vertex that is used to connect rainbow paths.

Step 1. Balance the number of vertices in A and B .

In this step, we claim that if \mathcal{G} is not a graph collection described in Theorem 1.6, then by moving vertices in $B \cap V_{bad}$ or deleting rainbow paths inside $\mathcal{G}[B]$, we can make the number of remaining vertices in B and A differ by σ . Denote $s := |V_{bad} \cap B|$. If $|B| - |A| - \sigma \leq 2s$, then move $\frac{|B| - |A| - \sigma}{2}$ vertices in $V_{bad} \cap B$ to A , we can get our desired result. Hence, it suffices to consider $|B| - |A| - \sigma > 2s$. We first move all vertices in $V_{bad} \cap B$ to A . Then $B \cap V_{bad} = \emptyset$. Assume that $\{Q_1, \dots, Q_t\}$ is a set of disjoint maximal rainbow paths in $\mathcal{G}[B]$. If $|E(Q_1)| + \dots + |E(Q_t)| \geq |B| - |A| - \sigma$, then there must exist a set of disjoint rainbow paths, say $\{Q'_1, \dots, Q'_{t'}\}$, such that $|E(Q'_1)| + \dots + |E(Q'_{t'})| = |B| - |A| - \sigma$. That is, $|A| - t' = |B| - (|V(Q'_1)| + \dots + |V(Q'_{t'})|) - \sigma$. Using Claim 4.5 (ii) to connect $Q'_1, \dots, Q'_{t'}$ into a single rainbow path P^1 , whose endpoints are in different parts. (In fact, by using Claim 4.5 we only get a rainbow path with two endpoints in B . But we usually want to find a rainbow path with endpoints in different parts. Hence we extend it by using an unused color of \mathcal{C}_2 and an unused vertex in A). Therefore, $|A \setminus V(P^1)| = |B \setminus V(P^1)| - \sigma$, as desired.

In what follows, it suffices to consider $|E(Q_1)| + \dots + |E(Q_t)| < |B| - |A| - \sigma = r - 2s - \sigma$. Now, we move all vertices of Q_1, \dots, Q_t to A . Let $\mathcal{C}' = \mathcal{C} \setminus \text{col}(\cup_{i \in [t]} Q_i)$. Hence

$$\begin{aligned} |A| &\leq \frac{n-r}{2} + s + 2(r-2s-\sigma) \leq \frac{n}{2} + 16\delta n, \\ |B| &\geq \frac{n+r}{2} - s - 2(r-2s-\sigma) \geq \frac{n}{2} - 16\delta n, \\ |\mathcal{C}'| &= |\mathcal{C}| - (|E(Q_1)| + \dots + |E(Q_t)|) \geq (1-8\delta)n. \end{aligned}$$

By the maximality of $\{Q_1, \dots, Q_t\}$, we know $G_i[B] = \emptyset$ for all $i \in \mathcal{C}'$. It follows from $\delta(\mathcal{G}) \geq \lceil \frac{n}{2} - 1 \rceil$ that $|A| \geq \lceil \frac{n}{2} - 1 \rceil$. Together with Theorem 4.3, we obtain that \mathcal{G} is one of the graph collections described in Theorem 1.6, as desired.

In this step, if we only move vertices in V_{bad} from B to A , then let P^1 be a null graph (i.e., no vertices); otherwise, assume P^1 is a rainbow path obtained in the above, whose length is at most $24\delta n$ and endpoints are $u_1 \in A \setminus V_{bad}$, $v_1 \in B \setminus V_{bad}$. Hence $|A \setminus V(P^1)| = |B \setminus V(P^1)| - \sigma$.

Step 2. Construct a series of disjoint rainbow P_3 such that all centers of them are exactly all vertices in V_{bad} .

Recall that $|X_A^2 \cup X_B^2| \leq |X_A \cup X_B| < \frac{1}{2}\sqrt{\delta}n$. Let $v \in X_Y^2$. If $v \in Z$, then v is moved from B to A in Step 1. Hence $v \in X_B$. Therefore, $v \in A_i \cup B_i$ for at least $(1-3\sqrt{\delta})|\mathcal{C}_2|$ colors $i \in \mathcal{C}_2$ and $v \notin B_i$ for at least $10\sqrt{\delta}|\mathcal{C}_2|$ colors $i \in \mathcal{C}_2$. Hence, $v \in A_i$ for at least $7\sqrt{\delta}|\mathcal{C}_2| - 24\delta n > 4|X_A^2 \cup X_B^2|$ colors $i \in \mathcal{C}_2 \setminus \text{col}(P^1)$. If $v \in Y$ (i.e., v has not been moved in Step 1), then v belongs to Y_i for at least $(3\sqrt{\delta} - 24\delta)n > 4|X_A^2 \cup X_B^2|$ colors $i \in \mathcal{C}_2 \setminus \text{col}(P^1)$. Hence, using colors in $\mathcal{C}_2 \setminus \text{col}(P^1)$, we can greedily choose $|X_A^2 \cup X_B^2|$ disjoint rainbow P_3 , such that all centers of them are exactly all vertices in $X_A^2 \cup X_B^2$. Moreover, if $v \in (X_A^2 \cup X_B^2) \cap Y$, then the rainbow P_3 with center in v has endpoints in $Z \setminus (V_{bad} \cup V(P^1))$.

Connect those rainbow P_3 with centers in Y (if exists) into a single rainbow path with two endpoints in different parts by Claim 4.5 (ii). Then connect the two resulted rainbow paths and P^1 by Claim 4.5 (i)-(ii), we obtain a rainbow path $P^2 = u_1 \dots v_2$ with length at most $|E(P^1)| + 4|X_A^2 \cup X_B^2| + 4 \leq 24\delta n + 2\sqrt{\delta}n + 4$ and $v_2 \in B \setminus V_{bad}$.

For vertices in X , we consider the following claim. Notice that the position (in A or B) of vertices in X will not work in our proof, so if $v \in X$ is a vertex moved from B to A , then we only need consider $v \in A$.

Claim 4.6. *Assume $X \setminus V(P^2) = \{x_1, \dots, x_s\}$ with $s \leq \sqrt{\epsilon}n$. If $x_i \in X \cap Y$, then there exist $c_i^1, c_i^2, c_i^3, c_i^4 \in \mathcal{C}_2 \setminus (\text{col}(P^2) \cup \{c_\ell^1, c_\ell^2, c_\ell^3, c_\ell^4 : \ell \in [i-1]\})$ and $x_i^1, x_i^2, x_i^3, x_i^4 \in Z \setminus (V(P^2) \cup V_{bad} \cup \{x_\ell^1, x_\ell^2, x_\ell^3, x_\ell^4 : \ell \in [i-1]\})$ such that $x_i x_i^k \in E(G_{c_i^k})$ for all $k \in [4]$.*

Proof of Claim 4.6. Suppose there exists an $i_0 \in [s]$ such that Claim 4.6 holds for all $i \in [i_0 - 1]$ but does not hold for i_0 . Without loss of generality, assume that $x_{i_0} \in X \cap A$. Hence for all but at most three $j \in \mathcal{C}_2 \setminus (\text{col}(P^2) \cup \{c_\ell^1, c_\ell^2, c_\ell^3, c_\ell^4 : \ell \in [i_0 - 1]\})$ we have $d_{G_j}(x_{i_0}, B) \leq 4(i_0 - 1) + 3 + |V(P^2)|$. Thus, for such a color j ,

$$\begin{aligned} d_{G_j}(x_{i_0}, A_j) &\geq d_{G_j}(x_{i_0}, A) - |A \setminus A_1| - |A_1 \Delta A_j| - |X_A \setminus X_A^2| \\ &\geq \delta(G_j) - d_{G_j}(x_{i_0}, B) - |A \setminus A_1| - |A_1 \Delta A_j| - |X_A \setminus X_A^2| \\ &\geq \frac{n}{2} - 1 - (4(i_0 - 1) + 3 + |V(P^2)|) - 2\epsilon n - 2\delta n - 4\delta n \\ &\geq \frac{n}{2} - 1 - (4\sqrt{\epsilon}n + 24\delta n + 2\sqrt{\delta}n + 4) - 2\epsilon n - 6\delta n \end{aligned}$$

$$\geq \left(\frac{1}{2} - 3\sqrt{\delta}\right)n.$$

Hence $x_{i_0} \in A_j \cup B_j$. Otherwise, there exists a new characteristic partition (A_j, B'_j, C'_j) of G_j with $B'_j = B_j \cup \{x_{i_0}\}$, which contradicts **(B3)**. By the choice of j , we know $x_{i_0} \in A_j \cup B_j$ for all but at most three $j \in \mathcal{C}_2 \setminus (\text{col}(P^2) \cup \{c_\ell^1, c_\ell^2, c_\ell^3, c_\ell^4 : \ell \in [i_0 - 1]\})$. It is routine to check that

$$\begin{aligned} & |\mathcal{C}_2 \setminus (\text{col}(P^2) \cup \{c_\ell^1, c_\ell^2, c_\ell^3, c_\ell^4 : \ell \in [i_0 - 1]\})| - 3 \\ & \geq |\mathcal{C}_2| - 24\delta n - 2\sqrt{\delta}n - 4 - 4\sqrt{\epsilon}n - 3 \\ & \geq |\mathcal{C}_2| - 3\sqrt{\delta}n. \end{aligned}$$

Therefore, $x_{i_0} \in C_j$ for at most $3\sqrt{\delta}n$ colors $j \in \mathcal{C}_2$, which implies $x_{i_0} \notin X$, a contradiction. \square

By Claim 4.6, for each $x_i \in X \cap Y$, there exists a rainbow path $Q_{x_i} = x_i^1 x_i x_i^2$ with $\text{col}(Q_{x_i}) = \{c_i^1, c_i^2\} \subseteq \mathcal{C}_2$ and $x_i^1, x_i^2 \in Z \setminus V_{\text{bad}}$. Furthermore, each color or endpoint in Q_{x_i} can be replaced by two colors or vertices not in P^2 . Connect rainbow paths in $\{Q_{x_i} : x_i \in X \cap Y\}$ by Claim 4.5 (i)-(ii) into a single rainbow path with endpoints in different parts (if $|X \cap Y| \geq 1$). Then using Claim 4.5 (i)-(ii) to connect P^2 and the resulted two rainbow paths into a single rainbow path P^3 with endpoints $u_1 \in A \setminus V_{\text{bad}}$ and $v_3 \in B \setminus V_{\text{bad}}$, whose length is at most $|E(P^2)| + 4|X| + 4 \leq 24\delta n + 2\sqrt{\delta}n + 4\sqrt{\epsilon}n + 8$.

Step 3. Select a rainbow matching inside $\{G_i[A \setminus V(P^3), B \setminus V(P^3)] : i \in \mathcal{C}_{\text{bad}} \setminus \text{col}(P^3)\}$.

Choose a maximum rainbow matching, say \tilde{M} , in $\{G_i[A \setminus V(P^3), B \setminus V(P^3)] : i \in \mathcal{C}_{\text{bad}} \setminus \text{col}(P^3)\}$. If \tilde{M} contains an edge with color j such that $G_j[A \setminus V(P^3 \cup \tilde{M}), B \setminus V(P^3 \cup \tilde{M})]$ contains no 2-matching, then delete all such edges from \tilde{M} and denote the resulted rainbow matching by M . Connect P^3 and all rainbow edges in M by Claim 4.5 (i), we obtain a rainbow path P^4 with endpoints $u_1 \in A \setminus V_{\text{bad}}$ and $v_4 \in B \setminus V_{\text{bad}}$, whose length is at most $|E(P^3)| + 12\delta n \leq 2\sqrt{\delta}n + 4\sqrt{\epsilon}n + 36\delta n + 8$.

Denote $\mathcal{C}'_{\text{bad}} := \mathcal{C}_{\text{bad}} \setminus \text{col}(P^4)$ and let $j \in \mathcal{C}'_{\text{bad}}$. Hence in G_j , all but at most one vertex in $Y \setminus V(P^3 \cup \tilde{M})$ is adjacent to at least $\frac{n}{2} - 1 - |V(P^3 \cup \tilde{M})| - 1 \geq (\frac{1}{2} - 3\sqrt{\delta})n$ vertices in Y . Therefore, G_j is $(3\sqrt{\delta}, K_{\lfloor \frac{n}{2} \rfloor} \cup K_{\lceil \frac{n}{2} \rceil})$ -extremal for all $j \in \mathcal{C}'_{\text{bad}}$.

Step 4. Construct two disjoint rainbow paths inside $\mathcal{G}[A \setminus V(P^4)]$ and $\mathcal{G}[B \setminus V(P^4)]$ by using unused colors in $\mathcal{C}_1 \cup \mathcal{C}'_{\text{bad}}$.

It is straightforward to check that

$$|B \setminus V(P^4)| - |A \setminus V(P^4)| = |B \setminus V(P^1)| - |A \setminus V(P^1)| = \sigma.$$

By the construction of P^4 , we know that if $w \in V(P^4)$ is a vertex not in $(V(P^1) \cap B) \cup V_{\text{bad}}$, then it is possible to avoid w when construct P^4 ; if $i \in \text{col}(P^4)$ is a color not in $\text{col}(P^1 - A) \cup \text{col}(M)$, then it is possible to avoid i when construct P^4 . That is, we can assume that $w \notin V(P^4)$ unless $w \in (V(P^1) \cap B) \cup V_{\text{bad}}$ and $i \notin \text{col}(P^4)$ unless $i \in \text{col}(P^1 - A) \cup \text{col}(M)$.

Furthermore, in the process of choosing rainbow paths inside $\mathcal{G}[B]$ (in Step 1), one may use colors in $\mathcal{C}_1 \cup \mathcal{C}'_{\text{bad}}$ before other colors and avoid vertices in $V_{\text{bad}} \cup V(M)$. This implies that either $\mathcal{C}_1 \cup \mathcal{C}'_{\text{bad}} \subseteq \text{col}(P^1)$ or $\text{col}(P^1 - A) \subseteq \mathcal{C}_1 \cup \mathcal{C}'_{\text{bad}}$. Denote $\tilde{\mathcal{C}} := (\mathcal{C}_1 \cup \mathcal{C}'_{\text{bad}}) \setminus \text{col}(P^1)$.

Claim 4.7. *Let $Q = x_1 \dots x_s$ be a rainbow path inside \mathcal{G} such that $s \leq 3\sqrt{\delta}n$, $V_{\text{bad}} \subseteq V(Q)$, $(\mathcal{C}_1 \cup \mathcal{C}_{\text{bad}}) \setminus \tilde{\mathcal{C}} \subseteq \text{col}(Q)$, $x_1 \in A \setminus V_{\text{bad}}$ and $x_s \in B \setminus V_{\text{bad}}$. If $|\tilde{\mathcal{C}}| - ||B \setminus V(Q)| - |A \setminus V(Q)||$ is a nonnegative even integer, then \mathcal{G} has a transversal Hamilton cycle.*

Proof of Claim 4.7. Using colors in $\tilde{\mathcal{C}}$, we can greedily choose two disjoint rainbow paths P^5 and P^6 in $\mathcal{G}[A \setminus V(Q)]$ and $\mathcal{G}[B \setminus V(Q)]$ with lengths $\frac{|\tilde{\mathcal{C}}| - (|B \setminus V(Q)| - |A \setminus V(Q)|)}{2}$ and $\frac{|\tilde{\mathcal{C}}| + (|B \setminus V(Q)| - |A \setminus V(Q)|)}{2}$ respectively, whose endpoints are $u_5, v_5 \in A \setminus V_{bad}$ and $u_6, v_6 \in B \setminus V_{bad}$. (Note that if a rainbow path has length 0, then it contains one vertex instead of the null graph.) Applying Claim 4.5 (i)-(ii) to connect Q, P^5 and P^6 in turn, we get a rainbow path P with endpoints $x_1 \in A \setminus V_{bad}$ and $v_6 \in B \setminus V_{bad}$. Clearly, $|V(P)| \leq 3\sqrt{\delta}n + 3\delta n + \eta n + 4$ and $|A \setminus V(P)| = |B \setminus V(P)|$.

Notice that $x_1, v_6 \notin V_{bad}$. Hence there exist $i_1, i_2 \in \mathcal{C}_2 \setminus \text{col}(P)$ such that $x_1 \in A_{i_1}$ and $v_6 \in B_{i_2}$. Let $W^* = A \setminus V(P)$, $T^* = B \setminus V(P)$, $\mathcal{C}^* = \mathcal{C} \setminus (\text{col}(P) \cup \{i_1, i_2\})$, $W^- = N_{G_{i_2}}(v_6) \cap W^*$ and $T^+ = N_{G_{i_1}}(x_1) \cap Z^*$. It is routine to check that $|W^*| = |T^*|$, $|\mathcal{C}^*| = |W^*| + |T^*| - 1$,

$$\begin{aligned} |W^-| &\geq d_{G_{i_2}}(v_6, A) - |V(P)| \geq d_{G_{i_2}}(v_6, A_{i_2}) - |A_1 \Delta A_{i_2}| - |X_A \setminus X_A^2| - |V(P)| \\ &\geq \left(\frac{1}{2} - 3\sqrt{\delta}\right)n - 2\delta n - 4\delta n - |V(P)| > \left(\frac{1}{2} - 2\eta\right)n. \end{aligned}$$

Similarly, $|T^+| > \eta n$. Applying Claim 4.4 yields that $\{G_i[W^*, T^*] : i \in \mathcal{C}^*\}$ contains a transversal path P' starting at $v' \in W^-$ and ending at $u' \in T^+$. Thus, $x_1 P v_6 v' P' u' x_1$ is a transversal Hamilton cycle inside \mathcal{G} (see Figure 5). \square




FIGURE 5.

If n is even and $|\tilde{\mathcal{C}}|$ is even, or n is odd and $|\tilde{\mathcal{C}}|$ is odd, then in view of Claim 4.7, there exists a transversal Hamilton cycle inside \mathcal{G} , a contradiction. Hence, it suffices to consider n is even and $|\tilde{\mathcal{C}}|$ is odd, or n is odd and $|\tilde{\mathcal{C}}|$ is even. Therefore, either $|\tilde{\mathcal{C}}| - (|B \setminus V(P^4)| - |A \setminus V(P^4)|)$ is a positive odd integer or $|\tilde{\mathcal{C}}| = 0$ and n is odd.

Claim 4.8. (i) If $|\tilde{\mathcal{C}}| \geq 1$, then $E(G_i[A]) = \emptyset$ for all $i \in \mathcal{C} \setminus \tilde{\mathcal{C}}$,
(ii) $E(G_i[A, B]) = \emptyset$ for all $i \in \tilde{\mathcal{C}}$.

Proof of Claim 4.8. (i) Since $|\tilde{\mathcal{C}}| \geq 1$, one has $\text{col}(P^1 - A) \subseteq \mathcal{C}_1 \cup \mathcal{C}'_{bad}$. Suppose that there exists an $i_0 \in \mathcal{C} \setminus \tilde{\mathcal{C}}$ such that $E(G_{i_0}[A]) \neq \emptyset$. Choose $w_1 w_2 \in E(G_{i_0}[A])$. Clearly, $w_1, w_2 \notin V(P^1) \cap B$. Recall that $w \notin V(P^4)$ unless $w \in (V(P^1) \cap B) \cup V_{bad}$ and $i_0 \notin \text{col}(P^4)$ unless $i_0 \in \text{col}(P^1 - A) \cup \text{col}(M)$. If $i_0 \in \text{col}(P^1 - A) \cup \text{col}(M)$, then let \tilde{P}^4 be a rainbow path obtained by deleting the edge with color i_0 from P^4 and connecting its two endpoints by Claim 4.5 (i) (if $i_0 \in \text{col}(M)$) or Claim 4.5 (ii) (if $i_0 \in \text{col}(P^1 - A)$). It is routine to check that $|\mathcal{C}_1 \cup \mathcal{C}'_{bad} \setminus \text{col}(\tilde{P}^4)| - (|B \setminus V(\tilde{P}^4)| - |A \setminus V(\tilde{P}^4)|)$ is a positive odd integer. Let $P^4 := \tilde{P}^4$ if $i_0 \in \text{col}(P^1 - A) \cup \text{col}(M)$. Therefore, it suffices to consider $i_0 \notin \text{col}(P^4)$.

• $w_1, w_2 \notin V_{bad}$. Then $w_1, w_2 \notin V(P^4)$. Connect w_2 with the endpoint u_1 of P^4 by Claim 4.5 (ii). Denote by Q_0 the resulted rainbow path (see Figure 6).

- Exactly one of w_1, w_2 lies in V_{bad} . Assume, without loss of generality, that $w_1 \notin V_{bad}$ and $w_2 \in V_{bad}$. Hence $w_1 \notin V(P^4)$ and $w_2 \in V_{bad} \cap A$. Thus, there exists a rainbow subpath $u_1 P^4 w_2^1 w_2^2 w_2$ of P^4 with $w_2^1 \in A \setminus V_{bad}$, $w_2^2 \in B \setminus V_{bad}$. Replace $w_2^1 w_2^2 w_2$ with $w_1 w_2$ in P^4 , and connect w_1 with the vertex in $N_{P^4}(w_2^1) \setminus \{w_2^2\}$ (if exists) by Claim 4.5 (i). Let Q_1 be the resulted rainbow path (see Figure 6).

- $w_1, w_2 \in V_{bad}$. By rearranging, one may assume w_1 and w_2 is connected by a rainbow path $w_1 w_1^1 w_2^1 w_2^2 w_2$ in P^4 , where $w_1^1, w_2^2 \in B \setminus V_{bad}$ and $w_2^1 \in A \setminus V_{bad}$. Replace $w_1 w_1^1 w_2^1 w_2^2 w_2$ with $w_1 w_2$ in P^4 . Denote by Q_2 the resulted rainbow path (see Figure 6).




FIGURE 6.

In each of the above cases, we get a rainbow path Q_i ($i \in \{0, 1, 2\}$) such that $|\tilde{\mathcal{C}}| - ||B \setminus V(Q_i)| - |A \setminus V(Q_i)||$ is a nonnegative even integer. It follows from Claim 4.7 that \mathcal{G} contains a transversal Hamilton cycle, a contradiction.

(ii) Suppose that $E(G_{i_1}[A, B]) \neq \emptyset$ for some $i_1 \in \tilde{\mathcal{C}}$. Choose $z_1 z_2 \in E(G_{i_1}[A, B])$ with $z_1 \in A$ and $z_2 \in B$. Notice that $col(P^1 - A) \subseteq \mathcal{C}_1 \cup \mathcal{C}'_{bad}$. If $z_i \in V_{bad} \cap Y$ for $i \in [2]$, then P^4 can be written as $u_1 P^4 z_i^0 z_i^1 z_i z_i^2 P^4 v_4$, where $z_i^0 \in Y \setminus V_{bad}$ and $z_i^1, z_i^2 \in Z \setminus V_{bad}$; if $z_2 \in V(P^1) \cap B$, then P^4 can be written as $u_1 P^4 z_2 z_2' P^4 v_4$, where $z_2 z_2' \in E(P^1) \cap E(G_{i_2})$ for some $i_2 \in \mathcal{C}$. Clearly, $i_2 \in \mathcal{C}_2$ if $z_2' \in A$ and $i_2 \in (\mathcal{C}_1 \cup \mathcal{C}'_{bad}) \setminus \tilde{\mathcal{C}}$ if $z_2' \in B$. Notice that $V(P^1) \cap B \cap V_{bad} = \emptyset$. Hence $z_2' \notin V_{bad}$.

- $z_1, z_2 \notin (V(P^1) \cap B) \cup V_{bad}$. Then $z_1, z_2 \notin V(P^4)$. Connect z_2 with the endpoint u_1 of P^4 by Claim 4.5 (i). Let Q_3 be the resulted rainbow path and $\tilde{\mathcal{C}}_3 = \tilde{\mathcal{C}} \setminus \{i_1\}$.

- Exactly one of z_1, z_2 lies in $(V(P^1) \cap B) \cup V_{bad}$. Without loss of generality, assume that $z_1 \notin (V(P^1) \cap B) \cup V_{bad}$ and $z_2 \in (V(P^1) \cap B) \cup V_{bad}$ (the another case is more easier since $z_1 \notin V(P^1) \cap B$). Hence $z_1 \notin V(P^4)$. If $z_2 \in V(P^1) \cap B$, then connect $u_1 P^1 z_2 z_1$ with $z_2' P^4 v_4$ into a single rainbow path by Claim 4.5 (i)-(ii). Let Q_4 be the resulted rainbow path, let $\tilde{\mathcal{C}}_4 = \tilde{\mathcal{C}} \setminus \{i_1\}$ if $z_2' \in A$ and $\tilde{\mathcal{C}}_4 = (\tilde{\mathcal{C}} \setminus \{i_1\}) \cup \{i_2\}$ if $z_2' \in B$. If $z_2 \in V_{bad}$, then connect $u_1 P^4 z_2^1$ with $z_1 z_2 z_2' P^4 v_4$ into a single rainbow path by Claim 4.5 (ii). Let Q_5 be the resulted rainbow path and $\tilde{\mathcal{C}}_5 = \tilde{\mathcal{C}} \setminus \{i_1\}$.

- $z_1, z_2 \in (V(P^1) \cap B) \cup V_{bad}$. Then $z_1 \in V_{bad}$. If $z_2 \in V_{bad}$, by rearranging, one may assume z_1 and z_2 is connected by a rainbow path $z_1 z_1^2 z_1^3 z_2^3 z_2^1 z_2$ in P^4 , where $z_1^2, z_2^3 \in B \setminus V_{bad}$ and $z_1^3, z_2^1 \in A \setminus V_{bad}$. Replace such a rainbow path with $z_1 z_2$ in P^4 . Let Q_6 be the resulted rainbow path and $\tilde{\mathcal{C}}_6 = \tilde{\mathcal{C}} \setminus \{i_1\}$. If $z_2 \in V(P^1) \cap B$, then delete vertices z_1^0, z_1^1, z_1, z_1^2 from P^4 and connect the resulted two components by Claim 4.5 (i). Denote the resulted rainbow path by \tilde{P}^4 . Then delete the edge $z_2 z_2'$ from \tilde{P}^4 . Applying Claim 4.5 (i)-(ii) again to connect $u_1 \tilde{P}^4 z_2 z_1 z_1^1$ with $z_2' \tilde{P}^4 v_4$ into a single rainbow path Q_7 . Let $\tilde{\mathcal{C}}_7 = \tilde{\mathcal{C}} \setminus \{i_1\}$ if $z_2' \in A$ and $\tilde{\mathcal{C}}_7 = (\tilde{\mathcal{C}} \setminus \{i_1\}) \cup \{i_2\}$ if $z_2' \in B$.

In each of the above cases, we get a rainbow path Q_i ($i \in [3, 7]$) such that $|\tilde{\mathcal{C}}_i| - ||B \setminus V(Q_i)| - |A \setminus V(Q_i)||$ is a nonnegative even integer. Applying Claim 4.7 yields that \mathcal{G} contains a transversal Hamilton cycle, a contradiction. Hence $E(G_i[A, B]) = \emptyset$ for all $i \in \tilde{\mathcal{C}}$. \square

If $|\tilde{\mathcal{C}}| = 0$, then n is odd. By a similar discussion as the proof of Claim 4.8, we can show that $E(G_i[B]) = E(G_i[V(P^1) \cap B])$ for all $i \in \mathcal{C}$. Then $|A| \geq \frac{n-1}{2}$. Recall that $|B| \geq |A|$. Hence $|A| = \frac{n-1}{2}$ and $|B| = \frac{n+1}{2}$. This implies that P^1 is a null graph. Thus, $G_i[B] = \emptyset$ for all $i \in \mathcal{C}$. Therefore, \mathcal{G} is the half-split graph collection and it does not contain transversal Hamilton cycles, as desired.

If $|\tilde{\mathcal{C}}| \geq 1$, then based on Claim 4.8 and the fact that $\delta(\mathcal{G}) \geq \frac{n}{2} - 1$, we know n is even, $|A| = \frac{n}{2}$ and $|B| = \frac{n}{2}$, and $G_i[A] = \emptyset$ for all but at most $\eta n + 3\delta n$ colors in \mathcal{C} . By Theorem 4.3, we know \mathcal{G} must be one of the collections described in Theorem 1.6, as desired.

Case 2. $|\mathcal{C}_2| < \eta n$.

In this case, each vertex in $Y \setminus (X \cup X_Y^1)$ belongs to Y_i for at least $(1 - 10\sqrt{\delta})|\mathcal{C}_1|$ colors $i \in \mathcal{C}_1$. By a similar discussion as Claim 4.6, we get the following claim.

Claim 4.9. *Assume $X = \{x_1, \dots, x_s\}$ with $s \leq \sqrt{\epsilon n}$. If $x_i \in X \cap Y$, then there exist $c_i^1, c_i^2, c_i^3, c_i^4 \in \mathcal{C}_1 \setminus \{c_\ell^1, c_\ell^2, c_\ell^3, c_\ell^4 : \ell \in [i-1]\}$ and $x_i^1, x_i^2, x_i^3, x_i^4 \in Y \setminus (V_{bad} \cup \{x_\ell^1, x_\ell^2, x_\ell^3, x_\ell^4 : \ell \in [i-1]\})$ such that $x_i x_i^k \in E(G_{c_i^k})$ for all $k \in [4]$.*

By Claim 4.9 and a similar discussion as Steps 1-3 in Case 1, the following hold:

- (C1) In $\{G_i[A] \cup G_i[B] : i \in \mathcal{C}_1\}$, there are $|V_{bad}|$ disjoint rainbow P_3 with centers in V_{bad} and endpoints in $(A \cup B) \setminus V_{bad}$. Let \mathbf{P} be a set consisting of those rainbow P_3 .
- (C2) For colors in \mathcal{C}_{bad} , there exists a maximal rainbow matching, say M , inside $\{G_i[A \setminus V(\mathbf{P})] \cup G_i[B \setminus V(\mathbf{P})] : i \in \mathcal{C}_{bad}\}$ such that for each edge in $M[Y]$ with color j we have $G_j[Y \setminus V(\mathbf{P} \cup M)]$ contains a 2-matching. Denote $\mathcal{C}'_{bad} := \mathcal{C}_{bad} \setminus \text{col}(M)$. Then G_j is $(3\sqrt{\delta}, K_{\lfloor \frac{n}{2} \rfloor, \lceil \frac{n}{2} \rceil})$ -extremal for all $j \in \mathcal{C}'_{bad}$.
- (C3) Let $q = |\mathcal{C}_2 \cup \mathcal{C}'_{bad}|$. In the graph collection $\{G_i[A \setminus V(\mathbf{P} \cup M), B \setminus V(\mathbf{P} \cup M)] : i \in \mathcal{C}_2 \cup \mathcal{C}'_{bad}\}$, there exists a transversal matching $M' = \{u_1 v_1, u_2 v_2, \dots, u_q v_q\}$.
- (C4) We can assume a vertex $v \notin V(\mathbf{P} \cup M \cup M')$ unless $v \in V_{bad}$; and a color $c \notin \text{col}(\mathbf{P})$ if $c \in \mathcal{C}_1$.

Next, we give the following claim.

Claim 4.10. *Assume that $Q_1 = z_1 \dots z_s$ and $Q_2 = z'_1 \dots z'_t$ are two disjoint rainbow paths inside \mathcal{G} such that $z_1, z'_1 \in A \setminus V_{bad}$, $z_s, z'_t \in B \setminus V_{bad}$ and $s + t < 4\eta n$. If $V_{bad} \subseteq V(Q_1 \cup Q_2)$ and $\mathcal{C}_2 \cup \mathcal{C}_{bad} \subseteq \text{col}(Q_1 \cup Q_2)$, then \mathcal{G} contains a transversal Hamilton cycle.*

Proof of Claim 4.10. Note that $z_1, z'_1 \in A \setminus V_{bad}$ and $z_s, z'_t \in B \setminus V_{bad}$. There exist $i_1, i_2, i_3, i_4 \in \mathcal{C}_1 \setminus \text{col}(Q_1 \cup Q_2)$ such that $z_1 \in A_{i_1}$, $z'_1 \in A_{i_2}$, $z_s \in B_{i_3}$ and $z'_t \in B_{i_4}$. Split $\mathcal{C}_1 \setminus (\text{col}(Q_1 \cup Q_2) \cup \{i_1, i_2, i_3, i_4\})$ into two parts $\mathcal{C}_a \cup \mathcal{C}_b$, where $|\mathcal{C}_a| = |A \setminus V(Q_1 \cup Q_2)| - 1$ and $|\mathcal{C}_b| = |B \setminus V(Q_1 \cup Q_2)| - 1$. Let $I_1 \cup I_2$ and $J_1 \cup J_2$ be equitable partitions of $A \setminus V(Q_1 \cup Q_2)$ and $B \setminus V(Q_1 \cup Q_2)$ with $|I_1| \geq |I_2|$ and $|J_1| \geq |J_2|$, respectively. Define

$$\mathcal{G}_a = \{G_i[I_1, I_2] : i \in \mathcal{C}_a\} \text{ and } \mathcal{G}_b = \{G_i[J_1, J_2] : i \in \mathcal{C}_b\}.$$

Notice that

$$|Y \setminus V(Q_1 \cup Q_2)| \geq \frac{n-r}{2} - |V(Q_1 \cup Q_2)| > \frac{n}{2} - 5\eta n,$$

and for $j \in [2]$ we have

$$\begin{aligned}
|N_{G_{i_1}}(z_1, I_j)| &\geq d_{G_{i_1}}(z_1, A \setminus V(Q_1 \cup Q_2)) - |I_{3-j}| \\
&\geq d_{G_{i_1}}(z_1, A_{i_1}) - |A_1 \Delta A_{i_1}| - |X_A \setminus X_A^1| - |V(Q_1 \cup Q_2)| - |I_{3-j}| \\
&\geq \left(\frac{1}{2} - 3\sqrt{\delta}\right)n - (2\delta + 4\delta + 8\eta)n - \lceil \frac{n+r}{4} \rceil > \frac{n}{4} - 5\eta n.
\end{aligned}$$

Similarly, we have $|N_{G_{i_2}}(z'_1, I_j)|, |N_{G_{i_3}}(z_s, J_j)|, |N_{G_{i_4}}(z'_t, J_j)| > \frac{n}{4} - 5\eta n$ for each $j \in [2]$. By Claim 4.4, there exists a transversal path Q_a inside \mathcal{G}_a with two endpoints $x_a \in N_{G_{i_1}}(z_1, I_1)$ and $y_a \in N_{G_{i_2}}(z'_1, I)$; and a transversal path Q_b inside \mathcal{G}_b with two endpoints $x_b \in N_{G_{i_3}}(z_s, J_1)$ and $y_b \in N_{G_{i_4}}(z'_t, J)$ (here $y_a \in I_2$ (resp. $y_b \in J_2$) if and only if $|I_1| = |I_2|$ (resp. $|J_1| = |J_2|$) is even). Thus, $z_1 Q_1 z_s x_b Q_b y_b z'_t Q_2 z'_1 y_a Q_a x_a z_1$ is a transversal Hamilton cycle inside \mathcal{G} (see Figure 7). \square




FIGURE 7.




FIGURE 8.

Now, we proceed by considering the value of q .

Subcase 2.1. q is even and $q \geq 2$.

Using Claim 4.5 (iii) to connect all rainbow paths in \mathbf{P} , M and $M' - u_q v_q$ into a single rainbow path P , whose length is at most $(4\sqrt{\epsilon} + 2\sqrt{\delta} + 9\delta + 3\eta)n$ and endpoints are $x \in A \setminus V_{bad}$, $y \in B \setminus V_{bad}$ (see Figure 8). Thus, P and $u_q v_q$ are two disjoint rainbow paths satisfying all conditions in Claim 4.10. Thus, \mathcal{G} contains a transversal Hamilton cycle, a contradiction.

Subcase 2.2. q is odd.

Suppose that $E(G_{i_1}[A, B]) \neq \emptyset$ for some $i_1 \in \mathcal{C}_1 \cup \text{col}(M)$. Choose $w_1 w_2 \in E(G_{i_1}[A, B])$ with $w_1 \in A$ and $w_2 \in B$. Based on **(C1)**, the edge $w_1 w_2$ can be extended to a rainbow path of length at most 3, say Q_1 , which has endpoints in $A \setminus V_{bad}$ and $B \setminus V_{bad}$ respectively. Next, by using Claim 4.5 (iii) and avoiding vertices and colors in Q_1 , one may connect all rainbow P_3 in \mathbf{P} with centers not in $\{w_1, w_2\}$, all rainbow edges in M except the possible edge with color i_1 and all rainbow edges in M' into a single rainbow path P^1 , whose length is at most $(4\sqrt{\epsilon} + 2\sqrt{\delta} + 9\delta + 3\eta)n$ and endpoints are $x_1 \in A \setminus V_{bad}$, $y_1 \in B \setminus V_{bad}$. Then P^1 and Q_1 satisfy all conditions in Claim 4.10. Therefore, \mathcal{G} contains a transversal Hamilton cycle, a contradiction. Hence $E(G_i[A, B]) = \emptyset$ for all $i \in \mathcal{C}_1 \cup \text{col}(M)$. Recall that $\delta(\mathcal{G}) \geq \frac{n}{2} - 1$. Then n is even and $G_i[A] = G_i[B] = K_{\frac{n}{2}}$ for all $i \in \mathcal{C}_1 \cup \text{col}(M)$.

If $q = 1$, then it is easy to see that \mathcal{G} contains no transversal Hamilton cycle, as desired. Next, we consider $q \geq 3$. Suppose that there exists an $i_2 \in \mathcal{C}_2 \cup \mathcal{C}'_{bad}$ such that $E(G_{i_2}[A]) \cup E(G_{i_2}[B]) \neq \emptyset$. Without loss of generality, assume that $z_1 z_2 \in E(G_{i_2}[A])$ and $u_1 v_1$ is the edge in M' with color i_2 . Based on **(C1)**, we can extend $z_1 z_2$ to a rainbow path Q_2 with two endpoints in $A \setminus V_{bad}$ and length at most 3. Using Claim 4.5 (iii) to connect Q_2 , all rainbow P_3 in \mathbf{P} with centers not in $\{z_1, z_2\}$, all rainbow edges in M and all rainbow edges in $M' - \{u_1 v_1, u_q v_q\}$ into a

single rainbow path P^2 , whose length is at most $(4\sqrt{\epsilon} + 2\sqrt{\delta} + 9\delta + 3\eta)n + 6$ and endpoints are $x_2 \in A \setminus V_{bad}$, $y_2 \in B \setminus V_{bad}$. Clearly, P^2 and $u_q v_q$ are two rainbow paths satisfying all conditions in Claim 4.10. Therefore, \mathcal{G} contains a transversal Hamilton cycle, a contradiction. Thus, for each $i \in \mathcal{C}_2 \cup \mathcal{C}'_{bad}$, $E(G_i[A]) \cup E(G_i[B]) = \emptyset$ and so G_i is a subgraph of $K_{\frac{n}{2}, \frac{n}{2}}$. It follows that \mathcal{G} is a spanning collection of \mathcal{H}_{n-q}^q with odd q .

We claim that \mathcal{H}_{n-q}^q contains no transversal Hamilton cycles if n is even and q is odd. Suppose not, let C be a transversal Hamilton cycle and give it an arbitrary direction to make it a directed cycle. We say an edge of C is 1-type if it comes from some G_i with $i \in \mathcal{C}_1 \cup \text{col}(M)$ and 2-type if it comes from some G_i with $i \in \mathcal{C}_2 \cup \mathcal{C}'_{bad}$. Note that the number of 2-type edges directed from A to B is equal to the number of 2-type edges directed from B to A . Hence, the total number of 2-type edges in C is even, which implies that q is even, a contradiction.

Subcase 2.3. $q = 0$.

Suppose that $\{G_i[A, B] : i \in \mathcal{C}\}$ contains a rainbow 2-matching. Let $\{w_1 w_2, z_1 z_2\}$ be such a rainbow 2-matching with colors c_1, c_2 . Based on **(C1)**, we can extend $w_1 w_2$ and $z_1 z_2$ to two disjoint rainbow paths Q_3 and Q_4 with length at most 3, each of which has one endpoint in $A \setminus V_{bad}$ and the other in $B \setminus V_{bad}$. By applying Claim 4.5 (iii) and avoiding vertices and colors in Q_4 , one may connect Q_3 , all rainbow P_3 in \mathbf{P} with centers outside $\{w_1, w_2, z_1, z_2\}$ and all rainbow edges in M with colors outside $\{c_1, c_2\}$ into a single rainbow path P^3 , whose length is at most $(4\sqrt{\epsilon} + 2\sqrt{\delta} + 9\delta + 3\eta)n + 6$ and endpoints are $x_3 \in A \setminus V_{bad}$, $y_3 \in B \setminus V_{bad}$. Clearly, P^3 and Q_4 are two rainbow paths satisfying all conditions in Claim 4.10. Therefore, \mathcal{G} contains a transversal Hamilton cycle, a contradiction. Hence $\{G_i[A, B] : i \in \mathcal{C}\}$ contains no rainbow 2-matching. Finally, we finish the proof of this case by proving the following claim.

Claim 4.11. *If $\mathcal{G}[A, B]$ has no rainbow 2-matching, then \mathcal{G} does not contain transversal Hamilton cycles. Furthermore, \mathcal{G} must be a graph collection described in Theorem 1.6 (i)(b) or Theorem 1.6 (ii) (c).*

Proof of Claim 4.11. Notice that each transversal Hamilton cycle contains at least 2 disjoint rainbow edges in $\mathcal{G}[A, B]$. Hence \mathcal{G} contains no transversal Hamilton cycles. Next, we characterize the structure of \mathcal{G} .

Assume that each vertex of A has at least one neighbor in B for all $G_i \in \mathcal{G}$. Since $\mathcal{G}[A, B]$ contains no rainbow 2-matching, there exists a fixed vertex $w \in B$ such that $N_{G_i}(v, B) = \{w\}$ for all $i \in \mathcal{C}$ and all $v \in A$. Therefore, $|A| \geq \lceil \frac{n}{2} - 1 \rceil$. If n is odd, then $d_{G_i}(w) = n - 1$ and $G_i[V \setminus \{w\}] = G_i[A] \cup G_i[B \setminus \{w\}] = 2K_{\frac{n-1}{2}}$ for all $i \in \mathcal{C}$, as desired. If n is even, then there exists an equitable partition $A' \cup B'$ of V and a vertex $w' \in A'$ such that $G_i[A'] = K_{\frac{n}{2}}$ and $E(G_i[A', B']) = E(G_i[\{w'\}, B'])$ for all $i \in \mathcal{C}$, as desired.

In what follows, it suffices to consider that there is a vertex $v \in A$ and a color $i \in \mathcal{C}$ such that $N_{G_i}(v, B) = \emptyset$. Hence $|B| \geq |A| \geq \lceil \frac{n}{2} - 1 \rceil + 1$. Therefore, n is even and $|A| = |B| = \frac{n}{2}$.

- There exists a color $j_1 \in \mathcal{C}$ and a vertex $v_1 \in A$ such that $|N_{G_{j_1}}(v_1) \cap B| \geq 2$. Since \mathcal{G} contains no rainbow 2-matching, we know $G_i[A \setminus \{v_1\}, B] = \emptyset$ for all $i \in \mathcal{C} \setminus \{j_1\}$. This implies that $E(G_i[A, B]) = E(G_i[\{v_1\}, B])$ for all $i \in \mathcal{C} \setminus \{j_1\}$. If $G_i[\{v_1\}, B] = \emptyset$ for all $i \in \mathcal{C} \setminus \{j_1\}$, then $G_i[A] = G_i[B] = K_{\frac{n}{2}}$ for all $i \in \mathcal{C} \setminus \{j_1\}$, as desired. If there exists some $j_2 \in \mathcal{C} \setminus \{j_1\}$ such that $G_{j_2}[\{v_1\}, B] \neq \emptyset$, then let $v_1 w \in E(G_{j_2}[\{v_1\}, B])$. Therefore, $N_{G_{j_1}}(v, B) \subseteq \{w\}$ for each $v \in A \setminus \{v_1\}$.

If $N_{G_{j_1}}(v, B) = \emptyset$ for all $v \in A \setminus \{v_1\}$, then $E(G_i[A, B]) = E(G_i[\{v_1\}, B])$ and $E(G_i[A]) = K_{\frac{n}{2}}$ for all $i \in \mathcal{C}$, as desired. If there exists $v_2 \in A \setminus \{v_1\}$ such that $N_{G_{j_1}}(v_2, B) = \{w\}$, then

$E(G_i[\{v_1\}, B]) \subseteq \{v_1w\}$ for all $i \in \mathcal{C} \setminus \{j_1\}$. That is, $E(G_{j_1}[A, B]) \subseteq \{v_1v : v \in B\} \cup \{vw : v \in A\}$, $E(G_i[A, B]) \subseteq \{v_1w\}$ and $G_i[A] = G_i[B] = K_{\frac{n}{2}}$ for all $i \in \mathcal{C} \setminus \{j_1\}$, as desired.

- $|N_{G_i}(v, B)| \leq 1$ for all $i \in \mathcal{C}$ and all $v \in A$. By symmetry, one may assume that $|N_{G_i}(v, A)| \leq 1$ for all $i \in \mathcal{C}$ and all $v \in B$. Let $|E(G_{j_3}[A, B])| = \max\{|E(G_j[A, B])| : j \in \mathcal{C}\}$. If $|E(G_{j_3}[A, B])| \geq 3$, then $G_i = G_i[A] \cup G_i[B] = 2K_{\frac{n}{2}}$ for all $i \in \mathcal{C} \setminus \{j_3\}$, as desired.

If $|E(G_{j_3}[A, B])| = 2$, then assume $E(G_{j_3}[A, B]) = \{v_1w_1, v_2w_2\}$, where v_1, w_1, v_2, w_2 are pairwise distinct. This implies $G_{j_3}[A] \in \{K_{\frac{n}{2}} - v_1v_2, K_{\frac{n}{2}}\}$ and $G_{j_3}[B] \in \{K_{\frac{n}{2}} - w_1w_2, K_{\frac{n}{2}}\}$. Recall that \mathcal{G} contains no rainbow 2-matching. Then either $E(G_{j_4}[A, B]) = \{v_1w_2, v_2w_1\}$ for a unique $j_4 \in \mathcal{C} \setminus \{j_3\}$ and $G_i[A, B] = \emptyset$ for all $i \in \mathcal{C} \setminus \{j_3, j_4\}$, or $E(G_i[A, B]) \subseteq \{v_1w_2\}$ for all $i \in \mathcal{C} \setminus \{j_3\}$. The former case implies $E(G_{j_4}[A, B]) = \{v_1w_2, v_2w_1\}$, $G_{j_4}[A] \in \{K_{\frac{n}{2}} - v_1v_2, K_{\frac{n}{2}}\}$, $G_{j_4}[B] \in \{K_{\frac{n}{2}} - w_1w_2, K_{\frac{n}{2}}\}$ and $G_i = G_i[A] \cup G_i[B] = 2K_{\frac{n}{2}}$ for all $i \in \mathcal{C} \setminus \{j_3, j_4\}$ (see Figure 1 (a)), as desired. The latter case implies $E(G_i[A, B]) \subseteq \{v_1w_2\}$ and $G_i[A] = G_i[B] = K_{\frac{n}{2}}$ for all $i \in \mathcal{C} \setminus \{j_3\}$ (see Figure 1 (b)), as desired.

If $|E(G_{j_3}[A, B])| \leq 1$, then $G_i[A] = G_i[B] = K_{\frac{n}{2}}$ for all $i \in \mathcal{C}$ and all edges in $\{G_i[A, B] : i \in \mathcal{C}\}$ must have a common vertex, as desired.

This completes the proof of Claim 4.11. \square

Case 3. $|\mathcal{C}_1| \geq \eta n$ and $|\mathcal{C}_2| \geq \eta n$.

In this case, each vertex in $Y \setminus V_{bad}$ lies in Y_i for at least $(1 - 10\sqrt{\delta})|\mathcal{C}_1 \cup \mathcal{C}_2|$ colors $i \in \mathcal{C}_1 \cup \mathcal{C}_2$. Denote $\cup_{k \in [2]}(X_A^k \cup X_B^k) = \{v_1, v_2, \dots, v_t\}$. Notice that $t < \frac{1}{2}\sqrt{\delta}n$. By the definition of X_Y^k , we obtain that for each $i \in [t]$, there exist $c_{4i-3}, c_{4i-2}, c_{4i-1}, c_{4i} \in (\mathcal{C}_1 \cup \mathcal{C}_2) \setminus \{c_1, c_2, \dots, c_{4i-5}, c_{4i-4}\}$ and $v_i^1, v_i^2, v_i^3, v_i^4 \in (A \cup B) \setminus (V_{bad} \cup \{v_\ell^1, v_\ell^2, v_\ell^3, v_\ell^4 : \ell \in [i-1]\})$ such that

- if $v_i \in X_Y^1$, then $c_{4i-4+j} \in \mathcal{C}_1$, $v_i^j \in Y \cap Y_{c_{4i-4+j}}$ and $v_i v_i^j \in E(G_{c_{4i-4+j}})$ for all $j \in [4]$,
- if $v_i \in X_Y^2$, then $c_{4i-4+j} \in \mathcal{C}_2$, $v_i^j \in Z \cap Z_{c_{4i-4+j}}$ and $v_i v_i^j \in E(G_{c_{4i-4+j}})$ for all $j \in [4]$.

Therefore, for each $i \in [t]$, there exists a rainbow path $v_i^1 v_i v_i^2$ with colors c_{4i-3} and c_{4i-2} . Connect those rainbow paths by using colors in \mathcal{C}_2 and Claim 4.5 (i)-(ii), one may get a rainbow path P^1 with endpoints $u_1 \in A \setminus V_{bad}$ and $w_1 \in B \setminus V_{bad}$, whose length is at most $4t \leq 2\sqrt{\delta}n$.

By a similar discussion as Claim 4.6, we obtain the following claim.

Claim 4.12. *Assume $X = \{x_1, \dots, x_s\}$ with $s \leq \sqrt{\epsilon}n$. Then for each $i \in [s]$, there exist $c_i^1, c_i^2, c_i^3, c_i^4 \in \mathcal{C}_2 \setminus (\text{col}(P^1) \cup \{c_\ell^1, c_\ell^2, c_\ell^3, c_\ell^4 : \ell \in [i-1]\})$ and $x_i^1, x_i^2, x_i^3, x_i^4 \in V \setminus (V(P^1) \cup V_{bad} \cup \{x_\ell^1, x_\ell^2, x_\ell^3, x_\ell^4 : \ell \in [i-1]\})$ such that: if $x_i \in Y$, then $x_i x_i^j \in E(G_{c_i^j})$, and either $x_i^j \in Y$ and $c_i^j \in \mathcal{C}_1$ for all $j \in [4]$, or $x_i^j \in Z$ and $c_i^j \in \mathcal{C}_2$ for all $j \in [4]$.*

By Claim 4.12, there exists a rainbow path $x_i^1 x_i x_i^2$ with colors c_i^1, c_i^2 for each $i \in [s]$. Connect all those rainbow paths and P^1 by Claim 4.5 (i)-(ii), we obtain a rainbow path P^2 with endpoints $u_1 \in A \setminus V_{bad}$ and $w_2 \in B \setminus V_{bad}$, whose length is at most $|E(P^1)| + 4\sqrt{\epsilon}n + 2 \leq (2\sqrt{\delta} + 4\sqrt{\epsilon})n + 2$.

Choose a maximal rainbow matching, say M , inside $\{G_i[A \setminus V(P^2)] \cup G_i[B \setminus V(P^2)] : i \in \mathcal{C}_{bad}\}$ such that for each edge in $M[Y]$ with color j we have $G_j[Y \setminus V(P^2)]$ contains a 2-matching. Denote $\mathcal{C}'_{bad} := \mathcal{C}_{bad} \setminus \text{col}(M)$. Note that G_j is $(3\sqrt{\delta}, K_{\lfloor \frac{n}{2} \rfloor, \lfloor \frac{n}{2} \rfloor})$ -extremal for all $j \in \mathcal{C}'_{bad}$. Hence, $\{G_i[A, B] : i \in \mathcal{C}'_{bad}\}$ contains a transversal matching, say M' . Connect P^2 and all rainbow edges in $M \cup M'$ by Claim 4.5 (i)-(ii), we obtain a rainbow path P^3 with endpoints $u_1 \in A \setminus V_{bad}$ and $w_3 \in B \setminus V_{bad}$, whose length is at most $|E(P^2)| + 12\delta n \leq (2\sqrt{\delta} + 4\sqrt{\epsilon} + 12\delta)n + 2$.

Since $u_1, w_3 \notin V_{bad}$, there exist $i_0, i_1 \in \mathcal{C}_1 \setminus \text{col}(P^3)$ such that $u_1 \in A_{i_0}$ and $w_3 \in B_{i_1}$. By the construction of P^3 , one may assume that $v \notin V(P^3)$ unless $v \in V_{bad}$, and $c \notin \text{col}(P^3)$ unless $c \in \mathcal{C}_{bad}$. In what follows, we proceed by considering the parity of $|\mathcal{C}_2 \setminus \text{col}(P^3)|$.

Subcase 3.1. $|\mathcal{C}_2 \setminus \text{col}(P^3)|$ is odd.

Split $A \setminus V(P^3)$ into I_1 and I_2 such that $|I_1| = \frac{|\mathcal{C}_2 \setminus \text{col}(P^3)| + 1}{2}$, and split $B \setminus V(P^3)$ into J_1 and J_2 such that $|J_1| = \frac{|\mathcal{C}_2 \setminus \text{col}(P^3)| + 1}{2}$. Let $\mathcal{G}_1 = \{G_i[I_1, J_1] : i \in \mathcal{C}_2 \setminus \text{col}(P^3)\}$. By Claim 4.4, there exists a transversal path Q inside \mathcal{G}_1 with endpoints $x \in I_1$ and $y \in J_1$. Since $x, y \notin V_{bad}$, there exist $j_0, j_1 \in \mathcal{C}_1$ such that $x \in A_{j_0}$ and $y \in B_{j_1}$.

Suppose that $|I_2| = |A \setminus V(P^3)| - |I_1| = k_1$ and $|J_2| = |B \setminus V(P^3)| - |J_1| = k_2$. We further split I_2 into I'_2 and I''_2 such that $|I'_2| = \lceil \frac{k_1}{2} \rceil$, split J_2 into J'_2 and J''_2 such that $|J'_2| = \lceil \frac{k_2}{2} \rceil$. Then split the color set $\mathcal{C}_1 \setminus (\text{col}(P^3) \cup \{j_0, j_1\})$ into two subsets \mathcal{C}_a and \mathcal{C}_b such that $|\mathcal{C}_a| = k_1 - 1$ and $|\mathcal{C}_b| = k_2 - 1$. Let $\mathcal{G}_a = \{G_i[I'_2, I''_2] : i \in \mathcal{C}_a\}$ and $\mathcal{G}_b = \{G_i[J'_2, J''_2] : i \in \mathcal{C}_b\}$. Applying Claim 4.4 again yields that there is a transversal path Q_a inside \mathcal{G}_a with endpoints $x_a \in N_{G_{i_0}}(u_1, I'_2)$, $y_a \in N_{G_{j_0}}(x, I_2)$; and a transversal path Q_b inside \mathcal{G}_b with endpoints $x_b \in N_{G_{i_1}}(w_3, J'_2)$, $y_b \in N_{G_{j_1}}(y, J_2)$. Thus, $u_1 P^3 w_3 x_b Q_b y_b y Q x y_a Q_a x_a u_1$ is a transversal Hamilton cycle in \mathcal{G} (see Figure 9), a contradiction.




FIGURE 9.

Subcase 3.2. $|\mathcal{C}_2 \setminus \text{col}(P^3)|$ is even.

In this subcase, we first prove the following two claims.

Claim 4.13. Assume that $Q_1 = z_1 \dots z_k$ and $Q_2 = z'_1 \dots z'_t$ are two disjoint rainbow paths inside \mathcal{G} such that $z_1, z'_1 \in A \setminus V_{bad}$, $z_k, z'_t \in B \setminus V_{bad}$ and each of them has length at most $3\sqrt{\delta}n$. If $V_{bad} \subseteq V(Q_1 \cup Q_2)$, $\mathcal{C}_{bad} \subseteq \text{col}(Q_1 \cup Q_2)$ and $|\mathcal{C}_2 \setminus \text{col}(Q_1 \cup Q_2)|$ is even, then \mathcal{G} contains a transversal Hamilton cycle.

Proof of Claim 4.13. Note that $z_1, z'_1 \in A \setminus V_{bad}$ and $z_k, z'_t \in B \setminus V_{bad}$. There exist $l_1, l_2, l_3 \in \mathcal{C}_1 \setminus \text{col}(Q_1 \cup Q_2)$ and $l_4 \in \mathcal{C}_2 \setminus \text{col}(Q_1 \cup Q_2)$ such that $z_1 \in A_{l_1}$, $z_k \in B_{l_2}$, $z'_1 \in A_{l_4}$ and $z'_t \in B_{l_3}$.

Split $A \setminus V(Q_1 \cup Q_2)$ into I_1 and I_2 such that $|I_1| = \frac{|\mathcal{C}_2 \setminus \text{col}(Q_1 \cup Q_2)|}{2}$, and split $B \setminus V(Q_1 \cup Q_2)$ into J_1 and J_2 such that $|J_1| = \frac{|\mathcal{C}_2 \setminus \text{col}(Q_1 \cup Q_2)|}{2}$. Let $\mathcal{G}_1 = \{G_i[I_1, J_1] : i \in \mathcal{C}_2 \setminus (\text{col}(Q_1 \cup Q_2) \cup \{l_4\})\}$. By Claim 4.4, there exists a transversal path Q inside \mathcal{G}_1 with endpoints $z''_1 \in N_{G_{l_4}}(z'_1) \cap J_1$ and $y \in I_1$. Since $y \notin V_{bad}$, there exists $l_5 \in \mathcal{C}_1 \setminus (\text{col}(Q_1 \cup Q_2) \cup \{l_1, l_2, l_3\})$ such that $y \in A_{l_5}$.

Assume that $|I_2| = k_1$ and $|J_2| = k_2$. We further split I_2 into I'_2 and I''_2 such that $|I'_2| = \lceil \frac{k_1}{2} \rceil$, and split J_2 into J'_2 and J''_2 such that $|J'_2| = \lceil \frac{k_2}{2} \rceil$. Split the color set $\mathcal{C}_1 \setminus (\text{col}(Q_1 \cup Q_2) \cup \{l_1, l_2, l_3, l_5\})$ into two subsets \mathcal{C}_a and \mathcal{C}_b such that $|\mathcal{C}_a| = k_1 - 1$ and $|\mathcal{C}_b| = k_2 - 1$. Let $\mathcal{G}_a = \{G_i[I'_2, I''_2] : i \in \mathcal{C}_a\}$ and $\mathcal{G}_b = \{G_i[J'_2, J''_2] : i \in \mathcal{C}_b\}$. By Claim 4.4, there exists a transversal path Q_a inside \mathcal{G}_a with endpoints $x_a \in N_{G_{l_1}}(z_1) \cap I'_2$, $y_a \in N_{G_{l_5}}(y) \cap I_2$, and a transversal path Q_b inside \mathcal{G}_b with endpoints $x_b \in N_{G_{l_2}}(z_k) \cap J'_2$, $y_b \in N_{G_{l_3}}(z'_t) \cap J_2$. Hence, $z_1 Q_1 z_k x_b Q_b y_b z'_t Q_2 z'_1 z''_1 Q y y_a Q_a x_a z_1$ is a transversal Hamilton cycle in \mathcal{G} . \square




FIGURE 10. All possible cases for $y_1, y_2 \in V_{bad}$. We use “1” (resp. “2”) to denote that the color of this edge lies in \mathcal{C}_1 (resp. \mathcal{C}_2).

Claim 4.14. $E(G_i[A, B]) = \emptyset$ for all $i \in \mathcal{C}_1 \cup \text{col}(M)$.

Proof of Claim 4.14. Suppose that there exists an edge $y_1y_2 \in E(G_{i_2}[A, B])$ for some $i_2 \in \mathcal{C}_1 \cup \text{col}(M)$, where $y_1 \in A$ and $y_2 \in B$. If $i_2 \in \text{col}(M)$, then assume $uv \in E(M[A]) \cap E(G_{i_2})$. Connect P^2 , all rainbow edges in $M - uv$ and all rainbow edges in M' in turn by Claim 4.5 (i)–(ii), we get a rainbow path \tilde{P}^3 with endpoints $\tilde{u}_1 \in A \setminus V_{bad}$ and $\tilde{w}_3 \in B \setminus V_{bad}$. Therefore, P^3 can be written as $uvv'\tilde{u}_1\tilde{P}^3\tilde{w}_3$, where the colors of vv' and $v'\tilde{u}_1$ are in \mathcal{C}_2 . Since $|\mathcal{C}_2 \setminus \text{col}(P^3)|$ is even, one has $|\mathcal{C}_2 \setminus \text{col}(\tilde{P}^3)|$ is even. Let $P^3 := \tilde{P}^3$ if $i_2 \in \text{col}(M)$. Hence we can assume $i_2 \notin \text{col}(P^3)$. Recall that $y_i \notin V(P^3)$ unless it is not in V_{bad} for each $i \in [2]$. Moreover, if $y_i \in V_{bad}$ for some $i \in [2]$, then P^3 can be written as $u_1P^3y_i^1y_i^2P^3w_3$. Assume $y_i^1y_i$ and $y_iy_i^2$ have colors j_{2i-1} and j_{2i} in P^3 , respectively.

If $y_1, y_2 \notin V_{bad}$, then P^3 and y_1y_2 are two disjoint rainbow paths satisfying all conditions in Claim 4.13. Hence, \mathcal{G} contains a transversal Hamilton cycle, a contradiction. Therefore, at least one of y_1 and y_2 lies in V_{bad} . Without loss of generality, assume that $y_2 \in V_{bad}$. Observe that either $j_3, j_4 \in \mathcal{C}_1$ and $y_2^1, y_2^2 \in B$, or $j_3, j_4 \in \mathcal{C}_2$ and $y_2^1, y_2^2 \in A$ holds.

Assume $y_1 \notin V_{bad}$. Delete $y_2^1y_2$ from P^3 , we get two rainbow paths $P' = u_1P^3y_2^1$ and $P'' = y_1y_2y_2^2P^3w_3$. Recall that $u_1, y_1 \in A \setminus V_{bad}$, $w_3 \in B \setminus V_{bad}$ and $y_2^1 \notin V_{bad}$. It is routine to check that P' can be extended to a rainbow path with one endpoint u_1 and the other endpoint $\hat{y} \in B \setminus V_{bad}$, where $\hat{y} = y_2^1$ if $y_2^1 \in B$. Therefore, there are two rainbow paths satisfying all conditions in Claim 4.13. Therefore, \mathcal{G} contains a transversal Hamilton cycle, a contradiction.

Finally, we consider $y_1, y_2 \in V_{bad}$. Then by rearranging, one may assume that $y_1^2P^3y_2^1$ is a rainbow path with length at most 3. Replace $y_1P^3y_2$ by the edge y_1y_2 with color i_2 . It is routine to check that $|\mathcal{C}_2 \setminus \text{col}(u_1P^3y_1y_2P^3w_3)|$ is odd (see Figure 10 for all possible cases). Thus, this can be attributed to the Subcase 3.1. Hence \mathcal{G} contains a transversal Hamilton cycle, a contradiction. \square

By Claim 4.14, one has $E(G_i[A, B]) = \emptyset$ for all $i \in \mathcal{C}_1 \cup \text{col}(M)$. Therefore, n is even and $G_i[A] = G_i[B] = K_{\frac{n}{2}}$ for all $i \in \mathcal{C}_1 \cup \text{col}(M)$. By a similar argument as Claim 4.14, we can show that $E(G_i[A] \cup G_i[B]) = \emptyset$ for all $i \in \mathcal{C}_2 \cup \mathcal{C}'_{bad}$. It follows that \mathcal{G} is a spanning collection of \mathcal{H}_{n-t}^t , where $t = |\mathcal{C}_2 \cup \mathcal{C}'_{bad}|$.

We claim that t is odd. Otherwise, t is even. Recall that $G_i[A] = G_i[B] = K_{\frac{n}{2}}$ for all $i \in \mathcal{C}_1 \cup \text{col}(M)$. Then there exist two rainbow paths, say $Q_1 := z_1^1 \dots z_{s_1}^1$ and $Q_2 := z_1^2 \dots z_{s_2}^2$, such that $V(Q_1 \cup Q_2) = V_{bad}$ and $\text{col}(Q_1 \cup Q_2) \subseteq \mathcal{C}_1 \cup \text{col}(M)$. For $i \in [2]$, it is easy to see that Q_i can be extended to a rainbow path $Q'_i := x^i z_1^i \dots z_{s_i}^i y^i z^i$, where $x^i, y^i \in A \setminus V_{bad}$, $z_i \in B \setminus V_{bad}$, and the colors of $x^i z_1^i$, $z_{s_i}^i y^i$, $y^i z^i$ are unused colors in $\mathcal{C}_1, \mathcal{C}_1, \mathcal{C}_2$, respectively. Since t is even, one has $|\mathcal{C}_2 \cup \mathcal{C}'_{bad} \setminus \text{col}(Q'_1 \cup Q'_2)|$ is even. By a similar argument as Claim 4.13, we know \mathcal{G} contains a transversal Hamilton cycle, a contradiction. Therefore, \mathcal{G} is a spanning collection of \mathcal{H}_{n-t}^t , where n is even and t is odd.

This completes the proof of Theorem 1.6. \square

5. PROOF OF THEOREM 4.3

In this section, we prove Theorem 4.3, which characterizes the structure of almost balanced graph collections that do not contain transversal Hamilton cycles. Based on the parity of n and the size of the vertex partition, we split the proof of Theorem 4.3 into four lemmas (see, Lemmas 5.1, 5.2, 5.6 and 5.8).

Lemma 5.1. *For every δ with $0 < \delta \ll 1$, there exists $n_0 \in \mathbb{N}$ satisfying the following for every odd integer $n \geq n_0$. Let \mathcal{C} be a set of n colors and $\mathcal{G} = \{G_i : i \in \mathcal{C}\}$ be a collection of graphs with common vertex set V of size n . Let $A \cup B$ be a partition of V with $|A| = \frac{n-1}{2}$, and let $\mathcal{C}' \cup \mathcal{C}''$ be a partition of \mathcal{C} with $|\mathcal{C}''| \leq \delta n$. Assume that $\delta(\mathcal{G}) \geq \frac{n-1}{2}$ and $G_i[B] = \emptyset$ for all $i \in \mathcal{C}'$. If \mathcal{G} does not contain transversal Hamilton cycles, then \mathcal{G} is the half-split graph collection.*

Proof. In order to show that \mathcal{G} is the half-split graph collection, it suffices to prove $G_i[B] = \emptyset$ for all $i \in \mathcal{C}''$. By adding colors to \mathcal{C}'' if necessary we may assume $|\mathcal{C}''| = \delta n > 0$ and $\mathcal{C}'' = \{i_1, i_2, \dots, i_{\delta n}\}$. Suppose that $G_{i_1}[B] \neq \emptyset$ and let $uv \in E(G_{i_1}[B])$. Choose a maximal rainbow matching inside $\{G_{i_j}[A, B \setminus \{u, v\}] : j \in [2, \delta n]\}$, say $M = \{e_{i_j} : 2 \leq j \leq s\}$ with $s \leq \delta n$ and $e_{i_j} \in E(G_{i_j})$, such that $G_{i_j}[A \setminus V(M), B \setminus (V(M) \cup \{u, v\})] \neq \emptyset$ for all $j \in [2, s]$. Notice that G_{i_j} is $(2\delta n, K_{\lfloor \frac{n}{2} \rfloor} \cup K_{\lceil \frac{n}{2} \rceil})$ -extremal for all $j \in [s+1, \delta n]$. We greedily select two rainbow paths P_A and P_B inside $\{G_{i_j}[A \setminus V(M)] \cup G_{i_j}[B \setminus (V(M) \cup \{u, v\})] : j \in [s+1, \delta n]\}$ with lengths $\lfloor \frac{\delta n - s}{2} \rfloor$ and $\lceil \frac{\delta n - s}{2} \rceil$ respectively, where $V(P_A) \subseteq A$ and $V(P_B) \subseteq B$.

We first consider $\delta n - s$ is even. It is easy to see that $G_i[A, B] = K_{\frac{n-1}{2}, \frac{n+1}{2}}$ for all $i \in \mathcal{C}'$. Using colors in \mathcal{C}' and edges between A and B , one may connect uv , all rainbow edges in M , P_A and P_B into a single rainbow path, say $P^1 := u \dots v_1$, where $v_1 \in B$. Therefore, $|A \setminus V(P^1)| = |B \setminus V(P^1)| + 1$. Observe that $\{G_i[A \setminus V(P^1), B \setminus V(P^1 - \{u, v_1\})] : i \in \mathcal{C} \setminus \text{col}(P^1)\}$ is a collection of complete bipartite graphs, therefore it contains a transversal path that connects u and v_1 . Hence \mathcal{G} has a transversal Hamilton cycle, a contradiction.

Now, assume that $\delta n - s$ is odd. Recall that $|A| = \frac{n-1}{2}$ and $\delta(\mathcal{G}) \geq \frac{n-1}{2}$. Hence there exists an edge $w_1 w_2 \in E(G_{i_1}[A \setminus V(M \cup P_A \cup P_B), B])$ with $w_1 \in A$ and $w_2 \in B$. Notice that one may avoid w_2 when choosing M and P_B . Using colors in \mathcal{C}' and edges between A and B , we can connect $w_1 w_2$, all rainbow edges in M , P_A and P_B into a single rainbow path P^2 , whose endpoints are in different parts. Therefore, $|A \setminus V(P^1)| = |B \setminus V(P^1)|$. Similarly, \mathcal{G} has a transversal Hamilton cycle, a contradiction. \square

Lemma 5.2. *For every δ with $0 < \delta \ll 1$, there exists $n_0 \in \mathbb{N}$ satisfying the following for every even integer $n \geq n_0$. Let \mathcal{C} be a set of n colors, and $\mathcal{G} = \{G_i : i \in \mathcal{C}\}$ be a collection of graphs with common vertex set V of size n . Let $A \cup B$ be a partition of V with $|A| = \frac{n}{2} - 1$, and let $\mathcal{C}' \cup \mathcal{C}''$ be a partition of \mathcal{C} with $|\mathcal{C}''| \leq \delta n$. Assume that $\delta(\mathcal{G}) \geq \frac{n}{2} - 1$ and $G_i[B] = \emptyset$ for all $i \in \mathcal{C}'$. If \mathcal{G} does not contain transversal Hamilton cycles, then either $G_i[B] = \emptyset$ for all but at most one $i \in \mathcal{C}$, or $E(G_i[B]) \subseteq \{uv\}$ for two fixed vertices $u, v \in B$ and all $i \in \mathcal{C}$.*

Proof. In order to prove this lemma, it suffices to show that $\{G_i[B] : i \in \mathcal{C}''\}$ contains no rainbow P_3 or $2P_2$. By adding colors to \mathcal{C}'' if necessary we may assume $|\mathcal{C}''| = \delta n > 0$ and $\mathcal{C}'' = \{i_1, i_2, \dots, i_{\delta n}\}$. Suppose that $\{G_i[B] : i \in \mathcal{C}''\}$ contains a rainbow path $u_1 u_2 u_3$ with $u_1 u_2 \in E(G_{i_1})$ and $u_2 u_3 \in E(G_{i_2})$. (The proof of the case that $\{G_i[B] : i \in \mathcal{C}''\}$ contains a rainbow $2P_2$ is similar).

Choose a maximal rainbow matching inside $\{G_{i_j}[A, B \setminus \{u_1, u_2, u_3\}] : j \in [3, \delta n]\}$, say $M = \{e_{i_j} : 3 \leq j \leq s\}$ with $s \leq \delta n$ and $e_{i_j} \in E(G_{i_j})$, such that $G_{i_j}[A \setminus V(M), B \setminus (V(M) \cup \{u_1, u_2, u_3\})] \neq \emptyset$ for all $j \in [3, s]$. Notice that G_{i_j} is $(2\delta n, K_{\lfloor \frac{n}{2} \rfloor} \cup K_{\lceil \frac{n}{2} \rceil})$ -extremal for all $j \in [s+1, \delta n]$. We greedily select two rainbow paths P_A and P_B inside $\{G_{i_j}[A \setminus V(M)] \cup G_{i_j}[B \setminus (V(M) \cup \{u_1, u_2, u_3\})] : j \in [s+1, \delta n]\}$ with lengths $\lfloor \frac{\delta n - s}{2} \rfloor$ and $\lceil \frac{\delta n - s}{2} \rceil$ respectively, where $V(P_A) \subseteq A$ and $V(P_B) \subseteq B$.

If $\delta n - s$ is even, then by a similar discussion as Lemma 5.1, it is easy to find a transversal Hamilton cycle inside \mathcal{G} , a contradiction.

Now, assume that $\delta n - s$ is odd. Recall that $|A| = \frac{n}{2} - 1$ and $\delta(\mathcal{G}) \geq \frac{n}{2} - 1$. Then there exists an edge $w_1 w_2 \in E(G_{i_1}[A \setminus V(M \cup P_A \cup P_B), B])$ with $w_1 \in A$ and $w_2 \in B$. Moreover, $G_i[A, B] = K_{\frac{n}{2}-1, \frac{n}{2}+1}$ for all $i \in \mathcal{C}'$. Notice that one may avoid w_2 when choosing M and P_B . If $w_2 \in \{u_2, u_3\}$, then without loss of generality, assume that $w_2 = u_2$. Using colors in \mathcal{C}' and edges between A and B , we can connect $w_1 u_2 u_3$, all rainbow edges in M , P_A and P_B in turn to get a single rainbow path P^1 , whose endpoints are in different parts. Therefore, $|A \setminus V(P^1)| = |B \setminus V(P^1)|$. If $w_2 \notin \{u_2, u_3\}$, then using colors in \mathcal{C}' and edges between A and B , one may connect $u_2 u_3, w_1 w_2$, all rainbow edges in M , P_A and P_B in turn to get a single rainbow path P^2 , whose endpoints are in B . Therefore, $|A \setminus V(P^2)| = |B \setminus V(P^2)| + 1$. In each of the above two cases, it is routine to check that \mathcal{G} has a transversal Hamilton cycle, a contradiction. \square

Lemma 5.3. *Let \mathcal{C} be a set of colors, and $\mathcal{G} = \{G_i[Y, B] : i \in \mathcal{C}\}$ be a collection of bipartite graphs with common bipartition (Y, B) such that $7|Y| < |B| \leq \frac{3}{5}|\mathcal{C}|$. If $\sum_{i \in \mathcal{C}} |E(G_i[Y, B])| \geq t|B||\mathcal{C}|$ for some integer t with $1 \leq t \leq |Y|$, then \mathcal{G} contains t disjoint rainbow star S_5 , each of them has center in Y and other vertices in B .*

Proof. We use induction to prove this result. If $t = 1$, then $\sum_{i \in \mathcal{C}} |E(G_i[Y, B])| \geq |B||\mathcal{C}|$. Suppose that \mathcal{G} does not contain any rainbow S_5 with center in Y and other vertices in B . Then each vertex $v \in Y$ satisfies either $N_{G_i}(v, B) \neq \emptyset$ for at most three $i \in \mathcal{C}$, or there exist three vertices $w, w', w'' \in B$ such that $N_{G_i}(v) \subseteq \{w, w', w''\}$ for all $i \in \mathcal{C}$. It follows that

$$\sum_{i \in \mathcal{C}} |E(G_i[Y, B])| = \sum_{v \in Y} \sum_{i \in \mathcal{C}} d_{G_i}(v, B) \leq \sum_{v \in Y} \max\{3|B|, 3|\mathcal{C}|\} = 3|Y||\mathcal{C}| < |B||\mathcal{C}|,$$

a contradiction. That is to say, our result holds for $t = 1$.

Assume the result holds for t , and we prove it for $t + 1$. Now, we give the following claim.

Claim 5.4. *There exists a vertex $w \in Y$ such that $d_{G_i}(w, B) \geq 4t + 4$ for at least $4t + 4$ colors $i \in \mathcal{C}$.*

Proof of Claim 5.4. Suppose not, then for each vertex $v \in Y$, we have $d_{G_i}(v, B) \geq 4t + 4$ for at most $4t + 3$ colors $i \in \mathcal{C}$. It implies that

$$\begin{aligned} \sum_{i \in \mathcal{C}} |E(G_i[Y, B])| &= \sum_{v \in Y} \sum_{i \in \mathcal{C}} d_{G_i}(v, B) \\ &= \sum_{v \in Y} \left(\sum_{\substack{i \in \mathcal{C} \\ d_{G_i}(v, B) \geq 4t+4}} d_{G_i}(v, B) + \sum_{\substack{i \in \mathcal{C} \\ d_{G_i}(v, B) \leq 4t+3}} d_{G_i}(v, B) \right) \\ &\leq \sum_{v \in Y} \left(\sum_{\substack{i \in \mathcal{C} \\ d_{G_i}(v, B) \geq 4t+4}} |B| + \sum_{\substack{i \in \mathcal{C} \\ d_{G_i}(v, B) \leq 4t+3}} (4t + 3) \right) \\ &\leq |Y|((4t + 3)|B| + |\mathcal{C}|(4t + 3)) \end{aligned}$$

$$\begin{aligned}
&\leq (4t+3)|Y|(|B|+|\mathcal{C}|) \\
&< 7(t+1)|Y||\mathcal{C}| \\
&< (t+1)|B||\mathcal{C}|,
\end{aligned}$$

a contradiction. \square

By Claim 5.4, there exists a vertex $w \in Y$ such that $d_{G_i}(w, B) \geq 4t+4$ for at least $4t+4$ colors $i \in \mathcal{C}$. Let $\mathcal{G}' = \{G_i[Y \setminus \{w\}, B] : i \in \mathcal{C}\}$. Then

$$\sum_{i \in \mathcal{C}} |E(G_i[Y \setminus \{w\}, B])| \geq (t+1)|B||\mathcal{C}| - |B||\mathcal{C}| = t|B||\mathcal{C}|.$$

By induction, there are t disjoint rainbow S_5 inside \mathcal{G}' , each of which has center in $Y \setminus \{w\}$ and other vertices in B . Let \mathbf{S} be a set consisting of those disjoint rainbow S_5 . Recall that $d_{G_i}(w, B) \geq 4t+4$ for at least $4t+4$ colors $i \in \mathcal{C}$. Hence there exist four colors $c^1, c^2, c^3, c^4 \in \mathcal{C} \setminus \text{col}(\mathbf{S})$ such that $d_{G_{c^j}}(w) \geq 4t+4$ for all $j \in [4]$. Thus, there are four vertices $w^1, w^2, w^3, w^4 \in B \setminus V(\mathbf{S})$ such that $ww^j \in E(G_{c^j})$ for all $j \in [4]$. Therefore, $\{G_{c^j}[\{w, w^1, w^2, w^3, w^4\}] : j \in [4]\}$ contains a rainbow S_5 with center w , which is disjoint with each rainbow star in \mathbf{S} , as desired. \square

Lemma 5.5. *Assume $0 < \frac{1}{n} \ll \delta \ll 1$ and $0 \leq \gamma \leq 3\delta$. Let \mathcal{C} be a set of n colors, and $\mathcal{G} = \{G_i : i \in \mathcal{C}\}$ be a collection of graphs with common vertex set V of size n such that $\delta(\mathcal{G}) \geq \frac{n}{2} - 1$. Let $A \cup B$ be a partition of V with $|A| = \lfloor \frac{n+1}{2} \rfloor + \gamma n$, and let $\mathcal{C}' \cup \mathcal{C}''$ be a partition of \mathcal{C} with $|\mathcal{C}''| \leq \delta n$. Define $Y = \{v \in A : d_{G_i}(v, B) \leq (1 - \delta^{\frac{1}{4}})|B| \text{ for at least } \delta^{\frac{1}{4}}|\mathcal{C}'| \text{ colors } i \in \mathcal{C}'\}$. Assume that $|Y| > \gamma n$, \mathcal{G} does not contain transversal Hamilton cycles and $G_i[B] = \emptyset$ for all $i \in \mathcal{C}'$.*

- (i) *If n is odd, then \mathcal{G} is the half-split graph collection.*
- (ii) *If n is even, then there exists a new partition $A' \cup B'$ of V with $|A'| = \frac{n}{2} - 1$ such that either $G_i[B'] = \emptyset$ for all but at most one $i \in \mathcal{C}$, or $E(G_i[B']) \subseteq \{v_1 v_2\}$ for fixed vertices $v_1, v_2 \in B'$ and all $i \in \mathcal{C}$.*

Proof. It is routine to check that

$$\begin{aligned}
\lceil \frac{n}{2} - 1 \rceil |B||\mathcal{C}'| &\leq \sum_{i \in \mathcal{C}'} |E(G_i[A, B])| \\
&\leq |Y|(1 - \delta^{\frac{1}{4}})|B|\delta^{\frac{1}{4}}|\mathcal{C}'| + |Y||B|(1 - \delta^{\frac{1}{4}})|\mathcal{C}'| + (|A| - |Y|)|B||\mathcal{C}'| \\
&= (|A| - \delta^{\frac{1}{2}}|Y|)|B||\mathcal{C}'|. \tag{2}
\end{aligned}$$

It implies that $|Y| \leq \frac{\gamma n + 1}{\sqrt{\delta}} \leq 4\sqrt{\delta}n$. Notice that Claim 4.4 and Claim 4.5 (i)-(ii) hold by setting $V_{\text{bad}} = Y$ and $\mathcal{C}_2 = \mathcal{C}'$.

Since $\delta(\mathcal{G}) \geq \frac{n}{2} - 1$, we have $|E(G_i[A, B])| \geq |B|\lceil \frac{n}{2} - 1 \rceil$ for each $i \in \mathcal{C}'$, i.e., there are at most $|B|(\lfloor \frac{n+1}{2} \rfloor + \gamma n) - |B|\lceil \frac{n}{2} - 1 \rceil = |B|(\gamma n + 1)$ non-edges between A and B in G_i . Hence $\sum_{i \in \mathcal{C}'} |E(G_i[Y, B])| \geq (|Y| - \gamma n - 1)|B||\mathcal{C}'|$. By Lemma 5.3, there exist $|Y| - \gamma n - 1$ disjoint rainbow P_3 inside $\{G_i[Y, B] : i \in \mathcal{C}'\}$ with centers in Y . Let \mathbf{P}_1 be a set consisting of those rainbow P_3 and Y_1 be a subset of Y consisting of the centers of them. Denote $Y_2 := Y \setminus Y_1$. Then $|Y_2| = \gamma n + 1$.

Recall that each vertex in Y is adjacent to at least $\lceil \frac{n}{2} - 1 \rceil - (1 - \delta^{\frac{1}{4}})|B| > 4|Y|$ vertices in A for at least $\delta^{\frac{1}{4}}|\mathcal{C}'| > \delta^{\frac{1}{4}}(1 - \delta)n > 4|Y|$ colors $i \in \mathcal{C}'$. Then there exist $\gamma n + 1$ disjoint rainbow P_3 inside $\{G_i[Y_2, A \setminus Y] : i \in \mathcal{C}' \setminus \text{col}(\mathbf{P}_1)\}$ with centers in Y_2 , and let \mathbf{P}_2 be a set consisting of those

rainbow P_3 . For each $y \in Y$, denote the rainbow P_3 in $\mathbf{P}_1 \cup \mathbf{P}_2$ with center y by $P_y := y^1 y y^2$ with colors c_y^1 and c_y^2 . Let $\mathbf{P} = \mathbf{P}_1 \cup \mathbf{P}_2 = \{P_y : y \in Y\}$.

It is routine to check that there exists a rainbow matching inside $\{G_i[A \setminus V(\mathbf{P}), B \setminus V(\mathbf{P})] : i \in \mathcal{C}''\}$, say M , such that $G_j[A \setminus V(\mathbf{P} \cup M), B \setminus V(\mathbf{P} \cup M)]$ contains a 3-matching for each $j \in \text{col}(M)$. Furthermore, G_j is $(13\sqrt{\delta}, K_{\lfloor \frac{n}{2} \rfloor} \cup K_{\lceil \frac{n}{2} \rceil})$ -extremal for all $j \in \mathcal{C}'' \setminus \text{col}(M)$. We proceed by considering the following three cases.

Case 1. n is odd and $|\mathcal{C}'' \setminus \text{col}(M)|$ is odd, or n is even and $|\mathcal{C}'' \setminus \text{col}(M)| \geq 2$ is even.

Greedy choose two disjoint rainbow paths P_A and P_B inside $\{G_i[A \setminus V(\mathbf{P} \cup M)] \cup G_i[B \setminus V(\mathbf{P} \cup M)] : i \in \mathcal{C}'' \setminus \text{col}(M)\}$ with lengths $\lfloor \frac{|\mathcal{C}'' \setminus \text{col}(M)| - 1}{2} \rfloor$ and $\lceil \frac{|\mathcal{C}'' \setminus \text{col}(M)| + 1}{2} \rceil$ respectively, such that $V(P_A) \subseteq A$ and $V(P_B) \subseteq B$. In view of Claim 4.5 (i)-(ii), by using colors in \mathcal{C}' , one may connect all rainbow paths in \mathbf{P}_2 and \mathbf{P}_1 in turn to get a single rainbow path P^1 with $2|Y_1| + 3|Y_2|$ vertices in A and $2|Y_1| + |Y_2|$ vertices in B , whose endpoints are in different parts. Therefore, $|B \setminus V(P^1)| - |A \setminus V(P^1)| = 2 - \sigma$. Applying Claim 4.5 (i) again to connect P^1 , all rainbow edges in M , P_A and P_B in turn, we get a single rainbow path P^2 with endpoints are in different parts. Clearly, $|A \setminus V(P^2)| = |B \setminus V(P^2)|$. Together with Claim 4.4, we know \mathcal{G} contains a transversal Hamilton cycle, a contradiction.

Case 2. n is odd and $|\mathcal{C}'' \setminus \text{col}(M)|$ is even, or n is even and $|\mathcal{C}'' \setminus \text{col}(M)|$ is odd.

Greedy choose two disjoint rainbow paths P_A and P_B inside $\{G_i[A \setminus V(\mathbf{P} \cup M)] \cup G_i[B \setminus V(\mathbf{P} \cup M)] : i \in \mathcal{C}'' \setminus \text{col}(M)\}$ with lengths $\lfloor \frac{|\mathcal{C}'' \setminus \text{col}(M)|}{2} \rfloor$ and $\lceil \frac{|\mathcal{C}'' \setminus \text{col}(M)|}{2} \rceil$ respectively, such that $V(P_A) \subseteq A$ and $V(P_B) \subseteq B$. Suppose that there exists an $i_1 \in \mathcal{C}' \cup \text{col}(M)$ such that $G_{i_1}[Y_2 \cup B] \neq \emptyset$. Choose $y_1 y_2 \in E(G_{i_1}[Y_2 \cup B])$. By Claim 4.5 (i)-(ii), we connect all rainbow P_3 in $\mathbf{P}_2 \setminus \{P_{y_1}, P_{y_2}\}$, \mathbf{P}_1 , all rainbow edges in M except the possible edge with color i_1 and P_A, P_B in turn to get a single rainbow path P^3 , whose endpoints are in different parts. Similar to Case 1, we have $|A \setminus V(P^3)| - |B \setminus V(P^3)| = 2|\{y_1, y_2\} \cap Y_2| - 1$.

If $y_1, y_2 \in Y_2$, then let $Q := y_1^1 y_1 y_2 y_2^2$ with $\text{col}(Q) = \{c_{y_1}^1, i_1, c_{y_2}^2\}$. If $y_1 \in Y_2$ and $y_2 \in B$, then one may assume $y_2 \notin V(P^3)$ and let $Q := y_1^1 y_1 y_2$ with $\text{col}(Q) = \{c_{y_1}^1, i_1\}$. If $y_1, y_2 \in B$, then we may assume $y_1, y_2 \notin V(P^3)$ and let $Q := y_1 y_2$ with color i_1 . Next, connect P^3 with the rainbow path Q by Claim 4.5 (i)-(ii), we get a rainbow path P^4 with endpoints y_1^1, v_3 (if $y_1, y_2 \in Y_2$) or u_3, y_3 (otherwise). It is routine to check that $|A \setminus V(P^4)| = |B \setminus V(P^4)|$. Together with Claim 4.4, one obtains that \mathcal{G} contains a transversal Hamilton cycle, a contradiction. Hence, $G_i[Y_2 \cup B] = \emptyset$ for all $i \in \mathcal{C}' \cup \text{col}(M)$. Recall that $\delta(G_i) \geq \lceil \frac{n}{2} - 1 \rceil$. This implies that $N_{G_i}(w) = A \setminus Y_2$ for all $i \in \mathcal{C}' \cup \text{col}(M)$ and all $w \in Y_2 \cup B$. Move vertices in Y_2 from A to B , we get $|A| = \lceil \frac{n}{2} - 1 \rceil$, $|B| = \lfloor \frac{n}{2} + 1 \rfloor$ and $G_i[B] = \emptyset$ for each $i \in \mathcal{C}' \cup \text{col}(M)$. By Lemmas 5.1 and 5.2, our desired result holds.

Case 3. n is even and $|\mathcal{C}'' \setminus \text{col}(M)| = 0$.

By a similar discussion as Case 2 and the proof of Lemma 5.2, we obtain that $\mathcal{G}[Y_2 \cup B]$ contains no rainbow P_3 or $2P_2$. Hence either $G_i[Y_2 \cup B] = \emptyset$ for all but at most one unique $i \in \mathcal{C}$ or $E(G_i[Y_2 \cup B]) \subseteq \{uv\}$ for fixed u, v and all $i \in \mathcal{C}$. Move vertices in Y_2 to B , we get $|A| = \frac{n}{2} - 1$, $|B| = \frac{n}{2} + 1$. Therefore, either $G_i[B] = \emptyset$ for all but at most one $i \in \mathcal{C}$, or $E(G_i[B]) \subseteq \{uv\}$ for all $i \in \mathcal{C}$. Recall that $\delta(\mathcal{G}) \geq \frac{n}{2} - 1$. In both cases, \mathcal{G} does not contain transversal Hamilton cycles, as desired. \square

Lemma 5.6. Assume $0 < \frac{1}{n} \ll \delta \ll 1$. Let \mathcal{C} be a set of n colors and $\mathcal{G} = \{G_i : i \in \mathcal{C}\}$ be a collection of graphs with common vertex set V of size n . Let $A \cup B$ be a partition of V with $|A| = \frac{n}{2}$, and let $\mathcal{C}' \cup \mathcal{C}''$ be a partition of \mathcal{C} with $|\mathcal{C}''| \leq \delta n$. Assume that $\delta(\mathcal{G}) \geq \frac{n}{2} - 1$ and

$G_i[B] = \emptyset$ for all $i \in \mathcal{C}'$. If \mathcal{G} does not contain transversal Hamilton cycles, then one of the following holds:

- \mathcal{G} is a spanning collection of \mathcal{H}_{n-t}^t for some odd integer t ,
- there exists a partition $A' \cup B'$ of V with $|A'| = \frac{n}{2} - 1$, such that either $G_i[B'] = \emptyset$ for all but at most one $i \in \mathcal{C}$, or $E(G_i[B']) \subseteq \{uv\}$ for fixed vertices $u, v \in B'$ and all $i \in \mathcal{C}$.

Proof. Let $Y = \{v \in A : d_{G_i}(v, B) \leq (1 - \delta^{\frac{1}{4}})|B| \text{ for at least } \delta^{\frac{1}{4}}|\mathcal{C}'| \text{ colors } i \in \mathcal{C}'\}$. Then

$$\begin{aligned} \left(\frac{n}{2} - 1\right)|B||\mathcal{C}'| &\leq \sum_{i \in \mathcal{C}'} |E(G_i[A, B])| \\ &= \sum_{v \in Y} \sum_{i \in \mathcal{C}'} d_{G_i}(v, B) + \sum_{v \in A \setminus Y} \sum_{i \in \mathcal{C}'} d_{G_i}(v, B) \\ &\leq |Y|(1 - \delta^{\frac{1}{4}})|B|\delta^{\frac{1}{4}}|\mathcal{C}'| + |Y||B|(1 - \delta^{\frac{1}{4}})|\mathcal{C}'| + \left(\frac{n}{2} - |Y|\right)|B||\mathcal{C}'| \\ &= \left(\frac{n}{2} - \beta|Y|\right)|B||\mathcal{C}'|. \end{aligned}$$

It follows that $|Y| \leq \frac{1}{\sqrt{\delta}} < \sqrt{\delta}n$. If $|Y| \geq 1$, then our result holds by Lemma 5.5. In what follows, we only consider $|Y| = 0$. It is straightforward to check that Claim 4.4 and Claim 4.5 (i)-(ii) hold by setting $V_{bad} = Y$ and $\mathcal{C}_2 = \mathcal{C}'$.

Choose a maximal rainbow matching inside $\{G_i[A, B] : i \in \mathcal{C}''\}$, say M , such that $G_j[A, B]$ contains a 2-matching for each $j \in \text{col}(M)$. Furthermore, G_j is $(2\sqrt{\delta}, K_{\lfloor \frac{n}{2} \rfloor} \cup K_{\lceil \frac{n}{2} \rceil})$ -extremal for all $j \in \mathcal{C}'' \setminus \text{col}(M)$. Greedily choose two disjoint rainbow paths P_A and P_B inside $\{G_i[A \setminus V(M)] \cup G_i[B \setminus V(M)] : i \in \mathcal{C}'' \setminus \text{col}(M)\}$ with lengths x_A and x_B respectively, such that $V(P_A) \subseteq A$ and $V(P_B) \subseteq B$. In view of Claim 4.5 (i)-(ii), by using colors in \mathcal{C}' , one may connect all rainbow edges in M , P_A and P_B into a single rainbow path P , whose endpoints are denoted by $u_1 \in A$ and $v_1 \in B$.

If $|\mathcal{C}'' \setminus \text{col}(M)|$ is even, then let $x_A = x_B = \frac{|\mathcal{C}'' \setminus \text{col}(M)|}{2}$. Thus, $|B \setminus V(P)| = |A \setminus V(P)|$. Together with Claim 4.4, we know \mathcal{G} contains a transversal Hamilton cycle, a contradiction. Hence $|\mathcal{C}'' \setminus \text{col}(M)|$ is odd. Now, we give the following claim.

Claim 5.7. (i) $G_i[A] \cup G_i[B] = \emptyset$ for all $i \in \mathcal{C}' \cup \text{col}(M)$,
(ii) $G_i[A, B] = \emptyset$ for all $i \in \mathcal{C}'' \setminus \text{col}(M)$.

Proof of Claim 5.7. We only give the proof of $G_i[A] = \emptyset$ for all $i \in \mathcal{C}' \cup \text{col}(M)$, the other two statements can be proved by similar arguments.

Suppose that $G_{i_1}[A] \neq \emptyset$ for some $i_1 \in \mathcal{C}' \cup \text{col}(M)$. Choose $w_1 w_2 \in E(G_{i_1}[A])$. If $i_1 \in \text{col}(M)$, then delete the edge with color i_1 from M . Hence we can assume that $i_1 \notin \text{col}(M)$ and $w_1, w_2 \notin V(P)$. Using Claim 4.5 (ii) to connect $w_1 w_2$ and the rainbow path P , we get a rainbow path P' with endpoints w_1 and v_1 . Let $x_A = \frac{|\mathcal{C}'' \setminus \text{col}(M)| - 1}{2}$ and $x_B = \frac{|\mathcal{C}'' \setminus \text{col}(M)| + 1}{2}$. Hence $|B \setminus V(P')| = |A \setminus V(P')|$. It follows from Claim 4.4 that \mathcal{G} contains a transversal Hamilton cycle, a contradiction. \square

Based on Claim 5.7, we know \mathcal{G} is a spanning collection of \mathcal{H}_{n-t}^t for some odd t , as desired. \square

Lemma 5.8. Assume $0 < \frac{1}{n} \ll \delta \ll 1$ and $0 \leq \gamma \leq 3\delta$. Let \mathcal{C} be a set of n colors, and $\mathcal{G} = \{G_i : i \in \mathcal{C}\}$ be a collection of graphs with common vertex set V of size n such that $\delta(\mathcal{G}) \geq \frac{n}{2} - 1$. Let $A \cup B$ be a partition of V with $|A| = \lfloor \frac{n+1}{2} \rfloor + \gamma n$, and let $\mathcal{C}' \cup \mathcal{C}''$ be a partition of \mathcal{C} with $|\mathcal{C}''| \leq \delta n$. Assume that \mathcal{G} does not contain transversal Hamilton cycles and $G_i[B] = \emptyset$ for all $i \in \mathcal{C}'$.

- (i) If n is odd, then \mathcal{G} is the half-split graph collection.
- (ii) If n is even, then one of the following holds:
- \mathcal{G} is a spanning collection of \mathcal{H}_{n-t}^t for some odd integer t ,
 - there exists a new partition $A' \cup B'$ of V with $|A'| = \frac{n}{2} - 1$ such that either $G_i[B'] = \emptyset$ for all but at most one $i \in \mathcal{C}$, or $E(G_i[B']) \subseteq \{v_1 v_2\}$ for two fixed vertices $v_1, v_2 \in B'$ and all $i \in \mathcal{C}$.

Proof. Define $Y = \{v \in A : d_{G_i}(v, B) \leq (1 - \delta^{\frac{1}{4}})|B| \text{ for at least } \delta^{\frac{1}{4}}|\mathcal{C}'| \text{ colors } i \in \mathcal{C}'\}$. In view of (2), we have $|Y| \leq 4\sqrt{\delta}n$. If $|Y| > \gamma n$, then our result holds by Lemma 5.5. In what follows, we only consider $|Y| \leq \gamma n$. Therefore, $|B| \leq |A| - 2|Y| - \sigma$ where $\sigma = 0$ if n is even and $\sigma = 1$ otherwise. Notice that Claim 4.4 and Claim 4.5 (i)-(ii) hold by setting $V_{bad} = Y$ and $\mathcal{C}_2 = \mathcal{C}'$.

Recall that each vertex in Y is adjacent to at least $\lceil \frac{n}{2} - 1 \rceil - (1 - \delta^{\frac{1}{4}})|B| > 5|Y|$ vertices in A for at least $\delta^{\frac{1}{4}}|\mathcal{C}'| > \delta^{\frac{1}{4}}(1 - \delta)n > 5|Y|$ colors $i \in \mathcal{C}'$. Then there exist $|Y|$ disjoint rainbow paths P_3 with centers in Y and endpoints in $A \setminus Y$ using colors in \mathcal{C}' . In the graph collection $\mathcal{G}[A \setminus Y]$, we extend those rainbow paths or choose other disjoint rainbow paths into a set of disjoint maximal rainbow paths. Let $\mathbf{P} = \{Q_1, Q_2, \dots, Q_t\}$ be a set consisting of all disjoint rainbow paths in the above, each of which has length s_i ($1 \leq i \leq t$).

Case 1. $|B| \geq |A| - (s_1 + \dots + s_t) - \sigma$.

In this case, there exists a set $\mathbf{P}' = \{Q'_1, Q'_2, \dots, Q'_\ell\}$ such that $|A \setminus V(\mathbf{P}')| - \sigma = |B| - \ell$, where $Y \subseteq V(\mathbf{P}')$ and the endpoints of Q'_i are not in Y for all $i \in [\ell]$. Assume $|E(Q'_i)| = s'_i$ for each $i \in [\ell]$. By Claim 4.5 (ii), we can connect all rainbow paths of \mathbf{P}' into a single rainbow path P^1 , whose endpoints are in different parts. Since $|A| - |B| = 2\gamma n + \sigma$, we have $\ell \leq s'_1 + \dots + s'_\ell = 2\gamma n + \sigma$. Therefore, $|E(P^1)| \leq 6\gamma n + 3\sigma$ and $|A \setminus V(P^1)| - \sigma = |B \setminus V(P^1)|$.

Choose a maximal rainbow matching inside $\{G_i[A \setminus V(P^1), B \setminus V(P^1)] : i \in \mathcal{C}'' \setminus \text{col}(P^1)\}$, say M , such that $G_j[A \setminus V(P^1 \cup M), B \setminus V(P^1 \cup M)]$ contains a 2-matching for each $j \in \text{col}(M)$. Furthermore, G_j is $(6(\gamma + \delta), K_{\lfloor \frac{n}{2} \rfloor} \cup K_{\lceil \frac{n}{2} \rceil})$ -extremal for all $j \in \mathcal{C}'' \setminus \text{col}(P^1 \cup M)$. In fact, when construct $P^1 - B - Y$, it is possible to use colors in $\mathcal{C}'' \setminus \text{col}(P^1 \cup M)$ before other colors. Hence either $\text{col}(P^1 - B - Y) \subseteq \mathcal{C}'' \setminus \text{col}(M)$ or $\mathcal{C}'' \setminus \text{col}(M) \subseteq \text{col}(P^1 - B - Y)$.

Based on Claim 4.5 (i), one may connect P^1 , all rainbow edges in M into a single rainbow path P^2 , whose endpoints are in different parts. Clearly, $|E(P^2)| \leq 6\gamma n + 3\sigma + 4\delta n$. Next, we are to choose two disjoint rainbow paths P_A and P_B inside $\{G_i[A \setminus V(P^1 \cup M)] \cup G_i[B \setminus V(P^1 \cup M)] : i \in \mathcal{C}'' \setminus \text{col}(P^1 \cup M)\}$ such that $V(P_A) \subseteq A$ and $V(P_B) \subseteq B$ respectively, whose lengths are determined by the parity of $|\mathcal{C}'' \setminus \text{col}(P^1 \cup M)|$. Clearly, if n and $|\mathcal{C}'' \setminus \text{col}(P^1 \cup M)|$ have the same parity, then by Claim 4.4, we know \mathcal{G} contains a transversal Hamilton cycle, a contradiction. Hence n and $|\mathcal{C}'' \setminus \text{col}(P^1 \cup M)|$ have different parity. By similar arguments as the proof of Theorem 1.6 (Step 4 in Case 1), we can show that the following claim.

Claim 5.9. (i) If $|\mathcal{C}'' \setminus \text{col}(P^1 \cup M)| = 0$, then $E(G_i[A]) = E(G_i[V(\mathbf{P}') \cap A])$ for all $i \in \mathcal{C}$.
(ii) $G_i[A, B] = \emptyset$ for all $i \in \mathcal{C}'' \setminus \text{col}(P^1 \cup M)$.

If $|\mathcal{C}'' \setminus \text{col}(P^1 \cup M)| = 0$, then n is odd. By Claim 5.9 (i), one has $|A| \geq |B| \geq \frac{n-1}{2}$. Hence $|A| = \frac{n+1}{2}$ and $|B| = \frac{n-1}{2}$. This implies that P^1 is a null graph. Thus, $G_i[A] = \emptyset$ for all $i \in \mathcal{C}$. Together with Lemma 5.1, our desired result holds.

If $|\mathcal{C}'' \setminus \text{col}(P^1 \cup M)| \geq 1$, then based on Claim 5.9 (ii), we know n is even, $|A| = |B| = \frac{n}{2}$ and $G_i[B] = \emptyset$ for all but at most δn colors in \mathcal{C} . By Lemma 5.6, our desired result holds.

Case 2. $|B| < |A| - (s_1 + \dots + s_t) - \sigma$.

In this case, $\sum_{i=1}^t s_i < 2\gamma n$ and so $\sum_{i=1}^t \frac{s_i}{2} \leq \gamma n - \frac{1}{2}$. Using Claim 4.5 (ii) and t vertices in B , one may connect all rainbow paths of \mathbf{P} into a single rainbow paths P^3 , whose endpoint are in different parts. Let $\tilde{A} = A \setminus V(P^3)$, $\tilde{B} = B \setminus V(P^3)$ and $\tilde{C} = C \setminus \text{col}(P^3)$. Then $|\tilde{A}| = |A| - (s_1 + \dots + s_t + t)$, $|\tilde{B}| = |B| - t$ and $|\tilde{C}| = |C| - (s_1 + \dots + s_t + 2t - 1)$. For convenience, we assume that $Q_i = u_0^i u_1^i \dots u_{s_i}^i$ for each $i \in [t]$.

Let w be an arbitrary vertex in \tilde{A} and c_1, c_2 be two colors in $\mathcal{C} \setminus \text{col}(\mathbf{P})$. Without loss of generality, assume that $d_{G_{c_1}}(w, A \setminus \tilde{A}) \leq d_{G_{c_2}}(w, A \setminus \tilde{A})$. Clearly, in G_{c_1} and G_{c_2} , w is adjacent to at least $\lceil \frac{n}{2} - 1 \rceil - (\lceil \frac{n-1}{2} \rceil - \gamma n) = \gamma n - 1 + \sigma$ vertices in $A \setminus \tilde{A}$ and it cannot adjacent to the pendant vertices of Q_i for all $i \in [t]$ (by the maximality of \mathbf{P}). Furthermore, if $u_j^i \in N_{G_{c_1}}(w)$ for some $i \in [t]$ and $j \in [s_i]$, then $u_{j-1}^i, u_{j+1}^i \notin N_{G_{c_2}}(w)$. Denote by t_{odd} the number of rainbow paths in \mathbf{P} with odd lengths. Thus,

$$\gamma n - 1 + \sigma \leq d_{G_{c_1}}(w, A \setminus \tilde{A}) \leq \sum_{i=1}^t \lceil \frac{s_i - 1}{2} \rceil = \sum_{i=1}^t \frac{s_i}{2} - \frac{1}{2} t_{\text{odd}} \leq \gamma n - \frac{1}{2} - \frac{1}{2} t_{\text{odd}}. \quad (3)$$

This implies that $t_{\text{odd}} \leq 1 - 2\sigma$. Therefore, n is even and $t_{\text{odd}} \leq 1$.

• $t_{\text{odd}} = 0$. Notice that $\sum_{i=1}^t \frac{s_i}{2}$ and γn are integers, then $d_{G_{c_1}}(w, A \setminus \tilde{A}) = \sum_{i=1}^t \frac{s_i}{2} = \gamma n - 1$. It follows that $N_{G_{c_1}}(w, A \setminus \tilde{A}) = \cup_{i \in [t]} \{u_1^i, u_3^i, \dots, u_{s_i-1}^i\}$. Hence, if $i \in [t]$ and j is even, then u_j^i cannot adjacent to w in G_{c_2} . Recall that $d_{G_{c_2}}(w, A \setminus \tilde{A}) \geq \gamma n - 1 = \sum_{i=1}^t \frac{s_i}{2}$. We have $N_{G_{c_1}}(w, A \setminus \tilde{A}) = N_{G_{c_2}}(w, A \setminus \tilde{A})$. Thus, for any G_i with $i \in \mathcal{C} \setminus \text{col}(\mathbf{P})$ and any $w \in \tilde{A}$, we have $N_{G_i}(w) \cap (A \setminus \tilde{A}) = \cup_{i \in [t]} \{u_1^i, u_3^i, \dots, u_{s_i-1}^i\}$.

Now, move all vertices in $\cup_{i \in [t]} \{u_1^i, u_3^i, \dots, u_{s_i-1}^i\}$ from A to B . Then $|A| = \frac{n}{2} + 1$, $|B| = \frac{n-1}{2} - 1$ and $G_i[A]$ is empty for all $i \in \mathcal{C} \setminus \text{col}(\mathbf{P})$. Notice that $|\text{col}(\mathbf{P})| = \sum_{i=1}^t s_i \leq 2\gamma n$. Together with Lemma 5.2, our desired result holds.

• $t_{\text{odd}} = 1$. Then all inequalities in (3) must be equalities. Thus, $d_{G_{c_1}}(w, A \setminus \tilde{A}) = \sum_{i=1}^t \frac{s_i}{2} - \frac{1}{2} = \gamma n - 1$ and therefore $d_{G_{c_1}}(w, V(Q_i)) = \lceil \frac{s_i-1}{2} \rceil$ for all $i \in [t]$. Without loss of generality, assume that Q_1 is the unique rainbow path with odd length. By a similar discussion as the case for $t_{\text{odd}} = 0$, we know for every $i \in \mathcal{C} \setminus \text{col}(\mathbf{P})$ and every $w \in \tilde{A}$, $N_{G_i}(w, \cup_{i \in [t] \setminus \{1\}} V(Q_i)) = \cup_{i \in [t] \setminus \{1\}} \{u_1^i, u_3^i, \dots, u_{s_i-1}^i\}$.

Notice that in each G_i with $i \in \mathcal{C} \setminus \text{col}(\mathbf{P})$, each vertex in \tilde{A} has $\frac{s_1-1}{2}$ neighbors in $V(Q_1) \setminus \{u_0^1, u_{s_1}^1\}$ and it cannot adjacent to two adjacent vertices in Q_1 with two different colors. Hence for each $i \in \mathcal{C} \setminus \text{col}(\mathbf{P})$, $N_{G_i}(w)$ must be one of the following sets:

$$\begin{aligned} & \{u_1^1, u_3^1, \dots, u_{s_1-2}^1\}, \quad \{u_2^1, u_4^1, \dots, u_{s_1-1}^1\}, \\ & \{u_1^1, u_3^1, \dots, u_{j-3}^1, u_j^1, u_{j+2}^1, \dots, u_{s_1-1}^1\} \text{ for some even integer } j \text{ with } 4 \leq j \leq s_1 - 1. \end{aligned}$$

It is routine to check that $N_{G_{c_1}}(w, V(Q_1)) = N_{G_{c_2}}(w, V(Q_1))$. If there are two vertices $v, v' \in A \setminus \tilde{A}$ and colors $i_1, i_2, i_3, i_4 \in \mathcal{C} \setminus \text{col}(\mathbf{P})$ such that $u_{j-1}^1, u_{j+1}^1 \in N_{G_{i_1}}(v) = N_{G_{i_2}}(v)$ and $u_j^1, u_{j+2}^1 \in N_{G_{i_3}}(v') = N_{G_{i_4}}(v')$, then we can find a longer rainbow path $u_0^1 u_1^1 \dots u_{j-1}^1 v u_{j+1}^1 u_j^1 v' u_{j+2}^1 \dots u_{s_1}^1$, where $u_{j-1}^1 v \in E(G_{i_1})$, $v u_{j+1}^1 \in E(G_{i_2})$, $u_j^1 v' \in E(G_{i_3})$ and $v' u_{j+2}^1 \in E(G_{i_4})$, a contradiction. Notice that for two different vertices v and v' , it is possible that $N_{G_i}(v, V(Q_1)) = \{u_1^1, u_3^1, \dots, u_{s_1-2}^1\}$ and $N_{G_i}(v', V(Q_1)) = \{u_1^1, u_3^1, \dots, u_{s_1-4}^1, u_{s_1-1}^1\}$ hold for all $i \in \mathcal{C} \setminus \text{col}(\mathbf{P})$. Hence $\cup_{i \in \mathcal{C} \setminus \text{col}(\mathbf{P})} N_{G_i}(\tilde{A}, V(Q_1))$ is contained in one of the following sets:

$$\begin{aligned} & \{u_1^1, u_3^1, \dots, u_{s_1-2}^1, u_{s_1-1}^1\}, \quad \{u_2^1, u_4^1, \dots, u_{s_1-1}^1\}, \\ & \{u_1^1, u_3^1, \dots, u_{j_0-3}^1, u_{j_0}^1, u_{j_0+2}^1, \dots, u_{s_1-1}^1\} \text{ for a fixed even integer } j_0 \text{ with } 4 \leq j_0 \leq s_1 - 1. \end{aligned}$$

Move all vertices in $\bigcup_{i \in \mathcal{C} \setminus \text{col}(\mathbf{P})} N_{G_i}(\tilde{A}, A \setminus \tilde{A})$ from A to B , then $G_i[A]$ is empty for all $i \in \mathcal{C} \setminus \text{col}(\mathbf{P})$. What's more, either $|A| = \frac{n}{2} + 1$ and $|B| = \frac{n}{2} - 1$, or $|A| = \frac{n}{2}$ and $|B| = \frac{n}{2}$. Together with Lemmas 5.2 and 5.6, our desired result holds.

This completes the proof of Lemma 5.8. \square

Combining Lemmas 5.1, 5.2, 5.6 and 5.8, we obtain Theorem 4.3.

6. CONCLUDING REMARKS

In this paper, we studied the Dirac-type conditions for transversal Hamilton paths and transversal Hamilton cycles in graph collections. As we know, there are many sufficient conditions to guarantee the existence of Hamilton cycles in a graph, such as Ore's condition [41], Pósa's condition [42] and so on. It is natural to consider the Ore-type condition or Pósa-type condition for transversal Hamilton cycles in a graph collection (which was already proposed in [36]). Furthermore, motivated by the stability results for transversal Hamilton cycles under Dirac-type condition [20], one may study the stability result for transversal Hamilton cycles under Ore-type condition or Pósa-type condition.

The *closure* of an n -vertex graph G , denoted $C(G)$, is the graph with vertex set $V(G)$ obtained from G by iteratively adding edges joining pairs of nonadjacent vertices whose degree sum is at least n , until no such pair remains. Bondy and Chvátal [8] showed that an n -vertex graph is Hamiltonian if and only if its closure is Hamiltonian. We propose the following question.

Question 6.1. *Let $\mathcal{G} = \{G_1, \dots, G_n\}$ be a collection of graphs with common vertex set V of size n . If $\{C(G_i) : i \in [n]\}$ contains a transversal Hamilton cycle, does \mathcal{G} contain a transversal Hamilton cycle?*

Transversal generalizations were recently considered for digraphs. Cheng, Han, Wang and Wang [17] established a minimum degree condition to guarantee the existence transversal tournament factors in digraph collections. Chakraborti, Kim, Lee and Seo [16] proved the existence of transversal Hamilton paths and cycles in tournament collections.

For a digraph D , define $\delta^+(D) = \{d_D^+(v) : v \in V(D)\}$ and $\delta^-(D) = \{d_D^-(v) : v \in V(D)\}$ to be the minimum out-degree and minimum in-degree of D respectively. The minimum semi-degree of D is $\delta^0(D) := \min\{\delta^+(D), \delta^-(D)\}$. Ghouila-Houri [27] proved that any n -vertex digraph D with $\delta^0(D) \geq \frac{n}{2}$ contains a directed Hamilton cycle. Chakraborti, Kim, Lee and Seo [16] proposed that it would be interesting to consider a transversal version of the above theorem. Woodall [46] showed that if D is an n -vertex digraph satisfying $d_D^+(u) + d_D^-(v) \geq n$ for all pairs of vertices $\{u, v\}$ with $\vec{uv} \notin A(D)$, then D contains a directed Hamilton cycle. Hence it is natural to consider the transversal version of this result.

REFERENCES

- [1] R. Aharoni, C. St. J. A. Nash-Williams, and S. Shelah, *A general criterion for the existence of transversals*, Proc. London Math. Soc. (3) **47** (1983), no. 1, 43–68. $\uparrow 1$
- [2] R. Aharoni and E. Berger, *Rainbow matchings in r -partite r -graphs*, Electron. J. Combin. **16** (2009), no. 1, Research Paper 119, 9. $\uparrow 2$
- [3] R. Aharoni, E. Berger, M. Chudnovsky, D. Howard, and P. Seymour, *Large rainbow matchings in general graphs*, European J. Combin. **79** (2019), 222–227. $\uparrow 2$
- [4] R. Aharoni, M. DeVos, S. González Hermosillo de la Maza, A. Montejano, and R. Šámal, *A rainbow version of Mantel's theorem*, Adv. Comb. (2020), Paper No. 2, 12. $\uparrow 1, 2$

- [5] R. Aharoni and D. Howard, *A rainbow r -partite version of the Erdős-Ko-Rado theorem*, *Combin. Probab. Comput.* **26** (2017), no. 3, 321–337. ↑2
- [6] M. Anastos and D. Chakraborti, *Robust Hamiltonicity in families of Dirac graphs*, arXiv preprint arXiv:2309.12607 (2023). ↑2
- [7] J. A. Bondy, *Large cycles in graphs*, *Discrete Math.* **1** (1971/72), no. 2, 121–132. ↑2
- [8] J. A. Bondy and V. Chvátal, *A method in graph theory*, *Discrete Math.* **15** (1976), no. 2, 111–135. ↑31
- [9] J. A. Bondy and M. Kouider, *Hamilton cycles in regular 2-connected graphs*, *J. Combin. Theory Ser. B* **44** (1988), no. 2, 177–186. ↑2
- [10] J. Böttcher, M. Schacht, and A. Taraz, *Proof of the bandwidth conjecture of Bollobás and Komlós*, *Math. Ann.* **343** (2009), no. 1, 175–205. ↑2
- [11] C. Bowtell, P. Morris, Y. Pehova, and K. Staden, *Universality for transversal Hamilton cycles*, arXiv preprint arXiv:2310.04138 (2023). ↑2
- [12] P. Bradshaw, *Transversals and bipancyclicity in bipartite graph families*, *Electron. J. Combin.* **28** (2021), no. 4, Paper No. 4.25, 20. ↑2
- [13] P. Bradshaw, K. Halasz, and L. Stacho, *From one to many rainbow Hamiltonian cycles*, *Graphs Combin.* **38** (2022), no. 6, Paper No. 188, 21. ↑2
- [14] Y. Büyükoçak, D. Gözüpek, S. Özkan, and M. Shalom, *On one extension of Dirac’s theorem on Hamiltonicity*, *Discrete Appl. Math.* **252** (2019), 10–16. ↑4
- [15] D. Chakraborti, S. Im, J. Kim, and H. Liu, *A bandwidth theorem for graph transversals*, arXiv preprint arXiv:2302.09637 (2023). ↑2
- [16] D. Chakraborti, J. Kim, H. Lee, and J. Seo, *Hamilton transversals in tournaments*, arXiv preprint arXiv:2307.00912v2 (2023). ↑31
- [17] Y. Cheng, J. Han, B. Wang, and G. Wang, *Rainbow spanning structures in graph and hypergraph systems*, *Forum Math. Sigma* **11** (2023), Paper No. e95, 20. ↑2, 31
- [18] Y. Cheng, J. Han, B. Wang, G. Wang, and D. Yang, *Rainbow Hamilton cycle in hypergraph systems*, arXiv preprint arXiv:2111.07079 (2021). ↑2
- [19] Y. Cheng and K. Staden, *Transversals via regularity*, arXiv preprint arXiv:2306.03595 (2023). ↑2, 6, 12
- [20] ———, *Stability of transversal Hamilton cycles and paths*, arXiv preprint arXiv:2403.09913 (2024). ↑3, 6, 10, 12, 31
- [21] Y. Cheng, G. Wang, and Y. Zhao, *Rainbow pancyclicity in graph systems*, *Electron. J. Combin.* **28** (2021), no. 3, Paper No. 3.24, 9. ↑2
- [22] R. Diestel, *Graph theory*, Fifth, Graduate Texts in Mathematics, vol. 173, Springer, Berlin, 2017. ↑5
- [23] G. A. Dirac, *Some theorems on abstract graphs*, *Proc. London Math. Soc. (3)* **2** (1952), 69–81. ↑2
- [24] P. Erdős, A. Hajnal, and E. C. Milner, *On sets of almost disjoint subsets of a set*, *Acta Math. Acad. Sci. Hungar.* **19** (1968), 209–218. ↑1
- [25] A. Ferber, J. Han, and D. Mao, *Dirac-type problem of rainbow matchings and Hamilton cycles in random graphs*, arXiv preprint arXiv:2211.05477 (2022). ↑2
- [26] L. Fu, L. Gao, J. Wang, and W. Yang, *Non-Hamiltonian graphs with large minimum degree*, *Bull. Malays. Math. Sci. Soc.* **47** (2024), no. 1, Paper No. 32, 11. ↑4
- [27] A. Ghouila-Houri, *Une condition suffisante d’existence d’un circuit hamiltonien*, *C. R. Acad. Sci. Paris* **251** (1960), 495–497. ↑31
- [28] R. J. Gould, *Advances on the Hamiltonian problem—a survey*, *Graphs Combin.* **19** (2003), no. 1, 7–52. ↑2
- [29] ———, *Recent advances on the Hamiltonian problem: Survey III*, *Graphs Combin.* **30** (2014), no. 1, 1–46. ↑2
- [30] P. Gupta, F. Hamann, A. Müyesser, O. Parczyk, and A. Sgueglia, *A general approach to transversal versions of Dirac-type theorems*, *Bull. Lond. Math. Soc.* **55** (2023), no. 6, 2817–2839. ↑2
- [31] A. Hajnal and E. Szemerédi, *Proof of a conjecture of P. Erdős*, *Combinatorial theory and its applications, I-III (Proc. Colloq., Balatonfüred, 1969)*, 1970, pp. 601–623. ↑2
- [32] F. Joos and J. Kim, *On a rainbow version of Dirac’s theorem*, *Bull. Lond. Math. Soc.* **52** (2020), no. 3, 498–504. ↑1, 2, 3
- [33] J. Komlós, G. N. Sárközy, and E. Szemerédi, *Spanning trees in dense graphs*, *Combin. Probab. Comput.* **10** (2001), no. 5, 397–416. ↑2

- [34] D. Kühn and D. Osthus, *The minimum degree threshold for perfect graph packings*, *Combinatorica* **29** (2009), no. 1, 65–107. ↑2
- [35] A. Kupavskii, *Rainbow version of the Erds matching conjecture via concentration*, *Comb. Theory* **3** (2023), no. 1, Paper No. 1, 19. ↑2
- [36] L. Li, P. Li, and X. Li, *Rainbow structures in a collection of graphs with degree conditions*, *J. Graph Theory* **104** (2023), no. 2, 341–359. ↑2, 7, 31
- [37] H. Lu, Y. Wang, and X. Yu, *Rainbow perfect matchings for 4-uniform hypergraphs*, *SIAM J. Discrete Math.* **36** (2022), no. 3, 1645–1662. ↑2
- [38] ———, *A better bound on the size of rainbow matchings*, *J. Combin. Theory Ser. A* **195** (2023), Paper No. 105700, 19. ↑2
- [39] C. McDiarmid and N. Yelov, *Hamilton cycles, minimum degree, and bipartite holes*, *J. Graph Theory* **86** (2017), no. 3, 277–285. ↑2
- [40] R. Montgomery, A. Myesser, and Y. Pehova, *Transversal factors and spanning trees*, *Adv. Comb.* (2022), Paper No. 3, 25. ↑2
- [41] O. Ore, *Note on Hamilton circuits*, *Amer. Math. Monthly* **67** (1960), 55. ↑2, 31
- [42] L. Pósa, *A theorem concerning Hamilton lines*, *Magyar Tud. Akad. Mat. Kutató Int. Közl* **7** (1962), 225–226. ↑2, 31
- [43] M. S. Rahman and M. Kaykobad, *On Hamiltonian cycles and Hamiltonian paths*, *Inform. Process. Lett.* **94** (2005), no. 1, 37–41. ↑2
- [44] Y. Tang, B. Wang, G. Wang, and G. Yan, *Rainbow Hamilton cycle in hypergraph system*, arXiv preprint arXiv:2302.00080 (2023). ↑2
- [45] I. M. Wanless, *Transversals in Latin squares: a survey*, *Surveys in combinatorics*, 2011, pp. 403–437. ↑1
- [46] D. R. Woodall, *Sufficient conditions for circuits in graphs*, *Proc. London Math. Soc. (3)* **24** (1972), 739–755. ↑31

(Y. CHENG) MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, OXFORD, UK

Email address: yangyang.cheng@maths.ox.ac.uk

(W. SUN) DATA SCIENCE INSTITUTE, SHANDONG UNIVERSITY, JINAN, 250100, CHINA

Email address: wtsun2018@sina.com

(G. WANG) SCHOOL OF MATHEMATICS, SHANDONG UNIVERSITY, JINAN, 250100, CHINA

Email address: ghwang@sdu.edu.cn

(L. WEI) SCHOOL OF MATHEMATICS, SHANDONG UNIVERSITY, JINAN, 250100, CHINA

Email address: lanwei@mail.sdu.edu.cn