

Adler function, Bjorken polarized Sum Rule – confirmation of elements of the $\{\beta\}$ -expansion and the diagrams.

S. V. Mikhailov^a

^a*Bogoliubov Laboratory of Theoretical Physics, JINR, 141980 Dubna, Russia*

E-mail: mikhs@theor.jinr.ru

ABSTRACT: Different ways exist to obtain the elements of the $\{\beta\}$ -expansion for renorm-group invariant quantities. Here we consider independent confirmation within the standard QCD of a number of our results [1] for the values of elements of this expansion for the nonsinglet Adler D_A -function, Bjorken polarized sum rules S^{Bjp} up to the order N⁴LO. We suggest an approach to estimate the results of high order QCD calculations using a smaller number of diagrams of the specific type. This type is based on a proposed generalization of Naive NonAbelianization.

KEYWORDS: Renormalization Group, QCD

July 22, 2024

Contents

1	Introduction	1
2	Test for $\{\beta\}$-expansion elements of $D_A(S^{Bjp})$ at $O(a_s^3)$	2
3	Test for $\{\beta\}$-expansion elements of D_A at $O(a_s^4)$	4
4	Reduction of $\{\beta\}$-expansion to a generalized NNA	7
4.1	Generalized NNA, illustration of β_0 -dominance	7
4.2	Conjecture of β_0 -dominance	9
5	Conclusion	10
A	QCD β-function	11
B	Numerical estimates of the elements $d_{3,4}[\cdot], c_{3,4}[\cdot]$ of $\{\beta\}$-expansion	12

1 Introduction

Here we will consider the results of $\{\beta\}$ -expansion [2] and discuss the reliability of the obtained elements of this expansion [1] and the relation of the latter with the corresponding Feynman diagrams. The elements of this expansion appear at the decomposition of the standard perturbative coefficients of any renormalization group invariants (RGIs); here these coefficients are d_n or c_n for the nonsinglet Adler D_A -function, or the Bjorken polarized sum rule (BpSR) S^{Bjp} respectively. We consider independent confirmation of our results for the decomposition of the nonsinglet Adler D_A -function for the values of elements $d_3[\cdot]$ ($c_3[\cdot]$) of $\{\beta\}$ -expansion and for some elements of $d_4[\cdot]$ [1]. The agreements are obtained using the already known results of N^{3,4}LO calculations [3] of some diagrams in the $\overline{\text{MS}}$ -scheme in pure QCD. In the case of the Adler nonsinglet D_A -function at the default choice of scale $Q^2 = \mu^2$,

$$D_A(a(\mu^2)) = 1 + \sum_{n \geq 1} a_s^n(\mu^2) d_n, \quad a_s(\mu^2) = \alpha_s(\mu^2)/(4\pi), \quad (1.1)$$

the decomposition of d_n looks like [1, 2],

$$a_s^2 d_2 = \beta_0 d_2[1] + d_2[0], \quad (1.2a)$$

$$a_s^3 d_3 = \beta_0^2 d_3[2] + \beta_1 d_3[0, 1] + \beta_0 d_3[1] + d_3[0], \quad (1.2b)$$

$$a_s^4 d_4 = \beta_0^3 d_4[3] + \beta_1 \beta_0 d_4[1, 1] + \beta_2 d_4[0, 0, 1] + \beta_0^2 d_4[2] + \beta_1 d_4[0, 1] + \beta_0 d_4[1] + d_4[0], \quad (1.2c)$$

... = ...

$$a_s^n d_n = \beta_0^{n-1} d_n[n-1] + \dots + d_n[0], \quad (1.2d)$$

where β_i are the coefficients of expansion of the QCD β -function (see Appendix A). The $\{\beta\}$ -expansion shows how the intrinsic charge renormalization manifests itself; thus all possible β_i -terms appear in each order n of perturbative expansion (PT). This property differ it from more or less formal expansions over powers of selected lagrangian parameters or Casimirs, say the number of active quark flavors n_f , or powers of β_0 [4], which is based on the same formal reasons. The procedure of calculating the expansion elements in (1.2) was based, in its turn, on the results of extended QCD (QCDe) presented in [5], which possesses a number of new degrees of freedom (d.o.f.). These d.o.f. are arbitrary number of different *fermion representations*, revealing themselves only in intrinsic loops. After decomposition as in (1.2), one can equate the contributions of these new d.o.f. to zero and return from QCDe to the standard QCD. Note here that now are at least two different approaches to obtain the values of the elements within the standard QCD that lead to different results. One of them, started with [6], is based on an interpretation of RG transformation, it got the name ‘‘Principle of Maximum Conformality’’ (PMC), see review [7] and references therein. The other approach, see [8], is based on an extended interpretation of Crewter-Broadharst-Kataev (CBK) relation [9]. So, some doubt appeared about how reliable the results for the elements obtained within QCDe in [1] are with respect to the standard QCD.

Here we confirm our results in order $O(a_s^3)$ and confirm them in part in order of $O(a_s^4)$ based on the calculation *within the standard* QCD and focusing on the diagrammatic origins of the elements of the $\{\beta\}$ -expansion. In addition in Sec.4 we propose a hierarchy of contributions to the coefficients $d_n(c_n)$ based on a certain generalization of Naive Non-abelianization (NNA) in the framework of the $\{\beta\}$ -expansion. The usage of this hierarchy can facilitate the laborious process of estimating physical quantities in high-lows calculations.

2 Test for $\{\beta\}$ -expansion elements of $D_A(S^{Bjp})$ at $O(a_s^3)$

Recall the elements of the $\{\beta\}$ -expansion for D_A up to the order $O(a_s^3)$ [1, 2, 10], where the first term $d_1 = 3C_F$:

$$d_2[1] = d_1 \left(\frac{11}{2} - 4\zeta_3 \right); \quad d_2[0] = d_1 \left(\frac{C_A}{3} - \frac{C_F}{2} \right); \quad (2.1)$$

$$d_3[2] = d_1 \left(\frac{302}{9} - \frac{76}{3}\zeta_3 \right); \quad d_3[0, 1] = d_1 \left(\frac{101}{12} - 8\zeta_3 \right); \quad (2.2a)$$

$$d_3[1] = d_1 \left[C_A \left(-\frac{3}{4} + \frac{80}{3}\zeta_3 - \frac{40}{3}\zeta_5 \right) - C_F (18 + 52\zeta_3 - 80\zeta_5) \right]; \quad (2.2b)$$

$$d_3[0] = d_1 \left[C_A^2 \left(\frac{523}{36} - 72\zeta_3 \right) + \frac{71}{3}C_A C_F - \frac{23}{2}C_F^2 \right]. \quad (2.2c)$$

These decompositions for d_3 were first obtained in [2] where one used the results for D_A with a single additional degree of freedom – the light gluino [11], then the elements of this decomposition were confirmed in [1] within QCDe with any numbers of fermion d.o.f. Below

we reproduce the same values of the elements $d_3[\cdot]$ by considering the results for specific diagrams for D_A calculated in *pure* QCD.

Let us start with the simplest case in (2.1). The first element $d_2[1]$ originates from the diagram where the single gluon line is dressed in a quark bubble, which is proportional to $T_R n_f \sim \beta_0$. Just this correspondence allows one to perform the decomposition unambiguously. The sequential dressing of this only one gluon line generates its anomalous dimension that is finally revealed as a set of β -elements of our interest also in the higher orders of PT. In the next order the 2nd loop dressing of only one gluon line generates




Figure 1. Diagrams for D_A in order a_s^3 with self-energy dressing of only one gluon line (red in color). One should add to these samples of diagrams all possible connections of gluon lines. **Left.** proportional to $C_F \cdot (T_R n_f \equiv x)^2, \sim \beta_0^2$. **Right.** proportional to $C_F^2 (T_R n_f \equiv x), \sim \beta_1$.

specific diagrams with two 1-loop bubbles (reducible), see Fig.1(Left) and with 2-loop one (irreducible), Fig.1(Right).

We reproduced in Fig.1 the diagrams taken from Fig.C1(IV1) and Fig.C1(IV2) in [3], and the abelian projection of the results corresponding to them was presented in Appendix C there. The result for the diagram in Fig.1(Left) independently confirms the value of the $d_3[2]$ term in (2.2a) and coincides also with the general expression for $d_n[n-1]$, which can be extracted from the results of [9, 12]. The diagram in the right panel contributes to the term $\beta_1 d_3[0,1]$ by means of anomalous dimension of a gluon in order of a_s^2 . The latter, in its turn, includes the unique Casimir $C_F \cdot (T_R n_f \equiv x)$, which is common with β_1 , see Eq.(A.2b). This diagrammatic correspondence unambiguously determines the element $d_3[0,1]$ within the standard QCD calculation. Further we will trace the contributions to d_n of the color structures starting with maximum numbers of quark loops: the evident case of a renormalon chain with only one gluon line contribution gives us $C_F \cdot x^{(n-1)}$, then one fermion loop less – $C_F(C_F x^{(n-2)})$, then $C_F(C_F^2 x^{(n-3)})$ and so on. In other words, these terms are a projection onto the abelian sector of the QCD calculation (proportional to QED) of D and S^{Bjp} . The abelian part of the structure of β -coefficients is well recognized in the first terms (underlined) of the corresponding formulas in Appendix A.

Now we will show that independent knowledge of $d_3[0,1]$ guarantees also independent determination of two remainder elements $d_3[1], d_3[0]$ of this order if the coefficient d_3 as a whole is already known. Using the definition $T_R n_f = x$, we introduce [13] the consequence of roots $x_0, x_1, \dots \sim x$ of the QCD β -function coefficients $\beta_0(x_0) = 0, \beta_1(x_1) = 0, \dots$, respectively,

$$x_0 = \frac{11}{4}C_A; \quad (2.3a)$$

$$\beta_0(x_0) = 0; \beta_1(x_0) = -C_A(7C_A + 11C_F); \beta_2(x_0) = C_A \left(\frac{11}{2}C_F^2 - \frac{1127}{24}C_A^2 - \frac{77}{3}C_A C_F \right); \quad (2.3b)$$

$$x_1 = \frac{17}{2} \frac{C_A^2}{5C_A + 3C_F}; \quad (2.3c)$$

$$\beta_0(x_1) = C_A \frac{7C_A + 11C_F}{5C_A + 3C_F}; \beta_1(x_1) = 0; \beta_2(x_1) = \dots \quad (2.3d)$$

Introducing a reduced $\tilde{d}_3(x) \equiv d_3(x) - \beta_0^2(x)d_3[2]$, depending on the argument x , we obtain from the $\{\beta\}$ -expansion for d_3 (Eq.(1.3) in [1]),

$$\tilde{d}_3(x_0) - \beta_1(x_0)d_3[0, 1] = d_3[0]; \quad (2.4a)$$

$$\frac{\tilde{d}_3(x_0) - \tilde{d}_3(x_1)}{\beta_0(x_1)} + \frac{\beta_1(x_0)}{\beta_0(x_1)}d_3[0, 1] = d_3[1], \quad (2.4b)$$

where $d_3(x)$ (and $d_4(x)$) are known from the original calculations [14]. So we get *independent confirmation* within QCD of all the results for $d_3[.]$ displayed in Eqs.(2.2).

It is clear that the same relations, like Eq.(2.4), is true also for the elements $c_3[.]$ in the $\{\beta\}$ -expansion of S^{Bjp} . The required value of $c_3[0, 1]$ can be obtained from $d_3[0, 1]$ based on the CBK relation [1, 15],

$$c_2[1] + d_2[1] = c_3[0, 1] + d_3[0, 1] = c_n \underbrace{[0, 0, \dots, 1]}_{n-1} + d_n \underbrace{[0, 0, \dots, 1]}_{n-1} = 3C_F \left(\frac{7}{2} - 4\zeta_3 \right). \quad (2.5)$$

Therefore, the knowledge of this $d_3[0, 1]$ makes it possible to restore both $d_3[.]$ and $c_3[.]$ only within QCD calculations and confirms the results obtained with the help of QCDe in [1], and within QCD with light gluino [2].

3 Test for $\{\beta\}$ -expansion elements of D_A at $O(a_s^4)$

Here we discuss independent tests to verify the elements of $\{\beta\}$ -expansion for $d_4(c_4)$ obtained in [1] on the basis of the results presented in [5] within QCDe. These results for the $d_4[.]$ elements are presented below,

$$d_4[3] = C_F \left(\frac{6131}{9} - 406\zeta_3 - 180\zeta_5 \right); \quad (3.1a)$$

$$d_4[1, 1] = C_F \left(385 - \frac{1940}{3}\zeta_3 + 144\zeta_3^2 + 220\zeta_5 \right); \quad (3.1b)$$

$$d_4[0, 0, 1] = C_F \left(\frac{355}{6} + 136\zeta_3 - 240\zeta_5 \right); \quad (3.1c)$$

$$d_4[2] = -C_F \left[C_F \left(\frac{6733}{8} + 1920\zeta_3 - 3000\zeta_5 \right) + C_A \left(\frac{20929}{144} - \frac{12151}{6}\zeta_3 + 792\zeta_3^2 + 1050\zeta_5 \right) \right]; \quad (3.1d)$$

$$d_4[1] = C_F \left[-C_F^2 \left(\frac{447}{2} - 42\zeta_3 - 4920\zeta_5 + 5040\zeta_7 \right) + C_A C_F \left(\frac{3301}{4} - 678\zeta_3 - 2280\zeta_5 + 2520\zeta_7 \right) + C_A^2 \left(\frac{16373}{36} - \frac{17513}{3}\zeta_3 + 2592\zeta_3^2 + 3030\zeta_5 - 420\zeta_7 \right) \right]; \quad (3.1e)$$

$$d_4[0, 1] = -C_F \left[C_A \left(\frac{139}{12} + \frac{1054}{3}\zeta_3 - 460\zeta_5 \right) + C_F \left(\frac{251}{4} + 144\zeta_3 - 240\zeta_5 \right) \right]; \quad (3.1f)$$

$$d_4[0] = \tilde{d}_4[0] + \delta d_4 = C_F^4 \left(\frac{4157}{8} + 96\zeta_3 \right) - C_A C_F^3 \left(\frac{2409}{2} + 432\zeta_3 \right) + C_A^2 C_F^2 \left(\frac{3105}{4} + 648\zeta_3 \right) + C_A^3 C_F \left(\frac{68047}{48} + \frac{8113}{2}\zeta_3 - 7110\zeta_5 \right) + \delta d_4; \quad (3.1g)$$

$$\delta d_4 = -16 \left[n_f \frac{d_F^{abcd} d_F^{abcd}}{d_F} (13 + 16\zeta_3 - 40\zeta_5) + \frac{d_A^{abcd} d_F^{abcd}}{d_F} (-3 + 4\zeta_3 + 20\zeta_5) \right]. \quad (3.1h)$$

Equation(3.1a) for $d_4[3]$ can also be independently confirmed by the partial result for $\beta_0^3 d_4[3]$ in Eq.(C.4) in [3], see Fig.C1(V1) there, which agrees with the results in [9, 12] too. Here we reproduce these “abelian” diagrams in Fig.2(see left panel) for reader’s convenience.

(i)The projection of the weighted sum $\beta_2 d_4[0, 0, 1] + \beta_1 \beta_0 d_4[1, 1]$ onto the Casimir $C_F^2 x^2$ is denote as $s_4 = \hat{P}_{C_F^2 x^2} (\beta_2 d_4[0, 0, 1] + \beta_1 \beta_0 d_4[1, 1])$, where $\hat{P}_{C_F^2 x^2}$ means the corresponding projector. The evident diagrammatic origins just of these terms in s_4 are presented in Fig.C1(V2,V3) [3] that is reproduced in Fig.2(Right) here, the corresponding partial results are presented in Eqs.(C5, C6) [3]. The results for each of these two diagrams, evidently related to the factors β_2 , $\beta_1 \beta_0$ due to gluon line renormalizations, contributes to both the terms in the sum s_4 . The sum of the partial results of the direct QCD calculation of these




Figure 2. The types of the “abelian” diagrams for D_A in order a_s^4 of the self-energy type of only one gluon line (red in color) also include the symmetric diagram where the inner loops are interchanged. **Left:** proportional to $C_F x^3$, $\sim \beta_0^3$ **Right:** proportional to $C_F^2 x^2$: $\Rightarrow \beta_2, \beta_1 \beta_0$

diagrams reads in the lhs¹ of the equation as

$$C_F^2 x^2 \left(\frac{63250}{27} - 2784\zeta_3 + 768\zeta_3^2 \right) = s_4 \equiv \hat{P}_{C_F^2 x^2} (\beta_2 d_4[0, 0, 1] + \beta_1 \beta_0 d_4[1, 1]) \quad (3.2)$$

$$= \left(\frac{44}{9} C_F x^2 \right) d_4[0, 0, 1] + (-4 C_F x) \left(-\frac{4}{3} x \right) d_4[1, 1].$$

¹up to the factor 4^{-4} in the lhs due to another definition of a_s , $a_s = \alpha_s/\pi$ in [3]

The abelian projections of the β -coefficients in the last equation correspond to the underlined terms in Eq.(A.2) (Appendix A). Substituting the corresponding elements $d_4[.]$ from Eqs.(3.1b,3.1c) into the rhs, one obtains the equality with the lhs; at the same time, the terms with ζ_5 that exist in each of the elements $d_4[0, 0, 1], d_4[1, 1]$ cancel in the sum s_4 . So the sum of these elements *is confirmed*. Recall that $\beta_2 d_4[0, 0, 1] + \beta_1 \beta_0 d_4[1, 1]$ corresponds to special kinds of dressing of only one gluon line in Fig. 2, but it does not exhaust all the contributions to $C_F^2 x^2$.

(ii) To obtain the completed projection on $C_F^2 x^2$, presented in the lhs of (3.3), see [14],

$$C_F^2 x^2 \left(\frac{5713}{27} - \frac{4648}{3} \zeta_3 + 192 \zeta_3^2 + \frac{4000}{3} \zeta_5 \right) = \hat{P}_{C_F^2 x^2} \left(\beta_2 d_4[0, 0, 1] + \beta_1 \beta_0 d_4[1, 1] + \beta_0^2 d_4[2] \right), \quad (3.3)$$

one should supplement s_4 with the contribution $\hat{P}_{C_F^2 x^2} (\beta_0^2 d_4[2])$ in the rhs. The diagrams forming this additional term are not of “the only one dressed gluon line” type. Equality (3.3) unambiguously determines the abelian part of the element $d_4[2]$ and confirms now Eq.(3.1d). Concluding we confirm the values of the elements $d_4[0, 0, 1], d_4[1, 1]$ and the abelian part of $d_4[2]$ based on the direct QCD calculations ²

(iii) In the next step we will consider the contribution to $C_F (C_F^2 x)$ that is known from [14], see the lhs of (3.4a),

$$\begin{aligned} C_F^3 x \left(\frac{2002}{3} + 792 \zeta_3 - 8000 \zeta_5 + 6720 \zeta_7 \right) &= \hat{P}_{C_F^3 x} \left(\beta_2 d_4[0, 0, 1] + \beta_1 d_4[0, 1] + \beta_0 d_4[1] \right) \\ &= \hat{P}_{C_F^3 x} \left[(2C_F^2 x) d_4[0, 0, 1] + (-4C_F x) d_4[0, 1] + \left(-\frac{4}{3} x \right) d_4[1] \right], \end{aligned} \quad (3.4a)$$

while its rhs is fulfilled by the weighted sum. Of course, the values of elements $d_4[.]$ in Eq.(3.1) satisfy this equality. Assuming that the value $d_4[0, 0, 1]$ is already confirmed from the condition (3.2) in item (i), one obtains an independent test for the partial sum $\hat{P}_{C_F^3 x} (\beta_1 d_4[0, 1] + \beta_0 d_4[1])$ of the remaining elements,

$$C_F^3 x (549 + 520 \zeta_3 - 7520 \zeta_5 + 6720 \zeta_7) = -\hat{P}_{C_F^3 x} \left[(4C_F x) d_4[0, 1] + \left(\frac{4}{3} x \right) d_4[1] \right]. \quad (3.4b)$$

To complete the confirmation for the 5-loop $d_4[.]$ elements, we need at least 3 independently calculated elements, e.g., $d_4[0, 0, 1], d_4[1, 1]$ and, say, $d_4[0, 1]$. In other words, taking into consideration the Casimir content of these elements, this parameterization consist of 4 independent scalar parameters³. So to obtain the important element $d_4[0](c_4[0])$, one should take $d_4(x_0)$ or $c_4(x_0)$, all the powers of β_0 cancel in the corresponding $\{\beta\}$ -expansion, see Eq.(1.2c,2.3b), and one arrives at

$$d_4[0] = d_4(x_0) - \beta_2(x_0) d_4[0, 0, 1] - \beta_1(x_0) d_4[0, 1], \quad (3.5a)$$

$$c_4[0] = c_4(x_0) - \beta_2(x_0) c_4[0, 0, 1] - \beta_1(x_0) c_4[0, 1]. \quad (3.5b)$$

²The corresponding values of $d_4[0, 0, 1], d_4[1, 1]$ presented in [8], and [16], [17] (Table 2 there) do not agree with the lhs in (3.2), and the abelian part of $d_4[2]$ does not agree with (3.3).

³a similar parameterization was considered by A.Kataev, see A. Kataev’s talk at the session of RAS&JINR on 03.04.2024, <https://indico.jinr.ru/event/4174/contributions/25722/>

It is important if the elements $d_4[0, 0, 1], d_4[0, 1]$ are already known, then the counter part elements $c_4[0, 0, 1], c_4[0, 1]$ are also known from the corollaries [1] of the CBK relation [1, 5] like Eq.(2.5).

Let us mention here a rather rough test for elements $d_{3,4}[0]$ that is related to a special choice of Casimirs C_A, C_F , see (A.3). The coefficients of perturbative expansion based on $SU_c(N)$ dynamics are the polynomials over the C_A, C_F , let us consider these Casimirs as variables, which are related by the condition $7C_A + 11C_F = 0$ (that formally corresponds to the value $N = \sqrt{11}/5$). This linear condition, see Eqs.(2.3b, 2.3d), leads to the evident nullification of the coefficients β_0, β_1 ,

$$\beta_0(x_0) = \beta_1(x_0) = \beta_0(x_1) = \beta_1(x_1) = 0. \quad (3.6)$$

This condition radically simplifies the color structure of elements and in virtue of (3.6) leads

at $7C_A + 11C_F = 0$:

$$d_3(x_0) = d_3[0] = C_A^3 (22607 + 313632\zeta_3) \frac{7}{15972}; \quad (3.7)$$

$$d_4(x_0) = d_4[0] + \beta_2(x_0)d_4[0, 0, 1] = C_A^4 \left(\frac{308178983}{351384} + \frac{7782229}{29282}\zeta_3 + \frac{22680}{121}\zeta_5 \right) + \delta d_4. \quad (3.8)$$

The Eqs.(2.2c) for $d_3[0]$, and (3.1c) for $d_4[0, 0, 1]$, (3.1g) for $d_4[0]$ as well as the corresponding results in [17] satisfy both the tests in Eqs.(3.7,3.8).

4 Reduction of $\{\beta\}$ -expansion to a generalized NNA

Here we consider a conjecture about the hierarchy of contributions to the perturbation coefficients d_n, c_n based on the $\{\beta\}$ -expansion, which generalizes the well-known NNA [18, 19]. It also generalizes the results of the β_0^n -expansion [4] and the $(\beta_0^n, \beta_1\beta_0^{n-2})$ expansion in [3]. This hierarchy rearranges the elements into a group that can provide the main contribution to the coefficients of the expansion $d_n(c_n)$. It was first introduced in [2] (Sec.4) and was discussed in [10, 20]. We will discuss the relations between the main term of this hierarchy, the topological structure of the diagrams corresponding to it in the abelian limit, and the conditions on the indices of the involved elements of the $\{\beta\}$ -expansion.

4.1 Generalized NNA, illustration of β_0 -dominance

Here we will use the ordering of contributions in powers of β_0 under the condition of “large β_0 ” (in QCD $\beta_0 \sim 10$). More precisely, the common factor β_0^{n-1} in any order n will be taken out of the expansion coefficient d_n to order all the elements over the inverse powers of $1/(\beta_0)^i$ [2]. Recall that the well-known BLM [21] trick that was a predecessor of the $\{\beta\}$ -expansion, assumes for its efficiency the condition $|\beta_0 d_2[1]| \gtrsim d_2[0]$, see the corresponding ratios in Eq.(4.1).

Then, we follow the “practical” observation $\beta_i = O(\beta_0^{i+1})$ that works at $n_f = 0 \div 5$, $i = 1 \div 4$, and invent the notation $b_i \stackrel{def}{=} \beta_i/\beta_0^{i+1} = O(1)$, see (A.5). Of course, the last equality should be broken down at higher orders i of pQCD due to the factorial growth of

β_i . So joining both these conditions, we take into account the terms with the coefficients b_i together with others in the same order of $1/(\beta_0)^i$, see the first brackets in Eqs.(4.2) and the first and the second brackets in Eqs.(4.3) below. We call this kind of rearrangement in the groups of the same $1/\beta_0$ -order in brackets the generalized NNA (gNNA). The expansion coefficients of the D_A -function and BpSR S^{Bjp} are presented below as the sums of these groups accompanied by their numerical estimates (at $n_f = 3$, NNA estimates are shown from below in red),

$$d_2 = \beta_0 \left[\underbrace{d_2[1]}_{\mathbf{2.77}} + \frac{\mathbf{0.15}}{\beta_0} d_2[0] \right] \approx \beta_0 [\mathbf{2.92}]; \quad c_2 = \beta_0 \left[\underbrace{c_2[1]}_{\mathbf{-8}} + \frac{\mathbf{1.6}}{\beta_0} d_2[0] \right] \approx \beta_0 [\mathbf{-6.4}] \quad (4.1)$$

$$d_3 = \beta_0^2 \left[\underbrace{\left(d_3[2] + b_1 d_3[0, 1] \right)}_{\mathbf{12.4}} + \frac{\mathbf{8.6}}{\beta_0} d_3[1] + \frac{\mathbf{32.6}}{\beta_0^2} d_3[0] \right] \approx \beta_0^2 [\mathbf{12.9}] \quad (4.2a)$$

$$c_3 = \beta_0^2 \left[\underbrace{\left(c_3[2] + b_1 c_3[0, 1] \right)}_{\mathbf{-25.6}} + \frac{\mathbf{-25.9}}{\beta_0} c_3[1] + \frac{\mathbf{-17.8}}{\beta_0^2} c_3[0] \right] \approx \beta_0^2 [\mathbf{-16}] \quad (4.2b)$$

$$d_4 = \beta_0^3 \left[\underbrace{\left(d_4[3] + b_1 d_4[1, 1] + b_2 d_4[0, 0, 1] \right)}_{\mathbf{8.7}} + \frac{\mathbf{24}}{\beta_0} (d_4[2] + b_1 d_4[0, 1]) + \frac{\mathbf{34.3}}{\beta_0^2} d_4[1] + \frac{\mathbf{-6.8}}{\beta_0^3} d_4[0] \right] \\ \approx \beta_0^3 [\mathbf{17.2}] \quad (4.3a)$$

$$c_4 = \beta_0^3 \left[\underbrace{\left(c_4[3] + b_1 c_4[1, 1] + b_2 c_4[0, 0, 1] \right)}_{\mathbf{-89.6}} + \frac{\mathbf{-112}}{\beta_0} (c_4[2] + b_1 c_4[0, 1]) + \frac{\mathbf{-95.5}}{\beta_0^2} c_4[1] + \frac{\mathbf{136.5}}{\beta_0^3} c_4[0] \right] \\ \approx \beta_0^3 [\mathbf{-61.7}] \quad (4.3b)$$

As it is seen, the leading $(1/\beta_0)^0$ -groups numerically dominate in all of the sums in (4.2,4.3), while the subleading groups for the most part mutually cancel one another, compare the lhs and the rhs there. The evident manifestation of gNNA vs NNA can be seen in the leading group of (4.3a). The numerical estimates of each of the elements $d_{3,4}[\cdot], c_{3,4}[\cdot]$ are shown in Appendix B, where the diagrammatically confirmed results are marked in red. In addition, just the result of this leading group has been independently confirmed here in Sec.3, see the rhs of Eq.(3.2), which was based on the calculations of the QCD diagrams presented in Fig.2. Therefore, the correspondence of gNNA and the total sum in (4.3a) does not even depend on the detailed results in Eqs.(3.1b,3.1c) obtained with the help of QCDe. On the other hand, the contributions of the suppressed in $1/\beta_0$ the so-called ‘‘scale invariant’’ terms $d_{3,4}[0]$ almost cancel with the other subleading groups. At the same time, these $d_i[0](c_i[0])$ do not correspond by themselves to the final sum in the rhs neither in value nor in sign

⁴. We recognize that our conjecture is based on observations of a few examples only, and the reason of the aforementioned mutual cancellations is not clear; nevertheless, it looks reasonable to formulate the conjecture for the next orders of pQCD.

4.2 Conjecture of β_0 -dominance

In the case of 6 loop [1], the leading gNNA group is presented in the first line in Eq.(4.4a) below. Following our conjecture, one needs to calculate *restricted types* of the corresponding diagrams *to estimate* the complete result d_5 based on (4.4a),

$$d_5 = \beta_0^4 \left[\overbrace{\left(d_5[4] + b_1 d_5[2, 1] + b_1^2 d_5[0, 2] + b_2 d_5[1, 0, 1] + b_3 d_5[0, 0, 0, 1] \right)}^{p(5-1)} \right] + \quad (4.4a)$$

$$\frac{1}{\beta_0} \left(d_5[3] + b_1 d_5[1, 1] + b_2 d_5[0, 0, 1] \right) + \frac{1}{\beta_0^2} \left(d_5[2] + b_1 d_5[0, 1] \right) + \quad (4.4b)$$

$$\frac{1}{\beta_0^3} d_5[1] + \frac{1}{\beta_0^4} d_5[0] \Big]. \quad (4.4c)$$

The number of elements of the leading group in Eq.(4.4a) is equal to the number of partitions p depending on perturbation order n , *e.g.*, here $-p(5-1) = \mathbf{5}$ [22], see the detailed discussion of this correspondence in [1], Sec.4. So taking a natural series for the argument of the function p , one obtains $p(1, 2, 3, 4, 5, 6, 7, \dots) = 1, 2, 3, \mathbf{5}, 7, 11, 15, \dots$. For the order n of pQCD the leading gNNA group consists of the number of partitions $p(n-1)$ elements that can be traced from the dressing of *only one gluon line of the diagram*. The indices of the elements $d_n[j_1, \dots, j_{n-1}]$ of the leading group satisfy the evident equality $1j_1 + 2j_2 + \dots + (n-1)j_{n-1} = n-1$, while for the different subleading groups the conditions for the indices are $1j_1 + 2j_2 + \dots + (n-1)j_{n-1} = n-2, n-3, \dots$, where the shift from the number $n-1$ in the rhs determines the order of $1/\beta_0$ suppression see, *e.g.*, the groups in (4.4b, 4.4c).

If the elements of the gNNA group in (4.4a) are already obtained, they can be easily checked following the topological types of “abelian” diagrams in Fig.3, where the chain of n




Figure 3. The types of “abelian” diagrams corresponding to gNNA in order a_s^5 with all possible interchanges of the inner loops. (a) $C_F x^4 \sim \beta_0^4$; $C_F x^3$: (b), (c), (d) $\Rightarrow \beta_3, \beta_0 \beta_2, \beta_0^2 \beta_1$; $C_F^2 x^2$: (e), (f) $\Rightarrow \beta_1 \beta_1, \beta_3$.

bubbles in Fig.3(a) definitely corresponds to only $d_n[n-1]$. The other diagrams – (b, c, d) contribute to different combinations of the β -coefficients admissible for the leading group

⁴The authors of PMC, see [7] and references therein, propagate another, important role of the “scale invariant” elements $d_n[0]$ within the PMC approach.

that has a projection on $C_F \cdot x^3$, while the diagrams $(e, f) \sim C_F^2 \cdot x^2$. So one has

$$\text{sum of diag. (b,c,d)} = P_{C_F x^3} (\beta_3 d_5[0, 0, 0, 1] + \beta_2 \beta_0 d_5[1, 0, 1] + \beta_1 \beta_0^2 d_5[2, 1]) \quad (4.5)$$

$$\text{sum of diag. (e,f)} = P_{C_F^2 x^2} (\beta_3 d_5[0, 0, 0, 1] + \beta_1^2 d_5[0, 2]) \quad (4.6)$$

It is clear that the diagrams similar those in Figs. 2, 3 for the leading gNNA group can also be easily identified for the twist 2 coefficient function C_{Bjp} of the observable S^{Bjp} .

To conclude, we present the leading gNNA for any order $n \geq 4$ as a weighted sum of $p(n-1)$ elements,

$$d_n = \beta_0^{n-1} \left[\overbrace{\left(d_n[n-1] + b_1 d_n[n-3, 1] + \dots + b_{(n-2)} d_n \underbrace{[0, \dots, 1]}_{n-1} \right)}^{p(n-1)} + \dots + \frac{d_n[0]}{\beta_0^{n-1}} \right]. \quad (4.7)$$

This group corresponds to the diagrams with only one dressed gluon line like those presented in Fig. 3. On the contrary, the subleading gNNA groups correspond to the diagrams with more than one number of (dressed) gluon lines. It should be noted that taking into account the first two terms in the leading group of (4.7) to improve the BLM was first proposed in [3].

Using another language based on the hierarchy of n_f orders, the application of gNNA requires taking into account in addition to the leading powers of n_f , a subseries of subleading orders also originating from different sources.

5 Conclusion

We have confirm our results for the elements of the $\{\beta\}$ -expansion in [1, 2]: in order of $O(a_s^3)$ – completely; in order of $O(a_s^4)$ we confirm two elements, $d_4[0, 0, 1]$, $d_4[1, 1]$, and the abelian part of $d_4[2]$ from [1], and for both the cases – basing on the known calculation [3] within the standard QCD only.

We proposed a conjecture of β_0 -dominance in d_n . The weighted sum of such elements $d_n[j_1, j_2, \dots, j_{n-1}]$ dominates, whose indices satisfy the condition: $1j_1 + 2j_2 + \dots + (n-1)j_{n-1} = n-1$, their contributions are not suppressed by powers of $1/\beta_0$. This condition one to one corresponds to allowing for the topological class of diagrams with renormalization of only one gluon line. The conjecture, in a lucky case, can significantly reduce the number of diagrams considered and make the laborious process of estimating physical quantities in high loops of calculations easier.

Acknowledgments

The author is thankful to A. L. Kataev for drawing my attention to ref. [3] and also for the constructive criticism of the paper, to N. Volchanskiy for the fruitful discussions. I am very grateful to K. Chetyrkin for the clarifying the important items of this consideration.

A QCD β -function

The well-known renormalization group equation for the QCD coupling constant $a_s(\mu^2) = \frac{\alpha_s(\mu^2)}{4\pi}$ reads

$$\mu^2 \frac{da_s(\mu^2)}{d\mu^2} = -\beta(a) = -a_s^2(\mu^2) \sum_{i \geq 0} \beta_i a_s^i(\mu^2). \quad (\text{A.1})$$

The explicit expressions for the first coefficients of the β -function expansion with the underlined abelian terms read

$$\beta_0 = -\frac{4}{3} \underline{T_R n_f} + \frac{11}{3} C_A; \quad (\text{A.2a})$$

$$\beta_1 = -4 \underline{C_F T_R n_f} + \frac{34}{3} C_A^2 - \frac{20}{3} C_A T_R n_f; \quad (\text{A.2b})$$

$$\begin{aligned} \beta_2 = & \frac{44}{9} \underline{C_F (T_R n_f)^2} + 2 \underline{C_F^2 T_R n_f} + \frac{2857}{54} C_A^3 - \frac{205}{9} C_F C_A T_R n_f - \frac{1415}{27} C_A^2 T_R n_f \\ & + \frac{158}{27} C_A (T_R n_f)^2; \end{aligned} \quad (\text{A.2c})$$

$$\begin{aligned} \beta_3 = & \frac{1232}{243} \underline{C_F (T_R n_f)^3} + \underline{(C_F T_R n_f)^2} \left(\frac{1352}{27} - \frac{704}{9} \zeta_3 \right) + 46 \underline{(C_F)^3 T_R n_f} + C_A^3 T_R n_f \left(-\frac{39143}{81} + \frac{136}{3} \zeta_3 \right) \\ & + C_A C_F \underline{T_R^2 n_f^2} \left(\frac{17152}{243} + \frac{448}{9} \zeta_3 \right) + C_A C_F^2 T_R n_f \left(-\frac{4204}{27} + \frac{352}{9} \zeta_3 \right) + \frac{424}{243} C_A T_R^3 n_f^3 \\ & + C_A^2 C_F T_R n_f \left(\frac{7073}{243} - \frac{656}{9} \zeta_3 \right) + C_A^2 T_R^2 n_f^2 \left(\frac{7930}{81} + \frac{224}{9} \zeta_3 \right) + C_A^4 \left(\frac{150653}{486} - \frac{44}{9} \zeta_3 \right) \\ & + n_f \frac{d_F^{abcd} d_A^{abcd}}{N_A} \left(\frac{512}{9} - \frac{1664}{3} \zeta_3 \right) + n_f^2 \frac{d_F^{abcd} d_F^{abcd}}{N_A} \left(-\frac{704}{9} + \frac{512}{3} \zeta_3 \right) \\ & + \frac{d_A^{abcd} d_A^{abcd}}{N_A} \left(-\frac{80}{9} + \frac{704}{3} \zeta_3 \right), \end{aligned} \quad (\text{A.2d})$$

with the $SU_c(N)$ -group fundamental fermion invariants

$$T_R = \frac{1}{2}; \quad C_F = T_R \frac{N^2 - 1}{N}; \quad C_A = N; \quad N_A = 2C_F C_A \equiv N^2 - 1; \quad (\text{A.3})$$

$$\begin{aligned} d^{abc} d^{abc} &= \frac{(N^2 - 4)}{N} N_A; \quad d_F^{abcd} d_A^{abcd} = \frac{N(N^2 + 6)}{48} N_A; \\ d_F^{abcd} d_F^{abcd} &= \frac{N^4 - 6N^2 + 18}{96N^2} N_A; \quad d_A^{abcd} d_A^{abcd} = \frac{N^2(N^2 + 36)}{24} N_A, \end{aligned} \quad (\text{A.4})$$

d_R is the dimension of the quark color representation, $d_R = 3$ in QCD and n_f denotes the number of active flavors. At $n_f = 3$,

$$\beta_0 = 9; \quad b_1 \approx 0.79; \quad b_2 \approx 0.88; \quad b_3 \approx 1.9. \quad (\text{A.5})$$

B Numerical estimates of the elements $d_{3,4}[\cdot], c_{3,4}[\cdot]$ of $\{\beta\}$ -expansion

We present the numerical estimates for the each of elements, discussed in Sec.4 for reader's convenience. For the expressions $d_3[\cdot]$ in Eq.(2.2) and similar ones $c_3[\cdot]$ [1] we have [23, 24]

$$d_3[2] \approx \mathbf{12.41}, \quad d_3[0, 1] \approx \mathbf{-4.799}, \quad d_3[1] \approx 293.55, \quad d_3[0] \approx -2295.84, \quad (\text{B.1a})$$

$$c_3[2] \approx \mathbf{-25.56}, \quad c_3[0, 1] \approx \mathbf{-0.43}, \quad c_3[1] \approx -159.84, \quad c_3[0] \approx 2242.51. \quad (\text{B.1b})$$

From the expressions for $d_4[\cdot]$ in Eq.(3.1) one has

$$d_4[3] \approx \mathbf{8.720}, \quad d_4[1, 1] \approx \mathbf{58.487}, \quad d_4[0, 0, 1] \approx \mathbf{-34.955}, \quad (\text{B.2a})$$

$$d_4[2] \approx 154.47, \quad d_4[0, 1] \approx 195.46, \quad d_4[1] \approx -550.54, \quad (\text{B.2b})$$

$$d_4[0] \approx -25105.61 + 20.5427 n_f \approx -25043.97 \text{ (at } n_f = 3) \quad (\text{B.2c})$$

$$c_4[3] \approx \mathbf{-89.63}, \quad c_4[1, 1] \approx -61.647, \quad c_4[0, 0, 1] \approx \mathbf{29.722}, \quad (\text{B.3a})$$

$$c_4[2] \approx -707.165, \quad c_4[0, 1] \approx -193.028, \quad c_4[1] \approx 11053.7, \quad (\text{B.3b})$$

$$c_4[0] \approx 6871.02 \text{ (at } n_f = 3) \quad (\text{B.3c})$$

The elements marked in red in (B.1,B.2,B.3) have been confirmed, at least in their abelian part, by the diagrammatical calculations in the standard QCD in Secs.2, 3. The widely debated “zero” element $d_4[0]$ reads $-98.069 + 0.0802 n_f$ in the case of the using the coupling constant α_s/π instead of a_s .

References

- [1] P. A. Baikov and S. V. Mikhailov, “The β -expansion for Adler function, Bjorken Sum Rule, and the Crewther-Broadhurst-Kataev relation at order $O(\alpha_s^4)$ ”, JHEP **09** (2022), 185 doi:10.1007/JHEP09(2022)185 [arXiv:2206.14063 [hep-ph]], Addendum, JHEP **03** (2023),053 doi:10.1007/JHEP03(2023)053
- [2] S. V. Mikhailov, “Generalization of BLM procedure and its scales in any order of pQCD: A Practical approach”, JHEP **2007**, 009 (2007), doi: 10.1088/1126-6708/2007/06/009, [hep-ph/0411397].
- [3] Ball, Patricia and Beneke, M. and Braun, Vladimir M., “Resummation of $(\beta_0\alpha_s)^n$ corrections in QCD: Techniques and applications to the τ hadronic width and the heavy quark pole mass”, Nucl. Phys. **B**, 452 (1995)563; [hep-ph/9502300]
- [4] C. N. Lovett-Turner and C. J. Maxwell, “All orders renormalon resummations for some QCD observables”, Nucl. Phys. B **452** (1995), 188-212, doi:10.1016/0550-3213(95)00383-4, [arXiv:hep-ph/9505224].
- [5] K. G. Chetyrkin, “Adler function, Bjorken Sum Rule and Crewther-Broadhurst-Kataev relation with generic fermion representations at order $O(\alpha_s^4)$,” Nucl. Phys. B **985** (2022), 115988 doi:10.1016/j.nuclphysb.2022.115988 [arXiv:2206.12948 [hep-ph]].
- [6] M. Mojaza, S. J. Brodsky and X. G. Wu, “Systematic All-Orders Method to Eliminate Renormalization-Scale and Scheme Ambiguities in Perturbative QCD,” Phys. Rev. Lett. **110** (2013), 192001 doi:10.1103/PhysRevLett.110.192001 [arXiv:1212.0049 [hep-ph]].

- [7] Di Giustino, S. J. Brodsky, P. G. Ratcliffe, X.-G. Wu, S.-Q. Wang, “High precision tests of QCD without scale or scheme ambiguities: The 40th anniversary of the Brodsky-Lepage-Mackenzie method”, *Prog. Part. Nucl. Phys.* **135**, 104092 (2024), [[2307.03951](#)].
- [8] G. Cvetič and A. L. Kataev, “Adler function and Bjorken polarized sum rule: Perturbation expansions in powers of the $SU(N_c)$ conformal anomaly and studies of the conformal symmetry limit,” *Phys. Rev. D* **94** (2016) no.1, 014006 doi:10.1103/PhysRevD.94.014006 [arXiv:1604.00509 [hep-ph]].
- [9] D. J. Broadhurst and A. L. Kataev, “Connections between deep inelastic and annihilation processes at next to next-to-leading order and beyond,” *Phys. Lett. B* **315** (1993) 179.
- [10] A. L. Kataev and S. V. Mikhailov, “Generalization of the Brodsky-Lepage-Mackenzie optimization within the $\{\beta\}$ -expansion and the principle of maximal conformality”, *Phys. Rev. D* **91**, 014007 (2015), doi: 10.1103/PhysRevD.91.014007, [arXiv:1408.0122].
- [11] K. G. Chetyrkin, “Corrections of order α_s^3 to $R(\text{had})$ in pQCD with light gluinos”, *Phys. Lett. B* **391** (1997) 402 [hep-ph/9608480].
- [12] D. J. Broadhurst, “Large N expansion of QED: Asymptotic photon propagator and contributions to the muon anomaly, for any number of loops”, *Z. Phys. C* **58** (1993), 339-346 doi:10.1007/BF01560355
- [13] S. V. Mikhailov, “On a realization of $\{\beta\}$ -expansion in QCD” *JHEP* **2017**, 169 (2017), doi: 10.1007/JHEP04(2017)169, [arXiv:1610.01305].
- [14] P. A. Baikov, K. G. Chetyrkin, J. H. Kuhn, “Adler Function, Bjorken Sum Rule, and the Crewther Relation to Order α_s^4 in a General Gauge Theory”, *Phys. Rev. Lett.* **104**, 132004 (2010), [arXiv:1001.3606].
- [15] A. L. Kataev and S. V. Mikhailov, “Conformal symmetry breaking effects in gauge theories: new perturbative expressions”, *Theor. Math. Phys.* **170** (2012) 139, doi: 10.1007/s11232-012-0016-7, [arXiv:1011.5248].
- [16] A. L. Kataev and V. S. Molokoedov, *Phys. Part. Nucl.* **54** (2023) no.5, 931-941 doi:10.1134/S1063779623050106 [arXiv:2211.10242 [hep-ph]].
- [17] I. O. Goriachuk and A. L. Kataev and V. S. Molokoedov, “The $\overline{\text{MS}}$ -scheme α_s^5 QCD contributions to the Adler function and Bjorken polarized sum rule in the Crewther-type two-fold $\{\beta\}$ -expanded representation”, *JHEP* **2022**, 028 (2022), doi: 10.1007/JHEP05(2022)028, [arXiv:2111.12060].
- [18] D. J. Broadhurst and A. G. Grozin, “Matching QCD and HQET heavy - light currents at two loops and beyond”, *Phys. Rev. D* **52**, 4082 (1995), [hep-ph/9410240].
- [19] M. Beneke and V. M. Braun, “Naive nonAbelianization and resummation of fermion bubble chains”, *Phys. Lett. B* **348** (1995), 513-520 doi:10.1016/0370-2693(95)00184-M [arXiv:hep-ph/9411229 [hep-ph]].
- [20] A. P. Bakulev, S. V. Mikhailov, N. G. Stefanis, “Higher-order QCD perturbation theory in different schemes: From FOPT to CIPT to FAPT”, *J. High Energ. Phys.* **06** (2010) 085 [[1004.4125](#)]
- [21] S. J. Brodsky, G. P. Lepage and P. B. Mackenzie, “On the Elimination of Scale Ambiguities in Perturbative Quantum Chromodynamics”, *Phys. Rev. D* **28**, 228 (1983).
- [22] “The On-Line Encyclopedia of Integer Sequences”, <https://oeis.org/A000041>.

- [23] A. L. Kataev and S. V. Mikhailov, “ β -expansion in QCD, its conformal symmetry limit: theory + applications,” Nucl. Part. Phys. Proc. **258-259** (2015), 45-50
doi:10.1016/j.nuclphysbps.2015.01.011 [arXiv:1410.0554 [hep-ph]].
- [24] A. L. Kataev, J. Phys. Conf. Ser. **608** (2015) no.1, 012078
doi:10.1088/1742-6596/608/1/012078 [arXiv:1411.2257 [hep-ph]].