# Multiscale modelling of animal movement with persistent dynamics

Théo Michelot\*

Dalhousie University

#### Abstract

Wild animals are commonly fitted with trackers that record their position through time, to learn about their behaviour. Broadly, statistical models for tracking data often fall into two categories: local models focus on describing small-scale movement decisions, and global models capture large-scale spatial distributions. Due to this dichotomy, it is challenging to describe mathematically how animals' distributions arise from their short-term movement patterns, and to combine data sets collected at different scales. We propose a multiscale model of animal movement and space use based on the underdamped Langevin process, widely used in statistical physics. The model is convenient to describe animal movement for three reasons: it is specified in continuous time (such that its parameters are not dependent on an arbitrary time scale), its speed and direction are autocorrelated (similarly to real animal trajectories), and it has a closed form stationary distribution that we can view as a model of long-term space use. We use the common form of a resource selection function for the stationary distribution, to model the environmental drivers behind the animal's movement decisions. We further increase flexibility by allowing movement parameters to be time-varying, e.g., to account for daily cycles in an animal's activity. We formulate the model as a state-space model and present a method of inference based on the Kalman filter. The approach requires discretising the continuous-time process, and we use simulations to investigate performance for various time resolutions of observation. The approach works well at fine resolutions, though the estimated stationary distribution tends to be too flat when time intervals between observations are very long.

### 1 Introduction

Wild animals are commonly tracked using satellite tags, typically resulting in two-dimensional locations collected at discrete time intervals. Various statistical methods have been developed to analyse such data and answer increasingly complex ecological questions (Hooten et al., 2017). Those methods can broadly be divided into two classes, depending on the spatio-temporal scale of focus.

<sup>\*</sup>Email: theo.michelot@dal.ca

Global models describe large-scale patterns of space use by animals, and local models focus on smallscale movement patterns. These roughly correspond to the distinction in mathematical biology between Eulerian approaches, which describe the temporal evolution of the spatial distribution of animals, and Lagrangian approaches, which describe the movement dynamics of a single animal (Turchin, 1998). Although the large-scale patterns emerge from small-scale decisions, most existing models ignore this mechanism. The main approach to bridge this gap has been to develop analytical and numerical methods to simulate the large-scale distribution that emerges from a fitted animal movement model (Barnett and Moorcroft, 2008; Whitehead and Jonsen, 2013; Potts and Börger, 2023; Signer et al., 2024). However, those methods do not provide an explicit model for the largescale distribution. Recently, it has been proposed that stationary processes could be used to define movement models that scale up to a known long-term distribution (Michelot et al., 2019a,b, 2020). Here, we present the underdamped Langevin diffusion process, a model from statistical physics, as a multiscale model of animal space use.

### 2 Model overview

#### 2.1 The underdamped Langevin process

Let  $(\mathbf{Z}_t)$  denote the continuous-time process for the location  $X_t$  and velocity  $V_t$  of an animal at time  $t \ge 0$ , i.e.,  $\mathbf{Z}_t = (X_t, V_t)^{\intercal}$ . The underdamped Langevin process is described by the system of SDEs,

$$\begin{cases} dX_t = V_t \ dt \\ dV_t = -\gamma V_t \ dt + \sigma^2 \nabla \log[\pi(X_t)] \ dt + \sqrt{2\gamma} \sigma \ dW_t \end{cases}, \tag{1}$$

with initial condition  $Z_0 = (x_0, v_0)^{\intercal}$ , where  $\pi$  is a differentiable function from  $\mathbb{R}$  to  $\mathbb{R}^+$ ,  $\nabla$  is the gradient with respect to location,  $(W_t)$  is a standard Wiener process, and  $\gamma > 0$  and  $\sigma > 0$  are parameters (Cheng et al., 2018). In Equation 1, the location is defined as the time integral of the velocity, and the velocity is a mean-reverting process pulled towards higher values of  $\pi$  (with some stochasticity). The location process  $(X_t)$  therefore combines directional persistence, because velocity is autocorrelated, and attraction to high values of  $\pi$ . We focus the mathematical exposition on the one-dimensional case because Equation 1 can be used directly to model each dimension of a multi-dimensional *isotropic* process (e.g., two or three dimensions for animal movement). Under mild regularity conditions, the marginal stationary distribution of the process  $(X_t)$  defined in Equation 1 is proportional to  $\pi$  (Cheng et al., 2018; Eberle et al., 2019). In later sections, we will use this property to define a multiscale model where an animal's movement decisions, described by the SDE, are directly linked to the emergent distribution.

In physics, Equation 1 is used to describe the motion of a particle under an external force measured by  $F = \nabla \log \pi$  (Risken, 1996, Chapter 10). In that context, the parameter  $\gamma$  is often called the friction coefficient, as larger values imply stronger reversion of the velocity to zero (Erdmann et al., 2000). Using the Fokker-Planck approach, the underdamped Langevin process has also been formulated through a partial differential equation for evolution of the probability density function of ( $Z_t$ ), known as the Kramers equation (Kramers, 1940; Erdmann et al., 2000; Hadeler et al., 2004; Gardiner, 2004).

Because  $(X_t)$  has  $\pi$  as its stationary distribution, this process has also been used to build Markov Chain Monte Carlo (MCMC) algorithms to sample from  $\pi$  (Chen et al., 2014). It is a special case of the general method of Ma et al. (2015, 2019) for specifying stochastic differential equations with a given stationary distribution, which they use to define a general family of samplers. In particular, Equation 1 defines an irreversible stochastic process, in contrast with more common reversible processes such as the *overdamped* Langevin diffusion (Roberts and Tweedie, 1996).

#### 2.2 A multiscale animal movement model

If the process  $(X_t)$  defined in Equation 1 describes the position of an animal, the function  $\pi$  represents its long-term distribution over space, analogous to species distribution models widely used in ecology (Aarts et al., 2012). Then,  $\pi$  may be determined by habitat suitability and other factors that affect the animal's space use (e.g., corridors it needs to travel through). We propose using the underdamped Langevin process as a multi-scale model of animal movement, which links the small-scale movement dynamics described by the stochastic differential equations (Equation 1) to the large-scale spatial distribution  $\pi$ . Michelot et al. (2019b) proposed a similar model based on the *overdamped* Langevin diffusion, but the underdamped process presented herein has the advantage of describing realistic small-scale movement dynamics, including a finite speed and directional persistence.

#### 2.2.1 Modelling the utilisation distribution

To ensure that it is positive, a natural choice for the utilisation distribution  $\pi$  is of the form

$$\pi(x) \propto \exp\left(\sum_{k=1}^{K} \beta_k \psi_k(x)\right),\tag{2}$$

where the  $\psi_k$  are known differentiable functions over space. Due to the linearity of the gradient operator, the gradient term in the SDE for the velocity process can then be written as a linear combination of the gradients of the functions  $\psi_k$ , i.e.,  $\nabla \log \pi(X_t) = \sum_{k=1}^K \beta_k \nabla \psi_k(X_t)$ . We suggest two possible choices for the functions  $\psi_k$ , either as spatial covariates that can be used to predict space use, or as basis functions in a non-parametric approach.

The log-linear form for the utilisation distribution  $\pi$  (Equation 2) is often called a resource selection function when the functions  $\psi_k$  are spatial covariates of interest (Johnson et al., 2006; Aarts et al., 2012). The  $\beta_k$  coefficients then measure the association between an animal's space use and the measured environmental features, and often interpreted in terms of strength of "selection" or "avoidance" (Fieberg et al., 2021). This approach can directly be integrated into the underdamped Langevin process: the velocity  $V_t$  tends to be "pulled" towards higher values (if  $\beta_k > 0$ ) or lower values (if  $\beta_k < 0$ ) of the covariate  $\psi_k$ . This provides an intuitive description of an animal's movement decisions, based on the local environmental features captured by the gradient.

Alternatively, if spatial covariates are not available, or if they do not fully capture patterns of space use, we can use a non-parametric model of  $\pi$  where the  $\psi_k$  are basis functions. In this context, the weights  $\beta_k$  cannot be interpreted separately, and only the resulting function  $\pi$  is of interest. This is for example similar to the approaches of Preisler et al. (2004), Russell et al. (2018) and Brückner et al. (2020). In the non-parametric approach, we assume that the vector of basis coefficients follows  $\beta \sim N(0, \Sigma)$ , which can be viewed as a prior distribution or as a smoothness penalty to constrain the shape of the function  $\pi$  (Wood, 2017). In the two-dimensional case of animal movement, one could for example use thin plate regression splines, for which both the basis functions  $\psi_k$  and the smoothness penalty matrix  $\Sigma$  can be obtained using the R package mgcv (Wood, 2003, 2017). A model could combine parametric and non-parametric terms, as is common in spatial modelling where penalised splines are used to capture patterns that cannot be explained by available covariates.

Note that, in practice, environmental covariates are almost always measured and stored over a discrete spatial grid (a "raster"). The corresponding functions  $\psi_k$  are then piecewise constant, and their gradients cannot be used directly in the model (because the slope is zero within each grid cell, and undefined at the boundaries). In this case, we instead consider the bilinear interpolation of the raster, for which the gradient can easily be computed analytically (see e.g., Michelot et al., 2019b). Other smoothing or interpolation methods may be considered.

#### 2.2.2 Parameter interpretation

The parameter  $\gamma$  is called the friction coefficient in physics, and larger values lead to less smoothlooking trajectories. Its inverse  $1/\gamma$ , sometimes called the relaxation time of the process (Risken, 1996), is the time interval it takes for the autocorrelation of the velocity to decrease by a factor e, such that  $3/\gamma$  can be interpreted as the time scale over which the autocorrelation of the velocity decreases by approximately 95% (Michelot and Blackwell, 2019). The parameter  $\sigma^2$  is most directly interpreted as the variance of the limiting distribution of the velocity process. As a result,  $\sqrt{\pi/2} \times \sigma$ is the mean speed of movement of the animal (Michelot and Blackwell, 2019). Figure 1 shows simulations from the underdamped Langevin process for different combinations of values for  $\gamma$  and  $\sigma$ , all with the same underlying stationary distribution  $\pi$ . It illustrates the flexibility of the model to capture a wide range of movement dynamics, all leading to the same large-scale emergent patterns. In Section 2.3, we extend this further by allowing  $\gamma$  and  $\sigma$  to depend on time-varying covariates, e.g., to account for cyclical patterns in an animal's activity.



Figure 1: Simulations from the underdamped Langevin process for different values of the movement parameters  $\gamma$  (columns) and  $\sigma$  (rows). The background colour shows the stationary distribution, which was identical for all simulations.

#### 2.2.3 Links to other movement models

The underdamped Langevin process includes many widely-used continuous-time movement models as special or limiting cases. The continuous-time correlated random walk (or integrated Ornstein-Uhlenbeck process) of Johnson et al. (2008) is obtained when  $\nabla \log \pi = 0$ , i.e., when the stationary distribution  $\pi$  is flat. The model presented here can therefore be viewed as an extension of that model to the case where movement is driven by habitat, similar to the approach of Russell et al. (2018). When both  $\gamma \to \infty$  and  $\sigma \to \infty$  such that  $\sigma^2/\gamma$  remains constant, the underdamped Langevin process reduces to the overdamped (first-order) Langevin process (proof in Appendix A). Michelot et al. (2019b) used the overdamped process for animal movement, but it has no directional persistence and infinite speed, making it a less realistic model for the fine-scale movement dynamics. Michelot et al. (2020) described a reversible stochastic process inspired by Markov chain Monte Carlo samplers as the basis for a multiscale movement model. Similarly to the overdamped Langevin process, that model did not include directional persistence, leading to random walk-like movement patterns.

### 2.3 Time-varying dynamics

The dynamics of an animal's movement often change through time, as a result of changes in its behavioural state (e.g., resting, foraging, exploring; Morales et al., 2004). To account for this, two common approaches are to allow for regime-switching through a latent continuous-time Markov chain (Blackwell, 1997; Michelot and Blackwell, 2019), or to specify the movement parameters as functions of time-varying covariates (Michelot et al., 2021). Here, we adopt the latter approach, which is relatively easy to implement, and can relate movement patterns to temporal variables. Following the varying-coefficient stochastic differential equation approach of Michelot et al. (2021), the movement parameters can be specified using a combination of linear, non-linear, and random effects of covariates. A common example would be to model the speed or directionality of movement with a cyclical effect of time of day.

As long as the movement parameters do not depend on space (e.g., do not depend on spatial covariates), the resulting process  $(X_t)$  still has the stationary distribution  $\pi$ . Indeed, it can be viewed as an infinite mixture of processes that all share the same stationary distribution. This provides a very flexible framework to relate an animal's time-varying movement dynamics to a explicit long-term emerging distribution.

It is in principle straightforward to apply a similar method to write the parameters of  $\pi$  as functions of time-varying covariates. Although this could be used to capture behaviour-dependent habitat preferences (similar to Nicosia et al., 2017), the time-varying function  $\pi$  would not be a stationary distribution for  $(X_t)$  any more, and the large-scale distribution of space use may not be available in closed form.

### 3 Statistical inference and implementation

Given observations from the location process  $(X_t)$ , we wish to estimate the dynamics of the velocity process, i.e., the movement parameters  $\gamma$  and  $\sigma$ , and the parameters of the stationary distribution  $\pi$ . Russell et al. (2018) proposed a similar model for the movements of ants, and implemented a Bayesian approach based on the Euler-Maruyama discretisation of the position and velocity processes. Brückner et al. (2020) used an alternative method to derive estimators of the force field and noise in underdamped systems, while correcting for discretisation and observation errors.

In this section, we propose another framework of inference based on the Kalman filter. The key challenge is that the velocity process is not observed directly, so we treat it as a latent variable in a state-space model formulation, similar to the approach of Johnson et al. (2008) for the integrated Ornstein-Uhlenbeck process.

### 3.1 Transition density of discretised process

We first consider a realisation from the joint process  $(\mathbf{Z}_t)$  (location and velocity), with values  $(\mathbf{z}_1, \mathbf{z}_2, \ldots, \mathbf{z}_n)$  at observation times  $t_1 < t_2 < \cdots < t_n$ , and we define  $\Delta_i = t_{i+1} - t_i$  as the *i*-th time interval. There is no general solution to the system of SDEs in Equation 1 because the dynamics of the process depend on the possibly complex function  $\pi$ , but an approximate solution can be found based on the assumption that the force term  $\nabla \log \pi(X_t)$  is constant over each time interval  $[t_i, t_{i+1})$ . Cheng et al. (2018) derived the transition density of a special case of this discretised underdamped Langevin process (for a particular value of the parameters  $\gamma$  and  $\sigma$ ), and we use similar calculations to find the following general formulas:

$$Z_{t_{i+1}} \mid \{Z_{t_i} = z_i\} \sim N(\mu_i, \ Q_i)\}$$

where

$$\boldsymbol{\mu}_{i} = \begin{pmatrix} x_{i} + \frac{v_{i}(1 - e^{-\gamma\Delta_{i}})}{\gamma} + \frac{\sigma^{2}\nabla\log\pi(x_{i})}{\gamma} \left(\Delta_{i} - \frac{1 - e^{-\gamma\Delta_{i}}}{\gamma}\right) \\ v_{i}e^{-\gamma\Delta_{i}} - \frac{\sigma^{2}\nabla\log\pi(x_{i})}{\gamma}(1 - e^{-\gamma\Delta_{i}}) \end{pmatrix}$$

and

$$\boldsymbol{Q}_{i} = \begin{pmatrix} \sigma^{2} \left( \frac{2\Delta_{i}}{\gamma} - \frac{e^{-2\gamma\Delta_{i}}}{\gamma^{2}} - \frac{3}{\gamma^{2}} + \frac{4e^{-\gamma\Delta_{i}}}{\gamma^{2}} \right) & \frac{\sigma^{2}}{\gamma} \left( 1 - 2e^{-\gamma\Delta_{i}} + e^{-2\gamma\Delta_{i}} \right) \\ \frac{\sigma^{2}}{\gamma} \left( 1 - 2e^{-\gamma\Delta_{i}} + e^{-2\gamma\Delta_{i}} \right) & \sigma^{2} (1 - e^{-2\gamma\Delta_{i}}) \end{pmatrix}$$

Over very short time intervals, the slope of  $\log \pi$  will typically be almost constant, leading to a good approximation, but the error will be greater at coarser time resolutions. We illustrate this in simulations in Section 4.

#### 3.2 Inference using state-space model formulation

In practice, the velocity process is not observed, and we may only observe the location at discrete times, so we take a state-space modelling approach. A state-space model jointly describes the dynamics of an unobserved state process, and the mechanism that connects the state to the observed data (Durbin and Koopman, 2012). In this context, the underlying state is the joint process for the location and velocity of the animal, and the observation are just the locations. The model can be rewritten in the form

$$\begin{array}{ll} \text{Observation Equation:} & X_{t_i} = \boldsymbol{A} \boldsymbol{Z}_{t_i} \\ \text{State Equation:} & \boldsymbol{Z}_{t_{i+1}} = \boldsymbol{T}_i \boldsymbol{Z}_{t_i} + \boldsymbol{B}_i \boldsymbol{h}_i + \boldsymbol{\eta}_i \end{array}$$

where  $\eta_i \sim N(\mathbf{0}, \mathbf{Q}_i)$  is the process error (with  $\mathbf{Q}_i$  as defined in Section 3.1),  $\mathbf{h}_i = \nabla \log \pi(X_{t_i})$ , and

$$\boldsymbol{A} = \begin{pmatrix} 1 & 0 \end{pmatrix}, \quad \boldsymbol{T}_i = \begin{pmatrix} 1 & \frac{1-e^{-\gamma\Delta_i}}{\gamma} \\ 0 & e^{-\gamma\Delta_i} \end{pmatrix}, \quad \text{and} \ \boldsymbol{B}_i = \begin{pmatrix} \frac{\sigma^2}{\gamma} \left( \Delta_i - \frac{1}{\gamma} (1-e^{-\gamma\Delta_i}) \right) \\ \frac{\sigma^2}{\gamma} (1-e^{-\gamma\Delta_i}) \end{pmatrix}.$$

The observation equation simply represents the fact that velocity is not observed. Note that, although the state-space formulation could in principle be used to account for measurement error with an additional term in the observation equation, this would not be sufficient to correct for the corresponding errors in  $\nabla \log \pi(X_t)$ . Therefore, we assume that locations are observed with negligible error, and leave the problem of noisy locations open. The state equation is directly derived from the transition density of the discretised underdamped Langevin process given in Section 3.1. For two-dimensional data such as animal movement tracks, the model takes a similar formulation, but where the matrices given above are combined into block matrices.

For a state-space model in this form, the likelihood can be evaluated using the Kalman filter (Durbin and Koopman, 2012, Section 7.2). We use maximum likelihood estimation, but other likelihood-based inference methods could also be implemented directly using this approach. Using large-sample approximations, we derive uncertainty estimates based on the inverse of the Hessian matrix of all model parameters. Model parameters can be estimated from multiple independent time series, e.g., movement tracks collected on different animals, by multiplying their likelihoods. If random effect and penalised terms are included in the model for the utilisation distribution  $\pi$  (e.g., spatial random field) or in the movement parameters (e.g., cyclical effect of time of day), the model can be fitted using the marginal likelihood approach described by Michelot et al. (2021).

### 4 Simulations

We assessed the performance of the method of inference presented in Section 3 using simulations. We defined the true stationary distribution as  $\pi(x) = \exp(\beta_1\psi_1(x) + \beta_2\psi_2(x) + \beta_3\psi_3(x))$  where  $\psi_1$ and  $\psi_2$  were generated at random on a raster grid, to resemble spatial environmental covariates, and where  $\psi_3$  was the squared distance to the centre of the study region. We used the coefficients  $(\beta_1, \beta_2, \beta_3) = (2, 5, -10)$ , representing positive "selection" for the two environmental variables, and a tendency to remain close to the centre (similar to home range behaviour of animals). For the simulation, the movement parameters were set to  $\gamma = 1$  and  $\sigma = 1$ . We simulated 100 trajectories from the discretised underdamped Langevin process on this map, using a very fine time resolution of  $\Delta = 0.01$  to avoid any discretisation error in the simulation. Then, we downsampled the data at random to (average) time intervals of  $\Delta \in \{0.02, 0.05, 0.1, 0.2, 0.5, 1, 2\}$ , and fitted the model on the thinned data.

The results are shown in Figure 2. All model parameters were recovered well at short time resolutions. The friction parameter  $\gamma$  displayed some moderate positive bias at coarse time resolutions, corresponding to a slight underestimation of the time scale of autocorrelation of the velocity process. The speed parameter  $\sigma$  was estimated well even for the longer time intervals. In the model for the stationary distribution  $\pi$ , the estimates for the coefficients  $\beta_1$  and  $\beta_2$  (measuring the association between the distribution and two artificial environmental variables) decreased as the time interval of observation increased; for very long intervals, they were estimated close to zero. This is because those parameters are estimated based on the association between the animal's movement and the gradient of the covariates at the start point of each observed step. As the time intervals get longer, the association becomes weaker, i.e., the coefficients are underestimated. Interestingly, the coefficient for the squared distance  $\psi_3$  did not seem to be biased even at coarse resolutions, although the estimation variance increased somewhat. This is probably because  $\psi_3$  does not change quickly over space, i.e., the assumption that its gradient does not change between two observed locations is approximately correct even for coarser data. As a result, even though  $\beta_1$  and  $\beta_2$  could not be recovered at long intervals, the overall shape of the utilisation distribution  $\pi$  was estimated quite well; the correlation between the true and estimated  $\pi$  (measured on a raster grid) was around 0.85 on average even at the coarsest resolution.



Figure 2: Results of simulation study. Parameter estimates from 100 simulations, for data thinned to different time resolutions  $\Delta$ . The bottom-left panel shows the correlation between the estimated and true utilisation distributions, measured on a raster grid. The other five panels show the five parameters of the model, with the true values shown as horizontal red lines.

### 5 Discussion

It is a common challenge to carry out inference for stochastic differential equations from discrete data, and most approaches have to use approximations such as the Euler-Maruyama discretisation (Iacus, 2008). Similarly, the approach we presented requires the assumption that the gradient of the stationary distribution  $\nabla \log \pi(X_{t_i})$  is approximately constant between two observation times  $t_i$ and  $t_{i+1}$ , and simulations revealed that this leads to bias for coarse temporal resolutions. It may be possible to use higher-order approximations to improve inference. We tried replacing  $\nabla \log \pi(X_{t_i})$  by  $(\nabla \log \pi(X_{t_i}) + \nabla \log \pi(X_{t_{i+1}}))/2$ , or by  $\nabla \log \pi((X_{t_i} + X_{t_i})/2)$ , to account for dependence on the shape of  $\pi$  along the whole step (rather than only at the start point), but this did not yield noticeably better results. It would be interesting to explore other methods of inference for second-order stochastic differential equations, such as that described by Pilipovic et al. (2024).

The Kramers equation is a partial differential equation describing the evolution of the joint probability density of the position and velocity through time, for the underdamped Langevin process (Hadeler et al., 2004). It could be used to simulate the spatial distribution of an animal over intermediate time steps, to learn about its transient movement dynamics (Potts and Painter, 2024). This could for example provide insights into the time scale over which animals explore their environment, and the effects of habitat fragmentation on their ability to efficiently travel across their range.

The underdamped Langevin diffusion process is a promising approach to analyse animal tracking data, with great flexibility to capture small-scale movement patterns, and with a direct link to the emerging utilisation distribution. Other stationary processes within the general class described by Ma et al. (2015, 2019) might provide even better descriptions of animal movement, and we hope that this paper will stimulate more work in this area.

### Acknowledgements

I want to thank Paul Blackwell for discussions on this topic, and for bringing Ma et al. (2019) to my attention.

### References

- Aarts, G., Fieberg, J., and Matthiopoulos, J. (2012). Comparative interpretation of count, presence–absence and point methods for species distribution models. *Methods in Ecology and Evolution*, 3(1):177–187.
- Barnett, A. H. and Moorcroft, P. R. (2008). Analytic steady-state space use patterns and rapid computations in mechanistic home range analysis. *Journal of Mathematical Biology*, 57:139–159.
- Blackwell, P. (1997). Random diffusion models for animal movement. *Ecological Modelling*, 100(1-3):87–102.
- Brückner, D. B., Ronceray, P., and Broedersz, C. P. (2020). Inferring the dynamics of underdamped stochastic systems. *Physical Review Letters*, 125(5):058103.
- Chen, T., Fox, E., and Guestrin, C. (2014). Stochastic gradient Hamiltonian Monte Carlo. In International Conference on Machine Learning, pages 1683–1691. PMLR.

- Cheng, X., Chatterji, N. S., Bartlett, P. L., and Jordan, M. I. (2018). Underdamped Langevin MCMC: A non-asymptotic analysis. In *Conference on Learning Theory*, pages 300–323. PMLR.
- Durbin, J. and Koopman, S. J. (2012). *Time series analysis by state space methods*, volume 38. OUP Oxford, 2nd edition.
- Eberle, A., Guillin, A., and Zimmer, R. (2019). Couplings and quantitative contraction rates for Langevin dynamics. *The Annals of Probability*, 47(4):1982 2010.
- Erdmann, U., Ebeling, W., Schimansky-Geier, L., and Schweitzer, F. (2000). Brownian particles far from equilibrium. The European Physical Journal B-Condensed Matter and Complex Systems, 15:105–113.
- Fieberg, J., Signer, J., Smith, B., and Avgar, T. (2021). A 'How to' guide for interpreting parameters in habitat-selection analyses. *Journal of Animal Ecology*, 90(5):1027–1043.
- Gardiner, C. W. (2004). Handbook of stochastic methods. Springer Berlin, 3rd edition.
- Hadeler, K. P., Hillen, T., and Lutscher, F. (2004). The Langevin or Kramers approach to biological modeling. *Mathematical Models and Methods in Applied Sciences*, 14(10):1561–1583.
- Hooten, M. B., Johnson, D. S., McClintock, B. T., and Morales, J. M. (2017). Animal movement: statistical models for telemetry data. CRC Press.
- Iacus, S. M. (2008). Simulation and inference for stochastic differential equations: with R examples, volume 486. Springer.
- Johnson, C. J., Nielsen, S. E., Merrill, E. H., McDonald, T. L., and Boyce, M. S. (2006). Resource selection functions based on use-availability data: theoretical motivation and evaluation methods. *The Journal of Wildlife Management*, 70(2):347–357.
- Johnson, D. S., London, J. M., Lea, M.-A., and Durban, J. W. (2008). Continuous-time correlated random walk model for animal telemetry data. *Ecology*, 89(5):1208–1215.
- Kramers, H. A. (1940). Brownian motion in a field of force and the diffusion model of chemical reactions. *Physica*, 7(4):284–304.
- Ma, Y.-A., Chen, T., and Fox, E. (2015). A complete recipe for stochastic gradient MCMC. Advances in Neural Information Processing Systems, 28.
- Ma, Y.-A., Fox, E. B., Chen, T., and Wu, L. (2019). Irreversible samplers from jump and continuous Markov processes. *Statistics and Computing*, 29:177–202.
- Michelot, T. and Blackwell, P. G. (2019). State-switching continuous-time correlated random walks. Methods in Ecology and Evolution, 10(5):637–649.

- Michelot, T., Blackwell, P. G., Chamaillé-Jammes, S., and Matthiopoulos, J. (2020). Inference in MCMC step selection models. *Biometrics*, 76(2):438–447.
- Michelot, T., Blackwell, P. G., and Matthiopoulos, J. (2019a). Linking resource selection and step selection models for habitat preferences in animals. *Ecology*, 100(1):e02452.
- Michelot, T., Glennie, R., Harris, C., and Thomas, L. (2021). Varying-coefficient stochastic differential equations with applications in ecology. *Journal of Agricultural, Biological and Environmental Statistics*, 26:446–463.
- Michelot, T., Gloaguen, P., Blackwell, P. G., and Étienne, M.-P. (2019b). The Langevin diffusion as a continuous-time model of animal movement and habitat selection. *Methods in Ecology and Evolution*, 10(11):1894–1907.
- Morales, J. M., Haydon, D. T., Frair, J., Holsinger, K. E., and Fryxell, J. M. (2004). Extracting more out of relocation data: building movement models as mixtures of random walks. *Ecology*, 85(9):2436–2445.
- Nicosia, A., Duchesne, T., Rivest, L.-P., and Fortin, D. (2017). A multi-state conditional logistic regression model for the analysis of animal movement. *The Annals of Applied Statistics*, 11(3):1537 – 1560.
- Pilipovic, P., Samson, A., and Ditlevsen, S. (2024). Strang splitting for parametric inference in second-order stochastic differential equations. arXiv preprint arXiv:2405.03606.
- Potts, J. R. and Börger, L. (2023). How to scale up from animal movement decisions to spatiotemporal patterns: An approach via step selection. *Journal of Animal Ecology*, 92(1):16–29.
- Potts, J. R. and Painter, K. J. (2024). Distinguishing between long-transient and asymptotic states in a biological aggregation model. *Bulletin of Mathematical Biology*, 86(3):1–16.
- Preisler, H. K., Ager, A. A., Johnson, B. K., and Kie, J. G. (2004). Modeling animal movements using stochastic differential equations. *Environmetrics*, 15(7):643–657.
- Risken, H. (1996). Fokker-Planck equation. Springer.
- Roberts, G. O. and Tweedie, R. L. (1996). Exponential convergence of Langevin distributions and their discrete approximations. *Bernoulli*, pages 341–363.
- Russell, J. C., Hanks, E. M., Haran, M., and Hughes, D. (2018). A spatially varying stochastic differential equation model for animal movement. *The Annals of Applied Statistics*, 12(2):1312 – 1331.
- Signer, J., Fieberg, J., Reineking, B., Schlaegel, U., Smith, B., Balkenhol, N., and Avgar, T. (2024).

Simulating animal space use from fitted integrated step-selection functions (iSSF). Methods in Ecology and Evolution, 15(1):43–50.

- Turchin, P. (1998). Quantitative analysis of movement: measuring and modeling population redistribution in animals and plants. Sinauer, Sunderland, MA.
- Whitehead, H. and Jonsen, I. D. (2013). Inferring animal densities from tracking data using Markov chains. *PloS one*, 8(4):e60901.
- Wood, S. N. (2003). Thin plate regression splines. Journal of the Royal Statistical Society Series B: Statistical Methodology, 65(1):95–114.

Wood, S. N. (2017). Generalized additive models: an introduction with R. Chapman and Hall/CRC.

## Appendices

### A Overdamped Langevin process as a limiting case

We first define  $\tilde{\sigma} = \sigma \sqrt{\varepsilon}$  and  $\tilde{\lambda} = \lambda \varepsilon$  for some  $\epsilon > 0$ , and we consider the situation where  $\varepsilon \to 0$ (i.e.,  $\sigma \to \infty$ , and  $\sigma^2/\lambda$  is fixed). The SDE for  $V_t$  (Equation 1) can be rewritten

$$dV_t = -\frac{\tilde{\lambda}}{\varepsilon} V_t \ dt + \frac{\tilde{\sigma}^2}{\varepsilon} \nabla \log[\pi(X_t)] \ dt + \frac{\sqrt{2\tilde{\lambda}\tilde{\sigma}}}{\varepsilon} \ dW_t$$

and, if we multiply both sides by  $\varepsilon$ ,

$$\varepsilon dV_t = -\tilde{\lambda} V_t dt + \tilde{\sigma}^2 \nabla \log[\pi(X_t)] dt + \sqrt{2\tilde{\lambda}} \tilde{\sigma} dW_t$$

In the limit  $\varepsilon \to 0$ , the left-hand side tends to zero, and this equation becomes

$$\tilde{\lambda} V_t \ dt = \tilde{\sigma}^2 \nabla \log[\pi(X_t)] \ dt + \sqrt{2\tilde{\lambda}}\tilde{\sigma} \ dW_t$$

We plug this into the equation  $dX_t = V_t dt$  (Equation 1), and we get

$$dX_t = \frac{\tilde{\sigma}^2}{\tilde{\lambda}} \nabla \log[\pi(X_t)] \ dt + \frac{\sqrt{2}\tilde{\sigma}}{\sqrt{\tilde{\lambda}}} \ dW_t.$$

This equation defines an overdamped (first-order) Langevin diffusion process, where the parameter  $\tilde{\sigma}/\sqrt{\tilde{\lambda}}$  measures the diffusion speed (Michelot et al., 2019b).