State Transfer and Readout Times for Trees of Diameter 4

Stephen Kirkland^{1 ‡} Christopher M. van Bommel²

June 24, 2024

Abstract

We consider the state transfer properties of continuous time quantum walks on trees of diameter 4. We characterize all pairs of strongly cospectral vertices in trees of diameter 4, finding that they fall into pairs of three different types. For each type, we construct an infinite family of diameter 4 trees for which there is pretty good state transfer between the pair of strongly cospectral vertices. Moreover, for two of those types, for each tree in the infinite family, we give an explicit sequence of readout times at which the fidelity of state transfer converges to 1. For strongly cospectral vertices of the remaining type, we identify a sequence of trees and explicit readout times so that the fidelity of state transfer between the strongly cospectral vertices approaches 1.

We also prove a result of independent interest: for a graph with the property that the fidelity of state transfer between a pair of vertices at time t_k converges to 1 as $k \to \infty$, then the derivative of the fidelity at t_k converges to 0 as $k \to \infty$.

Keywords: Quantum state transfer; Fidelity; Pretty good state transfer; Sensitivity.

MSC 2020: 81P45; 05C50; 15A18.

1 Introduction

Interacting qubits can be modelled by a graph, where the individual qubits are represented by vertices of the graph and the presence of interactions between qubits is represented by an edge between the corresponding vertices. This forms a natural extension of classical random walks to the quantum setting, which we refer to as continuous quantum walks. We will consider the propagation model of this quantum system corresponding to the adjacency matrix A of the corresponding graph; then the time evolution of the system is given by

$$U(t) = \exp(itA),$$

where t is a nonnegative real value representing the length of time under which the system has propagated. It follows that U(t) is both symmetric and unitary, and in particular that $|U(t)_{a,b}| \leq 1$ for any t. The *fidelity* at time t is defined as $|U(t)_{a,b}|^2$, and is interpreted as the probability of state transfer from a to b at time t. We say a graph has *perfect state transfer* between vertices a and b at time \hat{t} if $|U(\hat{t})_{a,b}| = 1$.

Perfect state transfer is an important task in the area of quantum information processing, yet the graphs in which this property can be found appear to be rare. Christandl et al. [3, 4] demonstrated that the path graph has perfect state transfer between its end vertices if and only if the length of the

¹Department of Mathematics, University of Manitoba, Winnipeg, MB, Canada. Stephen.Kirkland@umanitoba.ca

²Department of Mathematics and Statistics, University of Guelph, Guelph, ON, Canada. cvanbomm@uoguelph.ca

 $^{^{\}ddagger}$ Research supported in part by The Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant RGPIN-2019-05408.

path is 1 or 2; Stevanović [21] and Godsil [14] independently extended this result to exclude perfect state transfer between internal nodes of paths. Godsil [15] found that for each $k \in \mathbb{N}$, the number of connected graphs with valency at most k that admit perfect state transfer is finite. Coutinho and Liu [10] considered the quantum system corresponding to the Laplacian matrix of the graph, and demonstrated that the path on 2 vertices is the only tree that admits perfect state transfer in this model. Recently, Coutinho, Juliano, and Spier [9] demonstrated that the only trees that admit perfect state transfer with respect to the adjacency matrix are P_2 and P_3 .

In order to transmit quantum information across larger distances, we would like to find examples of perfect state transfer between more distant vertices. One way to construct graphs that admit perfect state transfer at larger distances is to take Cartesian products of graphs admitting perfect state transfer. In this way, we can take the hypercube Q_k as an example of a graph that admits perfect state transfer between vertices at distance k. However, such a construction is exponential (in k) in both the number of vertices and the number of edges. Other infinite families demonstrating perfect state transfer at larger distances also exhibit this exponential growth. In the laboratory setting, each interaction between qubits (i.e. each edge in the corresponding graph) has an associated cost, and so to reduce those costs, it is desirable to identify examples of graphs exhibiting perfect state transfer where the number of vertices and/or edges is subexponential in the distance k – to date, no such families of examples are known.

An alternate approach to finding ideal graphs to admit perfect state transfer at specific distances is to consider weighted graphs. Indeed, perfect state transfer can be exhibited on a path of any length provided that suitable weights are used. One such weighting can be obtained by taking the quotient graph of an associated hypercube. However, these and other examples (see [23, 24]), require large weights to be used, again requiring significant cost to achieve. A related investigation is whether perfect state transfer can be achieved on paths by adding weighted loops to the end vertices; this was raised by Casaccino et al. [2] and resolved in the negative by Kempton, Lippner, and Yau [19]. Coutinho et al. [6] investigated a generalization of this notion by decorating the ends of a path with arbitrary graphs; they demonstrated that perfect state transfer cannot be achieved on the path on 2 or 3 vertices with non-trivial decorations.

In light of the scarcity of small examples of perfect state transfer, a third approach is to make a small sacrifice to the fidelity of the transfer and consider pretty good state transfer (PGST), which requires a sequence of times t_k such that $\lim_{k\to\infty} |U(t_k)_{a,b}| = 1$. Pretty good state transfer between end-vertices of paths was studied by Godsil, Kirkland, Severini, and Smith [16], and determined to be achieved precisely on paths whose number of vertices was one less than one of the following: a prime, twice a prime, or a power of 2. Coutinho, Guo, and van Bommel [7] provided a family of paths in which pretty good state transfer was achieved between internal vertices, and van Bommel [22] subsequently completed the characterization by demonstrating this was the only such family. These results can be summarized as follows.

Theorem 1. There is pretty good state transfer on P_n between vertices a and b if and only if a+b=n+1 and:

- a) $n = 2^t 1$, where t is a positive integer;
- b) n = p 1, where p is an odd prime; or
- c) $n = 2^t p 1$, where t is a positive integer and p is an odd prime, and a is a multiple of 2^{t-1} .

Fan and Godsil [11] studied the double star graph $S_{k,\ell}$, i.e., the graph formed by joining two stars $K_{k,1}$ and $K_{\ell,1}$ by an edge between the centre vertices, and demonstrated the following.

Theorem 2. [11] Pretty good state transfer is admitted between vertices u and v of the double star graph $S_{k,\ell}$ if and only if:

a) $S_{k,\ell} \cong P_4$, and u and v are the two leaves or the two stems, or

- b) $k=2,\ \ell\neq 2,\ and\ u$ and v are the two leaves corresponding to $K_{1,2},\ or$
- c) $k = \ell > 2$, 1 + 4k is not a perfect square, and u and v are the two centre vertices.

Hou, Gu, and Tong [13] considered extended double stars $F_{k,\ell}$, which are constructed from two stars $K_{k+1,1}$ and $K_{\ell+1,1}$ by taking a leaf from each and identifying those vertices, and proved the following.

Theorem 3. [13] There is pretty good state transfer on $F_{k,\ell}$ if and only if:

- a) $k = \ell = 1$ (i.e. P_5), and pretty good state transfer occurs between the two end-vertices or the second and fourth vertices; or
- b) $k = \ell > 1$, and pretty good state transfer occurs between the two original centre vertices.

We observe in passing that the double stars are precisely the trees of diameter 3 while the extended double stars are a subfamily of the trees of diameter 4.

As noted above, the main result of [9] shows that no tree on 4 or more vertices can yield perfect state transfer. In view of that, the present paper considers pretty good state transfer on trees, with a focus on trees of diameter 4. While many results on PGST characterize circumstances under which that phenomena takes place, there is scant literature that addresses the issue of readout times. One of our novel contributions in the present work is, for certain families of trees, the explicit description of readout times at which the fidelity of state transfer is close to 1. By so doing we enhance the applicability of PGST. We also prove that for any graph, if t_k is a sequence of times such that $|U(t_k)_{a,b}| \to 1$ as $k \to \infty$, then the derivative of the fidelity at time t_k converges to 0 as $k \to \infty$. Consequently, for a graph exhibiting PGST between a and b the fidelity at t_k becomes less sensitive to small changes in the readout time as $k \to \infty$.

2 Preliminaries

We analyze the time evolution of quantum systems through the spectral decomposition of the adjacency matrix, which is given by

$$A = \sum_{\lambda} \lambda E_{\lambda},$$

where we sum over the eigenvalues λ of A and E_{λ} is the projection onto the λ -eigenspace. The time evolution of the system can be expressed as

$$U(t) = \sum_{\lambda} \exp(it\lambda) E_{\lambda}.$$

The perfect state transfer condition of $|U(\hat{t})_{a,b}| = 1$ is equivalent to $U(\hat{t})e_a = \gamma e_b$ for some $\gamma \in \mathbb{C}$ with $|\gamma| = 1$, where e_j denotes the standard basis vector which contains a 1 in row j and 0 in all other entries. Equivalently, we have $e^{it\lambda}E_{\lambda}e_a = \gamma E_{\lambda}e_b$ for each eigenvalue λ , which is equivalent to the following pair of properties:

- (a) For each eigenvalue λ , there is a $\sigma_{\lambda} \in \{-1, +1\}$ such that $E_{\lambda}e_a = \sigma_{\lambda}E_{\lambda}e_b$, and
- (b) For each eigenvalue λ_r , whenever $E_{\lambda_r}e_a \neq 0$, then $t(\lambda_1 \lambda_r) = k_r\pi$, where $k_r \in \mathbb{Z}$, and $k_r \equiv (1 \sigma_{\lambda_r})/2 \pmod{2}$.

If vertices a and b satisfy property (a) above, then we say they are *strongly cospectral*. We note that this condition implies the more well-known condition of being *cospectral*, which requires that $(E_{\lambda})_{a,a} = (E_{\lambda})_{b,b}$ for all eigenvalues λ . It follows that such vertices satisfy $A_{a,a}^k = A_{b,b}^k$ for every integer k, which counts the number of closed walks from each vertex. In particular, we have that

a and b necessarily have the same degree. We distinguish the eigenvalues for which $E_{\lambda}e_{a} \neq 0$ as forming the eigenvalue support of a, denoted Θ_{a} , since these are the only terms that affect the time evolution of the system at that vertex. Observe that if a,b are strongly cospectral vertices then Θ_{a} can be partitioned naturally as $\Theta_{a} = \sigma_{ab}^{+} \cup \sigma_{ab}^{-}$, where $\sigma_{ab}^{+} = \{\lambda \in \Theta_{a} | E_{\lambda}e_{a} = E_{\lambda}e_{b}\}$ and $\sigma_{ab}^{-} = \{\lambda \in \Theta_{a} | E_{\lambda}e_{a} = -E_{\lambda}e_{b}\}$.

Godsil [15] provided a characterization of the eigenvalues in the eigenvalue support of a vertex admitting perfect state transfer, which was expanded by Coutinho [5] as follows.

Theorem 4. [15, 5] Let X be a graph, and let $a, b \in V(X)$. There is perfect state transfer between a and b at time t if and only if all conditions below hold.

- (a) For each eigenvalue λ , there is a $\sigma_{\lambda} \in \{-1, +1\}$ such that $E_{\lambda}e_a = \sigma_{\lambda}E_{\lambda}e_b$.
- (b) There is an integer α , a square-free positive integer Δ (possibly equal to 1), so that for all λ_r in the support of a, there is an integer β_r giving

$$\lambda_r = \frac{\alpha + \beta_r \sqrt{\Delta}}{2}.$$

In particular, because λ_r is an algebraic integer, it follows that all β_r have the same parity as α .

(c) There is $g \in \mathbb{Z}$ so that, for all λ_r in the support of a, $(\beta_1 - \beta_r)/g = k_r$ with $k_r \in \mathbb{Z}$, and $k_r \equiv (1 - \sigma_{\lambda_r})/2 \pmod{2}$.

If the conditions hold, then the positive values of t for which perfect state transfer occurs are precisely the odd multiples of $\pi/(g\sqrt{\Delta})$.

We note the following useful consequence of strong cospectrality, due to Godsil and Smith [17].

Lemma 5. [17] If a and b are strongly cospectral vertices in a graph X, then any automorphism of X that fixes a also fixes b.

For pretty good state transfer, we have the following characterization provided by Banchi et al. [1].

Theorem 6. [1] Let a and b be vertices of a graph G. Then pretty good state transfer occurs between a and b if and only if both conditions below are satisfied:

- a) Vertices a and b are strongly cospectral, in which case for each $\lambda_r \in \Theta_a$, let $k_r \equiv (1 \sigma_{\lambda_r})/2 \pmod{2}$.
- b) If there is a set of integers $\{\ell_i\}$ such that

$$\sum_{\lambda_j \in \Theta_a} \ell_j \lambda_j = 0 \text{ and } \sum_{\lambda_j \in \Theta_a} \ell_j k_j \text{ is odd,}$$

then necessarily

$$\sum_{\lambda_j \in \Theta_a} \ell_j \neq 0.$$

Finally, we note that for trees, Coutinho, Juliano, and Spier demonstrated that a vertex can have at most one other vertex to which it is strongly cospectral [8].

3 Sensitivity of state transfer for strongly cospectral vertices

If there is perfect state transfer from one state to another at time \hat{t} , then necessarily the derivative of the fidelity is zero at \hat{t} , since \hat{t} corresponds to a local maximum for the fidelity. In this section, we obtain an expression for the derivative of the fidelity between strongly cospectral vertices, with particular interest in pairs of vertices exhibiting PGST. Observe that our result is not restricted to trees

Theorem 7. Suppose that a, b are strongly cospectral vertices in a graph on n vertices, and denote the fidelity at time t by f(t). For each eigenvalue λ of A, let s_{λ} denote the (a, a) entry of the corresponding orthogonal idempotent projection matrix E_{λ} . Then

$$\frac{df}{dt} = 2 \left[\sum_{\lambda_j, \lambda_\ell \in \sigma_{ab}^+} s_{\lambda_j} s_{\lambda_\ell} \lambda_\ell \sin(t(\lambda_j - \lambda_\ell)) + \sum_{\lambda_j, \lambda_\ell \in \sigma_{ab}^-} s_{\lambda_j} s_{\lambda_\ell} \lambda_\ell \sin(t(\lambda_j - \lambda_\ell)) + \sum_{\lambda_j, \lambda_\ell \in \sigma_{ab}^-} s_{\lambda_j} s_{\lambda_\ell} (\lambda_j - \lambda_\ell) \sin(t(\lambda_j - \lambda_\ell)) \right].$$
(1)

Proof. We have

$$\begin{split} f(t) &= \\ \left(\sum_{\lambda_j \in \sigma_{ab}^+} s_{\lambda_j} \cos(t\lambda_j) - \sum_{\lambda_j \in \sigma_{ab}^-} s_{\lambda_j} \cos(t\lambda_j) \right)^2 + \left(\sum_{\lambda_j \in \sigma_{ab}^+} s_{\lambda_j} \sin(t\lambda_j) - \sum_{\lambda_j \in \sigma_{ab}^-} s_{\lambda_j} \sin(t\lambda_j) \right)^2. \end{split}$$

Differentiating with respect to t yields

$$\frac{df}{dt} = 2 \left(\sum_{\lambda_j \in \sigma_{ab}^+} s_{\lambda_j} \cos(t\lambda_j) - \sum_{\lambda_j \in \sigma_{ab}^-} s_{\lambda_j} \cos(t\lambda_j) \right) \left(\sum_{\lambda_\ell \in \sigma_{ab}^-} s_{\lambda_\ell} \lambda_\ell \sin(t\lambda_\ell) - \sum_{\lambda_\ell \in \sigma_{ab}^+} s_{\lambda_\ell} \lambda_\ell \sin(t\lambda_\ell) \right) + 2 \left(\sum_{\lambda_j \in \sigma_{ab}^+} s_{\lambda_j} \sin(t\lambda_j) - \sum_{\lambda_j \in \sigma_{ab}^-} s_{\lambda_j} \sin(t\lambda_j) \right) \left(\sum_{\lambda_\ell \in \sigma_{ab}^+} s_{\lambda_\ell} \lambda_\ell \cos(t\lambda_\ell) - \sum_{\lambda_\ell \in \sigma_{ab}^-} s_{\lambda_\ell} \lambda_\ell \cos(t\lambda_\ell) \right).$$

Collecting terms and simplifying, we have

$$\frac{df}{dt} = 2 \left[\sum_{\lambda_j, \lambda_\ell \in \sigma_{ab}^+} s_{\lambda_j} s_{\lambda_\ell} \lambda_\ell \sin(t(\lambda_j - \lambda_\ell)) + \sum_{\lambda_j, \lambda_\ell \in \sigma_{ab}^-} s_{\lambda_j} s_{\lambda_\ell} \lambda_\ell \sin(t(\lambda_j - \lambda_\ell)) + \sum_{\lambda_j \in \sigma_{ab}^+, \lambda_\ell \in \sigma_{ab}^-} s_{\lambda_j} s_{\lambda_\ell} \lambda_\ell \sin(t(\lambda_\ell - \lambda_j)) + \sum_{\lambda_j \in \sigma_{ab}^-, \lambda_\ell \in \sigma_{ab}^-} s_{\lambda_j} s_{\lambda_\ell} \lambda_\ell \sin(t(\lambda_\ell - \lambda_j)) \right].$$

It now follows that

$$\frac{df}{dt} = 2 \left[\sum_{\lambda_j, \lambda_\ell \in \sigma_{ab}^+} s_{\lambda_j} s_{\lambda_\ell} \lambda_\ell \sin(t(\lambda_j - \lambda_\ell)) + \sum_{\lambda_j, \lambda_\ell \in \sigma_{ab}^-} s_{\lambda_j} s_{\lambda_\ell} \lambda_\ell \sin(t(\lambda_j - \lambda_\ell)) + \sum_{\lambda_j, \lambda_\ell \in \sigma_{ab}^-} s_{\lambda_j} s_{\lambda_\ell} (\lambda_j - \lambda_\ell) \sin(t(\lambda_j - \lambda_\ell)) \right],$$

as desired. \Box

Example 8. In this example, we illustrate the conclusion of Theorem 7 for P_3 , where a and b are taken to be leaves. We have $\sigma_{ab}^+ = \{\sqrt{2}, -\sqrt{2}\}, \sigma_{ab}^- = \{0\}$, with $s_{\sqrt{2}} = s_{-\sqrt{2}} = \frac{1}{4}$ and $s_0 = \frac{1}{2}$. We find readily that $f(t) = \frac{1}{4}(1 - \cos(\sqrt{2}t))^2$, and differentiating that expression with respect to t yields

$$\frac{df}{dt} = \frac{1}{\sqrt{2}}(1 - \cos(\sqrt{2}t))\sin(\sqrt{2}t).$$

Alternatively, we can find $\frac{df}{dt}$ via (1) as follows:

$$\begin{split} &\frac{df}{dt} = \\ &2\left(\frac{1}{4}\frac{1}{4}\sqrt{2}\sin(-2\sqrt{2}t) + \frac{1}{4}\frac{1}{4}(-\sqrt{2})\sin(2\sqrt{2}t) + \right. \\ &\left. \frac{1}{4}\frac{1}{2}\sqrt{2}\sin(\sqrt{2}t) + \frac{1}{4}\frac{1}{2}(-\sqrt{2})\sin(-\sqrt{2}t) \right) = \\ &2\left(\frac{\sqrt{2}}{4}\sin(\sqrt{2}t) - \frac{\sqrt{2}}{8}\sin(2\sqrt{2}t)\right) = \\ &\frac{\sqrt{2}}{2}(\sin(\sqrt{2}t) - \sin(\sqrt{2}t)\cos(\sqrt{2}t)) = \frac{1}{\sqrt{2}}(1 - \cos(\sqrt{2}t))\sin(\sqrt{2}t), \end{split}$$

as expected.

The following corollary provides useful information in the context of PGST.

Corollary 9. Suppose that there is PGST from a to b. Denote the fidelity from a to b at time t by f(t), and let $\tau_m, m \in \mathbb{N}$ denote a sequence of times such that $f(\tau_m) \to 1$ as $m \to \infty$. Then $\frac{df}{dt}\Big|_{\tau_m} \to 0$ as $m \to \infty$.

Proof. Since there is PGST from a to b, there is a sequence $\gamma_m \in \mathbb{C}$ such that: i) $|\gamma_m| = 1, m \in \mathbb{N}$; and ii) $\sum_{\lambda \in \Theta_a} e^{i\tau_m \lambda} E_{\lambda} e_a - \gamma_m e_b \to 0$ as $m \to \infty$. (This observation is essentially made in [1].) Writing $e_b = \sum_{\lambda \in \Theta_a} E_{\lambda} e_b$, we may rewrite ii) as $\sum_{\lambda \in \sigma_{ab}^+} (e^{i\tau_m \lambda} - \gamma_m) E_{\lambda} e_a + \sum_{\lambda \in \sigma_{ab}^-} (e^{i\tau_m \lambda} + \gamma_m) E_{\lambda} e_a \to 0$ as $m \to \infty$.

We now deduce that: iii) for each $\lambda \in \sigma_{ab}^+, e^{i\tau_m\lambda} - \gamma_m \to 0$ as $m \to \infty$; and iv) for each $\lambda \in \sigma_{ab}^-, e^{i\tau_m\lambda} + \gamma_m \to 0$ as $m \to \infty$. Hence, we find that if $\lambda_j, \lambda_\ell \in \sigma_{ab}^+$, then $e^{i\tau_m(\lambda_j - \lambda_\ell)} \to 1$ as $m \to \infty$. Similarly, $e^{i\tau_m(\lambda_j - \lambda_\ell)} \to 1$ as $m \to \infty$ whenever $\lambda_j, \lambda_\ell \in \sigma_{ab}^-$, while $e^{i\tau_m(\lambda_j - \lambda_\ell)} \to -1$ as $m \to \infty$ whenever $\lambda_j \in \sigma_{ab}^+, \lambda_\ell \in \sigma_{ab}^-$. In particular, as $m \to \infty$, we have $\sin(\tau_m(\lambda_j - \lambda_\ell)) \to 0$ if either $\lambda_j, \lambda_\ell \in \sigma_{ab}^+$, or $\lambda_j, \lambda_\ell \in \sigma_{ab}^-$, or $\lambda_j \in \sigma_{ab}^+, \lambda_\ell \in \sigma_{ab}^-$. The conclusion now follows from Theorem 7.

4 Algebraic and combinatorial structure of trees of diameter 4

We now consider the properties of trees of diameter 4 that will allow us to consider PGST and readout times. We first observe that if we delete the leaves of a tree with diameter 4, we obtain a star on at least 3 vertices. This allows us to describe the structure of a tree with diameter 4 as a centre vertex (which we denote by v henceforth) with some number of adjacent stems, each of which has a number of adjacent leaves. (For ease of notation, we will consider a leaf of the centre vertex to be a stem with 0 adjacent leaves.) We begin this section by classifying the pairs of strongly cospectral vertices of trees of diameter 4, as this is a necessary condition for PGST.

Theorem 10. Let T be a tree of diameter 4 described as follows: there is a $t \in \mathbb{N}$ and parameters $q_j, a_j, j = 1, ..., t$ such that $q_1 < q_2 < \cdots < q_t$, and for each j = 1, ..., t there are a_j stems that are adjacent to the central vertex v, and also adjacent to q_j leaves. Then vertices a and b of T are strongly cospectral if and only if one of the following holds:

- (a) a and b are the only neighbours of v with q_i leaves (i.e. $a_i = 2$),
- (b) a and b are the only leaves at distance 2 from v with no twin (i.e. $q_j = 1, a_j = 2$),
- (c) a and b are the only leaves of some stem, and the central vertex v has at least one leaf (i.e. $q_1 = 0, q_i = 2$).

Proof. Assume a and b are strongly cospectral. Then $\deg(a) = \deg(b)$ and a and b have the same number of vertices at distance 2. Suppose a is a neighbour of v so that the number of vertices at distance 2 from a is $\sum_{j=1}^t a_j - 1$, while $\deg(a) = q_{j_0} + 1$ for some j_0 . If b = v, then the number of vertices at distance 2 from b is $\sum_{j=1}^t a_j q_j$ and $\deg(b) = \sum_{j=1}^t a_j$. Suppose now that $\sum_{j=1}^t a_j - 1 = \sum_{j=1}^t a_j q_j$ and $q_{j_0} + 1 = \sum_{j=1}^t a_j$. From the first equation we find that $\sum_{j=1}^t a_j (q_j - 1) = -1$, so that necessarily $q_1 = 0$ and $a_1 = \sum_{j=2}^t a_j (q_j - 1) + 1$. But then the second equation yields $q_{j_0} + 1 = a_1 + \sum_{j=2}^t a_j = \sum_{j=2}^t a_j q_j + 1$. From this we deduce that $q_{j_0} \geq 1$ (i.e. $j_0 \geq 2$) and hence that $q_{j_0} + 1 \geq a_{j_0} q_{j_0} + 1$, with strict inequality if $t \geq 3$. We conclude that necessarily t = 2, in which case T has diameter 3, contrary to our hypothesis.

Suppose b is distance 2 from v, in particular, b is a leaf. Then a is also a leaf. Let w be the neighbour of b, then $\deg(w) = \deg(v)$. If $\deg(v) = 2$, then T is a double star, contracting that T has diameter 4. If $\deg(v) \geq 3$, then w is adjacent to another leaf, and there is an automorphism swapping this vertex and b fixing all other vertices, so by Lemma 5, a and b are not strongly cospectral, a contradiction.

Finally, suppose b is a neighbour of v. If there exists a third neighbour of v with the same number of leaves, then there is an automorphism swapping this vertex and its leaves with y and its leaves fixing all other vertices, so by Lemma 5, a and b are not strongly cospectral, a contradiction. Hence, if a is a neighbour of v, then (a) holds.

Now, suppose a is a leaf of T that is not a neighbour of v. Then b is also a leaf of T, and from the argument given above we may assume that b is not a neighbour of v. Since T has diameter 4, we have $v \neq a, b$. If a and b do not have a common neighbour, then the unique neighbours of a and b must have degree 2, otherwise there is an automorphism swapping a or b with its twin and fixing all other vertices, so by Lemma 5, a and b are not strongly cospectral, a contradiction. Moreover, if there is a third neighbour of v with degree 2, then there is an automorphism swapping this stem-leaf pair with b and its stem and fixing all other vertices, so by Lemma 5, a and b are not strongly cospectral, a contradiction. Hence, if a is a leaf with no twin, then (b) holds.

Finally, if a and b have a common neighbour w, then $\deg(w)=3$, otherwise there exists a leaf $z\neq a,b$ adjacent to w, and then there is an automorphism swapping b and z and fixing all other vertices, so by Lemma 5, a and b are not strongly cospectral, a contradiction. Suppose v is adjacent to no leaves. We observe T has eigenvectors α with $\alpha(a)=1$, $\alpha(b)=-1$, and zero otherwise; and β with $\beta(v)=-1$, $\beta(\ell)=\frac{1}{q_i}$ for each leaf ℓ of a stem with q_i leaves, and zero otherwise. Then α is an eigenvector corresponding to eigenvalue 0 for which $\alpha(a)=-\alpha(b)$, and β is an eigenvector corresponding to eigenvalue 0 for which $\beta(a)=\beta(b)$, which contradicts strong cospectrality. Hence, if a is a leaf with a twin, then a holds.

Conversely, we demonstrate that each case is strongly cospectral. Suppose a and b are not strongly cospectral. Without loss of generality, there exists an eigenvector α with $\alpha(a) = 0$ and $\alpha(b) = 1$ corresponding to some eigenvalue λ .

(a) Suppose a and b are the only neighbours of v with q_j leaves. First suppose a and b are leaves. It follows that $\alpha(v) = \lambda = 0$. Moreover, α must be 0 on every stem of T. But then $(A\alpha)(v) = 1 \neq 0$,

a contradiction. Therefore, there is no such eigenvector, so a and b are strongly cospectral. Now suppose a and b are not leaves. Considering a leaf of a, we must have $\lambda = 0$, but considering a leaf of b, we must have $\lambda \neq 0$, a contradiction. Therefore, there is no such eigenvector, so a and b are strongly cospectral.

- (b) Suppose a and b are the only leaves at distance 2 from v with no twin. Considering a, we have that α assigns its stem 0 and so $\alpha(v) = 0$. Considering b, we have that α assigns its stem λ and so $\alpha(v) = \lambda^2 1$. It follows that $\lambda = \pm 1$. Considering v, there must be a neighbour z of v with nonzero entry c. Then each other neighbour of z has entry $\pm \frac{c}{q_j} = \frac{c}{\lambda}$, so $q_j = 1$, contradicting that a and b are the only leaves with no twin. Therefore, there is no such eigenvector, so a and b are strongly cospectral.
- (c) Suppose a and b are the only leaves of some stem, and the central vertex has at least one leaf. Let s be their common neighbour. It follows that $\alpha(s) = \lambda = 0$ and $\alpha(v) = -1$. Since every stem is assigned 0 by α , there is a leaf ℓ of v with nonzero entry, but $(A\alpha)(\ell) = -1 \neq 0$. Therefore, there is no such eigenvector, so a and b are strongly cospectral.

We now analyze the eigenvalues of a tree of diameter 4.

Lemma 11. Let T be a tree of diameter 4 consisting of a central vertex v adjacent to a_j vertices each with q_j leaves, where $q_1 < q_2 < \cdots < q_t$ for some integer t. Then the roots of the equation

$$\sum_{j=1}^{t} \frac{a_j}{x^2 - q_j} = 1 \tag{2}$$

are the nonzero eigenvalues of T in the support of v. Moreover, the squares of those nonzero eigenvalues satisfy

$$0 \le q_1 < \lambda_1^2 < q_2 < \lambda_2^2 < \dots < q_t < \lambda_t^2.$$

Proof. Let w be an eigenvector of T in the support of v with eigenvalue x. Without loss of generality, we may assume w(v)=1. Let s be a neighbour of v with q_j leaves. Then for each leaf ℓ of s, we have $xw(\ell)=w(s)$ and $xw(s)=q_jw(\ell)+1$. It follows that $w(\ell)=\frac{1}{x^2-q_j}$ and $w(s)=\frac{x}{x^2-q_j}$. Finally, we must have $x=xw(v)=\sum_{s\sim v}w(s)=\sum_{s\sim v}\frac{x}{x^2-q_j}$. Dividing through by x and grouping vertices with the same number of leaves gives the desired relation. We note in passing that from the argument above, the eigenvector w is unique up to scalar multiple.

argument above, the eigenvector w is unique up to scalar multiple. Conversely, if λ satisfies $\sum_{j=1}^t \frac{a_j}{x^2-q_j} = 1$, then the vector w given by w(v) = 1, $w(s) = \frac{\lambda}{\lambda^2-q_j}$ for each neighbour s of v with q_j leaves, and $w(\ell) = \frac{1}{\lambda^2-q_j}$ for each leaf ℓ adjacent to s is an eigenvector with eigenvalue λ in the support of v.

It follows from the intermediate value theorem that for each $\ell=1,\ldots,t-1$ there is a root λ of (2) such that $q_{\ell}<\lambda^2< q_{\ell+1}$, and similarly there is a root λ such that $q_{\ell}<\lambda^2$. Now, suppose there exist eigenvalues λ,μ such that $q_{\ell}<\lambda^2<\mu^2< q_{\ell+1}$ for some $\ell\leq t$. Then we have

$$1 = \sum_{j=1}^{t} \frac{a_j}{\lambda^2 - q_j} > \sum_{i=1}^{k} \frac{a_j}{\mu^2 - q_j} = 1,$$

a contradiction. Moreover, if $\lambda^2 < q_1$, we have

$$1 = \sum_{j=1}^{t} \frac{a_j}{\lambda^2 - q_j} < 0,$$

a contradiction. The result follows.

Working with (2) for each of the eigenvalues in the support of v, we consider fixed eigenvalues and numbers of leaves, and solve for the parameters counting the number of stems with a given number of leaves.

Lemma 12. Let q_1, q_2, \ldots, q_t and $\lambda_1, \lambda_2, \ldots, \lambda_t$ be given and satisfy $0 \le q_1 < \lambda_1^2 < q_2 < \lambda_2^2 < \cdots < q_t < \lambda_t^2$. Consider the linear system in $\{a_j\}$ given by

$$1 = \sum_{j=1}^{t} \frac{a_j}{\lambda_i^2 - q_j}, 1 \le i \le t.$$

Then the solution is given by

$$a_j = \frac{\prod_{i=1}^t (\lambda_i^2 - q_j)}{\prod_{i \neq j} (q_i - q_j)}.$$

Proof. The coefficient matrix for the linear system is given by the Cauchy matrix

$$M = \left[\frac{1}{\lambda_h^2 - q_i}\right]_{1 \le h, i \le t}.$$

For each j = 1, ..., t, let M_j denote the matrix formed from M by replacing its j-th column with the all-ones vector.

We have that

$$\det(M) = \frac{\prod_{h=2}^{t} \prod_{i=1}^{h-1} (\lambda_h^2 - \lambda_i^2) (q_h - q_i)}{\prod_{h=1}^{t} \prod_{i=1}^{t} (\lambda_h^2 - q_i)}$$

(see [18], for example) and

$$\begin{split} \det(M_j) &= \frac{\prod_{h=1}^t (\lambda_h^2 - q_j)}{\prod_{i \neq j} (q_i - q_j)} \det(M) \sum_{g=1}^t \frac{\prod_{i \neq j} (\lambda_g^2 - q_i)}{\prod_{h \neq g} (\lambda_h^2 - \lambda_g^2)} \\ &= \frac{\prod_{h=1}^t (\lambda_h^2 - q_j)}{\prod_{i \neq j} (q_i - q_j)} \det(M) \sum_{g=1}^t \frac{(-1)^{g-1} \prod_{i \neq j} (\lambda_g^2 - q_i) \prod_{g \notin \{h, i\}} (\lambda_h^2 - \lambda_i^2)}{\prod_{h=2}^t \prod_{i=1}^{h-1} (\lambda_h^2 - \lambda_i^2)}. \end{split}$$

By a tedious calculation, we obtain

$$\det(M_j) = \frac{\prod_{h=1}^t (\lambda_h^2 - q_j)}{\prod_{i \neq j} (q_i - q_j)} \det(M)$$

and therefore by Cramer's Rule, we obtain

$$a_j = \frac{\det(M_j)}{\det(M)} = \frac{\prod_{h=1}^t (\lambda_h^2 - q_j)}{\prod_{i \neq j} (q_i - q_j)}$$

as claimed.

Remark 13. Lemma 11 identifies the nonzero eigenvalues in the support of the central vertex v. It is straightforward to determine the remaining eigenvalues of A as follows, and we subdivide the discussion into two cases.

Case 1: $q_1 \geq 1$. In this scenario the remaining eigenvalues consist of $\pm \sqrt{q_j}$ with multiplicity $a_j - 1, j = 1, \ldots, t$, as well as 0 with multiplicity $1 + \sum_{j=1}^t a_j(q_j - 1)$. We note that in this case the eigenvalue 0 is in the support of v.

Case 2: $q_1 = 0$. In this scenario the remaining eigenvalues are $\pm \sqrt{q_j}$ with multiplicity $a_j - 1, j = 1, \ldots, t$, as well as 0 with multiplicity $a_1 - 1 + \sum_{j=2}^t a_j(q_j - 1)$. In this case the eigenvalue 0 is not in the support of v.

5 PGST for trees of diameter 4

Theorem 10 characterizes the three scenarios in which a pair of vertices in a tree of diameter 4 exhibits strong cospectrality. In this section, we consider each of those scenarios and analyse suitable trees in which there is PGST between the strongly cospectral vertices. Specifically, we construct families of trees of diameter 4 with convenient sets of eigenvalues that allow us to demonstrate PGST and in addition, to determine explicit readout times that ensure high fidelity. Our results rely heavily on the machinery of Section 4.

We begin with the family with two types of stems: leaves adjacent, which corresponds to Theorem $10\ c)$.

Theorem 14. Suppose that $t=2, q_1=0, q_2=2,$ and that k>1 is an odd integer. Set $a_1=k^2, a_2=k^2-1$. Suppose further that there are sequences of positive odd integers μ_n, ν_n such that $\lim_{n\to\infty}\frac{\mu_n}{\nu_n}=\sqrt{2}$, and let $\delta_n=\nu_n\sqrt{2}-\mu_n$. Letting $f(\tau)$ denote the fidelity (at time τ) between leaves adjacent to a common stem, we have

$$f\left(\frac{\mu_n \pi}{\sqrt{2}}\right) = \left(1 - \frac{1}{2(k^2 - 1)} \left(1 - \cos\left(\frac{\delta_n \pi}{\sqrt{2}}\right)\right)\right)^2. \tag{3}$$

In particular, when n is odd, we have

$$f\left(\frac{(1+\sqrt{2})^n + (1-\sqrt{2})^n}{2\sqrt{2}}\pi\right) = \left(1 - \frac{1}{2k^2 - 1}\left(1 - \cos\left(\frac{(\sqrt{2}-1)^n}{\sqrt{2}}\pi\right)\right)\right)^2.$$
(4)

Proof. Our graph has a total of $4k^2 - 2$ vertices, and from the results in section 4 it follows that the eigenvalues of the adjacency matrix A are:

$$0^{(2k^2-2)},1,-1,\sqrt{2}k,-\sqrt{2}k,\sqrt{2}^{(k^2-2)},-\sqrt{2}^{(k^2-2)}$$

(here the superscripts denote multiplicities). Suppose for concreteness that vertices 1 and 2 are leaves adjacent to a common stem.

We have the following observations.

- 1. $\frac{1}{\sqrt{2}}(e_1 e_2)$ is a null vector, and this can be extended to a basis of the null space in which all remaining null vectors are 0 in the first two positions.
- 2. For each of the eigenvalues 1 and -1, there is a corresponding (1,-1) eigenvector.
- 3. For each of the eigenvalues $\sqrt{2}k$ and $-\sqrt{2}k$, there is a corresponding normalized eigenvector whose first entry is $\frac{1}{\sqrt{2(k^2-1)(4k^2-2)}}$.
- 4. For each eigenvector corresponding to eigenvalue $\sqrt{2}$, we may construct a corresponding eigenvector corresponding to $-\sqrt{2}$ by changing the signs of the entries associated with all vertices on one side of the bipartition.

For each eigenvalue λ with corresponding orthogonal eigenprojection matrix E_{λ} , let $s_{\lambda}=(E_{\lambda})_{1,1}$. From observations 1–3, we find that $s_0=\frac{1}{2}, s_1=s_{-1}=\frac{1}{4k^2-2}$, and $s_{\sqrt{2}k}=s_{-\sqrt{2}k}=\frac{1}{2(k^2-1)(4k^2-2)}$. From observation 4, we have $s_{\sqrt{2}}=s_{-\sqrt{2}}$, and from the fact that $e_1^T(\sum_{\lambda\in\Theta_1}E_{\lambda})e_1=1$, we have $s_{\sqrt{2}}+s_{-\sqrt{2}}=1-s_0-s_1-s_{-1}-s_{\sqrt{2}k}-s_{-\sqrt{2}k}$. Hence, $s_{\sqrt{2}}=s_{-\sqrt{2}}=\frac{k^2-2}{4(k^2-1)}$.

It now follows that

$$\begin{split} e_1^T e^{i\tau A} e_2 &= -\frac{1}{2} + \frac{1}{4k^2 - 2} (e^{i\tau} + e^{-i\tau}) + \frac{1}{2(k^2 - 1)(4k^2 - 2)} (e^{i\tau\sqrt{2}k} + e^{-i\tau\sqrt{2}k}) \\ &+ \frac{k^2 - 2}{4(k^2 - 1)} (e^{i\tau\sqrt{2}} + e^{-i\tau\sqrt{2}}) \\ &= -\frac{1}{2} + \frac{1}{2k^2 - 1} \cos(\tau) + \frac{1}{(k^2 - 1)(4k^2 - 2)} \cos(\tau\sqrt{2}k) + \frac{k^2 - 2}{2(k^2 - 1)} \cos(\tau\sqrt{2}). \end{split}$$

Consider $\tau = \frac{\mu_n \pi}{\sqrt{2}}$. From our hypotheses on μ_n, ν_n , we see that $\cos(\mu_n \pi) = -1, \cos(k\mu_n \pi) = -1$ and $\cos\left(\frac{\mu_n\pi}{\sqrt{2}}\right) = -\cos\left(\frac{\delta_n\pi}{\sqrt{2}}\right)$. The expression (3) now follows.

Next we consider a specific sequence of rational approximations to $\sqrt{2}$. Take

$$\alpha_n = \frac{(1+\sqrt{2})^n + (1-\sqrt{2})^n}{2},$$

$$\beta_n = \frac{(1+\sqrt{2})^n - (1-\sqrt{2})^n}{2\sqrt{2}},$$
(5)

and recall that $\frac{\alpha_n}{\beta_n}$ is the sequence of convergents associated with the continued fraction expansion for $\sqrt{2}$ (see sequences A001333 and A000129 in [20], for example). In particular, $\frac{\alpha_n}{\beta_n} \to \sqrt{2}$ as $n \to \infty$, and it is readily determined that α_n, β_n are both odd when n is odd. A straightforward computation shows that $\beta_n \sqrt{2} - \alpha_n = (-1)^{n+1} (\sqrt{2} - 1)^n$, and (4) now follows.

Example 15. We illustrate Theorem 14 with a small numerical example. Taking k=3, the corresponding tree has 34 vertices. Considering the convergent in (5) with n=3 we have $\alpha_3=7,\beta_3=$ 5 and a Matlab computation yields $f(7\pi/\sqrt{2}) \approx 0.99853...$ Taking n=5 we have $\alpha_5=41, \beta_5=29$ and another computation gives $f(51\pi/\sqrt{2}) \approx 0.99995...$

The following result also considers Theorem 10 c), in the case that t = 3. Observe that as in Theorem 14, the sequence of readout times $\frac{\alpha_n \pi}{\sqrt{2}}$, n odd, is advantageous.

Theorem 16. Let $k_2, k_3 \in \mathbb{N}$ with $2 < k_2 < k_3$. Suppose that T is a diameter 4 tree with t = 3 and parameters $q_1 = 0, q_2 = 2, q_3, a_1, a_2, a_3$, and that a, b are the only leaves of some stem. The eigenvalue support of a (and b) contains the values $\pm k_2\sqrt{2}$ and $\pm k_3\sqrt{2}$ if and only if i) $2k_2^2 < q_3 < 2k_3^2$, ii) $q_3|2k_2^2k_3^2$, iii) $(q_3-2)|2(k_2^2-1)(k_3^2-1)$, and iv) $a_1 = \frac{2k_2^2k_3^2}{q_3}$, $a_2 = \frac{2(k_2^2-1)(k_3^2-1)}{q_3-2}$, $a_3 = (q_3-1)(a_1-a_2-1)$. In the event that i)-iv) hold and one of k_2 and k_3 is even, there is no PGST from a to b. If

i)-iv) hold and both k_2 and k_3 are odd, then letting f(t) denote the a to b fidelity at time t, we have

$$f\left(\frac{\alpha_n \pi}{\sqrt{2}}\right) = \left(1 - \frac{q_3 - 1}{(2k_2^2 - 1)(2k_3^2 - 1)} \left(1 - \cos\left(\frac{(\sqrt{2} - 1)^n}{\sqrt{2}}\pi\right)\right)\right)^2,$$

where n is odd and α_n is as in (5).

Proof. Observe that the eigenvalue support of a consists of: $0, \pm \sqrt{2}$ and $\pm \lambda_j, j = 1, \ldots, 3$, where $0 < \lambda_1 < \lambda_2 < \lambda_3$ are the positive roots of the equation

$$\frac{a_1}{x^2} + \frac{a_2}{x^2 - 2} + \frac{a_3}{x^2 - q_3} = 1. ag{6}$$

From Lemma 12 we find that $a_1 = \frac{\lambda_1^2 \lambda_2^2 \lambda_3^2}{2q_3}$, $a_2 = \frac{(\lambda_1^2 - 2)(\lambda_2^2 - 2)(\lambda_3^2 - 2)}{-2(q_3 - 2)}$, $a_3 = \frac{(\lambda_1^2 - q_3)(\lambda_2^2 - q_3)(\lambda_3^2 - q_3)}{-q_3(2 - q_3)}$. Note also that necessarily $0 < \lambda_1^2 < 2 < \lambda_2^2 < q_3 < \lambda_3^2$.

Suppose now that the support of a contains $k_2\sqrt{2}$ and $k_3\sqrt{2}$; then it must be the case that $\lambda_2^2=2k_2^2, \lambda_3^2=2k_3^2$. Since $\lambda_1^2=\frac{2a_1q_3}{4k_2^2k_3^2}\in\mathbb{Q}$, we find that $\lambda_1^2\in\mathbb{N}$. As $0<\lambda_1^2<2$, we conclude that $\lambda_1=1$. It now follows by setting x=1 in (6) that necessarily $a_3=(q_3-1)(a_1-a_2-1)$. Since $a_1=\frac{\lambda_1^2\lambda_2^2\lambda_3^2}{2q_3}=\frac{2k_2^2k_3^2}{q_3}$, we see that $q_3|2k_2^2k_3^2$. Similarly, since $a_2=\frac{(\lambda_1^2-2)(\lambda_2^2-2)(\lambda_3^2-2)}{-2(q_3-2)}=\frac{2(k_2^2-1)(k_3^2-1)}{q_3-2}$, we see that $(q_3-2)|2(k_2^2-1)(k_3^2-1)$. This establishes the necessity of conditions i)-iv). Conversely, if conditions i)-iv) hold, it is readily checked that both $k_2\sqrt{2}$ and $k_3\sqrt{2}$ are solutions to the equation $a_1+a_2+a_3=1$.

Conversely, if conditions i)-iv) hold, it is readily checked that both $k_2\sqrt{2}$ and $k_3\sqrt{2}$ are solutions to the equation $\frac{a_1}{x^2} + \frac{a_2}{x^2-2} + \frac{a_3}{x^2-q_3} = 1$. Suppose henceforth that i)-iv) hold. Suppose that one of k_2, k_3 is even, and for concreteness we

Suppose henceforth that i)-iv) hold. Suppose that one of k_2, k_3 is even, and for concreteness we assume that in fact k_2 is even. Note that $(k_2 - 1) \times 0 + (-k_2) \times \sqrt{2} + (1) \times k_2 \sqrt{2} = 0, k_2 - 1$ is odd, and $(k_2 - 1) + (-k_2) + (1) = 0$. From Theorem 2 of [1], we deduce that there cannot be PGST from a to b.

Now suppose that both k_2 and k_3 are odd. The eigenvalues in the support of a and b are: $0, \pm 1, \pm \sqrt{2}, \pm k_2 \sqrt{2}$ and $\pm k_3 \sqrt{2}$. Note that $\sigma^- = \{0\}$. Then $(E_0)_{a,b} = -\frac{1}{2}, (E_{\pm 1})_{a,b} = \frac{q_3 - 1}{2(2k_2^2 - 1)(2k_3^2 - 1)}$ and

$$\begin{split} (E_{\sqrt{2}})_{a,b} + (E_{-\sqrt{2}})_{a,b} + (E_{k_2\sqrt{2}})_{a,b} + (E_{-k_2\sqrt{2}})_{a,b} + (E_{k_3\sqrt{2}})_{a,b} + (E_{-k_3\sqrt{2}})_{a,b} = \\ \frac{1}{2} - (E_1)_{a,b} - (E_{-1})_{a,b}. \end{split}$$

It now follows that

$$f\left(\frac{\alpha_n \pi}{\sqrt{2}}\right) = \left(1 - 2(E_1)_{a,b} \left(1 - \cos\left(\frac{(\sqrt{2} - 1)^n}{\sqrt{2}}\pi\right)\right)\right)^2 = \left(1 - \frac{q_3 - 1}{(2k_2^2 - 1)(2k_3^2 - 1)} \left(1 - \cos\left(\frac{(\sqrt{2} - 1)^n}{\sqrt{2}}\pi\right)\right)\right)^2.$$

Example 17. Here is an example that illustrates Theorem 16. Suppose that t = 3, $a_1 = 99$, $q_1 = 0$, $a_2 = 96$, $q_2 = 2$, $a_3 = 42$, $q_3 = 22$. This graph has a total of 1354 vertices, and the eigenvalues of the adjacency matrix (with corresponding multiplicities) are $0^{(1076)}$, ± 1 , $\pm \sqrt{2}^{(95)}$, $\pm 3\sqrt{2}$, $\pm \sqrt{22}^{(41)}$, $\pm 11\sqrt{2}$. If a and b are the only leaves of some stem, then we have

$$(E_0)_{a,b} = -\frac{1}{2}; (E_{\pm 1})_{a,b} = \frac{21}{8194}; (E_{\pm \sqrt{2}})_{a,b} = \frac{95}{384};$$

$$(E_{\pm 3\sqrt{2}})_{a,b} = \frac{1}{30464}; (E_{\pm \sqrt{22}})_{a,b} = 0; (E_{\pm 11\sqrt{2}})_{a,b} = \frac{11}{1295616};$$

It now follows that

$$U(\tau)_{a,b} = -\frac{1}{2} + \frac{21}{4097}\cos(\tau) + \frac{95}{192}\cos(\sqrt{2}\tau) + \frac{1}{15232}\cos(3\sqrt{2}\tau) + \frac{11}{647808}\cos(11\sqrt{2}\tau).$$

In particular for n odd and α_n given by (5), we have

$$f\left(\frac{\alpha_n \pi}{\sqrt{2}}\right) = \left(1 - \frac{21}{4097} \left(1 - \cos\left(\frac{(\sqrt{2} - 1)^n}{\sqrt{2}}\pi\right)\right)\right)^2.$$

For instance, $f\left(\frac{7\pi}{\sqrt{2}}\right) \approx 0.999873...$ and $f\left(\frac{41\pi}{\sqrt{2}}\right) \approx 0.999996...$

Our next result considers the scenario of Theorem 10 a).

Theorem 18. Suppose that $q \in \mathbb{N}$ and consider the diameter 4 tree with $t = 1, q_1 = q, a_1 = 2$. For each $\ell \in \mathbb{N} \cup \{0\}$, define

$$D_{\ell} = \frac{r_1^{2\ell}(2q+1-r_2)(r_1-1) - r_2^{2\ell}(2q+1-r_1)(r_2-1)}{2(r_1-r_2)},$$

where $r_1 = q + 1 + \sqrt{q^2 + 2q}$, $r_2 = q + 1 - \sqrt{q^2 + 2q}$. Set $\delta_{\ell} = \frac{r_2^{2\ell}(r_2 - 1)}{2} \left(\sqrt{\frac{q + 2}{q}} - 1 \right)$ and $\tau_{\ell} = \frac{D_{\ell}\pi}{\sqrt{q}}$. The stem-to-stem fidelity at time τ_{ℓ} is $\frac{1}{4}(1 + \cos(\delta_{\ell}\pi))^2$.

Proof. For the stems a,b we have $\sigma_{ab}^+ = \{\pm \sqrt{q+2}\}$, $\sigma_{ab}^- = \{\pm \sqrt{q}\}$ and for each of the corresponding unit eigenvectors, the entries corresponding to the stems are $\frac{1}{2}$ or $-\frac{1}{2}$. It follows that at any time τ , the stem-to-stem fidelity is $\frac{1}{4}(-\cos(\tau\sqrt{q})+\cos(\tau\sqrt{q+2}))^2$.

Define

$$N_{\ell} = \frac{r_1^{2\ell}(2q+3-r_2)(r_1-1) - r_2^{2\ell}(2q+3-r_1)(r_2-1)}{2(r_1-r_2)}.$$

We claim that if q is odd, then for each $\ell \in \mathbb{N}$, N_{ℓ} and D_{ℓ} are even and odd natural numbers, respectively. (It turns out that $\frac{N_{\ell}}{D_{\ell}}$ is a convergent arising from the continued fraction expansion for $\sqrt{\frac{q+2}{q}}$, hence our interest in N_{ℓ} and D_{ℓ} .)

To see the claim, set $x(1) = \begin{bmatrix} q+1 \\ 1 \end{bmatrix}, y(1) = \begin{bmatrix} q \\ 1 \end{bmatrix}$ and define the recurrences $x(j+1) = \begin{bmatrix} 2q+1 & q \\ 2 & 1 \end{bmatrix} x(j)$ and $y(j+1) = \begin{bmatrix} 2q+1 & q \\ 2 & 1 \end{bmatrix} y(j), j \in \mathbb{N}$. Evidently $x(j), y(j) \in \mathbb{N}^2$ for all $j \in \mathbb{N}$, and a straightforward induction proof shows that whenever j is odd, $e_1^T x(j)$ is even, while $e_2^T x(j), e_1^T y(j), e_2^T y(j)$ are odd. The eigenvalues of $\begin{bmatrix} 2q+1 & q \\ 2 & 1 \end{bmatrix}$ are readily seen to be r_1, r_2 , with corresponding eigenvectors $v_1 = \begin{bmatrix} r_1 - 1 \\ 2 \end{bmatrix}, v_2 = \begin{bmatrix} r_2 - 1 \\ 2 \end{bmatrix}$, respectively.

We can write

$$x(1) = \frac{(2q+3-r_2)(r_1-1)}{2(r_1-r_2)}v_1 - \frac{(2q+3-r_1)(r_2-1)}{2(r_1-r_2)}v_2$$

and

$$y(1) = \frac{(2q+1-r_2)(r_1-1)}{2(r_1-r_2)}v_1 - \frac{(2q+1-r_1)(r_2-1)}{2(r_1-r_2)}v_2$$

so as to deduce that

$$x(2\ell+1) = \frac{r_1^{2\ell}(2q+3-r_2)(r_1-1)}{2(r_1-r_2)}v_1 - \frac{r_2^{2\ell}(2q+3-r_1)(r_2-1)}{2(r_1-r_2)}v_2$$

and

$$y(2\ell+1) = \frac{r_1^{2\ell}(2q+1-r_2)(r_1-1)}{2(r_1-r_2)}v_1 - \frac{r_2^{2\ell}(2q+1-r_1)(r_2-1)}{2(r_1-r_2)}v_2.$$

In particular we find that for each $\ell \in \mathbb{N} \cup \{0\}$, $N_{\ell} = e_1^T x(2\ell+1)$ is an even natural number, while $D_{\ell} = e_1^T y(2\ell+1)$ is an odd natural number. This completes the proof of the claim.

In the case that q is even, an analogous argument shows that for each $\ell \in \mathbb{N}$, N_{ℓ} and D_{ℓ} are odd and even, respectively.

A computation shows that $D_{\ell}\sqrt{\frac{q+2}{q}}-N_{\ell}=\delta_{\ell}$. Hence, when q is odd, at time τ_{ℓ} we have $\cos(\tau_{\ell}\sqrt{q})=\cos(D_{\ell}\pi)=-1$ and $\cos(\tau_{\ell}\sqrt{q+2})=\cos(D_{\ell}\sqrt{\frac{q+2}{q}}\pi)=\cos((N_{\ell}+\delta_{\ell})\pi)=\cos(\delta_{\ell}\pi)$.

Similarly, when q is even, we have $\cos(\tau_{\ell}\sqrt{q}) = \cos(D_{\ell}\pi) = 1$ and $\cos(\tau_{\ell}\sqrt{q+2}) = \cos(D_{\ell}\sqrt{\frac{q+2}{q}}\pi) = \cos((N_{\ell} + \delta_{\ell})\pi) = -\cos(\delta_{\ell}\pi)$.

The conclusion now follows.

Remark 19. Observe that in Theorem 18, the case q = 1 yields the graph P_5 , which, according to Theorem 1, exhibits PGST between next-to-leaf vertices. Thus Theorem 18 provides an advantageous sequence of readout times for a family of trees that includes P_5 .

An analysis similar to that in Theorem 18 also shows that for P_5 the fidelity between end vertices at time $D_\ell \pi$ is equal to $\left(\frac{5}{6} + \frac{\cos(\delta_\ell \pi)}{6}\right)^2$. Hence that sequence of readout times also ensures good fidelity between the leaves of P_5 .

The following technical result will be useful in the proof of Theorem 21 below.

Lemma 20. Suppose that G is a graph with strongly cospectral vertices a,b and denote the eigenvalues in σ_{ab} by $\lambda_j, j=1,\ldots,m$. Let $0<\epsilon<\frac{1}{2}$ be given, and suppose that for some $\tau>0$ there are integers $n_j, j=1,\ldots,m$ such that $|\tau\lambda_j-n_j|<\frac{\epsilon}{2\pi}$ for each $\lambda_j\in\sigma_{ab}^+$ and $|\tau\lambda_j-n_j-\frac{1}{2}|<\frac{\epsilon}{2\pi}$ for each $\lambda_j\in\sigma_{ab}^-$. Then the a to b fidelity at time $2\pi\tau$ is bounded below by $1-2\epsilon$.

Proof. Since a and b are strongly cospectral, it follows that there are positive numbers $c_j, j = 1, ..., m$ with $\sum_{j=1}^{m} c_j = 1$ such that the fidelity f at time $2\pi\tau$ can be written as

$$f = \left| \sum_{\lambda_j \in \sigma_{ab}^+} c_j e^{2\pi\tau i \lambda_j} - \sum_{\lambda_j \in \sigma_{ab}^-} c_j e^{2\pi\tau i \lambda_j} \right|^2$$

$$= \left| \sum_{\lambda_j \in \sigma_{ab}^+} c_j + \sum_{\lambda_j \in \sigma_{ab}^-} c_j + \sum_{\lambda_j \in \sigma_{ab}^+} (c_j e^{2\pi\tau i \lambda_j} - 1) - \sum_{\lambda_j \in \sigma_{ab}^-} c_j (e^{2\pi\tau i \lambda_j} + 1) \right|^2$$

$$\geq \left| 1 - \sum_{\lambda_j \in \sigma_{ab}^+} c_j \left| e^{2\pi\tau i \lambda_j} - 1 \right| - \sum_{\lambda_j \in \sigma_{ab}^-} c_j \left| e^{2\pi\tau i \lambda_j} + 1 \right|^2,$$

since $\sum_{j=1}^{m} c_j = 1$ together with an application of the triangle inequality. If $\lambda_j \in \sigma_{ab}^+$ we have

$$|e^{2\pi\tau i\lambda_j} - 1| = \sqrt{2(1 - \cos(2\pi\tau\lambda_j))} \le \sqrt{2(1 - \cos(\epsilon))} \le \epsilon.$$

We find similarly that if $\lambda_j \in \sigma_{ab}^-$ then $|e^{2\pi\tau i\lambda_j} + 1| \le \epsilon$. The conclusion now follows.

We next attempt to localize an advantageous readout time in the context of Theorem 10 b).

Theorem 21. Suppose that t=2, $a_1=2$, $q_1=1$, and let a,b denote the leaves on the branches at v corresponding to q_1 . Suppose that for each eigenvalue λ in the support of a we have $\lambda^2 \in \mathbb{N}$. Then $q_2 > 2$, and the eigenvalues in the support of a are $\pm 1, \pm \sqrt{2}, \pm \sqrt{2q_2 - 1}$. In particular, there is PGST between a and b if and only if $2q_2 - 1$ is not a perfect square.

is PGST between a and b if and only if $2q_2-1$ is not a perfect square. Suppose $2q_2-1$ is not a perfect square and that $0<\epsilon<\frac{1}{2}$. There is a τ_0 such that $|\tau_0|\leq \frac{2^{24}3^{11}\pi^4(2q_2-1)^7}{\epsilon^3}+\pi$ such that the a to b fidelity is bounded below by $1-2\epsilon$.

Proof. Observe that $\pm 1 \in \sigma_{ab}^-$. Suppose that the branches at v not containing a or b have stems of degree q_2 . Then σ_{ab}^+ consists of 0, along with the roots of the following equation: $\frac{2}{x^2-1} + \frac{a_2}{x^2-q_2} = 1$. Denote those roots by $\pm \lambda_1, \pm \lambda_2$ where $0 < \lambda_1 < \lambda_2$, and observe that if $q_2 = 0$, then $0 < \lambda_1 < 1$, contrary to our hypothesis. Hence, $q_2 \geq 2$.

We have $2=\frac{(\lambda_1^2-1)(\lambda_2^2-1)}{q_2-1}$ and since $1<\lambda_1^2< q_2<\lambda_2^2$, we find that $2>\lambda_1^2-1$. As $\lambda_1^2\in\mathbb{N}$, it must be the case that $\lambda_1^2=2$. Substituting $x^2=2$ into the equation $\frac{2}{x^2-1}+\frac{a_2}{x^2-q_2}=1$, we find that $a_2=q_2-2$. It is now straightforward to determine that $x^2=2q_2-1$ yields the other root of the equation $\frac{2}{x^2-1}+\frac{q_2-2}{x^2-q_2}=1$. Hence $\sigma_{ab}^+=\{\pm\sqrt{2},\pm\sqrt{2q_2-1}\}$.

If $2q_2-1$ is not a perfect square, we find that $\pm 1, \pm \sqrt{2}, \pm \sqrt{2q_2-1}$ are independent over \mathbb{Q} , which ensures that there is PGST between a and b. Next, suppose that $2q_2-1$ is a perfect square, say with $\sqrt{2q_2-1}=k\in\mathbb{N}$. Consider the following integer parameters: $\ell_1=-k,\ell_1'=0,\ell_{\sqrt{2}}=\frac{k-1}{2},\ell_1'=\frac{k-1}{2},\ell_2'=0,\ell_1'=0$. Then the following equations are satisfied:

$$(1)\ell_1 + (-1)\ell'_1 + (\sqrt{2})\ell_{\sqrt{2}} + (-\sqrt{2})\ell'_{\sqrt{2}} + (k)\ell_k + (-k)\ell'_k + (0)\ell_0 = 0,$$

$$\ell_1 + \ell'_1 = -k \text{ (odd)},$$

$$\ell_1 + \ell'_1 + \ell_{\sqrt{2}} + \ell'_{\sqrt{2}} + \ell_k + \ell'_k + \ell_0 = 0.$$

By Theorem 2 of [1], PGST between a and b cannot hold.

Next we suppose that $2q_2-1$ is not a perfect square, and that $0<\epsilon<\frac{1}{2}$. According to Theorem 3.1 of [12], there is an integer r and integers n_1,n_2 such that $|r\sqrt{2}+\frac{\sqrt{2}}{2}-n_1|\leq \frac{\epsilon}{2\pi},|r\sqrt{2q_2-1}+\frac{\sqrt{2q_2-1}}{2}-n_2|\leq \frac{\epsilon}{2\pi}$. Evidently for such an r we have $|(2r+1)\pi\sqrt{2}-2\pi n_1|<\epsilon,|(2r+1)\pi\sqrt{2q_2-1}-2\pi n_1|<\epsilon$. Further, that result of [12] provides an upper bound on |r| that depends on: the degrees of the minimal polynomials of $\sqrt{2q_2-1}$ and $\sqrt{2}$, the Weil heights (see [25]) of $\sqrt{2q_2-1}$ and $\sqrt{2}$, and the dimension of $\mathbb{Q}(\sqrt{2q_2-1},\sqrt{2})$ over \mathbb{Q} . Computing those quantities we find that the minimal polynomials have degree 2, that the Weil heights of $\sqrt{2q_2-1}$ and $\sqrt{2}$ are $\sqrt{2q_2-1}$ and $\sqrt{2}$, respectively, and that the dimension of $\mathbb{Q}(\sqrt{2q_2-1},\sqrt{2})$ over \mathbb{Q} is 4. Using that information and then applying the upper bound of [12, Theorem 3.1], we find that

$$|r| \le 2^{-8} 6^8 (3(\sqrt{2q_2-1})^2)^3 (2\sqrt{2})^8 (2\sqrt{2q_2-1})^8 \left(\frac{\epsilon}{2\pi}\right)^{-3} = \frac{2^{23} 3^{11} \pi^3 (2q_2-1)^7}{\epsilon^3}.$$

Setting τ_0 equal to $\pi(2r+1)$ now yields the desired conclusion.

Remark 22. The coefficient $2^{24}3^{11}\pi^4$ in Theorem 21 is on the order of 10^{14} .

Evidently the readout time bound of Theorem 21 is quite large. The next example shows that under certain circumstances, a more precise statement can be made about readout times corresponding to good fidelity.

Example 23. Here we consider the family of trees dealt with in Theorem 21, and maintain the notation of that result. Note that the bound on τ_0 in Theorem 21 does little to inform us on the value of a suitable τ_0 . In this example we show how q_2 and the readout time can be chosen together in such a way as to provide a sequence of trees and readout times for which there is good fidelity of state transfer between the leaves a, b that are at distance 4. For this family of trees we find the following entries in the relevant eigenprojection matrices: $(E_{\pm 1})_{a,a} = \frac{1}{4}, (E_{\pm \sqrt{2}})_{a,a} = \frac{q_2 - 2}{8q_2 - 12}, (E_{\pm \sqrt{2}q_2 - 1})_{a,a} = \frac{1}{4(4q_2^2 - 8q_2 + 3)}, (E_0)_{a,a} = \frac{q_2}{4q_2 - 2}.$

Suppose that α_n, β_n are as in (5). It is straightforward to show (for example by induction) that α_n is odd for all $n \in \mathbb{N}$, while β_n is odd or even according as n is odd or even. Observe that $(\beta_n + \beta_{n+1})\sqrt{2} = \alpha_n + \alpha_{n+1} + (-1)^{n+1}(2 - \sqrt{2})(\sqrt{2} - 1)^n$. Now select $q_2 = \frac{(\beta_n + \beta_{n+1})^2 + 3}{2}$, so that $2q_2 - 1 = (\beta_n + \beta_{n+1})^2 + 2$. We find that

$$(\beta_n + \beta_{n+1})\sqrt{2q_2 - 1} = (\beta_n + \beta_{n+1})\sqrt{(\beta_n + \beta_{n+1})^2 + 2}$$

$$= (\beta_n + \beta_{n+1})^2 + 1 - \frac{1}{(\beta_n + \beta_{n+1})\sqrt{(\beta_n + \beta_{n+1})^2 + 2} + (\beta_n + \beta_{n+1})^2 + 1}.$$

Set $t_0 = (\beta_n + \beta_{n+1})\pi$. Letting f(t) denote the fidelity of ab state transfer at time t, we have

$$f(t_0) = \left| -\frac{1}{2}\cos(t_0) + \frac{q_2}{4q_2 - 2} + \frac{q_2 - 2}{4q_2 - 6}\cos(\sqrt{2}t_0) + \frac{1}{(4q_2 - 2)(2q_2 - 3)}\cos(\sqrt{2}q_2 - 1t_0) \right|^2$$

$$= \left| 1 - \frac{q_2 - 2}{4q_2 - 6}(1 - \cos(\sqrt{2}t_0)) - \frac{1}{(4q_2 - 2)(2q_2 - 3)}(1 - \cos(\sqrt{2}q_2 - 1t_0)) \right|^2.$$

From our observations above, we find that

$$\cos(\sqrt{2}t_0) = \cos((2-\sqrt{2})\pi(\sqrt{2}-1)^n),$$

and

$$\cos(\sqrt{2q_2 - 1}t_0) = \cos\left(\frac{\pi}{(\beta_n + \beta_{n+1})\sqrt{(\beta_n + \beta_{n+1})^2 + 2} + (\beta_n + \beta_{n+1})^2 + 1}\right).$$

For example, taking n = 2 we have $t_0 = 7\pi$, $q_2 = 26$ and $f(t_0) \approx 0.9759...$ With n = 3 we get $t_0 = 17\pi$, $q_2 = 146$ and $f(t_0) \approx 0.9979...$

Our final three examples serve to quantify Corollary 9 by providing the value of the sensitivity of the fidelity in some of the families of trees studied in this section.

Example 24. Here we compute the sensitivity of the fidelity with respect to the readout time in the setting of Theorem 18. Maintaining the notation of that result, we find that $\cos(\tau_{\ell}\sqrt{q}) = (-1)^q, \sin(\tau_{\ell}\sqrt{q}) = 0, \cos(\tau_{\ell}\sqrt{q+2}) = (-1)^{q-1}\cos(\delta_{\ell}\pi), \sin(\tau_{\ell}\sqrt{q+2}) = (-1)^{q-1}\sin(\delta_{\ell}\pi)$. Denoting the stems by a, b, we have $\sigma_{ab}^+ = \{\pm\sqrt{q}+2\}, \sigma_{ab}^- = \{\pm\sqrt{q}\}, \text{ and } s_{\lambda} = \frac{1}{4} \text{ for each eigenvalue } \lambda \text{ in the support of } a.$

Appealing to (1), it follows that

$$\frac{df}{dt}\Big|_{\tau_{\ell}} = \frac{1}{8} \{ -2\sqrt{q+2}\sin(2\tau_{\ell}\sqrt{q+2}) - 2\sqrt{q}\sin(2\tau_{\ell}\sqrt{q}) + 2(\sqrt{q+2} - \sqrt{q})\sin((\sqrt{q+2} - \sqrt{q})\tau_{\ell}) + 2(\sqrt{q+2} + \sqrt{q})\sin((\sqrt{q+2} + \sqrt{q})\tau_{\ell}) \}.$$

Substituting the various sine and cosine values, we find that

$$\frac{df}{dt}\Big|_{\tau_{\ell}} = \frac{1}{8} \{-2\sqrt{q+2}\sin(2\delta_{\ell}\pi) - 4\sqrt{q+2}\sin(\delta_{\ell}\pi)\}
= -\frac{\sin(\delta_{\ell}\pi)(1+\cos(\delta_{\ell}\pi))\sqrt{q+2}}{2}.$$

Example 25. We now consider the sensitivity of the end-to-end fidelity for P_5 (see Remark 19 above). For end vertices a,b we have $\sigma_{ab}^+ = \{0,\pm\sqrt{3}\}, \sigma_{ab}^- = \{\pm1\}$, and adopting the notation of Theorem $7, s_0 = \frac{1}{3}, s_{\pm\sqrt{3}} = \frac{1}{12}, s_{\pm1} = \frac{1}{4}$. Using (1) and the values of the sin and cos functions at τ_ℓ and $\sqrt{3}\tau_\ell$ found in Example 24, a computation reveals that the sensitivity of the end-to-end fidelity at time τ_ℓ is given by $-\frac{\sqrt{3}\sin(\delta_\ell\pi)(5+\cos(\delta_\ell\pi))}{18}$.

Example 26. Here we revisit Theorem 14, and compute the sensitivity of state transfer between leaves of the same stem at time $\tau = \frac{\alpha_n \pi}{\sqrt{2}}$ when n is odd. Denote the two leaves by a,b, and observe that $\sigma_{ab}^- = \{0\}$, while $\sigma_{ab}^+ = \{\pm 1, \pm \sqrt{2}, \pm \sqrt{2k}\}$. We have $s_0 = \frac{1}{2}, s_{\pm 1} = \frac{1}{4k-2}, s_{\pm \sqrt{2}} = \frac{k-2}{4(k-1)}, s_{\pm \sqrt{2k}} = \frac{1}{2(k-1)(4k-2)}$.

Set $\gamma_n = \frac{(\sqrt{2}-1)^n}{\sqrt{2}}$, so that $\frac{\alpha_n\pi}{\sqrt{2}} = \beta_n\pi - \gamma_n\pi$ (recall that β_n is odd). It now follows that $\sin(\sqrt{2}\tau) = 0$, $\cos(\sqrt{2}\tau) = -1$, $\sin(\sqrt{2k}\tau) = 0$, $\cos(\sqrt{2k}\tau) = -1$, $\sin(\tau) = \sin(\gamma_n\pi)$, $\cos(\tau) = -\cos(\gamma_n\pi)$. Referring to (1), we see that the first summation inside the brackets contributes $2s_0s_1\sin(\gamma_n\pi)$, while the second summation is zero. A long computation (in which many terms cancel or are zero) yields the fact that the third summation is equal to $2s_1^2\sin(2\gamma_n\pi) + 4s_1s_{\sqrt{2}}\sin(\gamma_n\pi) + 4s_1s_{\sqrt{2k}}\sin(\gamma_n\pi)$. Substituting the three summations into (1), we then find that the sensitivity of the fidelity at time τ is given by

$$4s_1(s_0\sin(\gamma_n\pi) + s_1\sin(2\gamma_n\pi) + 2s_{\sqrt{2}}\sin(\gamma_n\pi) + s_{\sqrt{2k}}\sin(\gamma_n\pi)),$$

which in turn can be written as

$$\frac{4\sin(\gamma_n\pi)}{4k-2} \left(\frac{1}{2} + \frac{2\cos(\gamma_n\pi)}{4k-2} + \frac{k-2}{2(k-1)} + \frac{2}{2(k-1)(4k-2)} \right)$$

This last expression can be simplified to yield

$$\frac{2\sin(\gamma_n\pi)(2k-2+\cos(\gamma_n\pi))}{(2k-1)^2}.$$

In Examples 24–26 we see that the readout times do not necessarily correspond to the optimal fidelity (i.e. local maxima) in their respective neighbourhoods. Nevertheless, each of those sequences of readout times provides a convenient, systematic approach to achieving high fidelity state transfer, suffers little impact from minor perturbations, and approaches optimal fidelity.

References

- [1] L. Banchi, G. Coutinho, C. Godsil, S. Severini. Pretty good state transfer in qubit chains: the Heisenberg Hamiltonian. *Journal of Mathematical Physics* 58, 032202 (2017).
- [2] A. Casaccino, S. Lloyd, S. Mancini, S. Severini. Quantum state transfer through a qubit network with energy shifts and fluctuations. *International Journal of Quantum Information* 7 (08), 1417–1427 (2009).
- [3] M. Christandl, N. Datta, A. Ekert, A.J. Landahl. Perfect state transfer in quantum spin networks. *Physical Review Letters* **92**, 187902 (2004).
- [4] M. Christandl, N. Datta, T.C. Dorlas, A. Ekert, A. Kay, A.J. Landahl. Perfect state transfer of arbitrary states in quantum spin networks. *Physical Review A* 71, 032312 (2005).
- [5] G. Coutinho. Quantum State Transfer in Graphs. Ph.D. Dissertation, University of Waterloo (2014).
- [6] G. Coutinho, C. Godsil, E. Juliano, C.M. van Bommel. Quantum walks do not like bridges. Linear Algebra and its Applications 652, 155–172 (2022).
- [7] G. Coutinho, K. Guo, C.M. van Bommel. Pretty good state transfer between internal nodes of paths. Quantum Information & Computation 17, 825–830 (2017).
- [8] G. Coutinho, E. Juliano, T. J. Spier. Strong cospectrality in trees. *Algebraic Combinatorics* 6 (4), 955–963 (2023).
- [9] G. Coutinho, E. Juliano, T. J. Spier. No perfect state transfer in trees with more than 3 vertices. Journal of Combinatorial Theory, Series B 168, 68–85 (2024).
- [10] G. Coutinho, H. Liu. No Laplacian perfect state transfer in trees. SIAM Journal on Discrete Mathematics 29(4), 2179–2188 (2015).

- [11] X. Fan, C. Godsil. Pretty good state transfer on double stars. *Linear Algebra and its Applications* 438(5), 2346–2358 (2013).
- [12] L. Fukshansky, N. Moshchevitin. On an effective variation of Kronecker's approximation theorem avoiding algebraic sets. *Proceedings of the American Mathematical Society* 146, 4151–4163 (2018).
- [13] H. Hou, R. Gu, M. Tong. Pretty good state transfer on 1-sum of star graphs. *Open Mathematics* 16.1, 1483–1489 (2018).
- [14] C. Godsil. State transfer on graphs. Discrete Mathematics 312(1), 129–147 (2012).
- [15] C. Godsil. When can perfect state transfer occur? *Electronic Journal of Linear Algebra* **23**, 877–890 (2012).
- [16] C. Godsil, S. Kirkland, S. Severini, J. Smith. Number-theoretic nature of communication in quantum spin systems. *Physical Review Letters* 109.5, 050502 (2012).
- [17] C. Godsil, J. Smith. Strongly cospectral vertices. Australasian Journal of Combinatorics 88(1), 1–21 (2024).
- [18] R. Horn, C. Johnson. Matrix Analysis, Second Edition. Cambridge University Press, 2013.
- [19] M. Kempton, G. Lippner, S.-T. Yau. Perfect state transfer on graph with a potential. *Quantum Information & Computation* 17, 303–327 (2017).
- [20] N. J. A. Sloane et al. The On-Line Encyclopedia of Integer Sequences. Available at https://oeis.org, 2024.
- [21] D. Stevanović. Applications of graph spectra in quantum physics. *Zbornik Radova. (Beograd)* 14(22), 85–111 (2011).
- [22] C.M. van Bommel. A complete characterization of pretty good state transfer on paths. *Quantum Information & Computation* **19**, 601–608 (2019).
- [23] L. Vinet, A. Zhedanov. Dual-1 Hahn polynomials and perfect state transfer. *Journal of Physics, Conference Series* 343.1, 12125 (2012).
- [24] L. Vinet, A. Zhedanov. How to construct spin chains with perfect state transfer. Physical Review A 85.1, 12323 (2012).
- [25] U. Zannier. Basics on the theory of heights and its applications to certain diophantine problems. Expositiones Mathematicae 36, 1–31 (2018).