

Note on entanglement and edge modes

J.S.Dowker¹

*Theory Group,
School of Physics and Astronomy,
The University of Manchester,
Manchester, England*

A recent numerical evaluation of the spherical universal log coefficient in the Maxwell free-energy and its decomposition into bulk and edge contributions via a bounded hyperbolic geometry mode calculation is shown to be equivalent to an existing compact formulation for p -forms, at $p = 1$, on a conically deformed sphere. The edge mode seems to be a ghost $(p - 1)$ -form. Some numbers are given.

¹ dowkeruk@yahoo.co.uk

1. Introduction

A recent article, [1], has reinvestigated the notion of 'edge modes' introduced in order to reconcile computed values of the entanglement entropy and the divergent log term in the Maxwell free-energy (for even dimensions).

The geometries involved were the d -sphere, for d even, and the static patch of d -de Sitter space with novel 'dynamic' boundary conditions imposed at a stretched horizon in order to justify an earlier, no brick-wall calculation.

Numerical evaluations of various quantities were performed using character integrals which incorporate the relevant degeneracies and eigenvalues. The 'bulk' contribution was obtained via an integral over the de Sitter static patch, as in [2]. The numbers for the log coefficient were presented up to $d = 16$.

The content of the present very short note is only to point out that these numbers can readily be computed without leaving the compact arena by working on the conically deformed d -sphere which is also conveniently relevant for determining the Rényi and entanglement entropies.

Such a calculation is expounded for p -forms in [3], where all the details can be found. It yields, after the usual mode properties and the ghosts-for ghosts sum have been incorporated, the total universal log coefficient, $\mathcal{F}(q)$, in the free-energy, as an explicit *rational* function of the deformation parameter, q , where $2\pi/q$ is the conical angle. The number, q , starts out as an orbifold integer but can be extended into the reals by sight.

As a simple check of the general formulae, for conformally invariant propagation by (minus) the de Rham Laplacian, *i.e.* for $p = (d - 2)/2$, the total is just the standard conformal anomaly as early computed by Cappelli and d'Appollonio, [4]. See also Raj, [5]. The Maxwell theory employed in [1] means a vector field, $p = 1$, for all dimensions, which limits the immediate conformal significance to $d = 4$.

The hyperbolic constituents can be extracted easily from the resulting expression for $\mathcal{F}(q)$. Thus, the value at $q = 1$, $\mathcal{F}(1)$, is the total value and the term independent of q is the 'shift' or 'correction', denoted here by Δ . The difference is then the bulk contribution, B .

2. Numerology and comments

The explicit expressions derived in [3] follow from a standard spectral argument which organises the p -form eigenvalues and degeneracies on the conically deformed

d -sphere into generating functions. These are the analogues of the characters employed in [1] and in works cited therein.

No reference to boundary conditions was, of course, needed in [3], ², in sharp contrast to the analysis in [1] where appropriate conditions have to be devised, and tested which is a consequence of working in a non-compact background.

The calculations in [3] can be taken, by machine, as far as desired regarding form order and dimension. It is unnecessary to give any specific expressions here. Evaluation at $q = 1$ gives the quantities for the full, round sphere which is the relevant geometry and, simply in order to extend the numbers listed ³ in [1], the $p = 1$ values for $d = 18$ and $d = 20$ are found to be, ⁴

	$d = 18$	$d = 20$
Bulk =	$-\frac{904237137092591}{3118343638410000}$	$-\frac{129858859323129841}{457357066966800000}$
Edge =	$-\frac{15587954101729}{62523180720000}$	$-\frac{12156987650908561}{49893498214560000}$
Total =	$-\frac{13453490783330599}{24946749107280000}$	$-\frac{447787476738749901}{2744142401800800000}$

In the conformally invariant case, the Rényi entropy can be derived from $\mathcal{F}(q)$ and thence, via the replica method, the entanglement entropy across a spherical surface. For convenience only, this terminology will be maintained for *any* p and d (even).

It turns out that $\mathcal{F}(q)$, as a function of q , is an extremum at the round sphere, $q = 1$, for *all* p and d . This important property is the mathematical reason why the entanglement entropy equals (minus) the conformal anomaly in the conformally invariant case. Moreover, it was noticed in [3] that, numerically, the p -form shift equalled the entanglement entropy for a $(p - 1)$ -form in $d - 2$ dimensions and, by the extremum property, this is minus the corresponding total quantity. In symbols

² Although the *periodic* deformed sphere was obtained by combining absolute and relative p -form mode sets on a π/q segment of the d -sphere.

³ The values were actually computed with no brick-wall.

⁴ The expressions in [3] are not geared for computation at large d . Some radical simplification is required but I could not find one for general p . The edge contribution for $d = 22$ is $-3930799044044576533/16464854410804800000$ and for $d = 24$, $-532598043781522096919/2272149908691062400000$. The values tend to a constant as $d \rightarrow \infty$.

this reads,

$$\Delta(p, d) = -\mathcal{F}(p - 1, d - 2),$$

which actually holds for all p and d . One therefore has, for the total quantity,

$$\mathcal{F}(p, d) = B(p, d) - \mathcal{F}(p - 1, d - 2), \quad \forall p, d. \quad (1)$$

A number of interpretations for the Maxwell shift discrepancy have been suggested in the literature, mostly involving 'edge' (codimension-2) modes of a ghost scalar field, of one sort or another. In the present formulation, equation (1) could be interpreted as saying that the contributing codimension-2 edge mode is a ghost $(p - 1)$ -form.

As a curiosity, this equation can be iterated to give the total value, essentially as a finite, alternating sum of bulk quantities.

References

1. Ball,A., Law,Y.T.A. and Wong,G. *Dynamical Edge Modes and Entanglement in Maxwell Theory*, arXiv:2403.14542.
2. Dowker,J.S., *Entanglement entropy on even spheres*, arXiv:1009.3854.
3. Dowker,J.S. *Rényi entropy and CT for p -forms on even dimensional spheres*, arXiv:1706.45074.
4. Cappelli,A. and D'Appollonio,G., *On the Trace Anomaly as a Measure of Degrees of Freedom Phys. Lett.* **487B** (2000) 87, hep-th 0005115 .
5. Raj,H.,*A note on sphere free energy of p -form gauge theory and Hodge duality*, *Class. Quant. Grav.* **34** (2017) 247001, arXiv:1611.02507.