

An Optimal Selection Problem Associated with the Poisson Process

Zakaria Derbazi
Queen Mary University of London

July 2, 2024

Abstract

Cowan and Zabczyk (1978) introduced a continuous-time generalisation of the best-choice (secretary) problem, where offers arrive at epochs of a homogeneous Poisson process. We expand their work to encompass the last-success problem under the Karamata-Stirling success profile. In this setting, the k th trial is a success with probability $p_k = \theta/(\theta + k - 1)$, where $\theta > 0$. In the best-choice setting ($\theta = 1$), the myopic strategy is optimal, and the proof hinges on verifying the monotonicity of certain critical roots. We extend this crucial result to the last-success case by exploiting a connection to the sign of the derivative in the first parameter of a quotient of Kummer's hypergeometric functions. Finally, we derive bounds and asymptotics of the critical roots, strengthening and improving the findings of Ciesielski and Zabczyk (1979).

1 Introduction

Cowan and Zabczyk (1978) considered a continuous-time extension of the classical secretary problem. In this scenario, the task is to select the best apartment within a fixed time horizon, T . Apartments are presented at epochs of a homogeneous Poisson process with a known intensity $\lambda > 0$. The decision-maker may accept or discard each presented apartment irrevocably. Each arrival can be ranked relative to previously seen apartments, and all permutations of ranks are assumed equally likely and independent of the arrival process.

The optimal strategy selects the k th apartment presented at time t only if the expected number of remaining arrivals in the time window $(T - t)$ falls below a predefined cutoff threshold a_k . To be precise, the condition is

$$\lambda(T - t) \leq a_k. \quad (1)$$

Each *cutoff* a_k ($k = 1, 2, \dots$) is the unique solution to the equation $h_k(x) = 0$, where $h_k(x)$ is formed by the difference between the probability of zero successes and exactly one success and is given by

$$h_k(x) := \frac{1}{k} + \sum_{n=1}^{\infty} \frac{x^n}{n!(k+n)} \left(1 - \sum_{j=1}^n \frac{1}{j+k-1} \right). \quad (2)$$

For a strategy to be optimal, the sequence of roots $(a_k)_{k \geq 0}$ must be increasing. Cowan and Zabczyk developed an ingenious argument to establish this property. They begin by establishing the uniqueness of the roots, leveraging the continuity of the function h_k and all of its derivatives,

along with the behaviour of the first and second derivatives, in the vicinity of the root a_k . Subsequently, they exploit a relationship between the change in probability of exactly one success across consecutive trials and the difference between the probabilities of one and zero successes to prove monotonicity (see p. 588–590 in [10]).

Cowan and Zabczyk’s optimal selection model can be categorised as a best-choice problem with a random number of observations, denoted N , with an a priori Poisson distribution. These observations arrive uniformly and independently at times in $[0, T]$. This formulation might appear similar to the best-choice problem introduced by Presman and Sonin (1972), in which N is a Poisson-distributed random variable. However, the key distinction lies in the role of time as a crucial decision factor [10, 5, 24, 15]. Consequently, Cowan and Zabczyk’s model is more accurately characterised as a best-choice problem with random observation times, where the number of items to be inspected has a Poisson prior distribution.

We aim to extend this framework to encompass last-success problems, where the decision-maker only observes a sequence of indicator variables. In this setting, the defining characteristic of the trials (and, consequently, the problem itself) is the sequence of success probabilities, which we refer to as the *success* profile or *record* profile. In the classical best-choice problem, the profile is given by the success probability $p_k = 1/k$ for the k th trial.

Our investigation is closely related to the theory of random records [14, 20]. Gaver (1976) played a pivotal role in linking this theory to optimal stopping problems. He examined the embedding of record times within a point process and analysed the problem of selecting the last record appearing between epochs of a Poisson process. Bruss (1988) further highlighted the strong relationship between optimal selection problems and optimal stopping related to record sequences. However, the initial work on last-success problems is attributed to Pfeifer (1989). He examined a sequence of record indicators and addressed the problem of selecting the last success within that sequence. For best-choice problems with a random number of trials, a brief overview of recent advancements is provided in [15].

A common strategy for solving last-success problems involves checking whether they fall within the *monotone case* of optimal stopping theory [8, 13]. Should this criterion be met, the optimal strategy is deduced via a nonincreasing sequence of cutoffs. These cutoffs, indexed by the number of arrivals, represent the earliest instant at which accepting the current offer (record) becomes optimal. The optimality of the myopic strategy is ensured whenever the sequence of cutoffs is nonincreasing.

Cowan and Zabczyk’s problem has led to further research on optimal selection related to Poisson arrivals. Bruss (1987) broadened the framework to accommodate inhomogeneous Poisson processes. Ano and Ando (2000) considered a random availability attached to each offer. Schwalko and Szajowski (2003) explored the variant of the problem of selecting the best or second-best apartment. Subsequently, Szajowski (2007) introduced a game-theoretic version of the problem, thus offering a distinct perspective on the optimal selection strategy. Considering the last-success version of the problem, Hofmann (1997) investigated Poisson arrivals with successes appearing according to Nevzorov’s F^α record profile. In this setting, success probabilities are given by $p_k = \alpha_k / (\alpha_1 + \dots + \alpha_k)$ for some positive α ’s. However, Hofmann fell short of demonstrating the crucial property of monotonicity of the roots. In scenarios where trials arrive based on an inhomogeneous Poisson process, the instances of stopping at the last success, the last ℓ th success, and any of the last ℓ successes were studied by Bruss (2000), Bruss and Paindaveine (2000), and Tamaki (2011) respectively.

Using the Karamata-Stirling profile to model the success probabilities in our analysis is motivated by two key factors. First, it relates to the well-known Ewens Sampling Formula (ESF) [11]. The number of distinct components within the ESF framework can be expressed as

a sum of independent indicator variables. Each of these variables follows the Karamata-Stirling profile, where the probability of success in the k th trial is given by

$$p_k = \frac{\theta}{(\theta + k - 1)}, \quad k \geq 1, \quad (3)$$

with *mutation* parameter $\theta > 0$. The case $\theta = 1$ corresponds to the classical best-choice problem. Second, the generating function of the number of successes in n Bernoulli trials under the Karamata-Stirling profile exhibits advantageous analytical properties. This characteristic allows us to strengthen and simplify the solution presented by Cowan and Zabczyk while revealing a connection between last-success problems with random observation times and hypergeometric functions. In this regard, we rely on Kummer's confluent hypergeometric function properties to demonstrate that the sequence of cutoffs (a_k) is increasing. This approach aligns with the recent work of [16, 15], who employed the Gaussian hypergeometric function to analyse other power series distributions.

The paper is structured as follows. Section 2 introduces the fundamental model based on a generic prior for the number of observations and a generic success profile. Subsequently, it derives the necessary condition for the optimality of the myopic strategy. Section 3 presents new findings related to Kummer's confluent hypergeometric function, focusing specifically on deriving an expression to determine the sign of the derivative in parameter of a quotient of Kummer's functions. Section 4 establishes the core optimality result, demonstrating the monotonicity of the cutoffs through a connection to the sign of the derivative in the first parameter of a quotient of Kummer's functions. Section 5 transitions to the scenario where observations are revealed at epochs of an inhomogeneous Poisson process. By a trivial time-change argument on the arrivals, we show that the optimal rule is similar to the homogeneous case. When the success probability is a function of the time parameter, the problem reduces to selecting the last arrival in an inhomogeneous Poisson process. This section concludes with an illustrative example that considers an asymptotic Karamata-Stirling profile as an example of a continuous-time success probability function. Subsequently, the optimal threshold and the classical optimal winning probability are derived. Finally, Section 6 generalises the two theorems of Ciesielski and Zabczyk (1979) to encompass the Karamata-Stirling profile. We improve the upper bound on the roots and derive their asymptotic values.

2 The Mathematical Model

Let $\{T_n, X_n\}_{n \geq 1}$ denote a marked Poisson process defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, where T_1, T_2, \dots are the chronologically ordered arrival epochs. The marks $\{X_n\}_{n \geq 1}$ form an i.i.d. sequence of Bernoulli random variables independent of $\{T_n\}_{n \geq 1}$ having a success profile $\mathbf{p} := (p_n)_{n \geq 1}$, that is $\mathbb{P}(X_n = 1) = p_n$. Without loss of generality, assume that $0 < p_k < 1$ for all $k \geq 1$.

The bivariate process $\{T_n, X_n\}_{n \geq 1}$ is adapted to the natural filtration $(\mathcal{F}_t)_{t \in [0, 1]} \subseteq \mathcal{F}$, where the σ -algebra \mathcal{F}_t encapsulates the complete knowledge of epochs and outcomes within the time interval $[0, t]$. The random record process of points T_n at which record marks occur ($X_n = 1$) can be regarded as a thinning of $\{T_n\}_{n \geq 1}$, where each mark T_n is independently retained with probability p_n [3].

Another way to model this process is to consider the marks X_1, X_2, \dots as a sequence of trials, whose number N is Poisson-distributed. The number of successes among trials $k, k + 1, \dots, n$

has a Poisson-binomial distribution with probability generating function

$$z \mapsto \prod_{j=k}^n (1 - p_j + zp_j).$$

A simple computation gives the probability of zero successes and the probability of one success from stage $k \geq 1$ as

$$s_0(k, n) := \prod_{j=k}^n (1 - p_j), \quad k < n, \quad (4)$$

$$s_1(k, n) := s_0(k, n) \sum_{j=k}^n \frac{p_j}{1 - p_j}, \quad k < n. \quad (5)$$

Without loss of generality, we apply a deterministic time-change to transform arrivals from $[0, T]$ to $[0, 1]$. Furthermore, given $N = n$, the trials arrive in continuous time at independent, uniformly distributed epochs U_1, U_2, \dots, U_n in $[0, 1]$. We order these arrivals chronologically so each arrival epoch T_k of the Poisson process in the original setting corresponds to the k th order statistic of U_1, \dots, U_n . We associate with each T_k , a trial X_k resulting in success with probability p_k , independently of anything else, and we let T_n undefined on the event $\{N < n\}$. The number of arrivals up to time t is a mixed binomial process defined as

$$N_t = \sum_{n=1}^N \mathbb{1}\{U_n \leq t\}, \quad t \in [0, 1].$$

We introduce the state variable $(t, k) \in [0, 1] \times \{0, 1, \dots\}$ to represent the number of arrivals observed up to any epoch t . We denote the occurrence of exactly k arrivals up to time t by the event $\{N_t = k\}$. When the k th trial occurs at time $T_k = t$ and the outcome is a success, we designate a transition to this *success* state as $(t, k)^\circ$.

By independence of the trials and the Markov property of mixed binomial processes [18], the distribution of the point process of epochs and outcomes of the trials on $(t, 1]$ both depend on \mathcal{F}_t only through N_t .

Let $\boldsymbol{\pi} = (\pi_1, \pi_2, \dots)$ be the prior distribution of the number of trials, where $\pi_n = \mathbb{P}(N = n)$. The posterior distribution factors as

$$\begin{aligned} \mathbb{P}(N = k + j \mid N_t = k) &= \frac{\binom{k+j}{k} \pi_{k+j} (1-t)^j t^k}{\sum_{n=k}^{\infty} \binom{n}{k} \pi_n (1-t)^{n-k} t^k} \\ &= f_k(t) \binom{k+j}{k} \pi_{k+j} (1-t)^j, \quad j \geq 0, \end{aligned} \quad (6)$$

where

$$f_k(t) = \left(\sum_{n=k}^{\infty} \binom{n}{k} \pi_n (1-t)^{n-k} \right)^{-1}$$

is a normalisation function. By (6), the conditional probability that there are no successes following state (t, k) is given by

$$\mathcal{S}_0(t, k) := f_k(t) \sum_{j=0}^{\infty} \pi_{k+j} \binom{k+j}{k} (1-t)^j s_0(k+1, k+j), \quad (7)$$

which is the *adapted* reward from stopping at a success state $(t, k)^\circ$. Analogously, the probability of exactly one success following state (t, k) is

$$\mathcal{S}_1(t, k) := f_k(t) \sum_{j=m}^{\infty} \pi_{k+j} \binom{k+j}{k} (1-t)^j s_1(k+1, k+j). \quad (8)$$

Similarly, this probability represents the adapted reward from stopping at the next success state $(t, k)^\circ$.

The following result recalls some properties of \mathcal{S}_1 and \mathcal{S}_0 that prove instrumental in the latter sections.

Lemma 1 (Lemma 2 in [15]). *For fixed $k \geq 1$, suppose $\pi_k > 0$. Then*

- (i) $\mathcal{S}_0(1, k) = 1$, $\mathcal{S}_1(1, k) = 0$,
- (ii) $\mathcal{S}_0(t, k)$ is nondecreasing in t , and is strictly increasing if $\sum_{j=k+1}^{\infty} \pi_j > 0$,
- (iii) There exists cutoffs $a_k \in [0, 1)$ such that $\text{sgn}(\mathcal{S}_1(t, k) - \mathcal{S}_0(t, k)) = \text{sgn}(a_k - t)$ for $t \in (0, 1]$.

We define a *stopping strategy* to be a random variable τ which takes values in the random set of times $\{T_1, \dots, T_N, 1\}$ and is adapted to \mathcal{F}_t for $t \in [0, 1]$, where $\{\tau = 1\}$ is interpreted as not stopping at all. Our analysis will focus solely on strategies stopping at success states $(T_k, k)^\circ$, where the objective is to stop at the last success. For $k \geq 1$, stopping at $(t, k)^\circ$ is regarded as a win if $T_k = t$, and no successes occur in $(t, 1)$.

With every ‘stopping set’ $A_k \subset [0, 1) \times \mathbb{N}$ we associate a *Markovian* stopping strategy which stops at the first success trial $(t, k)^\circ$ with (t, k) falling in A_k . The optimal strategy exists for arbitrary $\boldsymbol{\pi}$ and \boldsymbol{p} . Moreover, it is Markovian by a theorem on the exclusion of randomised stopping strategies (see Theorem 5.3 in [8]).

The *myopic* strategy τ^* is defined as the strategy stopping at the first state $(t, k)^\circ$ satisfying $\mathcal{S}_1(t, k) \leq \mathcal{S}_0(t, k)$. This strategy is Markovian with stopping set

$$A := \bigcup_{k=1}^{\infty} ([a_k, 1) \times \{k\}),$$

where the cutoffs (a_k) are defined in Lemma 1. Each cutoff a_k corresponds to the earliest time τ^* can accept a success trial with index k . If $(t, k) \notin A$, then stopping in the state $(t, k)^\circ$ is not optimal.

The so-called *monotone case* of the optimal stopping theory holds if A is ‘closed’, meaning that after entering A , the sequence of successes $(t, N_t)^\circ$ does not exit the set (see [8, 13]). This condition is sufficient for the optimality of τ^* . Since the sequence of successes is increasing in both components, this property of A is equivalent to the condition $a_1 \geq a_2 \geq \dots$, characterising the optimality of τ^* .

3 Hypergeometrics

Recall the definition of Kummer’s hypergeometric function

$$M(a, c, x) := \sum_{j=0}^{\infty} \frac{(a)_j}{(c)_j} \frac{x^j}{j!}, \quad (9)$$

viewed as a function of real parameters a, c and $x > 0$, where $c > a > 0$, its basic integral representation is given by

$$M(a, c, x) := \frac{\Gamma(c)}{\Gamma(a)\Gamma(c-a)} \int_0^1 e^{xu} u^{a-1} (1-u)^{c-a-1} du, \quad (10)$$

where Γ is the Gamma function, which is defined by Euler's integral of the second kind as follows

$$\Gamma(x) := \int_0^\infty u^{x-1} e^{-u} du \quad \text{for } x > 0. \quad (11)$$

Let D_x denote the derivative with respect to the variable (or the parameter) x . The following elementary results are foundational for the proofs presented in this section.

Lemma 2. *Consider two functions $f : I \rightarrow J$ and $g : I \rightarrow J$, where $I, J \subseteq (0, \infty)$. If f and g are differentiable, then it holds that*

$$\begin{aligned} \operatorname{sgn} D_x \log f(x) &= \operatorname{sgn} D_x f(x), \\ \operatorname{sgn} D_x \{f(x)g(x)\} &= \operatorname{sgn} D_x f(x) \quad \text{if } \operatorname{sgn} D_x f(x) = \operatorname{sgn} D_x g(x), \\ \operatorname{sgn} D_x \left\{ \frac{f(x)}{g(x)} \right\} &= \operatorname{sgn} D_x f(x) \quad \text{if } \operatorname{sgn} D_x f(x) = -\operatorname{sgn} D_x g(x). \end{aligned}$$

Proof. Given that f, g are strictly positive, it is evident that

$$\operatorname{sgn} D_x \log f(x) := \operatorname{sgn} \frac{D_x f(x)}{f(x)} = \operatorname{sgn} D_x f(x).$$

From this result and by definition of the log function, we have the elementary relations

$$\operatorname{sgn} D_x \{f(x)g(x)\} = \operatorname{sgn} D_x \log \{f(x)g(x)\} = \operatorname{sgn} \{D_x \log f(x) + D_x \log g(x)\},$$

and

$$\operatorname{sgn} D_x \{f(x)/g(x)\} = \operatorname{sgn} D_x \log \{f(x)/g(x)\} = \operatorname{sgn} \{D_x \log f(x) - D_x \log g(x)\}.$$

The assertions are readily demonstrated by a simple sign analysis. □

The next result provides a test for assessing the sign of expressions of the kind

$$D_x \frac{\Gamma(ax + x_1)}{\Gamma(ax + x_2)},$$

for arbitrary real numbers x_1, x_2 and a . We employ a lengthier but more instructive argument based on Euler's integral of the first kind. This approach circumvents the direct proof, which uses the logarithmic derivative of the Gamma function.

Lemma 3. *For a fixed $c \in \mathbb{R}$, the following sign identity holds:*

$$\operatorname{sgn} D_x \frac{\Gamma(x)}{\Gamma(x+c)} = -\operatorname{sgn}(c) \quad \text{where } x > 0 \text{ and } x+c > 0.$$

Proof. The derivative of the quotient of Gamma functions is given by

$$D_x \frac{\Gamma(x)}{\Gamma(x+c)} = \frac{D_x \Gamma(x) \Gamma(x+c) - D_x \Gamma(x+c) \Gamma(x)}{\Gamma^2(x+c)}.$$

We only need to check the derivative's numerator as the quotient's denominator is positive. Use Euler's representation of the Gamma function (11) to note that

$$\begin{aligned} & D_x \Gamma(x) \Gamma(x+c) - D_x \Gamma(x+c) \Gamma(x) \\ &= \int_0^\infty \frac{1}{u} e^{-u} \log(u) du \int_0^\infty v^{c-1} e^{-v} dv - \int_0^\infty v^{c-1} e^{-v} \log(v) dv \int_0^\infty \frac{1}{u} e^{-u} du \\ &= \iint_{\overline{\mathbb{R}}_+ \times \overline{\mathbb{R}}_+} \left\{ \frac{1}{uv} e^{-u-v} \right\} v^c \log(u) dudv - \iint_{\overline{\mathbb{R}}_+ \times \overline{\mathbb{R}}_+} \left\{ \frac{1}{uv} e^{-u-v} \right\} v^c \log(v) dudv \\ &= \iint_{\overline{\mathbb{R}}_+ \times \overline{\mathbb{R}}_+} \left\{ \frac{u}{v} e^{-u-v} \right\} v^c \log\left(\frac{u}{v}\right) dudv \\ &= \iint_{0 < u < v < \infty} \left\{ \frac{1}{uv} e^{-u-v} \right\} v^c \log\left(\frac{u}{v}\right) dudv + \iint_{0 < v < u < \infty} \left\{ \frac{1}{uv} e^{-u-v} \right\} v^c \log\left(\frac{u}{v}\right) dudv \\ &= \iint_{0 < u < v < \infty} \left\{ \frac{1}{uv} e^{-u-v} \right\} v^c \log\left(\frac{1}{uv}\right) dudv + \iint_{0 < u < v < \infty} \left\{ \frac{1}{uv} e^{-u-v} \right\} u^c \log\left(\frac{v}{u}\right) dudv \\ &= \iint_{0 < u < v < \infty} \left\{ \frac{1}{uv} e^{-u-v} \log\left(\frac{v}{u}\right) \right\} [u^c - v^c] dudv. \end{aligned}$$

The sign of the integrand when $u < v$ depends on the expression $u^c - v^c$ whose sign matches that of $-c$. This completes the proof. \square

Next, we proceed to the main result of this section, which is to derive an expression for the sign of derivative in the first parameter and the second parameter, respectively, of a quotient of Kummer's functions.

Theorem 4. *For $x > 0$, the following sign identities hold*

- (i) $\operatorname{sgn} D_a \left(\frac{M(a, c_1, x)}{M(a, c_2, x)} \right) = \operatorname{sgn}(c_1 - c_2)$, if $c_1 > a > 0$ and $c_2 > a > 0$,
- (ii) $\operatorname{sgn} D_c \left(\frac{M(a_1, c, x)}{M(a_2, c, x)} \right) = \operatorname{sgn}(a_2 - a_1)$, if $c > a_1 > 0$ and $c > a_2 > 0$.

Proof. We only prove the first assertion of the theorem and defer part (ii), which is potentially of broader interest, to the appendix. Use part one of Lemma 2 to reformulate the analysis to the derivative of the log of a quotient instead since

$$\operatorname{sgn} D_a \left(\frac{M(a, c_1, x)}{M(a, c_2, x)} \right) = \operatorname{sgn} D_a \log \left(\frac{M(a, c_1, x)}{M(a, c_2, x)} \right).$$

Now, let ϕ be this the integral part in Kummer's function representation (10)

$$\phi(a, c, x) := \int_0^1 e^{xu} \left(\frac{u}{1-u} \right)^{a-1} (1-u)^{c-2} du.$$

The logarithmic derivative in parameter a , where c_1, c_2 are constants satisfying $c_1 > a > 0$ and $c_2 > a > 0$, respectively, is

$$\begin{aligned}
D_a \log \left(\frac{M(a, c_1, x)}{M(a, c_2, x)} \right) &= D_a \log M(a, c_1, x) - D_a \log M(a, c_2, x) \\
&= D_a \left\{ \log \frac{\Gamma(c_1)}{\Gamma(c_2)} + \log \frac{\Gamma(c_2 - a)}{\Gamma(c_1 - a)} + \log \frac{\phi(a, c_1, x)}{\phi(a, c_2, x)} \right\} \\
&= D_a \log \frac{\Gamma(c_2 - a)}{\Gamma(c_1 - a)} + D_a \log \frac{\phi(a, c_1, x)}{\phi(a, c_2, x)}. \tag{12}
\end{aligned}$$

By the second part of Lemma 2, it is sufficient to show that the two logarithmic derivatives in (12) have the same sign. Using Lemma 3, the sign of the first quotient matches the sign of $c_2 - c_1$.

For the second quotient, check the sign of $D_a \{ \phi(a, c_1, x) / \phi(a, c_2, x) \}$. Again, we only need to check the numerator's sign after taking the derivative. For positive constants, $c_1 > a > 0$ and $c_2 > a > 0$, use the Eulerian integral representation (10) to express the derivative as a double integral in the following way

$$\begin{aligned}
&D_a \left\{ \frac{\phi(a, c_1, x)}{\phi(a, c_2, x)} \right\} \\
&= \int_0^1 e^{xu} \left(\frac{u}{1-u} \right)^{a-1} (1-u)^{c_1-2} \log \left(\frac{u}{1-u} \right) du \int_0^1 e^{xv} \left(\frac{v}{1-v} \right)^{a-1} (1-v)^{c_2-2} dv \\
&\quad - \int_0^1 e^{xu} \left(\frac{u}{1-u} \right)^{a-1} (1-u)^{c_1-2} du \int_0^1 e^{xv} \left(\frac{v}{1-v} \right)^{a-1} (1-v)^{c_2-2} \log \left(\frac{v}{1-v} \right) dv \\
&= \iint_{[0,1] \times [0,1]} e^{x(u+v)} \left(\frac{uv}{(1-u)(1-v)} \right)^{a-1} [(1-u)(1-v)]^{c_1-2} (1-v)^{c_2-c_1} \log \left(\frac{u}{1-u} \right) dudv \\
&\quad - \iint_{[0,1] \times [0,1]} e^{x(u+v)} \left(\frac{uv}{(1-u)(1-v)} \right)^{a-1} [(1-u)(1-v)]^{c_1-2} (1-v)^{c_2-c_1} \log \left(\frac{v}{1-v} \right) dudv \\
&= \iint_{[0,1] \times [0,1]} \left\{ \underbrace{e^{x(u+v)} \left(\frac{uv}{(1-u)(1-v)} \right)^{a-1} [(1-u)(1-v)]^{c_1-2}}_{\text{symmetric in } u,v} (1-v)^{c_2-c_1} \log \left(\frac{u(1-v)}{v(1-u)} \right) \right\} dudv \\
&= \iint_{0 < v < u < 1} \{ \dots \} (1-v)^{c_2-c_1} \log \left(\frac{u(1-v)}{v(1-u)} \right) dudv + \iint_{0 < u < v < 1} \{ \dots \} (1-v)^{c_2-c_1} \log \left(\frac{u(1-v)}{v(1-u)} \right) dudv \\
&= \iint_{0 < v < u < 1} \{ \dots \} (1-v)^{c_2-c_1} \log \left(\frac{u(1-v)}{v(1-u)} \right) dudv + \iint_{0 < v < u < 1} \{ \dots \} (1-u)^{c_2-c_1} \log \left(\frac{v(1-u)}{u(1-v)} \right) dudv \\
&= \iint_{0 < v < u < 1} \{ \dots \} \log \left(\frac{u(1-v)}{v(1-u)} \right) \{ (1-v)^{c_2-c_1} - (1-u)^{c_2-c_1} \} dudv.
\end{aligned}$$

In the given domain of integration where $u > v$, the expression $\log \left(\frac{u(1-v)}{v(1-u)} \right)$ is positive. This means that the sign of the last integrand depends solely on $(1-v)^{c_2-c_1} - (1-u)^{c_2-c_1}$. But, the sign of this expression is precisely that of $c_2 - c_1$ as $1-v > 1-u$.

Since the sign of both terms in the right-hand side of (12) coincides, we can conclude, through part two of Lemma 2, that the sign of $D_a \log \left(\frac{M(a, c_1, x)}{M(a, c_2, x)} \right)$ matches that of $c_2 - c_1$, which is the desired result.

□

4 The Poisson Prior

We specialise our investigation to the case where N follows a Poisson prior distribution $\boldsymbol{\pi} := (\pi_1, \pi_2, \dots)$ with intensity $\lambda > 0$, where

$$\pi_n = \frac{\lambda^n}{n!} e^{-\lambda} \quad n \geq 0.$$

This formulation recovers the original scenario where the underlying pacing process $(N_t, t \in [0, 1])$ is Poisson. By virtue of the stationarity of increments of the Poisson process, the posterior of $N - N_t$ given $N_t = k$ is again Poisson with a rescaled intensity

$$x = (1 - t)\lambda. \tag{13}$$

That is,

$$\mathbb{P}(N = k + j \mid N_t = k) = \frac{x^j}{j!} e^{-x}.$$

Based on this posterior distribution, the probability generating function of the number of successes following state (t, k) becomes

$$\begin{aligned} z \mapsto e^{-x} \sum_{j=0}^{\infty} \frac{(k + \theta z)_j}{(k + \theta)_j} \frac{x^j}{j!} &= e^{-x} M(k + \theta z; k + \theta; x) \\ &= M(\theta - \theta z; k + \theta; -x). \end{aligned}$$

where $M(a; c; x)$ is Kummer's hypergeometric function defined in (9). The last equality is a consequence of Kummer's transformation formula, where

$$M(a, c; x) = e^x M(c - a, c; -x).$$

The derivative of the generating function is given by

$$z \mapsto \theta e^{-x} D_a M(k + \theta z, \theta + k, x).$$

Explicit expressions for (7) and (8), which represent the probability of zero successes and the probability of exactly one success after state (t, k) , are as follows

$$\begin{aligned} \mathcal{S}_0(1 - x/\lambda, k) &= e^{-x} M(k; \theta + k; x), \\ \mathcal{S}_1(1 - x/\lambda, k) &= \theta e^{-x} D_a M(k; \theta + k; x). \end{aligned}$$

As \mathcal{S}_0 and \mathcal{S}_1 are power series in x , the cutoffs, defined by Lemma 1, can be recast in terms of a sole sequence of critical roots (γ_k) of equation $\mathcal{S}_0(1 - x/\lambda, k) - \mathcal{S}_1(1 - x/\lambda, k) = 0$, expressed through the equation

$$\theta D_a M(k; \theta + k; x) - M(k; \theta + k; x) = 0. \tag{14}$$

In the special case where $\theta = 1$, the equation reduces to that of Cowan and Zabczyk (2). Observe that the roots (γ_k) do not depend on the intensity λ , and are related to the cutoffs through the following identity

$$a_k = \left(1 - \frac{\gamma_k}{\lambda}\right)_+. \tag{15}$$

The optimality of the myopic strategy for all finite $\lambda > 0$ is equivalent to the condition $\gamma_k \uparrow$. That is, the roots must be nondecreasing with k . Notably, Cowan and Zabczyk posed and ingeniously solved this problem in the classical success profile setting ($\theta = 1$). The argument for uniqueness relies on showing that each derivative of $h_k(x) := \mathcal{S}_0(1-x/\lambda, k) - \mathcal{S}_1(1-x/\lambda, k)$ has a unique root. The uniqueness is achieved by exploiting the continuity of this function and its derivatives. They approach the problem inductively by starting at higher derivatives and continuing iteratively to lower ones until $h_k(x)$ is reached. They demonstrated the monotonicity of roots based on a relation between $h_{k+1}(x)$ and the derivative of an expression involving $\mathcal{S}_1(1-x/\lambda, k) - \mathcal{S}_1(1-x/\lambda, k+1)$. In contrast, our analysis of Cowan and Zabczyk's problem reveals a distinct structure characterised by a prominent connection to Kummer's hypergeometric function. The following theorem strengthens their original result by utilising a more general success profile and provides a more straightforward proof that capitalises on the properties of Kummer's hypergeometric function.

Theorem 5. *The roots γ_k are increasing. Therefore, the stopping rule τ^* which calls for stopping at state $(t, k)^\circ$ satisfying*

$$(1-t)\lambda \leq \gamma_k, \tag{16}$$

is optimal.

Proof. First, we need establish that the roots γ_k of equation (14) are increasing and hence $\gamma_{k+1} > \gamma_k$ for all $k \geq 1$. It is sufficient to show that

$$D_a \log(M(k, \theta + k, x)) < D_a \log(M(k, \theta + k + 1, x)), \quad x \in (0, 1).$$

But this holds by the first identity of Lemma 4. Consequently, by assertion (iii) of Lemma 1, it is optimal to stop at time $t \geq a_k$, where the cutoff a_k is given by (15). Using the definition of the cutoff, we obtain $t \geq 1 - \gamma_k/\lambda$. Rearranging this inequality yields the desired result. \square

Remark 6. In the setting where λ is random with a Gamma prior distribution, $(N_t, t \in [0, 1])$ becomes a negative binomial process (mixed Poisson). This scenario has been addressed under different guises in [14, 21, 19, 15].

One can extend Theorem 5 to the inhomogeneous case in a straightforward manner through a time-change argument, reducing the problem to its homogeneous counterpart. Nonetheless, the next section tackles the inhomogeneous case and presents an example that illustrates the reduction of the last-success problem in continuous time to the problem of stopping at the last arrival of a Poisson process.

5 The Inhomogeneous Case

We now turn to the case where the counting process (N_t) is an inhomogeneous Poisson process with known time-dependent intensity $\lambda(t)$, where the function

$$\Lambda(s) = \int_0^s \lambda(u) du, \tag{17}$$

is continuous and $s \in [0, 1]$.

In the best-choice case, Theorem 3 in [5] established the optimal strategy for selecting the best apartment, where arrivals are governed by (N_t) . The extension of this result to the last-success setting is immediate.

Theorem 7. Consider the last success problem with success profile (3), where trials arrive according to an inhomogeneous Poisson process $(N_t, t \in [0, 1])$ with mean arrivals given by (17). The optimal stopping strategy for selecting the last success is to stop at state $(t^*, k)^\circ$ such that

$$\Lambda(1) - \Lambda(t^*) \leq \gamma_k,$$

where γ_k is the root of equation (14).

Proof. Given $N_1 = n$, the trials arrive at independent epochs U_1, U_2, \dots in $[0, 1]$. These arrival times have a common distribution function F , where

$$F(s) = \frac{\Lambda(s)}{\Lambda(1)}. \quad (18)$$

The points $F(U_1), F(U_2), \dots$ are renewal times of a homogeneous Poisson process in $[0, 1]$ with known intensity $\Lambda(1)$. Therefore, by Theorem 5, the myopic strategy τ^* is optimal and the optimal rule is to stop at the success state $(t^*, k)^\circ$ satisfying

$$[1 - F(t^*)] \Lambda(1) \leq \gamma_k, \quad (19)$$

which is a straightforward generalisation of (16). Plugging (18) into (19) completes the proof. \square

5.1 The Case of the Poisson Process of Successes

Let $p(t)$ denote the success probability of a trial occurring at time t . A success at time t is independent of the arrival process (N_t) and previous outcomes. Our objective is to maximise the probability of stopping at the last arrival time that coincides with the occurrence of a success.

Fix a positive integer m and let $\delta_m := \{0 = t_0 < t_1 < \dots < t_m = 1\}$ be an partition of $[0, 1]$. Denote the length of the k th subinterval $[t_{k-1}, t_k]$ by Δt_k .

Let p_k be the probability of a success occurring in the interval $(t_{k-1}, t_k]$. From the independence of successes and the properties of the Poisson process it follows that

$$p_k = p(t_k)\lambda(t_k)\Delta t_k + o(\Delta t_k).$$

As $\|\delta_m\| := \max_{1 \leq i \leq m} \Delta t_i$ tends to 0, the well defined function

$$s(t) := \lim_{\Delta t \rightarrow 0} \frac{p(t)\lambda(t)\Delta t + o(\Delta t)}{\Delta t} = p(t)\lambda(t). \quad (20)$$

is referred to as the limiting *success intensity* by Bruss [6]. This function represents a random record process of points $\lambda(t)$ at which a record occurs with probability $p(t)$. This thinning of (N_t) preserves the Poisson process property, where each point is independently retained with probability $p(t)$. We denote this new thinned Poisson process of *successes* by $(P_t, t \in [0, 1])$. Each arrival generated by (P_t) , whose intensity is $s(t)$, corresponds to a success event.

Let

$$\begin{cases} m(t) := \mathbb{E}[P_t] = \int_0^t s(u)du, \text{ and} \\ M(t_0, t_1) := m(t_1) - m(t_0) = \int_{t_0}^{t_1} s(u)du, \end{cases}$$

denote respectively the expected number of arrivals of (P_t) up to time t and the expected number of arrivals in the time interval $[t_0, t_1]$, where $0 \leq t_0 < t_1 < 1$.

In the context of the last success problem in continuous time, the optimal stopping rule τ^* translates to stopping at the last arrival time of the inhomogeneous process (P_t) . This case is much simpler than Cowan and Zabczyk's problem, given that all the arrivals are successes. Consequently, it is optimal to stop at the next arrival of (P_t) , if any, which occurs after time t^* , where

$$t^* := \sup \left\{ 0, \sup \{t \in [0, 1] : M(t, 1) \geq 1\} \right\}, \text{ and } \sup\{\emptyset\} = -\infty.$$

The threshold t^* corresponds to the last time the expected number of future arrivals of (P_t) is at least 1. The optimal stopping rule τ^* can be readily extended to the problem of stopping at the last ℓ th success [7, 12]. This generalisation follows intuitively where stopping is optimal after time t^* given by

$$t^* := \sup \left\{ 0, \sup \{t \in [0, 1] : M(t, 1) \geq \ell\} \right\}. \quad (21)$$

The following example illustrates a problem involving a continuous-time Karamata-Stirling profile combined with an inhomogeneous arrival process.

Example. Suppose the mutation parameter of the Karamata-Stirling profile exhibits a linear relationship with the number of trials n , such that a finite *mutation rate* $\kappa \in (0, \infty)$ exists. In this case

$$\kappa := \lim_{n \rightarrow \infty} \frac{\theta}{n}. \quad (22)$$

To introduce a continuous-time adaptation of the Karamata-Stirling profile, we divide both the numerator and denominator of (3) by n and let $t = (k - 1)/n$, where $k = 1, \dots, n$, be the proportion of trials elapsed. As $n \rightarrow \infty$, we obtain an asymptotic Karamata-Stirling profile, denoted by $p(t)$, where

$$\begin{aligned} p(t) &:= \lim_{n \rightarrow \infty} \frac{\theta/n}{\theta/n + (k-1)/n} \\ &= \frac{\kappa}{\kappa + t}, \quad t \in [0, 1]. \end{aligned}$$

With this continuous-time Karamata-Stirling profile established, we shift our focus to analysing the arrival process governed by the following intensity function

$$\lambda(t) = (t + \kappa)^\alpha,$$

where $\alpha \in \mathbb{R}$ is a fixed number and κ is the mutation rate defined earlier in (22).

The success intensity (20) becomes

$$s(t) = \kappa(t + \kappa)^{-1+\alpha}.$$

When $\alpha = 0$, the original counting process (N_t) is homogeneous and (P_t) , the Poisson process of successes, is inhomogeneous as $s(t) = p(t)$. The expected number of arrivals (successes) between $0 \leq t_0 < t_1$ and $t_1 < 1$ is given by

$$M(t_0, t_1) = \kappa \log \left(\frac{t_1 + \kappa}{t_0 + \kappa} \right). \quad (23)$$

Using (21), the optimal rule for selecting the last ℓ th success, where $\ell \geq 1$ is to stop at the first arrival occurring after time t^* , where

$$t^* = \sup \left\{ 0, \sup \{t \in [0, 1] : t \leq (\kappa + 1)e^{-\ell/\kappa} - \kappa\} \right\}.$$

Next, recall that $P_1 - P_t$ has a Poisson distribution with intensity $M(t, 1)$, therefore use (23) to express the probability of ℓ arrivals as

$$\mathbb{P}[P_1 - P_t = \ell] = \frac{1}{\ell!} \left(-\kappa \log \left(\frac{t + \kappa}{1 + \kappa} \right) \right)^\ell \left(\frac{t + \kappa}{1 + \kappa} \right)^\kappa. \quad (24)$$

Setting $t = (\kappa + 1)e^{-\ell/\kappa} - \kappa$ in (24), we obtain the asymptotic optimal winning probability $e^{-\ell} \ell^\ell / \ell!$ for stopping at the last ℓ th success [7, 12]. This result translates to the ubiquitous winning probability $1/e$ in the last success setting ($\ell = 1$).

Now suppose that $\alpha \neq 0$. An easy calculation gives the expected number of arrivals of (P_t) between times $t_0 \in [0, t_1)$ and $t_1 \in (0, 1)$ as

$$M(t_0, t_1) = \frac{\kappa}{\alpha} \{ (t_1 + \kappa)^\alpha - (t_0 + \kappa)^\alpha \}.$$

Consequently, the optimal threshold becomes, in this case

$$t^* := \sup \left\{ 0, \sup \{t \in [0, 1] : t \leq \left((1 + \kappa)^\alpha - \frac{\alpha \ell}{\kappa} \right)^{\frac{1}{\alpha}} - \kappa \} \right\},$$

and the asymptotic optimal winning probability is once again $e^{-\ell} \ell^\ell / \ell!$.

In the next section, we revert to the homogenous Poisson setting with a discrete Karamata-Stirling profile to assess the asymptotic properties of the roots obtained in Section 4.

6 Asymptotics and Bounds

Unlike other priors, such as the negative binomial addressed in [15], the roots γ_k do not converge. There is no Poisson process with infinite prior, which ensures that only finitely many nonzero cutoffs exist for each $\lambda \in (0, \infty)$. [10, 9, 2, 5] observed that the analogues of γ_k 's do not accumulate, rather grow about linearly with k .

In the best-choice setting, where $\theta = 1$, [10] showed that $\gamma_k \sim k$ for large k and [9] derived the bounds

$$(e - 1)(k - 1) \leq \gamma_k \leq 4e[(e - 1)k + 1],$$

and asymptotics

$$\lim_{k \rightarrow \infty} \frac{\gamma_k}{k} = (e - 1).$$

In the next theorem, we not only extend these results to the last success setting, where $\theta > 0$, but also improve the roots' upper bound.

Theorem 8. *For $k \geq 1$, the roots (γ_k) of the fundamental equation (14) satisfy*

$$(e^{1/\theta} - 1)(k - 1) < \gamma_k < (e^{1/\theta} - 1)k + 1,$$

and

$$\lim_{k \rightarrow \infty} \frac{\gamma_k}{k} = (e^{1/\theta} - 1).$$

Before proving this theorem, we need an auxiliary result to assess the shape and properties of the following functions

$$\begin{cases} \varphi_0(n, k) & := \frac{(k)_n}{(k + \theta)_n}, \quad \varphi_0(0) = 1, \\ \varphi(n, k) & := \psi(n + k) - \psi(k), \quad \psi(0) = 0. \end{cases}$$

where $\theta > 0$, $k > 0$ are fixed and ψ is the log-derivative of the Gamma function.

Proposition 9. *For a fixed integer $k \geq 1$ and a fixed positive real number θ , it holds that*

- (i) $\varphi_0(\cdot, k)$ is log-convex and decreasing,
- (ii) $\varphi(\cdot, k)$ is concave and increasing.

Proof. Fix $k \geq 1$ and recall that if a sequence $\{\varphi_0(n, k)\}_{n \geq 1}$ is log-convex, then it is convex. For the first part of assertion (i), check that $\varphi_0^2(n, k) \leq \varphi_0(n + 1, k)\varphi_0(n - 1, k)$ holds. For fixed $\theta > 0$, it is easy to verify that

$$\begin{aligned} \frac{\varphi_0(n, k)}{\varphi_0(n + 1, k)} - \frac{\varphi_0(n - 1, k)}{\varphi_0(n, k)} &= \frac{(k)_n/(k + \theta)_n}{(k)_{n+1}/(k + \theta)_{n+1}} - \frac{(k)_{n-1}/(k + \theta)_{n-1}}{(k)_n/(k + \theta)_n} \\ &= \frac{\theta}{k + n} - \frac{\theta}{k + n - 1} < 0. \end{aligned}$$

For the monotonicity of φ_0 in the first parameter, note that

$$\begin{aligned} \frac{\varphi_0(n, k)}{\varphi_0(n + 1, k)} &= \frac{(k)_n/(k + \theta)_n}{(k)_{n+1}/(k + \theta)_{n+1}} \\ &= \frac{k + \theta + n}{k + n} > 1. \end{aligned}$$

Moving to assertion (ii). For the first part, it is sufficient to check that $\psi(\cdot)$ is concave. A straightforward calculation yields

$$\begin{aligned} 2\psi(n) - \psi(n - 1) - \psi(n + 1) &= \frac{1}{n - 1} - \frac{1}{n} \\ &= \frac{1}{n(n - 1)} > 0. \end{aligned}$$

The monotonicity of φ in the first parameter follows from observing that $\psi(n + 1) - \psi(n) = \frac{1}{n} > 1$. This completes the proof. \square

Proof of Theorem 8. Recall the derivative of the Pochhammer symbol of a positive real x , which is given by

$$(x)_n = (x)_n \{\psi(x + n) - \psi(x)\}.$$

Using this definition, the derivative in the first parameter of $M(k, k + \theta, x)$ becomes

$$\begin{aligned} D_a M(k, k + \theta; x) &= \sum_{n \geq 0} \frac{(k)_n}{(k + \theta)_n} \frac{x^n}{n!} \{\psi(k + n) - \psi(k)\} \\ &= \sum_{n \geq 0} \frac{x^n}{n!} \varphi_0(n) \varphi(n). \end{aligned}$$

We may rewrite equation (14) as $h_k(x) = 0$, where

$$\begin{aligned} h_k(x) &= \mathcal{S}_1(1 - x/\lambda, k) - \mathcal{S}_0(1 - x/\lambda, k) \\ &= \sum_{n \geq 0} \frac{x^n}{n!} e^{-x} \varphi_0(n, k) (\theta \varphi(n, k) - 1) \\ &= \mathbb{E}[\varphi_0(\xi_x, k) \{\theta \varphi(\xi_x - 1, k)\}]. \end{aligned} \tag{25}$$

where ξ_x is a Poisson random variable with mean x . By the first assertion of Proposition 9, $\varphi_0(\cdot, k)$ is decreasing. Hence, use (25) to observe that

$$\begin{aligned} h_k(x) &= \sum_{n \geq 0} \frac{x^n}{n!} e^{-x} \varphi_0(n, k) (\theta \varphi(n, k) - 1) \\ &\leq \varphi_0(0) \sum_{n \geq 0} \frac{x^n}{n!} e^{-x} (\theta \varphi(n, k) - 1) \\ &= \theta \mathbb{E}[\varphi(\xi_x, k)] - 1. \end{aligned}$$

But $\varphi(\cdot, k)$ is concave by assertion (ii) of the same proposition. Consequently, apply Jensen's inequality to obtain

$$\begin{aligned} h_k(x) &\leq \theta \varphi(\mathbb{E}[\xi_x], k) - 1 \\ &= \theta \varphi(x, k) - 1. \end{aligned} \tag{26}$$

We now proceed to establish bounds on this critical root of $h_k(x) = 0$. Let this unique root be denoted $\gamma_k > 0$. It follows by (26) that

$$\begin{aligned} 0 &= h_k(\gamma_k) \\ &\leq \theta \varphi(\gamma_k, k) - 1. \end{aligned}$$

Therefore,

$$\varphi(\gamma_k, k) \geq \frac{1}{\theta}. \tag{27}$$

Next, let m be a positive integer such that

$$\varphi(m, k) \geq \frac{1}{\theta} > \varphi(m-1, k). \tag{28}$$

By the Archimedean property of the reals and the monotonicity of φ , we can infer that

$$m-1 < \gamma_k \leq m.$$

It is easy to see by using (27) and (28) together with the monotonicity of φ that

$$\psi(k+m-1) - \psi(k) < \frac{1}{\theta} \leq \psi(\gamma_k+k) - \psi(k) \leq \psi(k+m) - \psi(k). \tag{29}$$

Moreover, by summing the following inequalities

$$\log(j+1) - \log(j) = \int_j^{j+1} \frac{dx}{x} < \frac{1}{j} < \int_{j-1}^j \frac{dx}{x} = \log(j) - \log(j-1),$$

from $j = k$ to $j = k + m - 1$, we obtain the classical inequality

$$\log\left(\frac{k+m}{k}\right) < \psi(k+m) - \psi(k) < \log\left(\frac{k+m-1}{k-1}\right). \quad (30)$$

Finally, combine (29) and (30) together with the monotonicity of the log function to form the inequality

$$\log\left(\frac{k+\gamma_k-1}{k}\right) < \frac{1}{\theta} < \log\left(\frac{k+\gamma_k-1}{k-1}\right).$$

This inequality ultimately yields the desired bounds

$$(e^{1/\theta} - 1)(k-1) < \gamma_k < (e^{1/\theta} - 1)k + 1. \quad (31)$$

For the second assertion of the Theorem, dividing each side of (31) by k and sending $k \rightarrow \infty$ completes the proof. \square

Appendix

Proof of part (ii) of Theorem 4. The logarithmic derivative in parameter c , where a_1, a_2 are constants satisfying $c > a_1 > 0$ and $c > a_2 > 0$ respectively is given by

$$D_c \log\left(\frac{M(a_1, c, x)}{M(a_1, c, x)}\right) = D_c \log \frac{\Gamma(c-a_2)}{\Gamma(c-a_1)} + \log D_c \frac{\phi(a, c_1, x)}{\phi(a, c_2, x)}. \quad (32)$$

By the elementary Lemma 2, it is sufficient to show that the signs of the two derivatives of quotients match. First, the sign of the derivative of a quotient of two Gamma functions coincides with $a_1 - a_2$ based on Lemma 3.

Next, we consider the second logarithmic derivative involving a quotient of ϕ 's. Use Lemma 2 once more to switch back to the derivative of the quotient $\phi(a_1, c, x)/\phi(a_2, c, x)$. Given that the denominator in the expression of the derivative of a quotient is positive, we restrict our investigation to the numerator, which we denote ϕ_c .

In much the same way as in the proof of part (i), consider the Eulerian integral representation (10) to express ϕ_c as a double integral, where $c > a_1 > 0$ and $c > a_2 > 0$ respectively

$$\begin{aligned} \phi_c &:= - \int_0^1 e^{xu}(1-u)^{c-2} \left(\frac{u}{1-u}\right)^{a_1-1} \log(1-u) du \int_0^1 e^{xv}(1-v)^{c-2} \left(\frac{v}{1-v}\right)^{a_2-1} dv \\ &\quad + \int_0^1 e^{xu}(1-u)^{c-2} \left(\frac{u}{1-u}\right)^{a_1-1} du \int_0^1 e^{xv}(1-v)^{c-2} \left(\frac{v}{1-v}\right)^{a_2-1} \log(1-v) dv \\ &= \iint_{[0,1] \times [0,1]} e^{x(u+v)} [(1-u)(1-v)]^{c-2} \left(\frac{uv}{(1-u)(1-v)}\right)^{a_1-1} \left(\frac{v}{1-v}\right)^{a_2-a_1} \log(1-v) dudv \\ &\quad - \iint_{[0,1] \times [0,1]} e^{x(u+v)} [(1-u)(1-v)]^{c-2} \left(\frac{uv}{(1-u)(1-v)}\right)^{a_1-1} \left(\frac{v}{1-v}\right)^{a_2-a_1} \log(1-u) dudv \end{aligned}$$

$$\begin{aligned}
&= \iint_{[0,1] \times [0,1]} \left\{ e^{x(u+v)} [(1-u)(1-v)]^{c-2} \left(\frac{uv}{(1-u)(1-v)} \right)^{a_1-1} \right\} \left(\frac{v}{1-v} \right)^{a_2-a_1} \log \left(\frac{1-v}{1-u} \right) dudv \\
&= \iint_{0 < v < u < 1} \{ \dots \} \left(\frac{v}{1-v} \right)^{a_2-a_1} \log \left(\frac{1-v}{1-u} \right) dudv + \iint_{0 < u < v < 1} \{ \dots \} \left(\frac{v}{1-v} \right)^{a_2-a_1} \log \left(\frac{1-v}{1-u} \right) dudv \\
&= \iint_{0 < v < u < 1} \{ \dots \} \left(\frac{v}{1-v} \right)^{a_2-a_1} \log \left(\frac{1-v}{1-u} \right) dudv + \iint_{0 < v < u < 1} \{ \dots \} \left(\frac{u}{1-u} \right)^{a_2-a_1} \log \left(\frac{1-u}{1-v} \right) dudv \\
&= \iint_{0 < v < u < 1} \{ \dots \} \left\{ \left(\frac{v}{1-v} \right)^{a_2-a_1} - \left(\frac{u}{1-u} \right)^{a_2-a_1} \right\} \log \left(\frac{1-v}{1-u} \right) dudv.
\end{aligned}$$

Mirroring the case of the derivative in parameter a , the term $\log \left(\frac{1-v}{1-u} \right)$ is positive in the domain of integration, where $u > v$. It follows that the sign of the last integrand coincides with the sign of

$$\left(\frac{v}{1-v} \right)^{a_2-a_1} - \left(\frac{u}{1-u} \right)^{a_2-a_1},$$

whose sign precisely matches that of $c_1 - c_2$. By the second part of Lemma 2, this concludes the proof as the sign of both terms in the right-hand side of (32) matches $c_1 - c_2$. \square

References

- [1] Ano, K. and Ando, M. (2000). A note on Bruss' stopping problem with random availability. In *IMS Lecture Notes - Monograph Series*, **35**, 71–82.
- [2] Berezovsky, B.A. and Gnedin, A. V. *The best choice problem* (in Russian). Nauka, 1984.
- [3] Browne, S. and Bunge, J. (1995). Random record processes and state dependent thinning. *Stochastic Process. Appl.* **55** 131–142.
- [4] Bruss, F.T. (1984) A Unified Approach to a Class of Best Choice Problems with an Unknown Number of Options. *Ann. Probab.* **12**, 882–889.
- [5] Bruss, F. T. (1987) On an optimal selection problem of Cowan and Zabczyk. *J. Appl. Probab.* **24**, 918–928.
- [6] Bruss, F. T. (2000) Sum the odds to one and stop. *Ann. Probab.* **28**, 1384–1391.
- [7] Bruss, F. T. and Paindaveine, D. (2000). Selecting a Sequence of Last Successes in Independent Trials. *J. Appl. Probab.* **37**, 389–399.
- [8] Chow, Y.S., Robbins, H. and Siegmund, D. *Great Expectations: The Theory of Optimal Stopping*, Houghton Mifflin Co., 1971.
- [9] Ciesielski, Z. and Zabczyk, J. (1979). A note on a selection problem. *Banach Center Publ.* **5** 47–51.
- [10] Cowan, R. and Zabczyk, J. (1978) An optimal selection problem associated with the Poisson process. *Theory Probab. Appl.* **23**, 584–592.
- [11] Crane, H. (2016) The ubiquitous Ewens sampling formula. *Statist. Sci.* **31**, 1–19.

- [12] Derbazi, Z. (2024). Sum the Probabilities to m and Stop. [arXiv:2406.07283](https://arxiv.org/abs/2406.07283).
- [13] Ferguson, T.S. *Optimal stopping and applications*, <https://www.math.ucla.edu/~tom/Stopping/Contents.html>, ebook, UCLA, 2006.
- [14] Gaver, D. P. (1976) Random record models. *J. Appl. Probab.* **13**, 538–547.
- [15] Gnedin, A. and Derbazi, Z. (2022), The Last-Success Problem with Random Observation Times, [arXiv:2207.05156](https://arxiv.org/abs/2207.05156)
- [16] Gnedin, A. and Derbazi, Z. (2022). Trapping the ultimate success, *Mathematics* **10**, 158.
- [17] Hofmann, G. *A Family of General Record Models*, Doctoral dissertation, The Ohio State University, 1997
- [18] Kallenberg, O. *Random Measures: Theory and Applications*. New York, Springer, 2017.
- [19] Kurushima, A. and Ano, K. (2003). A Poisson arrival selection problem for Gamma prior intensity with natural number parameter. *Sci. Math. Japon.* **57**, 217–231.
- [20] Nevzorov, V. *Records: Mathematical Theory* (Lecture 25), Translation of Mathematical Monographs **194**, AMS, 2001.
- [21] Orsingher, E. and Battaglia, F. (1979). Record values in Poisson and randomized Poisson models, *Publ. Inst. Stat. Univ. Paris* **24**, fasc. 3-4, 69–78.
- [22] Schwalko, A. and Szajowski, K. (2003). On Bruss' stopping problem with general gain function. *Game Theory Appl.* **9**, 161–171.
- [23] Szajowski, K. (2007). A game version of the Cowan-Zabczyk-Bruss' problem. *Statist. Probab. Lett.* **77** 1683–1689.
- [24] Szajowski, K. J. (2015). An apartment problem. *Math. Appl.* **42**, 193–206.
- [25] Yasuda, M. (1984). Asymptotic results for the best-choice problem with a random number of objects. *J. Appl. Probab.* **21**, 521–536.