Deducibility in the full Lambek calculus with weakening is HACк-complete

Vitor Greati^a, Revantha Ramanayake^{1a,b}

^aBernoulli Institute, University of Groningen, Nijenborgh 4, NL-9747 AG Groningen, Netherlands ^bCogniGron, University of Groningen, Nijenborgh 4, NL-9747 AG Groningen, Netherlands

Abstract

We show that the problem of deciding the consequence relation (deducibility) for the full Lambek calculus with weakening $(\mathbf{FL}_{\mathbf{w}})$ is complete for the class HACK of hyper-Ackermannian problems (i.e., $\mathbf{F}_{\omega^{\omega}}$ in the ordinal-indexed hierarchy of fast-growing complexity classes). Provability was already known to be PSPACE-complete. We prove that deducibility is HACK-complete even for the multiplicative fragment. Lower bounds are obtained via a novel reduction from reachability in lossy channel systems. Upper bounds are obtained via structural proof theory (forward proof search over sequent calculi) and well-quasi-order theory (length theorems for Higman's ordering on words over a finite alphabet).

Keywords: substructural logics, hyper-Ackermannian problems, Lambek calculus, Higman's Lemma, well-quasi-order theory, sequent calculi

1. Introduction

Substructural logics are obtained from intuitionistic and classical logic by removing familiar structural properties like exchange (e), contraction (c) and weakening (w), and extending with further axioms and language connectives. A prominent example is the family of *linear logics* where a propositional substructural base is extended with modalities ("exponentials") for weakening and contraction. The absence of structural properties leads to highly expressive and complex logics, and this is evident even in the most fundamental of substructural logics (see Table 1), namely the extensions of the full Lambek calculus **FL** by subsets of $\{(e), (c), (w)\}$. A significant gap in this table persisted at **FL**_w since the deducibility problem (provability of a formula from a finite set of formulas) was, thus far, only known to be decidable, a result obtained by Blok and van Alten [1] via the finite embeddability property for **FL**_w-algebras, the algebraic semantics of **FL**_w. We fill this gap by establishing completeness with respect to the fast-growing hyper-Ackermannian complexity class HACK, placing **FL**_w as the hardest for deducibility among the decidable basic substructural logics (taking the crown from **FL**_{ec}, where deducibility has Ackermannian complexity). Let **FL**^{\Sigma}_w denote the restriction of **FL**_w to connectives in Σ .

¹The financial support of the CogniGron research center, the Ubbo Emmius Funds (University of Groningen), and FWF project P33548 is acknowledged.

Theorem 1 (Main theorem). For $\{\otimes, \backslash, 0, 1\} \subseteq \Sigma \subseteq \Sigma_{\mathbf{FL}}$, deducibility in $\mathbf{FL}_{\mathbf{w}}^{\Sigma}$ is HACKcomplete. In particular, $\mathbf{FL}_{\mathbf{w}}$ and its multiplicative fragment $\mathbf{L}_{\mathbf{w}}$ are HACK-complete.

Logic	Deducibility	Provability
$\begin{array}{c} FL\\ FL_{e}\\ FL_{ec}\\ FL_{ew}\\ FL_{c}\\ FL_{w}\\ L_{w} \end{array}$	Undecidable [2] Undecidable [4] ACK-complete [5] 2-EXPTIME-complete [6] TOWER-complete [7] Undecidable [8] Decidable [1]. Complexity: this paper Decidable [9]. Complexity: this paper	PSPACE-complete [3]PSPACE-complete [3]ACK-complete [5]2-EXPTIME-complete [6]PSPACE-complete [3]Undecidable [8]PSPACE-complete [3]PSPACE-complete [3]PSPACE [3]

Table 1: The computational status of the basic substructural logics.

The above statement references Schmitz's ordinal-indexed fast-growing complexity classes $\{\mathbf{F}_{\alpha}\}_{\alpha}$ [10]. Important members of this family are \mathbf{F}_{2} , problems solvable in time expressed by an elementary function (such as polynomial and exponential); \mathbf{F}_{3} , the first class of nonelementary problems, known as TOWER; \mathbf{F}_{ω} , the first class of non-primitive recursive problems, known as ACK (short for 'Ackermann'); and $\mathbf{F}_{\omega^{\omega}}$, the first class of non-multiply recursive problems. The complexity of the latter cannot be expressed by a function defined by multiple nested recursion [11, §10], and it is known as HACK ('hyper-Ackermann').

An overview of the lower bound argument. Inspired by Urquhart's [5] lower bound for \mathbf{FL}_{ec} where acceptance for expansive additive counter machines is reduced to deducibility, we reduce reachability for lossy channel systems (LCS) to deducibility in \mathbf{FL}_{w} .

An LCS is a computational model whose configurations are tuples (q, u_1, \ldots, u_n) where q is the state and u_k is the kth channel's contents i.e., some word over a finite alphabet M. Its instructions take the following form: *read* letter **a** from the front of channel k in state q_i , remove it and transit to state q_j ; write letter **a** to the back of channel k in state q_i and transit to state q_j ; perform a lossy step by deleting some letters from any channel (the state remains unchanged). Chambart and Schnoebelen [12] show that deciding reachability of a configuration v from a configuration u is HACK-complete.

We code the set I of instructions of an LCS \mathcal{L} as a finite set $\mathcal{T}(\mathcal{L})$ of sequents, and simulate read and write instructions by cuts with these sequents, and lossy steps by left weakening. For instance, reachability of $(q_2, \mathbf{a}, \mathbf{b})$ from $(q_1, \mathbf{aa}, \mathbf{b})$ in \mathcal{L} is coded as deducibility of the following sequent from $\mathcal{T}(\mathcal{L})$:

$$Q_1, \#_1^s, A, A, \#_1^e, \#_2^s, B, \#_2^e \Rightarrow Q_2 \otimes (\#_1^s \otimes (A \otimes (\#_1^e \otimes (\#_2^s \otimes (B \otimes \#_2^e))))))$$

Reading upwards from the endsequent, the following deduction in $\mathbf{FL}_{\mathbf{w}}$ simulates an instruction $(q_i, c_1, \mathbf{a}, ?, q_j) \in \mathbf{I}$; i.e., read **a** from the front of channel c_1 in state q_i , remove it, then transit to q_j .

sequent in
$$\mathcal{T}(\mathcal{L})$$
 coding instruction

$$\frac{Q_{j}, \#_{1}^{s}, B, C, \#_{1}^{e} \Rightarrow v}{Q_{j} \otimes \#_{1}^{s}} \xrightarrow{Q_{j} \otimes \#_{1}^{s}, B, C, \#_{1}^{e} \Rightarrow v} (L\otimes)$$

$$\frac{Q_{i}, \#_{1}^{s}, A, B, C, \#_{1}^{e} \Rightarrow v}{Q_{i}, \#_{1}^{s}, A, B, C, \#_{1}^{e} \Rightarrow v} (cut)$$

. . .

Every computation in the LCS is shown to correspond to a deduction of its reachability encoding. The converse direction is far more challenging, requiring a careful analysis of deductions in $\mathbf{FL}_{\mathbf{w}}$. The starting point is a cut-elimination style argument that establishes a normal form (of "standard deductions") where the left premise of every cut is an element of the theory. The computation is read off the standard deduction, starting at the endsequent.

The upper bound argument. We define a forward proof procedure extending the construction in Balasubramanian et al. [13]. The idea is to start from the set D_0 of initial sequent instances and form an increasing chain $D_0 \subset D_1 \subset \ldots$ of finite sets of deducible sequents (each step corresponds to limited weakening and then a single rule application), taking care that each D_i contains only minimal elements under the Higman ordering on words (omit words that can be obtained from another by inserting letters); the ordering is a well-quasiorder (wqo) [14] so the chain must be finite. By construction, the bad sequence underlying the chain is controlled (no element in D_{i+1} is much larger than some element in D_i), and the complexity is dominated by the maximum length of a controlled bad sequence. This is a function in HACK as shown by Schmitz and Schnoebelen [6]. The argument applies to infinitely-many structural rule extensions of \mathbf{FL}_{w} .

Contributions. We show that (1) sequent-deducibility in fragments of $\mathbf{FL}_{\mathbf{w}}$ containing \otimes is HACK-hard and that (2) deducibility for any \mathcal{N}_2 -analytic structural rule extension [15] of any fragment of $\mathbf{FL}_{\mathbf{w}}$ is in HACK. Our main theorem follows from (1) and (2). Algebraizability of $\mathbf{FL}_{\mathbf{w}}$ implies that (3) the word problem and the quasiequational theory of the variety of integral zero-bounded full Lambek algebras are HACK-complete.

2. Noncommutative substructural logics with weakening

Let Σ be a propositional signature and fix a countable set \mathcal{P} of propositional variables. Let $\operatorname{Fm}_{\Sigma}(\mathcal{P})$ denote the set of formulas over Σ generated by \mathcal{P} . We consider subsignatures of $\Sigma_{\mathbf{FL}} := \{0, 1, \top, \bot, \otimes, \wedge, \vee, /, \setminus\}$, where $0, 1, \top$ and \bot are constants and the other connectives are binary. A subsignature is specified as $\Sigma_{\mathbf{FL}}^{\star_1 \cdots \star_k} := \{\star_1, \ldots, \star_k\} \subseteq \Sigma_{\mathbf{FL}}$. For example, $\Sigma_{\mathbf{FL}}^{\otimes \setminus} = \{\otimes, \setminus\}$. Sequences (or lists) of formulas will be written without delimiters, with commas separating the formulas. We write $\bigotimes \varphi_1, \ldots, \varphi_m$ as abbreviation for $\varphi_1 \otimes (\varphi_2 \otimes (\ldots (\varphi_{m-1} \otimes \varphi_m) \ldots)))$, where $m \geq 1$ and $\varphi_1, \ldots, \varphi_m$ is a list of formulas. If the constant 1 is present in the language, we allow m = 0 (empty fusion), the result being 1.

A sequent over $\operatorname{Fm}_{\Sigma}(\mathcal{P})$ is an expression $\Gamma \Rightarrow \Pi$, where Γ is a finite sequence of formulas and Π is an empty sequence or a sequence with a single formula. Given a set \mathcal{T} of sequents, let $\operatorname{subf}(\mathcal{T})$ denote the set of subformulas of the formulas appearing in it. We assume familiarity with the sequent calculus (see e.g., [16, Sec. 2.1.3]): a sequent calculus \mathbf{S} over $\operatorname{Fm}_{\Sigma}(\mathcal{P})$ is a collection of schematic sequent rules. Instantiations of their schematic variables lead to rule instances. Derivations in \mathbf{S} are finite rooted trees labelled with sequents over $\operatorname{Fm}_{\Sigma}(\mathcal{P})$ such that a node and its children are the conclusion and premises of a rule instance. For a finite set \mathcal{T} of sequents, a \mathcal{T} -deduction in \mathbf{S} is a derivation in which every leaf is a rule with no premises (axiom or initial sequent) or a sequent in \mathcal{T} . A proof in \mathbf{S} is a \emptyset -deduction in \mathbf{S} . Any $\mathbf{S}' \subseteq \mathbf{S}$ is called a subcalculus of \mathbf{S} , and any $\mathbf{S}'' \supseteq \mathbf{S}$ is a rule extension of \mathbf{S} . Structural rule extensions are extensions by structural rules (rules without logical connectives).

Subcalculi of the sequent calculus $\mathbf{FL}_{\mathbf{w}}$ displayed in Fig. 1 (*full Lambek calculus with weakening*) are induced by a subsignature of $\Sigma_{\mathbf{FL}}$ as follows. Let $\mathbf{FL}_{\mathbf{w}}^{\star_1\cdots\star_k}$ be the sequent calculus consisting of rules that mention no connective (i.e., rule (id) and structural rules), or that mentions a connective in $\{\star_1, \ldots, \star_k\}$. For instance, $\mathbf{FL}_{\mathbf{w}}^{\otimes \backslash}$ is the calculus over $\mathrm{Fm}_{\Sigma_{\mathbf{FL}}^{\otimes \backslash}}(\mathcal{P})$ consisting of the axioms, all structural rules and the rules mentioning \otimes and \backslash .

A sequent calculus **S** over $\operatorname{Fm}_{\Sigma}(\mathcal{P})$ canonically determines two consequence relations, one over sequents and another over formulas. The sequent-deduction relation $\vdash_{\mathbf{S}\Rightarrow}$ is such that, for all finite sets $\mathcal{T} \cup \{s\}$ of sequents over $\operatorname{Fm}_{\Sigma}(\mathcal{P}), \mathcal{T} \vdash_{\mathbf{S}\Rightarrow} s$ iff there is a \mathcal{T} -deduction of sin **S**. The formula-deduction relation $\vdash_{\mathbf{S}}$ is such that, for all finite sets $G \cup \{\varphi\}$ of formulas over $\operatorname{Fm}_{\Sigma}(\mathcal{P}), G \vdash_{\mathbf{S}} \varphi$ iff there is a $\{\Rightarrow \psi \mid \psi \in G\}$ -deduction of $\Rightarrow \varphi$ in **S**. Given a sequent calculus **S**, the problem of deciding $\vdash_{\mathbf{S}\Rightarrow}$ is called sequent-deducibility and the problem of deciding $\vdash_{\mathbf{S}}$ is called formula-deducibility or simply deducibility. We know that $\vdash_{\mathbf{FL}_{\mathbf{w}}}$ is the consequence relation of the full Lambek logic with weakening (see [16, 2.1.4] for the Hilbertcalculus presentation and residuated lattices semantics of these logics). The reader is also referred to [16, 2.1.4] for the notion of axiomatic extensions of $\vdash_{\mathbf{FL}_{\mathbf{w}}}$.

The following lemma says that $\vdash_{\mathbf{S}\Rightarrow}$ reduces to $\vdash_{\mathbf{S}}$ when \mathbf{S} is a subcalculus of $\mathbf{FL}_{\mathbf{w}}$ containing the rules for $\otimes, \backslash, 0$ and 1. For that, consider the translation $(\cdot)^{\rightarrow}$ of sequents over $\operatorname{Fm}_{\Sigma}(\mathcal{P})$ into formulas over $\operatorname{Fm}_{\Sigma\cup\{0,1,\otimes,\backslash\}}(\mathcal{P})$ defined $(\Gamma\Rightarrow\Pi)^{\rightarrow}:=\bigotimes\Gamma\backslash\Pi^*$, where Π^* is φ if $\Pi=\varphi$ and 0 if Π is empty.

Lemma 2. Let $\Sigma \supseteq \{\otimes, \setminus, 0, 1\}$ and $\mathbf{S} := \mathbf{FL}_{\mathbf{w}}^{\Sigma}$. For all finite sets of sequents \mathcal{T} and sequents s over the language of \mathbf{S} , $\mathcal{T} \vdash_{\mathbf{S}\Rightarrow} s$ iff $\{t^{\rightarrow} \mid t \in \mathcal{T}\} \vdash_{\mathbf{S}} s^{\rightarrow}$.

Proof. From left to right, let δ be a deduction witnessing $\mathcal{T} \vdash_{\mathbf{S}\Rightarrow} s$. We show how to convert δ into a witness of $\{t^{\rightarrow} \mid t \in \mathcal{T}\} \vdash_{\mathbf{S}} s^{\rightarrow}$ by structural induction. In the base case, s is either the result of an axiomatic rule or an element of \mathcal{T} . To translate $t \in \mathcal{T}$ into t^{\rightarrow} , repeatedly use the rules $(L\otimes)$ and $(R\setminus)$, and (L1) and (R0) if required. In the inductive step, assume $\Gamma \Rightarrow \Pi$ was derived by an application of a k-ary rule \mathbf{r} , whose premises are the sequents $\Gamma_1 \Rightarrow \Pi_1, \ldots, \Gamma_k \Rightarrow \Pi_k$. Note first that the rule $\underline{\Rightarrow \bigotimes \Gamma \setminus \Pi}_{\Gamma \Rightarrow \Pi}$ (FS) is derivable in presence of cut. By IH, the sequents $\Rightarrow \bigotimes \Gamma_1 \setminus \Pi_1^*, \ldots, \Rightarrow \bigotimes \Gamma_k \setminus \Pi_k^*$ are provable from $\mathcal{T}^{\rightarrow}$. Apply to these premises the derived rule (FS) thus obtaining the premises $\Gamma_1 \Rightarrow \Pi_1, \ldots, \Gamma_k \Rightarrow \Pi_k$ (maybe cuts with $\Rightarrow 1$ will be necessary to recover empty antecedents). Use cuts with $0 \Rightarrow$ whenever necessary to obtain from those the premises $\Gamma_1 \Rightarrow \Pi_1, \ldots, \Gamma_k \Rightarrow \Pi_k$. Apply the same rule \mathbf{r} to the latter, obtaining $\Gamma \Rightarrow \Pi$, from which $\Rightarrow \bigotimes \Gamma \setminus \Pi^*$ is easily derivable (if Γ is empty, use (L1)), and we are done.

From right to left, first transform all the leaves with translated sequents from \mathcal{T} into trees with leaves being either instances of axioms or the sequents themselves. This is easy

to do using rules (R\), (L \otimes), (L1) and (R0). Then it is enough to transform the root s^{\rightarrow} into s using (FS) as we did in the converse direction.

$$\begin{array}{c|c} \hline p \Rightarrow p & (\mathrm{id}) & \hline \Gamma, \bot, \Delta \Rightarrow \Pi & (\mathrm{L}\bot) & \hline \Gamma \Rightarrow \top & (\mathrm{R}\top) & \hline 0 \Rightarrow & (\mathrm{L0}) & \hline \Rightarrow 1 & (\mathrm{R1}) \\ \hline & \frac{\Gamma, \Delta \Rightarrow \Pi}{\Gamma, \varphi, \Delta \Rightarrow \Pi} & (\mathrm{lw}) & \frac{\Gamma \Rightarrow}{\Gamma \Rightarrow \varphi} & (\mathrm{rw}) & \frac{\Theta \Rightarrow \varphi}{\Gamma, \Theta, \Delta \Rightarrow \Pi} & (\mathrm{cut}) \\ \hline & \frac{\Gamma \Rightarrow}{\Gamma, \Theta, \Delta \Rightarrow \Pi} & (\mathrm{lw}) & \frac{\Gamma, \Delta \Rightarrow \Pi}{\Gamma, 1, \Delta \Rightarrow \Pi} & (\mathrm{L1}) & \frac{\Gamma, \varphi, \psi, \Delta \Rightarrow \Pi}{\Gamma, \varphi \otimes \psi, \Delta \Rightarrow \Pi} & (\mathrm{Le}) \\ \hline & \frac{\Gamma \Rightarrow \varphi}{\Gamma, \Delta \Rightarrow \varphi} & (\mathrm{Re}) & \frac{\Gamma, \varphi, \Delta \Rightarrow \Pi}{\Gamma, \varphi \lor \psi, \Delta \Rightarrow \Pi} & (\mathrm{Le}) & \frac{\Gamma \Rightarrow \varphi_i}{\Gamma \Rightarrow \varphi_1 \lor \varphi_2} & (\mathrm{Re}) \\ \hline & \frac{\Gamma, \varphi_i, \Delta \Rightarrow \Pi}{\Gamma, \varphi_1 \land \varphi_2, \Delta \Rightarrow \Pi} & (\mathrm{Le}) & \frac{\Gamma \Rightarrow \varphi}{\Gamma \Rightarrow \varphi \land \psi} & (\mathrm{Re}) & \frac{\Gamma \Rightarrow \varphi}{\Gamma \Rightarrow \varphi \land \psi} & (\mathrm{Re}) & \frac{\Gamma \Rightarrow \varphi}{\Delta, \psi, \Theta \Rightarrow \Pi} & (\mathrm{Le}) & \frac{\Gamma \Rightarrow \varphi}{\Delta, \psi/\varphi, \Gamma, \Theta \Rightarrow \Pi} & (\mathrm{Le}) \\ \hline & \frac{\Gamma, \varphi \Rightarrow \psi}{\Gamma \Rightarrow \psi/\varphi} & (\mathrm{Re}) & \frac{\Gamma \Rightarrow \varphi}{\Delta, \Gamma, \varphi \lor \psi, \Theta \Rightarrow \Pi} & (\mathrm{Le}) & \frac{\varphi, \Gamma \Rightarrow \psi}{\Gamma \Rightarrow \varphi \lor \psi} & (\mathrm{Re}) \\ \hline \end{array}$$

Figure 1: The sequent calculus $\mathbf{FL}_{\mathbf{w}}$.

Every \varnothing -deduction (i.e., every proof) in $\mathbf{FL}_{\mathbf{w}}^{\Sigma}$ can be converted to a proof without cuts (by cut elimination [16, 4.1.1]), so these calculi satisfy the *subformula property* (every formula that occurs in the proof is a subformula of the endsequent). However, not all cuts can be eliminated from a \mathcal{T} -deduction. Still, for suitable \mathcal{T} identified below, the applications of cut can be restricted to obtain a generalized subformula property.

Definition 3. A sequent is said to be regular if its antecedent contains only propositional variables and the succedent is not empty. A regular theory is a finite collection of regular sequents. For a regular theory \mathcal{T} , a standard cut (over \mathcal{T}) has as left premise a leaf that is a sequent from \mathcal{T} . A standard deduction (from \mathcal{T}) is one in which all cuts are standard. We write $\mathcal{T} \vdash_{\mathbf{FL}_{\mathbf{w}}^{\Sigma} \Rightarrow}^{\mathsf{std}} \Gamma \Rightarrow \varphi$ to denote that $\Gamma \Rightarrow \varphi$ has a standard deduction from \mathcal{T} in $\mathbf{FL}_{\mathbf{w}}^{\Sigma}$.

Lemma 4. Let $\Sigma \subseteq \Sigma_{\mathbf{FL}}$. For a regular theory \mathcal{T} and sequent $s, \mathcal{T} \vdash_{\mathbf{FL}_{\mathbf{w}}^{\Sigma} \Rightarrow} s$ iff $\mathcal{T} \vdash_{\mathbf{FL}_{\mathbf{w}}^{\Sigma} \Rightarrow}^{\mathsf{std}} s$. Moreover, every formula in a standard deduction in $\mathbf{FL}_{\mathbf{w}}^{\Sigma}$ is in $\mathsf{subf}(\mathcal{T} \cup \{s\})$.

Proof. We establish the claim for $\Sigma = \Sigma_{\mathbf{FL}}$. The reader will observe that removing rules means less cases to be checked, and thus the argument here applies to any $\Sigma \subseteq \Sigma_{\mathbf{FL}}$. Let us call the instantiation of the schematic-variable φ in an instance of the (cut) rule (cf. Fig. 1) as the *cut formula* of that instantiation.

For first assertion, the argument is based on the usual cut-elimination procedure for $\mathbf{FL}_{\mathbf{w}}$ [16, 4.1.1]; the difference is that we now have to deal with cuts with a premise in \mathcal{T} . We repeatedly eliminate topmost non-standard cuts, by primary induction on the grade of the cut (the length of the cut formula), and secondary induction on the cut-height (the sum of the number of sequents appearing in the proofs of the premises of the cut).

Consider a topmost non-standard cut with cut-height 2 (note that < 2 is not possible). The premises of the cut are instances of axiomatic rules or elements from \mathcal{T} . If no element from \mathcal{T} appears as a premise, the cut is eliminated in the usual way as for $\mathbf{FL}_{\mathbf{w}}$. If the left premise is from \mathcal{T} , nothing needs to be done since we admit standard cuts. If the right premise is from \mathcal{T} , the left premise can only be an instance either of (id) or (L \perp); i.e., the cut has one of these forms:

$$\frac{\overline{p \Rightarrow p} (\mathrm{id})}{\Gamma_1, p, \Gamma_2 \Rightarrow \varphi} \frac{\mathcal{T}}{(\mathrm{cut})} \quad \frac{\overline{\Delta_1, \bot, \Delta_2 \Rightarrow p} (\bot \mathrm{L})}{\Gamma_1, \Delta_1, \bot, \Delta_2, \Gamma_2 \Rightarrow \varphi} \frac{\mathcal{T}}{(\mathrm{cut})}$$

The cut above left is clearly eliminable since the second premise matches the conclusion; the cut above right is replaced by an instance of $(L\perp)$.

Inductive step. Assume the cut has cut-height ≥ 3 . If neither of the premises of cut are from \mathcal{T} , the usual cut-reduction steps apply [16, 4.1.1]. Else, if the topmost cut is standard, nothing needs to be done since we admit such cuts. The remaining case is that the right premise is from \mathcal{T} and the left premise is from some other rule. It follows that the topmost non-standard cut occurs as follows.

$$\frac{\stackrel{o_1}{\Gamma \Rightarrow p} \overline{\Delta_1, p, \Delta_2 \Rightarrow \Pi} \mathcal{T}}{\Delta_1, \Gamma, \Delta_2 \Rightarrow \Pi} (\text{cut})$$

Since the cut-height is ≥ 3 , we have that the left premise is not an instance of an axiomatic rule. Consider the last rule in δ_1 . We transform each case so the non-standard cut is replaced with a cut that is smaller under the induction measure. We indicate the transformations with \rightsquigarrow . Since the cut formula is a propositional variable (this is forced by the right premise being from the regular theory \mathcal{T}), the last rule in δ_1 cannot be a right-introduction rule. We provide some representative cases.

• The left premise of the cut under consideration is the conclusion of a standard cut:

• $(L\otimes)$

$$\frac{\delta_{2}}{\Gamma_{1},\varphi_{1},\varphi_{2},\Gamma_{2}\Rightarrow p} (L\otimes) \xrightarrow{\Delta_{1},p,\Delta_{2}\Rightarrow\Pi} \mathcal{T} \\
\frac{\Gamma_{1},\varphi_{1}\otimes\varphi_{2},\Gamma_{2}\Rightarrow p}{\Delta_{1},\Gamma_{1},\varphi_{1}\otimes\varphi_{2},\Gamma_{2},\Delta_{2}\Rightarrow\Pi} (cut)$$

$$\frac{\delta_{2}}{\frac{\Gamma_{1},\varphi_{1},\varphi_{2},\Gamma_{2}\Rightarrow p}{\Delta_{1},p,\Delta_{2}\Rightarrow\Pi} \mathcal{T} \\
\frac{\delta_{2}}{\frac{\Gamma_{1},\varphi_{1},\varphi_{2},\Gamma_{2}\Rightarrow p}{\Delta_{1},p,\Delta_{2}\Rightarrow\Pi} (cut)}$$

The second assertion follows by structural induction on standard deductions. The base case is obvious. Inductive step. The claim is immediate for a logical rule, (lw), and (rw), by the induction hypothesis applied to the deduction concluding the premise, and since every formula in the premise must be a subformula of a formula in the conclusion. It remains to check the case of a standard cut.

$$\frac{\overline{\Delta \Rightarrow \psi} \mathcal{T} \qquad \stackrel{\delta_1}{\Gamma_1, \psi, \Gamma_2 \Rightarrow \varphi}}{\Gamma_1, \Delta, \Gamma_2 \Rightarrow \varphi} \operatorname{cut}$$

The claim follows since ψ is a formula present in one of the sequents in \mathcal{T} .

The next lemma is a direct consequence of Lem. 4 and tells us that sequent-deducibility in $\mathbf{FL}_{\mathbf{w}}^{\Sigma'}$ reduces to the same problem over an expanded signature $\Sigma_{\mathbf{FL}} \supseteq \Sigma \supseteq \Sigma'$. In other words, $\vdash_{\mathbf{FL}_{\mathbf{w}}^{\Sigma}}$ is a *conservative expansion* of $\vdash_{\mathbf{FL}_{\mathbf{w}}^{\Sigma'}}$.

Lemma 5. Let $\Sigma' \subseteq \Sigma \subseteq \Sigma_{\mathbf{FL}}$. If $\mathcal{T} \cup \{s\}$ are sequents over $\operatorname{Fm}_{\Sigma'}(\mathcal{P})$, then $\mathcal{T} \vdash_{\mathbf{FL}_{\mathbf{w}}^{\Sigma'} \Rightarrow} s$ iff $\mathcal{T} \vdash_{\mathbf{FL}_{\mathbf{w}}^{\Sigma} \Rightarrow} s$.

Proof. The nontrivial direction is right-to-left, so suppose that $\mathcal{T} \vdash_{\mathbf{FL}_{\mathbf{w}}^{\Sigma} \Rightarrow} s$ is witnessed by a deduction δ . Obtain a standard deduction δ' from δ by Lem. 4. The new proof will have sequents over $\operatorname{Fm}_{\Sigma'}(\mathcal{P})$ only (by the subformula property proved also in Lem. 4), and thus must be a deduction in $\operatorname{FL}_{\mathbf{w}}^{\Sigma'}$.

3. Hyper-Ackermannian lower bounds

We obtain hyper-Ackermannian lower bounds for the sequent-deducibility problem in Σ -fragments of $\mathbf{FL}_{\mathbf{w}}$ where $\otimes \in \Sigma$, via a reduction from the reachability problem in lossy channel systems. The result extends to the problem of deducibility if Σ also contains \setminus (or /), 0 and 1, in view of Lem. 2. We begin by introducing lossy channel systems and their reachability problem, and then show how to encode it into sequent-deducibility in $\mathbf{FL}_{\mathbf{w}}^{\otimes}$.

3.1. Lossy channel systems

A channel system is a computational model with unbounded FIFO channels (queues), defined as follows.

Definition 6. A channel system (CS) is a structure $\mathcal{L} := \langle Q, C, M, I \rangle$, where

- 1. $\mathbf{Q} := \{q_1, \ldots, q_m\}$ is a finite set of states;
- 2. $C := \{c_1, \ldots, c_n\}$ is a finite set of channels;
- 3. $M := \{a_1, \ldots, a_k\}$ is a finite message alphabet; and
- 4. $I \subseteq Q \times C \times M \times \{!, ?\} \times Q$ is a finite set of instructions.

We denote by $\text{Conf}(\mathcal{L})$ the set $\mathbb{Q} \times (\mathbb{M}^*)^n$ of configurations of \mathcal{L} , where \mathbb{M}^* is the set of all finite sequences of elements in \mathbb{M} .

Definition 7. The perfect steps of a $CS\mathcal{L}$ are given by a binary relation $\rightarrow_{\mathcal{L} perf} \subseteq Conf(\mathcal{L}) \times Conf(\mathcal{L})$ such that $(q_i, u_1, \ldots, u_r, \ldots, u_n) \rightarrow_{\mathcal{L} perf} (q_j, u_1, \ldots, v_r, \ldots, u_n)$ if either

- 1. $u_r = \mathbf{a}_p v_r$ and $(q_i, c_r, \mathbf{a}_p, ?, q_j) \in \mathbf{I}$ (read and remove \mathbf{a}_p from the front of channel c_r and change state from q_i to q_j)
- 2. $v_r = u_r \mathbf{a}_p$ and $(q_i, c_r, \mathbf{a}_p, !, q_j) \in \mathbf{I}$ (write \mathbf{a}_p at the back of channel c_r and change state from q_i to q_j).

Let $\rightarrow^{?}_{\mathcal{L} \text{ perf}}$ and $\rightarrow^{!}_{\mathcal{L} \text{ perf}}$ denote the perfect steps given by items 1 and 2 respectively. Also let $\rightarrow^{*}_{\mathcal{L} \text{ perf}}$ denote the reflexive and transitive closure of $\rightarrow_{\mathcal{L} \text{ perf}}$.

Clearly, $u \to_{\mathcal{L}perf}^* v$ iff there is a finite sequence w_1, \ldots, w_η $(\eta \ge 1)$ of configurations such that $u = w_1, v = w_\eta$ and $w_i \to_{\mathcal{L}perf} w_{i+1}$ for each $i \ (1 \le i < \eta)$. This sequence is called a *perfect computation in* \mathcal{L} of length η and it witnesses that u reaches v in \mathcal{L} .

Reachability (as well as many other verification problems for CSs) is undecidable [17], but adding lossy behaviour makes it decidable. This behaviour is introduced in the operational semantics of a CS as follows.

Definition 8. The lossy semantics of a CS \mathcal{L} is given by a relation $\rightarrow_{\mathcal{L} \text{lossy}} \subseteq \text{Conf}(\mathcal{L}) \times \text{Conf}(\mathcal{L})$ that extends $\rightarrow_{\mathcal{L} \text{perf}}$ with all transitions (called lossy steps) of the form $(q_i, u_1, \ldots, u_r, \ldots, u_n) \rightarrow_{\mathcal{L} \text{lossy}}^l (q_i, u_1, \ldots, v_r, \ldots, u_n)$ such that $u_r = w_1 \mathbf{a}_p w_2$ and $v_r = w_1 w_2$. Let $\rightarrow_{\mathcal{L} \text{lossy}}^*$ be the reflexive and transitive closure of $\rightarrow_{\mathcal{L} \text{lossy}}$. A channel system with a lossy semantics is called a lossy channel system (LCS).

Intuitively, in an LCS the lossy steps permit any channel to lose a message (a symbol) from any position at any moment without changing the state. No instruction in the machine is required for this. The relation $\rightarrow_{\mathcal{L} \text{ lossy}}^*$ induces a notion of *lossy computation in* \mathcal{L} and reachability is defined analogously to perfect computations with $\rightarrow_{\mathcal{L} \text{ perf}}^*$. Since a step can be either read, write, or lossy, we use $\rightarrow_{\mathcal{L} \text{ lossy}}^?$, $\rightarrow_{\mathcal{L} \text{ lossy}}^!$ or $\rightarrow_{\mathcal{L} \text{ lossy}}^l$ to indicate precisely which step was performed in a lossy computation.

Definition 9 (Reachability problem in LCSs). Given a LCS \mathcal{L} and configurations $u, v \in Conf(\mathcal{L})$, does $u \to_{\mathcal{L} \text{ lossy }}^* v$?

Remark 10. The relation $\rightarrow_{\mathcal{L} \text{ lossy}}^*$ has an equivalent definition in the literature [18, Sec. 2] as $u \rightsquigarrow_{\mathcal{L}}^* v$ if, and only if, there are $u', v' \in \text{Conf}(\mathcal{L})$ such that $u \rightarrow_{\mathcal{L} \text{ lossy}}^{l*} u' \rightarrow_{\mathcal{L} \text{ perf}} v' \rightarrow_{\mathcal{L} \text{ lossy}}^{l*} v$ (the * denotes the reflexive transitive closure). That is, messages may be lost before and after performing a perfect step. It can be shown that $\rightarrow_{\mathcal{L} \text{ lossy}}^* = \rightarrow_{\mathcal{L} \text{ lossy}}^{l*} \circ \rightarrow_{\mathcal{L} \text{ perf}} \circ \rightarrow_{\mathcal{L} \text{ lossy}}^{l*} = \sim_{\mathcal{L}}^*$. The latter relation is shown [12, App. A] to have no impact on the complexity of reachability compared to the "write-lossy" mechanism employed in the proof of Thm. 11 [12] stated below.

Theorem 11 ([12, Thm. 5.5, Obs. 6.1, App. A]). Reachability in LCSs is HACK-complete.

3.2. Encoding reachability in LCSs into sequent-deducibility

We code each instruction of a lossy channel system into a sequent. The collection of these sequents is the *theory* associated to the system.

Definition 12. For an LCS $\mathcal{L} = \langle Q, C, M, I \rangle$, define these sets of propositional variables:

- 1. $\mathbf{Q} := \{Q_i \mid q_i \in \mathbf{Q}\}$ (state variables)
- 2. $\# := \{\#_i^s \mid c_i \in C\} \cup \{\#_i^e \mid c_i \in C\}$ (channel markers)
- 3. $\mathsf{M} := \{P_a \mid a \in \mathsf{M}\}$ (alphabet variables)

Moreover, define $Props(\mathcal{L}) := Q \cup \# \cup M$.

Definition 13 (Theory of an LCS). Given an LCS $\mathcal{L} = \langle Q, C, M, I \rangle$, the theory of \mathcal{L} , denoted by $\mathcal{T}(\mathcal{L})$, is defined as the union of the following finite sets of sequents:

1. $\mathcal{T}(\mathcal{L})^{!} := \{\#_{l}^{e}, Q_{i} \Rightarrow P_{a} \otimes (\#_{l}^{e} \otimes Q_{j}) \mid (q_{i}, c_{l}, a, !, q_{j}) \in \mathbf{I}\};$ 2. $\mathcal{T}(\mathcal{L})^{?} := \{\#_{l}^{s}, P_{a}, Q_{i} \Rightarrow \#_{l}^{s} \otimes Q_{j} \mid (q_{i}, c_{l}, a, ?, q_{j}) \in \mathbf{I}\};$ 3. $\mathcal{T}(\mathcal{L})^{\mathsf{Q}} := \bigcup_{\substack{Q \in \mathsf{Q} \\ R \in \mathsf{Props}(\mathcal{L}) \setminus \mathsf{Q}}} \{R, Q \Rightarrow Q \otimes R\} \cup \{Q, R \Rightarrow R \otimes Q\}.$

Note that $\mathcal{T}(\mathcal{L})$ is a regular theory in the sense of Def. 3.

Recall that $\bigotimes \varphi_1, \ldots, \varphi_m := \varphi_1 \otimes (\varphi_2 \otimes (\ldots (\varphi_{m-1} \otimes \varphi_m) \ldots))$, where $m \geq 1$ and $\varphi_1, \ldots, \varphi_m$ is a sequence of formulas. We now translate a reachability instance in a fixed LCS $\mathcal{L} = \langle \mathsf{Q}, \mathsf{C}, \mathsf{M}, \mathsf{I} \rangle$ to a sequent-deducibility instance in $\mathbf{FL}_{\mathbf{w}}^{\otimes}$. We begin by coding a configuration as a sequence.

Definition 14. Given $u := (q_i, u_1, \ldots, u_n) \in \text{Conf}(\mathcal{L})$, define the sequence $E(u) := Q_i, \#_1^s, U_1, \#_1^e, \ldots, \#_n^s, U_n, \#_n^e$, where, for $u_i = a_1 \cdots a_{k_i} \in M^*$, U_i is the sequence $P_{a_1}, \ldots, P_{a_{k_i}}$ of propositional variables.

Definition 15. For $u := (q_i, u_1, \ldots, u_n)$, $v := (q_j, v_1, \ldots, v_n) \in Conf(\mathcal{L})$, define the sequent

$$S_{u \to v} := E(u) \Rightarrow \bigotimes E(v).$$

Define also $S_{u\to v}^{\bowtie}$ as the set of sequents of the above form where the occurrence of Q_i in the antecedent may appear in any position (i.e., not necessarily at the front as in E(u)).

We note that sequents in standard deductions in $\mathbf{FL}_{\mathbf{w}}^{\otimes}$ from a regular theory have a nonempty succedent coming from the observation that every leaf has this property:

Lemma 16. Let \mathcal{T} be a regular theory. It is never the case that $\mathcal{T} \vdash_{\mathbf{FL}_{\mathbf{w}}^{\otimes}}^{\mathsf{std}} \Gamma \Rightarrow .$

Proof. It follows because no axiomatic rule has conclusion with empty succedent and no other rule allows to derive a sequent with empty succedent from one with nonempty succedent. \Box

The flattening $[\varphi]$ is the sequence of propositional variables obtained from a formula φ by replacing ' \otimes ' with ','. E.g., $[p \otimes ((q \otimes s) \otimes r)] = p, q, s, r$. The flattening $[\Gamma]$ of a sequence Γ of formulas is the concatenation of the flattening of the formulas in Γ in the order they appear in the sequence. Let $\#\Delta$ denote the subsequence of Δ having only variables from the set # (that is, only channel markers). The Q-free subsequence of a sequence of propositional variables is obtained by deleting all occurrences of state variables.

The following lemmas express that in standard deductions from a regular theory, in various situations, propositional variables occurring positively must also occur negatively. They abstract what is needed for the right-to-left direction of Lem. 20 below. The proofs are by structural induction on standard deductions.

Lemma 17. Let Γ be a sequence of formulas, none having a state variable. If $\mathcal{T}(\mathcal{L}) \vdash_{\mathbf{FL}_{\mathbf{w}}^{\otimes} \Rightarrow}^{\mathsf{std}} \Gamma \Rightarrow \varphi$, then the Q-free subsequence of $[\varphi]$ is a subsequence of $[\Gamma]$.

Proof. Structural induction on a standard deduction witnessing $\mathcal{T}(\mathcal{L}) \vdash_{\mathbf{FL}_{w}^{\otimes}} \Gamma \Rightarrow \varphi$. Base case. For elements from $\mathcal{T}(\mathcal{L})$ it holds vacuously since each sequent contains a state variable in the antecedent; for an identity sequent the claim is immediate. Inductive step. Suppose that the last rule instance applied is (lw), (R \otimes) or (L \otimes). Applying the IH to its premise(s), we have that the Q-free subsequence of [φ] is a subsequence of the flattening of the premise antecedent(s), and hence also of [Γ]. The last rule instance cannot be a standard cut as the antecedent of each element in $\mathcal{T}(\mathcal{L})$ contain a state variable hence Γ would contains a state variable, contradicting the hypotheses. The rule (rw) cannot occur due to Lem. 16.

Lemma 18. Let Γ be a sequence of formulas, $u \in Conf(\mathcal{L})$ and L a nonempty subsequence of E(u). Then $\mathcal{T}(\mathcal{L}) \vdash_{\mathbf{FL}_{\mathbf{w}}^{\otimes} \Rightarrow}^{\mathsf{std}} \Gamma \Rightarrow \bigotimes L$ implies that the variables in #L appear at least once in $[\Gamma]$.

Proof. Induction on the structure of a standard deduction witnessing $\mathcal{T}(\mathcal{L}) \vdash_{\mathbf{FL}_{\mathbf{w}}^{\otimes} \Rightarrow} \Gamma \Rightarrow \bigotimes L$. Base case. Straightforward: identity sequent $p \Rightarrow p$ (trivial) and an element from $\mathcal{T}(\mathcal{L})$ (by inspection, it contains no channel variables or the same channel variable occurs in the antecedent and succedent). Inductive step. Consider the last rule instance in the

standard deduction. For $(\mathbb{R}\otimes)$, $(\mathbb{L}\otimes)$ and $(\mathbb{I}w)$, we have by IH that each variable in #L appears in $[\Gamma']$ for the antecedent Γ' of some premise of the rule instance. Those variables will be carried to the conclusion antecedent. If the last rule applied is a standard cut, its right premise has the form $\Gamma_1, \varphi, \Gamma_2 \Rightarrow \bigotimes L$ where φ is the cut formula. By IH, every variable in #L appears in $[\Gamma_1, \varphi, \Gamma_2]$. Moreover, since the left premise is from $\mathcal{T}(\mathcal{L})$, any channel variable occurring in the cut formula will occur in its antecedent, and hence also in $[\Gamma]$. The case of (rw) does not need to be considered in view of Lem. 16.

Lemma 19. Let Γ be a sequence of formulas and φ be a formula. If $\mathcal{T}(\mathcal{L}) \vdash_{\mathbf{FL}_{\mathbf{w}}^{\otimes} \Rightarrow}^{\mathsf{std}} \Gamma \Rightarrow Q_i \otimes \varphi$ or $\mathcal{T}(\mathcal{L}) \vdash_{\mathbf{FL}_{\mathbf{w}}^{\otimes} \Rightarrow}^{\mathsf{std}} \Gamma \Rightarrow Q_i$, then $[\Gamma]$ must contain a state variable.

Proof. Structural induction on a derivation witnessing $\mathcal{T}(\mathcal{L}) \vdash_{\mathbf{SL}_{w} \Rightarrow} \Gamma \Rightarrow Q_i \otimes \varphi$. Base case. The case of an initial sequent is trivial, and the antecedent of any sequent from $\mathcal{T}(\mathcal{L})$ contains a state variable. Inductive step. Suppose that the last rule is $(\mathbb{R}\otimes)$. The left premise has Q_i as succedent, and hence by IH, its antecedent will contain a state variable that is carried down to the conclusion. The argument is analogous for (lw) and (L \otimes). If the last rule instance is a standard cut, its left premise is a sequent in $\mathcal{T}(\mathcal{L})$. As already observed, the antecedent of such a sequent contains a state variable and that will be carried down to the conclusion. The case of (rw) does not need to be considered in view of Lem. 16.

The case of $\mathcal{T}(\mathcal{L}) \vdash_{\mathbf{FL}_{\mathbf{w}}^{\otimes} \Rightarrow} \Gamma \Rightarrow Q_i$ is similar.

We are ready to reduce reachability in lossy channel systems to sequent-deducibility.

Lemma 20. For all LCS \mathcal{L} , given $u, v \in \mathsf{Conf}(\mathcal{L})$, $u \to_{\mathcal{L} \text{ lossy}}^* v$ iff $\mathcal{T}(\mathcal{L}) \vdash_{\mathbf{FL}_{\mathbf{w}}^{\otimes}}^{\mathsf{sd}} s$ for some $s \in S_{u \to v}^{\bowtie}$.

Proof. In the left-to-right direction, we work by induction on the length $k \geq 1$ of a computation witnessing $u \to_{\mathcal{L} \text{lossy}}^* v$. Base case. If k = 1, u = v, and $S_{u \to v}$ is provable by (id) and (R \otimes). Inductive step. The computation decomposes as $u \to_{\mathcal{L} \text{lossy}}^{\sigma} v' \to_{\mathcal{L} \text{lossy}}^* v$ with length k + 1. As $v' \to_{\mathcal{L} \text{lossy}}^* v$ is witnessed by a computation of length k, IH yields $\mathcal{T}(\mathcal{L}) \vdash_{\mathbf{FL}_{w}^{\otimes} \Rightarrow} s'$ for some $s' \in S_{v' \to v}^{\otimes}$. We extend the latter deduction downwards to s, using a case analysis on σ to determine the form of s' and s:

1. if $\sigma = ?$, by effect of $(q_i, c_l, a, ?, q_j) \in I$: then $u_l = av'_l$, all the other channels have the same content, and the state transits from q_i to q_j . Thus s' has the following form, possibly following repeated standard cuts with sequents in $\mathcal{T}(\mathcal{L})^{\mathbb{Q}}$.

$$\#_1^s, U_1, \#_1^e, \dots, \#_{l-1}^s, U_{l-1}, \#_{l-1}^e, \#_l^s, Q_j, V_l', \#_l^e, \dots, \#_n^s, U_n, \#_n^e \Rightarrow \bigotimes E(v)$$

Apply (L \otimes) and a standard cut with $\#_l^s, P_a, Q_i \Rightarrow \#_l^s \otimes Q_j$ to obtain s.

2. if $\sigma = l$: the state is unchanged, and $v'_l = z_1 z_2$, for $u = z_1 a z_2$. Thus s' has the following form, possibly following repeated standard cuts with sequents in $\mathcal{T}(\mathcal{L})^{\mathsf{q}}$.

 $Q_{j}, \#_{1}^{s}, U_{1}, \#_{1}^{e}, \dots, \#_{l}^{s}, Z_{1}, Z_{2}, \#_{l}^{e}, \dots, \#_{n}^{s}, U_{n}, \#_{n}^{e} \Rightarrow \bigotimes E(v)$ By (lw) $Q_{j}, \#_{1}^{s}, U_{1}, \#_{1}^{e}, \dots, \#_{l}^{s}, Z_{1}, P_{a}, Z_{2}, \#_{l}^{e}, \dots, \#_{n}^{s}, U_{n}, \#_{n}^{e} \Rightarrow \bigotimes E(v)$ 3. if $\sigma = !$, by effect of $(q_i, c_l, a, !, q_j) \in I$: by definition of this step, $v'_l = u_l a$, the content in other channels of u and v' is unchanged, and the state transits from q_i to q_j . Thus s' has the following form modulo the position of Q_j . In any case, the s' can be moved to the displayed position through repeated standard cuts with sequents in $\mathcal{T}(\mathcal{L})^{\mathbb{Q}}$.

$$\#_1^s, U_1, \#_1^e, \dots, \#_l^s, U_l, P_a, \#_l^e, Q_j, \#_{l+1}^s, \dots, \#_n^s, U_n, \#_n^e \Rightarrow \bigotimes E(v)$$

It suffices now to apply $(L\otimes)$ twice and a standard cut with $\#_l^e, Q_i \Rightarrow P_a \otimes \#_l^e \otimes Q_j$ to obtain the sequent s: $\#_1^s, U_1, \#_1^e, \ldots, \#_l^s, U_l, \#_l^e, Q_j, \#_{l+1}^s, \ldots, \#_n^s, U_n, \#_n^e \Rightarrow \bigotimes E(v)$.

For the right-to-left direction, we establish property $\mathcal{P}(\delta) :=:$ "for all $u, v \in \mathsf{Conf}(\mathcal{L})$, if δ is a standard deduction witnessing $\mathcal{T}(\mathcal{L}) \vdash_{\mathbf{FL}_{\mathbf{w}}^{\otimes} \Rightarrow} s$ for some $s \in S_{u \to v}^{\bowtie}$, then $u \to_{\mathcal{L}}^{*}_{\mathsf{lossy}} v$ ". Induction on the structure of δ .

Base case. No initial sequent nor any sequent from $\mathcal{T}(\mathcal{L})$ is an image of an encoding of a configuration since it does not contain any pair $\{\#_i^s, \#_i^e\}$ of channel markers.

Inductive step. Let $u = (q_i, u_1, \ldots, u_n)$ and $v = (q_j, v_1, \ldots, v_n)$. By cases on the last rule applied in δ . It cannot be $(L\otimes)$ since the antecedent of s is a sequence of propositional variables. Then it could be:

- 1. (rw): Not possible by Lem. 16 since δ is a standard deduction.
- 2. (R \otimes): Write s as $\Gamma_1, \Gamma_2 \Rightarrow Q_j \otimes \varphi_2$. The premises then are $\Gamma_1 \Rightarrow Q_j$ and $\Gamma_2 \Rightarrow \varphi_2$. By Lem. 19, Γ_1 must have a formula with a state variable, and thus Γ_2 lacks a state variable. If Γ_1 contained anything more, it must contain a channel marker. By Lem. 17, Γ_2 must contain all the channel markers to fulfil the requirement that $[\varphi_2]$ must be a subsequence of $[\Gamma_2]$. Hence Γ_1 is the singleton sequence Q_l . Now, for $Q_l \Rightarrow Q_j$ to be provable, we must have l = j. Also, $[\varphi_2]$ a subsequence of $[\Gamma_2]$ implies that the channel portions in the succedent are subsequences of the corresponding ones in the antecedent. Thus, u can reach v by losing messages. Observe: the IH is not employed.
- 3. (lw): We claim that the weakening introduces a variable into the channel portion. This suffices since the premise would then be the image of a configuration that, by IH, reaches v, and is itself reachable from u by a lossy step. To establish the claim, first note that the sequents involved in an application of (lw) have the same succedent, in this case ⊗ E(v). Let Γ' be the premise antecedent. By Lem. 18, we have that all channel variables appear in Γ'. Since these symbols appear only once in conclusion antecedent, weakening could not have introduced them. If the weakening introduced the state variable, then Γ' would have no state variables; that is impossible due to Lem. 19 since the succedent does contain a state variable. The remaining possibility is that weakening introduced a variable in the channel portion, so the claim is proved.
- 4. Standard cut with a sequent in $\mathcal{T}(\mathcal{L})^{?}$: The deduction has the form

$$\frac{\#_{l}^{s}, P_{a}, Q_{i} \Rightarrow \#_{l}^{s} \otimes Q_{p}}{\#_{l}^{s}, U_{1}, \#_{1}^{e}, \dots, \#_{l}^{s} \otimes Q_{p}, U_{l}, \#_{l}^{e}, \dots, \#_{n}^{s}, U_{n}, \#_{n}^{e} \Rightarrow \bigotimes E(v)}{\#_{1}^{s}, U_{1}, \#_{1}^{e}, \dots, \#_{l}^{s}, P_{a}, Q_{i}, U_{l}, \#_{l}^{e}, \dots, \#_{n}^{s}, U_{n}, \#_{n}^{e} \Rightarrow \bigotimes E(v)}$$
(cut)
12

s

Consider the last rule in δ_1 . It cannot be a (standard) cut as that would have introduced a state variable on its own (i.e., unfused) into the conclusion. If it was (R \otimes) then the left premise has the form $\Gamma_1 \Rightarrow Q_j$ and the second premise has the form $\Gamma_2 \Rightarrow \varphi_2$. By Lem. 19, $\#_l^s \otimes Q_p \in \Gamma_1$, and thus Γ_2 consists only of propositional variables. Since $\#_l^s$ is not in Γ_2 but φ_2 contains all channel markers, Lem. 18 gives a contradiction. So (R \otimes) was not the last rule applied. Neither can it be (rw) due to Lem. 16.

We are left with two alternatives for the last rule in δ_1 : (lw) and (L \otimes). If (lw), arguing as in the previous item, the introduced variable must be in the channel portion by Lem. 18 and Lem. 19. Iterating the entire argument up to this point (implicit induction), after multiple applications of (lw) we must ultimately encounter (L \otimes) (we have ruled out all the other possibilities). The situation is therefore the following.

$$\frac{\delta'_{1}}{\#_{1}^{s}, P_{a}, Q_{i} \Rightarrow \#_{l}^{s} \otimes Q_{p}} \mathcal{T}(\mathcal{L})^{?} = \frac{\frac{\#_{1}^{s}, U'_{1}, \#_{1}^{e}, \dots, \#_{l}^{s}, Q_{p}, U'_{l}, \#_{l}^{e}, \dots, \#_{n}^{s}, U'_{n}, \#_{n}^{e} \Rightarrow \bigotimes E(v)}{\#_{1}^{s}, U_{1}, \#_{1}^{e}, \dots, \#_{l}^{s} \otimes Q_{p}, U'_{l}, \#_{l}^{e}, \dots, \#_{n}^{s}, U'_{n}, \#_{n}^{e} \Rightarrow \bigotimes E(v)} (L\otimes) \\ \frac{\#_{1}^{s}, U_{1}, \#_{1}^{e}, \dots, \#_{l}^{s} \otimes Q_{p}, U_{l}, \#_{l}^{e}, \dots, \#_{n}^{s}, U_{n}, \#_{n}^{e} \Rightarrow \bigotimes E(v)}{\#_{1}^{s}, U_{1}, \#_{1}^{e}, \dots, \#_{l}^{s} \otimes Q_{p}, U_{l}, \#_{l}^{e}, \dots, \#_{n}^{s}, U_{n}, \#_{n}^{e} \Rightarrow \bigotimes E(v)} (L\otimes) \\ (Uw)^{*} \\ \frac{\#_{1}^{s}, U_{1}, \#_{1}^{e}, \dots, \#_{l}^{s}, P_{a}, Q_{i}, U_{l}, \#_{l}^{e}, \dots, \#_{n}^{s}, U_{n}, \#_{n}^{e} \Rightarrow \bigotimes E(v)}{(uv)^{*}} \\ (cut) \\ \frac{W_{1}^{s}, W_{1}, W_{1}^{e}, \dots, W_{l}^{s}, P_{a}, Q_{i}, U_{l}, \#_{l}^{e}, \dots, \#_{n}^{s}, U_{n}, \#_{n}^{e} \Rightarrow \bigotimes E(v)}{(uv)^{*}} \\ (cut) \\ \frac{W_{1}^{s}, W_{1}, W_{1}^{e}, \dots, \#_{l}^{s}, P_{a}, Q_{i}, U_{l}, \#_{l}^{e}, \dots, \#_{n}^{s}, W_{n}, W_{n}^{e} \Rightarrow \bigotimes E(v)}{(uv)^{*}} \\ (cut) \\ \frac{W_{1}^{s}, W_{1}, W_{1}^{e}, \dots, W_{n}^{s}, W_{n}, W_{n}^{e} \Rightarrow \bigotimes E(v)}{(uv)^{*}} \\ (uv)^{*} \\ (uv)^{*}$$

From the IH, we obtain that $(q_p, u'_1, \ldots, u'_l, \ldots, u'_n) \rightarrow^*_{\mathcal{L} \text{ lossy }} v$. Since $(q_i, u_1, \ldots, au_l, \ldots, u_n) \rightarrow^?_{\mathcal{L} \text{ lossy }} (q_p, u_1, \ldots, u_l, \ldots, u_n) \rightarrow^*_{\mathcal{L} \text{ lossy }} (q_p, u'_1, \ldots, u'_n)$, we are done.

5. Standard cut with a sequent in $\mathcal{T}(\mathcal{L})^{!}$: In this case the deduction has the form

Consider the last rules in δ_1 . Arguing as in the $\mathcal{T}(\mathcal{L})^?$ case, it is a sequence of (lw) rules introducing variables in channel portions, then (L \otimes). The latter premise is

$$\#_1^s, U_1', \#_1^e, \dots, \#_l^s, U_l', P_a, \#_l^e \otimes Q_p, \dots, \#_n^s, U_n', \#_n^e \Rightarrow \bigotimes E(v)$$

Since no state variable occurs by itself in the antecedent, the above sequent was not derived by a (standard) cut. Nor was it by (R \otimes), since its left premise would contain $\#_l^e \otimes Q_p$ hence depriving the right premise antecedent of $\#_l^e$ leading to a contradiction with Lem. 18. Nor was it (rw) due to Lem. 16. Hence we encounter once again a sequence of (lw) rules followed by an application of (L \otimes), resulting in the sequent

$$\#_1^s, U_1'', \#_1^e, \dots, \#_l^s, U_l'', P_a, \#_l^e, Q_p, \dots, \#_n^s, U_n'', \#_n^e \Rightarrow \bigotimes E(v)$$

By IH, we obtain that $(q_p, u''_1, \ldots, u''_l a, \ldots, u''_n) \to^*_{\mathcal{L} \text{ lossy }} v$. Since $(q_i, u_1, \ldots, u_l, \ldots, u_n) \to^!_{\mathcal{L} \text{ lossy }} (q_p, u_1, \ldots, u_l a, \ldots, u_n) \to^*_{\mathcal{L} \text{ lossy }} (q_p, u''_1, \ldots, u''_l a, \ldots, u''_n)$, we conclude $u \to^*_{\mathcal{L} \text{ lossy }} v$.

6. Standard cut with $Q_i, R \Rightarrow R \otimes Q_i \in \mathcal{T}(\mathcal{L})^{\mathbb{Q}}$ (sequent $R, Q_i \Rightarrow Q_i \otimes R$ is similar):

As in the previous cases, the last rule applied in δ_1 could not have been a standard cut nor (rw). Suppose it is (R \otimes). Writing the endsequent of δ_1 as $\Gamma_3, \Gamma_4 \Rightarrow Q_j \otimes \varphi_2$, the left premise would have the form $\Gamma_3 \Rightarrow Q_i$ where Γ_3 must contain $R \otimes Q_i$ (Lem. 19) and hence also Γ_1 . If Γ_1 is non-empty then it must contain a channel marker and hence in the right premise $\Gamma_4 \Rightarrow \varphi_2$, the Γ_4 would lack a channel marker contradicting Lem. 18. If Γ_1 is empty then R must be a channel marker and once again Γ_4 would lack a channel marker leading to a contradiction. Therefore the rule is not (R \otimes).

The remaining possibility is a sequence of (lw) followed by $(L\otimes)$. The premise of the latter is in the image of an encoding, thus the IH applies and we are done.

Theorem 21. Let $\Sigma \subseteq \Sigma_{\mathbf{FL}}$. If $\otimes \in \Sigma$, then sequent-deducibility in $\mathbf{FL}_{\mathbf{w}}^{\Sigma}$ is HACK-hard. Moreover, if $\{\otimes, \backslash, 0, 1\} \subseteq \Sigma$, then deducibility is HACK-hard.

Proof. The first assertion follows from Lem. 20, plus the fact that the reduction established there is polynomial-time, and Lem. 5. The second one follows from the first and Lem. 2. \Box

4. Hyper-Ackermannian upper bounds

We give an algorithm to decide the relation $\vdash_{\mathbf{S}\Rightarrow}$ for regular theories, for a sequent calculus **S** containing the rule (lw) and satisfying a generalized subformula property as defined below. In what follows, given a finite $\Omega \subseteq \operatorname{Fm}_{\Sigma}(\mathcal{P})$, an Ω -sequent is a sequent in which only formulas in Ω appear.

Definition 22. A sequent calculus that is a structural rule extension of $\mathbf{FL}_{\mathbf{w}}^{\Sigma}$ ($\Sigma \subseteq \Sigma_{\mathbf{FL}}$) is called amenable provided it satisfies the generalized subformula property: for every regular theory \mathcal{T} , it is the case that $\mathcal{T} \vdash_{\mathbf{S} \Rightarrow} s$ iff there is a \mathcal{T} -deduction of s where only $\mathsf{subf}(\mathcal{T} \cup \{s\})$ -sequents appear (such a deduction is called analytic).

 $\mathbf{FL}_{\mathbf{w}}^{\Sigma}$ is amenable for any $\Sigma \subseteq \Sigma_{\mathbf{FL}}$; Lem. 4 gives the generalized subformula property. Indeed, any *analytic structural rule* extension of an amenable calculus is amenable (see results for the substructural hierarchy, Ciabattoni et al. [15]).

4.1. Just enough on well-quasi-order theory

A quasi-ordered set (qo-set) is a structure $\mathbf{W} := \langle W, \preccurlyeq_{\mathbf{W}} \rangle$, where W is a set and $\preccurlyeq_{\mathbf{W}} \subseteq W \times W$ is reflexive and transitive (a quasi order, for short). Abusing notation, we write $a \in \mathbf{W}$ for $a \in W$. We denote by \mathbb{N} the set of natural numbers; by \mathbb{N}^+ the set $\mathbb{N} \setminus \{0\}$; and, given $\ell \in \mathbb{N} \cup \{\omega\}$, we write $\mathbb{N}_{<\ell}$ for the set $\{a \in \mathbb{N} \mid a < \ell\}$. Given a qo-set \mathbf{W} , a bad sequence over \mathbf{W} (of length ℓ) is a sequence $(a_i)_{i \in \mathbb{N}_{<\ell}}$ of elements of \mathbf{W} such that,

for all i < j, $a_i \not\preccurlyeq_{\mathbf{W}} a_j$. A qo-set \mathbf{W} is a *well-quasi-ordered set (wqo-set)* if every bad sequence over it is finite (see [14, Sec. 2] for equivalent definitions). Examples include \mathbb{N}^k under component-wise ordering (Dickson's Lemma [19]), and sequences under the subword embedding ordering:

Theorem 23 (Higman's Lemma [14]). For any finite set Ω , $\langle \Omega^*, \preccurlyeq_* \rangle$ is a wqo, where, recall, Ω^* is the set of all finite sequences of elements from Ω and $w_1 \preccurlyeq_* w_2$ iff w_1 is obtainable from w_2 by deleting some elements.

Bad sequences over a wqo, though finite, do not in general have a maximum length. For example, $(1,0), (0,k), (0,k-1), \ldots, (0,0)$ is a bad sequence on \mathbb{N}^2 under the componentwise ordering with length k + 2 for every $k \in \mathbb{N}$. Nevertheless, a maximum length can be ensured by limiting the size of each element in the sequence to some (fixed) function of the size of the preceding element. To achieve this, the wqo is enriched with more structure. A normed (well-)quasi-ordered set (n(w)qo-set) [20] is a structure $\mathbf{W} := \langle W, \preccurlyeq_{\mathbf{W}}, \llbracket \cdot \rrbracket_{\mathbf{W}} \rangle$, where $\langle W, \preccurlyeq_{\mathbf{W}} \rangle$ is a (w)qo-set and $\llbracket \cdot \rrbracket_{\mathbf{W}} : W \to \mathbb{N}$ is a proper norm, meaning that, for all $n \in \mathbb{N}$, $\{a \in W : \llbracket a \rrbracket_{\mathbf{W}} < n\}$ is finite. It is easy to check that the following is a nwqo.

Definition 24. Given a finite set Ω , let $\Omega^* := \langle \Omega^*, \preccurlyeq_*, \llbracket \cdot \rrbracket_* \rangle$, where $\llbracket w \rrbracket_*$ is the length of w (and $\langle \Omega^*, \preccurlyeq_* \rangle$ is as in Thm. 23).

Indeed, the *disjoint sum* of these nwqos, defined below, is also an nwqo [20].

Definition 25. Given a finite set Ω and $m \in \mathbb{N}$, let $m \cdot \Omega^* := \langle m \cdot \Omega^*, \preccurlyeq_{m \cdot \Omega^*}, \llbracket \cdot \rrbracket_{m \cdot \Omega^*} \rangle$. Here, $m \cdot \Omega^* := \{1, \ldots, m\} \times \Omega^*$ is the disjoint sum of m copies of Ω^* , and $(i, w_1) \preccurlyeq_{m \cdot \Omega^*} (j, w_2)$ iff i = j and $w_1 \preccurlyeq_* w_2$. Also, define $\llbracket (i, w) \rrbracket_{m \cdot \Omega^*} := \llbracket w \rrbracket_*$.

A control function is a mapping $g : \mathbb{N} \to \mathbb{N}$ that is strictly increasing (for all $a, b \in \mathbb{N}$, if a < b, then g(a) < g(b)) and strictly inflationary (a < g(a) for all $a \in \mathbb{N}$).

For an nwqo-set \mathbf{W} , control function g, and initial parameter $t \in \mathbb{N}$, a (g, t)-controlled bad sequence over \mathbf{W} is a bad sequence $a_0, \ldots, a_{\ell-1}$ over \mathbf{W} where $\llbracket a_i \rrbracket_{\mathbf{W}} < g^i(t)$ for all $0 \leq i < \ell$. The (g, t)-controlled bad sequences have a maximum length [21] using König's Lemma. Denote by $L^g_{\mathbf{W}}(t)$ the maximum length of a (g, t)-controlled bad sequence over \mathbf{W} . We call $L^g_{\mathbf{W}} : \mathbb{N} \to \mathbb{N}$ the length function of \mathbf{W} (for g). Bounds for (g, t)-controlled bad sequences over \mathbf{W} reduce to finding bounds for $L^g_{\mathbf{W}}$ ('length theorems'), expressed in terms of the ordinal-indexed extended Grzegorczyk hierarchy $\{\mathfrak{F}_{\alpha}\}_{\alpha}$, from which the hierarchy $\{\mathbf{F}_{\alpha}\}_{\alpha}$ of fast-growing complexity classes are defined (the reader is referred to [10, Sec. 2] for detailed definitions). What matters here is that to classify a problem in \mathbf{F}_{α} , it suffices to show that the complexity of every instance is upper bounded by a function in \mathfrak{F}_{β} for some $\beta < \alpha$. Schmitz and Schnoebelen obtained the following length theorem for Higman's lemma.

Theorem 26 ([6, Thm. 5.3]). For any finite set Ω , $m \in \mathbb{N}$, and primitive recursive control function g, the length function $L^g_{m:\Omega^*}$ is upper bounded by a function in $\mathfrak{F}_{\omega^{|\Omega|-1}}$.

A *reflection* is a map between nqo-sets that preserves bad sequences. Hence, its existence implies the reverse transfer of the wqo-property and upper bounding of length theorem. In the next subsection, we obtain a length theorem for a nwqo over sequents in this way.

Definition 27 (Reflection). Let $\mathbf{W}_1 := \langle W_1, \preccurlyeq_1, \llbracket \cdot \rrbracket_1 \rangle$ and $\mathbf{W}_2 := \langle W_2, \preccurlyeq_2, \llbracket \cdot \rrbracket_2 \rangle$ be non-sets. A reflection between \mathbf{W}_1 and \mathbf{W}_2 is a mapping $f : W_1 \to W_2$ such that, for all $a, b \in W_1$, if $f(a) \preccurlyeq_2 f(b)$, then $a \preccurlyeq_1 b$ and, for all $a \in W_1$, $\llbracket f(a) \rrbracket_2 \leq \llbracket a \rrbracket_1$. We write $\mathbf{W}_1 \hookrightarrow^f \mathbf{W}_2$ (sometimes omitting the superscript f).

Lemma 28 ([6, p. 446]). Whenever $\mathbf{W}_1 \hookrightarrow \mathbf{W}_2$, if \mathbf{W}_2 is a nwqo, then so is \mathbf{W}_1 ; and if g is a control function, then $L^g_{\mathbf{W}_1}(t) \leq L^g_{\mathbf{W}_2}(t)$ for all $t \in \mathbb{N}$.

4.2. A well-quasi-order over sequents based on weakening

Observe that the antecedent of the premise of the left-weakening rule (lw) is obtainable from the antecedent of the conclusion by deleting letters, and the succedents are identical. This motivates an nwqo over sequents, for which we obtain a length theorem via a reflection into a disjoint sum of Higman's orderings. In what follows, let Σ be an arbitrary signature.

Definition 29. For finite $\Omega \subseteq \operatorname{Fm}_{\Sigma}(\mathcal{P})$, let $\mathbf{W}_{\Sigma}(\Omega) := \langle \operatorname{Seq}_{\Sigma}(\Omega), \leadsto_{\Sigma}^{\Omega}, \llbracket \cdot \rrbracket \rangle$, where

- $\operatorname{Seq}_{\Sigma}(\Omega)$ is the set of all Ω -sequents over $\operatorname{Fm}_{\Sigma}(\mathcal{P})$;
- $\rightsquigarrow_{\Sigma}^{\Omega} \subseteq \mathsf{Seq}_{\Sigma}(\Omega) \times \mathsf{Seq}_{\Sigma}(\Omega)$ is such that $s_1 \rightsquigarrow_{\Sigma}^{\Omega} s_2$ iff s_2 is obtained by successive applications of (lw) starting from s_1 ;
- $\llbracket \cdot \rrbracket : \operatorname{Seq}_{\Sigma}(\Omega) \to \mathbb{N}$ is such that $\llbracket \Gamma \Rightarrow \Pi \rrbracket := |\Gamma|$ (i.e., the length of the antecedent).

Theorem 30. For all finite $\Omega \subseteq \operatorname{Fm}_{\Sigma}(\mathcal{P})$, $\mathbf{W}_{\Sigma}(\Omega)$ is a nwqo; and if g is primitive recursive, then $L^{g}_{\mathbf{W}_{\Sigma}(\Omega)}$ is upper bounded by a function in $\mathfrak{F}_{\omega^{|\Omega|-1}}$.

Proof. Fix an enumeration $\varphi_1, \ldots, \varphi_m$ of the formulas in Ω . Then the mapping $f : Seq_{\Sigma}(\Omega) \to (|\Omega| + 1) \cdot \Omega^*$ such that $f(\Gamma \Rightarrow) := (0, \Gamma)$ and $f(\Gamma \Rightarrow \varphi_j) := (j, \Gamma)$ is a reflection; hence the result follows from Thm. 26 and Lem. 28.

4.3. Decision procedure for amenable sequent calculi

Fix an amenable sequent calculus **S** (see Def. 22) over a signature Σ .

Definition 31. For \mathcal{X} a formula, a sequent, or a sequent rule, let $\langle \mathcal{X} \rangle$ be the length of the written representation of \mathcal{X} . If X is a finite set of formulas, of sequents, or of sequent rules, let $\langle X \rangle := \sum_{\mathcal{X} \in X} \langle \mathcal{X} \rangle$ and $\langle \max X \rangle := \max_{\mathcal{X} \in X} \langle \mathcal{X} \rangle$. In particular, $\langle \max S \rangle := \max_{r \in S} \langle r \rangle$ i.e., the largest among the representation lengths of the rules in the calculus.

The construction in this subsection extends the commutative setting that appears in Balasubramanian et al. [13]. For a finite set D of sequents, define $\llbracket_{\max}D\rrbracket := \max_{s \in D} \llbracket s \rrbracket$ ($\llbracket s \rrbracket$ is the length of the antecedent of s, see Def. 29).

Definition 32. Let Ω be a finite set of formulas closed under subformulas and \mathcal{T} be a regular theory containing only Ω -sequents. Define D_0 as the set of minimal elements with respect to $\sim \stackrel{\Omega}{\Sigma}$ (i.e., not obtainable from another element by repeated (lw)) from the following finite set: the union of \mathcal{T} with the set of all instances of initial sequents in **S** that satisfy the following:

- a) formula-variables are instantiated to elements of Ω ;
- b) succedent-variables are instantiated to an element in Ω or as empty;
- c) sequence-variables are instantiated to the empty sequence.

Define $D_{i+1} := D_i \cup \partial D_i$ $(i \ge 0)$, where ∂D_i is the set of Ω -sequents s satisfying:

- 1. $s_1 \cdots s_k/s$ is a rule instance of **S** such that, for all $1 \leq j \leq k$, there is $s_j' \in D_i$ with $s_j' \rightsquigarrow_{\Sigma}^{\Omega} s_j;$
- 2. the antecedent of s has length $\leq (\langle \mathbf{S} \rangle \llbracket_{\max} D_i \rrbracket) \cdot \langle \mathbf{S} \rangle$;
- 3. there is no $s' \in D_i$ such that $s' \rightsquigarrow_{\Sigma}^{\Omega} s$.

The above construction yields a chain $D_0 \subseteq D_1 \subseteq \ldots$ of sets of sequents. The following shows that the chain stabilizes at a finite index. Indeed, if that were not the case, Def. 32 (3) forces the existence of an infinite bad sequence $(s_i)_{i < \omega}$ from $s_i \in D_i \setminus D_{i-1}$ $(i < \omega)$, and that is impossible.

Theorem 33. D_{i+1} is computable from D_i and there is $N \in \mathbb{N}$ such that $D_N = D_{N+i}$ for all $i \geq 0$.

Proof. Each D_i is computable because (a) there are only finitely-many rules in \mathbf{S} , (b) finitelymany formulas in Ω , (c) Def. 32 (2) restricts the length of antecedents (thus there are only finitely-many rule instances to consider), and (d) the relation $\rightsquigarrow_{\Sigma}^{\Omega}$ is computable.

If there is some N such that $D_N = D_{N+1}$ then, by the construction of these sets, $D_N = D_{N+i}$ for every $i \ge 0$. Suppose that no such N exists. It follows that $D_0 \subset D_1 \subset D_2 \subset \ldots$. Choose any $s_i \in D_i \setminus D_{i-1}$ for each $i \in \mathbb{N}$. By Def. 32 (3), $(s_i)_{i < \omega}$ is an infinite bad sequence over $\mathbf{W}_{\Sigma}(\Omega)$, contradicting that the latter is a wqo (Thm. 30).

We now establish (Thm. 35) that every deducible sequent is obtainable by weakening from a sequent at the stabilization point. First, a technical lemma.

Lemma 34. If $s' \rightsquigarrow_{\Sigma}^{\Omega} s$ with $s = L_1 \cdots L_m \Rightarrow \Pi$, then there are sequences L'_1, \ldots, L'_m such that $s' = L'_1 \cdots L'_m \Rightarrow \Pi$ and $L'_i \preccurlyeq_* L_i$ for all $1 \le i \le m$.

Proof. By induction on a deduction δ witnessing $s' \rightsquigarrow_{\Sigma}^{\Omega} L_1 \cdots L_m \Rightarrow \Pi$. If it has a single node, the involved sequents are the same, so take $L'_i := L_i$. In the inductive step, assume $s = L_1 \cdots L_i^1 \varphi L_i^2 \cdots L_m \Rightarrow \Pi$ was obtained by weakening from $s'' = L_1 \cdots L_i^1 L_i^2 \cdots L_m \Rightarrow \Pi$, where $L_i = L_i^1 L_i^2$. Then, by the (IH), $s' = L'_1 \cdots L'_i \cdots L'_m \Rightarrow \Pi$ with $L'_j \preccurlyeq_* L_j$ for all $1 \le j \le m$. In particular, $L'_i \preccurlyeq_* L_i^1 L_i^2 \preccurlyeq_* L_i^1 \varphi L_i^2$, and we are done.

Theorem 35. Let $\mathcal{T} \cup \{s\}$ be a finite set of Ω -sequents such that \mathcal{T} is a regular theory (as in Def. 32). Then $\mathcal{T} \vdash_{\mathbf{S}\Rightarrow} s$ iff there is $M \in \mathbb{N}$ such that $s' \rightsquigarrow_{\Sigma}^{\Omega} s$ for some $s' \in D_M$.

Proof. The left-to-right direction is the non-trivial one. Since **S** is amenable, $\mathcal{T} \vdash_{\mathbf{S}\Rightarrow} s$ is witnessed by an analytic deduction δ . We argue by induction on the structure of δ .

Base case. Then δ is a single node, so the node is a sequent in \mathcal{T} or an instance of an initial sequent. So s is in D_0 , or s is obtainable from the node by applications of (lw).

Inductive step. The last rule instance in δ is an instance $s_1 \cdots s_k/s$ of \mathbf{r} . Since each s_i is an Ω -sequent by virtue of the deduction being analytic, by applying IH for each $1 \leq i \leq k$, there is $N_i \in \mathbb{N}$ with $s'_i \rightsquigarrow_{\Sigma}^{\Omega} s_i$ for some $s'_i \in D_{N_i}$. Let $M := \max_i N_i$, which gives us $s'_i \in D_M$ for all $1 \leq i \leq k$. Hence Def. 32 (1) is satisfied for s. The situation is the following.

We now show by cases that there is $s' \in D_{M+1}$ such that $s' \rightsquigarrow_{\Sigma}^{\Omega} s$.

If Def. 32 (2) is satisfied, then $s \in D_{M+1}$, or (due to Def. 32 (3)) there is some $s' \in D_{M+1}$ such that $s' \rightsquigarrow_{\Sigma}^{\Omega} s$. In either case we are done, so assume that Def. 32 (2) fails. Thus s has antecedent of length $> (\langle \mathbf{S} \rangle \llbracket_{\max} D_M \rrbracket) \cdot \langle \mathbf{S} \rangle$. Thus there are sequence-variables $\Gamma_1, \ldots, \Gamma_m$ (m > 0) in the conclusion of \mathbf{r} instantiated with sequences of length $> \langle \mathbf{S} \rangle \llbracket_{\max} D_M \rrbracket$ (if m = 0then the antecedent would be instantiations of formula-variables and hence its length would be $\leq \langle \mathbf{S} \rangle \llbracket_{\max} D_0 \rrbracket$). Assume wlog that $\Gamma_1, \ldots, \Gamma_{m'}$ with $m' \leq m$ appear in the premises of \mathbf{r} , while the other variables appear exclusively in the conclusion.

Write $s_i = L_{i1} \cdots L_{iu_i} \Rightarrow \prod_i$ and $s'_i = L'_{i1} \cdots L'_{iu_i} \Rightarrow \prod_i$ where each L_i is an instantiation of a sequence-variable or a singleton sequence of a formula, such that $L'_{ij} \preccurlyeq L_{ij}$ for each $1 \leq j \leq u_i$ (the existence of such sequences is guaranteed by Lem. 34). The length of $L'_{i1} \cdots L'_{iu_i}$ is $\leq [max D_M]$ by the fact that $s'_i \in D_M$, hence each L'_{ij} has length $\leq [max D_M]$.

We construct a new instance I' of \mathbf{r} , with premises s''_i instead of s_i , and a conclusion s' whose antecedent has size satisfying Def. 32(2), and $s' \rightsquigarrow_{\Sigma}^{\Omega} s$:

Construct I' by consideration of where each Γ_j $(1 \leq j \leq m')$ occurs. For Γ_j occurring in a single premise, instantiate it with the sequence L'_{jl} instead of the L_{jl} that was used before.

For each Γ_j $(1 \leq j \leq m')$ occurring in multiple premises, we need a single instantiation that can be used for each occurrence (reflecting the fact that we are dealing with additive rules here). Without loss of generality, assume that s_1, \ldots, s_l are all the premises where Γ_j occurs. Assume that Γ_j was instantiated with the sequence L. Let $L_{1j_1}, \ldots, L_{lj_l}$ be the sequences in these premises that instantiate Γ_j (each is equal to L). We want a subsequence of L reachable by weakening from each $L'_{1j_1}, \ldots, L'_{lj_l}$ (these need not be identical): start with L and cross off each of the sequences $L'_{1j_1}, \ldots, L'_{lj_l}$ (letter by letter, respecting the order). Use the sequence crossed-off to instantiate Γ_j . It has size $\leq k \cdot [\lceil \max D_M \rceil] < \langle \mathbf{S} \rangle \cdot [\lceil \max D_M \rceil]$. Finally, for sequence-variables that appear only in the conclusion of the rule, i.e., $\Gamma_{m'+1}, \ldots, \Gamma_m$, instantiate each of these with the empty sequence.

The new instantiation I' has premises s''_i where $s'_i \rightsquigarrow_{\Sigma}^{\Omega} s''_i$ for each $1 \leq i \leq k$, and its conclusion is $s' \rightsquigarrow_{\Sigma}^{\Omega} s$ since the instantiation of each Γ_j is a subsequence of the original instantiation. Moreover, the number of variables in the conclusion of \mathbf{r} is $\langle \mathbf{S} \rangle$ and each variable instantiation under I' has size $\langle \mathbf{S} \rangle [\![\mathsf{max} D_M]\!]$. Hence it follows that $s' \in D_{M+1}$ or there is some $s'_0 \in D_M$ with $s'_0 \rightsquigarrow_{\Sigma}^{\Omega} s'$. In either case, the claim is proved.

4.4. Complexity analysis

The fast-growing complexity classes are closed under exponentiation so the distinction between space and time (also determinism and non-determinism) is unimportant [10]. We undertake a space analysis of the algorithm based on Thm. 33 and Thm. 35. It suffices to write down each D_i in turn at the same location, computing each rule instance in turn; when the stabilization index D_N is reached, check whether $s' \rightsquigarrow_{\Sigma}^{\Omega} s$ for some $s' \in D_N$. Each $D_i \subseteq D_N$, and writing down a rule instance takes space bounded by the size of an element from $D_N^{(S)} \times D_N$. So the space requirements are an elementary function in $|D_N|$ and the size of an element from D_N . We upper bound each of these.

Lemma 36. Let \mathcal{T} be a regular theory, s be a sequent and $\Omega := \mathsf{subf}(\mathcal{T} \cup \{s\})$.

- 1. $|\Omega| \leq \langle s \rangle + \langle \mathcal{T} \rangle;$
- 2. if $s' \in D_i$ then $[\![s']\!] < (\langle \mathcal{T} \rangle + 1) \langle \mathbf{S} \rangle^{2i+1}$, and $[\![\max D_i]\!] < (\langle \mathcal{T} \rangle + 1) \langle \mathbf{S} \rangle^{2i+1}$;

3. if
$$s' \in D_i$$
 then $\langle s' \rangle < 4(\langle \mathcal{T} \rangle + \langle s \rangle)^2 \langle \mathbf{S} \rangle^{2N+1}$

- 4. for all $i \ge 0$, $|D_i| < E(\langle \mathcal{T} \rangle + \langle s \rangle, N)$ for E an elementary function (i.e., in \mathfrak{F}_2).
- *Proof.* 1. It is easy to see by structural induction that $|\mathsf{subf}(\varphi)| \leq \langle \varphi \rangle$ for all formulas φ . Then, for any sequent $t := \psi_1, \ldots, \psi_m \Rightarrow \Pi$, we have $|\mathsf{subf}(t)| \leq |\mathsf{subf}(\Pi)| + \sum_{i=1}^{m} |\mathsf{subf}(\psi_i)| \leq \langle \Pi \rangle + \sum_{i=1}^{m} \langle \psi_i \rangle \leq \langle t \rangle$. The result then follows because $|\Omega| \leq |\mathsf{subf}(s)| + \sum_{t \in \mathcal{T}} |\mathsf{subf}(t)| \leq \langle s \rangle + \langle \mathcal{T} \rangle$.
- 2. By induction on $i \geq 0$. Note that if $s' \in D_0$, since D_0 is a subset of the union of \mathcal{T} and instantiations from Ω of initial sequents in \mathbf{S} , $[\![s']\!] < [\![_{\max}\mathcal{T}]\!] + \langle \mathbf{S} \rangle \leq ([\![_{\max}\mathcal{T}]\!] + 1) \langle \mathbf{S} \rangle \leq (\langle \mathcal{T} \rangle + 1) \langle \mathbf{S} \rangle$. Inductive step: if $s' \in D_{i+1}$, then $[\![s']\!] < \langle \mathbf{S} \rangle^2 [\![_{\max}D_i]\!] < \langle \mathbf{S} \rangle^2 (\langle \mathcal{T} \rangle + 1) \langle \mathbf{S} \rangle^{2i+1} = (\langle \mathcal{T} \rangle + 1) \langle \mathbf{S} \rangle^{2(i+1)+1}$.

Since $\llbracket \max D \rrbracket = \max_{s \in D} \llbracket s \rrbracket$, it follows that $\llbracket \max D_i \rrbracket < (\langle \mathcal{T} \rangle + 1) \langle \mathbf{S} \rangle^{2i+1}$

3. It is enough to observe that $\langle s' \rangle$ is upper bounded by the sum of the sizes of each formula in the antecedent, plus 1 for each comma, plus 1 to account for the sequent symbol, plus the size of the succedent. This gives us $\langle s' \rangle \leq [\![s']\!] \langle \Omega \rangle + [\![s']\!] + 1 + \langle \Omega \rangle \leq 4([\![s']\!] + 1) \langle \Omega \rangle$. From the previous items, the latter is $\leq 4((\langle \mathcal{T} \rangle + 1) \langle \mathbf{S} \rangle^{2N+1} + 1)(\langle s \rangle + \langle \mathcal{T} \rangle) \leq 4((\langle \mathcal{T} \rangle + \langle s \rangle) \langle \mathbf{S} \rangle^{2N+1})(\langle s \rangle + \langle \mathcal{T} \rangle)$ where we have used that $\langle s \rangle \geq 2$ and $\langle \mathbf{S} \rangle \geq 1$. 4. Let $\eta, \rho \in \mathbb{N}$. Then define $\begin{bmatrix} \eta \\ \rho \end{bmatrix}_{\mathsf{seq}} := (\sum_{l=0}^{\rho} \eta^l) \cdot (\eta+1) \leq (\eta+1)^{\rho+1} \cdot (\eta+1)$, the number of distinct sequents over η formulas and having norm $\leq \rho$. Note that $|D_0| \leq \begin{bmatrix} \langle s \rangle + \langle \mathcal{T} \rangle \\ \langle s \rangle + \langle \mathcal{T} \rangle \end{bmatrix}_{\mathsf{seq}}$ and $|D_{i+1}| \leq \begin{bmatrix} |\Omega| \\ \langle \mathbf{S} \rangle^2 \llbracket_{\mathsf{max}} D_i \rrbracket \end{bmatrix}_{\mathsf{seq}} < \begin{bmatrix} \langle s \rangle + \langle \mathcal{T} \rangle \\ \langle \mathbf{S} \rangle^2 (\langle \mathcal{T}+1) \rangle \langle \mathbf{S} \rangle^{2N+1} \end{bmatrix}_{\mathsf{seq}} \leq \begin{bmatrix} \langle s \rangle + \langle \mathcal{T} \rangle \\ \langle \mathbf{S} \rangle^2 (\langle s \rangle + \langle \mathcal{T} \rangle) \langle \mathbf{S} \rangle^{2N+1} \end{bmatrix}_{\mathsf{seq}}$. Thus, for all $i \geq 0$, $|D_i| < E(\langle s \rangle + \langle \mathcal{T} \rangle, N)$ for E an elementary function (i.e. in \mathfrak{F}_2).

Theorem 37. Deciding whether $\mathcal{T} \vdash_{\mathbf{S}\Rightarrow} s$ for a regular theory \mathcal{T} is in HACK.

Proof. Lem. 36 (4) shows that $|D_N| < E_1(\langle \mathcal{T} \rangle + \langle s \rangle, N)$, for an elementary function E_1 . Also, Lem. 36 (3) shows that that $s' \in D_N$ implies that its size $\langle s' \rangle$ is bounded by $E_2(\langle \mathcal{T} \rangle + \langle s \rangle, N)$, for an elementary function E_2 . It remains to upper bound N, and it is this that forces the fast-growing complexity. Let $\Omega := \operatorname{subf}(\mathcal{T} \cup \{s\})$. In the proof of Thm. 33, we extracted a bad sequence $(s_i)_{i \leq N}$ over $\mathbf{W}_{\Sigma}(\Omega)$ from the chain of sets $D_0 \subset \ldots \subset D_N$ by choosing any $s_i \in D_i \setminus D_{i-1}$. We claim that this bad sequence is (g, n)-controlled for the control function $g(x) := \langle \mathbf{S} \rangle^2 x$ and $n := (\langle \mathcal{T} \rangle + 1) \langle \mathbf{S} \rangle$, allowing us to conclude that $N \leq L^g_{\mathbf{W}_{\Sigma}(\Omega)}(n) \leq L^g_{\mathbf{W}_{\Sigma}(\Omega)}((\langle \mathcal{T} \rangle + \langle s \rangle) \langle \mathbf{S} \rangle)$. By induction on *i*. Base case: $s_0 \in D_0$, so Lem. 36(2) yields $[[s_0]] < (\langle \mathcal{T} \rangle + 1) \langle \mathbf{S} \rangle$. Inductive step: $[[s_{i+1}]] < (\langle \mathcal{T} \rangle + 1) \langle \mathbf{S} \rangle^{2(i+1)+1} = \langle \mathbf{S} \rangle^2 (\langle \mathcal{T} \rangle + 1) \langle \mathbf{S} \rangle^{2i+1} = \langle \mathbf{S} \rangle^2 g^i(\langle \mathcal{T} \rangle + 1) \langle \mathbf{S} \rangle) = g^{i+1}(n)$, so we are done. So the space to run the algorithm is a composition of elementary functions with a function in $\mathfrak{F}_{\omega^{|\Omega|-1}}$ by Thm. 30, thus it is bounded by a function in $\mathfrak{F}_{\omega^{|\Omega|-1}}$. So, for this particular $|\Omega|$, the problem is in $\mathbf{F}_{\omega^{|\Omega|}}$. As Ω varies with the input, we wish to eliminate its dependence. Upper bounding over all Ω , we have that the problem is in $\mathbf{F}_{\omega^{\omega}} = \text{HACK}$.

Corollary 38. Any subcalculus of $\mathbf{FL}_{\mathbf{w}}$ containing (lw) has deducibility in HACK. This holds in particular for $\mathbf{FL}_{\mathbf{w}}^{\Sigma}$ for any $\Sigma \subseteq \Sigma_{\mathbf{FL}}$.

5. Final considerations

Combining Sec. 3 and Sec. 4, we finally obtain the promised result:

Theorem 1 (Main theorem). For $\{\otimes, \backslash, 0, 1\} \subseteq \Sigma \subseteq \Sigma_{\mathbf{FL}}$, deducibility in $\mathbf{FL}_{\mathbf{w}}^{\Sigma}$ is HACKcomplete. In particular, $\mathbf{FL}_{\mathbf{w}}$ and its multiplicative fragment $\mathbf{L}_{\mathbf{w}}$ are HACK-complete.

Complexity of the word problem in integral FL-algebras. Let V be an equational class of algebras (a variety) over a signature Σ . The word problem of V [16, Sec. 4.4.2] asks whether, given fixed finite sets of variables X (generators) and equations E over $\operatorname{Fm}_{\Sigma}(X)$, the quasiequation $\& E \Longrightarrow e$ is valid in V, where e is an equation over $\operatorname{Fm}_{\Sigma}(X)$. Note that deciding this problem allows for one algorithm per pair (X, E), as opposed to deciding the quasiequational theory of V, which asks for a single algorithm that applies to every (X, E). In view of the algebraizability of $\vdash_{\mathbf{FL}_{\mathbf{w}}}$ [16, Sec. 2.6] w.r.t. the variety of integral zero-bounded FL-algebras $\mathsf{FL}_{\mathbf{w}}$, our results imply that the word problem and the quasiequational theory of FL-algebras $\mathsf{FL}_{\mathbf{w}}$, our results imply that the word problem and the quasiequational theory of end $\mathcal{T}(\mathcal{L})^{\rightarrow}$ (cf. Def. 13) is a finite set of formulas over $\operatorname{Fm}_{\Sigma_{\mathbf{FL}}}(\mathsf{Props}(\mathcal{L}))$, we have that the word problem in this variety is HACK-hard. The quasiequational theory is in HACK in view of the proof search procedure in the previous section. Since the word problem reduces to the quasiequational theory, we obtain that both problems are HACK-complete. This also applies to integral FL-algebras and integral residuated lattices since the constant 0 does not play any essential role in the arguments.

Non-existence of deduction theorem. Our lower bounds imply that fragments of $\mathbf{FL}_{\mathbf{w}}$ covered by Thm. 21 have no deduction theorem (DT); else deducibility would reduce to provability, yet provability in these logics is PSPACE.

Upper bound for axiomatic extensions. The obtained upper-bounds apply to deducibility for \mathcal{N}_2 -analytic structural rule extensions of $\mathbf{FL}_{\mathbf{w}}$, and to the corresponding axiomatic extensions (refer to the substructural hierarchy [15]).

Undecidability of deducibility in \mathbf{FL} . The encoding in Sec. 3.2 offers a new proof of the undecidability of the deducibility problem in \mathbf{FL} (remove lossiness; reachability in channel systems is undecidable). This might be useful to prove undecidability of deducibility in some axiomatic extensions of \mathbf{FL} .

References

- W. J. Blok, C. J. V. Alten, On the finite embeddability property for residuated ordered groupoids, Transactions of the American Mathematical Society 357 (10) (2005) 4141–4157.
- [2] P. Jipsen, C. Tsinakis, A Survey of Residuated Lattices, Springer US, Boston, MA, 2002, pp. 19–56. doi:10.1007/978-1-4757-3627-4_3.
- [3] R. Horčík, K. Terui, Disjunction property and complexity of substructural logics, Theoretical Computer Science 412 (31) (2011) 3992–4006. doi:https://doi.org/10.1016/j.tcs.2011.04.004.
- [4] P. Lincoln, J. Mitchell, A. Scedrov, N. Shankar, Decision problems for propositional linear logic, Annals of Pure and Applied Logic 56 (1) (1992) 239-311. doi:https://doi.org/10.1016/0168-0072(92)90075-B.
- [5] A. Urquhart, The complexity of decision procedures in relevance logic II, The Journal of Symbolic Logic 64 (4) (1999) 1774–1802.
- [6] S. Schmitz, P. Schnoebelen, Multiply-recursive upper bounds with Higman's Lemma, in: L. Aceto, M. Henzinger, J. Sgall (Eds.), Automata, Languages and Programming, Springer Berlin Heidelberg, Berlin, Heidelberg, 2011, pp. 441–452.
- H. Tanaka, Tower-complete problems in contraction-free substructural logics, in: B. Klin, E. Pimentel (Eds.), 31st EACSL Annual Conference on Computer Science Logic, CSL 2023, February 13-16, 2023, Warsaw, Poland, Vol. 252 of LIPIcs, Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2023, pp. 34:1–34:19.
- [8] K. Chvalovský, R. Horčík, Full Lambek calculus with contraction is undecidable, The Journal of Symbolic Logic 81 (2) (2016) 524–540.
- [9] N. Galatos, H. Ono, Cut elimination and strong separation for substructural logics: An algebraic approach, Annals of Pure and Applied Logic 161 (9) (2010) 1097–1133. doi:10.1016/j.apal.2010.01.003.
- S. Schmitz, Complexity hierarchies beyond elementary, ACM Trans. Comput. Theory 8 (1) (feb 2016). doi:10.1145/2858784.
- [11] R. Péter, Recursive Functions, Academic Press, 1967.
- [12] P. Chambart, P. Schnoebelen, The ordinal recursive complexity of lossy channel systems, in: 2008 23rd Annual IEEE Symposium on Logic in Computer Science, 2008, pp. 205–216. doi:10.1109/LICS.2008.47.
- [13] A. R. Balasubramanian, T. Lang, R. Ramanayake, Decidability and complexity in weakening and contraction hypersequent substructural logics, in: 36th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2021, Rome, Italy, June 29 - July 2, 2021, IEEE, 2021, pp. 1–13. doi:10.1109/LICS52264.2021.9470733.
- [14] G. Higman, Ordering by divisibility in abstract algebras, Proceedings of the London Mathematical Society s3-2 (1) (1952) 326-336. doi:https://doi.org/10.1112/plms/s3-2.1.326.
- [15] A. Ciabattoni, N. Galatos, K. Terui, Algebraic proof theory: Hypersequents and hypercompletions, Ann. Pure Appl. Logic 168 (3) (2017) 693-737. doi:10.1016/j.apal.2016.10.012.
- [16] N. Galatos, P. Jipsen, T. Kowalski, H. Ono, Residuated lattices: an algebraic glimpse at substructural logics, Vol. 151 of Studies in Logic and the Foundations of Mathematics, Elsevier, 2007.
- [17] D. Brand, P. Zafiropulo, On communicating finite-state machines, J. ACM 30 (2) (1983) 323–342. doi:10.1145/322374.322380.
- [18] P. Schnoebelen, Verifying lossy channel systems has nonprimitive recursive complexity, Information Processing Letters 83 (5) (2002) 251–261. doi:10.1016/S0020-0190(01)00337-4.
- [19] L. E. Dickson, Finiteness of the odd perfect and primitive abundant numbers with n distinct prime factors, American Journal of Mathematics 35 (4) (1913) 413–422.
- [20] S. Schmitz, P. Schnoebelen, Algorithmic Aspects of WQO Theory, lecture (Aug. 2012).
- [21] D. Figueira, S. Figueira, S. Schmitz, P. Schnoebelen, Ackermannian and primitive-recursive bounds with Dickson's Lemma, in: Proceedings - Symposium on Logic in Computer Science, 2011, pp. 269– 278. doi:10.1109/LICS.2011.39.