

Horseshoe-type Priors for Independent Component Estimation *

Jyotishka Datta
Department of Statistics
Virginia Tech

Nicholas G. Polson
Booth School of Business
University of Chicago[†]

September 4, 2024

Abstract

Independent Component Estimation (ICE) has many applications in modern day machine learning as a feature engineering extraction method. Horseshoe-type priors are used to provide scalable algorithms that enables both point estimates via expectation-maximization (EM) and full posterior sampling via Markov Chain Monte Carlo (MCMC) algorithms. Our methodology also applies to flow-based methods for nonlinear feature extraction and deep learning. We also discuss how to implement conditional posteriors and envelope-based methods for optimization. Through this hierarchy representation, we unify a number of hitherto disparate estimation procedures. We illustrate our methodology and algorithms on a numerical example. Finally, we conclude with directions for future research.

Key words: Deep Learning, ICE, factor models, Bayes, Flow models.

*This is a work in progress and has not been peer-reviewed. There may be inadvertent errors.

[†]Jyotishka Datta is Assistant Professor of Statistics at Virginia Tech. Nicholas G. Polson is Professor of Econometrics and Statistics at the Booth School of Business, University of Chicago.

1 Introduction

Linear independent component analysis (ICA) is central to blind source separation and has many fields of application, e.g. signal processing, medical imaging, machine learning and many others. ICA can be viewed as a feature extraction problem where a random vector \mathbf{x} with coordinates that admit a representation as linear combination of independent latent variables s with an unknown mixing matrix \mathbf{A} , namely $x = \mathbf{A}s$. One needs to estimate \mathbf{A} or the unmixing matrix $\mathbf{W} = \mathbf{A}^{-1}$ (Comon et al., 1991; Comon, 1994; Bell and Sejnowski, 1995; Hyvärinen and Pajunen, 1999). ICA can be viewed of as a refinement or an extension of principle component analysis (PCA) or factor analysis, in that the PCA only requires uncorrelated components, not independent.

Feature engineering (a.k.a. nonlinear factor analysis) is a fundamental problem in many modern day machine learning applications ranging from medical imaging to signal processing. A key aspect of feature engineering is decomposing high-dimensional data into independent latent factors. Bhadra et al. (2024) presents a statistical view of high-dimensional deep learning. In this formulation, the output Y and input X connected via a statistical model:

$$P(\mathbf{Y} | \mathbf{W}, \mathbf{X}) = P(\mathbf{Y} | \mathbf{a}), \mathbf{a} = \mathbf{W}\mathbf{X},$$

where, \mathbf{a} is linear factors to be extracted, connected via a hierarchical model with no error in second stage of hierarchy. Linear independent components analysis (ICA) can be useful here as it separates the observed data into a lower-dimensional array of independent sources (a.k.a. features) to offer a high-dimensional probabilistic structure. Thus, ICA falls into the class of high-dimensional data reduction methods and help one identify pivotal data features as well as yield optimal predictive outcomes. A key feature of popular statistical or machine learning models is that they necessitate an approach that concurrently discerns both the foundational independent features and their associated mixing weights, colloquially termed as nonlinear mixing maps.

Our primary objective is to unify an array of algorithmic approaches to allow for both MAP optimization and fully Bayesian estimation. Method of moments estimators are commonplace in this literature and have lead to tensor methods. Building on the seminal work of (MacKay, 1992a, 1996) who showed that the (Bell and Sejnowski, 1995) algorithm for signal processing for a non-linear feedforward network can be viewed as a maximum likelihood algorithm for the optimisation of a linear generative model. Our contribution then is three-fold:

- (i) Develop full Bayesian methods for independent component estimation (ICE).
- (ii) Unify existing models using Gaussian scale mixture representation.
- (iii) Develop Nonlinear deep ICE.

The hallmark of ICA methods is the use of a super-Gaussian distribution over hidden states. Surprisingly, this leads to identification where traditional Gaussian modeling assumptions do not. A linear mixing of Gaussian distribution is itself Gaussian, so de-mixing is impossible. Hence the source distribution must have heavy tails (a.k.a. super-Gaussian). We show that the easiest way to achieve this is via Horseshoe-type priors which are heavy-tailed by construction. Horseshoe priors have been a default class of priors for many problems, see

Generative models have achieved many successful applications ranging from machine learning to image processing. They can be applied in the context of ICA. To fix notation let the data generating model be constructed as follows: data \mathbf{y} is generated from a baseline source distribution, $\mathbf{s} \sim p(\mathbf{s})$ via a latent state x which is a ridge function of \mathbf{s} , namely $g(\mathbf{A}\mathbf{s} + \mathbf{b})$ for an activation

function $g(\cdot)$ and then,

$$\mathbf{x} = g(\mathbf{A}\mathbf{s} + \mathbf{b}), \text{ where } \mathbf{s} \sim p(\mathbf{s}).$$

The mixing matrix \mathbf{A} provides ‘feature extraction’ – a central problem in machine learning. We consider below the special case $g = 1(\cdot)$, the identity map, but g could be a deep learner with many layers. Training high-dimensional generative models such as LLMs (large language models) is extremely costly and novel scalable algorithms to help in this task are an active area of interest. We refer the readers to [Polson and Sokolov \(2020\)](#); [Tran et al. \(2020\)](#) for a comprehensive review of the computational aspects of deep learning.

Our work builds on the seminal work of [MacKay \(1992a, 1996\)](#) who interprets ICA as a latent variable modeling strategy. [MacKay \(1996\)](#) provides instances of many statistical models (e.g. mixture models, hidden Markov models, factor analysis or Helmholtz machines) that are generative models with a layer of latent variables that are usually modeled with simple, separable distributions. Across each of these modeling frameworks, learning the latent variables is tantamount to describing the observables in terms of independent components, and hence, it is natural to expect that ICA should also admit a generative latent variable representation. With this interpretation, [MacKay \(1992a\)](#) shows that the original blind separation algorithm of [Bell and Sejnowski \(1995\)](#) can be viewed as a maximum a posteriori (MAP) estimator from the marginalised likelihood, where the trick is to write the ICA as a latent variable model with a separable distribution on the hidden states. In this representation, the latent variables are assumed be mutually independent and non-Gaussian, and are called independent components. The key insight then that underlies our unifying approach is that the components of the source distribution can be modeled with scale mixtures of normals ([West, 1987](#); [Barndorff-Nielsen et al., 1982](#)). We show that MacKay’s original model and its extensions are simply scale mixtures with Pólya-Gamma mixing ([Polson et al., 2013](#)) and are thus Horseshoe-type priors ([Polson and Scott, 2012a, 2013](#)).

On the algorithmic side, we show that a number of hitherto disjoint algorithms can be unified as envelope optimization methods (see [Geman and Yang, 1995](#); [Polson and Scott, 2016](#)). For example, the ICA updating scheme due to [Bell and Sejnowski \(1995\)](#) is equivalent to a maximum likelihood approach, see ([MacKay, 1992a, 1996](#)). Auxiliary variable methods allow for both EM and MCMC algorithms to be developed across a wide spectrum of source distributions. Such methods can lead to faster convergence ([Ono and Miyabe, 2010](#)) and can incorporate methods for fast convergence such as Nesterov acceleration and block-coordinate descent, thus providing an alternative to traditional stochastic gradient descent methods.

From a historical perspective, Independent Component Analysis arises as a possible solution to the ‘Blind source separation’ problem, a classical problem in signal processing. In the formulation by ([Herault and Jutten, 1986](#); [Bell and Sejnowski, 1995](#)), algorithms for blind source separation attempt to recover source signals \mathbf{s} from observations \mathbf{x} , where $\mathbf{x} = \mathbf{V}\mathbf{s}$, are linear mixtures with unknown weights \mathbf{V} . This is done by finding a square matrix \mathbf{W} which is the inverse of the mixing matrix \mathbf{V} , up to permutation and change of scale. For example, ([Bell and Sejnowski, 1994, 1995](#)) take an algorithmic approach summarized as a linear mapping $\mathbf{a} = \mathbf{W}\mathbf{x}$ where \mathbf{a} is an estimate of the source signals. This is done by adjusting the unmixing matrix \mathbf{W} to maximize the entropy of the outputs.

The algorithm proceeds iteratively using a gradient ascent method on the log-likelihood of the estimated sources, where the update rule for W is:

$$\nabla W \propto (I - 2\phi(\mathbf{a})\mathbf{a}^T) \mathbf{W}$$

where, I is the identity matrix, and $\phi(\mathbf{a})$ is a non-linear function applied component-wise to \mathbf{a} .

Common choices for f are the logistic sigmoid function or the hyperbolic tangent (i.e., a nonlinear map $\mathbf{z}_i = \phi_i(\mathbf{a}_i)$ where $\phi(\cdot) = -\tanh(\cdot)$).

A Bayesian approach (Févotte et al., 2004; Févotte and Godsill, 2006; Donnat et al., 2019; MacKay, 1992a) has a two-fold advantage. First, rather than casting problem as a method of moments approach (kurtosis), a Bayesian approach finds the solution by a suitable regularization with a heavy-tailed prior (thus imposing constraint on higher order moments). A particular suitable class is super-Gaussian priors (Palmer et al., 2006) via scale mixtures of normals (West, 1987). MacKay (1992a) uses a heavy-tailed source distribution towards this. The advantage is that mixture and envelope methods can be used to extract features/factors and provide fast scalable algorithms for ICA. In this paper, we unify the existing Bayesian algorithms and provide new ones.

Assuming a source distribution, $\mathbf{s} \sim p(\mathbf{s})$ acts as a regularization penalty and allows us to unify existing procedures as MAP estimators. For example, MacKay (1996) shows that if $\phi_i(a_i) = -\tanh(a_i)$ then this is equivalent to a source distribution of the form $p_i(s_i) \propto 1/\cosh(s_i) = 1/(e^{s_i} + e^{-s_i})$. We argue in §2 that this can be written as a Gaussian scale mixture using a Pólya-Gamma mixing density. Moreover, (MacKay, 1992b) suggests adding a gain β and considering a heavy-tailed source distribution of the form $p_i(s_i) \propto 1/\cosh^{1/\beta}(\beta s_i)$. Again we show that this can be represented as a Normal scale mixture thus leading to classes of auxiliary variable methods for inference and optimisation. In the limit as $\beta \rightarrow \infty$ this becomes $p_i(s_i) \propto \exp(-|s_i|)$ the double exponential (a.k.a. Lasso) prior, and as $\beta \rightarrow 0$, $p_i(s_i)$ would converge to a zero-mean Gaussian with variance $1/\beta$.

A related goal is to bring together the modeling of source distributions under the encompassing paradigm of Gaussian scale mixtures (Andrews and Mallows, 1974; West, 1987; Carlin and Polson, 1991). This latent variable or parameter expansion approach offers fast, scalable algorithms akin to the auxiliary variable methods proposed by (Ono and Miyabe, 2010). Furthermore, it obviates dependence on approximative techniques like variational Bayes (VB). Our methodology bears significance for deep Bayesian models wherein latent variable distributions are forged as superpositions of nonlinear mappings, akin to deep learning constructs (Polson and Sokolov, 2017; Polson and Rovcková, 2018).

The rest of our paper is outlined as follows. The next subsection describes connections with previous work. Section 3 provides a discussion of deep Bayesian generative models and provides the latent variable representation for McKay’s algorithm. Section 4 provides a unifying framework for super-Gaussian source distributions. Section 5 discusses envelope methods to provide fast optimisation methods. Section 6.1 illustrates our methodology. Finally, Section 7 concludes with directions for future research.

2 Connections with Previous Work

As Auddy and Yuan (2023) point out, the mixing matrix \mathbf{A} can be identified up to permutation and scaling under the assumption of at most one Gaussian component in § due to a theorem by Comon (1994). The review paper Auddy et al. (2024) demonstrates that this can also be connected with the Kruskal identifiability theorem (Kruskal, 1977) that shows a remarkable difference between matrix and tensors: while a matrix of rank ρ can be decomposed in many ways as a sum of ρ rank-one matrices, but higher-order tensors typically admit unique decomposition into rank-one tensors. In the context of PCA versus ICA, Auddy et al. (2024) observe that the the principal components are singular vectors of covariance matrix, while the independent components are ‘singular vector’s of fourth order cumulant tensors, and hence, PCA can not reconstruct independent random variables with mean zero, unit variance but non-zero excess kurtosis, but ICA can. There have been many

approaches for solving the ICA problem including but not limited to (MacKay, 1992a; Comon, 1994; Bell and Sejnowski, 1995; MacKay, 1996; Deco and Obradovic, 1996; Hyvärinen and Pajunen, 1999; Hastie and Tibshirani, 2002; Ono and Miyabe, 2010; Samworth and Yuan, 2012; Auddy and Yuan, 2023) as well as hierarchical Bayesian approaches such as MacKay (1992a); Févotte et al. (2004); Févotte and Godsill (2006); Karklin and Lewicki (2005); Asaba et al. (2018). We refer the readers to Hyvärinen et al. (2001); Nordhausen and Oja (2018) for comprehensive reviews.

Samworth and Yuan (2012) proposes a nonparametric maximum likelihood approach for estimating the unmixing weights and the marginal distributions by projecting the empirical distributions on the space of log-concave univariate distributions. They also address the question of identifiability of the unmixing matrices, first studied by (Comon, 1994), and in their formulation identifiability up to scaling and permutation is guaranteed by requiring that not more than one of the univariate log-concave projections is Gaussian. This is also connected to Kruskal’s identifiability conditions (Kruskal, 1977) for a tensor product $T = \sum_{r=1}^R A \otimes B \otimes C$, in terms of the Kruskal ranks of the individual matrices: $\kappa_A + \kappa_B + \kappa_C \geq 2R + 2$. Bhaskara et al. (2014) provide a robust version of the identifiability theorem for approximate recovery and point uses in several latent variable models. We refer the readers to the comprehensive review by Auddy et al. (2024) on tensor methods for high-dimensional data analysis.

Camuto et al. (2021) combine a linearly independent component analysis (ICA) with nonlinear bisective feature maps (from flow-based methods). For non-square ICA, they can assume the number of sources is less than the data dimensionality – thus achieving better unsupervised latent factor discovering than other ICA flow-based methods. Dinh et al. (2014) discuss NICE deep generative models with nonlinear invertible neural networks, building on the work of Deco and Obradovic (1996), Obradovic and Deco (1998), Comon et al. (1991) and Pearlmutter and Parra (1996) and Malthouse (1998). Recent work includes Bayesian ICA models of (Donnat et al., 2019). Other related approaches include auto-encoder and sparsity models.

Nonlinear ICA models have been proposed by Hyvärinen and Pajunen (1999), although identification can be challenging. Khemakhem et al. (2020) provides recent identification results, providing mild conditions under which the joint distribution encompassing both observed and latent variables within Variational Autoencoders (VAEs) are identifiable and estimable, thus establishing a connection between VAEs and nonlinear Independent Component Analysis (ICA). The source ξ is a feature vector that needs to be learned, see, for example, Olshausen and Field (1996, 1997) for sparse coding of natural images, where an image is modeled as a natural superposition using an over-complete basis set where the amplitudes are given sparsity-inducing prior distributions. This is based on the intuition of Barlow’s principle of redundancy reduction (Barlow, 1989, 2001).

Other popular approaches for dimension reduction include Sliced Inverse Regression (Li, 1991) that finds a low-dimensional projection of the data that captures the most relevant information for explaining the variation in the data, specifically designed for non-linear relationships in the data, Unlike traditional linear dimensionality reduction methods like Principal Component Analysis (PCA). Lopes et al. (2012) introduces a sequential online strategy for efficient posterior simulation. Finally, as noted by Brillinger (2012) and Naik and Tsai (2000), the mixing matrix can be consistently estimated through PLS, regardless of the activation function’s nonlinearity, albeit with a proportionality constant. While the assumption by Brillinger (2012) of Gaussian input \mathbf{X} is necessary for applying Stein’s lemma, we note that this outcome extends to scale-mixtures of Gaussians.

NICE (non-linear independent component estimation) Dinh et al. (2014) provide a deep learning framework called the for high-dimensional density estimation, followed by the real NVP (Dinh et al., 2016) transformations for unsupervised learning. The real NVP method learns a stable and

invertible bijective function or map between samples $x \sim p_X$ and latent space $s \sim p_S$ or, $\theta \sim p_\theta$. For example, [Trippe and Turner \(2018\)](#) utilize normalizing flows as likelihoods for conditional density estimation for complex densities. [Jimenez Rezende and Mohamed \(2015\)](#) provide an approximation framework using a series of parametric transformations for complex posteriors. A new method for Monte Carlo integration called Neural Importance Sampler was provided by [Müller et al. \(2019\)](#) based on the NICE framework by parametrizing the proposal density by a collection of neural networks.

Invertible Neural Network: An important concept in the context of generative models is an invertible neural network or INNs ([Dinh et al., 2016](#)). Loosely speaking, an INN is a one-to-one function with a forward mapping $f: \mathbb{R}^d \mapsto \mathbb{R}^d$, and its inverse $g = f^{-1}$. [Song et al. \(2019\)](#) provides the ‘MintNet’ algorithm to construct INNs by using simple building blocks of triangular matrices, leading to efficient and exact Jacobian calculation. On the other hand, [Behrmann et al. \(2021\)](#) show that common INN method suffer from exploding inverses and provide conditions for stability of INNs. For image representation, [Jacobsen et al. \(2018\)](#) introduce a deep invertible network, called the i-revnet, that retains all information from input data up until the final layer.

Flow transformation models: Here $\mathbf{Y} = h(\mathbf{X})$ where $h(\cdot)$ is typically modeled as an invertible neural network (INN), with both $p_Y(\cdot)$ and $p_F(\cdot)$ as Gaussian densities,

$$\begin{aligned} p(\mathbf{Y}, \mathbf{X} \mid \xi) &= p(\mathbf{Y} \mid \mathbf{X}, \xi) p(\mathbf{X} \mid \xi) \\ &= p_y(\mathbf{Y} \mid h^{-1}(\mathbf{X}), \mathbf{s}) p_F(h^{-1}(\mathbf{X})) \left| \frac{\partial h^{-1}}{\partial \mathbf{X}} \right| \end{aligned}$$

where the determinant of Jacobian is easy to compute. These models can be thought of as latent factor models. Flow-based methods can construct a nonlinear ICA where the dimensionality of the latent space is equal to the data as in an auto-encoder approach, see [Camuto et al. \(2021\)](#).

Latent Factor Model: Another interesting class of models contain latent factors that are driven with INNs. These models take the form

$$\mathbf{Y} = \mathbf{F}^T \xi + \epsilon \tag{1}$$

$$\mathbf{X} = h(\mathbf{F}) \tag{2}$$

$$\mathbf{F} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p), \xi \sim p(\mathbf{s}) \tag{3}$$

where (\mathbf{Y}, \mathbf{X}) are observed data and ϵ is the mean zero Gaussian noise. Here $h(\cdot)$ is an invertible neural network (INN) and \mathbf{F} are the latent factors. This is essentially a flow transformation model and therefore, we can estimate h and \mathbf{s} using the loss function:

$$L(h, \xi) = \sum_{i=1}^N \left\{ \lambda \|y_i - h^{-1}(x_i)^T \mathbf{s}\|^2 + \|h^{-1}(x_i)\|^2 - \log \left| \frac{\partial h^{-1}}{\partial x} \right| (x_i) \right\} - \log p(\mathbf{s})$$

An iterative two-step minimization procedure to learn h and \mathbf{s} is given by: For $t = 1, 2, \dots$

$$(i) \hat{h}^{(t)} = \hat{h}^{(t-1)} - \eta \nabla L(h, \hat{\xi}^{(t-1)})$$

$$(ii) \hat{\xi}^{(t)} = \arg \min_{\xi} L(\hat{h}^{(t)}, \xi), \text{ or draw samples from the posterior } \propto \exp(-L(\hat{h}^{(t)}, \xi)).$$

HINT (Hierarchical Invertible Neural Transport): [Kruse et al. \(2021\)](#); [Detommaso et al. \(2019\)](#) provide the algorithm for posterior sampling. In this formulation, the function T moves in the

normalizing direction: a complicated and irregular data distribution $p_w(\mathbf{w})$ towards the simpler, more regular or ‘normal’ form, of the base measure $p_z(z)$. Let $\mathbf{w} := [\mathbf{y}, \mathbf{x}] \in \mathbb{R}^{m+d}$ and $T(\mathbf{w}) := [T^y(\mathbf{y}), T^x(\mathbf{x}, \mathbf{y})] \in \mathbb{R}^{m+d}$. The inverse function $S^{-1} = T$ is denoted as $S(\mathbf{z}) := [S^y(\mathbf{z}_y), S^x(\mathbf{z}_x, \mathbf{z}_y)]$ where $S^y = (T^y)^{-1}$ where we assume that $\mathbf{z} \sim \mathcal{N}(0, I_{m+d})$. As $p_z = p_{z_y} p_{z_x|z_y}$ and S^y pushes forward the base density p_{z_y} to p_y , it can be shown that $S^x(\cdot, z_y)$ pushes forward the base density $p_{z_x|z_y}(\cdot|z_y)$ to the posterior density $p_{x|y}(\cdot|y)$, when $z_y = T^y(y)$. To sample from $p_{x|y}$, we simply sample $z_x \sim \mathcal{N}(0, I_d)$ and calculate $x = S^x(z_x, z_y) = S^x(z_x, T^y(y))$, since $p_{z_x|z_y} = p_{z_x}$.

Two popular approaches for generative models that rely on latent space representation of the input data, albeit using substantially different architecture) are VAE (variational auto-encoders) and GAN (generative adversarial networks), but the iterative algorithms are only approximate based on a Kullback–Leiber divergence based approximation. On the theoretical side, [Wang et al. \(2023\)](#) provide exact proximal algorithms based on EM and MCMC algorithms. There are many directions for future work, particularly extensions to fields where traditional statistical methods dominate such as spatial or spatiotemporal data analysis.

3 Deep Independent Components Estimation (DICE)

3.1 The generative model and the likelihood function

We begin with linear independent component models. The goal of ICA is to attempt to recover source signals \mathbf{S} from observations \mathbf{X} which are linear mixtures (with unknown coefficients \mathbf{A}) of the source signals, i.e., $\mathbf{X} = \mathbf{A}\mathbf{S}$, where $\mathbf{W} = \mathbf{A}^{-1}$. This can be interpreted as a Bayesian hierarchical model with a degenerate first stage. We follow [MacKay \(1996\)](#)’s formulation here. We observe data as N observations $\mathbf{x}^{(n)}$, $1 \leq n \leq N$ which are linear mixtures of sources $\mathbf{s}^{(n)}$. Let $\mathbf{s}^{(n)} \in \mathbb{R}^d$, $1 \leq n \leq N$ denote the set of sources, that are independently distributed with marginal density $p_i(s_i^{(n)})$.

$$\mathbf{x}^{(n)} = \mathbf{A}\mathbf{s}^{(n)} \quad \text{with} \quad p(\mathbf{s}^{(n)}) = \prod_{i=1}^d p_i(s_i^{(n)}).$$

where \mathbf{A} is the mixing matrix. We wish to estimate $\mathbf{W} = \mathbf{A}^{-1}$. As stated before, the sources $\mathbf{s}^{(n)}$ have an independent components distribution and can be viewed as latent variables. The joint probability of the observed and the hidden latent variables can be written as:

$$\begin{aligned} p(\{\mathbf{x}^{(n)}\}_{n=1}^N, \{\mathbf{s}^{(n)}\}_{n=1}^N | \mathbf{A}) &= \prod_{n=1}^N p(\mathbf{x}^{(n)} | \mathbf{s}^{(n)}, \mathbf{A}) p(\mathbf{s}^{(n)}) \\ &= \prod_{n=1}^N \delta \left(\prod_i (x_i^{(n)} - \sum_j A_{ij} s_j^{(n)}) \right) \prod_j p_j(s_j^{(n)}) \end{aligned} \quad (4)$$

[MacKay \(1996\)](#) points out that it is straightforward to replace $\delta(x_j^{(n)} - A_{ji} s_i^{(n)})$ with a probability distribution over $x_j^{(n)}$ with mean $A_{ji} s_i^{(n)}$, but we need to assume \mathbf{x} is generated without noise to get the [Bell and Sejnowski \(1995\)](#) algorithm.

To access a wider range of probability densities, we can introduce auxiliary variables λ where,

$$p(\mathbf{s}^{(n)}) = \int p(\mathbf{s}^{(n)} | \lambda) p(\lambda) d\lambda.$$

Gaussian scale mixtures (West, 1987; Bhadra et al., 2016) encompasses a wide range of commonly used prior distributions in Bayesian literature, and is also a source of constructing newer priors for handling data with sparsity or other structures. In fact, it is easy to see that the existing prior distributions for Bayesian ICA, viz., student's t (Févotte et al., 2004), Jeffrey's prior $p(s_i) \propto 1/|s_i|$ (Févotte and Godsill, 2006), or Laplace (Asaba et al., 2018), are all Gaussian scale mixtures. We will argue later in the section that one can recover MacKay (1992a)'s hyperbolic secant distribution as a Gaussian scale mixture, too. This has many theoretical and practical advantages. For example, in designing computational algorithms, we can develop EM and MCMC algorithms using the joint posterior $p(\mathbf{s}^{(n)}, \lambda | \mathbf{x})$. We also know a great deal about the behavior of Gaussian scale mixtures due to the works of Barndorff-Nielsen et al. (1982) and others.

To obtain the maximum likelihood estimator, we first observe that the likelihood can be obtained as a product of the following factors from (4), for $n = 1, \dots, N$.

$$p(\mathbf{x}^{(n)} | \mathbf{A}, \mathbf{H}) = \int p(\mathbf{x}^{(n)} | A, \mathbf{s}^{(n)}) p(\mathbf{s}^{(n)}) d\mathbf{s}^{(n)} = \int \prod_i \delta(x_i^{(n)} - A_{ij} s_j^{(n)}) \prod_i p_i(s_i^{(n)}) d\mathbf{s}^{(n)}$$

We use the summation convention like (MacKay, 1996), i.e., $A_{ij} s_j^{(n)} \equiv \sum_j A_{ij} s_j^{(n)}$. Now, using the elementary fact that $\int \delta(x - As) f(s) ds = |A|^{-1} f(x/A)$, we obtain the log likelihood for a single term:

$$p(\mathbf{x} | A) = \frac{1}{|A|} \prod_i p_i(A_{ij}^{-1} x_j) = |W| \prod_i p_i(W_{ij} x_j) |W|$$

$$\Rightarrow \log p(\mathbf{x}^{(n)} | A) = \log |W| + \sum_i \log p_i(W_{ij} x_j) \doteq \log |W| + \sum_i \log p_i(a_i), \text{ where } a_i \equiv W_{ij} x_j.$$

We can then optimize the log-likelihood using any gradient or envelope methods. The terms $\phi_i(a_i) = d \log p_i(a_i) / da_i$ are key terms here that indicate the gradient direction for maximum likelihood. MacKay (1996) shows how to interpret the heuristic algorithms of Bell and Sejnowski (1995) as a MAP estimation procedure from this marginal likelihood. The above formulation assumes \mathbf{x} is generated without noise, as in the original Bell-Sejnowski formulation (Bell and Sejnowski, 1995). MacKay (1992a) comments that an analogous algorithm results if we replace the $\delta(x_j^{(n)} - \sum_i A_{ji} s_i^{(n)})$ term with a suitable probability density over \mathbf{x} with mean $\sum_i A_{ji} s_i^{(n)}$ and sufficiently small noise variance.

We mention two key points in MacKay (1996) here (Section 2.4 point 2,3):

- (i) Employing a tanh nonlinearity of the form $\phi_i(a_i) = -\tanh(a_i)$ implicitly assumes a probability distribution for latent variables, $p_i(s_i) \propto \frac{1}{\cosh(s_i)} \propto \frac{1}{e^{s_i} + e^{-s_i}}$. This distribution exhibits heavier tails compared to the Gaussian distribution, offering a broader range of potential tail behaviors.
- (ii) Alternatively, by incorporating a tanh non-linearity with a gain parameter β , denoted as $\phi_i(a_i) = -\tanh(\beta a_i)$, the associated probabilistic model varies with β . The resulting distribution is expressed as $p_i(s_i) \propto 1/[\cosh(\beta s_i)]^{1/\beta}$. As β becomes large, the non-linearity converges to a step function, resulting in a Laplace density $p_i(s_i) \propto \exp(-|s|)$. Conversely, as β approaches zero, $p_i(s_i)$ tends towards a Gaussian distribution with a mean of zero and a variance of $\frac{1}{\beta}$.

Remark 1. Time versus Transfer Domain: It is worthwhile to note that separation in transfer domain is equivalent to separation in the time domain due to the one-to-one mapping between the models in time and

transfer domain. Using the notations used in (Févotte and Godsill, 2006; Févotte et al., 2004), the linear instantaneous model in time domain is given by the following model where observations at time t are noisy combination of sources at t , and signals of length n :

$$\mathbf{x}_t = \mathbf{A}s_t + \boldsymbol{\epsilon}_t, \quad t = 0, \dots, N - 1, \quad (5)$$

where, $\mathbf{x}_t = (x_{1t}, x_{2t}, \dots, x_{dt})^T$ are the observation vectors, $\mathbf{s}_t = (s_{1t}, s_{2t}, \dots, s_{nt})^T$ are the sources and $\boldsymbol{\epsilon}_t = (\epsilon_{1t}, \epsilon_{2t}, \dots, \epsilon_{dt})$ are the additive noises. The goal is to estimate \mathbf{s} and \mathbf{A} . We assume that there is a basis $n \times n$ matrix Φ such that sources have a sparse representation on it. The equivalent model on the transfer domain is then,

$$\mathbf{x}^{(n)} = \mathbf{A}\mathbf{s}^{(n)} + \boldsymbol{\epsilon}^{(n)}, \quad n = 0, \dots, N - 1, \quad (6)$$

where, n is the coefficient of the basis in the decomposition. Separation using (5) and (6) are equivalent since Φ is a basis matrix.

3.2 Exponential family representation and the posterior

We show the general derivation for the posterior under the aforementioned hierarchical model for the observables and the latent variables. To reduce the notational clutter, we drop the suffix and adopt summation convention as needed. The key insight here is that the posterior can be easily recovered from an exponential family representation. First note that the likelihood $p(\mathbf{x}|\mathbf{s})$ derived above together with the source distribution $p(\mathbf{s})$ can be combined to form a posterior of \mathbf{s} given \mathbf{x} is given by:

$$\begin{aligned} p(\mathbf{s}|\mathbf{x}) &= \frac{1}{(2\pi\sigma^2)^{q/2}} \exp \left\{ -\frac{1}{2\sigma^2} (\mathbf{x} - \mathbf{A}\mathbf{s})' (\mathbf{x} - \mathbf{A}\mathbf{s}) \right\} p(\mathbf{s}) \\ &= h(\mathbf{s}) \exp \left\{ \boldsymbol{\eta}' \mathbf{s} - \frac{1}{2\sigma^2} \mathbf{x}' \mathbf{x} - \log f(\mathbf{x}) \right\} \\ &= h(\mathbf{s}) \exp \{ \boldsymbol{\eta}' \mathbf{s} - \psi(\boldsymbol{\eta}) \}. \end{aligned}$$

where, the different components or known functions are given by:

$$\boldsymbol{\eta} = \frac{1}{\sigma^2} \mathbf{A}' \mathbf{x} \quad (\text{Canonical parameter})$$

$$\psi(\boldsymbol{\eta}) = \frac{1}{2\sigma^2} \mathbf{x}' \mathbf{x} + \log f(\mathbf{x}) = \frac{\sigma^2}{2} \boldsymbol{\eta}' \mathbf{W} \mathbf{W}' \boldsymbol{\eta} + \log f(\sigma^2 \mathbf{W}' \boldsymbol{\eta}) \quad (\text{Cumulant function}),$$

$$\text{where, } f(\mathbf{x}) = \int h(\mathbf{s}) \exp \left\{ \boldsymbol{\eta}' \mathbf{s} - \frac{1}{2\sigma^2} \mathbf{x}' \mathbf{x} \right\} d\mathbf{s}, \quad \text{and}$$

$$h(\mathbf{s}) = \frac{1}{(2\pi\sigma^2)^{q/2}} \exp \left\{ -\frac{1}{2\sigma^2} \mathbf{s}' \mathbf{A}' \mathbf{A} \mathbf{s} \right\} p(\mathbf{s}).$$

Therefore, we can calculate the posterior mean of \mathbf{s} given \mathbf{x} as:

$$\begin{aligned} \mathbb{E}(\mathbf{s} | \mathbf{x}) &= \frac{\partial \psi(\boldsymbol{\eta})}{\partial \boldsymbol{\eta}} = \mathbf{W} \mathbf{x} + \sigma^2 \mathbf{W} \frac{\partial}{\partial \mathbf{x}} \log f(\mathbf{x}) = \frac{\int h(\mathbf{s}) \exp\{\boldsymbol{\eta}' \mathbf{s}\} \mathbf{s} d\mathbf{s}}{\int h(\mathbf{s}) \exp\{\boldsymbol{\eta}' \mathbf{s}\} d\mathbf{s}} \\ \frac{\partial}{\partial \mathbf{x}} \log f(\mathbf{x}) &= \frac{1}{\sigma^2} \left(\mathbf{A} \frac{\int h(\mathbf{s}) \exp\{\boldsymbol{\eta}' \mathbf{s}\} \mathbf{s} d\mathbf{s}}{\int h(\mathbf{s}) \exp\{\boldsymbol{\eta}' \mathbf{s}\} d\mathbf{s}} - \mathbf{x} \right). \end{aligned}$$

We will use this later when deriving a full MCMC algorithms for uncertainty in DICE models.

4 Super-Gaussian Source Distributions

As discussed earlier, it is well known that one needs to use a heavy-tailed super-Gaussian distribution as prior for the sources \mathbf{s} to identify \mathbf{A} . In addition, as [Févotte et al. \(2004\)](#) pointed out, while over-determined ($d \geq n$) case easy: there are many efficient approaches, esp. within Independent Component Analysis, the the general linear instantaneous case, with mixtures possibly noisy and under-determined ($d \leq n$) is challenging. A common approach in such situation is sparsity-Based Blind Source Separation (BSS), especially for under-determined mixtures. The basic premise here is that one exploits source sparsity assumptions: only a few expansion coefficients of sources are significantly different from zero. For example, recommendations in the literature include Student's t or Jeffrey's prior on \mathbf{s} ([Févotte et al., 2004](#); [Févotte and Godsill, 2006](#)) or Laplace [Asaba et al. \(2018\)](#) attempts to do these with commonly used priors. We take a Bayesian shrinkage approach: $\mathbf{s} \sim p(\mathbf{s})$ acts as a regularization penalty and allows us to unify existing procedures as MAP estimators.

4.1 MacKay source distribution: hyperbolic secant

[MacKay \(1996\)](#) shows that $\phi_i(a_i) = -\tanh(a_i)$ is equivalent to a source distribution of the form $p_i(s_i) \propto 1/\cosh(s_i) = 1/(e^{s_i} + e^{-s_i})$. This can be written as a normal scale mixture using a Pólya-Gamma mixing density.

$$s \mid \tau \sim \mathcal{N}(0, 1/4\tau), \tau \sim PG(1, 0), \text{ since } \frac{1}{\cosh(s)} = \int_0^\infty e^{-2\tau t^2} p(\tau) d\tau. \quad (7)$$

due to the following equation:

$$\frac{1}{\cosh(s)} = \int_0^\infty e^{-2\tau t^2} p(\tau) d\tau. \quad (8)$$

Therefore, the location parameter \mathbf{s} is also a normal scale mixture:

$$\mathbf{s} \mid v, \tau \sim \mathcal{N}\left(0, \frac{v^2}{4\tau}\right), v \sim p(v), \tau \sim PG(1, 0) \quad (9)$$

To see this, recall the Pólya-Gamma density is defined as:

Definition 2. Suppose $b \geq 0$, and $z \geq 0$. The Pólya-Gamma distribution $PG(\beta)$ is defined as the density p_{PG} on \mathbb{R}^+ that has the following Laplace transform:

$$\cosh^{-\beta} \left(\sqrt{\frac{t}{2}} \right) = \int_0^\infty \exp(-tx) p_{PG}(x \mid \beta) dx.$$

By reparameterizing $t \mapsto 2\tau^2$, we get the normal scale mixture results. We note that, this is connected to the Jacobi theta distributions studied by [Biane et al. \(2001\)](#); [Devroye \(2009\)](#): J^* has a Jacobi distribution if,

$$J^* \stackrel{D}{=} \frac{2}{\pi^2} \sum_{k=1}^{\infty} \frac{e_k}{(k - \frac{1}{2})^2}, e_k \sim \mathcal{Exp}(1),$$

and the moment generating function is:

$$\mathbb{E}(e^{-tJ^*}) = \frac{1}{\cosh \sqrt{2t}}.$$

The $\cosh(\cdot)$ priors have a connection with the popular horseshoe priors:

$$\beta_j \sim \mathcal{N}(0, \lambda_j^2 \tau^2 \sigma^2), \lambda_j \sim p(\lambda_j) d\lambda_j, \tau \sim p(\tau) d\tau.$$

The horseshoe prior places a standard half-Cauchy distribution over λ_j . This induces an unstandardized unit hyperbolic secant distribution over $\xi_j = \log(\lambda_j)$ (see [Polson and Scott, 2012b](#), section 5).

$$p_{HS}(\xi_j) = \frac{1}{\pi} \frac{1}{\cosh(\xi_j)}.$$

Such log-scale shrinkage priors can unify commonly used continuous shrinkage priors, as shown in ([Schmidt and Makalic, 2018](#)). This also provides a link with the non-Gaussian tail behaviour for blind source separation and that of sparse signal extraction. A similar framework to horseshoe and optimal rates of reconstruction is an area of future research.

Other proposals include the scaled cosh ([MacKay, 1996](#)): $p(s) \sim 1/\cosh^{1/\beta}(\beta s)$, which has the interesting property that,

$$p_i(s_i|\beta) \begin{cases} \exp(-|s_i|) & \beta \rightarrow \infty \text{ (Bayesian Lasso)} \\ \mathcal{N}(0, 1/\beta) & \beta \rightarrow 0. \text{ (Bayesian Ridge)}. \end{cases}$$

That is, in the limit $\beta \rightarrow 0$ or $\rightarrow \infty$, we get Lasso or its Bayesian analog ([Tibshirani, 1996](#); [Park and Casella, 2008](#)), as shown in ([MacKay, 1996](#)).

4.2 Gibbs' sampling strategies

The general scheme for Gibbs sampling follows the following steps. First, from we can write the updates as:

$$p(\mathbf{s} | \tau, \mathbf{A}, \mathbf{x}) = \prod_i p(s_i | \tau_i, \mathbf{A}, \mathbf{x})$$

where $p(s)$ is $\mathcal{N}(\mu_s, \Sigma_s)$ and $\mu_s = \frac{1}{\sigma^2} \sum_s A^T x_i$ and $\Sigma_s = (\frac{1}{\sigma^2} A^T A + \text{diag}(\tau_i)^{-1})^{-1}$. Then, sampling would follow the steps iteratively: (1) update \mathbf{A} : the rows of A are independent a posterior, and (2) update τ : $p(\tau | \mathbf{s})$. We can also link the posterior of \mathbf{A} , viz. , $p(\mathbf{A} | \mathbf{x}, \sigma) = \int p(\mathbf{a}, \tau | \mathbf{x}, \sigma) d\tau$, with the MacKay update equation:

$$\Delta(A^{-1}) \propto A - \tanh(A^{-1}x) \cdot x.$$

The full MCMC algorithm can be derived as follows: Let i be observation index, k the row index of \mathbf{A} , j the column index of \mathbf{A} . $\mathbf{X} = \mathbf{A}^T + \epsilon$ where \mathbf{X} is $n \times d$ and \mathbf{A} is $d \times d$. For the i -th

observation, $\mathbf{x}_i = \mathbf{A}s_i$.

$$\begin{aligned} s_{ij} \mid \mathbf{x}, \mathbf{A}, \tau, s_{ij'} &\sim \mathcal{N}\left(\frac{(\mathbf{X}\mathbf{A} - \xi\mathbf{A}^T\mathbf{A} + S \cdot \text{diag}(\mathbf{A}^T\mathbf{A}))_{ij}}{\text{diag}(\mathbf{A}^T\mathbf{A})_{jj} + 4\tau_{ij}^2\sigma^2}, \frac{\sigma^2}{\text{diag}(\mathbf{A}^T\mathbf{A})_{jj} + 4\tau_{ij}^2\sigma^2}\right) \\ \mathbf{A}_{kj} \mid x, s, \tau, \mathbf{A}_{kj'} &\sim \mathcal{N}\left(\frac{(\mathbf{X}^T\xi - \mathbf{A}\xi^T S + \mathbf{A} \cdot \text{diag}(\xi^T\xi))_{kj}}{\text{diag}(\xi^T\xi)_{jj} + (\sigma/\sigma_2)^2}, \frac{\sigma^2}{\text{diag}(\xi^T\xi)_{jj} + (\sigma/\sigma_2)^2}\right) \\ \tau_{ij} \mid \mathbf{X}, \xi, \mathbf{A} &\sim PG(1, |2s_{ij}|). \end{aligned}$$

Jeffreys prior This corresponds to the improper source distribution $p(s) \sim 1/|s|$. See (Févotte and Godsill, 2006). Févotte and Godsill (2006) derive the EM algorithm for MAP estimation under the Jeffrey's prior ($p(s) \propto 1/|s|$) using the following Normal scale mixture representation:

$$\frac{1}{|s_{ik}|} = \int \mathcal{N}(s_{ik} \mid 0, v_{ik}) \frac{1}{v_{ik}} dv_{ik}$$

The prior $p(v_{ik}) = 1/v_{ik}$ can be thought of as a limiting case of the Inverse Gamma prior, as both parameters of inverse gamma approaches zero, as $p(v) \propto v^{-(a+1)}e^{b/v} \rightarrow 1/v$ as $a, b \rightarrow 0$. It then follows that the prior $p(s_{ik}) = 1/|s_{ik}|$ can be derived as a limiting case of the t -prior in Févotte et al. (2004).

Student's t prior (Févotte et al., 2004) In the case of noise, $\mathbf{x}_j = \mathbf{A}_j\mathbf{s}_j + \boldsymbol{\epsilon}_j$, $j = 1, \dots, N$ we still have a scale mixture of Gaussians framework – and EM algorithm can be found. For example, Févotte et al. (2004) provides MCMC algorithms by iterating the conditionals where each s_i has a $t(\alpha_i, \lambda_i)$ prior, with d.f. α_i and scale λ_i , which can be written as a Normal scale mixture of Inverse Gamma distribution. Specifically, Févotte et al. (2004) uses the representation:

$$p(s_{ij} \mid v_{ij}) = \mathcal{N}(s_{ij} \mid 0, v_{ij}), \text{ and } p(v_{ij} \mid \alpha_i, \lambda_i) = \mathcal{IG}(v_{ij} \mid \alpha_i/2, 2/\alpha_i\lambda_i^2).$$

The Gibbs sampler for Févotte et al. (2004) would be as follows:

- Let $\Sigma_{s_j} = (\frac{1}{\sigma^2}\mathbf{A}^T\mathbf{A} + \text{diag}(v_j)^{-1})^{-1}$, $\mu_{s_j} = \frac{1}{\sigma^2}\Sigma_{s_j}\mathbf{A}^T\mathbf{x}_j$, then,

$$\mathbf{s} \mid \mathbf{A}, \sigma, \mathbf{v}, \alpha, \lambda \sim \prod_{j=1}^K \mathcal{N}(\mathbf{s}_j \mid \mu_{s_j}, \Sigma_{s_j}).$$

- Let $\Sigma_a = \sigma^2(\sum_{j=1}^K S_j^T S_j)^{-1}$, $\mu_a = \frac{1}{\sigma^2}\Sigma_a \sum_j S_j^T \mathbf{x}_j$, then,

$$\mathbf{a} \mid s, \sigma, \mathbf{v}, \boldsymbol{\alpha}, \boldsymbol{\lambda} \sim \mathcal{N}(\mathbf{a} \mid \mu_a, \Sigma_a).$$

- Let $\gamma_\sigma = dN/2$ and $\beta_\sigma = 2/\sum_j \|\mathbf{x}_j - \mathbf{A}s_j\|_F^2$.

$$\sigma \sim \sqrt{\mathcal{IG}(\gamma_\sigma, \beta_\sigma)}.$$

- Let $\gamma_{v_i} = (\alpha_i + 1)/2$, and $\beta_{v_i,t} = 2/(s_{it}^2 + \alpha_i\lambda_i^2)$.

$$\mathbf{v} \mid \sim \prod_{j=1}^K \prod_{i=1}^n \mathcal{IG}(v_{i,t} \mid \gamma_{v_i}, \beta_{v_i,t}).$$

- Finally, λ can be drawn from $\mathcal{G}(\gamma_{\lambda_i}, \beta_{\lambda_i})$, where $\gamma_{\lambda_i} = \alpha_i N/2$, and $\beta_{\lambda_i} = 2/(\alpha_i S_i)$, and since the conditional distribution of α is not available in a closed form but known up to a constant, one can use any approximation given that the individual values of the hyperparameter are not very critical for the algorithm.

Other source distributions There have been a number of other source distributions analysed in the literature. For example, finite mixtures of the form

$$p(s) = \prod_{i=1}^S \sum_{\lambda_i=1}^{M_i} p(s_i(\lambda_i))p(\lambda_i)$$

where $p(\lambda) \sim \text{Gamma}(r, c)$. See the algorithm for constrained EM for ICA ([Hinton et al., 2001](#)).

5 Envelope Optimization Methods

5.1 Mixtures and Envelopes: MCMC and Optimisation

We have two approaches at our disposal: mixtures and envelopes. Their advantages include avoiding approximate estimation techniques, simulation-based inference and prediction and envelope methods for optimisation via Bayes MAP procedures. These two approaches correspond to the two following representations:

$$p(x) = \int p(x, \lambda) d\lambda \quad (\text{mixture})$$

$$p(x) = \sup_{\lambda > 0} \{p(x, \lambda)\} \quad (\text{envelope}).$$

Here, λ is an auxiliary variable ([Geman and Yang, 1995](#)). Before we proceed, let us introduce a few basic notations and results. First, for a closed convex function $\theta(x) : \mathbb{R}^n \mapsto \bar{R}$, the convex conjugate is defined as $\theta^*(\lambda) = \sup_x \{\lambda^T x - \theta(x)\}$, and by the Fenchel-Moreau theorem, the dual relationship holds, i.e. $\theta(x) = \sup_{\lambda} \{\lambda^T x - \theta^*(\lambda)\}$ ([Polson and Scott, 2016](#)). The next result needed is Theorem 2 from ([Polson and Scott, 2016](#)), stated below:

Theorem 3 ([Polson and Scott, 2016](#)). *Let $p(x) \propto \exp\{-\phi(x)\}$ is symmetric in x , and $\theta(x) = \phi(\sqrt{2x})$ for $x > 0$, with completely monotone derivative. Then $p(x)$ admits both normal scale mixture and envelope representation.*

$$\exp -\phi(x) \propto \int_{\mathbb{R}^+} \mathcal{N}(x | 0, \lambda^{-1}) p_I(\lambda) d\lambda \propto \sup_{\lambda \geq 0} \{\mathcal{N}(x | 0, \lambda^{-1}) p_V(\lambda)\}, \quad (10)$$

where, $p_V(\lambda)$, the variational prior is $p_V(\lambda) \propto \lambda^{-\frac{1}{2}} \exp\{\theta^*(\lambda)\}$, and any optimal value of λ would either lie in the subdifferential of $\theta(x^2/2)$ or satisfy: $\hat{\lambda}(x) = \phi'(x)/x$.

The envelope approach generalises the ([Ono and Miyabe, 2010](#)) approach. This leads to an iterative algorithm from the recursions:

$$x^{(t+1)} = \arg_x \min_{\lambda} \left\{ (y - x)^2 + \lambda^{(t)} x^2 \right\},$$

$$\lambda^{(t+1)} = \phi'(x^{(t+1)})/x^{(t+1)}, \quad t = 1, 2, \dots$$

Consider now the envelope representation:

$$p(x) = \sup_{\lambda > 0} p(x, \lambda)$$

where λ is an auxiliary variable (Geman and Yang, 1995). The key result for $1/\cosh$ is given in Polson and Scott (2016):

$$\frac{1}{\cosh(x)} \propto \int_0^\infty \mathcal{N}(x|0, \lambda^{-1}) \cdot p_I(\lambda) d\lambda = \sup_{\lambda \geq 0} \{\mathcal{N}(x|0, \lambda^{-1}) \cdot p_V(\lambda)\} \quad (11)$$

where $p_I(\lambda) \sim 4 \cdot PG(1, 0)$ is the density of Pólya-Gamma distribution. This establishes the link with Proximal Algorithms, sub-differentials or EM if differentiable, and can be used to speed up ICA algorithms. The gradient-based nature avoids matrix inversion and auxiliary variables dynamically adjusts step-sizes, i.e. it lets you perform large step sizes at the beginning, leading to a fast convergence.

An example is given below:

$$\log \cosh(x/2) = \inf_{\lambda \geq 0} \left\{ \frac{\lambda}{2} x^2 - \theta^*(\lambda) \right\} \quad (12)$$

$$\hat{\lambda}(x) = \mathbb{E}_{PG}(\lambda|x) = \frac{1}{2x} \tanh\left(\frac{x}{2}\right) \quad (13)$$

where, we have utilized the fact that $\log \cosh$ is a scale mixture of Pólya-Gamma distributions. We have a sum of $\log \cosh$:

$$\sum_{n=1}^N \sum_{k=1}^K \log \cosh(w_{ij} x_j).$$

For statistical properties of the $1/\pi \cosh(x)$ function as a probability density and the $\log \cosh$ as an alternative loss function, we refer the readers to (Saleh and Saleh, 2022).

Bouchard (2007) obtains an upper bound in two steps: first by bounding sum of exponentials by product of sigmoids and then using a standard quadratic bound for the logarithm of sigmoids. For a vector \mathbf{x} of size n , we have:

$$\log \sum_{i=1}^n e^{x_i} \leq \alpha + \sum_{i=1}^n \log(1 + e^{x_i - \alpha}), \text{ for any } \alpha \in \mathbb{R}. \quad (14)$$

Then one applies the standard quadratic tight bound for $\log(1 + e^x)$ (Jaakkola and Jordan, 1997), given by:

$$\log(1 + e^x) \leq \lambda(\xi)(x^2 - \xi^2) + \frac{1}{2}(x - \xi) + \log(1 + e^\xi), \forall \xi \in \mathbb{R},$$

where $\lambda(\xi) = \frac{1}{2\xi} \left(\frac{1}{1+e^{-\xi}} - \frac{1}{2} \right)$ ($= \frac{1}{4\xi} \tanh(\frac{\xi}{2})$). Equality holds when $x = \xi$. Applying this to (14), one gets,

$$\log \sum_{i=1}^n e^{x_i} \leq \alpha + \sum_{i=1}^n \lambda(\xi_i) \{ (x_i - \alpha)^2 - \xi_i^2 \} + \frac{1}{2} (x_i - \xi_i - \alpha) + \log(1 + e^{\xi_i}). \quad (15)$$

5.2 Auxiliary-Function-Based EM Algorithm

The EM algorithm alternates between an E -step (expectation) and an M -step (maximisation) defined by

$$\begin{aligned} \text{E step : } Q(W|W^{(g)}) &= \int \log p(\mathbf{W}|\lambda, \mathbf{x}) p(\lambda|\mathbf{W}^{(g)}, \mathbf{x}) d\lambda \\ \text{M step : } W^{(g+1)} &= \arg \max_{\mathbf{W}} Q(\mathbf{W}|\mathbf{W}^{(g)}) \end{aligned}$$

From before, the log-likelihood is given by the following objective function:

$$l(\mathbf{W}) = \log |\mathbf{W}| - \sum_i \hat{E} [\phi(W_{ij}x_j)] \quad (16)$$

where \hat{E} denote the sample average. When $p(s) \propto 1/\cosh(s)$, then $\phi(s) = \log \cosh(s)$.

[MacKay \(1996\)](#) directly solves the optimization problem with natural gradient descent algorithm. We now exploit the auxiliary variable scale mixture of normals representation and show that this faster algorithm is equivalent to that of [Ono and Miyabe \(2010\)](#). The complete-data posterior is given by:

$$\begin{aligned} p(\mathbf{W}, \boldsymbol{\lambda}|\mathbf{x}) &= |\mathbf{W}|^N \cdot \prod_{n=1, i=1}^{N, S} \frac{1}{\cosh(W_{ij}x_j^{(n)})} \\ &\propto |\mathbf{W}|^N \cdot \prod_{n=1, i=1}^{N, S} \int_0^\infty e^{-\frac{1}{2}(W_{ij}x_j^{(n)})^2 \lambda_{ni}} \sqrt{\lambda_{ni}} \cdot p_I(\lambda_{ni}) d\lambda_{ni}. \end{aligned}$$

E-step: Since the conditional expectation of λ_{ni} is

$$\hat{\lambda}_{ni} = \frac{\phi'(W_{ij}x_j^{(n)})}{W_{ij}x_j^{(n)}} = \frac{\tanh(W_{ij}x_j^{(n)})}{W_{ij}x_j^{(n)}},$$

We calculate the conditional log posterior (up to a scale $1/N$) as

$$Q(\mathbf{W}|\mathbf{W}^{(t)}) = \log |\mathbf{W}| - \sum_i \hat{E} \left[\frac{1}{2} \hat{\lambda}_{ni} (W_{ij}x_j)^2 \right] \quad (17)$$

$$= \log |\mathbf{W}| - \sum_i W_i' \hat{E} \left[\frac{\tanh(W_{ij}^{(t)} x_j)}{2W_{ij}^{(t)} x_j} x x' \right] W_i. \quad (18)$$

$$= \log |\mathbf{W}| - \sum_i W_i' Z_i^{(t)} W_i. \quad (19)$$

where

$$Z_i^{(t)} := \hat{E} \left[\frac{\tanh(W_{ij}^{(t)} x_j)}{2W_{ij}^{(t)} x_j} x x' \right] = \hat{E} \left[\frac{\phi'(W_{ij}^{(t)} x_j)}{2W_{ij}^{(t)} x_j} x x' \right].$$

M-step: Maximizing $Q(\mathbf{W}|\mathbf{W}^{(t)})$ with respect to \mathbf{W} is equivalent to solving the following system of equations

$$\frac{1}{2}Z_i^{(t)}W_i - \frac{\partial}{\partial W_i} \log |\mathbf{W}| = 0, \forall i \quad (20)$$

which reduces to

$$W'_j Z_i^{(t)} W_i = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases} \quad (21)$$

Now we show that Ono's auxiliary-function-based optimization is exactly the same as our latent variable EM described above.

According to Theorem 1, [Ono and Miyabe \(2010\)](#),

$$\phi(s) \leq \frac{\phi'(r)}{2r}|s|^2 + F(r), \forall r \text{ with } F(r) = \phi(r) - \frac{r\phi'(r)}{2} \quad (22)$$

Here equality holds if and only if $r = |s|$. Thus, the objective function (16) satisfies

$$l(\mathbf{W}) = \arg \max_{\mathbf{R}} Q(\mathbf{W}, \mathbf{R}) \quad (23)$$

$$= \arg \max_{\mathbf{R}} \left\{ \log |\mathbf{W}| - \sum_i \hat{E} \left[\frac{\phi'(r_i)}{2r_i} |W_{ij}x_j|^2 + F(r_i) \right] \right\} \quad (24)$$

Then we can maximize $J(\mathbf{W}, \mathbf{R})$ in terms of \mathbf{W} and \mathbf{R} alternatively, the variables being iteratively updated as below:

Step 1: Given $\mathbf{W}^{(t)}$, $\mathbf{R}^{(t+1)} = \arg \max_{\mathbf{R}} J(\mathbf{W}^{(t)}, \mathbf{R})$.

The solution is $r_i^{(t+1)} = \hat{E} [W_{ij}^{(t)} x_j]$, which results in

$$J(\mathbf{W}, \mathbf{R}^{(t+1)}) = \log |\mathbf{W}| - \sum_i W'_i Z_i^{(t)} W_i + C^{(t)} \quad (25)$$

where $C^{(t)}$ is a constant independent of \mathbf{W} . This is essentially the same of E-step above.

Step 2: Given $\mathbf{R}^{(t+1)}$, $\mathbf{W}^{(t+1)} = \arg \max_{\mathbf{W}} J(\mathbf{W}, \mathbf{R}^{(t+1)})$. That is,

$$\mathbf{W}^{(t+1)} = \arg \max_{\mathbf{W}} \left\{ \log |\mathbf{W}| - \sum_i W'_i Z_i^{(t)} W_i \right\} \quad (26)$$

This is the same as M-step.

6 Applications

6.1 Numerical Experiments

We fix $n = 500$, $d = 4$, $\epsilon \sim \mathcal{N}(0, \sigma^2)$, $v_{ij} \sim \mathcal{N}(0, \sigma^2)$. The data generating process can be described as follows:

$$\begin{aligned}
v &\sim \mathcal{N}(0, \sigma_2^2), \quad v \in \mathbb{R}^{d \times d} \\
\tau &\sim PG(b = 1, c = 0), \quad \tau \in \mathbb{R}^{n \times d} \\
s &\sim \mathcal{N}\left(0, 1/\sqrt{4\tau}\right), \quad s \in \mathbb{R}^{n \times d} \\
x &\sim \mathcal{N}(0, \sigma^2) + s \cdot v^T, \quad x \in \mathbb{R}^{n \times d}
\end{aligned} \tag{27}$$

We set $\sigma = 0.01$ and $\sigma_2 = 1$ for the first experiment and compare Mackay's original method with the EM algorithm described here. The densities of \hat{s} and s for the candidate methods are shown in Fig. 1, and the correlations are shown in Figure 2. EM and MacKay's algorithms seem to have similar performance in terms of recovering signals.




Figure 1: Comparison of the densities for \hat{s} and s




Figure 2: Correlations between \hat{s} and s

Now, we scale the first τ by 100x and setting $\sigma = 0.1$ in the data generating process (27), and repeat the comparisons. Now, both EM algorithm and MacKay's algorithm has difficulty in identifying the first signal but can identify the rest and performs quite similarly. Figures 3 and 4 show the density estimates and the correlations.




Figure 3: Comparison of the densities for \hat{s} and s




Figure 4: Correlations between \hat{s} and s

7 Discussion

High dimensional feature extraction has been a central tool in many modern day applications. Pattern matching methods such as deep learning compose nonlinear maps to find features. Non-linear ICA methods can be notoriously hard to learn and recently combinations of flow-based methods and ICA have been used. We have developed a framework for optimization and posterior simulation using auxiliary-based methods. Our proposed method utilize the normal scale

mixture representation of the $1/\cosh$ function with Pólya-Gammamixing density. This representation makes it possible to implement both MCMC for fully Bayes inference as well as EM for MAP estimation. This representation also establishes connections with extant literature such as (MacKay, 1992a), (Ono and Miyabe, 2010) for auxiliary methods, and Bayesian schemes such as (Févotte et al., 2004), and others. This is a fruitful area for future research and implementation of fully Bayesian methods.

References

- Andrews, D. F. and Mallows, C. L. (1974). Scale mixtures of normal distributions. *J. Roy. Statist. Soc. Ser. B*, 36:99–102.
- Asaba, K., Saito, S., Horii, S., and Matsushima, T. (2018). Bayesian independent component analysis under hierarchical model on independent components. In *2018 Asia-Pacific Signal and Information Processing Association Annual Summit and Conference (APSIPA ASC)*, pages 959–962. IEEE.
- Auddy, A., Xia, D., and Yuan, M. (2024). Tensor methods in high dimensional data analysis: Opportunities and challenges. *arXiv preprint arXiv:2405.18412*.
- Auddy, A. and Yuan, M. (2023). Large dimensional independent component analysis: Statistical optimality and computational tractability. *arXiv preprint arXiv:2303.18156*.
- Barlow, H. (2001). Redundancy reduction revisited. *Network: computation in neural systems*, 12(3):241.
- Barlow, H. B. (1989). Unsupervised learning. *Neural computation*, 1(3):295–311.
- Barndorff-Nielsen, O., Kent, J., and Sørensen, M. (1982). Normal variance-mean mixtures and z distributions. *International Statistical Review*, 50:145–159.
- Behrmann, J., Vicol, P., Wang, K.-C., Grosse, R., and Jacobsen, J.-H. (2021). Understanding and mitigating exploding inverses in invertible neural networks. In *International Conference on Artificial Intelligence and Statistics*, pages 1792–1800. PMLR.
- Bell, A. and Sejnowski, T. J. (1994). A non-linear information maximisation algorithm that performs blind separation. *Advances in neural information processing systems*, 7.
- Bell, A. J. and Sejnowski, T. J. (1995). An information-maximization approach to blind separation and blind deconvolution. *Neural computation*, 7(6):1129–1159.
- Bhadra, A., Datta, J., Polson, N. G., Sokolov, V., and Xu, J. (2024). Merging two cultures: Deep and statistical learning. *Wiley Interdisciplinary Reviews: Computational Statistics*, 16(2):e1647.
- Bhadra, A., Datta, J., Polson, N. G., and Willard, B. (2016). Global-Local Mixtures. *arXiv preprint arXiv:1604.07487*.
- Bhaskara, A., Charikar, M., and Vijayaraghavan, A. (2014). Uniqueness of tensor decompositions with applications to polynomial identifiability. In *Conference on Learning Theory*, pages 742–778. PMLR.
- Biane, P., Pitman, J., and Yor, M. (2001). Probability laws related to the jacobi theta and riemann zeta functions, and brownian excursions. *Bulletin of the American Mathematical Society*, 38(4):435–465.

- Bouchard, G. (2007). Efficient bounds for the softmax function and applications to approximate inference in hybrid models. In *NIPS 2007 workshop for approximate Bayesian inference in continuous/hybrid systems*, volume 6.
- Brillinger, D. R. (2012). A Generalized Linear Model With “Gaussian” Regressor Variables. In Guttorp, P. and Brillinger, D., editors, *Selected Works of David Brillinger*, Selected Works in Probability and Statistics, pages 589–606. Springer, New York, NY.
- Camuto, A., Willetts, M., Roberts, S., Holmes, C., and Rainforth, T. (2021). Towards a theoretical understanding of the robustness of variational autoencoders. In *International Conference on Artificial Intelligence and Statistics*, pages 3565–3573. PMLR.
- Carlin, B. P. and Polson, N. G. (1991). Inference for nonconjugate bayesian models using the gibbs sampler. *Canadian Journal of statistics*, 19(4):399–405.
- Comon, P. (1994). Independent component analysis, a new concept? *Signal processing*, 36(3):287–314.
- Comon, P., Jutten, C., and Herault, J. (1991). Blind separation of sources, part ii: Problems statement. *Signal processing*, 24(1):11–20.
- Deco, G. and Obradovic, D. (1996). *An information-theoretic approach to neural computing*. Springer Science & Business Media.
- Detommaso, G., Kruse, J., Ardizzone, L., Rother, C., Köthe, U., and Scheichl, R. (2019). Hint: Hierarchical invertible neural transport for general and sequential bayesian inference. *stat*, 1050:25.
- Devroye, L. (2009). On exact simulation algorithms for some distributions related to jacobi theta functions. *Statistics & Probability Letters*, 79(21):2251–2259.
- Dinh, L., Krueger, D., and Bengio, Y. (2014). Nice: Non-linear independent components estimation. *arXiv preprint arXiv:1410.8516*.
- Dinh, L., Sohl-Dickstein, J., and Bengio, S. (2016). Density estimation using real nvp. *arXiv preprint arXiv:1605.08803*.
- Donnat, C., Tozzi, L., and Holmes, S. (2019). Constrained bayesian ica for brain connectome inference. *arXiv preprint arXiv:1911.05770*.
- Févotte, C. and Godsill, S. J. (2006). Blind separation of sparse sources using jeffrey’s inverse prior and the em algorithm. In *International Conference on Independent Component Analysis and Signal Separation*, pages 593–600. Springer.
- Févotte, C., Godsill, S. J., and Wolfe, P. J. (2004). Bayesian approach for blind separation of underdetermined mixtures of sparse sources. In *Independent Component Analysis and Blind Signal Separation: Fifth International Conference, ICA 2004, Granada, Spain, September 22-24, 2004. Proceedings 5*, pages 398–405. Springer.
- Geman, D. and Yang, C. (1995). Nonlinear image recovery with half-quadratic regularization. *IEEE transactions on Image Processing*, 4(7):932–946.
- Hastie, T. and Tibshirani, R. (2002). Independent components analysis through product density estimation. *Advances in neural information processing systems*, 15.

- Herault, J. and Jutten, C. (1986). Space or time adaptive signal processing by neural network models. In *AIP conference proceedings*, volume 151, pages 206–211. American Institute of Physics.
- Hinton, G. E., Welling, M., Teh, Y. W., and Osindero, S. (2001). A new view of ica. In *Proceedings of 3rd International Conference on Independent Component Analysis and Blind Signal Separation (ICA'01)*, pages 746–751.
- Hyvärinen, A., Karhunen, J., and Oja, E. (2001). *Independent component analysis, adaptive and learning systems for signal processing, communications, and control*, volume 1. John Wiley & Sons, Inc.
- Hyvärinen, A. and Pajunen, P. (1999). Nonlinear independent component analysis: Existence and uniqueness results. *Neural networks*, 12(3):429–439.
- Jaakkola, T. S. and Jordan, M. I. (1997). A variational approach to bayesian logistic regression models and their extensions. In *Sixth International Workshop on Artificial Intelligence and Statistics*, pages 283–294. PMLR.
- Jacobsen, J.-H., Smeulders, A., and Oyallon, E. (2018). i-revnet: Deep invertible networks. *arXiv preprint arXiv:1802.07088*.
- Jimenez Rezende, D. and Mohamed, S. (2015). Variational inference with normalizing flows. *arXiv e-prints*, pages arXiv–1505.
- Karklin, Y. and Lewicki, M. S. (2005). A hierarchical bayesian model for learning nonlinear statistical regularities in nonstationary natural signals. *Neural computation*, 17(2):397–423.
- Khemakhem, I., Kingma, D., Monti, R., and Hyvarinen, A. (2020). Variational autoencoders and nonlinear ica: A unifying framework. In *International Conference on Artificial Intelligence and Statistics*, pages 2207–2217. PMLR.
- Kruse, J., Detommaso, G., Köthe, U., and Scheichl, R. (2021). Hint: Hierarchical invertible neural transport for density estimation and bayesian inference. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 35, pages 8191–8199.
- Kruskal, J. B. (1977). Three-way arrays: rank and uniqueness of trilinear decompositions, with application to arithmetic complexity and statistics. *Linear algebra and its applications*, 18(2):95–138.
- Li, K.-C. (1991). Sliced inverse regression for dimension reduction. *Journal of the American Statistical Association*, 86(414):316–327.
- Lopes, H. F., Polson, N. G., and Carvalho, C. M. (2012). Bayesian statistics with a smile: A resampling–sampling perspective.
- MacKay, D. J. (1992a). Bayesian interpolation. *Neural computation*, 4(3):415–447.
- MacKay, D. J. (1992b). Information-based objective functions for active data selection. *Neural computation*, 4(4):590–604.
- MacKay, D. J. (1996). Maximum likelihood and covariant algorithms for independent component analysis.
- Malthouse, E. C. (1998). Limitations of nonlinear pca as performed with generic neural networks. *IEEE Transactions on neural networks*, 9(1):165–173.

- Müller, T., McWilliams, B., Rousselle, F., Gross, M., and Novák, J. (2019). Neural importance sampling. *ACM Transactions on Graphics (ToG)*, 38(5):1–19.
- Naik, P. and Tsai, C.-L. (2000). Partial least squares estimator for single-index models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 62(4):763–771.
- Nordhausen, K. and Oja, H. (2018). Independent component analysis: A statistical perspective. *Wiley Interdisciplinary Reviews: Computational Statistics*, 10(5):e1440.
- Obradovic, D. and Deco, G. (1998). Information maximization and independent component analysis: Is there a difference? *Neural Computation*, 10(8):2085–2101.
- Olshausen, B. A. and Field, D. J. (1996). Emergence of simple-cell receptive field properties by learning a sparse code for natural images. *Nature*, 381(6583):607–609.
- Olshausen, B. A. and Field, D. J. (1997). Sparse coding with an overcomplete basis set: A strategy employed by v1? *Vision research*, 37(23):3311–3325.
- Ono, N. and Miyabe, S. (2010). Auxiliary-function-based independent component analysis for super-gaussian sources. In *International Conference on Latent Variable Analysis and Signal Separation*, pages 165–172. Springer.
- Palmer, J. A., Kreutz-Delgado, K., and Makeig, S. (2006). Super-gaussian mixture source model for ica. In *International Conference on Independent Component Analysis and Signal Separation*, pages 854–861. Springer.
- Park, T. and Casella, G. (2008). The bayesian lasso. *Journal of the American Statistical Association*, 103(482):681–686.
- Pearlmutter, B. A. and Parra, L. C. (1996). A context-sensitive generalization of ica. *Advances in neural information processing systems*, 151.
- Polson, N. and Sokolov, V. (2020). Deep learning: Computational aspects. *Wiley Interdisciplinary Reviews: Computational Statistics*, 12(5):e1500.
- Polson, N. G. and Rovcková, V. (2018). Posterior concentration for sparse deep learning. *Advances in Neural Information Processing Systems*, 31.
- Polson, N. G. and Scott, J. G. (2012a). On the half-Cauchy prior for a global scale parameter. *Bayesian Analysis*, 7(4):887–902.
- Polson, N. G. and Scott, J. G. (2012b). On the Half-Cauchy Prior for a Global Scale Parameter. *Bayesian Analysis*, 7(4):887–902.
- Polson, N. G. and Scott, J. G. (2013). Data augmentation for non-Gaussian regression models using variance-mean mixtures. *Biometrika*, 100(2):459–471.
- Polson, N. G. and Scott, J. G. (2016). Mixtures, envelopes and hierarchical duality. *Journal of the Royal Statistical Society. Series B*, 78:701–727.
- Polson, N. G., Scott, J. G., and Windle, J. (2013). Bayesian inference for logistic models using pólya–gamma latent variables. *Journal of the American statistical Association*, 108(504):1339–1349.

- Polson, N. G. and Sokolov, V. (2017). Deep learning: A bayesian perspective. *Bayesian Analysis*, 12(4):1275–1304.
- Saleh, R. A. and Saleh, A. (2022). Statistical properties of the log-cosh loss function used in machine learning. *arXiv preprint arXiv:2208.04564*.
- Samworth, R. J. and Yuan, M. (2012). Independent component analysis via nonparametric maximum likelihood estimation. *The Annals of Statistics*, 40(6):2973–3002.
- Schmidt, D. F. and Makalic, E. (2018). Log-scale shrinkage priors and adaptive bayesian global-local shrinkage estimation. *arXiv preprint arXiv:1801.02321*.
- Song, Y., Meng, C., and Ermon, S. (2019). Mintnet: Building invertible neural networks with masked convolutions. *Advances in Neural Information Processing Systems*, 32.
- Tibshirani, R. (1996). Regression Shrinkage and Selection via the Lasso. *Journal of the Royal Statistical Society. Series B*, 58:267–288.
- Tran, M.-N., Nguyen, N., Nott, D., and Kohn, R. (2020). Bayesian deep net glm and glmm. *Journal of Computational and Graphical Statistics*, 29(1):97–113.
- Trippe, B. L. and Turner, R. E. (2018). Conditional density estimation with bayesian normalising flows. *arXiv preprint arXiv:1802.04908*.
- Wang, Y., Polson, N., and Sokolov, V. O. (2023). Data augmentation for bayesian deep learning. *Bayesian Analysis*, 18(4):1041–1069.
- West, M. (1987). On scale mixtures of normal distributions. *Biometrika*, 74(3):646–648.