

A Strongly Normalising System of Dependent Types for Transparent and Opaque Probabilistic Computation

Francesco A. Genco*

LUCI Lab, Department of Philosophy, University of Milan

June 26, 2024

Keywords. probabilistic computation; non-deterministic computation; opacity; type theory; dependent types

1 Introduction

We define an extension of λ -calculus with dependents types that enables us to encode transparent and opaque probabilistic programs and prove a strong normalisation result for it by a *reducibility* technique. While transparent non-deterministic programs are formalised by rather usual techniques, opaque non-deterministic programs are formalised by introducing in the syntax oracle constants, the behaviour of which is governed by oracular functions. The generality of these functions and the fact that their values are determined by the form of the whole term inside which the relative oracle occurs also enable us to simulate learning-like behaviours.

2 Kinds, types and terms

Definition 2.1 (Typeless Terms). The terms of the calculus are defined by the following grammar:

$$\begin{aligned}
 x, y, z \dots & ::= x_0 \mid x_1 \mid x_2 \dots \\
 o & ::= o_0 \mid o_1 \mid o_2 \dots \\
 t, s, u, v \dots & ::= x \mid o \mid ot \mid o\nu \mid (ot)\nu \mid \\
 & \quad \lambda x : \varphi. t \mid ts \mid \\
 & \quad t(p)s \mid t\nu \mid \\
 & \quad \langle t, s \rangle \mid t\pi_j \mid [\kappa] \mid [\kappa]^p \mid \\
 & \quad [t, [\kappa_1 / \dots / \kappa_n], s] \mid [t, [\kappa_1 / \dots / \kappa_n], s]^p \mid
 \end{aligned}$$

*Work funded by the PRIN project n.2020SSKZ7R BRIO (Bias, Risk and Opacity in AI).

where $n \in \mathbb{N}$, $j \in \{0, 1\}$ and $p \in \{i \mid i \in \mathbb{Q}, 0 \leq i \leq 1\}$

Definition 2.2 (Kindless Type constructors). The terms of the calculus are defined by the following grammar:

$$\begin{aligned} \alpha &::= \alpha_0 \mid \alpha_1 \mid \alpha_2 \dots \\ \varphi &::= \alpha \mid \lambda x : \varphi_1. \varphi_2 \mid \varphi t \mid \forall x : \varphi_1. \varphi_2 \mid \oplus \varphi \mid \Sigma \varphi \end{aligned}$$

The standard part of the grammar for terms includes variables $x_0, x_1 \dots$ for terms, applications and λ -abstractions. Nondeterministic choice terms of the form $t \langle p \rangle s$ are supposed to represent programs that nondeterministically reduce to either t , with probability p , or to s , with probability $1 - p$. The terms of the form $[\kappa]$, where κ is a possibly empty list t_1, \dots, t_n of terms, denote computations represented as reductions of t_i in t_{i+1} for $1 \leq i \leq n$. If a term representing a computation is of the form $[\kappa]^p$, then it also indicates that the computation has probability p of developing as specified by the list κ of terms.

Instead of defining a grammar of types directly, we define, as per standard practice, a grammar of type constructors, that is, syntactic objects that are supposed to evaluate to types. Thus we can handle smoothly the definition of types that depend on terms. According to the grammar, a type constructor can be a type constructor variable $\alpha_1, \alpha_2 \dots$, a λ -abstraction of a type constructor—where the bound variable is a variable for terms—or the application of a type constructor to a term. Type constructor applications are supposed to be evaluated in order to reduce the type constructor to a proper type, which is a type constructor that does not contain occurrences of λ but only occurrences of variables $\alpha_1, \alpha_2 \dots$ and of \forall quantifiers. When we form a type constructor, we can also generalise a type constructor by the \forall quantifier—where the bound variable is a variable for terms. As opposed to the occurrences of λ , the occurrences of these quantifiers are not supposed to disappear during the evaluation of the type constructor. Indeed, λ in a type constructor of the form $(\lambda x : \varphi_1. \forall y : \varphi_1 \alpha xy) t$ is meant to have the resulting type depend on the term t given as an argument to the type constructor, while \forall is meant to stay as a first-order quantifier occurring also in the type resulting from the evaluation of the constructor: $(\lambda x : \varphi_1. \forall y : \varphi_1 \alpha xy) t$ evaluates to $\forall y : \varphi_1 \alpha ty$, which is a type. Notice that the variables $\alpha, \alpha_1, \alpha_2, \dots$ are meant to represent first-order predicates. That is, what we would normally write as $\forall x. \forall y. P(x, y)$ is here written as $\forall x : \varphi_1. \forall y : \varphi_2. \alpha xy$ where α is meant as a type constructor variable corresponding to the predicate symbol P . As specified by the deductive system below, the kind of α will determine its arity.

Type constructors encode functions just as terms do, with the only difference that terms take terms as inputs and yield terms as outputs while type constructors take terms as inputs and yield types as outputs. Therefore, in order to guarantee that their rewriting system is well-behaved, we introduce a system to constrain their usage and obtain a terminating system. Just as we assign types to terms, we assign kinds to type constructors.

The grammar of kinds is rather simple: we have a base case, the constant $*$ denoting the kind of types, and an abstraction Π that binds term variables.¹ For instance

¹Notice that x can never occur as a sub-kind of a kind $\Pi x : \varphi. \Phi$. Hence, quantified kinds of dependent type constructors can be seen as implication kinds of the form $\varphi \Rightarrow \Phi$ (similarly to what we will do for types $\forall x : \varphi_1. \varphi_2$ where x does not occur in φ_2). For instance, all the information provided in the kind of the type constructor $(\forall x : \varphi. \alpha x) : (\Pi x : \varphi. *)$ is also contained in the kind $\varphi \Rightarrow *$ since x does not occur in $*$.

Definition 2.3 (Kinds). The kinds of the calculus are defined by the following grammar:

$$\Phi ::= * \mid \Pi x : \varphi. \Phi$$

We assume that the notions of *free variables*, *bound variables* and substitution are defined as usual, see, for instance, [Sørensen and Urzyczyn, 2006].

Definition 2.4 (Variables occurring in a kind). The term variables occurring in a kind Φ are all the term variables that appear in Φ but not immediately to the right of a Π .

Definition 2.5 (Subconstructors and variables occurring in a type constructor). The subconstructors of a type constructor are inductively defined as follows.

- $\alpha x_1 \dots x_n$ (possibly for $n = 0$) is a subconstructor of $\alpha x_1 \dots x_n$.
- $\lambda x : \varphi_1. \varphi_2$ and all subconstructors of φ_2 are subconstructors of $\lambda x : \varphi_1. \varphi_2$.
- φt and all subconstructors of φ are subconstructors of φt .
- $\forall x : \varphi_1. \varphi_2$ and all subconstructors of φ_2 are subconstructors of $\forall x : \varphi_1. \varphi_2$.
- $\oplus \varphi$ and all subconstructors of φ are subconstructors of $\oplus \varphi$.
- $\Sigma \varphi$ and all subconstructors of φ are subconstructors of $\Sigma \varphi$.

The constructor variables occurring in a type constructor φ are all the constructor variables that appear in φ .

The term variables occurring in a type constructor φ are all the term variables that appear in φ but not immediately to the right of a λ or \forall .

Definition 2.6 (Subterms and variables occurring in a term). The subterms of a term are inductively defined as follows.

- x is a subterm of x , o is a subterm of o .
- $\lambda x : \varphi. t$ and all subterms of t are subconstructors of $\lambda x : \varphi. t$.
- ts and all subterms of t and s are subconstructors of ts .
- $t(p)s$ and all subterms of t and s are subconstructors of $t(p)s$.
- $t\nu$ and all subterms of t are subconstructors of $t\nu$.
- $\langle t, s \rangle$ and all subterms of t and s are subconstructors of $\langle t, s \rangle$.
- $t\pi_j$ and all subterms of t are subconstructors of $t\pi_j$.
- $[\kappa]^p$ (where p might not occur) and all subterms of κ are subconstructors of $[\kappa]^p$.
- $[t, [\kappa_1 / \dots / \kappa_n], s]^p$ (where p might not occur) and all subterms of $t, \kappa_1, \dots, \kappa_n$ and s are subconstructors of $[t, [\kappa_1 / \dots / \kappa_n], s]^p$.

The term variables occurring in a term t are all the term variables that appear in t but not immediately to the right of a λ or \forall .

Definition 2.7 (Binder scope, bound variables, free variables). The *scope* of the displayed occurrence of $\forall x : A$ in $\forall x : A. \varphi$ is φ . The *scope* of the displayed occurrence of $\lambda x : A$ in $\lambda x : A. \varphi$ is φ . The *scope* of the displayed occurrence of $\lambda x : A$ in $\lambda x : A. t$ is t .

An occurrence of the variable $x : A$ in φ is *bound in* φ if it occurs inside the scope of an occurrence of $\lambda x : A$ or $\forall x : A$ in φ . An occurrence of the variable $x : A$ in t is *bound in* t if it occurs inside the scope of an occurrence of $\lambda x : A$ or $\forall x : A$ in t .

A variable occurrence in φ is *free* if it is not bound in φ . A variable occurrence in t is *free* if it is not bound in t .

Definition 2.8 (Substitution). For any two terms t and s and term variable x , the term substitution $t[s/x]$ denotes the term obtained by uniformly and simultaneously replacing all free occurrences of x in t by an occurrence of s .

For any type constructor φ , term t , and term variable x , the type constructor substitution $\varphi[t/x]$ denotes the type constructor obtained by uniformly and simultaneously replacing all free occurrences of x in φ by an occurrence of t .

For any two type constructors φ and ψ , and type constructor variable α , the type constructor substitution $\varphi[\psi/\alpha]$ denotes the type constructor obtained by uniformly and simultaneously replacing all occurrences of α in φ by an occurrence of ψ .

Notice that—since a type constructor variable α is never bound by an occurrence of Π, \forall or λ —type constructor substitutions, such as $[\psi/\alpha]$, are neither employed in deductions nor while normalising type constructors. We only introduce these substitutions as technical devices to define type constructor contexts.

Definition 2.9 (Term contexts and type constructor contexts). A term context $\mathcal{C}[]_1 \dots []_n$ is a term in which, for each $i \in \{1, \dots, n\}$, a designated term variable x_i occurs exactly once. By $\mathcal{C}[]_1 \dots [t]_m \dots []_n$ with $1 \leq m \leq n$ we denote the term obtained by applying the following substitution: $(\mathcal{C}[]_1 \dots []_n)[t/x_m]$.

A type constructor context $\mathcal{T}[]_1 \dots []_n$ is a type constructor in which, for each $i \in \{1, \dots, n\}$, a designated type constructor variable α_i occurs exactly once. By $\mathcal{T}[]_1 \dots [\varphi]_m \dots []_n$ with $1 \leq m \leq n$ we denote the type constructor obtained by applying the following substitution: $(\mathcal{T}[]_1 \dots []_n)[\varphi/\alpha_m]$.

3 The calculus

Notation. We use Γ, Δ to denote lists of type and kind assignments for term and type constructor variables, respectively. We use s, t, u, v as metavariables for terms, A, B, C, D, E, F, T for types, φ, ψ, ξ for type constructors, and Φ, Ψ, Ξ for kinds. We use x, y, z for term variables and α, β, γ for constructor variables. We use κ to denote a list of terms. We add subscripts when necessary.

Definition 3.1 (Oracle constants and oracular functions). Each oracle constant o has either arity 0 or 1, and is associated to a type T and to an oracular function $f_o(,)$.

If o_0 is a 0-ary oracle constant, then $T = \Sigma A$ for any type A and $f_{o_0}(,)$ is any function that accepts as first input a term context $\mathcal{C}[]_1 \dots []_n$ with any number n of holes and as second input a non-zero number $m \leq n$. The output of the function is a closed term of type A that does not contain any oracle constant.

If o_1 is a 1-ary oracle constant, then $T = \forall x : A. \Sigma B$ and $f_{o_1}(,)$ is any function that accepts as first input a term context $\mathcal{C}[]_1 \dots []_n$ with any number n of holes and as second input a non-zero number $m \leq n$. The output of the function is a closed term of type $B[t/x]$ that does not contain any oracle constant, where t is the term to which the m th hole of $\mathcal{C}[]_1 \dots []_n$ is applied.

Oracular functions are used to determine the reducti of oracle constants. In particular, as we will see in Table 5, if the oracular function application $f_o(\mathcal{C}[]_1 \dots []_n, m)$ has as value the term u , then we use u as reductum of o —possibly applied to some argument—when it occurs in the m th hole of the context $\mathcal{C}[]_1 \dots []_n$. Hence, for instance, if o does not accept arguments, then $\mathcal{C}[o]_1 \dots [o]_n$ will reduce to a term of the form $\mathcal{C}[\dots]_1 \dots [u]_m \dots [\dots]_n$ if $f_o(\mathcal{C}[]_1 \dots []_n, m) = u$. Since the values of f_o

$$\frac{\Gamma_1 \vdash \Phi : \square}{\Gamma_1, \alpha : \Phi \vdash \alpha : \Phi} \quad \frac{\Gamma_2 \vdash A : *}{\Gamma_2, x : A \vdash x : A}$$

where α does not occur in Γ_1 and x does not occur in Γ_2

$$\vdash * : \square \quad \frac{\Gamma, x : A \vdash \Phi : \square}{\Gamma \vdash (\Pi x : A. \Phi) : \square} \quad \frac{\Gamma, x : A \vdash B : *}{\Gamma \vdash (\forall x : A. B) : *}$$

$$\frac{\Gamma \vdash A : *}{\Gamma \vdash \oplus A : *} \quad \frac{\Gamma \vdash A : *}{\Gamma \vdash \Sigma A : *} \quad \frac{\Gamma \vdash A : * \quad \Gamma \vdash B : *}{\Gamma \vdash A \wedge B : *}$$

Figure 1: Formation and axiom rules for kinds and types

$$\frac{\Gamma, x : A \vdash \varphi : \Phi}{\Gamma \vdash (\lambda x : A. \varphi) : (\Pi x : A. \Phi)} \quad \frac{\Gamma \vdash \varphi : (\Pi x : A. \Phi) \quad \Gamma \vdash t : A}{\Gamma, \vdash (\varphi t) : \Phi[t/x]}$$

Figure 2: Kind assignment rules

depend on a context representing the whole term in which o occurs, there is no need to specify whether o is applied to some argument or not, this information is already contained in the first argument of the function. So, in case o does accept an argument, then the term $\mathcal{C}[os_1]_1 \dots [os_n]_n = \mathcal{D}[o]_1 \dots [o]_n$ will reduce to a term of the form $\mathcal{C}[\dots]_1 \dots [u]_m \dots [\dots]_n$ if $f_o(\mathcal{D}[\]_1 \dots [\]_n, m) = u$. Thus, in general, we use applications $f_o(\mathcal{C}[\]_1 \dots [\]_n, m)$ of oracular functions in which $1 \leq m \leq n$.

Oracular functions are defined with respect to contexts with possibly several holes because thus we can encode, by using oracle constants, the behaviour of functions implementing learning mechanisms. Indeed, when we reduce a term of the form $\mathcal{C}[os_1]_1 \dots [os_n]_n = \mathcal{D}[o]_1 \dots [o]_n$ to a term of the form $\mathcal{C}[u_1]_1 \dots [u_n]_n$ since $f_o(\mathcal{D}[\]_1 \dots [\]_n, i) = u_i$ for $1 \leq i \leq n$, we have that the values produced by the different terms os_i can be determined by f_o as if they were produced through a learning process.

Definition 3.2 (Conversion equivalence \equiv_β). Let us define the conversion relation for type constructors as follows:

$$\mathcal{T}[(\lambda x : \varphi_1. \varphi_2)t] \mapsto \mathcal{T}[\varphi_2[t/x]]$$

where $\mathcal{T}[\]$ is a type constructor context. Let us, moreover, define \mapsto^* as the reflexive and transitive closure of \mapsto . For any two type constructors φ and φ' , $\varphi \equiv_\beta \varphi'$ if, and only if, either $\varphi \mapsto^* \varphi'$ or $\varphi' \mapsto^* \varphi$.

Since types in our system are constructed by also employing terms, a simple way to guarantee that a type is well-formed is to build it by applying rules similar to those used to construct terms. The rules in Tables 1, 2 and 3 precisely enable us to do so. In particular, the first three rules in Table 1 make it possible, starting from the atomic kind $*$, to construct complex kinds and, starting from an atomic type, to construct

$$\begin{array}{c}
\frac{\Gamma \vdash \Psi : \square \quad \Gamma \vdash \Phi : \square}{\Gamma, \alpha : \Psi \vdash \Phi : \square} \quad \frac{\Gamma \vdash A : * \quad \Gamma \vdash \Phi : \square}{\Gamma, x : A \vdash \Phi : \square} \\
\frac{\Gamma \vdash \Psi : \square \quad \Gamma \vdash \varphi : \Phi}{\Gamma, \alpha : \Psi \vdash \varphi : \Phi} \quad \frac{\Gamma \vdash A : * \quad \Gamma \vdash \varphi : \Phi}{\Gamma, x : A \vdash \varphi : \Phi} \\
\frac{\Gamma \vdash \Psi : \square \quad \Gamma \vdash t : A}{\Gamma, \alpha : \Psi \vdash t : A} \quad \frac{\Gamma \vdash A : * \quad \Gamma \vdash t : B}{\Gamma, x : A \vdash t : B}
\end{array}$$

Figure 3: Weakening rules

complex types by introducing the universal quantifier Π for kinds and \forall for types. Once we have thus constructed a kind Φ , we can derive an axiom-shaped sequent in which a type constructor variable α of kind Φ is added to the context and the same type constructor variable $\alpha : \Phi$ occurs on the right-hand side of the sequent. Similarly, for types, when we have formed a type A , we can derive the axiom-shaped sequent with the variable term $x : A$ both in the context and on the right-hand side of the sequent.

By using these rules we can obtain complex kinds and types from atomic ones, but we can only assign them to atomic objects: type constructor variables such as α and term variables such as x . The kind assignment rules, in Table 2, precisely enable us to form complex type constructors while we assign them a suitable kind. The type assignment rules, in Table 4, enable us to do the same thing but for terms: we can form complex terms by starting from term variables, such as x , while we assign them a suitable type. In simply typed λ -calculus, we only have type assignment rules. Indeed, types in simply typed λ -calculus can be defined without reference to terms and hence there is no need for a mechanism to handle by levels—in a way suitable for a type theory—the definition of types and terms. It is, indeed, the reciprocal dependence of terms from types and of types from terms that makes it useful to handle also the construction of types, and not only of terms, by derivation rules. The presented system, in order to orderly stratify the interdependent sets of types and terms, uses the following idea: types that do not depend on terms can be directly obtained, terms can be directly obtained possibly using previously obtained types—those of the variables in the context, obtained by type formation and weakening rules—and, finally, types that do depend on terms can be obtained by applying type constructors to previously obtained terms. In order to construct a type constructor we must first construct the atomic kinds of its type constructor variables, and then construct the type constructor itself by kind assignment rules. Since kinds, just as types in simply typed λ -calculus, do not depend on terms, they can be directly obtained by kind formation rules and kind assignment rules without resorting to kind constructors.

The fact that we are able to derive type constructors, which are not terms, makes it necessary to have kinds to restrict their behaviour. We cannot use types because otherwise we would mix up terms and type constructors. The presence of kinds makes it, obviously, necessary to also have rules for constructing kinds.

The weakening rules, in Table 3, simply enable us to build up the context from which we construct a kind—first line of the table—derive the kind assignment of a type constructor—second line—and derive the type assignment of a term—third line.

The conversion rules, Table 4, enable us to evaluate type constructor applications

during the typing phase. This is required since type constructors are meant, not surprisingly, to produce types that can then be assigned to terms. At the same time, type constructors produce types by being applied to arguments and by being evaluated, not unlike terms produce outputs. This generates an apparent contradiction: we need the type constructor to provide us with a type to be used while typing, but the type constructor can only this type during its evaluation, which happens after the typing phase is completed. The adopted solution consists in enabling us to evaluate type constructor applications during the typing phase by the evaluation rules. While we deductively construct our term, we can also construct its type by forming the required type constructors, by applying them and by evaluating them, and we can do all this inside the derivation through which we are constructing our term. Thus we do not need to wait for the dynamic phase of evaluation in order to obtain the type we need to assign to our term.

Let us now define, in the standard way, the arrow type \rightarrow as a special case of the product type \forall .

Notation. If the variable x does not occur in B , then we write $\forall x : A.B$ as $A \rightarrow B$. We then define $\neg A$ as $A \rightarrow \perp$ and $A_1 \wedge \dots \wedge A_n \rightarrow C$ as $A_1 \rightarrow \dots \rightarrow A_n \rightarrow C$. We, moreover, abbreviate by A^n a conjunction $A \wedge \dots \wedge A$ in which A occurs n times. Finally, by $\langle t_1, \dots, t_n \rangle$ we denote the term $\langle t_1, \langle t_2, \langle t_3, \dots, \langle t_{n-1}, t_n \rangle \dots \rangle \rangle$ containing $n - 1$ pairs nested on the right, and by π_i^* for $i \in \mathbb{N}$ the sequence of projections π_0, π_1 that selects the i th element of $\langle t_1, \dots, t_n \rangle$.

Definition 3.3 (Derivations, kinding and typing). Derivations are inductively defined as follows:

- $\vdash * : \square$ is a derivation with conclusion $\vdash * : \square$
- if $\delta_1, \dots, \delta_n$ are derivations with conclusions S_1, \dots, S_n and a rule application of the form

$$\frac{S_1 \quad \dots \quad S_n}{S}$$

is an instance of one of the rules in Tables 1, 2, 3, 4, then

$$\frac{\delta_1 \quad \dots \quad \delta_n}{S}$$

is a derivation with conclusion S .

For any kindless type constructor φ and kind Φ , if there exists a derivation with conclusion $\Gamma \vdash \varphi : \Phi$, we say that the type constructor φ has kind Φ and we express it formally by $\varphi : \Phi$.

If $\varphi : *$, then φ is a type.

For any typeless term t and type $A : *$, if there exists a derivation with conclusion $\Gamma \vdash t : A$, we say that the term t has type A and we express it formally by $t : A$.

4 The evaluation relation and its basic properties

By Definition 3.2, while defining typing derivations, we already defined the evaluation relation for type constructors. Indeed, we need to be able to statically evaluate type constructors if we wish to construct terms and assign them a type which is already in

normal form. Nevertheless, we still need to define the evaluation relation that enables us to formalise the actual dynamic process of execution of a term. This relation, unlike the one between type constructors, formalises a probabilistic notion of computation and, therefore, is a ternary relation $t \mapsto^p t'$ intuitively indicating that the term t evaluates to the term t' with a probability of p , where $p \in \mathbb{Q}$ and $0 \leq p \leq 1$.

The reduction rules for evaluating terms are shown in Table 5. Just as evaluation rules for type constructors, These rules can be applied at any depth inside a term.² That is, if a rule of the form $t \mapsto_p t'$ is displayed in Table 5, then, for any term s that contains at least one occurrence of x , $s[t/x] \mapsto_p s[t'/x]$.

We now show that \mapsto preserves kinds and that \mapsto^p preserves typability.

Theorem 4.1 (Subject reduction). *For any type constructor $\varphi : \Phi$, if $\varphi \mapsto \varphi'$, then $\varphi' : \Phi$ and all free term variables of φ' occur free also in φ .*

For any term $t : A$, if $t \mapsto_p t'$, then $t' : A'$ —where A' has the same propositional and quantificational structure as A but might contain different terms—and all free term variables of t' occur free also in t .

Proof. We start by proving the part of the statement concerning type constructors. The only possible evaluation rule applicable to a type constructor is the following:

$$\mathcal{T}[(\lambda x : \varphi_1.\varphi_2)t] \mapsto \mathcal{T}[\varphi_2[t/x]]$$

Now, suppose that $\varphi_2 : \Phi_2$. Then the kind of $\lambda x : \varphi_1.\varphi_2$ must be $\Pi x : \varphi_1.\Phi_2$. This is clear by simple inspection of the rules in Table 2 for assigning kinds. As a consequence, for any $t : \varphi_1$, the kind of $(\lambda x : \varphi_1.\varphi_2)t$ is Φ_2 . The kind of $\varphi_2[t/x]$ is Φ_2 too since the substitution replaces $x : \varphi_1$ by $t : \varphi_1$ inside $\varphi_2 : \Phi_2$. Therefore, $(\lambda x : \varphi_1.\varphi_2)t$ and $\varphi_2[t/x]$ always have the same kind and—since replacing a subconstructor with a kind by a subconstructor with the same kind inside a constructor leaves the kind of the constructor unchanged—also the constructors $\mathcal{T}[(\lambda x : \varphi_1.\varphi_2)t]$ and $\mathcal{T}[\varphi_2[t/x]]$ have the same kind.

This reduction, moreover, does not free any variable:

- All variables bound in φ_2 before the reduction, remain bound after it.
- All variables bound by occurrences of λ occurring in $\mathcal{T}[\]$ before the reduction, remain bound after it.
- All occurrences of x in φ_2 are replaced by t during the reduction.

We prove now the part of the statement concerning terms. We reason by cases on the reduction rule that has the reduction $t \mapsto_p t'$ as an instance.

- $\mathcal{C}[(\lambda x : A.s)t] \mapsto_1 \mathcal{C}[s[t/x]]$.

Suppose that $s : B$. Then the type of $\lambda x : A.s$ must be $\forall x : A.B$. This is clear by simple inspection of the rules in Table 4 for assigning types. As a consequence, for any $t : A$, the type of $(\lambda x : A.s)t$ is B . The type of $s[t/x]$ is B too since the substitution replaces $x : A$ by $t : A$ inside $s : B$. Therefore, $(\lambda x : A.s)t$ and $s[t/x]$ always have the same type and—since replacing a subterm with a type by a subterm with the same type inside a term leaves the type of the term unchanged—also the terms $\mathcal{C}[(\lambda x : A.s)t]$ and $\mathcal{C}[s[t/x]]$ have the same type.

This reduction does not free any variable:

- All variables bound in s before the reduction, remain bound after it.

²Notice that we use the words *reduction* and *evaluation* interchangeably.

- All variables bound by occurrences of λ occurring in $\mathcal{C}[\]$ before the reduction, remain bound after it.
- All occurrences of x in s are replaced by t during the reduction.

- $\mathcal{C}[\langle t_0, t_1 \rangle \pi_i] \mapsto_1 \mathcal{C}[t_i]$.

Suppose that $t_0 : A_0$ and $t_1 : A_1$. Then the type of $\langle t_0, t_1 \rangle$ must be $A_1 \wedge A_2$. This is clear by simple inspection of the rules in Table 4 for assigning types. As a consequence, the type of $\langle t_0, t_1 \rangle \pi_i$ is the type of t_i , that is, A_i . Since replacing a subterm with a type by a subterm with the same type inside a term leaves the type of the term unchanged, also the terms $\mathcal{C}[(\lambda x : A.s)t]$ and $\mathcal{C}[s[t/x]]$ have the same type.

This reduction does not free any variable:

- All variables bound in t_i before the reduction, remain bound after it.
- All variables bound by occurrences of λ occurring in $\mathcal{C}[\]$ before the reduction, remain bound after it.

- $\mathcal{C}[(t_0(p)t_1)\nu] \mapsto_q \mathcal{C}[t_i]$.

Suppose that $t_0 : A_0$ and $t_1 : A_1$. Then the type of $t_0(p)t_1$ must be $\oplus A$ where $A = A_0 = A_1$. This is clear by simple inspection of the rules in Table 4 for assigning types. As a consequence, the type of $(t_0(p)t_1)\nu$ is the type of t_i , that is, $A_i = A$. Since replacing a subterm with a type by a subterm with the same type inside a term leaves the type of the term unchanged, also the terms $\mathcal{C}[(t_0(p)t_1)\nu]$ and $\mathcal{C}[t_i]$ have the same type.

This reduction does not free any variable:

- All variables bound in t_i before the reduction, remain bound after it.
- All variables bound by occurrences of λ occurring in $\mathcal{C}[\]$ before the reduction, remain bound after it.

- $\mathcal{C}[\mathcal{C}'[o_0\nu]_1 \dots [o_0\nu]_n] \mapsto_1 \mathcal{C}[\mathcal{C}'[s_1]_1 \dots [s_n]_n]$.

Suppose that the types of s_1, \dots, s_n are A_1, \dots, A_n . Then we must have that $A_1 = \dots = A_n$ and that the type of $o_0\nu$ is $A = A_1 = \dots = A_n$. The type of o_0 must then be ΣA . This is clear by Definition 3.1 and by inspection of the rules in Table 4 for assigning types. As a consequence, the type of $o_0\nu$ and the type of each s_1, \dots, s_n is the same, that is, $A = A_1 = \dots = A_n$. Since replacing a subterm with a type by a subterm with the same type inside a term leaves the type of the term unchanged, also the terms $\mathcal{C}[\mathcal{C}'[o_0\nu]_1 \dots [o_0\nu]_n]$ and $\mathcal{C}[\mathcal{C}'[s_1]_1 \dots [s_n]_n]$ have the same type.

This reduction does not free any variable:

- All variables occurring in s_1, \dots, s_n are bound according to Definition 3.1.
- All variables bound by occurrences of λ occurring in $\mathcal{C}[\mathcal{C}'[\]_1 \dots [\]_n]$ before the reduction, remain bound after it.

- $\mathcal{C}[\mathcal{C}'[(o_1 t_1)\nu]_1 \dots [(o_1 t_n)\nu]_n] \mapsto_1 \mathcal{C}[\mathcal{C}'[s_1]_1 \dots [s_n]_n]$.

Suppose that the types of s_1, \dots, s_n are B_1, \dots, B_n . Then we must have that $B_1 = \dots = B_n$ and that the type of $(o_1 t_1)\nu, \dots, (o_1 t_n)\nu$ is $B = B_1 = \dots = B_n$. The type of o_1 must then be of the form $\forall x : A. \Sigma B$ and we must have that $t_1 : A, \dots, t_n : A$. This is clear by Definition 3.1 and by inspection of

the rules in Table 4 for assigning types. As a consequence, the type of each $(o_1 t_1)\nu, \dots, (o_1 t_n)\nu$ and the type of each s_1, \dots, s_n is the same, that is, $B = B_1 = \dots = B_n$. Since replacing a subterm with a type by a subterm with the same type inside a term leaves the type of the term unchanged, also the terms $\mathcal{C}[\mathcal{C}'[o_0\nu]_1 \dots [o_0\nu]_n]$ and $\mathcal{C}[\mathcal{C}'[s_1]_1 \dots [s_n]_n]$ have the same type.

This reduction does not free any variable:

- All variables occurring in s_1, \dots, s_n are bound according to Definition 3.1.
- All variables bound by occurrences of λ occurring in $\mathcal{C}[\mathcal{C}'[\]_1 \dots [\]_n]$ before the reduction, remain bound after it.

□

We obviously cannot prove that a term only has one normal form. Indeed, a term of the form $s(p)t$, for $s \neq t$, is *supposed* to have at least two normal forms: the normal form of s and that of t . Hence, we will simply prove that type constructors have a unique normal form. In Section 5, we will show that each type constructor has a normal form that can be reached in a finite number of evaluation steps. We will actually prove much more than that, but the rest will be discussed in due time. Right now, we prove Theorem 4.4, which guarantees that all reduced forms of a term can be evaluated to the same normal form. First, though, we need to prove a couple of lemmata that enable us to handle, inside the proof of Theorem 4.4, the substitutions generated by the reduction of applications.

Lemma 4.2. *For any two variables x and y such that $x \neq y$, term t in which y does not occur free and type constructor or term θ , $(\theta[t/x])[s[t/x]/y] = (\theta[s/y])[t/x]$.*

Proof. The proof is by induction on the number of symbol occurrences in θ . We reason by cases on the structure of θ .

- $\theta = \alpha$. Since $(\alpha[t/x])[s[t/x]/y] = \alpha[s[t/x]/y] = \alpha = \alpha[t/x] = (\alpha[s/y])[t/x]$ we have that the statement holds.
- $\theta = \forall z : A.\psi$. Considering that we can always rename bound variables, we suppose that $x \neq z \neq y$. Then

$$((\forall z : A.\psi)[t/x])[s[t/x]/y] = (\forall z : (A[t/x]).\psi[t/x])[s[t/x]/y] = \forall z : ((A[t/x])[s[t/x]/y]).(\psi[t/x])[s[t/x]/y]$$

By inductive hypothesis, $(\psi[t/x])[s[t/x]/y] = (\psi[s/y])[t/x]$ and $(A[t/x])[s[t/x]/y] = (A[s/y])[t/x]$, which implies that $\forall z : ((A[t/x])[s[t/x]/y]).(\psi[t/x])[s[t/x]/y] = \forall z : ((A[s/y])[t/x]).(\psi[s/y])[t/x]$. From this and the following equalities:

$$\forall z : ((A[s/y])[t/x]).(\psi[s/y])[t/x] = (\forall z : (A[s/y]).(\psi[s/y]))[t/x] = ((\forall z : A.\psi)[s/y])[t/x]$$

—also due to the fact that $x \neq z \neq y$ —we can conclude that the statement holds in this case as well.

- $\theta = \oplus\psi$. We have that

$$((\oplus\psi)[t/x])[s[t/x]/y] = (\oplus\psi[t/x])[s[t/x]/y] = \oplus(\psi[t/x])[s[t/x]/y]$$

By inductive hypothesis, $(\psi[t/x])[s[t/x]/y] = (\psi[s/y])[t/x]$, which implies that $\oplus(\psi[t/x])[s[t/x]/y] = \oplus(\psi[s/y])[t/x]$. From this and the fact that $\oplus(\psi[s/y])[t/x] = (\oplus(\psi[s/y]))[t/x] = ((\oplus\psi)[s/y])[t/x]$ we can conclude that the statement holds in this case as well.

- $\theta = \psi_1 \wedge \psi_2$. We have that

$$\begin{aligned} ((\psi_1 \wedge \psi_2)[t/x])[s[t/x]/y] &= (\psi_1[t/x] \wedge \psi_2[t/x])[s[t/x]/y] \\ &= (\psi_1[t/x])[s[t/x]/y] \wedge (\psi_2[t/x])[s[t/x]/y] \end{aligned}$$

By inductive hypothesis, $(\psi_i[t/x])[s[t/x]/y] = (\psi_i[s/y])[t/x]$, which implies that

$$(\psi_1[t/x])[s[t/x]/y] \wedge (\psi_2[t/x])[s[t/x]/y] = (\psi_1[s/y])[t/x] \wedge (\psi_2[s/y])[t/x]$$

From this and the fact that

$$(\psi_1[s/y])[t/x] \wedge (\psi_2[s/y])[t/x] = (\psi_1[s/y] \wedge \psi_2[s/y])[t/x] = ((\psi_1 \wedge \psi_2)[s/y])[t/x]$$

we can conclude that that the statement holds in this case as well.

- $\theta = \lambda z : A.\psi$. Considering that we can always rename bound variables, we suppose that $x \neq z \neq y$. Then

$$((\lambda z : A.\psi)[t/x])[s[t/x]/y] = (\lambda z : (A[t/x]).\psi[t/x])[s[t/x]/y] = \lambda z : ((A[t/x])[s[t/x]/y]).(\psi[t/x])[s[t/x]/y]$$

By inductive hypothesis, $(\psi[t/x])[s[t/x]/y] = (\psi[s/y])[t/x]$ and $(A[t/x])[s[t/x]/y] = (A[s/y])[t/x]$, which implies that $\lambda z : ((A[t/x])[s[t/x]/y]).(\psi[t/x])[s[t/x]/y] = (\lambda z : (A[s/y]).(\psi[s/y]))[t/x]$. From this and the following equalities: $\lambda z : ((A[s/y])[t/x]).(\psi[s/y])[t/x] = (\lambda z : (A[s/y]).(\psi[s/y]))[t/x] = ((\lambda z : A.\psi)[s/y])[t/x]$ —also due to the fact that $x \neq z \neq y$ —we can conclude that the statement holds in this case as well.

- $\theta = \psi u$. We have that

$$((\psi u)[t/x])[s[t/x]/y] = (\psi[t/x]u[t/x])[s[t/x]/y] = (\psi[t/x])[s[t/x]/y](u[t/x])[s[t/x]/y]$$

By inductive hypothesis, $(\psi[t/x])[s[t/x]/y] = (\psi[s/y])[t/x]$ and $(u[t/x])[s[t/x]/y] = (u[s/y])[t/x]$, which implies that

$$(\psi[t/x])[s[t/x]/y](u[t/x])[s[t/x]/y] = (\psi[s/y])[t/x](u[s/y])[t/x]$$

From this and the fact that

$$(\psi[s/y])[t/x](u[s/y])[t/x] = (\psi[s/y]u[s/y])[t/x] = ((\psi u)[s/y])[t/x]$$

we can conclude that that the statement holds in this case as well.

- $\theta = z$. We have three subcases: $x \neq z \neq y$, $z = x$ and $z = y$.

If $x \neq z \neq y$ then $(z[t/x])[s[t/x]/y] = z = (z[s/y])[t/x]$.

If $\theta = z = x$ then $(x[t/x])[s[t/x]/y] = t[s[t/x]/y] = t$ since, by assumption, y does not occur free in t . But we also have that $(x[s/y])[t/x] = x[t/x]$ since $x \neq y$ and, finally, $x[t/x] = t$. Hence, indeed, $(x[t/x])[s[t/x]/y] = (x[s/y])[t/x]$ as desired since $x = \theta$.

If $\theta = z = y$ then $(y[t/x])[s[t/x]/y] = y[s[t/x]/y] = s[t/x]$ since, by assumption, $y \neq x$. But we also have that $(y[s/y])[t/x] = s[t/x]$. Hence, indeed, $(y[t/x])[s[t/x]/y] = (y[s/y])[t/x]$ as desired since $y = \theta$.

- $\theta = \langle u_1, u_2 \rangle$. We have that

$$\begin{aligned} (\langle u_1, u_2 \rangle[t/x])[s[t/x]/y] &= \langle u_1[t/x], u_2[t/x] \rangle[s[t/x]/y] \\ &= \langle (u_1[t/x])[s[t/x]/y], (u_2[t/x])[s[t/x]/y] \rangle \end{aligned}$$

By inductive hypothesis, $(u_i[t/x])[s[t/x]/y] = (u_i[s/y])[t/x]$, which implies that

$$\langle (u_1[t/x])[s[t/x]/y], (u_2[t/x])[s[t/x]/y] \rangle = \langle (u_1[s/y])[t/x], (u_2[s/y])[t/x] \rangle$$

From this and the fact that

$$\langle (u_1[s/y])[t/x], (u_2[s/y])[t/x] \rangle = \langle u_1[s/y], u_2[s/y] \rangle[t/x] = \langle (u_1, u_2)[s/y] \rangle[t/x]$$

we can conclude that that the statement holds in this case as well.

- $\theta = u\pi_0$ and $t = u\pi_1$. We have that

$$((u\pi_i)[t/x])[s[t/x]/y] = ((u[t/x]\pi_i)[s[t/x]/y] = ((u[t/x])[s[t/x]/y])\pi_i$$

By inductive hypothesis, $(u[t/x])[s[t/x]/y] = (u[s/y])[t/x]$, which implies that

$$((u[t/x])[s[t/x]/y])\pi_i = ((u[s/y])[t/x])\pi_i$$

From this and the fact that

$$((u[s/y])[t/x])\pi_i = ((u[s/y])\pi_i)[t/x] = ((u\pi_i)[s/y])[t/x]$$

we can conclude that that the statement holds in this case as well.

- $\theta = \lambda z : A.u$. Considering that we can always rename bound variables, we suppose that $x \neq z \neq y$. Then

$$((\lambda z : A.u)[t/x])[s[t/x]/y] = (\lambda z : (A[t/x]).u[t/x])[s[t/x]/y] = \lambda z : ((A[t/x])[s[t/x]/y]).(u[t/x])[s[t/x]/y]$$

By inductive hypothesis, $(u[t/x])[s[t/x]/y] = (u[s/y])[t/x]$ and $(A[t/x])[s[t/x]/y] = (A[s/y])[t/x]$, which implies that $\lambda z : ((A[t/x])[s[t/x]/y]).(u[t/x])[s[t/x]/y] = (\lambda z : (A[s/y]).(u[s/y]))[t/x]$. From this and the following equalities: $\lambda z : ((A[s/y])[t/x]).(u[s/y])[t/x] = (\lambda z : (A[s/y]).(u[s/y]))[t/x] = ((\lambda z : A.u)[s/y])[t/x]$ —also due to the fact that $x \neq z \neq y$ —we can conclude that the statement holds in this case as well.

- $\theta = vu$. We have that

$$((vu)[t/x])[s[t/x]/y] = (v[t/x]u[t/x])[s[t/x]/y] = (v[t/x])[s[t/x]/y](u[t/x])[s[t/x]/y]$$

By inductive hypothesis, $(v[t/x])[s[t/x]/y] = (v[s/y])[t/x]$ and $(u[t/x])[s[t/x]/y] = (u[s/y])[t/x]$, which implies that

$$(v[t/x])[s[t/x]/y](u[t/x])[s[t/x]/y] = (v[s/y])[t/x](u[s/y])[t/x]$$

From this and the fact that

$$(v[s/y])[t/x](u[s/y])[t/x] = (v[s/y]u[s/y])[t/x] = ((vu)[s/y])[t/x]$$

we can conclude that that the statement holds in this case as well.

- $\theta = u_1\langle p \rangle u_2$. We have that

$$\begin{aligned} ((u_1\langle p \rangle u_2)[t/x])[s[t/x]/y] &= ((u_1[t/x]\langle p \rangle (u_2[t/x]))[s[t/x]/y]) \\ &= \langle p \rangle (u_2[t/x])[s[t/x]/y] \end{aligned}$$

By inductive hypothesis, $(u_i[t/x])[s[t/x]/y] = (u_i[s/y])[t/x]$, which implies that

$$((u_1[t/x]\langle p \rangle (u_2[t/x]))[s[t/x]/y]) = ((u_1[s/y])[t/x]\langle p \rangle (u_2[s/y])[t/x])$$

From this and the fact that

$$((u_1[s/y])[t/x]\langle p \rangle (u_2[s/y])[t/x]) = ((u_1[s/y]\langle p \rangle (u_2[s/y]))[t/x]) = ((u_1\langle p \rangle u_2)[s/y])[t/x]$$

we can conclude that that the statement holds in this case as well.

- $\theta = u\nu$. We have that

$$((u\nu)[t/x])[s[t/x]/y] = ((u[t/x]\nu)[s[t/x]/y] = ((u[t/x])[s[t/x]/y])\nu$$

By inductive hypothesis, $(u[t/x])[s[t/x]/y] = (u[s/y])[t/x]$, which implies that

$$((u[t/x])[s[t/x]/y])\nu = ((u[s/y])[t/x])\nu$$

From this and the fact that

$$((u[s/y])[t/x])\nu = ((u[s/y])\nu)[t/x] = ((u\nu)[s/y])[t/x]$$

we can conclude that that the statement holds in this case as well. \square

Lemma 4.3. *For any type constructor χ and term t that does not contain free variables which are bound in χ , if $\chi \mapsto \chi'$ then $\chi[t/x] \mapsto \chi'[t/x]$.*

Proof. We reason by induction on the number of symbol occurrences in χ .

In the base case, $\chi = \alpha$. Since α cannot be reduced to any type constructor, the statement is trivially satisfied.

Suppose now that all type constructors with less than n symbols verify the statement, we show that also a type constructor χ with n symbols does. We reason by cases on the structure of χ .

- $\chi = \forall y : A.\psi$. First, if x is bound in χ then $\chi[t/x] = \chi$, $\chi'[t/x] = \chi'$ and the statement is tautologous. We hence suppose that $x \neq y$. Now, $\forall y : A.\psi \mapsto \forall y : A.\psi'$ if, and only if, $\psi \mapsto \psi'$ and $\forall y : A.\psi$ cannot be reduced in any other way. By inductive hypothesis, if $\psi \mapsto \psi'$ then $\psi[t/x] \mapsto \psi'[t/x]$. Hence, we have that $\psi[t/x] \mapsto \psi'[t/x]$ enables us to obtain $\forall y : (A[t/x]).(\psi[t/x]) \mapsto \forall y : (A[t/x]).(\psi'[t/x])$. Now, since $x \neq y$ and since t does not contain free variables which are bound in χ , we have that $\forall y : (A[t/x]).(\psi[t/x]) = (\forall y : A.\psi)[t/x]$ and $\forall y : (A[t/x]).(\psi'[t/x]) = (\forall y : A.\psi')[t/x]$ and thus that the statement holds for $\chi = \forall y : A.\psi$.
- $\chi = \oplus\psi$. Now, $\oplus\psi \mapsto \oplus\psi'$ if, and only if, $\psi \mapsto \psi'$ and $\oplus\psi$ cannot be reduced in any other way. By inductive hypothesis, if $\psi \mapsto \psi'$ then $\psi[t/x] \mapsto \psi'[t/x]$. But since $\psi[t/x] \mapsto \psi'[t/x]$ implies that $\oplus(\psi[t/x]) \mapsto \oplus(\psi'[t/x])$ and since $\oplus(\psi[t/x]) = (\oplus\psi)[t/x]$ and $\oplus(\psi'[t/x]) = (\oplus\psi')[t/x]$, we have that the statement holds for $\chi = \oplus\psi$.
- $\chi = \psi_1 \wedge \psi_2$. Now, $\psi_1 \wedge \psi_2$ can only reduce in one step to terms of the form $\psi'_1 \wedge \psi_2$ or $\psi_1 \wedge \psi'_2$ such that $\psi_i \mapsto \psi'_i$ for $i \in \{1, 2\}$. Moreover, $\psi_i \mapsto \psi'_i$ implies that $\psi_1 \wedge \psi_2 \mapsto \psi'_1 \wedge \psi_2$ and that $\psi_1 \wedge \psi_2 \mapsto \psi_1 \wedge \psi'_2$. Moreover, by inductive hypothesis, if $\psi_i \mapsto \psi'_i$ then $\psi_i[t/x] \mapsto \psi'_i[t/x]$. Therefore, if $\chi \mapsto \chi'$, then
 - either $\chi' = \psi'_1 \wedge \psi_2$ and $\chi[t/x] = (\psi_1 \wedge \psi_2)[t/x] = \psi_1[t/x] \wedge \psi_2[t/x] \mapsto \psi'_1[t/x] \wedge \psi_2[t/x] = (\psi'_1 \wedge \psi_2)[t/x] = \chi'[t/x]$,
 - or $\chi' = \psi_1 \wedge \psi'_2$ and $\chi[t/x] = (\psi_1 \wedge \psi_2)[t/x] = \psi_1[t/x] \wedge \psi_2[t/x] \mapsto \psi_1[t/x] \wedge \psi'_2[t/x] = (\psi_1 \wedge \psi'_2)[t/x] = \chi'[t/x]$

In both cases, the statement holds for $\chi = \psi_1 \wedge \psi_2$.

- $\chi = \lambda y : A.\psi$. First, if x is bound in χ then $\chi[t/x] = \chi$, $\chi'[t/x] = \chi'$ and the statement is tautologous. Hence, we suppose that x is free in χ . Now, it is clear that $\lambda y : A.\psi \mapsto \lambda y : A.\psi'$ if, and only if, $\psi \mapsto \psi'$ and that $\lambda y : A.\psi$ cannot be reduced in any other way. Therefore we know that $\chi \mapsto \chi'$ implies that $\psi \mapsto \psi'$. By inductive hypothesis, moreover, $\psi \mapsto \psi'$ implies that $\psi[t/x] \mapsto \psi'[t/x]$. Since we assumed that x is free in χ we have, as a consequence, that $\lambda y : (A[t/x]).(\psi[t/x]) = (\lambda y : A.\psi)[t/x]$ and $\lambda y : (A[t/x]).(\psi'[t/x]) = (\lambda y : A.\psi')[t/x]$. But then from $\psi[t/x] \mapsto \psi'[t/x]$ —again from the fact that $\lambda y : A.\theta \mapsto \lambda y : A.\theta'$ if, and only if, $\theta \mapsto \theta'$ —we can obtain $(\lambda y : A.\psi)[t/x] = \lambda y : (A[t/x]).(\psi[t/x]) \mapsto \lambda y : (A[t/x]).(\psi'[t/x]) = (\lambda y : A.\psi')[t/x]$, which proves that the statement holds in this case as well.
- $\chi = \psi s$. Now, either ψs reduces to $\psi's$ —in which case we can prove the statement by induction just like in the first case considered—or we have that $\chi = \psi s = (\lambda y : A.\theta)s \mapsto \theta[s/y]$. In this case we have to prove that $((\lambda y : A.\theta)s)[t/x] \mapsto (\theta[s/y])[t/x]$. As in the previous case, we suppose that x is free in χ because otherwise $\chi[t/x] = \chi$, $\chi'[t/x] = \chi'$ and the statement is tautologous. But if x is free in $\chi = \psi s = (\lambda y : A.\theta)s$ then $x \neq y$. Thus, $((\lambda y : A.\theta)s)[t/x] = (\lambda y : A.(\theta[t/x]))(s[t/x]) \mapsto (\theta[t/x])[(s[t/x])/y]$. Since— with the help of the assumption that y does not occur in t and of the fact that $x \neq y$ —Lemma 4.2 guarantees us that $(\theta[t/x])[(s[t/x])/y] = (\theta[s/y])[t/x]$, we have that the statement holds also in this case. □

Theorem 4.4 (Confluence for type constructors). *For any $\varphi : \Phi$, if $\varphi \mapsto^* \varphi_1$ and $\varphi \mapsto^* \varphi_2$, then there exists a type constructor φ_3 for which $\varphi_1 \mapsto^* \varphi_3$ and $\varphi_2 \mapsto^* \varphi_3$.*

Proof. We are going to prove the following statement:

If $\varphi \mapsto \varphi_1$ and $\varphi \mapsto \varphi_2$, then we can always reduce φ_1 and φ_2 in such a way to show that $\varphi_1 \mapsto \varphi_3$ and $\varphi_2 \mapsto \varphi_3$.

Then a simple argument by diagram chase—see, for instance, [Sørensen and Urzyczyn, 2006, Chap. 1, Sec. 1.4]—yields the statement.

Let us reason by induction on the number of symbol occurrences in φ and by cases on the reductions $\varphi \mapsto \varphi_1$ and $\varphi \mapsto \varphi_2$.

In the base case, $\varphi = \alpha$, which is impossible since α cannot be reduced.

Suppose now that all type constructors with less than n symbols verify the statement, we show that also a type constructor φ with n symbols does.

- $\varphi = \forall x : A.\psi$.

Now, clearly, $\forall x : A.\psi$ can only reduce to terms of the form $\forall x : A.\psi \mapsto \forall x : A.\psi'$ such that $\psi \mapsto \psi'$ and, moreover, $\psi \mapsto \psi'$ implies that $\forall x : A.\psi \mapsto \forall x : A.\psi'$. Therefore, any two reductions $\forall x : A.\psi = \varphi \mapsto \varphi_1$ and $\forall x : A.\psi = \varphi \mapsto \varphi_2$ can be represented as follows:

$$\varphi = \forall x : A.\psi \mapsto \forall x : A.\psi_1 = \varphi_1 \qquad \varphi = \forall x : A.\psi \mapsto \forall x : A.\psi_2 = \varphi_2$$

But this means that

$$\psi \mapsto \psi_1 \qquad \psi \mapsto \psi_2$$

which, by inductive hypothesis, implies that there exists a ψ_3 for which

$$\psi_1 \mapsto \psi_3 \qquad \psi_2 \mapsto \psi_3$$

This, in turn, implies that

$$\oplus\psi_1 \mapsto \oplus\psi_3 \quad \oplus\psi_2 \mapsto \oplus\psi_3$$

which gives us what we were looking for if we take $\oplus\psi_3$ as our φ_3 . Hence, $\varphi = \forall x : A.\psi$ verifies the statement.

- $\varphi = \oplus\psi$.

Now, clearly, $\oplus\psi$ can only reduce to terms of the form $\oplus\psi \mapsto \oplus\psi'$ such that $\psi \mapsto \psi'$ and, moreover, $\psi \mapsto \psi'$ implies that $\oplus\psi \mapsto \oplus\psi'$. Therefore, any two reductions $\oplus\psi = \varphi \mapsto \varphi_1$ and $\oplus\psi = \varphi \mapsto \varphi_2$ can be represented as follows:

$$\varphi = \oplus\psi \mapsto \oplus\psi_1 = \varphi_1 \quad \varphi = \oplus\psi \mapsto \oplus\psi_2 = \varphi_2$$

But this means that

$$\psi \mapsto \psi_1 \quad \psi \mapsto \psi_2$$

which, by inductive hypothesis, implies that there exists a ψ_3 for which

$$\psi_1 \mapsto \psi_3 \quad \psi_2 \mapsto \psi_3$$

This, in turn, implies that

$$\oplus\psi_1 \mapsto \oplus\psi_3 \quad \oplus\psi_2 \mapsto \oplus\psi_3$$

which gives us what we were looking for if we take $\oplus\psi_3$ as our φ_3 . Hence, $\varphi = \oplus\psi$ verifies the statement.

- $\varphi = \psi \wedge \chi$. Now, $\psi \wedge \chi$ can only reduce in one step to terms of the form $\psi' \wedge \chi$ or $\psi \wedge \chi'$ such that $\psi \mapsto \psi'$ and, respectively, $\chi \mapsto \chi'$. Moreover, $\psi \mapsto \psi'$ and $\chi \mapsto \chi'$ imply that $\psi \wedge \chi \mapsto \psi' \wedge \chi$ and $\psi \wedge \chi \mapsto \psi \wedge \chi'$. By inductive hypothesis, ψ and χ verify the statement. Therefore, if both reductions of $\varphi = \psi \wedge \chi$ act exclusively inside ψ or exclusively inside χ , the statement clearly holds for $\varphi = \psi \wedge \chi$ as well. Suppose then that $\psi \wedge \chi = \varphi \mapsto \varphi_1 = \psi' \wedge \chi$ and $\psi \wedge \chi = \varphi \mapsto \varphi_2 = \psi \wedge \chi'$. Since then we must have that $\chi \mapsto \chi'$ and $\psi \mapsto \psi'$, we also have that $\psi' \wedge \chi \mapsto \psi' \wedge \chi'$ and $\psi \wedge \chi' \mapsto \psi' \wedge \chi'$. Taking $\psi' \wedge \chi'$ as φ_3 gives us that the statement is verified by φ as well.
- $\varphi = \lambda x : A.\psi$. By inductive hypothesis, if $\psi \mapsto \psi_1$ and $\psi \mapsto \psi_2$, then there exists a ψ_3 for which $\psi_1 \mapsto \psi_3$ and $\psi_2 \mapsto \psi_3$. Since $\lambda x : A.\psi \mapsto \lambda x : A.\psi'$ if, and only if, $\psi \mapsto \psi'$ and since $\lambda x : A.\psi$ cannot reduce in any other way, we have, by an argument analogous to the one used in the first case considered, that also $\varphi = \lambda x : A.\psi$ verifies the statement.
- $\varphi = \psi t$. By inductive hypothesis, if $\psi \mapsto \psi_1$ and $\psi \mapsto \psi_2$, then there exists a ψ_3 for which $\psi_1 \mapsto \psi_3$ and $\psi_2 \mapsto \psi_3$. Therefore, if both reductions of ψt exclusively act inside ψ , we have that that $\psi t \mapsto \psi_1 t$ and $\psi t \mapsto \psi_2 t$, and then $\psi_1 t \mapsto \psi_3 t$ and $\psi_2 t \mapsto \psi_3 t$. If, otherwise, $\varphi = (\lambda x : A.\chi)t \mapsto \chi[t/x] = \varphi_1$ and $\varphi = (\lambda x : A.\chi)t \mapsto (\lambda x : A.\chi')t = \varphi_2$, we take φ_3 to be $\chi'[t/x]$. Indeed, clearly, $\varphi_2 = (\lambda x : A.\chi')t \mapsto \chi'[t/x] = \varphi_3$. Moreover, by Lemma 4.3 and by possibly renaming the variables which are bound in χ , if $\chi \mapsto \chi'$ then $\varphi_1 = \chi[t/x] \mapsto \chi'[t/x] = \varphi_3$. In conclusion, $\varphi = \psi t$ verifies the statement in all cases.

□

5 Strong normalisation

We generalise the proof of strong normalisation by Tait's reducibility technique presented in [Girard et al., 1989, Chapter 6].

Definition 5.1 (Normal and strongly normalising). A term t is normal if there is no term t' and real number p such that $t \mapsto_p t'$. A type constructor φ is normal if there is no type constructor φ' such that $\varphi \mapsto \varphi'$.

A term t (type constructor φ) is strongly normalising if there exists no infinite sequence of reduction steps $t \mapsto_{p_1} t_1 \mapsto_{p_2} t_2 \mapsto_{p_3} t_3 \dots$ ($\varphi \mapsto \varphi_1 \mapsto \varphi_2 \mapsto \varphi_3 \dots$)

We define now, for any type $A : *$ and kind $\Phi : \square$, the set Red_A of reducible terms of type A and Red_Φ of reducible type constructors of kind Φ .

Definition 5.2 (Reducibles and sets of reducibles). For any type $A : *$ and kind $\Phi : \square$, the set Red_A of reducibles of type A and the set Red_Φ of reducibles of kind Φ are defined as follows:

- for $\Phi = *$, Red_* contains all strongly normalising type constructors of kind $*$.
- for $\Phi = \Pi x : B. \Psi$, $\text{Red}_{\Pi y : B. \Psi}$ contains all type constructors $\varphi : \Pi y : B. \Psi$ such that, for any term $s : B$, $\varphi s \in \text{Red}_{\Psi[s/x]}$.
- for $A = P$ atomic type, Red_P contains all strongly normalising terms of type A .
- for $A = B \wedge C$, $\text{Red}_{B \wedge C}$ contains all terms $t : B \wedge C$ such that $t\pi_0 \in \text{Red}_B$ and $t\pi_1 \in \text{Red}_C$.
- for $A = \forall y : B. C$, $\text{Red}_{\forall y : B. C}$ contains all terms $t : (\forall y : B. C)$ such that, for any term $s \in \text{Red}_B$, $ts \in \text{Red}_{C[s/y]}$.
- for $A = \oplus B$, $\text{Red}_{\oplus B}$ contains all terms $t : \oplus B$ such that $t\nu \in \text{Red}_B$.

The set Red of all reducibles is defined as follows:

$$\left(\bigcup_{A:*} \text{Red}_A \right) \cup \left(\bigcup_{\Psi:\square} \text{Red}_\Psi \right)$$

Clearly, the definition of the set $\text{Red}_{\Pi y : B. \Psi}$ of all reducibles of kind $\Pi y : B. \Psi$ is modelled after the definition of the set $\text{Red}_{\forall y : B. C}$ of all reducibles of type $\forall y : B. C$. Nevertheless, since during the evaluation of type constructors we do not reduce the terms to which they are applied, there is no need to mention whether or not the arguments of type constructors of kind $\Pi y : B. \Psi$ must be reducible in order to obtain a reducible type constructor application. As we will show, all terms are reducible, but this does not influence in any way the evaluation termination proof for type constructors.

Definition 5.3 (Neutral terms and type constructors). A term or type constructor is neutral if it is in **none** of the following forms:

$$\langle s, t \rangle \quad \lambda x. t \quad s(p)t \quad \lambda x. \varphi$$

That is, t is neutral if, and only if, each redex occurring in $ts, t\pi_i, t\nu$ either occurs in t or occurs in s ; and φ is neutral if, and only if, each redex occurring in φt occurs in φ .

Let us introduce three conditions that we will show to hold for the sets of reducible terms.

- CSN * If $t \in \text{Red}_A$, then t is strongly normalising.
 □ If $\varphi \in \text{Red}_\Phi$, then φ is strongly normalising.
- CEval * If $t : A, t' : A', t \in \text{Red}_A$ and $t \mapsto_p t'$, then $t' \in \text{Red}_{A'}$.
 □ If $\varphi \in \text{Red}_\Phi$ and $\varphi \mapsto \varphi'$, then $\varphi' \in \text{Red}_\Phi$.
- CAntiEval * If
 – $t : A$ is neutral and
 – for each $t' : A'$ such that $t \mapsto_p t'$, we have that $t' \in \text{Red}_{A'}$
 then $t \in \text{Red}_A$.
 □ If
 – φ is neutral and
 – for each φ' such that $\varphi \mapsto \varphi'$, we have that $\varphi' \in \text{Red}_\Phi$
 then $\varphi \in \text{Red}_\Phi$.

Where A' is a type with the same propositional and quantificational structure as A but that possibly contains a term which is the one-step reduced form of the corresponding term occurring in A .

Notice that CEval states that Red is closed under evaluation (the relation \mapsto_p on terms and \mapsto on type constructors), and CAntiEval states that Red is closed under those instances of anti-evaluation (the inverse of \mapsto_p and \mapsto) that yield neutral terms.

Proposition 5.1. *For any type A and kind Φ , Red_A and Red_Φ verify CSN, CEval and CAntiEval.*

Proof. We prove the statement by induction on the number of symbol occurrences in A and in Φ . In the base case for types, $A = P$ is an atomic type.

- CSN. Since Red_P only contains strongly normalising terms by definition, CSN is trivially verified.
- CEval. Since all elements of Red_P are strongly normalising by definition and since a strongly normalising term can only reduce to strongly normalising terms, CEval is verified.
- CAntiEval. We suppose that $t : P$ is neutral and that, for each t' such that $t \mapsto_p t'$, we have $t' \in \text{Red}_{P'}$. We need to prove that $t \in \text{Red}_P$. Any sequence of reductions starting from t must be of the form $t \mapsto_{p_1} t_1 \mapsto_{p_1} \dots$ where $t_1 \in \text{Red}_{P_1}$ and thus t_1 is strongly normalising. Hence, any sequence of reductions starting from t must be finite. But this means that $t : P$ is strongly normalising, which implies, by definition of Red_P , that $t \in \text{Red}_P$. Hence, also CAntiEval is verified.

In the base case for kinds, $\Phi = *$ is an atomic kind.

- CSN. Since Red_* only contains strongly normalising elements by definition, CSN is trivially verified.
- CEval. Since all elements of Red_* are strongly normalising by definition and since a strongly normalising constructor can only reduce to strongly normalising constructors, CEval is verified.

- CAntiEval. We suppose that $\varphi : *$ is neutral and that, for each φ' such that $\varphi \mapsto \varphi'$, we have $\varphi' \in \text{Red}_*$. We need to prove that $\varphi \in \text{Red}_*$. Any sequence of reductions starting from φ must be of the form $\varphi \mapsto \varphi_1 \mapsto \dots$ where $\varphi_1 \in \text{Red}_*$ and thus φ_1 is strongly normalising. Hence, any sequence of reductions starting from φ must be finite. But this means that $\varphi : *$ is strongly normalising, which implies, by definition of Red_* , that $\varphi \in \text{Red}_*$. Hence, also CAntiEval is verified.

Suppose now that, for any type F and kind Ψ containing less than n symbols, Red_F and Red_Ψ verify CSN, CEval and CAntiEval. We prove that this holds also for any type A and kind Φ containing n symbols.

Suppose that $A = B \wedge C$.

- CSN. If $t \in \text{Red}_{B \wedge C}$, then $t\pi_0 \in \text{Red}_B$. Hence, and by induction hypothesis (CSN), $t\pi_0$ is strongly normalising. Suppose, by reasoning indirectly, that an infinite reduction sequence of the form $t \mapsto_{p_1} t_1 \mapsto_{p_2} t_2 \dots$ exists, then also an infinite reduction sequence of the form $t\pi_0 \mapsto_{p_1} t_1\pi_0 \mapsto_{p_2} t_2\pi_0 \dots$ exists. Which is absurd since $t\pi_0$ is strongly normalising. Hence, t is strongly normalising and CSN is verified.
- CEval. Since $t \in \text{Red}_{B \wedge C}$, we have, by definition of $\text{Red}_{B \wedge C}$, that $t\pi_0 \in \text{Red}_B$ and $t\pi_1 \in \text{Red}_C$. Moreover, if $t \mapsto_p t'$, then $t\pi_0 \mapsto_p t'\pi_0$ and $t\pi_1 \mapsto_p t'\pi_1$. From $t\pi_0 \in \text{Red}_B$, $t\pi_1 \in \text{Red}_C$, $t\pi_0 \mapsto_p t'\pi_0$ and $t\pi_1 \mapsto_p t'\pi_1$, by induction hypothesis (CEval), we obtain that $t'\pi_0 \in \text{Red}_{B'}$ and $t'\pi_1 \in \text{Red}_{C'}$. From this and by definition of $\text{Red}_{B \wedge C}$, we have that $t' \in \text{Red}_{B' \wedge C'}$, and thus that CEval is verified.
- CAntiEval. We suppose that $t : B \wedge C$ is neutral and that, for each t' such that $t \mapsto_p t'$, we have $t' \in \text{Red}_{B' \wedge C'}$. We need to prove that $t \in \text{Red}_{B \wedge C}$. Since t is neutral, $t\pi_0$ can only reduce to a term of the form $t'\pi_0$ such that $t \mapsto_p t'$. Since the local assumptions guarantee us that $t' \in \text{Red}_{B' \wedge C'}$, we also know that $t'\pi_0 \in \text{Red}_{B'}$. But then, since $t\pi_0$ is neutral and can only reduce to terms in $\text{Red}_{B'}$, by inductive hypothesis (CAntiEval), we have that $t\pi_0 \in \text{Red}_B$. Since the same argument guarantees that $t\pi_1 \in \text{Red}_C$, we have that $t \in \text{Red}_{B \wedge C}$ and thus that CAntiEval is verified.

Suppose that $A = \forall y : B.C$.

- CSN. If $t \in \text{Red}_{\forall y : B.C}$, then $tx \in \text{Red}_{C[x/y]}$ for any $x : B$. Indeed, by inductive hypothesis (CAntiEval), from $x : A$ being normal and neutral we can infer that $x \in \text{Red}_A$. By induction hypothesis (CSN), $tx \in \text{Red}_{C[x/y]}$ implies that tx is strongly normalising. Suppose now, by reasoning indirectly, that an infinite reduction sequence of the form $t \mapsto_{p_1} t_1 \mapsto_{p_2} t_2 \dots$ exists, then also an infinite reduction sequence of the form $tx \mapsto_{p_1} t_1x \mapsto_{p_2} t_2x \dots$ exists. Which is absurd since tx is strongly normalising. Hence, t is strongly normalising and CSN is verified.
- CEval. Since $t \in \text{Red}_{\forall y : B.C}$, we have, by definition of $\text{Red}_{\forall y : B.C}$, that $ts \in \text{Red}_{C[s/y]}$ for any $s \in \text{Red}_B$. Moreover, if $t \mapsto_p t'$, then $ts \mapsto_p t's$. From $ts \in \text{Red}_{C[s/y]}$ and $ts \mapsto_p t's$, by induction hypothesis (CEval), we obtain that $t's \in \text{Red}_{C'[s/y]}$. Since s is a generic element of Red_B , from $t's \in \text{Red}_{C'[s/y]}$ and by definition of $\text{Red}_{\forall y : B.C'}$, we have that $t' \in \text{Red}_{\forall y : B.C'}$, and thus that CEval is verified.
- CAntiEval. We suppose that $t : (\forall y : B.C)$ is neutral and that, for each t' such that $t \mapsto_p t'$, we have $t' \in \text{Red}_{\forall y : B.C'}$. We need to prove that $t \in \text{Red}_{\forall y : B.C}$. In

order to show this, it is enough to show that $ts \in \text{Red}_{C[s/y]}$ for any $s \in \text{Red}_B$. Since, by induction hypothesis (CEval), $s \in \text{Red}_B$ implies that s is strongly normalising, we know that the longest sequence of reductions starting from s is finite and we can reason by induction on its length $l(s)$.

If $l(s) = 0$, since t is neutral, then ts can only reduce to terms of the form $t's$ such that $t \mapsto_p t'$. Since the local assumptions guarantee us that $t' \in \text{Red}_{\forall y:B.C'}$, we also know that $t's \in \text{Red}_{C[s/y]}$. But then, since ts is neutral and can only reduce to terms in $\text{Red}_{C[s/y]}$, by inductive hypothesis (CAntiEval), we have that $ts \in \text{Red}_{C[s/y]}$ as desired.

Suppose now that $ts \in \text{Red}_{C[s/y]}$ for any $s \in \text{Red}_B$ such that $l(s) < n$, we show that this holds also for any $s \in \text{Red}_B$ such that $l(s) = n$.

Since t is neutral, ts can either reduce to a term of the form $t's$ such that $t \mapsto_p t'$ or of the form ts' such that $s \mapsto_p s'$. Since the local assumptions guarantee us that $t' \in \text{Red}_{\forall y:B.C'}$, we also know that $t's \in \text{Red}_{C[s/y]}$. We moreover have that $l(s') < l(s)$, and hence, by local inductive hypothesis, we know that $ts' \in \text{Red}_{C[s'/y]}$. But then, since ts is neutral and can only reduce to terms in Red (either in $\text{Red}_{C[s'/y]}$ or in $\text{Red}_{C[s/y]}$), by global inductive hypothesis (CAntiEval), we have that $ts \in \text{Red}_{C[s/y]}$.

Since we just showed that $ts \in \text{Red}_{C[s/y]}$ for any $s \in \text{Red}_B$, we have that $t \in \text{Red}_{\forall y:B.C}$, which means that CAntiEval is verified.

Suppose that $A = \oplus B$.

- CSN. If $t \in \text{Red}_{\oplus B}$, then $t\nu \in \text{Red}_B$. Hence, and by induction hypothesis (CSN), $t\nu$ is strongly normalising. Suppose, by reasoning indirectly, that an infinite reduction sequence of the form $t \mapsto_{p_1} t_1 \mapsto_{p_2} t_2 \dots$ exists, then also an infinite reduction sequence of the form $t\nu \mapsto_{p_1} t_1\nu \mapsto_{p_2} t_2\nu \dots$ exists. Which is absurd since $t\nu$ is strongly normalising. Hence, t is strongly normalising and CSN is verified.
- CEval. Since $t \in \text{Red}_{\oplus B}$, we have, by definition of $\text{Red}_{\oplus B}$, that $t\nu \in \text{Red}_B$. Moreover, if $t \mapsto_p t'$, then $t\nu \mapsto_p t'\nu$. From $t\nu \in \text{Red}_B$ and $t\nu \mapsto_p t'\nu$, by induction hypothesis (CEval), we obtain that $t'\nu \in \text{Red}_{B'}$. From this and by definition of $\text{Red}_{\oplus B}$, we have that $t' \in \text{Red}_{\oplus B'}$, and thus that CEval is verified.
- CAntiEval. We suppose that $t : \oplus B$ is neutral and that, for each t' such that $t \mapsto_p t'$, we have $t' \in \text{Red}_{\oplus B'}$. We need to prove that $t \in \text{Red}_{\oplus B}$. Since t is neutral, $t\nu$ can only reduce to a term of the form $t'\nu$ such that $t \mapsto_p t'$. Since the local assumptions guarantee us that $t' \in \text{Red}_{\oplus B'}$, we also know that $t'\nu \in \text{Red}_{B'}$. But then, since $t\nu$ is neutral and can only reduce to terms in $\text{Red}_{B'}$, by inductive hypothesis (CAntiEval), we have that $t\nu \in \text{Red}_B$. Hence, by definition of $\text{Red}_{\oplus B}$, we have that $t \in \text{Red}_{\oplus B}$ and thus that CAntiEval is verified.

Suppose that $\Phi = \Pi y : B. \Xi$.

- CSN. If $\varphi \in \text{Red}_{\Pi y:B.\Xi}$, then $\varphi x \in \text{Red}_{\Xi[x/y]}$ for any $x : B$. By induction hypothesis (CSN), $\varphi x \in \text{Red}_{\Xi[x/y]}$ implies that φx is strongly normalising. Suppose now, by reasoning indirectly, that an infinite reduction sequence of the form $\varphi \mapsto \varphi_1 \mapsto \varphi_2 \dots$ exists, then also an infinite reduction sequence of the form $\varphi x \mapsto \varphi_1 x \mapsto \varphi_2 x \dots$ exists. Which is absurd since φx is strongly normalising. Hence, t is strongly normalising and CSN is verified.

- CEval. Since $\varphi \in \text{Red}_{\Pi y : B. \Xi}$, we have, by definition of $\text{Red}_{\Pi y : B. \Xi}$, that $\varphi s \in \text{Red}_{\Xi[s/y]}$ for any $s : B$. Moreover, if $\varphi \mapsto \varphi'$, then $\varphi s \mapsto \varphi' s$. From $\varphi s \in \text{Red}_{\Xi[s/y]}$ and $\varphi s \mapsto \varphi' s$, by induction hypothesis (CEval), we obtain that $\varphi' s \in \text{Red}_{\Xi[s/y]}$. Since s is a generic term of type B , from $\varphi' s \in \text{Red}_{\Xi[s/y]}$ and by definition of $\text{Red}_{\Pi y : B. \Xi}$, we have that $\varphi' \in \text{Red}_{\Pi y : B. \Xi}$, and thus that CEval is verified.
- CAntiEval. We suppose that $\varphi : (\Pi y : B. \Xi)$ is neutral and that, for each φ' such that $\varphi \mapsto_p \varphi'$, we have $\varphi' \in \text{Red}_{\Pi y : B. \Xi}$. We need to prove that $\varphi \in \text{Red}_{\Pi y : B. \Xi}$. In order to show this, it is enough to show that $\varphi s \in \text{Red}_{\Xi[s/y]}$ for any $s : B$. Since φ is neutral, then φs can only reduce to terms of the form $\varphi' s$ such that $\varphi \mapsto_p \varphi'$. Now, the local assumptions guarantee us that $\varphi' \in \text{Red}_{\Pi y : B. \Xi}$. Therefore, we also know that $\varphi' s \in \text{Red}_{\Xi[s/y]}$. But then, since φs is neutral and can only reduce to terms in $\text{Red}_{\Xi[s/y]}$, by inductive hypothesis (CAntiEval), we have that $\varphi s \in \text{Red}_{\Xi[s/y]}$ as desired. Considering that we just showed that $\varphi s \in \text{Red}_{\Xi[s/y]}$ for any $s : B$, we have that $\varphi \in \text{Red}_{\Pi y : B. \Xi}$, which means that CAntiEval is verified. □

We now show that the term constructors corresponding to logical introduction rules preserve reducibility.

Lemma 5.2 (Pairing lemma). *If $s \in \text{Red}_A$ and $t \in \text{Red}_B$ then $\langle s, t \rangle \in \text{Red}_{A \wedge B}$.*

Proof. In order to prove the statement, we need to show that $\langle s, t \rangle \pi_0 \in \text{Red}_A$ and $\langle s, t \rangle \pi_1 \in \text{Red}_B$.

Since we have that $s \in \text{Red}_A$ and $t \in \text{Red}_B$ and because of CSN, we know that all sequences of reductions starting from s and t are of finite length. Hence, we can prove the statement by induction on $l(s) + l(t)$ where the function $l(\)$ has as value the length of the longest sequence of reductions starting from its argument.

Let us consider $\langle s, t \rangle \pi_0 : A$. In one step, it can only reduce to one of the following terms:

- $s \in \text{Red}_A$
- $\langle s', t \rangle \pi_0$ where $s \mapsto_p s'$. Since, by CEval, $s' \in \text{Red}_{A'}$ and since $l(s') < l(s)$, we can apply the inductive hypothesis to $\langle s', t \rangle \pi_0$ and conclude that $\langle s', t \rangle \pi_0 \in \text{Red}_{A'}$.
- $\langle s, t' \rangle \pi_0$ where $t \mapsto_p t'$. Since, by CEval, $t' \in \text{Red}_{B'}$ and since $l(t') < l(t)$, we can apply the inductive hypothesis to $\langle s, t' \rangle \pi_0$ and conclude that $\langle s, t' \rangle \pi_0 \in \text{Red}_A$.

Hence, in one step, $\langle s, t \rangle \pi_0$ can only reduce to terms in Red (either in Red_A or in $\text{Red}_{A'}$). By CAntiEval we then know that it is an element of Red_A .

Since we can conclude by an analogous argument that $\langle s, t \rangle \pi_1 \in \text{Red}_B$, we have that $\langle s, t \rangle \in \text{Red}_{A \wedge B}$, as desired. □

Lemma 5.3 (Abstraction lemma). *If, for any $s \in \text{Red}_A$, $t[s/x] \in \text{Red}_{B[s/x]}$, then $\lambda x. t \in \text{Red}_{\forall x : A. B}$.*

Proof. In order to prove the statement, we need to show that, for any $s \in \text{Red}_A$, $(\lambda x. t)s \in \text{Red}_{B[s/x]}$. Since, by Proposition 5.1 (CSN), $s \in \text{Red}_A$ and $t \in \text{Red}_{\forall x : A. B}$ imply that s and t are strongly normalising, we can reason by induction on $l(s) + l(t)$, where $l(\)$ has as value the length of the longest sequence of reductions starting from its argument.

Let us consider the possible outcomes of a one-step reduction of $(\lambda x. t)s$:

- $t[s/x]$, which is assumed to be an element of $\text{Red}_{B[s/x]}$.
- $(\lambda x.t')s$. Since $t \mapsto_p t'$, $l(s) + l(t') < l(s) + l(t)$. We can then use the induction hypothesis to conclude that $(\lambda x.t')s \in \text{Red}'_B[s/x]$.
- $(\lambda x.t)s'$. Since $s \mapsto_p s'$, we also have that $l(s') + l(t) < l(s) + l(t)$. We can then use the induction hypothesis to conclude that $(\lambda x.t)s' \in \text{Red}_B[s'/x]$.

Since all terms to which $(\lambda x.t)s : B[s/x]$ can reduce in one step are elements of Red (of $\text{Red}_{B[s/x]}$, of $B'[s/x]$ or of $B[s'/x]$), by Proposition 5.1 (CEval), we know that $(\lambda x.t)s \in \text{Red}_{B[s/x]}$. Since, moreover, s is a generic element of Red_B , by definition of $\text{Red}_{\forall x:B.C}$, we can conclude, as desired, that $(\lambda x.t) \in \text{Red}_{\forall x:A.B}$. \square

Lemma 5.4 (Nondeterministic choice lemma). *If $s, t \in \text{Red}_A$ then $s(p)t \in \text{Red}_{\oplus A}$.*

Proof. In order to prove the statement, we need to show that $(s(p)t)\nu \in \text{Red}_A$.

Since we have that $s, t \in \text{Red}_A$ and because of CSN, we know that all sequences of reductions starting from s and t are of finite length. Hence, we can prove the statement by induction on $l(s) + l(t)$ where the function $l(\cdot)$ has as value the length of the longest sequence of reductions starting from its argument.

Let us consider $(s(p)t)\nu : A$. In one step, it can only reduce to one of the following terms:

- $s \in \text{Red}_A$
- $t \in \text{Red}_A$
- $(s'(p)t)\nu$ where $s \mapsto_p s'$. Since, by CEval, $s' \in \text{Red}_{A'}$ and since $l(s') < l(s)$, we can apply the inductive hypothesis to $(s'(p)t)\nu$ and conclude that $(s'(p)t)\nu \in \text{Red}_{A'}$.
- $(s(p)t')\nu$ where $t \mapsto_p t'$. Since, by CEval, $t' \in \text{Red}_{A'}$ and since $l(t') < l(t)$, we can apply the inductive hypothesis to $(s(p)t')\nu$ and conclude that $(s(p)t')\nu \in \text{Red}_{A'}$.

Hence, in one step, $(s(p)t)\nu$ can only reduce to terms in Red (either in Red_A or in $\text{Red}_{A'}$). By CAntiEval we then know that it is an element of Red_A .

We have then that $s(p)t \in \text{Red}_{\oplus A}$, as desired. \square

Lemma 5.5 (Abstraction lemma for type constructors). *If, for any $s : A$, $\varphi[s/x] \in \text{Red}_{\Xi[s/x]}$, then $\lambda x.\varphi \in \text{Red}_{\Pi x:A.\Xi}$.*

Proof. In order to prove the statement, we need to show that, for any $s : A$, $(\lambda x.\varphi)s \in \text{Red}_{\Xi[s/x]}$. Since, by Proposition 5.1 (CSN), $\varphi \in \text{Red}_{\Pi x:A.\Xi}$ implies that φ is strongly normalising, we can reason by induction on $l(\varphi)$, where $l(\cdot)$ has as value the length of the longest sequence of reductions starting from its argument.

Let us consider the possible outcomes of a one-step reduction of $(\lambda x.\varphi)s$:

- $\varphi[s/x]$, which is assumed to be an element of $\text{Red}_{\Xi[s/x]}$.
- $(\lambda x.\varphi')s$. Since $\varphi \mapsto_p \varphi'$, we have also $l(\varphi') < l(\varphi)$. We can then use the induction hypothesis to conclude that $(\lambda x.\varphi')s \in \text{Red}_{\Xi[s/x]}$.

Since all type constructors to which $(\lambda x.\varphi)s : \Xi[s/x]$ can reduce in one step are elements of $\text{Red}_{\Xi[s/x]}$, by Proposition 5.1 (CEval), we know that $(\lambda x.\varphi)s \in \text{Red}_{\Xi[s/x]}$. Since, moreover, s is a generic term of type B , by definition of $\text{Red}_{\Pi x:B.\Xi}$, we can conclude, as desired, that $(\lambda x.\varphi) \in \text{Red}_{\Pi x:A.\Xi}$. \square

Theorem 5.6 (Reducibility theorem). *For any type $A : *$ and term $t : A$, $t \in \text{Red}_A$.*

Proof. We prove the following stronger statement:

For any type $F : *$, term $t : F$ with free variables $x_1 : A_1, \dots, x_n : A_n$ and sequence of terms $s_1 \in \text{Red}_{A_1}, \dots, s_n \in \text{Red}_{A_n}$, we have that $t[s_1/x_1, \dots, s_n/x_n] \in \text{Red}_{F[s_1/x_1, \dots, s_n/x_n]}$.

Let us denote by $[\overline{s}/\overline{x}]$ the substitution $[s_1/x_1, \dots, s_n/x_n]$. The proof is by induction on the number of symbol occurrences in t .

- $t = x_i$ for some $1 \leq i \leq n$. Since $t[\overline{s}/\overline{x}] = s_i$ and $s_i \in \text{Red}_{A_i}$ by assumption, we obviously have $t[\overline{s}/\overline{x}] \in \text{Red}_{F[\overline{s}/\overline{x}]}$ where $F[\overline{s}/\overline{x}] = A_i$.
- $t = \langle u, v \rangle$. By induction hypothesis, we have that $u[\overline{s}/\overline{x}] \in \text{Red}_{F_1[\overline{s}/\overline{x}]}$ and $v[\overline{s}/\overline{x}] \in \text{Red}_{F_2[\overline{s}/\overline{x}]}$ for suitable types F_1, F_2 . By Lemma 5.2, we know that $\langle u[\overline{s}/\overline{x}], v[\overline{s}/\overline{x}] \rangle \in \text{Red}_{F_1[\overline{s}/\overline{x}] \wedge F_2[\overline{s}/\overline{x}]}$. But $\langle u[\overline{s}/\overline{x}], v[\overline{s}/\overline{x}] \rangle = \langle u, v \rangle[\overline{s}/\overline{x}]$ and $F_1[\overline{s}/\overline{x}] \wedge F_2[\overline{s}/\overline{x}] = (F_1 \wedge F_2)[\overline{s}/\overline{x}]$, and hence $\langle u, v \rangle[\overline{s}/\overline{x}] \in \text{Red}_{(F_1 \wedge F_2)[\overline{s}/\overline{x}]}$ as desired.
- $t = u\pi_0$ and $t = u\pi_1$. By induction hypothesis, $u[\overline{s}/\overline{x}] \in \text{Red}_{(F_1 \wedge F_2)[\overline{s}/\overline{x}]}$ for any sequence $\overline{u} = s_1, \dots, s_n$ such that $s_1 \in \text{Red}_{A_1}, \dots, s_n \in \text{Red}_{A_n}$. By definition of $\text{Red}_{(F_1 \wedge F_2)[\overline{s}/\overline{x}]}$ and since $(F_1 \wedge F_2)[\overline{s}/\overline{x}] = F_1[\overline{s}/\overline{x}] \wedge F_2[\overline{s}/\overline{x}]$, this means that $(u[\overline{s}/\overline{x}])\pi_i \in \text{Red}_{F_i[\overline{s}/\overline{x}]}$ for $i \in \{0, 1\}$. But since $(u[\overline{s}/\overline{x}])\pi_i = (u\pi_i)[\overline{s}/\overline{x}]$, we have that $(u\pi_i)[\overline{s}/\overline{x}] \in \text{Red}_{F_i[\overline{s}/\overline{x}]}$, as desired.
- $t = \lambda y. u$. By induction hypothesis, for suitable types F_1 and F_2 , we have that $u[\overline{s}/\overline{x}, v/y] \in \text{Red}_{F_2[\overline{s}/\overline{x}, v/y]}$ for any $v \in \text{Red}_{F_1}$. By Lemma 5.3, this implies that $\lambda y. (u[\overline{s}/\overline{x}]) \in \text{Red}_{\forall y : F_1. (F_2[\overline{s}/\overline{x}])}$. But the assumption that \overline{x} are all free in $t = \lambda y. u$ implies that $y \notin \overline{x}$, which in turn implies that $\lambda y. (u[\overline{s}/\overline{x}]) = (\lambda y. u)[\overline{s}/\overline{x}]$ and that $\forall y : F_1. (F_2[\overline{s}/\overline{x}]) = (\forall y : F_1. F_2)[\overline{s}/\overline{x}]$. Hence, $(\lambda y. u)[\overline{s}/\overline{x}] \in \text{Red}_{(\forall y : F_1. F_2)[\overline{s}/\overline{x}]}$, as desired.
- $t = uv$. By induction hypothesis, $u[\overline{s}/\overline{x}] \in \text{Red}_{(\forall y : F_1. F_2)[\overline{s}/\overline{x}]}$ and $v[\overline{s}/\overline{x}] \in \text{Red}_{F_1[\overline{s}/\overline{x}]}$ for any sequence $\overline{s} = s_1, \dots, s_n$ such that $s_1 \in \text{Red}_{A_1}, \dots, s_n \in \text{Red}_{A_n}$. Since $(\forall y : F_1. F_2)[\overline{s}/\overline{x}] = \forall y : (F_1[\overline{s}/\overline{x}]).(F_2[\overline{s}/\overline{x}])$ and by definition of $\forall y : (F_1[\overline{s}/\overline{x}]).(F_2[\overline{s}/\overline{x}])$, this means that $(u[\overline{s}/\overline{x}])(v[\overline{s}/\overline{x}]) \in \text{Red}_{F_2[\overline{s}/\overline{x}]}$. But since $(u[\overline{s}/\overline{x}])(v[\overline{s}/\overline{x}]) = (uv)[\overline{s}/\overline{x}]$, we have that $(uv)[\overline{s}/\overline{x}] \in \text{Red}_{F_2[\overline{s}/\overline{x}]}$, as desired.
- $t = u(p)v$. By induction hypothesis, we have that $u[\overline{s}/\overline{x}], v[\overline{s}/\overline{x}] \in \text{Red}_{F_1[\overline{s}/\overline{x}]}$ for a suitable type F_1 . By Lemma 5.4, we know that $u[\overline{s}/\overline{x}](p)v[\overline{s}/\overline{x}] \in \text{Red}_{\oplus F_1[\overline{s}/\overline{x}]}$. But $u[\overline{s}/\overline{x}](p)v[\overline{s}/\overline{x}] = (u(p)v)[\overline{s}/\overline{x}]$ and $\oplus F_1[\overline{s}/\overline{x}] = (\oplus F_1)[\overline{s}/\overline{x}]$. Hence, $(u(p)v)[\overline{s}/\overline{x}] \in \text{Red}_{(\oplus F_1)[\overline{s}/\overline{x}]}$ as desired.
- $t = u\nu$. By induction hypothesis, $u[\overline{s}/\overline{x}] \in \text{Red}_{(\oplus F_1)[\overline{s}/\overline{x}]}$ for any sequence $\overline{u} = s_1, \dots, s_n$ such that $s_1 \in \text{Red}_{A_1}, \dots, s_n \in \text{Red}_{A_n}$. Since $(\oplus F_1)[\overline{s}/\overline{x}] = \oplus F_1[\overline{s}/\overline{x}]$ and by definition of $\text{Red}_{\oplus F_1[\overline{s}/\overline{x}]}$, this means that $(u[\overline{s}/\overline{x}])\nu \in \text{Red}_{F_1[\overline{s}/\overline{x}]}$. But since $(u[\overline{s}/\overline{x}])\nu = (u\nu)[\overline{s}/\overline{x}]$, we have that $(u\nu)[\overline{s}/\overline{x}] \in \text{Red}_{F_1[\overline{s}/\overline{x}]}$, as desired. □

Theorem 5.7 (Reducibility theorem for type constructors). *For any kind $\Phi : \square$ and type constructor $\varphi : \Phi$, $\varphi \in \text{Red}_\Phi$.*

Proof. We prove the following stronger statement:

For any kind $\Phi : \square$ and type constructor $\varphi : \Phi$ with free variables $x_1 : A_1, \dots, x_n : A_n$ and sequence of terms $s_1 : A_1, \dots, s_n : A_n$, we have that $\varphi[s_1/x_1, \dots, s_n/x_n] \in \text{Red}_{\Phi[s_1/x_1, \dots, s_n/x_n]}$.

Let us denote by $[\bar{s}/\bar{x}]$ the substitution $[s_1/x_1, \dots, s_n/x_n]$. The proof is by induction on the number of symbol occurrences in φ .

- $\varphi = \alpha$. Now, if $\varphi[\bar{s}/\bar{x}] = \alpha$ then $\varphi[\bar{s}/\bar{x}] = \varphi$. Moreover, since $\alpha : \Phi$ is normal and neutral we can infer by CAntiEval that $\alpha \in \text{Red}_\Phi$. Therefore, $\varphi \in \text{Red}_\Phi$.
- $\varphi = \lambda y.\psi$. By induction hypothesis, for a suitable kind Φ and type F , we have that $\psi[\bar{s}/\bar{x}, v/y] \in \text{Red}_{\Phi[\bar{s}/\bar{x}, v/y]}$ for any $v : F$. By Lemma 5.5, this implies that $\lambda y.(\psi[\bar{s}/\bar{x}]) \in \text{Red}_{\Pi y : F.(\Phi[\bar{s}/\bar{x}])}$. But the assumption that \bar{x} are all free in $\varphi = \lambda y.\psi$ implies that $y \notin \bar{x}$, which in turn implies that $\lambda y.(\psi[\bar{s}/\bar{x}]) = (\lambda y.\psi)[\bar{s}/\bar{x}]$ and that $\Pi y : F.(\Phi[\bar{s}/\bar{x}]) = (\Pi y : F.\Phi)[\bar{s}/\bar{x}]$. Hence, $(\lambda y.\psi)[\bar{s}/\bar{x}] \in \text{Red}_{(\Pi y : F.\Phi)[\bar{s}/\bar{x}]}$, as desired.
- $\varphi = \psi v$. By induction hypothesis, $\psi[\bar{s}/\bar{x}] \in \text{Red}_{(\forall y : F.\Phi)[\bar{s}/\bar{x}]}$ and $v[\bar{s}/\bar{x}] : F[\bar{s}/\bar{x}]$ for any sequence $\bar{s} = s_1, \dots, s_n$ such that $s_1 \in \text{Red}_{A_1}, \dots, s_n \in \text{Red}_{A_n}$. Since $(\Pi y : F.\Phi)[\bar{s}/\bar{x}] = \Pi y : (F[\bar{s}/\bar{x}]).(\Phi[\bar{s}/\bar{x}])$, by definition of $\Pi y : (F[\bar{s}/\bar{x}]).(\Phi[\bar{s}/\bar{x}])$, we have that $(\psi[\bar{s}/\bar{x}])(v[\bar{s}/\bar{x}]) \in \text{Red}_{\Phi[\bar{s}/\bar{x}]}$. But since $(\psi[\bar{s}/\bar{x}])(v[\bar{s}/\bar{x}]) = (\psi v)[\bar{s}/\bar{x}]$, we obtain $(\psi v)[\bar{s}/\bar{x}] \in \text{Red}_{\Phi[\bar{s}/\bar{x}]}$, as desired.

□

Corollary (Strong normalisation). *For any type $A : *$ and term $t : A$, t is strongly normalising. For any kind $\Phi : \square$ and type constructor $\varphi : \Phi$, φ is strongly normalising.*

Proof. We just need to apply Theorems 5.6 and 5.7 with $s_i = x_i$ for each $i \in \{1 \dots, n\}$. The result then follows by CSN. □

6 Conclusion

References

- [Girard et al., 1989] Girard, J.-Y., Lafont, Y., and Taylor, P. (1989). *Proofs and Types*. Cambridge University Press.
- [Sørensen and Urzyczyn, 2006] Sørensen, M. H. and Urzyczyn, P. (2006). *Lectures on the Curry-Howard isomorphism*. Elsevier.

$$\frac{\Gamma, x : A \vdash t : B}{\Gamma \vdash (\lambda x : A.t) : (\forall x : A.B)} \quad \frac{\Gamma \vdash t : (\forall x : A.B) \quad \Gamma \vdash s : A}{\Gamma \vdash ts : B[s/x]}$$

$$\frac{\Gamma \vdash t : \perp \quad \Gamma \vdash P : *}{\Gamma \vdash \text{efq}(t) : P} \quad \text{where } P \text{ does not contain } \forall$$

$$\frac{\Gamma \vdash t : A \quad \Gamma \vdash s : B}{\Gamma \vdash \langle s, t \rangle : A \wedge B} \quad \frac{\Gamma \vdash t : A_0 \wedge A_1}{\Gamma \vdash t\pi_i : A_i} \quad \text{where } i \in \{0, 1\}$$

$$\frac{\Gamma \vdash t : A \quad \Gamma \vdash s : A}{\Gamma \vdash t(p)s : \oplus A} \quad \frac{\Gamma \vdash A : *}{\Gamma \vdash o : \Sigma A}$$

where o is an oracle constant and A is the type associated to it by definition

$$\frac{\Gamma \vdash t : \oplus A}{\Gamma \vdash t\nu : A} \quad \frac{\Gamma \vdash o : \Sigma A}{\Gamma \vdash o\nu : A}$$

$$\frac{\Gamma \vdash o_1 : \forall x : A. \Sigma B \quad \Gamma \vdash t : A}{\Gamma \vdash o_1 t : \Sigma(B[t/x])} \quad \frac{\Gamma \vdash o_0 \nu : A}{\Gamma \vdash \langle o_0 \nu, \dots, o_0 \nu \rangle : A^n}$$

$$\frac{\Gamma \vdash o_1 : \forall x : A. \Sigma B \quad \Gamma \vdash t_1 : A \quad \dots \quad \Gamma \vdash t_n : A}{\Gamma \vdash \langle (o_1 t_1)\nu, \dots, (o_1 t_n)\nu \rangle : B[t/x]^n}$$

where o_1 is a 1-ary oracle constant, o_0 is a 0-ary oracle constant,

$$\frac{\Gamma \vdash t : C \quad \Gamma \vdash D : *}{\Gamma \vdash t : D} \quad \text{where } C \equiv_\beta D \text{ according to Definition 3.2}$$

Figure 4: Type assignment and conversion rules

$$\begin{aligned}
& (\lambda x : A.t)s \mapsto_1 t[s/x] \\
& \langle t_0, t_1 \rangle \pi_i \mapsto_1 t_i \quad \text{for } i \in \{0, 1\} \\
& (t\langle p \rangle s)\nu \mapsto_p t \quad (t\langle p \rangle s)\nu \mapsto_{1-p} s
\end{aligned}$$

$$\mathcal{C}[o_0\nu]_1 \dots [o_0\nu]_n \mapsto_1 \mathcal{C}[s_1]_1 \dots [s_n]_n$$

where o_0 is a 0-ary oracle constant,
and $f_{o_0}(\mathcal{C}[\]_1 \dots [\]_n, i) = s_i$ for $1 \leq i \leq n$

$$\mathcal{C}[(o_1 t_1)\nu]_1 \dots [(o_1 t_n)\nu]_n \mapsto_1 \mathcal{C}[s_1]_1 \dots [s_n]_n$$

where o_1 is a 1-ary oracle constant,
 $\mathcal{C}[(o_1 t_1)\nu]_1 \dots [(o_1 t_n)\nu]_n = \mathcal{D}[o_1]_1 \dots [o_1]_n$,
and $f_{o_1}(\mathcal{D}[\]_1 \dots [\]_n, i) = s_i$ for $1 \leq i \leq n$

Figure 5: Evaluation reductions