

Traveling waves in reaction-diffusion-convection equations with combustion nonlinearity

Pavel Drábek¹ and Michaela Zahradníková^{✉2}

¹Department of Mathematics and NTIS, Faculty of Applied Sciences, University of West Bohemia,
Univerzitní 8, 301 00 Plzeň, Czech Republic

²Department of Mathematics, Faculty of Science, University of South Bohemia,
Braníšovská 1760, 370 05 České Budějovice, Czech Republic

Abstract. This paper concerns the existence and properties of traveling wave solutions to reaction-diffusion-convection equations on the real line. We consider a general diffusion term involving the p -Laplacian and combustion-type reaction term. We extend and generalize the results established for $p = 2$ to the case of singular and degenerate diffusion. Our approach allows for non-Lipschitz reaction as well. We also discuss the shape of the traveling wave profile near equilibria, assuming power-type behavior of the reaction and diffusion terms.

Keywords: reaction-diffusion-convection equation, combustion-type reaction, traveling waves.

2020 Mathematics Subject Classification: Primary 35K57, 35C07; Secondary 34A12, 35K65.

1 Introduction

In this paper, we study the existence of traveling wave solutions for the reaction-diffusion-convection equation

$$v_t = [D(v)|v_x|^{p-2}v_x]_x + (H(v))_x + g(v), \quad x \in \mathbb{R}, t \geq 0. \quad (1.1)$$

Here $p > 1$, $D \in C^1(0, 1)$ is a diffusion coefficient with $D > 0$ in $(0, 1)$, $H \in C^1[0, 1]$ represents a nonlinear convective flux function, and the reaction term $g \in C[0, 1]$ satisfies

$$g(v) = 0 \text{ in } [0, \theta], \quad g(v) > 0 \text{ in } (\theta, 1), \quad g(1) = 0 \quad (1.2)$$

for some $\theta \in (0, 1)$. Let us emphasize that the diffusion coefficient D might degenerate or have a singularity at one or both endpoints. A traveling wave solution (t.w.s. for short) is a solution of the form $v(x, t) = u(x - ct)$, $c \in \mathbb{R}$, connecting the equilibria 0 and 1. Besides the unknown profile u , the constant speed c also needs to be determined.

Reaction-diffusion equations with and without convective effects are used to describe a variety of phenomena in biology, chemistry and physics. We refer to [7] for an overview of classical applications in the case $p = 2$, involving different types of reaction terms. Nonlinearity g satisfying (1.2) arises in combustion models with an ignition temperature assumption, in which the reaction starts only after the temperature reaches the threshold value θ , cf. [3]. The diffusion

[✉] Corresponding author. Orcid ID: 0000-0003-3303-6969. Email: zahram05@prf.jcu.cz

coefficient D is usually assumed to be strictly positive in $[0, 1]$. However, modeling of certain dispersal phenomena suggests also coefficients with degenerations or singularities at equilibria, see, e.g., [9], [12]. More recently, p -Laplacian-type diffusion has also been considered in literature, see [1, 6, 8] and their references. The p -Laplacian operator itself appears for instance in models derived from the power-type Darcy's law, cf. [2].

If H is constant, i.e., no convective effects are present, equation (1.1) possesses a unique nonincreasing t.w.s. with a positive wave speed c^* . For $p = 2$, this result was proved in [11] assuming $g \in C[0, 1]$ and $D \in C^1[0, 1]$ strictly positive in $[0, 1]$. Furthermore, if stronger regularity conditions on g are imposed, the t.w.s. is strictly monotone on \mathbb{R} . In our previous work [4], we proved the existence of a nonincreasing solution to (1.1) without convection in an even more general setting. In particular, we considered a discontinuous coefficient D with finitely many jumps in $(0, 1)$ and with possible singularities and/or degenerations at equilibria 0 and 1. This required a new concept of non-classical non-smooth solution, but the existence result remained the same also for $p > 1$.

In [10], the authors investigate the effect of convection on the existence of t.w.s. in the case $p = 2$ and $D > 0$ in $[0, 1]$. They derive existence and non-existence results based on whether, in some sense, H prevails over or is weak compared to the terms D and g . Our paper focuses on the same phenomena but we consider arbitrary $p > 1$ as well as a more general diffusion coefficient D (possibly degenerate or singular at 0 and 1) and a non-Lipschitz reaction g . The main contribution of our paper consists in showing how the results from [10] extend to our more general case. We also note that our method of proof differs from that in [10].

The paper is organized as follows. In Section 2, we present the definition of possibly non-smooth traveling wave profile u and our main existence and nonexistence results. In Section 3, we first discuss monotonicity property of the wave profile, which then allows us to reduce the second-order problem for the unknown profile u to a first-order one on a bounded interval. Proofs of the main results are provided in Sections 4 (nonexistence) and 5 (existence). Auxiliary lemmas referenced in Sections 3 and 5 can be found in the Appendices A and B, respectively. Finally, Section 6 is dedicated to discussions concerning the asymptotic properties of solutions, while assuming power-type behavior of the terms D and g .

2 Main results

Without loss of generality, we assume $H(0) = 0$ and write

$$H(u) := \int_0^u h(s) \, ds.$$

Here $h(u) = \frac{d}{du}H(u)$, $u \in [0, 1]$, is the convective velocity. When looking for traveling wave solutions $v(x, t) = u(\xi)$, $\xi := x - ct$, the partial differential equation (1.1) becomes a boundary value problem for the second-order ordinary differential equation on the real line

$$\begin{cases} (D(u)|u'|^{p-2}u')' + (c + h(u))u' + g(u) = 0, \\ u(-\infty) = 1, \quad u(+\infty) = 0 \end{cases} \quad (2.1)$$

where $u' := \frac{du}{d\xi}$ stands for the derivative with respect to the wave coordinate ξ . Since the equation in (2.1) is autonomous, its solutions are invariant under translations. Therefore, we can always normalize solutions of (2.1) as $u(0) = \theta$.

Before presenting the main results of this paper, we first provide the definition of solution to (2.1) on \mathbb{R} . This concept accounts for situations when the profile u reaches one or both equilibria 0 and 1.

Definition 2.1. A continuous function $u : \mathbb{R} \rightarrow [0, 1]$ is a solution of (2.1) if

- (a) $u \in C^1(I_u)$, $I_u := \{\xi \in \mathbb{R} : 0 < u(\xi) < 1\}$, and the equation in (2.1) holds at every point of I_u ;
- (b) the function $\xi \mapsto D(u(\xi))|u'(\xi)|^{p-2}u'(\xi)$ is continuous on \mathbb{R} and $D(u(\xi))|u'(\xi)|^{p-2}u'(\xi) \rightarrow 0$ as $u(\xi) \rightarrow 0$ and $u(\xi) \rightarrow 1$;
- (c) (boundary conditions) $u(\xi) \rightarrow 1$ as $\xi \rightarrow -\infty$ and $u(\xi) \rightarrow 0$ as $\xi \rightarrow +\infty$.

Remark 2.2. At the beginning of Section 3, we prove that I_u is an open interval, bounded or unbounded, and that $u'(\xi) < 0$ for all $\xi \in I_u$. If $p = 2$, $D \in C^1[0, 1]$ with $D > 0$ in $[0, 1]$ and g is a Lipschitz function on $[0, 1]$, then $I_u = \mathbb{R}$ and $u \in C^2(\mathbb{R})$ is a classical solution, cf. [10]. Note that in this case we have $u'(\xi) \rightarrow 0$ as $\xi \rightarrow \pm\infty$.

On the other hand, if $p \neq 2$ and $D \in C^1(0, 1)$ is singular or degenerate at 0 and/or 1, then I_u might be an interval of finite length, i.e., $I_u = (\xi_1, \xi_2)$, $\xi_1, \xi_2 \in \mathbb{R}$. Moreover, the derivative $u'(\xi_i)$, $i = 1, 2$, need not exist. We provide detailed discussion of these cases in Section 6 based on asymptotic properties of D and g which yield different shapes of traveling wave profile.

Remark 2.3. Notice that it follows from Definition 2.1 (a) that the function

$$\xi \mapsto D(u(\xi))|u'(\xi)|^{p-2}u'(\xi)$$

belongs to $C^1(I_u)$. In particular, the function $\xi \mapsto |u'(\xi)|^{p-2}u'(\xi)$ also belongs to $C^1(I_u)$ due to our assumption $D \in C^1(0, 1)$, $D > 0$ in $(0, 1)$.

In what follows, we denote

$$h_m := \min_{u \in [0, 1]} h(u)$$

and p' stands for the exponent conjugate to p , i.e., $p' = \frac{p}{p-1}$. Furthermore, we assume

$$\int_0^1 D^{p'-1}(u)g(u) \, du < +\infty. \quad (2.2)$$

Theorem 2.4 (Nonexistence). *Let*

$$H(\theta) \geq \theta h_m + \left(p' \int_0^1 D^{p'-1}(u)g(u) \, du \right)^{\frac{1}{p'}}. \quad (2.3)$$

Then the boundary value problem (2.1) has no solution for any $c > -h_m$. If strict inequality holds in (2.3), there is no solution for any $c \geq -h_m$.

In Section 4, we prove that $c \geq -h(0)$ is a necessary condition for the existence of solutions. An immediate consequence is the following corollary, which addresses the nonexistence of solution for any real value of c .

Corollary 2.5. *If strict inequality holds in (2.3) and $h_m = h(0)$, then (2.1) has no solution for any $c \in \mathbb{R}$.*

Let

$$k = k(p) = \begin{cases} \frac{1}{2^{p'-1}-1} & \text{if } 1 < p < 2, \\ 1 & \text{if } p = 2, \\ \frac{p'}{p'-1 + \frac{1+p'(p'-1)^{\frac{1}{p'-2}} + (p'-1)^{\frac{p'}{p'-2}}}{\left(1+(p'-1)^{\frac{1}{p'-2}}\right)^{p'}}} & \text{if } p > 2. \end{cases} \quad (2.4)$$

Then $k = k(p)$ is a continuous function in $(1, +\infty)$ satisfying

$$\lim_{p \rightarrow 1^+} k(p) = 0 \quad \text{and} \quad \lim_{p \rightarrow +\infty} k(p) = \frac{1}{2}.$$

Theorem 2.6 (Existence). *Let*

$$H(1) \leq h_m + \left(k(p) \int_0^1 D^{p'-1}(u)g(u) \, du \right)^{\frac{1}{p'}}. \quad (2.5)$$

Then there exists a unique $c = c^ > -h_m$ such that the boundary value problem (2.1) has a unique (up to translation) solution $u = u(\xi)$. Moreover, the solution u is strictly decreasing on I_u and c^* satisfies*

$$c^* < \frac{1}{\theta} \left[\left(p' \int_0^1 D^{p'-1}(u)g(u) \, du \right)^{\frac{1}{p'}} - H(\theta) \right] - h_m. \quad (2.6)$$

Clearly, if $h_m \leq 0$ it follows from the above theorem that the unique wave speed c^* is positive. The following result concerns the existence of a positive wave speed in the case $h_m > 0$.

Theorem 2.7 (Positive wave speed c). *If $h(u) > 0$, $u \in [0, 1]$, and*

$$H(1) \leq \left(k(p) \int_0^1 D^{p'-1}(u)g(u) \, du \right)^{\frac{1}{p'}}, \quad (2.7)$$

then $c^ > 0 > -h_m$.*

Remark 2.8. The expression for $k(p)$ given in (2.4) is an optimal value of the constant in Theorems 2.6 and 2.7. For $p = 2$ it coincides with the estimates derived in [10] by a different approach.

3 Reduction to a first order problem

In this section, we establish our main tool for investigating the existence and nonexistence of solutions to (2.1). In particular, we show that (2.1) can be transformed into a first-order boundary value problem. First, we prove that each solution to (2.1) is a decreasing function in I_u .

Proposition 3.1. *Let u be a solution of (2.1). There exist $-\infty \leq \xi_1 < \xi_2 \leq +\infty$ such that $u \equiv 1$ in $(-\infty, \xi_1]$, $u \equiv 0$ in $[\xi_2, +\infty)$ and $u'(\xi) < 0$ for any $\xi \in (\xi_1, \xi_2)$.*

Proof. First, we show that the derivative of a solution to (2.1) does not vanish in the set $I_u = \{\xi \in \mathbb{R} : 0 < u(\xi) < 1\}$. Indeed, let $\xi_0 \in I_u$ be such that $0 < u(\xi_0) \leq \theta$. If $u'(\xi_0) = 0$ then it follows from Lemma A.1 that the boundary conditions in (2.1) are not satisfied, a contradiction. Now consider $\xi_0 \in I_u$, $\theta < u(\xi_0) < 1$ with $u'(\xi_0) = 0$. Then

$$(D(u(\xi))|u'(\xi)|^{p-2}u'(\xi))' \Big|_{\xi=\xi_0} = -g(u(\xi_0)) < 0.$$

It follows from Lemma A.2 that ξ_0 must be the point of strict local maximum of u and therefore $\lim_{\xi \rightarrow -\infty} u(\xi) \neq 1$, again a contradiction.

Next we prove that $u'(\xi) < 0$ for all $\xi \in I_u$, i.e., the solution cannot “switch” from 0 to 1 and back again finitely many times (while still satisfying the boundary conditions). To this end, we observe that $c > -H(1)$ is a necessary condition for the existence of solution to (2.1). Indeed, integrating the equation in (2.1) we obtain

$$\begin{aligned} D(u(\xi))|u'(\xi)|^{p-2}u'(\xi) - D(u(\hat{\xi}))|u'(\hat{\xi})|^{p-2}u'(\hat{\xi}) + c(u(\xi) - u(\hat{\xi})) \\ + H(u(\xi)) - H(u(\hat{\xi})) + \int_{\hat{\xi}}^{\xi} g(u(\zeta)) d\zeta = 0, \quad \xi, \hat{\xi} \in \mathbb{R}. \end{aligned}$$

Passing to the limits $\xi \rightarrow +\infty$, $\hat{\xi} \rightarrow -\infty$ and taking into account parts (b) and (c) of Definition 2.1 yields

$$c + H(1) - H(0) = \int_{-\infty}^{+\infty} g(u(\zeta)) d\zeta.$$

Since $H(0) = 0$ and the integral on the left-hand side is positive, we conclude that $c > -H(1)$.

Suppose that there exist $\underline{\xi}, \bar{\xi} \in \mathbb{R}$ such that $u(\underline{\xi}) = 0$, $u(\bar{\xi}) = 1$ and $u'(\xi) > 0$ for all $\xi \in (\underline{\xi}, \bar{\xi})$. Integrating the equation in (2.1) from $\underline{\xi}$ to $\bar{\xi}$ and employing the same arguments as above, we arrive at

$$c + H(1) = - \int_{\underline{\xi}}^{\bar{\xi}} g(u(\zeta)) d\zeta < 0,$$

i.e., $c < -H(1)$, a contradiction.

Therefore, there exist $-\infty \leq \xi_1 < \xi_2 \leq +\infty$ such that $u \equiv 1$ in $(-\infty, \xi_1]$, $u \equiv 0$ in $[\xi_2, +\infty)$ and $u'(\xi) < 0$ for any $\xi \in (\xi_1, \xi_2)$. This concludes the proof. \square

In particular, it follows from Proposition 3.1 that $I_u = (\xi_1, \xi_2)$ is an open interval, bounded or unbounded.

Following substitutions from [6, p. 174], we set

$$-w(u) := D(u)|u'|^{p-2}u'. \tag{3.1}$$

Since $u'(\xi) < 0$ for all $\xi \in (\xi_1, \xi_2)$, we have $w = w(u) > 0$ in $(0, 1)$ and w satisfies

$$\frac{1}{p'D^{p'-1}(u)} \frac{d}{du} w^{p'}(u) - (c + h(u)) \left(\frac{w(u)}{D(u)} \right)^{p'-1} + g(u) = 0, \quad u \in (0, 1).$$

Put

$$y(u) := w^{p'}(u) > 0.$$

Then $y = y(u)$ solves the equation

$$y'(u) = p' \left[(c + h(u))(y^+(u))^{\frac{1}{p'}} - f(u) \right], \quad u \in (0, 1), \tag{3.2}$$

where $y' = \frac{dy}{du}$, $y^+(u) := \max\{y(u), 0\}$ and

$$f(u) := D^{p'-1}(u)g(u). \quad (3.3)$$

In terms of y , part (b) of Definition 2.1 translates to

$$y(0) := \lim_{u \rightarrow 0^+} y(u) = 0, \quad y(1) := \lim_{u \rightarrow 1^-} y(u) = 0. \quad (3.4)$$

It follows from (3.1) that

$$\frac{\partial \xi}{\partial u} = - \left(\frac{D(u)}{w(u)} \right)^{p'-1}$$

and therefore, since $u(0) = \theta$,

$$\xi(u) = - \int_{\theta}^u \left(\frac{D(s)}{w(s)} \right)^{p'-1} ds = - \int_{\theta}^u \frac{D^{p'-1}(s)}{y^{\frac{1}{p}}(s)} ds, \quad u \in (0, 1). \quad (3.5)$$

Since $\xi = \xi(u)$ maps $(0, 1)$ onto (ξ_1, ξ_2) , we have

$$\xi_1 = - \int_{\theta}^1 \frac{D^{p'-1}(s)}{y^{\frac{1}{p}}(s)} ds \quad \text{and} \quad \xi_2 = \int_0^{\theta} \frac{D^{p'-1}(s)}{y^{\frac{1}{p}}(s)} ds. \quad (3.6)$$

It follows from the above calculations that the existence of a monotone solution to (2.1) implies the existence of a positive solution to (3.2), (3.4) which, in addition, satisfies (3.6). Proceeding as in [5, Proposition 3.3] where $h \equiv 0$, it can be shown that these problems are equivalent. We thus have the following assertion.

Proposition 3.2. *Let (2.2) hold. A function $u : \mathbb{R} \rightarrow [0, 1]$ is a unique solution (up to translation) of (2.1) in the sense of Definition 2.1 if and only if $y : [0, 1] \rightarrow \mathbb{R}$ is a unique positive solution of (3.2), (3.4).*

4 Proof of nonexistence results

Due to Proposition 3.2, to prove the nonexistence results, it suffices to show that the first-order boundary value problem (3.2), (3.4) does not admit positive solutions for the given values of c . First, we notice that

$$c \geq -h(0) \quad (4.1)$$

is a necessary condition for the existence of a positive solution of (3.2), (3.4). Indeed, if $c < -h(0)$ then, by the continuity of h , there exists $\delta > 0$ such that $c < -h(u)$ for all $u \in [0, \delta]$. Integrating the equation (3.2) over $[0, \delta]$ and taking into account $y_c(0) = 0$ together with $c + h(u) < 0$ in $[0, \delta]$, we arrive at

$$y_c(\delta) = p' \int_0^{\delta} (c + h(u))(y_c^+(u))^{\frac{1}{p}} du < 0,$$

a contradiction with the positivity of solution $y_c = y_c(u)$.

Proof of Theorem 2.4. We proceed by contradiction and assume that $c > -h_m$ and $y_c = y_c(u)$ is a positive solution of (3.2), (3.4). Integrating the equation (3.2) over $(\theta, 1)$ and using (3.4) yields

$$y_c(\theta) = -p' \int_{\theta}^1 (c + h(\sigma))(y_c(\sigma))^{\frac{1}{p}} d\sigma + p' \int_{\theta}^1 f(\sigma) d\sigma < p' \int_{\theta}^1 f(\sigma) d\sigma, \quad (4.2)$$

where f is given by (3.3). On the other hand, since $f \equiv 0$ on $(0, \theta)$ the equation (3.2) is separable on $(0, \theta)$. Using (3.4) we obtain

$$y_c^{\frac{1}{p'}}(\theta) = c\theta + H(\theta). \quad (4.3)$$

It follows from (4.2), (4.3) and the condition (2.3) that

$$\left(p' \int_0^1 f(\sigma) d\sigma\right)^{\frac{1}{p'}} > y_c^{\frac{1}{p'}}(\theta) = c\theta + H(\theta) > -h_m\theta + H(\theta) \geq \left(p' \int_0^1 f(\sigma) d\sigma\right)^{\frac{1}{p'}},$$

a contradiction.

Assuming strict inequality in (2.3) and $c \geq -h_m$, we would arrive at

$$\left(p' \int_0^1 f(\sigma) d\sigma\right)^{\frac{1}{p'}} \geq y_c^{\frac{1}{p'}}(\theta) = c\theta + H(\theta) \geq -h_m\theta + H(\theta) > \left(p' \int_0^1 f(\sigma) d\sigma\right)^{\frac{1}{p'}},$$

again a contradiction. This concludes the proof. \square

5 Proof of existence results

Proof of Theorem 2.6. We first prove the statement of Theorem 2.6 assuming that $h_m = 0$, i.e., we will show that if

$$H(1) \leq \left(k(p) \int_0^1 D^{p'-1}(u)g(u) du\right)^{\frac{1}{p'}},$$

then there exists a unique positive value $c = c^*$ for which (2.1) admits a solution. This result can then be applied to the case of a more general $h \in C[0, 1]$ with $h_m \neq 0$ by means of a suitable shift, as discussed at the end of this section.

Thanks to the equivalence established in Proposition 3.2, we proceed by investigating the initial value problem

$$\begin{cases} y'_c(u) = p' \left[(c + h(u))(y_c^+(u))^{\frac{1}{p}} - f(u) \right], & u \in (0, 1), \\ y_c(1) = 0, \end{cases} \quad (5.1)$$

where f is given by (3.3). Let $c \geq 0$. Since $c + h(u) \geq 0$ for all $u \in [0, 1]$, the function

$$y \mapsto (c + h(u))(y^+)^{\frac{1}{p}}, \quad u \in [0, 1],$$

satisfies one-sided Lipschitz condition and it follows from [5, Lemma 4.1 and Lemma 4.3], where we replace c by $c + h(u)$, that (5.1) has a unique global solution $y_c = y_c(u)$ defined on $[0, 1]$. Our aim is to show that there exists $c > 0$ such that $y_c(u) > 0$ if $u \in (0, 1)$ and $y_c(0) = 0$.

First, let us observe that $f(u) > 0$ in $(\theta, 1)$ implies that

$$y_c(u) > 0 \quad \text{for } u \in (\theta, 1), \quad (5.2)$$

and

$$y_c(\theta) = -p' \int_{\theta}^1 (c + h(\sigma))(y_c(\sigma))^{\frac{1}{p}} d\sigma + p' \int_{\theta}^1 f(\sigma) d\sigma < p' \int_0^1 f(\sigma) d\sigma. \quad (5.3)$$

According to Lemma B.2, for any $p > 1$ we have

$$y_0^{\frac{1}{p'}}(\theta) > H(\theta). \quad (5.4)$$

In particular, $y_0(\theta) > 0$ and hence there exists $0 < \delta \leq \theta$ such that $y_0(u) > 0$ for $u \in (\theta - \delta, \theta)$. Since $f \equiv 0$ on $(0, \theta)$, $y_0 = y_0(u)$ solves the equation

$$y_0'(u) = p'h(u)(y_0(u))^{\frac{1}{p}}, \quad u \in (\theta - \delta, \theta).$$

Separating variables, we obtain for $u \in (\theta - \delta, \theta)$

$$y_0^{\frac{1}{p'}}(\theta) - y_0^{\frac{1}{p'}}(u) = H(\theta) - H(u),$$

i.e.,

$$y_0^{\frac{1}{p'}}(u) - H(u) = y_0^{\frac{1}{p'}}(\theta) - H(\theta) > 0$$

by (5.4). It follows that $\delta = \theta$ and

$$y_0^{\frac{1}{p'}}(u) > 0 \quad \text{for all } u \in [0, \theta].$$

Therefore,

$$y_0(u) > 0 \quad \text{for all } u \in [0, 1]. \tag{5.5}$$

Set

$$c^* := \sup\{c > 0 : y_c(u) > 0 \text{ for all } u \in (0, 1)\}.$$

It follows from (5.2), (5.5) and the continuous dependence of the solution to (5.1) on the parameter c that the set $\{c > 0 : y_c(u) > 0 \text{ for all } u \in (0, 1)\}$ is nonempty and $c^* > 0$. If $c^* = +\infty$ then there exist $c_n \rightarrow +\infty$ and corresponding $y_{c_n} = y_{c_n}(u) > 0$, $u \in (0, 1)$, which satisfy

$$y_{c_n}'(u) = p'(c_n + h(u))(y_{c_n}(u))^{\frac{1}{p}}, \quad u \in (0, \theta).$$

Separating variables yields

$$(y_{c_n}(u))^{\frac{1}{p'}} = (y_{c_n}(\theta))^{\frac{1}{p'}} + c_n(u - \theta) + H(u) - H(\theta), \quad u \in (0, \theta), \tag{5.6}$$

and from (5.3) we get

$$y_{c_n}(\theta) < p' \int_0^1 f(\sigma) d\sigma < +\infty.$$

Therefore, the right-hand side in (5.6) tends to $-\infty$, a contradiction. Hence $0 < c^* < +\infty$.

Next we prove that $y_{c^*}(u) > 0$, $u \in (0, 1)$, $y_{c^*}(0) = 0$. Indeed, by the continuous dependence of (5.1) on the parameter c and the definition of c^* , the solution $y_{c^*} = y_{c^*}(u)$ must vanish somewhere in the interval $[0, \theta]$. Let $\eta \in [0, \theta]$ be the largest zero of y_{c^*} . It follows from the comparison argument that solutions of (5.1) decrease (not strictly) with c . This can be easily shown as in [5, Lemma 4.5 and Corollary 4.6] by replacing c with $c + h(u)$. If $\eta > 0$ then for $c < c^*$ we have $y_c(u) \geq 0$ on $(0, \eta)$ and hence from

$$y_c'(u) = p'(c + h(u))(y_c(u))^{\frac{1}{p}}, \quad u \in (0, \eta),$$

we again deduce

$$0 \leq (y_c(u))^{\frac{1}{p'}} = (y_c(\eta))^{\frac{1}{p'}} + c(u - \eta) + H(u) - H(\eta). \tag{5.7}$$

Since for $c \rightarrow c^*$ we have $y_c(\eta) \rightarrow y_{c^*}(\eta) = 0$ by continuous dependence on parameter, for any fixed $u \in (0, \eta)$ there exists $c < c^*$, $(c^* - c)$ sufficiently small, such that

$$(y_c(\eta))^{\frac{1}{p'}} + c(u - \eta) + H(u) - H(\eta) < 0$$

($h_m = 0$ implies that H is nondecreasing), which contradicts (5.7). Hence $\eta = 0$ and we have $y_{c^*}(u) > 0$ in $(0, 1)$, $y_{c^*}(0) = 0$. By the uniqueness for the initial value problem (5.1), this solution is also a unique solution of the boundary value problem (3.2), (3.4) with $c = c^*$.

Finally, we show that positive solutions of (5.1) do not vanish at 0 for values of $c \neq c^*$. Assume by contradiction that there exists $\hat{c} \neq c^*$ such that $y_{\hat{c}} = y_{\hat{c}}(u) > 0$ solves (5.1) in $(0, 1)$, $y_{\hat{c}}(0) = 0$. The definition of c^* yields $\hat{c} < c^*$. Separating variables in the equation in (5.1) on $(0, \theta)$, we obtain

$$y_{\hat{c}}^{\frac{1}{p'}}(\theta) = \hat{c}\theta + H(\theta)$$

and also

$$y_{c^*}^{\frac{1}{p'}}(\theta) = c^*\theta + H(\theta). \quad (5.8)$$

Hence $y_{\hat{c}}(\theta) < y_{c^*}(\theta)$. On the other hand, the comparison argument applied to (5.1) yields $y_{\hat{c}}(u) \geq y_{c^*}(u)$, $u \in [0, 1]$. This follows from [5, Corollary 4.6] with $c_1 = \hat{c} + h(u)$ and $c_2 = c^* + h(u)$. In particular, $y_{\hat{c}}(\theta) \geq y_{c^*}(\theta)$, a contradiction.

It follows from (5.8) together with (5.3) that

$$c^* = \frac{1}{\theta} \left(y_{c^*}^{\frac{1}{p'}}(\theta) - H(\theta) \right) < \frac{1}{\theta} \left[\left(p' \int_0^1 D^{p'-1}(u)g(u) du \right)^{\frac{1}{p'}} - H(\theta) \right],$$

i.e., (2.6) holds. This concludes the proof for $h_m = 0$.

If $h_m \neq 0$, we can consider a new convective velocity $\tilde{h}(u) := h(u) - h_m$, $u \in [0, 1]$. Then $\tilde{h}_m := \min_{u \in [0, 1]} \tilde{h}(u) = 0$ and $\tilde{H}(u) = \int_0^u \tilde{h}(s) ds = H(u) - h_m u$ is a nondecreasing function. Setting $\tilde{c} := c + h_m$, the equation in (2.1) becomes

$$(D(u)|u'|^{p-2}u')' + (\tilde{c} + \tilde{h}(u))u' + g(u) = 0$$

and we can apply the above reasoning to prove the existence of a unique positive value \tilde{c}^* assuming that

$$\tilde{H}(1) \leq \left(k(p) \int_0^1 D^{p'-1}(u)g(u) du \right)^{\frac{1}{p'}}.$$

Since $\tilde{H}(1) = H(1) - h_m$, we immediately see that condition (2.5) yields a unique value $c^* = \tilde{c}^* - h_m > -h_m$ corresponding to the problem with convective velocity h and the estimate (2.6) holds. \square

Proof of Theorem 2.7. For $h_m > 0$ we can carry out the proof exactly as in the case $h_m = 0$, see the proof of Theorem 2.6. In particular, statements concerning the initial value problem (5.1) remain valid and the positivity of h justifies the use of Lemma B.2. Therefore, if (2.7) holds we conclude that $c^* > 0 > -h_m$. \square

6 Asymptotic analysis of the wave profile

In this section, we discuss asymptotic behavior of the solution $u = u(\xi)$ to (2.1) as $\xi \rightarrow \pm\infty$. Our aim is to determine whether the solution attains 0 and/or 1 (or neither of them). To this end, we study the convergence of the integrals from (3.6), and hence the boundedness of the interval (ξ_1, ξ_2) . For technical reasons and for the sake of brevity, we assume power-type behavior of D and g near equilibria 0 and 1.

In what follows, we consider $H(u) > 0$, $u \in (0, 1]$, and profiles with $c^* > 0$. For notational brevity, we use the following notation: for $s_0 \in \mathbb{R}$ we write

$$\phi_1(s) \sim \phi_2(s) \text{ as } s \rightarrow s_0 \text{ if and only if } \lim_{s \rightarrow s_0} \frac{\phi_1(s)}{\phi_2(s)} \in (0, +\infty).$$

6.1 Asymptotics near 0

Let us assume that $D(u) \sim u^\alpha$ as $u \rightarrow 0+$ for some $\alpha \in \mathbb{R}$. Thanks to $f \equiv 0$ in $[0, \theta]$, we have

$$y_{c^*}^{\frac{1}{p}}(u) = c^* u + H(u), \quad u \in (0, \theta),$$

and due to the assumption $H \in C^1[0, 1]$, $H(u) > 0$ together with $c^* > 0$, we have $y_{c^*}^{\frac{1}{p}}(u) \sim u$ as $u \rightarrow 0+$. Let us recall that

$$\xi_2 = \int_0^\theta \frac{D^{p'-1}(s)}{y_{c^*}^{1/p}(s)} ds.$$

Since

$$\int_0^u \frac{D^{p'-1}(s)}{y_{c^*}^{1/p}(s)} ds \sim \int_0^u \frac{s^{\alpha(p'-1)}}{s^{p'-1}} ds = \int_0^u s^{\frac{\alpha-1}{p-1}} ds \quad \text{as } u \rightarrow 0+, \quad (6.1)$$

we conclude that the following two cases occur:

- (a) $\xi_2 = +\infty$ if and only if $p + \alpha \leq 2$;
- (b) $\xi_2 < +\infty$ if and only if $p + \alpha > 2$,

see Figure 6.1 for geometric interpretation.




Figure 6.1: Visualization of cases (a) and (b), leading to ξ_2 finite or infinite

Observe that for any $\alpha > 1$, the profile $u = u(\xi)$ is always right compactly supported, i.e., $u \equiv 0$ in $[\xi_2, +\infty)$, $\xi_2 \in \mathbb{R}$. If $\alpha = 0$ and $1 < p \leq 2$, the profile does not attain 0 for any finite ξ . This result is consistent with that from [10] for $p = 2$ and $D \in C^1[0, 1]$ strictly positive in $[0, 1]$.

In case (b), we can also study the one-sided derivative $u'(\xi_2-)$. In particular, differentiating (3.5) yields

$$\frac{d\xi}{du} = -\frac{d}{du} \int_{\frac{1}{2}}^u \frac{D^{p'-1}(s)}{y_{c^*}^{1/p}(s)} ds = -\frac{D^{p'-1}(u)}{y_{c^*}^{1/p}(u)}, \quad u \in (0, 1).$$

Since $D(u) \sim u^\alpha$, $y(u) \sim u^{p'}$ as $u \rightarrow 0+$, we have

$$\left. \frac{d\xi}{du} \right|_{u=0+} \sim -u^{\frac{\alpha-1}{p-1}} \rightarrow \begin{cases} 0 & \text{if } \alpha > 1 \\ \text{const.} < 0 & \text{if } \alpha = 1 \\ -\infty & \text{if } \alpha < 1 \end{cases} \quad \text{as } u \rightarrow 0+.$$

From an inverse perspective, we obtain the following classification for the profile $u = u(\xi)$:

$$u'(\xi_2-) = \begin{cases} -\infty & \text{if } \alpha > 1, \\ \text{const.} < 0 & \text{if } \alpha = 1, \\ 0 & \text{if } \alpha < 1. \end{cases}$$

Therefore, if $p + \alpha > 2$ and $\alpha < 1$ we have $u'(\xi_2-) = u'(\xi_2+) = 0$.

6.2 Asymptotics near 1

Let us assume that $D(u) \sim (1-u)^\beta$ and $g(u) \sim (1-u)^\gamma$ as $u \rightarrow 1-$ for some $\beta \in \mathbb{R}$, $\gamma > 0$. Since the equation (3.2) is not separable on $(\theta, 1)$, the asymptotic analysis becomes more involved than in the previous case. However, we can apply the same reasoning as in [4, Section 5.1], where we investigated asymptotic properties of solutions in the absence of convection. In fact, this technique yields the same results also when $h(u) \geq 0$ instead of $h \equiv 0$. Replacing c by $c + h(u)$ in [4, Section 5.1], we obtain the same classification of solutions as in Theorems 5.1 and 5.2 therein. In our current notation, these theorems read as follows.

Theorem 6.1. *Let $D(u) \sim (1-u)^\beta$, $g(u) \sim (1-u)^\gamma$ as $u \rightarrow 1-$ where $\beta \in \mathbb{R}$ and $\gamma > 0$ are such that*

$$-1 < \gamma + \frac{\beta}{p-1} \leq \frac{1}{p-1}$$

for given $p > 1$. If

$$\frac{\gamma - \beta + 1}{p} < 1,$$

then $\xi_1 > -\infty$. If

$$\frac{\gamma - \beta + 1}{p} \geq 1,$$

then $\xi_1 = -\infty$.

Theorem 6.2. *Let $D(u) \sim (1-u)^\beta$, $g(u) \sim (1-u)^\gamma$ as $u \rightarrow 1-$ where $\beta \in \mathbb{R}$ and $\gamma > 0$ are such that*

$$\gamma + \frac{\beta}{p-1} > \frac{1}{p-1}$$

for given $p > 1$. If $\gamma < 1$ then $\xi_1 > -\infty$. If $\gamma \geq 1$ then $\xi_1 = -\infty$.

Remark 6.3. To visualize conditions from Theorems 6.1, 6.2, we introduce the following sets:

$$\mathcal{M}_1^1 := \{(\gamma, \beta) \in \mathbb{R}^2 : \gamma > 0, -1 < \gamma + \frac{\beta}{p-1} \leq \frac{1}{p-1}, \gamma - \beta + 1 < p\},$$

$$\mathcal{M}_1^2 := \{(\gamma, \beta) \in \mathbb{R}^2 : \gamma > 0, -1 < \gamma + \frac{\beta}{p-1} \leq \frac{1}{p-1}, \gamma - \beta + 1 \geq p\},$$

$$\mathcal{M}_1^3 := \{(\gamma, \beta) \in \mathbb{R}^2 : \gamma > 0, \gamma + \frac{\beta}{p-1} > \frac{1}{p-1}, \gamma < 1\},$$

$$\mathcal{M}_1^4 := \{(\gamma, \beta) \in \mathbb{R}^2 : \gamma > 0, \gamma + \frac{\beta}{p-1} > \frac{1}{p-1}, \gamma \geq 1\}.$$

Then $\xi_1 > -\infty$ if and only if $(\gamma, \beta) \in \mathcal{M}_1^1 \cup \mathcal{M}_1^3$ and $\xi_1 = -\infty$ if and only if $(\gamma, \beta) \in \mathcal{M}_1^2 \cup \mathcal{M}_1^4$. See Figure 6.2 for the case $p = 2$.




Figure 6.2: Visualization of the sets \mathcal{M}_1^1 , \mathcal{M}_1^2 , \mathcal{M}_1^3 and \mathcal{M}_1^4 for $p = 2$

Moreover, one can show exactly as in [5, Remark 6.4] that if $\xi_1 \in \mathbb{R}$, i.e., $(\gamma, \beta) \in \mathcal{M}_1^1 \cup \mathcal{M}_1^3$, then $u'(\xi_1) = 0$.

Acknowledgement

Pavel Drábek was supported by the Grant Agency of the Czech Republic (GAČR) under Grant No. 22-18261S.

A Auxiliary lemmas for Proposition 3.1

Lemma A.1. *Let $u \in C^1(\mathbb{R})$ be a solution of the initial value problem*

$$\begin{cases} (D(u)|u|^{p-2}u)' = -(c + h(u))u', \\ u(\xi_0) = u_0 \in (0, 1), u'(\xi_0) = 0. \end{cases} \quad (\text{A.1})$$

Then u does not verify part (c) of Definition 2.1.

Proof. Integrating the equation in (A.1) and using the initial conditions yields

$$D(u(\xi))|u'(\xi)|^{p-2}u'(\xi) = c(u_0 - u(\xi)) + H(u_0) - H(u(\xi)), \quad \xi \in \mathbb{R}. \quad (\text{A.2})$$

Put

$$\mathcal{S}_p(\nu) := |\nu|^{p-2}\nu \text{ for } \nu \neq 0, \mathcal{S}_p(0) = 0, \quad p > 1.$$

Since $\mathcal{S}_{p'}$ is the inverse function to \mathcal{S}_p , equation (A.2) is for $D(u(\xi)) \neq 0$ equivalent to

$$u'(\xi) = \mathcal{S}_{p'} \left(\frac{1}{D(u(\xi))} [c(u_0 - u(\xi)) + H(u_0) - H(u(\xi))] \right). \quad (\text{A.3})$$

If $1 < p \leq 2$ then, due to $D \in C^1(0, 1)$, $H \in C^1[0, 1]$ and $p' > 2$, the right-hand side of (A.3) is locally Lipschitz continuous in u . Hence $u(\xi) = u_0$, $\xi \in \mathbb{R}$, is a unique solution of (A.1) in \mathbb{R} , and therefore does not verify part (c) of Definition 2.1.

If $p > 2$, i.e., $1 < p' < 2$, then the right-hand side of (A.3) is not Lipschitz continuous only at one point $u = u_0$, but it is one-sided Lipschitz continuous there due to the fact that $\mathcal{S}_{p'}(\nu) = |\nu|^{p'-2}\nu$ satisfies one-sided Lipschitz condition. Therefore, either $u(\xi) = u_0$, $\xi \in (-\infty, \xi_0]$ is a unique solution of (A.1) in $(-\infty, \xi_0]$, or $u(\xi) = u_0$, $\xi \in [\xi_0, +\infty)$, is a unique solution of (A.1) in $[\xi_0, +\infty)$. In either case, part (c) of Definition 2.1 is not satisfied. \square

Lemma A.2. *Let u be a solution of (2.1) and let $\xi_0 \in \mathbb{R}$ be such that $u(\xi_0) \in (0, 1)$, $u'(\xi_0) = 0$ and*

$$(D(u(\xi))|u'(\xi)|^{p-2}u'(\xi))' \Big|_{\xi=\xi_0} < 0.$$

Then u has a strict local maximum at ξ_0 .

Proof. Let us recall that it follows from Remark 2.3 that both functions $\xi \mapsto D(u(\xi))|u'(\xi)|^{p-2}u'(\xi)$ and $\xi \mapsto |u'(\xi)|^{p-2}u'(\xi)$ are continuously differentiable in a small neighborhood of ξ . We have

$$0 > (D(u(\xi))|u'(\xi)|^{p-2}u'(\xi))' \Big|_{\xi=\xi_0} = \frac{dD}{du} \Big|_{u=u(\xi_0)} \underbrace{|u'(\xi_0)|^p}_{=0} + D(u(\xi_0)) (|u'(\xi)|^{p-2}u'(\xi))' \Big|_{\xi=\xi_0}.$$

Since $D(u(\xi_0)) > 0$, we get $(|u'(\xi)|^{p-2}u'(\xi))' \Big|_{\xi=\xi_0} < 0$, and therefore, $|u'(\xi)|^{p-2}u'(\xi)$ is strictly decreasing in ξ_0 and equal to 0 at $\xi = \xi_0$. Since the power $\mathcal{S}_p(\nu) = |\nu|^{p-2}\nu$ is strictly increasing, $u'(\xi)$ is strictly decreasing at $\xi = \xi_0$. Hence ξ_0 is the point of strict local maximum of u . \square

B Technical lemmas for Section 5

Lemma B.1 (Technical inequalities). *Let $a > 0$, $b > 0$. Then*

(i) *for $r \geq 2$ we have*

$$a^r + ra^{r-1}b + b^r \leq (a+b)^r;$$

(ii) *for $1 < r < 2$ we have*

$$a^r + ra^{r-1}b + b^r \leq \hat{k}(r)(a+b)^r,$$

where

$$\hat{k}(r) = \frac{1 + r(r-1)^{\frac{1}{r-2}} + (r-1)^{\frac{r}{r-2}}}{\left(1 + (r-1)^{\frac{1}{r-2}}\right)^r}.$$

Proof. We put $t = \frac{b}{a} > 0$ and write the inequality in an equivalent form

$$f(t) := \frac{1 + rt + t^r}{(1+t)^r} \leq \hat{k}(r),$$

where we set $\hat{k}(r) = 1$ for $r \geq 2$. Then the optimal choice for $\hat{k}(r)$ would be $\hat{k}(r) = \max_{t \geq 0} f(t)$, if this maximum exists. Indeed, it does. Namely, f is a continuously differentiable function on $[0, +\infty)$ satisfying $f(0) = 1 = \lim_{t \rightarrow +\infty} f(t)$. An elementary calculation yields that $t_1 = (r-1)^{\frac{1}{r-2}}$ is the only stationary point of f in $(0, +\infty)$.

Part (i). It is clear that equality holds for $r = 2$. Let $r > 2$. Then $f(1) = \frac{2+r}{2^r} < 1$. Hence $t_1 = (r-1)^{\frac{1}{r-2}}$ is the point of global minimum of f , $0 < f(t_1) \leq f(1) < 1$ and therefore $\max_{t \geq 0} f(t) = f(0) = 1$.

Part (ii). Let $1 < r < 2$. Then $f(1) = \frac{2+r}{2^r} > 1$ and hence t_1 is the point of global maximum of f in $[0, +\infty)$ with

$$\hat{k}(r) = f(t_1) = \frac{1 + r(r-1)^{\frac{1}{r-2}} + (r-1)^{\frac{r}{r-2}}}{\left(1 + (r-1)^{\frac{1}{r-2}}\right)^r}.$$

□

Lemma B.2 (Inequality (5.4)). *Assume that $h(u) \geq 0$ in $[0, 1]$ and let $y_0 = y_0(u)$ be a solution of the initial value problem (5.1) with $c = 0$. If*

$$H^{p'}(1) \leq k(p) \int_0^1 f(u) \, du, \quad (\text{B.1})$$

where $k = k(p)$ is given by (2.4), then

$$y_0^{\frac{1}{p'}}(\theta) > H(\theta).$$

Proof. We proceed by contradiction, that is, we assume that

$$y_0^{\frac{1}{p'}}(\theta) \leq H(\theta).$$

Since $f > 0$ on $(\theta, 1)$, it follows from the equation in (5.1) that $y_0(u) > 0$ for all $u \in (\theta, 1)$. Set $z(u) := y_0^{\frac{1}{p'}}(u)$. Then $z(u) > 0$ in $(\theta, 1)$, $z(1) = 0$,

$$z(\theta) \leq H(\theta) \quad (\text{B.2})$$

and $z = z(u)$ satisfies the equation

$$[z^{p'}(u)]' = p'h(u)z^{p'-1}(u) - p'f(u), \quad u \in (\theta, 1), \quad (\text{B.3})$$

or, equivalently,

$$z'(u) = h(u) - \frac{f(u)}{z^{p'-1}(u)}, \quad u \in (\theta, 1). \quad (\text{B.4})$$

Integrating (B.3) and using the mean value theorem, we obtain

$$\begin{aligned} z^{p'}(\theta) &= z^{p'}(1) - p' \int_{\theta}^1 h(\sigma)z^{p'-1}(\sigma) \, d\sigma + p' \int_{\theta}^1 f(\sigma) \, d\sigma \\ &= -p'z^{p'-1}(\tau_0)(H(1) - H(\theta)) + p' \int_0^1 f(\sigma) \, d\sigma \end{aligned} \quad (\text{B.5})$$

for some $\tau_0 \in (\theta, 1)$. From (B.4) we have

$$z(\tau_0) - z(\theta) = \int_{\theta}^{\tau_0} h(\sigma) \, d\sigma - \int_{\theta}^{\tau_0} \frac{f(\sigma)}{z^{p'-1}(\sigma)} \, d\sigma < H(\tau_0) - H(\theta)$$

and hence

$$z(\tau_0) < z(\theta) + H(1) - H(\theta) \quad (\text{B.6})$$

thanks to the monotonicity of H (in particular, $h \geq 0$ implies that H is nondecreasing). It follows from (B.5), (B.6) together with (B.2)

$$H^{p'}(\theta) > -p'(H(\theta) + [H(1) - H(\theta)])^{p'-1}(H(1) - H(\theta)) + p' \int_0^1 f(\sigma) \, d\sigma. \quad (\text{B.7})$$

Next we proceed separately for $p = 2$, $1 < p < 2$ and $p > 2$.

Case 1: $p = 2$. Since $p' = 2$, (B.7) becomes

$$H^2(\theta) > -2(H(\theta) + [H(1) - H(\theta)])(H(1) - H(\theta)) + 2 \int_0^1 f(\sigma) d\sigma.$$

Reorganizing the terms in the above inequality and using (B.1), we obtain

$$\begin{aligned} 0 &> -H^2(\theta) - 2H(\theta)(H(1) - H(\theta)) - (H(1) - H(\theta))^2 - (H(1) - H(\theta))^2 + 2 \int_0^1 f(\sigma) d\sigma \\ &= -H^2(1) - (H(1) - H(\theta))^2 + 2 \int_0^1 f(\sigma) d\sigma > 2 \left(\int_0^1 f(\sigma) d\sigma - H^2(1) \right) \geq 0, \end{aligned}$$

a contradiction.

Case 2: $1 < p < 2$. Since $p' > 2$, we use the well-known inequality

$$(a + b)^r \leq 2^{r-1}(a^r + b^r), \quad a, b > 0, \quad r > 1,$$

with $a = H(\theta)$, $b = H(1) - H(\theta)$, $r = p' - 1$ in (B.7) and obtain

$$\begin{aligned} H^{p'}(\theta) &> -p'(H(\theta) + [H(1) - H(\theta)])^{p'-1}(H(1) - H(\theta)) + p' \int_0^1 f(\sigma) d\sigma \\ &\geq -p'2^{p'-2} \left(H^{p'-1}(\theta) + [H(1) - H(\theta)]^{p'-1} \right) (H(1) - H(\theta)) + p' \int_0^1 f(\sigma) d\sigma. \end{aligned}$$

Hence

$$\begin{aligned} 0 &> -H^{p'}(\theta) - p'2^{p'-2}H^{p'-1}(\theta)(H(1) - H(\theta)) - p'2^{p'-2}(H(1) - H(\theta))^{p'} + p' \int_0^1 f(\sigma) d\sigma \\ &= -H^{p'}(\theta) - p'H^{p'-1}(\theta)(H(1) - H(\theta)) - (H(1) - H(\theta))^{p'} \\ &\quad + (1 - p'2^{p'-2})(H(1) - H(\theta))^{p'} + (p' - p'2^{p'-2})H^{p'-1}(\theta)(H(1) - H(\theta)) + p' \int_0^1 f(\sigma) d\sigma \end{aligned}$$

and, using the inequality from Lemma B.1 (i) with $a = H(\theta)$, $b = H(1) - H(\theta)$ and $r = p'$,

$$\begin{aligned} 0 &> -(H(\theta) + (H(1) - H(\theta)))^{p'} + (1 - p'2^{p'-2})(H(1) - H(\theta))^{p'} \\ &\quad + (p' - p'2^{p'-2})H^{p'-1}(\theta)(H(1) - H(\theta)) + p' \int_0^1 f(\sigma) d\sigma. \end{aligned}$$

Then $0 \leq H(\theta) \leq H(1)$ implies

$$0 > -H^{p'}(1) + (1 - p'2^{p'-2})H^{p'}(1) + (p' - p'2^{p'-2})H^{p'}(1) + p' \int_0^1 f(\sigma) d\sigma$$

and from (B.1) we conclude

$$0 > -p'(2^{p'-1} - 1)H^{p'}(1) + p' \int_0^1 f(\sigma) d\sigma \geq 0,$$

a contradiction.

Case 3: $p > 2$. Since $1 < p' < 2$, we now use the well-known inequality

$$(a + b)^r \leq a^r + b^r, \quad a, b > 0, \quad 0 < r < 1,$$

with $a = H(\theta)$, $b = H(1) - H(\theta)$, $r = p' - 1$ in (B.7) and obtain

$$0 > -H^{p'}(\theta) - p' \left(H^{p'-1}(\theta) + [H(1) - H(\theta)]^{p'-1} \right) (H(1) - H(\theta)) + p' \int_0^1 f(\sigma) \, d\sigma,$$

i.e.,

$$0 > -H^{p'}(\theta) - p' H^{p'-1}(\theta)(H(1) - H(\theta)) - p' (H(1) - H(\theta))^{p'} + p' \int_0^1 f(\sigma) \, d\sigma,$$

or equivalently

$$0 > -H^{p'}(\theta) - p' H^{p'-1}(\theta)(H(1) - H(\theta)) - (H(1) - H(\theta))^{p'} \\ - (p' - 1) (H(1) - H(\theta))^{p'} + p' \int_0^1 f(\sigma) \, d\sigma.$$

For $a > 0$, $b > 0$, $r \in (1, 2)$ we have $a^r + ra^{r-1}b + b^r \leq \hat{k}(r)(a+b)^r$ by the technical Lemma B.1 (ii). We apply it with $a = H(\theta)$, $b = H(1) - H(\theta)$, $r = p'$:

$$0 > -\hat{k}(p') (H(\theta) + (H(1) - H(\theta)))^{p'} - (p' - 1) (H(1) - H(\theta))^{p'} + p' \int_0^1 f(\sigma) \, d\sigma.$$

But (B.1) yields

$$0 > -(\hat{k}(p') + (p' - 1))H^{p'}(1) + p' \int_0^1 f(\sigma) \, d\sigma \geq 0,$$

a contradiction. □

References

- [1] A. Audrito and J.L. Vázquez, *The Fisher-KPP problem with doubly nonlinear diffusion*, J. Differ. Equ. **263** (2017), no. 11, 7647–7708.
- [2] J. Benedikt, P. Girg, L. Kotrla and P. Takáč, *Origin of the p -Laplacian and A. Missbach*, Electron. J. Differ. Equ. **2018** (2018), no. 16, 1–17.
- [3] H. Berestycki, B. Nicolaenko and B. Scheurer, *Traveling wave solutions to combustion models and their singular limits*, SIAM J. Math. Anal. **33** (1985), no. 6, 1207–1242.
- [4] P. Drábek and M. Zahradníková. *Traveling waves for unbalanced bistable equations with density dependent diffusion*, Electron. J. Differ. Equ. **2021** (2021), no. 76, 1–21.
- [5] P. Drábek and M. Zahradníková. *Travelling waves for generalized Fisher-Kolmogorov equation with discontinuous density dependent diffusion*, Math. Meth. Appl. Sci. **46** (2023), no. 11, 12064–12086.
- [6] R. Enguiça, A. Gavioli and L. Sanchez, *A class of singular first order differential equations with applications in reaction-diffusion*, Discrete Contin. Dyn. Syst. **33** (2013), 173–191.
- [7] B.H. Gilding and R. Kersner, *Travelling Waves in Nonlinear Diffusion-Convection Reaction*. Birkhäuser Verlag, Basel, 2004.
- [8] A. Hamydy, *Travelling wave for absorption-convection-diffusion equations*, Electron. J. Differ. Equ. **2006** (2006), no. 86, 1–9.

- [9] J.R. King and P.M. McCabe, *On the Fisher–KPP Equation with Fast Nonlinear Diffusion*, Proceedings: Mathematical, Physical and Engineering Sciences **459** (2003), no. 2038, 2529–46.
- [10] L. Malaguti and C. Marcelli, *The influence of convective effects on front-propagation in certain diffusive models*, in Mathematical Modelling & computing in biology and medicine, 5th ESMTB Conference, Milan, Jul 2–6, 2002, ed. V. Capasso, Milan Res. Cent. Ind. Appl. Math. MIRIAM Proj., **1** Società Editrice Esculapio, Bologna, 2003. 362–367.
- [11] L. Malaguti, C. Marcelli and S. Matucci, *A unifying approach to travelling wavefronts for reaction-diffusion equations arising from genetics and combustion models*, Dynamic Systems and Applications **12** (2003), 333–354.
- [12] F. Sánchez-Garduño and P.K. Maini, *Existence and uniqueness of a sharp travelling wave in degenerate non-linear diffusion Fisher-KPP equation*, J. Math. Biol. **33** (1994), 163–192.