Study of semileptonic $B \rightarrow DP\ell^+\nu_\ell$ decays based on the SU(3) flavor symmetry

Ru-Min Wang^{1,†}, Yi-Jie Zhang¹, Meng-Yuan Wan¹, Xiao-Dong Cheng^{2,§}, Yuan-Guo Xu^{1, \sharp}

¹College of Physics and Communication Electronics, JiangXi Normal University, NanChang, JiangXi 330022, China

²College of Physics and Electronic Engineering, Xinyang Normal University, Xinyang, Henan 464000, China

[†]ruminwang@sina.com [§]chengxd@mails.ccnu.edu.cn [#]yuanguoxu@jxnu.edu.cn

Decays $B \to DP\ell^+\nu_\ell$ ($\ell = e, \mu, \tau$) with the non-resonance, the charmed vector resonances, the charmed scalar resonances and the charmed tensor resonances are calculated by using the SU(3) flavor symmetry. Firstly, the decay amplitudes of different modes are related by the SU(3) flavor symmetry. Then, relevant experiential data are used to constrain nonperturbative coefficients in the non-resonant and various resonant $B \to DP\ell^+\nu_\ell$ decays. Finally, using the constrained nonperturbative coefficients, the branching ratios of not-yet-measured $B \to D^*P\ell^+\nu_\ell$ decays with the non-resonant and various charmed resonant contributions are predicted. Many branching ratios are predicted for the first time. We find that $B \to D\eta'\ell^+\nu_\ell$, $B_s \to D_s\eta'\ell^+\nu_\ell$ decays only receive the non-resonant contributions, $B \to D_s K\ell^+\nu_\ell$, $B_s \to DK\ell^+\nu_\ell$, $B \to D\eta\ell^+\nu_\ell$ and $B_s \to D_s\eta\ell^+\nu_\ell$ decays receive both non-resonant and charmed tensor resonant contributions, $B^+ \to D^-\pi^+\ell^+\nu_\ell$ decays receive the non-resonant, the charmed scalar resonant and the charmed tensor resonant contributions, and other $B \to D\pi\ell^+\nu_\ell$ decays receive all four kinds of contributions. These results can be used to test the SU(3) flavor symmetry approach in the four-body semileptonic decays by the future LHCb and Belle-II experiments.

I. Introduction

Semileptonic *B* decays play a key role in testing the Standard Model and understanding the heavy quark dynamics. Some three-body semileptonic *B* decays, such as $B \to D\ell^+\nu_\ell$ and $B \to D^*\ell^+\nu_\ell$, have been well understood. Nevertheless, other decays, like $B \to D_0\ell^+\nu_\ell$, $B \to D_2^*\ell^+\nu_\ell$ and $B \to DP\ell^+\nu_\ell$ decays, received less attention. Some branching ratios of the $B \to DP\ell'^+\nu_{\ell'}$ ($\ell' = e, \mu$) decays have been measured, and the experimental data from the Particle Data Group (PDG) within 2σ errors are [1]

$$\mathcal{B}(B^+ \to D^- \pi^+ \ell'^+ \nu_{\ell'})_T = (4.4 \pm 0.8) \times 10^{-3},\tag{1}$$

$$\mathcal{B}(B^+ \to D^- \pi^+ \ell'^+ \nu_{\ell'})_{\overline{D}_0} = (2.5 \pm 1.0) \times 10^{-3}, \tag{2}$$

$$\mathcal{B}(B^+ \to D^- \pi^+ \ell'^+ \nu_{\ell'})_{\overline{D}_2^*} = (1.53 \pm 0.32) \times 10^{-3}, \tag{3}$$

$$\mathcal{B}(B^0 \to D^0 \pi^- \ell'^+ \nu_{\ell'})_T = (4.1 \pm 1.0) \times 10^{-3}, \tag{4}$$

$$\mathcal{B}(B^0 \to D^0 \pi^- \ell'^+ \nu_{\ell'})_{D_0} = (3.0 \pm 2.4) \times 10^{-3}, \tag{5}$$

$$\mathcal{B}(B^0 \to D^0 \pi^- \ell'^+ \nu_{\ell'})_{D_2^*} = (1.21 \pm 0.66) \times 10^{-3}, \tag{6}$$

$$\mathcal{B}(B^+ \to D_s^- K^+ \ell'^+ \nu_{\ell'})_T = (3.0^{+2.8}_{-2.4}) \times 10^{-4},\tag{7}$$

where $\mathcal{B}_{T,M}$ denote the total and M resonant branching ratios. For $\mathcal{B}(B^+ \to D^- \pi^+ \ell'^+ \nu_{\ell'})_T$ and $\mathcal{B}(B^0 \to D^0 \pi^- \ell'^+ \nu_{\ell'})_T$ given in Eq. (1) and Eq. (4), the PDG reports decays via the D_0 and D_2^* resonances. With the experiment running, more relevant decays could be measured at LHCb and Belle-II. Present measurements of the $B \to DP\ell'^+\nu_{\ell'}$ decays give us an opportunity to test theoretical approaches of the $B \to DP\ell^+\nu_{\ell}$ decays and to predict many non-measured decays, which can be further tested by LHCb and Belle-II in the near future.

Theoretically, semileptonic decays are relatively simple, since the weak and strong dynamics can be separated in these decays. All the strong dynamics in the initial and final hadrons is included in the hadronic transition form factors. The analytic structure of the $B \rightarrow DP$ form factors is more complicated than ones of the $B \rightarrow D$ or $B \rightarrow P$ form factors [2]. Due to our poor understanding of hadronic interactions, the $B \rightarrow DP$ evaluations of the form factors are difficult. In the absence of reliable calculations, the symmetry analysis can provide very useful information about the decays. SU(3) flavor symmetry is one of popular symmetry approaches. SU(3) flavor symmetry has been widely used to study hadron decays, including *b*-hadron decays [3–16], *c*-hadron decays [15–31], and light hadron decays [15, 32–37]. SU(3) flavor breaking effects due to the mass difference of u, d, s quarks have also been studied, for instance, in Refs. [38–50].

Some four-body semileptonic decays $B/D \to PP\ell^+\nu_\ell$ and $B \to D^{(*)}P\ell^+\nu_\ell$ have been studied, for instance, in Refs. [2, 51–64]. In this work, we will study the $B \to DP\ell^+\nu_\ell$ decays with the non-resonant, the charmed vector resonant, the charmed scalar resonant and the charmed tensor resonant contributions by the SU(3) flavor symmetry based on the experimental data. Firstly, we will construct the hadronic amplitude relations or the form factor relations between different decay modes. Then, we will use the available data to extract the hadronic amplitudes or the form factors. Finally, we will predict the not-yet-measured modes for further tests in experiments.

This paper is organized as follows. The non-resonant contributions of the $B \to DP\ell\nu_{\ell}$ decays are discussed in Sec. II. The charmed vector resonant, the charmed scalar resonant and the charmed tensor resonant contributions of $B \to DP\ell^+\nu_{\ell}$ decays are presented in Sec. III. Finally, summary is given in Sec. IV.

II. Non-resonant $B \to DP\ell^+\nu_\ell$ decays

A. Meson multiplets

Since the SU(3) flavor analysis is based on the SU(3) flavor group, we give relevant meson multiplets first. Bottom pseudoscalar triplet B_i , charm pseudoscalar triplet D_i , charm scalar triplet D_{0i} and charm vector triplet D_i^* under the SU(3) flavor symmetry of u, d, s quarks are

$$B_{i} = \left(B^{+}(\bar{b}u), B^{0}(\bar{b}d), B^{0}_{s}(\bar{b}s)\right), \qquad D_{i} = \left(\overline{D}^{0}(\bar{c}u), \ D^{-}(\bar{c}d), \ D^{-}_{s}(\bar{c}s)\right), \tag{8}$$

$$D_{0i} = \left(\overline{D}_0^0(\bar{c}u), D_0^-(\bar{c}d), D_{s0}^-(\bar{c}s)\right), \qquad D_i^* = \left(\overline{D}^{*0}(\bar{c}u), D^{*-}(\bar{c}d), D_s^{*-}(\bar{c}s)\right), \qquad (9)$$

where i = 1, 2, 3 for u, d, s quarks. Note that the structures of scalar D_0 mesons are not known well, and they might be four-quark states, $D_{0i}^{4q} = \left(\overline{D}_0^0(\bar{c}u\bar{d}d), D_0^-(\bar{c}d\bar{u}u), D_{s0}^-(\bar{c}s(\bar{u}u + \bar{d}d)/\sqrt{2})\right)$ [65].

The charm tensor triplet D_{2i}^* is

$$D_{2i}^* = \left(\overline{D}_2^*(2460)^0, \ D_2^*(2460)^-, \ D_{s2}^*(2573)^-\right).$$
(10)

Note that D_{0i} , D_i^* and D_{2i}^* will be used for the resonances in Sec. III, and they are given here together. In addition, charm axial vector mesons D'_1/D_1 have been studied in Ref. [66] for $B \to D^*P\ell^+\nu_\ell$ decays, D'_1/D_1 do not contribute to $B \to DP\ell^+\nu_\ell$ decays, so relevant multiplets are not listed here. Light pseudoscalar octets and singlets P_i^i is

$$P_{j}^{i} = \begin{pmatrix} \frac{\pi^{0}}{\sqrt{2}} + \frac{\eta_{8}}{\sqrt{6}} + \frac{\eta_{1}}{\sqrt{3}} & \pi^{+} & K^{+} \\ \pi^{-} & -\frac{\pi^{0}}{\sqrt{2}} + \frac{\eta_{8}}{\sqrt{6}} + \frac{\eta_{1}}{\sqrt{3}} & K^{0} \\ K^{-} & \overline{K}^{0} & -\frac{2\eta_{8}}{\sqrt{6}} + \frac{\eta_{1}}{\sqrt{3}} \end{pmatrix},$$
(11)

with j = 1, 2, 3 for u, d, s quarks. The η and η' are mixtures of $\eta_1 = \frac{u\bar{u} + d\bar{d} + s\bar{s}}{\sqrt{3}}$ and $\eta_8 = \frac{u\bar{u} + d\bar{d} - 2s\bar{s}}{\sqrt{6}}$ with the mixing angle θ_P

$$\begin{pmatrix} \eta \\ \eta' \end{pmatrix} = \begin{pmatrix} \cos\theta_P & -\sin\theta_P \\ \sin\theta_P & \cos\theta_P \end{pmatrix} \begin{pmatrix} \eta_8 \\ \eta_1 \end{pmatrix}.$$
 (12)

And $\theta_P = [-20^\circ, -10^\circ]$ from the PDG [1] will be used in our numerical analysis.

B. Decay amplitudes

The four-body semileptonic $B \to DP\ell^+\nu_\ell$ decays are generated by $\bar{b} \to \bar{c}\ell^+\nu_\ell$ transition, and the effective Hamiltonian is

$$\mathcal{H}_{eff}(\bar{b} \to \bar{c}\ell^+ \nu_\ell) = \frac{G_F}{\sqrt{2}} V_{cb} \bar{b} \gamma^\mu (1 - \gamma_5) \bar{c} \ \bar{\nu}_\ell \gamma_\mu (1 - \gamma_5) \ell, \tag{13}$$

where G_F is the Fermi constant, and V_{cb} is the CKM matrix element. Decay amplitudes of the $B \to DP\ell^+\nu_\ell$ decays can be written as

$$\mathcal{A}(B \to DP\ell^+\nu_\ell) = \langle D(k_1)P(k_2)\ell^+(q_1)\nu_\ell(q_2)|\mathcal{H}_{eff}(\bar{b} \to \bar{c}\ell^+\nu_\ell)|B(p_B)\rangle$$

$$= \frac{G_F}{\sqrt{2}}V_{cb}L_\mu H^\mu,$$
(14)

where $L_{\mu} = \bar{\nu_{\ell}} \gamma_{\mu} (1 - \gamma_5) \ell$ is the leptonic charged current, and $H^{\mu} = \langle D(k_1) P(k_2) | \bar{c} \gamma^{\mu} (1 - \gamma_5) \bar{b} | B(p_B) \rangle$ is the hadronic matrix element. Usually, the hadronic matrix element H^{μ} can be obtained in terms of the form factors $F_{\perp,t,0,\parallel}$ of the $B \to DP$ transitions, which are similar to ones of $B \to PP$ transitions [64]. Nevertheless, the calculations of the $F_0, F_t, F_{\perp}, F_{\parallel}$ form factors are very difficult. In this work, the hadronic matrix element will be related by the SU(3) flavor symmetry.

Since the leptonic charged current L_{μ} is invariant under the SU(3) flavor symmetry, the hadronic matrix element H^{μ} can be parameterized by the SU(3) flavor symmetry. Then the decay amplitudes of the non-resonant $B \to DP\ell^+\nu_{\ell}$ decays in Eq. (14) can be transformed as

$$\mathcal{A}(B \to DP\ell^+\nu_\ell)_N = \frac{G_F}{\sqrt{2}} V_{cb}^* H(B \to DP)_N \ \bar{\nu}_\ell \gamma_\mu (1 - \gamma_5)\ell, \tag{15}$$

$$H(B \to DP)_N = c_{01} B^i P^j_i D_j + c_{02} B^i D_i P^k_k,$$
(16)

where $c_{01,02}$ are the nonperturbative coefficients under the SU(3) flavor symmetry. The c_{02} term is suppressed by the Okubo-Zweig-Iizuka (OZI) rule [67–69], and it only appears in the decays with η, η' final states.

Decay modes	Non-resonant decay amplitudes	Decay modes	Non-resonant decay amplitudes
$B^+ \to D^- \pi^+ \ell^+ \nu_\ell$	c_{01}	$B^0 \to D^- \pi^0 \ell^+ \nu_\ell$	$-\frac{1}{\sqrt{2}}c_{01}$
$B^+ \to D_s^- K^+ \ell^+ \nu_\ell$	c_{01}	$B^0 \to D^- \eta \ell^+ \nu_\ell$	$\frac{c_{01}cos\theta_P}{\sqrt{6}} - \frac{(c_{01}+3c_{02})sin\theta_P}{\sqrt{3}}$
$B^+ \to \overline{D}{}^0 \pi^0 \ell^+ \nu_\ell$	$\frac{1}{\sqrt{2}}c_{01}$	$B^0 \to D^- \eta' \ell^+ \nu_\ell$	$\frac{c_{01}sin\theta_P}{\sqrt{6}} + \frac{(c_{01}+3c_{02})cos\theta_P}{\sqrt{3}}$
$B^+ \to \overline{D}{}^0 \eta \ell^+ \nu_\ell$	$\frac{c_{01}cos\theta_P}{\sqrt{6}} - \frac{(c_{01}+3c_{02})sin\theta_P}{\sqrt{3}}$	$B^0_s \to \overline{D}^0 K^- \ell^+ \nu_\ell$	c_{01}
$B^+ \to \overline{D}{}^0 \eta' \ell^+ \nu_\ell$	$\frac{c_{01}sin\theta_P}{\sqrt{6}} + \frac{(c_{01}+3c_{02})cos\theta_P}{\sqrt{3}}$	$B_s^0 \to D^- \overline{K}^0 \ell^+ \nu_\ell$	c_{01}
$B^0 \to \overline{D}{}^0 \pi^- \ell^+ \nu_\ell$	c_{01}	$B_s^0 \to D_s^- \eta \ell^+ \nu_\ell$	$-rac{2c_{01}cos heta_P}{\sqrt{6}} - rac{(c_{01}+3c_{02})sin heta_P}{\sqrt{3}}$
$B^0 \to D_s^- K^0 \ell^+ \nu_\ell$	c_{01}	$B_s^0 \to D_s^- \eta' \ell^+ \nu_\ell$	$-\frac{2c_{01}sin\theta_P}{\sqrt{6}}+\frac{(c_{01}+3c_{02})cos\theta_P}{\sqrt{3}}$

TABLE I: SU(3) IRA amplitudes for $B \to DP\ell^+\nu$ decays due to $\bar{b} \to \bar{c}\ell^+\nu$.

The idiographic hadronic amplitudes of the non-resonant $B \to DP\ell^+\nu_\ell$ decays are given in Tab. I. From Tab. I, one can see that, if ignoring the OZI suppressed c_{02} term, all hadronic amplitudes can be related by only one nonperturbative coefficient c_{01} .

The differential branching ratios of the nonresonant $B \to DP\ell^+\nu_\ell$ decays are [64]

$$\frac{d\mathcal{B}(B \to DP\ell^+\nu_\ell)_N}{dq^2 dk^2} = \frac{1}{2}\tau_B |\mathcal{N}|^2 \beta_\ell (3-\beta_\ell) |H_N|^2,\tag{17}$$

with

$$|\mathcal{N}|^{2} = G_{F}^{2} |V_{cb}|^{2} \frac{\beta_{\ell} q^{2} \sqrt{\lambda}}{3 \times 2^{10} \pi^{5} m_{B}^{3}}, \quad \text{with} \quad \beta_{\ell} = 1 - \frac{m_{\ell}^{2}}{q^{2}}.$$

$$|H_{N}|^{2} = |F_{0}|^{2} + \frac{2}{3} (|F_{\parallel}|^{2} + |F_{\perp}|^{2}) + \frac{3m_{\ell}^{2}}{q^{2}(3 - \beta_{\ell})} |F_{t}|^{2}, \qquad (18)$$

where $\tau_M(m_M)$ is lifetime (mass) of M particle. The ranges of integration are given by $(m_D + m_P)^2 \leq k^2 \leq (m_B - m_\ell)^2$ and $m_\ell^2 \leq q^2 \leq (m_B - \sqrt{k^2})^2$. If we ignore $|F_t|^2$ term since it is proportional to m_ℓ^2 and it is small when $\ell = e, \mu, |H_N|^2$ is only include the hadronic part. Noted that although $|F_t|^2$ term might be large when $\ell = \tau$, it is difficult to estimate its contribution in this work, so we still ignore it. Then H_N , which only includes hadronic part, follow the relationship of the SU(3) flavor symmetry in the Tab. I.

C. Numerical results

For the non-resonant $B \to DP\ell^+\nu_\ell$ decays, no any process has been measured until now. However, as given in Eq. (7), $\mathcal{B}(B^+ \to D_s^- K^+ \ell'^+ \nu_{\ell'})_T$ has been measured. The $B^+ \to D_s^- K^+ \ell'^+ \nu_{\ell'}$ mode can decay via the non-resonance and the D_2^* tensor meson resonance. In the later analysis of Sec. III, the contributions of D_2^* tensor meson resonance are far less than the experimental data given in Eq. (7). So we think that the non-resonant contributions are dominant in the $B^+ \to D_s^- K^+ \ell'^+ \nu_{\ell'}$ decays, *i.e.*, $\mathcal{B}(B^+ \to D_s^- K^+ \ell'^+ \nu_{\ell'})_N \approx \mathcal{B}(B^+ \to D_s^- K^+ \ell'^+ \nu_{\ell'})_T$. The experimental data of $\mathcal{B}(B^+ \to D_s^- K^+ \ell'^+ \nu_{\ell'})_T$ are used to determine c_{01} in the non-resonant $B \to DP\ell^+ \nu_{\ell'}$ decays (Due to poor relevant experimental data, the OZI suppressed c_{02} term is ignored). Then many other branching ratios of the non-resonant $B \to DP\ell^+ \nu_{\ell}$ decays can be predicted by using the data of $\mathcal{B}(B^+ \to D_s^- K^+ \ell'^+ \nu_{\ell'})_T$, which are listed in the second column of Table II.

From the second column of Table II, one can see that many branching ratio central values of the non-resonant $B \to DP\ell'^+\nu_{\ell'}$ decays, such as $\mathcal{B}(B^+ \to D^-\pi^+\ell'^+\nu_{\ell'})_N$, $\mathcal{B}(B^+ \to \overline{D}^0\pi^0\ell'^+\nu_{\ell'})_N$, $\mathcal{B}(B^+ \to \overline{D}^0\eta\ell'^+\nu_{\ell'})_N$, $\mathcal{B}(B^0 \to \overline{D}^0\pi^-\ell'^+\nu_{\ell'})_N$, $\mathcal{B}(B^0 \to D^-\pi^0\ell'^+\nu_{\ell'})_N$, $\mathcal{B}(B^0 \to D^-\eta\ell'^+\nu_{\ell'})_N$, $\mathcal{B}(B_s^0 \to \overline{D}^0K^-\ell'^+\nu_{\ell'})_N$ $\mathcal{B}(B_s^0 \to D^-\overline{K}^0\ell'^+\nu_{\ell'})_N$ and $\mathcal{B}(B_s^0 \to D_s^-\eta\ell'^+\nu_{\ell'})_N$, are on the orders of 10^{-4} , which could be measured by the LHCb and Belle II experiments. Nevertheless, other decays, for examples, the non-resonant $B^+ \to \overline{D}^0\eta'\ell'^+\nu_{\ell'}$, $B^0 \to D^-\eta'\ell'^+\nu_{\ell'}$, $B_s^0 \to D_s^-\eta'\ell'^+\nu_{\ell'}$, and all $B \to DP\tau^+\nu_{\tau}$ decays, are strongly suppressed by the narrow phase spaces, their branching ratio central values are on the orders of $\mathcal{O}(10^{-5} - 10^{-7})$, and they might not be observed by the experiments in the near future.

TABLE II: Branching ratios for $B \to DP\ell^+\nu_\ell$ decays due to $\bar{b} \to \bar{c}\ell^+\nu_\ell$ within 2σ errors. The unit is 10^{-3} for the branching ratios. \mathcal{B}_N denotes the resonant branching ratios, $\mathcal{B}_{[R]}$ denotes the *R* resonant branching ratios, ^e denotes experimental data within 2σ errors, and [†]denotes the results obtained by considering the resonance width effects.

Decay modes	\mathcal{B}_N	$\mathcal{B}_{[D^*]}$	$\mathcal{B}_{[D_0]}$	$\mathcal{B}_{[D_2^*]}$
$B^+ \to D^- \pi^+ \ell'^+ \nu_{\ell'}$	0.64 ± 0.52		$2.64 \pm 0.86 [D_0^0]$ 2.5+1.0 ^e	$1.33 \pm 0.12 [D_2^{*0}]$ 1.53 ± 0.32^{e}
$B^+ \to D_s^- K^+ \ell'^+ \nu_{\ell'}$	$0.32 {\pm} 0.26 \\ 0.30 {+} 0.28 e \\ -0.24$			$[5.66 \times 10^{-15}, 2.15 \times 10^{-7}]_{[D_2^{*0}]}$
$B^+ \to \overline{D}{}^0 \pi^0 \ell'^+ \nu_{\ell'}$	0.32 ± 0.26	$35.10\pm 2.68_{[D^{*0}]}$ $31.22\pm 2.25^{\dagger}_{[D^{*0}]}$	$1.34\pm0.44_{[D_0^0]}$	$0.70\pm 0.06_{[D_2^{*0}]}$
$B^+ \to \overline{D}^0 \eta \ell'^+ \nu_{\ell'}$	0.11 ± 0.10			$(4.36 \pm 1.22) \times 10^{-3} {}_{[D_2^{*0}]}$
$B^+ \to \overline{D}^0 \eta' \ell'^+ \nu_{\ell'}$	0.038 ± 0.033			
$B^0\to \overline{D}{}^0\pi^-\ell'^+\nu_{\ell'}$	0.60 ± 0.49	$33.67 \pm 2.11_{[D^{*}]}$ $30.14 \pm 1.72^{\dagger}_{[D^{*}]}$	2.48 ± 0.81 [D_0^-] 3.0 ± 2.4^e	$\frac{1.28 \pm 0.13}{[D_2^{*}]}$ 1.21 $\pm 0.66^{e}$
$B^0 \to D^s K^0 \ell'^+ \nu_{\ell'}$	0.30 ± 0.24			$[4.07 \times 10^{-14}, 5.30 \times 10^{-6}]_{[D_2^{*-}]}$
$B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'}$	0.30 ± 0.24	$\frac{15.29 \pm 1.21_{[D^*-]}}{13.69 \pm 1.05_{[D^*-]}^{\dagger}}$	$1.23\pm0.40_{[D_0^-]}$	$0.62 \pm 0.06_{[D_2^{*-}]}$
$B^0 \to D^- \eta \ell'^+ \nu_{\ell'}$	0.10 ± 0.09			$(3.54 \pm 1.40) \times 10^{-3}$
$B^0 \to D^- \eta' \ell'^+ \nu_{\ell'}$	0.035 ± 0.031			• • •
$B^0_s \to \overline{D}^0 K^- \ell'^+ \nu_{\ell'}$	0.41 ± 0.33			$1.23 \pm 0.18_{[D_{22}^{*-}]}$
$B^0_s \to D^- \overline{K}{}^0 \ell'^+ \nu_{\ell'}$	0.40 ± 0.33			$1.11 \pm 0.16_{[D_{-2}^{*-}]}$
$B^0_s \to D^s \eta \ell'^+ \nu_{\ell'}$	0.15 ± 0.13			$(1.67 \pm 0.59) \times 10^{-2} [D_{22}^{*-}]$
$B_s^0 \to D_s^- \eta' \ell'^+ \nu_{\ell'}$	0.095 ± 0.081			• • • • • • • • • • • • • • • • • • • •
$B^+ \to D^- \pi^+ \tau^+ \nu_\tau$	0.091 ± 0.074		$0.35 \pm 0.12_{[D_0^0]}$	$(8.62 \pm 1.82) \times 10^{-2} {}_{[D_2^{*0}]}$
$B^+ \to D_s^- K^+ \tau^+ \nu_\tau$	0.022 ± 0.017		•••	$[3.54 \times 10^{-16}, 1.45 \times 10^{-8}]_{[D_2^{*0}]}$
$B^+ \to \overline{D}{}^0 \pi^0 \tau^+ \nu_\tau$	0.047 ± 0.038	$8.59 \pm 0.66_{[D^{*0}]}$ $7.63 \pm 0.55_{[D^{*0}]}^{\dagger}$	$0.18\pm 0.06_{[D_0^0]}$	$(4.52 \pm 0.95) \times 10^{-2} {}_{[D_2^{*0}]}$
$B^+ \to \overline{D}^0 \eta \tau^+ \nu_{\tau}$	0.0085 ± 0.0072			$(2.81 \pm 1.00) \times 10^{-4} {}_{[D_2^{*0}]}$
$B^+ \to \overline{D}{}^0 \eta' \tau^+ \nu_{\tau}$	0.00086 ± 0.00074		• • •	
$B^0 \to \overline{D}^0 \pi^- \tau^+ \nu_\tau$	0.086 ± 0.070	$8.22 \pm 0.52 [D^{*-}]$ $7.34 \pm 0.42^{\dagger}_{[D^{*-}]}$	$0.32\pm0.11_{[D_0^-]}$	$(8.34 \pm 1.78) \times 10^{-2} {}_{[D_2^{*}]}$
$B^0 \to D_s^- K^0 \tau^+ \nu_\tau$	0.020 ± 0.016			$[2.51 \times 10^{-15}, 3.20 \times 10^{-7}]_{[D_2^{*}]}$
$B^0 \to D^- \pi^0 \tau^+ \nu_{\tau}$	0.043 ± 0.035	$3.73 \pm 0.30_{[D^*-]}$ $3.34 \pm 0.26^{\dagger}_{[D^*-]}$	$0.16\pm0.06_{[D_0^-]}$	$(4.06 \pm 0.87) \times 10^{-2} {}_{[D_2^{*}]}$
$B^0 \to D^- \eta \tau^+ \nu_{\tau}$	0.0077 ± 0.0066			$(2.34 \pm 1.03) \times 10^{-4} {}_{[D_2^{*-}]}$
$B^0 \to D^- \eta' \tau^+ \nu_\tau$	0.00077 ± 0.00067			
$B_s^0 \to \overline{D}^0 K^- \tau^+ \nu_\tau$	0.040 ± 0.033			$(7.86 \pm 1.99) \times 10^{-2} {}_{[D_{s2}^{*-}]}$
$B_s^0 \to D^- \overline{K}{}^0 \tau^+ \nu_\tau$	0.039 ± 0.032			$(7.10 \pm 1.79) \times 10^{-2} [D_{\circ 2}^{*-}]$
$B_s^0 \to D_s^- \eta \tau^+ \nu_\tau$	0.011 ± 0.009			$(1.04 \pm 0.44) \times 10^{-3} [D_{+2}^{*-}]$
$B_s^0 \to D_s^- \eta' \tau^+ \nu_\tau$	0.0020 ± 0.0017			

III. Decays $B \to DP\ell^+\nu_\ell$ with the D^*, D_0, D_2^* resonances

For the $B \to DP\ell^+\nu_\ell$ decays with the resonances, in the case of the decay widths of the resonance states are very narrow, the resonant branching ratios respect a simple factorization relation

$$\mathcal{B}(B \to D_J \ell^+ \nu_\ell, D_J \to DP) = \mathcal{B}(B \to D_J \ell^+ \nu_\ell) \times \mathcal{B}(D_J \to DP), \tag{19}$$

where $D_J = D^*, D_0, D_2^*$. And this result is also a good approximation for wider resonances. Eq. (19) will be used in our analysis for resonant $B \to D_J (\to DP) \ell^+ \nu_\ell$ decays. Relevant $\mathcal{B}(B \to D_J \ell^+ \nu_\ell)$ and $\mathcal{B}(D_J \to DP)$ also can be obtained by the SU(3) flavor symmetry.

A. Semileptonic three-body $B \to D_J \ell^+ \nu_\ell$ decays

The helicity amplitudes of the decays $B \to D_J \ell^+ \nu_\ell$ can be written as

$$\mathcal{M}(B \to D_J \ell^+ \nu_\ell) = \frac{G_F}{\sqrt{2}} V_{cb} \sum_{mm'} g_{mm'} L_m^{\lambda_\ell \lambda_\nu} H_{m'}^{\lambda_M}, \qquad (20)$$

with

$$L_m^{\lambda_\ell \lambda_\nu} = \epsilon_\alpha(m) \bar{\nu_\ell} \gamma^\alpha (1 - \gamma_5) \ell, \qquad (21)$$

$$H_{m'}^{\lambda_{D_J}} = \epsilon_{\beta}^*(m') \langle D_J(p, \epsilon^*) | \bar{c} \gamma^{\beta} (1 - \gamma_5) b | B(p_B) \rangle, \qquad (22)$$

where the particle helicities $\lambda_{D_J} = 0$ for D/D_0^* , $\lambda_{D_J} = 0, \pm 1$ for $D_J = D^*/D_2^*$, $\lambda_\ell = \pm \frac{1}{2}$, $\lambda_\nu = \pm \frac{1}{2}$, and $\epsilon_\mu(m)$ is the polarization vectors of the virtual W with $m = 0, t, \pm 1$.

The matrix elements can be parameterized via the form factors of the $B \rightarrow D_J$ transitions [70, 71]

$$\langle D(p) | \bar{c} \gamma_{\mu} b | B(p_B) \rangle = f_1^{BD}(q^2) \Big((p+p_B)_{\mu} - \frac{m_B^2 - m_D^2}{q^2} q_{\mu} \Big) + f_0^{BD}(q^2) \frac{m_B^2 - m_D^2}{q^2} q_{\mu}, \tag{23}$$

$$\langle D_{0}^{*}(p) | \bar{c}\gamma_{\mu}\gamma_{5}b | B(p_{B}) \rangle = -i \left[f_{1}^{BD_{0}}(q^{2}) \left((p+p_{B})_{\mu} - \frac{m_{B}^{2} - m_{D_{0}}^{2}}{q^{2}}q_{\mu} \right) + f_{0}^{BD_{0}}(q^{2}) \frac{m_{B}^{2} - m_{D_{0}}^{2}}{q^{2}}q_{\mu} \right], \quad (24)$$

$$\langle D^{*}(p,\varepsilon^{*}) | \bar{c}\gamma_{\mu}(1-\gamma_{5})b | B(p_{B}) \rangle = \frac{2V^{BD^{*}}(q^{2})}{m_{B} + m_{D^{*}}} \epsilon_{\mu\nu\alpha\beta} \varepsilon^{*\nu} p_{B}^{\alpha} p^{\beta}$$

$$-i \left[\varepsilon_{\mu}^{*}(m_{B} + m_{D^{*}})A_{1}^{BD^{*}}(q^{2}) - (p_{B} + p)_{\mu}(\varepsilon^{*}.p_{B}) \frac{A_{2}^{BD^{*}}(q^{2})}{m_{B} + m_{D^{*}}} \right]$$

$$+iq_{\mu}(\varepsilon^{*}.p_{B}) \frac{2m_{D^{*}}}{q^{2}} [A_{3}^{BD^{*}}(q^{2}) - A_{0}^{BD^{*}}(q^{2})], \quad (25)$$

$$\langle D_{2}(p,\varepsilon^{*}) | \bar{c}\gamma_{\mu}(1-\gamma_{5})b | B(p_{B}) \rangle = \frac{2iV^{BD_{2}}(q^{2})}{2} \epsilon_{\mu\nu\alpha\beta} \varepsilon^{*\nu} p_{D}^{\alpha} p^{\beta}$$

$$\langle D_2(p,\varepsilon^*) | \bar{c}\gamma_{\mu}(1-\gamma_5)b | B(p_B) \rangle = \frac{(1-\gamma)}{m_B + m_{D_2}} \epsilon_{\mu\nu\alpha\beta} e^{*\nu} p_B^{\alpha} p^{\beta} + 2m_{D_2} \frac{e^* \cdot q}{q^2} q_{\mu} A_0^{BD_2}(q^2) + (m_B + m_{D_2}) \Big(e^*_{\mu} - \frac{e^* \cdot q}{q^2} q_{\mu} \Big) A_1^{BD_2}(q^2) - \frac{e^* \cdot q}{m_B + m_{D_2}} \Big((p_B + p)_{\mu} - \frac{m_B^2 - m_{D_2}^2}{q^2} q_{\mu} \Big) A_2^{BD_2}(q^2),$$
(26)

where $s = q^2$ $(q = p_B - p_M)$, ε^* is the polarization of vector meson, and $e^{*\nu} \equiv \frac{\varepsilon^{*\mu\nu} \cdot p_{B\mu}}{m_B}$, $f_{+,0}^{BD}$, $f_{1,0}^{BD_0}$. The hadronic helicity amplitudes can be written as

$$H_{\pm}^{D} = 0, \qquad \qquad H_{0}^{D} = \frac{2m_{B}|\vec{p}_{D}|}{\sqrt{q^{2}}}f_{\pm}^{BD}(q^{2}), \qquad \qquad H_{t}^{D} = \frac{m_{B}^{2} - m_{D}^{2}}{\sqrt{q^{2}}}f_{0}^{BD}(q^{2}), \qquad (27)$$

for $B \to D\ell^+\nu_\ell$ decays,

$$H_{\pm}^{D_0} = 0, \qquad \qquad H_0^{D_0} = \frac{i2m_B |\vec{p}_{D_0}|}{\sqrt{q^2}} f_1^{BD_0}(q^2), \qquad \qquad H_t^{D_0} = \frac{i(m_B^2 - m_{D_0}^2)}{\sqrt{q^2}} f_0^{BD_0}(q^2), \qquad (28)$$

for $B \to D_0 \ell^+ \nu_\ell$ decays,

$$H_{\pm}^{D^*} = (m_B + m_{D^*}) A_1^{BD^*}(q^2) \mp \frac{2m_B |\vec{p}_{D^*}|}{(m_B + m_{D^*})} V^{BD^*}(q^2),$$
(29)

$$H_0^{D^*} = \frac{1}{2m_{D^*}\sqrt{q^2}} \left[(m_B^2 - m_{D^*}^2 - q^2)(m_B + m_{D^*})A_1^{BD^*}(q^2) - \frac{4m_B^2|\vec{p}_{D^*}|^2}{m_B + m_{D^*}}A_2^{BD^*}(q^2) \right],$$
(30)

$$H_t^{D^*} = \frac{2m_B |\vec{p}_{D^*}|}{\sqrt{q^2}} A_0^{BD^*}(q^2), \tag{31}$$

for $B \to D^* \ell^+ \nu_\ell$ decays, and

$$H_{\pm}^{D_{2}^{*}} = \frac{2|\vec{p}_{D_{2}^{*}}|}{2\sqrt{2}m_{D_{2}^{*}}} \left[(m_{B} + m_{D_{2}^{*}})A_{1}^{BD_{2}^{*}}(q^{2}) \mp \frac{2m_{B}|\vec{p}_{D_{2}^{*}}|}{(m_{B} + m_{D_{2}^{*}})}V^{BD_{2}^{*}}(q^{2}) \right],$$
(32)

$$H_0^{D_2^*} = \frac{|\vec{p}_{D_2^*}|}{\sqrt{2}m_{D_2^*}} \frac{1}{2m_{D_2^*}\sqrt{q^2}} \left[(m_B^2 - m_{D_2^*}^2 - q^2)(m_B + m_{D_2^*})A_1^{BD_2^*}(q^2) - \frac{4m_B^2|\vec{p}_{D_2^*}|^2}{m_B + m_{D_2^*}}A_2^{BD_2^*}(q^2) \right], \quad (33)$$

$$H_t^{D_2^*} = \frac{|\vec{p}_{D_2^*}|}{\sqrt{2}m_{D_2^*}} \frac{2m_B |\vec{p}_{D_2^*}|}{\sqrt{q^2}} A_0^{BD_2^*}(q^2), \tag{34}$$

for $B \to D_2^* \ell^+ \nu_\ell$ decays, where $|\vec{p}_{D_J}| \equiv \sqrt{\lambda(m_B^2, m_{D_J}^2, q^2)}/2m_B$ with $\lambda(a, b, c) = a^2 + b^2 + c^2 - 2ab - 2ac - 2bc$. The differential branching ratios of the $B \to D_J \ell^+ \nu_\ell$ decays are [72]

$$\frac{d\mathcal{B}(B \to D_J \ell^+ \nu_\ell)}{dq^2} = \frac{\tau_B G_F^2 |V_{cb}|^2 \lambda^{1/2} (q^2 - m_\ell^2)^2}{24(2\pi)^3 m_B^3 q^2} \mathcal{H}_{\text{total}},\tag{35}$$

with

$$\mathcal{H}_{\text{total}} = \left(\mathcal{H}_U + \mathcal{H}_L\right) \left(1 + \frac{m_\ell^2}{2q^2}\right) + \frac{3m_\ell^2}{2q^2} \mathcal{H}_S, \tag{36}$$

$$\mathcal{H}_U = |H_+^{D_J}|^2 + |H_-^{D_J}|^2, \quad \mathcal{H}_L = |H_0^{D_J}|^2, \quad \mathcal{H}_S = |H_t^{D_J}|^2, \quad (37)$$

where $\lambda \equiv \lambda(m_B^2, m_{D_J}^2, q^2)$ and $m_\ell^2 \le q^2 \le (m_B - m_{D_J})^2$.

Now one can obtain the branching ratios of the $B \to D_J \ell^+ \nu_\ell$ decays by the relevant form factors, which depend on the different methods. In this work, we use the SU(3) flavor symmetry to obtain the relations of the hadronic helicity amplitudes, and the same relations are also true for the form factors. In terms of the SU(3) flavor symmetry, the hadronic helicity amplitudes can be parameterized as

$$H_{m'}^{\lambda_{D_J}} = C_0^{D_J} B_i (D_J)^i, (38)$$

where $C_0^{D_J}$ are the nonperturbative coefficients of the $B \to D_J \ell^+ \nu_\ell$ decays under the SU(3) flavor symmetry. For the charmed four-quark states D_0^{4q} , $H_{m'}^{\lambda_{D_0}} = C_0^{4q,D_0} B_i (D_J)_j^{ij}$. And the hadronic helicity amplitude relations for the $B \to D_J \ell^+ \nu_\ell$ decays are summarized in Tab. III.

The relations in Tab. III will be used for the form factors F(0), which are $f_1^{BD}(0)$ in the $B \to D\ell^+\nu_\ell$ decays, $V^{BD^*}(0)$ in the $B \to D^*\ell^+\nu_\ell$ decays, $f_1^{BD_0}(0)$ in the $B \to D_0\ell^+\nu_\ell$ decays, and $A_1^{BD_2^*}(0)$ in $B \to D_2^*\ell^+\nu_\ell$ decays. The form factors F(0) are determined by the relevant experimental data. Other form factors $F_i(0)$ can be expressed as $r_i \times F(0)$, and the values of the ratios $r_i = \frac{F_i(0)}{F(0)}$ are taken from Ref. [73] for the $B \to D/D^*\ell^+\nu_\ell$ decays, from Ref.

Decay moeds	SU(3) hadronic amplitudes	Decay moeds	SU(3) hadronic amplitudes
$B^+ \to \overline{D}^0 \ell^+ \nu_\ell$	C_0^D	$B^+ \to \overline{D}^{*0} \ell^+ \nu_\ell$	$C_0^{D^*}$
$B^0 \to D^- \ell^+ \nu_\ell$	C_0^D	$B^0 \to D^{*-} \ell^+ \nu_\ell$	$C_0^{D^*}$
$B_s^0 \to D_s^- \ell^+ \nu_\ell$	C_0^D	$B_s^0 \to D_s^{*-} \ell^+ \nu_\ell$	$C_0^{D^*}$
$B^+ \to \overline{D}_0^0 \ell^+ \nu_\ell$	$C_0^{D_0}, C_0^{4q,D_0}$	$B^+ \to \overline{D}_2^{*0} \ell^+ \nu_\ell$	$C_{0}^{D_{2}^{*}}$
$B^0 \to D_0^- \ell^+ \nu_\ell$	$C_0^{D_0}, C_0^{4q,D_0}$	$B^0 \to D_2^{*-} \ell^+ \nu_\ell$	$C_{0}^{D_{2}^{*}}$
$B_s^0 \to D_{s0}^- \ell^+ \nu_\ell$	$C_0^{D_0}, \ \sqrt{2}C_0^{4q,D_0}$	$B_s^0 \to D_{s2}^{*-} \ell^+ \nu_\ell$	$C_0^{D_2^*}$

TABLE III: SU(3) IRA amplitudes for $B \to D_J \ell^+ \nu$ decays due to $\bar{b} \to \bar{c} \ell^+ \nu$.

TABLE IV: The experimental data and the SU(3) flavor symmetry predictions of the branching ratios of $B \to D/D^* \ell^+ \nu_{\ell}$ decays within 2σ errors. $\mathcal{B}(B \to D/D^* \ell^+ \nu_{\ell})$ is in unit of 10^{-2} . ^adenotes that the experimental data are not used to constrain the parameter $C_0^{D^*}$.

Branching ratios	Exp. data $[1]$	Our predictions	Branching ratios	Exp. data $[1]$	Our predictions
$\mathcal{B}(B^+ \to \overline{D}^0 \ell'^+ \nu_{\ell'})$	2.30 ± 0.18	2.34 ± 0.14	$\mathcal{B}(B^+ \to \overline{D}^{*0} \ell'^+ \nu_{\ell'})$	5.58 ± 0.44	5.41 ± 0.27
$\mathcal{B}(B^0\to D^-\ell'^+\nu_{\ell'})$	2.24 ± 0.18	2.19 ± 0.13	$\mathcal{B}(B^0\to D^{*-}\ell'^+\nu_{\ell'})$	4.97 ± 0.24	4.97 ± 0.24
$\mathcal{B}(B^0_s \to D^s \ell'^+ \nu_{\ell'})$		2.20 ± 0.14	$\mathcal{B}(B^0_s \to D^{*-}_s \ell'^+ \nu_{\ell'})$		4.99 ± 0.28
$\mathcal{B}(B^0_s \to D^s \mu^+ \nu_\mu)$	2.44 ± 0.46	2.20 ± 0.14	$\mathcal{B}(B_s^0 \to D_s^{*-} \mu^+ \nu_\mu)$	5.30 ± 1.0	4.98 ± 0.28
$\mathcal{B}(B^+ \to \overline{D}^0 \tau^+ \nu_\tau)$	0.77 ± 0.50	0.68 ± 0.04	$\mathcal{B}(B^+ \to \overline{D}^{*0} \tau^+ \nu_\tau)$	1.88 ± 0.40^a	1.35 ± 0.07
$\mathcal{B}(B^0\to D^-\tau^+\nu_\tau)$	1.05 ± 0.46	0.64 ± 0.04	$\mathcal{B}(B^0 \to D^{*-} \tau^+ \nu_\tau)$	1.58 ± 0.18^a	1.21 ± 0.06
$\mathcal{B}(B^0_s \to D^s \tau^+ \nu_\tau)$		0.63 ± 0.04	$\mathcal{B}(B^0_s \to D^{*-}_s \tau^+ \nu_\tau)$		1.20 ± 0.07

TABLE V: The experimental data and the SU(3) flavor symmetry predictions of the branching ratios of the $B \to D_0/D_2^* \ell^+ \nu_\ell$ decays within 2σ errors. ${}^{2q(4q)}$ denote the two(four) quark state predictions. $\mathcal{B}(B \to D_0/D_2^* \ell'^+ \nu_{\ell'})$ is in unit of 10^{-3} , and $\mathcal{B}(B \to D_0/D_2^* \tau^+ \nu_{\tau})$ is in unit of 10^{-4} .

Branching ratios	Exp. data $[74]$	Our predictions	Branching ratios	Exp. data	Our predictions
$\mathcal{B}(B^+ \to \overline{D}_0^0 \ell'^+ \nu_{\ell'})$	4.2 ± 1.6	3.98 ± 1.30	$\mathcal{B}(B^+ \to \overline{D}^0_2 \ell'^+ \nu_{\ell'})$	2.9 ± 0.6 [74]	3.20 ± 0.30
$\mathcal{B}(B^0\to D_0^-\ell'^+\nu_{\ell'})$	3.9 ± 1.4	3.71 ± 1.21	$\mathcal{B}(B^0\to D_2^-\ell'^+\nu_{\ell'})$	2.7 ± 0.6 [74]	2.99 ± 0.29
$\mathcal{B}(B^0_s\to D^{s0}\ell'^+\nu_{\ell'})$		$4.43 \pm 1.54^{2q}, \ 8.84 \pm 3.08^{4q}$	$\mathcal{B}(B^0_s \to D^{s2} \ell'^+ \nu_{\ell'})$		2.72 ± 0.27
$\mathcal{B}(B^+ \to \overline{D}_0^0 \tau^+ \nu_\tau)$		5.23 ± 1.85	$\mathcal{B}(B^+ \to \overline{D}_2^0 \tau^+ \nu_\tau)$		2.15 ± 0.47
$\mathcal{B}(B^0 \to D_0^- \tau^+ \nu_\tau)$		4.86 ± 1.72	$\mathcal{B}(B^0 \to D_2^- \tau^+ \nu_\tau)$		1.97 ± 0.43
$\mathcal{B}(B^0_s \to D^{s0}\tau^+\nu_\tau)$		$6.75 \pm 2.40^{2q}, \ 13.50 \pm 4.80^{4q}$	$\mathcal{B}(B_s^0 \to D_{s2}^- \tau^+ \nu_\tau)$	•••	1.73 ± 0.38

[75] for the $B \to D_0 \ell^+ \nu_\ell$ decays and from Ref. [76] for the $B \to D_2^* \ell^+ \nu_\ell$ decays. Taking the $B \to D^* \ell^+ \nu_\ell$ decays as an example, there are four form factors $V^{BD^*}(0)$ and $A_{0,1,2}^{BD^*}(0)$ in the $B \to D^* \ell^+ \nu_\ell$ decays, $A_{0,1,2}^{BD^*}(0)$ are expressed by $r_{0,1,2} \times V^{BD^*}(0)$, and the values of $r_{0,1,2} = \frac{A_{0,1,2}^{BD^*}(0)}{V^{BD^*}(0)}$ are taken from Ref. [73], and then there is only one parameter $V^{BD^*}(0)$ in the $B \to D^* \ell^+ \nu_\ell$ decays, and it can be determined by the experimental data of the $B \to D^* \ell^+ \nu_\ell$ decays.

Now we give our branching ratio predictions of the semileptonic $B \to D_J \ell^+ \nu_\ell$ decays under the SU(3) flavor symmetry. If not specially specified, the theoretical input parameters, such as the lifetimes, the masses, and the experimental data within the 2σ error bars from PDG [1] will be used in our numerical analysis.

Theoretically, exclusive semileptonic $B \to D/D^* \ell^+ \nu_\ell$ are well understood. Although the $B \to D\ell^+ \nu_\ell$ decays are not used for the four-body semileptonic decay branching ratios, there are five experimental data in the $B \to D\ell^+ \nu_\ell$ decays, which could be used to test the SU(3) flavor symmetry approach, so we present their results here. The experimental data of the $B \to D\ell^+\nu_\ell$ decays are listed in the second column of Tab. IV, which are used to constrain the only one free parameter $f_1^{BD}(0)$. We obtain that $f_1^{BD}(0) = 0.66 \pm 0.05$, which agrees with 0.67 given in Ref. [73]. Then one can predict the branching ratios of the $B_s^0 \to D_s^- \ell^+ \nu_\ell$ decays in terms of the constrained $f_1^{BD}(0)$, which are listed in the third column of Tab. IV.

For the $B \to D^* \ell^+ \nu_\ell$ decays, there are also five measured modes, and they are listed in the fifth column of Tab. IV. $\mathcal{B}(B^+ \to \overline{D}^{*0} \tau^+ \nu_\tau)$ and $\mathcal{B}(B^0 \to D^{*-} \tau^+ \nu_\tau)$ are not used to constrain the only free parameter $V^{BD^*}(0)$. We obtain that $V^{BD^*}(0) = 0.65 \pm 0.05$ from three experimental data of $\mathcal{B}(B_{(s)} \to D^* \ell'^+ \nu_{\ell'})$, which is smaller than 0.76 given in Ref. [73]. Then one can predict other branching ratios of the $B \to D^* \ell^+ \nu_\ell$ decays, which are listed in the last column of Tab. IV. One can see that our prediction and experimental data of $\mathcal{B}(B^+ \to \overline{D}^{*0} \tau^+ \nu_\tau)$ intersect within 2σ error ranges, nevertheless, our prediction of $\mathcal{B}(B^0 \to D^{*-} \tau^+ \nu_\tau)$ is slightly smaller than its data, and they will agree within 3σ error ranges.

For the $B \to D_0 \ell^+ \nu_\ell$ decays, only two decay modes have been measured, and they are listed in the second column of Tab. V, which are used to constrain the parameter $f_1^{BD_0}(0)$. Our constrained $f_1^{BD_0}(0) = 0.38 \pm 0.09$, which is obviously larger than 0.27 ± 0.03 given in Ref. [75]. Our branching ratio predictions of the semileptonic $B \to D_0 \ell^+ \nu_\ell$ decays are listed in the third column of Tab. V. The branching ratio predictions of $B_s^0 \to D_{s0}^- \ell^+ \nu_\ell$ are different between the two quark state and the four quark state, the prediction with four quark state are 2 times of one with the two quark state. In the later analysis of $B \to DP\ell\nu_\ell$ with D_0 resonances, the results of the two quark state will be used.

For the $B \to D_2^* \ell^+ \nu_\ell$ decays, only $\mathcal{B}(B^+ \to \overline{D}_2^0 \ell'^+ \nu_{\ell'})$ and $\mathcal{B}(B^0 \to D_2^- \ell'^+ \nu_{\ell'})$ have been measured, and they are listed in the fifth column of Tab. V. We obtain $A_1^{BD_2^*}(0) = 0.46 \pm 0.06$ from two measured branching ratios, which are consistent with $0.63^{+0.11}_{-0.12}$ within 2σ errors given in Ref. [76]. Then one can predict the branching ratios of the semileptonic $B \to D^* \ell^+ \nu_\ell$ decays, which are listed in the last column of Tab. IV. Decays $B_s^0 \to D_{s2}^- \ell^+ \nu_\ell$ have been calculated by the QCD sum rule approach for different scale parameter $\mu = 2/3/4$ GeV [77], for examples, $\mathcal{B}(B_s^0 \to D_{s2}^- \ell'^+ \nu_{\ell'}) = (3.07 \pm 1.40) \times 10^{-3}$ and $\mathcal{B}(B_s^0 \to D_{s2}^- \tau^+ \nu_{\tau}) = (1.03 \pm 0.61) \times 10^{-3}$ for $\mu = 4$ GeV. The predictions of $\mathcal{B}(B_s^0 \to D_{s2}^- \ell'^+ \nu_{\ell'})$ in Ref. [77] are consistent with ours, nevertheless, the predictions of $\mathcal{B}(B_s^0 \to D_{s2}^- \tau^+ \nu_{\tau})$ in Ref. [77] are smaller than ours.

Until now, the SU(3) flavor symmetry predictions of the three-body semileptonic $B \to D_J \ell^+ \nu_\ell$ decays are quite coincident with their experimental data within 2σ errors. The SU(3) flavor breaking effects mainly come from different masses of u, d, and s quarks. Since $m_{u,d}$ are much smaller than m_s , the SU(3) breaking effects due to a non-zero m_s dominate the SU(3) breaking effects [50]. If considering the SU(3) flavor breaking effects due to a non-zero m_s , the nonperturbative coefficients of the $B_s^0 \to D_s^-/D_s^{*-}/D_{s0}^-/D_{s2}^{*-}\ell^+\nu_\ell$ decays are different from those of the $B^+ \to \overline{D}^0/\overline{D}^{*0}/\overline{D}_0^0/\overline{D}_2^{*0}\ell^+\nu_\ell$ and $B^0 \to D^-/D^{*-}/D_0^-/D_2^{*-}\ell^+\nu_\ell$ decays. As given in Tab. IV, decays $B_s^0 \to D_s^-\mu^+\nu_\mu$ and $B_s^0 \to D_s^{*-}\mu^+\nu_\mu$ have been measured, so one can estimate the SU(3) flavor breaking effects due to a nonzero m_s in the $B \to D/D^*\ell^+\nu_\ell$ decays. Comparing our SU(3) flavor symmetry predictions and their experimental measurements of $B_s^0 \to D_s^-/D_s^{*-}\mu^+\nu_\mu$ decays within 2σ errors, one can find the SU(3) breaking contributions to $\mathcal{B}(B_s^0 \to D_s^-\ell^+\nu_\ell)$ and $\mathcal{B}(B_s^0 \to D_s^{*-}\ell^+\nu_\ell)$ should be less than roughly 23% and 20% of their experimental central values, respectively. After the $B_s^0 \to D_{s0}^-/D_{s2}^{*-}\ell^+\nu_\ell$ decays are measured, one can estimate the SU(3) flavor breaking effects in the $B \to D_0/D_2^*\ell^+\nu_\ell$ decays.

Non-leptonic two-body $D_J \rightarrow DP$ decays

Non-leptonic two-body $D_J \to DP$ decays via strong or electromagnetic interaction are presented in this section. Due to parity conservation, only $D_J = D_0/D^*/D_2^*$ resonances are considered for the $B \to DP\ell^+\nu_\ell$ decays [78]. In terms of the SU(3) flavor symmetry, the decay amplitudes of the strong or electromagnetic $D_J \to DP$ decays can be parameterized as

$$A(D_J \to DP) = a_{01}^{D_J} (D_J)_i P_j^i D^j + a_{02}^{D_J} (D_J)_i D^i P_j^j,$$
(39)

where $a_{01,02}^{D_J}$ are the nonperturbative coefficients, and all D_J are two-quark state. $a_{02}^{D_J}$ are OZI suppressed and it will be ignored in later numerical analysis. The decay amplitudes for each $D_2^* \to DP$ decay are summarized in Tab. VI. The decay amplitudes for $D^*/D_0 \to DP$ can be obtained by replacing $a_{01,02}^{D_2^*}$ listed in Tab. VI with $a_{01,02}^{D^*,D_0}$ only if their phase spaces are allowed.

Then the branching ratios of the $D_J \to DP$ decays can be written as [79]

В.

$$\mathcal{B}(D_0 \to DP) = \frac{\tau_{D_0} p_c(m_{D_0}, m_D, m_P)}{8\pi m_{D_0}^2} |A(D_0 \to DP)|^2, \tag{40}$$

$$\mathcal{B}(D^* \to DP) = \frac{\tau_{D^*} p_c^3(m_{D^*}, m_D, m_P)}{6\pi m_{D^*}^2} |A(D^* \to DP)|^2, \tag{41}$$

$$\mathcal{B}(D_2^* \to DP) = \frac{\tau_{D_2^*} p_c^5(m_{D_2^*}, m_D, m_P)}{60\pi m_{D_2^*}^2} |A(D_2^* \to DP)|^2, \tag{42}$$

where the center of mass momentum $p_c(m_{D_J}, m_D, m_P) \equiv \frac{\sqrt{\lambda(m_{D_J}^2, m_D^2, m_P^2)}}{2m_{D_J}}$.

Four decay modes of the $D^* \to D\pi$ decays have been measured, and the data within 2σ errors are [1]

$$\mathcal{B}(D^{*0} \to D^0 \pi^0) = (64.7 \pm 1.8)\%, \qquad \mathcal{B}(D^{*+} \to D^0 \pi^+) = (67.7 \pm 1.0)\%, \\ \mathcal{B}(D^{*+} \to D^+ \pi^0) = (30.7 \pm 1.0)\%, \qquad \mathcal{B}(D_s^{*+} \to D_s^+ \pi^0) = (5.8 \pm 1.4)\%.$$
(43)

Isospin violating decays $D_s^{*+} \to D_s^+ \pi^0$ and later $D_{s0}^0 \to D_s^0 \pi^0$, which might decay via $\eta - \pi^0$ mixing [80], are not considered in this work. The experimental data of $\mathcal{B}(D^{*0} \to D^0 \pi^0)$ and $\mathcal{B}(D^{*+} \to D^0 \pi^+, D^+ \pi^0)$ will be used to obtain $\mathcal{B}(B \to D \pi \ell \nu_\ell)$ with the D^* resonances.

In the $D_0 \to DP$ decays, only $D_0 \to D\pi$ decays have been seen but have no data. Since $D_0 \to D\eta$, $D_0 \to D_s K$ and $D_{s0} \to DK$ are not allowed by the phase spaces, we assume $\mathcal{B}(D_0 \to D\pi) = 1$ to obtain four branching ratios of the $D_0^0 \to D^0 \pi^0$, $D^+ \pi^-$ and $D_0^+ \to D^+ \pi^0$, $D^0 \pi^+$ decays. And they are

$$\mathcal{B}(D_0^0 \to D^0 \pi^0) = (33.62 \pm 0.03)\%, \qquad \mathcal{B}(D_0^0 \to D^+ \pi^-) = (66.38 \pm 0.03)\%, \\ \mathcal{B}(D_0^+ \to D^+ \pi^0) = (33.18 \pm 0.01)\%, \qquad \mathcal{B}(D_0^+ \to D^0 \pi^+) = (66.82 \pm 0.01)\%.$$
(44)

The branching ratios of $D_0 \to D\pi$ decays are the same as those in Eq. (44) when considering D_0 as the four quark state.

For the $D_2^* \to DP$ decays, since there is no experimental data of the branching ratios, we can not constrain $a_{01}^{D_2^*}$ directly. Nevertheless, $\frac{\mathcal{B}(D_2^* \to D\pi)}{\mathcal{B}(D_2^* \to D^*\pi)} = 1.52 \pm 0.14$ within 2σ errors are measured [1]. The SU(3) relation of the decay amplitudes of the $D_2^* \to D^*P$ decays are given in Ref. [66]. Using $\frac{\mathcal{B}(D_2^* \to D\pi)}{\mathcal{B}(D_2^* \to D^*\pi)} = 1.52 \pm 0.14$ and assuming $\mathcal{B}(D_2^{*0} \to D\pi) + \mathcal{B}(D_2^{*0} \to D^*\pi) \leq 1$, $\mathcal{B}(D_2^{*+} \to D\pi) + \mathcal{B}(D_2^{*+} \to D^*\pi) \leq 1$ and $\mathcal{B}(D_{s2}^{*+} \to DK) + \mathcal{B}(D_{s2}^{*+} \to D^*K) \leq 1$,

Decay modes	Coupling vertexs	Decay modes	Coupling vertexs
$D_2^{*0}(2460) \to D^0 \pi^0$	$a_{01}^{D_2^*}/\sqrt{2}$	$D_2^{*+}(2460) \to D^+ \pi^0$	$-a_{01}^{D_2^*}/\sqrt{2}$
$D_2^{*0}(2460) \to D^0 \eta$	$a_{01}^{D_2^*} \left(\frac{\cos \theta_P}{\sqrt{6}} - \frac{\sin \theta_P}{\sqrt{3}} \right) - \sqrt{3} a_{02}^{D_2^*} \sin \theta_P$	$D_2^{*+}(2460) \to D^+\eta$	$a_{01}^{D_2^*}\left(\frac{\cos\theta_P}{\sqrt{6}}-\frac{\sin\theta_P}{\sqrt{3}}\right)-\sqrt{3}a_{02}^{D_2^*}\sin\theta_P$
$D_2^{*0}(2460) \to D^+\pi^-$	$a_{01}^{D_2^*}$	$D_2^{*+}(2460) \to D^0 \pi^+$	$a_{01}^{D_2^*}$
$D_2^{*0}(2460) \to D_s^+ K^-$	$a_{01}^{D_2^*}$	$D_2^{*+}(2460) \to D_s^+ \overline{K}^0$	$a_{01}^{D_2^*}$
$D_{s2}^{*+}(2573) \to D^0 K^+$	$a_{01}^{D_2^*}$		
$D_{s2}^{*+}(2573) \to D^+ K^0$	$a_{01}^{D_2^*}$		
$D_{s2}^{*+}(2573) \to D_s^+ \eta$	$-a_{01}^{D_2^*}\left(\frac{2cos\theta_P}{\sqrt{6}} + \frac{sin\theta_P}{\sqrt{3}}\right) - \sqrt{3}a_{02}^{D_2^*}sin\theta_P$		

TABLE VI: The decay amplitudes for the $D_2^* \to DP$ decays under the SU(3) flavor symmetry.

TABLE VII: The predictions of the $D_2^* \to DP$ decays within 2σ errors.

Decay modes	Branching ratios $(\times 10^{-2})$	Decay widthes (MeV)	Decay widthes from $[81]$ (MeV)
$D_2^{*0}(2460) \to D^0 \pi^0$	20.18 ± 1.77	9.59 ± 1.06	$4.14^{+1.82}_{-1.57}$
$D_2^{*0}(2460) \to D^0 \eta$	0.13 ± 0.03	0.06 ± 0.02	
$D_2^{*0}(2460) \to D^+ \pi^-$	38.51 ± 3.41	18.30 ± 2.03	$7.91^{+3.49}_{-3.00}$
$D_2^{*0}(2460) \to D_s^+ K^-$	$[1.82\times 10^{-15}, 6.56\times 10^{-6}]$	$[8.77\times10^{-14}, 3.00\times10^{-6}]$	
$D_2^{*+}(2460) \to D^+ \pi^0$	19.40 ± 1.63	9.33 ± 0.92	
$D_2^{*+}(2460) \to D^+\eta$	0.11 ± 0.04	$(5.45 \pm 2.15) \times 10^{-2}$	
$D_2^{*+}(2460) \to D^0 \pi^+$	39.88 ± 3.33	19.18 ± 1.90	
$D_2^{*+}(2460) \rightarrow D_s^+ \overline{K}^0$	$[1.26\times 10^{-14}, 1.80\times 10^{-6}]$	$[6.01\times 10^{-13}, 8.28\times 10^{-5}]$	
$D_{s2}^{*+}(2573) \to D^0 K^+$	42.32 ± 5.20	7.52 ± 0.95	$3.35^{+1.48}_{-1.27}$
$D_{s2}^{*+}(2573) \to D^+ K^0$	38.12 ± 4.72	6.79 ± 0.86	$3.04^{+1.34}_{-1.15}$
$D_{s2}^{*+}(2573) \to D_s^+ \eta$	0.58 ± 0.19	$(10.23 \pm 3.43) \times 10^{-2}$	

one can constrain the nonperturbative coefficients $a_{01}^{D_2^*}$, and we obtain that $|a_{01}^{D_2^*}| = 25.14 \pm 1.47$. Then the branching ratios of the $D_2^* \to DP$ decays can be predicted, which are summarized in the second column of Tab. VII. In addition, their decay width predictions and previous width predictions are also given in the forth and fifth columns of Tab. VII, respectively. Our width predictions are about 1 time larger than ones in Ref. [81].

C. Numerical results of the resonant $B \to DP\ell^+\nu_\ell$ decays

In terms of $\mathcal{B}(B \to D_J \ell^+ \nu_\ell)$ given in Tabs. V-IV and $\mathcal{B}(D_J \to DP)$ given in Eqs. (43-44) and Tab. VII, after considering the further experimental bounds of the resonant $B \to D\pi \ell^+ \nu_\ell$ decays given in Eqs. (2-3) and (5-6), one can obtain the branching ratio predictions of the resonant $B \to DP\ell^+ \nu_\ell$ decays, and they are listed in the third, forth and fifth columns of Tab. II for the D^* , D_0 and $D^*_{(s)2}$ resonances, respectively. Corresponding experimental data with 2σ errors are also listed in Tab. II for the convenience of comparison. Note that, since the vector resonances are also considered in this work, $\mathcal{B}(B^+ \to D^- \pi^+ \ell'^+ \nu_{\ell'})_T$ and $\mathcal{B}(B^0 \to D^0 \pi^- \ell'^+ \nu_{\ell'})_T$ in Eq. (1) and Eq. (4) are not used for our results. Many resonant branching ratios are predicted for the first time.

One can see that the the vector meson D^* resonances give the dominant contributions in the $B^+ \to \overline{D}^0 \pi^0 \ell^+ \nu_\ell$, $B^0 \to \overline{D}^0 \pi^- \ell^+ \nu_\ell$ and $B^0 \to D^- \pi^0 \ell^+ \nu_\ell$ decays, largely because of its proximity to the $D\pi$ threshold. Please note that decay amplitude of the $B^+ \to D^- \pi^+ \ell^+ \nu_\ell$ decays is larger than ones of the $B^+ \to D^0 \pi^0 \ell^+ \nu_\ell$ decays by factor $\sqrt{2}$, nevertheless, the latter branching ratios are much larger than the former ones, since the most dominant resonance D^{*0} cannot decay into $D^- \pi^+$ on its mass-shell [1]. In previous studies, $\mathcal{B}(B^+ \to \overline{D}^0 \pi^0 \ell'^+ \nu_{\ell'})_{D^{*0}} = 34.9 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7 \times 10^{-3}$ [51], $\mathcal{B}(B^0 \to D^- \pi^0 \ell'^+ \nu_{\ell'})_{D^{*-}} = 16.7$

As for the scalar meson D_0 resonances and the tensor meson D_2^* resonances, the experimental upper limit of $\mathcal{B}(B^+ \to D^- \pi^+ \ell'^+ \nu_{\ell'})_{D_0^0}$ gives further constraint on the $\mathcal{B}(B \to D\pi\ell'^+ \nu_{\ell'})_{D_0}$ predictions, and the experimental lower limit of $\mathcal{B}(B^+ \to D^- \pi^+ \ell'^+ \nu_{\ell'})_{D_2^{*0}}$ gives further constraint on the $\mathcal{B}(B \to D\pi\ell'^+ \nu_{\ell'})_{D_2^*}$ predictions. Our predictions for $\mathcal{B}(B^0 \to \overline{D}^0 \pi^- \ell'^+ \nu_{\ell'})_{D_0^-, D_2^{*-}}$ are more precise than their experimental measurements. The contributions of the D_0 and D_2^* resonances are in the same order of magnitude in the $B^+ \to D^- \pi^+ \ell'^+ \nu_{\ell'}$ and $B^0 \to \overline{D}^0 \pi^- \ell'^+ \nu_{\ell'}$ decays. But the contributions of the D_0 resonances are larger than ones of the D_2^* resonances in the $B^+ \to \overline{D}^0 \pi^0 \ell^+ \nu_{\ell}$, $B^0 \to D^- \pi^0 \ell^+ \nu_{\ell}$, $B^+ \to D^- \pi^+ \tau^+ \nu_{\tau}$ and $B^0 \to \overline{D}^0 \pi^- \tau^+ \nu_{\tau}$ decays.

The Belle II experiment has reported the branching ratios of the $B \to D\eta \ell'^+ \nu_{\ell'}$ decays with quite large errors, $\mathcal{B}(B^0 \to D^- \eta \ell'^+ \nu_{\ell'})_T = (4.0 \pm 4.0) \times 10^{-3}$ and $\mathcal{B}(B^+ \to D^0 \eta \ell'^+ \nu_{\ell'})_T = (4.0 \pm 4.0) \times 10^{-3}$ [74], which are not used for our predictions. From our predictions, one can see that the non-resonant branching ratios are dominant in the $B \to D\eta \ell'^+ \nu_{\ell'}$ decays. And our predictions of $\mathcal{B}(B \to D\eta \ell'^+ \nu_{\ell'})_N$ lie in the range of experimental data with 1σ error.

In addition, the interference terms between the non-resonant, the vector resonant, the scalar resonant and the tensor resonant contributions exist, and they might not be ignored if more than one kind of contributions are important in the decays, and they will be studied in our succeeding work.

All current experimental data of $\mathcal{B}(B \to DP\ell^+\nu_\ell)$ can be explained by the SU(3) flavor symmetry approach. For the SU(3) flavor breaking effects in the $B \to DP\ell^+\nu_\ell$ decays, which are in the similar situation to the $B \to D^*P\ell^+\nu_\ell$ decays. The SU(3) flavor breaking effects due to a nonzero *s* quark mass dominate the SU(3) breaking effects. As given in Tab. II, the dominate SU(3) flavor breaking effects might appear in the non-resonant and the charmed tensor resonant $B \to DP\ell^+\nu_\ell$ decays. Nevertheless, there is only the data for $\mathcal{B}(B_{u,d} \to D\pi\ell^+\nu_\ell)_{D_2^*}$, and there is not any data for $\mathcal{B}(B_s \to DK\ell^+\nu_\ell)_{D_{s_2}^*}$. Or there is only the data of $\mathcal{B}(B^+ \to D_s^-K^+\ell^+\nu_\ell)_N$, and there is no any data of $\mathcal{B}(B_{u,d,s} \to DK\ell^+\nu_\ell)_N$. Therefore, we can not directly judge how large the possible SU(3) breaking effects are in the $B \to DP\ell^+\nu_\ell$ decays.

Although the widths of all resonances are narrow, following Refs. [51, 53], the width effects of D^* mesons are analyzed. After considering the width effects of D^* mesons, the decay branching ratios of $B \to D\pi \ell^+ \nu_{\ell}$ are [51, 52]

$$\mathcal{B}(B \to D^* \pi \ell^+ \nu_\ell)_{D^*} = \frac{1}{\pi} \int_{(m_{D^*} - n\Gamma_{D^*})^2}^{(m_{D^*} + n\Gamma_{D^*})^2} dt_V \int_{m_\ell^2}^{(m_B - \sqrt{t_V})^2} dq^2 \frac{\sqrt{t_V} d\mathcal{B}(B \to D^*(\lambda)\ell^+ \nu_\ell, t_V)/dq^2 \ \mathcal{B}(D^*(\lambda) \to D\pi, t_V)\Gamma_{D^*}}{(t_V - m_{D^*}^2)^2 + m_{D^*}^2 \Gamma_{D^*}^2},$$
(45)

where $d\mathcal{B}(B \to D^*(\lambda)\ell^+\nu_\ell, t_V)/dq^2$ and $\mathcal{B}(D^*(\lambda) \to D\pi, t_V)$ are obtained from to Eq. (35) and Eq. (41) by replacing $m_{D^*} \to \sqrt{t_V}$, respectively. There are two nonperturbative coefficients $V^{BD^*}(0)$ in $d\mathcal{B}(B \to D^*(\lambda)\ell^+\nu_\ell, t_V)/dq^2$ and $a_{01}^{D^*}$ in $\mathcal{B}(D^*(\lambda) \to D\pi, t_V)$. $V^{BD^*}(0) = 0.65 \pm 0.05$ from the data of $\mathcal{B}(B \to D^*\ell^+\nu_\ell)$ listed in Tab. IV, and $a_{01}^{D^*} = 8.42 \pm 0.38$ from the data of $\mathcal{B}(D^{*+} \to D\pi)$ given in Eq. (43). $\Gamma_{D^{*0}} = (56.00 \pm 5.74) \ KeV$, which is obtained

via the experimental data of $\mathcal{B}(D^{*0} \to D^0 \pi^0)$ in Eq. (43) and the SU(3) flavor symmetry in $D^* \to D\pi$. Following Refs. [51], choosing n = 3, the results of $\mathcal{B}(B \to D\pi \ell^+ \nu_\ell)_{D^*}$ are obtained, and they are listed in Tab. by denoting \dagger . One can see that the results obtained by considering the D^* width effects are slightly smaller than ones obtained by the narrow width approximation.

IV. Summary

The semileptonic $B \to DP\ell^+\nu_\ell$ decays with the non-resonances, the vector resonances, the scalar resonances and the tensor resonances have been investigated by using the SU(3) flavor symmetry based on the relevant experimental data. The amplitude relations of the non-resonant $B \to DP\ell^+\nu_\ell$ decays, the semileptonic $B \to D/D^*/D_0/D_2^*\ell^+\nu_\ell$ decays and the non-leptonic $D^*/D_0/D_2^* \to DP$ decays have been obtained, and then the resonant branching ratios have been obtained by the narrow width approximation after considering the resonant experimental data of the $B \to DP\ell'^+\nu_{\ell'}$ decays. Our main results can be summarized as follows.

For the non-resonant $B \to DP\ell^+\nu_\ell$ decays, the central values of $\mathcal{B}(B^+ \to D^-\pi^+\ell'^+\nu_{\ell'})_N$, $\mathcal{B}(B^+ \to \overline{D}^0\pi^0\ell'^+\nu_{\ell'})_N$, $\mathcal{B}(B^+ \to \overline{D}^0\eta\ell'^+\nu_{\ell'})_N$, $\mathcal{B}(B^0 \to \overline{D}^0\pi^-\ell'^+\nu_{\ell'} \mathcal{B}(B^0 \to D_s^-K^0\ell'^+\nu_{\ell'})_N$, $\mathcal{B}(B^0 \to D^-\pi^0\ell'^+\nu_{\ell'})_N$, $\mathcal{B}(B^0 \to D^-\eta\ell'^+\nu_{\ell'})_N$, $\mathcal{B}(B^0_s \to \overline{D}^0K^-\ell'^+\nu_{\ell'} \mathcal{B}(B^0_s \to D^-\overline{K}^0\ell'^+\nu_{\ell'})_N$ and $\mathcal{B}(B^0_s \to D_s^-\eta\ell'^+\nu_{\ell'})_N$ are on the orders of 10^{-4} , which could be measured by the LHCb and Belle II experiments. Other non-resonant decay branching ratios of $B^+ \to \overline{D}^0\eta'\ell'^+\nu_{\ell'}$, $B^0 \to D^-\eta'\ell'^+\nu_{\ell'}$, $B^0_s \to D^-_s\eta'\ell'^+\nu_{\ell'}$ and all $B \to DP\tau^+\nu_\tau$ decays are strongly suppressed by the narrow phase spaces.

For the charmed vector resonant $B \to DP\ell^+\nu_\ell$ decays, they give the dominant contributions in the $B^+ \to \overline{D}^0 \pi^0 \ell^+ \nu_\ell$, $B^0 \to \overline{D}^0 \pi^- \ell^+ \nu_\ell$ and $B^0 \to D^- \pi^0 \ell^+ \nu_\ell$ decays. Nevertheless, since the resonance D^{*0} can not decay into $D^- \pi^+$, the total branching ratios of the $B^+ \to D^- \pi^+ \ell^+ \nu_\ell$ decays are much smaller than ones of the $B^+ \to \overline{D}^0 \pi^0 \ell^+ \nu_\ell$, $B^0 \to \overline{D}^0 \pi^- \ell^+ \nu_\ell$ and $B^0 \to D^- \pi^0 \ell^+ \nu_\ell$ decays.

As for the charmed scalar and tensor resonant $B \to DP\ell^+\nu_\ell$ decays, our predictions of $\mathcal{B}(B^0 \to \overline{D}^0 \pi^- \ell'^+ \nu_{\ell'})_{D_0^-, D_2^{*-}}$ are more precise than their experimental measurements. The contributions of the D_0 and D_2^* resonances are in the same order of magnitude in the $B^+ \to D^- \pi^+ \ell'^+ \nu_{\ell'}$ and $B^0 \to \overline{D}^0 \pi^- \ell'^+ \nu_{\ell'}$ decays. But the contributions of the D_0 resonances are larger than ones of D_2^* resonances in the $B^+ \to \overline{D}^0 \pi^0 \ell^+ \nu_\ell$, $B^0 \to D^- \pi^0 \ell^+ \nu_\ell$, $B^+ \to D^- \pi^+ \tau^+ \nu_\tau$ and $B^0 \to \overline{D}^0 \pi^- \tau^+ \nu_\tau$ decays.

Although only approximate predictions can be obtained by the SU(3) flavor symmetry, they are still useful for understanding these decays. Until now, our predictions of the $B \to DP\ell^+\nu_\ell$ decays are quite coincident with present experimental data, and they could be tested in future experiments, such as LHCb and Belle II.

ACKNOWLEDGEMENTS

The work was supported by the National Natural Science Foundation of China (No. 12175088 and No. 12365014).

References

- [1] R. L. Workman et al. [Particle Data Group], PTEP 2022, 083C01 (2022).
- [2] T. Feldmann, D. Van Dyk and K. K. Vos, JHEP 10, 030 (2018) [arXiv:1807.01924 [hep-ph]].
- [3] X. G. He, Eur. Phys. J. C 9, 443-448 (1999) [arXiv:hep-ph/9810397 [hep-ph]].
- [4] X. G. He, Y. K. Hsiao, J. Q. Shi, Y. L. Wu and Y. F. Zhou, Phys. Rev. D 64, 034002 (2001) [arXiv:hep-ph/0011337 [hep-ph]].
- [5] H. K. Fu, X. G. He and Y. K. Hsiao, Phys. Rev. D 69, 074002 (2004) [arXiv:hep-ph/0304242 [hep-ph]].
- [6] Y. K. Hsiao, C. F. Chang and X. G. He, Phys. Rev. D 93, no.11, 114002 (2016) [arXiv:1512.09223 [hep-ph]].
- [7] X. G. He and G. N. Li, Phys. Lett. B 750, 82-88 (2015) [arXiv:1501.00646 [hep-ph]].
- [8] M. He, X. G. He and G. N. Li, Phys. Rev. D 92, no.3, 036010 (2015) [arXiv:1507.07990 [hep-ph]].
- [9] N. G. Deshpande and X. G. He, Phys. Rev. Lett. 75, 1703-1706 (1995) [arXiv:hep-ph/9412393 [hep-ph]].
- [10] M. Gronau, O. F. Hernandez, D. London and J. L. Rosner, Phys. Rev. D 50, 4529-4543 (1994) [arXiv:hep-ph/9404283 [hep-ph]].
- [11] M. Gronau, O. F. Hernandez, D. London and J. L. Rosner, Phys. Rev. D 52, 6356-6373 (1995) [arXiv:hep-ph/9504326 [hep-ph]].
- [12] S. Shivashankara, W. Wu and A. Datta, Phys. Rev. D 91, no.11, 115003 (2015) [arXiv:1502.07230 [hep-ph]].
- [13] S. H. Zhou, Q. A. Zhang, W. R. Lyu and C. D. Lü, Eur. Phys. J. C 77, no.2, 125 (2017) [arXiv:1608.02819 [hep-ph]].
- [14] H. Y. Cheng, C. W. Chiang and A. L. Kuo, Phys. Rev. D 91, no.1, 014011 (2015) [arXiv:1409.5026 [hep-ph]].
- [15] R. M. Wang, Y. G. Xu, C. Hua and X. D. Cheng, Phys. Rev. D 103, no.1, 013007 (2021) [arXiv:2101.02421 [hep-ph]].
- [16] R. M. Wang, X. D. Cheng, Y. Y. Fan, J. L. Zhang and Y. G. Xu, J. Phys. G 48, no.8, 085001 (2021) [arXiv:2008.06624 [hep-ph]].
- [17] Y. Grossman and D. J. Robinson, JHEP 04, 067 (2013) [arXiv:1211.3361 [hep-ph]].
- [18] D. Pirtskhalava and P. Uttayarat, Phys. Lett. B 712, 81-86 (2012) [arXiv:1112.5451 [hep-ph]].
- [19] M. J. Savage and R. P. Springer, Phys. Rev. D 42, 1527-1543 (1990).
- [20] M. J. Savage, Phys. Lett. B 257, 414-418 (1991).
- [21] G. Altarelli, N. Cabibbo and L. Maiani, Phys. Lett. B 57, 277-280 (1975).
- [22] C. D. Lü, W. Wang and F. S. Yu, Phys. Rev. D 93, no.5, 056008 (2016) [arXiv:1601.04241 [hep-ph]].
- [23] C. Q. Geng, Y. K. Hsiao, Y. H. Lin and L. L. Liu, Phys. Lett. B 776, 265-269 (2018) [arXiv:1708.02460 [hep-ph]].
- [24] C. Q. Geng, Y. K. Hsiao, C. W. Liu and T. H. Tsai, Phys. Rev. D 97, no.7, 073006 (2018) [arXiv:1801.03276 [hep-ph]].
- [25] C. Q. Geng, Y. K. Hsiao, C. W. Liu and T. H. Tsai, JHEP 11, 147 (2017) [arXiv:1709.00808 [hep-ph]].
- [26] C. Q. Geng, C. W. Liu, T. H. Tsai and S. W. Yeh, Phys. Lett. B 792, 214-218 (2019) [arXiv:1901.05610 [hep-ph]].
- [27] W. Wang, Z. P. Xing and J. Xu, Eur. Phys. J. C 77, no.11, 800 (2017) [arXiv:1707.06570 [hep-ph]].
- [28] D. Wang, Eur. Phys. J. C 79, no.5, 429 (2019) [arXiv:1901.01776 [hep-ph]].
- [29] D. Wang, P. F. Guo, W. H. Long and F. S. Yu, JHEP 03, 066 (2018) [arXiv:1709.09873 [hep-ph]].
- [30] S. Müller, U. Nierste and S. Schacht, Phys. Rev. D 92, no.1, 014004 (2015) [arXiv:1503.06759 [hep-ph]].
- [31] Y. Qiao, Y. X. Liu, Y. G. Xu and R. M. Wang, [arXiv:2404.03857 [hep-ph]].
- [32] R. M. Wang, M. Z. Yang, H. B. Li and X. D. Cheng, Phys. Rev. D 100, no.7, 076008 (2019) [arXiv:1906.08413 [hep-ph]].
- [33] Y. G. Xu, X. D. Cheng, J. L. Zhang and R. M. Wang, J. Phys. G 47, no.8, 085005 (2020) [arXiv:2001.06907 [hep-ph]].
- [34] H. M. Chang, M. González-Alonso and J. Martin Camalich, Phys. Rev. Lett. 114, no.16, 161802 (2015) [arXiv:1412.8484

[hep-ph]].

- [35] P. Zenczykowski, Phys. Rev. D 73, 076005 (2006) [arXiv:hep-ph/0512122 [hep-ph]].
- [36] P. Zenczykowski, Nucl. Phys. B Proc. Suppl. 167, 54-57 (2007) [arXiv:hep-ph/0610191 [hep-ph]].
- [37] N. Cabibbo, Phys. Rev. Lett. 10, 531-533 (1963).
- [38] G. Martinelli, M. Naviglio, S. Simula and L. Vittorio, Phys. Rev. D 106, no.9, 093002 (2022) [arXiv:2204.05925 [hep-ph]].
- [39] R. M. Wang, Y. Qiao, Y. J. Zhang, X. D. Cheng and Y. G. Xu, [arXiv:2301.00090 [hep-ph]].
- [40] M. Imbeault and D. London, Phys. Rev. D 84, 056002 (2011) [arXiv:1106.2511 [hep-ph]].
- [41] Y. L. Wu and Y. F. Zhou, Phys. Rev. D 72, 034037 (2005) [arXiv:hep-ph/0503077 [hep-ph]].
- [42] H. Y. Cheng and C. W. Chiang, Phys. Rev. D 86, 014014 (2012) [arXiv:1205.0580 [hep-ph]].
- [43] A. Dery, M. Ghosh, Y. Grossman and S. Schacht, JHEP 03 (2020), 165 [arXiv:2001.05397 [hep-ph]].
- [44] S. Sasaki and T. Yamazaki, Phys. Rev. D 79 (2009), 074508 [arXiv:0811.1406 [hep-ph]].
- [45] G. S. Yang and H. C. Kim, Phys. Rev. C 92, 035206 (2015) [arXiv:1504.04453 [hep-ph]].
- [46] T. N. Pham, Phys. Rev. D 87 (2013) no.1, 016002 [arXiv:1210.3981 [hep-ph]].
- [47] C. Q. Geng, Y. K. Hsiao, C. W. Liu and T. H. Tsai, Eur. Phys. J. C 78 (2018) no.7, 593 [arXiv:1804.01666 [hep-ph]].
- [48] R. Flores-Mendieta, E. E. Jenkins and A. V. Manohar, Phys. Rev. D 58 (1998), 094028 [arXiv:hep-ph/9805416 [hep-ph]].
- [49] D. Xu, G. N. Li and X. G. He, Int. J. Mod. Phys. A 29 (2014), 1450011 [arXiv:1307.7186 [hep-ph]].
- [50] X. G. He, G. N. Li and D. Xu, Phys. Rev. D **91**, no.1, 014029 (2015) [arXiv:1410.0476 [hep-ph]].
- [51] H. Y. Cheng, C. Y. Cheung, W. Dimm, G. L. Lin, Y. C. Lin, T. M. Yan and H. L. Yu, Phys. Rev. D 48, 3204-3220 (1993) [arXiv:hep-ph/9305340 [hep-ph]].
- [52] S. Y. Tsai and Y. K. Hsiao, [arXiv:2107.03634 [hep-ph]].
- [53] A. Le Yaouanc, J. P. Leroy and P. Roudeau, Phys. Rev. D 99, no.7, 073010 (2019) [arXiv:1806.09853 [hep-ph]].
- [54] Y. J. Shi and U. G. Meißner, Eur. Phys. J. C 81, no.5, 412 (2021) [arXiv:2103.12977 [hep-ph]].
- [55] C. S. Kim, J. Lee and W. Namgung, Phys. Rev. D 60, 094019 (1999) [arXiv:hep-ph/9904470 [hep-ph]].
- [56] C. S. Kim, J. Lee and W. Namgung, Phys. Rev. D 59, 114005 (1999) [arXiv:hep-ph/9811395 [hep-ph]].
- [57] N. N. Achasov, A. V. Kiselev and G. N. Shestakov, Phys. Rev. D 102, no.1, 016022 (2020) [arXiv:2005.06455 [hep-ph]].
- [58] J. Wiss, eConf C070805, 34 (2007) [arXiv:0709.3247 [hep-ex]].
- [59] S. Faller, T. Feldmann, A. Khodjamirian, T. Mannel and D. van Dyk, Phys. Rev. D 89, no.1, 014015 (2014) [arXiv:1310.6660 [hep-ph]].
- [60] W. Wang, Phys. Lett. B 759, 501-506 (2016) [arXiv:1602.05288 [hep-ph]].
- [61] N. N. Achasov, A. V. Kiselev and G. N. Shestakov, Phys. Rev. D 104, no.1, 016034 (2021) [arXiv:2106.10670 [hep-ph]].
- [62] E. J. Gustafson, F. Herren, R. S. Van de Water, R. van Tonder and M. L. Wagman, [arXiv:2311.00864 [hep-ph]].
- [63] C. Hambrock and A. Khodjamirian, Nucl. Phys. B 905, 373-390 (2016) [arXiv:1511.02509 [hep-ph]].
- [64] P. Böer, T. Feldmann and D. van Dyk, JHEP 02, 133 (2017) [arXiv:1608.07127 [hep-ph]].
- [65] H. Y. Cheng and W. S. Hou, Phys. Lett. B 566, 193-200 (2003) [arXiv:hep-ph/0305038 [hep-ph]].
- [66] M. Y. Wan, Y. G. Xu, Q. L. Jia, Y. X. Liu and Y. J. Zhang, [arXiv:2403.14929 [hep-ph]].
- [67] S. Okubo, Phys. Lett. 5, 165-168 (1963).
- [68] H. J. Lipkin, Nucl. Phys. B 291, 720-730 (1987).
- [69] H. J. Lipkin and B. s. Zou, Phys. Rev. D 53, 6693-6696 (1996).
- [70] H. Y. Cheng and X. W. Kang, Eur. Phys. J. C 77 (2017) no.9, 587 [erratum: Eur. Phys. J. C 77 (2017) no.12, 863] [arXiv:1707.02851 [hep-ph]].
- [71] H. Y. Cheng, C. K. Chua and C. W. Hwang, Phys. Rev. D 69 (2004), 074025 [arXiv:hep-ph/0310359 [hep-ph]].
- [72] M. A. Ivanov, J. G. Körner, J. N. Pandya, P. Santorelli, N. R. Soni and C. T. Tran, Front. Phys. (Beijing) 14 (2019) no.6, 64401 [arXiv:1904.07740 [hep-ph]].

- [73] E. J. Totten, M. J. Irwin and P. A. Whitelock, Mon. Not. Roy. Astron. Soc. 314, 630-640 (2000) [arXiv:astro-ph/0001113 [astro-ph]].
- [74] F. Abudinén et al. [Belle-II], Phys. Rev. D 107, no.7, 072002 (2023) [arXiv:2205.06372 [hep-ex]].
- [75] R. C. Verma, J. Phys. G 39, 025005 (2012) [arXiv:1103.2973 [hep-ph]].
- [76] L. Chen, Y. W. Ren, L. T. Wang and Q. Chang, Eur. Phys. J. C 82, no.5, 451 (2022) [arXiv:2112.08016 [hep-ph]].
- [77] K. Azizi, H. Sundu and S. Sahin, Eur. Phys. J. C 75, no.5, 197 (2015) [arXiv:1411.3100 [hep-ph]].
- [78] C. S. Kim, J. Lee and W. Namgung, Phys. Rev. D 59, 114006 (1999) [arXiv:hep-ph/9811396 [hep-ph]].
- [79] H. Y. Cheng and C. K. Chua, Phys. Rev. D 102, no.5, 053006 (2020) [arXiv:2007.02558 [hep-ph]].
- [80] P. L. Cho and M. B. Wise, Phys. Rev. D 49, 6228-6231 (1994) [arXiv:hep-ph/9401301 [hep-ph]].
- [81] Z. G. Wang, Eur. Phys. J. C 74, no.10, 3123 (2014) [arXiv:1406.2632 [hep-ph]].
- [82] C. S. Kim, G. L. Castro and S. L. Tostado, Phys. Rev. D 95, no.7, 073003 (2017) [arXiv:1702.01704 [hep-ph]].