

Two-loop Quarkonium Hamiltonian in Non-annihilation Channel

Go Mishima,¹ Yukinari Sumino,¹ and Hiromasa Takaura²

¹*Department of Physics, Tohoku University, Sendai, 980-8578 Japan*

²*Center for Gravitational Physics and Quantum Information,
Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan*

(Dated: July 2, 2024)

We calculate the two-loop heavy quarkonium Hamiltonian within potential-NRQCD effective field theory in the non-annihilation channel. This calculation represents the first non-trivial step towards determining the N⁴LO Hamiltonian in the weak coupling regime. To cope with the large amount of calculations, we make use of the β expansion and differential equation for master integrals and systematize the procedure.

PACS numbers:

The heavy quarkonium system provides an ideal laboratory for investigating theoretical and phenomenological aspects of QCD in depth [1–3]. In particular, analyses of quarkonium systems in the weak coupling regime have enabled precise examinations of QCD, in the case that their relevant physical scales are much larger than the typical QCD scale $\Lambda_{\text{QCD}} \sim 300$ MeV. For instance, these analyses have led to the determinations of fundamental physical constants such as the charm quark mass, bottom quark mass, and strong coupling constant α_s [4, 5]. Moreover, physics studies in the top-antitop quark threshold region in future e^+e^- collider experiments are expected to be major targets of high energy experiments, prompting enormous efforts towards their high-precision theoretical studies [6, 7]. The quarkonium Hamiltonian within the potential Nonrelativistic QCD (pNRQCD) effective field theory (EFT) [8] has played a principal role in these precision analyses. The Hamiltonian has been known up to the next-to-next-to-next-to-leading order (N³LO) accuracy for more than a decade [9–11], and it took roughly a decade to calculate the N³LO Hamiltonian after the N²LO Hamiltonian was completed [12]. It is now timely to start calculations at N⁴LO accuracy.

In this paper, as a first non-trivial partial calculation of the N⁴LO quarkonium Hamiltonian, we calculate the Hamiltonian at the two-loop level, or at $\mathcal{O}(\alpha_s^3)$, and up to $\mathcal{O}(\beta^0)$ [relative $\mathcal{O}(\alpha_s^2\beta^2)$ compared to the LO $\sim \alpha_s/\beta^2$ contribution]. Here, β denotes the velocity of the heavy quark or antiquark in the center-of-mass (c.m.) frame and is a small expansion parameter of the quarkonium system. To reduce the labor of the calculation, we compute only the contributions from the non-annihilation channel of the quark and antiquark. Our motivation is to establish a calculational procedure for such a complex calculation.

We consider the SU(3) color gauge theory with n_h heavy quark flavors (each with the same mass m) and n_l massless quark flavors. (For convenience we call this

theory as QCD.) We calculate the scattering amplitude between a heavy quark Q and a heavy antiquark \bar{Q}' of different flavors (with mass m). In this case only diagrams in the non-annihilation channel contribute. We evaluate this amplitude in the β expansion ($1/m$ expansion) in the c.m. frame, which is valid in the non-relativistic region $\beta \ll 1$. Until now, no analytic evaluation of the full relativistic scattering amplitude at the two-loop level has been available. Hence, we aim to devise a method to systematically evaluate this series expansion. We also calculate the same scattering amplitude in pNRQCD EFT. Then we determine the singlet Hamiltonian in the latter theory by matching the two amplitudes (“direct matching”).

Let us first highlight the characteristic features of our calculational method compared to the previous one. The calculation of the N³LO Hamiltonian [9] was, so to speak, done “manually,” utilizing the expansion-by-regions (EBR) technique [13]. This technique was used to separate contributions from the hard (H) and soft (S) regions from those of the potential (P) and ultra-soft (US) regions, where only the H and S contributions constitute the Hamiltonian of the EFT. This is not sufficient to determine the Hamiltonian unless we perform the matching off-shell. In the EFT, there are different choices for the operator basis of the Hamiltonian that are equivalent on-shell, and depending on which one is chosen, the off-shell effects within the loop change. Therefore, it is necessary to identify and manually add the operators (“off-shell operators” [9]) that compensate for these effects and reproduce correctly the S -matrix elements.

However, it is highly demanded to systematize the calculational procedure to handle the vast amount of calculations required at higher orders. In particular, identifying the off-shell operators in the above manual procedure seems difficult to systematize at higher orders. In contrast, we systematize the calculation of the β expansion with the aid of the differential equation satisfied by the

master integrals, which enables calculation of contributions from the whole regions manageable.

Our procedure does not separate contributions from the S and P regions, although contributions from the P region eventually cancel out in the calculation of the Hamiltonian. This approach avoids the need to identify the off-shell operators, at the cost of calculating the P contributions which eventually cancel. (The direct matching procedure incorporates the off-shell operators and the cancellation of the P contributions simultaneously.) In short, our method is adapted to systematically handle the large amount of calculations.

We calculate the QCD two-loop on-shell scattering amplitude for $Q(\vec{p}) + \bar{Q}'(-\vec{p}) \rightarrow Q(\vec{p}') + \bar{Q}'(-\vec{p}')$ in the color singlet channel and in the c.m. frame as follows, where

$$\vec{p}' = \vec{p} + \vec{k}, \quad |\vec{p}'|^2 = |\vec{p}|^2. \quad (1)$$

We first project the amplitude on to a spinor basis. Feynman integrals are regularized by dimensional regularization, where the number of the space-time dimensions is set to $4 - 2\epsilon$. The spinor basis can be easily expressed by a two-component spinor basis using the Pauli matrices in dimensional regularization, satisfying

$$\{\sigma^i, \sigma^j\} = 2\delta^{ij}\mathbb{I}, \quad \delta^{ii} = 3 - 2\epsilon, \quad \text{tr}\mathbb{I} = 2. \quad (2)$$

We adopt a 21-dimensional basis $\{\Lambda_1, \dots, \Lambda_{21}\}$, where the first five elements are chosen as

$$\begin{aligned} \Lambda_1 &= \mathbb{I} \otimes \mathbb{I}, \\ \Lambda_2 &= \sigma^a \sigma^b \otimes \sigma^a \sigma^b, \\ \Lambda_3 &= \sigma^a \sigma^b \sigma^c \sigma^d \otimes \sigma^a \sigma^b \sigma^c \sigma^d, \\ \Lambda_4 &= \frac{1}{m^2} \left(\vec{\sigma} \cdot \vec{k} \sigma^a \otimes \vec{\sigma} \cdot \vec{k} \sigma^a \right), \\ \Lambda_5 &= \frac{1}{m^2} \left(\vec{\sigma} \cdot \vec{p}' \vec{\sigma} \cdot \vec{p} \otimes \mathbb{I} + \mathbb{I} \otimes \vec{\sigma} \cdot \vec{p}' \vec{\sigma} \cdot \vec{p} \right). \end{aligned} \quad (3)$$

m denotes the pole mass of the heavy quarks, and the amplitude is renormalized in the on-shell scheme. The coefficients of the spinor basis are expressed by scalar integrals by the projection.

Next we express the scalar integrals by master integrals using the integration-by-parts identities [14]. We use the program `Kira` [15–17] (also `LiteRed` [18] and `FIRE` [19] for cross checks) for this reduction, by which all the coefficients are expressed by 149 master integrals. Up to this stage the obtained expression is exact.

We expand the master integrals in β . This is done by solving the differential equation satisfied by the master integrals [20] in series expansions in β , that is, expansions in p and k . We can interpret the β expansions in the language of the EBR technique, where non-zero contributions originate from seven regions: HH, HS, HP, SS, SP, PP, and PUS regions. We can group them into four regions as HH, HS+HP, SS+SP+PP and PUS. Then the differential equation is satisfied independently by each of

the contributions from these four regions.[29] We determine the boundary condition for the solution corresponding to each region by evaluating the LO term of the expansion using the EBR technique. Moreover, in many cases, the boundary condition can be fixed by simply demanding regularity of the solution at $u = \vec{k}^2 - 4\vec{p}^2 = 0$ in the physical sheet. Once we have the β expansions of the master integrals, it is straightforward to expand the scattering amplitude in β . This procedure is much more efficient and economical than to calculate each master integral or each diagram by only the EBR technique.

The structure of the QCD $Q\bar{Q}'$ scattering amplitude is fairly complicated, even after the expansion in β . Although the amplitude is regular at $u = 0$ in the physical sheet, it has singularities at $u = 0$ in the second and other Riemann sheets by analytical continuation. [An example of such a structure is $\log(4\vec{p}^2/\vec{k}^2)/(\vec{k}^2 - 4\vec{p}^2)$.] The regularity of the amplitude at $u = 0$ in the physical sheet follows from the flavor conservation of QCD. Furthermore, the contributions from the P region include non-elementary functions of p/k (counted as order one in the β expansion).

The calculational procedure for the scattering amplitude for the same process in pNRQCD EFT is similar.[30] The amplitude is projected on to the two-component spinor basis. The scalar integrals are reduced to master integrals of the EFT using the integration-by-parts identities. After matching the amplitude to that of QCD, we readily obtain the two-loop Hamiltonian in the β expansion. The Hamiltonian up to $\mathcal{O}(\beta^0)$ consists only of the five operators $\Lambda_1, \dots, \Lambda_5$ of the spinor basis before expansion in ϵ , where the coefficients of the other operators vanish.

We have performed the following cross checks for our results before expanding them in ϵ . (1) We checked that the QCD scattering amplitude is regular at $u = 0$ in the physical sheet, even though this is not so obvious from the obtained expression, which has singularities at $u = 0$ in the second and other Riemann sheets and contains non-elementary functions of p/k . (2) The two-loop Hamiltonian is regular in \vec{p} but singular in \vec{k} . Each term of the coefficient of Λ_i takes the form $P_1(\vec{p})V(k)P_2(\vec{p}')$, where P_j is a homogeneous polynomial and $V(k)$ is proportional to k^{a+be} with $a \in \{-2, -1, 0\}$, $b \in \{-4, -2, 0\}$. This form is expected to originate from the hard (H) and soft (S) contributions according to the EBR technique and is consistent with the concept of the EFT construction. This means that all the singularities at $u = 0$ in the second and other Riemann sheets cancel between the scattering amplitudes of QCD and pNRQCD EFT. Additionally, the functions of p/k (P contributions) also cancel out.

After expanding our results in ϵ , we have performed the following cross checks. (3) We reproduced the known two-loop static QCD potential $V_{\text{QCD}}^{(2 \text{ loop})}$ [12] and two-loop

$1/(mk)$ potential $V_{1/(mk)}^{(2 \text{ loop})}$ [21] as part of the two-loop Hamiltonian. We also reproduced the known one-loop part of the $N^3\text{LO}$ Hamiltonian [9] at an intermediate stage of the calculation. (4) We evaluated the coefficients of the expansion in ϵ of each QCD master integral by numerical integrations (in part using FIESTA [22, 23]) in the non-relativistic region. We compared them with the expansion of the master integral in β and ϵ obtained analytically and checked consistency. (5) Some of the ϵ expansions of the master integrals are known analytically [24, 25]. We expanded the analytical expressions in β and found agreement with our results.

Let us present our final result. We expand the coefficients of $\Lambda_1, \dots, \Lambda_5$ in ϵ while we ignore any $\mathcal{O}(\epsilon)$ contributions in Λ_i . (Namely, we simply take the limit $\epsilon \rightarrow 0$ for Λ_i .) The Hamiltonian is given in the form

$$H = \frac{16\pi^2 C_F}{k^2} \sum_{i=1}^4 \sum_{j=1}^3 \sum_{n, \ell \geq 0} \left(\frac{\alpha_s(k)}{4\pi} \right)^j \times C_{\{i,j,n,2\ell\}} \left(\frac{k}{m} \right)^n \frac{(p^2)^\ell + (p'^2)^\ell}{2m^{2\ell}} O_i, \quad (4)$$

where

$$p^2 = |\vec{p}|^2, \quad p'^2 = |\vec{p}'|^2, \quad k = |\vec{k}| = |\vec{p}' - \vec{p}|. \quad (5)$$

We absorb $\log(\mu/k)$ terms originating from the running of α_s by expressing the Hamiltonian by $\alpha_s(k)$, the strong coupling constant in the modified minimal subtraction ($\overline{\text{MS}}$) scheme of the theory with n_l flavors only, renormalized at $\mu = k$. [31][32] The spinor basis in three dimensions is defined as

$$O_1 = \mathbb{I} \otimes \mathbb{I}, \quad O_2 = \vec{S}^2, \quad O_3 = \frac{i}{k^2} \vec{S} \cdot (\vec{p} \times \vec{k}), \quad O_4 = \sigma^a \otimes \sigma^a - \frac{3}{k^2} (\vec{k} \cdot \vec{\sigma}) \otimes (\vec{k} \cdot \vec{\sigma}), \quad (6)$$

with

$$\vec{S} = \frac{\vec{\sigma}}{2} \otimes \mathbb{I} + \mathbb{I} \otimes \frac{\vec{\sigma}}{2}. \quad (7)$$

The Wilson coefficients are separated into finite and divergent parts as

$$C_{\{i,j,n,2\ell\}} = C_{\{i,j,n,2\ell\}}^{\text{fin}} + C_{\{i,j,n,2\ell\}}^{\text{div}}. \quad (8)$$

Two-loop non-zero finite Wilson coefficients up to $\mathcal{O}(\beta^0)$ are given by

$$C_{\{1,3,0,0\}}^{\text{fin}} = V_{\text{QCD}}^{(2 \text{ loop})}(k) \cdot (4\pi k^2) / [C_F \alpha_s(k)^3], \quad (9)$$

$$C_{\{1,3,1,0\}}^{\text{fin}} = V_{1/(mk)}^{(2 \text{ loop})}(k) \cdot (4\pi mk) / [C_F \alpha_s(k)^3], \quad (10)$$

$$C_{\{1,3,2,0\}}^{\text{fin}} = L_{m/\mu} \left(-\frac{208}{9} - \frac{260\pi^2}{9} + \frac{40n_l}{27} \right) + L_{k/m} \left(-\frac{106\pi^2}{9} - \frac{2152}{9} + \frac{3013n_l}{162} - \frac{17n_h}{6} \right) + L_{k/m}^2 \left(-\frac{1825}{18} + \frac{109n_l}{27} \right) + \frac{27\pi^4}{16} + \frac{1285\zeta(3)}{6} - \frac{11}{3}\pi^2 \log(2) + \frac{99541\pi^2}{1944} + \frac{41395}{162} + \frac{8}{27}n_h n_l + n_h \left(\frac{277\pi^2}{324} - \frac{1375}{108} \right) + n_l \left(\frac{139}{81} + \frac{220\pi^2}{81} \right), \quad (11)$$

$$C_{\{1,3,0,2\}}^{\text{fin}} = L_{k/\mu} \left(32\pi^2 + 416 - \frac{80n_l}{3} \right) - \frac{100n_l^2}{81} + n_l \left(\frac{52\zeta(3)}{3} + \frac{1901}{27} - \frac{4\pi^2}{9} \right) - \frac{9\pi^4}{4} - 114\zeta(3) + \frac{266\pi^2}{3} - \frac{7919}{18}, \quad (12)$$

$$C_{\{2,3,2,0\}}^{\text{fin}} = \frac{352}{27}\pi^2 L_{m/\mu} + L_{k/m}^2 \left(38 - \frac{5n_l}{3} \right) + L_{k/m} \left(\frac{584}{9} - \frac{182n_l}{27} - 4\pi^2 \right) + \frac{100n_l^2}{243} + n_l \left(-\frac{52\zeta(3)}{9} - \frac{1777}{81} - \frac{10\pi^2}{9} \right) - \frac{3\pi^4}{4} - \frac{1181\zeta(3)}{6} - \frac{121}{9}\pi^2 \log(2) - \frac{1133\pi^2}{81} + \frac{10771}{54} + n_h \left(\frac{770}{81} - \frac{16\pi^2}{27} \right), \quad (13)$$

$$C_{\{3,3,2,0\}}^{\text{fin}} = L_{k/m}^2 (42 - 2n_l) + L_{k/m} \left(\frac{1010}{3} - \frac{242n_l}{9} \right) + n_l \left(-26\zeta(3) - \frac{4591}{54} - \frac{4\pi^2}{3} \right) - \frac{27\pi^4}{8} + \frac{161\zeta(3)}{3} + \frac{56}{9}\pi^2 \log(2) + \frac{820\pi^2}{27} + \frac{7823}{12} + \frac{50n_l^2}{27} + n_h \left(\frac{1010}{27} - \frac{34\pi^2}{9} \right), \quad (14)$$

$$C_{\{4,3,2,0\}}^{\text{fin}} = L_{k/m}^2 \left(\frac{17}{4} - \frac{n_l}{6} \right) + L_{k/m} \left(\frac{583}{18} - \frac{121n_l}{54} \right) + n_l \left(-\frac{13\zeta(3)}{9} - \frac{2257}{324} - \frac{\pi^2}{9} \right) - \frac{3\pi^4}{16} + \frac{31\zeta(3)}{18} + \frac{14}{27}\pi^2 \log(2) + \frac{545\pi^2}{162} + \frac{15739}{216} + \frac{25n_l^2}{243} + n_h \left(\frac{505}{162} - \frac{17\pi^2}{54} \right), \quad (15)$$

where $L_{a/b}$ represents $\log(a^2/b^2)$. Note that $V_{\text{QCD}}^{(2 \text{ loop})}$ and $V_{1/(mk)}^{(2 \text{ loop})}$ do not include the heavy-quark-loop contributions, which are included in other Wilson coefficients.

Two-loop non-zero divergent Wilson coefficients are

given by

$$C_{\{1,3,1,0\}}^{\text{div}} = -\frac{136\pi^2}{3\epsilon}, \quad (16)$$

$$C_{\{1,3,2,0\}}^{\text{div}} = L_{k/\mu} \left(\frac{44}{9\epsilon} - \frac{8n_l}{27\epsilon} \right) - \frac{22}{9\epsilon^2} + \frac{130\pi^2}{9\epsilon} + \frac{104}{9\epsilon} + n_l \left(\frac{4}{27\epsilon^2} - \frac{20}{27\epsilon} \right), \quad (17)$$

$$C_{\{1,3,0,2\}}^{\text{div}} = L_{k/\mu} \left(-\frac{88}{\epsilon} + \frac{16n_l}{3\epsilon} \right) + \frac{44}{\epsilon^2} - \frac{16\pi^2}{\epsilon} - \frac{208}{\epsilon} + n_l \left(\frac{40}{3\epsilon} - \frac{8}{3\epsilon^2} \right), \quad (18)$$

$$C_{\{2,3,2,0\}}^{\text{div}} = -\frac{176\pi^2}{27\epsilon}. \quad (19)$$

In summary we computed the quarkonium Hamiltonian at the two-loop level in the non-annihilation channel. The obtained Hamiltonian has an expected form as resulting from integrating the H and S modes. This shows that the singular structure of the scattering amplitude at $u = 0$ in the second and other Riemann sheets as well as non-elementary functions of p/k originating from the P region are reproduced correctly by the EFT and canceled in the calculation of the Hamiltonian. The developed calculational procedure would also be useful to compute various Wilson coefficients at high orders relevant for quarkonium observables. In particular, the obtained Hamiltonian will play a major role, for instance, in calculating the fine and hyperfine splittings of the quarkonium.

For a consistent calculation of physical observables at N⁴LO accuracy using the Hamiltonian, it is sometimes required to include higher-order terms of β and ϵ at the tree and one-loop levels than those given in the literature. We provide them as well as the expression of the two-loop Hamiltonian for general color factors in the Supplementary Material [28].

Acknowledgement

The works of Y.S. and H.T., respectively, were supported in part by JSPS KAKENHI Grant Numbers JP23K03404 and by JP19K14711 and JP23K13110. H.T. is the Yukawa Research Fellow supported by Yukawa Memorial Foundation.

-
- [1] N. Brambilla *et al.* [Quarkonium Working Group], doi:10.5170/CERN-2005-005 [arXiv:hep-ph/0412158 [hep-ph]].
- [2] N. Brambilla, S. Eidelman, B. K. Heltsley, R. Vogt, G. T. Bodwin, E. Eichten, A. D. Frawley, A. B. Meyer, R. E. Mitchell and V. Papadimitriou, *et al.* Eur. Phys. J. C **71** (2011), 1534 doi:10.1140/epjc/s10052-010-1534-9 [arXiv:1010.5827 [hep-ph]].

- [3] Y. Sumino, [arXiv:1411.7853 [hep-ph]].
- [4] Y. Kiyo, G. Mishima and Y. Sumino, Phys. Lett. B **752** (2016), 122-127 [erratum: Phys. Lett. B **772** (2017), 878-878] doi:10.1016/j.physletb.2015.11.040 [arXiv:1510.07072 [hep-ph]].
- [5] C. Peset, A. Pineda and J. Segovia, JHEP **09** (2018), 167 doi:10.1007/JHEP09(2018)167 [arXiv:1806.05197 [hep-ph]].
- [6] M. Beneke, Y. Kiyo and K. Schuller, [arXiv:1312.4791 [hep-ph]].
- [7] M. Vos, G. Abbas, M. Beneke, S. Bilokin, M. J. Costa, S. De Curtis, K. Fujii, J. Fuster, I. Garcia Garcia and P. Gomis, *et al.* [arXiv:1604.08122 [hep-ex]].
- [8] N. Brambilla, A. Pineda, J. Soto and A. Vairo, Rev. Mod. Phys. **77** (2005), 1423 doi:10.1103/RevModPhys.77.1423 [arXiv:hep-ph/0410047 [hep-ph]].
- [9] B. A. Kniehl, A. A. Penin, V. A. Smirnov and M. Steinhauser, Nucl. Phys. B **635** (2002), 357-383 doi:10.1016/S0550-3213(02)00403-0 [arXiv:hep-ph/0203166 [hep-ph]].
- [10] C. Anzai, Y. Kiyo and Y. Sumino, Phys. Rev. Lett. **104** (2010), 112003 doi:10.1103/PhysRevLett.104.112003 [arXiv:0911.4335 [hep-ph]].
- [11] A. V. Smirnov, V. A. Smirnov and M. Steinhauser, Phys. Rev. Lett. **104** (2010), 112002 doi:10.1103/PhysRevLett.104.112002 [arXiv:0911.4742 [hep-ph]].
- [12] Y. Schroder, DESY-THESIS-1999-021.
- [13] M. Beneke and V. A. Smirnov, Nucl. Phys. B **522** (1998), 321-344 doi:10.1016/S0550-3213(98)00138-2 [arXiv:hep-ph/9711391 [hep-ph]].
- [14] K. G. Chetyrkin and F. V. Tkachov, Nucl. Phys. B **192** (1981), 159-204 doi:10.1016/0550-3213(81)90199-1
- [15] P. Maierhöfer, J. Usovitsch and P. Uwer, Comput. Phys. Commun. **230** (2018), 99-112 doi:10.1016/j.cpc.2018.04.012 [arXiv:1705.05610 [hep-ph]].
- [16] J. Klappert, F. Lange, P. Maierhöfer and J. Usovitsch, Comput. Phys. Commun. **266** (2021), 108024 doi:10.1016/j.cpc.2021.108024 [arXiv:2008.06494 [hep-ph]].
- [17] R. H. Lewis, *Computer Algebra System Fermat*, <https://home.bway.net/lewis>.
- [18] R. N. Lee, J. Phys. Conf. Ser. **523** (2014), 012059 doi:10.1088/1742-6596/523/1/012059 [arXiv:1310.1145 [hep-ph]].
- [19] A. V. Smirnov, Comput. Phys. Commun. **189** (2015), 182-191 doi:10.1016/j.cpc.2014.11.024 [arXiv:1408.2372 [hep-ph]].
- [20] T. Gehrmann and E. Remiddi, Nucl. Phys. B **580** (2000), 485-518 doi:10.1016/S0550-3213(00)00223-6 [arXiv:hep-ph/9912329 [hep-ph]].
- [21] B. A. Kniehl, A. A. Penin, M. Steinhauser and V. A. Smirnov, Phys. Rev. D **65** (2002), 091503 doi:10.1103/PhysRevD.65.091503 [arXiv:hep-ph/0106135 [hep-ph]].
- [22] A. V. Smirnov, N. D. Shapurov and L. I. Vysotsky, Comput. Phys. Commun. **277** (2022), 108386 doi:10.1016/j.cpc.2022.108386 [arXiv:2110.11660 [hep-ph]].
- [23] M. Borinsky, Ann. Inst. H. Poincaré D Comb. Phys. Interact. **10** (2023) no.4, 635-685 doi:10.4171/aihpd/158 [arXiv:2008.12310 [math-ph]].
- [24] V. A. Smirnov, Phys. Lett. B **524** (2002), 129-

- 136 doi:10.1016/S0370-2693(01)01382-X [arXiv:hep-ph/0111160 [hep-ph]].
- [25] R. Bonciani, A. Ferroglia, P. Mastrolia, E. Remiddi and J. J. van der Bij, Nucl. Phys. B **681** (2004), 261-291 [erratum: Nucl. Phys. B **702** (2004), 364-366] doi:10.1016/j.nuclphysb.2004.08.003 [arXiv:hep-ph/0310333 [hep-ph]].
- [26] A. G. Grozin, P. Marquard, J. H. Piclum and M. Steinhauser, Nucl. Phys. B **789** (2008), 277-293 doi:10.1016/j.nuclphysb.2007.08.012 [arXiv:0707.1388 [hep-ph]].
- [27] A. G. Grozin, M. Hoeschele, J. Hoff, M. Steinhauser, M. Hoeschele, J. Hoff and M. Steinhauser, JHEP **09** (2011), 066 doi:10.1007/JHEP09(2011)066 [arXiv:1107.5970 [hep-ph]].
- [28] See Supplemental Material associated to this Letter.
- [29] The contributions from the S and P regions mix with each other through the prescription in the EBR technique [13], which subtracts the pinch singularities in the S region and compensates them as regularized contributions in the P region. As a result, the contributions from HS, HP, SS, SP and PP regions individually do not satisfy the differential equation.
- [30] We have not calculated the contributions from the US regions on the EFT side and simply assumed that they are the same as the PUS contribution of QCD. This is expected since in QCD the PUS contribution does not mix with other contributions and therefore is well defined.
- [31] In expressing the Hamiltonian, it is customary to rewrite the coupling constant of the full theory $\alpha_s^{(n_h+n_l)}(\mu)$ by that of the theory with n_l flavors only $\alpha_s(\mu) \equiv \alpha_s^{(n_l)}(\mu)$. We include the $\mathcal{O}(\epsilon)$ term in this decoupling relation [1, 2], which simplifies the result slightly since the one-loop Hamiltonian includes the $1/\epsilon$ pole.
- [32] We do not include the $\mathcal{O}(\epsilon)$ correction to the running formula

$$\alpha_s(k) = \alpha_s(\mu) - \frac{\alpha_s(\mu)^2}{4\pi} \left(11 - \frac{2n_l}{3} \right) \log \left(\frac{k^2}{\mu^2} \right) + \dots$$

Supplemental: More details of Hamiltonian

We present the tree-level and one-loop Hamiltonians before expanding in ϵ , up to $\mathcal{O}(\beta^2)$ and $\mathcal{O}(\beta)$ [$\mathcal{O}(\beta^4)$ and $\mathcal{O}(\alpha_s\beta^3)$ relative to the LO $\sim \alpha_s/\beta^2$], respectively. These are in general necessary ingredients to calculate physical observables at the N⁴LO accuracy. We also present the two-loop Hamiltonian after expansion in ϵ and retaining the color factors.

Tree-level and one-loop Hamiltonians

The Hamiltonian is given in the form

$$H = \frac{C_F \bar{\mu}^{-2\epsilon}}{k^2} \sum_{i=1}^6 \sum_{j=1}^3 \sum_{n,\ell \geq 0} (g_R^2 \bar{\mu}^{2\epsilon})^j W_{\{i,j,n,2\ell\}} \left(\frac{k}{m}\right)^n \frac{(p^2)^\ell + (p'^2)^\ell}{2m^{2\ell}} \Lambda_i, \quad (20)$$

where

$$k = |\vec{k}|, \quad p^2 = |\vec{p}|^2, \quad p'^2 = |\vec{p}'|^2, \quad \vec{p}' = \vec{p} + \vec{k}. \quad (21)$$

$g_R = \sqrt{4\pi\alpha_s^{(n_h+n_l)}}(\mu)$ denotes the renormalized gauge coupling constant in the $\overline{\text{MS}}$ scheme of the full theory (with n_h heavy quark flavors and n_l massless quark flavors); $\bar{\mu}^2 = \mu^2 e^{\gamma_E}/(4\pi)$. The spinor basis is defined in dimensional regularization as

$$\begin{aligned} \Lambda_1 &= \mathbb{I} \otimes \mathbb{I}, \\ \Lambda_2 &= \sigma^a \sigma^b \otimes \sigma^a \sigma^b, \\ \Lambda_3 &= \sigma^a \sigma^b \sigma^c \sigma^d \otimes \sigma^a \sigma^b \sigma^c \sigma^d, \\ \Lambda_4 &= \frac{1}{m^2} \left(\vec{\sigma} \cdot \vec{k} \sigma^a \otimes \vec{\sigma} \cdot \vec{k} \sigma^a \right), \\ \Lambda_5 &= \frac{1}{m^2} \left(\vec{\sigma} \cdot \vec{p}' \vec{\sigma} \cdot \vec{p} \otimes \mathbb{I} + \mathbb{I} \otimes \vec{\sigma} \cdot \vec{p}' \vec{\sigma} \cdot \vec{p} \right), \\ \Lambda_6 &= \frac{1}{m^4} \left(\vec{\sigma} \cdot \vec{p}' \vec{\sigma} \cdot \vec{p} \otimes \vec{\sigma} \cdot \vec{p}' \vec{\sigma} \cdot \vec{p} \right). \end{aligned} \quad (22)$$

We present the Wilson coefficients before expansion in ϵ . We list only those coefficients which are non-zero.

At tree level and up to $\mathcal{O}(\beta^4)$ relative to LO, they are given by

$$\begin{aligned} W_{\{1,1,0,0\}} &= -1, & W_{\{1,1,0,2\}} &= \frac{1}{2}, & W_{\{1,1,0,4\}} &= -\frac{7}{16}, \\ W_{\{4,1,0,0\}} &= -\frac{1}{4}, & W_{\{4,1,0,2\}} &= \frac{1}{4}, & W_{\{5,1,0,0\}} &= -\frac{3}{4}, & W_{\{5,1,0,2\}} &= \frac{3}{4}, \\ W_{\{6,1,0,0\}} &= -\frac{1}{16}. \end{aligned} \quad (23)$$

At one loop and up to $\mathcal{O}(\alpha_s\beta^3)$ relative to LO, the Wilson coefficients are given by

$$W_{\{1,2,0,0\}} = m^{-2\epsilon} \cdot \frac{2}{3}(\epsilon - 1)n_h iI_H - \bar{\mu}^{-2\epsilon} \cdot 2\delta_1 Z_g + k^{-2\epsilon} \left(-\frac{(\epsilon - 1)(8\epsilon - 11)C_A}{2\epsilon - 3} - \frac{2(\epsilon - 1)n_l}{2\epsilon - 3} \right) iI_S^a, \quad (24)$$

$$W_{\{1,2,1,0\}} = k^{-2\epsilon} \left((\epsilon - 1)C_A - \frac{1}{2}(2\epsilon - 1)C_F \right) iI_S^b, \quad (25)$$

$$W_{\{1,2,2,0\}} = m^{-2\epsilon} \left(-\frac{(\epsilon - 1)(96\epsilon^3 - 100\epsilon^2 + 12\epsilon - 29)C_A}{24(2\epsilon - 1)(2\epsilon + 1)} + \frac{(\epsilon - 1)(96\epsilon^4 + 44\epsilon^3 - 96\epsilon^2 + 37\epsilon + 6)C_F}{6(2\epsilon - 1)(2\epsilon + 1)(2\epsilon + 3)} - \frac{2}{15}\epsilon(\epsilon - 1)n_h \right) iI_H + k^{-2\epsilon} \left(\frac{1}{24}(-48\epsilon^2 + 104\epsilon - 61)C_A + \frac{1}{3}(\epsilon - 1)(8\epsilon - 7)C_F \right) iI_S^a, \quad (26)$$

$$W_{\{1,2,0,2\}} = m^{-2\epsilon} \left(\frac{2(\epsilon - 1)(2\epsilon^2 - 1)C_A}{2\epsilon - 1} - \frac{4(\epsilon - 1)\epsilon(2\epsilon + 1)C_F}{2\epsilon - 1} + \frac{1}{3}(1 - \epsilon)n_h \right) iI_H + \bar{\mu}^{-2\epsilon} \delta_1 Z_g + k^{-2\epsilon} \left(\frac{(\epsilon - 1)n_l}{2\epsilon - 3} - \frac{(40\epsilon^2 - 95\epsilon + 51)C_A}{6(2\epsilon - 3)} \right) iI_S^a, \quad (27)$$

$$W_{\{1,2,1,2\}} = k^{-2\epsilon} \left(\frac{1}{2}(\epsilon - 2)C_A + \frac{1}{4}(-\epsilon - 1)C_F \right) iI_S^b, \quad (28)$$

$$W_{\{1,2,3,0\}} = k^{-2\epsilon} \left(\frac{1}{8}(2\epsilon^2 - 6\epsilon + 5)C_A + \frac{1}{16}(-6\epsilon^2 + 15\epsilon - 14)C_F \right) iI_S^b, \quad (29)$$

$$W_{\{2,2,2,0\}} = m^{-2\epsilon} \left(\frac{1}{8}(1 - \epsilon)C_A + \frac{(\epsilon - 1)\epsilon C_F}{2(2\epsilon + 1)} \right) iI_H - k^{-2\epsilon} \cdot \frac{1}{8}C_A iI_S^a, \quad (30)$$

$$W_{\{2,2,3,0\}} = -k^{-2\epsilon} \cdot \frac{\epsilon C_F}{16(\epsilon - 1)} iI_S^b, \quad (31)$$

$$W_{\{4,2,0,0\}} = m^{-2\epsilon} \left(-\frac{(\epsilon - 1)(2\epsilon^2 - 1)C_A}{2(2\epsilon - 1)} + \frac{(\epsilon - 1)\epsilon(2\epsilon + 1)C_F}{2\epsilon - 1} + \frac{1}{6}(\epsilon - 1)n_h \right) iI_H - \bar{\mu}^{-2\epsilon} \frac{\delta_1 Z_g}{2} + k^{-2\epsilon} \left(-\frac{(\epsilon - 1)(4\epsilon - 5)C_A}{4(2\epsilon - 3)} - \frac{(\epsilon - 1)n_l}{2(2\epsilon - 3)} \right) iI_S^a, \quad (32)$$

$$W_{\{4,2,1,0\}} = k^{-2\epsilon} \left(\frac{\epsilon C_A}{8} - \frac{(2\epsilon^2 - 7\epsilon + 4)C_F}{16(\epsilon - 1)} \right) iI_S^b, \quad (33)$$

$$W_{\{5,2,0,0\}} = m^{-2\epsilon} \left(-\frac{(\epsilon - 1)(2\epsilon^2 - 1)C_A}{2\epsilon - 1} + \frac{2(\epsilon - 1)\epsilon(2\epsilon + 1)C_F}{2\epsilon - 1} + \frac{1}{2}(\epsilon - 1)n_h \right) iI_H - \bar{\mu}^{-2\epsilon} \frac{3\delta_1 Z_g}{2} + k^{-2\epsilon} \left(-\frac{(24\epsilon^2 - 49\epsilon + 21)C_A}{4(2\epsilon - 3)} - \frac{3(\epsilon - 1)n_l}{2(2\epsilon - 3)} \right) iI_S^a, \quad (34)$$

$$W_{\{5,2,1,0\}} = k^{-2\epsilon} \left(\frac{1}{4}(3\epsilon - 2)C_A + \frac{1}{4}(4 - 3\epsilon)C_F \right) iI_S^b. \quad (35)$$

The master integrals of the hard and soft regions in the expansion-by-regions technique are given by

$$iI_H = (4\pi)^{\epsilon-2} \Gamma(\epsilon - 1), \quad iI_S^a = -\frac{2^{4\epsilon-5} \pi^{\epsilon-\frac{1}{2}}}{\sin(\pi\epsilon) \Gamma\left(\frac{3}{2} - \epsilon\right)}, \quad iI_S^b = \frac{16^{\epsilon-1} \pi^\epsilon}{\cos(\pi\epsilon) \Gamma(1 - \epsilon)}, \quad (36)$$

(after factoring out the dimensionful parameters). The one-loop counter term for the gauge coupling constant reads

$$\delta_1 Z_g = \frac{2(n_h + n_l) - 11C_A}{96\pi^2\epsilon}. \quad (37)$$

The color factors are given by $C_F = 4/3$ and $C_A = 3$ for the SU(3) gauge group. It is straightforward to expand the above Wilson coefficients in ϵ .

Two-loop Hamiltonian for general gauge group

We expand the coefficients[3] of $\Lambda_1, \dots, \Lambda_5$ in ϵ while we ignore any $\mathcal{O}(\epsilon)$ contributions in Λ_i . (Namely, we simply take the limit $\epsilon \rightarrow 0$ for Λ_i .) The Hamiltonian is given in the form

$$H = \frac{16\pi^2 C_F}{k^2} \sum_{i=1}^4 \sum_{j=1}^3 \sum_{n,\ell \geq 0} \left(\frac{\alpha_s(k)}{4\pi} \right)^j C_{\{i,j,n,2\ell\}} \left(\frac{k}{m} \right)^n \frac{(p^2)^\ell + (p'^2)^\ell}{2m^{2\ell}} O_i. \quad (38)$$

$\alpha_s(k)$ denotes the strong coupling constant in the $\overline{\text{MS}}$ scheme of the theory with n_l flavors only, renormalized at $\mu = k$. [4] The spinor basis in three dimensions is defined as

$$O_1 = \mathbb{I} \otimes \mathbb{I}, \quad O_2 = \vec{S}^2, \quad O_3 = \frac{i}{k^2} \vec{S} \cdot (\vec{p} \times \vec{k}), \quad O_4 = \sigma^a \otimes \sigma^a - \frac{3}{k^2} (\vec{k} \cdot \vec{\sigma}) \otimes (\vec{k} \cdot \vec{\sigma}), \quad (39)$$

with

$$\vec{S} = \frac{\vec{\sigma}}{2} \otimes \mathbb{I} + \mathbb{I} \otimes \frac{\vec{\sigma}}{2}. \quad (40)$$

The Wilson coefficients are separated into finite and divergent parts as

$$C_{\{i,j,n,2\ell\}} = C_{\{i,j,n,2\ell\}}^{\text{fin}} + C_{\{i,j,n,2\ell\}}^{\text{div}}. \quad (41)$$

2-loop non-zero finite Wilson coefficients are given by

$$\begin{aligned} C_{\{1,3,0,0\}}^{\text{fin}} &= V_{\text{QCD}}^{(2 \text{ loop})}(k) [C_F \alpha_s(k)^3 / (4\pi k^2)]^{-1} \\ &= -\frac{100n_l^2}{81} + n_l \left(\left(\frac{28\zeta(3)}{3} + \frac{899}{81} \right) C_A + \left(\frac{55}{6} - 8\zeta(3) \right) C_F \right) \\ &\quad + \left(-\frac{22\zeta(3)}{3} - \frac{4343}{162} - 4\pi^2 + \frac{\pi^4}{4} \right) C_A^2, \end{aligned} \quad (42)$$

$$\begin{aligned} C_{\{1,3,1,0\}}^{\text{fin}} &= V_{1/(mk)}^{(2 \text{ loop})}(k) [C_F \alpha_s(k)^3 / (4\pi mk)]^{-1} \\ &= \left(-\frac{32}{3} \pi^2 C_A C_F - \frac{16}{3} \pi^2 C_A^2 \right) \log \left(\frac{\mu^2}{k^2} \right) + n_l \left(\frac{49\pi^2 C_A}{18} - \frac{4\pi^2 C_F}{9} \right) \\ &\quad + \left(\frac{130\pi^2}{9} - \frac{32}{3} \pi^2 \log(2) \right) C_A C_F + \left(-\frac{101\pi^2}{9} - \frac{16}{3} \pi^2 \log(2) \right) C_A^2, \end{aligned} \quad (43)$$

$$\begin{aligned}
C_{\{1,3,2,0\}}^{\text{fin}} &= \log^2\left(\frac{k^2}{m^2}\right) \left(-\frac{11C_A C_F}{9} + \frac{13C_{Anl}}{9} - \frac{193C_A^2}{18} - \frac{2C_F n_l}{9}\right) \\
&+ \log\left(\frac{k^2}{m^2}\right) \left(-2\pi^2 C_F^2 + \left(\frac{146}{9} + \frac{13\pi^2}{9}\right) C_A C_F - \frac{17C_{Anh}}{18} + \frac{637C_{Anl}}{54}\right. \\
&\quad \left.+ \left(-\frac{304}{9} - \frac{14\pi^2}{9}\right) C_A^2 - \frac{340C_F n_l}{27}\right) \\
&+ \log\left(\frac{m^2}{\mu^2}\right) \left(\left(\frac{416}{9} - \frac{13\pi^2}{9}\right) C_A C_F + \frac{40C_{Anl}}{9} - \frac{80C_F n_l}{9} - 4\pi^2 C_F^2\right. \\
&\quad \left.+ \left(-\frac{208}{9} - \frac{16\pi^2}{9}\right) C_A^2\right) \\
&+ \log(2) \left(22\pi^2 C_A C_F - 9\pi^2 C_A^2 - 6\pi^2 C_F^2\right) + \left(\frac{103\zeta(3)}{2} - \frac{407}{6} + \frac{13\pi^2}{24} + \frac{3\pi^4}{16}\right) C_A^2 \\
&+ \left(-77\zeta(3) + \frac{2369}{9} - \frac{713\pi^2}{54}\right) C_A C_F + \left(33\zeta(3) - \frac{946}{9} + \frac{6023\pi^2}{108}\right) C_F^2 + \frac{8n_h n_l}{27} \\
&+ \left(\frac{1387}{108} - \frac{41\pi^2}{36}\right) C_{Anh} + \left(\frac{173\pi^2}{54} - \frac{346}{9}\right) C_{Fnh} + \left(\frac{28\pi^2}{27} - \frac{173}{27}\right) C_{Anl} \\
&+ \left(\frac{424}{27} - \frac{8\pi^2}{27}\right) C_F n_l, \tag{44}
\end{aligned}$$

$$\begin{aligned}
C_{\{1,3,0,2\}}^{\text{fin}} &= \log\left(\frac{k^2}{\mu^2}\right) \left(\left(\frac{416}{9} + \frac{32\pi^2}{9}\right) C_A^2 - \frac{80C_{Anl}}{9}\right) + \left(\frac{28\zeta(3)}{3} + \frac{1571}{81} - \frac{4\pi^2}{27}\right) C_{Anl} \\
&+ \left(\frac{55}{6} - 8\zeta(3)\right) C_F n_l - \frac{100n_l^2}{81} + \left(-\frac{38\zeta(3)}{3} - \frac{7919}{162} + \frac{266\pi^2}{27} - \frac{\pi^4}{4}\right) C_A^2, \tag{45}
\end{aligned}$$

$$\begin{aligned}
C_{\{2,3,2,0\}}^{\text{fin}} &= \log^2\left(\frac{k^2}{m^2}\right) \left(\frac{38C_A^2}{9} - \frac{5C_{Anl}}{9}\right) + \log\left(\frac{m^2}{\mu^2}\right) \left(\frac{8}{3}\pi^2 C_A C_F + \frac{4}{3}\pi^2 C_F^2\right) \\
&+ \log\left(\frac{k^2}{m^2}\right) \left(-\frac{46C_A C_F}{9} - \frac{82C_{Anl}}{27} + \left(\frac{256}{27} - \frac{4\pi^2}{9}\right) C_A^2 + \frac{16C_F n_l}{9}\right) \\
&+ \log(2) \left(12\pi^2 C_F^2 - \frac{238}{9}\pi^2 C_A C_F + \frac{71}{9}\pi^2 C_A^2\right) + \left(\frac{16\pi^2}{27} - \frac{538}{81}\right) C_{Anh} \\
&+ \left(\frac{596}{27} - \frac{16\pi^2}{9}\right) C_{Fnh} + \frac{100n_l^2}{243} + \left(-18\zeta(3) - \frac{10}{3} - \frac{239\pi^2}{9}\right) C_F^2 \\
&+ \left(-\frac{295\zeta(3)}{18} + \frac{11651}{486} - \frac{127\pi^2}{27} - \frac{\pi^4}{12}\right) C_A^2 + \left(-\frac{13\zeta(3)}{3} - \frac{70}{27} + \frac{170\pi^2}{9}\right) C_A C_F \\
&+ \left(-\frac{28\zeta(3)}{9} - \frac{503}{243} - \frac{10\pi^2}{27}\right) C_{Anl} + \left(\frac{8\zeta(3)}{3} - \frac{637}{54}\right) C_F n_l, \tag{46}
\end{aligned}$$

$$\begin{aligned}
C_{\{3,3,2,0\}}^{\text{fin}} &= \log\left(\frac{k^2}{m^2}\right) \left(\frac{56C_A C_F}{3} - \frac{70C_{Anl}}{9} + \frac{262C_A^2}{9} - \frac{8C_F n_l}{3}\right) \\
&+ \log^2\left(\frac{k^2}{m^2}\right) \left(\frac{14C_A^2}{3} - \frac{2C_{Anl}}{3}\right) + \log(2) \left(\frac{8}{3}\pi^2 C_A C_F + \frac{8}{3}\pi^2 C_A^2 - 16\pi^2 C_F^2\right) \\
&+ \left(3\zeta(3) + \frac{5999}{108} - \frac{4\pi^2}{9} - \frac{3\pi^4}{8}\right) C_A^2 + \left(\frac{10\pi^2}{9} - \frac{298}{27}\right) C_{Anh} + \left(\frac{476}{9} - \frac{16\pi^2}{3}\right) C_{Fnh} \\
&+ \frac{50n_l^2}{27} + \left(-14\zeta(3) - \frac{827}{54} - \frac{4\pi^2}{9}\right) C_{Anl} + \left(-4\zeta(3) + \frac{590}{9} + \frac{8\pi^2}{3}\right) C_A C_F \\
&+ \left(12\zeta(3) - \frac{1055}{36}\right) C_F n_l + \left(24\zeta(3) - 62 + \frac{40\pi^2}{3}\right) C_F^2, \tag{47}
\end{aligned}$$

$$\begin{aligned}
C_{\{4,3,2,0\}}^{\text{fin}} &= \log\left(\frac{k^2}{m^2}\right) \left(\frac{17C_A C_F}{9} - \frac{35C_A n_l}{54} + \frac{149C_A^2}{54} - \frac{2C_F n_l}{9}\right) \\
&+ \log^2\left(\frac{k^2}{m^2}\right) \left(\frac{17C_A^2}{36} - \frac{C_A n_l}{18}\right) + \log(2) \left(\frac{2}{9}\pi^2 C_A C_F + \frac{2}{9}\pi^2 C_A^2 - \frac{4}{3}\pi^2 C_F^2\right) \\
&+ \left(-\frac{\zeta(3)}{18} + \frac{12299}{1944} + \frac{\pi^2}{18} - \frac{\pi^4}{48}\right) C_A^2 + \left(\frac{5\pi^2}{54} - \frac{149}{162}\right) C_A n_h + \left(\frac{119}{27} - \frac{4\pi^2}{9}\right) C_F n_h \\
&+ \frac{25n_l^2}{243} + \left(-\frac{7\zeta(3)}{9} - \frac{1367}{972} - \frac{\pi^2}{27}\right) C_A n_l + \left(-\frac{\zeta(3)}{3} + \frac{331}{54} + \frac{2\pi^2}{9}\right) C_A C_F \\
&+ \left(\frac{2\zeta(3)}{3} - \frac{445}{216}\right) C_F n_l + \left(2\zeta(3) - \frac{29}{6} + \frac{10\pi^2}{9}\right) C_F^2. \tag{48}
\end{aligned}$$

2-loop non-zero divergent Wilson coefficients are given by

$$C_{\{1,3,1,0\}}^{\text{div}} = -\frac{8\pi^2 C_A^2}{3\epsilon} - \frac{16\pi^2 C_A C_F}{3\epsilon}, \tag{49}$$

$$\begin{aligned}
C_{\{1,3,2,0\}}^{\text{div}} &= \log\left(\frac{k^2}{\mu^2}\right) \left(-\frac{88C_A C_F}{9\epsilon} - \frac{8C_A n_l}{9\epsilon} + \frac{44C_A^2}{9\epsilon} + \frac{16C_F n_l}{9\epsilon}\right) \\
&+ \left(\frac{44}{9\epsilon^2} + \frac{13\pi^2}{18\epsilon} - \frac{208}{9\epsilon}\right) C_A C_F + \left(-\frac{22}{9\epsilon^2} + \frac{8\pi^2}{9\epsilon} + \frac{104}{9\epsilon}\right) C_A^2 + \frac{2\pi^2 C_F^2}{\epsilon} \\
&+ \left(\frac{4}{9\epsilon^2} - \frac{20}{9\epsilon}\right) C_A n_l + \left(\frac{40}{9\epsilon} - \frac{8}{9\epsilon^2}\right) C_F n_l, \tag{50}
\end{aligned}$$

$$\begin{aligned}
C_{\{1,3,0,2\}}^{\text{div}} &= \log\left(\frac{k^2}{\mu^2}\right) \left(\frac{16C_A n_l}{9\epsilon} - \frac{88C_A^2}{9\epsilon}\right) + \left(\frac{44}{9\epsilon^2} - \frac{16\pi^2}{9\epsilon} - \frac{208}{9\epsilon}\right) C_A^2 \\
&+ \left(\frac{40}{9\epsilon} - \frac{8}{9\epsilon^2}\right) C_A n_l, \tag{51}
\end{aligned}$$

$$C_{\{2,3,2,0\}}^{\text{div}} = -\frac{4\pi^2 C_A C_F}{3\epsilon} - \frac{2\pi^2 C_F^2}{3\epsilon}. \tag{52}$$

-
- [1] A. G. Grozin, P. Marquard, J. H. Piclum and M. Steinhauser, Nucl. Phys. B **789** (2008), 277-293 doi:10.1016/j.nuclphysb.2007.08.012 [arXiv:0707.1388 [hep-ph]].
- [2] A. G. Grozin, M. Hoeschele, J. Hoff, M. Steinhauser, M. Hoeschele, J. Hoff and M. Steinhauser, JHEP **09** (2011), 066 doi:10.1007/JHEP09(2011)066 [arXiv:1107.5970 [hep-ph]].
- [3] Λ_6 is $\mathcal{O}(\beta^4)$ by itself, and it contributes only to the tree-level Hamiltonian within the accuracy orders of our current interest.
- [4] It is customary to express the Hamiltonian in terms of the coupling constant of the theory with n_l flavors only $\alpha_s(\mu) \equiv \alpha_s^{(n_l)}(\mu)$. We include the $\mathcal{O}(\epsilon)$ term in the decoupling relation [1, 2] to rewrite $\alpha_s^{(n_h+n_l)}(\mu)$ by $\alpha_s(\mu)$.
-